

NUCLEIC ACIDS AND METHODS FOR DETECTING PATHOGENS AND BENEFICIAL MICROORGANISMS

Field of the invention

The present invention relates to nucleic acids and methods for detecting pathogens and beneficial microorganisms.

Background of the invention

Disease in lawn grasses or turfgrasses develops from an interaction among a susceptible plant, an environment favorable for disease development, and a pathogenic organism (fungi and bacteria). Such organisms may also develop on decorative grasses, plants and crops; indeed, they may appear on any suitable organic substrate. Thus, treatment of a diseased substrate, especially turfgrass, usually consists in applying a treatment agent that will either kill the pathogen or keep it from growing.

However, the first step in disease management, and especially turfgrass disease management, should always consist in identifying the causative pathogenic agent. The knowledge of the causative pathogenic agent of a disease is important to select the most appropriate treatment agent (fungicides, bactericides etc.). Indeed, it is important to have identified the disease correctly, so that an appropriate fungicide or bactericide can be selected. Inadequate treatment will not cure the disease and may have a severe effect on the soil and other beneficial organisms. Using the wrong treatment is cost ineffective and may involve the risk of exacerbating the disease, as well as causing other unwanted side effects.

Classical methods for the identification of the causative pathogenic agent essentially rely on the symptoms which can be observed on the individual plant and on the turf stand, as well as on the pathogen structures, which can be found in the vicinity of the diseased turfgrass.

However, these methods may require a long time to be implemented, since they often involve the isolation and the culture of the pathogen in a laboratory. Besides, differentiating closely related pathogen species can be difficult.

Accordingly, molecular biology methods have been developed which circumvent these difficulties. One of the most popular fungal detection methods relies on the Polymerase chain reaction (PCR) amplification of the internal transcribed spacers (1, 2) and the 5.8S rRNA gene (ITS1 -5.8S-ITS2) from the fungal rRNA operon (Goodwin et al. (1995) Plant Pathology 44:384-391; Ranjard et al. (2001) Applied and Environmental Microbiology 67:4479-4487).

For bacterial detection, PCR amplification and sequencing of the 16S rRNA gene is used (Nam HR, Lee HM, Lee Y. Isolation of quinupristin/dalfopristin-resistant *Streptococcus agalactiae* from asymptomatic Korean women. J Microbiol. 2008 Feb; 46(1):108-11. doi: 10.1007/s12275-007-0217-1. PubMed PMID: 18337702).

However, sequencing of the region of a specific primer pair is often necessary for the identification of a given pathogenic species, which renders this method cumbersome where the identity of pathogen is unknown and is sought for. Furthermore, sequencing of mixtures of pathogens is expensive and laborious. Species specific primers to identify certain species out of mixtures are not available for all microorganisms. Accordingly, these methods are not used in routine for determining the anti-pathogenic agent most adapted to treat a given turfgrass disease.

Japanese patent application No. 2008005760 discloses 458 probes for detecting molds that can be found in food. These 458 probes are designed for detecting molds by a hybridization-based method involving the use of a microarray.

Patent application WO 2009/147017 A1, of the same applicant, discloses nucleic acids and methods for detecting pathogenic fungi in turfgrass. This invention relates to turfgrass but focuses on AFLP and T-RFLP as methods for identification. This method has the limit that often many different fungal species, pathogenic and non-pathogenic, are present in turfgrass. Many different species in a complex sample make it difficult or even impossible identifying numerical underrepresented species. The disease causing pathogenic species, however, is not necessarily the most abundant species.

As a consequence, current methods of choice for the detection of plant pathogens and beneficial microorganisms are time-consuming, not sensitive enough, not specific enough, and/or not broad enough to detect the spectrum of microorganisms.

It is therefore an objective of the present invention to provide specific nucleic acids that enable identification of the pathogens and beneficial microorganisms present in a sample.

It is also an objective of the present invention to provide a single Microarray-Chip containing up to 100 spots with these specific nucleic acids. These nucleic acids are specific for e.g. 90 known fungi and e.g. 10 bacteria species.

A further object of the present invention is a method that enables to test in a single sample for up to 100 different species. Since DNA sequences of very closely related species are identical, this method is also able to include detection of those microorganisms. This method thus makes it possible to screen very fast for many microorganisms. Furthermore, the method is adaptable for high-throughput. The method

according to the invention is thus very sensitive and enables detection of pathogens that are present at low levels.

Summary of the invention

The present invention arises from the identification, by the inventors, of a selection of nucleotide sequences specific to a conserved region within the gene of rRNA and its intergenic regions of fungi and bacteria generally relevant for turfgrasses' health, but also for other organic substrates.

Thus, the present invention relates to the use of a set of nucleic acids comprising or consisting of:

(i) the sequences selected from the group consisting of:

- GGGTTGCTTGCTTGCGAGCTCC (SEQ ID N°1);
- CTGGTGTGTTGGACTCGCCTTAAAC (SEQ ID N°2);
- GCACATATTTTGGCCTTTGTATCAGG (SEQ ID N°3);
- TCCGCCAGGGAARACCAAACTCT (SEQ ID N°4);
- CGGAGCGCGGGCCGTCGCG (SEQ ID N°5);
- ACTTATACCCAAAACGTTGCCTCG (SEQ ID N°6);
- CATTATCGAGTTTACGCTCCATAAC (SEQ ID N°7);
- GCGCAGCTATTAGATCTACGGTG (SEQ ID N°8);
- GCAAGGCTGGAGTATTTTATTACCCT (SEQ ID N°9);
- CCGTGGCCTTGTTGCCACGCCC (SEQ ID N°10);
- ATTGGGGCCTTGTTGCCACACCC (SEQ ID N°11);
- TTTTGGCCTTTGTCCAGTTGCGG (SEQ ID N°12);
- GATTGTCGCCCCCCTCCTGG (SEQ ID N°13);
- GACCTTATTCAAACCTTTTTTTCAGTT (SEQ ID N°14);
- GCTGTTGGGGACCGGCTCACCCG (SEQ ID N°15);
- TGACCGTTGTACGAGACGACTTTAT (SEQ ID N°16);
- CTTGGTGTGGGAGCTGCAGTCC (SEQ ID N°17);
- CCATTGCGTAGTAGTAAAACCC (SEQ ID N°18);
- AACGCGCTTCGTTCCGAGGCTT (SEQ ID N°19);
- TTCAACCCTCAAGCCCCCGGGTTTG (SEQ ID N°20);
- CGAAGTAGTGATATTCCGCATCGG (SEQ ID N°21);
- GTTAGGGGGTCCCCTCTCCGG (SEQ ID N°22);
- CTCGGTCTCGAGCCGCCGG (SEQ ID N°23);
- ACACCCCATTTGAACCTATTTATTTTYAA (SEQ ID N°24);
- ACATCTCGCGCTTTGCATTCAGAA (SEQ ID N°25);

- CGATTTTGGGGGGTGGCTAGTGC (SEQ ID N°26);
- CTCTGAGTACGAAAAGAACCTGAAA (SEQ ID N°27);
- TGCTGGCTCTTCTAGAGTCGGCTC (SEQ ID N°28);
- GGCCAGACGACAGCCATAAACC (SEQ ID N°29);
- 5 - CGCCGGTGGACTACCTAAACTCT (SEQ ID N°30);
- CGTCGTCGCTGTTGCAAGGAC (SEQ ID N°31);
- TCGGGCAACGGAACCAGGCGC (SEQ ID N°32);
- GTTGCCTCGGCGGGCACGGC (SEQ ID N°33);
- CCAGTTATATAGGCACCCAATAAGCC (SEQ ID N°34);
- 10 - CACCAAACCAGCTTGGGAAACCTT (SEQ ID N°35);
- GTCACGTGGTCTTGGTTTTGAA (SEQ ID N°36);
- CGCAACCGGGAGCCGCGGCGCGG (SEQ ID N°37);
- TGGACTGGCTTCGGCTAGACTGG (SEQ ID N°38);
- ACCATAGCTCAGTTGCTTGGCTTTT (SEQ ID N°39);
- 15 - TGGCGAATGTTTGGACTTCGGTCT (SEQ ID N°40);
- GCGCAAGCTGGGGTGGRCGAG (SEQ ID N°41);
- GAACTTGTGTCTCTGCGGCGCG (SEQ ID N°42);
- AACTGGTGAACCGTAGCTGTGTGGT (SEQ ID N°43);
- CCGTACATTAACTTGACTTTCTTCC (SEQ ID N°44);
- 20 - GGAACCTCCACCCTTGAATACACTG (SEQ ID N°45);
- CGACCCCTTTTATAATTACCCAAC (SEQ ID N°46);
- CTAAAAAACCCCTCATAACCTTTTTTTT (SEQ ID N°47);
- CTCCTAAAACCCAATATCTTATTTTAAAG (SEQ ID N°48);
- CAATACTGCCATCTTGTTTTTGAAGG (SEQ ID N°49);
- 25 - ATACTTGCCATCTTTTGAAGG (SEQ ID N°50);
- CCTAAAAACCCCTTATCA (SEQ ID N°51);
- CAACCTTTTTTGGAGTATTCTAATGAT (SEQ ID N°52);
- ATACTGCCATCTTATTKAAGGGAGAC (SEQ ID N°53);
- CAGGGCTATCCCCCTGCCAGG (SEQ ID N°54);
- 30 - GGAGGGTTGCGCACTTTGTGCGTG (SEQ ID N°55);
- GACGCCCTGTTTTCGGATAGGG (SEQ ID N°56);
- TGGCGTGCGTTTGCTTGCGCTTC (SEQ ID N°57);
- CTATACTCCGAGAACGAAAGTTTTTGG (SEQ ID N°58);
- TAGTAGTGTGTGTRGCACGTTGTC (SEQ ID N°59);
- 35 - TGTTTTTGTTTTGTGGAAATACGCTGTTT (SEQ ID N°60);
- GTAGAATTTTGCTGCTCTTGGGCG (SEQ ID N°61);

- CTGTGTAGTCAGGGATGGAATGTGC (SEQ ID N°62);
- TGAACGCATCATGTTGCTTCGGG (SEQ ID N°63);
- CTGGCTTTTGTGTTTGGATTTGGAGGT (SEQ ID N°64);
- CAGCGACAACCGACTCTAAGTTCA (SEQ ID N°65);
- 5 - CTCGTGAAACACATGAAGTCTGAG (SEQ ID N°66);
- TCGGCGCCCCAGGAGAAATCCT (SEQ ID N°67);
- AGTCCATGTCCGCAATGGCAGG (SEQ ID N°68);
- AACACATACCTCTCGTTACAGGGTC (SEQ ID N°69);
- GAGCTGCTCTTCGGGGCCTTGTAT (SEQ ID N°70);
- 10 - CTTGTATGCGCGCCAGAGAATATCA (SEQ ID N°71);
- TATTGGGCGTCCGCGGGGGA (SEQ ID N°72);
- GCGGAGTTCACGAGCCCTCAC (SEQ ID N°73);
- GCGCCTTGTCTCTCGCGAGAC (SEQ ID N°74);
- AGCTGGATCTCAGTGTTATGCTTGG (SEQ ID N°75);
- 15 - GTGCCTCTCGGGGCTTCTGCCG (SEQ ID N°76);
- CCTGTGTAGTAATGCTTAGCTTACAC (SEQ ID N°77);
- CTACGGAGGGGTGGCTGCGTTG (SEQ ID N°78);
- CTACGGAGGGGTGGCTGCGTTG (SEQ ID N°79);
- ACAGCTCTGAGCAAAAATTCAAATG (SEQ ID N°80);
- 20 - GTCCAATGTAGGCGCAGCGTAA (SEQ ID N°81);
- ATTAGCTGGAACCTCTTGTGGACC (SEQ ID N°82);
- ATTAGCTGGAACCTCTTGTGGACC (SEQ ID N°83);
- CTCATTAAACAATTTTTCTTATAAAGATTG (SEQ ID N°84);
- GTCATTGCACTCAGGTAGACGTAACA (SEQ ID N°85);
- 25 - CTTTTTCTTTTGAAAAGGTTGACG (SEQ ID N°86);
- ACAGACAATTTTATTGAACACTTTTT (SEQ ID N°87);
- TTTGAAGAGTTGGCGGATCGGTAT (SEQ ID N°88);
- ACGGACAATTTTATTTAACACTTTTTG (SEQ ID N°89);
- GCCCTTAAAKCAGTGGCGGACC (SEQ ID N°90);
- 30 - TAATACATGCAAGTCGAGCGGACAGAT (SEQ ID N°91);
- GGGTACTTACCTAATACGTGAGTAT (SEQ ID N°92);
- GAAGGGTARTGTCTTAATACGGCATT (SEQ ID N°93);
- GGAAGGGYAGTGTGTTAATAGCAC (SEQ ID N°94);
- ATGACTGTCCATCGCATGGTGGAT (SEQ ID N°95);
- 35 - CTGCGGATCGCATGGTCTGCG (SEQ ID N°96);
- AATACTTCTCCTCGCATGGGGAGG (SEQ ID N°97);

- ATGACACGACCGCATGGTCTGT (SEQ ID N°98);
 - CACCGGAAACGGCAGAGATGTCG (SEQ ID N°99); and
 - GAGTGTGTAAGGATGGCGAA (SEQ ID N°100);

(iii) the sequences having at least 80% identity with SEQ ID N°1, SEQ ID N°2, SEQ

5 ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9,

SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15,
 SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21,
 SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27,
 SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33,
 SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39,
 SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45,
 SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51,
 SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57,
 SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63,
 SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69,

15 SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75,
 SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81,
 SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87,
 SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93,
 SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99
 and SEQ ID N°100, or

(iii) the complementary sequences of (i) or (ii),

for the detection by hybridization techniques of at least one pathogen and/or one
 beneficial microorganism in a sample.

25 It also relates to a method for detecting by hybridization techniques at least one
 pathogen and/or one beneficial microorganism in a sample, said method comprising the
 use of a set of nucleic acids comprising or consisting of:

(i) the sequences selected from the group consisting of SEQ ID N°1, SEQ ID N°2,

30 SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID
 N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID

N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID
 N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID
 N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID
 N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID
 N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID
 N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID

N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100,

(ii) the sequences having at least 80% identity with SEQ ID N°1, SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100, or

(iii) the complementary sequences of (i) or (ii).

In one embodiment, the invention also relates to the use of at least one nucleic acid comprising or consisting of:

(i) a sequence selected from the group consisting of SEQ ID N°1, SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33,

5	SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 or SEQ ID N°100,	or
10	(ii) a sequence having at least 80% identity with SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 or SEQ ID N°100,	
15	(iii) a complementary sequence of (i) or (ii),	
20	for the detection by hybridization techniques of at least one pathogen and/or one beneficial microorganism in a sample.	
25	In a particular embodiment, the present invention also relates to a method for detecting by hybridization techniques at least one pathogen and/or one beneficial	
30		
35		

microorganism in a sample, said method comprising the use of at least one nucleic acid

comprising or consisting of:

(i) a sequence selected from the group consisting of SEQ ID N°1, SEQ ID N°2, SEQ

ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9,

SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15,

SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21,

SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27,

SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33,

SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39,

SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45,

SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51,

SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57,

SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63,

SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69,

SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75,

SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81,

SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87,

SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93,

SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 or

SEQ ID N°100,

(ii) a sequence having at least 80% identity with SEQ ID N°2, SEQ ID N°3, SEQ ID

N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10,

SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16,

SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22,

SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28,

SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34,

SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40,

SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46,

SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52,

SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58,

SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64,

SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70,

SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76,

SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82,

SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88,

SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94,

35

30

25

20

15

10

5

SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 or SEQ ID N°100,
or

(iii) the complementary sequences of (i) or (ii).

The present invention also relates to a Microarray-Chip comprising a set of nucleic acids
comprising or consisting of:

(i) the sequences SEQ N°1 SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5,
SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID
N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID
N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID
N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID
N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID
N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID
N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID
N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID
N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID
N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID
N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID
N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID
N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID
N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID
N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID
N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100,

(ii) the sequences having at least 80% identity with SEQ ID N°1, SEQ ID N°2, SEQ
ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9,
SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15,
SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21,
SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27,
SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33,
SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39,
SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45,
SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51,
SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57,
SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63,
SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69,
SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75,
SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81,

SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100, or

(iii) the complementary sequences of (i) or (ii).

In a particular embodiment, the present invention also relates to a Microarray-Chip

comprising at least one nucleic acid comprising or consisting of:

(i) a sequence selected from the group consisting of: SEQ ID N°1, SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9,

5

SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100,

25

(ii) a sequence having at least 80% identity with SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10,

SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64,

30

35

SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70,
 SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76,
 SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82,
 SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88,
 5 SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94,
 SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 or SEQ ID N°100,
 or

(iii) a complementary sequence of (i) or (ii).

The present invention also relates to a detection method of pathogens and/or
 10 beneficial microorganisms in an organic sample, comprising the steps of:

- a) DNA isolation from the sample and from standards comprising DNA of
 said pathogens and/or beneficial microorganisms;
- b) amplification of the isolated DNAs of step a) by PCR;
- c) hybridization of the PCR products obtained on a Microarray-Chip as
 15 defined above;
- d) detection of the bound DNAs; and
- e) deduction therefrom if the sample contains pathogens and/or beneficial
 microorganisms.

The present invention also relates to a kit for the detection of pathogens and/or
 20 beneficial microorganisms in an organic sample comprising:

- (i) the nucleic acids of the following sequences: SEQ N°1 SEQ ID N°2, SEQ ID
 N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ
 ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID
 N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19,
 25 SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ
 ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID
 N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35,
 SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ
 ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID
 30 N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51,
 SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ
 ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID
 N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67,
 SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ
 35 ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID
 N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83,

SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100,

- (ii) the nucleic acids of sequences having at least 80% identity with SEQ ID N°1, SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100, or

- (iii) the nucleic acids of complementary sequences of (i) or (ii).

The present invention also relates to a method of treating a diseased turfgrass comprising the steps of:

- a) detecting by a method combining PCR and Microarray-Chip the absence or the presence of nucleic acids from at least one pathogenic fungi and/or bacteria in a sample of soil in which the diseased turfgrass is growing, or in a sample of the diseased turfgrass, with at least one nucleic acid as defined above, in particular with the set of nucleic acids as defined above;
- b) if nucleic acids from one or more pathogenic fungi and/or bacteria have been detected in step a), selecting one or more antifungal and/or antibacterial agents which target the one or more pathogenic fungi and/or bacteria from which nucleic acids have been detected; and

- c) applying the selected one or more antifungal and/or antibacterial agents of step b) to the diseased turfgrass.

Detailed description of the invention

5 Nucleic acids, as intended herein can be of any type, however it is preferred that they are DNA.

The table 1 below provides details on the nucleic acid sequences that can be used for the identification of the assigned species. It gives the reference SEQ ID, the name of the oligonucleotide, the corresponding species and which type of the 2 PCRs is needed to amplify them (ITS: genomic region comprising of genes and intergenic regions encoding the ribosomal RNA in fungi; 16S rRNA: genomic region encoding the 16S ribosomal RNA in bacteria).

Table 1:

SEQ ID N°	Oligo ID	Species	Nucleic acid sequences	Region
1	ARME	Armillaria mellea	GGGTTGCTTGCTTGCGAGCTCC	ITS
2	ASPH	Ascochyta phleina	CTGGTGTTTGGACTCGCCTTAAAC	ITS
3	BISO	Bipolaris sorokiniana	GCACATATTTTGCCTTTGTATCAGG	ITS
4	BLGR	Blumeria graminis	TCCGCCAGGGAARACCAAACTCT	ITS
5	CEZE	Cercospora zebrina	CGGAGCGCGGGCCGTCGCG	ITS
6	CLPU	Claviceps purpurea	ACTTATACCCAAAACGTTGCCTCG	ITS
7	COTR	Colletotrichum trifolii	CATTATCGAGTTTACGCTCCATAAC	ITS
8	COFU	Corticium fuciforme	GCGCAGCTATTAGATCTACGGTG	ITS
9	CUAF	Curvularia affinis	GCAAGGCTGGAGTATTTTATTACCCT	ITS
10	DRDI	Drechslera dictyoides	CCGTGGCCTTGTTGCCACGCCC	ITS
11	DRPH	Drechslera phlei	ATTGGGGCCTTGTTGCCACACCC	ITS
12	DRPO	Drechslera poae	TTTTGCGCTTTGTCCAGTTGCGG	ITS
13	DRSI	Drechslera siccans	GATTGTCGCCCCCCTCCTGG	ITS
14	DRTR	Drechslera tritici-repentis	GACCTTATTCAAACCTTTTTTTCAGTT	ITS
15	EPTY	Epichloe typhina	GCTGTTGGGGACCGGCTCACCCG	ITS
16	EXTU	Exserohilum turcicum	TGACCGTTGTACGAGACGACTTTAT	ITS
17	FUCU	Fusarium culmorum	CTTGGTGTTGGGAGCTGCAGTCC	ITS
18	FUPO	Fusarium poae	CCATTGCGTAGTAGTAAAACCC	ITS
19	GAGR	Gaeumannomyces graminis	AACGCGCTTCGTTGCGAGGCTT	ITS

20	FUPR	Gibberella intermedia	TTCAACCCTCAAGCCCCCGGGTTTG	ITS
21	GLCA	Gliocladium catenulatum	CGAAGTAGTGATATTCCGCATCGG	ITS
22	GLGR	Glomerella graminicola	GTTAGGGGGTCCCCCTCTCCGG	ITS
23	KACA	Kabatiella caulivora	CTCGGTCTCGAGCCGCCGG	ITS
24	LEKO	Leptosphaeria korrae	ACACCCCATTTGAACCTATTTATTTTYAA	ITS
25	LEAU	Leptosphaerulina australis	ACATCTCGCGCTTTGCATTCAGAA	ITS
26	MAPH	Macrophomina phaseolina	CGATTTTGGGGGGTGGCTAGTGC	ITS
27	MAPO	Magnaporthe poae	CTCTGAGTACGAAAAGAACCTGAAA	ITS
28	MAOR	Marasmius oreades	TGCTGGCTCTTCTAGAGTCGGCTC	ITS
29	MIBO	Microdochium bolleyi	GGCCAGACGACAGCCATAAACC	ITS
30	MINI	Microdochium nivale	CGCCGGTGGACTACCTAAACTCT	ITS
31	MITR	Microsphaera trifolii	CGTCGTGCTGTTGCGAAGGAC	ITS
32	MYRO	Myrothecium roridum	TCGGGCAACGGAACCAGGCGC	ITS
33	NESP	Neotyphodium sp.	GTTGCCTCGGCGGGCACGGC	ITS
34	OPHE	Ophiosphaerella herpotricha	CCAGTTATATAGGCACCCAATAAGCC	ITS
35	LENO	Ophiosphaerella narmari	CACCAAACCAGCTTGGGAAACCTT	ITS
36	PETR	Peronospora trifoliorum	GTCACGTGGTCTTGGTTTTGAA	ITS
37	PHSP	Phyllachora vulgata	CGCAACCGGGAGCCGCGGCGCGG	ITS
38	PLBR	Phytophthora brassicae	TGGACTGGCTTCGGCTAGACTGG	ITS
39	PHCA	Phytophthora cactorum	ACCATAGCTCAGTTGCTTGGCTTTT	ITS
40	PHCI	Phytophthora citricola	TGGCGAATGTTTGGACTTCGGTCT	ITS
41	PHDR	Phytophthora drechsleri	GCGCAAGCTGGGGTGGRCGAG	ITS
42	PHFR	Phytophthora fragariae	GAAC TTGTGTCTCTGCGGCGCG	ITS
43	PHME	Phytophthora megasperma	AACTGGTGAACCGTAGCTGTGTGGT	ITS
44	PHNI	Phytophthora nicotianae	CCGTACATTAACTTGACTTTCTTCC	ITS
45	PSME	Pseudopeziza medicaginis	GGAACCTCCACCCTTGAATACACTG	ITS
46	PUCO1	Puccinia coronata	CGACCCCTTTTATAATTACCCAAC	ITS
47	PUCO2	Puccinia coronata	CTAAAAAACCCCTCATAACCTTTTTT	ITS

48	PUGRA	<i>Puccinia graminis</i>	CTCCTAAAACCCAATATCTTATTTTTTAAG	ITS
49	PUPO	<i>Puccinia poae-nemoralis</i>	CAATACTGCCATCTTGTTTTGAAGG	ITS
50	PUPA	<i>Puccinia poarum</i>	ATACTTGCCATCTTTTTGGAAGG	ITS
51	PURE	<i>Puccinia recondita</i>	CCTAAAAACCCCCCTTATCA	ITS
52	PUSO	<i>Puccinia sorghi</i>	CAACCTTTTTGGAGTATTCTAATGAT	ITS
53	PUST	<i>Puccinia striiformis</i>	ATACTGCCATCTTATTKAAGGGAGAC	ITS
54	PUTR	<i>Puccinia triticina</i>	CAGGGCTATCCCCCTGCCAGG	ITS
55	PYLY	<i>Pyrenochaeta lycopersici</i>	GGAGGGTTGCGCACTTTGTGCGTG	ITS
56	PYAP	<i>Pythium aphanidermatuma</i>	GACGCCCTGTTTTCGGATAGGG	ITS
57	PYDE	<i>Pythium debaryanum</i>	TGGCGTGCGTTTGCTTGCGCTTC	ITS
58	PHGR	<i>Pythium graminicola</i>	CTATACTCCGAGAACGAAAGTTTTTG	ITS
59	PYIR	<i>Pythium irregulare</i>	TAGTAGTGTGTGTRGCACGTTGTC	ITS
60	PYMA	<i>Pythium mastophorum</i>	TGTTTTTGTTTTGTGGAAATACGCTGTTT	ITS
61	PYSU	<i>Pythium sulcatum</i>	GTAGAATTTTGCTGCTCTTGGGCG	ITS
62	PYUL	<i>Pythium ultimum</i>	CTGTGTAGTCAGGGATGGAATGTGC	ITS
63	RACO	<i>Ramularia collycygni</i>	TGAACGCATCATGTTGCTTCGGG	ITS
64	RHCE	<i>Rhizoctonia cerealis</i>	CTGGCTTTTGTTTTGGATTTGGAGGT	ITS
65	RHFR	<i>Rhizoctonia fragariae</i>	CAGCGACAACCGACTCTAAGTTCA	ITS
66	RHOR	<i>Rhynchosporium orthosporum</i>	CTCGTGAAACACATGAAGTCTGAG	ITS
67	RHSE	<i>Rhynchosporium secalis</i>	TCGGCGCCCCAGGAGAAATCCT	ITS
68	SCBO	<i>Sclerotinia borealis</i>	AGTCCATGTCCGCAATGGCAGG	ITS
69	SCHO	<i>Sclerotinia homoeocarpa</i>	AACACATACCTCTCGTTACAGGGTC	ITS
70	SCSC	<i>Sclerotinia sclerotiorum</i>	GAGCTGCTCTTCGGGGCCTTGTAT	ITS
71	SCTR	<i>Sclerotinia trifoliorum</i>	CTTGTATGCGCGCCAGAGAATATCA	ITS
72	SEMA	<i>Septoria macropoda</i>	TATTGGGCGTCCGCGGGGGA	ITS
73	SETR	<i>Septoria tritici</i>	GCGGAGTTCACGAGCCCTCAC	ITS
74	STSA	<i>Stemphylium sarcinaeforme</i>	GCGCCTTGTCTCTCGCGAGAC	ITS
75	THCU	<i>Thanatephorus cucumeris</i>	AGCTGGATCTCAGTGTTATGCTTGG	ITS
76	CHEL	<i>Thielaviopsis basicola</i>	GTGCCTCTCGGGGCTTCTGCCG	ITS
77	CHTH	<i>Thielaviopsis populi</i>	CCTGTGTAGTAATGCTTAGCTTACAC	ITS
78	TICA	<i>Tilletia caries</i>	CTACGGAGGGGTGGCTGCGTTG	ITS
79	TICO	<i>Tilletia controversa</i>	CTACGGAGGGGTGGCTGCGTTG	ITS

80	TRHA	Trichoderma hamatum	ACAGCTCTGAGCAAAAATTCAAAATG	ITS
81	TYIN	Typhula incarnate	GTCCAATGTAGGCGCAGCGTAA	ITS
82	TYIS1	Typhula ishkariensis	ATTAGCTGGAACCTCTTGTGGACC	ITS
83	TYIS2	Typhula ishkariensis	ATTAGCTGGAACCTCTTGTGGACC	ITS
84	URDA	Uromyces dactylidis	CTCATTAACAATTTTTCTTATAAAGATTG	ITS
85	URTR	Uromyces trifolii-repentis	GTCATTGCACTCAGGTAGACGTAACA	ITS
86	USMA	Ustilago maydis	CTTTTTCTTTTGGAAAAGGTTGACG	ITS
87	USNU	Ustilago nuda	ACAGACAATTTTATTGAACACTTTTT	ITS
88	USST	Ustilago striiformis	TTTGAAGAGTTGGCGGATCGGTAT	ITS
89	USTR	Ustilago tritici	ACGGACAATTTTATTTAACACTTTTG	ITS
90	VEDA	Verticillium dahliae	GCCCTTAAAKCAGTGGCGGACC	ITS
91	BASU	Bacillus amyloliquefaciens	TAATACATGCAAGTCGAGCGGACAGAT	16S rRNA
92	PSCH	Pseudomonas chlororaphis	GGGTACTTACCTAATACGTGAGTAT	16S rRNA
93	SEEN	Serratia entomophila	GAAGGGTARTGTCTTAATACGGCATT	16S rRNA
94	SEPL	Serratia plymuthica	GGAAGGGYAGTGTGTTAATAGCAC	16S rRNA
95	STAL	Streptomyces albidoflavus	ATGACTGTCCATCGCATGGTGGAT	16S rRNA
96	STGR1	Streptomyces graminofaciens	CTGCGGATCGCATGGTCTGCG	16S rRNA
97	STGR2	Streptomyces griseoviridis	AATACTTCTCCTCGCATGGGGAGG	16S rRNA
98	STRI	Streptomyces rimosus	ATGACACACGACCGCATGGTCTGT	16S rRNA
99	STSC	Streptomyces scabiei	CACCGGAAACGGCCAGAGATGGTCG	16S rRNA
100	XATR	Xanthomonas translucens	GAGTGTGGTAGAGGATGGCGGAA	16S rRNA

Nucleotide sequences are written using the nucleotide base code (IUPAC) (Cornish-Bowden, A. Nucl Acid Res 13, 3021-3030 (1985)). In nucleotide sequences letters G, A, T, C stand for Guanine, Adenine, Thymine and Cytosine, M stands for A and C, R stands for A and G, W stands for A and T, Y stands for C and T, S stands for C and G, K stands for G and T, H stands for A and C and T, V stands for A and C and G, D stands for A and G and T, B stands for C and G and T, N stands for A and C and G and T.

As intended herein, the nucleic acids comprising or consisting of sequences having at least 80% identity with SEQ ID N°1 to 100, as previously mentioned, are nucleic acid sequences 80 to 100% identical to the nucleic acid sequences of SEQ ID N°1 to 100. In particular, these nucleic acid sequences are 90 to 100% identical, more particularly 95

to 100%, most preferably 95, 96, 97, 98, 99 or 100% identical to the nucleic acid sequences of SEQ ID N°1 to 100.

By nucleic acids having at least, for example, 95% identity with a query nucleic acid sequence of the present invention, it is intended that the nucleic acid sequence is identical to the query sequence except that the sequence may include up to five nucleic acid alterations per each 100 nucleic acids of the query nucleic acid sequence. In other words, to obtain a nucleic acid having a sequence at least 95% identical to a query nucleic acid sequence, up to 5% (5 of 100) of the nucleic acids in the subject sequence may be inserted, deleted, or substituted with another nucleic acid.

In the context of the present application, the percentage of identity is calculated using a global alignment (*i.e.* the two sequences are compared over their entire length). Methods for comparing the identity of two or more sequences are well known in the art. The « needle » program, which uses the Needleman-Wunsch global alignment algorithm (Needleman and Wunsch, 1970 J. Mol. Biol. 48:443-453) to find the optimum alignment (including gaps) of two sequences when considering their entire length, may for example be used.

In particular, the maximum length of the nucleic acids according to the invention is less than or equal to 500 nucleotides, preferably less than or equal to 300, more preferably less than or equal to 200, still more preferably less than or equal to 100, even more preferably less than or equal to 50, most preferably less than or equal to 35; in a particular embodiment the nucleotide sequence length is from 19 to 31 nucleotides.

In the context of the invention, by "at least one nucleic acid", it is meant, for example, 1, 2, 3, 10, 20, 30, 40, 50 or more (up to 100) nucleic acids comprising or consisting of sequences chosen from the sequences SEQ N°1 to 100, sequences at least 80% identical to SEQ ID N°1 to 100 and complementary sequences thereof. It can for example be at least 40 nucleic acids, at least 50, 60, 70, 80 or 90 nucleic acids comprising or consisting of sequences chosen from said sequences. It has to be noted that any of the sequences above mentioned in (i), (ii) and (iii) can be chosen and combined together.

In a particular embodiment of the uses, methods and Microarray-Chips according to the invention, at least 40 nucleic acids, and more particularly at least 50 nucleic acids are used.

"Stringent conditions" can be easily defined by the person skilled in the art using common knowledge. If necessary, guidance for defining such conditions can be found in numerous textbooks, such as Tijssen, 1993, Laboratory Techniques in Biochemistry and Molecular Biology - Hybridization with Nucleic Acid Probes, Part I, Chapter 2 "Overview of principles of hybridization and the strategy of nucleic acid probe assays", Elsevier, New

York. In particular, stringent conditions according to the invention can be constituted using a hybridization reaction requiring an optimized combination of hybridization buffer and hybridization temperature, depending on the ingredients of the hybridization buffer. Such determination of hybridization conditions falls within the routine work of the person skilled in the art.

By way of example, which has not to be considered as limiting the present invention, hybridization can be carried out at 55°C according to the ALERE DNA Array protocol and by using the hybridization buffers mentioned therein (Hybridisation Kit, ALERE Technologies GmbH, Germany, order numbers 245200100 and 24520K100).

As intended herein, "one pathogen" is a microorganism of any type that causes disease in its host. The host may be an animal, a plant, or even another microorganism. In particular, the pathogen is selected from bacteria and/or fungi. More particularly, fungi and/or bacteria are selected from the group consisting of : *Armillaria mellea*, *Ascochyta phleina*, *Bipolaris sorokiniana*, *Blumeria graminis*, *Cercospora zebrina*, *Claviceps purpurea*, *Colletotrichum trifolii*, *Corticium fuciforme*, *Curvularia affinis*, *Drechslera dictyoides*, *Drechslera phlei*, *Drechslera poae*, *Drechslera siccans*, *Drechslera tritici-repentis*, *Epichloe typhina*, *Exserohilum turcicum*, *Fusarium culmorum*, *Fusarium poae*, *Gaeumannomyces graminis*, *Gibberella intermedia*, *Gliocladium catenulatum*, *Glomerella graminicola*, *Kabatiella caulivora*, *Leptosphaeria korrae*, *Leptosphaerulina australis*, *Macrophomina phaseolina*, *Magnaporthe poae*, *Marasmius oreades*, *Microdochium bolleyi*, *Microdochium nivale*, *Microsphaera trifolii*, *Myrothecium roridum*, *Neotyphodium sp.*, *Ophiosphaerella herpotricha*, *Ophiosphaerella narmari*, *Peronospora trifoliorum*, *Phyllachora vulgata*, *Phytophthora brassicae*, *Phytophthora cactorum*, *Phytophthora citricola*, *Phytophthora drechleri*, *Phytophthora fragariae*, *Phytophthora megasperma*, *Phytophthora nicotianae*, *Pseudopeziza medicaginis*, *Puccinia coronata*, *Puccinia coronata*, *Puccinia graminis*, *Puccinia poae-nemoralis*, *Puccinia poarum*, *Puccinia recondite*, *Puccinia sorghi*, *Puccinia striiformis*, *Puccinia tritici*, *Pyrenochaeta lycopersici*, *Pythium aphanidermatum*, *Pythium debaryanum*, *Pythium graminicola*, *Pythium irregulare*, *Pythium mastophorum*, *Pythium sulcatum*, *Pythium ultimum*, *Ramularia collo-cygni*, *Rhizoctonia cerealis*, *Rhizoctonia fragariae*, *Rhynchosporium orthosporum*, *Rhynchosporium secalis*, *Sclerotinia borealis*, *Sclerotinia homoeocarpa*, *Sclerotinia sclerotiorum*, *Sclerotinia trifoliorum*, *Septoria macropoda*, *Septoria tritici*, *Stemphylium sarcinaeforme*, *Thanatephorus cucumeris*, *Thielaviopsis basicola*, *Thielaviopsis populi*, *Tilletia caries*, *Tilletia controversa*, *Trichoderma hamatum*, *Typhula incarnate*, *Typhula ishikariensis*, *Typhula ishikariensis*, *Uromyces dactylidis*, *Uromyces trifolii-repentis*, *Ustilago maydis*, *Ustilago nuda*, *Ustilago striiformis*, *Ustilago tritici*,

Verticillium dahliae; *Bacillus amyloliquefaciens*, *Pseudomonas chlororaphis*, *Serratia entomophila*, *Serratia plymuthica*, *Streptomyces albidoflavus*, *Streptomyces graminofaciens*, *Streptomyces griseoviridis*, *Streptomyces rimosus*, *Streptomyces scabiei*, *Xanthomonas translucens*.

It has to be understood that a fungus and/or bacterium being part of the above mentioned list but which could have synonymous names and/or which could have been renamed, are also included. In addition, closely related species of those mentioned in the above list are also in the frame of the present invention. By closely related species, it is meant species having an identical sequence to one or more nucleotide(s) of SEQ ID NOs 1 to 100, as described above, within their conserved regions.

In a particular embodiment, the present invention relates to the use of a set of nucleic acids comprising or consisting of:

(i) the sequences SEQ N°1, SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100,

(ii) the sequences comprising or consisting of sequences having at least 80% identity with SEQ N°1, SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID

N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100, or

(iv) the complementary sequences of (i) or (ii),

for the detection of all of the fungus and bacteria mentioned in the above list.

As intended herein, "one beneficial microorganism" is any organism that benefits the growing process, including insects, arachnids, other animals, plants, bacteria, fungi, viri, and nematodes. In particular, the beneficial microorganism is a plant growth promoting rhizobacteria.

As intended herein, the sample in which nucleic acids are to be detected can be of any type of organic substrate liable to contain nucleic acids. In particular, it can be a turfgrass, soil, natural fertilizer, seed or crop sample. More particularly, the natural fertilizer can be of animal origin such as animal dung or animal manure, or of plant or herbal origin such as leaves, stalk, root, seed, fruit body or blossom.

Where the sample is a turfgrass sample, it can be a sample obtained from the turf grass as a whole or from a sample of a part of the turfgrass, such as the root, seeds or soil. In particular, the sample can be a turfgrass root or seed or a blade of the turfgrass sample.

In particular, turfgrasses to be considered within the frame of the present invention are notably described in the Compendium of Turfgrass Diseases, Third Edition (2005) issued by the American Phytopathological Society.

More particularly, the turfgrass sample is selected from the group consisting of the *Festaceae*, *Aveneae*, *Triticeae*, *Chlorideae*, *Zoysieae*, *Paniceae* and *Andropogoneae* Tribe.

Where the sample is a soil sample, it is preferably taken directly under the diseased area or in the vicinity of the diseased area, preferably turfgrass.

The sample can be obtained directly from turfgrass, fertilizer, seed, crop or soil, or be obtained after treatment steps, such as grinding, freezing or extraction, in particular nucleic acid extraction, steps.

In a particular embodiment, the sample in which nucleic acids are to be detected is not of human origin.

As intended in the present invention, the at least one nucleic acid as defined above or the set of nucleic acid as defined above is to be used in an hybridization technique, in particular a nucleic acid hybridization-based detection method. Examples of such methods are Microarray-Chip, Southern blot analysis, Northern blot analysis, or Polymerase chain reactions (PCR) wherein the nucleic acids can be used as a primer, probe or template for hybridization. Depending on the method of choice, the nucleic acid is labeled or unlabeled. Examples of such labels are biotin, streptavidin, fluorescent chemicals, radioactive isotopes, Digoxigenin and many others which allow for the detection of bound DNA.

Nucleic acid hybridization-based detection methods are well known to a person skilled in the art.

Microarray-Chips, for instance, are well described in "Microarray Technology in Practice" by Steve Russell (2008, ISBN-10: 012372516X, ISBN-13: 978-0123725165) and are used in numerous publications (Oh S, et al., Genomic diversity of Escherichia isolates from diverse habitats. PLoS One. 2012;7(10):e47005.doi: 10.1371/journal.pone.0047005. Epub 2012 Oct 8).

Polymerase chain reaction (PCR), Southern blot analysis and Northern blot analysis, for instance, are well described in Molecular Cloning: A Laboratory Manual, 4th Edition (2012, ISBN-10: 1936113422 | ISBN-13: 978-1936113422).

For polymerase chain reactions as detection method of choice, at least one nucleic acid as defined above or the set of nucleic acid as defined above, is used to convey specificity of the PCR, wherein the nucleic acid can be labeled or unlabeled and be used as primer or probe.

For Southern or Northern blot analysis as detection method of choice, at least one nucleic acid as defined above or the set of nucleic acid as defined above is used during hybridization wherein the nucleic acid can be labeled or unlabeled.

For Microarray-Chip as detection method of choice, at least one nucleic acid as defined above or the set of nucleic acid as defined above is used as probe on the Microarray-Chip, wherein the nucleic acid can be labeled or unlabeled.

The Microarray-Chip hybridization methods above can be combined with general PCRs to increase sensitivity or specificity of the assay. At least one nucleic acid sequence

as defined above or the set of nucleic acid as defined above will be part of the final PCR product. Such amplification methods, using PCRs, and variations thereof, are well known to the person skilled in the art.

In a particular embodiment of the present invention, the hybridization technique
5 can thus comprise a PCR combined with a Microarray-Chip, wherein the PCR increased the sensitivity, reduces background and/or adds detection label to the PCR product.

If nucleotide sequences from fungi or isolated DNA from fungi are to be amplified by PCR, the PCR comprises primers targeting a sequence within the genome (Internal Transcribed Spacer (ITS)) of the one or more fungi comprising at least one nucleic acid
10 sequence described above. In particular, one primer targets the 18S rDNA/ITS1 region or the ITS2/28S rDNA region of the one or more fungi. The fungal rRNA operon comprises the 18S rRNA gene, the Internal Transcribed Spacer 1 (ITS1), the 5.8S rRNA gene, the Internal Transcribed Spacer 2 (ITS2), and the 28S rRNA gene. The 18S rDNA/ITS1 and ITS2/28S rDNA regions thus relate to the regions of the genome of the one or more fungi
15 in the vicinity respectively of the junction of the 18S rRNA gene and ITS1, and of the junction of ITS2 and the 28S rRNA gene. As will be clear to anyone of skill in the art, to design one nucleic acid as a forward primer, the primer which targets the 18S rDNA/ITS1 region or the ITS2/28S rDNA region will be a reverse primer, and vice versa. The sequences of the 18S rDNA/ITS1 and ITS2/28S rDNA regions are well known to one of
20 skill in the art and can usually be accessed from public sequence databases. Where sequences of the 18S rDNA/ITS1 and ITS2/28S rDNA regions would not be publicly available for a particular fungus species, they can be routinely sequenced. Besides, it is well within the common knowledge of anyone skilled in the art to select primers within the known sequences.

By way of example of a primer targeting the 18S rDNA/ITS1 region, one can cite the so-called "ITS1 primer" of sequence TCCGTAGGTGAACCTGCGG (SEQ ID N°101). Conversely, by way of example of a primer targeting the ITS2/28S rDNA region one can cite the so-called "ITS4 primer" of sequence TCCTCCGCTTATTGATATGC (SEQ ID N°
102). Still other examples are available from WO 2009/147017 A1, White et al. 1990,
30 Gardes & Bruns 1993.

If nucleotide sequences from bacteria or isolated DNA from bacteria are to be amplified by PCR, the PCR comprises primers targeting a sequence within the genome (16S rRNA) of the one or more bacteria comprising at least one nucleic acid sequence described above. However, it is preferred that the primer targets the gene for the 16S
35 rRNA. As will be clear to anyone skilled in the art, to design one nucleic acid as a forward primer, the other primer will be a reverse primer, and vice versa. The sequences of the

16S rRNA gene are well known to one skilled in the art and can usually be accessed from public sequence databases. Where sequences of the 16S rRNA gene would not be publicly available for a particular bacterial species, they can be routinely sequenced. Besides, it is well within the common knowledge of anyone skilled in the art to select primers within the known sequences.

By way of example of a primer targeting the 16S rRNA, one can cite the so-called "EUB8m_f primer" of sequence AGAGTTTGATCMTGGCTCAG (SEQ ID N°103) and "EUB1088_r primer" of sequence CTCGTTGCGGGACTTAACC (SEQ ID N°104), wherein one primer acts as forward primer and the other as reverse primer.

As intended herein the primers to be used may be unmodified or modified nucleic acids, in particular DNA. Where the primers are modified nucleic acids they can notably be labeled nucleic acids, in particular biotin labeled, fluorescently labeled or radioactively labeled nucleic acids.

Nucleic acids may need to be isolated and purified from organic samples (e.g. turfgrass, soil, crops, plants, fertilizers, seeds etc.). Methods for the isolation of nucleic acids are well known for the person skilled in the art and may depend on the method for detection used for detection and source of organic sample. By way of example, methods for nucleic acid isolation are described in the scientific literature (e.g. Michele K. Nishiguchi, Phaedra Doukakis, Mary Egan et al., "DNA Isolation Methods", Methods and Tools in Biosciences and Medicine, Techniques in molecular systematics and evolution, ed. by Rob DeSalle et al.© 2002 Birkhäuser Verlag Basel/Switzerland). Still other examples are many commercially available methods or kits (e.g. "DNeasy® Plant Mini Kit (QIAGEN 69104) or "PowerSoil® DNA Isolation Kit (MO BIO laboratories Inc, Cat no. 12888-100).

As mentioned above, the present invention also relates to a Microarray-Chip comprising at least one nucleic acid according to the present invention.

In particular, the Microarray-Chip comprises the all set of nucleic acids according to the invention, and mentioned above.

According to an embodiment of the present invention, the Microarray-Chip according to the invention is used for the detection of at least one pathogen and/or beneficial microorganism in a sample, the said pathogen, beneficial microorganism and sample being as defined above.

In particular, the Microarray-Chip according to the invention is used for the detection of all of the fungi and bacteria mentioned in the above list.

As mentioned above, the present invention also relates to a detection method of pathogens and/or beneficial microorganisms in an organic sample.

In particular, this method is for detection of all of the fungi and bacteria defined in the above mentioned list, and two PCRs are conducted in its step b), the first directed to fungus DNAs and the second directed to bacterial DNAs.

In a particular embodiment, the two PCRs are combined into one single reaction tube.

Experimental procedure

DNA is isolated from organic sample (e.g. turfgrass, soil, crops, plant parts etc.) and from standards (isolated bacteria, isolated fungus etc.). Methods are known for a skilled person. Examples are described in the scientific literature (e.g. Michele K. Nishiguchi, Phaedra Doukakis, Mary Egan et al., "DNA Isolation Methods", Methods and Tools in Biosciences and Medicine, Techniques in molecular systematics and evolution, ed. by Rob DeSalle et al. © 2002 Birkhäuser Verlag Basel/Switzerland) and many commercially methods are available, too. (e.g. "DNeasy[®] Plant Mini Kit (QIAGEN 69104) or "PowerSoil[®] DNA Isolation Kit (MO BIO laboratories Inc, Cat no. 12888-100).

A Fungi-PCR (ITS) can be conducted. By means of a fungi specific, but species unspecific PCR, the rDNA internal transcribed spacer sequence (ITS) is amplified. Example for such a primer pair are the ITS1 (TCCGTAGGTGAACCTGCGG) (SEQ ID N°101) and ITS4 (TCCTCCGCTTATTGATATGC) (SEQ ID N°102), targeting the 18S rDNA/ITS1 and ITS2/28S rDNA region, as described elsewhere (White et al. 1990, Gardes & Bruns 1993). Other primer pairs, amplifying the same region, are applicable, too, with this method. A person skilled in the art will be able to design and/or use such primers that target this region in all organisms.

A Bacterial-PCR (16S rRNA) can also be conducted. By means of a bacterial specific, but species unspecific PCR, the bacterial 16S rDNA is amplified. Example for such a primer pair is "EUB8m_f primer" of sequence AGAGTTTGATCMTGGCTCAG (SEQ ID N°103) and "EUB1088_r primer" of sequence CTCGTTGCGGGACTTAACC (SEQ ID N°104). Other primer pairs, amplifying the same region or the whole 16S rRNA, are applicable, too, with this method. A person skilled in the art will be able to design and/or use such primers that target this region in all organisms.

The two PCRs can be combined into a single reaction tube. The person skilled in the art will be able to do this.

Eventually the PCR product has to be labelled. Many methods and dyes exist to do so (fluorescent dye, biotin, isotopes, Digoxigenin labelling). The PCR product can be labelled during PCR (e.g. labelled primers) or the PCR product is labelled after amplification.

The PCR product is purified by means of standard methods (e.g. QIAGEN PCR purification Kit).

The individual or combined purified PCR products are hybridized to the DNA on a Microarray-Chip according to the present invention and as described above, and stringently washed in order to remove unbound DNA (e.g. Hybridisation Kit, ALERE Technologies GmbH, Germany, order numbers 245200100 and 24520K100).

Bound DNA is detected by a detection method specific to the label that was added to the PCR product during amplification.

Such detection methods are well known by the person skilled in the art. For example, fluorescent labels can be measured using apparatus capable to excite the fluorochrom and to measure the emitted signal using fluorescent sensitive cameras. Biotin labels can be detected using Streptavidin linked with horseradish peroxidase (HRP). HRP catalyzes the conversion substrates into coloured products. The presence of HRP and thus of biotin is then measured by the detection of processed substrate. Examples of substrates are 3,3'-Diaminobenzidine or 3,3',5,5'-Tetramethylbenzidine. Methods for product measurement depend on the substrate and may be chemiluminescent signals sensitive cameras or optical cameras measuring the optical spectrum, including the human eye. Specific examples of such apparatus are ATR3 ArrayTube™ Reader (for Microarrays, by Identibac, Alere) or ChemiDoc™ MP imager (for southern and northern blots by Biorad) and Rotor Gene Q (for PCR by QIAGEN).

It is then possible to deduce therefrom if the sample contains pathogens and/or beneficial microorganisms and to identify the species present in the sample.

In particular, this method can be used for screening of organic samples before its use for absence of pathogens and can detect pathogens already before the pathogen emerges and causes disease.

As mentioned above, the present invention also relates to a detection kit for the detection of pathogens and/or beneficial microorganisms in an organic sample.

As mentioned above, the present invention also relates to a method of treating a diseased turfgrass.

Thereby, thanks to the present invention, analysis of turfgrass, soils, natural fertilizers (e.g. absence of pathogens), seeds, and crops and detection of causative pathogen in grass showing the disease is possible.

The invention will be further illustrated by the following example.

Example 1

Microarray tubes were covalently linked with nucleic acid sequences according to the invention (SEQ ID NOs 1 to 100) (provided by Alere Technologies GmbH, Jena, Germany). The nucleic acid sequences according to the invention and above described were spotted in duplicates onto the microarray in order to generate a reproducible result.

5 Each spot contained 0.02 μ mol of nucleotide and the nucleotides were immobilized onto the array by a 3'C7-amino modification.

DNA from isolated fungi, turfgrass or soil was extracted using the commercial kit PowerSoil® DNA Isolation Kit (from MO BIO laboratories Inc., order number 12888-100) according to the manufacturer's instructions. DNA from samples consisting of turfgrass

10 was isolated using maximal 0.3 g of soil.

Amplification of fungal rDNA genomic region (ITS) was performed using the primer pairs ITS1 and ITS4 that contained a biotin label at their 5' end. Primers were purchased from Microsynth (Balgach, Switzerland). PCR products were purified using the "PCR purification kit" from Qiagen (#28104) to remove residual primers, buffer and nucleotides.

15 The resulting purified biotinylated PCR products were hybridized to the microarray tubes. The Hybridization protocol and reagents were from Alere (Hybridisation Kit, ALERE Technologies GmbH, Germany, order numbers 245200100). The hybridization temperature was increased (from 55 to 61°C) to increase specificity. After hybridization and washing off unbound DNA, bound, biotinylated PCR products were detected using

20 horseradish peroxidase labeled streptavidin and horseradish peroxidase substrate from Alere (Hybridisation Kit, ALERE Technologies GmbH, Germany, order numbers 24520K100).

The signal intensities on the microarray tubes and location to the spotted chip were measured with the ATR3 (Alere tube reader version 3). Intensity measurements produced

25 a semi-quantitative value of the bound PCR-product. The analysis software (IconoClust, Standard version, Build: 3.6r0), developed by Alere, processed signal intensities and background signals received from a given spot and allocated signal intensities to the nucleic acid present at a given spot. Each spot produced signal intensities between approximately 0 and +1 (Arbitrary units), calculated from control spots and signal intensity.

30 Results are provided in Table 2 below.

Table 2: Signal intensities detected in different samples (A-E: individual turfgrass samples, F: mix of isolated fungal cultures: *Bipolaris sorokiniana* strain 583, *Sclerotinia minor* strain 454, *Pyrenochaeta lycopersici* strain 45, expected to bind to probes BISO, SCSC and PYLY, *S.minor* is very closely related to *S.sclerotiorum* and expected to bind

35

the probe. Signal intensities >0.07 are written in bold. Correspondance between Oligo ID and species can be found in Table 1.

Oligo ID	A	B	C	D	E	F
ARME	0.00	-0.01	0.01	0.00	0.00	0.00
ASPH	0.65	0.84	0.28	0.35	0.62	0.06
BASU	0.05	0.01	0.00	-0.01	0.00	-0.01
BISO	0.00	0.00	0.00	-0.01	0.12	0.83
BLGR	0.01	0.00	0.00	0.06	0.00	0.00
CEZE	0.00	0.00	0.00	-0.01	0.01	0.00
CHEL	0.00	0.00	-0.01	0.01	-0.01	-0.01
CHTH	0.00	0.01	0.01	0.01	0.01	0.01
CLPU	0.00	0.01	0.00	0.01	0.00	0.00
COFU	0.00	0.01	0.01	0.00	0.00	0.00
COTR	0.00	0.00	0.00	0.01	0.00	0.00
CUAF	0.00	0.00	0.00	0.18	0.00	0.00
DRDI	0.02	-0.01	-0.01	0.01	-0.01	0.00
DRPH	0.00	0.00	0.01	-0.01	0.00	0.00
DRPO	0.00	0.01	0.00	-0.01	-0.01	0.01
DRSI	0.00	0.01	-0.01	-0.01	0.00	0.00
DRTR	0.00	0.02	0.00	-0.01	-0.01	0.00
EPTY	0.00	0.00	0.01	0.00	0.00	0.01
EXTU	0.00	0.01	0.01	0.01	-0.01	0.00
FUCU	0.00	0.01	0.00	0.00	0.00	-0.01
FUPO	0.01	0.00	-0.01	-0.01	0.01	0.00
FUPR	0.01	0.86	0.60	0.79	0.24	0.01
GAGR	0.00	0.00	0.00	0.01	0.00	0.00
GLCA	0.01	0.01	0.00	0.02	0.01	0.00
GLGR	-0.01	-0.01	-0.01	-0.01	0.00	0.00
KACA	0.00	0.00	0.00	0.00	0.00	-0.01
LEAU	0.03	0.81	0.04	0.05	0.22	0.00
LEKO	-0.01	0.01	0.01	0.00	0.00	0.00
LENO	0.00	0.00	0.01	0.01	-0.01	-0.01
MAOR	0.00	0.00	-0.01	0.00	0.00	-0.01
MAPH	0.01	0.00	0.00	-0.01	0.01	0.01
MAPO	0.00	0.01	0.00	0.00	0.00	0.00
MIBO	0.00	0.05	0.00	-0.01	0.00	-0.01
MINI	0.53	0.01	-0.01	-0.01	0.00	0.00
MITR	0.01	0.01	0.00	0.00	0.01	0.00
MYRO	0.00	0.00	0.01	-0.01	0.01	0.00
NESP	0.00	0.01	0.00	0.00	0.00	0.00
OPHE	0.00	0.00	0.00	-0.01	0.00	-0.01
PETR	0.00	0.01	0.00	-0.01	0.00	0.00
PHCA	0.00	0.01	-0.01	0.01	0.00	0.00
PHCI	-0.01	0.01	0.00	0.01	0.01	0.00
PHDR	-0.01	0.00	-0.01	0.02	0.00	0.00
PHFR	0.00	-0.01	0.00	-0.01	-0.01	-0.01
PHGR	0.00	0.38	0.01	0.00	-0.01	-0.01
PHME	0.07	-0.01	0.00	0.00	0.00	0.00
PHNI	0.00	0.00	0.00	0.01	0.00	0.00
PHSP	0.00	0.01	0.00	-0.01	0.00	0.00
PLBR	0.01	0.00	0.00	0.02	0.00	-0.01
PSCH	0.01	0.00	-0.01	0.01	-0.01	-0.01
PSME	0.00	0.00	0.01	0.01	0.01	0.00
PUCO1	0.00	0.01	0.00	0.00	0.01	0.00

PUCO2	0.00	0.01	0.00	-0.01	0.00	0.00
PUGRA	-0.01	0.01	0.00	0.00	0.00	0.00
PUPA	-0.01	0.01	-0.01	0.00	-0.01	0.00
PUPO	0.00	0.01	0.00	-0.01	0.00	0.00
PURE	-0.01	0.01	0.01	0.00	0.00	0.00
PUSO	0.00	0.01	0.01	-0.01	0.01	0.00
PUST	0.00	0.01	0.01	0.01	-0.01	0.00
PUTR	0.00	0.00	0.00	0.01	0.00	0.00
PYAP	0.00	0.00	0.01	0.01	0.01	0.00
PYDE	0.01	0.01	0.01	0.01	-0.01	0.00
PYIR	0.00	0.00	0.01	0.00	0.00	0.01
PYLY	0.00	0.01	0.01	0.01	0.01	0.07
PYMA	0.01	0.00	0.01	0.00	0.00	0.00
PYSU	0.01	0.00	0.00	-0.01	-0.01	-0.01
PYUL	-0.01	0.00	0.00	0.00	0.00	0.00
RACO	0.00	0.00	0.01	-0.01	0.00	0.00
RHCE	0.00	-0.01	0.00	0.12	0.13	-0.01
RHFR	0.02	0.01	0.02	0.00	0.00	0.00
RHOR	0.00	0.01	0.00	0.01	0.01	0.00
RHSE	-0.01	0.01	-0.01	0.00	0.01	-0.01
SCBO	0.00	0.00	0.02	0.01	-0.01	0.03
SCHO	0.05	0.00	0.00	0.00	0.01	0.01
SCSC	0.00	0.01	0.00	0.01	-0.01	0.36
SCTR	0.07	0.01	0.00	0.01	0.02	0.01
SEEN	-0.01	0.00	-0.01	-0.01	0.00	0.00
SEMA	0.00	0.00	0.00	0.00	0.00	0.00
SEPL	-0.01	0.00	-0.01	-0.01	-0.01	0.00
SETR	0.00	0.01	0.00	0.01	0.00	0.00
STAL	0.00	0.00	0.01	0.01	0.00	0.00
STGR1	0.01	0.01	0.00	0.01	0.00	0.00
STGR2	-0.01	0.01	-0.01	-0.01	0.00	0.00
STRI	0.00	0.02	0.00	0.00	-0.01	0.00
STSA	0.01	0.00	0.00	0.00	0.01	0.00
STSC	0.00	0.00	0.02	0.01	0.01	0.01
THCU	0.00	0.01	0.00	-0.01	-0.01	0.00
TICA	0.00	0.01	0.00	0.00	0.00	0.00
TICO	0.00	0.00	0.00	-0.01	-0.01	0.00
TRHA	-0.01	0.00	0.26	0.01	0.00	0.00
TYIN	0.00	-0.01	0.00	0.00	0.00	0.00
TYIS1	0.00	0.00	0.01	0.01	0.01	0.00
TYIS2	0.00	-0.01	-0.01	-0.01	-0.01	-0.01
URDA	0.00	0.01	0.01	0.01	0.01	0.00
URTR	0.00	-0.01	-0.01	0.01	0.00	0.00
USMA	0.00	0.00	0.00	0.01	-0.01	0.01
USNU	-0.01	0.00	0.00	0.01	-0.01	0.00
USST	0.01	0.00	0.00	0.01	0.00	0.00
USTR	0.00	0.01	0.00	-0.01	0.00	0.00
VEDA	0.14	0.63	0.56	0.79	0.78	0.01
XATR	-0.01	0.00	0.01	0.00	0.00	0.00
Biotin-Control	0.85	0.85	0.85	0.87	0.87	0.85

Conclusion:

The present invention allows identification of microorganisms from a large selection of fungi and bacterial species.

The method according to the invention is faster, has a broader spectrum of microorganisms that can be identified and is more specific than conventional methods.

5

10

15

20

25

30

35

CLAIMS

1. The use of a set of nucleic acids comprising or consisting of:

(i) the sequences selected from the group consisting of:

- GGGTTGCTTGCTTGCGAGCTCC (SEQ ID N° 1);
- CTGGTGTGGACTCGCCTTAAAC (SEQ ID N° 2);
- GCACATATTTTGCCTTTGTATCAGG (SEQ ID N° 3);
- TCCGCCAGGGAARACCAAACTCT (SEQ ID N° 4);
- CGGAGCGCGGGCCGTCGCG (SEQ ID N° 5);
- ACTTATACCCAAAACGTTGCCTCG (SEQ ID N° 6);
- CATTATCGAGTTTACGCTCCATAAC (SEQ ID N° 7);
- GCGCAGCTATTAGATCTACGGTG (SEQ ID N° 8);
- GCAAGGCTGGAGTATTTTATTACCCT (SEQ ID N° 9);
- CCGTGGCCTTGTTGCCACGCCC (SEQ ID N° 10);
- ATTGGGGCCTTGTTGCCACACCC (SEQ ID N° 11);
- TTTTGCGCTTTGTCCAGTTGCGG (SEQ ID N° 12);
- GATTCGTCGCCCCCCTCCTGG (SEQ ID N° 13);
- GACCTTATTCAAACCTTTTTTTTTCAGTT (SEQ ID N° 14);
- GCTGTTGGGGACCGGCTCACCCG (SEQ ID N° 15);
- TGACCGTTGTCACGAGACGACTTTAT (SEQ ID N° 16);
- CTTGGTGTGGGAGCTGCAGTCC (SEQ ID N° 17);
- CCATTGCGTAGTAGTAAAACCC (SEQ ID N° 18);
- AACGCGCTTCGTTCCGAGGCTT (SEQ ID N° 19);
- TTCAACCCTCAAGCCCCCGGGTTTG (SEQ ID N° 20);
- CGAAGTAGTGATATTCCGCATCGG (SEQ ID N° 21);
- GTTAGGGGGTCCCCTCTCCGG (SEQ ID N° 22);
- CTCGGTCTCGAGCCGCCGG (SEQ ID N° 23);
- ACACCCCATTTGAACCTATTTATTTTYAA (SEQ ID N° 24);
- ACATCTCGCGCTTTGCATTCAGAA (SEQ ID N° 25);
- CGATTTTGGGGGGTGGCTAGTGC (SEQ ID N° 26);
- CTCTGAGTACGAAAAGAACCTGAAA (SEQ ID N° 27);
- TGCTGGCTCTTCTAGAGTCGGCTC (SEQ ID N° 28);
- GGCCAGACGACAGCCATAAACC (SEQ ID N° 29);
- CGCCGGTGGACTACCTAAACTCT (SEQ ID N° 30);
- CGTCGTCGCTGTTGCAAGGAC (SEQ ID N° 31);

- TCGGGCAACGGAACCAGGCGC (SEQ ID N°32);
- GTTGCCCTCGGCGGGCACGGC (SEQ ID N°33);
- CCAGTTATATAGGCACCCAATAAGCC (SEQ ID N°34);
- CACCAAACCAGCTTGGGAAACCTT (SEQ ID N°35);
- 5 - GTCACGTGGTCTTGGTTTTGAA (SEQ ID N°36);
- CGCAACCGGGAGCCGCGGCGCGG (SEQ ID N°37);
- TGGACTGGCTTCGGCTAGACTGG (SEQ ID N°38);
- ACCATAGCTCAGTTGCTTGGCTTTT (SEQ ID N°39);
- TGGCGAATGTTTGGACTTCGGTCT (SEQ ID N°40);
- 10 - GCGCAAGCTGGGGTGGRCGAG (SEQ ID N°41);
- GAACTTGTGTCTCTGCGGCGCG (SEQ ID N°42);
- AACTGGTGAACCGTAGCTGTGTGGT (SEQ ID N°43);
- CCGTACATTAACTTGACTTTCTTCC (SEQ ID N°44);
- GGAACTCCACCCTTGAATACACTG (SEQ ID N°45);
- 15 - CGACCCCTTTTATAATTCACCCAAC (SEQ ID N°46);
- CTAAAAAACCCCTCATAACCTTTTTTTT (SEQ ID N°47);
- CTCCTAAAACCCAATATCTTATTTTAAAG (SEQ ID N°48);
- CAATACTGCCATCTTGTTTTTGAAGG (SEQ ID N°49);
- ATACTTGCCATCTTTTTGGAAGG (SEQ ID N°50);
- 20 - CCTAAAAACCCCCCTTATCA (SEQ ID N°51);
- CAACCTTTTTGGAGTATTCTAATGAT (SEQ ID N°52);
- ATACTGCCATCTTATTKAAGGGAGAC (SEQ ID N°53);
- CAGGGCTATCCCCCTGCCAGG (SEQ ID N°54);
- GGAGGGTTGCGCACTTTGTGCGTG (SEQ ID N°55);
- 25 - GACGCCCTGTTTTCGGATAGGG (SEQ ID N°56);
- TGGCGTGCGTTTGCTTGCGCTTC (SEQ ID N°57);
- CTATACTCCGAGAACGAAAGTTTTTGG (SEQ ID N°58);
- TAGTAGTGTGTGTRGCACGTTGTC (SEQ ID N°59);
- TGTTTTTGTGTTTGTGGAAATACGCTGTTT (SEQ ID N°60);
- 30 - GTAGAATTTTGCTGCTCTTGGGCG (SEQ ID N°61);
- CTGTGTAGTCAGGGATGGAATGTGC (SEQ ID N°62);
- TGAACGCATCATGTTGCTTCGGG (SEQ ID N°63);
- CTGGCTTTTGTGTTTGGATTTGGAGGT (SEQ ID N°64);
- CAGCGACAACCGACTCTAAGTTCA (SEQ ID N°65);
- 35 - CTCGTGAAACACATGAAGTCTGAG (SEQ ID N°66);
- TCGGCGCCCCAGGAGAAATCCT (SEQ ID N°67);

- AGTCCATGTCCGCAATGGCAGG (SEQ ID N°68);
 - AACACATACCTCTCGTTACAGGGTC (SEQ ID N°69);
 - GAGCTGCTCTTCGGGGCCTTGTAT (SEQ ID N°70);
 - CTTGTATGCGCGCCAGAGAATATCA (SEQ ID N°71);
 - 5 - TATTGGGCGTCCGCGGGGGA (SEQ ID N°72);
 - GCGGAGTTCACGAGCCCTCAC (SEQ ID N°73);
 - GCGCCTTGTCTCTCGCGAGAC (SEQ ID N°74);
 - AGCTGGATCTCAGTGTTATGCTTGG (SEQ ID N°75);
 - GTGCCTCTCGGGGCTTCTGCCG (SEQ ID N°76);
 - 10 - CCTGTGTAGTAATGCTTAGCTTACAC (SEQ ID N°77);
 - CTACGGAGGGGTGGCTGCGTTG (SEQ ID N°78);
 - CTACGGAGGGGTGGCTGCGTTG (SEQ ID N°79);
 - ACAGCTCTGAGCAAAAATTCAAAATG (SEQ ID N°80);
 - GTCCAATGTAGGCGCAGCGTAA (SEQ ID N°81);
 - 15 - ATTAGCTGGAACCTCTTGTGGACC (SEQ ID N°82);
 - ATTAGCTGGAACCTCTTGTGGACC (SEQ ID N°83);
 - CTCATTAAACAATTTTTTCTTATAAAGATTG (SEQ ID N°84);
 - GTCATTGCACTCAGGTAGACGTAACA (SEQ ID N°85);
 - CTTTTTCTTTTGGAAAAGGTTGACG (SEQ ID N°86);
 - 20 - ACAGACAATTTTATTGAACACTTTTT (SEQ ID N°87);
 - TTTGAAGAGTTGGCGGATCGGTAT (SEQ ID N°88);
 - ACGGACAATTTTATTTAACACTTTTG (SEQ ID N°89);
 - GCCCTTAAAKCAGTGGCGGACC (SEQ ID N°90);
 - TAATACATGCAAGTCGAGCGGACAGAT (SEQ ID N°91);
 - 25 - GGGTACTTACCTAATACGTGAGTAT (SEQ ID N°92);
 - GAAGGGTARTGTCTTAATACGGCATT (SEQ ID N°93);
 - GGAAGGGYAGTGTGTTAATAGCAC (SEQ ID N°94);
 - ATGACTGTCCATCGCATGGTGGAT (SEQ ID N°95);
 - CTGCGGATCGCATGGTCTGCG (SEQ ID N°96);
 - 30 - AATACTTCTCCTCGCATGGGGAGG (SEQ ID N°97);
 - ATGACACACGACCGCATGGTCTGT (SEQ ID N°98);
 - CACCGGAAACGGCCAGAGATGGTTCG (SEQ ID N°99); and
 - GAGTGTGGTAGAGGATGGCGGAA (SEQ ID N°100);
- (ii) the sequences having at least 80% identity with SEQ N°1 SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID

N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100, or

(iii) the complementary sequences of (i) or (ii),

for the detection by hybridization techniques of at least one pathogen and/or one beneficial microorganism in a sample.

2. The use according to claim 1, wherein the at least one pathogen is selected from bacteria and/or fungi.

3. The use according to claim 2 wherein the at least one fungus and/or bacterium is selected from the group consisting of : *Armillaria mellea*, *Ascochyta phleina*, *Bipolaris sorokiniana*, *Blumeria graminis*, *Cercospora zebrine*, *Claviceps purpurea*, *Colletotrichum trifolii*, *Corticium fuciforme*, *Curvularia affinis*, *Drechslera dictyoides*, *Drechslera phlei*, *Drechslera poae*, *Drechslera siccans*, *Drechslera tritici-repentis*, *Epichloe typhina*, *Exserohilum turcicum*, *Fusarium culmorum*, *Fusarium poae*, *Gaeumannomyces graminis*, *Gibberella intermedia*, *Gliocladium catenulatum*, *Glomerella graminicola*, *Kabatiella caulivora*, *Leptosphaeria korrae*, *Leptosphaerulina australis*, *Macrophomina phaseolina*, *Magnaporthe poae*, *Marasmius oreades*, *Microdochium bolleyi*, *Microdochium nivale*, *Microsphaera trifolii*, *Myrothecium roridum*, *Neotyphodium sp.*, *Ophiosphaerella herpotricha*, *Ophiosphaerella narmari*, *Peronospora trifoliorum*, *Phyllachora vulgata*, *Phytophthora brassicae*, *Phytophthora cactorum*,

Phytophthora citricola, *Phytophthora drechsleri*, *Phytophthora fragariae*,
Phytophthora megasperma, *Phytophthora nicotianae*, *Pseudopeziza*
medicaginis, *Puccinia coronata*, *Puccinia coronate*, *Puccinia graminis*, *Puccinia*
poae-nemoralis, *Puccinia poarum*, *Puccinia recondite*, *Puccinia sorghi*, *Puccinia*
striiformis, *Puccinia tritici*, *Pyrenochaeta lycopersici*, *Pythium*
aphanidermatum, *Pythium debaryanum*, *Pythium graminicola*, *Pythium*
irregulare, *Pythium mastophorum*, *Pythium sulcatum*, *Pythium ultimum*,
Ramularia collo-cygni, *Rhizoctonia cerealis*, *Rhizoctonia fragariae*,
Rhynchosporium orthosporum, *Rhynchosporium secalis*, *Sclerotinia borealis*,
Sclerotinia homoeocarpa, *Sclerotinia sclerotiorum*, *Sclerotinia trifoliorum*,
Septoria macropoda, *Septoria tritici*, *Stemphylium sarcinaeforme*,
Thanatephorus cucumeris, *Thielaviopsis basicola*, *Thielaviopsis populi*, *Tilletia*
caries, *Tilletia controversa*, *Trichoderma hamatum*, *Typhula incarnate*, *Typhula*
ishikariensis, *Typhula ishikariensis*, *Uromyces dactylidis*, *Uromyces trifolii-*
repentis, *Ustilago maydis*, *Ustilago nuda*, *Ustilago striiformis*, *Ustilago tritici*,
Verticillium dahliae; *Bacillus amyloliquefaciens*, *Pseudomonas chlororaphis*,
Serratia entomophila, *Serratia plymuthica*, *Streptomyces albidoflavus*,
Streptomyces graminofaciens, *Streptomyces griseoviridis*, *Streptomyces*
rimosus, *Streptomyces scabiei*, *Xanthomonas translucens*.

4. The use according to claim 1 for the detection of all of the fungi and bacteria defined in claim 3.
5. The use according to claim 1, wherein the at least one beneficial microorganism is a plant growth promoting rhizobacteria.
6. The use according to any one of the preceding claims, wherein the sample is a turfgrass; soil; natural fertilizer, in particular a natural fertilizer of animal origin such as animal dung or animal manure, or of plant or herbal origin such as leaves, stalk, root, seed, fruit body or blossom; seed or crop sample.
7. The use according to claim 6 wherein the sample is a turfgrass root or seed or a blade of turfgrass sample.

8. The use according to claim 6 or 7, wherein the turfgrass is selected from the group consisting of the *Festaceae*, *Aveneae*, *Triticeae*, *Chlorideae*, *Zoysieae*, *Paniceae* and *Andropogoneae* Tribe.

9. The use according to any one of the preceding claims wherein the hybridization technique is chosen from Microarray-Chip, Southern blotting, Northern blotting, PCR.

10. The use according to any one of the preceding claims wherein the hybridization technique comprises a PCR combined with a Microarray-Chip, wherein the PCR increased the sensitivity, reduces background and/or adds detection label to the PCR product.

11. A Microarray-Chip comprising a set of nucleic acids comprising or consisting of :

- (i) the sequences SEQ N°1, SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100,
- (ii) the sequences having at least 80% identity with SEQ N°1, SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ

ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100, or

(iii) the complementary sequences of (i) or (ii).

12. Use of a Microarray-Chip according to claim 11 for the detection of at least one pathogen and/or beneficial microorganism in a sample.

13. The use according to claim 12, wherein the at least one pathogen is selected from bacteria and/or fungus.

14. The use according to claim 13 wherein the at least one fungus and/or bacterium is selected from the group consisting of : *Armillaria mellea*, *Ascochyta phleina*, *Bipolaris sorokiniana*, *Blumeria graminis*, *Cercospora zebrina*, *Claviceps purpurea*, *Colletotrichum trifolii*, *Corticium fuciforme*, *Curvularia affinis*, *Drechslera dictyoides*, *Drechslera phlei*, *Drechslera poae*, *Drechslera siccans*, *Drechslera tritici-repentis*, *Epichloe typhina*, *Exserohilum turcicum*, *Fusarium culmorum*, *Fusarium poae*, *Gaeumannomyces graminis*, *Gibberella intermedia*, *Gliocladium catenulatum*, *Glomerella graminicola*, *Kabatiella caulivora*, *Leptosphaeria korrae*, *Leptosphaerulina australis*, *Macrophomina phaseolina*, *Magnaporthe poae*, *Marasmius oreades*, *Microdochium bolleyi*, *Microdochium nivale*, *Microsphaera trifolii*, *Myrothecium roridum*, *Neotyphodium sp.*, *Ophiosphaerella herpotricha*, *Ophiosphaerella narmari*, *Peronospora trifoliorum*, *Phyllachora vulgata*, *Phytophthora brassicae*, *Phytophthora cactorum*, *Phytophthora citricola*, *Phytophthora drechsleri*, *Phytophthora fragariae*,

Phytophthora megasperma, *Phytophthora nicotianae*, *Pseudopeziza medicaginis*,
Puccinia coronata, *Puccinia coronata*, *Puccinia graminis*, *Puccinia poae-nemoralis*,
Puccinia poarum, *Puccinia recondite*, *Puccinia sorghi*, *Puccinia striiformis*, *Puccinia*
trititica, *Pyrenochaeta lycopersici*, *Pythium aphanidermatum*, *Pythium debaryanum*,
5 *Pythium graminicola*, *Pythium irregulare*, *Pythium mastophorum*, *Pythium sulcatum*,
Pythium ultimum, *Ramularia collo-cygni*, *Rhizoctonia cerealis*, *Rhizoctonia fragariae*,
Rhynchosporium orthosporum, *Rhynchosporium secalis*, *Sclerotinia borealis*,
Sclerotinia homoeocarpa, *Sclerotinia sclerotiorum*, *Sclerotinia trifoliorum*, *Septoria*
macropoda, *Septoria tritici*, *Stemphylium sarcinaeforme*, *Thanatephorus cucumeris*,
10 *Thielaviopsis basicola*, *Thielaviopsis populi*, *Tilletia caries*, *Tilletia controversa*,
Trichoderma hamatum, *Typhula incarnate*, *Typhula ishikariensis*, *Typhula*
ishikariensis, *Uromyces dactylidis*, *Uromyces trifolii-repentis*, *Ustilago maydis*, *Ustilago*
nuda, *Ustilago striiformis*, *Ustilago tritici*, *Verticillium dahliae*; *Bacillus*
amyloliquefaciens, *Pseudomonas chlororaphis*, *Serratia entomophila*, *Serratia*
15 *plymuthica*, *Streptomyces albidoflavus*, *Streptomyces graminofaciens*, *Streptomyces*
griseoviridis, *Streptomyces rimosus*, *Streptomyces scabiei*, *Xanthomonas translucens*.

15. The use according to claim 12 for the detection of all of the fungi and bacteria
defined in claim 14.

16. The use according to claim 12, wherein the at least one beneficial microorganism
is a plant growth promoting rhizobacterium.

17. The use according to any one of claims 12 to 16, wherein the sample is a
25 turfgrass; soil; natural fertilizer, in particular a natural fertilizer of animal origin such as
animal dung or animal manure, or of plant or herbal origin such as leaves, stalk, root,
seed, fruit body or blossom; seed or crop sample.

18. The use according to claim 17 wherein the sample is a turfgrass root or seed or a
30 blade of turfgrass sample.

19. The use according to claim 17 or 18, wherein the turfgrass is selected from the
group consisting of the *Festaceae*, *Aveneae*, *Triticeae*, *Chlorideae*, *Zoysieae*,
Paniceae and *Andropogoneae* Tribe.

20. A detection method of pathogens and/or beneficial microorganisms in an organic sample, comprising the steps of:

- a) DNA isolation from the sample and from standards comprising DNA of said pathogens and/or beneficial microorganisms;
- b) amplification of the isolated DNAs of step a) by PCR;
- c) hybridization of the PCR products obtained, on a Microarray-Chip as defined in claim 11;
- d) detection of the bounded DNAs; and
- e) deduction therefrom if the sample contains pathogens and/or beneficial microorganisms.

21. A method according to claim 20, wherein the pathogens to be detected are selected from bacteria and/or fungi.

22. A method according to claim 21, wherein fungi and/or bacteria are selected from the group consisting of: *Armillaria mellea*, *Ascochyta phleina*, *Bipolaris sorokiniana*, *Blumeria graminis*, *Cercospora zebrina*, *Claviceps purpurea*, *Colletotrichum trifolii*, *Corticium fuciforme*, *Curvularia affinis*, *Drechslera dictyoides*, *Drechslera phlei*, *Drechslera poae*, *Drechslera siccans*, *Drechslera tritici-repentis*, *Epichloe typhina*, *Exserohilum turcicum*, *Fusarium culmorum*, *Fusarium poae*, *Gaeumannomyces graminis*, *Gibberella intermedia*, *Gliocladium catenulatum*, *Glomerella graminicola*, *Kabatiella caulivora*, *Leptosphaeria korrae*, *Leptosphaerulina australis*, *Macrophomina phaseolina*, *Magnaporthe poae*, *Marasmius oreades*, *Microdochium bolleyi*, *Microdochium nivale*, *Microsphaera trifolii*, *Myrothecium roridum*, *Neotyphodium sp.*, *Ophiosphaerella herpotricha*, *Ophiosphaerella narmari*, *Peronospora trifoliorum*, *Phyllachora vulgata*, *Phytophthora brassicae*, *Phytophthora cactorum*, *Phytophthora citricola*, *Phytophthora drechsleri*, *Phytophthora fragariae*, *Phytophthora megasperma*, *Phytophthora nicotianae*, *Pseudopeziza medicaginis*, *Puccinia coronata*, *Puccinia coronata*, *Puccinia graminis*, *Puccinia poae-nemoralis*, *Puccinia poarum*, *Puccinia recondite*, *Puccinia sorghi*, *Puccinia striiformis*, *Puccinia tritici*, *Pyrenochaeta lycopersici*, *Pythium aphanidermatum*, *Pythium debaryanum*, *Pythium graminicola*, *Pythium irregulare*, *Pythium mastophorum*, *Pythium sulcatum*, *Pythium ultimum*, *Ramularia collo-cygni*, *Rhizoctonia cerealis*, *Rhizoctonia fragariae*, *Rhynchosporium orthosporum*, *Rhynchosporium secalis*, *Sclerotinia borealis*, *Sclerotinia homoeocarpa*, *Sclerotinia sclerotiorum*, *Sclerotinia trifoliorum*, *Septoria macropoda*, *Septoria tritici*, *Stemphylium sarcinaeforme*, *Thanatephorus cucumeris*, *Thielaviopsis basicola*,

Thielaviopsis populi, *Tilletia caries*, *Tilletia controversa*, *Trichoderma hamatum*,
Typhula incarnate, *Typhula ishikariensis*, *Typhula ishikariensis*, *Uromyces dactylidis*,
Uromyces trifolii-repentis, *Ustilago maydis*, *Ustilago nuda*, *Ustilago striiformis*, *Ustilago*
tritici, *Verticillium dahliae*; *Bacillus amyloliquefaciens*, *Pseudomonas chlororaphis*,
5 *Serratia entomophila*, *Serratia plymuthica*, *Streptomyces albidoflavus*, *Streptomyces*
graminofaciens, *Streptomyces griseoviridis*, *Streptomyces rimosus*, *Streptomyces*
scabiei, *Xanthomonas translucens*.

23. A method according to claim 20 for detection of all of the fungi and bacteria
10 defined in claim 22 wherein two PCRs are conducted in step b), the first directed to
fungus DNAs and the second directed to bacterial DNAs.

24. A method according to claim 23 wherein the two PCRs are combined into one
15 single reaction tube.

25. A method according to claim 20 wherein the beneficial microorganisms are plant
growth promoting rhizobacteria.

26. A method according to any one of claims 20 to 24 for screening samples before its
20 use for absence of pathogens.

27. A kit for the detection of pathogens and/or beneficial microorganisms in an organic
sample comprising:

- (i) the nucleic acids of the following sequences: SEQ N°1 SEQ ID N°2, SEQ ID
25 N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ
ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID
N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19,
SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ
ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID
30 N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35,
SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ
ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID
N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51,
SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ
35 ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID
N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67,

5	SEQ ID N°68, SEQ ID N°70, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100,	(iii)	the nucleic acids of sequences having at least 80% identity with SEQ ID N°1, SEQ ID N°2, SEQ ID N°3, SEQ ID N°4, SEQ ID N°5, SEQ ID N°6, SEQ ID N°7, SEQ ID N°8, SEQ ID N°9, SEQ ID N°10, SEQ ID N°11, SEQ ID N°12, SEQ ID N°13, SEQ ID N°14, SEQ ID N°15, SEQ ID N°16, SEQ ID N°17, SEQ ID N°18, SEQ ID N°19, SEQ ID N°20, SEQ ID N°21, SEQ ID N°22, SEQ ID N°23, SEQ ID N°24, SEQ ID N°25, SEQ ID N°26, SEQ ID N°27, SEQ ID N°28, SEQ ID N°29, SEQ ID N°30, SEQ ID N°31, SEQ ID N°32, SEQ ID N°33, SEQ ID N°34, SEQ ID N°35, SEQ ID N°36, SEQ ID N°37, SEQ ID N°38, SEQ ID N°39, SEQ ID N°40, SEQ ID N°41, SEQ ID N°42, SEQ ID N°43, SEQ ID N°44, SEQ ID N°45, SEQ ID N°46, SEQ ID N°47, SEQ ID N°48, SEQ ID N°49, SEQ ID N°50, SEQ ID N°51, SEQ ID N°52, SEQ ID N°53, SEQ ID N°54, SEQ ID N°55, SEQ ID N°56, SEQ ID N°57, SEQ ID N°58, SEQ ID N°59, SEQ ID N°60, SEQ ID N°61, SEQ ID N°62, SEQ ID N°63, SEQ ID N°64, SEQ ID N°65, SEQ ID N°66, SEQ ID N°67, SEQ ID N°68, SEQ ID N°69, SEQ ID N°70, SEQ ID N°71, SEQ ID N°72, SEQ ID N°73, SEQ ID N°74, SEQ ID N°75, SEQ ID N°76, SEQ ID N°77, SEQ ID N°78, SEQ ID N°79, SEQ ID N°80, SEQ ID N°81, SEQ ID N°82, SEQ ID N°83, SEQ ID N°84, SEQ ID N°85, SEQ ID N°86, SEQ ID N°87, SEQ ID N°88, SEQ ID N°89, SEQ ID N°90, SEQ ID N°91, SEQ ID N°92, SEQ ID N°93, SEQ ID N°94, SEQ ID N°95, SEQ ID N°96, SEQ ID N°97, SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100, , or	(iiii)	the nucleic acids of complementary sequences of (i) or (ii).
25	SEQ ID N°98, SEQ ID N°99 and SEQ ID N°100, , or				
30	28. A method of treating a diseased turfgrass comprising the steps of:				
35	a) detecting by a method combining PCR and Microarray-Chip the absence or the presence of nucleic acids from at least one pathogenic fungus and/or bacterium in a sample of soil in which the diseased turfgrass is growing, or in a sample of the diseased turfgrass, with the set of nucleic acids as defined in claim 1;				

b) if nucleic acids from one or more pathogenic fungi and/or bacteria have been detected in step a), selecting one or more antifungal and/or antibacterial agents which target the one or more pathogenic fungi and/or bacteria from which nucleic acids have been detected; and

5 c) applying the selected one or more antifungal and/or antibacterial agents of step b) to the diseased turfgrass.