

PROCESS FOR PURIFYING ADIPIC ACID BY CRYSTALLIZATION

The present invention relates to a process for purifying adipic acid by crystallization or recrystallization from at least one carboxylic acid.

5 Adipic acid is one of the two base materials for preparing polyamide 6-6. For the applications of polyamide 6-6 it is necessary to have a very high purity, and this purity must exist already at the stage of the precursors, especially at the adipic acid stage.

10 Depending on the process by which adipic acid is prepared the impurities it contains are obviously different. The present process can be applied to adipic acid originating from various processes of synthesis. In fact, one of the most troublesome and sometimes most
15 costly impurities is formed by the presence of traces of the catalyst employed during the preparation of adipic acid.

 However, in the description which follows, the process will be applied more particularly to adipic
20 acid obtained from the double hydroxycarbonylation of butadiene or from the oxidation of cyclohexane.

 The first hydroxycarbonylation of butadiene leads to a mixture of pentenoic acids, principally 3-pentenoic acid. The second hydroxycarbonylation
25 affects the pentenoic acids obtained in the first reaction and leads to adipic acid which also includes a certain amount of 2-methylglutaric acid, 2-ethylsuccinic acid and other compounds originating

from the first hydroxycarbonylation reaction, such as gamma-valerolactone, unconverted pentenoic acids, and methylbutenoic acid. It also includes traces of the catalyst employed in the second hydroxycarbonylation
5 reaction, usually iridium and/or rhodium.

The direct oxidation of cyclohexane to adipic acid is generally carried out in the presence of cobalt, and in this process the adipic acid obtained contains traces of cobalt catalyst.

10 Since adipic acid is of low solubility in water when cold but is much more soluble when hot, this solvent is generally employed for the crystallization of the said acid.

However, owing to the very high purities
15 which are increasingly required for adipic acid, especially insofar as trace metals are concerned, one or even a number of recrystallizations from water often prove to be inadequate.

Besides the trouble which can be caused by
20 the presence of trace metals to the various uses of adipic acid, the intrinsic value of certain catalysts, such as iridium or rhodium, bearing in mind the very large tonnages of adipic acid, means that it is essential to recover them as thoroughly as possible in
25 the context of an economically viable industrial process.

The present invention consists in an improved process for crystallization or recrystallization of

adipic acid, characterized in that the said crystallization or recrystallization is carried out in at least one carboxylic acid having a melting point of less than 20°C.

5 The carboxylic acids employed in the present process are, more particularly, aliphatic carboxylic acids which are saturated or which contain an ethylenic unsaturation.

 They are preferably linear or branched
10 monocarboxylic acids having 2 to 6 carbon atoms.

 As nonlimitative examples of such monocarboxylic acids mention may be made of acetic acid, propionic acid, butanoic acids, pentanoic acids, hexanoic acids and pentenoic acids.

15 Acetic acid and pentenoic acids are preferred, acetic acid owing to its availability and to its use in the synthesis of adipic acid from cyclohexane, and pentenoic acids because they are an intermediate in the preparation of adipic acid from
20 butadiene.

 The purity of the adipic acid recrystallized in this way may be improved further when recrystallization is carried out in the presence of carbon monoxide.

25 The carbon monoxide can make up at least part of the atmosphere above the solution in the crystallization or recrystallization reactor (or reactor headspace) or can create within the said

reactor a pressure which is greater than the atmospheric pressure.

In practice, the process will therefore be operated under an absolute pressure of from 0 bar (preferably at least 0.5 bar) to 50 bars of carbon monoxide, the upper limit not being critical in nature but being representative of industrial apparatus which is not excessively expensive.

The crude adipic acid subjected to recrystallization according to the present process is usually an adipic acid which has already undergone one or more purification treatments, in particular by crystallization from water, by refining or else by distillation, to give it a minimum purity of approximately 95%.

Generally, the adipic acid recrystallized by the process of the invention has a purity of from 95 to 99.95%.

The recrystallization consists in taking the adipic acid to be purified and dissolving it in the minimum amount of hot aliphatic carboxylic acid, i.e. usually at a temperature from 80 to 250°C, optionally under an at least partial pressure or atmosphere of carbon monoxide, and in then inducing crystallization of the dissolved adipic acid by cooling the solution, optionally after having seeded the solution using crystals of pure adipic acid.

Generally, the quantity of carboxylic acid

employed is that which leads to a saturated solution of adipic acid at the chosen temperature. By way of indication, at 90°C the saturated solution in 3-pentenoic acid contains approximately 33% of adipic acid by weight per weight.

The catalyst content of the adipic acid can also be reduced when recrystallization is carried out in the presence of a strong protic acid.

By strong protic acid is meant in the present text an inorganic protic acid having a pKa of less than 1.

As nonlimitative examples of such strong protic acids mention may be made of hydroiodic acid, hydrobromic acid, hydrochloric acid, nitric acid and sulphuric acid.

The quantity of strong protic acid can vary from 0 mol to 100 mol per mole of catalyst metal present in the adipic acid. Preferably, the quantity of protic acid varies from 0 mol (more preferably from 1 mol) to 50 mol per mole of catalyst metal.

The process of the invention likewise embraces the crystallization of adipic acid from reaction mixtures in which it is present.

It is thus possible, for example, to crystallize adipic acid from the mixture obtained by hydroxycarbonylation of pentenoic acid with water and carbon monoxide. This reaction mixture can be mixed with the carboxylic acid in the presence or absence of

the carbon monoxide employed for the hydroxycarbonylation reaction, and the whole mixture can be kept at a temperature from 80 to 250°C as indicated above for the recrystallization.

5 Crystallization can also be carried out in the hydroxycarbonylation reactor by allowing the reaction mixture to cool, preferably under carbon monoxide pressure. This variant can be employed in particular when hydroxycarbonylation is conducted in a
10 carboxylic acid or when it is conducted in 3-pentenoic acid with a degree of conversion of the latter which is incomplete.

 Since the hydroxycarbonylation reaction is conducted in the presence of carbon monoxide, it is not
15 generally necessary to add this compound for the crystallization, although this possibility is not excluded if appropriate.

 Similarly, since the promoter used in the hydroxycarbonylation reaction may be hydroiodic acid or
20 hydrobromic acid, it may not be necessary to add the strong protic acid. However, if desired, the quantity of strong protic acid present in the reaction mixture may be supplemented. As for the recrystallization of adipic acid, it is also possible to operate in the
25 absence of strong protic acid, although this variant is not preferred.

 The recrystallization according to the invention can be carried out a number of times in

succession on the adipic acid in order to reduce further the content of catalyst metal. It is also possible to follow a crystallization or recrystallization according to the invention by one or
5 more recrystallizations from water.

The examples which follow illustrate the invention.

EXAMPLE 1

A glass bulb is charged with 5.2 g of adipic
10 acid, containing 31.2 μ g of Co (0.0006% by weight per weight of adipic acid), and 7.5 ml of acetic acid. The adipic acid has been prepared by direct oxidation of cyclohexane in the presence of Co acetate and has been purified by recrystallization from water. It does not
15 contain any measurable quantities of organic impurities.

The open bulb is placed in a 125 ml autoclave, which is then closed.

The headspace is filled cold with nitrogen
20 (approximately 1 bar).

The temperature is raised to 185°C and is maintained at this level for approximately 30 minutes.

After cooling the autoclave and purging it with nitrogen, the adipic acid is filtered off and the
25 autoclave is rinsed with a few ml of acetic acid.

The adipic acid filtered off is washed with 2 times 5 ml of acetic acid and then 3 times 8 ml of

acetic acid.

The adipic acid is dried overnight in an oven (60°C). The cobalt present in the final adipic acid is assayed by inductively coupled plasma in conjunction
5 with mass spectrometry (ICP/mass). 0.00008% of Co by weight per weight is found.

EXAMPLE 2

Example 1 is repeated with the same charges and under the same operating conditions but with the
10 addition to the reaction mixture employed of 10 molar equivalents of HCl per molar equivalent of Co present in the adipic acid employed.

The same treatment as for Example 1 gives a dry final adipic acid containing 0.000009% of Co by
15 weight per weight.

EXAMPLES 3 TO 8

Example 1 is repeated under the same operating conditions but with the addition to the reaction mixture employed of HI (with a molar ratio of
20 HI to Ir as indicated in Table 1 below), with the use of 3-pentenoic acid (P3) as the recrystallization solvent, and with the use of an adipic acid (AdOH) containing iridium. The adipic acid was prepared by hydroxycarbonylation of 3-pentenoic acid in the
25 presence of a catalyst based on Ir and was purified by recrystallization from water. It contains no measurable

quantities of organic impurities.

Following recrystallization, the adipic acid is washed with 3-pentenoic acid saturated with adipic acid and then with water saturated with adipic acid.

5 Table 1 below summarizes the conditions under which the examples were carried out (Tp = temperature) and the initial and final contents of Ir (initial Ir and final Ir), expressed in micrograms per gram, of the adipic acid employed.

Example	AdOH in g	P3 in g	Temp in °C	Duration in min	CO in bars	Molar ratio HI:Ir	Initial Ir	Final Ir
Ex. 3	2.7	3.9	185	20	30	10	5.0	0.76
Ex. 4	1.43	2.0	90	120	30	10	5.0	2.6
Ex. 5	5.2	7.8	185	30	30	20	2.2	0.42
Ex. 6	6.2	9.3	185	30	1	20	2.2	0.56
Ex. 7	2.9	4.5	185	1200	1	10	0.46	0.18
Ex. 8	8.05	13	185	40	1	20	8.0	1.3

Table 1

EXAMPLES 9 TO 12

Examples 3 to 8 are repeated but using an adipic acid (AdOH) containing rhodium. The adipic acid was prepared by hydroxycarbonylation of 3-pentenoic acid in the presence of a catalyst based on Rh and was purified by recrystallization from water. It contains no measurable quantities of organic impurities.

Following recrystallization, the adipic acid is washed with 3-pentenoic acid saturated in adipic acid and then with water saturated in adipic acid.

Table 2 below summarizes the conditions under which the examples were carried out (Tp = temperature) and the initial and final contents of Rh (initial Rh and final Rh), expressed in micrograms per gram, of the adipic acid employed.

Example	AdOH in g	P3 in g	Tp in °C	Duration in min	CO in bars	Molar ratio HI:Rh	Initial Rh	Final Rh
Ex. 9	4.99	7.5	190	30	30	10	9.5	3.2
Ex. 10	5.02	7.5	190	30	1	13	9.5	3.1
Ex. 11	5.04	7.5	190	30	0 (1 bar of argon)	12	9.5	3.8
Ex. 12	5.05	7.5	190	600	1	14	9.5	0.46

Table 2

EXAMPLE 13

In Example 13, the recrystallization described for Examples 3 to 8 is repeated using the recrystallized adipic acid obtained in Example 7.

5 Table 3 below summarizes the conditions under which the example was carried out and the initial and final contents of Ir (initial Ir and final Ir), expressed in micrograms per gram, of the adipic acid employed.

10 COMPARATIVE TEST 1

 An adipic acid obtained by hydroxycarbonylation of 3-pentenoic acid in the presence of iridium and HI is recrystallized from water. This adipic acid has already undergone one
15 crystallization and still contains 0.00022% of iridium.

 The recrystallization is carried out conventionally by dissolving adipic acid in the minimum amount of water at approximately 95°C and then by gradual cooling of the resulting solution, followed by
20 filtration and, finally, by washing of the adipic acid filtered off with 2 times 5 ml of water and with 3 times 8 ml of water.

 The adipic acid is dried overnight in an oven (60°C). The iridium present in the final adipic acid is
25 assayed. 0.00022% of Ir by weight per weight is found). Therefore, the iridium content of the adipic acid has not been successfully reduced.

Example	AdOH in g	P3 in g	Tp in °C	Duration in min	CO in bars	Molar ratio HI:Rh	Initial Rh	Final Rh
Ex. 13	1.26	2.0	185	30	1	30	0.18	0.14

Table 3

EXAMPLE 14

The hydroxycarbonylation of 3-pentenoic acid is carried out with the following charges:

- 77.7 g of 3-pentenoic acid
- 5 - 88.1 mg of $[\text{IrCl}(\text{COD})]_2$
- 169.6 mg of a 57% by weight aqueous solution of HI
- 5.7 g of water, injected continuously.

The carbon monoxide pressure at the reaction
10 temperature (185°C) is maintained at 22 bars and the reaction is stopped after a period of 1 h 5 minutes.

After gradual cooling under the carbon monoxide pressure of 22 bars, the crystallized adipic acid is drawn off and filtered. Assay by gas
15 chromatography and by high-performance liquid chromatography of the compounds of the filtrate is then used to determine a degree of conversion of the 3-pentenoic acid of 56.6%.

The adipic acid crystallized in this way
20 contains 0.000122% of iridium.

This adipic acid is recrystallized from 3-pentenoic acid under a carbon monoxide atmosphere (circulation bubble by bubble).

After recrystallization, the adipic acid is
25 washed with 3-pentenoic acid saturated with adipic acid and then with water saturated with adipic acid.

The recrystallized and dried adipic acid has an iridium content of 0.00002% by weight per weight.

CLAIMS

1. Process for crystallizing or
recrystallizing adipic acid, characterized in that the
said crystallization or recrystallization is carried
5 out in at least one carboxylic acid having a melting
point of less than 20°C and in that the adipic acid
which is subjected to the recrystallization has a
minimum purity of 95%.
2. Process according to claim 1,
10 characterized in that the carboxylic acid employed is
selected from aliphatic carboxylic acids which are
saturated or which contain an ethylenic unsaturation
and, preferably, from linear or branched monocarboxylic
acids having 2 to 6 carbon atoms.
- 15 3. Process according to either of claims 1
and 2, characterized in that the carboxylic acid
employed is selected from acetic acid, propionic acid,
butanoic acids, pentanoic acids, hexanoic acids and
pentenoic acids.
- 20 4. Process according to one of claims 1 to
3, characterized in that the crystallization or
recrystallization is carried out in the presence of
carbon monoxide.
- 25 5. Process according to claim 4,
characterized in that the carbon monoxide makes up at
least part of the atmosphere above the solution in the
crystallization or recrystallization reactor or creates

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED



within the said reactor a pressure which is greater than the atmospheric pressure.

6. Process according to either of claims 4 and 5, characterized in that it is carried out under an absolute pressure of from 0.5 bar to 50 bars of carbon monoxide.

7. Process according to one of claims 1 to 6, characterized in that the crude adipic acid subjected to recrystallization has a purity of from 95 to 99.95%.

8. Process according to one of claims 1 to 7, characterized in that recrystallization is carried out in the presence of a strong protic acid.

9. Process according to claim 8, characterized in that the strong protic acid is selected from hydroiodic acid, hydrobromic acid, hydrochloric acid, nitric acid and sulphuric acid.

10. Process according to one of claims 1 to 9, characterized in that the quantity of strong protic acid varies from 0 to 100 mol per mole of catalyst metal present in the adipic acid and, preferably, varies from 0 to 50 mol per mole of catalyst metal.

This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

CLAIMS

1. Process for crystallizing or
recrystallizing adipic acid which makes it possible to
reduce the content of trace metals, characterized in
5 that the said crystallization or recrystallization is
carried out in at least one carboxylic acid having a
melting point of less than 20°C and in that the adipic
acid which is subjected to the recrystallization has a
minimum purity of 95%.
- 10 2. Process according to claim 1,
characterized in that the carboxylic acid employed is
selected from aliphatic carboxylic acids which are
saturated or which contain an ethylenic unsaturation
and, preferably, from linear or branched monocarboxylic
15 acids having 2 to 6 carbon atoms.
3. Process according to either of claims 1
and 2, characterized in that the carboxylic acid
employed is selected from acetic acid, propionic acid,
butanoic acids, pentanoic acids, hexanoic acids and
20 pentenoic acids.
4. Process according to one of claims 1 to
3, characterized in that the crystallization or
recrystallization is carried out in the presence of
carbon monoxide.
- 25 5. Process according to claim 4,
characterized in that the carbon monoxide makes up at

AMENDED SHEET





least part of the atmosphere above the solution in the crystallization or recrystallization reactor or creates within the said reactor a pressure which is greater than the atmospheric pressure.

5 6. Process according to either of claims 4 and 5, characterized in that it is carried out under an absolute pressure of from 0.5 bar to 50 bars of carbon monoxide.


10 7. Process according to one of claims 1 to 6, characterized in that the crude adipic acid subjected to recrystallization has a purity of from 95 to 99.95%.

15 8. Process according to one of claims 1 to 7, characterized in that recrystallization is carried out in the presence of a strong protic acid.

9. Process according to claim 8, characterized in that the strong protic acid is selected from hydroiodic acid, hydrobromic acid, hydrochloric acid, nitric acid and sulphuric acid.

20 10. Process according to one of claims 1 to 9, characterized in that the quantity of strong protic acid varies from 0 to 100 mol per mole of catalyst metal present in the adipic acid and, preferably, varies from 0 to 50 mol per mole of catalyst metal.

AMENDED SHEET



PCT WORLD ORGANISATION FOR INTELLECTUAL PROPERTY
International Office
INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International patent classification⁶ : C07C 55/14, 51/43</p>	<p>A1</p>	<p>(11) International publication number: WO 97/46510</p> <p>(43) International publication date: 11 December 1997 (11.12.97)</p>
<p>(21) International application number: PCT/FR97/00939</p> <p>(22) International filing date: 29 May 1997 (29.05.97)</p> <p>(30) Data relating to the priority: 96/07.171 4 June 1996 (04.06.96) FR</p> <p>(71) Applicant (for all designated States except US): RHÔNE-POULENC FIBER AND RESIN INTERMEDIATES <i>124a</i> [FR/FR]: 25, quai Paul Doumer, F-92408 Courbevoie Cedex (FR). <i>8 Nov 95</i></p> <p>(72) Inventors: and</p> <p>(75) Inventors/Applicants (US only): Eric B. HENRIET [FR/FR]: 89, rue du Pensionnat, F-69003 Lyon (FR). Philippe LECONTE [FR/FR]: 43, rue Sainte-Beuve, F-69330 Meyzieu (FR). Carl PATOIS [FR/FR]: 2, avenue Jules Jusserand, F-69003 Lyon (FR). Robert PERRON [FR/FR]: La Pecolière, F-69390 Charly (FR).</p> <p>(74) Representative: Noël VIGNALLY; Rhône-Poulenc Chimie, Industrial Property Directorate, C.R.I.T. - Carrières, P.O. Box 62, F-69192 Saint-Fons Cedex (FR).</p>	<p>(81) Designated States: BR, BY, CA, CN, CZ, JP, KR, MX, PL, RO, RU, SG, SK, UA, US, VN, European Patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p><i>PCT/13/306</i></p> <p>Published With the International Search Report.</p>	

As printed

(54) Title: METHOD FOR PURIFYING ADIPIC ACID BY CRYSTALLIZATION

(54) Titre: PROCÉDÉ DE PURIFICATION DE L'ACIDE ADIPIQUE PAR CRISTALLISATION

(57) Abstract

The invention discloses a method for purifying adipic acid by crystallization or recrystallization in at least one carboxylic acid. More precisely, it discloses an improvement in adipic acid crystallization or recrystallization, characterised in that the said crystallization or recrystallization is effected in at least one carboxylic acid with a melting point below 20 °C. The presence of carbon monoxide during crystallization or recrystallization can have a favourable effect on the adipic acid purity, in particular by reducing the content of metal catalyst traces of the adipic acid. The adipic acid purity can also be improved when the crystallization or recrystallization is effected in presence of a strong proton acid.

(57) Abrégé

La présente invention concerne un procédé de purification de l'acide adipique par cristallisation ou recristallisation dans au moins un acide carboxylique. Plus précisément elle concerne un perfectionnement à la cristallisation ou à la recristallisation de l'acide adipique, caractérisé en ce que ladite cristallisation ou recristallisation est effectuée dans au moins un acide carboxylique ayant un point de fusion inférieur à 20 °C. La présence de monoxyde de carbone pendant la cristallisation ou la recristallisation peut avoir un effet avantageux sur la pureté de l'acide adipique, en particulier pour réduire la teneur de l'acide adipique en traces de catalyseur métallique. La pureté de l'acide adipique peut aussi être améliorée lorsque la cristallisation ou la recristallisation est effectuée en présence d'un acide protonique fort.

ONLY FOR INFORMATION

Codes used to identify the PCT member States on the flyleaves of the brochures in which international applications made under the PCT are published.

AL	Albania	LI	Liechtenstein
AM	Armenia	LK	Sri Lanka
AT	Austria	LR	Liberia
AU	Australia	LS	Lesotho
AZ	Azerbaijan	LT	Lithuania
BA	Bosnia-Herzegovina	LU	Luxembourg
BB	Barbados	LV	Latvia
BE	Belgium	MC	Monaco
BF	Burkina Faso	MD	Republic of Moldova
BG	Bulgaria	MG	Madagascar
BJ	Benin	MK	Former Yugoslav Republic of Macedonia
BR	Brazil	ML	Mali
BY	Belarus	MN	Mongolia
CA	Canada	MR	Mauritania
CF	Central African Republic	MW	Malawi
CG	Congo	MX	Mexico
CH	Switzerland	NE	Niger
CI	Ivory Coast	NL	Netherlands
CN	China	NO	Norway
CU	Cuba	NZ	New Zealand
CZ	Czech Republic	PL	Poland
DE	Germany	PT	Portugal
DK	Denmark	RO	Romania
EE	Estonia	RU	Russian Federation
ES	Spain	SD	Sudan
FI	Finland	SE	Sweden
FR	France	SG	Singapore
GA	Gabon	SI	Slovenia
GB	United Kingdom	SK	Slovakia
GE	Georgia	SN	Senegal
GH	Ghana	SZ	Swaziland
GN	Guinea	TD	Chad
GR	Greece	TG	Togo
HU	Hungary	TJ	Tajikistan
IE	Ireland	TM	Turkmenistan
IL	Israel	TR	Turkey
IS	Iceland	TT	Trinidad and Tobago
IT	Italy	UA	Ukraine
JP	Japan	UG	Uganda
KE	Kenya	US	United States of America
KG	Kyrgyzstan	UZ	Uzbekistan
KP	Democratic People's Republic of Korea	VN	Vietnam
KR	Republic of Korea	YU	Yugoslavia
KZ	Kazakhstan	ZW	Zimbabwe
LC	Saint Lucia		