

SUBSTANTIALLY ANIMAL PROTEIN-FREE RECOMBINANT FURIN AND METHODS FOR PRODUCING SAME

This application claims priority of U.S. Provisional Application No. 61/018,152, filed December 31, 2007, which is incorporated by reference herein in its entirety.

FIELD OF THE INVENTION

[0001] The present invention generally relates to recombinant furin (rFurin) and methods for producing rFurin. More specifically, the invention relates to substantially animal protein-free rFurin and methods for producing substantially animal protein-free rFurin.

BACKGROUND OF THE INVENTION

[0002] Active or mature proteins are usually present in very low amounts in living organisms. Therefore, their pro-proteins or pro-enzymes are preferably activated *in vitro* by contacting them with activation enzymes (e.g. proteases). Pro-proteins (or protein precursors) are inactive proteins that become active by one or more posttranslational modifications and, in particular, by the cleavage of a pro-peptide from the pro-protein. Examples of pro-proteins include, for example, pro-insulin, prothrombin, pro-von Willebrand Factor (pro-VWF), and the like.

[0003] Von Willbrand factor (VWF) is a blood glycoprotein involved in coagulation. VWF is deficient or defective in von Willebrand disease and is involved in a large number of other diseases, including thrombotic thrombocytopenic purpura, Heyde's syndrome, and possibly hemolytic-uremic syndrome. VWF is a glycoprotein circulating in plasma as a series of multimers ranging in size from about 500 to 20,000 kD. Multimeric forms of VWF are composed of 250 kD polypeptide subunits linked together by disulfide bonds. VWF mediates the initial platelet adhesion to the sub-endothelium of the damaged vessel wall, and it is believed that only the larger multimers of VWF exhibit hemostatic activity. VWF multimers having large molecular masses are stored in the Weibel-Pallade bodies of endothelial cells and are liberated upon stimulation. Liberated VWF is then further processed by plasma proteases to result in low molecular weight forms of VWF.

[0004] In humans the removal of the pro-peptide is almost complete, whereas, in mammalian cell lines with a high level of recombinant VWF expression, this process is not very efficient. Therefore, cell culture supernatants from such recombinant cell lines usually comprise a mixture of mature VWF and VWF precursors, like pro-VWF. In order to obtain mature VWF, it is therefore necessary to convert the VWF precursors, in particular pro-VWF, into mature VWF. This process is usually achieved by cleaving the pro-peptide with a protease.

[0005] Current conventional methods produce mature VWF by either incubating its pro-form with proteases in a liquid phase, whereby the maturation itself (i.e., the cleavage of the pro-peptide from the pro-protein) occurs in an unbound state in free solution, or as described, for example, in WO 00/49047, by immobilizing the protease on a solid carrier, which is contacted and incubated with a preparation comprising pro-VWF (see e.g. WO 00/49047). However, these methods comprise various disadvantages over the methods according to the present invention.

[0006] Industrially, VWF and, in particular, recombinant VWF (rVWF) is synthesized and expressed together with recombinant Factor VIII (rFVIII) in a genetically engineered Chinese Hamster ovary (CHO) cell line. The function of the co-expressed rVWF is to stabilize rFVIII in the cell culture process. rVWF is synthesized in the cell in the pro-form, containing a large pro-peptide attached to the N-terminus. Upon maturation in the endoplasmic reticulum and Golgi apparatus, the pro-peptide is cleaved by the action of the cellular protease furin and the mature protein is secreted as a homopolymer of identical subunits, consisting of dimers of the expressed protein. However, the maturation is typically incomplete, leading to a product comprising a mixture of pro-VWF and mature VWF.

[0007] Previous publications have shown that pro-VWF can be converted to mature VWF by *in vitro* treatment with furin or furin-like proteases (Schlokot et al., *Biotechnol. Appl. Biochem.* 24: 257-267, 1996; Preininger et al., *Cytotechnology* 30: 1-15, 1999; and EP 0775750A). In particular, EP 0775750A suggests the co-expression of furin and VWF recombinantly so that the maturation of VWF may occur *in situ*.

[0008] Recombinant furin (rFurin) transforms pro-rVWF (pro-recombinant von Willebrand factor) to rVWF by cleaving the Arg741-Ser742 peptide bond. This

maturation step is part of a rVWF production process for the treatment of von Willebrand Disease Type B and part of the manufacturing process for recombinant Factor VIII – half life (rFVIII-HL). Furin belongs to the family of the pro-protein convertases and is dependent on calcium (Ca^{2+}). Furin specifically cleaves the C-terminal peptide bond of arginine within a specific sequence, containing arginine at positions –1 and –4. This sequence can be found in numerous human proteins, showing that furin plays a major role in the maturation of a number of human pro-proteins.

[0009] The production of activated proteins is of high clinical and diagnostic importance. For example, active or mature proteins, like mature VWF, may be used to control blood coagulation. The present invention provides improved recombinant furin (rFurin) which is substantially animal protein-free rFurin for the subsequent production of activated proteins. More specifically, the present invention provides substantially animal protein-free rFurin for transforming pro-VWF into mature VWF.

SUMMARY

[0010] The present disclosure describes recombinant furin (rFurin), which is substantially animal protein-free recombinant furin (rFurin), and methods for producing same. Such rFurin is substantially free of other proteins which may normally be associated with the production of rFurin, such as serum proteins and host cell proteins. This rFurin allows for the subsequent production of mature proteins with high specific activity and high purity without side effects associated with protein contaminant in the rFurin preparation. More specifically, this rFurin allows for the production of mature VWF with high specific activity and high purity. Accordingly, described herein are methods for selection and adaptation of recombinant host cells to chemically-defined medium, expression of rFurin which is secreted into cell culture supernatant, and purification of rFurin after cell removal. The present invention refers to subject matter as defined in the attached set of claims.

[0011] The substantially animal protein-free rFurin of the present disclosure includes preparations or compositions of rFurin comprising host cell protein in a concentration which ranges between about 0.1 to 0.6 ng protein or less/Unit furin activity or between about 2 and 11 μg protein or less/mL and essentially lacking contaminating proteins from serum in the culture medium. In one example, the substantially animal

protein-free rFurin encompasses preparations of rFurin comprising contaminating host cell DNA in a concentration between about 0 to 0.4 pg DNA or less/Unit furin activity or between about 0 and 24 ng DNA or less/mL and essentially lacking contaminating proteins from serum in the culture medium.

[0012] Described herein are compositions comprising substantially animal protein free recombinant furin at an activity of at least 10000 U furin/mL and host cell protein at a concentration less than about 11 μ g protein/mL. Such compositions may also comprise host cell protein at a concentration less than about 1.0 ng protein/U furin activity. In one example, the host cell protein is from CHO cell.

[0013] In another example, there are disclosed compositions comprising substantially animal protein free recombinant furin at an activity of at least 10000 U furin/mL and host cell DNA at a concentration less than about 14 ng DNA/mL. In various aspects, such compositions also comprise host cell DNA at a concentration less than about 0.5 pf DNA/U furin activity. In one example, the host cell DNA is from a CHO cell.

[0014] Disclosed herein are compositions comprising substantially animal protein free recombinant furin at a specific furin activity of at least about 100 U/ μ g and host cell protein at a concentration less than about 11 μ g protein/mL. Such compositions may also comprise host cell protein at a concentration less than about 1.0 ng protein/U furin activity. In one aspect, the host cell protein is from a CHO cell.

[0015] Further disclosed herein are compositions comprising substantially animal protein free recombinant furin at a specific furin activity of at least about 100 U/ μ g and host cell DNA at a concentration less than about 14 ng DNA/mL. Such compositions may also comprise host cell DNA at a concentration less than about 0.5 pg DNA/U furin activity. In one aspect, the host cell DNA is from a CHO cell.

[0016] Also disclosed herein are methods of making a composition comprising substantially animal protein-free recombinant furin described herein. Such methods comprise the step of adapting the host cells to growth in medium with increasingly lower concentrations of serum until all serum is removed from the medium. In another example, the methods comprise the step of transferring the host cell from growth in medium comprising serum to growth in serum-free medium. In another example, the host cell is a CHO cell.

[0017] Also disclosed are methods of using a composition comprising substantially animal protein-free recombinant furin described herein. Such uses comprise the step of contacting a pro-protein with the composition under conditions to cleave a pro-peptide from the pro-protein to form a mature protein. The rFurin can be used in the formation of any mature protein from a pro-protein that is cleaved by furin. In one example, the mature protein is von Willebrand factor. In another example, the mature protein is Factor VIII. In addition, it is contemplated that the rFurin disclosed herein is useful for both *in vitro* and *in vivo* processing of any pro-protein that it cleaves.

BRIEF DESCRIPTION OF THE DRAWING

[0018] A further illustration of the invention is given with reference to the accompanying drawings, which are set out below in Figures 1-18.

[0019] Figure 1 depicts the expressed active rFurin protease construct in one embodiment of the invention. The rFurin construct is truncated at the C-terminal end at AA 577 to remove the Cys-rich transmembrane and cytosol domains.

[0020] Figure 2 depicts a pedigree of the generation of the CHO/rFurin clone #488-3.

[0021] Figure 3 shows a pedigree of the generation of the CHO/rFurin clone #289-20.

[0022] Figure 4 sets out a comparison of the graphical distribution of the rFurin producers in the cell populations of PMCB#01 and PMCB#04. 80.74% of the cells in PMCB#04 express rFurin; 74.06% of the cells in PMCB#01 express rFurin.

[0023] Figure 5 shows a "Doehlert Matrix" where five temperatures were combined with three pH values, resulting in seven combinations of temperature and pH.

[0024] Figure 6 shows a surface plot analysis of the data in reference to the volumetric productivity. The coordinates of the data in Figure 6 are marked as points. The surface shows the assumed correlation of the single data.

[0025] Figure 7 shows a contour plot which illustrates the influence of temperature and pH on the volumetric productivity. The dots indicate the conditions (pH / temp.) which had been tested experimentally.

[0026] Figure 8 shows a surface plot which is a three-dimensional illustration demonstrating the strong influence of the temperature and the weak influence of the pH on the volumetric productivity.

[0027] Figure 9 shows a surface plot which illustrates the modeled correlation three-dimensionally; it demonstrates the quadratic relationship and shows clearly a maximum for the growth rate at 36.5°C.

[0028] Figure 10 sets out an analysis of the data in reference to specific productivity. There is a similar correlation of the specific productivity with temperature and pH as seen for the volumetric productivity.

[0029] Figure 11 shows that by decreasing the temperature from 37°C to 35.1°C, the volumetric productivity could be increased from approx. 200 kU/L/d to 540 kU/L/d.

[0030] Figure 12 shows SDS-page and silver-stain for rFurin. The band patterns of the Capto-MMC eluates of campaign ORFU06002 and ORFU07002 correlate to a high degree; all samples show a prominent Furin band at approx. 60kDa. A trend to slightly lower molecular weight of the Furin bands is visible in samples of campaign ORFU06002 from batches MMC01 to MMC08 (Figure 12, lanes 1-8).

[0031] Figure 13 shows the Western blot analysis of samples using a monoclonal anti-Furin antibody.

[0032] Figure 14 shows the specific band patterns for rFurin from isoelectric focusing (IEF) and subsequent Western blotting of rFurin samples of campaign ORFU06002.

[0033] Figure 15 shows the specific band patterns for rFurin from isoelectric focusing (IEF) and subsequent Western blotting of rFurin samples of campaign ORFU07002.

[0034] Figure 16 shows Western blot results for rFurin from isoelectric focusing (IEF) and subsequent Western blotting of rFurin samples of campaign ORFU07002.

[0035] Figure 17 shows Furin Reverse Phase HPLC for samples from campaign ORFU06002 (Capto-MMC eluates). Samples were tested with C4 RP-HPLC in order to establish a fingerprint pattern for rFurin.

[0036] Figure 18 shows Furin Reverse Phase HPLC for samples from campaign ORFU07002 (Capto-MMC eluates). Samples were tested with C4 RP-HPLC in order to establish a fingerprint pattern for rFurin.

DETAILED DESCRIPTION OF THE INVENTION

[0037] The present invention relates to the development and production of a recombinant host cell line that is capable of growing in serum-free medium and secreting active recombinant furin (rFurin) into the cell culture supernatant. The host cell line selected for transfection of a plasmid encoding recombinant furin is in one aspect the same as used for expression of recombinant Factor VIII and recombinant VWF. The resulting rFurin is then purified so that is substantially free of animal protein.

[0038] Furin, also known as PACE, PACE4, PC1/PC3, PC2, PC4 and PC5/PC6, belongs to the group of the subtilisin-like serine proteases, which play an important role in the cleavage of proproteins, especially in the secretory synthesis (Van de Ven et al., *Crit. Rev. Oncogen.* 4:115-136, 1993). Pro-proteins are post-translationally, intracellularly processed to their mature form by the endogenous protease in the Golgi apparatus. The protease cleavage site comprises a recognition sequence which is characterized by the amino acid sequence Arg-X-Lys/Arg-Arg. The protease furin cleaves proproteins specifically after this consensus sequence (Hosaka et al., *J. Biol. Chem.* 266:12127-12130, 1991).

[0039] The DNA and amino acid sequence of human and murine furin, as well as further proteins with subtilisin-like protease function have been identified (Roebroek et al., *Mol. Biol. Rep.* 11: 117-125, 1986; Roebroek et al., *EMBO J.* 5:2197-2202, 1986; Barr et al., *DNA Cell Biol.* 10:319-328, 1991; Van den Ouweland et al., *Nucleic Acids Res.* 17:7101-7102, 1989; Van den Ouweland et al., *Nucleic Acids Res.* 18:664, 1990; Smeekens et al., 1990, *J. Biol. Chem.* 265:2997-3000; Smeekens et al., *Proc. Natl. Acad. Sci. USA.* 88: 340-344, 1991; Kiefer et al., *DNA Cell Biol.* 10: 757, 1991; Nakayama et al., *J. Biol. Chem.* 267:5897-5900, 1992; and Hatsuzawa et al., *J. Biol. Chem.* 265: 22075-22078, 1990). The human furin gene encodes a protein consisting of 794 amino acids, certain functions being allocatable to individual characteristic regions: a catalytic center, a middle domain, a cystine-rich

region, a transmembrane domain, and a cytoplasmatic domain (Van de Ven et al., *Crit. Rev. Oncogen.* 4:115-136, 1993). In one aspect, the human furin polypeptide is set out in GenBank Accession No: EAX02111 (National Center for Biotechnology Information, U.S. National Library of Medicine, Bethesda, MD). However, the worker of ordinary skill in the art will appreciate that any protein having furin biological activity, i.e., the ability to cleave pro-proteins (e.g., pro-VWF to produce mature VWF) can be produced by the methods described herein.

[0040] Intact furin is incorporated into the membrane system of the Golgi apparatus where it is functionally active (Bresnahan et al., *J. Cell Biol.* 111:2851-2859, 1990). A truncated form of the over-expressed native furin of 75-80 kD could be detected in cell supernatant as secreted protein (Wise et al., *Proc. Natl. Acad. Sci. USA* 87: 9378-9382, 1990). This naturally secreted furin is known as "shed furin" (Vidricaire et al., *Biochem. Biophys. Res. Comm.* 195:1011-1018, 1993) and is cleaved N-terminally of the transmembrane portion (Vey et al., *J. Cell Biol.* 127:1829-1842, 1994).

[0041] Furin truncated by genetic engineering, in which the encoding part of the transmembrane and cytoplasmatic domains has been deleted, can also be expressed and secreted correspondingly. Such N-terminal deletions have been described for amino acids 714-794 (Leduc et al., *J. Biol. Chem.* 267:14304-14308, 1992; Molloy et al., *J. Biol. Chem.* 267:16396-16402, 1992); for amino acids 716-794 ("Sol-PACE") (Wasley et al., *J. Biol. Chem.* 268:8458-8465, 1993; and Rehemtulla et al., *Blood* 79:2349-2355, 1992); and for amino acids 705-794 (Hatsuzawa et al., *J. Biol. Chem.* 267:16094-16099, 1992). Furin mutants additionally comprising a deletion of the cystine-rich region have also been described (Hatsuzawa et al., *J. Biochem.* 101:296-301, 1992; Creemers et al., *J. Biol. Chem.* 268:21826-21834, 1993).

[0042] The endoproteolytic activity of furin and its selectivity for basic amino acids was first determined in experiments with pro-von Willebrand factor (pro-vWF). Pro-vWF consists of a propolypeptide with 741 amino acids and mature von Willebrand factor (vWF) with 2050 amino acids (Verweij et al., *EMBO J.* 5:1839-1847, 1986). The liberation of mature vWF from pro-vWF results from a proteolytic cleavage after Arg763. Transfection of pro-vWF cDNA in eukaryotic expression vectors results in the production of equimolar amounts of the 360 kD pro-vWF and of the 260 kD

mature vWF in the cell culture supernatant. vWF is probably processed into its mature form in transfected cells, by endogenously occurring furin (Wise et al., *Proc. Natl. Acad. Sci. USA* 87:9378-9382, 1990, Van de Ven et al., *Mol. Biol. Rep.* 14:265-275, 1990).

[0043] Among additional pro-proteins which are cleaved by furin or by subtilisin-like enzymes, respectively, are a series of hormones and growth factors (e.g., proactivin A, hepatocyte-growth factor), plasma proteins (albumin, factor VII, factor IX, factor X), receptors (insulin pro-receptor), viral proteins (e.g. HIV-1 gp160, influenza virus haemagglutinin) as well as bacterial proteins (diphtheria toxin, anthrax toxin) (Decroly et al., *J. Biol. Chem.* 269:12240-12247, 1994; Stieneke-Grober et al., *EMBO J.* 11:2407-2414, 1992; Barr, *Cell* 66:1-3, 1991, Wasley et al., *J. Biol. Chem.* 268:8458-8465, 1993; Klimpel et al., *Proc. Natl. Acad. Sci. USA* 89:10277-10281, 1992; Tsuneoka et al., *J. Biol. Chem.* 268:26461-26465, 1993; Bresnahan et al., *J. Cell. Biol.* 111:2851-2859, 1990; Hosaka et al., *J. Biol. Chem.* 266:12127-12130, 1991; and Vey et al., *J. Cell. Biol.* 127:1829-1842, 1994). The rFurin of the present invention is contemplated for use in cleaving these pro-proteins as well.

[0044] By co-expression of the nucleic acid sequences encoding intact furin and a pro-protein in eukaryotic cell cultures, an increased processing of the pro-proteins has been achieved *in vivo*. This has been demonstrated, e.g., for pro-factor IX (Wasley et al., *J. Biol. Chem.* 268:8458-8465, 1993) and for pro-vWF (WO 91/06314; Van de Ven et al., *Mol. Bio. Rep.* 14:265-275, 1990; and Rehemtulla et al., *Blood* 79:2349-2355, 1992). The present invention contemplates that the rFurin of the invention is useful for both *in vitro* and *in vivo* processing of any pro-protein that it cleaves.

[0045] Beside the co-expression of intact furin with pro-proteins, truncated furin has been expressed together with pro-proteins. Furin deletion mutants have been demonstrated as enzymatically active when co-expressed *in vivo* and as secreted; the enzymatic activity of such deletion mutants could be detected *inter alia* in the processing of pro-factor IX (Wasley et al., *J. Biol. Chem.* 268:8458-8465, 1993) and pro-vWF (Rehemtulla et al., *Blood* 79: 2349-2355, 1992). Co-expression experiments with furin deletion mutants have shown that the transmembrane and the cytoplasmic parts of the protein are not essential to the catalytic function (Rehemtulla et al., *Proc. Natl. Acad. Sci. USA* 89: 8235-8239, 1992).

[0046] WO 91/06314 discloses the recombinant expression of furin in prokaryotic and eukaryotic cells, the preparation of furin fusion proteins, deletion mutants and fragments, the purification of recombinantly prepared furin, and the potential use of purified furin for the processing of proproteins *in vitro* in general. WO 92/09698 describes the expression of PACE (furin), the co-expression with inactive precursors of proteins, such as, e.g., pro-vWF, as well as the preparation of fusion proteins. Stieneke-Grober et al. (*EMBO J.* 11:2407-2414, 1992) describe the *in vitro* cleavage of influenza virus HA protein by means of purified furin. Decroly et al. (*J. Biol. Chem.* 269:12240-12247, 1994) describe the *in vitro* cleavage of HIV gp160 by means of furin.

[0047] In experiments with C-terminally shortened furin, the cleavage of pro-albumin and complement Pro-C3 (Oda et al., *Biochem. Biophys. Res. Commun.* 189:1353-1361, 1992), anthrax toxin (Klimpel et al., *Proc. Natl. Acad. Sci. USA* 89:10277-10281, 1992), diphtheria toxin (Tsuneoka et al., *J. Biol. Chem.* 268: 26461-26465, 1993) and pro-factor IX (Wasley et al., *J. Biol. Chem.* 268:8458-8468, 1993, Bristol et al., *Biochemistry* 33:14136-14143, 1994) has been carried out successfully *in vitro*.

[0048] The rFurin of the present invention, therefore, is contemplated for use in the *in vivo* and *in vitro* processing of pro-proteins as described above. In one aspect, the rFurin of the invention is especially useful in the *in vitro* processing of pro-VWF and pro-factor IX. However, its use is not to be construed as limited to the processing of said proteins. In a further aspect the rFurin of the invention is particularly useful in the *in vitro* processing of recombinant pro-proteins.

[0049] A further aspect of the present invention is the co-culturing of cells which express pro-vWF and rFurin. Thus, pro-vWF in the cell culture supernatant is cleaved *in vitro* into its active form by rFurin which is also present in the cell culture supernatant. Processed vWF is subsequently isolated from the culture and purified, as discussed in U.S. Patent No. 6,210,929, incorporated herein by reference. For co-culturing, all the common expression systems can be used, and various systems for expressing pro-vWF and rFurin may be combined with each other. In one aspect, an expression system is used in which both pro-vWF and rFurin are expressed in host cells of the same origin.

[0050] The term "host cell" is used to refer to a cell which has been transformed, or is capable of being transformed with a nucleic acid sequence and then of expressing a selected gene of interest. The term includes the progeny of the parent cell, whether or not the progeny is identical in morphology or in genetic make-up to the original parent, so long as the selected gene is present.

[0051] The invention includes any host cells or hosts known in the art for recombinant protein production. Therefore, the cells in the present invention can be derived from any source. In one aspect, the invention includes eukaryotic and prokaryotic host cells. In another aspect, the invention includes plant cells, animal cells, fish cells, amphibian cells, avian cells, insect cells, and yeast cells. In one aspect, exemplary yeast cells include *Pichia*, e.g. *P. pastoris*, and *Saccharomyces* e.g. *S. cerevisiae*, as well as *Schizosaccharomyces pombe*, *Kluyveromyces*, *K. Zactis*, *K. fragilis*, *K. bulgaricus*, *K. wickerhamii*, *K. waltii*, *K. drosophilarum*, *K. thermotolerans*, and *K. marxianus*; *K. yarrowia*; *Trichoderma reesia*, *Neurospora crassa*, *Schwanniomyces*, *Schwanniomyces occidentalis*, *Neurospora*, *Penicillium*, *Totipocladium*, *Aspergillus*, *A. nidulans*, *A. niger*, *Hansenula*, *Candida*, *Kloeckera*, *Torulopsis*, and *Rhodotorula*. Exemplary insect cells include *Autographa californica* and *Spodoptera frugiperda*, and *Drosophila*.

[0052] In a further aspect, the host cells are mammalian cells, including primary epithelial cells (e.g., keratinocytes, cervical epithelial cells, bronchial epithelial cells, tracheal epithelial cells, kidney epithelial cells and retinal epithelial cells) and established cell lines and their strains (e.g., 293 embryonic kidney cells, BHK cells, HeLa cervical epithelial cells and PER-C6 retinal cells, MDBK (NBL-1) cells, 911 cells, CRFK cells, MDCK cells, CHO cells, BeWo cells, Chang cells, Detroit 562 cells, HeLa 229 cells, HeLa S3 cells, Hep-2 cells, KB cells, LS 180 cells, LS 174T cells, NCI-H-548 cells, RPMI 2650 cells, SW-13 cells, T24 cells, WI-28 VA13, 2RA cells, WISH cells, BS-C-1 cells, LLC-MK₂ cells, Clone M-3 cells, 1-10 cells, RAG cells, TCMK-1 cells, Y-1 cells, LLC-PK₁ cells, PK(15) cells, GH₁ cells, GH₃ cells, L2 cells, LLC-RC 256 cells, MH₁C₁ cells, XC cells, MDOK cells, VSW cells, and TH-1, B1 cells, or derivatives thereof), fibroblast cells from any tissue or organ (including but not limited to heart, liver, kidney, colon, intestines, esophagus, stomach, neural tissue (brain, spinal cord), lung, vascular tissue (artery, vein, capillary), lymphoid tissue (lymph gland, adenoid, tonsil, bone marrow, and blood), spleen, and fibroblast

and fibroblast-like cell lines (e.g., Chinese hamster ovary (CHO) cells, TRG-2 cells, IMR-33 cells, Don cells, GHK-21 cells, citrullinemia cells, Dempsey cells, Detroit 551 cells, Detroit 510 cells, Detroit 525 cells, Detroit 529 cells, Detroit 532 cells, Detroit 539 cells, Detroit 548 cells, Detroit 573 cells, HEL 299 cells, IMR-90 cells, MRC-5 cells, WI-38 cells, WI-26 cells, MiCl₁ cells, CV-1 cells, COS-1 cells, COS-3 cells, COS-7 cells, Vero cells, DBS-FrhL-2 cells, BALB/3T3 cells, F9 cells, SV-T2 cells, M-MSV-BALB/3T3 cells, K-BALB cells, BLO-11 cells, NOR-10 cells, C.sub.3H/IOTI/2 cells, HSDM₁C₃ cells, KLN205 cells, McCoy cells, Mouse L cells, Strain 2071 (Mouse L) cells, L-M strain (Mouse L) cells, L-MTK (Mouse L) cells, NCTC clones 2472 and 2555, SCC-PSA1 cells, Swiss/3T3 cells, Indian muntjac cells, SIRC cells, C.sub.II cells, and Jensen cells, or derivatives thereof).

[0053] Exemplary mammalian cells include varieties of CHO, BHK, HEK-293, NS0, YB2/3, SP2/0, and human cells such as PER-C6 or HT1080, as well as VERO, HeLa, COS, MDCK, NIH3T3, Jurkat, Saos, PC-12, HCT 116, L929, Litk-, WI38, CV1, TM4, W138, Hep G2, MMT, a leukemic cell, an embryonic stem cell or a fertilized egg cell. In one aspect of the invention, an exemplary host cell is a CHO cell. In a further aspect of the invention, the medium is used to culture CHO cells in suspension.

[0054] Host cells can be engineered to express a protein in a variety of ways known in the art, including but not limited to insertion of exogenous nucleic acid encoding the desired protein, optionally as part of an expression vector, insertion of an exogenous expression control sequence such that it causes increased expression of the host cell's endogenous gene encoding the desired protein, or activation of the host cell's endogenous expression control sequence(s) to increase expression of endogenous gene encoding the desired protein.

[0055] Cultures of host cells can be prepared according to any methods known in the art, and methods of growing such host cells and recovering recombinant protein produced by the cells, whether from the cells or culture medium, are known in the art. Such culturing methods may involve addition of chemical inducers of protein production to the culture medium. Exemplary host cells and procedures are described below.

[0056] A nucleic acid encoding a furin polypeptide is inserted into an appropriate expression vector using standard molecular biology techniques. In one aspect, the nucleic acid encodes the human furin polypeptide as set out in GenBank Accession No: EAX02111 (National Center for Biotechnology Information, U.S. National Library of Medicine, Bethesda, MD), however the worker of ordinary skill in the art will appreciate that any protein having furin biological activity, i.e., the ability to cleave pro-VWF to produce mature VWF, can be produced by the methods described herein. In a further aspect, a C-terminally truncated, fully secreted rFurin was designed by deleting nucleotides encoding amino acids 578 to 794 comprising the cystine-rich, the transmembrane, and the cytoplasmic domains. In an even further aspect, a tail of amino acids may be added to aid in purification processes. In yet another aspect, a tail of 10 histidine residues was added after amino acid 577, with or without interjacent four glycine residues serving as a flexible linker.

[0057] Expression vectors optionally may include a promoter, one or more enhancer sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader or signal sequence for polypeptide secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and/or a selectable marker element. Each of these sequences is discussed below.

[0058] Optionally, the vector may contain a "tag"-encoding sequence, i.e., an oligonucleotide sequence located at the 5' or 3' end of the furin polypeptide coding sequence; the oligonucleotide molecule encodes polyHis (such as hexaHis), or another "tag" such as FLAG, HA (hemagglutinin influenza virus) or myc for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for detection or affinity purification of the furin polypeptide from the host cell.

[0059] Suitable vectors include, but are not limited to, cosmids, plasmids, or modified viruses, but it will be appreciated that the vector system must be compatible with the selected host cell. In one aspect, the vector is a plasmid. In a further aspect, the plasmid is pUC-based cloning vector. Other vectors that can be used in the invention include expression vectors, replication vectors, probe generation vectors, sequencing vectors, and retroviral vectors. Vectors contemplated by the

invention include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, phagemid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transfected with virus expression vectors (e.g., Cauliflower Mosaic Virus, CaMV; Tobacco Mosaic Virus, TMV) or transformed with bacterial expression vectors (e.g., Ti or pBR322 plasmid); or even animal cell systems.

[0060] Mammalian expression vectors typically comprise an origin of replication, a suitable promoter, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, the SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide the required expression control elements. Exemplary eukaryotic vectors include pcDNA3, pWLneo, pSV2cat, pOG44, PXTI, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL, and pVITRO3.

[0061] Nucleic acid can be transferred into host cells by any means known in the art, e.g. through liposome-mediated transfer, receptor-mediated transfer (ligand-DNA complex), electroporation, microinjection of DNA, cell fusion, DEAE-dextran, calcium chloride, calcium phosphate precipitation, microparticle bombardment, infection with viral vectors, lipofection, transfection, or homologous recombination.

[0062] The term "transformed" or "transfected" as used herein refers to a host cell modified to contain an exogenous polynucleotide, which can be integrated into the chromosome of the host cell or maintained as an episomal element. It is contemplated that in certain aspects of the methods provided, the host cell is transfected in a "transfection step." The method may comprise multiple transfection steps. In addition, other methods known in the art for introducing exogenous polynucleotides into a host cell, including for example, electroporation and cell fusion which are not technically "transformation" are within the definition of the term "transformation" for purposes of this description.

[0063] The invention also provides methods for culturing, i.e. growing, host cells under conditions that result in rFurin protein expression. Such methods include the step of recovering the rFurin produced by the host cells from the culture medium. In

an exemplary aspect, the host cells are grown in a chemically defined, serum-free medium. Because serum is a biochemically undefined material, contains many components which have not been fully identified, differs from lot to lot, and is frequently contaminated with microorganisms, such as viruses and mycoplasma, the presence of serum in the recombinant production of rFurin is undesirable. Furthermore, the presence of animal proteins in serum in the culture media can require lengthy purification procedures.

[0064] The invention therefore provides a biochemically defined culture medium, essentially free from animal protein, for culturing cells recombinantly transfected with a human furin gene. The components of the medium are mostly inorganic, synthetic or recombinant and as such are not obtained directly from any animal source.

[0065] The cell culture medium of the present invention may comprise one or more replacement compounds and can comprise one or more replacement compounds which can be metal binding compounds and/or can comprise one or more complexes comprising one or more replacement compounds. In some embodiments, the medium can comprise one or more complexes, said complex comprising one or more transition elements or salts or ions thereof complexed one or more replacement compounds which can be metal-binding compounds. In some embodiments, the medium is capable of supporting the culture of cells *in vitro* and permits transfection of cells cultured therein.

[0066] According to one aspect of the invention, a transition element is preferably selected from the group consisting of scandium, titanium, vanadium, chromium, manganese, iron, cobalt, nickel, copper, zinc, yttrium, zirconium, niobium, molybdenum, technetium, rubidium, rhodium, palladium, silver, cadmium, lanthanum, hafnium, tantalum, tungsten, rhenium, osmium, iridium, platinum, gold, mercury, and actinium, or salts or ions thereof, and is preferably an iron salt. Suitable iron salts include, but are not limited to, FeCl_3 , $\text{Fe}(\text{NO}_3)_3$ or FeSO_4 or other compounds that contain Fe^{+++} or Fe^{++} ions.

[0067] Metal binding compounds in the medium include any macromolecules which can interact with or bind with transition elements and facilitate their uptake by cells. Such interaction/binding can be covalent or non-covalent in nature. The metal-binding compound used in this aspect of the invention is preferably selected

from the group consisting of a polyol, a hydroxypyridine derivative, 1,3,5-N,N',N''-tris(2,3-dihydroxybenzoyl)amino-methylbenzene, ethylenediamine-N,N'-tetramethylenephosphonic acid, trisuccin, an acidic saccharide (e.g., ferrous gluconate), a glycosaminoglycan, diethylenetriaminepentaacetic acid, nicotinic acid-N-oxide, 2-hydroxy-nicotinic acid, mono-, bis-, or tris-substituted 2,2'-bipyridine, a hydroxamate derivative (e.g. acetohydroxamic acid), an amino acid derivative, deferoxamine, ferrioxamine, iron basic porphine and derivatives thereof, DOTA-lysine, a texaphyrin, a sapphyrin, a polyaminocarboxylic acid, an .alpha.-hydroxycarboxylic acid, a polyethylenecarbamate, ethyl maltol, 3-hydroxy-2-pyridine, and IRC011. In one aspect, the metal-binding compound is a polyol such as sorbitol or dextran, and particularly sorbitol. In a related aspect, the metal-binding compound is a hydroxypyridine derivative, such as 2-hydroxypyridine-N-oxide, 3-hydroxy-4-pyrrone, 3-hydroxypyrid-2-one, 3-hydroxypyrid-2-one, 3-hydroxypyrid-4-one, 1-hydroxypyrid-2-one, 1,2-dimethyl-3-hydroxypyrid-4-one, 1-methyl-3-hydroxypyrid-2-one, 3-hydroxy-2(1H)-pyridinone, ethyl maltol or pyridoxal isonicotinyl hydrazone. The metal binding compounds of the present invention can also bind divalent cations such as Ca^{++} and Mg^{++} .

[0068] The culture medium of the present invention may comprise one or more ingredients selected from the group consisting of adenine, ethanolamine, D-glucose, heparin, a buffering agent, hydrocortisone, insulin, linoleic acid, lipoic acid, phenol red, phosphoethanolamine, putrescine, sodium pyruvate, tri-iodothyronine, thymidine, L-alanine, L-arginine, L-asparagine, L-aspartic acid, L-cysteine, L-glutamic acid, L-glutamine, glycine, L-histidine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, L-valine, N-acetyl-cysteine, biotin, choline chloride, D- Ca^{++} -pantothenate, folic acid, i-inositol, niacinamide, pyridoxine, riboflavin, thiamine, vitamin B₁₂, Pluronic F68, recombinant insulin, a calcium salt, CuSO_4 , FeSO_4 , FeCl_3 , $\text{Fe}(\text{NO}_3)_3$, KCl, a magnesium salt, a manganese salt, sodium acetate, NaCl, NaHCO_3 , Na_2HPO_4 , Na_2SO_4 , a selenium salt, a silicon salt, a molybdenum salt, a vanadium salt, a nickel salt, a tin salt, ZnCl_2 , ZnSO_4 or other zinc salts, wherein each ingredient is added in an amount which supports *in vitro* cell culture.

[0069] In another aspect, the culture medium of the invention may optionally further comprise one or more supplements selected from the group consisting of one

or more cytokines, soy peptone, one or more yeast peptides and one or more plant peptides (most preferably one or more of rice, aloe vera, soy, maize, wheat, pea, squash, spinach, carrot, potato, sweet potato, tapioca, avocado, barley, coconut and/or green bean, and/or one or more other plants), e.g., see international application no. PCT/US97/18255, published as WO 98/15614.

[0070] The culture medium of the present invention may also optionally include one or more buffering agents to maintain an optimal pH. Suitable buffering agents include, but are not limited to, N-[2-hydroxyethyl]-piperazine-N'-[2-ethanesulfonic acid] (HEPES), MOPS, MES, phosphate, bicarbonate and other buffering agents suitable for use in cell culture applications. A suitable buffering agent is one that provides buffering capacity without substantial cytotoxicity to the cells cultured. The selection of suitable buffering agents is within the ambit of ordinary skill in the art of cell culture.

[0071] The above-described media components when admixed together in solution form a complete culture medium of the present invention. A complete medium is suitable for use in the culture of a variety of mammalian cells, as described in more detail herein. Based on the information obtained herein, and knowledge possessed by those of ordinary skill in the art, one of ordinary skill in the art can obtain operative media formulations without undue experimentation.

[0072] Initially and prior to adaptation for growth in a chemically defined serum-free medium, host cells may be grown in standard media well known to one of ordinary skill in the art. The media will usually contain all nutrients necessary for the growth and survival of the cells. Suitable media for culturing eukaryotic cells are, Roswell Park Memorial Institute (RPMI) medium 1640 (RPMI 1640), Minimal Essential Medium (MEM), and/or Dulbecco's Modified Eagle Medium (DMEM), DMEM/F12, and ExCell 325 medium, all of which may be supplemented with serum and/or growth factors as indicated by the particular cell line being cultured. Notably, however, the invention provides that the serum in the media is then removed from the culture to obtain host cells that can grow in serum-free medium. Thus, the invention provides optimal media for culturing host cells under serum-free conditions for maximal production of rFurin. In a further aspect, the cells are grown in serum-free medium in suspension culture. Recipes for the various media of the invention are provided in the Examples herein.

[0073] In one aspect, an antibiotic or other compound useful for selective growth of transformed cells is added as a supplement to the media. The compound to be used will be dictated by the selectable marker element present on the plasmid with which the host cell was transformed. Selectable markers that confer resistance to particular drugs that are ordinarily toxic to an animal cell can be used in the methods and compositions of the invention. For example, where the selectable marker element is kanamycin resistance, the compound added to the culture medium will be kanamycin. Other compounds for selective growth include ampicillin, tetracycline, geneticin, neomycin, zeomycin (zeo); puromycin (PAC); Blasticidin S (BlaS), and GPT. Additional selectable markers are known in the art and useful in the compositions and methods of the invention.

[0074] Metabolic enzymes that confer cell survival or induce cell death under prescribed conditions can also be used in the methods and compositions of the inventions. Examples include, but are not limited to: dihydrofolate reductase (DHFR); herpes simplex virus thymidine kinase (TK), hypoxanthine-guanine phosphoribosyltransferase (HGPRT), and adenine phosphoribosyltransferase (APRT), which are genes which can be employed in cells lacking TK, HGPRT or APRT, respectively. The worker of ordinary skill in the art will appreciate, however, that a rFurin product of the invention will be essentially free of these added proteins.

[0075] The medium can be used to culture any host cells or hosts known in the art for recombinant protein production. In one aspect of the invention, an exemplary host cell is a CHO cell. In a further aspect of the invention, the medium is used to culture CHO cells in suspension.

[0076] When the recombinant protein of interest is secreted into the medium by the host cells, the medium can be harvested periodically, so that the same host cells can be used through several harvest cycles. Culture medium may be added in a batch process, e.g. where culture medium is added once to the cells in a single batch, or in a fed batch process in which small batches of culture medium are periodically added. Medium can be harvested at the end of culture or several times during culture. Continuously perfused production processes are also known in the art, and involve continuous feeding of fresh medium into the culture, while the same volume is continuously withdrawn from the reactor. Perfused cultures generally achieve higher cell densities than batch cultures and can be maintained for weeks or

months with repeated harvests. Thus, chemostat cultures and batch reseeded cultures are both suitable for the manufacturing of rFurin, as are other culture methods known in the art.

[0077] A variety of culture systems are known in the art, including T-flasks, spinner and shaker flasks, roller bottles and stirred-tank bioreactors. Roller bottle cultivation is generally carried out by seeding cells into roller bottles that are partially filled (e.g., to 10-30% of capacity) with medium and slowly rotated, allowing cells to attach to the sides of the bottles and grow to confluency. The cell medium is harvested by decanting the supernatant, which is replaced with fresh medium. Anchorage-dependent cells can also be cultivated on microcarrier, e.g. polymeric spheres, that are maintained in suspension in stirred-tank bioreactors. Alternatively, cells can be grown in single-cell suspension.

[0078] The amount of rFurin produced by a host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis, SDS-polyacrylamide gel electrophoresis, non-denaturing gel electrophoresis, High Performance Liquid Chromatography (HPLC) separation, immunoprecipitation, ELISA, and/or activity assays such as DNA binding gel shift assays. The invention also contemplates that specific productivity (expressed as amount of protein/cell/day) of rFurin can be evaluated using standard methods as known in the art and as described herein.

[0079] "Substantially animal protein-free rFurin" is defined as encompassing preparations of rFurin comprising host cell protein in a concentration which ranges from between about 0.1 to 0.6 ng protein or less/Unit furin activity or between about 2 and 11 μ g protein or less/mL and essentially lacking contaminating proteins from serum in the culture medium. In one aspect, the substantially animal protein-free rFurin encompasses preparations of rFurin comprising host cell DNA in a concentration which ranges from between about 0 to 0.4 pg DNA or less/Unit furin activity or between about 0 and 24 ng DNA or less/mL and essentially lacking contaminating proteins from serum in the culture medium. In one aspect, host cells expressing rFurin are grown in a chemically-defined, serum-free medium. Alternatively, the cells may be grown in medium with serum and purified according to methods provided herein.

[0080] Host cells expressing rFurin are cultured in suspension in a medium free of animal (including human) derived substances under chemostat conditions. The cells are removed by filtration and the rFurin containing cell culture supernatant is concentrated by ultrafiltration and purified by ion exchange chromatography to result in a solution of rFurin with an activity of at least about 1000 Units/ml, of at least about 2000 Units/ml, of at least about 3000 Units/ml, of at least about 4000 Units/ml, of at least about 5000 Units/ml, of at least about 6000 Units/ml, of at least about 7000 Units/ml, of at least about 8000 Units/ml, of at least about 9000 Units/ml, of at least about 10000 Units/ml, of at least about 15000 Units/ml, of at least about 20000 Units/ml, of at least about 25000 Units/ml, of at least about 30000 Units/ml, of at least about 35000 Units/ml, of at least about 40000 Units/ml, of at least about 45000 Units/ml, of at least about 50000 Units/ml, of at least about 55000 Units/ml, of at least about 60000 Units/ml, of at least about 65000 Units/ml, of at least about 70000 Units/ml, of at least about 75000 Units/ml, of at least about 80000 Units/ml, of at least about 85000 Units/ml, of at least about 90000 Units/ml, of at least about 95000 Units/ml, of at least about 100000 Units/ml, of at least about 120000 Units/ml, of at least about 140000 Units/ml, of at least about 160000 Units/ml, of at least about 180000 Units/ml, of at least about 200000 Units/ml, and of at least about 500000 Units/ml, and up to more than 500000 U/ml.

[0081] In another aspect the purified solution of recombinant furin of the invention has a specific activity of at least about 10 U/ μ g protein, at least about 20 U/ μ g protein, at least about 30 U/ μ g protein, at least about 40 U/ μ g protein, at least about 50 U/ μ g protein, at least about 60 U/ μ g protein, at least about 70 U/ μ g protein, at least about 80 U/ μ g protein, at least about 90 U/ μ g protein, at least about 100 U/ μ g protein, at least about 120 U/ μ g protein, at least about 140 U/ μ g protein, at least about 160 U/ μ g protein, at least about 180 U/ μ g protein, at least about 200 U/ μ g protein, at least about 250 U/ μ g protein, at least about 300 U/ μ g protein, at least about 350 U/ μ g protein, at least about 400 U/ μ g protein, at least about 450 U/ μ g protein, at least about 500 U/ μ g protein, at least about 550 U/ μ g protein, at least about 600 U/ μ g protein, at least about 650 U/ μ g protein, at least about 700 U/ μ g protein, at least about 750 U/ μ g protein, at least about 800 U/ μ g protein, at least about 850 U/ μ g protein, at least about 900 U/ μ g protein, at least about 950 U/ μ g protein, and at least about 1000 U/ μ g protein.

[0082] In another embodiment, the purified solution of rFurin in the invention contains host cell protein at a concentration of less than about 20.0 $\mu\text{g/ml}$, less than about 19.0 $\mu\text{g/ml}$, less than about 18.0 $\mu\text{g/ml}$, less than about 17.0 $\mu\text{g/ml}$, less than about 16.0 $\mu\text{g/ml}$, less than about 15.0 $\mu\text{g/ml}$, less than about 14.0 $\mu\text{g/ml}$, less than about 13.0 $\mu\text{g/ml}$, less than about 12.0 $\mu\text{g/ml}$, less than about 11.0 $\mu\text{g/ml}$, less than about 10.5 $\mu\text{g/ml}$, less than about 10.0 $\mu\text{g/ml}$, less than about 9.5 $\mu\text{g/ml}$, less than about 9.0 $\mu\text{g/ml}$, less than about 8.5 $\mu\text{g/ml}$, less than about 8.0 $\mu\text{g/ml}$, less than about 7.5 $\mu\text{g/ml}$, less than about 7.0 $\mu\text{g/ml}$, less than about 6.5 $\mu\text{g/ml}$, less than about 6.0 $\mu\text{g/ml}$, less than about 5.5 $\mu\text{g/ml}$, less than about 5.0 $\mu\text{g/ml}$, less than about 4.5 $\mu\text{g/ml}$, less than about 4.0 $\mu\text{g/ml}$, less than about 4.0 $\mu\text{g/ml}$, less than about 3.5 $\mu\text{g/ml}$, less than about 3.0 $\mu\text{g/ml}$, less than about 2.5 $\mu\text{g/ml}$, less than about 2.0 $\mu\text{g/ml}$, less than about 1.5 $\mu\text{g/ml}$, less than about 1.0 $\mu\text{g/ml}$, less than about 0.5 $\mu\text{g/ml}$, less than about 0.4 $\mu\text{g/ml}$, less than about 0.3 $\mu\text{g/ml}$, less than about 0.2 $\mu\text{g/ml}$, less than about 0.1 $\mu\text{g/ml}$, and about 0 $\mu\text{g/ml}$.

[0083] In another aspect, the purified solution of rFurin in the invention contains host cell protein at a concentration of less than about 1.0 ng protein/U rFurin, less than about 0.95 ng protein/U rFurin, less than about 0.90 ng protein/U rFurin, less than about 0.85 ng protein/U rFurin, less than about 0.80 ng protein/U rFurin, less than about 0.75 ng protein/U rFurin, less than about 0.70 ng protein/U rFurin, less than about 0.65 ng protein/U rFurin, less than about 0.60 ng protein/U rFurin, less than about 0.55 ng protein/U rFurin, less than about 0.50 ng protein/U rFurin, less than about 0.45 ng protein/U rFurin, less than about 0.40 ng protein/U rFurin, less than about 0.35 ng protein/U rFurin, less than about 0.30 ng protein/U rFurin, less than about 0.25 ng protein/U rFurin, less than about 0.20 ng protein/U rFurin, less than about 0.15 ng protein/U rFurin, less than about 0.10 ng protein/U rFurin, less than about 0.05 ng protein/U rFurin, less than about 0.04 ng protein/U rFurin, less than about 0.03 ng protein/U rFurin, less than about 0.02 ng protein/U rFurin, less than about 0.01 ng protein/U rFurin, and about 0 ng protein/U rFurin.

[0084] The examples herein below demonstrate the invention using CHO host cells to produce rFurin, however, the worker of ordinary skill will realize that any host cell type can be similarly adapted for producing the rFurin of the invention. CHO cells have been widely used in the production of recombinant proteins, and

engineered CHO cells (those in which a CHO cell line is transfected with a product gene and a selectable marker gene) are routinely grown in culture medium containing serum. However, the use of serum poses a number of problems. Serum is an expensive commodity, which is not readily available in amounts required for commercial production. Serum is also a biochemically undefined material and contains many components which have not been fully identified nor their actions determined. Thus serum will differ from batch to batch, possibly requiring testing to determine levels of the various components and their effect on the cells.

[0085] In addition, serum is frequently contaminated with microorganisms such as viruses and mycoplasma many of which may be harmless, but still represent an additional unknown factor. Furthermore, the presence of animal proteins in culture media can require lengthy purification procedures. In particular, the presence of bovine antibodies in bovine serum albumin (BSA) makes purification of the desired antibodies expressed by the recombinant CHO cell line extremely difficult. Removal of bovine antibody from medium prior to use is possible, but this removal and the additional product testing required after removal adds greatly to the cost of production of the product. Consequently, there are benefits in using a culture medium devoid of animal components which will support cellular growth, especially of CHO cells. While CHO cells do not readily grow in serum-free conditions, the present invention provides rFurin grown in CHO cells under serum-free conditions.

[0086] Engineered CHO cells are also difficult to grow in suspension. It is highly desirable to achieve growth in suspension when using the cells to express a product like rFurin. For the production of such a biological protein on a commercial scale, it is desirable to be able to support growth in fermenters of a considerable size. A suitable medium is also required to support the cells so that they may grow in large production conditions. Such suitable media are set out in the Examples herein. The worker of ordinary skill in the art will appreciate that any methods of culturing cells in the art can be used in culturing the host cells comprising rFurin as set out in the invention. Non-limiting examples of culture methods are provided in the Examples herein.

[0087] The invention also provides purification methods that are carried out after cells are grown in serum-free medium to remove CHO cell proteins from rFurin. The worker of ordinary skill in the art will appreciate that any methods of protein

purification known in the art can be used in the purification of rFurin from the culture medium. Non-limiting examples of purification methods are provided in the Examples herein. Accordingly, rFurin which is essentially substantially free from all animal source protein can be produced. The substantially animal protein-free rFurin is optionally stored frozen until use.

EXAMPLES

[0088] Additional aspects and details of the invention will be apparent from the following examples, which are intended to be illustrative rather than limiting. Example 1 describes the construction of a rFurin expression plasmid and host cell transfection; Example 2 describes the processes of adapting the rFurin-expressing CHO cell clones to growth in serum-free conditions; Example 3 describes a process of optimization for manufacturing rFurin in animal-protein free medium; Example 4 describes the purification of rFurin; Example 5 sets out the downstream processing (concentration and purification) and analysis of the large scale production of rFurin; and Example 6 demonstrates the safety, sterility, and stability testing that is performed to determine and maintain the quality of the host cell bank.

EXAMPLE 1:

CONSTRUCTION OF A RECOMBINANT FURIN EXPRESSION PLASMID AND HOST CELL TRANSFECTION

[0089] A detailed description of the furin progenitor plasmids used to construct a rFurin expression plasmid designated #556 is set out in Table 1. Expressed under control of a constitutive cytomegalovirus (CMV) promoter, the mature rFurin contains the catalytic domain, the P domain, and a small portion of the cystine-rich domain whereas regions located C-terminal to amino acid 577 are removed leading to a fully secreted active protease.

[0090] A description of the construction of the DHFR-vector used as the selection plasmid is depicted in Table 2. For the development of stably expressing CHO/rFurin cell clones designated # 488-3 and # 289-20, CHO cells lacking a functional endogenous DHFR gene were co-transfected with plasmids # 556 and # 73 employing calcium phosphate co-precipitation. Clones secreting high levels of

rFurin were selected in several rounds of subcloning and amplification using the DHFR/MTX selection system.

Table 1. Description of furin plasmid generation

Plasmid	Description	Comments
# 177	Original plasmid obtained from Wim J.M. van de Ven (University of Leuven, Belgium). A pUC18-based plasmid containing the 4.0 kbp <i>EcoRI</i> fragment of the human furin cDNA (2.385 kbp) with additional sequences of the 5'- and 3'-untranslated regions (UTR).	Carried out in the University of Leuven, Belgium.
# 180	Human full-length furin expression vector.	The 2.8 kbp <i>SmaI</i> / <i>AvrII</i> fragment of plasmid # 177, comprising the complete furin cDNA (2.385 kbp) and in addition ~50 bp of the 5'- UTR and ~400 bp of the 3'-UTR, was cloned into the expression vector # 55.
# 55	Eukaryotic expression plasmid from Clontech (Palo Alto, CA, USA) which has been modified to contain a multiple cloning site instead of the β -galactosidase cDNA. The plasmid provides a human cytomegalovirus immediate early (CMV IE) gene promoter and enhancer, the RNA splicing signals from the SV40 genome consisting of the late viral protein gene 16s/19s splice donor and acceptor sequences, and the SV40 polyadenylation signal. The original vector pCMV β is a pUC19 derivative containing the <i>E. coli</i> ~ 3.4 kbp β -galactosidase cDNA inserted into the <i>NotI</i> site.	The β -galactosidase <i>NotI</i> -cassette was removed and a multiple cloning site was inserted instead having amongst other restriction sites also the unique sites for <i>SmaI</i> and <i>AvrII</i> .
# 229	Human furin lacking the C-terminal sequences from position 578 to 794 spanning the cysteine-rich, the trans-membrane and the cytoplasmic regions. After glycine 577, 4 additional glycines as 'spacer' and 10 histidine residues were introduced.	Between the <i>SmaI</i> and <i>AvrII</i> sites, an appropriate reannealed oligodesoxynucleotide-linker was inserted coding for 4 glycines, 10 histidines followed by a stop codon.
# 378	Derivative of plasmid # 229.	The 12 bps coding for 4

Plasmid	Description	Comments
	Truncated furin after glycine 577 containing 10 histidines but without 4 glycines.	glycines located on a <i>SauI</i> / <i>Hind III</i> fragment were removed by PCR. The modified fragment was then religated into the plasmid-backbone.
# 556	Derivative of plasmid # 378. Truncated furin after glycine 577 which is devoid of any additional heterologous sequences.	Deletion of 30 bps coding for the 10 histidine residues by PCR. The modified <i>SauI</i> / <i>Hind III</i> fragment was religated into the plasmid-backbone.

Table 2. Description of DHFR plasmid generation

Plasmid	Description	Comments
# 29	The original plasmid named pAdd26SV(A)-3 was obtained from H.J. Hauser (GBF, Braunschweig, Germany). This plasmid contains the full length murine dihydrofolate reductase (DHFR) cDNA behind an adenovirus major late promoter.	Constructed outside Baxter.
# 73	Murine DHFR-cDNA under control of the SV40 early promoter.	The <i>PstI</i> fragment comprising the DHFR cDNA and the SV40 polyadenylation signal of plasmid # 29 was cloned into the <i>PstI</i> site of plasmid # 53. Due to the cloning strategy, plasmid # 73 contains two polyadenylation signals.
# 53	Eukaryotic expression vector from Clontech (Palo Alto, CA, USA) which has been modified to contain a multiple cloning site instead of the β -galactosidase cDNA. The plasmid provides a simian virus 40 (SV40) early gene promoter and enhancer, the RNA splicing signals from the SV40 genome consisting of the late viral protein gene 16s/19s splice donor and acceptor sequences, and the SV40 polyadenylation signal. The original vector pSV40 β is a pUC19 derivative containing the <i>E. coli</i> ~ 3.4 kbp β -galactosidase cDNA inserted into the <i>NotI</i> site.	The β -galactosidase <i>NotI</i> -cassette was removed and a multiple cloning site was inserted instead having amongst other restriction sites also a unique site for <i>PstI</i> .

[0091] Transfected CHO cells were grown in DHFR-medium, composed of DMEM NUT MIX F12 (1:1) without hypoxanthine, thymidine and glycine supplemented with Hepes, L-glutamine, Penicillin-Streptomycin, and with 5% or 10% dialyzed, gamma-irradiated FBS (= 5% DHFR, 10% DHFR). Dialyzed and gamma-irradiated FBS was purchased from Life Technologies with full documentation of certificates of analysis, origin and irradiation. The preparation of gamma-Trypsin solution (1mg/ml) was performed at Baxter in Orth, Austria, by the department of Media Preparation/PCC.

[0092] A pedigree of the generation of the CHO/rFurin clone # 488-3 is depicted in Figure 2. Clone CHO/rFurin # 488-3 was obtained from initial clones which underwent two rounds of subcloning in 10% DHFR selection medium before entering amplification in selection medium supplemented with 100 nM MTX in which one round of subcloning in medium containing unchanged MTX concentration was performed. Clone # 448-3 was expanded for freezing. The CHO/rFurin clone # 289-20 was likewise prepared and expanded. However, clone #289-20 is a successor clone derived from clone #488-3. A pedigree of the generation of the CHO/rFurin clone # 289-20 is depicted in Figure 3.

[0093] , Furin activity was measured in the conditioned medium of clones, which were cultured for 24 hrs in serum-free DHFR medium. Cell clones which showed high furin activity (U/10⁶ cells per 24 hrs) were selected. Selected high producer clones were expanded for the preparation of freeze stock ampules, and used for splitting for the next cloning round. Isolation and identification of high producer cell clones was performed. Cell densities were analyzed using the Casy cell-counter. Furin expression levels of up to 200-300 U/10⁶ cells per 24 hrs were achieved for clone # 488-3. Furin expression levels of up to 400 U/10⁶ cells per 24 hrs were achieved for clone # 289-20.

EXAMPLE 2:

ADAPTING THE RECOMBINANT FURIN EXPRESSING CELL CLONES TO GROWTH IN SERUM-FREE CONDITIONS

[0094] The strategy for cell line adaptation and selection is to adapt the cell line to a serum- and protein-free cell line in either gradually in a step-wise dilution or abruptly. The purpose of this study was to find a CHO cell population growing under

serum-free conditions, which was stably producing rFurin. The CHO cell clone #488-3 was used as starting material. The rFurin expressing cell clone CHO #488-3 was changed over to serum-free conditions in three parallel conducted adaptations as set out in detail below.

[0095] The serum depletion process started in spinner flasks with use of microcarriers to find a means to hold back cells in the phase of adaptation, since in that phase cells usually show slow growth. By using this method, it was possible to avoid, during subsequent media changes, diluting the cells to such concentrations where growth could be inhibited.

[0096] Three variants of an in-house developed medium, BAP, BAS, and BCS, (as shown in Table 3) were used during the course of this study.

Table 3: In-house media formulations of BAP, BAS and BCS

Components	JDE Item No.	Concentration [g/kg]	BAP	BAS	BCS
DMEM/F12	0200437	11.745	X	X	X
L-Glutamine	0200444	0.600	X	X	X ¹
Phenol red sodium salt	0200425	0.008	X	X	X
Putrescine dihydrochloride	0200233	0.0036	---	---	X
Iron (II) sulfate heptahydrate	0200231	0.0006	---	---	X
Ethanolamine	0200426	0.00153	X	X	X
Synperonic F68	0200172	0.250	X	X	X ²
Soy peptone	0200171	2.50	X	---	---
Sodium bicarbonate	0301012	2.00	X	X	X
WFI (water for injection)	F124	ad 1 kg	X	X	X

¹ Concentration of L-Glutamine in BCS: 0.900 g/kg

² Concentration of Synperonic F68 in BCS: 1.00 g/kg

[0097] Depending on the purpose of the respective experiment these media variants were provided with different supplements, as listed in Table 4.

Table 4: Media and their supplements

	Put ¹ (mg/l)	Glut ² (mg/l)	Synp ³ (mg/l)	Fe ⁴ (mg/l)	Zn ⁵ (mg/l)
ExCell 325PF CHO	---	600	---	---	1.0 or 5.0

BAS	3.6	300	750	0.6	5.0
BCS	---	---	---	---	1.0 or 5.0

¹ Putrescine dihydrochloride

² L-Glutamine

³ Synperonic F68

⁴ Iron(II) sulfate heptahydrate

⁵ Zinc(II) sulfate heptahydrate

[0098] The following tables give an overview of media and reagents, which were used in the course of this study. Table 5 summarizes media and reagents which were used for the establishment of the pre-master cell bank clones PMCB#01 and the PMCB#04.

Table 5: Media and Reagents Used for the Establishment of PMCB#01 and PMCB#04

	Description	Lot Number
MEDIA	10% DHFR medium	#061005/09
	BAP + 5% FBS	M/MAB-05/009, M/MAB-05/013
	BAS+Put+Glut+Synp+Fe+5% FBS	M/MAB-06/019 ⁴
	BAS+Put+Glut+Synp+Fe	M/MAB-06/012 ¹ , M/MAB-06/017, M/MAB-06/031, M/MAB-06/036, M/MAB-06/044, M/MAB-06/048 ⁴ , M/MAB-06/058 ⁴ , M/MAB-06/061 ⁴
	BAS+Put+Glut+Synp+Fe+Zn	M/MAB-06/063, M/MAB-06/069, M/MAB-06/072, M/MAB-06/074 ⁴ , M/MAB-06/076 ⁴ , M/MAB-06/079 ⁴
	BCS+Zn	M/CLD-06/001
REAGENTS	Gamma-Trypsin Solution 1mg/ml	GT_04002_1
	N1 Buffer pH 7.3	N1_05001_2
	Dimethylsulfoxide (Baxter Mat.No.: 330000000001 / JDE 0200407)	219102
	Cytopore 2 Carrier	Cyt2_24_06_001 ¹
	Na ₂ HCO ₃ - Solution	R/CBL/05/005 ¹

¹ only applied to PMCB#01

⁴ only applied to PMCB#04

[0099] Table 6 summarizes media and reagents which were used in the course of sub-cloning and establishing of the corresponding evaluation cell banks (ECBs) of the sub-clones #488-3/CJ06-19/5F10 (5F10) and #488-3/CJ06-19/1E8 (1E8).

Table 6: Lot numbers of media and reagents used for the establishment of subclones 5F10 and 1E8

	Description	Lot Number
MEDIA	10% DHFR Anzuchtmedium	#061005/09
	BAP+5% FBS	M/MAB-05/009, M/MAB-05/013
	ExCell+5% FBS+Glut	M/MAB-05/017
	ExCell+Glut	M/MAB-05/016, M/MAB-05/022, M/MAB-06/025, M/MAB-06/035, M/MAB-06/054, M/MAB-06/057, M/MAB-06/060
	ExCell+Glut+Zn	M/MAB-06/067, M/MAB-06/071, M/MAB-06/104, M/MAB-06/142
	ExCell+Glut / BAS+Put+Glut+Synp+Fe (1:1)	M/MAB-06/062
	ExCell+Glut / BAS+Put+Glut+Synp+Fe+Zn (1:1)	M/MAB-06/070
	ExCell+Glut / BCS (1:1)	M/MAB-06/077
	ExCell+BCS+Glut+Zn	M/MAB-06/090, M/MAB-06/103, M/MAB-06/106
	BCS+Zn	M/MAB-06/105, M/MAB-06/110, M/MAB-06/116, M/MAB-06/143, M/MAB-06/129, M/MAB-06/149
REAGENTS	Gamma-Trypsin Solution 1mg/ml	GT_04002_1
	Trypsin-Inhibitor Solution 1mg/ml	TI_04001_1
	N1 Buffer pH 7.3	N1_05001_2
	Dimethylsulfoxid (Baxter Mat.No.: 33000000001 / JDE 0200407)	219102
	Na ₂ HCO ₃ - Solution	R/CBL/05005

[0100] The lot numbers of all supplements, which were added to the media, are referenced in the appropriate manufacturing protocol of the corresponding medium. Other media additives are listed in Table 7.

Table 7: Other Media Additives

Description	Manufacturer	Catalog No.	Lot No.	Provided by
Aqua Bidest	Fresenius	B230673	SBV 093	Manufacturer
Aqua Bidest	Fresenius	B230673	TDV 252	Manufacturer
Aqua Bidest	Fresenius	B230673	TKV 261	Manufacturer
Aqua Bidest	Fresenius	B230673	UCV 282	Manufacturer
Cytopore 2 Carrier	Pharmacia Biotech	17-1271-03	249933	Pilot Plant II
DMEM NUT MIX F12 ²	Gibco	041-90163M	3097428	Recombinant Cell Lines
ExCell 325PF CHO	JRH	14340	4N0597	Manufacturer
ExCell 325PF CHO	JRH	14340	5L0191	Manufacturer
ExCell 325PF CHO	SAFC Biosciences	14340C	5M0775	Manufacturer
FBS ^{1,2}	Gibco	10603-017	3092829A	Recombinant Cell Lines
FBS ¹	JRH	12303	1A0348	Manufacturer
Glutamin ²	Gibco	25030-024	3096452	Recombinant Cell Lines
Glutamine	Sigma	G-5763	117 H 00655	Manufacturer
Hepes-Puffer ²	Gibco	15630-056	3094125	Recombinant Cell Lines
Paraformaldehyde	Sigma	P-6148	043 K 0653	Manufacturer
Paraformaldehyde	Sigma	P-6148	045 K 0703	Manufacturer
Paraformaldehyde	Sigma	P-6148	118 H 0987	Manufacturer
Penic./Streptomycin ²	Gibco	15140-122	1276184	Recombinant Cell Lines
Putrescine	Sigma	P5780	22 K 2615	Manufacturer
Synperonic F68	Serva	35724	01137	Manufacturer

¹ animal derived materials: for certificates refer to appendix

² ingredients of "10% DHFR medium", only used in the beginning of experiment SF05-80

[0101] Adaptation to serum-free conditions was performed in T-flasks or in spinner flasks in conjunction with cell retention (centrifugation and the like) by weaning off the cells from fetal bovine serum (FBS).

[0102] Suspension cultures in T-flasks were incubated at $36 \pm 2^\circ\text{C}$ and 7.5 ± 1.0 % CO₂. The culture in spinner flasks was performed in Techne and Bellco spinners without carriers operating at $36 \pm 2^\circ\text{C}$ with 80 rpm and 130 rpm, respectively.

[0103] The subcloning of the CHO/rFurin cell clone #488-3, subsequent to its adaptation to serum-free medium conditions, resulted in a CHO cell clone, growing in suspension in serum-free in-house medium, stably producing rFurin in a large amount. The procedure was based on the limited dilution method. Briefly, the cell suspension was diluted so that 100 μl of the suspension contains a cell. The wells of a 96-well plate were filled with 100 μl of this suspension. In theory, each well

contains one cell for clonal development. As these single cells start to grow, clones develop. Thus, every newly generated cell can be traced back to the first original cell in the well. The clones were expanded in 24-well plates, then in T25 flasks, then in T75 flasks, and then in T175 flasks.

[0104] During culture, in process controls (IPC) were performed to monitor growth conditions and to measure rFurin expression. Cell densities during culture were measured using the Casy instrument. The Nucleo counter instrument was applied for the detection of cell nuclei after CTX extraction. Determination of cell density and viability after thawing was performed with trypan blue exclusion method using a hemacytometer. Cell density and viability was also analyzed by using an automated trypan blue exclusion method performed by the Cedex instrument.

[0105] The supernatants of the cell cultures were used to determine the amount and activity of expressed rFurin. Fluorescence activated cell sorting (FACS) analysis was used to see the ratio of producer to non-producer cells in a given cell population. Morphology and growth behavior of cells were determined by optical control. Additionally, the supernatant of the cell cultures was also examined to monitor medium conditions, such as the determination of the pH value and the residual concentration of glucose, glutamine, lactate, and ammonium. These analyses were performed by means of a NOVA instrument.

[0106] In the course of this study, two pre-master cell banks (PMCBs) PMCB#01 and PMCB#04 were produced and two cell clone lines were established as set out below. Evaluation cell banks (ECB) (see Table 15) were also generated.

Preparation and QC testing of PMCB#01

[0107] One vial of the CHO/rFurin #488-3 (ECB#01) was thawed in BAS medium containing additives (Put, Glut, Synp and Fe) and 5% FBS. The cells were cultured cells for five days in a T175. The cells were then adapted to serum-free conditions as set out below.

[0108] Cells were prepared in 150 ml growth medium (BAS with Put, Glut, Synp and Fe) containing 5% FBS, 0.2 g/l. Cytopore 2 carriers were used for the adaptation to serum-free conditions. Having inoculated the 5-day old cell suspension resulting in a starting cell density of about 2.0×10^5 cells/ml, the cells attached to the porous carriers within the first hours. Thus, the cells were kept in the

spinner flask, while the growth medium was exchanged to reduce the serum concentration in two steps, from 5% to 3.8% to 0% on day 05. During the following culture period the cells got adapted to the serum-free medium conditions. The cells detached from the surface of the carriers and continued their growth in suspension. The cell density and the viability of the suspension cells increased continuously. Every two to three days the suspension cells were split in a ratio of 1:2.

[0109] An evaluation cell bank (ECB) was then prepared. On culture day 28, cells having a viability of greater than 60% were transferred to a new experiment. The culture was grown in 200-300 ml BAS medium (with Put, Glut, Synp and Fe) in a Belco spinner without carriers. According to the determined CASY cell density, the suspension culture was split every two to three days to a starting cell density of about 2.0×10^5 cells / ml. After 16 days, when the cell culture reached a viability of greater than 80%, the evaluation cell bank (ECB) consisting of six vials of cells was produced.

[0110] One vial of the ECB was thawed in a new experiment. The cells were grown for four days in a T175 in BAS medium with Put, Glut, Synp, Fe and Zn. Cells were then transferred to a Belco spinner containing up to 600 ml BAS medium with Put, Glut, Synp, Fe and Zn. Again, every two to three days the suspension culture was split to a starting cell density of about 2.0×10^5 cells / ml. On culture day 13, 143 ml of the cell suspension were removed for the preparation of the PMCB#01, consisting of 20 vials. The cells were expanded and quality control tests were performed on PMCB#01.

Preparation and QC testing of PMCB#04

[0111] One vial of the CHO/rFurin #488-3 (ECB#01) was thawed in BAS medium containing containing Put, Glut, Synp, Fe and 5% FBS. Half of the six day old culture was transferred to a new experiment for adaptation to serum-free conditions.

[0112] Here, the adaptation to serum-free conditions was not performed step by step, but rather was carried out abruptly. A Belco spinner flask was prepared with BAS medium (with addition of Put, Glut, Synp and Fe) containing no FBS and no carriers. The cells were inoculated with a starting cell density of about 2.5×10^5 cells/ml. Due to the sudden serum-free conditions, the doubling time of the cells decreased to a rather low level. To avoid diluting the cells to such a concentration

where growth could be possibly inhibited, the medium was changed by spinning down the cell suspension. The cell pellet was resuspended in fresh growth medium. Culture splits were performed when the cell density was greater than 4.0×10^5 cells/ml. After having reached a minimum viability of about 50% at culture day 15, the cells started to recover and, from culture day 32 and on, their viability increased to between 85–90%. On culture day 61, the adapted cells were frozen as an ECB consisting of 15 vials.

[0113] One vial of the ECB was thawed in a new experiment in serum-free BAS medium comprising Put, Glut, Synp and Fe. After the culture was cultured in T175 flasks for seven days, it was transferred to a Belco Spinner, where the cells grew for two further days. Then, the addition of Zn was tested. About 100 ml of the centrifuged cell suspension were resuspended in serum-free BAS medium containing Put, Glut, Synp, Fe and Zn. The suspension was cultured in a Belco Spinner. According to the determined CASY cell density, every two to three days the suspension culture was split to a starting cell density of about 2.0×10^5 cells / ml.

[0114] For inoculum preparation and to be able to produce a large amount of a homogenous cell suspension for the generation of the PMCB#04, the cell culture was scaled up to 1000 ml in a Belco Spinner. On culture day 02, 465 ml of the cell suspension were used to produce the PMCB#04, consisting of 20 ampules. The cells were expanded and quality control tests were performed on PMCB#04.

[0115] Figure 4 sets out a comparison of the graphical distribution of the rFurin producers in the cell populations of PMCB#01 and PMCB#04. 80.74% of the cells in PMCB#04 express rFurin. 74.06% of the cells in PMCB#01 express rFurin.

[0116] CHO/rFurin #488-3 subclones CJ06-19/5F10 and CJ06-19/1E8 were then generated. An ampule of the CHO/rFurin clone #488-3 was thawed and the culture was passaged in a T175 flask. On culture day 17, three different media were tested, BAP medium (developed by Baxter), CD-CHO (provided by Gibco) and ExCell 325PF CHO (provided by JRH), each of them contained 5% FBS. After four days of growth in T175 flasks, cells grown in ExCell medium were adapted to serum-free conditions.

[0117] The cells were weaned off from serum in small steps. The whole procedure ranged over three experiments. First, the anchorage-dependent cells were initially

cultured in T175 flasks in ExCell 325PF CHO containing 5% FBS. The serum was then slowly reduced to 0.5% on culture day 13. During the next 13 culture days two splits were performed, wherein the serum concentration was further reduced to 0.25% and the viability decreased to lesser than 70%. The cells lost their anchorage-dependent behavior, showed more and more spherical shape, and started to grow in suspension.

[0118] The last step of serum reduction occurred in a Techne Spinner in ExCell 325PF CHO medium containing 0.25% FBS and no carriers. After a culture period of 23 days, the serum concentration reached 0%. Cells were split and cultured to keep all relevant parameters in their given range. The cell density moved between 0.15 to 0.9×10^6 cells/ml. The viability dropped to less than 40% during the first week, but then returned to values of greater than 90%. On culture day 42, the adapted cells were frozen in the ECB / CJ06-20 consisting of 20 vials.

[0119] The cells were then subcloned. A cell suspension was diluted with preconditioned ExCell 325PF CHO in such a way, that 100 μ l of the suspension theoretically contained 0.5-1.0 cells. In the subcloning experiment, five 96-well plates were filled with 100 μ l of this cell suspension per well. The day after the seeding of the cells, the wells were searched for single cells under the microscope. Wells containing one cell were marked and observed further. Addition and exchange of preconditioned ExCell 325PF CHO were performed when necessary. When the cell died or in the absence of cell division during the next two weeks, the relevant well was excluded from the experiment.

[0120] Two single cells showed growth. The evolved clones, having reached an appropriate size, were transferred into a well of a 24-well plate. Here, the exchange of preconditioned ExCell 325PF CHO was also performed according to the growth and the requirements of the culture. The subclones CJ06-19/5F10 and CJ06-19/1E8 were transferred into a T25 flask on culture day 07 and day 10, respectively.

[0121] The ECBs were prepared in ExCell medium. These ECBs present the source material for further investigations concerning the two subclones, such as a re-adaptation from expensively purchased media to a more economical formulation, self-developed by Baxter. ECB of sub clone CJ06-19/5F10, CJ06-42, was expanded in ExCell 325PF CHO medium from a T25, to a T75, to a T175, and then into a

Bellco Spinner. The ECB / CJ06-63 consisting of 10 vials were frozen from a culture out of the T175 on day 21. ECB of subclone CJ06-19/1E8, CJ06-43, was also expanded in ExCell 325PF CHO medium. The culture was expanded from a T25, to a T75, to a T175, and then into a Bellco Spinner. The ECB / CJ06-64 consisting of 10 vials were frozen from a culture out of the T175 on day 18.

[0122] The ECBs (Subclones 5F10 and 1E8) were adapted to BCS medium as set out below. The subclone CJ06-19/5F10 of ECB clone CJ06-19 was thawed in experiment CJ06-66, and then by adding BCS medium to ExCell medium in an increasing volume, the cells were weaned from ExCell medium and acquired the ability to grow in BCS medium. The subclone CJ06-19/1E8 of ECB clone CJ06-19 was simultaneously adapted to BCS medium in a similar manner.

[0123] Table 8 shows all serum-free cell banks which were prepared in the course of this study.

Table 8: Summary of serum-free cell banks of rFurin expressing CHO cell clone #488-3

	Test No.:	Date	Name	Cell density / Amp	Amp.	Medium	Thawing contr.
PMCB#01	DE06-02	10.03.06	ECB / DE06-02	1.0×10^7	6	BAS+Put+Glut+Synp+Fe	ok
	CJ06-73	12.05.06	PMCB#01	1.5×10^7	20	BAS+Put+Glut+Synp+Fe+Zn	ok
PMCB#04	CJ06-51	11.04.06	ECB / CJ06-51	1.2×10^7	15	BAS+Put+Glut+Synp+Fe	ok
	SK06-65	12.05.06	PMCB#04	1.5×10^7	20	BAS+Put+Glut+Synp+Fe+Zn	ok
Sub Clones	CJ06-20	27.02.06	ECB / CJ06-20	1.1×10^7	20	ExCell + Glut	n.d.
	CJ06-63	14.04.06	Klon:CJ06-19/5F10 ECB / CJ06-63	1.0×10^7	10	ExCell + Glut	ok
	SK06-58	10.05.06	Klon:CJ06-19/5F10 ECB / SK06-58	1.0×10^7	5	ExCell+Glut / BAS+Put+Glut+Synp+Fe (1:1)	ok
	SK06-83	19.06.06	ECB / SK06-83	1.5×10^7	15	ExCell+Glut / BAS+Put+Glut+Synp+Fe+Zn (1:1)	ok
	SK06-151	03.11.06	Klon:CJ06-19/5F10 ECB / SK06-151	1.0×10^7	5	BCS+Zn	n.d.
	CJ06-64	14.04.06	Klon:CJ06-19/1E8 ECB / CJ06-64	1.0×10^7	10	ExCell + Glut	ok
	SK06-59	10.05.06	Klon:CJ06-19/1E8 ECB / SK06-59	1.0×10^7	5	ExCell+Glut / BAS+Put+Glut+Synp+Fe (1:1)	ok

SK06-84	19.06.06	ECB / SK06-84	1.5×10^7	15	ExCell+Glut / BCS (1:1)	ok
SK06-152	03.11.06	Klon:CJ06-19/1E8 ECB / SK06-152	1.0×10^7	7	BCS+Zn	n.d.

[0124] A comparison of PMCB#01 and PMCB#04 and subclones 5F10 and 1E8 is set out in Table 9.

Table 9: Comparison of PMCB#01, PMCB#04 and Subclones 5F10 and 1E8

Cell population	Experiments	Evaluated period	Average values of							
			Cell density x [$\times 10^6$ cells/ml]	Viability [%]	Furin act. [U/ml]	FACS [%]	μ [d ⁻¹]	g [h]	q _p [U/(10 ⁶ ·d)]	Q _p [U/(ml·d)]
PMCB#01	CJ06-69	29.04.-17.05.06	1.16	84.4	274.87	71.30	0.494	35.94	123.0	77.1
PMCB#04	SK06-49, SK06-63	26.04.-17.05.06	1.10	94.4	287.03	72.02	0.614	28.52	207.7	96.8
Clone 5F10	CJ06-66, SK06-52	21.04.-17.05.06	0.59	94.3	105.57	83.30	0.523	37.43	135.0	40.8
Clone 1E8	CJ06-67, SK06-53	21.04.-17.05.06	0.59	94.8	95.55	88.34	0.512	39.01	114.4	35.7

[0125] The cell populations cultured in the experiments CJ06-69 as well as SK06-49 and SK06-63 represent the precursor cultures of the cell lines for the PMCBs. These cells were grown in BAS medium (with the addition of Put, Glut, Synp, Fe and Zn), and were frozen as PMCB#01 and PMCB#04, respectively.

[0126] The subclones 5F10 and 1E8 were grown in the commercially available medium ExCell 325PF CHO provided by JRH. Because the ability of growing in medium based on the BAS formulation was preferred, PMCB#01 and PMCB#04 were chosen for further production of rFurin.

[0127] As presented in Table 10, the cells of PMCB#04 showed better growth behavior in comparison to PMCB#01. Viabilities and growth rates were greater in PMCB#04, and generation doubling times were lower. The cell-specific as well as the volumetric production ratio of rFurin was also greater throughout the evaluated period of time. In a further experiment, the data concerning the viabilities was confirmed. One ampule of each of the PMCBs was thawed into a T175 flask containing BCS medium with Zn as additive. As shown in Table 10 the viability of PMCB#04 was again greater than that of PMCB#01.

Table 10: Thawing experiments of PMCB#01 and PMCB#04

Cell population	experiment	cell density [x 10 ⁷ cells / vial]	viability [%]
PMCB#01	CJ06-77 [75]	1.05	68.7
PMCB#04	SK06-67 [76]	0.99	77.4

[0128] Thus, PMCB#04 was chosen as the source material for the future production of a rFurin Master Cell Bank (MCB) and all further working cell banks (WCBs). The PMCB#04 consisting of 20 vials was established in compliance with the current Good Tissue Practice regulations. Quality Control (QC) testing was performed in accordance to requests of the ICH-Guideline Q5D. PMCB#04 was chosen for producing rFurin in manufacturing processes.

[0129] The growth medium is free of human or animal derived substances and is self-developed. Having the cells removed by filtration, the rFurin containing supernatant is concentrated by ultrafiltration. After purification by exchange chromatography, the activity of the rFurin solution is aimed to be at least 200 Units rFurin/ml.

[0130] Testing for mycoplasma, viruses, extraneous agents, sterility, and expression efficiency was carried out. The criteria for selection are high Furin activity (e.g., Furin protein, ELISA) and homogeneity of the cell population in immunofluorescence (FACS), performed on different subclones in comparison to initial clones #488-3 and #289-20, respectively. In addition, the impurity profiles have to be compared qualitatively (e.g. by UV-peak patterns after RP HPLC or by SDS-PAGE/Coomassie techniques).

EXAMPLE 3:

OPTIMIZATION FOR MANUFACTURING RECOMBINANT FURIN IN ANIMAL PROTEIN-FREE MEDIUM

[0131] This example describes the development and optimization process for the culture of the rFurin expressing CHO clone #488-3. Specific medium optimization with regard to amino acids, glucose, and NaHCO₃ concentration was carried out, which resulted in increased cell growth rates and higher productivities of the

fermentation process. Optimization for inline controlled process parameters was carried out with the optimized medium formulation for pO₂ (10%, 20% and 50%), and a factorial experiment was carried out to determine optimum pH (range 7.1-7.3) and temperature (range 35.1°C-37.9°C), which resulted in a significant yield improvement for CHO clone #488-3 when fermentation was carried out at lower temperatures between 35°-36°C.

[0132] As possible production modes, chemostat cultures and batch reseeded cultures were compared, indicating that both process types are suitable for the manufacturing of rFurin, and give comparable yields with identical parameter settings. Specific experiments were carried out to investigate the influence of agitation types and rates in different bioreactor setups. It was shown that the specific growth rates and expression rates, cell densities and therefore volumetric productivities are strongly influenced by variations in the bioreactor setup, and that under conditions of increased agitation rates the yields could be significantly increased. Altogether, the influence of the main parameters of the rFurin upstream process are well characterized with regard to their effect, and a high yielding process could be transferred to the pilot plant for preclinical and clinical manufacturing. The details are set out below.

[0133] A subclone # 488-03 was developed in-house which was adapted to a serum- and insulin-free medium. The following experiments were carried out in a FBS-free and insulin-free medium (BACD-medium) as the basic medium formulation (see Table 11).

Table 11: Components of the basic medium (BACD-medium)

Component	Concentration [g/kg]
DMEM/F12 (1:1)	11.76
L-Glutamine	0.6
Ethanolamine	0.00153
Synperonic	0.25
Putrescine.2HCl	0.0036
FeSO ₄ .7H ₂ O	0.0006
CuSO ₄ .5H ₂ O	0.00000125
NaHCO ₃	2.0

[0134] In order to improve cell growth and to provide optimal conditions for cell propagation, an amino acid analysis of the supernatant of a chemostat culture was performed (see Table 12). As a result, three essential amino acids were added to the medium, namely methionine (10 mg/L), leucine (40 mg/L) and phenylalanine (10 mg/L). Fermentation was performed in a 1.5 L bioreactor at 37°C, pH 7.15 and a pO₂ of 20%.

Table 12: Amino acid analysis by HPLC

Amino acids	BACD_24_06_014_026	FUR_06/10_M04_K06	FUR_06/10_M04_K07
	Peak area [%]	relative peak area [%]	relative peak area [%]
Aspartic acid	100.00	24.04	9.88
Glutamic acid	100.00	57.78	24.89
Serine	100.00	14.37	15.35
Asparagine H ₂ O	100.00	15.03	19.17
Glycine	100.00	103.55	90.58
Glutamine	100.00	19.26	23.14
Histidine	100.00	45.59	45.32
Threonine	100.00	63.09	61.39
Arginine	100.00	72.22	74.25
Alanine	100.00	1513.36	1518.46
Proline	100.00	58.54	55.36
Tyrosine	100.00	46.37	47.63
Cystine	100.00	53.43	56.39
Valine	100.00	42.75	42.11
Methionine	100.00	13.23	11.51
Isoleucine	100.00	48.39	48.79
Leucine	100.00	23.58	20.63
Lysine HCl	100.00	41.85	40.65
Phenylalanine	100.00	34.99	33.00
Tryptophan %:	100	69.1	68.8

BACD_24_06_014_026...
FUR_06/10_M04_K06...

basic medium (s. materials and methods)
Fermentation lot, 6th day of culture

FUR_06/10_M04_K07...

Fermentation lot, 7th day of culture

The peak area of the respective amino acid in the HPLC-diagram of the basic medium was set to 100%. By comparison to the peak area obtained on day 6 and 7 of the chemostat culture the decrease of the respective amino acid was determined.

[0135] Due to low glutamine concentrations in the supernatant of the culture, glutamine was added to the medium (300 mg/L) to give a final concentration of glutamine at 900 mg/L. After the addition of glutamine to the medium the growth rate increased from 0.55 d⁻¹ to 0.67 d⁻¹ (see Tables 13 and 14). By the addition of these three amino acids mentioned above (namely methionine (10 mg/L), leucine (40 mg/L) and phenylalanine (10 mg/L)) to the medium, the growth rate of the cells could be increased again (0.69 d⁻¹) (see Table 15). The volumetric productivity went up to approximately 267 kU/L/d (= + 13%) and the specific productivity showed an increase of 16%. The supplementation of the medium with glutamine, methionine, leucine and phenylalanine showed a positive effect on cell growth, and volumetric and specific activity, and was therefore retained for further medium preparation.

Table 13: Fermentation data of the chemostat culture FUR_06/10-M04 from day 6 to day 9 of culture

Experiment	glc	gln	pH	D	CC	μ	Furin	P	qP
FUR_06/10-M04	[g/L]	[g/L]	NOVA	[1/d]	[1 E6/mL]	[1/d]	[IU/mL]	[kIU/L/d]	[IU/C/d]*10 ⁶
Day 6-9	1.58	0.10	7.16	0.494	2.35	0.548	306.55	151	64

Table 14: Fermentation data of the chemostat culture FUR_06/10-M04 after the supplementation of the medium with of 300 mg/L glutamine on day 12

Experiment	glc	gln	pH	D	CC	μ	Furin	P	qP
FUR_06/10-M04	[g/L]	[g/L]	NOVA	[1/d]	[1 E6/mL]	[1/d]	[IU/mL]	[kIU/L/d]	[IU/C/d]*10 ⁶
Day 15-21	1.47	0.35	7.16	0.668	2.27	0.673	352.68	236	104

Table 15: Fermentation data of the chemostat culture FUR_06/10-M04 after the addition of methionine (10 mg/L), leucine (40 mg/L) and phenylalanine (10 mg/L) on day 22 to the medium

Experiment	glc	gln	pH	D	CC	μ	Furin	P	qP
FUR_06/10-M04	[g/L]	[g/L]	NOVA	[1/d]	[1 E6/mL]	[1/d]	[IU/mL]	[kIU/L/d]	[IU/C/d]*10 ⁶

Day 26-29	1.37	0.38	7.13	0.713	2.20	0.695	373.80	267	121
-----------	------	------	------	-------	------	-------	--------	-----	-----

[0136] To check whether the supplemented amounts of the amino acids in the medium were sufficient, another amino acid analysis of a chemostat culture (10L) was performed. The culture had been supplied with 300 mg/L glutamine and the respective amounts of the three amino acids. The samples were drawn on day 7 and 15 of culture, and the culture conditions were similar to those mentioned above. The results (see Table 16 where only the data for methionine, leucine and phenylalanine are shown) confirmed that sufficient amounts of the supplemented amino acids were present in the fermentation broth.

Table 16: Relative amount of methionine, leucine and phenylalanine in a chemostat culture (FUR 06/17_F04) supplemented with glutamine (300 mg/L), methionine (10 mg/L), leucine (40 mg/L) and phenylalanine (10 mg/L)

Amino acid	Peak area [%]	relative peak area [%]	relative peak area [%]
	BACD-24-06-030-057	FUR 06/17_F04-K07	FUR 06/17_F04-K15
Methionine	100.0	47.5	61.7
Leucine	100.0	50.3	65.4
Phenylalanine	100.0	50.1	65.2

[0137] Reduction of the NaHCO_3 concentration and increase of the glucose concentration. The influence of high dissolved CO_2 concentrations in the cell culture on growth and productivity was investigated. Due to large scale production in a bioreactor, a greater CO_2 concentration in the cell culture can be expected than in 2.5–32 L bioreactors. Therefore, two fermentation runs were carried out in parallel, one run with a CO_2 concentration of approximately 7.5% and the other one with a CO_2 concentration of approximately 12%. The CO_2 concentration was adjusted by varying the CO_2 fraction in the head space flow. The CO_2 concentration in the cell culture was measured by analyzing drawn samples with the NOVA instrument. The fermentation was carried out at 37°C, at a pH of 7.15 and with a $p\text{O}_2$ of 20%.

[0138] A CO_2 -concentration of 11-12% had a negative influence on cell growth and productivity. At this CO_2 -concentration, a growth rate of 0.29 d^{-1} was reached over an interval of 12 days at a cell count of 1.1×10^6 cells/mL and a dilution rate of

0.30 d⁻¹. In the fermentation run with approx. 7.5% CO₂, a growth rate of 0.52 d⁻¹ was reached at a cell count of 1.49x10⁶ cells/mL and a dilution rate of 0.53 d⁻¹ in the same interval. At high CO₂ concentrations, the viability was reduced to 86.1%, compared to 95.9% at 7.5% CO₂. Additionally, the volumetric productivity was reduced to approximately 36% and the specific productivity to 50%. Due to the high CO₂ concentration, the specific glucose uptake rate was decreased as well (-39%). The negative influence of an increased CO₂ concentration (11-12%) on cell growth and productivity was quite obvious and, therefore, it is optimal to carry out the fermentation at 7.5% CO₂.

[0139] As set out above, experiments showed that when the concentration of CO₂ was increased to 12% in the 1,000 L bioreactor, a strong decrease in the performance of the CHO-Furin clone 488-3 can be expected. Therefore, it was decided to reduce the NaHCO₃ concentration in the medium from 2 g/L to 1.5 g/L. A lesser amount of NaHCO₃ in the medium also decreased the buffer capacity of the medium and, therefore, two fermentation runs (10L) were compared, one with 3.15 g/L glucose and 2 g/L NaHCO₃ in the medium (FUR_06/24_F01) and another one with 4.65 g/L glucose and 1.5 g/L NaHCO₃ (FUR_06/26_F04).

[0140] To simulate the conditions of a large scale bioreactor (1,000 L), the concentration of dissolved CO₂ during fermentation was adjusted to 7-8% in the fermenter with 2 g/L NaHCO₃ and to 6-7% in the fermenter with 1.5 g/L NaHCO₃ by constant CO₂ gassing in the head space. The growth rates of both cultures were similar (0.58 and 0.56 d⁻¹) and the cultures showed comparable volumetric productivities and viabilities. However, the specific glucose uptake rate was slightly higher in the culture with a lower NaHCO₃ concentration (0.83 mg/10⁶ cells/d vs. 0.67 mg/10⁶ cells/d). Therefore, a glucose concentration of 4.65 g/L was considered to be reasonable and was retained in further medium preparation.

[0141] Investigation of the Synperonic F68 concentration in the Furin medium. The regular concentration of Synperonic F68 in the cell culture medium was set at 0.25 g/L. The purpose of Synperonic F68 in the medium is to protect the cells from damage due to submerged oxygenation. Therefore, one experiment was carried out in 2 x 10L bioreactors, where an increased Synperonic F68 concentration of 1.0 g/L vs. the regular concentration of 0.25 g/L was investigated. The fermentation was carried out at 35.8°C at a pH of 7.30 and with a pO₂ of 20%.

[0142] With increasing Pluronic concentration (Synperonic F68), a slightly higher specific growth rate and cell density could be achieved. Thus, due to increased specific productivity, a proportionally greater volumetric productivity of 365 vs. 278 kU/L/d could be achieved.

[0143] Supernatants from these experiments were collected and filtered with a depth filter (Cuno Cart.Z08P4A30SP 4 discs) followed by a membrane filter (Pall Fluorodyne II KA2DFLP2). The filtered supernatants were concentrated by ultrafiltration (Sartorius 30S1463901E--SG PSU 10KD, 0.2m²) and diafiltrated against the Furin diafiltration buffer. The sterile filtered concentrate (Sartorius Sartobran P 523130748-00 0.45/0.2μ) was purified on the Capto MMC column. Results from this purification experiments revealed that the increased Synperonic F68 has no detrimental influence on quality and/or purity of the purified rFurin. No elevated Synperonic F68 concentrations could be found in the eluate of the Capto MMC column, which were <0.15 mg/mL in both eluates.

[0144] Despite these results the Synperonic F68 concentration in the rFurin medium was kept at the original 0.25g/L. However, in case of scale up problems due to submerged oxygenation, an increasing Synperonic concentration of up to 1.0 g/L could be considered, with the potential of increased yields without detrimental influence of the final Furin product.

[0145] Final Medium Composition for rFurin Production in Fermentation. Based on the medium optimization experiments, the following medium composition is shown to be optimal for the fermentation of CHO clone 488-3 in bioreactor cultures for the production of rFurin.

Table 17: Components of the basic medium and optimized fermentation medium

Component	Basic Medium Concentration [g/kg]	Optimized Medium Concentration [g/kg]
DMEM/F12 (1:1)	11.76	11.76
L-Glutamine	0.6	0.9
D-Glucose	-	1.5
Ethanolamine	0.00153	0.00153
Synperonic	0.25	0.25

Putrescine.2HCl	0.0036	0.0036
Methionine	-	0.01
Leucine	-	0.04
Phenylalanine	-	0.01
FeSO ₄ .7H ₂ O	0.0006	0.0006
CuSO ₄ .5H ₂ O	0.00000125	0.00000125
NaHCO ₃	2.0	1.5

[0146] The influence of different pO₂ set points on growth rate and productivity was also investigated. Three different pO₂ set points were tested, i.e. 10%, 20% and 50%. The experiments were performed in 1.5 L bioreactors with gassing via head space. All fermentation runs were carried out at 35.1°C at pH 7.20. Comparison of the fermentation runs at 10, 20 and 50% pO₂ showed a slight increase of the growth rates (0.59, 0.62 and 0.65 d⁻¹) with increasing pO₂ set points. Likewise, the volumetric productivity was greatest at 50% pO₂. The mean cell counts of the different fermentation runs were very similar. Thus, the increase in the volumetric productivity was the result of the increasing growth rate.

[0147] Consequently, a pO₂ set point of 50% seemed to influence the growth rate positively, and as a result the volumetric productivity was approx. 4% higher than at standard conditions (= 20% pO₂). No effect on viability was observed. Therefore, a setpoint of pO₂= 20% and a range for regulation between 10%-50% could be confirmed to be suitable for fermentation of the CHO clone #488-3.

[0148] Optimization of temperature and pH to maximize the volumetric productivity. The influence of pH and temperature on the performance of the CHO-Furin clone was investigated. By using a 'design of experiments method', different temperatures were combined with different pH values to ascertain the conditions which resulted in maximum volumetric productivity. Five temperatures were combined with three pH values according to the "Doehlert Matrix", resulting in seven combinations of temperature and pH, as shown in Figure 5.

[0149] The combination of 36.5°C and pH 7.20 was chosen as the center point, which was applied to two fermentation lots. Table 18 sets out the culture conditions used in the different fermentation lots.

Table 18: Experimental setup of the Doehlert Matrix: Culture conditions of the different fermentation lots

Fermentation lot	Temp. [°C]	pH
FUR_06/43-B07	35.1	7.20
FUR_06/43-B05	35.8	7.10
FUR_06/40-B03	35.8	7.30
FUR_06/40-B01	36.5	7.20
FUR_06/43-B02	36.5	7.20
FUR_06/40-B08	37.2	7.10
FUR_06/40-B04	37.2	7.30
FUR_06/40-B06	37.9	7.20

Table 19: Mean values of the fermentation data of the fermentations FUR_06/40-B01, -B03, -B04, -B06, -B08 and FUR_06/43-B02, -B05, -B07

Fermentation lot	Temp.	pH	CC	μ	Furin	P ¹⁾	qP ¹⁾	Viability
	[°C]		[10 ⁶ /mL]	[1/d]	[IU/mL]	[kIU/L/d]	[U/10 ⁶ /d]	[%]
FUR_06/43-B07	35.1	7.20	1.74	0.595	930	543	312	97.7
FUR_06/40-B03	35.8	7.30	1.64	0.632	621	379	232	97.6
FUR_06/43-B05	35.8	7.10	1.73	0.647	683	429	250	98.5
FUR_06/40-B01	36.5	7.20	1.78	0.696	439	309	174	97.9
FUR_06/43-B02	36.5	7.20	1.66	0.684	454	310	186	96.8
FUR_06/40-B04	37.2	7.30	1.71	0.642	284	184	108	96.8
FUR_06/40-B08	37.2	7.10	1.68	0.652	338	215	128	98.3
FUR_06/40-B06	37.9	7.20	1.52	0.473	143	65	43	96.1

¹⁾ the mean volumetric and the mean specific productivity over 5 culture days were calculated by forming the the mean value of the single productivities

[0150] The data were analyzed statistically with the Response Surface Methodology (RSM), using the "Minitab" software. RSM explores the relationships between several explanatory variables and one or more response variables. The main idea of RSM is to use a set of designed experiments to obtain an optimal response. The analysis focused on the volumetric and specific productivity as well as on the growth rate.

[0151] Analysis of the data in reference to the volumetric productivity. As a first step, a surface plot was created (Figure 6). The coordinates of the data in Figure 6 are marked as points. The surface shows the assumed correlation of the single data.

[0152] The surface plot supposes a linear correlation between the parameters temperature/pH and the responding volumetric productivity. The chart indicates an increase of the volumetric productivity with decreasing temperature. The influence of the pH is considered to be weak. Subsequent calculations showed a linear correlation of the data (calculation not shown). By variance analysis, an equation was generated which describes the correlation between pH, temperature and volumetric productivity:

$$P = 7693.1 - 162.4 \cdot \text{Temp} - 202.8 \cdot \text{pH}$$

[0153] Based on the mathematical model a contour plot was generated (Figure 7) which illustrates the influence of temperature and pH on the volumetric productivity. The dots indicate the conditions (pH / temp.) which had been tested experimentally.

[0154] The contour plot shows that the area, where maximum volumetric productivity can be expected, is at 35.1°C and a pH of approx. 7.10. Both values are at the edge of the experimental design which means that the real maximum could be found even below those values. Furthermore the Contour plot shows that the influence of the pH on the volumetric productivity is marginal and slightly higher at low pH values.

[0155] The surface plot (Figure 8) which is a three-dimensional illustration of the mathematical model, gives the same result as the contour plot: the strong influence of the temperature and the weak influence of the pH on the volumetric productivity.

[0156] Analysis of the data in reference to the growth rate μ . The correlation of the growth rate with temperature and pH is described in a quadratic equation. Again, a strong influence of the temperature and a weak influence of the pH can be found. Strong variation of the growth rate hampers the design of a mathematical model. Analysis of variance and regression gave following equation:

$$P = -103.539 + 5.706 \cdot \text{Temp} - 0.079 \cdot \text{Temp}^2 - 0.233 \cdot \text{pH} - 0.020 \cdot \text{pH}^2$$

[0157] The influence of the pH on the growth rate is statistically not ascertainable, as indicated by the high P-value for the pH-terms which was calculated to be 0.99 (calculation not shown).

[0158] The surface plot (Figure 9) illustrates the modeled correlation three-dimensionally, demonstrates the quadratic relationship, and shows a maximum growth rate at 36.5°C.

[0159] Analysis of the data shows a similar correlation of the specific productivity with temperature and pH as seen for the volumetric productivity (Figure 10). The correlation is described by a linear equation. The influence of the pH is again low as proved by variance analysis (calculation not shown):

$$qP = 4261.40 - 93.35 \cdot \text{temp} - 93.77 \cdot \text{pH}$$

[0160] In summary, the experiments for optimization of temperature and pH and subsequent analysis of the data gave a quite clear result. The greatest volumetric productivity, which is considered as the most important parameter for process optimization, was achieved by culture at the lowest temperature (35.1°C) and lowest pH (7.10). The influence of pH is statistically hardly significant, and a pH value between 7.10 and 7.30 would give very similar results. By decreasing the temperature from 37°C to 35.1°C, the volumetric productivity could be raised from approx. 200 kU/L/d to 540 kU/L/d which is 2.7 times greater (Figure 11). Based on these results, the new set points for temperature and pH, for the culture of the CHO-Furin clone in the chemostat mode, were determined as 35.1°C and 7.15.

[0161] Comparison of chemostat mode to batch refeed mode. The culture of the CHO-Furin clone in the chemostat mode at a low temperature (35.1°C) resulted in high yields in small scale experiments (1.5L, 2.5L, 10L and 32L) and was considered to be the appropriate culture method for the large scale culture for production of rFurin. However, initial fermentation runs in the large scale (1200 L bioreactor in the PP1 facility) showed a strong decrease of the growth rate at the chemostat mode. As an alternative, to get higher growth rates, the batch refeed mode was set up in the large scale. Experiments in the 10L bioreactor scale were carried out to compare growth and productivity in the chemostat mode with the batch refeed mode. The cells were cultured at 36.5°C at a pH of 7.15 and a pO_2 of 20%. The batch was split to a cell count of $0.6\text{--}0.7 \times 10^6$ cells/mL every second culture day.

[0162] The chemostat- and the batch refeed-fermentations were run in parallel, using the same cell culture as inoculum. The cells were cultured in the chemostat mode until day 6. Then one fermentation run was switched to the batch refeed mode

(FUR_06_50-F03) while the other one was continued in the chemostat mode (FUR_06_51-F04). Culture in the batch refeed mode resulted in a mean cell count of 2.22×10^6 cells/mL at the end of the batch, with a growth rate of 0.64 d^{-1} (Table 20). In the chemostat mode, a growth rate of 0.50 d^{-1} was obtained with an average cell count of 1.67×10^6 cells/mL. Due to the higher cell count and growth rate in the batch refeed mode, the volumetric productivity was greater as well (238 vs. 197 kU/L/d).

[0163] The data indicate that the batch refeed mode is a preferable culture method for the CHO-Furin clone in the 10 L scale, which results in even slightly higher volumetric productivities than in the chemostat mode (at 36.5°C and a pH of 7.15). However, in the batch refeed mode, harvest- and further downstream processes are restricted to certain intervals, while in the chemostat mode, harvesting can be performed continuously. Thus, each method has its advantages. No optimization experiments for the parameters of the batch refeed mode were performed like for the optimal cell counts at the end of a batch or for the best split ratio.

Table 20: Mean values of the fermentation data from the fermentation runs FUR_06_51-F04 (Chemostat) and Fur_06_50-F03 (Batch Refeed)

Ferment. lot	Vol.	Mode	D	μ	CC	Furin	P
			[1/d]	[1/d]	[1 E6/mL]	[IU/mL]	[kU/L/d]
FUR_06_51-F04	10 L	chemostat	0.494	0.497	1.67	401.4	198
FUR_06_50-F03	10 L	batch refeed	-	0.637	2.22 (end) ¹⁾	539.8	238 ²⁾

¹⁾ Mean value of the cell counts at the end of each batch (2 batches)

²⁾ Mean value of the productivities at the end of each batch (2 batches)

[0164] Scale Up Investigations. Comparison of Rushton type- versus ball type-impellers at different agitation rates. The influence of the agitator type on growth rate and productivity was investigated. The standard setting used in the bioreactors for experiments with the CHO-Furin clone comprised a Rushton impeller and four baffle plates. The 1,000 L bioreactor for rFurin production was equipped with three pairs of ball impellers without any baffle plates. Therefore, an experiment was carried out to test three different settings in 10L bioreactors: one reactor was equipped with

the standard setting, another reactor with two pairs of ball impellers and no baffle plates, and a third reactor was assembled like the standard one but the agitation rate was reduced from 170 rpm to 90 rpm.

[0165] Table 21 gives an overview of the bioreactor set-ups. The agitation rate of 60 rpm with ball impellers gave a similar tip speed as the rushton impeller at 90 rpm (see Table 21). However, the tip speeds cannot be equated with each other due to the different geometry of the impellers (shape, diameter).

Table 21: 10L Bioreactor Set-ups for impeller / agitation rate experiment

Fermentation lot	Impeller type / diam.	Levels	Baffle plates	Agitation rate	Tip speed
FUR_06/37_F03	Rushton 80 mm	2	4	170	0.712 m s ⁻¹
FUR_06/35_F01	Rushton 80 mm	2	4	90	0.377 m s ⁻¹
FUR_06/38_F04	Ball 140 mm, 39.5 mm diameter	2	-	60	0.440 m s ⁻¹

[0166] The fermentation conditions were as follows: 37°C, pH 7.15, pO₂ of 20% and pCO₂ of 6-7%. (Medium: 4,65 g/L Glc, 1,5 g/L NaHCO₃). Comparison of the fermentation data showed that the performance in the bioreactor with the standard set-up, i.e. equipped with the rushton impeller and agitated at a rate of 170 rpm, was higher than in the bioreactors with the other set-ups (Table 22). The use of ball impellers at 60 rpm resulted in a somewhat lower growth rate (0.57 vs. 0.61d⁻¹), a lower volumetric (197 vs. 227 kU/L/D) compared to the rushton impeller at 170 rpm. The viability was hardly affected though. Using a rushton-impeller stirred bioreactor at an agitation rate of 90 rpm instead of 170 rpm clearly diminished the growth rate (0.54 vs. 0.61d⁻¹), volumetric productivity (175 vs. 227 kU/L/D) and cell specific productivity (115 vs. 138 U/10⁶ cells/day).

[0167] The data from this experiment showed that both, a ball impeller at 60 rpm and Rushton impellers at a reduced agitation rate, result in a declined performance of the CHO-Furin clone compared to the Rushton impellers at 170 rpm. Nevertheless, the results also revealed that it is possible to use a ball impeller for culture of the CHO-Furin clone. Among the set-ups which were tested, the set-up with the rushton impeller at 170 rpm plus four baffle plates displayed the roughest conditions. Notably, however, no decrease of cell viability was observed under these

conditions, which indicates a high mechanical resistance of the investigated CHO-Furin clone.

Table 22: Mean values of the fermentation data of three fermentation runs of the CHO-Furin clone in bioreactors with different set-ups (impeller type, baffle plates)

Fermentation lot	Set-up	CC	D	Furin	μ	P	qP	Viability
		[10 ⁶ /mL]	[d ⁻¹]	[IU/mL]	[d ⁻¹]	[kU/L/d]	[U/10 ⁶ /d]	[%]
FUR_06/37_F03	Rushton 170 rpm	1.65	0.628	361.37	0.614	227	138	95.4
FUR_06/38_F04	Ball impeller 60 rpm	1.52	0.600	328.10	0.568	197	130	94.1
FUR_06/35_F01	Rushton 90 rpm	1.53	0.574	305.31	0.544	175	115	94.3

[0168] Comparison of different agitation rates with pitched blade impellers in 32 L bioreactors. For the GMP run campaign ORFURFB07002 in the PP1 the bioreactor (working volume 950L) was equipped with a pitched blade impeller type (2 pcs., d = 700mm) instead of the previously used ball type impellers.

[0169] 2 day batch reseeded cycles were carried out during this campaign. In order to assess this change compared to the previous campaigns, a 32L bioreactor was set up with a similar type of pitched blade impellers (1 piece, d=140mm), and fermentations were carried out using cell suspension from the PP1 facility and medium to ensure comparable materials to be used.

[0170] A similar batch reseeded process was carried out, where a 2-3 consecutive two day batch cycles were started with a starting cell density of 0.5x10E06 cells/mL. The experiment was carried out at 2 different agitation rates, i.e. 55rpm and 120 rpm). Culture conditions were identical with the culture conditions used in the PP1: pH-SP 7.15, Temp-SP 35.5°C, pO2-SP 20%. Data from the last batch were compared after adaptation to the respective agitation conditions.

Table 23: Growth rate, cell count and vol. productivity of the batches of run FUR_07/06-F07 with pitched blade impellers in the 32L bioreactor

FUR_07/06-F07I	Agitation rate	CC ¹⁾	Split ²⁾	Furin	μ	P ³⁾
Intervall	[rpm]	[10 ⁶ /ml]	1: x	[IU/mL]	[d ⁻¹]	[kU/L/d]
Batch K02-K04	120	1.95	3.89	1197.8	0.680	445
K10-K12	50	1.50	3.13	812.0	0.570	276

- 1) cell count at the end of each batch
- 2) theoretical split ratio to be applied in a 2 batch reseeded process: split ratio = $e^{(2 \times \mu)}$
- 3) calculation of volumetric productivity: $P = (1 - (1/\text{split ratio})) \times \text{product titer} / 2 \text{ days}$

[0171] The experiment demonstrated again the effect of agitation conditions on specific growth rates of the CHO clone #488-3. The increasing growth rates led to greater final cell densities at the end of the batch, when applying the same starting cell densities. Volumetric productivities increased from 276 kU/L/D to 445 kU/L/D (+61%), whereas the final cell density increased by 30% (1.95 vs. 1.5x10E06 cells/mL). These results indicate that the increased agitation rate had a positive impact on cell-specific productivity.

[0172] It also might be concluded that fermentation runs, which resulted in average productivities of around 200kU/L/D were mainly a result of the applied low agitation rates of 20 rpm, and not due to the impeller design itself. Here it could be demonstrated, that pitched blade impellers can result in yields >400kU/L/D, if the agitation rates are adjusted accordingly.

EXAMPLE 4:

PURIFICATION OF RECOMBINANT FURIN (rFURIN)

[0173] This example provides methods for the filtering and purifying rFurin. The collected cell culture supernatants were first filtered on depth filters (Cuno Zetaplus filters) to get them cell-free and particle-free, followed by membrane filtration at 0.45 μm PVDF filters (PALL Fluorodyne II). The filtered cell culture supernatant containing the rFurin was then concentrated by ultrafiltration on 10 K PES UF cassettes from Sartorius (Sartocon PESU 10kDa) with concentration factors ranging

from 10–50. The furin concentrates (with furin activity ranging from 290–1700 Units/ml) were then stored in aliquots at <-60°C.

[0174] In an effort to get a more homogeneous rFurin preparation for use in the VWF maturation process, experiments were conducted to partially purify the rFurin from the cell culture supernatant. A partially purified rFurin reagent is easier to use for characterization and release testing. It also results in a reduced presence of CHO host cell proteins in the VWF maturation process.

[0175] A purification procedure on an anion exchange resin was developed that required a loading conductivity of <5 mS/cm (RT) for efficient binding of rFurin. The elution was then performed as a step procedure at an ionic strength of approximately 500 - 300 mM NaCl and during the screening phase, a gradient elution up to 300 mM NaCl was applied (see overview in Table 24). The original pH of the buffers of 6.7 (RT) was increased to 7.5 to improve binding of rFurin during loading, in particular at high protein loading and high liner flow rates (see summary of relevant buffers in Table 25). The purification experiments were performed on EMD TMAE (Merck) and CaptoQ (GE Healthcare) anion exchange resins that differed in the stability of the packed and the maximum flow rate to operate. The analytical data summarized in Table 26 show that rFurin can be concentrated from the cell culture supernatant up to 362 fold with yields ranging from 20–71%. The rFurin activities in the eluate pools was between 639 Units/ml and 27651 Units/ml depending on the load applied. The CHO impurity level in the eluate was found in a range between 10–134 ng CHO protein/Unit rFurin and reduction rates up to 12.3 with a slightly better performance for CHO reduction found for the CaptoQ resin.

[0176] In summary, purification of rFurin by anion exchange chromatography proved to be an option for concentration, which is important for storage of the rFurin reagent at <-60°C. In addition, a slight CHO reduction by a factor from 3-12 is important in reducing the concentration of CHO protein in the preparation of VWF.

Table 24: Approximate procedure for the purification of rFurin on anion exchange resin. For EMD TMAE (Merck) the flow rate applied was 150/75 cm/h, for CaptoQ (GE Healthcare) the flow rate was 600/300.

Step	Buffer	CV	Flow rate [cm/h]	comment
------	--------	----	---------------------	---------

Step	Buffer	CV	Flow rate [cm/h]	comment
Equilibration	FEQ buffer	10	150 - 600	
Load	CCS diluted with EQ1	up to 1200		Conductivity of the load $\leq 5\text{mS/cm}$;
Wash 1	EQ buffer	up to 50		Optional; at high column loading this step was omitted
Wash 2	Mix of EQ buffer and FEL buffer	Approx. 10		
Step Elution	30% FEL/ 70%FEQ	14	75 - 300	
Gradient elution	100% FEQ – 30%FEL/70%FEQ	14	75 - 300	Optional: used for initial screening experiments
Post conditioning	Various buffers (FEL, NaCl buffer, CIP buffer)	15	n.d.	

Table 25: Buffers for the purification of furin on anion exchange resin

Buffer	formulation	
Equilibration buffer (FEQ)	50 mM HEPES, 1 mM CaCl_2 , pH= 6.7 – 7.5 at RT	At high loadings the pH of the buffers were increased to improve binding
Elution buffer (FEL)	50 mM HEPES, 1 mM CaCl_2 , 1 M NaCl, pH=6.7 – 7.5 at RT	
NaCl buffer	2 M NaCl	
CIP buffer	0.5 M NaOH	

Table 26: Summary of relevant purification experiments of rFurin on anion exchange resin. The total protein content was determined using a Bradford assay. The CHO reduction rate is calculated as CHO protein loaded divided by total CHO protein found in the eluate pool.

Run ID	Load	Eluate		Conc. Factor	yield	CHO		comment
	Column load	Furin						
	U Furin/ml resin	U/ml	U/mg protein	Vol load/Vol eluate	% Furin activity	ng/ U Furin	reduction	
011	2705	639	n.d.	32	36	134	3.6	TMAE; RT, elution

Run ID		Load	Eluate		Conc. Factor	yield	CHO		comment
		Column load	Furin						
		U Furin/ml resin	U/ml	U/mg protein	Vol load/Vol eluate	% Furin activity	ng/ U Furin	reduction	
									500 mM NaCl
013		4016	2259	n.d.	33	57	51	3.0	TMAE; RT, elution 400 mM NaCl
015		4403	2007	n.d.	41	46	50	3.8	TMAE; RT, elution 300 mM NaCl
018		10635	2584	n.d.	65	37	41	6.5	TMAE; RT, elution 300 mM NaCl
021		8684	3550	n.d.	126	20	59	5.5	TMAE; RT, elution 300 mM NaCl
024		4151	903	n.d.	19	54	38	6.3	CaptoQ; RT; elution 300 mM NaCl
025	E1	15696	6185	n.d.		45	10	12.3	CaptoQ; RT; gradient elution
	E2		1068	n.d.		12	n.d.	n.d.	
026	E1	31316	10237	14138		51	8	10.1	CaptoQ; 4°C; gradient elution
	E2		1868	n.d.		17	n.d.	n.d.	
028		32635	27651	10011	288	68	19	3.6	TMAE; 4°C; elution 300 mM NaCl
029C		23682	12642	7789	176	71	14.4	6.9	CaptoQ; RT; elution 300 mM NaCl
029T		23806	14922	2671	362	30	65.1	3.3	TMAE; RT, elution 300 mM NaCl
033		20625	6843	4568	n.d.	35	n.d.	4.6	CaptoQ; RT; elution 300 mM NaCl; load UF/DF conc.
035		26244	13209	6967	n.d.	78	n.d.	1.9	TMAE; RT, elution 300 mM NaCl; load UF/DF conc.

EXAMPLE 5:

CONCENTRATION, PURIFICATION, AND ANALYSIS OF rFURIN

[0177] This example describes other methods used in the concentration and purification (i.e., downstream processing) of large-scale rFurin. Such processing methods include ultrafiltration, diafiltration, and capto-MMC chromatography, that was carried out in the production of substantially animal protein-free rFurin. It also

describes methods of analyzing protein concentration, specific activity, and contamination by host cell protein and DNA.

[0178] Ultrafiltration. The supernatant (approx. 800–1200 kg in the Chemostat campaigns, and approx. 550–700 kg in the RFB-campaigns) was separated from the cells and concentrated to a final volume of 35–45 L by ultrafiltration. The parameters and setpoints of the Ultrafiltration/Diafiltration System (UFS) during the concentration step are listed in Table 27.

Table 27: Operating Parameters and Setpoints for the Concentration step

Parameter	Setpoint
Temperature of the filtered harvest	10–15 °C
Filter Area	14 m ²
Feed Pressure	0.9–1.5 bar
Retentate Pressure	0.7 – 1.2 bar
Permeate Pressure	0 – 0.1 bar
Transmembrane Pressure	0.7 – 1.0 bar
Specific Cross Flow Rate (Feed flow rate)	300–600 L/h/m ²
Specific Permeate Flow Rate	15–40 L/h/m ²
Concentration factor	20–30
Processing Time	3–4 h

[0179] Diafiltration. Immediately after finishing the concentration step, diafiltration of the retentate was initiated. The parameters and setpoints of the UFS during the diafiltration step are listed in Table 28.

Table 28: Operating Parameters and Setpoints for the Diafiltration Step

Parameter	Setpoint
Filter Area	14 m ²
Feed Pressure	0.9 – 1.5 bar

Parameter	Setpoint
Retentate Pressure	0.7 – 1.2 bar
Permeate Pressure	0 – 0.1 bar
Transmembrane Pressure	0.7 – 1.1 bar
Specific Cross Flow Rate (Feed flow rate)	300–600 L/h/m ²
Specific Permeate Flow Rate	7–20 L/h/m ²
Target Conductivity	< 2 mS/cm
Diafiltration rate ($v_{\text{permeate}}/v_{\text{retentate}}$)	4–5,5
Processing Time ¹⁾	0.75 – 1.25 h

[0180] When the conductivity of the retentate has fallen below 2 mS/cm (determined with the inline conductivity probe of the UFS), this process step was finished. This low conductivity is required to ensure a quantitative binding of the rFurin to the chromatographic gel in the subsequent purification step. The diafiltered product was transferred and the pH-value of the diafiltered product was adjusted to 6.0 by adding a 1 M acetic acid solution. The product was stored at room temperature (RT) before applying to the Capto-MMC column.

[0181] Capto-MMC Chromatography. The Capto-MMC gel, a multimodal cation exchanger, was used to bind rFurin and to eliminate the vast majority of contaminants from the diafiltrated product. After equilibration of the chromatography gel, the diafiltered product is loaded to the column. A 0.22 µm filter capsule was installed to perform an online filtration of the diafiltered product. The further chromatographic steps are listed and detailed in Table 29.

Table 29: Capto-MMC Chromatography Steps

	Chromatography step	Buffer / Product	Linear flow rate	Quantity
			cm/h	CV ¹⁾
1	Acid wash	ES4 online diluted to 0,2M acetic acid	300	1
2	Equilibration	EP2	150	37
3	Absorption	35 – 45 L diafiltered product	150	

	Chromatography step	Buffer / Product	Linear flow rate	Quantity
			cm/h	CV ¹⁾
4	Equilibration	EP2	150	30
5	Washing	WPF	150	10
6	Gradient Elution	EL1 / MMC eluate	150	15
7	Elution II	EL2 / MMC eluate II	150	10
8	Regeneration	ES4	150	10
9	Regeneration	SHD	150	10

¹⁾ CV = column volume: volume of the packed resin in the chromatography column

[0182] In the downstream processing, no issues were observed. The downstream steps (Filtration, UF, DF) could be carried out at the same conditions used for the harvests from the Chemostat culture. Mean total rFurin activities yields of 91 % (Campaign ORFU06002), 66 % (Campaign ORFU07001) and 84 % (Campaign ORFU07002) could be achieved. The only minor change is given by the fact, that the starting volume for the UF-step is somewhat lower compared to continuous culture. But this change has no effect on the quality of the purified product. (See data on total activity, Host Cell DNA, Host Cell Protein; see Table 30).

[0183] The mean values of all three campaigns for the Host Cell Protein and Host Cell DNA content reveal rather low mean maximum values of 8.55 µg/ml (ranging from 2.2-10.4 µg/ml) and 13.48 ng/ml (ranging from 0.0-23.9 ng/ml), respectively (see Table 30). The chromatographic step was able to reduce the specific contamination to low mean maximum values of 0.35 ng CHO Protein/U rFurin Activity (ranging from 0.13-0.52 ng CHO Protein/U) and 0.148 pg Host Cell DNA/U rFurin Activity (ranging from 0.0- 365 pg DNA/U rFurin Activity).

Table 30: Capto-MMC Chromatography impurity results – Campaign comparison

campaign	CHO-Protein Content	CHO-Protein / rFurin Activity	Host Cell DNA Content	Host Cell DNA / rFurin Activity
	µg/mL	ng/U	ng/mL	pg/U
ORFU06002	8.55	0.22	1.66	0.059
ORFU07001	3.50	0.35	2.54	0.031

ORFU07002	2.36	0.026	13.48	0.148
-----------	------	-------	-------	-------

[0184] Furin Characterization. The quality and functionality of rFurin was assessed by following biochemical / biophysical methods (Table 31).

Table 31: Analytical methods for rFurin

Method	Aim	Comment
Fluorescent substrate cleavage	Enzyme activity	Low molecular weight substrate
Furin use test	Enzyme activity, maturation efficacy	rVWF substrate
SDS-PAGE	Protein composition	Western blot assay
IEF	Protein composition	Coomassie staining, Western blot assay
RP-HPLC	Protein composition	Protein fingerprint

[0185] Furin Activity Assay. The purified rFurin batches (Capto-MMC eluate pools) were tested for enzymatic activity of Furin. The substrate is a short synthetic peptide containing the dibasic recognition sequence attached to a fluorescent amino-methyl coumarin (AMC) group, that is released after cleavage (BOC-RVRR-AMC). The released fluorogenic group can be detected by excitation at 380nm and subsequent measurement of the emitted light at 435nm. One activity unit is defined as the release of 1 pMol of AMC per minute at 30°C.

[0186] Depending on the fermentation and purification efficacy the measured values of rFurin activity were in the range of about 10000 U/ml up to more than about 100000 U/ml, with a mean value of approx. 69000 U/ml (Table 32, Table 32, and Table 34). An increase of rFurin activity for the RFB mode campaigns was noticed, especially when comparing the mean values of the Chemostat campaign ORFU06002 (47737 U/ml) with the mean values of RFB campaigns ORFU07001 (77653 U/ml) and ORFU07002 (93178 U/ml). (Overall, rFurin activity ranged from about 10000 to greater than 100000 U/ml; all data not shown).

[0187] The specific activity of Furin is expressed as the activity U/μg protein (see Tables 32-34). The mean specific activity for campaign ORFU06002 was 269U/μg

protein, increasing to 500U/μg for ORFU07001 and 563U/μg for ORFU07002, respectively. (Overall, specific activity ranged from 124-620 U/μg protein; data not shown). Thus, specific activity doubled for the two RFB campaigns, a result of the higher enzymatic activity of the RFB rFurin compared to the batches produced in Chemostat mode.

Table 32: Campaign ORFU06002 Capto-MMC Eluate Pool

Furin-Lot Chemostat Mode	Furin Activity	Protein Content	Specific rFurin Activity
	[U/ml]	[μg/ml]	[U/μg]
ORFUCHR06002MMC01	28231	148	190.8
ORFUCHR06002MMC02	16307	162	124.3
ORFUCHR06002MMC03	44854	174	257.8
ORFUCHR06002MMC04	39102	148	264.2
ORFUCHR06002MMC05	71871	236	304.5
ORFUCHR06002MMC06	44292	142	311.9
ORFUCHR06002MMC07	68728	155	443.4
ORFUCHR06002MMC08	68510	271	252.8
Mean Value	47737	180	269

Table 33: Campaign ORFU07001 Capto-MMC Eluate Pool

Furin-Lot Repeated Fed-Batch	Furin Activity	Protein Content	Specific rFurin Activity
	[U/ml]	[μg/ml]	[U/μg]
ORFUCHR07001MMC01	77090	151	510.5
ORFUCHR07001MMC02	68450	112	611.2
ORFUCHR07001MMC03	87420	231	378.4
Mean Value	77653	165	500

Table 34: Campaign ORFU07002 Capto-MMC Eluate Pool

Furin-Lot Repeated Fed-Batch	Furin Activity	Protein Content	Specific rFurin Activity
	[U/ml]	[μg/ml]	[U/μg]
ORFUCHR07002MMC01	91200	162	563.0
ORFUCHR07002MMC02	96400	160	602.5
ORFUCHR07002MMC03	95020	173	549.2
ORFUCHR07002MMC04	85690	190	451.0
ORFUCHR07002MMC05	102670	174	590.1
ORFUCHR07002MMC06	88090	142	620.4
Mean Value	93178	167	563

[0188] **Furin-Use-Test Activity.** The Furin-Use-Test is designed to quantify the efficacy of rFurin to process pro-VWF to mature rVWF. The maturation efficacy is expressed as the amount of Furin units required for the maturation of 1 VWF Antigen unit (U Furin / U VWF). The substrate is a proVWF/VWF preparation that has been purified at Pilot scale according to the current manufacturing procedure but omitting the Furin maturation and the final purification step on Superose 6. The rVWF substrate concentration was 100 U Ag/ml (F8HL_24_01UF02-R).

[0189] Four dilutions of the sample were tested in a 1 ml Eppendorf Tube to cover specific Furin concentrations of, e.g., 0.25–2.0 U/U VWF. The reaction and dilution buffer was 100 mM HEPES, 1 mM CaCl₂, pH7.0 and the maturation experiment was performed for 16 hours at 37°C under slight agitation. After the incubation the enzymatic reaction is stopped by addition of SDS-sample buffer and heating the samples for 5 minutes at > 90°C. The samples are then analysed by SDS-PAGE on 8% gels using silver staining of the separated polypeptides. The specific Furin activity required to get a maturation efficacy of > 95% as evaluated by visual inspection of the gels (the proVWF band should represent less than 5% compared to the mature VWF band) is then reported and used as a quality attribute of the rFurin tested.

Table 35: Campaign ORFU06002 Capto-MMC Eluate Pool

Furin-Lot Chemostat Mode	Furin Activity	Use-test Activity	Maturation Degree
	[U/ml]	[U/U]	[%]
ORFUCHR06002MMC01	28231	0.7	>95
ORFUCHR06002MMC02	16307	0.4	>95
ORFUCHR06002MMC03	44854	1.1	>95
ORFUCHR06002MMC04	39102	1.3	>95
ORFUCHR06002MMC05	71871	1.0	>95
ORFUCHR06002MMC06	44292	1.0	>95
ORFUCHR06002MMC07	68728	1.0	>95
ORFUCHR06002MMC08	68510	1.0	>95
Mean Value	47737	0.58	>95

Table 36: Campaign ORFU07001 Capto-MMC Eluate Pool

Furin-Lot Repeated Fed-Batch	Furin Activity	Use-test Activity	Maturation Degree
	[U/ml]	[U/U]	[%]
ORFUCHR07001MMC01	77090	0.5	>95
ORFUCHR07001MMC02	68450	0.5	>95
ORFUCHR07001MMC03	87420	0.5	>95

Mean Value	77653	0.5	>95
------------	-------	-----	-----

Table 37: Campaign ORFU07002 Capto-MMC Eluate Pool

Furin-Lot Repeated Fed-Batch	Furin Activity [U/ml]	Use-test Activity [U/U]	Maturation Degree [%]
ORFUCHR07002MMC01	91200	0.5	>95
ORFUCHR07002MMC02	96400	0.5	>95
ORFUCHR07002MMC03	95020	0.5	>95
ORFUCHR07002MMC04	85690	0.3	>95
ORFUCHR07002MMC05	102670	0.6	>95
ORFUCHR07002MMC06	88090	1.0	>95
Mean Value	93178	0.6	>95

[0190] Good and consistent maturation activity for proVWF was found for all tested rFurin batches. The mean values of the campaigns are below 1.0 U Furin/U VWF:Ag and the maturation degree exceeds 95% in all cases. (Tables 35-37). Maturation activities of all rFurin batches are comparable and, referring to the calculated mean values of the campaigns ORFU06002, ORFU07001 and ORFU07002, almost no differences can be found between rFurin produced in Chemostat mode and RFB-mode.

[0191] Furin SDS-PAGE and Silver Stain. 8% SDS-PAGE with silver staining and Western blot analysis was performed for all rFurin batches to ensure consistent quality and visualize the degree of impurity. As seen in Figure 12, the band patterns of the Capto-MMC eluates of campaign ORFU06002 and ORFU07002 correlate to a high degree; all samples show a prominent Furin band at approx. 60kDa. A trend to slightly lower molecular weight of the Furin bands is visible in samples of campaign ORFU06002 from batches MMC01 to MMC08 (Figure 12, lanes 1-8). Samples of those batches were deglycosylated with the effect that this trend was not visible anymore in subsequent SDS-PAGE with silvers staining (data not shown), supporting the assumption that during campaign ORFU06002, rFurin was glycosylated slightly different or to a lesser degree in course of ongoing fermentation. This trend was not noticed in RFB campaign ORFU07002, suggesting constant glycosylation of the rFurin during the whole campaign. The impurities in batches from ORFU06002 and ORFU07002 are almost completely the same; however, samples from the RFB-mode campaign ORFU07002 show less intense bands in the 40kDa region, polypeptides that were particularly strong enriched in samples of campaign ORFU06002.

[0192] SDS-PAGE Western Blot. Western blot analysis for all samples was performed using a monoclonal anti-Furin antibody (Figure 13). The prominent band at ~60kDa can be identified as the Furin band and is found in all samples. Comparability of the samples is very high. Slight variations in band intensity are due to the different Furin concentration in the samples. Overall, SDS-PAGE analysis underlines the comparability of rFurin produced in Chemostat and RFB mode.

[0193] To support conventional SDS-PAGE, Isoelectric Focusing (IEF) was performed on samples of campaigns ORFU06002 and ORFU07002. IEF was performed between pH 7.0 and pH 3.0, using vertical IEF. The polypeptides were visualized with Coomassie staining and Western blot analysis. IEF and subsequent Western blotting of rFurin samples of campaign ORFU06002 provided specific band patterns for Furin (see Figure 14). Up to seven separate bands in the region of pH 4.5 to pH 5.5 can be identified, with at least five bands present in all samples.

[0194] IEF of samples of campaign ORFU07002 was carried out using Coomassie staining and Western blot analysis for visualization. Coomassie staining reveals the specific band pattern with all samples showing at minimum five separate bands in the range of pH 4.5 to pH 5.5, and up to eight bands can be identified (see Figure 15, Lane 3).

[0195] Western blot analysis (see Figure 16) corroborates these results, with some samples showing not all the bands visible in the Coomassie gel (see Figure 17), probably due to incomplete transfer of the proteins from the gel to the blotting membrane.

[0196] Furin Reverse Phase HPLC. Samples from campaigns ORFU06002 and ORFU07003 (Capto-MMC eluates) were tested with C4 RP-HPLC in order to establish a fingerprint pattern for rFurin. The peak patterns of the samples of the two campaigns were compared (see Figure 17 and Figure 18).

[0197] Chromatograms of all tested samples show a characteristic main peak at a retention time of approx. 13min., which can be assigned to Furin. The peak heights correlate well with the Furin concentration in the samples. Other protein impurities can be seen as minor peaks in the range of 8min to 17min. All samples from campaign ORFU07002 show significantly less and smaller peaks from impurities than those from ORFU06002. This fact is well in accordance with the results of

SDS-PAGE, as in the RFB mode campaign ORFU07002 a smaller number and decreased amount of impurities were found.

[0198] Analytical data obtained from the above discussed characterization methods prove very good comparability of the rFurin produced in both Chemostat mode and RFB mode. No major differences in rFurin quality could be detected from the data available, however rFurin batches produced in RFB mode showed higher specific activity and less impurities compared to production in Chemostat mode. The chromatographic step accounts for the very low Host Cell Protein and Host Cell DNA content in the Bulk Drug Substance (BDS). At least a complete production process consisting of a RFB-fermentation and the whole down-stream processing (Filtration, UF/DF, Capto-MMC chromatography) is maybe easier to be implemented for the commercial production of rFurin, e.g. in Single Use Bioreactor (SUB) systems.

EXAMPLE 6:

SAFETY, STERILITY, AND STABILITY TESTING

[0199] This example describes the safety, sterility, and stability testing that is performed to determine and maintain the quality of the CHO cell bank. Testing on sterility/mycoplasma has to be performed in accordance to requests of the ICH-Guideline Q5D. The quality of the cell bank has to be checked by determination of average viability and cell density of the thawed cells and subsequent growth rate of the cultures.

[0200] Cells were tested for viral safety (Table 38), genetic stability (Table 39), and identity (Table 40). Cells were found to be sterile, free of mycoplasma, free from extraneous agents, free from retroviruses, negative for MVM virus, negative for adventitious viruses, negative for rodent viruses, free from porcine and bovine viruses, and free from Cache Valley virus (CVV).

[0201] Cell banks were examined with regard to the Furin producer/non-producer ratio by FACS analysis. They are also tested for their long term stability (ability to produce Furin over a period of time). Further, the secreted Furin produced under serum-free conditions has to be investigated with respect to generated isoforms. All data should indicate that stable growth and Furin production can be achieved using these cell banks.

[0202] Cells from the MCB/WCB should show stable growth and Furin expression over the entire production process.

Table 38. Viral Safety Program

Test	MCB	WCB	MEPC
Adventitious viruses in vitro (CHO, A9, Vero, MRC-5 Cells)	x	x	x
Adventitious viruses in vivo (in suckling mice, adult mice, guinea pigs, and embryonated eggs)	x	(x)	x
Rodent viruses MAP test with LCMV challenge	x	---	---
Rodent viruses HAP	x	---	---
Rodent viruses MMV (in vitro assay)	x	---	x
Retroviruses EM (Transmission electron microscopic examination)	x	---	x
Retroviruses Retroviral Infectivity Assay with Pert Endpoint	x	---	x
Retroviruses Ex S+L- (in vitro detection of xenotropic retroviruses by Mink S+L-Focus)	x	---	x
Retroviruses/ Co cultivation Detection of infectious retroviruses by co-cultivation with HEK 293 cells (5 passages)	---	---	x
Bovine virus: Detection of viral contaminants in Bovine serum according CMPC & US 9CFR requirements	x	---	---
Porcine virus: In vitro assay, detection of porcine viral contaminants using PPK indicator cells according to	x	---	---

Test	MCB	WCB	MEPC
9 CFR			
PCR Bovine polyomavirus	x	---	x
Real time PCR of Cache Valley Virus (CVV)	x	---	x

(x) viral testing has not to be performed on the WCB from which the MEPC have been prepared

Table 39. Genetic Stability Program

Test	MCB	WCB	MEPC
Productivity: Furin activity assay	x	x	x
Sequencing	x	---	x
quantitative PCR (gene copy number)	x	---	x
Southern Blot	x	---	x
Northern Blot	x	---	x

Table 40. Identity Program

Test	MCB	WCB	MEPC
Isoenzyme analysis	x	x	x

[0203] The present invention has been described in terms of particular embodiments found or proposed to comprise preferred modes for the practice of then invention. It will be appreciated by those of ordinary skill in the art that, in light of the present disclosure, numerous modifications and changes can be made in the particular embodiments exemplified without departing from the intended scope of the invention. Therefore, it is intended that the appended claims cover all such equivalent variations which come within the scope of the invention as claimed.

WHAT IS CLAIMED IS:

1. A composition comprising substantially animal protein-free recombinant furin lacking transmembrane and cytoplasmic domains and comprising a specific activity of at least 400 U/ μ g protein, the composition comprising host cell protein at a concentration less than 1.0 ng protein/U furin activity or host cell DNA at a concentration less than 0.5 pg DNA/U furin activity, and essentially lacking contaminating proteins from serum.
2. The composition of claim 1 comprising host cell protein at a concentration less than 0.8 ng protein/U furin activity.
3. The composition of claim 1 comprising recombinant furin at an activity of at least 450 U/ μ g.
4. The composition of claim 1 comprising recombinant furin at an activity of at least 500 U/ μ g.
5. The composition of claim 1 comprising recombinant furin at an activity of at least 550 U/ μ g.
6. The composition of claim 1 comprising host cell protein at a concentration less than 0.6 ng protein/U furin activity.
7. The composition of claim 1 comprising host cell protein at a concentration less than 0.4 ng protein/U furin activity.
8. The composition of claim 1 comprising host cell DNA at a concentration less than 0.4 pg DNA/U furin activity.
9. The composition of claim 1 comprising host cell DNA at a concentration from between 0 to 0.4 pg DNA/U furin activity.
10. A method of making the composition of claim 1 comprising growing a host cell transformed or transfected with a polynucleotide encoding secreted recombinant furin lacking transmembrane and cytoplasmic domains in serum-free medium in a repeated fed batch process; binding the secreted furin on a multimodal cation exchange resin to remove host cell protein, host cell DNA, or a combination thereof; and recovering the secreted recombinant furin from the multimodal cation exchange resin.
11. The method of claim 10 comprising adapting the host cell to grow in medium with increasingly lower concentrations of serum until all serum is removed from the medium.

12. The method of claim 10 comprising transferring the host cell from growth in medium comprising serum to growth in serum-free medium.
13. The method of claim 11 or 12, wherein the host cell is a Chinese Hamster Ovary (CHO) cell.
14. A method of processing a pro-protein comprising a furin cleavage site to cleave a pro-peptide from the pro-protein to form a mature protein comprising contacting the pro-protein comprising the furin cleavage site with the composition of claim 1.
15. The method of claim 14, wherein the mature protein is von Willebrand Factor.
16. The method of claim 14, wherein the mature protein is Factor VIII.
17. The composition of any one of claims 1-9, wherein the host cell is a Chinese Hamster Ovary (CHO) cell.