

DESCRIPTION

TITLE OF THE INVENTION

CARBOXYMETHYL PIPERIDINE DERIVATIVE

5

TECHNICAL FIELD

[0001]

The present invention relates to carboxymethyl piperidine derivatives useful as medicaments.

10

[0002]

More particularly, the present invention relates to carboxymethyl piperidine derivatives or pharmaceutically acceptable salts thereof which have substance P/neurokinin 1 (NK₁) receptor antagonist activity, and which are useful as agents for the prevention or treatment of cancer-chemotherapy-induced nausea and vomiting (CINV) and so on.

15

BACKGROUND ART

[0003]

CINV occurs when the vomiting center located in the lateral reticular formation of the medulla oblongata receives a stimulus. The area postrema and the solitary nucleus of the medulla oblongata contain NK₁ receptors, and the NK₁ receptors are believed to be closely involved in vomiting.

20

Administration of an antineoplastic agent facilitates the serotonin secretion from the enterochromaffin (EC) cells in the digestive tract, and serotonin directly stimulates the vomiting center through 5-hydroxytryptamine₃ (5-HT₃) receptors in the digestive tract. Also, when serotonin stimulates the vomiting center through the chemoreceptor trigger zone (CTZ) located in the area postrema of the fourth ventricle,

25

nausea and vomiting occur. Substance P, like serotonin, is found in the EC cells in the digestive tract, and its secretion is promoted by administration of an antineoplastic agent. Recently, it has been revealed that substance P induces vomiting through the NK₁ receptors in the CTZ or by binding to the NK₁ receptors in the central nervous system, and therefore NK₁ receptors have been attracting attention as the target for developing antiemetic agents (Non-patent literature 1).

[0004]

Aprepitant is the first selective NK₁ receptor antagonist in the world which was approved as a preventive agent for nausea and vomiting associated with administration of antineoplastic agents. Regarding the mechanism of action of aprepitant, it is believed that aprepitant selectively inhibits the binding of substance P and the NK₁ receptors in the central nervous system, which is one of the pathways that induce CINV, and thus prevents CINV. Aprepitant has been launched as a preventive agent for CINV (Non-patent literature 2).

[0005]

It is known that aprepitant is metabolized by cytochrome P450 (CYP) 3A4. Also, aprepitant is known to have a dose-dependent inhibitory effect on CYP3A4, a CYP3A4-inducing effect and a CYP2C9-inducing effect. Accordingly, aprepitant may cause the drug-drug interactions with drugs that inhibit or induce CYP3A4 or with drugs that are metabolized by CYP3A4 or CYP2C9. For example, it is reported that the inhibitory effect of aprepitant on CYP3A4 sometimes inhibits the metabolism of dexamethasone and that the dose should be thus adjusted when dexamethasone is combined with aprepitant (Non-patent literature 3).

Therefore, when aprepitant is used, sufficient care should be directed to the drug-drug interactions based on the inhibitory effect of aprepitant on CYP3A4.

[0006]

For the above reasons, a novel NK₁ receptor antagonist with fewer drug-drug

interactions is required in the prevention or treatment of CINV.

[0007]

Compounds with an NK₁ receptor antagonist activity such as casopitant, netupitant, ezlopitant, rolapitant, vestipitant, vofopitant and so on, are known.

5 However, casopitant is reported to have an inhibitory effect on CYP3A4 and cause the drug-drug interactions due to the effect (Non-patent literature 4). Clinical trials on casopitant, as a preventive agent for cancer-chemotherapy-induced nausea and vomiting, had been conducted in the U.S. and Europe; however, its development was discontinued after the application. Netupitant is currently under development as a
10 preventive agent for cancer-chemotherapy-induced nausea and vomiting,; however, netupitant is reported to have an inhibitory effect on CYP3A4 and cause the drug-drug interactions due to the effect (Non-patent literature 5). Clinical trials on ezlopitant, as a preventive agent for cancer-chemotherapy-induced nausea and vomiting, had been conducted in the U.S.; however, its development was discontinued. Clinical trials on
15 vofopitant, as a preventive agent for cancer-chemotherapy-induced nausea and vomiting, had been conducted in Europe; however, its development was discontinued.

Many of the above compounds resulted in the discontinuance.

[0008]

Pyridine derivatives claiming to be having NK₁ receptor antagonist activity are
20 described in Patent literature 1 to 16. And, prodrugs of pyridine derivatives are described in Patent literature 17 and 18.

However, carboxymethyl piperidine derivatives of the present invention are not described in the above literatures.

25 **CITATION LIST**

Patent literature

[0009]

Patent literature 1: U.S. Patent No. 6,479,483

Patent literature 2: U.S. Patent No. 6,770,637

Patent literature 3: U.S. Patent No. 7,939,533

Patent literature 4: European Patent No. 1,103,545

5 Patent literature 5: U.S. Patent No. 7,211,579

Patent literature 6: U.S. Patent Publication No.2006/0030600

Patent literature 7: U.S. Patent No. 6,576,762

Patent literature 8: U.S. Patent No. 6,225,316

Patent literature 9: U.S. Patent No. 7,683,056

10 Patent literature 10: U.S. Patent No. 8,344,005

Patent literature 11: International publication No. WO2011/ 054773

Patent literature 12: U.S. Patent Publication No. 2007/0071813

Patent literature 13: U.S. Patent Publication No. 2003/0083345

Patent literature 14: U.S. Patent Publication No. 2003/0004157

15 Patent literature 15: U.S. Patent No. 6,849,624

Patent literature 16: U.S. Patent No. 6,297,375

Patent literature 17: U.S. Patent No. 6,593,472

Patent literature 18: U.S. Patent No. 8,426,450

Non-Patent literature

20 [0010]

Non-patent literature 1: P. J. Hesketh et al., European Journal of Cancer, 2003, Vol. 39, pp. 1074-1080

Non-patent literature 2: Toni M. Dando et al., Drugs, 2004, Vol. 64, No. 7, pp. 777-794

25 Non-patent literature 3: Jacqueline B. McCrea et al., CLINICAL PHARMACOLOGY & THERAPEUTICS, 2003, Vol. 74, No. 1, pp. 17-24

Non-patent literature 4: Stefano Zamuner et al., British Journal of Clinical

Pharmacology, 2010, Vol. 70, No. 4, pp. 537-546

Non-patent literature 5: Corinna Lanzarotti et al., Support Care Cancer, 2013, Vol. 21, No. 10, pp. 2783-2791

5 SUMMARY OF THE INVENTION

Problem to be solved by the Invention

[0011]

A problem of the present invention is to provide a new compound which has NK₁ receptor antagonist activity, whose CYP3A4 inhibitory activity is reduced compared to aprepitant, and which are useful for the prevention or treatment of cancer-chemotherapy-induced nausea and vomiting. A problem of the present invention is preferably to provide the above compound whose central transportation property and long-acting medicinal effect is excellent.

Means for solving the Problem

15 [0012]


The present invention relates to a compound represented by the following formula (I) or a pharmaceutically acceptable salt thereof.

[0013]

That is, the present invention relates to the following [1] to [12] and the like.

20 [1] A compound represented by the formula (I):


[Chem.1]



wherein

ring A is a group represented by the formula:


[Chem.2]



;

ring B is a group represented by the formula:

[Chem.3]




5

;

with the proviso that bonds with (*) are bonding site to the formula:

[Chem.4]




;

bonds with (**) are bonding site to the ring A;

10 bonds with (***) are bonding site to the formula:

[Chem.5]



;


R₁ is C₁₋₆ alkyl or C₁₋₆ alkoxy;R₂ and R₃ are each independently a hydrogen atom or methyl;

15 n is 0, 1, 2, 3, 4 or 5;

or a pharmaceutically acceptable salt thereof.

[2] The compound represented by the formula (Ia) according to the above [1]:

[Chem.6]




wherein

ring A, R_1 and n have the same meaning as described in the above [1];

- 5 ring B^b is a group represented by the formula:


[Chem.7]



;

wherein, bond with $(*)$ is a bonding site to the formula:

[Chem.8]



10


;

 $(**)$ and $(***)$ have the same meaning as described in the above [1] ;

or a pharmaceutically acceptable salt thereof.

[3] The compound represented by the formula (Ib) according to the above [2]:

[Chem.9]




15

wherein

ring A and ring B^b have the same meaning as described in the above [2];

with the proviso that bond with (*) is a bonding site to the formula:

[Chem.10]



;


- 5 (**) and (***) have the same meaning as described in the above [2] ;

R^{1a}, R^{1b}, R^{1c}, R^{1d} and R^{1e} are each independently any one of a hydrogen atom, methyl or methoxy;

or a pharmaceutically acceptable salt thereof.

[4] The compound according to the above [1], wherein ring B is the following formula:

10 [Chem.11]



wherein, (*), (**) and (***) have the same meaning as described in the above [1] ;


or a pharmaceutically acceptable salt thereof.

[5] The compound according to the above [4], wherein R₁ is methyl, and n is 0 or 1, or a

- 15 pharmaceutically acceptable salt thereof.

[6] The compound represented by the following formula according to the above [2]:


[Chem.12]



or a pharmaceutically acceptable salt thereof.

[7] The compound represented by the following formula according to the above [2]:

[Chem.13]




5

or a pharmaceutically acceptable salt thereof.

[8] The compound represented by the following formula according to the above [2]:


[Chem.14]



10 or a pharmaceutically acceptable salt thereof.

[9] The compound represented by the following formula according to the above [2]:


[Chem.15]



or a pharmaceutically acceptable salt thereof.

[10] The compound represented by the following formula according to the above [1]:

[Chem.16]



5

or a pharmaceutically acceptable salt thereof.

[11] A pharmaceutical composition comprising as an active ingredient a compound according to any one of the above [1] to [10], or a pharmaceutically acceptable salt thereof.

- 10 [12] The pharmaceutical composition according to the above [11], for use in the prevention or treatment of cancer-chemotherapy-induced nausea and vomiting.

Effect of the Invention

[0014]

- 15 The compounds of the present invention have an excellent NK₁ receptor antagonist activity. And, CYP3A4 inhibitory of the compounds of the present invention is reduced compared to aprepitant. Furthermore, the preferable compounds of the present invention excel in central transportation property and long-acting

medicinal effect.

Therefore, the compounds of the present invention or pharmaceutically acceptable salts thereof are useful as an agent for the prevention or treatment of cancer-chemotherapy-induced nausea and vomiting.

5 **Brief Description of the Drawings**

[0015]

[Figure 1] Figure 1 shows the effect on cisplatin-induced acute and delayed emetic response. In the figure, each bar chart shows a value of control group (Control), the group intravenously administered with 3 mg/kg of the compound of Example 1 (Ex. No 1, 3 mg/kg, iv) and the group orally administered with 1 mg/kg of the compound of Example 1 (Ex. No 1, 1 mg/kg, po) in the acute phase, and a value of control group, the group intravenously administered with 3 mg/kg of the compound of Example 1 (Ex. No 1, 3 mg/kg, iv) and the group orally administered with 1 mg/kg of the compound of Example 1 (Ex. No 1, 1 mg/kg, po) in the delayed phase from the left respectively. The vertical axes show the number of retching and vomiting (Retches + Vomites) (the mean+standard error of 5 examples of control group, the mean+standard error of 5 examples of the group intravenously administered, and the mean+standard error of 5 examples of the group orally administered).

Mode for Carrying out the Invention

20 [0016]

In the present invention, each term has the following meaning unless otherwise specified.

[0017]

The term "C₁₋₆ alkyl" means a straight-chained or a branched alkyl group having 1 to 6 carbon atoms, and for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl can be illustrated.

The term "C₁₋₆ alkoxy" means a straight-chained or a branched alkoxy group

having 1 to 6 carbon atoms, and for example, methoxy, ethoxy, propoxy, isopropoxy can be illustrated.

[0018]

In the compounds represented by the formula (I) of the present invention, the
5 symbol "R₁" means a substituent of the piperidine ring.

[0019]

The present invention is further illustrated below.

[0020]

In the case where the compounds represented by the formula (I) of the present
10 invention contain one or more asymmetric carbon atoms, all stereoisomers in the *R*- or *S*-configuration at each of asymmetric carbons and their mixtures are included in the present invention. In such cases, racemic compounds, racemic mixtures, individual enantiomers and mixtures of diastereomers are included in the scope of the present invention. In the case where the compounds represented by the formula (I) of the
15 present invention have the cis-trans isomers, all cis-trans isomers are included in the present invention.

[0021]

In the present invention, stereochemical determination can be also determined according to well-known methods in the art. For example, see also "Tokuron NMR
20 rittai kagaku", Kodansha, 2012, p. 59.

[0022]

A compound represented by the formula (I) of the present invention can be converted into pharmaceutically acceptable salts thereof according to a general method. As such salts, acid additive salts and salts with a base can be illustrated.

25 [0023]

As the acid additive salt, an acid additive salt with a mineral acid such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, nitric acid,

phosphoric acid and the like, and an acid additive salt with an organic acid such as formic acid, acetic acid, trifluoroacetic acid, methanesulfonic acid, benzenesulfonic acid, *p*-toluenesulfonic acid, propionic acid, citric acid, succinic acid, tartaric acid, fumaric acid, butyric acid, oxalic acid, malonic acid, maleic acid, lactic acid, malic acid, carbonic acid, benzoic acid, glutamic acid, aspartic acid and the like can be illustrated.

[0024]

As the salt with a base, a salt formed with inorganic base such as a lithium salt, a sodium salt, a potassium salt, a calcium salt, a magnesium salt and the like, and a salt formed with organic base such as N-methyl-D-glucamine, N,N'-dibenzylethylenediamine, triethylamine, piperidine, morpholine, pyrrolidine, arginine, lysine, choline and the like.

[0025]

In the present invention, a pharmaceutically acceptable salt also includes a solvate thereof with a pharmaceutically acceptable solvent such as water, ethanol or the like.

[0026]


In an embodiment of a compound represented by the formula (I) of the present invention, *n* is 0, 1 or 2.

[0027]

A compound represented by the formula (I) of the present invention can be also prepared, for example, by a method described below or a similar method thereto, or a method described in literatures or a similar method thereto.

Scheme 1

[Chem.17]



In the formula, L^1 and L^2 are each independently a leaving group such as a chlorine atom, a bromine atom, an iodine atom, a trifluoromethanesulfonyloxy group or the like, PG^1 is a protective group, and ring A, ring B, R_1 , R_2 , R_3 and n have the same meanings as defined above.

[0028]

Process 1

Compound (4) can be also prepared by conducting coupling reaction of Compound (2) with Compound (3) in an inert solvent in the presence of a base and a palladium catalyst.

[0029]

Process 2

Compound (6) can be also prepared by conducting condensation reaction of Compound (4) with Compound (10) in an inert solvent in the presence of a base.

[0030]

Process 3

Compound (I) can be also prepared by allowing Compound (6) to react with


Compound (7), and conducting deprotection in an inert solvent in the presence or absence of a base.

As the inert solvent, for example, N, N-dimethylformamide, N-methylpyrrolidone, dimethylsulfoxide, diethyl ether, tetrahydrofuran, 1, 4-dioxane, 1, 2-dimethoxyethane, benzene, toluene, xylene, and a mixed solvent thereof can be illustrated. As the base, potassium carbonate, sodium carbonate, cesium carbonate, sodium hydroxide, potassium hydroxide, lithium hydroxide, potassium fluoride, cesium fluoride, triethylamine, pyridine, N, N-diisopropylethylamine, 2, 6-lutidine, and 1, 8-diazabicyclo[5, 4, 0]-7-undecene can be illustrated. The reaction temperature is usually at 0°C to reflux temperature. The reaction time is usually from 30 minutes to 7 days, varying based on a used starting material, solvent and reaction temperature or the like. The above reaction can also be conducted by using a microwave reactor (Biotage). When a microwave reactor is used, the reaction is conducted at pressure range: 1 to 30 bar, power range: 1 to 400 W, reaction temperature: room temperature to 300°C, and reaction time: a minute to 1 day, varying based on a used starting material, solvent and model.

[0031]

Scheme 2

[Chem.18]



In the formula, PG¹ and PG² represent a protective group, and R₁, R₂, and n have the same meanings as defined above.

[0032]

Process 4

Compound (10) can also be prepared by conducting olefination reaction of Compound (9) with Compound (8) in an inert solvent in the presence of a base. As the inert solvent, for example, N, N-dimethylformamide, N-methylpyrrolidone, dimethylsulfoxide, diethyl ether, tetrahydrofuran, 1, 4-dioxane, 1, 2-dimethoxyethane, benzene, toluene, xylene, and a mixed solvent thereof can be illustrated. As the base, sodium hydride, sodium methoxide, potassium carbonate, cesium carbonate, potassium *tert*-butoxide, triethylamine, N, N-diisopropylethylamine, and 1, 8-diazabicyclo[5, 4, 0]-7-undecene can be illustrated. The reaction temperature is usually at -78°C to reflux temperature. The reaction time is usually from 30 minutes to 7 days, varying based on a used starting material, solvent and reaction temperature or the like. The reaction can also be conducted by using a microwave reactor (Biotage).

[0033]

Process 5

Compound (7a) can also be prepared by conducting olefin reduction such as catalytic reduction method of Compound (10) and so on, and conducting deprotection. The catalytic reduction method can be conducted, for example, by allowing Compound (10) to react by using a catalyst under a hydrogen atmosphere in a solvent. As the solvent, for example, methanol, ethanol, ethyl acetate, tetrahydrofuran, acetic acid and the like can be illustrated. As the catalyst, for example, palladium-carbon powder, rhodium-carbon powder, platinum-carbon powder, platinum-carbon powder doped with vanadium can be illustrated. The reaction temperature is usually at room temperature to reflux temperature. The reaction time is usually from 30 minutes to 7 days, varying based on a used starting material, solvent and reaction temperature or the like.

[0034]

The above-mentioned schemes are exemplary for preparing compounds

represented by the formula (I) of the present invention and synthetic intermediates thereof. The above schemes can be changed or modified into schemes which a person ordinarily skilled in the art can easily understand.

[0035]

5 In the above schemes, when a protective group is necessary based on variation of functional group, operations of introduction and deprotection can also be conducted optionally in combination according to a general method.

[0036]

10 Compounds represented by the formula (I) of the present invention and intermediates thereof can also be isolated and purified, if required, according to conventional isolation and purification techniques well known to a person ordinarily skilled in the art in the relevant field, such as solvent extraction, crystallization, recrystallization, chromatography, preparative high performance liquid chromatography or the like.

15 [0037]

 The compounds of the present invention have an excellent NK₁ receptor antagonist activity, and thus can also be used as an agent for the prevention or treatment of various diseases mediated by NK₁ receptor. For example, the compounds of the present invention are useful as antiemetic agent, especially useful as preventive agent of
20 cancer-chemotherapy (for example, cisplatin)-induced gastrointestinal symptom (for example, nausea and vomiting). Preferable compounds of the present invention are not only useful for acute cancer-chemotherapy-induced nausea and vomiting but also delayed cancer-chemotherapy-induced nausea and vomiting.

[0038]

25 In an embodiment, the compounds of the present invention have an excellent NK₁ receptor antagonist activity, and thus can also be used as an agent for the prevention of postoperative nausea and vomiting (PONV), nausea and vomiting

associated with radiotherapy, morphine-induced vomiting or motion sickness, and the treatment of schizophrenia, social phobia, anxiety and depression, alcoholism, irritable bowel syndrome, ulcerative colitis, coughing, asthma, atopic dermatitis, psoriasis, pruritus, pain, migraine, tinnitus, benign prostatic hyperplasia, overactive bladder or
5 urinary incontinence.

[0039]

Pharmaceutical compositions of the present invention can be administered in various dosage forms depending on their usages. As such dosage forms, for example, powders, granules, fine granules, dry syrups, tablets, capsules, injections, liquids,
10 ointments, suppositories and poultices can be illustrated, which are administered orally or parenterally.

[0040]

Pharmaceutical compositions of the present invention can be prepared by using a compound represented by the formula (I) or a pharmaceutically acceptable salt thereof
15 and at least one of a pharmaceutical additive. These pharmaceutical compositions can be formulated by admixing, diluting or dissolving with appropriate pharmaceutical additives such as excipients, disintegrants, binders, lubricants, diluents, buffers, tonicity agents, preservatives, wetting agents, emulsifying agents, dispersing agents, stabilizing agents, solubilizing agents and the like, according to a conventional formulation
20 procedure depending upon their dosage forms.

[0041]

When a pharmaceutical composition of the present invention is used in the prevention or treatment, the dosage of a compound represented by the formula (I) or a pharmaceutically acceptable salt thereof as the active ingredient is appropriately decided
25 to depend on the age, sex, body weight, degree of disorders and treatment of each patient and the like. The dosage for an adult can be decided within the range of, for example, 1 to 1000 mg per day, 0.1 to 500 mg per day, 0.1 to 100 mg per day, or 0.1 to

50 mg per day in the case of oral administration, and the daily dose can be divided into one, two, three or four times per day and administered. And, the dosage for an adult can be decided within the range of, for example, 1 to 1000 mg per day, 0.1 to 500 mg per day, 0.1 to 100 mg per day, or 0.1 to 50 mg per day in the case of parenteral administration, and the daily dose can be divided into one, two, three or four times per day and administered.

[0042]

When a pharmaceutical composition of the present invention is used in the prevention of cancer-chemotherapy-induced nausea and vomiting, this pharmaceutical can also be administered before administration of antineoplastic agents. For example, the pharmaceutical can be administered immediately before administration to before an hour and a half of the administration in chemotherapy, and after the second day, the pharmaceutical can also be administered in the morning.

[0043]

In an embodiment, a compound represented by the formula (I) of the present invention or a pharmaceutically acceptable salt thereof can also be used in combination with any other medicament other than NK_1 receptor antagonists. As such other medicaments used in combination, for example, corticosteroid and 5-HT₃ receptor antagonist antiemetic agent can be illustrated.

[0044]

When a compound represented by the formula (I) of the present invention or a pharmaceutically acceptable salt thereof are used in combination with the other medicament, it can be administered as a formulation comprising together with there active ingredients or as formulations each of which is separately formulated from each active ingredient. When separately formulated, these formulations can be administered separately or concurrently.

Furthermore, the dosage of the compound represented by the formula (I) of the

present invention can be reduced depending on the dosage of the other medicaments used in combination.

EXAMPLES

5 [0045]

The present invention is further illustrated in more detail by way of the following Reference Examples, Examples and Test Examples. However, the present invention is not limited thereto.

[0046]

10 Reference Example 1

(3-Methylpiperidin-4-yl)acetic acid ethyl ester

To a suspension of sodium hydride (60%, 0.17 g) in tetrahydrofuran (5 mL) was added triethyl phosphonoacetate (1.04 g) under ice-cooling, and the mixture was stirred at room temperature for 1 hour. To the reaction mixture was added a solution of
15 3-methyl-4-oxopiperidine-1-carboxylic acid *tert*-butyl ester (0.50 g) in tetrahydrofuran (5 mL) at room temperature, and the mixture was stirred at 50°C for 2.5 hours. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under
20 reduced pressure. The obtained residue was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate) to give 4-ethoxycarbonylmethylene-3-methylpiperidine-1-carboxylic acid *tert*-butyl ester (0.65 g). To a solution of the obtained compound (0.65 g) in ethanol (12 mL) was added 10% palladium on carbon (250 mg, wet) at room temperature, and the mixture was stirred at room temperature
25 under a hydrogen gas atmosphere for 18 hours. The reaction mixture was diluted with ethyl acetate and the resulting mixture was stirred at room temperature for 15 minutes. The reaction mixture was filtered through a Celite pad, and the filtrate was concentrated

under reduced pressure to give 4-ethoxycarbonylmethyl-3-methylpiperidine-1-carboxylic acid *tert*-butyl ester (0.64 g). To a solution of the obtained compound (0.64 g) in ethyl acetate (10 mL) was added 4 mol/L hydrochloric acid (ethyl acetate solution, 10 mL) at room temperature, and the mixture was stirred at room temperature for 39
5 hours. The reaction mixture was concentrated under reduced pressure, and to the residue was added a saturated aqueous sodium hydrogen carbonate solution. The resulting mixture was extracted with a mixed solvent of dichloromethane-isopropyl alcohol (dichloromethane/isopropyl alcohol=3/1) twice. The combined organic layer was dried over anhydrous sodium sulfate, and the solvent was removed under reduced
10 pressure to give the title compound (0.39 g).

[0047]

Reference Examples 2 and 3

The compounds of Reference Examples 2 to 3 were prepared in a similar manner to that described in Reference Example 1 using the corresponding starting
15 materials.

[0048]

Reference Example 4

(6-Chloro-4-iodopyridin-3-yl)carbamic acid *tert*-butyl ester

To a solution of (6-chloropyridin-3-yl)carbamic acid *tert*-butyl ester (5.0 g) and
20 N, N, N', N'-tetramethylethane-1,2-diamine (7.7 g) in diethyl ether (120 mL) was added dropwise *n*-butyllithium (2.65 mol/L *n*-hexane solution, 25 mL) at -78°C under an argon gas atmosphere. After the mixture was stirred at -10°C for 2 hours, a solution of iodine (11.4 g) in diethyl ether (40 mL) was added dropwise at -78°C. The resulting mixture was stirred at room temperature for 1 day. To the reaction mixture was added
25 a saturated aqueous ammonium chloride solution, and the resulting mixture was extracted with diethyl ether. The organic layer was washed with 10% aqueous sodium pyrosulfite and brine, and dried over anhydrous magnesium sulfate, and the solvent was

removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate) to give the title compound (2.59 g).

[0049]

5 Reference Example 5

(6-Chloro-4-iodopyridin-3-yl)methylcarbamic acid *tert*-butyl ester

To a solution of (6-Chloro-4-iodopyridin-3-yl)carbamic acid *tert*-butyl ester (2.59 g) in N,N-dimethylformamide (30 mL) was added sodium hydride (60%, 0.32 g) under ice-cooling, and the mixture was stirred at room temperature for 30 minutes. To
10 the mixture was added iodomethane (2.60 g) under ice-cooling and the mixture was stirred at room temperature overnight. To the reaction mixture was added a saturated aqueous sodium hydrogen carbonate solution, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure.
15 The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate) to give the title compound (2.66 g).

[0050]

Reference Example 6

20 (6-Chloro-4-iodopyridin-3-yl)methylamine

To a solution of (6-Chloro-4-iodopyridin-3-yl)methylcarbamic acid *tert*-butyl ester (2.66 g) in dichloromethane (10 mL) was added trifluoroacetic acid (8.23 g) under ice-cooling, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was concentrated under reduced pressure, and to the residue was added a
25 saturated aqueous sodium carbonate solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure to give the

title compound (1.89 g).

[0051]

Reference Example 7

[6-Chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]methylanine

- 5 To a mixed solution of (6-Chloro-4-iodopyridin-3-yl)methylanine (1.89 g) and 4-fluoro-2-methylphenyl boronic acid (1.30 g) in 1,2-dimethoxyethane (20 mL)-water (20 mL) were added palladium (II) acetate (0.16 g), triphenylphosphine (0.37 g) and sodium carbonate (3.73 g) at room temperature, and the mixture was stirred at 90°C overnight. The reaction mixture was cooled to room temperature and water was added.
- 10 The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate) to give the title compound (1.56 g).

15 [0052]

Reference Example 8

(6-Chloro-4-*ortho*-tolylpyridin-3-yl)methylanine

- To a mixed solution of (6-Chloro-4-iodopyridin-3-yl)methylanine (0.70 g) and 2-methylphenyl boronic acid (0.42 g) in 1,2-dimethoxyethane (10 mL)-water (10 mL)
- 20 were added palladium (II) acetate (0.058 g), triphenylphosphine (0.14 g) and sodium carbonate (1.38 g) at room temperature, and the mixture was stirred at 90°C overnight. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was
- 25 removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate) to give the title compound (0.54 g).

[0053]

Reference Example 9

2-(3,5-Bistrifluoromethylphenyl)-N-[6-chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide

- 5 To a solution of 2-(3,5-bistrifluoromethylphenyl)-2-methylpropionic acid (0.66 g) in dichloromethane (10 mL) was added oxalyl chloride (0.56 g) and N,N-dimethylformamide (2 drops) at room temperature, and the mixture was stirred at the same temperature for 1 hour. The reaction mixture was concentrated under reduced pressure to give the residue. Under an argon gas atmosphere, to a solution of [6-
10 Chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]methylamine (0.50 g) in tetrahydrofuran (10 mL) was added potassium bis(trimethylsilyl)amide (0.500 mol/L toluene solution, 5.0 mL) under ice-cooling, and the mixture was stirred at room temperature for 30 minutes. To the reaction mixture was added dropwise a solution of the above residue in tetrahydrofuran (5 mL) under ice-cooling, and the mixture was
15 stirred at room temperature for 2 hours. To the reaction mixture was added 1.0 mol/L aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel
20 (eluent: *n*-hexane/ethyl acetate) to give the title compound (1.03 g).

[0054]

Reference Example 10

2-(3,5-Bistrifluoromethylphenyl)-N-(6-chloro-4-*ortho*-tolylpyridin-3-yl)-N-methylisobutylamide

- 25 To a solution of 2-(3,5-bistrifluoromethylphenyl)-2-methylpropionic acid (0.77 g) in dichloromethane (12 mL) was added oxalyl chloride (0.65 g) and N,N-dimethylformamide (2 drops) at room temperature, and the mixture was stirred at the

same temperature for 1 hour. The reaction mixture was concentrated under reduced pressure to give the residue. Under an argon gas atmosphere, to a solution of (6-Chloro-4-*ortho*-tolylpyridin-3-yl)methylamine (0.54 g) in tetrahydrofuran (12 mL) was added potassium bis(trimethylsilyl)amide (0.500 mol/L toluene solution, 6.0 mL) under ice-cooling, and the mixture was stirred at room temperature for 30 minutes. To the reaction mixture was added dropwise a solution of the above residue in tetrahydrofuran (6 mL) under ice-cooling, and the mixture was stirred at room temperature for 1 hours. To the reaction mixture was added 1.0 mol/L aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate) to give the title compound (1.0 g).

[0055]

15 Reference Example 11

[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester

A suspension of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.79 g), (piperidin-4-yl)acetic acid ethyl ester (0.38 g), and potassium carbonate (0.41 g) in dimethyl sulfoxide (4.5 mL) was stirred at 180°C under microwave irradiation for 1 hour. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate=100/0-60/40). The obtained material was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl

acetate=100/0-60/40) to give the title compound (0.38 g).

[0056]

Reference Example 12

(5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-*ortho*-tolyl-
5 3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)acetic acid ethyl ester

A suspension of 2-(3,5-bistrifluoromethylphenyl)-N-(6-chloro-4-*ortho*-tolylpyridin-3-yl)-N-methylisobutylamide (0.50 g), (piperidin-4-yl)acetic acid ethyl ester (0.25 g), and potassium carbonate (0.27 g) in dimethyl sulfoxide (3.0 mL) was stirred at 180°C under microwave irradiation for 2 hours. The reaction mixture was
10 cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=100/0-50/50) to give the title compound (0.35 g).

15 [0057]

Reference Example 13

5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methyl phenyl)-3,3-dimethyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester

20 A solution of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.05 g) and (3,3-dimethylpiperidin-4-yl)acetic acid ethyl ester (0.094 g) in 1-methyl-2-pyrrolidone (0.5 mL) was stirred at 180°C under microwave irradiation for 3 hours. The reaction mixture was cooled to room temperature and water was added. The resulting mixture
25 was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on

silica gel (eluent: *n*-hexane/ethyl acetate=100/0-50/50) to give the title compound (0.043 g).

[0058]

Reference Example 14

- 5 5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methyl phenyl)-3-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester

A solution of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.05 g) and (3-methylpiperidin-4-yl)acetic acid ethyl ester (0.087 g) in 1-methyl-2-pyrrolidone (0.5 mL) was stirred at 180°C under microwave irradiation for 1 hour. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude
10 product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=100/0-50/50) to give the title compound (0.040 g).

[0059]

Reference Example 15

- 20 [5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methyl phenyl)-3-methoxyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester

A solution of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.05 g) and (3-methoxypiperidin-4-yl)acetic acid ethyl ester (0.094 g) in 1-methyl-2-pyrrolidone (0.5 mL) was stirred at
25 180°C under microwave irradiation for 1 hour. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium

sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=100/0-60/40) to give the title compound (0.025 g).

[0060]

5 Reference Example 16

2-Methyl-2-piperidin-4-ylpropionic acid ethyl ester

To a solution of lithium diisopropylamide (1.09 mol/L tetrahydrofuran/*n*-hexane solution, 30.0 mL) in tetrahydrofuran (40 mL) was added dropwise a solution of ethyl isobutylate in tetrahydrofuran (20 mL) at -78°C under an argon gas atmosphere, and the mixture was stirred at the same temperature for 1 hour. To the reaction mixture was added a solution of 4-oxopiperidine-1-carboxylic acid benzyl ester (5.83 g) in tetrahydrofuran (50 mL) at -78°C, and the mixture was stirred at room temperature overnight. To the reaction mixture was added a saturated aqueous ammonium chloride solution, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=90/10-10/90) to give 4-(1-ethoxycarbonyl-1-methylethyl)-4-hydroxypiperidine-1-carboxylic acid benzyl ester (6.13 g). To a solution of the obtained compound (6.13 g) in toluene (100 mL) was added (methoxycarbonylsulfamoyl)triethylammonium hydroxide inner salt (5.00 g) at room temperature, and the mixture was stirred at 90°C for 2 hours. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution, water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=100/0-50/50) to give 4-(1-ethoxycarbonyl-1-

methylethyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid benzyl ester (5.85 g). Under a hydrogen gas atmosphere, a mixture of the obtained compound (5.85 g) and Pearlman's catalyst (0.600 g) in methanol (65 mL) and ethyl acetate (65 mL) was stirred at room temperature overnight. The reaction mixture was filtered through a Celite pad, and the
 5 filtrate was concentrated under reduced pressure to give the title compound (3.18 g).

[0061]

Reference Example 17

2-Piperidin-4-ylpropionic acid ethyl ester

To a suspension of sodium hydride (60%, 0.78 g) in N,N-dimethylformamide
 10 (20 mL) was added 2-(diethoxyphosphoryl)propionic acid ethyl ester (4.44 g) under ice-cooling, and the mixture was stirred at the same temperature for 30 minutes. To the reaction mixture was added a solution of 4-oxopiperidine-1-carboxylic acid benzyl ester (3.50 g) in N,N-dimethylformamide (10 mL) under ice-cooling, and the mixture was stirred at room temperature for 1 hour. To the reaction mixture was added water, and
 15 the resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=100/0-50/50) to give 4-(1-ethoxycarbonylethylidene)piperidine-1-carboxylic acid benzyl ester (4.60 g). Under a
 20 hydrogen gas atmosphere, a mixture of the obtained compound (4.60 g) and 10% palladium on carbon (500 mg, wet) in methanol (50 mL) was stirred at room temperature for 2 hours. The reaction mixture was filtered through a Celite pad, and the filtrate was concentrated under reduced pressure to give the title compound (2.85 g).

[0062]

25 Reference Example 18

(4-Methylpiperidin-4-yl)acetic acid ethyl ester

To a solution of 4-hydroxymethyl-4-methyl piperidine-1-carboxylic acid *tert*-

butyl ester (0.67 g) and triethylamine (0.44 g) in dichloromethane (15 mL) was added methanesulfonyl chloride (0.40 g) at room temperature, and the mixture was stirred at the same temperature for 3 hours. The reaction mixture was washed with water and brine, and dried over anhydrous magnesium sulfate. The solvent was removed under reduced pressure to give 4-methanesulfonyloxymethyl-4-methylpiperidine-1-carboxylic acid *tert*-butyl ester (0.85 g). To a solution of the obtained compound (0.85 g) in *N,N*-dimethylformamide (6 mL) was added sodium cyanide (0.27 g) at room temperature, and the mixture was stirred at 50°C for 5 hours and at 80°C for 13 hours. To the reaction mixture were added sodium cyanide (0.22 g) and sodium iodide (0.02 g), and the mixture was stirred at 120°C for 8 hours. To the reaction mixture was added sodium cyanide (0.72 g), and the mixture was stirred at 140°C for 17 hours. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with a mixed solvent of *n*-hexane and ethyl acetate (*n*-hexane/ethyl acetate=1/4). The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=80/20-50/50) to give 4-cyanomethyl-4-methylpiperidine-1-carboxylic acid *tert*-butyl ester (0.53 g). A mixture of the obtained compound (0.53 g) and concentrated hydrochloric acid (12 mL) was stirred at 110°C for 47 hours. The reaction mixture was cooled to room temperature, and to the mixture were added water (24 mL), aqueous sodium hydroxide solution (2.0 mol/L, 45 mL) and di-*tert*-butyl dicarbonate (0.51 g), and the resulting mixture was stirred at room temperature for 21 hours. The solvent was removed under reduced pressure, and to the residue were added water and hydrochloric acid (2.0 mol/L, 3 mL), and the resulting mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was removed under reduced pressure to give 4-carboxymethyl-4-methylpiperidine-1-carboxylic acid *tert*-butyl ester (0.34 g). To a

solution of the obtained compound (0.34 g) in N,N-dimethylformamide (5 mL) were added potassium carbonate (0.27 g) and ethyl iodide (0.41 g) at room temperature, and the resulting mixture was stirred at the same temperature for 22 hours. To the reaction mixture was added water, and the resulting mixture was extracted with diethyl ether.

- 5 The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=100/0-60/40) to give 4-ethoxycarbonylmethyl-4-methylpiperidine-1-carboxylic acid *tert*-butyl ester (0.28 g). To a solution of the obtained compound
 10 (0.28 g) in 1, 4-dioxane (5 mL) was added hydrogen chloride (4.0 mol/L 1, 4-dioxane solution, 5.0 mL) at room temperature, and the mixture was stirred at the same temperature for 26 hours. The solvent was removed under reduced pressure, and to the residue was added saturated aqueous sodium hydrogen carbonate, and the resulting mixture was extracted with a mixed solvent of dichloromethane/isopropyl alcohol
 15 (dichloromethane/isopropyl alcohol=3/1) twice. The combined organic layer was dried over anhydrous sodium sulfate, and the solvent was removed under reduced pressure to give the title compound (0.17 g).

[0063]

Reference Example 19

20 6-Chloro-3-nitropyridine-2-carbonitrile

- To a solution of 2, 6-dichloro-3-nitropyridine (2.50 g) in N-methylpyrrolidone (25 mL) was added copper (I) cyanide (2.32 g) at room temperature, and the mixture was stirred at 180°C for 1 hour. The reaction mixture was cooled to room temperature, and to the mixture were added ethyl acetate and water. The mixture was filtered through
 25 a Celite pad. The filtrate was washed with brine, and the separated aqueous layer was re-extracted with ethyl acetate. The combined organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed

under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=90/10-70/30) to give the title compound (0.90 g).

[0064]

5 Reference Example 20

3-Amino-6-chloropyridine-2-carbonitrile

To a solution of 6-chloro-3-nitropyridine-2-carbonitrile (0.32 g) and concentrated hydrochloric acid (1.2 mL) in ethanol (3.6 mL) was added iron powder (0.34 g) at room temperature, and the mixture was heated at reflux for 30 minutes. The
10 reaction mixture was cooled to room temperature, and basified by addition of saturated aqueous sodium hydrogen carbonate solution. To the mixture was added ethyl acetate, and the resulting mixture was filtered through a Celite pad, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and brine, dried over anhydrous magnesium sulfate. The solvent was removed under reduced pressure to
15 give the title compound (0.24 g).

[0065]

Reference Example 21

3-Amino-4-bromo-6-chloropyridine-2-carbonitrile

To a solution of 3-amino-6-chloropyridine-2-carbonitrile (0.24 g) in *N,N*-
20 dimethylformamide (8 mL) was added *N*-bromosuccinimide (0.37 g) at room temperature, and the mixture was stirred at the same temperature overnight. To the reaction mixture was added saturated aqueous sodium thiosulfate solution, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under
25 reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate=75/25-50/50) to give the title compound (0.30 g).

[0066]

Reference Example 22

3-Amino-6-chloro-4-(4-fluoro-2-methylphenyl)pyridine-2-carbonitrile

A mixture of 3-amino-4-bromo-6-chloropyridine-2-carbonitrile (0.15 g), 4-fluoro-2-methylphenylboronic acid (0.08 g), tetrakis(triphenylphosphine)palladium(0) (0.07 g), sodium carbonate (0.20 g), 1, 2-dimethoxyethane (3.2 mL) and water (0.8 mL) was stirred at 100°C under microwave irradiation for 1 hour. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate=50/50-0/100) to give the title compound (0.14 g).

[0067]

15 Reference Example 23

2-(3,5-Bistrifluoromethylphenyl)-N-[6-chloro-2-cyano-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]isobutylamide

To a solution of 2-(3,5-bistrifluoromethylphenyl)-2-methylpropionic acid (0.31 g) in dichloromethane (2.6 mL) were added oxalyl chloride (0.26 g) and N,N-dimethylformamide (2 drops) at room temperature, and the mixture was stirred at the same temperature for 1 hour. The reaction mixture was concentrated under reduced pressure to give the residue. To a solution of 3-amino-6-chloro-4-(4-fluoro-2-methylphenyl)pyridine-2-carbonitrile (0.14 g) in tetrahydrofuran (5 mL) was added sodium bis(trimethylsilyl)amide (1.0 mol/L tetrahydrofuran solution, 1.1 mL) under ice-cooling, and the mixture was stirred at the same temperature for 30 minutes. To the reaction mixture was added dropwise a solution of the above residue in tetrahydrofuran (2.0 mL) under ice-cooling, and the mixture was stirred at room temperature for 30

minutes. To the reaction mixture was added a saturated aqueous sodium hydrogen carbonate solution, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate=85/15-40/60) to give the title compound (0.21 g).

[0068]

Reference Example 24

2-(3,5-Bistrifluoromethylphenyl)-N-[6-chloro-2-cyano-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide

To a solution of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-2-cyano-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]isobutylamide (0.21 g) in N,N-dimethylformamide (2.4 mL) was added sodium hydride (60 %, 0.018 g) under ice-cooling, and the mixture was stirred at the same temperature for 5 minutes. To the reaction mixture was added iodomethane (0.11 g) under ice-cooling and the mixture was stirred at room temperature overnight. To the reaction mixture was added water, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=90/10-50/50) to give the title compound (0.09 g).

[0069]

Reference Example 25

[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-6'-cyano-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester

A suspension of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-2-cyano-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.03 g), piperidin-4-yl

acetic acid ethyl ester (0.05 g) and potassium carbonate (0.02 g) in dimethyl sulfoxide (1.0 mL) was stirred at 100°C under microwave irradiation for 1 hour. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate=80/20-20/80) to give the title compound (0.02 g).

[0070]

10 Reference Example 26

2-[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]-2-methylpropionic acid ethyl ester

A mixture of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.15 g), 2-methyl-2-piperidin-4-yl propionic acid ethyl ester (0.28 g) and N-methylpyrrolidone (1.5 mL) was stirred at 190°C under microwave irradiation for 1 hour. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=100/0-60/40) to give the title compound (0.14 g).

[0071]

Reference Example 27

25 2-[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]propionic acid ethyl ester

A suspension of 2-(3,5-bis(trifluoromethyl)phenyl)-N-[6-chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.48 g), 2-piperidin-4-yl propionic acid ethyl ester (0.83 g) and potassium carbonate (0.25 g) in N-methylpyrrolidone (3.6 mL) was stirred at 190°C under microwave irradiation for 1 hour. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=100/0-50/50) to give the title compound (0.51 g).

[0072]

Reference Examples 28 and 29

N-[4-[(*S*)-2-((*S*)-4-Benzyl-2-oxo-oxazolidin-3-yl)-1-methyl-2-oxo-ethyl]-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl]-2-(3,5-bis(trifluoromethyl)phenyl)-N-methylisobutylamide (Reference Examples 28) and N-[4-[(*R*)-2-((*S*)-4-Benzyl-2-oxo-oxazolidin-3-yl)-1-methyl-2-oxo-ethyl]-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl]-2-(3,5-bis(trifluoromethyl)phenyl)-N-methylisobutylamide (Reference Examples 29)

A mixture of 2-[5'-{[2-(3,5-bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]propionic acid ethyl ester (0.50 g), aqueous sodium hydroxide solution (1.0 mol/L, 2.0 mL), tetrahydrofuran (2 mL) and methanol (6 mL) was stirred at 140°C under microwave irradiation for 2 hours. The reaction mixture was cooled to room temperature and hydrochloric acid (1.0 mol/L, 3.0 mL) was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography

on silica gel (eluent: *n*-hexane/ethyl acetate=50/50-0/100) to give 2-[5'-{[2-(3,5-bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]propionic acid (0.42 g). To a solution of (*S*)-4-benzylloxazolidin-2-one (0.06 g) in tetrahydrofuran (4 mL) was added dropwise *n*-butyllithium (2.65 mol/L in *n*-hexane solution, 0.12 mL) at -78°C under an argon gas atmosphere, and the mixture was stirred at the same temperature for 30 minutes to give lithium solution of (*S*)-4-benzylloxazolidin-2-one. Under an argon gas atmosphere, to a solution of 2-[5'-{[2-(3,5-bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]propionic acid (0.20 g) and triethylamine (0.04 g) in diethyl ether (4 mL) was added pivaloyl chloride (0.04 g) under ice-cooling, and the mixture was stirred at the same temperature for 1 hour. To the reaction mixture was added dropwise the above lithium solution at -78°C, and the mixture was stirred under ice-cooling for 1 hour. To the reaction mixture was added a saturated aqueous ammonium chloride solution, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=90/10-10/90). Then, the obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=100/0-50/50). Then, the obtained crude product was purified by preparative thin-layer chromatography (silica gel thickness: 0.5 mm, eluent: *n*-hexane/ethyl acetate=2/1) to give Reference Examples 28 (0.09 g) and Reference Examples 29 (0.09 g). In the above chromatography, the compound of Reference Examples 28 was in the high polarity side, and the compound of Reference Examples 29 was in the low polarity side.

[0073]

Reference Example 30

2-[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-6'-cyano-4'-(4-fluoro-2-methyl-phenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]propionic acid ethyl ester

A suspension of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-2-cyano-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.06 g), 2-piperidin-4-yl propionic acid ethyl ester (0.10 g) and potassium carbonate (0.03 g) in dimethyl sulfoxide (1.0 mL) was stirred at 180°C under microwave irradiation for 2 hours. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate=90/10-65/35) to give the title compound (0.05 g).

[0074]

15 Reference Example 31

5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methyl phenyl)-4-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester

A mixture of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.05 g), (4-methylpiperidin-4-yl)acetic acid ethyl ester (0.09 g) and N-methylpyrrolidone (0.5 mL) was stirred at 190°C under microwave irradiation for 30 minutes. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=90/10-60/40) to give the title compound (0.03 g).

[0075]

Reference Example 32

[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-6'-cyano-4'-(4-fluoro-2-methylphenyl)-4-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester

A suspension of 2-(3,5-bistrifluoromethylphenyl)-N-[6-chloro-2-cyano-4-(4-fluoro-2-methylphenyl)pyridin-3-yl]-N-methylisobutylamide (0.06 g), (4-methylpiperidin-4-yl)acetic acid ethyl ester (0.09 g) and potassium carbonate (0.03 g) in dimethyl sulfoxide (1.0 mL) was stirred at 180°C under microwave irradiation for 1 hour. The reaction mixture was cooled to room temperature and water was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on aminopropylated silica gel (eluent: *n*-hexane/ethyl acetate=90/10-65/35) to give the title compound (0.06 g).

[0076]

Example 1

[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid

To a mixture of [5'-{[2-(3,5-bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester (0.38 g) in tetrahydrofuran (6 mL), methanol (6 mL) and water (2 mL) was added lithium hydroxide monohydrate (0.12 g) at room temperature, and the mixture was stirred at room temperature overnight. The reaction mixture was neutralized by the addition of 2 mol/L hydrochloric acid (1.5 mL), and the resulting mixture was concentrated under reduced pressure. To the residue was added water, and the resulting mixture was extracted with ethyl acetate. The organic

layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure to give the title compound (0.37 g).

[0077]

Example 2

5 (5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-*ortho*-tolyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)acetic acid

To a mixture of [5'-{[2-(3,5-bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-*ortho*-tolyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester (0.30 g) in tetrahydrofuran (4 mL), methanol (2 mL) and
10 water (2 mL) was added lithium hydroxide monohydrate (0.084 g) at room temperature, and the mixture was stirred at room temperature for 3 hours. The reaction mixture was neutralized by the addition of 2 mol/L hydrochloric acid (1.1 mL), and the resulting mixture was concentrated under reduced pressure. To the residue was added water, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed
15 with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure to give the title compound (0.29 g).

[0078]

Example 3

5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methyl phenyl)-3,3-dimethyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid
20

To a mixture of [5'-{[2-(3,5-bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,3-dimethyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester (0.043 g) in tetrahydrofuran (1 mL), methanol (0.5 mL) and water (0.5 mL) was added lithium hydroxide
25 monohydrate (0.012 g) at room temperature, and the mixture was stirred at room temperature for 3 hours. The reaction mixture was neutralized by the addition of 2 mol/L hydrochloric acid (0.14 mL). Water was added to the mixture, and the resulting

mixture was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure to give the title compound (0.033 g).

[0079]

5 Example 4

5'-{[2-(3,5-Bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methyl phenyl)-3-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid

To a mixture of [5'-{[2-(3,5-bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester (0.040 g) in tetrahydrofuran (1 mL), ethanol (0.5 mL) and water (0.5 mL) was added lithium hydroxide monohydrate (0.011 g) at room temperature, and the mixture was stirred at room temperature for 3 hours. The reaction mixture was neutralized by the addition of 2 mol/L hydrochloric acid (0.13 mL). Water was added to the mixture, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure to give the title compound (0.035 g).

[0080]

Example 5

20 [5'-{[2-(3,5-Bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methyl phenyl)-3-methoxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid

To a mixture of [5'-{[2-(3,5-bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3-methoxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester (0.025 g) in tetrahydrofuran (1 mL), ethanol (0.5 mL) and water (0.5 mL) was added lithium hydroxide monohydrate (0.006 g) at room temperature, and the mixture was stirred at room temperature for 3 hours. The reaction mixture was neutralized by the addition of 2

mol/L hydrochloric acid (0.075 mL). Water was added to the mixture, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure to give the title compound (0.021 g).

5 [0081]

Example 6

[5'-{[2-(3,5-Bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-6'-cyano-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid

To a mixture of [5'-{[2-(3,5-bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-6'-cyano-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester (0.02 g) in tetrahydrofuran (0.30 mL), methanol (0.15 mL) and water (0.15 mL) was added lithium hydroxide monohydrate (0.007 g) at room temperature, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was neutralized by the addition of acetic acid. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure to give the title compound (0.02 g).

[0082]

Example 7

20 2-[5'-{[2-(3,5-Bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]-2-methylpropionic acid

A mixture of 2-[5'-{[2-(3,5-bis(trifluoromethyl)phenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]-2-methylpropionic acid ethyl ester (0.14 g), aqueous sodium hydroxide solution (1.0 mol/L, 0.60 mL), tetrahydrofuran (0.6 mL) and methanol (1.8 mL) was stirred at 140°C under microwave irradiation for 4.5 hours. To the reaction

mixture was added aqueous sodium hydroxide solution (2.0 mol/L, 0.50 mL), and the mixture was stirred at 140°C under microwave irradiation for 2 hours. The reaction mixture was cooled to room temperature and hydrochloric acid (1.0 mol/L, 2.0 mL) was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure to give the title compound (0.11 g).

[0083]

Example 8

2-[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-6'-cyano-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]propionic acid

To a mixture of 2-[5'-{[2-(3,5-bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-6'-cyano-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]propionic acid ethyl ester (0.03 g) in tetrahydrofuran (0.4 mL), methanol (0.2 mL) and water (0.2 mL) was added lithium hydroxide monohydrate (0.008 g) at room temperature, and the mixture was stirred at room temperature overnight. The reaction mixture was neutralized by the addition of acetic acid. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate/methanol=50/50/0-0/100/0-0/90/10) to give the title compound (0.02 g).

[0084]

Example 9

5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-4-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid

To a mixture of [5'-{[2-(3,5-bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-4-methyl-3,4,5,6-

tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester (0.03 g), tetrahydrofuran (0.375 mL), methanol (0.375 mL) and water (0.15 mL) was added lithium hydroxide monohydrate (0.02 g) at room temperature, and the mixture was stirred at room temperature for 3 hours and then at 50°C for 4 hours. The reaction mixture was cooled to room temperature and hydrochloric acid (2.0 mol/L, 0.2 mL) was added. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, and dried over anhydrous sodium sulfate, and the solvent was removed under reduced pressure to give the title compound (0.025 g).

[0085]

10 Example 10

[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-6'-cyano-4'-(4-fluoro-2-methylphenyl)-4-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid

To a mixture of [5'-{[2-(3,5-bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-6'-cyano-4'-(4-fluoro-2-methylphenyl)-4-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]acetic acid ethyl ester (0.06 g), tetrahydrofuran (0.50 mL), methanol (0.25 mL) and water (0.25 mL) was added lithium hydroxide monohydrate (0.02 g) at room temperature, and the mixture was stirred at room temperature for 1 hour, and then stirred at 50°C for 3 hours. The reaction mixture was cooled to room temperature and neutralized by the addition of acetic acid. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate/methanol=20/80/0-0/100/0-0/90/10) to give the title compound (0.03 g).

[0086]

Example 11

(*S*)-2-[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]propionic acid

To a mixture of N-[4-[(*S*)-2-((*S*)-4-Benzyl-2-oxo-oxazolidin-3-yl)-1-methyl-2-oxo-ethyl]-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl]-2-(3,5-bistrifluoromethylphenyl)-N-methylisobutylamide (0.08 g) in tetrahydrofuran (1.5 mL) and water (0.5 mL) were added lithium hydroxide monohydrate (0.008 g) and hydrogen peroxide solution (30%, 0.06 mL) under ice-cooling, and the mixture was stirred at the same temperature overnight. To the reaction mixture was added aqueous sodium sulfite solution (10%, 0.75 mL), and the resulting mixture was stirred for 30 minutes. The solvent was removed under reduced pressure. The residue was acidified with hydrochloric acid and the resulting mixture was extracted with ethyl acetate. The aqueous layer was re-extracted with ethyl acetate. The combined organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=90/10-10/90) to give the title compound (0.014 g).

[0087]

Example 12

(*R*)-2-[5'-{[2-(3,5-Bistrifluoromethylphenyl)-2-methylpropionyl]methylamino}-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl]propionic acid

To a mixture of N-[4-[(*R*)-2-((*S*)-4-Benzyl-2-oxo-oxazolidin-3-yl)-1-methyl-2-oxo-ethyl]-4'-(4-fluoro-2-methylphenyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl]-2-(3,5-bistrifluoromethylphenyl)-N-methylisobutylamide (0.04 g) in tetrahydrofuran (0.75 mL) and water (0.25 mL) were added lithium hydroxide monohydrate (0.004 g) and hydrogen peroxide solution (30%, 0.03 mL) under ice-cooling, and the mixture was stirred under the same temperature overnight. To the reaction mixture was added aqueous sodium sulfite solution (10%, 0.375 mL), and the resulting mixture was stirred









for 30 minutes. The solvent was removed under reduced pressure. The residue was acidified with hydrochloric acid and the resulting mixture was extracted with ethyl acetate. The aqueous layer was re-extracted with ethyl acetate. The combined organic layer was washed with brine, and dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The obtained crude product was purified by column chromatography on silica gel (eluent: *n*-hexane/ethyl acetate=90/10-10/90) to give the title compound (0.004 g).

[0088]

Tables 1 to 7 show the chemical structures of the above compounds of Reference Examples 1 to 32, and the chemical structures and the physical properties of the above compounds of Examples 1 to 12. The abbreviations in these Tables: "Ref No.", "Ex No.", "Str.", "Physical data", "¹H-NMR", "DMSO-d₆" and "CDCl₃" represent Reference Example number, Example number, chemical structure, physical property, hydrogen nuclear magnetic resonance spectrum, dimethylsulfoxide-d₆ and chloroform-d₁, respectively. And, "MS" and "ESI_APCI" represent mass spectrometry and measurement of Electrospray ionization-Atmospheric pressure chemical ionization, respectively.








[0089]

[Table 1]

Ref. No.	Str.	Ref. No.	Str.
1		2	
3		4	
5		6	
7		8	

[0090]

[Table 2]

Ref. No.	Str.	Ref. No.	Str.
9		10	
11		12	
13		14	
15			






[0091]

[Table 3]

Ref. No.	Str.	Ref. No.	Str.
16		17	
18		19	
20		21	
22		23	
24		25	
26		27	







[0092]

[Table 4]


Ex. No.	Str.	Physical data
1		$^1\text{H-NMR } \delta \text{ ppm (DMSO-}d_6\text{): 1.00-1.60 (9H, m), 1.60-1.80 (2H, m), 1.80-2.00 (1H, m), 2.00-2.40 (7H, m), 2.70-2.90 (2H, m), 4.20-4.40 (2H, m), 6.66 (1H, s), 6.90-7.20 (3H, m), 7.60-7.80 (2H, m), 7.87 (1H, s), 8.03 (1H, s), 12.09 (1H, brs)}$ MS (ESI_APCI, m/z): 640 (M+H) ⁺
2		$^1\text{H-NMR } \delta \text{ ppm (DMSO-}d_6\text{): 1.00-1.60 (9H, m), 1.60-1.80 (2H, m), 1.80-2.00 (1H, m), 2.00-2.40 (7H, m), 2.70-2.90 (2H, m), 4.20-4.40 (2H, m), 6.65 (1H, s), 7.00-7.40 (4H, m), 7.60-7.85 (2H, m), 7.86 (1H, s), 8.03 (1H, s), 12.09 (1H, brs)}$ MS (ESI_APCI, m/z): 622 (M+H) ⁺
3		$^1\text{H-NMR } \delta \text{ ppm (DMSO-}d_6\text{): 0.70-2.80 (25H, m), 3.90-4.10 (1H, m), 4.20-4.50 (1H, m), 6.67 (1H, s), 6.90-7.30 (3H, m), 7.60-7.80 (2H, m), 7.82 (1H, s), 8.03 (1H, s), 12.13 (1H, brs)}$ MS (ESI_APCI, m/z): 668 (M+H) ⁺
4		$^1\text{H-NMR } \delta \text{ ppm (DMSO-}d_6\text{): 0.75-3.20 (23H, m), 3.80-4.40 (2H, m), 6.60-6.70 (1H, m), 6.90-7.20 (3H, m), 7.60-7.90 (3H, m), 8.03 (1H, s), 12.10 (1H, brs)}$ MS (ESI_APCI, m/z): 654 (M+H) ⁺
5		$^1\text{H-NMR } \delta \text{ ppm (DMSO-}d_6\text{): 0.75-3.50 (23H, m), 4.10-4.35 (1H, m), 4.40-4.70 (1H, m), 6.60-6.80 (1H, m), 6.90-7.30 (3H, m), 7.60-7.95 (3H, m), 8.03 (1H, s), 12.16 (1H, brs)}$ MS (ESI_APCI, m/z): 670 (M+H) ⁺

[0094]

[Table 6]

Ex. No.	Str.	Physical data
6		$^1\text{H-NMR}$ δ ppm (CDCl_3) : 1.20–1.75 (8H, m), 1.80–1.90 (2H, m), 2.00–2.75 (9H, m), 2.80–3.00 (2H, m), 4.20–4.40 (2H, m), 6.55–6.65 (1H, m), 6.80–7.25 (3H, m), 7.55–7.65 (2H, m), 7.70–7.80 (1H, m); MS (ESI/APCI, m/z) : 665 ($\text{M}+\text{H}$) $^+$
7		$^1\text{H-NMR}$ δ ppm ($\text{DMSO}-d_6$) : 1.04 (6H, s), 1.10–1.85 (11H, m), 2.00–2.40 (4H, m), 2.60–2.80 (2H, m), 4.30–4.50 (2H, m), 6.65 (1H, s), 6.90–7.25 (3H, m), 7.60–7.80 (2H, m), 7.87 (1H, s), 8.03 (1H, s), 12.15 (1H, brs); MS (ESI/APCI, m/z) : 668 ($\text{M}+\text{H}$) $^+$
8		$^1\text{H-NMR}$ δ ppm (CDCl_3) : 1.15–1.95 (14H, m), 2.05–2.30 (3H, m), 2.30–2.40 (1H, m), 2.40–2.70 (3H, m), 2.80–2.95 (2H, m), 4.25–4.45 (2H, m), 6.55–6.65 (1H, m), 6.80–7.25 (3H, m), 7.60–7.75 (2H, m), 7.70–7.80 (1H, m); MS (ESI/APCI, m/z) : 679 ($\text{M}+\text{H}$) $^+$
9		$^1\text{H-NMR}$ δ ppm ($\text{DMSO}-d_6$) : 1.07 (3H, s), 1.10–1.70 (11H, m), 2.00–2.40 (7H, m), 3.35–3.55 (2H, m), 3.55–3.75 (2H, m), 6.65 (1H, s), 6.80–7.30 (3H, m), 7.55–7.85 (2H, m), 7.88 (1H, s), 8.02 (1H, s), 12.02 (1H, brs); MS (ESI/APCI, m/z) : 654 ($\text{M}+\text{H}$) $^+$
10		$^1\text{H-NMR}$ δ ppm (CDCl_3) : 1.15–1.75 (13H, m), 2.05–2.70 (8H, m), 3.40–3.55 (2H, m), 3.65–3.75 (2H, m), 6.55–6.65 (1H, m), 6.80–7.25 (3H, m), 7.55–7.65 (2H, m), 7.75–7.80 (1H, m); MS (ESI/APCI, m/z) : 679 ($\text{M}+\text{H}$) $^+$
11		$^1\text{H-NMR}$ δ ppm ($\text{DMSO}-d_6$) : 1.04 (3H, d, $J=6.8\text{Hz}$), 1.10–1.90 (12H, m), 2.05–2.40 (6H, m), 2.65–2.85 (2H, m), 4.25–4.45 (2H, m), 6.65 (1H, s), 6.85–7.25 (3H, m), 7.60–7.80 (2H, m), 7.87 (1H, s), 8.02 (1H, s), 12.10 (1H, brs); MS (ESI/APCI, m/z) : 654 ($\text{M}+\text{H}$) $^+$

[Table 7]

Ex. No.	Str.	Physical data
12		¹ H-NMR δ ppm (CDCl ₃) : 1.10–2.70 (21H, m), 2.75–2.95 (2H, m), 4.20–4.45 (2H, m), 6.46 (1H, s), 6.80–7.10 (3H, m), 7.65 (2H, s), 7.76 (1H, s), 7.97 (1H, s); MS (ESI/APCI, m/z) : 654 (M+H) ⁺

[0096]

Test Example 1

5 Affinity for human NK₁ receptor(1) Preparation of human NK₁ receptor expression vector

PCR was performed using human adult normal tissue-derived brain cDNA (BioChain) as the template, with the forward primer of SEQ ID NO:1 and the reverse primer of SEQ ID NO:2, using a PCR enzyme, PrimeSTAR Max DNA Polymerase or PrimeSTAR GXL DNA Polymerase (registered trademark, Takara Bio). The amplified product was inserted into a plasmid (pCR-BluntII-TOPO (registered trademark), Life Technologies) using Zero Blunt PCR Cloning Kit (registered trademark, Life Technologies). By a general method, *Escherichia coli* (One Shot TOP10 competent cells, Life Technologies) was transformed by the plasmid into which the amplified product had been inserted. The *Escherichia coli* cells were cultured on an LB agar medium containing 50 µg/mL kanamycin for a day. After the culture, a colony was selected and cultured in an LB medium containing 50 µg/mL kanamycin. After the culture, the plasmid was purified using Quantum Prep Plasmid Miniprep Kit (Bio-Rad). The plasmid was double digested for about two hours using restriction enzymes, *Xho*I and *Hind*III (New England Biolabs). Then, electrophoresis using 1% agarose gel was performed, and the fragment that was cleaved was collected and purified using TaKaRa RICOCHIP (Takara Bio). Separately, a plasmid was also purified from *Escherichia*

coli that had been transformed by a vector (pcDNA3.1(-) (registered trademark), Life Technologies), and the plasmid was double digested for about two hours using restriction enzymes, *Xho*I and *Hind*III (New England Biolabs). Then, electrophoresis using 1% agarose gel was performed, and the vector that was cleaved was collected and purified using TaKaRa RICOCHIP (Takara Bio). The fragment cut out of pCR-Blunt-II and the pcDNA3.1(-) vector treated with the restriction enzymes were ligated using DNA Ligation Kit <Mighty Mix> (Takara Bio). By a general method, *Escherichia coli* (One Shot TOP10 competent cells, Life Technologies) was transformed by the plasmid obtained by the ligation. The *Escherichia coli* cells were cultured on an LB agar medium containing 50 µg/mL ampicillin for a day. After the culture, a colony was selected and cultured in an LB medium containing 50 µg/mL ampicillin, and then the plasmid was purified using Quantum Prep Plasmid Miniprep Kit (Bio-Rad). The protein-encoding nucleotide sequence (SEQ ID NO:3) of the obtained plasmid was completely identical to the nucleotide sequence (NM_001058.3) of human tachykinin receptor 1 (TACR1, NK1R) registered on a known database (NCBI). Therefore, it was confirmed that the cloned gene sequence was the nucleotide sequence of human NK₁ receptor and that the amino acid sequence which would be translated was human NK₁ receptor. The pcDNA3.1(-) (registered trademark) into which the nucleotide sequence of SEQ ID NO:3 was inserted was used as the human NK₁ receptor expression plasmid.

[0097]

(2) Preparation of human NK₁ receptor-expressing cells

(2-1) Culture of 293T cells

Using a liquid D-MEM (Dulbecco's Modified Eagle Medium) medium (low glucose, containing L-glutamine, Wako Pure Chemical Industries) supplemented with an antibiotic penicillin-streptomycin solution (Life Technologies, final penicillin concentration of 100 U/mL and final streptomycin concentration of 100 µg/mL) and fetal bovine serum (final concentration of 10%), 293T cells (RIKEN) were cultured in

an incubator under the condition of 5% CO₂ gas at 37°C.

(2-2) Subculture of 293T cells

Almost confluent cells were washed with PBS (Phosphate Buffered Saline, Wako Pure Chemical Industries), detached using 0.05% trypsin-EDTA (Life Technologies) and suspended in the liquid medium. The cell suspension was diluted with the above liquid medium in such a manner that the spread ratio became 1:10, and then the cells were cultured.

(2-3) Preparation for human NK₁ receptor-expressing cells

Confluent cells were washed with PBS, detached using 0.05% trypsin-EDTA (Life Technologies) and suspended in a liquid D-MEM medium (low glucose, containing L-glutamine, Wako Pure Chemical Industries) supplemented with fetal bovine serum (final concentration of 10%). The cell suspension was diluted with the liquid medium, and the cells were seeded into the wells of a poly-D-lysine-coated 96-well microplate (BD Biocoat (registered trademark), Nippon Becton Dickinson) at a density of 5×10^4 cells/well and a liquid medium volume of 100 µL/well. After seeding, the cells were cultured in an incubator under the condition of 5% CO₂ gas at 37°C for about four to five hours, and the cells to be transfected with the human NK₁ receptor expression plasmid were thus prepared.

(2-4) Transfection of human NK₁ receptor expression plasmid into 293T cells

For the transfection of the human NK₁ receptor expression plasmid, Lipofectamine 2000 (registered trademark, Life Technologies) was used. The human NK₁ receptor expression plasmid was diluted with Opti-MEM (registered trademark) I Reduced-Serum Medium (Life Technologies) to a concentration resulting in 0.2 µg/25 µL/well. At the same time, Lipofectamine 2000 (registered trademark, Life Technologies) was diluted with Opti-MEM (registered trademark) I Reduced-Serum Medium (Life Technologies) to a concentration resulting in 0.4 µL/25 µL/well and incubated at room temperature for five minutes. After five minutes, to form a complex

of human NK₁ receptor expression plasmid/Lipofectamine 2000, the diluted human NK₁ receptor expression plasmid and the diluted Lipofectamine 2000 were mixed and incubated at room temperature for 20 to 25 minutes. After the incubation, 50 µL/well of the complex solution was added to the cells to be transfected with the human NK₁ receptor expression plasmid, and the cells were cultured in an incubator under the condition of 5% CO₂ gas at 37°C for about 48 hours. The cells that were cultured for 48 hours were used for the assays as the human NK₁ receptor-expressing cells.

[0098]

(3) Measurement of binding affinity to human NK₁ receptor

10 (3-1) Preparation of membrane fraction from human NK₁ receptor-expressing cells

Human NK₁ receptor-expressing cells were prepared in a 175cm² culture flask (Nippon Becton Dickinson). The formation of a complex of the human NK₁ receptor expression plasmid and Lipofectamine 2000 was performed by calculating the culture area ratio and increasing the scale of the method described in the above 2-4 by the ratio.

- 15 The human NK₁ receptor-expressing cells were collected in a buffer solution for the membrane fraction preparation (50 mM Tris (Wako Pure Chemical), 120 mM sodium chloride (Wako Pure Chemical Industries), 5 mM potassium chloride (Wako Pure Chemical Industries), 1 mM ethylenediaminetetraacetic acid (Sigma), 0.002 mg/mL chymostatin (Peptide Institute), 0.04 mg/ bacitracin (Wako Pure Chemical Industries),
- 20 0.005 mg/mL phosphoramidon (Peptide Institute) and 0.5 mM phenylmethylsulfonyl fluoride (Wako Pure Chemical Industries), pH7.4) and centrifuged at 1,880 g for 10 minutes, and the cell sediment was suspended in the buffer solution for the membrane fraction preparation. After freezing and thawing the cells once, the cells were homogenized using a Dounce-type homogenizer (cooled on ice, 1000 rpm, 20 times).
- 25 The homogenized cell suspension was centrifuged at 20,000 rpm for 10 minutes, and the supernatant was removed to obtain cell sediment. The cell sediment was suspended again in the buffer solution for the membrane fraction preparation and homogenized

using a Dounce-type homogenizer (cooled on ice, 1000 rpm, 30 times). The cell suspension was centrifuged at 20,000 rpm for 10 minutes, and the supernatant was removed to obtain cell sediment. The same homogenization and centrifugation were repeated again, and final cell sediment was obtained. The final cell sediment was suspended in a buffer solution for the receptor binding test (50 mM Tris (Wako Pure Chemical Industries), 3 mM manganese chloride (Wako Pure Chemical Industries), 0.002 mg/mL chymostatin (Peptide Institute), 0.04 mg/ bacitracin (Wako Pure Chemical Industries) and 0.02% bovine serum albumin (Sigma), pH 7.4), and the protein concentration was measured using BCA Protein Assay Kit (Pierce).

10 (3-2) Receptor binding test

The buffer solution for the receptor binding test was dispensed to the wells of a 96-well assay plate (Greiner) at 22.5 μ L/well. DMSO solutions of a test compound, which were prepared at an 80-time higher concentration using 100% dimethyl sulfoxide (DMSO), were added to the wells at 2.5 μ L/well (final concentrations of 1 nM to 100 nM), and the solutions were mixed. As a radiolabeled ligand, 125 I-substance P (Substance P, [125 I]Tyr⁸-, PerkinElmer) was used. 125 I-substance P was diluted with the buffer solution for the receptor binding test to a concentration resulting in 125 pmol/25 μ L/well and added to the 96-well assay plate, and the solutions were mixed. The membrane fraction prepared from the human NK₁ receptor-expressing cells was diluted with the buffer solution for the receptor binding test to a concentration resulting in 8 to 10 μ g/well, suspended until the suspension became in such a homogenous state that the suspension could flow through a 27G injection needle smoothly and then added to the 96-well assay plate at 150 μ L/well. Then, the plate was incubated at room temperature for 60 minutes while shaking the plate. The reaction solutions were suction-filtered through a multiscreen 96-well filter plate (Millipore) which had been pre-treated with 0.3% polyethyleneimine, and the reaction was terminated by washing with a washing solution (50 mM Tris and 0.02% bovine serum albumin, pH 7.4) four times. The

bottom of the microplate was dried at 60°C, and then 100 µL/well of MicroScint 20 (PerkinElmer) was dispensed to the wells. The top of the plate was sealed with TopSeal A (PerkinElmer), and the plate was shaken for 5 to 10 minutes. Then, the radioactivities were measured with TopCount NXT (registered trademark)

- 5 (PerkinElmer). The radioactivity of each well was calculated by subtracting the radioactivity of the well to which 10 µM aprepitant was added (non-specific binding). The binding rate (%) of ¹²⁵I-substance P = (the radioactivity of the group to which the test compound was added) / (the radioactivity of the group to which the vehicle was added) x100 was calculated. Using analysis software, GraphPad Prism (GraphPad
- 10 Software), the binding rate (%) was plotted against the concentration of the test compound and linearly approximated, and the concentration required for 50% inhibition, IC₅₀, was calculated. These results were shown in Table 8. In the table, Ex. No. means the Example number, and IC₅₀ (nM) is the concentration required for 50% inhibition.

[0099]

15 (4) Result

[Table 8]

E x . N o .	I C ₅₀ (nM)
1	2 . 1 1
2	2 . 5 3
3	4 . 6 1
4	3 . 5 4
5	3 . 2 9
6	2 . 6 0
7	4 . 6 5
8	2 . 2 5
9	2 . 8 3
1 0	3 . 3 2
1 1	1 . 3 2
1 2	7 . 4 4

As shown in Table 8, it was demonstrated that the compounds of the present invention exhibit a high binding affinity for human NK₁ receptor.

[0100]

Test Example 2

Inhibitory effect on human NK₁ receptor

(1) Preparation of human NK₁ receptor-expressing cells

- 5 Human NK₁ receptor-expressing cells were prepared by the same methods as those described in 2-3 and 2-4 of Test Example 1.

[0101]

(2) Study on inhibitory effect on increase in intracellular calcium concentration

- 10 The human NK₁ receptor-expressing cells were washed with 300 μ L/well of a washing solution (20 mM HEPES/Hank's Balanced Salt Solution (HBSS) pH 7.3). A fluorescent calcium indicator (Fluo-4 Direct Calcium Assay Kit, Life Technologies, containing 0.42 mM probenecid and 0.1% bovine serum albumin, prepared according to the protocol of the product) was added to the wells at 150 μ L/well, and the plate was incubated at 37°C for 30 minutes in an incubator. Then, DMSO solutions of a test
- 15 compound, which were prepared at an 80-time higher concentration using 100% dimethyl sulfoxide (DMSO), were added to the wells at 2.5 μ L/well (final concentrations of 0.1, 1 and 10 μ M), and the solutions were mixed. Then, the plate was further incubated at 37°C for 30 minutes in an incubator. After 30 minutes, the intracellular calcium concentrations were measured immediately.

- 20 The intracellular calcium concentrations were each measured as a fluorescent signal using FDSS (registered trademark) 7000 (Hamamatsu Photonics). A substance P (Peptide Institute, Inc.) solution which was prepared at 0.4 μ M or 4 μ M using an assay buffer (20 mM HEPES/Hank's Balanced Salt Solution (HBSS) pH 7.3, containing 0.1% bovine serum albumin) was added automatically to each well at 50 μ L/well (final
- 25 concentration of 0.1 or 1 μ M) 10 seconds after starting reading, and the fluorescent signal was measured up to 120 seconds.

The intracellular calcium concentration (%) of the cells to which a test

compound was added was calculated by the equation below, where the fluorescent signal of the group to which the vehicle (DMSO) was added was regarded as 100%, and the fluorescent signal before the addition of substance P was regarded as 0%.

Intracellular calcium concentration (%) = (Fluorescent signal of test compound addition group) / (Fluorescent signal of vehicle addition group) x100

The intracellular calcium concentration (%) calculated was regarded as the remaining agonist activity of substance P (Substance P-Response Remaining: SPRR). These results were shown in Table 9. In the table, Ex. No. means the Example number. SPRR (%) is the value obtained when the concentration of substance P was 1 μ M and the concentration of the compound was 0.1 μ M.

[0102]

(3) Results

[Table 9]

E x . N o .	S P R R (%)
1	2 8
2	5 4
3	1 3
4	1 6
5	4 9
6	7 3
7	6
8	6 9
9	6
1 1	7

As shown in Table 9, it was demonstrated that the compounds of the present invention exhibit a human NK₁ receptor antagonist activity.

[0103]

Test Example 3

Inhibitory effect on CYP3A4

A dimethyl sulfoxide (DMSO) solution of a test compound with a

concentration 1000 times higher than the evaluation concentration was prepared, and a reaction solution was prepared by diluting the solution. Enzyme reaction was performed by incubating in a potassium phosphate buffer solution (pH 7.4) containing 1 nM to 20 μ M test compound, 3.2 mM magnesium chloride, 0.2 pmol human CYP3A4 (BD Biosciences), 0.5 mM reduced nicotinamide adenine dinucleotide phosphate (NADPH) and 3 μ M Luciferin-IPA (Promega) at 37°C for 10 minutes. The volume of the reaction solution was 50 μ L/well. The 30-minute pre-incubation group was incubated at 37°C for 30 minutes before adding the substrate, the Luciferin-IPA solution (12.5 μ L/well). At the end of the enzyme reaction, 50 μ L/well of a Luciferin detection reagent (Promega) was added to the wells, and the plate was left at room temperature for 20 minutes. Then, the emission intensities were measured with Infinite M1000 (TECAN). The enzyme activities (%) relative to the value of the group to which the test compound was not added were calculated. A dose-response curve was drawn using analysis software, GraphPad Prism (GraphPad Software), and the concentration of each compound that exhibited 50% inhibition, IC_{50} , was calculated. As a comparative example, aprepitant, which is an NK_1 receptor antagonist, was tested in the same manner.

[0104]

IC_{50} values of the 30-minute pre-incubation groups using the test compounds were measured by the above measurement method, and the results are shown in Table 10. In the table, Ex. No. means the Example number, and IC_{50} (μ M) is the concentration required for 50% inhibition.

[Table 10]

E x . N o .	I C ₅₀ (μM)
1	6 . 2
2	1 3
3	3 . 7
4	4 . 8
5	5 . 5
6	2 . 6
7	4 . 1
8	1 . 4
9	6 . 3
1 0	3 . 8
1 1	4 . 9
1 2	4 . 6
Aprepitant	0 . 0 2

It was demonstrated that the CYP3A4-inhibitory activities of the compounds of the present invention are reduced as compared to that of aprepitant. Therefore, it is expected that the compounds of the present invention have fewer drug-drug interactions based on the inhibitory effect on CYP3A4 than aprepitant.

[0105]

Test Example 4

Effect on foot-tapping

(1) Effect on foot-tapping

A male gerbil (Japan SLC) was anesthetized with isoflurane, and 0.3 mg/kg of a test compound was administered from the jugular vein. After four hours, GR73632 (5 pmol/5 μl), which is an NK₁ receptor agonist, was administered into the cerebral ventricle at the part 1 mm lateral to and 4.5 mm below the bregma in the head, under anesthesia with isoflurane. After the administration, the gerbil was moved to an observation cage, and the foot-tapping period during five minutes after the recovery of the righting reflex was measured. The foot-tapping inhibition rate (%) of each test compound was calculated by the following equation.

Foot-tapping inhibition rate (%) = {1 - (Foot-tapping period when test compound was administered) / (Foot-tapping period when solvent was administered)} x100

[0106]

(2) Measurement of drug concentrations

After foot-tapping was finished, laparotomy was performed immediately under anesthesia with ether, and a blood sample was taken from the abdominal vena cava. At the same time, the brain was extracted. Through a quantitative analysis using liquid chromatography-mass spectrometry (LC/MS), the concentrations of the test compound in the plasma and the brain were measured.

[0107]

(3) Results

The effects on foot-tapping were measured by the above test method, and the results are shown in Table 11. In the table, Ex. No. means the Example number. Inhibition (%) is the foot-tapping inhibition rate, and Conc. (nM) is the drug concentration in the brain.

[Table 11]

E x . N o .	I n h i b i t i o n (%)	C o n c . (nM)
1	1 0 0	2 7 5
2	8 1	1 6 7
3	1 0 0	2 6 2
4	1 0 0	1 7 9
5	9 7	1 1 1
6	1 0 0	2 4 3
8	1 0 0	2 1 3
1 0	1 0 0	2 1 9

The compounds of the present invention were penetrated into the central nervous system and exhibited an excellent NK₁ receptor antagonist activity also in vivo.

[0108]

Test Example 5

20 Ferret pharmacokinetic test

(1) Methods

The test compound solution was prepared by dissolving the test compound in a

vehicle (50% N,N-dimethylformamide (Wako Pure Chemical Industries), 30% propylene glycol (Wako Pure Chemical Industries) and 4% 2-hydroxypropyl- β -cyclodextrin (Wako Pure Chemical Industries)). Under anesthesia with isoflurane, 0.3 mg/kg of the test compound was intravenously administered to a male ferret (Marshall BioResources Japan) from the jugular vein. In the case of oral administration, 1 mg/kg of the test compound was orally administered to an awake animal. After the administration of the test compound, blood samples were taken sequentially from the brachial cephalic vein, and the concentrations of the test compound in the plasma were measured by a quantitative analysis using liquid chromatography-mass spectrometry (LC/MS).

[0109]

(2) Results

The pharmacokinetic test in a ferret was tested by the above test method, and the results are shown in Table 12 and Table 13. In the tables, Ex. No. means the Example number. CL_{tot} and V_{ss} are the total body clearance and the steady-state volume of distribution, based on the plasma concentrations in the case of intravenous administration, respectively. C_{max}, AUC and BA are the maximum plasma test compound concentration, the area under the plasma test compound concentration-time curve and the bioavailability, in the case of oral administration, respectively.

[Table 12]

Ex. No.	CL _{tot} (mL/min/kg)	V _{ss} (mL/kg)
1	0.75	336

[Table 13]

Ex. No.	C _{max} (ng/mL)	AUC (ng·min/mL)	BA (%)
1	1,270	863,023	61

As shown in Table 12 and Table 13, the compound of the present invention exhibited an excellent pharmacokinetics.

[0110]

Test Example 6

5 Effect on cisplatin-induced acute and delayed emetic response

(1) Methods

(1-1) Intravenous administration test

The test compound solution was prepared by dissolving the test compound in a vehicle (a mixture of 50% N,N-dimethylformamide (Wako Pure Chemical Industries),
10 30% propylene glycol (Wako Pure Chemical Industries) and 4% 2-hydroxypropyl- β -cyclodextrin (Wako Pure Chemical Industries)). The vehicle only was administered to the control group.

Under anesthesia with isoflurane, 3 mg/kg of the test compound was intravenously administered to a male ferret (Marshall BioResources Japan) from the
15 jugular vein. Cisplatin, which was heated to 40-50°C, was intraperitoneally administered at 5 mg/kg one hour after the drug administration. The ferret was observed for 72 hours from immediately after the cisplatin administration, and the number of retching (periodic abdominal contraction without vomiting of the gastric content) and vomiting was counted.

20 (1-2) Oral administration test

The test compound solution was prepared by dissolving the test compound in a vehicle (1% N-methyl-2-pyrrolidone, 0.5% Tween 80, 20% polyethylene glycol 400 and 78.5% physiological saline). The solvent only was administered to the control group.

25 The test compound was orally administered to a male ferret (Marshall BioResources Japan). The oral administration was performed every 24 hours at 1 mg/kg three times in total. Cisplatin, which was heated to 40-50°C, was

intraperitoneally administered at 5 mg/kg one hour after the first oral administration of the test compound. The ferret was observed for 72 hours from immediately after the cisplatin administration, and the number of retching (periodic abdominal contraction without vomiting of the gastric content) and vomiting was counted.

5 [0111]

(2) Results

The results are shown in Fig. 1. In the control group, an increase in the number of retching and vomiting was observed in the acute phase (up to 24 hours after the cisplatin administration) and in the delayed phase (24 hours to 72 hours after the
10 cisplatin administration). In the group to which the compound of Example 1 was intravenously administered and in the group to which the compound was orally administered, the inhibition of the number of retching and vomiting was observed in the acute phase and in the delayed phase.

It was demonstrated that the compound of the present invention has a long-
15 acting medicinal effect and an inhibitory effect on the cisplatin-induced acute and delayed emetic responses.

[0112]

Test Example 7

Evaluation of hERG current

20 (1) Method

A dimethyl sulfoxide (DMSO) solution of the test compound with a concentration 1000 times higher than the evaluation concentration (10 μ M) was prepared, and a solution with the a final application concentration was prepared by diluting the solution. The hERG current was measured by a whole-cell method using a
25 patch clamp system, where a cover glass on which hERG channel-expressing human embryonic kidney (HEK) 293 cells were seeded was placed on a perfusion bath and a perfusion solution was caused to flow. The change in the hERG channel-derived

current caused by a pulse protocol (holding potential of -80 mV, depolarization pulse of +20 mV for 1.9 seconds, repolarization pulse of -50 mV for 2 seconds, stimulated at intervals of 15 seconds) of data acquisition/analysis software, pCLAMP9 (Axon Instruments, Inc.), was measured. The measurement conditions were a flow rate of about 1.5 mL/min and a temperature of about 33°C. Two wave forms just before applying the test compound and two wave forms immediately after the application for 10 minutes were analyzed, and the statistical analysis was performed. The value before the application of the test compound was regarded as 100%, and the change rate based on the value was determined.

(2) Result

The effect of the test compound on hERG current was evaluated by the above method (the result is shown in Table 14). In the table, Ex. No. means the Example number, and the value is the average $\pm$ the standard error.

[Table 14]

Ex. No.	% (n=3)
1	93.9 $\pm$ 2.7

As shown in Table 14, the compound of the present invention did not cause any change in the hERG current with a statistical significance compared to the vehicle control (0.1% DMSO).

INDUSTRIAL APPLICABILITY

[0113]

The compounds of the present invention or pharmaceutically acceptable salts thereof have an excellent NK₁ receptor antagonist activity, and thus are also useful as an agent for the prevention or treatment of cancer-chemotherapy-induced nausea and vomiting.

SEQUENCE LISTING FREE TEXT

[0114]

<Sequence listing 1>

SEQ ID NO:1 is the sequence of forward primer which was used for DNA amplification
5 of SEQ ID NO:3.

<Sequence listing 2>

SEQ ID NO:2 is the sequence of reverse primer which was used for DNA amplification
of SEQ ID NO:3.


<Sequence listing 3>

10 SEQ ID NO:3 is the DNA sequence of protein expression site, which was amplified by
using primers of SEQ ID NO:1 and 2 for expressing human tachykinin receptor 1.

CLAIMS

1. A compound represented by the formula (I):


[Chem.1]



5 wherein

ring A is a group represented by the formula:


[Chem.2]



;

ring B is a group represented by the formula:


10 [Chem.3]



;

with the proviso that bonds with (*) are bonding site to the formula:

[Chem.4]




;

15 bonds with (**) are bonding site to the ring A;

bonds with (***) are bonding site to the formula:

[Chem.5]



;

R_1 is C_{1-6} alkyl or C_{1-6} alkoxy;


R_2 and R_3 are each independently a hydrogen atom or methyl;

n is 0, 1, 2, 3, 4 or 5;

5 or a pharmaceutically acceptable salt thereof.

2. The compound represented by the formula (Ia) according to claim 1:

[Chem.6]




10 wherein

ring A, R_1 and n have the same meaning as described in claim 1;

ring B^b is a group represented by the formula:


[Chem.7]



;

15 wherein, bond with (*) is a bonding site to the formula:

[Chem.8]




;

(**) and (***) have the same meaning as described in claim 1 ;

or a pharmaceutically acceptable salt thereof.

5 3. The compound represented by the formula (Ib) according to claim 2:

[Chem.9]




wherein

ring A and ring B^b have the same meaning as described in claim 2;

10 with the proviso that bond with (*) is a bonding site to the formula:

[Chem.10]



;


(**) and (***) have the same meaning as described in claim 2 ;

R^{1a}, R^{1b}, R^{1c}, R^{1d} and R^{1e} are each independently any one of a hydrogen atom, methyl or

15 methoxy;


or a pharmaceutically acceptable salt thereof.

4. The compound according to claim 1, wherein ring B is the following formula:
[Chem.11]




- 5 wherein, (*), (**) and (***) have the same meaning as described in claim 1 ;
or a pharmaceutically acceptable salt thereof.
5. The compound according to claim 4, wherein R₁ is methyl, and n is 0 or 1, or a pharmaceutically acceptable salt thereof.

6. A compound according to claim 1 represented by the following formula:
[Chem.12]



or a pharmaceutically acceptable salt thereof.


7. A compound according to claim 1 represented by the following formula:
[Chem.13]



or a pharmaceutically acceptable salt thereof.

8. A compound according to claim 1
represented by the following formula:


5 [Chem.14]



or a pharmaceutically acceptable salt thereof.

9. A compound according to claim 1
represented by the following formula:


10 [Chem.15]



or a pharmaceutically acceptable salt thereof.

10. A compound according to claim 1 represented by the following formula:

[Chem.16]



5 or a pharmaceutically acceptable salt thereof.

11. A pharmaceutical composition comprising as an active ingredient a compound according to any one of claims 1 to 10, or a pharmaceutically acceptable salt thereof.