

METHOD OF SYNTHESIS


Technical Field of the Invention

The present invention resides in the field of chemical synthesis. More specifically, the present invention relates to novel methods useful in the
5 synthesis of a positron emission tomography (PET) tracer, and novel intermediates useful in said method.

Description of Related Art

WO 94/27591 describes certain substituted guanidines and their use for treatment and/or prophylaxis of neurological conditions such as epilepsy, neurodegenerative
10 conditions and/or nerve cell death resulting from e.g. hypoxia, hypoglycemia, brain or spinal chord ischemia, brain or spinal chord trauma. WO 94/27591 describes that the substituted guanidines can be prepared by the reaction of an amine, typically an amine salt such as an amine hydrochloride, with a preformed alkyl or aryl cyanamide (Safer *et al* 1948 J Org Chem; 13: 924) or the corresponding N-
15 substituted alkyl or aryl cyanamide.

WO 2004/007440 and WO 2006/136846 describe radiolabelled guanidine derivatives and their use for imaging central nervous system (CNS) receptors and teach synthesis of these radiolabelled derivatives from precursor compounds. For example, WO 2006/136846 teaches a compound of Formula (A):



or a salt or solvate thereof, wherein:

R¹ is hydrogen or C₁₋₄alkyl;


R² and R⁴ are each independently selected from C₁₋₄ alkyl, [¹¹C]-C₁₋₄alkyl, and [¹⁸F]-C₁₋₄ fluoroalkyl provided that at least one of R² and R⁴ is [¹¹C]-C₁₋₄alkyl or

$[^{18}\text{F}]$ -C₁₋₄ fluoroalkyl; and,

R³ is halo.


WO 2006/136846 teaches that the above compound of Formula (A) is synthesised by reaction of a suitable source of ¹¹C or ¹⁸F with a precursor

5 compound of Formula (B):




wherein one of R² or R⁴ is hydrogen, and the other is hydrogen, C₁₋₄ alkyl, or a thiol protecting group such as benzyl; R¹ is hydrogen or C₁₋₄alkyl, and R³ is halo.

10 WO 2006/136846 also teaches that the method to obtain the precursor compound of Formula (B) above wherein R² is hydrogen is based on that disclosed by Hu *et al* (J. Med. Chem. 1997; 40(26): 4281-9), wherein a compound of Formula (C):




15 or a salt or solvate thereof, wherein R³ is halo and P¹ is a thiol protecting group; is reacted with a compound of Formula (D):



wherein R¹ is hydrogen or C₁₋₄ alkyl and R⁴ is as defined for the desired compound of Formula (B).

This method has also been recently reported by Robins *et al* (2010 Bioorg Med Chem Lett; 20: 1749-51) as a successful way to obtain the following ^{18}F -labelled S-fluoroalkyl diarylguanidines:



- 5 However, the above-described method of preparing the radiolabelling precursor of Formula (B) suffers from a number of problems. First of all, tin chloride is used in a reduction step used in the preparation of the compound of Formula (C) from sulfonylchloride starting material. Residual tin complicates the workup by gel forming tin hydroxides at pH >2. Furthermore, following this reduction
- 10 step, a benzyl protective group is introduced at the thiol and this group has to be removed at the end of the synthesis, requiring use of AlCl_3 followed by flash chromatography. There is therefore a need for a method to prepare this radiolabelling precursor that overcomes these problems.

In addition, and not described in the prior art, the present inventors have found

15 that the compound of Formula (B) wherein R^2 is hydrogen decomposes to form a disulfide impurity, even under what would be considered "inert" conditions, which complicates subsequent radiolabelling to obtain the corresponding compound of Formula (A). There is therefore an additional need for alternative strategies to obtain said compound of Formula (A) that do not suffer from this

20 disadvantage.

Summary of the Invention

The present invention provides a method to prepare a radiolabelled guanidine derivative. A novel intermediate and its method of synthesis are also provided.


In the method of the invention, use of a step comprising iodine oxidation has

25 the advantage that the iodine oxidises any tin present as a consequence of the

initial tin chloride reduction step. No protecting groups are required in the method of the invention, thereby eliminating the need for a deprotection step. In addition, the method of the invention overcomes the problem observed by the present inventors with formation of a disulphide impurity. The method of the invention therefore overcomes a number of problems associated with known methods for the preparation of radiolabelled guanidine derivatives.

Detailed Description of the Invention

In one aspect, the present invention provides a method to obtain a positron emission tomography (PET) tracer of Formula I:



wherein:

X¹ is an X group selected from C₁₋₄ alkyl or halo;


Y¹ is a Y group selected from hydrogen or C₁₋₄ alkyl;

Z¹ is a Z group which is C₁₋₄ alkyl; and,

Q is [¹¹C]C₁₋₄ alkyl- or [¹⁸F]-C₁₋₄fluoroalkyl-;

wherein said method comprises:

(a) providing a compound of Formula II:




wherein:

X^{11} and X^{12} are the same and are both an X group as defined for X^1 ;

Y^{11} and Y^{12} are the same and are both a Y group as defined for Y^1 ; and,

5 Z^{11} and Z^{12} are the same and are both a Z group as defined for Z^1 ;

(b) reducing said compound of Formula II with a reducing agent to obtain a compound of Formula III:



X^{21} is an X group as defined for X^1 ;

10 Y^{21} is a Y group as defined for Y^1 ; and,

Z^{21} is a Z group as defined for Z^1 ;

(c) reacting the compound of Formula III as obtained in step (b), with either $[^{11}\text{C}]\text{C}_{1-4}$ alkyl- LG^1 or $[^{18}\text{F}]\text{-C}_{1-4}$ fluoroalkyl- LG^2 , wherein LG^1 and LG^2 are independently halo, or the group $-\text{O}-\text{SO}_2-\text{R}^1$ wherein R^1 represents an

optionally-substituted C₆₋₁₀ aryl, an optionally-substituted C₁₋₄ alkyl, or C₁₋₄ fluoroalkyl, wherein said reacting is carried out in a suitable solvent.

The term "PET tracer" refers to a chemical compound that comprises a radionuclide that undergoes positron emission decay (also known as positive beta decay), and is therefore detectable using PET imaging. The most commonly used radionuclides for PET are ¹⁸F and ¹¹C.

The term "alkyl", alone or in combination, means a straight-chain or branched-chain alkyl radical having the general formula C_nH_{2n+1}. Examples of such radicals include methyl, ethyl, and isopropyl.

10 The term "halogen" or "halo" in the context of the present invention means a substituent selected from fluorine, chlorine, bromine or iodine.

The term "reducing agent" (also commonly referred to as a "reductant" or "reducer") is the element in an oxidation-reduction reaction that donates an electron to another species. For the present invention, non-limiting examples of suitable reducing agents for use in step (b) in the above-defined method include: sodium borohydride (NaBH₄), zinc in hydrochloric acid, zinc in acetic acid, magnesium in hydrochloric acid, sodium hydrogentelluride (NaTeH) in ethanol, lithium aluminium hydride (LiAlH₄) in tetrahydrofuran, indium in ammonium chloride, and sodium hydride (NaH). A preferred reducing agent is NaBH₄. In a preferred embodiment, the reducing agent is bound to a solid phase such as a resin in the form of particles such as beads.

The term "fluoroalkyl" refers to an alkyl as defined above that comprises a fluorine atom in place of a hydrogen. Specifically, the term fluoroalkyl as used herein is taken to mean [¹⁸F]fluoroalkyl, and as such the fluorine atom comprised therein is radioactive ¹⁸F. Preferably, said fluoroalkyl includes a single ¹⁸F atom, most preferably at the terminal end of the chemical group.


The term "leaving group" refers to a moiety suitable for nucleophilic substitution and is a molecular fragment that departs with a pair of electrons in heterolytic bond cleavage.

The term “aryl” refers to a monovalent aromatic hydrocarbon having a single ring (i.e. phenyl) or fused rings (i.e. naphthalene).

A chemical group defined herein as “optionally substituted” may either have no substituents or may include one or more substituents. Preferred substituents include C₁₋₄ alkyl, C₁₋₄ haloalkyl, halo, and nitro, wherein alkyl is as defined above, “haloalkyl” is an alkyl as defined above that comprises a halo, halo is as defined above, and “nitro” refers to the group -NO₂. Accordingly, examples of preferred -O-SO₂-R¹ groups for the present invention include: toluenesulfonic acid, nitrobenzenesulfonic acid, benzenesulfonic acid, trifluoromethanesulfonic acid, fluorosulfonic acid, and perfluoroalkylsulfonic acid.

The reaction step (c) with [¹¹C]C₁₋₄ alkyl-LG¹ or [¹⁸F]-C₁₋₄ fluoroalkyl-LG² is an alkylation reaction carried out in a suitable solvent. A “suitable solvent” is one in which the reactants are readily soluble and readily react to result in the desired product. Such a suitable solvent may be selected from the group comprising N,N-dimethylformamide (DMF), acetone, dichloromethane (DCM), chloroform, dimethylsulphoxide (DMS), methanol, ethanol, propanol, isopropanol, tetrahydrofuran (THF), or acetonitrile.

In a preferred embodiment of the method of the invention said compound of Formula II is a compound of Formula IIa:



wherein:

X^{11} and X^{12} are the same and are both an X group as suitably and preferably defined herein;

Y^{11} and Y^{12} are the same and are both a Y group as suitably and preferably defined herein; and,

5 Z^{11} and Z^{12} are the same and are both a Z group as suitably and preferably defined herein.


Preferably for the method of the invention said X group is halo, most preferably chloro.

10 Preferably for the method of the invention said Y group is C_{1-4} alkyl, most preferably methyl.

Preferably for the method of the invention said Z group is methyl.


In a preferred embodiment for the intermediates and the product of the method of the invention, said X group is chloro, said Y group is methyl and said Z group is methyl.

15 In summary, the compound of Formula II provided in step (a) of the method of the invention is obtained by reacting a compound of Formula IV:



wherein X^{31} and X^{32} are the same and are both an X group as suitably and preferably defined herein;

20 with a compound of Formula V:




(V)

wherein Y^{41} is a Y group as suitably and preferably defined herein, and Z^{41} is a Z group as suitably and preferably defined herein.

The coupling of the compound of Formula IV with the cyanamide of Formula V
 5 may be performed without solvent, or in the presence of a high boiling non-protic solvent such as chlorobenzene, toluene, or xylene. This reaction may be effected at elevated temperature, for example 50 to 200°C, preferably at around 130°C.

The above-defined compound of Formula IV is obtained by iodine oxidation of a
 10 compound of Formula VI:










(VI)

wherein X^{51} is an X group as suitably and preferably defined herein.

Table 1 below illustrates a known method to obtain a compound of Formula III alongside the methods used in the present invention:

15

Prior Art	Present Invention
 Formula VI	 Formula VI
↓ add thiol protecting group 	↓ I₂  Formula IV
+  Formula V	+  Formula V



In Table 1:

each X, Y and Z group is as suitably and preferably defined herein for X, Y and Z, respectively; and,

- 5 P^1 is a thiol protecting group.

Protecting groups are well known to those skilled in the art. For thiol groups suitable protecting groups are benzyl, trityl, and 4-methoxybenzyl. The use of further protecting groups are described in 'Protective Groups in Organic Synthesis', Theodorora W. Greene and Peter G. M. Wuts, (Fourth Edition,

10 John Wiley & Sons, 2007).

The cyanamide starting material of Formula V in Table 1 can be prepared according to the method described by Hu *et al* (1997 J Med Chem; 40: 4281-4289), by reaction of cyanogen bromide with the primary amine in diethyl ether, or by alkylation of an arylcyanamide with sodium hydride or alkyl halide in

15 tetrahydrofuran. The nitrobenzenesulfonyl chloride starting material illustrated

in Table 1 is commercially available. The first step in Table 1, common to the prior art method and the method of the present invention, is reduction of the nitrobenzenesulfonyl chloride starting material to form the aminobenzenethiol intermediate of Formula VI. In the second step of the prior art method, a

5 protecting group is placed onto the thiol group, which is removed at the end of the synthesis using known methods. For example, where the thiol protecting group is benzene, AlCl_3 and flash chromatography may be used for its removal. The disulfide of Formula IV is obtained in the method of the invention by iodine oxidation of the aminobenzenethiol intermediate of Formula VI. This step has

10 the advantage that the iodine additionally oxidises any tin present. In the prior art method, residual tin complicates the workup because the tin hydroxides form gels at $\text{pH} > 2$. The method used in the present invention overcomes this problem as oxidised tin does not form gels, thereby facilitating extractive work up of the product. The method is furthermore advantageous because use of

15 the disulfide intermediate of Formula IV circumvents the need to protect the thiol group. The method used in the present invention to obtain a compound of Formula III illustrated in Table I above may be regarded as another aspect of the present invention.

The present inventors have observed that the compound of Formula III

20 decomposes to form the disulfide of Formula II, even under what would be considered "inert" conditions. This problem is effectively overcome by storage of the disulfide compound of Formula II instead of the compound of Formula III. In the method to obtain a PET tracer of Formula I the compound of Formula II is reduced by step (b) immediately before the radiolabelling step (c). In order to

25 facilitate this, steps (b) and (c) are preferably carried out in the same vessel. A further advantage of this strategy over the prior art methods is that it is not necessary to include a base in the reaction step (c). Inclusion of a base in the reaction is required by the methods taught by the prior art to deprotonate the thiol in order to allow reaction with the radiolabelled synthon. Suitable bases taught in

30 the prior art methods include inorganic bases such as potassium carbonate, potassium hydroxide, or sodium hydride, or organic bases such as a trialkylamine, for example triethylamine, diisopropylethylamine, or dimethylaminopyridine. In the

method of the present invention, once the compound of Formula II is reduced in step (b) it can be reacted directly in step (c) with $[^{11}\text{C}]\text{C}_{1-4}$ alkyl- LG^1 or $[^{18}\text{F}]\text{-C}_{1-4}$ fluoroalkyl- LG^2 to obtain the PET tracer of Formula I without any requirement to add a base.

- 5 $[^{11}\text{C}]\text{C}_{1-4}$ alkyl- LG^1 provided in step (c) of the method can be prepared using methods well-known in the art of radiochemistry. For example, $[^{11}\text{C}]\text{methyl iodide}$ can be prepared by reduction of $[^{11}\text{C}]\text{carbon dioxide}$ with lithium aluminium hydride followed by reaction with hydroiodic acid. $[^{11}\text{C}]\text{carbon dioxide}$ is usually produced by the $^{14}\text{N}(\rho,\alpha)^{11}\text{C}$ reaction from nitrogen gas containing trace amounts of oxygen.
- 10 $[^{11}\text{C}]\text{methyl triflate}$ can be prepared from $[^{11}\text{C}]\text{methyl iodide}$, or by gas phase reaction of $[^{11}\text{C}]\text{methyl bromide}$ prepared from $[^{11}\text{C}]\text{methane}$. All these methods are described in more detail in "Aspects on the Synthesis of ^{11}C -Labelled Compounds", Chapter 3 of Handbook of Radiopharmaceuticals (2003 Welch & Redvanly eds. pp 141-194). A preferred $[^{11}\text{C}]\text{C}_{1-4}$ alkyl- LG^1 is selected from
- 15 $[^{11}\text{C}]\text{methyl-}\text{LG}^1$ or $[^{11}\text{C}]\text{ethyl-}\text{LG}^1$, and LG^1 is preferably iodo.

- $[^{18}\text{F}]\text{-C}_{1-4}$ fluoroalkyl- LG^2 provided in step (c) of the method can be prepared by radiolabelling alkyl dihalides or sulfonates using $[^{18}\text{F}]\text{fluoride}$. $[^{18}\text{F}]\text{Fluoride}$ is typically obtained as an aqueous solution which is a product of the irradiation of an $[^{18}\text{O}]\text{-water}$ target. It has been widespread practice to carry out various steps in
- 20 order to convert $[^{18}\text{F}]\text{Fluoride}$ into a reactive nucleophilic reagent, such that it is suitable for use in nucleophilic radiolabelling reactions. These steps include the elimination of water from $[^{18}\text{F}]\text{-fluoride ion}$ and the provision of a suitable counterion (Handbook of Radiopharmaceuticals 2003 Welch & Redvanly eds. ch. 6 pp 195-227). Suitable counterions include large but soft metal ions such as
- 25 rubidium or caesium, potassium complexed with a cryptand such as KryptofixTM, or tetraalkylammonium salts. A preferred $[^{18}\text{F}]\text{-C}_{1-4}$ fluoroalkyl- LG^2 is $[^{18}\text{F}]\text{-fluoroethyl-}\text{LG}^2$ wherein LG^2 is preferably a sulfonate, most preferably tosylate.

- Preferably for the method to obtain said PET tracer of Formula I, the reducing step (b) and the reacting step (c) are carried out in immediate sequence. The
- 30 term "in immediate sequence" should be understood to mean that the reacting step (c) is carried out as soon as possible, i.e. directly, after the reducing step

(b), such that there is as little time as practically possible between the two steps and ideally no time between the two steps. In this way, any decomposition of the compound of Formula III to form the disulfide is minimised, thereby facilitating the radiolabelling reaction.

5 In a particularly preferred embodiment, the above-described method to obtain said PET tracer of Formula I is automated. PET tracers, and [^{18}F]-tracers in particular, are now often conveniently prepared on an automated radiosynthesis apparatus. There are several commercially-available examples of such apparatus, including TracerlabTM and FastlabTM (both from GE Healthcare Ltd). Such apparatus
10 commonly comprises a “cassette”, often disposable, in which the radiochemistry is performed, which is fitted to the apparatus in order to perform a radiosynthesis. The cassette normally includes fluid pathways, a reaction vessel, and ports for receiving reagent vials as well as any solid-phase extraction cartridges used in post-radiosynthetic clean up steps. The present invention therefore provides in
15 another aspect a cassette for carrying out these steps wherein said cassette comprises:

- (i) a vessel containing the compound of Formula II as suitably and preferably defined herein;
- (ii) means for reacting said compound of Formula II with a reducing agent
20 to form a compound of Formula III, wherein said reducing agent and said compound of Formula III are as suitably and preferably defined herein; and,
- (iii) means for reacting said compound of Formula III with either [^{11}C]C₁₋₄ alkyl-LG¹ or [^{18}F]-C₁₋₄ fluoroalkyl-LG² to obtain a PET tracer of
25 Formula I, wherein LG¹, LG² and said PET tracer of Formula I are as suitably and preferably defined herein.

The means for reacting said compound of Formula II with said reducing agent may be a vessel containing the reducing agent in solution (or in a soluble) form, wherein the reducing agent is passed through the vessel containing the compound
30 of Formula II in order to effect the reduction. Alternatively, said means may be a

vessel in which the reducing agent is bound to a solid phase, wherein the compound of Formula II is passed through the vessel containing the reducing agent in order to effect the reduction. The suitable and preferred embodiments described herein for the reduction step (b) and the reaction step (c) also apply to
 5 the method as carried out on the cassette of the invention.

The cassette may additionally comprise:

- (i) an ion-exchange cartridge for removal of excess [^{11}C]C₁₋₄ alkyl-LG¹ or [^{18}F]-C₁₋₄ fluoroalkyl-LG².

The PET tracer of Formula I obtained by the method of the invention is useful as a
 10 radioligand for the NMDA receptor and can be used in an *in vivo* diagnostic or imaging method such as positron emission tomography (PET). A PET tracer of Formula I as defined above, or a salt or solvate thereof, may be used to image the NMDA receptor in healthy human volunteers. Because the PET tracer of Formula I is useful for *in vivo* imaging of NMDA receptors it thus also has utility in the
 15 diagnosis of NMDA-mediated disorders, such as stroke, brain or spinal chord trauma, epilepsy, Alzheimer's disease, or Huntington's disease.

Brief Description of the Examples

Example 1 describes the synthesis of 1,1'-(5,5'-disulfanediy)bis(2-chloro-5,1-phenylene))bis(3-methyl-3-(3-(methylthio)phenyl)guanidine).

20 Example 2 describes the synthesis of 3-(2-chloro-5-mercaptophenyl)-1-methyl-1-(3-(methylthio)phenyl)guanidine.

Example 3 describes the synthesis of 3-(2-chloro-5-((2-fluoroethyl)thio)phenyl)-1-methyl-1-(3-(methylthio)phenyl)guanidine.

Example 4 describes the synthesis of 3-(2-chloro-5-((2-fluoroethyl)thio)phenyl)-
 25 1-methyl-1-(3-(methylthio)phenyl)guanidine using resinbound borohydride.

List of Abbreviations used in the Examples

DCM dichloromethane


	EtOH	ethanol
	g	gram(s)
	HPLC	high performance liquid chromatography
	M	molar
5	MeOH	methanol
	mg	milligram(s)
	mL	millilitre(s)
	mmol	millimole(s)
	NMR	nuclear magnetic resonance
10	RT	room temperature

Examples

Unless otherwise specified, the intermediates and reagents used in the examples were purchased from Sigma Aldrich, Merck, or Alfa Aesar.

Example 1: Synthesis of 1,1'-(5,5'-disulfanediylbis(2-chloro-5,1-phenylene))bis(3-methyl-3-(3-(methylthio)phenyl)guanidine)

1(a) Synthesis of 5,5'-disulfanediylbis(2-chlorobenzenaminium) chloride




Tin(II) chloride (33.32g, 175.74mmol) was dissolved in 30% hydrochloric acid (99.7mL) and 4-chloro-3-nitrobenzene-1-sulfonyl chloride (5.00g, 19.553mmol) was added before submerging the flask in a 125°C preheated oil bath. After 3 hours all solid material had dissolved and the reaction mixture was allowed to cool to RT, which caused spontaneous crystallization.

The crystals (contaminated with Tin) were filtered off, dissolved in water (250mL) and portions of iodine solution (50mg/mL) were added until HPLC analysis confirmed that all the 2-chloro-5-mercaptopbenzenaminium chloride had been converted to 5,5'-disulfanediybis(2-chlorobenzenaminium) chloride.

The solution was filtered, and water (400mL) was added to the filtrate, followed by stirring and neutralization using NaOH solution (~1mL, 10%). The solution was extracted with diethyl ether (4 x 150mL), dried with magnesium sulphate (anhydrous) and filtered. To the ether solution was added HCl (dry, 1M in diethyl ether, 10mL), the solution was filtered and the filtrate was dried under vacuum to give 5,5'-disulfanediybis(2-chlorobenzenaminium) chloride as an off-white powder (21,47g, 63%).

¹H NMR (400 MHz, CDCl₃): δ 7.154 (d, J = 8.3 Hz, 2H), δ 6.865 (d, J = 2.2 Hz, 2H), δ 6.783 (dd, J₁ = 2.2 Hz, J₂=8.3 Hz, 2H), δ 4.080 (broad s, 4H)

1(b) Synthesis of 1,1'-(5,5'-disulfanediybis(2-chloro-5,1-phenylene))bis(3-methyl-3-(3-(methylthio)phenyl)guanidine)




A mixture of 5,5'-disulfanediybis(2-chlorobenzenaminium) chloride (1.0g, 2.6mmol) and N-methyl-N-(3-(methylthio)phenyl)cyanamide (1.83g, 10.3 mmol) was heated to 130°C. This thick slow-stirring melt was left for 17h (HPLC yield

after 1 hour ~80%), then allowed to cool to RT. The solid was dissolved in DCM (25mL), extracted with water (3 x 200mL) and the combined aqueous phases back-extracted with DCM (50mL). The aqueous phase was neutralised with NaHCO₃ and extracted with diethyl ether (3 x 150mL). The combined organic phases were dried with magnesium sulphate (~5g), filtered and concentrated to dryness under reduced pressure to give 1,1'-(5,5'-disulfanediylbis(2-chloro-5,1-phenylene))bis(3-methyl-3-(3-(methylthio)phenyl)guanidine) (1.16g, 1.7mmol, 67%, HPLC purity 94.8%) as off-white powder.

- ¹H NMR (400 MHz, CDCl₃): δ 7.300 (t, J = 7.9 Hz, 2H), δ 7.298 (d, J = 8.3 Hz, 2H), δ 7.175 (t, J = 1.9 Hz, 2H), δ 7.122 (ddd, J₁ = 1.0, J₂ = 1.8 Hz, J₃ = 7.9 Hz, 2H), δ 7.108 (d, J = 2.3 Hz, 2H), δ 7.054 (ddd, J₁ = 1.0, J₂ = 2.2 Hz, J₃ = 7.9 Hz, 2H), δ 7.049 (dd, J₁ = 2.3, J₂ = 8.3 Hz, 2H), δ 3.893 (broad s, 4H), δ 3.338 (s, 6H), δ 2.494 (s, 6H).

Example 2: Synthesis of 3-(2-chloro-5-mercaptophenyl)-1-methyl-1-(3-(methylthio)phenyl)guanidine




- 1,1'-(disulfanediylbis(2-chloro-5,1-phenylene))bis(3-methyl-3-(3-(methylthio)phenyl)guanidine) (3.1g, 4.6mmol), obtained as described in Example 1, was suspended in EtOH (62mL) and then sodium borohydride (0.5g, 13.8mmol) was added in portions. The reaction solution was left to stir overnight under an inert atmosphere. The reaction was cooled to 10 degrees before quenching with hydrochloric acid in ether (2M) and concentrated to dryness under reduced pressure to give a creamy solid. The solid was purified by column chromatography (CHCl₃->7% MeOH in CHCl₃) to give 3-(2-chloro-5-

mercaptophenyl)-1-methyl-1-(3-(methylthio)phenyl)guanidine (2.5g, 6,7mmol, 73.6%) as a white foam.


¹H NMR (500 MHz, CDCl₃): δ 9.686 (s, 1H), δ 8.531 (s, 2H), δ 7.132 (t, J = 7.8 Hz, 1H), δ 7.032 (d, J = 8.4 Hz, 1H), δ 7.022 (d, J = 7.8 Hz, 1H), δ 7.014 (s, 1H), δ 6.922 (s, 1H), δ 6.892 (d, J = 8.4 Hz, 1H), δ 6.855 (d, J = 7.8 Hz, 1H), δ 3.740 (s, 1H), δ 3.656 (s, 3H), δ 2.460 (s, 3H).

Example 3: Synthesis of 3-(2-chloro-5-((2-fluoroethyl)thio)phenyl)-1-methyl-1-(3-(methylthio)phenyl)guanidine



- 10 1,1'-(disulfanediy)bis(2-chloro-5,1-phenylene))bis(3-methyl-3-(3-(methylthio)phenyl)guanidine) (3.0g, 4.45mmol), obtained as described in Example 1, was dissolved in ethanol (120mL) and 2-fluoroethyl 4-methylbenzenesulfonate (2.1g, 9.8mmol) was added. The clear solution was heated to 60°C under inert atmosphere and sodium borohydride (0.8g, 22.3mmol) was added in portions. The reaction was heated for 45 minutes before being concentrated to dryness under reduced pressure. The crude was purified by column chromatography (CHCl₃→2% MeOH in CHCl₃) to give 3-(2-chloro-5-((2-fluoroethyl)thio)phenyl)-1-methyl-1-(3-(methylthio)phenyl)guanidine (2.2g, 5.6 mmol, 62.9%) as a clear slowly crystalizing oil.
- 20 ¹H NMR (400MHz, CDCl₃): δ7.28-7.35 (m, 2H), δ7.05-7.25 (m, 4H), δ6.90-6.95 (dd, J₁=3.18Hz, J₂=8.33Hz 1H), δ4.6 (t, J=6.75Hz, 1H), δ4.5 (t, J=6.75Hz, 1H), δ3.9 (broad s, 1H), δ3.4 (s, 3H), δ3.2 (t, J=6.75Hz, 1H), δ3.15 (t, J=6.75Hz, 1H), δ2.52 (s, 3H) δ1.65 (broad s, 1H)

Example 4: Synthesis of 3-(2-chloro-5-((2-fluoroethyl)thio)phenyl)-1-methyl-1-(3-(methylthio)phenyl)guanidine using resinbound Borohydride




- 1,1'-(disulfanediy)bis(2-chloro-5,1-phenylene))bis(3-methyl-3-(3-
 5 (methylthio)phenyl)guanidine) (0.15 g, 0.22 mmol), obtained as described in
 Example 1, and 2-fluoroethyl 4-methylbenzenesulfonate (0.10 g, 0.45 mmol)
 was dissolved in ethanol (96%, 10 mL), and solid phase supported borohydride
 (0.45g, ~1.11mmol) was added. The resulting reaction suspension was heated
 under inert atmosphere to 60°C for 17 hours. The suspension was filtered and
 10 the solid phase material was washed with ethanol (96%, 2*5mL). The
 combined organic phases were concentrated under reduced pressure to give a
 moist crude that was redissolved in dichloromethane (10mL), dried with
 magnesium sulfate (~0.3 g, 2.492 mmol), filtered and concentrated to dryness
 to give 3-(2-chloro-5-((2-fluoroethyl)thio)phenyl)-1-methyl-1-(3-
 15 (methylthio)phenyl)guanidine (0.13 g, 0.32 mmol, 70.7 % yield) as a clear oil.

^1H NMR (400MHz, CDCl_3): δ 7.28-7.35 (m, 2H), δ 7.05-7.25 (m, 4H), δ 6.90-6.95
 (dd, $J_1=3.18\text{Hz}$, $J_2=8.33\text{Hz}$ 1H), δ 4.6 (t, $J=6.75\text{Hz}$, 1H), δ 4.5 (t, $J=6.75\text{Hz}$, 1H),
 δ 3.9 (broad S, 1H), δ 3.4 (s, 3H), δ 3.2(t, $J=6.75\text{Hz}$, 1H), δ 3.15 (t, $J=6.75\text{Hz}$,
 1H), δ 2.52 (S, 3H) δ 1.65 (broad S, 1H)

Claims

- 1) A method to obtain a positron emission tomography (PET) tracer of Formula I:



- 5 wherein:

X^1 is an X group selected from C_{1-4} alkyl or halo;


Y^1 is a Y group selected from hydrogen or C_{1-4} alkyl;

Z^1 is a Z group which is C_{1-4} alkyl; and,

Q is $[^{11}C]C_{1-4}$ alkyl- or $[^{18}F]-C_{1-4}$ fluoroalkyl-;

- 10 wherein said method comprises:

(a) providing a compound of Formula II:




wherein:

X^{11} and X^{12} are the same and are both an X group as defined for X^1 ;

Y^{11} and Y^{12} are the same and are both a Y group as defined for Y^1 ; and,

Z^{11} and Z^{12} are the same and are both a Z group as defined for Z^1 ;

- 5 (b) reducing said compound of Formula II with a reducing agent to obtain a compound of Formula III:




X^{21} is an X group as defined for X^1 ;

Y^{21} is a Y group as defined for Y^1 ; and,

- 10 Z^{21} is a Z group as defined for Z^1 ;

- (c) reacting the compound of Formula III as obtained in step (b), with either $[^{11}\text{C}]\text{C}_{1-4}\text{ alkyl-LG}^1$ or $[^{18}\text{F}]\text{-C}_{1-4}\text{ fluoroalkyl-LG}^2$, wherein LG^1 and LG^2 are independently halo, or the group $-\text{O-SO}_2\text{-R}^1$ wherein R^1 represents an optionally-substituted C_{6-10} aryl, an optionally-substituted C_{1-4} alkyl, or C_{1-4} fluoroalkyl, wherein said reacting is carried out in a suitable solvent.
- 15

- 2) The method as defined in Claim 1 wherein said compound of Formula II is a compound of Formula IIa:



wherein:

X^{11} and X^{12} are the same and are both an X group as defined in Claim 1;


5 Y^{11} and Y^{12} are the same and are both a Y group as defined in Claim 1; and,

Z^{11} and Z^{12} are the same and are both a Z group as defined in Claim 1.

- 3) The method as defined in either Claim 1 or Claim 2 wherein said X group is halo.
- 10 4) The method as defined in any one of Claims 1-3 wherein said Y group is C_{1-4} alkyl.
- 5) The method as defined in any one of Claims 1-4 wherein said Z group is methyl.
- 6) The method as defined in any one of Claims 1-5 wherein said X group is
15 chloro, said Y group is methyl and said Z group is methyl.
- 7) The method as defined in any one of Claims 1-6 wherein said reducing agent used in step (b) is selected from sodium borohydride ($NaBH_4$), zinc in hydrochloric acid, zinc in acetic acid, magnesium in hydrochloric acid, sodium hydrogentelluride ($NaTeH$) in ethanol, lithium aluminium hydride


(LiAlH₄) in tetrahydrofuran, indium in ammonium chloride, and sodium hydride (NaH).

- 8) The method as defined in any one of Claims 1-7 wherein in said reacting step (c) the compound of Formula III is reacted with [¹⁸F]-C₁₋₄ fluoroalkyl-LG².
- 9) The method as defined in any one of Claims 1-8 wherein said reducing step (b) and said reacting step (c) are carried out in immediate sequence.
- 10) The method as defined in any one of Claims 1-9 wherein said reducing step (b) and said reacting step (c) are carried out in the same vessel.
- 11) The method as defined in any one of Claims 1-10 wherein said method is automated.
- 12) A method for the preparation of the compound of Formula III as defined in any one of Claims 1 and 3-6, said method comprising reacting a compound of Formula IV:



wherein X³¹ and X³² are the same and are both an X group as defined in any one of Claims 1, 3 and 6;

with a compound of Formula V:



wherein Y^{41} is a Y group as defined in any one of Claims 1, 4 and 6, and Z^{41} is a Z group as defined in any one of Claims 1, 5 and 6;

to obtain a compound of Formula II as defined in any one of Claims 1-6;

5 and,

reducing said compound of Formula II using a reducing agent as defined in either Claim 1 or Claim 7 to obtain said compound of Formula III.

13) A cassette for carrying out the method as defined in Claim 11, wherein said cassette comprises:

10 (i) a vessel containing the compound of Formula II as defined in any one of Claims 1-6;

(ii) means for reacting said compound of Formula II with a reducing agent to form a compound of Formula III as defined in any one of Claims 1 and 3-6; and,

15 (iii) means for reacting said compound of Formula III with either $[^{11}\text{C}]\text{C}_{1-4}$ alkyl- LG^1 or $[^{18}\text{F}]\text{-C}_{1-4}$ fluoroalkyl- LG^2 to obtain a PET tracer of Formula I, wherein LG^1 , LG^2 , and said PET tracer of Formula I are as defined in Claim 1.

14) The cassette as defined in Claim 12 which further comprises:

20 (iv) an ion-exchange cartridge for removal of excess $[^{11}\text{C}]\text{C}_{1-4}$ alkyl- LG^1 or $[^{18}\text{F}]\text{-C}_{1-4}$ fluoroalkyl- LG^2 .