

COMBINATION COMPOSITIONS COMPRISING A BETA-LACTAMASE INHIBITOR AND USES THEREOF

CROSS-REFERENCE

[0001] This patent application claims the benefit of US Provisional Application No. 62/773,063, filed November 29, 2018; US Provisional Application No. 62/777,643, filed December 10, 2018; US Provisional Application No. 62/796,524, filed January 24, 2019; US Provisional Application No. 62/828,354, filed April 2, 2019; and US Provisional Application No. 62/832,118, filed April 10, 2019; each of which is incorporated herein by reference in their entirety.

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

[0002] This invention was made with government support under Grant No. R01AI111539, Grant No. R43AI109879, Grant No. R44AI109879, and Contract No. HHSN272201600029C, awarded by the National Institutes of Health (NIH). The government has certain rights in the invention.

FIELD OF INVENTION

[0003] The present invention relates to pharmaceutical compositions containing boron-containing compounds and their use as inhibitors of beta-lactamase enzymes and as antibacterial agents in combination with a beta-lactam antibiotic.

BACKGROUND OF THE INVENTION

[0004] Antibiotics are the most effective drugs for curing bacterial-infectious diseases clinically. They have a wide market due to their advantages of good antibacterial effect with limited side effects. Among them, the beta-lactam class of antibiotics (for example, penicillins, cephalosporins, and carbapenems) are widely used because they have a strong bactericidal effect and low toxicity.



[0005] To counter the efficacy of the various beta-lactams, bacteria have evolved to produce variants of beta-lactam deactivating enzymes called beta-lactamases, and in the ability to share this tool inter- and intra-species. These beta-lactamases are categorized as “serine” or “metallo” based, respectively, on presence of a key serine or zinc in the enzyme active site. The rapid spread of this mechanism of bacterial resistance can severely limit beta-lactam treatment options in the hospital and in the community.

[0006] There is a need for new orally-delivered antibacterial agents to treat resistant gram-negative infections in both the community and hospital setting.

SUMMARY OF THE INVENTION

[0007] Disclosed herein is a pharmaceutical composition comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof.




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) ceftibuten.

[0008] In some embodiments of a pharmaceutical composition, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0009] In some embodiments of a pharmaceutical composition, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0010] In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated for oral administration. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as an emulsion. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a microemulsion. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a Self-Emulsifying Drug Delivery System (SEDDS). In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a hydrophilic solubilizer. In some embodiments of a pharmaceutical composition, the hydrophilic solubilizer is a hydrophilic polymer. In some embodiments of a pharmaceutical composition, the hydrophilic polymer is a polyethylene glycol. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a surfactant. In some embodiments of a pharmaceutical composition, the surfactant is a polyoxyethylene stearate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises polyethylene glycol, caprylic/capric glycerides, and tocopheryl polyethylene glycol succinate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises propylene glycol, polyethylene glycol, and tocopheryl polyethylene glycol succinate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a tablet. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a capsule. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a powder for reconstitution. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a solution.

[0011] In some embodiments of a pharmaceutical composition, the powder for reconstitution is reconstituted with a liquid carrier to form an oral suspension.

[0012] Also disclosed herein is a pharmaceutical composition comprising:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) ceftibuten.

[0013] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) ceftibuten.



[0014] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or ((R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
- (ii) ceftibuten.

[0015] In some embodiments of a method of treating a bacterial infection, the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.

[0016] In some embodiments of a method of treating a bacterial infection, the method comprises administering to the subject in need thereof:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

- (ii) ceftibuten.

[0017] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





[0018] In some embodiments of a method of treating a bacterial infection, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0019] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and ceftibuten are formulated for oral administration. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and ceftibuten are administered sequentially. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and ceftibuten are administered concurrently.

[0020] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R¹ and R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl); each alkyl being optionally substituted with halogen, -OH, -O(C₁-C₆ alkyl), -NH₂, -NHC(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -COOH, or -COO(C₁-C₆ alkyl);


or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

(ii) ceftibuten.

[0021] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





[0022] In some embodiments of a method of treating a bacterial infection, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0023] In some embodiments of a method of treating a bacterial infection, the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and ceftibuten are formulated for oral administration. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered sequentially. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered concurrently.

[0024] Also disclosed herein is a pharmaceutical composition comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

- (ii) cefixime.

[0025] In some embodiments of a pharmaceutical composition, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0026] In some embodiments of a pharmaceutical composition, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0027] In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated for oral administration. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as an emulsion. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a microemulsion. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a Self-Emulsifying Drug Delivery System (SEDSS). In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a hydrophilic solubilizer. In some embodiments of a pharmaceutical composition, the hydrophilic solubilizer is a hydrophilic polymer. In some embodiments of a pharmaceutical composition, the hydrophilic polymer is a polyethylene glycol. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a surfactant. In some embodiments of a pharmaceutical composition, the surfactant is a polyoxyethylene stearate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises polyethylene glycol, caprylic/capric glycerides, and tocopheryl polyethylene glycol succinate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises propylene glycol, polyethylene glycol, and tocopheryl polyethylene glycol succinate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a capsule. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a tablet. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a powder for reconstitution. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a solution.

[0028] In some embodiments of a pharmaceutical composition, the powder for reconstitution is reconstituted with a liquid carrier to form an oral suspension.

[0029] Also disclosed herein is a pharmaceutical composition comprising:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) cefixime.

[0030] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) cefixime.



[0031] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or ((R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
- (ii) cefixime.

[0032] In some embodiments of a method of treating a bacterial infection, the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.

[0033] In some embodiments of a method of treating a bacterial infection, the method comprises administering to the subject in need thereof:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R¹ and R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl); each alkyl being optionally substituted with halogen, -OH, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, or -COO(C₁-C₆ alkyl);


or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

- (ii) cefixime.

[0034] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





[0035] In some embodiments of a method of treating a bacterial infection, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0036] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefixime are formulated for oral administration. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefixime are administered sequentially. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefixime are administered concurrently.

[0037] A method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R¹ and R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl); each alkyl being optionally substituted with halogen, -OH, -O(C₁-C₆ alkyl), -NH₂, -NHC(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -COOH, or -COO(C₁-C₆ alkyl);


or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

(ii) cefixime.

[0038] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





[0039] In some embodiments of a method of treating a bacterial infection, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0040] In some embodiments of a method of treating a bacterial infection, the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefixime are formulated for oral administration. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered sequentially. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered concurrently.

[0041] Disclosed herein is a pharmaceutical composition comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:


each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);

or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and
(ii) cefditoren pivoxil.

[0042] In some embodiments of a pharmaceutical composition, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0043] In some embodiments of a pharmaceutical composition, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0044] In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated for oral administration. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as an emulsion. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a microemulsion. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a Self-Emulsifying Drug Delivery System (SEDSS). In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a hydrophilic solubilizer. In some embodiments of a pharmaceutical composition, the hydrophilic solubilizer is a hydrophilic polymer. In some embodiments of a pharmaceutical composition, the hydrophilic polymer is a polyethylene glycol. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a surfactant. In some embodiments of a pharmaceutical composition, the surfactant is a polyoxyethylene stearate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises polyethylene glycol, caprylic/capric glycerides, and tocopheryl polyethylene glycol succinate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises propylene glycol, polyethylene glycol, and tocopheryl polyethylene glycol succinate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a tablet. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a capsule. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a powder for reconstitution. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a solution.

[0045] In some embodiments of a pharmaceutical composition, the powder for reconstitution is reconstituted with a liquid carrier to form an oral suspension.

[0046] Also disclosed herein is a pharmaceutical composition comprising:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) cefditoren pivoxil.

[0047] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) cefditoren pivoxil.



[0048] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or ((R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
- (ii) cefditoren pivoxil.

[0049] In some embodiments of a method of treating a bacterial infection, the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.

[0050] In some embodiments of a method of treating a bacterial infection, the method comprises administering to the subject in need thereof:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R¹ and R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl); each alkyl being optionally substituted with halogen, -OH, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, or -COO(C₁-C₆ alkyl);


or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

- (ii) cefditoren pivoxil.

[0051] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





[0052] In some embodiments of a method of treating a bacterial infection, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0053] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefditoren pivoxil are formulated for oral administration. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefditoren pivoxil are administered sequentially. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefditoren pivoxil are administered concurrently.

[0054] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) cefditoren pivoxil.

[0055] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





[0056] In some embodiments of a method of treating a bacterial infection, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0057] In some embodiments of a method of treating a bacterial infection, the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefditoren pivoxil are formulated for oral administration. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered sequentially. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered concurrently.

[0058] Disclosed herein is a pharmaceutical composition comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

- (ii) cefpodoxime proxetil.

[0059] In some embodiments of a pharmaceutical composition, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0060] In some embodiments of a pharmaceutical composition, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0061] In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated for oral administration. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as an emulsion. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a microemulsion. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a Self-Emulsifying Drug Delivery System (SEDDS). In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a hydrophilic solubilizer. In some embodiments of a pharmaceutical composition, the hydrophilic solubilizer is a hydrophilic polymer. In some embodiments of a pharmaceutical composition, the hydrophilic polymer is a polyethylene glycol. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a surfactant. In some embodiments of a pharmaceutical composition, the surfactant is a polyoxyethylene stearate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises polyethylene glycol, caprylic/capric glycerides, and tocopheryl polyethylene glycol succinate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition further comprises propylene glycol, polyethylene glycol, and tocopheryl polyethylene glycol succinate. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a tablet. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a capsule. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a powder for reconstitution. In some embodiments of a pharmaceutical composition, the pharmaceutical composition is formulated as a solution. In some embodiments of a pharmaceutical composition, the powder for reconstitution is reconstituted with a liquid carrier to form an oral suspension.

[0062] Also disclosed herein is a pharmaceutical composition comprising:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) cefpodoxime proxetil.

[0063] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) cefpodoxime proxetil.



[0064] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or ((R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
- (ii) cefpodoxime proxetil.

[0065] In some embodiments of a method of treating a bacterial infection, the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.

[0066] In some embodiments of a method of treating a bacterial infection, the method comprises administering to the subject in need thereof:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R¹ and R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl); each alkyl being optionally substituted with halogen, -OH, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, or -COO(C₁-C₆ alkyl);


or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

- (ii) cefpodoxime proxetil.

[0067] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





[0068] In some embodiments of a method of treating a bacterial infection, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0069] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefpodoxime proxetil are formulated for oral administration. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefpodoxime proxetil are administered sequentially. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefpodoxime proxetil are administered concurrently.

[0070] Also disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene)COOH, or $-(C_1$ - C_6 alkylene)COO(C_1 - C_6 alkyl); and

(ii) cefpodoxime proxetil.

[0071] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0072] In some embodiments of a method of treating a bacterial infection, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0073] In some embodiments of a method of treating a bacterial infection, the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefpodoxime proxetil are formulated for oral administration. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered sequentially. In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered concurrently.

BRIEF DESCRIPTION OF THE DRAWINGS

[0074] **FIG. 1A** shows kill curves over 24 h for ceftibuten alone or in combination with Compound 2, vs. comparators in *E. coli* ESBL4 (CTX-M-15, TEM-1 producing).

[0075] **FIG. 1B** shows kill curves over 24 h for ceftibuten alone or in combination with Compound 2, vs. comparators in *K. pneumoniae* BAA 1705 (KPC-producing).

[0076] **FIG. 1C** shows kill curves over 24 h for ceftibuten alone or in combination with Compound 2, vs. comparators *E. cloacae* ECL01 (p99, TEM-1, ACT-1 producing).

[0077] **FIG. 1D** shows kill curves over 24 h for ceftibuten alone or in combination with Compound 2, vs. comparators in *E. coli* VER (OXA-48 producing).

[0078] **FIG. 2** shows the dissolution of 150 mg Compound 1-ethanolate capsule.

[0079] **FIG. 3** shows the dissolution of 100 mg Compound 1-ethanolate/100 mg ceftibuten combination capsule.

[0080] **FIG. 4** shows the cumulative % inhibited across 193 isolates of serine-beta-lactamase expressing Enterobacteriaceae.

[0081] **FIG. 5A** Log CFU of Bacteria in Kidneys Following Administration of Ceftibuten, Ceftibuten/Compound 2 (1:1), and Amoxicillin/Clavulanate (2:1) - *E. coli* expressing CTX-M-15.

[0082] **FIG. 5B** Log CFU of Bacteria in Kidneys Following Administration of Ceftibuten, Ceftibuten/Compound 2 (1:1), and Amoxicillin/Clavulanate (2:1) - *E. coli* expressing TEM-1 + CTX-M-15.

[0083] **FIG. 5C** Log CFU of Bacteria in Kidneys Following Administration of Ceftibuten, Ceftibuten/Compound 2 (1:1), and Amoxicillin/Clavulanate (2:1) - *E. coli* expressing KPC-2 + SHV-12.

[0084] **FIG. 6A** Log CFU of Bacteria in Bladders Following Administration of Ceftibuten, Ceftibuten/Compound 2 (1:1), and Amoxicillin/Clavulanate (2:1) - *E. coli* expressing CTX-M-15.

[0085] **FIG. 6B** Log CFU of Bacteria in Bladders Following Administration of Ceftibuten, Ceftibuten/Compound 2 (1:1), and Amoxicillin/Clavulanate (2:1) - *E. coli* expressing TEM-1 + CTX-M-15.

[0086] **FIG. 6C** Log CFU of Bacteria in Bladders Following Administration of Ceftibuten, Ceftibuten/Compound 2 (1:1), and Amoxicillin/Clavulanate (2:1) - *E. coli* expressing KPC-2 + SHV-12.

[0087] **FIG. 7A** Log CFU of Bacteria in Urine Following Administration of Ceftibuten, Ceftibuten/Compound 2 (1:1), and Amoxicillin/Clavulanate (2:1) - *E. coli* expressing CTX-M-15.

[0088] **FIG. 7B** Log CFU of Bacteria in Urine Following Administration of Ceftibuten, Ceftibuten/Compound 2 (1:1), and Amoxicillin/Clavulanate (2:1) - *E. coli* expressing TEM-1 + CTX-M-15.

[0089] FIG. 7C Log CFU of Bacteria in Urine Following Administration of Ceftributen, Ceftributen/Compound 2 (1:1), and Amoxicillin/Clavulanate (2:1) - *E. coli* expressing KPC-2 + SHV-12.

[0090] FIG. 8 Log CFU of Bacteria in Murine Thighs Following Administration of Ceftributen and Ceftributen/Compound 2 in a dose fractionation study of Compound 2 - *E. coli* expressing AmpC + CTX-M-15 + SHV + TEM.

[0091] FIG. 9 Log CFU of Bacteria in Murine Thighs Following Administration of Ceftributen and Ceftributen/Compound 2 in a dose fractionation study of Compound 2 - *E. coli* expressing AmpC + CTX-M-15 + TEM.



[0092] FIG. 10 shows the dissolution of 25/75 PEG1500/TPGS 250 mg Compound 1 Capsule.

[0093] FIG. 11 shows dissolution of 20/20/60 PG/PEG400/TPGS 250 mg Compound 1 Capsule.

DETAILED DESCRIPTION

Compounds

[0094] Disclosed herein are compounds of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);

or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl).

[0095] In some embodiments of a compound of formula (I), R^1 is hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl). In some embodiments of a compound of formula (I), R^1 is hydrogen or C_1 - C_6 alkyl. In some embodiments of a compound of formula (I), R^1 is hydrogen. In some embodiments of a compound of

formula (I), R¹ is C₁-C₆ alkyl. In some embodiments of a compound of formula (I), R¹ is -CO(C₁-C₆ alkyl).


[0096] In some embodiments of a compound of formula (II), each R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl). In some embodiments of a compound of formula (II), each R² is independently hydrogen or C₁-C₆ alkyl. In some embodiments of a compound of formula (II), each R² is independently hydrogen. In some embodiments of a compound of formula (II), each R² is independently C₁-C₆ alkyl. In some embodiments of a compound of formula (II), each R² is independently -CO(C₁-C₆ alkyl).

[0097] In some embodiments of a compound of formula (II), two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, -OH, -COOH, or -(C₁-C₆ alkylene)COOH. In some embodiments of a compound of formula (II), two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, -COOH, or -(C₁-C₆ alkylene)COOH. In some embodiments of a compound of formula (II), two R² are taken together to form a 5- or 6-membered heterocycloalkyl optionally substituted with oxo, -COOH, or -(C₁-C₆ alkylene)COOH. In some embodiments of a compound of formula (II), two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -COOH, or -COO(C₁-C₆ alkyl). In some embodiments of a compound of formula (II), two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, -OH, or -COOH. In some embodiments of a compound of formula (II), the 4- to 8-membered heterocycloalkyl formed when two R² are taken together contains 1 to 4 heteroatoms selected from oxygen and boron. In some embodiments of a compound of formula (II), the 5- or 6-membered heterocycloalkyl formed when two R² are taken together contains 1 to 4 heteroatoms selected from oxygen and boron.


[0098] In some embodiments of a compound of formula (II), two R² are taken together to form a moiety derived from an alpha-hydroxy carboxylic acid or a beta-hydroxy carboxylic acid. In some embodiments of a compound of formula (II), two R² are taken together to form a moiety derived from an alpha-hydroxy carboxylic acid. In some embodiments of a compound of formula (II), two R² are taken together to form a moiety derived from a beta-hydroxy carboxylic acid.

Compound 1


[0099] ((2-Ethylbutanoyl)oxy)methyl (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylate is as shown in the structure below:




. In some embodiments, ((2-ethylbutanoyl)oxy)methyl (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylate is also referred to as Compound 1. In some embodiments, Compound 1 exists in equilibrium as shown below:




. In some embodiments, Compound 1 exists in an equilibrium between the “closed” cyclic form (as shown

above) and the “open” acyclic form:  ((R)-(2-(3-(((2-ethylbutanoyl)oxy)methoxy)carbonyl)-2-hydroxyphenyl)-1-propionamidoethyl)boronic acid) which


exists in an equilibrium with  . In some embodiments, Compound 1 associates into intramolecular dimers, trimers, and any combinations thereof. In some embodiments, Compound 1 is in the form of a pharmaceutically acceptable salt. In some embodiments, Compound 1 is in the form of a pharmaceutically acceptable solvate. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein. In some embodiments, Compound 1 is in the form of a pharmaceutically acceptable salt and solvate. In some embodiments, Compound 1 is converted in vivo to release Compound 2.

Compound 1-ethanolate


[0100] In some embodiments, Compound 1 exists in solid form as a covalently bound solvate. In some embodiments, Compound 1 exists in solid form as a covalently bound ethanolate. In some



embodiments, Compound 1 in solid form is:
 ((2-ethylbutanoyl)oxy)methyl (R)-2-ethoxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylate. In some embodiments, Compound 1-ethanolate exists in equilibrium as shown below:




embodiments, the Compound 1-ethanolate converts to Compound 1 in contact with water:



Compound 2

[0101] (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid is as shown in the structure below:





“open” acyclic form:

((R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid). In some embodiments, Compound 2 associates into intramolecular dimers, trimers, and any combinations thereof.

Pharmaceutical Compositions

[0102] Disclosed herein are pharmaceutical compositions comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:



each R¹ and R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl); each alkyl being optionally substituted with halogen, -OH, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, or -COO(C₁-C₆ alkyl);

or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

- (ii) ceftibuten, cefexime, cefditoren pivoxil, cefpodoxime proxetil, amoxicillin, cefaclor, cephalixin, cefdinir, or cefuroxime.

[0103] Disclosed herein is a pharmaceutical composition comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:



each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);

or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) ceftibuten.

[0104] Disclosed herein is a pharmaceutical composition comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:


each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) cefixime.

[0105] Disclosed herein is a pharmaceutical composition comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:





formula (II);

wherein:


each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);

or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and


(ii) cefpodoxime proxetil.

[0106] Disclosed herein is a pharmaceutical composition comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



formula (I);



formula (II);


wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) cefditoren pivoxil.

[0107] In some embodiments, the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:




[0108] In some embodiments, the compound of formula (I); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0109] Disclosed herein is a pharmaceutical composition comprising:


- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



- (ii) ceftibuten.

[0110] Disclosed herein is a pharmaceutical composition comprising:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




- (ii) cefixime.

[0111] Disclosed herein is a pharmaceutical composition comprising:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a


pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



(ii) cefpodoxime proxetil.

[0112] Disclosed herein is a pharmaceutical composition comprising:

(i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



(ii) cefditoren pivoxil.

[0113] In some embodiments, the compositions described herein are administered via any of the accepted modes of administration for agents that serve similar utilities including, but not limited to, orally, subcutaneously, intravenously, intranasally, topically, transdermally, intraperitoneally, intramuscularly, intrapulmonarily, vaginally, rectally, or intraocularly. In some embodiments, the administration is oral administration. In some embodiments, the administration is intravenous administration. In some embodiments, the administration is intramuscular administration.

[0114] A summary of pharmaceutical compositions described herein can be found, for example, in Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Lieberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Dekker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999), herein incorporated by reference for such disclosure.

[0115] In some embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable excipient. The term "pharmaceutically acceptable excipient", as used herein, means one or more compatible solid or encapsulating substances, which are suitable for administration to a mammal. The term "compatible", as used herein, means that the components of the composition are capable of being commingled with the subject compound, and with each other, in a manner such that there is no interaction, which would substantially reduce the pharmaceutical efficacy of the composition under ordinary use situations. In some embodiments, the pharmaceutically acceptable

excipient is of sufficiently high purity and sufficiently low toxicity to render them suitable for administration preferably to an animal, preferably mammal, being treated.

[0116] The pharmaceutical compositions described herein include, but are not limited to, dispersions, solutions, liquids, gels, syrups, elixirs, slurries, suspensions, self-emulsifying dispersions, Self-Emulsifying Drug Delivery System (SED DS), liposomal dispersions, powders for reconstitution, powders, delayed release formulations, extended release formulations, pulsatile release formulations, immediate release formulations, controlled release formulations, fast melt formulations, tablets, capsules, pills, dragees, and effervescent formulations. In some embodiments, the pharmaceutical composition is formulated as a capsule. In some embodiments, the pharmaceutical composition is formulated as a tablet. In some embodiments, the pharmaceutical composition is formulated as liquid for injection.

Self-Emulsifying Drug Delivery System

[0117] Self-emulsifying drug delivery systems (SED DS) are isotropic mixtures of oil (or lipid), surfactant (with or without co-surfactant) and optionally a co-solvent which spontaneously emulsify when exposed to an aqueous medium with gentle agitation. SED DS have most commonly been studied to improve bioavailability of poorly water soluble drugs via oral administration.

[0118] Microemulsions arising from SMED DS (self-microemulsifying drug delivery system) are thermodynamically stable while regular emulsions are kinetically stable. According to the lipid formulation classification system (LFCS), SMED DS are characterized by a higher content of water-soluble components. These systems can achieve smaller-sized droplet dispersions and optical clarity, which is a desirable characteristic for improving currently existing ophthalmic emulsion formulations. SNED DS (self-nanoemulsifying drug delivery system) and their resultant nanoemulsions share many of the advantageous characteristics of SMED DS and microemulsions, but with the limitation of being only kinetically stable dispersions.

[0119] In some embodiments, the formulation does not comprise an oil. In some embodiments, the formulation comprises water-soluble surfactants and cosolvents. In some embodiments, the formulation is classified as a Type IV system by C. Pouton in *European Journal of Pharmaceutical Sciences*, 29 (2006), pp 278-287. (see table below).

[0120] The lipid formulation classification system:

Formulation type	Material	Characteristics	Advantages
Type I	Oils without surfactants (e.g., tri-, di-, and monoglycerides)	Nondispersing requires digestion	Generally recognized as safe (GRAS) status; simple; and excellent capsule compatibility
Type II	Oils and water insoluble surfactants	SED DS formed without water-soluble components	Unlikely to lose solvent capacity on dispersion

Formulation type	Material	Characteristics	Advantages
Type III	Oils, surfactants, and cosolvents (both water-insoluble and water-soluble excipients)	SEDDS/SMEDDS formed with water-soluble components	Clear or almost clear dispersion, drug absorption without digestion
Type IV	Water-soluble surfactants and cosolvents	Formulation disperses typically to form a micellar solution	Formulation has good solvent capacity for many drugs

[0121] In some embodiments, the pharmaceutical composition is formulated as an emulsion. In some embodiments, the pharmaceutical composition is formulated as a microemulsion.

Hydrophilic Solubilizer

[0122] In some embodiments, the pharmaceutical composition comprises a hydrophilic solubilizer.

In some embodiments, a suitable hydrophilic solubilizer is a hydrophilic polymer that include a variety of pharmaceutically acceptable hydrophilic agents that participate in the formation of the microemulsion, permit the accomplishment of the high levels of solubilized active ingredient, and are chemically compatible with the capsular material of the dosage form. In some embodiments, the hydrophilic solubilizer hydrophilic polymer.

[0123] In general, suitable hydrophilic polymers (i.e., two or more repeating monomer units) include, but are not limited to, pharmaceutically acceptable and water soluble polymers such as polyethylene glycols, methoxypolyethylene glycols, polyvinyl alcohols, polyvinyl pyrrolidones, and the like. The hydrophilic polymer can also include combinations or mixtures of pharmaceutically acceptable and water soluble polymers as well.

[0124] As used herein, “polyethylene glycol” or “PEG” means a liquid or solid polymer of the general formula $H(OCH_2CH_2)_nOH$, wherein n is at least 4. In certain embodiments, the hydrophilic polymer is a polyethylene glycol or a mixture of polyethylene glycols. Polyethylene glycols that can be used can include a wide range of molecular weights. In general, suitable polyethylene glycols that can be used with the invention include those from about PEG 400 to about PEG 8000, preferably PEG 400 to about PEG 1500, most preferably PEG 1000. Polyethylene glycols that can be used include, but are not limited to, PEG-400, PEG-600, PEG-1000, PEG-1450, PEG-1500, PEG-3350, or PEG-4600.

[0125] In some embodiments, the pharmaceutical composition includes one PEG or, alternatively, a mixture of two or more of the aforementioned polyethylene glycols. Representative mixtures include PEG-400/PEG-1000, PEG-400/PEG-1450, PEG-600/PEG-1000, PEG-600/PEG-1450.

[0126] The amount of hydrophilic polymer, e.g., polyethylene glycol to be used in the composition can vary provided a microemulsion is formed. In general, the amount of hydrophilic polymer is present in an amount from about 10% to about 95% per total composition.

Surfactants

[0127] In some embodiments, the pharmaceutical composition includes at least one surfactant. The use of a surfactant provides benefits in regard to dissolution or delivery stability. Suitable surfactants include, but are not limited to, nonionic, anionic and cationic surfactants, and combinations thereof.

[0128] Examples of suitable anionic surfactants that can be used, include, but are not limited to, sodium laurylsulfate or sodium dodecylsulfate. Examples of suitable cationic surfactants that can be used include, but are not limited to, cetyl trimethyl ammonium bromide (C-TAB). Examples of nonionic surfactants that can be used include, but are not limited to, polyoxyethylene stearates, such as polyoxyl 40 stearate (e.g., MYRJ® 52).

[0129] In addition to the above suitable surfactants for use in pharmaceutical composition include, but are not limited to, polyoxyethylene stearates, polyoxyethylene castor oil, polyoxyethylene sorbitan fatty acid esters (sorbitans), saturated polyglycolized glycerides, fatty acid esters of polyethylene glycol, hydroxylated lecithins, medium chain monoglycerides, medium chain fatty acid esters, polyethylene/propylene glycol copolymers, polyethylene glycol stearate, d- α -tocopheryl polyethylene glycol succinate, poloxyl stearate (e.g., Myrj® 52) and poloxyl castor oil. Polyoxyethylene sorbitan fatty acid esters (polysorbates) are non-ionic surfactants (detergents) that may consist of a mixture of fatty acids. Commercially available examples are Tween® 20 (polyoxyethylene (20) sorbitan monolaurate), Tween® 40 (polyoxyethylene (20) sorbitan monopalmitate), and Tween® 80 (polyoxyethylene (20) sorbitan monooleate).

[0130] Examples of other useful surfactants are saturated polyglycolized glycerides consisting of mono-, di-, or triglycerides; di-fatty acid esters of polyethylene glycol, e.g., Gelucire® 44/14; hydroxylated lecithins, e.g., Centrolene® A; medium chain monoglycerides, e.g., glyceryl monocaprylate (Imwitor® 308, Capmul® MCM C-8); caprylic/capric glycerides (Imwitor® 742); medium chain monoglycerides and diglycerides, e.g., glyceryl caprylate/caprate (Capmul® MCM); polyethylene/propylene glycol copolymers; block copolymers of ethylene oxide and propylene oxide (e.g., Poloxamer 188, Pluronic® F-68); ethoxylated castor oil (e.g., Cremophor® EL); and ethoxylated hydroxystearic acid (e.g., Solutol® HS 15). Some surfactants are solid or semisolid at room temperature, e.g., Poloxamer 188, glyceryl monocaprylate, Gelucire® 44/14, and any combinations thereof. In some embodiments, the surfactant is macroglycerol ricinoleate (Kolliphor EL® or Cremophor EL®), caprylocaproyl polyoxyl-8 glyceride (Labrasol®), polyoxyethylene hydrogenated castor oil 60 (HCO-60), polysorbate 80 (Tween®-80), polyoxyethylene sorbitan trioleate (Tween®-85), polyoxyethylene glyceryl trioleate (tagot-TO), or any combinations thereof. Additional surfactants are those found in The Handbook of Pharmaceutical Excipients, 2nd Ed., published by The Pharmaceutical Press, London and American Pharmaceutical Association (1994), a common text in the field, which is hereby incorporated by reference in its entirety.

[0131] In certain embodiments, the surfactant is a polyoxyl stearate. In a further embodiment, the polyoxyl stearate is polyoxyl 40 stearate (MYRJ® 52).

[0132] In some embodiments, the formulation comprises an emulsifier.

[0133] Exemplary emulsifiers used in lipid-based formulations:

Common name/type	Examples
Low HLB (<10) emulsifier	
Phosphatidylcholine and phosphatidylcholine/solvent mixtures	Phosphatidylcholine, phosphatidylcholine in propylene glycol, phosphatidylcholine in medium chain triglycerides, and phosphatidylcholine in safflower oil/ethanol
Unsaturated polyglycolized glycerides	Oleoyl macrogolglycerides, linoleoyl macrogolglycerides
Sorbitan esters	Sorbitan monooleate, sorbitan monostearate, sorbitan monolaurate, and sorbitan monopalmitate
High HLB (>10) emulsifier	
Polyoxyethylene sorbitan esters	Polysorbate 20, polysorbate 40, polysorbate 60, and polysorbate 80
Polyoxyl castor oil derivatives	Polyoxyl 35 castor oil, polyoxyl 40 hydrogenated castor oil
Polyoxyethylene polyoxypropylene block copolymer	Poloxamer 188, poloxamer 407
Saturated polyglycolized glycerides	Lauroyl macrogolglycerides, stearyl macrogolglycerides
PEG-8 caprylic/capric glycerides	Caprylocaproyl macrogolglycerides
Vitamin E derivative	Tocopherol PEG succinate

[0134] The amount of surfactant used in the pharmaceutical composition, when present, varies provided the amount is sufficient to participate in the formation and/or stabilization of the microemulsion. In general, the amount of surfactant if used is present in an amount from about 0.1% to about 60%—depending upon the particular surfactant employed. In some embodiments, the surfactant is present in an amount from about 0.1% to about 50%. In some embodiments, the surfactant is present in an amount from about 10% to about 60%. In some embodiments, the surfactant is present in an amount from about 20% to about 60%. In some embodiments, the surfactant is present in an amount from about 30% to about 60%. In some embodiments, the surfactant is present in an amount from about 40% to about 60%.

Lipid

[0135] In some embodiments, the pharmaceutical composition includes a lipid. In some embodiments, the lipid is a long- or medium-chain triglyceride oils with different degrees of saturation. In some embodiments, the lipid is a monoglyceride. In some embodiments, the lipid is a

diglyceride. In some embodiments, the lipid is propylene glycol monocaprylate (Capryol®), caprylic acid, capric acid, lauric acid, myristic acid, palmitic acid, stearic acid, oleic acid, ethyl oleate, soybean oil, glyceryl caprylate/caprate (Campul®) glyceryl behenate (Compritol® 888 ATO), glyceryl palmitostearate (Precirol® ATO 5), glyceryl monostearate (Geleol™), glyceryl monolinoleate (Maisine™ 35-1), glyceryl monooleate, (Peceol™), medium-chain triglycerides (Labrafac™ Lipophile WL1349), propylene glycol monolaurate (Lauroglycol™ 90), oleoyl macrogol-6 glycerides (Labrafil® M1944CS), polyglyceryl-3 dioleate (Plurol Oleique® CC 497), diethylene glycol monoethyl ether (Transcutol® HP), or any combinations thereof.

[0136] The amount of lipid used in the pharmaceutical composition, when present, varies provided the amount is sufficient to participate in the formation and/or stabilization of the microemulsion. In general, the amount of lipid if used is present in an amount from about 0.1% to about 50%—depending upon the particular lipid employed.

Solvents

[0137] In some embodiments, the pharmaceutical composition includes a solvent and/or a co-solvent. In some embodiments, the solvent/co-solvent is an alcohol (such as ethanol, benzyl alcohol, alkane diols and triols, glycol ethers, propylene glycol (PG), glycerol, tetraglycol, or polyethylene glycols), pyrrolidine derivatives, 2-pyrrolidone, iriacetin, or any combination thereof.

Tablets

[0138] Pharmaceutical preparations for oral use are obtained by mixing one or more solid excipient with one or more of the compounds described herein, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, for example, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methylcellulose, microcrystalline cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose; or others such as: polyvinylpyrrolidone (PVP or povidone) or calcium phosphate. If desired, disintegrating agents are added, such as the cross-linked croscarmellose sodium, polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. In some embodiments, dyestuffs or pigments are added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Capsule

[0139] Pharmaceutical preparations that are administered orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In

soft capsules, the active compounds are dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In some embodiments, stabilizers are added.

[0140] In some embodiments, the pharmaceutical composition is encapsulated. In some embodiments, the pharmaceutical composition is encapsulated into discrete units. In some embodiments, the pharmaceutical composition is enclosed in a capsule. In some embodiments, the capsule is formed using materials which include, but are not limited to, natural or synthetic gelatin, pectin, casein, collagen, protein, modified starch, polyvinylpyrrolidone, acrylic polymers, cellulose derivatives, or combinations thereof. In some embodiments, the capsule is coated. In some embodiments, the coating covering the capsule includes, but is not limited to, immediate release coatings, protective coatings, enteric or delayed release coatings, sustained release coatings, barrier coatings, seal coatings, or combinations thereof. In some embodiments, a capsule herein is hard or soft. In some embodiments, the capsule is seamless. In some embodiments, the shape and size of the capsule also vary. Examples of capsule shapes include, but are not limited to, round, oval, tubular, oblong, twist off, or a non-standard shape. The size of the capsule may vary according to the volume of the lipid-based formulation. In some embodiments, the size of the capsule is adjusted based on the volume of the lipid-based formulation. Hard or soft gelatin capsules may be manufactured in accordance with conventional methods as a single body unit comprising the standard capsule shape. A single-body soft gelatin capsule typically may be provided, for example, in sizes from 3 to 22 minims (1 minims being equal to 0.0616 ml) and in shapes of oval, oblong or others. The gelatin capsule may also be manufactured in accordance with conventional methods, for example, as a two-piece hard gelatin capsule, sealed or unsealed, typically in standard shape and various standard sizes, conventionally designated as (000), (00), (0), (1), (2), (3), (4), and (5). The largest number corresponds to the smallest size.

Powder for reconstitution

[0141] In some embodiments, the pharmaceutical composition for oral administration is formulated as a powder for reconstitution. In some embodiments, the powder for reconstitution is reconstituted with a liquid carrier to form a liquid formulation suitable for oral administration. In some embodiments, the powder for reconstitution is reconstituted with a liquid carrier to form a suspension.

[0142] Liquid carrier suitable for the powder formulations to be reconstituted into an oral formulation described herein are selected for a particular oral liquid formulation (solution, suspension, etc.) as well as other qualities such as clarity, toxicity, viscosity, compatibility with excipients, chemical inertness, palatability, odor, color and economy. Exemplary liquid carrier include water, ethyl alcohol, glycerin, propylene glycol, syrup (sugar or other sweetener based, e.g., Ora-Sweet® SF sugar-free flavored syrup), juices (apple, grape, orange, cranberry, cherry, tomato and the like), other beverages (tea, coffee, soft drinks, milk and the like), oils (olive, soybean, corn, mineral, castor and the like), and combinations or mixtures thereof. Certain liquid vehicles, e.g., oil and water, can be combined together to form emulsions. In some embodiments, water is used for as a liquid carrier.

[0143] Buffering agents maintain the pH of the liquid formulation. Non-limiting examples of buffering agents include, but are not limited to sodium bicarbonate, potassium bicarbonate, magnesium hydroxide, magnesium lactate, magnesium gluconate, aluminum hydroxide, aluminum hydroxide/sodium bicarbonate co precipitate, mixture of an amino acid and a buffer, a mixture of aluminum glycinate and a buffer, a mixture of an acid salt of an amino acid and a buffer, and a mixture of an alkali salt of an amino acid and a buffer. Additional buffering agents include citric acid, sodium citrate, sodium tartarate, sodium acetate, sodium carbonate, sodium polyphosphate, potassium polyphosphate, sodium pyrophosphate, potassium pyrophosphate, disodium hydrogenphosphate, dipotassium hydrogenphosphate, trisodium phosphate, tripotassium phosphate, sodium acetate, potassium metaphosphate, magnesium oxide, magnesium hydroxide, magnesium carbonate, magnesium silicate, calcium acetate, calcium glycerophosphate, calcium chloride, calcium hydroxide, calcium lactate, calcium carbonate, calcium bicarbonate, and other calcium salts. Some buffering agents also impart effervescent qualities when a powder is reconstituted in a solution.

[0144] In further embodiments, the powder for reconstitution described herein comprises additional excipients including, but not limited to, glidants, flavoring agents, coloring agents and thickeners. Additional excipients such as bulking agents, tonicity agents and chelating agents are within the scope of the embodiments.

[0145] Glidants are substances that improve flowability of a powder. Suitable glidants include, but are not limited to, calcium phosphate tribasic, calcium silicate, cellulose (powdered), colloidal silicon dioxide, magnesium silicate, magnesium trisilicate, silicon dioxide, starch, talc and the like.

[0146] In another embodiment, the powder for reconstitution described herein comprises a flavoring agent or flavorant to enhance the taste or aroma of the formulation in liquid form. Suitable natural or synthetic flavoring agents can be selected from standard reference books, for example Fenaroli's Handbook of Flavor Ingredients, 3rd edition (1995). Non-limiting examples of suitable natural flavors, some of which can readily be simulated with synthetic agents or combinations thereof, include almond, anise, apple, apricot, bergamot, blackberry, blackcurrant, blueberry, cacao, caramel, cherry, cinnamon, clove, coffee, coriander, cranberry, cumin, dill, eucalyptus, fennel, fig, ginger, grape, grapefruit, guava, hop, lemon, licorice, lime, malt, mandarin, molasses, nutmeg, mixed berry, orange, peach, pear, peppermint, pineapple, raspberry, rose, spearmint, strawberry, tangerine, tea, vanilla, wintergreen, etc.

[0147] In further embodiments, the powder for reconstitution described herein comprises a coloring agent for identity and/or aesthetic purposes. Suitable coloring agents illustratively include FD&C Red No. 3, FD&C Red No. 20, FD&C Red No. 40, FD&C Yellow No. 6, FD&C Blue No. 2, D&C Green No. 5, D&C Orange No. 5, caramel, ferric oxide and mixtures thereof.

[0148] In further embodiments, the powder for reconstitution described herein comprises a thickener. Thickeners impart viscosity or weight to the resultant liquid forms from the formulation described herein. Exemplary thickeners include dextrin, cellulose derivatives (ethylcellulose, hydroxyethyl

cellulose, methylcellulose, hypromellose, and the like) starches, pectin, polyethylene glycol, polyethylene oxide, trehalose and certain gums (xanthan gum, locust bean gum, etc.).

[0149] In certain embodiments, delivery systems for pharmaceutical compounds may be employed, such as, for example, liposomes and emulsions. In certain embodiments, compositions provided herein can also include an mucoadhesive polymer, selected from among, for example, carboxymethylcellulose, carbomer (acrylic acid polymer), poly(methylmethacrylate), polyacrylamide, polycarbophil, acrylic acid/butyl acrylate copolymer, sodium alginate and dextran.

[0150] In some embodiments, the pharmaceutical composition comprises dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, titanium dioxide, and triacetin.

[0151] In some embodiments, the pharmaceutical composition comprises colloidal silicon dioxide, crospovidone, low substituted hydroxy propyl cellulose, magnesium stearate and mannitol. The capsule shell contains the following inactive ingredients: ferric oxide black, ferric oxide red, gelatin, potassium hydroxide, propylene glycol, shellac, sodium lauryl sulfate, and titanium dioxide.

[0152] In some embodiments, the pharmaceutical composition comprises magnesium stearate, microcrystalline cellulose, and sodium starch glycolate inside a capsule. In some embodiments, the capsule shell contains gelatin, sodium lauryl sulfate, titanium dioxide, and polysorbate 80. In some embodiments, the capsule shell contains benzyl alcohol, sodium propionate, edetate calcium disodium, butylparaben, propylparaben, and methylparaben.

[0153] In some embodiments, the pharmaceutical composition comprises colloidal silicon dioxide, strawberry flavor, sodium benzoate, sucrose, and xanthan gum.

In some embodiments, the pharmaceutical composition comprises cherry flavoring, polysorbate 80, silicon dioxide, simethicone, sodium benzoate, sucrose, titanium dioxide, and xanthan gum.

[0154] Additional excipients are contemplated in the powder for reconstitution described herein. These additional excipients are selected based on function and compatibility with the powder formulation described herein and may be found, for example in *Remington: The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, PA: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*, (Easton, PA: Mack Publishing Co 1975); Lieberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms* (New York, NY: Marcel Decker 1980); and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed (Lippincott Williams & Wilkins 1999), herein incorporated by reference in their entirety.

[0155] In some embodiments, the pharmaceutically acceptable excipient is able to form a complex with the boron compounds such as alcohols (for example, but not limited to, ethanol, propylene glycol, benzyl alcohol, glycerin, fatty alcohol, sugars, alcohol polymers, cellulose based polymers, mono and diglycerides, or cyclodextrins), amines (meglumine, triethanolamine, eudragit, chitosan) and carboxylate/carboxy acids (acetic acid, organic acids, EDTA, polymer based acrylate polymers).

Boronic Ester Exchange

[0156] In some embodiments, the $-OR^1$ or $-OR^2$ in the compounds of Formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof react with an $-OH$ containing excipient or $-OH$ containing solvent present in the pharmaceutical composition to form a new boronic ester derived from the $-OH$ containing excipient or $-OH$ containing solvent.

[0157] In some embodiments, the $-OH$ containing excipient or $-OH$ containing solvent is a polymer (for example: polyvinyl alcohol), a monosaccharide (for example: glucose, galactose, fructose, or xylose), a disaccharide (for example: sucrose, lactose, maltose, or trehalose), a polyol (for example: sorbitol or mannitol), an oligosaccharide (for example: maltodextrin, raffinose, stachyose, or fructo-oligosaccharides), alcohols (for example: ethanol, propylene glycol, benzyl alcohol, glycerol, fatty alcohol, sugars, or alcohol polymers), cellulose based polymers, or cyclodextrins.

Method of Administration

[0158] In some embodiments, the pharmaceutical compositions described herein are administered orally.

[0159] In some embodiments, the pharmaceutical composition is administered once a day. In some embodiments, the pharmaceutical composition is administered twice a day. In some embodiments, the pharmaceutical composition is administered three times a day. In some embodiments, the pharmaceutical composition is administered four times a day.

[0160] In some embodiments, the pharmaceutical composition is administered once every 24 hours. In some embodiments, the pharmaceutical composition is administered once every 12 hours. In some embodiments, the pharmaceutical composition is administered once every 8 hours.

[0161] In some embodiments, the pharmaceutical composition is administered for about 1 to about 10 days. In some embodiments, the pharmaceutical composition is administered for about 5 to about 10 days. In some embodiments, the pharmaceutical composition is administered for about 5 days. In some embodiments, the pharmaceutical composition is administered for about 6 days. In some embodiments, the pharmaceutical composition is administered for about 7 days. In some embodiments, the pharmaceutical composition is administered for about 8 days. In some embodiments, the pharmaceutical composition is administered for about 9 days. In some embodiments, the pharmaceutical composition is administered for about 10 days.

[0162] In some embodiments, the pharmaceutical composition comprises about 50 mg to about 3200 mg of the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof. In some embodiments, the pharmaceutical composition comprises about 50 mg to about 800 mg of the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof. In some embodiments, the pharmaceutical composition comprises about 75 mg to about 1200 mg of

the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof. In some embodiments, the pharmaceutical composition comprises about 100 mg to about 1600 mg of the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof. In some embodiments, the pharmaceutical composition comprises about 150 mg to about 2400 mg of the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof.

[0163] In some embodiments, the pharmaceutical composition comprises about 50 mg, about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, about 300 mg, about 325 mg, about 350 mg, about 375 mg, about 400 mg, about 425 mg, about 450 mg, about 475 mg, about 500 mg, about 525 mg, about 550 mg, about 575 mg, about 600 mg, about 625 mg, about 650 mg, about 675 mg, about 700 mg, about 725 mg, about 750 mg, about 775 mg, about 800 mg, about 825 mg, about 850 mg, about 875 mg, about 900 mg, about 925 mg, about 950 mg, about 975 mg, about 1000 mg, about 1100 mg, about 1200 mg, about 1300 mg, about 1400 mg, about 1500 mg, about 1600 mg, about 1700 mg, about 1800 mg, about 1900 mg, about 2000 mg, about 2100 mg, about 2200 mg, about 2300 mg, about 2400 mg, about 2500 mg, about 2600 mg, about 2700 mg, about 2800 mg, about 2900 mg, about 3000 mg, about 3100 mg, or about 3200 mg of the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof.

[0164] In some embodiments, the pharmaceutical composition comprises about 100 to about 1000 mg of ceftibuten. In some embodiments, the pharmaceutical composition comprises about 100 to about 500 mg of ceftibuten. In some embodiments, the pharmaceutical composition comprises about 100, about 200, about 250, about 300, about 350, about 400, about 450, about 500, about 550, about 600, about 650, about 700, about 750, about 800, about 850, about 900, about 950, or about 1000 mg of ceftibuten. In some embodiments, the pharmaceutical composition comprises about 100 mg of ceftibuten. In some embodiments, the pharmaceutical composition comprises about 200 mg of ceftibuten. In some embodiments, the pharmaceutical composition comprises about 400 mg of ceftibuten.

[0165] In some embodiments, the pharmaceutical composition comprises between about 4:1 and about 1:4 weight ratio of the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof to ceftibuten.

[0166] In some embodiments, the daily dose of ceftibuten is about 100 mg to about 1000 mg. In some embodiments, the daily dose of ceftibuten is about 100, about 200, about 250, about 300, about 350, about 400, about 450, about 500, about 550, about 600, about 650, about 700, about 750, about 800, about 850, about 900, about 950, or about 1000 mg. In some embodiments, the daily dose of ceftibuten is about 100 mg. In some embodiments, the daily dose of ceftibuten is about 200 mg. In some embodiments, the daily dose of ceftibuten is about 400 mg.

[0167] In some embodiments, the pharmaceutical composition comprises about 100 to about 1000 mg of cefixime. In some embodiments, the pharmaceutical composition comprises about 100 to about 500 mg of cefixime. In some embodiments, the pharmaceutical composition comprises about 100, about 200, about 250, about 300, about 350, about 400, about 450, about 500, about 550, about 600, about 650, about 700, about 750, about 800, about 850, about 900, about 950, or about 1000 mg of cefixime. In some embodiments, the pharmaceutical composition comprises about 100 mg of cefixime. In some embodiments, the pharmaceutical composition comprises about 200 mg of cefixime. In some embodiments, the pharmaceutical composition comprises about 400 mg of cefixime.

[0168] In some embodiments, the daily dose of cefixime is about 100 mg to about 1000 mg. In some embodiments, the daily dose of cefixime is about 100, about 200, about 250, about 300, about 350, about 400, about 450, about 500, about 550, about 600, about 650, about 700, about 750, about 800, about 850, about 900, about 950, or about 1000 mg. In some embodiments, the daily dose of cefixime is about 100 mg. In some embodiments, the daily dose of cefixime is about 200 mg. In some embodiments, the daily dose of cefixime is about 400 mg.

[0169] In some embodiments, the pharmaceutical composition comprises between about 4:1 and about 1:4 weight ratio of the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof to cefixime.

[0170] In some embodiments, the pharmaceutical composition comprises about 100 to about 1000 mg of cefpodoxime proxetil. In some embodiments, the pharmaceutical composition comprises about 100 to about 500 mg of cefpodoxime proxetil. In some embodiments, the pharmaceutical composition comprises about 100, about 200, about 250, about 300, about 350, about 400, about 450, about 500, about 550, about 600, about 650, about 700, about 750, about 800, about 850, about 900, about 950, or about 1000 mg of cefpodoxime proxetil. In some embodiments, the pharmaceutical composition comprises about 100 mg of cefpodoxime proxetil. In some embodiments, the pharmaceutical composition comprises about 200 mg of cefpodoxime proxetil. In some embodiments, the pharmaceutical composition comprises about 400 mg of cefpodoxime proxetil.

[0171] In some embodiments, the pharmaceutical composition comprises between about 4:1 and about 1:4 weight ratio of the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof to cefpodoxime proxetil.

[0172] In some embodiments, the daily dose of cefpodoxime proxetil is about 100 mg to about 1000 mg. In some embodiments, the daily dose of cefpodoxime proxetil is about 100, about 200, about 250, about 300, about 350, about 400, about 450, about 500, about 550, about 600, about 650, about 700, about 750, about 800, about 850, about 900, about 950, or about 1000 mg. In some embodiments, the daily dose of cefpodoxime proxetil is about 200 mg.

[0173] In some embodiments, the pharmaceutical composition comprises about 100 to about 1000 mg of cefditoren pivoxil. In some embodiments, the pharmaceutical composition comprises about 100

to about 500 mg of cefditoren pivoxil. In some embodiments, the pharmaceutical composition comprises about 100, about 200, about 250, about 300, about 350, about 400, about 450, about 500, about 550, about 600, about 650, about 700, about 750, about 800, about 850, about 900, about 950, or about 1000 mg of cefditoren pivoxil. In some embodiments, the pharmaceutical composition comprises about 100 mg of cefditoren pivoxil. In some embodiments, the pharmaceutical composition comprises about 200 mg of cefditoren pivoxil. In some embodiments, the pharmaceutical composition comprises about 400 mg of cefditoren pivoxil.

[0174] In some embodiments, the pharmaceutical composition comprises between about 4:1 and about 1:4 weight ratio of the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof to cefditoren pivoxil.

[0175] In some embodiments, the daily dose of cefditoren pivoxil is about 100 mg to about 1000 mg. In some embodiments, the daily dose of cefditoren pivoxil is about 100, about 200, about 250, about 300, about 350, about 400, about 450, about 500, about 550, about 600, about 650, about 700, about 750, about 800, about 850, about 900, about 950, or about 1000 mg. In some embodiments, the daily dose of cefditoren pivoxil is about 200 mg. In some embodiments, the daily dose of cefditoren pivoxil is about 400 mg. In some embodiments, the daily dose of cefditoren pivoxil is about 800 mg.

Methods of Treatment

[0176] The present disclosure also provides methods for inhibiting bacterial growth, by, e.g., reducing bacterial resistance to a beta-lactam antibiotic, such methods comprising contacting a bacterial cell culture, or a bacterially infected cell culture, tissue, or organism, with a compound disclosed herein, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof. In some embodiments, the bacteria to be inhibited by administration of a compound disclosed herein, a pharmaceutically acceptable salt, a solvate thereof, or a pharmaceutically acceptable salt and solvate thereof are bacteria that are resistant to beta-lactam antibiotics. The term “resistant” is well-understood by those of ordinary skill in the art (see, e.g. Payne et al., *Antimicrobial Agents and Chemotherapy* 38 767-772 (1994), Hanaki et al., *Antimicrobial Agents and Chemotherapy* 30 1120-1126 (1995)).

[0177] These methods are useful for inhibiting bacterial growth in a variety of contexts. In certain other embodiments, a compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is administered to a mammal, including a human to prevent the growth of beta-lactam resistant bacteria in vivo. The method according to this embodiment comprises administering a therapeutically effective amount of a beta-lactamase inhibitor for a therapeutically effective period of time to a mammal, including a human. Preferably, the beta-lactamase inhibitor is administered in the form of a pharmaceutical composition as described herein. In some embodiments, an antibiotic is co-administered with the beta-lactamase inhibitor. In some embodiments, the antibiotic is a beta-lactam antibiotic. In some embodiments, the beta-lactam antibiotic is ceftibuten. In some embodiments, the beta-lactam antibiotic is cefixime. In some

embodiments, the beta-lactam antibiotic is cefpodoxime proxetil. In some embodiments, the beta-lactam antibiotic is cefditoren pivoxil.

[0178] In some embodiments, the beta-lactam antibiotic is amoxicillin. In some embodiments, the beta-lactam antibiotic is cefaclor. In some embodiments, the beta-lactam antibiotic is cephalixin. In some embodiments, the beta-lactam antibiotic is cefdinir. In some embodiments, the beta-lactam antibiotic is cefuroxime.

[0179] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:



- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) ceftibuten.

[0180] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
- (ii) ceftibuten.

[0181] In some embodiments of a method of treating a bacterial infection, the method comprises administering to the subject in need thereof:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:



each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);

or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

(ii) cefibuten.

[0182] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:

each R¹ and R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl); each alkyl being optionally substituted with halogen, -OH, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, or -COO(C₁-C₆ alkyl);

or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

(ii) cefixime.

[0183] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) cefixime.



[0184] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and

(ii) cefixime.

[0185] In some embodiments of a method of treating a bacterial infection, the method comprises administering to the subject in need thereof:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:



each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);

or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) cefixime.

[0186] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:

- each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);
- or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and
- (ii) cefixime.

[0187] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:



- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) cefpodoxime proxetil.

[0188] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
- (ii) cefpodoxime proxetil.

[0189] In some embodiments of a method of treating a bacterial infection, the method comprises administering to the subject in need thereof:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:



each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);

or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) cefpodoxime proxetil.

[0190] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);

or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) cefpodoxime proxetil.

[0191] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:



- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
- (ii) cefditoren pivoxil.

[0192] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:

- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
- (ii) cefditoren pivoxil.

[0193] In some embodiments of a method of treating a bacterial infection, the method comprises administering to the subject in need thereof:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:



wherein:


each R¹ and R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl); each alkyl being optionally substituted with halogen, -OH, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, or -COO(C₁-C₆ alkyl);


or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

- (ii) cefditoren pivoxil.

[0194] Disclosed herein is a method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:





formula (II);


wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) cefditoren pivoxil.

[0195] In some embodiments of a method of treating a bacterial infection, the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0196] In some embodiments of a method of treating a bacterial infection, the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



[0197] In some embodiments, the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.

[0198] In some embodiments, the infection that is treated or prevented comprises a bacteria that includes *Achromobacter xylosoxidans*, *Bacillus cereus*, *Bacillus anthracis*, *Elizabethkingia meningoseptica*, *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Pseudomonas acidovorans*, *Pseudomonas alcaligenes*, *Pseudomonas putida*, *Stenotrophomonas maltophilia*, *Burkholderia cepacia*,

Aeromonas hydrophilia, *Escherichia coli*, *Citrobacter freundii*, *Salmonella typhimurium*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella enteritidis*, *Shigella dysenteriae*, *Shigella flexneri*, *Shigella sonnei*, *Enterobacter cloacae*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Klebsiella oxytoca*, *Serratia marcescens*, *Francisella tularensis*, *Morganella morganii*, *Proteus mirabilis*, *Proteus vulgaris*, *Providencia alcalifaciens*, *Providencia rettgeri*, *Providencia stuartii*, *Acinetobacter baumannii*, *Acinetobacter calcoaceticus*, *Acinetobacter haemolyticus*, *Yersinia enterocolitica*, *Yersinia pestis*, *Yersinia pseudotuberculosis*, *Yersinia intermedia*, *Bordetella pertussis*, *Bordetella parapertussis*, *Bordetella bronchiseptica*, *Haemophilus influenzae*, *Haemophilus parainfluenzae*, *Haemophilus haemolyticus*, *Haemophilus parahaemolyticus*, *Haemophilus ducreyi*, *Pasteurella multocida*, *Pasteurella haemolytica*, *Branhamella catarrhalis*, *Helicobacter pylori*, *Campylobacter fetus*, *Campylobacter jejuni*, *Campylobacter coli*, *Borrelia burgdorferi*, *Vibrio cholerae*, *Vibrio parahaemolyticus*, *Legionella pneumophila*, *Listeria monocytogenes*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Kingella*, *Moraxella*, *Gardnerella vaginalis*, *Bacteroides fragilis*, *Bacteroides distasonis*, *Bacteroides 3452A* homology group, *Bacteroides vulgatus*, *Bacteroides ovalus*, *Bacteroides thetaiotaomicron*, *Bacteroides uniformis*, *Bacteroides eggerthii*, *Bacteroides splanchnicus*, *Clostridium difficile*, *Mycobacterium tuberculosis*, *Mycobacterium avium*, *Mycobacterium intracellulare*, *Mycobacterium leprae*, *Corynebacterium diphtheriae*, *Corynebacterium ulcerans*, *Streptococcus pneumoniae*, *Streptococcus agalactiae*, *Streptococcus pyogenes*, *Enterococcus faecalis*, *Enterococcus faecium*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Staphylococcus intermedius*, *Staphylococcus hyicus* subsp. *hyicus*, *Staphylococcus haemolyticus*, *Staphylococcus hominis*, or *Staphylococcus saccharolyticus*.

[0199] In some embodiments, the infection that is treated or prevented comprises a bacteria that includes *Elizabethkingia meningoseptica*, *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Stenotrophomonas maltophilia*, *Escherichia coli*, *Citrobacter freundii*, *Salmonella typhimurium*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella enteritidis*, *Shigella dysenteriae*, *Shigella flexneri*, *Shigella sonnei*, *Enterobacter cloacae*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Klebsiella oxytoca*, *Serratia marcescens*, *Acinetobacter calcoaceticus*, *Acinetobacter haemolyticus*, *Yersinia enterocolitica*, *Yersinia pestis*, *Yersinia pseudotuberculosis*, *Yersinia intermedia*, *Haemophilus influenzae*, *Haemophilus parainfluenzae*, *Haemophilus haemolyticus*, *Haemophilus parahaemolyticus*, *Helicobacter pylori*, *Campylobacter fetus*, *Campylobacter jejuni*, *Campylobacter coli*, *Vibrio cholerae*, *Vibrio parahaemolyticus*, *Legionella pneumophila*, *Listeria monocytogenes*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Moraxella*, *Bacteroides fragilis*, *Bacteroides vulgatus*, *Bacteroides ovalus*, *Bacteroides thetaiotaomicron*, *Bacteroides uniformis*, *Bacteroides eggerthii*, or *Bacteroides splanchnicus*.

[0200] In certain embodiments, the combinations described herein are useful in the treatment of bacterial infections.

[0201] In certain embodiments, the combinations described herein are useful in the treatment of acute bacterial exacerbations of chronic bronchitis (ABECB), acute bacterial otitis media, pharyngitis, or tonsillitis. In certain embodiments, the combinations described herein are useful in the treatment of pneumonia, urinary tract infections, enteritis, or gastroenteritis.

[0202] In certain embodiments, the combinations described herein are useful in the treatment of otitis media, strep throat, pneumonia, urinary tract infections, gonorrhea, or Lyme disease.

[0203] In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered sequentially. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered concurrently. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered in the same pharmaceutical composition. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered in separate pharmaceutical composition. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are provided in separate containers. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are provided in a single container. In some embodiments, the container is a bottle.

[0204] In some embodiments, the method comprises administering about 400 mg ceftibuten and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0205] In some embodiments, the method comprises administering about 600 mg ceftibuten and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0206] In some embodiments, the method comprises administering about 800 mg ceftibuten and about 200 mg to about 3200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0207] In some embodiments, the method comprises administering about 200 mg ceftibuten and about 50 mg to about 800 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0208] In some embodiments, the method comprises administering about 300 mg ceftibuten and about 75 mg to about 1200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0209] In some embodiments, the method comprises administering about 400 mg ceftibuten and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0210] In some embodiments, the method comprises administering about 600 mg ceftibuten and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0211] In some embodiments, the method comprises administering about 800 mg ceftibuten and about 200 mg to about 3200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0212] In some embodiments, the method comprises administering about 200 mg ceftibuten and about 50 mg to about 800 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0213] In some embodiments, the method comprises administering about 300 mg ceftibuten and about 75 mg to about 1200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0214] In some embodiments, the method comprises administering about 400 mg ceftibuten and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0215] In some embodiments, the method comprises administering about 600 mg ceftibuten and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0216] In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered sequentially. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered concurrently. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof

and cefixime are administered in the same pharmaceutical composition. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered in separate pharmaceutical composition. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are provided in separate containers. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are provided in a single container. In some embodiments, the container is a bottle.

[0217] In some embodiments, the method comprises administering about 400 mg cefixime and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0218] In some embodiments, the method comprises administering about 600 mg cefixime and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0219] In some embodiments, the method comprises administering about 800 mg cefixime and about 200 mg to about 3200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0220] In some embodiments, the method comprises administering about 200 mg cefixime and about 50 mg to about 800 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0221] In some embodiments, the method comprises administering about 300 mg cefixime and about 75 mg to about 1200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0222] In some embodiments, the method comprises administering about 400 mg cefixime and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0223] In some embodiments, the method comprises administering about 600 mg cefixime and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0224] In some embodiments, the method comprises administering about 800 mg cefixime and about 200 mg to about 3200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0225] In some embodiments, the method comprises administering about 200 mg cefixime and about 50 mg to about 800 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0226] In some embodiments, the method comprises administering about 300 mg cefixime and about 75 mg to about 1200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0227] In some embodiments, the method comprises administering about 400 mg cefixime and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0228] In some embodiments, the method comprises administering about 600 mg cefixime and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0229] In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered sequentially. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered concurrently. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered in the same pharmaceutical composition. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered in separate pharmaceutical composition. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are provided in separate containers. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are provided in a single container. In some embodiments, the container is a bottle.

[0230] In some embodiments, the method comprises administering about 400 mg cefpodoxime proxetil and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0231] In some embodiments, the method comprises administering about 600 mg cefpodoxime proxetil and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0232] In some embodiments, the method comprises administering about 800 mg cefpodoxime proxetil and about 200 mg to about 3200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0233] In some embodiments, the method comprises administering about 200 mg cefpodoxime proxetil and about 50 mg to about 800 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0234] In some embodiments, the method comprises administering about 300 mg cefpodoxime proxetil and about 75 mg to about 1200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0235] In some embodiments, the method comprises administering about 400 mg cefpodoxime proxetil and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0236] In some embodiments, the method comprises administering about 600 mg cefpodoxime proxetil and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0237] In some embodiments, the method comprises administering about 800 mg cefpodoxime proxetil and about 200 mg to about 3200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0238] In some embodiments, the method comprises administering about 200 mg cefpodoxime proxetil and about 50 mg to about 800 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0239] In some embodiments, the method comprises administering about 300 mg cefpodoxime proxetil and about 75 mg to about 1200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0240] In some embodiments, the method comprises administering about 400 mg cefpodoxime proxetil and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0241] In some embodiments, the method comprises administering about 600 mg cefpodoxime proxetil and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0242] In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered sequentially. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered concurrently. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered in the same pharmaceutical composition. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered in separate pharmaceutical composition. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are provided in separate containers. In some embodiments, the compound of formula (I) or (II), a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are provided in a single container. In some embodiments, the container is a bottle.

[0243] In some embodiments, the method comprises administering about 400 mg cefditoren pivoxil and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0244] In some embodiments, the method comprises administering about 600 mg cefditoren pivoxil and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0245] In some embodiments, the method comprises administering about 800 mg cefditoren pivoxil and about 200 mg to about 3200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0246] In some embodiments, the method comprises administering about 200 mg cefditoren pivoxil and about 50 mg to about 800 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0247] In some embodiments, the method comprises administering about 300 mg cefditoren pivoxil and about 75 mg to about 1200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0248] In some embodiments, the method comprises administering about 400 mg cefditoren pivoxil and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically

acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0249] In some embodiments, the method comprises administering about 600 mg cefditoren pivoxil and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0250] In some embodiments, the method comprises administering about 800 mg cefditoren pivoxil and about 200 mg to about 3200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0251] In some embodiments, the method comprises administering about 200 mg cefditoren pivoxil and about 50 mg to about 800 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0252] In some embodiments, the method comprises administering about 300 mg cefditoren pivoxil and about 75 mg to about 1200 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0253] In some embodiments, the method comprises administering about 400 mg cefditoren pivoxil and about 100 mg to about 1600 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0254] In some embodiments, the method comprises administering about 600 mg cefditoren pivoxil and about 150 mg to about 2400 mg of a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0255] In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered sequentially. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered concurrently. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered in the same pharmaceutical composition. In some embodiments, (R)-2-hydroxy-3-

propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered in separate pharmaceutical composition. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are provided in separate containers. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are provided in a single container. In some embodiments, the container is bottle.

[0256] In some embodiments, the method comprises administering about 400 mg ceftibuten and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0257] In some embodiments, the method comprises administering about 600 mg ceftibuten and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0258] In some embodiments, the method comprises administering about 800 mg ceftibuten and about 200 mg to about 3200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0259] In some embodiments, the method comprises administering about 200 mg ceftibuten and about 50 mg to about 800 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0260] In some embodiments, the method comprises administering about 300 mg ceftibuten and about 75 mg to about 1200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0261] In some embodiments, the method comprises administering about 400 mg ceftibuten and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0262] In some embodiments, the method comprises administering about 600 mg ceftibuten and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0263] In some embodiments, the method comprises administering about 800 mg ceftibuten and about 200 mg to about 3200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0264] In some embodiments, the method comprises administering about 200 mg ceftibuten and about 50 mg to about 800 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0265] In some embodiments, the method comprises administering about 300 mg ceftibuten and about 75 mg to about 1200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0266] In some embodiments, the method comprises administering about 400 mg ceftibuten and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0267] In some embodiments, the method comprises administering about 600 mg ceftibuten and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0268] In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-

hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered sequentially. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered concurrently. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered in the same pharmaceutical composition. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered in separate pharmaceutical composition. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are provided in separate containers. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are provided in a single container. In some embodiments, the container is bottle.

[0269] In some embodiments, the method comprises administering about 400 mg cefixime and

[0270] In some embodiments, the method comprises administering about 400 mg cefixime and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0271] In some embodiments, the method comprises administering about 600 mg cefixime and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0272] In some embodiments, the method comprises administering about 800 mg cefixime and about 200 mg to about 3200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0273] In some embodiments, the method comprises administering about 200 mg cefixime and about 50 mg to about 800 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0274] In some embodiments, the method comprises administering about 300 mg cefixime and about 75 mg to about 1200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0275] In some embodiments, the method comprises administering about 400 mg cefixime and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0276] In some embodiments, the method comprises administering about 600 mg cefixime and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0277] In some embodiments, the method comprises administering about 800 mg cefixime and about 200 mg to about 3200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0278] In some embodiments, the method comprises administering about 200 mg cefixime and about 50 mg to about 800 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0279] In some embodiments, the method comprises administering about 300 mg cefixime and about 75 mg to about 1200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0280] In some embodiments, the method comprises administering about 400 mg cefixime and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-

benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0281] In some embodiments, the method comprises administering about 600 mg cefixime and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0282] In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered sequentially. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered concurrently. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered in the same pharmaceutical composition. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered in separate pharmaceutical composition. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are provided in separate containers. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are provided in a single container. In some embodiments, the container is bottle.

[0283] In some embodiments, the method comprises administering about 400 mg cefpodoxime proxetil and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0284] In some embodiments, the method comprises administering about 600 mg cefpodoxime proxetil and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0285] In some embodiments, the method comprises administering about 800 mg cefpodoxime proxetil and about 200 mg to about 3200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0286] In some embodiments, the method comprises administering about 200 mg cefpodoxime proxetil and about 50 mg to about 800 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0287] In some embodiments, the method comprises administering about 300 mg cefpodoxime proxetil and about 75 mg to about 1200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0288] In some embodiments, the method comprises administering about 400 mg cefpodoxime proxetil and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0289] In some embodiments, the method comprises administering about 600 mg cefpodoxime proxetil and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0290] In some embodiments, the method comprises administering about 800 mg cefpodoxime proxetil and about 200 mg to about 3200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0291] In some embodiments, the method comprises administering about 200 mg cefpodoxime proxetil and about 50 mg to about 800 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-

benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0292] In some embodiments, the method comprises administering about 300 mg cefpodoxime proxetil and about 75 mg to about 1200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0293] In some embodiments, the method comprises administering about 400 mg cefpodoxime proxetil and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0294] In some embodiments, the method comprises administering about 600 mg cefpodoxime proxetil and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0295] In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered sequentially. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered concurrently. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered in the same pharmaceutical composition. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered in separate pharmaceutical composition. In some embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are provided in separate containers. In some

embodiments, (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are provided in a single container. In some embodiments, the container is bottle.

[0296] In some embodiments, the method comprises administering about 400 mg cefditoren pivoxil and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0297] In some embodiments, the method comprises administering about 600 mg cefditoren pivoxil and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0298] In some embodiments, the method comprises administering about 800 mg cefditoren pivoxil and about 200 mg to about 3200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 24 hours.

[0299] In some embodiments, the method comprises administering about 200 mg cefditoren pivoxil and about 50 mg to about 800 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0300] In some embodiments, the method comprises administering about 300 mg cefditoren pivoxil and about 75 mg to about 1200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0301] In some embodiments, the method comprises administering about 400 mg cefditoren pivoxil and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0302] In some embodiments, the method comprises administering about 600 mg cefditoren pivoxil and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-

hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0303] In some embodiments, the method comprises administering about 800 mg cefditoren pivoxil and about 200 mg to about 3200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 12 hours.

[0304] In some embodiments, the method comprises administering about 200 mg cefditoren pivoxil and about 50 mg to about 800 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0305] In some embodiments, the method comprises administering about 300 mg cefditoren pivoxil and about 75 mg to about 1200 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0306] In some embodiments, the method comprises administering about 400 mg cefditoren pivoxil and about 100 mg to about 1600 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

[0307] In some embodiments, the method comprises administering about 600 mg cefditoren pivoxil and about 150 mg to about 2400 mg of (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; both administered every 8 hours.

Certain Definitions

[0308] Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of embodiments described herein, certain preferred methods, devices, and materials are now described.

[0309] As used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to “an excipient” is a reference to one or more excipients and equivalents thereof known to those skilled in the art, and so forth.

[0310] The term “about” is used to indicate that a value includes the standard level of error for the device or method being employed to determine the value.

[0311] The use of the term “or” is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and to “and/or.”

[0312] The terms “comprise,” “have” and “include” are open-ended linking verbs. Any forms or tenses of one or more of these verbs, such as “comprises,” “comprising,” “has,” “having,” “includes” and “including,” are also open-ended. For example, any method that “comprises,” “has” or “includes” one or more steps is not limited to possessing only those one or more steps and also covers other unlisted steps.

[0313] “Optional” or “optionally” may be taken to mean that the subsequently described structure, event or circumstance may or may not occur, and that the description includes instances where the events occurs and instances where it does not.

[0314] As used herein, the term “therapeutic” means an agent utilized to treat, combat, ameliorate, prevent or improve an unwanted condition or disease of a patient.

[0315] “Administering” when used in conjunction with a therapeutic means to administer a therapeutic systemically or locally, as directly into or onto a target tissue, or to administer a therapeutic to a patient whereby the therapeutic positively impacts the tissue to which it is targeted. “Administering” a pharmaceutical composition may be accomplished by injection, topical administration, and oral administration or by other methods alone or in combination with other known techniques.

[0316] The term “animal” as used herein includes, but is not limited to, humans and non-human vertebrates such as wild, domestic and farm animals. As used herein, the terms “patient,” “subject” and “individual” are intended to include living organisms in which certain conditions as described herein can occur. Examples include humans, monkeys, cows, sheep, goats, dogs, cats, mice, rats, and transgenic species thereof. In a preferred embodiment, the patient is a primate. In certain embodiments, the primate or subject is a human. In certain instances, the human is an adult. In certain instances, the human is child. In further instances, the human is 12 years of age or younger. In certain instances, the human is elderly. In other instances, the human is 60 years of age or older. Other examples of subjects include experimental animals such as mice, rats, dogs, cats, goats, sheep, pigs, and cows.

[0317] By “pharmaceutically acceptable,” it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the composition and not deleterious to the recipient thereof.

[0318] The term “pharmaceutical composition” means a composition comprising at least one active ingredient, whereby the composition is amenable to investigation for a specified, efficacious outcome in a mammal (for example, without limitation, a human). Those of ordinary skill in the art will

understand and appreciate the techniques appropriate for determining whether an active ingredient has a desired efficacious outcome based upon the needs of the artisan.

[0319] A “therapeutically effective amount” or “effective amount” as used herein refers to the amount of active compound or pharmaceutical agent that elicits a biological or medicinal response in a tissue, system, animal, individual or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes one or more of the following: (1) preventing the disease; for example, preventing a disease, condition or disorder in an individual that may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease, (2) inhibiting the disease; for example, inhibiting a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., arresting further development of the pathology and/or symptomatology), and (3) ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., reversing the pathology and/or symptomatology).

[0320] The terms “treat,” “treated,” “treatment,” or “treating” as used herein refers to both therapeutic treatment in some embodiments and prophylactic or preventative measures in other embodiments, wherein the object is to prevent or slow (lessen) an undesired physiological condition, disorder or disease, or to obtain beneficial or desired clinical results. For the purposes described herein, beneficial or desired clinical results include, but are not limited to, alleviation of symptoms; diminishment of the extent of the condition, disorder or disease; stabilization (i.e., not worsening) of the state of the condition, disorder or disease; delay in onset or slowing of the progression of the condition, disorder or disease; amelioration of the condition, disorder or disease state; and remission (whether partial or total), whether detectable or undetectable, or enhancement or improvement of the condition, disorder or disease. Treatment includes eliciting a clinically significant response without excessive levels of side effects. Treatment also includes prolonging survival as compared to expected survival if not receiving treatment. A prophylactic benefit of treatment includes prevention of a condition, retarding the progress of a condition, stabilization of a condition, or decreasing the likelihood of occurrence of a condition. As used herein, “treat,” “treated,” “treatment,” or “treating” includes prophylaxis in some embodiments. As used herein, “treat,” “treated,” “treatment,” or “treating” does not include prophylaxis in some embodiments.

[0321] The term “emulsion” as used herein refers to a colloidal dispersion comprising water and organic components including hydrophobic (lipophilic) organic components. Generally, a traditional emulsion is comprised of oil droplets (>about 200 nm) dispersed in water, resulting in a milky white liquid which is not stable.

[0322] The term “microemulsion,” as used herein, refers to a dispersion comprising water and organic components including hydrophobic (lipophilic) organic components, wherein the droplets or

particles formed from the organic components have an average maximum dimension of less than about 200 nm.

[0323] “Alkyl” refers to a substituted straight-chain, or branched-chain saturated hydrocarbon monoradical having from one to about ten carbon atoms, from one to six carbon atoms, or from one to four carbon atoms. Examples include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, isopentyl, neopentyl, tert-amyl and hexyl, and longer alkyl groups, such as heptyl, octyl, and the like. Whenever it appears herein, a numerical range such as “C₁-C₆ alkyl” means that the alkyl group consists of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, 4 carbon atoms, 5 carbon atoms or 6 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated. In some embodiments, the alkyl is a C₁-C₁₀ alkyl, a C₁-C₉ alkyl, a C₁-C₈ alkyl, a C₁-C₇ alkyl, a C₁-C₆ alkyl, a C₁-C₅ alkyl, a C₁-C₄ alkyl, a C₁-C₃ alkyl, a C₁-C₂ alkyl, or a C₁ alkyl.

[0324] “Heterocycloalkyl” refers to a stable 3- to 24-membered partially or fully saturated ring radical comprising 2 to 23 carbon atoms and from one to 8 heteroatoms selected from the group consisting of boron, nitrogen, oxygen, phosphorous, and sulfur. In some embodiments, the heterocycloalkyl comprises 1 to 4 heteroatoms selected from boron and oxygen. Unless stated otherwise specifically in the specification, the heterocycloalkyl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused (when fused with an aryl or a heteroaryl ring, the heterocycloalkyl is bonded through a non-aromatic ring atom) or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heterocycloalkyl radical may be optionally oxidized; the nitrogen atom may be optionally quaternized. Representative heterocycloalkyls include, but are not limited to, heterocycloalkyls having from two to fifteen carbon atoms (C₂-C₁₅ heterocycloalkyl), from two to ten carbon atoms (C₂-C₁₀ heterocycloalkyl), from two to eight carbon atoms (C₂-C₈ heterocycloalkyl), from two to six carbon atoms (C₂-C₆ heterocycloalkyl), from two to five carbon atoms (C₂-C₅ heterocycloalkyl), or two to four carbon atoms (C₂-C₄ heterocycloalkyl). In some embodiments, the heterocycloalkyl is a 3- to 6-membered heterocycloalkyl. In some embodiments, the heterocycloalkyl is a 5- to 6-membered heterocycloalkyl. In some embodiments, the heterocycloalkyl is a 4- to 8-membered heterocycloalkyl. Examples of such heterocycloalkyl radicals include, but are not limited to, aziridinyl, azetidiny, dioxolanyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazoliny, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholinyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, quinuclidinyl, thiazolidinyl, tetrahydrofuryl, trithianyl, tetrahydropyranyl, thiomorpholinyl, thiamorpholinyl, 1-oxo-thiomorpholinyl, 1,1-dioxo-thiomorpholinyl, 1,3-dihydroisobenzofuran-1-yl, 3-oxo-1,3-dihydroisobenzofuran-1-yl,

methyl-2-oxo-1,3-dioxol-4-yl, and 2-oxo-1,3-dioxol-4-yl. In some embodiments, the heterocycloalkyl is 1,3,2-dioxaborolane, 1,3,2-dioxaborinane, or 1,3,2-dioxaborocane.

EXAMPLES

Example 1. Experimental Method for Beta-Lactamase Enzyme Assays.

Part 1. Isolation of Beta-Lactamases.

[0325] *E. coli* BL21(DE3) bacterial cells carrying expression plasmids for the individual beta-lactamases (SHV-5, KPC-2, p99AmpC, CTX-M-15, CMY-2, IMP-1, NDM-1, VIM-2, OXA-1, and OXA 48; expressed as native untagged proteins), were grown in 1 L of Superbroth (Teknova Inc. Hollister, CA) supplemented with 100 µg/mL kanamycin selection and 1x 5052 (0.5 % glycerol, 0.05 % glucose and 0.2 % α-lactose) at 35 °C for 18-20 hours. Cells were harvested by centrifugation (4,000 x g, 4 °C, 20 min) and resuspended in 50 mL of 10 mM HEPES pH 7.5 (1/20 of the initial volume). The cells were lysed by sonication (5 pulses of 45 seconds) at 45 W on ice. The lysates were clarified by centrifugation at 10,000 x g for 40 minutes at 4 °C. Samples were diluted 5-fold in 50 mM sodium acetate pH 5.0, stored overnight at 4 °C, after which they were centrifuged at 10,000 x g for 30 minutes to clarify, and filtered through 0.45 µm filters. The samples were loaded onto a 5 mL Capto S sepharose cation exchange column (GE Healthcare) pre-equilibrated with 50 mM sodium acetate pH 5.0. The column was washed with 5 column volumes of 50 mM sodium acetate pH 5.0 to wash out unbound protein and a linear gradient of NaCl (0 to 500 mM) was used to elute the protein (over 16 CV) from the column. Fractions were assayed for beta-lactamase activity using Centa (Calbiochem, Gibbstown, NJ) or Nitrocefin (EMD Millipore chemicals, Darmstadt, Germany) as a reporter beta-lactamase substrate for activity in the isolated fractions. Active fractions were pooled, concentrated and further purified by gel filtration chromatography on a Superdex 75 prep grade gel filtration column (GE Healthcare, Piscataway, NJ) pre-equilibrated in 50 mM Hepes pH 7.5, 150 mM NaCl. Active fractions were pooled, concentrated, quantitated by BCA protein determination (Thermo Scientific, Rockford, IL), dialyzed into PBS and frozen at -80 °C in 20 % glycerol until use.

[0326] For VIM-2 metallo beta-lactamase, the procedure was identical with the following exceptions, first the protein was not pH adjusted to pH 5 with 50 mM sodium acetate, second, the chromatography step was changed to a 5 ml Q sepharose anion exchange column pre-equilibrated with 50 mM Hepes pH 7.5, and elution of the protein was achieved by a linear gradient of NaCl (0-600 mM). Finally, the VIM-2 purification required a second run (3rd step) on the Q sepharose anion exchange column to achieve acceptable purity (>90%).

Part 2. Inhibition of Diverse Beta-Lactamases.

[0327] To determine the level of inhibition of beta-lactamase enzymes, Compound 2 was diluted in PBS at pH 7.4 to yield concentrations ranging from 100 to 0.00005 µM in 96-well microtiter plates. An equal volume of diluted enzyme stock was added, and the plates were incubated at 37 °C for 15

min. Nitrocefin was used as substrate for p99 AmpC, CMY-2, IMP-1, VIM-2 and OXA-1 and OXA-48, and dispensed into each well at a final concentration of 100 μ M. Absorbance at 486 nm was immediately monitored for 10 min using a Biotek Powerwave XS2 microplate spectrophotometer using the GEN5 software package (Biotek Instruments, Winooski VT). In an analogous fashion, imipenem was used as substrate for KPC-2, and Cefotaxime was used for CTX-M-15 and SHV-5, while changes in absorbance upon hydrolysis of the beta-lactam ring were monitored at 300 nm and 260 nm respectively in UV-transparent 96-well microtiter assay plates. Maximum rates of hydrolysis were compared to those in control wells (without inhibitors), and percentages of enzyme inhibition were calculated for each concentration of inhibitor. The concentration of inhibitor needed to reduce the initial rate of hydrolysis of substrate by 50% (IC_{50}) was calculated as the residual activity of beta-lactamase at 486 nm using GraFit version 7 kinetics software package (Erithacus Software, Surrey, UK).

[0328] Using the methodology described above, Compound 2 was evaluated for the ability to inhibit beta-lactamase enzymes from all four Ambler classifications (A through D). The results of these assays are summarized in Table 1 for representative enzymes across different subtypes (note SHV-5 represents an Ambler Class A Extended Spectrum beta-Lactamases, KPC-2 exemplifies a Class A carbapenemase, AmpC represents a chromosomal Class C, OXA-1 and OXA-48 represent Class D oxacillinases, and VIM-2, NDM-1, and IMP-1 represent Class B zinc-dependent metallo-beta-lactamases also possessing carbapenemase activity).

[0329] Steady state kinetic parameters for reversible inhibition of beta-lactam hydrolysis by Compound 2 were determined with a subset of enzymes (Table 2). The onset of inhibition, or rate of covalent complex formation (k_2/K_i), was determined by monitoring the progress of beta-lactam hydrolysis by the respective enzymes in the presence of increasing concentrations of Compound 2. Assays (200 μ l final volume) were performed in PBS in triplicate using 96-well microtiter plates. Cephalothin was used as the substrate for p99 AmpC and CTX-M-15 (50 μ M for p99 AmpC; 75 μ M for CTX-M-15) and reactions were initiated by addition of enzyme (0.2 nM p99 AmpC; 3 nM CTX-M-15). The decrease in the absorbance at λ 260 nm was recorded continuously on a BioTek Powerwave XS2 microplate reader. The concentrations of Compound 2 tested were: 20 μ M, 10 μ M, 5 μ M, 2.5 μ M, 1.25 μ M, 0.625 μ M, 0.313 μ M, 0.156 μ M, 0.0781 μ M, 0.0391 μ M, 0.0195 μ M and 0 μ M. SHV-5 reactions were initiated with 40 nM enzyme, using cefotaxime (100 μ M) as the substrate. The concentrations of Compound 2 were as above, and the reaction progress monitored continuously by measuring the reduction in absorbance at λ 260 nm. For KPC-2, imipenem (75 μ M) was used as the substrate, and reactions initiated with 3 nM enzyme. Reaction progress was monitored by measuring the reduction in absorbance at λ 300 nm in continuous fashion. The concentrations of Compound 2 tested were: 10 μ M, 5 μ M, 2.5 μ M, 1.25 μ M, 0.625 μ M, 0.313 μ M, 0.156 μ M, 0.0781 μ M, 0.0391 μ M, 0.0195 μ M, 0.0098 and 0 μ M. Time courses for reaction progress were fit to the following equation, using Prism 7.04 (GraphPad Software, LaJolla, USA) to obtain k_{obs} :

$$A_i = A_0 + v_s t + (v_0 - v_s) \left[\frac{1 - e^{-k_{\text{obs}} t}}{k_{\text{obs}}} \right]$$

[0330] In the above equation, A_i is the observed absorbance, A_0 the initial absorbance, v_0 the initial velocity, v_s the steady state velocity and t , is time. k_2/K_i was subsequently derived using the following equation, where $[I]$ is the concentration of Compound 2, $[S]$ the substrate concentration and K_m the Michaelis constant for the substrate:

$$k_{\text{obs}} = k_{-2} + \frac{k_2}{K_i} \left[\frac{[I]}{1 + \frac{[S]}{K_m}} \right]$$

[0331] Off-rates for Compound 2 with the various beta-lactamases were determined by jump dilution experiments, performed in triplicate. Enzyme and inhibitor were incubated at room temperature for 10 minutes. Enzyme:inhibitor complexes were then diluted 800-fold into reaction buffer (50 mM Hepes, pH 7.0, 0.1 mg/ml BSA), and 20 μ L of the diluted reaction immediately added to 180 μ L of 110 μ M nitrocefin in a 96-well microtiter plate, and the absorbance at λ 486 nm measured continuously on a BioTek Powerwave XS2 microplate reader. The resulting progress curves were fit to a single exponential, from which k_{off} was derived. Half-life was determined using:

$$t_{1/2} = \frac{0.693}{k_{\text{off}}}$$

[0332] To determine K_i , plots of the inhibitor concentration versus inverse the initial velocity were fit to a linear equation, and the observed K_i corrected for substrate concentration and affinity by:

$$K_i = \frac{K_i^{\text{observed}}}{1 + \frac{[S]}{K_m}}$$

Table 1. Enzyme inhibition of Compound 2 against representative beta-lactamase enzymes from Ambler Classes A, B, C and D.

Amber Class	Enzyme	IC ₅₀ (μ M)
Class A (serine-based)	CTX-M-15	0.018
	KPC-2	0.08
	SHV-5	0.368
Class C (serine-based)	p99 AmpC	0.014
	CMY-2	0.014
Class D (serine-based)	OXA-1	0.066
	OXA-48	0.317
Class B (metallo-based)	VIM-2	9.04
	NDM-1	38.1
	IMP-1	>100

Table 2. Kinetic parameters of beta-lactamase inhibition by Compound 2.

Enzyme	k_2/K_i ($10^4 \text{ M}^{-1} \text{ s}^{-1}$)	k_2 (10^{-4} s^{-1})	$t_{1/2}$ (min)	K_i (μM)
KPC-2	2.9 ± 0.07	2.5 ± 0.1	46 ± 2.0	0.11
CTX-M-15	4.8 ± 0.9	4.5 ± 0.1	26 ± 0.6	0.01
SHV-5	1.1 ± 0.16	12.7 ± 0.07	6.5 ± 0.04	0.041
p99 AmpC	6.0 ± 0.6	24 ± 0.5	5.0 ± 0.1	0.02

Example 2: In vitro Antibacterial Assays Demonstrating Potentiation of Beta-Lactam Antibiotics by Beta-Lactamase Inhibition.

[0333] To determine the ability of test compounds to potentiate the antibacterial activity of various oral cephalosporins against beta-lactamase expressing Enterobacteriaceae, classic cell-based broth microdilution MIC assays were employed. A panel of 40 *E. coli* and *K. pneumoniae* with elevated cephalosporin MICs were used. These isolates were molecularly characterized and known to produce various Ambler Class A, Extended Spectrum beta-lactamase (ESBL). The assay was conducted in Cation Adjusted Mueller Hinton Broth (CAMHB, BD # 212322, BD Diagnostic Systems, Sparks, MD). Bacteria strains were grown for 3-5 hours in CAMHB broth. Ceftibuten (#SML0037-50MG, Sigma, St Louis, MO), cefixime (#1097658, USP, Rockville, MD), cefdinir (#C7118-1G, Sigma, St Louis, MO), cephalexin (#15085, MP Biomedicals, Solon, OH), cefpodoxime (#1098027, Rockville, MD) were added either alone to a microtiter plate in 2-fold serial dilutions in CAMHB or in combination with beta-lactamase inhibitors (BLI) at fixed concentration. Compound 2 was added to each test cephalosporin at fixed concentrations of 1 $\mu\text{g/mL}$, 2 $\mu\text{g/mL}$, and 4 $\mu\text{g/mL}$ and clavulanic acid was added at 4 $\mu\text{g/mL}$. Once the test articles are added, the plates can be inoculated according to CLSI broth microdilution method. After inoculation, the plates are incubated for 16-20 hours at 37 °C then the Minimal Inhibitory Concentration (MIC) of the test compound is determined visually. Comparison of the MIC data from antibiotic alone with the BLI-cephalosporin combination serves as a measure of the potentiation.

[0334] Table 3 lists the MIC data for ceftibuten titrated against a set of forty ESBL-producing strains of Enterobacteriaceae, alone and in combination with Compound 2 or clavulanic acid at the fixed concentrations listed. Table 4 provides the beta-lactamase content for this set of strains.

Table 3. Minimum Inhibitory Concentrations of ceftibuten (CTB) in combination with Compound 2 (fixed at 1, 2, or 4 $\mu\text{g/mL}$) or clavulanic acid (4 $\mu\text{g/mL}$) in a panel of ESBL-producing Enterobacteriaceae.

Species	Strain	Compound 2 (Fixed Concentration)			Clavulanic acid	CTB alone
		1 $\mu\text{g/mL}$	2 $\mu\text{g/mL}$	4 $\mu\text{g/mL}$	4 $\mu\text{g/mL}$	
<i>E. coli</i>	1924	0.125	0.25	0.125	0.5	0.5
<i>E. coli</i>	2150	0.125	0.125	0.06	0.5	0.5

Species	Strain	Compound 2 (Fixed Concentration)			Clavulanic acid	CTB alone
		1 µg/mL	2 µg/mL	4 µg/mL	4 µg/mL	
<i>E. coli</i>	2806	0.06	0.125	0.125	0.25	2
<i>E. coli</i>	3174	0.125	0.25	0.25	0.5	0.5
<i>E. coli</i>	3327	0.25	0.25	0.25	1	8
<i>E. coli</i>	BAS 1	1	1	0.5	>32	>32
<i>E. coli</i>	ESBL 2	0.125	0.125	0.125	4	8
<i>E. coli</i>	ESBL4	0.25	0.125	0.25	0.5	>32
<i>E. coli</i>	ESBL5	0.25	0.125	0.125	0.5	>32
<i>E. coli</i>	SI-AIRT-4	0.06	0.06	0.06	0.5	0.5
<i>E. coli</i>	Si-FDL-GES	0.125	0.125	0.125	0.25	0.5
<i>E. coli</i>	SI-LP377	0.125	0.125	0.125	0.125	4
<i>E. coli</i>	SI-M004	0.25	0.125	0.125	0.5	32
<i>E. coli</i>	SI-NO36	0.06	0.125	0.06	0.25	0.5
<i>E. coli</i>	SI-PBLII	0.125	0.125	0.125	0.5	0.5
<i>E. coli</i>	SI-V502	0.25	0.25	0.125	0.5	0.5
<i>E. coli</i>	25922	0.125	0.06	0.06	0.5	0.25
<i>K. oxytoca</i>	169219	0.03	0.0149	0.0149	0.0149	4
<i>K. oxytoca</i>	176877	0.5	0.125	0.25	0.125	4
<i>K. oxytoca</i>	496 #2	0.125	0.125	0.125	0.125	8
<i>K. pneumoniae</i>	3151	1	0.25	0.25	0.06	>32
<i>K. pneumoniae</i>	115468	0.06	0.06	0.06	0.25	8
<i>K. pneumoniae</i>	153239	0.06	0.125	0.06	0.06	0.25
<i>K. pneumoniae</i>	170375	0.06	0.06	0.06	0.06	32
<i>K. pneumoniae</i>	304487	0.25	0.25	0.25	0.125	8
<i>K. pneumoniae</i>	319478	0.125	0.25	0.125	0.06	>32
<i>K. pneumoniae</i>	329633	0.06	0.06	0.06	0.03	16
<i>K. pneumoniae</i>	11/23 LF#2	0.25	0.25	0.125	0.25	2
<i>K. pneumoniae</i>	ESBL 10	0.25	0.25	0.25	0.125	16
<i>K. pneumoniae</i>	ESBL 7 #1	0.25	0.25	0.125	0.125	32
<i>K. pneumoniae</i>	ESBL 8 #1	0.25	0.125	0.06	0.03	16
<i>K. pneumoniae</i>	ESBL 8 #2	0.125	0.125	0.06	0.06	16
<i>K. pneumoniae</i>	ESBL7 #2	4	2	1	>32	>32
<i>K. pneumoniae</i>	KI (KC1)	0.125	0.06	0.125	0.06	1
<i>K. pneumoniae</i>	KP 3	0.25	0.25	0.125	0.125	2
<i>K. pneumoniae</i>	KP 4	0.25	0.125	0.125	2	32
<i>K. pneumoniae</i>	KP 8	0.06	0.03	0.03	0.06	0.25
<i>K. pneumoniae</i>	KPN 508	0.125	0.125	0.06	0.06	32
<i>K. pneumoniae</i>	SI-KP NO30	0.06	0.06	0.06	0.06	0.25
<i>K. pneumoniae</i>	BAA 1705	0.5	0.5	0.25	8	32

Table 4. Beta-Lactamases expressed in test set of Enterobacteriaceae strains.

Species	Strain	Enzyme
<i>E. coli</i>	1924	TEM-12
<i>E. coli</i>	2150	SHV-3, TEM-1
<i>E. coli</i>	2806	SHV-12, TEM-1
<i>E. coli</i>	3174	TEM-1, CTXM-15
<i>E. coli</i>	3327	TEM-1, SHV-12
<i>E. coli</i>	BAS 1	none detected
<i>E. coli</i>	ESBL 2	TEM-1
<i>E. coli</i>	ESBL4	CTX-M15, TEM-1
<i>E. coli</i>	ESBL5	CTX-M15, TEM-1

Species	Strain	Enzyme
<i>E. coli</i>	SI-AIRT-4	TEM-1
<i>E. coli</i>	SI-FDL-GES	GES-5
<i>E. coli</i>	SI-LP377	CTX-M2
<i>E. coli</i>	SI-M004	SHV-2, GES-12
<i>E. coli</i>	SI-NO36	SHV-11, TEM-1
<i>E. coli</i>	SI-PBLII	CTX-M2, TEM-1
<i>E. coli</i>	SI-V502	TEM-29, CTX-M15
<i>E. coli</i>	25922	None detected
<i>K. oxytoca</i>	169219	TEM-1, CTXM-15
<i>K. oxytoca</i>	176877	SHV-105
<i>K. oxytoca</i>	496 #2	SHV-105, TEM-1
<i>K. pneumoniae</i>	3151	SHV-12, TEM-1
<i>K. pneumoniae</i>	115468	TEM-1
<i>K. pneumoniae</i>	153239	SHV-11, TEM-1
<i>K. pneumoniae</i>	170375	SHV-12, CTX-M15 and TEM-1
<i>K. pneumoniae</i>	304487	SHV-12, TEM-1
<i>K. pneumoniae</i>	319478	SHV-12, CTX-M3 and TEM-1
<i>K. pneumoniae</i>	329633	SHV-5, TEM-1
<i>K. pneumoniae</i>	11/23 LF#2	SHV-11, GES-5
<i>K. pneumoniae</i>	ESBL 10	SHV-12, TEM-1
<i>K. pneumoniae</i>	ESBL 7 #1	SHV-12, TEM-1
<i>K. pneumoniae</i>	ESBL 8 #1	TEM-1, SHV-12
<i>K. pneumoniae</i>	ESBL 8 #2	SHV-12, TEM-1
<i>K. pneumoniae</i>	ESBL7 #2	SHV-12, CTX-M15, TEM-1
<i>K. pneumoniae</i>	KI (KC1)	TEM-10
<i>K. pneumoniae</i>	KP 3	SHV-1, TEM-1
<i>K. pneumoniae</i>	KP 4	SHV-2a, SHV-5, TEM-1
<i>K. pneumoniae</i>	KP 8	TEM-10, TEM-1, CTX-M15
<i>K. pneumoniae</i>	KPN 508	SHV-7, TEM-1
<i>K. pneumoniae</i>	SI-KP NO30	TEM-26b, SHV-60, SHV-26

[0335] Table 5 lists MIC data for cefdinir in combination with Compound 2 or clavulanic acid in this panel of ESBL Enterobacteriaceae.

Table 5. Minimum Inhibitory Concentrations of cefdinir (CDR) in combination with Compound 2 (fixed at 1, 2, or 4 µg/mL) or clavulanic acid (4 µg/mL), in a panel of ESBL Enterobacteriaceae.

Species	Strain	Compound 2 (Fixed Concentration)			Clavulanic acid 4 µg/mL	CDR alone
		1 µg/mL	2 µg/mL	4 µg/mL		
<i>E. coli</i>	1924	0.5	0.25	0.25	0.5	2
<i>E. coli</i>	2150	1	0.5	0.5	0.5	>32
<i>E. coli</i>	2806	0.125	0.125	0.25	0.25	4
<i>E. coli</i>	3174	0.25	0.25	0.125	0.25	1
<i>E. coli</i>	3327	0.5	0.5	0.5	0.5	16
<i>E. coli</i>	BAS 1	1	1	2	16	>32
<i>E. coli</i>	ESBL 2	0.5	0.5	0.25	>32	>32
<i>E. coli</i>	ESBL4	2	1	0.5	1	>32
<i>E. coli</i>	ESBL5	8	2	2	1	>32
<i>E. coli</i>	SI-AIRT-4	0.25	0.25	0.25	0.25	0.25

Species	Strain	Compound 2 (Fixed Concentration)			Clavulanic acid	CDR alone
		1 µg/mL	2 µg/mL	4 µg/mL	4 µg/mL	
<i>E. coli</i>	SI-FDL-GES	0.5	0.5	0.5	1	>32
<i>E. coli</i>	SI-LP377	4	1	1	1	>32
<i>E. coli</i>	SI-M004	0.5	0.5	0.5	0.5	32
<i>E. coli</i>	SI-NO36	0.125	0.125	0.25	0.25	4
<i>E. coli</i>	SI-PBLII	0.25	0.25	0.25	0.5	2
<i>E. coli</i>	SI-V502	1	0.5	0.5	1	2
<i>E. coli</i>	25922	0.125	0.125	0.125	0.25	0.5
<i>K. oxytoca</i>	169219	1	0.5	0.5	0.25	>32
<i>K. oxytoca</i>	176877	2	2	1	2	8
<i>K. oxytoca</i>	496 #2	0.25	0.25	0.06	0.06	16
<i>K. pneumoniae</i>	3151	2	1	0.5	0.125	>32
<i>K. pneumoniae</i>	115468	0.5	0.25	0.125	0.06	8
<i>K. pneumoniae</i>	153239	0.5	0.25	0.25	0.25	2
<i>K. pneumoniae</i>	170375	8	4	2	1	>32
<i>K. pneumoniae</i>	304487	1	1	2	0.5	8
<i>K. pneumoniae</i>	319478	4	2	2	0.5	>32
<i>K. pneumoniae</i>	329633	0.25	0.25	1	0.25	16
<i>K. pneumoniae</i>	11/23 LF#2	0.25	0.5	0.25	0.5	2
<i>K. pneumoniae</i>	ESBL 10	1	1	0.5	0.5	16
<i>K. pneumoniae</i>	ESBL 7 #1	0.5	1	0.5	0.5	32
<i>K. pneumoniae</i>	ESBL 8 #1	0.25	0.125	0.25	0.06	16
<i>K. pneumoniae</i>	ESBL 8 #2	0.5	0.25	0.25	0.125	16
<i>K. pneumoniae</i>	ESBL7 #2	>32	32	16	>32	>32
<i>K. pneumoniae</i>	KI (KC1)	0.125	0.125	0.125	0.125	1
<i>K. pneumoniae</i>	KP 3	0.5	0.25	0.5	1	4
<i>K. pneumoniae</i>	KP 4	0.5	0.5	0.125	0.125	8
<i>K. pneumoniae</i>	KP 8	0.25	0.125	0.125	0.125	0.5
<i>K. pneumoniae</i>	KPN 508	0.5	0.25	0.5	0.125	32
<i>K. pneumoniae</i>	SI-KP NO30	0.125	0.25	0.125	0.125	0.5
<i>K. pneumoniae</i>	BAA 1705	4	2	1	>32	>32

[0336] Table 6 lists MIC data for cefalexin in combination with Compound 2 or clavulanic acid in this panel of ESBL Enterobacteriaceae.

Table 6. Minimum Inhibitory Concentrations of cefalexin (LEX) in combination with Compound 2 (fixed at 1, 2, or 4 µg/mL) or clavulanic acid (4 µg/mL), in a panel of ESBL Enterobacteriaceae.

Species	Strain	Compound 2 (Fixed Concentration)			Clavulanic acid	LEX alone
		1 µg/mL	2 µg/mL	4 µg/mL	4 µg/mL	
<i>E. coli</i>	1924	0.5	0.25	0.25	0.5	2
<i>E. coli</i>	2150	1	0.5	0.5	0.5	>32
<i>E. coli</i>	2806	0.125	0.125	0.25	0.25	4
<i>E. coli</i>	3174	0.25	0.25	0.125	0.25	1
<i>E. coli</i>	3327	0.5	0.5	0.5	0.5	16
<i>E. coli</i>	BAS 1	1	1	2	16	>32
<i>E. coli</i>	ESBL 2	0.5	0.5	0.25	>32	>32

Species	Strain	Compound 2 (Fixed Concentration)			Clavulanic acid	LEX alone
		1 µg/mL	2 µg/mL	4 µg/mL	4 µg/mL	
<i>E. coli</i>	ESBL4	2	1	0.5	1	>32
<i>E. coli</i>	ESBL5	8	2	2	1	>32
<i>E. coli</i>	SI-AIRT-4	0.25	0.25	0.25	0.25	0.25
<i>E. coli</i>	Si-FDL-GES	0.5	0.5	0.5	1	>32
<i>E. coli</i>	SI-LP377	4	1	1	1	>32
<i>E. coli</i>	SI-M004	0.5	0.5	0.5	0.5	32
<i>E. coli</i>	SI-NO36	0.125	0.125	0.25	0.25	4
<i>E. coli</i>	SI-PBLII	0.25	0.25	0.25	0.5	2
<i>E. coli</i>	SI-V502	1	0.5	0.5	1	2
<i>E. coli</i>	25922	0.125	0.125	0.125	0.25	0.5
<i>K. oxytoca</i>	169219	1	0.5	0.5	0.25	>32
<i>K. oxytoca</i>	176877	2	2	1	2	8
<i>K. oxytoca</i>	496 #2	0.25	0.25	0.06	0.06	16
<i>K. pneumoniae</i>	3151	2	1	0.5	0.125	>32
<i>K. pneumoniae</i>	115468	0.5	0.25	0.125	0.06	8
<i>K. pneumoniae</i>	153239	0.5	0.25	0.25	0.25	2
<i>K. pneumoniae</i>	170375	8	4	2	1	>32
<i>K. pneumoniae</i>	304487	1	1	2	0.5	8
<i>K. pneumoniae</i>	319478	4	2	2	0.5	>32
<i>K. pneumoniae</i>	329633	0.25	0.25	1	0.25	16
<i>K. pneumoniae</i>	11/23 LF#2	0.25	0.5	0.25	0.5	2
<i>K. pneumoniae</i>	ESBL 10	1	1	0.5	0.5	16
<i>K. pneumoniae</i>	ESBL 7 #1	0.5	1	0.5	0.5	32
<i>K. pneumoniae</i>	ESBL 8 #1	0.25	0.125	0.25	0.06	16
<i>K. pneumoniae</i>	ESBL 8 #2	0.5	0.25	0.25	0.125	16
<i>K. pneumoniae</i>	ESBL7 #2	>32	32	16	>32	>32
<i>K. pneumoniae</i>	KI (KC1)	0.125	0.125	0.125	0.125	1
<i>K. pneumoniae</i>	KP 3	0.5	0.25	0.5	1	4
<i>K. pneumoniae</i>	KP 4	0.5	0.5	0.125	0.125	8
<i>K. pneumoniae</i>	KP 8	0.25	0.125	0.125	0.125	0.5
<i>K. pneumoniae</i>	KPN 508	0.5	0.25	0.5	0.125	32
<i>K. pneumoniae</i>	SI-KP NO30	0.125	0.25	0.125	0.125	0.5
<i>K. pneumoniae</i>	BAA 1705	4	2	1	>32	>32

[0337] Table 7 lists MIC data for cefixime in combination with Compound 2 or clavulanic acid in this panel of ESBL Enterobacteriaceae.

Table 7. Minimum Inhibitory Concentrations of cefixime (CFM) in combination with Compound 2 (fixed at 1, 2, or 4 µg/mL) or clavulanic acid (4 µg/mL), in a panel of ESBL Enterobacteriaceae.

Species	Strain	Compound 2 (Fixed Concentration)			Clavulanic acid	CFM alone
		1 µg/mL	2 µg/mL	4 µg/mL	4 µg/mL	
<i>E. coli</i>	1924	0.5	0.5	0.25	1	2
<i>E. coli</i>	2150	0.5	0.5	0.25	1	>32
<i>E. coli</i>	2806	0.25	0.125	0.125	1	32
<i>E. coli</i>	3174	0.06	0.125	0.06	0.5	0.5
<i>E. coli</i>	3327	1	0.5	0.5	2	>32
<i>E. coli</i>	BAS 1	8	2	1	>32	>32

Species	Strain	Compound 2 (Fixed Concentration)			Clavulanic acid	CFM alone
		1 µg/mL	2 µg/mL	4 µg/mL	4 µg/mL	
<i>E. coli</i>	ESBL 2	0.25	0.25	0.125	32	>32
<i>E. coli</i>	ESBL4	0.5	0.25	0.25	1	>32
<i>E. coli</i>	ESBL5	1	1	0.5	1	>32
<i>E. coli</i>	SI-AIRT-4	0.125	0.125	0.06	0.25	0.5
<i>E. coli</i>	Si-FDL-GES	0.25	0.125	0.125	0.5	2
<i>E. coli</i>	SI-LP377	0.5	0.25	0.125	0.5	>32
<i>E. coli</i>	SI-M004	4	4	1	1	>32
<i>E. coli</i>	SI-NO36	0.5	0.25	0.5	0.5	1
<i>E. coli</i>	SI-PBLII	0.25	0.25	0.25	1	1
<i>E. coli</i>	SI-V502	0.25	0.5	0.25	1	2
<i>E. coli</i>	25922	0.125	0.06	0.06	0.5	1
<i>K. oxytoca</i>	169219	0.25	0.125	0.125	0.125	>32
<i>K. oxytoca</i>	176877	1	0.5	0.5	0.25	>32
<i>K. oxytoca</i>	496 #2	0.25	0.125	0.06	0.03	32
<i>K. pneumoniae</i>	3151	16	8	2	0.125	>32
<i>K. pneumoniae</i>	115468	2	0.5	0.5	0.125	>32
<i>K. pneumoniae</i>	153239	0.125	0.5	0.25	0.25	0.5
<i>K. pneumoniae</i>	170375	1	1	1	0.5	>32
<i>K. pneumoniae</i>	304487	1	1	0.5	0.25	>32
<i>K. pneumoniae</i>	319478	2	2	1	0.5	>32
<i>K. pneumoniae</i>	329633	1	0.5	0.5	0.03	>32
<i>K. pneumoniae</i>	11/23 LF#2	4	2	1	2	>32
<i>K. pneumoniae</i>	ESBL 10	2	1	0.5	0.125	>32
<i>K. pneumoniae</i>	ESBL 7 #1	4	4	1	0.25	>32
<i>K. pneumoniae</i>	ESBL 8 #1	2	0.5	0.25	0.06	>32
<i>K. pneumoniae</i>	ESBL 8 #2	2	1	0.5	0.03	>32
<i>K. pneumoniae</i>	ESBL7 #2	8	8	1	>32	>32
<i>K. pneumoniae</i>	KI (KC1)	1	1	0.25	0.125	32
<i>K. pneumoniae</i>	KP 3	4	4	2	0.5	>32
<i>K. pneumoniae</i>	KP 4	2	0.5	0.25	0.125	>32
<i>K. pneumoniae</i>	KP 8	0.25	0.125	0.125	0.06	>32
<i>K. pneumoniae</i>	KPN 508	4	1	0.5	0.25	>32
<i>K. pneumoniae</i>	SI-KP NO30	0.5	0.25	0.25	0.06	>32
<i>K. pneumoniae</i>	BAA 1705	0.5	0.5	0.25	>32	>32

[0338] Summary MIC data (MIC₅₀, the concentration inhibiting growth of 50% of the strains in the panel, and MIC₉₀, the concentration inhibiting growth of 90% of the strains) are shown in Table 8. This data indicates that Compound 2, in combination with Cefibuten, provides superior antibacterial activities across the range of ESBL-producing Enterobacteriaceae.

Table 8. Summary of Minimum Inhibitory Concentrations of cephalosporins in combinations with Compound 2 (fixed at 1, 2, or 4 µg/mL) or clavulanic acid (4 µg/mL) for 50% (MIC₅₀, µg/mL) and 90% (MIC₉₀, µg/mL) of tested strains of Enterobacteriaceae.

Antibiotic		Inhibitor (fixed concentration)				
		Compd 2 (1 µg/mL)	Compd 2 (2 µg/mL)	Compd 2 (4 µg/mL)	Clavulanic Acid (4 µg/mL)	No BLI
Ceftibuten	MIC ₅₀	0.125	0.125	0.125	0.25	8
	MIC ₉₀	0.5	0.25	0.25	2	>32
Cefixime	MIC ₅₀	1	0.5	0.25	0.5	>32
	MIC ₉₀	4	4	1	2	>32
Cefdinir	MIC ₅₀	0.5	0.5	0.5	0.5	16
	MIC ₉₀	4	2	2	2	>32
Cephalexin	MIC ₅₀	8	8	8	16	>32
	MIC ₉₀	32	32	16	32	>32
Cefpodoxime	MIC ₅₀	1	1	1	0.5	>32
	MIC ₉₀	8	2	2	4	>32

Example 3. In vivo efficacy in a murine septicemia infection model.

[0339] In vivo efficacy studies with Compound 1 were conducted (Dr. W. Weiss, U. North Texas Health Sciences Center) in the lethal murine model of septicemia, using a *K. pneumoniae* strain (UNT-023) expressing KPC-2 (in analogy to the procedure described in Endimiani, A., et al., *Antimicrob. Agents Chemother.* **2011**, 55(1), pp 82-85). The inoculum for the strain was titrated to produce 100% mortality in five days after intraperitoneal inoculation. Mice were administered compounds 1 h post-inoculation. Dosing groups (five animals per group) compared ceftibuten alone administered by oral gavage at 8-128 mg/kg versus ceftibuten plus Compound 1 given by oral gavage in a 1:1 ratio (2-32 mg/kg). Table 9 shows that Compound 1 demonstrated striking rescue of ceftibuten in this lethal septicemia model, with the 50% effective dose (EC₅₀) values shown. The results confirm that Compound 1, orally-delivered in combination with ceftibuten, is efficacious against Class A beta-lactamase producing Enterobacteriaceae.

Table 9. Proof of Concept – In vivo rescue of Ceftibuten by Compound 1 in a lethal murine septicemia model of infection – *K. pneumoniae* (UNT-023, KPC-2 producing).

Dosing Group (Oral)	MIC (µg/mL) vs. <i>K. pneumoniae</i> UNT-023 (KPC-2)	ED ₅₀ (mg/kg PO)	95% Confidence Interval
CTB alone	>128	>128	--
CTB + Compound 1	0.5	12.9	9.8-17.2

Example 4: In vivo efficacy in a murine urinary tract infection model.

[0340] In vivo efficacy studies with Compound 2 were conducted (Dr. W. Weiss, U. North Texas Health Sciences Center) in a murine model of pyelonephritis, using an *E. coli* strain (UNT-204-1) expressing TEM-1 and CTX-M-15 beta-lactamases (performed according to the procedure described in Weiss, W.J., et al., *Antimicrob. Agents Chemother.* **2018**, 62(1), e01439-17). To establish the ascending UTI model, female C3H/HeJ mice were placed on 5% glucose water for 6 days and then transurethrally infected with approximately $9 \log_{10}$ CFU of each bacterial isolate to establish kidney infection. Treatment was initiated on day 5 following inoculation, and dosing groups (five animals per group) compared ceftibuten alone (3-300 mg/kg), ceftibuten plus Compound 2 in a 1:1 ratio (3-300 mg/kg), and amoxicillin-clavulanate in a 2:1 ratio (10-300 mg/kg amoxicillin, 5-150 mg/kg clavulanate), administered by subcutaneous injection twice daily for 3 days. Animals were euthanized 2 h following the last dose of test article on post-inoculation day 7, and bacterial counts was determined in kidneys, bladder, and urine, and compared across dosing groups and untreated controls. Table 10 shows that Compound 2 demonstrated striking rescue of ceftibuten in this model. In the kidneys, ceftibuten (100 mg/kg) plus Compound 2 (100 mg/kg) reduced the bacterial load by an average of $3.17 \log_{10}$ CFU, compared with a reduction of $2.11 \log_{10}$ CFU for ceftibuten alone (100 mg/kg), or $1.55 \log_{10}$ CFU reduction for amoxicillin-clavulanate (100 mg/kg + 50 mg/kg, respectively). Similar improvements in reduction of bacterial counts in bladder and urine were observed for the combination of Compound 2 with ceftibuten. The results show that Compound 2, delivered in combination with ceftibuten, is efficacious against Class A beta-lactamase producing Enterobacteriaceae in a model of urinary tract infections.

[0341] In vivo efficacy study of ceftibuten with and without Compound 2 was conducted (Dr. W. Weiss, U. North Texas Health Sciences Center) in the murine ascending UTI model with three strains of *E. coli* expressing TEM-1 + CTX-M-15, CTX-M-15, and KPC-2 + SHV-12. In the first part of the study, the MICs of ceftibuten and ceftibuten with Compound 2 at a fixed concentration of 4 $\mu\text{g/mL}$ were determined against the test strains. Ceftibuten MICs were reduced from 8-64 $\mu\text{g/mL}$ to <0.06-0.12 $\mu\text{g/mL}$ with Compound 2.

[0342] The animals were prepared according to the procedure described in Weiss, W.J., et al., *Antimicrob. Agents Chemother.* **2018**, 62(1), e01439-17. Treatments of ceftibuten alone, ceftibuten with Compound 2 at various dose ratios, and amoxicillin-clavulanate (2:1) (used as a comparator) were initiated four days post-infection and administered subcutaneously, every 12 hours for 3 days. Bacterial burdens in kidneys, bladders, and urine 18 hours after the final dose were quantitated by serial dilution plating and compared to those of untreated controls.

[0343] At Day 7 post-infection, mean bacterial titers for the three bacterial strains were 6.5 to 7.1 $\log_{10}$ CFU/g in kidney, 5.7 to 6.8 $\log_{10}$ CFU/g in bladder, and 5.9 to 7.2 $\log_{10}$ CFU/mL in urine for the untreated controls. Administration of ceftibuten alone from 1 to 300 mg/kg resulted in a dose response. There were minimal CFU reductions in the kidneys observed and up to 2 $\log_{10}$ CFU lower

titers in the bladder and urine at ceftibuten doses of 100 and 300 mg/kg. The addition of Compound 2 to ceftibuten treatment resulted in increased efficacy, with bacterial titers that were up to 2 log₁₀ CFU/g lower in kidneys (FIG. 5A, FIG. 5B, and FIG. 5C), 3.2 log₁₀ CFU/g lower in bladders (FIG. 6A, FIG. 6B, and FIG. 6C), and up to 4 log₁₀ CFU/mL lower in urine (FIG. 7A, FIG. 7B, and FIG. 7C) than ceftibuten alone at doses of 3, 10, 30, 100, and 300 mg/kg.

[0344] In this UTI study with *E. coli* isolates expressing ESBLs or a combination of ESBLs and KPC carbapenemase, co-administration of Compound 2 (1:1) further reduced the bacterial titers in kidneys, bladders, and urine when compared to ceftibuten alone. The results demonstrate that Compound 2 rescues ceftibuten activity in a UTI model against uropathogenic strains of *E. coli* expressing TEM-1 + CTX-M-15, CTX-M-15, or KPC-2 + SHV-12.

Table 10. In vivo rescue of Ceftibuten (CTB) by Compound 2 in a murine model of pyelonephritis infection – *E. coli* (UNT-204-1, TEM-1 plus CTX-M-15 producing).

Test Article	Dose Group (mg/kg, SC, BID, 3 days)	Log ₁₀ Change in CFU/Sample vs. untreated controls, Day 7		
		Kidneys	Bladder	Urine
CTB alone	300	-2.14	-1.83	-3.37
	100	-2.11	-2.26	-2.54
	30	-1.57	-2.08	-2.28
	10	-1.55	-1.51	-0.59
	3	-0.44	-0.03	-1.24
CTB + Compound 2	300:300	-2.99	-1.93	-3.99
	100:100	-3.17	-2.75	-3.53
	30:30	-2.93	-2.65	-3.67
	10:10	-2.59	-2.12	-3.71
	3:3	-2.37	-1.77	-3.42
Amoxicillin-Clavulanate	300:150	-3.01	-1.62	-2.41
	100:50	-1.55	-1.17	-1.34
	30:15	-1.48	-0.96	-0.28
	10:5	-0.77	-0.87	-0.46

Example 5. Time-Kill Kinetic Analyses for Compound 2.

[0345] Compound 2 was assessed for its ability to rescue ceftibuten to cidal activity (≤ 3 log₁₀ killing) using time-kill experiments. Activity was determined against four beta-lactamase expressing Enterobacteriaceae (1 KPC, 1 p99, 1 CTX-M-15, and 1 OXA-48). Ceftibuten was diluted in macrodilution format either alone or with Compound 2 fixed at 4 µg/mL over a concentration range dictated by its MIC in that particular strain. Macrodilution tubes were incubated with shaking and bacterial amounts were quantified at 0, 2, 4, 6, 8, and 24 hours by limiting dilutions. Amoxicillin/clavulanic acid (2:1) was used as a comparator oral antibacterial. The data for all 4 strains are depicted in FIG. 1A, FIG. 1B, FIG. 1C, and FIG. 1D, and show that Compound 2 was able to rescue ceftibuten to a cidal endpoint through 24 hours. Similar responses were observed in all

strains tested, even in those resistant to carbapenems. The results demonstrate that Compound 2 returns the beta-lactam ceftibuten to full cidal activity in time-kill experiments.

Example 6. Synergy and Antagonism of Compound 2 plus Ceftibuten.

[0346] The potential for synergy or antagonism between Compound 2 and a range of antibacterial agents was evaluated by measuring fractional inhibitory concentration (FIC) values in vitro using a broth microdilution “checkerboard” panel, in which the combination agents were tested both alone and together at varying concentration ratios. In this study, ceftibuten was titrated in the presence and absence of Compound 2 (fixed at 4 µg/mL) and compared to linezolid, rifampin, levofloxacin, nitrofurantoin and trimethoprim-sulfamethoxazole against six Enterobacteriaceae strains (5 expressing beta-lactamases and 1 wild-type). In addition, ceftibuten/Compound 2 was tested in combination with metronidazole against three representative anaerobic strains of bacteria, and in combination with fluconazole in three representative strains of yeast. Results are shown in Tables 11-13, wherein FICI ≤ 0.5 = Synergy; $>0.5 - 4$ = additive/indifferent; >4 = antagonism. Importantly, antagonism was not observed between ceftibuten/Compound 2 and other agents tested.

Table 11. Fractional Inhibitory Concentration Index (FICI) for Ceftibuten/Compound 2 Combined with Antibacterial Agents Linezolid (LZD), Levofloxacin (LVX), Rifampin (RIF), Nitrofurantoin (NIT), or Trimethoprim-Sulfamethoxazole (SXT).

Species	Strain ID	Enzyme Content	FICI				
			LZD	RIF	LVX	NIT	SXT
<i>E. coli</i>	ATCC 25922	None	0.48	1.34	1.33	1.03	1.22
<i>K. pneumoniae</i>	ATCC BAA 1705	KPC-2	1.00	1.15	0.71	1.00	1.00
<i>K. pneumoniae</i>	ATCC 700603	SHV-18	2.00	0.74	1.12	1.03	1.03
<i>E. coli</i>	VER	OXA-48	0.50	1.12	1.21	1.08	1.47
<i>E. coli</i>	ESBL 4	CTX-M-15	0.50	0.86	1.10	0.72	1.33
<i>E. cloacae</i>	SI-ECL01	p99	0.42	0.58	0.98	0.96	1.05

Table 12. Fractional Inhibitory Concentration Index (FICI) for Ceftibuten/Compound 2 Combined with Metronidazole in Anaerobes.

Species	Strain ID	Enzyme Content	FICI of Ceftibuten/Compound 2 with metronidazole
<i>B. fragilis</i>	ATCC 25285	None	0.55
<i>C. difficile</i>	ATCC 700057	None	1.00
<i>B. thetaiotaomicron</i>	ATCC 29741	None	1.00

Table 13. Fractional Inhibitory Concentration Index (FICI) for Ceftibuten/Compound 2 Combined with Fluconazole in Representative Strains of Yeast.

Species	Strain ID	FICI of Ceftibuten/Compound 2 with Fluconazole
<i>Candida albicans</i>	ATCC 90028	0.88
<i>Candida tropicalis</i>	ATCC 750	0.85
<i>Candida krusei</i>	ATCC 6258	0.75

Example 7. Impact of concentration on the ability of Compound 2 to rescue ceftibuten in Enterobacteriaceae expressing Class A, C, and D beta-lactamase enzymes.

[0347] Minimum inhibitory concentrations (MICs) were determined by broth microdilution method following CLSI guidelines. Tested strains were ceftibuten-resistant Enterobacteriaceae expressing ESBL (n=20), KPC (n=20), OXA-48 (n=20) and/or Class C (n=20) beta-lactamases. For each isolate, one microtiter plate was prepared and formatted as a checkerboard with the BLI (Compound 2 or clavulanic acid) titrated down the plate and ceftibuten titrated across the plate. Each agent was also tested alone. MICs were determined as the well with the lowest concentration of ceftibuten without visible growth at each concentration of BLI.

Table 14. Ceftibuten MIC ($\mu\text{g/mL}$) summary for each enzyme-expressing subset of isolates with Compound 2 or clavulanic Acid (CLA) at 4 $\mu\text{g/mL}$.

	BLI	ESBL	KPC	OXA-48	Class C	All
CTB MIC₅₀	Compound 2	0.25	0.25	0.125	0.25	0.25
	CLA	0.5	8	8	2	4
CTB MIC₉₀	Compound 2	0.5	1	0.5	1	1
	CLA	2	32	>32	>32	32

Example 8. Pharmacokinetic Studies of Compound 1 in Rats.

[0348] The single dose plasma pharmacokinetic parameters were evaluated for Compound 2 following oral gavage dosing of Compound 1 in fasted and non-fasted Sprague-Dawley rats, as well as following intravenous bolus dosing of Compound 2. The results are summarized in Table 15.

Table 15. Mean plasma pharmacokinetic parameters for Compound 2 following intravenous administration of Compound 2, or oral administration of Compound 1, in Male Sprague-Dawley Rats.

Group	Number of rats (n)	C _{max} (ng/mL)	AUC _∞ (ng*h/mL)	F (%)
Compound 2 5 mg/kg IV	10	7,095	3,711	NA
Compound 1 5 mg/kg PO, fasted	10	1,441	1,735	70
Compound 1 5 mg/kg PO, non-fasted	10	574	2,263	91

Abbreviations: C_{max} = maximum plasma concentration; AUC_∞ = area under the plasma concentration time curve from time 0 to infinity; F = oral bioavailability corrected for the molecular weight difference between Compound 1 and Compound 2; NA = not applicable.

Example 9. Pharmacokinetic Studies of Compound 1 in Dogs.

[0349] The single dose plasma pharmacokinetic parameters were evaluated for Compound 2 following oral gavage dosing of Compound 1 in fasted and non-fasted male Beagle dogs, as well as following intravenous bolus dosing of Compound 2. The results are summarized in Table 16.

Table 16. Mean plasma pharmacokinetic parameters for Compound 2 following intravenous administration of Compound 2, or oral administration of Compound 1, in male Beagle dogs.

Group	Number of dogs (n)	C _{max} (ng/mL)	AUC _∞ (ng*h/mL)	F (%)
Compound 2 5 mg/kg IV	10	22,404	25,467	NA
Compound 1 5 mg/kg PO, fasted	5	3,540	9,875	58
Compound 1 5 mg/kg PO, non-fasted	15	3,960	10,939	64

Abbreviations: C_{max} = maximum plasma concentration; AUC_∞ = area under the plasma concentration time curve from time 0 to infinity; F = oral bioavailability corrected for the molecular weight difference between Compound 1 and Compound 2; NA = not applicable.

Example 10. Pharmacokinetic Studies of Compound 1 in Monkeys.

[0350] The single dose plasma pharmacokinetic parameters were evaluated for Compound 2 following oral gavage dosing of Compound 1 in fasted and non-fasted male Cynomolgus monkeys, as well as following intravenous bolus dosing of Compound 2. The results are summarized in Table 17.

Table 17. Mean plasma pharmacokinetic parameters for Compound 2 following intravenous administration of Compound 2, or oral administration of Compound 1, in male Cynomolgus monkeys.

Group	Number of monkeys (n)	C _{max} (ng/mL)	AUC _∞ (ng*h/mL)	F (%)
Compound 2 5 mg/kg IV	10	38,575	38,399	NA
Compound 1 5 mg/kg PO, fasted	10	4,794	10,224	42
Compound 1 5 mg/kg PO, non-fasted	15	6,186	15,392	63

Abbreviations: C_{max} = maximum plasma concentration; AUC_∞ = area under the plasma concentration time curve from time 0 to infinity; F = oral bioavailability corrected for the molecular weight difference between Compound 1 and Compound 2; NA = not applicable.

Example 11. Single Ascending Dose Pharmacokinetics in Rats.

[0351] The single dose pharmacokinetics and dose proportionality of Compound 2 were evaluated in non-fasted male Sprague-Dawley rats following oral gavage of Compound 1, as well as intravenous bolus administration of Compound 2. The dosing formulation for Compound 1 was prepared in Solutol HS-15:Water (20:80 v/v) for oral dosing (homogeneous and chemically-stable solution up to a concentration of 60 mg/mL; physically- and chemically-stable emulsion at concentrations ranging from 61-200 mg/mL). This study included dose concentrations ranging from 2-120 mg/mL. The data are shown in Table 18.

Table 18. Mean plasma pharmacokinetic parameters for Compound 2 following intravenous administration of Compound 2, or oral administration of Compound 1, in male Sprague-Dawley rats (n = 5/group), using Solutol HS-15:Water (20:80) Formulation.

Group	Calculated dose of Compound 2 (mg/kg)	C _{max} (μg/mL)	AUC _∞ (μg*h/mL)	F (%)
Compound 2 10 mg/kg IV	10	23.7	6.7	NA
Compound 1	6.7	1.25	2.42	53

10 mg/kg PO ^a				
Compound 1 100 mg/kg PO ^a	67.1	8.90	36.1	80
Compound 1 300 mg/kg PO ^a	201	19.2	90.3	67
Compound 1 600 mg/kg PO ^a	403	17.7	129	58
Compound 1 1000 mg/kg PO ^b	671	19.6	240	64

Abbreviations: C_{max} = maximum plasma concentration; AUC_∞ = area under the plasma concentration time curve from time 0 to infinity; F = oral bioavailability corrected for the molecular weight difference between Compound 1 and Compound 2; NA = not applicable; ^aDose volume = 5 mL/kg; ^bDose volume = 10 mL/kg

Example 12. Single Ascending Dose Pharmacokinetics in Dogs.

[0352] The single dose pharmacokinetics and dose proportionality of Compound 2 were evaluated in non-fasted Beagle dogs following oral gavage of Compound 1, as well as intravenous bolus administration of Compound 2. The dosing formulation for Compound 1 was prepared in Solutol HS-15:Water (10:90 v/v) which provided dosing formulations of Compound 1 as a solution (homogeneous and chemically-stable up to a concentration of 30 mg/mL) and as a emulsion (physically- and chemically-stable at concentrations ranging from 31-100 mg/mL) with dose concentrations ranging from 2-60 mg/mL. The data are shown in Table 19.

Table 19. Mean plasma pharmacokinetic parameters for Compound 2 following intravenous administration of Compound 2, or oral administration of Compound 1, in male Beagle dogs (n = 4/group), using Solutol HS-15:Water (10:90) Formulation.

Group	Calculated dose of Compound 2 (mg/kg)	C _{max} (μg/mL)	AUC _∞ (μg*h/mL)	F (%)
Compound 2 10 mg/kg IV	10	54.5	39.3	NA
Compound 1 10 mg/kg PO, non-fasted	6.7	6.40	26.4	100
Compound 1 30 mg/kg PO, non-fasted	20.1	19.2	68.2	86
Compound 1 100 mg/kg PO, non-fasted	67.1	58.4	228	86
Compound 1 300 mg/kg PO, non-fasted	201	57.1	298	38 ^a

Compound 1 100 mg/kg PO, fasted	67.1	74.4	199	75
------------------------------------	------	------	-----	----

Abbreviations: $C_{\max}$ = maximum plasma concentration; AUC_{∞} = area under the plasma concentration time curve from time 0 to infinity; F = oral bioavailability corrected for the molecular weight difference between Compound 1 and Compound 2; NA = not applicable; ^aValue lower due to 3 of 4 dogs exhibiting emesis, single dog without emesis exhibited F = 79%.

Example 13. Single Ascending Dose Pharmacokinetics in Cynomolgus Monkeys.

[0353] The single dose pharmacokinetics and dose proportionality of Compound 2 were evaluated in non-fasted Cynomolgus monkeys following oral gavage of Compound 1, as well as intravenous bolus administration of Compound 2. The dosing formulation for Compound 1 was prepared in Solutol HS-15:Water (20:80 v/v) for oral dosing (homogeneous and chemically-stable solution up to a concentration of 60 mg/mL; physically- and chemically-stable emulsion at concentrations ranging from 61-200 mg/mL). This study included dose concentrations ranging from 2-120 mg/mL. The data are shown in Table 20.

Table 20. Mean plasma pharmacokinetic parameters for Compound 2 following intravenous administration of Compound 2, or oral administration of Compound 1, in male cynomolgus monkeys (n = 4/group), using Solutol HS-15:Water (20:80) Formulation.

Group	Calculated dose of Compound 2 (mg/kg)	$C_{\max}$ ($\mu\text{g/mL}$)	AUC_{∞} ($\mu\text{g}\cdot\text{h/mL}$)	F (%)
Compound 2 1 mg/kg IV	1	26.0	10.0	NA
Compound 1 10 mg/kg PO, non-fasted	6.7	12.4	64.0	95
Compound 1 30 mg/kg PO, non-fasted	20.1	20.3	131	65
Compound 1 100 mg/kg PO, non-fasted	67.1	61.4	445	66
Compound 1 300 mg/kg PO, non-fasted	201	86.9	1107	55
Compound 1 600 mg/kg PO, non-fasted	403	90.9	1667	41
Compound 1 1000 mg/kg PO, non-fasted	671	97.3	1555	23

Compound 1 100 mg/kg PO, fasted	67.1	62.3	461	69
------------------------------------	------	------	-----	----

Abbreviations: $C_{\max}$ = maximum plasma concentration; AUC_{∞} = area under the plasma concentration time curve from time 0 to infinity; F = oral bioavailability corrected for the molecular weight difference between Compound 1 and Compound 2; NA = not applicable;

Example 14: Assessment of metabolic stability of Compounds 1 and 2 in liver S9, intestinal S9, and Plasma across species.

[0354] The stability of both Compound 1 and Compound 2 in intestinal S9, liver S9 and plasma was determined for five species: human, cynomolgus monkey, beagle dog, Sprague-Dawley® rat and CD-1® mouse, using UPLC/MS-MS methods. Based on the disappearance rate of Compound 1, the half-life was determined in each matrix. Cleavage rates for Compound 1 were compared to those of cefpodoxime proxetil, an orally bioavailable cephalosporin class antibiotic that also undergoes biotransformation in vivo. The data is shown in Table 21.

Table 21. Metabolic stability of Compounds 1 and 2 in liver S9, intestinal S9, and plasma from five species.

Test Article	Test conc.	Species	Plasma Half-life (min)	Liver S9 Half-life (min)	Intestinal S9 Half-life (min)
Compound 1	3 μ M	Human	10.7	1.0	2.9
		Monkey	22.0	0.8	11.2
		Dog	43.9	0.5	49.0
		Rat	1.6	2.3	3.9
		Mouse	4.6	1.2	1.1
Compound 2	3 μ M	Human	>120	>120	>120
		Monkey	>120	>120	>120
		Dog	>120	>120	>120
		Rat	>120	>120	>120
		Mouse	>120	>120	>120
Cefpodoxime proxetil	3 μ M	Human	4.8	3.4	35.6
		Monkey	12.4	4.3	16.9
		Dog	26.2	4.7	>120
		Rat	0.8	10.2	7.4
		Mouse	0.6	2.4	2.5

Example 15: Proposed clinical study of Compound 1 or Compound 1-ethanolate in combination with Ceftibuten for the treatment of ESBL-producing Enterobacteriaceae infections.

[0355] For possible combinations of Compound 1 and Ceftibuten, a Phase I drug-drug interaction study (DDI) will be conducted to ensure safety and appropriate pharmacokinetics of the various combinations. Such studies would be conducted in two parts 1) a randomized, crossover drug-drug interaction study (Part 1) and 2) a repeat-dose safety and PK study (Part 2). Part 1 would consist of a single dose of the given combination, and Part 2 would be conducted over 7-10 days of repeat doses of the given combination and dosing regimen. Following completion of these DDI studies, epithelial lung fluid and renal impairment studies would be conducted, using doses and regimens derived from the results from the DDI studies.

[0356] A compound of formula (I) or (II) and Ceftibuten will be dosed orally. The repeat dose study will be conducted with dosing every 8, 12, or 24 hours. Proposed drug ratios, doses, and dosing regimens are outlined as follows:

[0357] For dosing every 24 hours:

Ceftibuten	Compound of formula (I) or (II)
about 400 mg	about 100 mg to about 1600 mg
about 600 mg	about 150 mg to about 2400 mg
about 800 mg	about 200 mg to about 3200 mg

[0358] For dosing every 12 hours:

Ceftibuten	Compound of formula (I) or (II)
about 200 mg	about 50 mg to about 800 mg
about 300 mg	about 75 mg to about 1200 mg
about 400 mg	about 100 mg to about 1600 mg
about 600 mg	about 150 mg to about 2400 mg
about 800 mg	about 200 mg to about 3200 mg

[0359] For dosing every 8 hours:

Ceftibuten	Compound of formula (I) or (II)
about 200 mg	about 50 mg to about 800 mg
about 300 mg	about 75 mg to about 1200 mg
about 400 mg	about 100 mg to about 1600 mg
about 600 mg	about 150 mg to about 2400 mg

Example 16. Excipient Screening Study

[0360] The solubility of Compound 1-ethanolate was evaluated in a series of pharmaceutically acceptable excipients that are commonly used to prepare self-emulsifying drug delivery systems (SEDDS) for oral drug delivery. This class of formulation results in a solid solution of drug substance and excipients that enable the delivery of the drug substance and prevents precipitation upon contact with aqueous environments (i.e. the stomach).

[0361] The indicated quantity of Compound 1-ethanolate was weighed into a 3 mL screw top glass vial outfitted with a magnetic stir bar. The vial was transferred to a controlled temperature heating block and 100 μ L aliquots of each excipient was added with stirring. After each addition, the sample was evaluated for dissolution of the drug substance and additional aliquots of excipient were added until a clear, colorless solution was obtained. Table 22 summarizes the solubility screening for 25 excipients.

Table 22: Solubility of Compound 1-Ethanolate in SEDDS Excipients

Vial #	Excipient	Compound 1-ethanolate (mg)	Excipient (μ L)	Temp	Solubility		
1	Gelucire 44/14 (Lauroyl polyoxyl-32 glycerides)	113.8	400	50°C	>	284.5	mg/mL
2	Polysorbate 80 (Polyoxyethylene (20) sorbitan monooleate)	95.4	400	50°C	>	238.5	mg/mL
3	Myrj 52 (PEG-40 Stearate)	110.9	400	50°C	>	277.3	mg/mL
4	Kolliphor EL (Polyoxyl 35 hydrogenated castor oil)	112.8	300	50°C	>	376	mg/mL
5	Kolliphor RH (PEG-40 Hydrogenated Castor Oil)	103.1	600	50°C	~	171.8	mg/mL
6	Poloxamer 188	112.7	500	65°C	>	225.4	mg/mL
7	Span 20 (Sorbitan monolaurate)	113.6	500	50°C	>	227.2	mg/mL
8	TPGS	68.6	300	50°C	>	228.7	mg/mL
9	Labrasol (Caprylocaproyl polyoxyl-8 glycerides)	117	200	50°C	>	585	mg/mL
10	Labrafil M1944CS (Oleoyl polyoxyl-6 glycerides)	112.1	800	50°C	>	140.1	mg/mL
11	Labrafil M2125CS (Linoleoyl polyoxyl-6 glycerides)	121.2	800	50°C	>	151.5	mg/mL
12	Peceol (Glycerol monooleate)	124	300	50°C	>	413.3	mg/mL
13	Lauroglycol (Propylene Glycol Monolaurate)	145.9	500	50°C	>	291.8	mg/mL
14	Plurol Oleique CC497 (Polyglyceryl-3 dioleate)	112.4	300	50°C	>	374.7	mg/mL
15	Maisine CC (glyceryl monolinoleate)	110.4	300	50°C	>	368	mg/mL
16	Propylene Glycol	121.4	100	50°C	>	1214	mg/mL

17	Glycerol	111.6	2000	50°C	<	55.8	mg/mL
18	PEG-3350	111.8	400	65°C	>	279.5	mg/mL
19	PEG-1500	114.6	300	50°C	>	382	mg/mL
20	PEG-400	99.7	200	50°C	>	498.5	mg/mL
21	Ethanol	113.6	200	50°C	>	568	mg/mL
22	Triacetin (glycerin triacetate)	92.6	1100	50°C	>	84.18	mg/mL
23	Phosal 50 PG (50% PC and propylene glycol)	83.8	200	50°C	>	419	mg/mL
24	Capmul MCM (Caprylic/capric mono- & diglycerides)	70.2	200	50°C	>	351	mg/mL
25	Span 80	82.9	300	50°C	>	276.3	mg/mL

Example 17: Preparation of a Compound 1-ethanolate SEDDS Formulation.

[0362] A 1:1 mixture by weight of PEG 1500 and Myrj 52 was prepared by weighing 5.0 g of Myrj 52 into a clean 50 mL beaker. A magnetic stir bar was added and 5.0 g of molten PEG 1500 was weighed into the beaker. This mixture was stirred on a ~55 °C hot plate until a uniform solution was obtained. A 6.5 mL aliquot of this excipient blend was added to a scintillation vial containing 2.002 g of Compound 1-ethanolate (~280 mg/mL Compound 1-ethanolate). The blend was then stirred at ~55 °C on a hot plate using a magnetic stir bar until a clear, homogeneous solution was obtained.

Example 18: Preparation and Dissolution of a Compound 1-ethanolate HPMC Capsules.

[0363] The 280 mg/mL Compound 1-ethanolate PEG 1500/Myrj 52 stock formulation (from example 17) was re-melted on a ~55 °C hot plate with stirring. The resulting clear solution was filled into a Size 0 white, opaque HPMC capsule by adding 0.55 mL of the liquid to the capsule body using a positive displacement pipette. The capsule was capped and allowed to solidify at room temperature before transferring to 2-8 °C for storage. The dissolution profile was collected for the 150 mg Compound 1-ethanolate capsule by using USP Apparatus II conditions (50 RPM, 450 mL pH 2 water at 38 °C) with manual time pulls. The samples were removed at t = 0, 5, 10, 15, 20, 30, 45 and 60 minutes and filtered through a 0.22 µm syringe filter and analyzed using HPLC. The dissolution profile for the 150 mg capsule of Compound 1-ethanolate is shown in FIG. 2.

Example 19: Preparation and Dissolution of a Compound 1-ethanolate / Cefitibuten Fixed Dose Combination HPMC Capsules.

[0364] The 280 mg/mL Compound 1-ethanolate PEG 1500/Myrj 52 stock formulation (from example 17) was re-melted on a ~55 °C hot plate with stirring. The resulting clear solution was filled into a Size 2 white, opaque HPMC capsule by adding 0.353 mL of the liquid to the capsule body using a positive displacement pipette. The capsule was capped and allowed to solidify at room

temperature before transferring to 2-8 °C for storage. The Compound 1-ethanolate capsule was placed into the body of a Size 00 white, opaque HPMC capsule and 100 mg of ceftibuten was filled around the smaller capsule and the larger capsule was capped to yield a combination product in a single size 00 capsule. The dissolution profile was collected for the 100 mg Compound 1-ethanolate /100 mg ceftibuten capsule by using USP Apparatus II conditions (50 RPM, 450 mL pH 8 PBS bugger at 38 °C) with manual time pulls. The samples were removed at t = 0, 5, 10, 15, 20, 30, 45 and 60 minutes and filtered through a 0.22 µm syringe filter and analyzed using HPLC. The dissolution profile for the combination product is shown in FIG. 3.

Example 20: In vitro Antibacterial Assays Demonstrating Potentiation of Beta-Lactam Antibiotics by Beta-Lactamase Inhibition.

[0365] Broth microdilution minimum inhibitory concentration assays were performed according to CLSI methods with amoxicillin, cefaclor, cefalexin, cefdinir, cefditoren, cefixime, cefpodoxime, ceftibuten, and cefuroxime alone or in combination with Compound 2 fixed at 4 mg/L. Levofloxacin and amoxicillin-clavulanate were also tested as comparators.

[0366] One hundred representative isolates of Enterobacteriaceae expressing Class A ESBL (n=25), Class A KPC (n=25), Class C (n=25), and Class D OXA-48 (n=25) enzymes were used. beta-lactamase genes were verified using polymerase chain reaction (PCR) while expression of these genes was determined phenotypically.

[0367] Commercially available oral beta-lactam antibiotics were titrated (tested range 0.016 to 32 mg/L) across 96-well microtiter plates and were mixed with either cation adjusted Mueller Hinton broth (CAMHB), or Compound 2 supplemented CAMHB at 4 mg/L. Levofloxacin and amoxicillin-clavulanate (2:1 ratio) were also tested (range 0.016 to 32 mg/L and 0.06 to 128 mg/L, respectively). Bacterial inocula at a final concentration of 2-5x10⁵ CFU/mL were used. Microtiter plates were incubated aerobically at 37°C for 18-20 hours and MICs were read visually. Table 23a and 23b show the MIC₅₀ and MIC₉₀ results:

Table 23a. MIC₅₀ and MIC₉₀ Summary for combinations of Compound 2 and beta-lactam antibiotics.

Beta-Lactam	BLI	All		ESBL		KPC	
		MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀
Ceftibuten	-	16	≥64	4	≥64	16	≥64
	Compound 2	0.125	1	0.125	0.25	0.125	1
Amoxicillin	Clavulanate	64	≥256	16	≥64	≥64	≥256

		All		ESBL		KPC	
Beta-Lactam	BLI	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀
Levofloxacin	-	16	≥64	8	32	≥64	≥64
Amoxicillin	-	≥256	≥256	≥256	≥256	≥64	≥256
	Compound 2	64	≥256	64	≥256	≥64	≥256
Cefaclor	-	≥64	≥64	≥64	≥64	≥64	≥64
	Compound 2	2	≥64	1	2	4	≥64
Cefalexin	-	≥64	≥64	≥64	≥64	≥64	≥64
	Compound 2	8	≥64	8	16	16	≥64
Cefdinir	-	≥64	≥64	32	≥64	≥64	≥64
	Compound 2	0.5	16	0.5	1	0.5	16
Cefditoren	-	≥64	≥64	≥64	≥64	≥64	≥64
	Compound 2	0.5	2	0.5	1	0.5	4
Cefexime	-	≥64	≥64	≥64	≥64	≥64	≥64
	Compound 2	0.5	2	0.5	2	0.5	2
Cefpodoxime	-	≥64	≥64	≥64	≥64	≥64	≥64
	Compound 2	1	8	0.5	2	1	8
Cefuroxime	-	≥64	≥64	≥64	≥64	≥64	≥64
	Compound 2	8	≥64	8	16	16	≥64

Table 23b. MIC₅₀ and MIC₉₀ Summary for combinations of Compound 2 and beta-lactam antibiotics vs. Class C and Class D beta-lactamase producing strains.

		OXA		Class C	
Beta-lactam	BLI	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀
Ceftibuten	-	16	≥64	16	≥64
	Compound 2	0.125	1	0.125	1

		OXA		Class C	
Beta-lactam	BLI	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀
Amoxicillin	Clavulanate	≥256	≥256	32	≥256
Levofloxacin	-	32	≥64	4	32
Amoxicillin	-	≥256	≥256	≥256	≥256
	Compound 2	≥256	≥256	64	≥256
Cefaclor	-	≥64	≥64	≥64	≥64
	Compound 2	4	≥64	2	32
Cefaxelin	-	≥64	≥64	≥64	≥64
	Compound 2	16	≥64	8	≥64
Cefdinir	-	≥64	≥64	32	≥64
	Compound 2	2	32	0.25	2
Cefditoren	-	≥64	≥64	≥64	≥64
	Compound 2	1	4	0.5	0.2
Cefexime	-	≥64	≥64	≥64	≥64
	Compound 2	0.5	2	0.5	4
Cefpodoxime	-	≥64	≥64	≥64	≥64
	Compound 2	1	8	0.5	8
Cefuroxime	-	≥64	≥64	≥64	≥64
	Compound 2	8	≥64	8	≥64

Example 21: In vitro comparison of oral ceftibuten/Compound 2 with IV therapeutics against MDR Enterobacteriaceae

[0368] Broth microdilution minimum inhibitory concentration (MIC) assays were carried out according to CLSI guidelines. BLIs were fixed at a concentration of 4 mg/L (Compound 2, tazobactam, and avibactam) or 8 mg/L (vaborbactam). The antibacterial activity of Compound 2 in combination with ceftibuten was compared to ceftibuten alone, meropenem, piperacillin/tazobactam, ceftazidime/avibactam, meropenem/vaborbactam, tobramycin, and tigecycline in 193 strains of Enterobacteriaceae expressing Class A ESBL (N=33), Class A KPC (N=77), Class D OXA-48 (39), and Class C (44) enzymes. Beta-lactamase genes were verified using polymerase chain reaction while gene expression was determined phenotypically. MIC results were interpreted using CLSI M100 Ed. 29 (2019) or EUCAST v9.0 Clinical Breakpoints (2019). Tables 24a-24d and FIG. 4 summarize the results.

Table 24a: MICs of Ceftibuten/Compound 2 and comparators in ESBL-expressing *Enterobacteriaceae* (n = 33).

Test Article			Number of strains at MIC value (mg/L)									
	MIC ₅₀ mg/L	MIC ₉₀ mg/L	≤ 0.06	0.125	0.25	0.5	1	2	4	8	16	≥32
Ceftibuten	4	≥ 32	0	2	3	2	4	2	8	5	1	6
Ceftibuten + Compound 2 (4 mg/L)	0.125	0.25	12	15	4	1	1	0	0	0	0	0
Ceftazidime + avibactam (4 mg/L)	0.5	0.5	2	4	10	15	1	0	0	0	1	0
Meropenem + vaborbactam (8 mg/L)	≤ 0.06	≤ 0.06	31	2	0	0	0	0	0	0	0	0
Piperacillin + tazobactam (4 mg/mL)	16	≥ 32	0	0	0	0	0	5	7	3	3	15
Tobramycin	≥ 32	≥ 32	0	0	0	3	5	2	1	1	4	17
Tigecycline	0.5	2	0	4	10	11	3	3	2	0	0	0
Meropenem	≤ 0.06	0.125	29	1	0	0	1	1	0	1	0	0

Strain set composed of *C. freundii* (1), *E. coli* (11), *K. oxytoca* (3), and *K. pneumoniae* (18)

Table 24b: MICs of Ceftibuten/ Compound 2 and comparators in KPC-expressing *Enterobacteriaceae* (n = 77).

Test Article			Number of strains at MIC value (mg/L)									
	MIC ₅₀ mg/L	MIC ₉₀ mg/L	≤ 0.06	0.125	0.25	0.5	1	2	4	8	16	≥32
Ceftibuten	16	≥ 32	0	0	0	2	2	4	7	13	24	25
Ceftibuten + Compound 2 (4 mg/L)	0.25	1	10	20	17	22	4	0	2	1	0	1
Ceftazidime + avibactam (4 mg/L)	1	8	0	0	2	10	27	19	7	10	1	1
Meropenem + vaborbactam (8 mg/L)	≤ 0.06	2	40	6	6	7	9	5	3	1	0	0
Piperacillin + tazobactam (4 mg/mL)	≥ 32	≥ 32	0	0	0	0	0	0	1	0	0	76
Tobramycin	≥ 32	≥ 32	0	0	1	3	0	2	0	3	3	65
Tigecycline	1	2	0	0	4	19	34	15	5	0	0	0

Meropenem	≥ 32	≥ 32	1	0	0	1	0	3	4	11	13	44
-----------	------	------	---	---	---	---	---	---	---	----	----	----

Strain set composed of *E. cloacae* (4), *E. coli* (4), *K. pneumoniae* (67), and *K. oxytoca* (2)

Table 24c: MICs of Ceftibuten/ Compound 2 and comparators in OXA-48-expressing Enterobacteriaceae (n = 39).

Test Article			Number of strains at MIC value (mg/L)									
	MIC ₅₀ mg/L	MIC ₉₀ mg/L	≤ 0.06	0.125	0.25	0.5	1	2	4	8	16	≥32
Ceftibuten	16	≥ 32	1	0	0	1	0	0	2	4	13	18
Ceftibuten + Compound 2 (4 mg/L)	0.125	2	8	15	6	2	3	1	1	2	0	1
Ceftazidime + avibactam (4 mg/L)	0.5	2	0	1	9	13	10	4	1	0	0	1
Meropenem + vaborbactam (8 mg/L)	2	8	0	2	1	9	3	11	5	4	2	2
Piperacillin + tazobactam (4 mg/mL)	≥ 32	≥ 32	0	0	0	0	0	0	1	0	1	37
Tobramycin	16	≥ 32	0	0	0	3	4	6	1	1	6	18
Tigecycline	0.5	2	0	1	4	16	11	4	1	2	0	0
Meropenem	2	8	1	1	4	6	6	10	3	4	2	2

Strain set composed of *E. cloacae* (1), *E. coli* (15), *K. pneumoniae* (22), and *M. morgannii* (1)

Table 24d: MICs of Ceftibuten/ Compound 2 and comparators in AmpC/CMY-expressing Enterobacteriaceae (n = 44).

Test Article			Number of strains at MIC value (mg/L)									
	MIC ₅₀ mg/L	MIC ₉₀ mg/L	≤ 0.06	0.125	0.25	0.5	1	2	4	8	16	≥32
Ceftibuten	≥ 32	≥ 32	1	0	0	0	1	5	3	7	7	20
Ceftibuten + Compound 2 (4 mg/L)	0.25	2	13	6	8	8	2	3	3	0	1	0
Ceftazidime + avibactam (4 mg/L)	0.5	2	2	2	6	12	14	4	3	1	0	0
Meropenem + vaborbactam (8 mg/L)	≤ 0.06	0.25	37	1	3	1	2	0	0	0	0	0
Piperacillin + tazobactam (4 mg/mL)	≥ 32	≥ 32	0	1	1	0	3	5	5	4	1	24

Tobramycin	≥ 32	≥ 32	0	0	0	2	2	0	3	3	6	28
Tigecycline	1	4	0	4	6	8	10	10	4	2	0	0
Meropenem	≤ 0.06	0.5	34	2	2	2	1	0	3	0	0	0

Strain set composed of *C. freundii* (2), *E. aerogenes* (5), *E. cloacae* (5), *E. coli* (8), *K. oxytoca* (2), *K. pneumoniae* (10), *P. mirabilis* (1), *Salmonella spp.* (2), and *S. marcescens* (9)

Example 22: In vivo PK/PD in neutropenic mouse thigh infection model

[0369] The in vivo efficacy study of ceftibuten with and without Compound 2 was conducted in the neutropenic mouse thigh infection model (Dr. David Nicolau, Hartford Hospital, Hartford, CT) utilizing a total of 21 strains of Enterobacteriaceae expressing various beta-lactamases (11 ESBL, 1 ESBL+OXA, 5 KPC, 3 OXA-48 and 1 AmpC). In MIC testing, Compound 2 fixed at 4 mg/L reduced ceftibuten MICs from ≥32 µg/mL to ≤2 µg/mL. All strains were demonstrated to grow in neutropenic mouse thigh in the presence of a humanized dosing regimen of ceftibuten administered three times at 300 mg over 24 hrs.

[0370] Dose fractionation studies were conducted to determine the PK/PD index for Compound 2 that best described its efficacy in combination with ceftibuten. Six strains were examined. Mice were administered a human simulated regimen (HSR) of ceftibuten 300 mg q8h with Compound 2 at 1.2, 4 or 12 mg/kg/day (2 strains) or 4, 12 or 38 mg/kg/day (4 strains). The principle behind these assays is shown in Table 25 below. If a minimally efficacious dose of Compound 2 given as a single dose demonstrates maximal efficacy, then C_{max} is the predominant driver of efficacy. If 4 divided doses demonstrate maximal efficacy, then time above threshold is favored. If all dose fractionations give similar efficacy, then exposure (AUC) is the likely driver of BLI efficacy.

Table 25. Interpretative efficacy criteria for dose fractionation studies.

	Efficacy Outcome		
	C _{max}	Time above threshold	AUC
Single dose	+++	+	+++
Two divided doses	++	++	+++
Four divided doses	+	+++	+++

+++ , maximum efficacy; ++ , intermediate efficacy; + , minimum efficacy

[0371] As shown in FIG. 8 and FIG. 9 with *E. coli* 617 expressing AmpC, TEM-1, CTXM-15, SHV-5 and SHV-1, and *E. coli* C11-23 expressing AmpC, TEM-1 and CTXM-15, respectively, rescue of ceftibuten activity by Compound 2 was similar regardless of the manner in which the Compound 2 dose was fractionated. The results thus support AUC as the PK/PD driver for BLI efficacy.

[0372] Dose ranging studies were conducted to assess the ability of various doses of Compound 2 to potentiate the in vivo bactericidal activity of humanized ceftibuten exposures against these 21 beta-lactamase-producing Enterobacteriaceae strains and 1 wild-type Enterobacteriaceae strain. All treatment groups were administered the ceftibuten 300 mg q8h human-simulated regimen alone, or in combination with Compound 2 q8h at a ceftibuten:Compound 2 ratio of 10:1, 3.16:1, 1:1, 1:3.16 or 1:10 (approximately corresponding to ceftibuten 300:Compound 2@30, 95, 300, 950 or 3,000 mg, respectively). Compound 2 rescued ceftibuten activity in all 21 beta-lactamase-expressing strains. Net bacterial stasis was achieved at a Compound 2 median exposure of 8.98 mg*h/L with an interquartile range of 3.06-18.56 mg*h/L.

Table 26. Free Compound 2 exposure indexed to combination MIC ($fAUC_{0-24}/MIC$) required to achieve net bacterial stasis for ceftibuten 300 mg q8h. $fAUC_{0-24}$ units of mg*h/L.

Isolate (MIC, mg/L) ^a	Enzyme(s)	Compound 2 Exposure Required to Achieve Stasis	
		$fAUC_{0-24}/MIC$ (mg*h/L) ^a	R ²
EC 614 (1)	OXA-48	0.72	0.816
EC 617 (0.5)	AmpC, CTX-M-15, SHV-5, SHV-1, TEM-1	7.61	0.825
EC 636 (0.125)	CTX-M-15	18.89	0.972
EC 639 (0.125)	CTX-M-15, TEM-1	18.23	0.961
EC C11-23 (0.5)	AmpC, CTX-M-15, TEM-1	0.13	0.915
ECL 138 (1)	KPC	5.92	0.810
ECL 139 (1)	AmpC (p99), CTX-M-3, TEM-1	193.55	0.851
ECL 150 (0.25)	CTX-M-15	43.22	0.586
KP 630 (0.5)	CTX-M-15, SHV-WT, TEM-WT	1.53	0.929
KP 631 (0.5)	CTX-M-15, SHV-WT, TEM-WT	14.91	0.831
KP 774 (1)	KPC	0.55	0.921
KP 776 (1)	KPC	30.90	0.816
KP 780 (2)	KPC	8.98	0.843
KP 783 (2)	CMY-2, TEM-1	4.59	0.872
KP 785 (0.25)	OXA-204	0.10	0.843
KP 786 (0.5)	OXA-48	17.10	0.949
KP 787 (1)	KPC	6.43	0.634
KP 813 (0.125)	CTX-M-15, OXA-48, SHV-12, TEM- 1	8.95	0.913
KP 814 (0.125)	SHV-12, TEM-1	15.82	0.950
KP 816 (0.125)	CTX-M-3, SHV-12, TEM-1	28.63	0.930
KP 819 (0.25)	SHV-12	13.78	0.930
Median (interquartile range)		8.98 (3.06–18.56)	

^a Based on the MICs of ceftibuten/Compound 2 combination at a fixed Compound 2 concentration of 4 mg/L.

Example 23. Preparation of 560 mg/mL Compound 1 Propylene Glycol/Peg 400/ Imwitor® 742/TPGS SEDDS Formulation.

[0373] A 20/45/10/25 volume % stock solution of propylene glycol (PG), PEG 400, Imwitor® 742 and TPGS was prepared by pipetting 2.0 mL of propylene glycol and 4.5 mL of PEG 400 into a clean 20 mL scintillation vial using a positive displacement pipette. A magnetic stir bar was added and the vial was stirred at ~60°C on a stir plate. The Imwitor® 742 and TPGS were melted in a ~60°C oven and thoroughly mixed per the manufacturer's instructions before addition to the PG/PEG 400 solution. The Imwitor® 742 was added by pipetting 1.0 mL of molten excipient using a positive displacement pipette. Once a uniform solution was obtained, the TPGS was added by pipetting 2.5 mL using a positive displacement pipette. This mixture was stirred on a ~60°C hot plate until a uniform solution was obtained. A 0.5 mL aliquot of this stock solution was added to a glass vial containing 700 mg of Compound 1-ethanolate and blend was then stirred at ~60°C on a hot plate using a magnetic stir bar until a clear, homogeneous solution was obtained. The final solution was assayed by HPLC against a Compound 1 standard and found to be 563.2 mg/mL. A 50 mg capsule dose of this formulation was prepared by pipetting 89 µL into a Size 2 HPMC white, opaque capsule.

Example 24. Preparation of 350 mg/mL Compound 1 Peg 3350/ Imwitor® 742/TPGS SEDDS Formulation.

[0374] A 55/20/25 volume % stock solution of PEG 3350, Imwitor® 742 and TPGS was prepared by pipetting 5.5 mL of PEG 3350 into a clean 20 mL scintillation vial using a positive displacement pipette. A magnetic stir bar was added and the vial was stirred at ~60°C on a stir plate. The Imwitor® 742 and TPGS were melted in a ~60°C oven and thoroughly mixed per the manufacturer's instructions before addition to the PEG 3350. The Imwitor® 742 was added by pipetting 2.0 mL of molten excipient using a positive displacement pipette. Once a uniform solution was obtained, the TPGS was added by pipetting 2.5 mL using a positive displacement pipette. This mixture was stirred on a ~60°C hot plate until a uniform solution was obtained. A 0.5 mL aliquot of this stock solution was added to a glass vial containing 700 mg of Compound 1-ethanolate and blend was then stirred at ~60°C on a hot plate using a magnetic stir bar. After ~30 minutes the formulation was filtered through a 0.45µm PTFE syringe filter to remove undissolved VNR-7145. The resulting clear, homogeneous solution was assayed by HPLC against a Compound 1 standard and found to be 355.1 mg/mL. A 50 mg capsule dose of this formulation was prepared by pipetting 141 µL into a Size 2 HPMC white, opaque capsule.

Example 25. Preparation of 400 mg/mL Compound 1 Propylene Glycol/Peg 400/ TPGS SEDDS Formulation.

[0375] A 20/20/60 volume % stock solution of PEG 3350, Imwitor® 742 and TPGS was prepared by pipetting 5.5 mL of PEG 3350 into a clean 20 mL scintillation vial using a positive displacement pipette. A magnetic stir bar was added and the vial was stirred at ~60°C on a stir plate. The Imwitor® 742 and TPGS were melted in a ~60°C oven and thoroughly mixed per the manufacturer's instructions before addition to the PEG 3350. The Imwitor® 742 was added by pipetting 2.0 mL of molten excipient using a positive displacement pipette. Once a uniform solution was obtained, the TPGS was added by pipetting 2.5 mL using a positive displacement pipette. This mixture was stirred on a ~60°C hot plate until a uniform solution was obtained. A 0.5 mL aliquot of this stock solution was added to a glass vial containing 700 mg of Compound 1-ethanolate and blend was then stirred at ~60°C on a hot plate using a magnetic stir bar. After ~30 minutes the formulation was filtered through a 0.45µm PTFE syringe filter to remove undissolved Compound 1-ethanolate. The resulting clear, homogeneous solution was assayed by HPLC against a Compound 1 standard and found to be 355.1 mg/mL. A 50 mg capsule dose of this formulation was prepared by pipetting 141 µL into a Size 2 HPMC white, opaque capsule.

Example 26. Preparation of 350 mg/mL Compound 1 Peg 1500/ TPGS SEDDS Formulation.

[0376] A 25/75 volume % stock solution of PEG 1500 and TPGS was prepared by first melting the excipients in a ~60°C oven and thoroughly mixed per the manufacturer's instructions. The vehicle stock was prepared by pipetting 5.0 mL of PEG 1500 and 15.0 mL of PEG 1500 into a clean 20 mL scintillation vial using a positive displacement pipette. This mixture was stirred on a ~60°C hot plate until a uniform solution was obtained. A ~5 mL batch of 350 mg/mL Compound 1 was prepared by combining 2.1 g of Compound 1-ethanolate and 2.6 g of the vehicle stock into a glass vial outfitted with a magnetic stir bar and mixing at ~60°C on a hot plate until a clear, homogeneous solution was obtained. The final solution was assayed by HPLC against a Compound 1 standard and found to be 346.6 mg/mL. A 250 mg capsule dose of this formulation was prepared by pipetting 622 µL into a Size 00 HPMC white, opaque capsule. The capsules were capped and allowed to solidify at room temperature before transferring to 2-8°C for storage. The dissolution profile was collected for the 250 mg Compound 1 capsule by using USP Apparatus II conditions (50 RPM, 450 mL pH 2 water at 38°C) with manual time pulls. The samples were removed at t = 0, 5, 10, 15, 20, 30, 45 and 60 minutes and filtered through a 0.22 µm syringe filter and analyzed using HPLC. The dissolution profile for the 250 mg capsule of Compound 1 is shown in FIG. 10.

Example 27. Preparation and Dissolution of 250 mg Compound 1 HPMC Capsules.

[0377] A 20/20/60 volume % stock solution of propylene glycol (PG), PEG 400, and TPGS was prepared by pipetting 4.0 mL of propylene glycol and 4.4 mL of PEG 400 into a clean 50 mL beaker using a positive displacement pipette. A magnetic stir bar was added and the beaker was stirred at



~60°C on a stir plate. The TPGS was melted in a ~60°C oven and thoroughly mixed per the manufacturer's instructions before addition to the PG/PEG 400 solution. The TPGS was added by pipetting 12.0 mL using a positive displacement pipette. This vehicle stock was stirred on a ~60°C hot plate until a uniform solution was obtained. A ~20 mL batch of 400 mg/mL Compound 1 was prepared by combining 8.71 g of Compound 1-ethanolate and 12.3 g of the vehicle stock into a 100 mL round bottom flask outfitted with a magnetic stir bar and mixing at ~60°C on a hot plate until a clear, homogeneous solution was obtained. The final solution was assayed by HPLC against a Compound 1 standard and found to be 401.7 mg/mL. A 250 mg capsule dose of this formulation was prepared by pipetting 622 µL into a Size 0 HPMC white, opaque capsule. The capsules were capped and allowed to solidify at room temperature before transferring to 2-8°C for storage. The dissolution profile was collected for the 250 mg Compound 1 capsule by using USP Apparatus II conditions (75 RPM, 900 mL pH 2 water at 38°C) with manual time pulls. The samples were removed at t = 0, 5, 10, 15, 20, 30, 45 and 60 minutes and filtered through a 0.22 µm syringe filter and analyzed using HPLC. The dissolution profile for the 250 mg capsule of Compound 1 is shown in FIG. 11.

CLAIMS

WHAT IS CLAIMED:

1. A pharmaceutical composition comprising:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

- (ii) ceftibuten.

2. The pharmaceutical composition of claim 1, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





3. The pharmaceutical composition of claim 1, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



4. The pharmaceutical composition of any one of claims 1-3, wherein the pharmaceutical composition is formulated for oral administration.
5. The pharmaceutical composition of any one of claims 1-4, wherein the pharmaceutical composition is formulated as an emulsion.
6. The pharmaceutical composition of any one of claims 1-4, wherein the pharmaceutical composition is formulated as a microemulsion.
7. The pharmaceutical composition of any one of claims 1-4, wherein the pharmaceutical composition is formulated as a Self-Emulsifying Drug Delivery System (SEDDS).
8. The pharmaceutical composition of any one of claims 1-7, further comprising a hydrophilic solubilizer.
9. The pharmaceutical composition of claim 8, wherein the hydrophilic solubilizer is a hydrophilic polymer.
10. The pharmaceutical composition of claim 9, wherein the hydrophilic polymer is a polyethylene glycol.
11. The pharmaceutical composition of any one of claims 1-10, further comprising a surfactant.
12. The pharmaceutical composition of claim 11, wherein the surfactant is a polyoxyethylene stearate.
13. The pharmaceutical composition of any one of claims 1-11, further comprising a pharmaceutically acceptable excipient.
14. The pharmaceutical composition of any one of claims 1-7, further comprising polyethylene glycol, caprylic/capric glycerides, and tocopheryl polyethylene glycol succinate.
15. The pharmaceutical composition of any one of claims 1-7, further comprising propylene glycol, polyethylene glycol, and tocopheryl polyethylene glycol succinate.
16. The pharmaceutical composition of any one of claims 1-15, wherein the pharmaceutical composition is formulated as a capsule.
17. The pharmaceutical composition of any one of claims 1-15, wherein the pharmaceutical composition is formulated as a tablet.
18. The pharmaceutical composition of any one of claims 1-15, wherein the pharmaceutical composition is formulated as a powder for reconstitution.
19. The pharmaceutical composition of claim 18, wherein the powder for reconstitution is reconstituted with a liquid carrier to form an oral suspension.

20. The pharmaceutical composition of any one of claims 1-15, wherein the pharmaceutical composition is formulated as a solution.
21. A pharmaceutical composition comprising:
- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
 - (ii) ceftibuten.
22. A method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:
- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
 - (ii) ceftibuten.
23. A method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:
- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
 - (ii) ceftibuten.
24. The method of claim 23 wherein the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.
25. The method of claim 23 or 24, wherein the method comprises administering to the subject in need thereof:
- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:






formula (II);


wherein:

- each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);
- or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and
- (ii) ceftibuten.

26. The method of claim 25, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





27. The method of claim 25, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



28. The method of any one of claims 25-27, wherein the compound of formula (I) or (II) and ceftibuten are formulated for oral administration.
29. The method of any one of claims 25-28, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and ceftibuten are administered sequentially.
30. The method of any one of claims 25-28, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and ceftibuten are administered concurrently.

31. A method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) ceftibuten.



32. The method of claim 31, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



33. The method of claim 31, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:




34. The method of any one of claims 31-33, wherein the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.
35. The method of any one of claims 31-34, wherein the compound of formula (I) or (II) and ceftibuten are formulated for oral administration.
36. The method of any one of claims 31-35, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered sequentially.
37. The method of any one of claims 31-35, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and ceftibuten are administered concurrently.
38. A pharmaceutical composition comprising:
- a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

- each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);
- or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and
- cefexime.

39. The pharmaceutical composition of claim 38, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





40. The pharmaceutical composition of claim 38, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



41. The pharmaceutical composition of any one of claims 38-40, wherein the pharmaceutical composition is formulated for oral administration.
42. The pharmaceutical composition of any one of claims 38-41, wherein the pharmaceutical composition is formulated as an emulsion.
43. The pharmaceutical composition of any one of claims 38-41, wherein the pharmaceutical composition is formulated as a microemulsion.
44. The pharmaceutical composition of any one of claims 38-41, wherein the pharmaceutical composition is formulated as a Self-Emulsifying Drug Delivery System (SEDDS).
45. The pharmaceutical composition of any one of claims 38-43, further comprising a hydrophilic solubilizer.
46. The pharmaceutical composition of claim 45, wherein the hydrophilic solubilizer is a hydrophilic polymer.
47. The pharmaceutical composition of claim 46, wherein the hydrophilic polymer is a polyethylene glycol.
48. The pharmaceutical composition of any one of claims 38-47, further comprising a surfactant.
49. The pharmaceutical composition of claim 48, wherein the surfactant is a polyoxyethylene stearate.
50. The pharmaceutical composition of any one of claims 38-49, further comprising a pharmaceutically acceptable excipient.
51. The pharmaceutical composition of any one of claims 38-44, further comprising polyethylene glycol, caprylic/capric glycerides, and tocopheryl polyethylene glycol succinate.
52. The pharmaceutical composition of any one of claims 38-44, further comprising propylene glycol, polyethylene glycol, and tocopheryl polyethylene glycol succinate.


53. The pharmaceutical composition of any one of claims 38-52, wherein the pharmaceutical composition is formulated as a capsule.
54. The pharmaceutical composition of any one of claims 38-52, wherein the pharmaceutical composition is formulated as a tablet.
55. The pharmaceutical composition of any one of claims 38-52, wherein the pharmaceutical composition is formulated as a powder for reconstitution.
56. The pharmaceutical composition of claim 55, wherein the powder for reconstitution is reconstituted with a liquid carrier to form an oral suspension.
57. The pharmaceutical composition of any one of claims 38-52, wherein the pharmaceutical composition is formulated as solution.
58. A pharmaceutical composition comprising:
- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
 - (ii) cefixime.
59. A method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:
- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
 - (ii) cefixime.
60. A method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:
- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
 - (ii) cefixime.
61. The method of claim 60, wherein the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.
62. The method of claim 60 or 61, wherein the method comprises administering to the subject in need thereof:
- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof.




wherein:

- each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);
- or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and
- (ii) cefixime.

63. The method of claim 62, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





64. The method of claim 62, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



65. The method of any one of claims 62-64, wherein the compound of formula (I) or (II) and cefixime are formulated for oral administration.


66. The method of any one of claims 62-65, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefixime are administered sequentially.
67. The method of any one of claims 62-65, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefixime are administered concurrently.
68. A method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:
- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

- each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);
- or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and
- (ii) cefixime.



69. The method of claim 68, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



70. The method of claim 68, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



71. The method of any one of claims 68-70, wherein the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.
72. The method of any one of claims 68-71, wherein the compound of formula (I) or (II) and cefixime are formulated for oral administration.
73. The method of any one of claims 68-72, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered sequentially.
74. The method of any one of claims 68-72, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefixime are administered concurrently.
75. A pharmaceutical composition comprising:
- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:


- each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);
- or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) cefditoren pivoxil.

76. The pharmaceutical composition of claim 75, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



77. The pharmaceutical composition of claim 75, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





78. The pharmaceutical composition of any one of claims 75-77, wherein the pharmaceutical composition is formulated for oral administration.
79. The pharmaceutical composition of any one of claims 75-78, wherein the pharmaceutical composition is formulated as an emulsion.
80. The pharmaceutical composition of any one of claims 75-78, wherein the pharmaceutical composition is formulated as a microemulsion.
81. The pharmaceutical composition of any one of claims 75-78, wherein the pharmaceutical composition is formulated as a Self-Emulsifying Drug Delivery System (SEDDS).
82. The pharmaceutical composition of any one of claims 75-81, further comprising a hydrophilic solubilizer.
83. The pharmaceutical composition of claim 82, wherein the hydrophilic solubilizer is a hydrophilic polymer.
84. The pharmaceutical composition of claim 83, wherein the hydrophilic polymer is a polyethylene glycol.
85. The pharmaceutical composition of any one of claims 75-84, further comprising a surfactant.
86. The pharmaceutical composition of claim 85, wherein the surfactant is a polyoxyethylene stearate.
87. The pharmaceutical composition of any one of claims 75-86, further comprising a pharmaceutically acceptable excipient.

88. The pharmaceutical composition of any one of claims 75-81, further comprising polyethylene glycol, caprylic/capric glycerides, and tocopheryl polyethylene glycol succinate.
89. The pharmaceutical composition of any one of claims 75-81, further comprising propylene glycol, polyethylene glycol, and tocopheryl polyethylene glycol succinate.
90. The pharmaceutical composition of any one of claims 75-89, wherein the pharmaceutical composition is formulated as a capsule.
91. The pharmaceutical composition of any one of claims 75-89, wherein the pharmaceutical composition is formulated as a tablet.
92. The pharmaceutical composition of any one of claims 75-89, wherein the pharmaceutical composition is formulated as a powder for reconstitution.
93. The pharmaceutical composition of claim 92, wherein the powder for reconstitution is reconstituted with a liquid carrier to form an oral suspension.
94. The pharmaceutical composition of any one of claims 75-89, wherein the pharmaceutical composition is formulated as a solution.
95. A pharmaceutical composition comprising:
- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
 - (ii) cefditoren pivoxil.
96. A method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:
- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
 - (ii) cefditoren pivoxil.
97. A method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:
- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
 - (ii) cefditoren pivoxil.
98. The method of claim 97, wherein the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.

99. The method of claim 97 or 98, wherein the method comprises administering to the subject in need thereof:

- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

- (ii) cefditoren pivoxil.



100. The method of claim 99, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



101. The method of claim 99, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



102. The method of any one of claims 99-101, wherein the compound of formula (I) or (II) and cefditoren pivoxil are formulated for oral administration.
103. The method of any one of claims 99-102, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefditoren pivoxil are administered sequentially.
104. The method of any one of claims 99-103, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefditoren pivoxil are administered concurrently.
105. A method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:
 - (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof;




wherein:


each R¹ and R² is independently hydrogen, C₁-C₆ alkyl, or -CO(C₁-C₆ alkyl); each alkyl being optionally substituted with halogen, -OH, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, or -COO(C₁-C₆ alkyl);

or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and



- (ii) cefditoren pivoxil.
106. The method of claim 105, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



107. The method of claim 105, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



108. The method of any one of claims 105-107, wherein the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.
109. The method of any one of claims 105-108, wherein the compound of formula (I) or (II) and cefditoren pivoxil are formulated for oral administration.
110. The method of any one of claims 105-109, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered sequentially.
111. The method of any one of claims 105-109, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefditoren pivoxil are administered concurrently.
112. A pharmaceutical composition comprising:
- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);


or two R² are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, -OH, C₁-C₆ alkyl, -O(C₁-C₆ alkyl), -NH₂, -NHC₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂, -COOH, -COO(C₁-C₆ alkyl), -(C₁-C₆ alkylene)COOH, or -(C₁-C₆ alkylene)COO(C₁-C₆ alkyl); and

(ii) cefpodoxime proxetil.

113. The pharmaceutical composition of claim 112, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:





114. The pharmaceutical composition of claim 112, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



115. The pharmaceutical composition of any one of claims 112-114, wherein the pharmaceutical composition is formulated for oral administration.
116. The pharmaceutical composition of any one of claims 112-115, wherein the pharmaceutical composition is formulated as an emulsion.
117. The pharmaceutical composition of any one of claims 112-115, wherein the pharmaceutical composition is formulated as a microemulsion.
118. The pharmaceutical composition of any one of claims 112-115, wherein the pharmaceutical composition is formulated as a Self-Emulsifying Drug Delivery System (SEDSS).
119. The pharmaceutical composition of any one of claims 112-118, further comprising a hydrophilic solubilizer.
120. The pharmaceutical composition of claim 119, wherein the hydrophilic solubilizer is a hydrophilic polymer.
121. The pharmaceutical composition of claim 120, wherein the hydrophilic polymer is a polyethylene glycol.

122. The pharmaceutical composition of any one of claims 112-121, further comprising a surfactant.
123. The pharmaceutical composition of claim 122, wherein the surfactant is a polyoxyethylene stearate.
124. The pharmaceutical composition of any one of claims 112-123, further comprising a pharmaceutically acceptable excipient.
125. The pharmaceutical composition of any one of claims 112-118, further comprising polyethylene glycol, caprylic/capric glycerides, and tocopheryl polyethylene glycol succinate.
126. The pharmaceutical composition of any one of claims 112-118, further comprising propylene glycol, polyethylene glycol, and tocopheryl polyethylene glycol succinate.
127. The pharmaceutical composition of any one of claims 112-126, wherein the pharmaceutical composition is formulated as a capsule.
128. The pharmaceutical composition of any one of claims 112-126, wherein the pharmaceutical composition is formulated as a tablet.
129. The pharmaceutical composition of any one of claims 112-126, wherein the pharmaceutical composition is formulated as a powder for reconstitution.
130. The pharmaceutical composition of claim 129, wherein the powder for reconstitution is reconstituted with a liquid carrier to form an oral suspension.
131. The pharmaceutical composition of any one of claims 112-126, wherein the pharmaceutical composition is formulated as a solution.
132. A pharmaceutical composition comprising:
 - (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
 - (ii) cefpodoxime proxetil.
133. A method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:
 - (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid, or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and
 - (ii) cefpodoxime proxetil.
134. A method of treating a bacterial infection in a subject in need thereof, the method comprising delivering to the subject:


- (i) (R)-2-hydroxy-3-propionamido-3,4-dihydro-2H-benzo[e][1,2]oxaborinine-8-carboxylic acid or (R)-3-(2-borono-2-propionamidoethyl)-2-hydroxybenzoic acid; and
- (ii) cefpodoxime proxetil.
135. The method of claim 134, wherein the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.
136. The method of claim 134 or 135, wherein the method comprises administering to the subject in need thereof:
- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:

- each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);
- or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and
- (ii) cefpodoxime proxetil.



137. The method of claim 136, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



138. The method of claim 136, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



139. The method of any one of claims 134-138, wherein the compound of formula (I) or (II) and cefpodoxime proxetil are formulated for oral administration.
140. The method of any one of claims 134-139, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefpodoxime proxetil are administered sequentially.
141. The method of any one of claims 134-139, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof; and cefpodoxime proxetil are administered concurrently.
142. A method of treating a bacterial infection in a subject in need thereof, the method comprising administering to the subject:
- (i) a compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof:




wherein:


- each R^1 and R^2 is independently hydrogen, C_1 - C_6 alkyl, or $-CO(C_1$ - C_6 alkyl); each alkyl being optionally substituted with halogen, $-OH$, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, or $-COO(C_1$ - C_6 alkyl);
- or two R^2 are taken together to form a 4- to 8-membered heterocycloalkyl optionally substituted with oxo, halogen, $-OH$, C_1 - C_6 alkyl, $-O(C_1$ - C_6 alkyl), $-NH_2$, $-NHC_1$ - C_6 alkyl, $-N(C_1$ - C_6 alkyl) $_2$, $-COOH$, $-COO(C_1$ - C_6 alkyl), $-(C_1$ - C_6 alkylene) $COOH$, or $-(C_1$ - C_6 alkylene) $COO(C_1$ - C_6 alkyl); and

(ii) cefpodoxime proxetil.

143. The method of claim 142, wherein the compound of formula (I) or (II), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



144. The method of claim 143, wherein the compound of formula (I), or a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof is:



145. The method of any one of claims 142-144, wherein the bacterial infection is caused by carbapenem-resistant Enterobacteriaceae (CRE) or extended-spectrum beta-lactamase (ESBL) producing gram-negative bacteria.
146. The method of any one of claims 142-145, wherein the compound of formula (I) or (II) and cefpodoxime proxetil are formulated for oral administration.
147. The method of any one of claims 142-146, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered sequentially.
148. The method of any one of claims 142-146, wherein the compound of formula (I) or (II); a pharmaceutically acceptable salt, a solvate, or a pharmaceutically acceptable salt and solvate thereof and cefpodoxime proxetil are administered concurrently.