12. The composition of any one of claims 1 to 11, wherein the mineral is selected from the group consisting of sodium, magnesium, potassium, calcium, zinc, manganese, copper, molybdenum, selenium and chromium or combination thereof.

13. The composition of any one of claims 1 to 12, wherein the vitamin is selected from the group consisting of thiamine, riboflavin, niacin, vitamin C, vitamin D, vitamin E, vitamin B, vitamin B12, pantothenic acid, biotin and folate or combination thereof.

14. The composition of any one of claims 1 to 13, wherein the second active ingredient further comprises taurine.

15. The composition of any one of claims 1 to 14, wherein the second active ingredient further comprises a component selected from, the group consisting of carnitine, glucosamine, taurine, sodium citrate, flaxseed oil and micellular casein or combination thereof.

16. The composition of any one of claims 1 to 15, wherein the second active ingredient consists essentially of sodium chloride, magnesium gluconate, potassium chloride, selenium, zinc, sulphate, manganese sulphate, copper sulphate, chromium picolinate, sodium molybdate, potassium iodide, calcium lactate, sodium bicarbonate, L-glutamine, L-lysine, L-arginine, L-carnitine, vitamin C, vitamin D, vitamin B12, niacin, panthothenic acid, riboflavin, pyridoxine, thiamine, folate, biotin, taurine, dextrose, inulin, fructose, dextrose and glycine.

17. The composition of any one of claims 1 to 16, wherein the first active ingredient and the second active ingredient are present at a ratio of 5:1, 4:1, 3:1, 2:1, 1:1, 1:2, 1:3, 1:4 or 1:5.

18. A composition comprising a first active ingredient and a second active ingredient, wherein the first active ingredient comprises an antiseptic and the second active ingredient comprises an admixture of the carbohydrates dextrose, fructose and inulin, the amino acids glycine, arginine, glutamine and lysine, at least one vitamin, and at least one mineral, for use in enhancing wound healing and/or reducing pain and inflammation in a subject~~A method for enhancing wound healing and/or pain and inflammation, said method comprising the steps of administering to the subject a therapeutically effective amount of a composition comprising a first active ingredient~~

~~and a second active ingredient, wherein the first active ingredient comprises an antiseptic and the second active ingredient comprises an admixture of the carbohydrates dextrose, fructose and inulin, the amino acids glycine, arginine, glutamine and lysine, at least one vitamin, and at least one mineral~~.

19. The ~~method~~ composition for use of claim 18, wherein said administration is topical administration.

20. A composition comprising a first active ingredient and a second active ingredient, wherein the first active ingredient comprises an antiseptic and the second active ingredient comprises an admixture of the carbohydrates dextrose, fructose and inulin, the amino acids glycine, arginine, glutamine and lysine, at least one vitamin, and at least one mineral, for use in ~~A method for~~ treating a wound in a subject~~, comprising the steps of applying to the wound a therapeutically effective amount of a first active ingredient and a second active ingredient, wherein the first active ingredient comprises an antiseptic, and the second active ingredient comprises an admixture of the carbohydrates dextrose, fructose and inulin, the amino acids glycine, arginine, glutamine and lysine, at least one vitamin, and at least one mineral~~.

21. The composition for use ~~method~~ of any one of claims 18 to 20, wherein the second active ingredient comprises dextrose in an amount of 800 mg to 1200mg, fructose in an amount of from 200mg to 300mg, and inulin in amount of 400 to 600 mg, per 3.5 grams of the second active ingredient.

22. The composition for use ~~method~~ of any one of claims 18 to 21, wherein the second active ingredient comprises glutamine in an amount of 100 mg to 200 mg, lysine in an amount of 20 mg to 100 mg, arginine in an amount of 10 mg to 30 mg, per 3.5 grams of second active ingredient, and wherein the ratio of glycine to arginine is in the range of from 19:1 to 18:1 by weight.

23. The composition for use ~~method~~ according to any one of claims 18 to 22, wherein said antiseptic is a bis-biguanide compound.

24. The composition for use ~~method~~ according to claim 23, wherein said bis-biguanide compound is chlorhexidine or a chlorhexidine salt.

25. The composition for use ~~method~~ according to claim 24, wherein the chlorhexidine is chlorhexidine-digluconate.

26. The composition for use ~~method~~ according to any one of claims 18 to 25, wherein the first active ingredient is a 0.01% to 3% bis-biguanide compound.

27. The composition for use ~~method~~ according to any one of claim 18 to 22 wherein said antiseptic is iodine.

28. The composition for use ~~method~~ according to any one of claim 18 to 22 wherein said antiseptic is an antibacterial dye.

29. The composition for use ~~method~~ according to any one of claims 18 to 22 wherein said antiseptic is a halogenated phenol derivative.

30. The composition for use ~~method~~ according to any one of claims 18 to 22 wherein said antiseptic is a quinolone derivative.

31. The composition for use ~~method~~ according to any one of claims 18 to 30, wherein the mineral is selected from the group consisting of sodium, magnesium, potassium, calcium, zinc, manganese, copper, molybdenum, selenium and chromium or combination thereof.

32. The composition for use ~~method~~ according to any one of claims 18 to 31, wherein the vitamin is selected from the group consisting of thiamine, riboflavin, niacin, vitamin C, vitamin D, vitamin E, vitamin B, vitamin B12, pantothenic acid, biotin and folate or combination thereof.

33. The composition for use ~~method~~ according to any one of claims 18 to 32, wherein the second active ingredient further comprises a component selected from, the group consisting of carnitine, glucosamine, taurine, sodium citrate, flaxseed oil and micellular casein or combination thereof.

34. The composition for use ~~method~~ according to any one of claims 18 to 33, wherein the second active ingredient consists essentially of sodium chloride, magnesium gluconate, potassium chloride, selenium, zinc, sulphate, manganese sulphate, copper sulphate, chromium picolinate, sodium molybdate, potassium iodide, calcium lactate, sodium bicarbonate, L-glycine, L-glutamine, L-lysine, L-arginine, L-carnitine, vitamin C, vitamin D, vitamin B12, niacin, panthothenic acid, riboflavin, pyridoxine, thiamine, folate, biotin, taurine, dextrose, inulin, fructose, dextrose and glycine.

35. The composition for use ~~method~~ according to any one of claims 18 to 35, wherein the first active ingredient and the second active ingredient are present at a ratio of 5:1, 4:1, 3:1, 2:1, 1:1, 1:2, 1:3, 1:4 or 1:5, preferably 1:1.

36. The composition for use ~~method~~ of any one of claims 18 to 36, wherein the wound is the result of infection or inflammation.

37. The composition for use ~~method~~ of any one of claims 18 to 37, wherein the wound is selected from the group consisting of surgical wound, traumatic mouth ulcer, non-healing area of the mouth, inflammatory lesion and apthous ulcer.

38. A base composition for combining with a first active ingredient comprising an antiseptic to form a composition for enhancing wound healing and/or reduce pain and inflammation in a subject, the base composition comprising a second active ingredient comprising an admixture of the carbohydrates dextrose, fructose and inulin, the amino acids glycine, arginine, glutamine and lysine, at least one vitamin, and at least one mineral.

39. The base composition of claim 38, wherein the second active ingredient comprises dextrose in an amount of 800 mg to 1200mg, fructose in an amount of from 200mg to 300mg, and inulin in amount of 400 to 600 mg, per 3.5 grams of the second active ingredient.

40. The base composition of claim 38 or 39, wherein the second active ingredient comprises glutamine in an amount of 100 mg to 200 mg, lysine in an amount of 20 mg to 100 mg, arginine in an amount of 10 mg to 30 mg, per 3.5 grams of second