

DESCRIPTION

SATURATED-RING-FUSED DIHYDROPYRIMIDINONE OR
DIHYDROTRIAZINONE COMPOUNDS AND PHARMACEUTICAL USE THEREOF

5

TECHNICAL FIELD

[0001]

The present invention relates to saturated-ring-fused
dihydropyrimidinone or dihydrotriazinone compounds, or
10 pharmaceutically acceptable salts thereof, having ROR γ
antagonist activity, pharmaceutical compositions comprising
the same, and pharmaceutical use thereof.

BACKGROUND ART

15 [0002]

ROR γ (i.e., Retinoid-related Orphan Receptor gamma) is
a nuclear receptor which is important for the
differentiation and activation of Th17 cells. ROR γ t is
also known as a splicing variant of ROR γ (Non patent
20 literature 1). ROR γ and ROR γ t differ only in their N-
terminal domains and share the same ligand-binding domain
and DNA-binding domain. It is reported that ROR γ is
expressed in other tissues besides Th17 cells (Non Patent
Literature 1).

25 [0003]

Inhibition of ROR γ can inhibit the differentiation and activation of Th17 cells. IL-17 produced in Th17 cells is involved in the induction of a variety of chemokines, cytokines, metalloproteases, and other inflammatory mediators and the migration of neutrophil, and therefore, inhibition of IL-17 may lead to inhibit such induction and migration (Non Patent Literatures 2 and 3). It is known that Th17 cells are involved in autoimmune diseases (such as rheumatoid arthritis, psoriasis, inflammatory bowel disease (such as Crohn's disease and ulcerative colitis), multiple sclerosis, systemic lupus erythematosus (SLE), Behcet's disease, sarcoidosis, Harada disease, ankylosing spondylitis, uveitis, polymyalgia rheumatica, type I diabetes, graft-versus-host disease, alopecia areata, and vitiligo), allergic diseases, dry eye, fibrosis (such as lung fibrosis and primary biliary cirrhosis), and cancers (such as malignant melanoma and prostate cancer).

ROR γ in adipose tissues is related to the regulation of adipogenesis and inhibition of ROR γ can ameliorate insulin resistance (Non Patent Literature 4). It is known that adipose tissues are involved in metabolic diseases (such as hepatic steatosis).

It is also known that IL-17 and Th17 cells are involved in ischemia, cardiomyopathy, hypertension, and periodontitis.

[0004]

For example, as for rheumatoid arthritis, it is reported that administration of anti-IL-17 antibody can ameliorate swelling and joint destruction associated with collagen-induced arthritis (Non Patent Literature 5). It is also reported that swelling and joint destruction associated with collagen-induced arthritis can be ameliorated in experiments using IL-17-deficient mice (Non Patent Literature 6).

As for psoriasis, it is reported that administration of anti-IL-17 antibody is effective in treating psoriasis in clinical trials (Non Patent Literature 7). Anti IL-17 antibodies have been placed on the market for use in psoriasis (Non Patent Literature 8).

As for inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, adoptive transfer of T cells derived from ROR γ -KO mice does not increase IL-17 in the mucosa in a colitis model induced by the adoptive transfer of T cells, thereby the onset of colitis can be inhibited (Non Patent Literature 9). It is also reported that an anti-IL-23 antibody, an antibody against IL-23 which activates Th17 cells, was effective in treating Crohn's disease in clinical trials (Non Patent Literature 20).

As for multiple sclerosis, the disease state of a

mouse experimental autoimmune encephalomyelitis model which is an animal model of multiple sclerosis can be inhibited in RORγ-KO mice (Non Patent Literature 10). It is also reported that an anti-IL-17A antibody can ameliorate MRI observation in relapsing remitting multiple sclerosis in clinical trials (Non Patent Literature 21).

As for systemic lupus erythematosus, it is reported that administration of anti-IL-17 antibody can inhibit onset of GBM nephritis model in RORγt-KO mice which is an animal model of glomerulonephritis (Non Patent Literature 11). Administration of anti-IL-17 antibody potentially inhibits nephritis associated with SLE as well (Non Patent Literature 12).

As for ankylosing spondylitis, it is reported that administration of anti-IL-17 antibody is effective in treating ankylosing spondylitis (Non Patent Literature 13).

As for uveitis, it is reported that administration of anti-IL-17 antibody is effective in treating uveitis associated with Behcet's disease, sarcoidosis, and Harada disease (Non Patent Literature 7).

As for polymyalgia rheumatica, efficacy of anti-IL-17 antibody is currently assessed in clinical trials for polymyalgia rheumatica.

As for type I diabetes, administration of anti-IL-17 antibody can inhibit progression of disease states in a NOD

mouse model which is a type I diabetes model (Non Patent Literature 14). Efficacy of anti-IL-17A antibody is currently assessed in clinical trials (Non Patent Literature 22).

5 As for graft-versus-host disease, it is reported that transfection of ROR γ -KO-mouse-derived cells can ameliorate survival rates and rejections in a host in a mouse transplant model (Non Patent Literature 19).

As for alopecia areata, efficacy of anti-IL-17A
10 antibody is currently assessed in clinical trials (Non Patent Literature 25).

As for vitiligo, increases of IL-17 and Th17 cells are recognized in patient sera and pathological tissues, respectively (Non Patent Literature 34).

15 As for allergic diseases such as asthma, attenuated eosinophilic pulmonary inflammation, the reduced number of CD4⁺ lymphocytes, and the decrease of Th2 cytokines/chemokines levels are exhibited in ROR γ -KO mice in an OVA-sensitized model, which then allergic reactions
20 can be inhibited (Non Patent Literature 15). Efficacy of anti-IL-17A antibody is currently assessed in clinical trials for atopic dermatitis (Non Patent Literature 23). Efficacy of anti-IL-23 antibody is currently assessed in clinical trials for asthma (Non Patent Literature 24).

25 As for dry eye, it is reported that Th17 cells

increase in an animal model of dry eye, and efficacy of anti-IL-17 antibody is currently assessed in clinical trials for dry eye patients (Non Patent Literature 16).

As for fibrosis, administration of anti-IL-17 antibody
5 can inhibit inflammation and fibrosis in lung and extend survival of animals in a bleomycin-induced lung fibrosis model which is an animal model of lung fibrosis (Non Patent Literature 17).

As for primary biliary cirrhosis, it is reported that
10 Th17 cells increase in the lesion area of patients with primary biliary cirrhosis, and efficacy of anti-IL-23 antibody is currently assessed in clinical trials (Non Patent Literature 18).

As for malignant melanoma, efficacy of anti-IL-17
15 antibody is currently assessed in clinical trials (Non Patent Literatures 26 and 27).

As for prostate cancer, it is recognized that anti-IL-17 antibody treatment decreased the formation of micro-invasive prostate cancer in Pten-null mice (Non Patent
20 Literature 28).

As for insulin resistance, the insulin resistance induced by feeding high-fat diets can be inhibited in RORγ KO mice (Non Patent Literature 4).

As for hepatic steatosis, it is recognized that anti-
25 IL-17 antibody ameliorated steatosis on pathological

tissues in an alcoholic liver-disease model (Non Patent Literature 29).

As for non-alcoholic fatty liver disease, it is recognized that anti-IL-17 antibody treatment improved
5 liver function, attenuated hepatic lipid accumulation, suppressed Kupffer cells activation, and decreased pro-inflammatory cytokines levels in a high fat diet-induced non-alcoholic fatty liver disease model (Non Patent Literature 30).

10 As for ischemia and cardiomyopathy, it is reported that IL-17A contributes to myocardial ischemia/reperfusion injury by regulating cardiomyocyte apoptosis and neutrophil infiltration. It is recognized that anti-IL-17A antibody treatment or IL-17A knockout reduced infarct size, improved
15 cardiac function, and thus, ameliorated ischemia/reperfusion injury (Non Patent Literature 31).

As for hypertension, it is reported that treatment with antibody against IL-17A or IL-17RA suppressed increased blood pressure by administration of angiotensin
20 II (Non Patent Literature 32).

As for periodontitis, increase of Th17 cells or IL-17 was recognized in an experimental periodontitis model. It is reported that treatment with ROR γ antagonist, GSK805, or anti-IL-17A antibody diminished bone loss in the model (Non
25 Patent Literature 33).

[0005]

On the basis of these findings, ROR γ antagonists are deemed to be beneficial for preventing or treating autoimmune diseases, allergic diseases, dry eye, fibrosis, cancers (such as malignant melanoma and prostate cancer), metabolic disease, ischemia, cardiomyopathy, hypertension, and periodontal disease.

[0006]

[Non Patent Literature 1] JETTEN, "Retinoid-related orphan receptors (RORs): critical roles in development, immunity, circadian rhythm, and cellular metabolism", Nucl. Recept. Signal., 7: e003 (2009).

[Non Patent Literature 2] KOENDERS, et al., "Potential new targets in arthritis therapy: interleukin (IL)-17 and its relation to tumour necrosis factor and IL-1 in experimental arthritis", Ann. Rheum. Dis., 65: iii29-33 (2006).

[Non Patent Literature 3] SCHMIDT-WEBER, et al., "Th17 cells in the big picture of immunology", J. Allergy Clin. Immunol., 120: 247-54 (2007).

[Non Patent Literature 4] MEISSBURGER, et al., "Adipogenesis and insulin sensitivity in obesity are regulated by retinoid-related orphan receptor gamma", EMBO Mol. Med., 3: 637-51 (2011).

[Non Patent Literature 5] KELCHTERMANS et al.,

"Effector mechanisms of interleukin-17 in collagen-induced arthritis in the absence of interferon- γ and counteraction by interferon- γ ", *Arthritis Res. Ther.*, 11(4): R122 (2009).

[Non Patent Literature 6] NAKAE, et al., "Suppression
5 of Immune Induction of Collagen-Induced Arthritis in IL-17-Deficient Mice", *J. Immunol.*, 171: 6173-6177 (2003).

[Non Patent Literature 7] HUEBER et al., "Effects of AIN457, a Fully Human Antibody to Interleukin-17A, on Psoriasis, Rheumatoid Arthritis, and Uveitis", *Sci. Transl.*
10 *Med.*, 2(52): 52ra72 (2010).

[Non Patent Literature 8] SANFORD et al., "Secukinumab: first global approval", *Drugs*, 75(3): 329-338 (2015).

[Non Patent Literature 9] LEPPKES, et al., "ROR γ -Expressing Th17 Cells Induce Murine Chronic Intestinal
15 Inflammation via Redundant Effects of IL-17A and IL-17F", *Gastroenterology*, 136(1): 257-267 (2009).

[Non Patent Literature 10] IVANOV et al., "The Orphan Nuclear Receptor ROR γ t Directs the Differentiation Program of Proinflammatory IL-17+ T Helper Cells", *Cell*,
20 126(6): 1121-1133 (2006).

[Non Patent Literature 11] STEINMETZ et al., "The Th17-Defining Transcription Factor ROR γ t Promotes Glomerulonephritis", *J. Am. Soc. Nephrol.*, 22(3): 472-483 (2011).

25 [Non Patent Literature 12] CRISPIN et al.,

"Interleukin-17-producing T cells in lupus", Curr. Opin. Rheumatol., 22(5): 499-503 (2010).

[Non Patent Literature 13] BAETEN et al., "Anti-interleukin-17A monoclonal antibody secukinumab in
5 treatment of ankylosing spondylitis: a randomised, double-blind, placebo-controlled trial", Lancet, 382(9906): 1705-1713 (2013).

[Non Patent Literature 14] EMAMAULLEE et al., "Inhibition of Th17 Cells Regulates Autoimmune Diabetes in
10 NOD Mice", Diabetes, 58: 1302-1311 (2009).

[Non Patent Literature 15] TILLEY, et al., "Retinoid-Related Orphan Receptor γ Controls Immunoglobulin
Production and Th1/Th2 Cytokine Balance in the Adaptive
Immune Response to Allergen", J. Immunol., 178: 3208-3218
15 (2007).

[Non Patent Literature 16] U.S. NATIONAL INSTITUTES
OF HEALTH, "The Effects of a Single Intravenous
Administration of Secukinumab (AIN457) or Canakinumab
(ACZ885) in Dry Eye Patients", ClinicalTrials.gov
20 information for Clinical Trials Identifier NCT01250171
(December 4, 2012).

[Non Patent Literature 17] MI et al., "Blocking IL-17A Promotes the Resolution of Pulmonary Inflammation and
Fibrosis Via TGF- β 1-Dependent and -Independent Mechanisms",
25 J. Immunol., 187: 3003-3014 (2011).

[Non Patent Literature 18] U.S. NATIONAL INSTITUTES
OF HEALTH, "A Study of Efficacy and Safety of Ustekinumab
in Patients With Primary Biliary Cirrhosis (PBC) Who Had an
Inadequate Response to Ursodeoxycholic Acid",
5 ClinicalTrials.gov information for Clinical Trials
Identifier NCT01389973 (April 2, 2015).

[Non Patent Literature 19] FULTON et al.,
"Attenuation of Acute Graft-versus-Host Disease in the
Absence of the Transcription Factor ROR γ t", J. Immunol.,
10 189(4): 1765-1772 (2012).

[Non Patent Literature 20] Brian G Feagan et al.,
"Induction therapy with the selective interleukin-23
inhibitor risankizumab in patients with moderate-to-severe
Crohn's disease: a randomised, double-blind, placebo-
15 controlled phase 2 study", The Lancet, 389(10080): 1699-
1709 (2017).

[Non Patent Literature 21] Eva Havrdova et al.,
"Activity of secukinumab, an anti-IL-17A antibody, on brain
lesions in RRMS: results from a randomized, proof-of-
20 concept study", J. Neurol., 263(7): 1287-1295 (2016).

[Non Patent Literature 22] U.S. NATIONAL INSTITUTES
OF HEALTH, "Study of Secukinumab in Patients With Newly-
diagnosed Type 1 Diabetes Mellitus", ClinicalTrials.gov
information for Clinical Trials Identifier NCT02044848.

25 [Non Patent Literature 23] U.S. NATIONAL INSTITUTES

OF HEALTH, "Secukinumab for Treatment of Atopic Dermatitis",
ClinicalTrials.gov information for Clinical Trials
Identifier NCT02594098.

[Non Patent Literature 24] U.S. NATIONAL INSTITUTES
5 OF HEALTH, "Efficacy and Safety of BI 655066/ABBV-066
(Risankizumab) in Patients With Severe Persistent Asthma",
ClinicalTrials.gov information for Clinical Trials
Identifier NCT02443298.

[Non Patent Literature 25] U.S. NATIONAL INSTITUTES
10 OF HEALTH, "A Study of Secukinumab for the Treatment of
Alopecia Areata", ClinicalTrials.gov information for
Clinical Trials Identifier NCT02599129.

[Non Patent Literature 26] U.S. NATIONAL INSTITUTES
OF HEALTH, "An Open-Label, Proof-of-Concept Study of
15 Ixekizumab in the Treatment of Pyoderma Gangrenosum",
ClinicalTrials.gov information for Clinical Trials
Identifier NCT03137160.

[Non Patent Literature 27] U.S. NATIONAL INSTITUTES
OF HEALTH, "Single-arm Study to Assess a Potential Effect
20 of Anti-IL-17 (Secukinumab) in the Treatment of Pyoderma
Gangrenosum", ClinicalTrials.gov information for Clinical
Trials Identifier NCT02733094.

[Non Patent Literature 28] Q Zhang et al.,
"Targeting Th17-IL-17 Pathway in Prevention of Micro-
25 Invasive Prostate Cancer in a Mouse Model", Prostate,

77(8): 888-899 (2017).

[Non Patent Literature 29] W Shi et al., "Anti-IL-17 Antibody Improves Hepatic Steatosis by Suppressing Interleukin-17-Related Fatty Acid Synthesis and Metabolism", Clin. Dev. Immunol., Volume 2013, Article ID 253046 (2013).

[Non Patent Literature 30] R Xu et al., "Neutralization of interleukin-17 attenuates high fat diet-induced non-alcoholic fatty liver disease in mice", Acta Biochim. Biophys. Sin. (Shanghai), 45(9): 726-733 (2013).

[Non Patent Literature 31] Y.H. Lial et al., "Interleukin-17A Contributes to Myocardial Ischemia/Reperfusion Injury by Regulating Cardiomyocyte Apoptosis and Neutrophil Infiltration", J. Am. Coll. Cardiol. 59(4): 420-429 (2012).

[Non Patent Literature 32] M.A. Saleh et al., "Inhibition of Interleukin 17-A but not Interleukin-17F Signaling Lowers Blood Pressure and Reduces End-organ Inflammation in Angiotensin II-induced Hypertension", JACC Basic Transl. Sci. 1(7): 606-616 (2016).

[Non Patent Literature 33] N. Dutzan et al., "A dysbiotic microbiome triggers TH17 cells to mediate oral mucosal immunopathology in mice and humans", Sci. Transl. Med. 10(463): eaat0797 (2018).

[Non Patent Literature 34] R Speeckaert et al., "The many faces of interleukin-17 in inflammatory skin

diseases", Br. J. Dermatol. 175(5): 892-901 (2016).

SUMMARY OF INVENTION


[0007]

5 The present invention provides saturated-ring-fused dihydropyrimidinone or dihydrotriazinone compounds, or pharmaceutically acceptable salts thereof, having ROR_Y antagonist activity, pharmaceutical compositions comprising the same, and their medical use. One aspect of the present
10 invention includes the following illustrative embodiments.

[0008]

[Item 1]

A compound of Formula [I]:



15 wherein R¹ is

- (1) C₁₋₈ alkyl,
- (2) halo-C₁₋₈ alkyl,

(3) C₃₋₈ cycloalkyl optionally substituted with the same or different 1 to 3 substituents selected from Group A¹, or

(4) C₃₋₈ cycloalkyl-C₁₋₄ alkyl wherein the C₃₋₈ cycloalkyl moiety may be optionally substituted with the same or
5 different 1 to 3 substituents selected from Group A¹,

Group A¹ is

- (1) halogen,
- (2) C₁₋₄ alkyl, and
- (3) halo-C₁₋₄ alkyl,

10 X¹ is

- (1) a bond, or
- (2) -O-,

R² is

- (1) hydrogen, or
- 15 (2) halogen,

R³ is

- (1) hydrogen, or
- (2) -Y³-COO-R³⁰,

Y³ is

- 20 (1) C₁₋₈ alkylene,
- (2) C₃₋₈ cycloalkylene,
- (3) bridged C₅₋₈ cycloalkylene, or
- (4) C₆₋₁₄ arylene,

R³⁰ is

- 25 (1) hydrogen, or

(2) C₁₋₄ alkyl,

X² is

(1) =C(R⁴)-, or

(2) =N-,

5 R⁴ is

(1) hydrogen, or

(2) C₁₋₄ alkyl,

X³ is

(1) -C(R⁵)(R⁶)-,

10 X⁴ is

(1) a bond, or

(2) -C(R⁷)(R⁸)-,

X⁵ is

(1) -C(R⁹)(R¹⁰)-,

15 (2) -N(R¹¹)-, or

(3) -O-,

R⁵ and R⁶ are each independently

(1) hydrogen,

(2) C₁₋₄ alkyl,

20 (3) halo-C₁₋₄ alkyl,

(4) cyano-C₁₋₄ alkyl, or

(5) C₁₋₄ alkyl substituted with one substituent selected from the group consisting of -O-R⁵¹, -CO-R⁶¹, -COO-R⁵², -N(R⁷¹)(R⁷²), -CO-N(R⁷³)(R⁷⁴), -N(R⁷⁵)-CO-R⁶², -N(R⁷⁶)-COO-R⁵³,

25 and -O-S(O)₂-R⁶³,

R^7 , R^8 , R^9 , and R^{10} are each independently

- (1) hydrogen,
- (2) halogen,
- (3) cyano,
- 5 (4) hydroxy,
- (5) C_{1-4} alkyl,
- (6) halo- C_{1-4} alkyl,
- (7) cyano- C_{1-4} alkyl,
- (8) C_{1-4} alkoxy, or
- 10 (9) C_{1-4} alkyl substituted with one substituent selected from the group consisting of $-O-R^{51}$, $-CO-R^{61}$, $-COO-R^{52}$, $-N(R^{71})(R^{72})$, $-CO-N(R^{73})(R^{74})$, $-N(R^{75})-CO-R^{62}$, $-N(R^{76})-COO-R^{53}$, and $-O-S(O)_2-R^{63}$,

R^{51} , R^{52} , and R^{53} are each independently

- 15 (1) hydrogen,
- (2) C_{1-4} alkyl, or
- (3) C_{6-14} aryl- C_{1-4} alkyl,

R^{61} , R^{62} , and R^{63} are each independently

- (1) C_{1-4} alkyl,
- 20 R^{71} , R^{72} , R^{73} , R^{74} , R^{75} , and R^{76} are each independently

(1) hydrogen, or

(2) C_{1-4} alkyl,

R^{11} is

- (1) $-CO-R^{111}$, or
- 25 (2) $-COO-R^{112}$,

R^{111} is

(1) C_{1-4} alkyl,


R^{112} is

(1) C_{1-4} alkyl, or a pharmaceutically acceptable salt thereof.

[0009]

[Item 2]

The compound according to Item 1, having a structure of Formula [II]:



10

wherein each variable is defined as defined in Item 1, or a pharmaceutically acceptable salt thereof.

[0010]

[Item 3]

15 The compound according to Item 1 or 2, wherein X^2 is =N-, or a pharmaceutically acceptable salt thereof.

[0011]

[Item 4]

The compound according to Item 1 or 2, wherein X^2 is $=C(R^4)-$ and R^4 is hydrogen, or a pharmaceutically acceptable salt thereof.

5 [0012]

[Item 5]

The compound according to any one of Items 1 to 4, wherein R^3 is hydrogen, or a pharmaceutically acceptable salt thereof.

10 [0013]

[Item 6]

The compound according to any one of Items 1 to 4, wherein R^3 is $-Y^3-COO-R^{30}$,

Y^3 is

- 15 (1) C_{1-8} alkylene,
(2) C_{3-8} cycloalkylene, or
(3) bridged C_{5-8} cycloalkylene,

R^{30} is hydrogen or C_{1-4} alkyl, or a pharmaceutically acceptable salt thereof.

20 [0014]

[Item 7]

The compound according to any one of Items 1 to 6, wherein R^2 is halogen, or a pharmaceutically acceptable salt thereof.

25 [0015]

[Item 8]

The compound according to any one of Items 1 to 7, wherein R^1 is C_{1-8} alkyl and X^1 is a bond, or a pharmaceutically acceptable salt thereof.

5 [0016]

[Item 9]

The compound according to any one of Items 1 to 8, wherein R^5 and R^6 are each independently hydrogen or C_{1-4} alkyl, or a pharmaceutically acceptable salt thereof.

10 [0017]

[Item 10]

The compound according to any one of Items 1 to 9, wherein X^4 is a bond or $-C(R^7)(R^8)-$ and both of R^7 and R^8 are hydrogen, or a pharmaceutically acceptable salt thereof.

15 [0018]









[Item 11]









The compound according to any one of Items 1 to 10, wherein X^5 is $-C(R^9)(R^{10})-$ or $-O-$ and both of R^9 and R^{10} are hydrogen, or a pharmaceutically acceptable salt thereof.

20 [0019]

[Item 12]

The compound according to Item 1, selected from the following compound group:





or a pharmaceutically acceptable salt thereof.

[0020]

[Item 13]

A pharmaceutical composition comprising a compound according to any one of Items 1 to 12 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

[0021]

[Item 14]

An ROR γ antagonist comprising a compound according to any one of Items 1 to 12 or a pharmaceutically acceptable salt thereof.

[0022]

[Item 15]

A therapeutic or preventive agent for a disease selected from the group consisting of autoimmune diseases, allergic diseases, dry eye, fibrosis, cancers, metabolic disease, ischemia, cardiomyopathy, hypertension, and periodontal disease, comprising a compound according to any one of Items 1 to 12 or a pharmaceutically acceptable salt thereof.

[0023]

[Item 16]

A method of antagonizing ROR γ , comprising administering a therapeutically effective amount of a compound according to any one of Items 1 to 12 or a

pharmaceutically acceptable salt thereof to a mammal.

[0024]

[Item 17]

A method of treating or preventing a disease selected
5 from the group consisting of autoimmune diseases, allergic
diseases, dry eye, fibrosis, cancers, metabolic disease,
ischemia, cardiomyopathy, hypertension, and periodontal
disease, comprising administering a therapeutically
effective amount of a compound according to any one of
10 Items 1 to 12 or a pharmaceutically acceptable salt thereof
to a mammal.

[0025]

[Item 18]

Use of a compound according to any one of Items 1 to
15 12 or a pharmaceutically acceptable salt thereof in the
manufacture of an ROR γ antagonist.

[0026]

[Item 19]

Use of a compound according to any one of Items 1 to
20 12 or a pharmaceutically acceptable salt thereof in the
manufacture of a therapeutic or preventive agent for a
disease selected from the group consisting of autoimmune
diseases, allergic diseases, dry eye, fibrosis, cancers,
metabolic disease, ischemia, cardiomyopathy, hypertension,
25 and periodontal disease.

[0027]

[Item 20]

A compound according to any one of Items 1 to 12 or a
pharmaceutically acceptable salt thereof for use in an RORy
5 antagonist.

[0028]

[Item 21]

A compound according to any one of Items 1 to 12 or a
pharmaceutically acceptable salt thereof for use in
10 treating or preventing a disease selected from the group
consisting of autoimmune diseases, allergic diseases, dry
eye, fibrosis, cancers, metabolic disease, ischemia,
cardiomyopathy, hypertension, and periodontal disease.

[0029]

15 [Item 22]

A commercial package comprising a pharmaceutical
composition according to Item 13 and a package insert
concerning the pharmaceutical composition describing that
the pharmaceutical composition can be used for treating or
20 preventing a disease selected from the group consisting of
autoimmune diseases, allergic diseases, dry eye, fibrosis,
cancers, metabolic disease, ischemia, cardiomyopathy,
hypertension, and periodontal disease.

[0030]

25 [Item 23]

A kit comprising a pharmaceutical composition according to Item 13 and a package insert concerning the pharmaceutical composition describing that the pharmaceutical composition can be used for treating or preventing a disease selected from the group consisting of autoimmune diseases, allergic diseases, dry eye, fibrosis, cancers, metabolic disease, ischemia, cardiomyopathy, hypertension, and periodontal disease.

10 DESCRIPTION OF EMBODIMENTS

[0031]

Definitions of terms used herein are shown as follows.

[0032]

A wavy line as follows:



15

in a partial structure shows a binding site.

[0033]

The term "halogen" includes fluoro, chloro, bromo, and iodo. A preferable "halogen" is fluoro, chloro, or bromo.

20 [0034]

The term "C₁₋₄ alkyl" means a straight- or branched-chain saturated hydrocarbon group with 1 to 4 carbon atoms. The "C₁₋₄ alkyl" group includes methyl, ethyl, n-propyl,

isopropyl, n-butyl, isobutyl, sec-butyl, and tert-butyl.


[0035]

The term "C₁₋₈ alkyl" means a straight- or branched-chain saturated hydrocarbon group with 1 to 8 carbon atoms.

5 The "C₁₋₈ alkyl" group includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, 1,1-dimethylpropyl, 1-ethylpropyl, n-hexyl, isohexyl, 3,3-dimethylbutyl, 2,2-dimethylbutyl, 1,1-dimethylbutyl, 2-ethylbutyl, n-heptyl,
10 5-methylhexyl, 4-methylhexyl, 4,4-dimethylpentyl, 3,3-dimethylpentyl, 3,4-dimethylpentyl, 2,3-dimethylpentyl, 3-ethylpentyl, 2-ethylpentyl, heptan-4-yl, n-octyl, 6-methylheptyl, 5,5-dimethylhexyl, 4,5-dimethylhexyl, 4-ethylhexyl, 3-ethylhexyl, 2-propylpentyl, and octan-4-yl.

15 [0036]

The term "C₁₋₈ alkylene" means a divalent group derived from straight- or branched-chain saturated hydrocarbon with 1 to 8 carbon atoms. The "C₁₋₈ alkylene" group includes, for example, the following groups:



[0037]

The term "halo-C₁₋₄ alkyl" means the "C₁₋₄ alkyl" group substituted with 1 to 5 halogen atoms independently selected from the group of the term "halogen". The "halo-C₁₋₄ alkyl" group includes, for example, monofluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, 3-fluoropropyl, 3-chloropropyl, 1,1-difluoropropyl, 3,3,3-trifluoropropyl, 4-fluorobutyl, and 4,4,4-trifluorobutyl.

[0038]

The term "halo-C₁₋₈ alkyl" means the "C₁₋₈ alkyl" group substituted with 1 to 9 halogen atoms independently selected from the group of the term "halogen". The "halo-

C₁₋₈ alkyl" group includes, for example, monofluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, 3-fluoropropyl, 3-chloropropyl, 1,1-difluoropropyl, 3,3,3-trifluoropropyl, 4-fluorobutyl, 4,4,4-trifluorobutyl, 5-fluoropentyl, 5,5,5-trifluoropentyl, 4,4,5,5,5-pentafluoropentyl, 3,3,4,4,5,5,5-heptafluoropentyl, 6-fluorohexyl, 6,6,6-trifluorohexyl, 7-fluoroheptyl, 7,7,7-trifluoroheptyl, 8-fluorooctyl, 8,8,8-trifluorooctyl, and 7,7,8,8,8-pentafluorooctyl.

[0039]

The term "cyano-C₁₋₄ alkyl" means the "C₁₋₄ alkyl" group substituted with one cyano group. The "cyano-C₁₋₄ alkyl" group includes, for example, cyanomethyl, 1-cyanoethyl, 2-cyanoethyl, 2-cyanopropyl, 3-cyanopropyl, 4-cyanobutyl, and 2-cyano-2-methylpropyl.

[0040]

The term "C₁₋₄ alkoxy" means those which the "C₁₋₄ alkyl" group binds to an oxygen atom and the group binds to another group via the oxygen atom. The "C₁₋₄ alkoxy" group includes methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, and tert-butoxy.

[0041]

The term "C₃₋₈ cycloalkyl" means a monocyclic saturated


hydrocarbon group with 3 to 8 carbon atoms. The "C₃₋₈ cycloalkyl" group includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

[0042]

5 The term "C₃₋₈ cycloalkyl-C₁₋₄ alkyl" means the "C₁₋₄ alkyl" group substituted with one cycloalkyl group selected from the group of the "C₃₋₈ cycloalkyl". The "C₃₋₈ cycloalkyl-C₁₋₄ alkyl" group includes, for example, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cycloheptylmethyl, cyclooctylmethyl, 2-cyclopropylethyl, 2-cyclobutylethyl, 2-cyclopentylethyl, 2-cyclohexylethyl, 2-cycloheptylethyl, 2-cyclooctylethyl, 1-cyclopropylethyl, 1-cyclobutylethyl, 1-cyclopentylethyl, 1-cyclohexylethyl, 1-cycloheptylethyl, 1-cyclooctylethyl, 3-cyclopropylpropyl, 3-cyclobutylpropyl, 3-cyclopentylpropyl, 3-cyclohexylpropyl, 3-cycloheptylpropyl, and 3-cyclooctylpropyl.


[0043]

20 The term "C₃₋₈ cycloalkylene" means a divalent group derived from a monocyclic saturated hydrocarbon group with 3 to 8 carbon atoms. The "C₃₋₈ cycloalkylene" group includes, for example, the following groups:



[0044]

The term "bridged C₅₋₈ cycloalkylene" means a divalent group derived from a bridged cyclic saturated hydrocarbon group with 5 to 8 carbon atoms. The "bridged C₅₋₈ cycloalkylene" group includes, for example, the following groups:



[0045]

The term "C₆₋₁₄ aryl" means an aromatic hydrocarbon group with 6 to 14 carbon atoms. The "C₆₋₁₄ aryl" group includes, for example, phenyl, naphthyl, anthryl, indenyl, azulenyl, fluorenyl, phenanthryl, and pentalenyl.


[0046]

The term "C₆₋₁₄ aryl-C₁₋₄ alkyl" means the "C₁₋₄ alkyl" group substituted with one aryl group selected from the group of the "C₆₋₁₄ aryl". The "C₆₋₁₄ aryl-C₁₋₄ alkyl" group includes, for example, benzyl, phenethyl, 3-phenylpropyl, 4-phenylbutyl, naphthalen-1-ylmethyl, naphthalen-2-ylmethyl,

anthracen-1-ylmethyl, anthracen-2-ylmethyl, and anthracen-9-ylmethyl.

[0047]


The term "C₆₋₁₄ arylene" means a divalent group derived from an aromatic hydrocarbon group with 6 to 14 carbon atoms. The "C₆₋₁₄ arylene" group includes, for example, the following groups:



[0048]


As for the term "substituted", for example, the phrase "C₃₋₈ cycloalkyl optionally substituted with the same or different 1 to 3 substituents selected from Group A¹" in R¹ means unsubstituted C₃₋₈ cycloalkyl or a group where any of replaceable hydrogen atoms in C₃₋₈ cycloalkyl are substituted with the same or different 1 to 3 substituents selected from Group A¹, i.e., the group consisting of (1) halogen, (2) C₁₋₄ alkyl, and (3) halo-C₁₋₄ alkyl. Such a

substituted C₃₋₈ cycloalkyl group includes, for example, the following groups:



[0049]

5 The term "a compound of Formula [I]" herein may also be referred to as "Compound [I]". In one embodiment, Compound [I] is a compound of Formula [II]:



wherein each variable is defined as defined above. The
10 term "a compound of Formula [II]" herein may also be referred to as "Compound [II]".


[0050]

Embodiments of partial structures and substituents of Compound [I] and Compound [II] are illustrated as below,
15 but each partial structure or substituent of Compound [I]



and Compound [II] is not limited to those embodiments; Compound [I] and Compound [II] include any combinations of two or more embodiments optionally selected from the embodiments in each partial structure or substituent.

5 [0051]









The following partial structure of Compound [I]:




means either of the following partial structures:











10 A preferable structure is any of the following partial structures:



A more preferable one is any of the following partial structures:




A further preferable one is any of the following partial structures:





[0052]

The following partial structure of Compound [II]:




means either of the following partial structures:




A preferable structure is any of the following partial


5 structures:



A more preferable one is any of the following partial structures:




A further preferable one is any of the following partial structures:




[0053]


Another embodiment of the following partial structure of Compound [I]:




includes the following partial structure:




Such a partial structure means either of the following
5 partial structures:




a preferable one is any of the following partial
structures:



A more preferable one is any of the following partial structures:



A further preferable one is any of the following partial structures:



[0054]

R¹ is preferably C₁₋₈ alkyl, or C₃₋₈ cycloalkyl substituted with the same or different 1 to 3 substituents selected from Group A¹, and is more preferably C₁₋₈ alkyl.

[0055]


Group A¹ is preferably halogen and C₁₋₄ alkyl.

[0056]

X¹ is preferably a bond.

[0057]

5 A partial structure of -X¹-R¹ is preferably any one of the following structures:



[0058]

R² is preferably halogen and more preferably chloro.

10 [0059]

R³ is preferably -Y³-COO-R³⁰.

[0060]

Y³ is preferably C₁₋₈ alkylene, C₃₋₈ cycloalkylene, or bridged C₅₋₈ cycloalkylene, and more preferably C₃₋₈ cycloalkylene or bridged C₅₋₈ cycloalkylene.

15


[0061]

C₁₋₈ alkylene in Y³ is preferably any one of the following groups:




20 [0062]

C_{3-8} cycloalkylene in Y^3 is preferably the following group:




[0063]

5 Bridged C_{5-8} cycloalkylene in Y^3 is preferably any one of the following groups:



[0064]

C_{6-14} arylene in Y^3 is preferably the following group:



10

[0065]

R^{30} is preferably hydrogen or ethyl, and more preferably hydrogen.

[0066]

15 R^4 is preferably hydrogen or methyl, and more preferably hydrogen.

[0067]

R^5 and R^6 are preferably each independently hydrogen,

C₁₋₄ alkyl, cyano-C₁₋₄ alkyl, or C₁₋₄ alkyl substituted with one substituent selected from the group consisting of -O-R⁵¹, -COO-R⁵², -N(R⁷¹)(R⁷²), -CO-N(R⁷³)(R⁷⁴), -N(R⁷⁵)-CO-R⁶², and -O-S(O)₂-R⁶³ and are more preferably each independently
 5 hydrogen or C₁₋₄ alkyl.

[0068]

In one embodiment, R⁵ and R⁶ are selected from the following options:

- (A) both are hydrogen;
- 10 (B) one is hydrogen and the other is C₁₋₄ alkyl, preferably methyl;
- (C) both are C₁₋₄ alkyl, preferably methyl for both;
- (D) one is hydrogen and the other is cyano-C₁₋₄ alkyl, preferably cyanomethyl; or
- 15 (E) one is hydrogen and the other is C₁₋₄ alkyl, preferably methyl or ethyl, substituted with one substituent selected from the group consisting of -O-R⁵¹, -COO-R⁵², -N(R⁷¹)(R⁷²), -CO-N(R⁷³)(R⁷⁴), -N(R⁷⁵)-CO-R⁶², and -O-S(O)₂-R⁶³.

[0069]

20 R⁷ and R⁸ are preferably each independently hydrogen, halogen, cyano, C₁₋₄ alkyl, halo-C₁₋₄ alkyl, C₁₋₄ alkoxy, or C₁₋₄ alkyl substituted with -O-R⁵¹, and more preferably, both of them are hydrogen.

[0070]

25 In one embodiment, R⁷ and R⁸ are selected from the

following options:

(A) both are hydrogen;

(B) both are halogen, preferably fluoro for both;

(C) both are C₁₋₄ alkyl, preferably methyl for both; or

5 (D) one is hydrogen and the other is cyano, C₁₋₄ alkyl (preferably methyl, ethyl, or isopropyl), halo-C₁₋₄ alkyl (preferably trifluoromethyl), C₁₋₄ alkoxy (preferably methoxy), or C₁₋₄ alkyl, preferably methyl, substituted with one -O-R⁵¹.

10 [0071]

R⁹ and R¹⁰ are preferably each independently hydrogen or C₁₋₄ alkyl, and more preferably, both of them are hydrogen.

[0072]

15 In one embodiment, R⁹ and R¹⁰ are selected from the following options:

(A) both are hydrogen; or

(B) both are methyl.

[0073]

20 In one embodiment, X² is =N- and R³ is hydrogen.

[0074]

In another embodiment, X² is =C(R⁴)- and R³ is -Y³-COO-R³⁰.

[0075]

25 The term "pharmaceutically acceptable salt" may be any

salts without excess toxicity known in the art. Specifically, it includes, for example, salts with inorganic acids, salts with organic acids, salts with inorganic bases, and salts with organic bases. Various
5 forms of pharmaceutically acceptable salts are well known in the art and are listed, for example, in the following references:

(a) Berge et al., J. Pharm. Sci., 66, p1-19 (1977);

(b) Stahl et al., "Handbook of Pharmaceutical Salts:
10 Properties, Selection, and Use" (Wiley-VCH, Weinheim, Germany, 2002);

(c) Paulekuhn et al., J. Med. Chem., 50, p6665-6672 (2007).

According to known methods, Compound [I] may be reacted with an inorganic acid, organic acid, inorganic
15 base, or organic base to give each pharmaceutically acceptable salt thereof.

[0076]

Such salts with inorganic acids include, for example, salts with hydrofluoric acid, hydrochloric acid,
20 hydrobromic acid, hydroiodic acid, nitric acid, phosphoric acid, and sulfuric acid. Preferable salts include salts with hydrochloric acid, nitric acid, sulfuric acid, phosphoric acid, and hydrobromic acid.

Such salts with organic acids include, for example,
25 salts with acetic acid, adipic acid, alginic acid, 4-

aminosalicylic acid, anhydromethylenecitric acid, benzoic acid, benzenesulfonic acid, camphor acid, camphor-10-sulfonic acid, carbonic acid, citric acid, edetic acid, ethane-1,2-disulfonic acid, dodecylsulfonic acid, 5 ethanesulfonic acid, fumaric acid, glucoheptonic acid, gluconic acid, glucuronic acid, glucoheptonic acid, glycollylarsanilic acid, hydroxynaphthoic acid, 2-hydroxy-1-ethanesulfonic acid, lactic acid, lactobionic acid, malic acid, maleic acid, mandelic acid, methanesulfonic acid, 10 methylsulfuric acid, methylnitric acid, methylenebis(salicylic acid), galactaric acid, naphthalene-2-sulfonic acid, 2-naphthoic acid, 1,5-naphthalenedisulfonic acid, oleic acid, oxalic acid, pamoic acid, pantothenic acid, pectic acid, picric acid, propionic acid, 15 polygalacturonic acid, salicylic acid, stearic acid, succinic acid, tannic acid, tartaric acid, teoclic acid, thiocyanic acid, trifluoroacetic acid, p-toluenesulfonic acid, undecanoic acid, asparaginic acid, and glutamic acid. Preferable salts include salts with oxalic acid, maleic acid, 20 citric acid, fumaric acid, lactic acid, malic acid, succinic acid, tartaric acid, acetic acid, trifluoroacetic acid, benzoic acid, glucuronic acid, oleic acid, pamoic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, and 2-hydroxy-1-ethanesulfonic acid.

25 [0077]

Such salts with inorganic bases include, for example, salts with lithium, sodium, potassium, magnesium, calcium, barium, aluminum, zinc, bismuth, and ammonium. Preferable salts include salts with sodium, potassium, calcium, magnesium, and zinc.

Such salts with organic bases include, for example, salts with arecoline, betaine, choline, clemizole, ethylenediamine, N-methylglucamine, N-benzylphenethylamine, tris(hydroxymethyl)methylamine, arginine, and lysine. Preferable salts include salts with tris(hydroxymethyl)methylamine, N-methylglucamine, and lysine.

[0078]

A preferable "pharmaceutically acceptable salt" includes hydrochloride and sodium salt.

[0079]

Compound [I] or a pharmaceutically acceptable salt thereof may exist in a solvate form.

The term "solvate" means Compound [I] or a pharmaceutically acceptable salt thereof coordinate with a solvent molecule and includes a hydrate. Such a solvate is preferably a pharmaceutically acceptable solvate and includes hydrates, ethanolates, and solvates with dimethylsulfoxide of Compound [I] or a pharmaceutically acceptable salt thereof.

Specifically, such a solvate includes a hemihydrate, monohydrate, dihydrate, or monoethanolate of Compound [I], or a monohydrate of a hydrochloride salt of Compound [I] or a 2/3 ethanolate of a dihydrochloride salt thereof. Such a
5 solvate may be obtained according to known methods.

[0080]

Compound [I] or a pharmaceutically acceptable salt thereof may exist in its tautomeric form. Such Compound [I] or a pharmaceutically acceptable salt thereof may exist
10 in each tautomeric form or in the form of a mixture of its tautomers.

[0081]

Compound [I] or a pharmaceutically acceptable salt thereof may have stereoisomers recognized as cis/trans
15 isomers. Such Compound [I] or a pharmaceutically acceptable salt thereof may exist in its cis or trans form, or in the form of a mixture of its cis and trans isomers.

[0082]

Compound [I] or a pharmaceutically acceptable salt
20 thereof may have one or more asymmetric carbon atoms. Such Compound [I] or a pharmaceutically acceptable salt thereof may exist in a single enantiomeric form or a single diastereomeric form, or in the form of a mixture of its enantiomers or diastereomers.

25 [0083]


Compound [I] or a pharmaceutically acceptable salt thereof may exist in its atropisomeric form. Such Compound [I] or a pharmaceutically acceptable salt thereof may exist in each atropisomeric form or in the form of a mixture of its atropisomers.

[0084]

Compound [I] or a pharmaceutically acceptable salt thereof may simultaneously comprise multiple structural features responsible for the above isomers. Compound [I] or a pharmaceutically acceptable salt thereof may comprise the above isomers in any ratios.


[0085]

Formulae, chemical structures, or compound names herein without stereochemistry specified include any of the above isomers available, unless otherwise specified. For example, the following structure:



includes, unless otherwise specified, all of:

(1) a racemate of the following two (i.e., S- and R-) enantiomers:



(2) the S-enantiomer; and

(3) the R-enantiomer.

[0086]

5 A diastereomeric mixture may be separated into each diastereomer by conventional methods such as chromatography and crystallization. Each diastereomer may also be prepared with a stereochemically-single starting material or by synthetic methods with stereoselective reactions.

10 [0087]

An enantiomeric mixture may be separated into each single enantiomer by methods well known in the art.

For example, an enantiomeric mixture may be reacted with a substantially pure enantiomer that is known as a
 15 chiral auxiliary to form a diastereomeric mixture, followed by separation from the diastereomeric mixture by ordinary methods such as fractional crystallization and chromatography to give a single diastereomer with an enhanced isomeric ratio or a substantially pure single

diastereomer. Then, the separated diastereomer may be converted into a desired enantiomer by removal of the added chiral auxiliary in a cleavage reaction.

An enantiomeric mixture may also be directly separated
5 into each enantiomer by chromatography methods with a chiral stationary phase well known in the art. Alternatively, either of enantiomers may be obtained with a substantially-pure optically-active starting material or by stereoselective synthesis, i.e., asymmetric induction, for
10 a prochiral intermediate with a chiral auxiliary or asymmetric catalyst.

[0088]

Absolute configurations may be determined by X-ray crystallography for crystalline products or intermediates.
15 Crystalline products or intermediates derivatized with a reagent with a known configuration and an asymmetric center may optionally be used in the determination.

[0089]

Compound [I] or a pharmaceutically acceptable salt
20 thereof may be labelled with an isotope atom such as ^2H , ^3H , ^{14}C , and ^{35}S .

For example, any hydrogen atoms of Compound [I] include protium ^1H (H), deuterium ^2H (D), and tritium ^3H (T). For example, when C_{1-8} alkyl of R^1 is ethyl, the ethyl group
25 includes $-\text{CD}_2\text{CD}_3$ and $-\text{CT}_2\text{CT}_3$ besides $-\text{CH}_2\text{CH}_3$.

[0090]

Compound [I], or a pharmaceutically acceptable salt thereof is preferably Compound [I], or a pharmaceutically acceptable salt thereof, substantially purified. More
5 preferable one is Compound [I], or a pharmaceutically acceptable salt thereof, having 80% or more of purity.

[0091]

According to known methods in the art of pharmaceutical formulations, a pharmaceutical composition
10 herein may be prepared by, for example, mixing Compound [I] or a pharmaceutically acceptable salt thereof with at least one or more pharmaceutically acceptable carrier(s) in an appropriate amount. The content (also referred to as "a therapeutically effective amount" herein) of Compound [I]
15 or a pharmaceutically acceptable salt thereof in the pharmaceutical composition varies depending on dosage forms and doses and is, for example, 0.1 to 100% by weight of the composition.

[0092]

20 A dosage form of Compound [I] or a pharmaceutically acceptable salt thereof includes an oral preparation such as tablets, capsules, granules, powders, lozenges, syrups, emulsions, and suspensions and a parenteral preparation such as external preparations, suppositories, injections,
25 eye drops, nasal preparations, and pulmonary preparations.

[0093]

The term "pharmaceutically acceptable carrier" includes various conventional organic or inorganic carrier substances for formulation materials such as excipients, disintegrants, binders, fluidizers, and lubricants in solid formulations; solvents, solubilizing agents, suspending agents, tonicity agents, buffers, and soothing agents in liquid formulations; and bases, emulsifying agents, wetting agents, stabilizers, stabilizing agents, dispersants, plasticizers, pH regulators, absorption promoters, gelators, preservatives, fillers, solubilizers, solubilizing agents, and suspending agents in semisolid formulations. A preserving agent, an antioxidant agent, a colorant, or a sweetening agent may also be optionally used as an additive.

15 [0094]

Such an "excipient" includes, for example, lactose, white soft sugar, D-mannitol, D-sorbitol, cornstarch, dextrin, microcrystalline cellulose, crystalline cellulose, carmellose, carmellose calcium, sodium carboxymethyl starch, low substituted hydroxypropyl cellulose, and gum arabic.

[0095]

Such a "disintegrant" includes, for example, carmellose, carmellose calcium, carmellose sodium, sodium carboxymethyl starch, croscarmellose sodium, crospovidone, low substituted hydroxypropyl cellulose, hydroxypropyl

methylcellulose, and crystalline cellulose.

[0096]

Such a "binder" includes, for example, hydroxypropyl cellulose, hydroxypropyl methylcellulose, povidone, 5 crystalline cellulose, white soft sugar, dextrin, starch, gelatin, carmellose sodium, and gum arabic.

[0097]

Such a "fluidizer" includes, for example, light anhydrous silicic acid and magnesium stearate.

10 [0098]

Such a "lubricant" includes, for example, magnesium stearate, calcium stearate, and talc.

[0099]

Such a "solvent" includes, for example, purified water, 15 ethanol, propyleneglycol, macrogol, sesame oil, corn oil, and olive oil.

[0100]

Such a "solubilizing agent" includes, for example, propyleneglycol, D-mannitol, benzyl benzoate, ethanol, 20 triethanolamine, sodium carbonate, and sodium citrate.

[0101]

Such a "suspending agent" includes, for example, benzalkonium chloride, carmellose, hydroxypropyl cellulose, propyleneglycol, povidone, methylcellulose, and glyceryl 25 monostearate.

[0102]

Such a "tonicity agent" includes, for example, glucose, D-sorbitol, sodium chloride, and D-mannitol.

[0103]

5 Such a "buffer" includes, for example, sodium hydrogen phosphate, sodium acetate, sodium carbonate, and sodium citrate.

[0104]

10 Such a "soothing agent" includes, for example, benzyl alcohol.

[0105]

Such a "base" includes, for example, water, animal or vegetable oils such as olive oil, corn oil, arachis oil, sesame oil, and castor oil, lower alcohols such as ethanol, propanol, propylene glycol, 1,3-butylene glycol, and phenol, higher fatty acids and esters thereof, waxes, higher alcohols, polyalcohols, hydrocarbons such as white petrolatum, liquid paraffin, and paraffin, hydrophilic petrolatum, purified lanolin, absorptive ointment, hydrous lanolin, hydrophilic ointment, starch, pullulan, gum arabic, tragacanth gum, gelatin, dextran, cellulose derivatives such as methylcellulose, carboxymethylcellulose, hydroxyethyl cellulose, and hydroxypropyl cellulose, synthetic polymers such as carboxyvinyl polymers, sodium polyacrylate, polyvinyl alcohol, and polyvinylpyrrolidone,

15
20
25

propylene glycol, macrogol such as macrogol 200 to 600, and a combination of any two or more of them.

[0106]

Such a "preserving agent" includes, for example, ethyl
5 parahydroxybenzoate, chlorobutanol, benzyl alcohol, sodium dehydroacetate, and sorbic acid.

[0107]

Such an "antioxidant agent" includes, for example, sodium sulfite and ascorbic acid.

10 [0108]

Such a "colorant" includes, for example, food dye such as Food Red No. 2 and No. 3, and Food Yellow No. 4 and No. 5, and β -carotene.

[0109]

15 Such a "sweetening agent" includes, for example saccharin sodium, dipotassium glycyrrhizate, and aspartame.

[0110]

A pharmaceutical composition herein may be administered orally or parenterally such as locally,
20 rectally, intravenously, intramuscularly, and subcutaneously to human as well as mammals other than human such as mice, rats, hamsters, guinea pigs, rabbits, cats, dogs, pigs, cattle, horses, sheep, and monkeys. A dose may vary depending on subjects to be administered, diseases,
25 symptoms, dosage forms, routes of administration, etc. For

example, in oral administration to an adult patient, the dose of Compound [I], the active ingredient, ranges generally from about 0.01 mg to about 1 g per day, which may be administered once or several times in a divided amount.

[0111]

A kit such as kits for administration, treatment, and/or prevention, a package such as packaged goods, and a set and/or case of medicine which comprises a pharmaceutical composition comprising Compound [I] or a pharmaceutically acceptable salt thereof as the active ingredient or active agent and a written matter concerning the composition indicating that the composition may or should be used for treatment and/or prevention are also useful. Such a kit, package, and set of medicine may comprise one or more containers filled with the pharmaceutical composition or one or more active ingredients and other drugs or medicines (or ingredients) used for the composition. Examples of such a kit, package, and set of medicine include commercial kits, commercial packages, and commercial medicine set for appropriate use in the treatment and/or prevention of intended diseases. The written matter comprised in such a kit, package, and set of medicine includes a cautionary note or package insert in the form designated by the government

organization that regulates manufactures, use, or sales of pharmaceutical or biological products which ensures an approval by the government organization on manufactures, use, or sales of products concerning administration to humans. The kit, package, and set of medicine may include packaged products as well as structures configured for appropriate administration steps and configured so as to be able to achieve more preferable medical treatment and/or prevention including treatment and/or prevention of intended diseases.

[0112]

Compound [I] or a pharmaceutically acceptable salt thereof has RORy antagonism and is useful for an RORy antagonist.

[0113]

The term "having RORy antagonist activity", "having RORy antagonism", or "antagonizing RORy" means that the function of RORy is antagonized, preferably specifically antagonized, to disappear or reduce its activity, and includes, for example, antagonizing, preferably specifically antagonizing, the function of RORy according to the conditions described in Test Example 1 below.

The term "RORy antagonist" means any substances that antagonize the function of RORy, preferably any substances that specifically antagonize the function of RORy.

The term "ROR γ " is preferably "human ROR γ ".

[0114]

Compound [I] or a pharmaceutically acceptable salt thereof has ROR γ antagonism, and is expected to be effective against diseases that involve the function of ROR γ .

Specifically, Compound [I] or a pharmaceutically acceptable salt thereof is expected to be useful for treating or preventing a disease selected from the group consisting of autoimmune diseases, allergic diseases, dry eye, fibrosis, cancers, metabolic disease, ischemia, cardiomyopathy, hypertension, and periodontal disease.

[0115]

The term "autoimmune diseases" means a generic name of diseases where an immune system of a subject overreacts to and attacks even normal cells and tissues thereof to cause symptoms, and includes, specifically, rheumatoid arthritis, psoriasis, inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, multiple sclerosis, systemic lupus erythematosus (SLE), Behcet's disease, sarcoidosis, Harada disease, ankylosing spondylitis, uveitis, polymyalgia rheumatica, type I diabetes, graft-versus-host disease, alopecia areata, and vitiligo.

The term "allergic diseases" means diseases derived from the condition where an immune reaction excessively

occurs against a certain antigen, and includes, specifically, atopic dermatitis, allergic rhinitis such as pollen allergy, allergic conjunctivitis, allergic gastroenteritis, asthma such as bronchial asthma and
5 infantile asthma, food allergy, medication allergy, and hives.

The term "fibrosis" means a condition with increased fibroconnective tissues, and includes, specifically, lung fibrosis and primary biliary cirrhosis.

10 The term "cancer" includes malignant melanoma and prostate cancer.

The term "metabolic disease" means a disease caused by abnormality of metabolic turnover or a disease which includes metabolic abnormality as an element that
15 constitutes pathogenesis, and includes, for example, diabetes such as type I diabetes and type II diabetes, hepatic steatosis, and non-alcoholic fatty liver disease.

[0116]

The term "treating" used herein also includes
20 ameliorating symptoms, preventing from becoming severe, maintaining remission, preventing exacerbation, and preventing relapse.

The term "preventing" used herein means suppressing pathogenesis of symptoms.

25 [0117]

As long as an embodiment disclosed herein is compatible with another embodiment disclosed in another portion of the description, any two or more combinations of these embodiments are also intended to be included in the invention.

[0118]

A general method of preparing Compound [I] or a pharmaceutically acceptable salt thereof is illustrated as below. A method of preparing Compound [I] or a pharmaceutically acceptable salt thereof, however, is not intended to be limited thereto. Salts of each compound in the general method may be selected from the above "pharmaceutically acceptable salt" unless otherwise specified.

Each compound obtained in each step may be isolated and/or purified by known methods such as distillation, recrystallization, and column chromatography, if necessary, but each reaction may optionally proceed to a sequential step without isolation and/or purification.

The room temperature herein means a temperature under no control, and includes 1°C to 40°C as one embodiment.

[0119]


Abbreviations used herein are defined as follows.

IPA: Isopropyl alcohol

25 Hex.: n-Hexane

- DMSO: Dimethyl sulfoxide
- NOE: Nuclear overhauser effect
- DsPhSO₃N₃: p-Dodecylbenzenesulfonylazide
- DMEAD: Di-2-methoxyethyl azodicarboxylate
- 5 TBAI: Tetrabutylammonium iodide
- PPTS: Pyridinium p-toluenesulfonate
- THF: Tetrahydrofuran
- WSC · HCl: 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
- 10 DMAP: Dimethylaminopyridine
- LDA: Lithium diisopropylamide
- DMF: N,N-Dimethylformamide
- DIBAL-H: Diisobutylaluminum hydride
- TFA: Trifluoroacetic acid
- 15 NaHMDS: Sodium bis(trimethylsilyl)amide
- HMDS: Bis(trimethylsilyl)amine
- TEMPO: 2,2,6,6-Tetramethylpiperidin-1-oxyl
- TBAF: Tetrabutylammonium fluoride
- [0120]
- 20 [Preparation method 1]: Preparation of Compound [I-1] or a salt thereof

Compound [I] wherein X² is =C(R⁴)-, or a salt thereof, may be obtained by, for example, the following Preparation method 1.



In the scheme, R^1 , R^2 , R^3 , R^4 , X^1 , X^3 , X^4 , and X^5 are defined as defined above, and

L^1 is halogen, for example, selected from chloro, bromo, and iodo.

[0121]

(Step 1-1)

Compound [13] or a salt thereof may be prepared by reaction of Compound [11] with Compound [12] or a salt thereof in a solvent in the presence of an organometallic reagent and Lewis acid.

Such a solvent includes, for example, ether solvents such as tetrahydrofuran; hydrocarbon solvents such as toluene; and a mixed solvent of any of them. A preferable solvent herein is tetrahydrofuran.

Such an organometallic reagent includes, for example,

n-butyllithium and tert-butyllithium. A preferable organometallic reagent herein is n-butyllithium.

Such Lewis acid includes a boron trifluoride-diethyl ether complex.

5 The reaction temperature herein ranges, for example, from -102°C to -69°C, preferably from -78°C to -70°C.

Compound [11] is commercially available or may be prepared by known methods from commercially available products.

10 Compound [12] or a salt thereof may be prepared by, for example, any of Preparation methods 1A to 1R below.

[0122]

(Step 1-2)

Compound [14] or a salt thereof may be prepared by
15 reduction of Compound [13] or a salt thereof in a solvent in the presence of a metal reagent and an acid.

Such a metal reagent includes, for example, zinc and iron. A preferable metal reagent herein is zinc.

Such an acid includes, for example, acetic acid,
20 trifluoroacetic acid, hydrochloric acid, and sulfuric acid. A preferable acid herein is acetic acid or hydrochloric acid.

Such a solvent includes, for example, ether solvents such as tetrahydrofuran; alcohol solvents such as methanol;
25 water; and a mixed solvent of any of them. A preferable

solvent herein is tetrahydrofuran, methanol, or water.

The reaction temperature herein ranges, for example, from 0°C to 80°C, preferably from room temperature to 80°C.

Compound [14] or a salt thereof may also prepared by
5 hydrogenation of Compound [13] or a salt thereof in a solvent in the presence of a catalytic amount of palladium. Such a solvent includes, for example, ether solvents such as tetrahydrofuran; alcohol solvents such as ethanol; ester solvents such as ethyl acetate; and a mixed solvent of any
10 of them. A preferable solvent herein is tetrahydrofuran, ethanol, or ethyl acetate. The reaction temperature herein is room temperature.

[0123]

(Step 1-3)

15 Compound [16] or a salt thereof may be prepared by reaction of Compound [14] or a salt thereof with Compound [15] or a salt thereof in a solvent.

Such a solvent includes, for example, hydrocarbon solvents such as toluene; ether solvents such as
20 tetrahydrofuran; halogenated solvents such as dichloromethane; and a mixed solvent of any of them. A preferable solvent herein is toluene, tetrahydrofuran, or dichloromethane.

The reaction temperature herein ranges, for example,
25 from 0°C to 80°C, preferably from 0°C to room temperature.

The reaction may also be carried out with optional addition of triethylamine.

[0124]

(Step 1-4)

5 Compound [I-1] or a salt thereof may be prepared by oxidation of Compound [16] or a salt thereof in a solvent in the presence of an oxidizing agent, followed by cyclization.

Such a solvent includes, for example, halogenated
10 solvents such as chloroform; ester solvents such as ethyl acetate; nitrile solvents such as acetonitrile; ether solvents such as cyclopentyl methyl ether; carboxylic acid solvents such as acetic acid; and a mixed solvent of any of them. A preferable solvent herein is dichloromethane,
15 chloroform, cyclopentyl methyl ether, or acetic acid.

Such an oxidizing agent includes, for example, 2-azaadamantan-N-oxyl, 2,2,6,6-tetramethylpiperidin-1-oxyl radical, and Dess-Martin reagent. The reaction may also be carried out with optional addition of a co-oxidizing agent
20 such as (diacetoxyiodo)benzene and sodium hypochlorite. A preferable oxidizing agent herein is a mixture of 2,2,6,6-tetramethylpiperidin-1-oxyl radical and (diacetoxyiodo)benzene.

An acid in the cyclization includes hydrochloric acid,
25 trifluoroacetic acid, and p-toluenesulfonic acid. A


preferable acid herein is trifluoroacetic acid.

The reaction temperature herein ranges, for example, from 0°C to 80°C, preferably from 0°C to room temperature.


[0125]

- 5 [Preparation method 1A]: Preparation of Compound [I-1A] or a salt thereof

Compound [I] wherein X^2 is $=C(R^4)-$ and R^4 is hydrogen (Compound [I-1A]):



- 10 wherein R^1 , R^2 , R^3 , X^1 , X^3 , X^4 , and X^5 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12a] or a salt thereof obtained in Preparation method 1A as follows, instead of Compound [12] or a salt thereof in Preparation method 1.



15

In the scheme, X^3 , X^4 , and X^5 are defined as defined above.

[0126]

(Step 1A-1)

Compound [A2] or a salt thereof may be prepared by oxidation of Compound [A1] or a salt thereof in a solvent.

Such a solvent includes, for example, ester solvents such as ethyl acetate; hydrocarbon solvents such as toluene; sulfoxide solvents such as dimethyl sulfoxide; ether solvents such as tetrahydrofuran; and halogenated solvents such as chloroform. A preferable solvent herein is chloroform or dichloromethane.

The oxidizing agent herein includes, for example, 2,2,6,6-tetramethylpiperidin-1-oxyl radical, dimethyl sulfoxide, a sulfur trioxide-pyridine complex, iodoxybenzoic acid, pyridinium chlorochromate, and Dess-Martin reagent. A preferable oxidizing agent herein is 2,2,6,6-tetramethylpiperidin-1-oxyl radical.

The reaction temperature herein ranges, for example, from -78°C to room temperature, preferably from 0°C to room temperature.

The reaction may also be carried out with optional addition of (diacetoxyiodo)benzene.

[0127]

(Step 1A-2)

Compound [A3] or a salt thereof may be prepared by reaction of Compound [A2] or a salt thereof with hydroxylamine hydrochloride in a solvent.

Such a solvent includes, for example, alcohol solvents

such as ethanol; hydrocarbon solvents such as toluene; halogenated solvents such as dichloromethane; ether solvents such as tetrahydrofuran; amide solvents such as dimethylformamide; nitrile solvents such as acetonitrile; 5 water; and a mixed solvent of any of them. A preferable solvent herein is ethanol, toluene, tetrahydrofuran, or water.

The reaction temperature herein ranges from room temperature to 120°C.

10 The reaction may also be carried out with optional addition of sodium acetate.

[0128]

(Step 1A-3)

Compound [12a] or a salt thereof may be prepared by 15 cyclization of Compound [A3] or a salt thereof in a solvent in the presence of an oxidizing agent.

Such an oxidizing agent includes, for example, (diacetoxyiodo)benzene, sodium hypochlorite, chloramine T, and N-chlorosuccinimide. A preferable oxidizing agent 20 herein is (diacetoxyiodo)benzene or sodium hypochlorite.

When (diacetoxyiodo)benzene is used for the oxidizing agent, an acid is used for an additive. Such an acid includes trifluoroacetic acid. A solvent used herein includes, for example, alcohol solvents such as methanol; 25 halogenated solvents such as dichloromethane; and a mixed


solvent of any of them. A preferable solvent herein is methanol or dichloromethane. The reaction temperature herein ranges from 0°C to room temperature.

When aqueous sodium hypochlorite solution is used for the oxidizing agent, a base is used for an additive. Such a base includes, for example, triethylamine and pyridine. A preferable base herein is triethylamine. A solvent used herein includes, for example, halogenated solvents such as dichloromethane; alcohol solvents such as ethanol; nitrile solvents such as acetonitrile; ether solvents such as tert-butyl methyl ether; and a mixed solvent of any of them. A preferable solvent herein is dichloromethane. The reaction temperature herein ranges, for example, from 0°C to room temperature and is preferably room temperature.


[0129]

[Preparation method 1B]: Preparation of Compound [I-1B] or a salt thereof

Compound [I] wherein X^2 is $=C(R^4)-$, R^4 is hydrogen, X^4 is a bond, and X^5 is $-O-$ (Compound [I-1B]):



wherein R^1 , R^2 , R^3 , X^1 , and X^3 are defined as defined above,
 or a salt thereof, may be obtained by, for example, using
 Compound [12b] or a salt thereof obtained in Preparation
 method 1B as follows, instead of Compound [12] or a salt
 5 thereof in the above Preparation method 1.



In the scheme, X^3 is defined as defined above, and

G^1 is a protective group of carboxy, for example,
 selected from methyl, ethyl, isopropyl, tert-butyl, or
 10 benzyl.

[0130]

(Step 1B-1)

Compound [B3] or a salt thereof may be prepared by
 reaction of Compound [B1] with Compound [B2] or a salt
 15 thereof in a solvent or without any solvents in the
 presence of a catalyst.

Such a solvent includes, for example, halogenated
 solvents such as dichloromethane; ester solvents such as
 ethyl acetate; ether solvents such as diethyl ether;

hydrocarbon solvents such as benzene; and a mixed solvent of any of them. Herein, no solvent or dichloromethane is preferable.

Such a catalyst includes, for example, rhodium (II) acetate dimer dihydrate, indium (III) chloride, and iron (III) chloride. A preferable catalyst herein is rhodium (II) acetate dimer dihydrate.

The reaction temperature herein is room temperature.

[0131]

10 (Step 1B-2)

Compound [B4] or a salt thereof may be prepared by reduction of Compound [B3] or a salt thereof in a solvent.

Such a solvent includes, for example, ether solvents such as tetrahydrofuran; halogenated solvents such as dichloromethane; hydrocarbon solvents such as toluene; and a mixed solvent of any of them. A preferable solvent herein is tetrahydrofuran, dichloromethane, or toluene.

A reducing agent used herein includes, for example, diisobutylaluminum hydride, and lithium aluminum hydride. A preferable reducing agent herein is diisobutylaluminum hydride.

The reaction temperature herein ranges, for example, from -78°C to room temperature, preferably from -78°C to 0°C.

25 [0132]

(Step 1B-3)

Compound [B5] or a salt thereof may be prepared from Compound [B4] or a salt thereof in a similar manner to Step 1A-2.

5 [0133]


(Step 1B-4)

Compound [12b] or a salt thereof may be prepared from Compound [B5] or a salt thereof in a similar manner to Step 1A-3.


10 [0134]

[Preparation method 1C]: Preparation of Compound [I-1C] or a salt thereof

Compound [I] wherein X^2 is $=C(R^4)-$, R^4 is hydrogen, X^4 is a bond, X^5 is $-C(R^9)(R^{10})-$, and both of R^9 and R^{10} are
15 hydrogen (Compound [I-1C]):



wherein R^1 , R^2 , R^3 , X^1 , and X^3 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12c] or a salt thereof obtained in Preparation
20 method 1C as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1.



In the scheme, X^3 and G^1 are defined as defined above.


[0135]

[Preparation method 1D]: Preparation of Compound [I-1D] or

5 a salt thereof

Compound [I] wherein X^2 is $=\text{C}(\text{R}^4)-$, R^4 is hydrogen, X^3 is $-\text{C}(\text{R}^5)(\text{R}^6)-$, R^5 is hydrogen, R^6 is C_{1-4} alkyl, X^4 is $-\text{C}(\text{R}^7)(\text{R}^8)-$, R^7 and R^8 are each independently hydrogen or C_{1-4} alkyl, X^5 is $-\text{C}(\text{R}^9)(\text{R}^{10})-$, and both of R^9 and R^{10} are


10 hydrogen (Compound [I-1D]):



wherein R^{6D} is C_{1-4} alkyl,

R^{7D} and R^{8D} are each independently hydrogen or C_{1-4} alkyl, and

R^1 , R^2 , R^3 , and X^1 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12d] or a salt thereof obtained in Preparation method 1D as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1.




In the scheme, G^1 , R^{6D} , R^{7D} , and R^{8D} are defined as defined above.


[0136]

[Preparation method 1E]: Preparation of Compound [I-1E] or a salt thereof

Compound [I] wherein X^2 is $=C(R^4)-$, R^4 is hydrogen, and
 5 X^5 is $-O-$ (Compound [I-1E]):



wherein R^1 , R^2 , R^3 , X^1 , X^3 , and X^4 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12e] or a salt thereof obtained in
 10 Preparation method 1E as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1.



In the scheme, X^3 and X^4 are defined as defined above.

[0137]

(Step 1E-1)

Compound [E3] may be prepared by reaction of Compound [E1] with Compound [E2] in a solvent in the presence of a
5 base.

Such a solvent includes, for example, ether solvents such as tetrahydrofuran; hydrocarbon solvents such as toluene; alcoholic solvents such as methanol; amide solvents such as dimethylformamide; sulfoxide solvents such
10 as dimethylsulfoxide; and a mixed solvent of any of them. A preferable solvent herein is tetrahydrofuran.

Such a base includes, for example, sodium hydride, sodium hydroxide, sodium tert-butoxide, sodium bis(trimethylsilyl)amide, lithium diisopropylamide, and n-
15 butyllithium. A preferable base herein is sodium hydride.

The reaction temperature herein ranges, for example, from 0°C to 140°C, preferably from 0°C to room temperature.

[0138]

(Step 1E-2)

20 Compound [E4] may be prepared by treatment of Compound [E3] in a solvent in the presence of an acid.

Such a solvent includes, for example, ether solvents such as tetrahydrofuran; halogen solvents such as dichloromethane; alcoholic solvents such as methanol;
25 water; and a mixed solvent of any of them. A preferable

solvent herein is tetrahydrofuran or water.

Such an acid includes, for example, hydrochloric acid, sulfuric acid, acetic acid, phosphoric acid, boron trifluoride-diethyl ether complex, trimethylsilyl iodide, iodine, and ion-exchange resin. A preferable acid herein is hydrochloric acid.

The reaction temperature herein ranges, for example, from 0°C to 120°C and is preferably 60°C.

[0139]

10 (Step 1E-3)

Compound [E5] or a salt thereof may be prepared from Compound [E4] in a similar manner to Step 1A-2.

[0140]


(Step 1E-4)

15 Compound [12e] or a salt thereof may be prepared from Compound [E5] or a salt thereof in a similar manner to Step 1A-3.

[0141]


[Preparation method 1F]: Preparation of Compound [I-1F] or a salt thereof

20 Compound [I] wherein X^2 is $=C(R^4)-$, R^4 is hydrogen, X^3 is $-C(R^5)(R^6)-$, R^5 is hydrogen, R^6 is C_{1-4} alkyl, X^4 is a bond, X^5 is $-C(R^9)(R^{10})-$, and R^9 and R^{10} are each independently hydrogen or C_{1-4} alkyl (Compound [I-1F]):



wherein R^{9F} and R^{10F} are each independently hydrogen or C_{1-4} alkyl, and

R^1 , R^2 , R^3 , R^{6D} , and X^1 are defined as defined above, or
 5 a salt thereof, may be obtained by, for example, using
 Compound [12f] or a salt thereof obtained in Preparation
 method 1F as follows, instead of Compound [12] or a salt
 thereof in the above Preparation method 1.




In the scheme, $\text{R}^{6\text{D}}$, $\text{R}^{9\text{F}}$, and $\text{R}^{10\text{F}}$ are defined as defined above.


[0142]

- 5 [Preparation method 1G]: Alternative preparation of Compound [I-1A] or a salt thereof

Compound [I] wherein X^2 is $=\text{C}(\text{R}^4)-$ and R^4 is hydrogen (Compound [I-1A]):



wherein R^1 , R^2 , R^3 , X^1 , X^3 , X^4 , and X^5 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12a] or a salt thereof obtained in Preparation method 1G as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1.



In the scheme, R^{G1} and R^{G2} each independently C_{1-4} alkyl, and X^3 , X^4 , and X^5 are defined as defined above.

(Step 1G-1)

Compound [G3] or a salt thereof may be prepared by reaction of Compound [G1] or a salt thereof with Compound [G2] in a solvent in the presence of a base.

5 Such a solvent includes, for example, ether solvents such as tetrahydrofuran; hydrocarbon solvents such as toluene; sulfoxide solvents such as dimethyl sulfoxide; and a mixed solvent of any of them. A preferable solvent herein is tetrahydrofuran.

10 Such a base includes sodium hydride and lithium diisopropylamide.

The reaction temperature herein ranges, for example, from -78°C to 110°C, preferably from -78°C to 65°C.

The reaction may also be carried out with optional
15 addition of 18-crown-6-ether.

[0144]

(Step 1G-2)

Compound [G5] or a salt thereof may be prepared by protection of a carbonyl group of Compound [G3] or a salt
20 thereof with Compound [G4] in a solvent in the presence of an acid.

Such a solvent includes, for example, hydrocarbon solvents such as toluene; halogenated solvents such as dichloromethane; nitrile solvents such as acetonitrile; and
25 a mixed solvent of any of them. A preferable solvent

herein is toluene.

Such an acid includes, for example, p-toluenesulfonic acid and pyridinium p-toluenesulfonate. A preferable acid herein is p-toluenesulfonic acid.

5 The reaction temperature herein ranges, for example, from room temperature to 120°C, preferably from 100°C to 120°C.

[0145]

(Step 1G-3)

10 Compound [G6] or a salt thereof may be prepared by reduction of Compound [G5] or a salt thereof in a solvent.

Such a solvent includes, for example, ether solvents such as tetrahydrofuran; hydrocarbon solvents such as toluene; halogenated solvents such as dichloromethane; and
15 a mixed solvent of any of them. A preferable solvent herein is tetrahydrofuran or toluene.

A reducing agent used herein includes, for example, lithium aluminum hydride and diisobutylaluminum hydride. A preferable reducing agent herein is diisobutylaluminum
20 hydride.

The reaction temperature herein ranges, for example, from -78°C to 65°C, preferably from -78°C to room temperature.

[0146]

25 (Step 1G-4)

Compound [G8] or a salt thereof may be prepared by Mitsunobu reaction of Compound [G6] or a salt thereof with Compound [G7] in a solvent.

Such a solvent includes, for example, ether solvents
5 such as tetrahydrofuran; halogenated solvents such as dichloromethane; hydrocarbon solvents such as toluene; and a mixed solvent of any of them. A preferable solvent herein is tetrahydrofuran or dichloromethane.

A reagent used in Mitsunobu reaction includes, for
10 example, a mixture of triphenylphosphine or tributylphosphine with diethyl azodicarboxylate or dipiperidineamide azodicarboxylate. A preferable reagent used in Mitsunobu reaction is a mixture of triphenylphosphine with diethyl azodicarboxylate.

15 The reaction temperature herein ranges, for example, from 0°C to 80°C, preferably from 0°C to room temperature.

[0147]

(Step 1G-5)

Compound [G9] or a salt thereof may be prepared by
20 removal of a phthaloyl group of Compound [G8] or a salt thereof in a solvent.

Such a solvent includes, for example, alcohol solvents
such as ethanol; halogenated solvents such as
dichloromethane; ether solvents such as diethyl ether; and
25 a mixed solvent of any of them. A preferable solvent

herein is ethanol or dichloromethane.

A reagent used for removal of the phthaloyl group includes, for example, methylhydrazine, hydrazine, and ethanolamine. A preferable reagent used for removal of the
5 phthaloyl group is methylhydrazine or hydrazine.

The reaction temperature herein ranges, for example, from 0°C to 100°C, preferably from room temperature to 100°C.

[0148]

10 (Step 1G-6)

Compound [12a] or a salt thereof may be prepared by removal of an acetal group of Compound [G9] or a salt thereof in a solvent in the presence of an acid, followed by intramolecular cyclization in the presence of a base.

15 Such a solvent includes, for example, alcohol solvents such as methanol; ether solvents such as tetrahydrofuran; halogenated solvents such as dichloromethane; and a mixed solvent of any of them. A preferable solvent herein is methanol or tetrahydrofuran.

20 Such an acid includes, for example, hydrochloric acid, acetic acid, and p-toluenesulfonic acid. A preferable acid herein is hydrochloric acid or p-toluenesulfonic acid.


Such a base includes, for example, potassium carbonate, sodium acetate, and triethylamine. A preferable base
25 herein is potassium carbonate.

The reaction temperature herein ranges, for example, from 0°C to 120°C, preferably from 0°C to room temperature.

[0149]


[Preparation method 1H]: Preparation of Compound [I-1H] or
5 a salt thereof

Compound [I] wherein X^2 is $=C(R^4)-$, R^4 is hydrogen, X^3 is $-C(R^5)(R^6)-$, R^5 is hydrogen, R^6 is C_{1-4} alkyl, X^4 is $-C(R^7)(R^8)-$, R^7 is C_{1-4} alkyl, R^8 is hydrogen, X^5 is $-C(R^9)(R^{10})-$, and both of R^9 and R^{10} are hydrogen (Compound
10 [I-1H]):



wherein R^{7H} is C_{1-4} alkyl, and

R^1 , R^2 , R^3 , R^{6D} , and X^1 are defined as defined above, or
a salt thereof, may be obtained by, for example, using
15 Compound [12h] or a salt thereof obtained in Preparation
method 1H as follows, instead of Compound [12] or a salt
thereof in the above Preparation method 1.



In the scheme, R^{6D} , R^{7H} , and G^1 are defined as defined above.


[0150]

[Preparation method 1I]: Preparation of Compound [I-1I] or


5 a salt thereof

Compound [I] wherein X^2 is $=\text{C}(\text{R}^4)-$, R^4 is hydrogen, X^3 is $-\text{C}(\text{R}^5)(\text{R}^6)-$, R^5 is hydrogen, R^6 is C_{1-4} alkyl, X^4 is $-\text{C}(\text{R}^7)(\text{R}^8)-$, R^7 is C_{1-4} alkyl, R^8 is hydrogen, X^5 is $-\text{C}(\text{R}^9)(\text{R}^{10})-$, and both of R^9 and R^{10} are hydrogen (Compound

10 [I-1I]):



wherein R^1 , R^2 , R^3 , R^{6D} , R^{7H} , and X^1 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12i] or a salt thereof obtained in
 5 Preparation method 1I as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1.




In the scheme, $\text{R}^{6\text{D}}$, $\text{R}^{7\text{H}}$, and G^1 are defined as defined above.

[0151]

[Preparation method 1J]: Preparation of Compound [I-1J] or


5 a salt thereof

Compound [I] wherein X^2 is $=\text{C}(\text{R}^4)-$ and R^4 is C_{1-4} alkyl (Compound [I-1J]):



wherein R^{4J} is C_{1-4} alkyl, and

R^1 , R^2 , R^3 , X^1 , X^3 , X^4 , and X^5 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12j] or a salt thereof obtained in Preparation method 1J as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1.



In the scheme, X^3 , X^4 , X^5 , and R^{4J} are defined as defined above.


[0152]

[Preparation method 1K]: Preparation of Compound [I-1K] or a salt thereof

Compound [I] wherein X^2 is $=C(R^4)-$, R^4 is hydrogen, X^3

is $-C(R^5)(R^6)-$, R^5 is hydrogen, and R^6 is:

- (1) halo- C_{1-4} alkyl,
- (2) cyano- C_{1-4} alkyl, or
- (3) C_{1-4} alkyl substituted with one substituent selected from the group consisting of $-O-R^{51}$, $-CO-R^{61}$, $-COO-R^{52}$, $-N(R^{71})(R^{72})$, $-CO-N(R^{73})(R^{74})$, $-N(R^{75})-CO-R^{62}$, $-N(R^{76})-COO-R^{53}$, and $-O-S(O)_2-R^{63}$ (Compound [I-1K]):




wherein R^{6K} is:

- (1) halo- C_{1-4} alkyl,
- (2) cyano- C_{1-4} alkyl, or
- (3) C_{1-4} alkyl substituted with one substituent selected from the group consisting of $-O-R^{51}$, $-CO-R^{61}$, $-COO-R^{52}$, $-N(R^{71})(R^{72})$, $-CO-N(R^{73})(R^{74})$, $-N(R^{75})-CO-R^{62}$, $-N(R^{76})-COO-R^{53}$, and $-O-S(O)_2-R^{63}$, and

R^1 , R^2 , R^3 , X^1 , X^4 , and X^5 are defined as defined above, or a part of salts thereof, may be obtained by, for example, using Compound [12k] or a salt thereof obtained in Preparation method 1K as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1, or by

converting a benzyl ether moiety of the resulting compound into various substituents.




In the scheme, X^4 , X^5 , and G^1 are defined as defined above.

5 [0153]


[Preparation method 1M]: Preparation of Compound [I-1M] or a salt thereof

Compound [I] wherein X^2 is $=\text{C}(\text{R}^4)-$, R^4 is hydrogen, X^3 is $-\text{C}(\text{R}^5)(\text{R}^6)-$, both of R^5 and R^6 are hydrogen, X^4 is a bond, and X^5 is $-\text{C}(\text{R}^9)(\text{R}^{10})-$ (Compound [I-1M]):

10



wherein R^1 , R^2 , R^3 , R^9 , R^{10} , and X^1 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12m] or a salt thereof obtained in Preparation method 1M as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1.



In the scheme, R^9 and R^{10} are defined as defined above.


[0154]

[Preparation method 1N]: Preparation of Compound [I-1N] or a salt thereof

Compound [I] wherein X^2 is $=C(R^4)-$, R^4 is hydrogen, X^3 is $-C(R^5)(R^6)-$, R^5 is hydrogen, R^6 is:

- (1) halo- C_{1-4} alkyl,
- (2) cyano- C_{1-4} alkyl, or
- (3) C_{1-4} alkyl substituted with one substituent selected from the group consisting of $-O-R^{51}$, $-CO-R^{61}$, $-COO-R^{52}$, $-N(R^{71})(R^{72})$, $-CO-N(R^{73})(R^{74})$, $-N(R^{75})-CO-R^{62}$, $-N(R^{76})-COO-R^{53}$, and $-O-S(O)_2-R^{63}$, and

X^4 is a bond (Compound [I-1N]):




wherein R^1 , R^2 , R^3 , R^{6K} , X^1 , and X^5 are defined as defined above, or a part of salts thereof, may be obtained by, for example, using Compound [12n] or a salt thereof obtained in Preparation method 1N as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1, or by converting a benzyl ether moiety of the resulting compound into various substituents.

[0155]


[Preparation method 1P]: Preparation of Compound [I-1P] or

5 a salt thereof

Compound [I] wherein X^2 is $=C(R^4)-$, R^4 is hydrogen, X^4 is a bond, and X^5 is $-N(R^{11})-$ (Compound [I-1P]):



wherein R^1 , R^2 , R^3 , R^{11} , X^1 , and X^3 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12p] or a salt thereof obtained in Preparation method 1P as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1.



In the scheme, R^{11} , X^3 , and G^1 are defined as defined above.

[0156]

10 (Step 1P-1)

Compound [P2] or a salt thereof may be prepared from Compound [P1] or a salt thereof in a similar manner to Step 1B-2.

[0157]

15 (Step 1P-2)

Compound [P3] or a salt thereof may be prepared from

Compound [P2] or a salt thereof in a similar manner to Step 1A-2.

[0158]


(Step 1P-3)

5 Compound [12p] or a salt thereof may be prepared from Compound [P3] or a salt thereof in a similar manner to Step 1A-3.

[0159]

[Preparation method 1Q]: Preparation of Compound [I-1Q] or
10 a salt thereof

Compound [I] wherein X^2 is $=C(R^4)-$, R^4 is hydrogen, X^3 is $-C(R^5)(R^6)-$, R^5 and R^6 are each independently C_{1-4} alkyl, X^4 is a bond, X^5 is $-C(R^9)(R^{10})-$, and both of R^9 and R^{10} are hydrogen (Compound [I-1Q]):




[I-1Q]

15 wherein R^{5Q} is C_{1-4} alkyl, and

R^1 , R^2 , R^3 , R^{6D} , and X^1 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12q] or a salt thereof obtained in Preparation
20 method 1Q as follows, instead of Compound [12] or a salt

thereof in the above Preparation method 1.




In the scheme, R^{5Q} and R^{6D} are defined as defined above.


[0160]

- 5 [Preparation method 1R]: Preparation of Compound [I-1R] or a salt thereof

Compound [I] wherein X² is =C(R⁴)-, R⁴ is hydrogen, X⁴ is -C(R⁷)(R⁸)-, both of R⁷ and R⁸ are hydrogen, X⁵ is -C(R⁹)(R¹⁰)-, and both of R⁹ and R¹⁰ are hydrogen (Compound [I-1R]):



wherein R^1 , R^2 , R^3 , X^1 , and X^3 are defined as defined above, or a salt thereof, may be obtained by, for example, using Compound [12r] or a salt thereof obtained in Preparation method 1R as follows, instead of Compound [12] or a salt thereof in the above Preparation method 1.




In the scheme, X^3 and G^1 are defined as defined above.

[0161]

[Preparation method 1S]: Preparation of Compound [11] used in Preparation method 1


Compound [11]:



[11]

wherein R^1 , R^2 , X^1 , and L^1 are defined as defined above,

may be prepared by, for example, cross-coupling reaction of Compound [S1] shown as follows, when X^1 is a bond, R^1 is C_{1-8} alkyl, optionally substituted C_{3-8} cycloalkyl, or optionally substituted C_{3-8} cycloalkyl- C_{1-4} alkyl (i.e.,
 5 Compound [11s] or [11t]).



In the scheme, R^2 and L^1 are defined as defined above,
 10 L^2 is halogen (e.g., iodo) or trifluoromethanesulfonyloxy,
 R^{1S} is C_{1-8} alkyl, optionally substituted C_{3-8} cycloalkyl, or optionally substituted C_{3-8} cycloalkyl- C_{1-4} alkyl,
 15 R^{W1} is boronic acid, boronic acid ester, or trifluoroborate,

R^{W2} is zinc, or zinc halide,


R^{1T} is C_{2-8} alkyl, or optionally substituted C_{3-8} cycloalkyl- C_{2-4} alkyl, and

R^{100T} is trimethylsilyl or a straight- or branched-
5 chain saturated hydrocarbon with 1 to 6 carbon atoms.

[0162]

Such a cross-coupling reaction includes methods described in literatures such as F. Diederich, P. J. Stang (1998). Metal-catalyzed Cross-coupling Reactions, Weinheim,
10 Germany, Wiley-VCH, which includes Suzuki coupling, Negishi coupling, and Sonogashira coupling.

Compound [S1] is preferably a compound wherein L^1 is bromo and L^2 is iodo, more preferably a compound shown as follows.



As for Compound [S2], Compound [S3], and Compound [S4],
for example, commercially available products such as isobutylboronic acid, 1-hexylboronic acid pinacol ester, potassium (3,3-dimethylbutyl)trifluoroborate, butylzinc
20 bromide, and cyclohexylacetylene may be used, or these compounds may be prepared from corresponding commercially available compounds with chloro, bromo, or iodo, such as 1-chloro-3,3-dimethyl-butane and bromomethyl-cyclohexane, according to known methods.

Compound [S2] wherein R^{W1} is boronic acid may be prepared by preparation of a Grignard reagent from commercially available compounds such as R^1 -Br and magnesium, followed by reaction with, for example,
5 trimethyl borate or triisopropyl borate.

Compound [S2] wherein R^{W1} is boronic acid ester may be prepared by, for example, reaction of a boronic acid compound with pinacol.

Compound [S2] wherein R^{W1} is trifluoroborate may be
10 prepared by, for example, reaction of a boronic acid compound with potassium hydrogen fluoride.

Compound [S3] may be prepared from, for example, commercially available compounds such as R^1 -I and zinc.

An activating agent for zinc includes iodine,
15 trimethylsilyl chloride, and 1,2-dibromoethane, and these agents can be used alone or in combination with any two or more of them. A preferable activating agent is trimethylsilyl chloride or 1,2-dibromoethane.

As for Compound [S4], commercially available products
20 such as 3,3-dimethyl-1-butyne, cyclohexylacetylene, and phenylacetylene may be used.

Compound [11t] may be obtained by catalytic hydrogenation of an alkynylene compound obtained in Sonogashira reaction with a catalyst such as palladium
25 carbon, platinum carbon, and rhodium-alumina to convert

into an alkyl compound.


The solvent in each step includes tetrahydrofuran, N,N-dimethylformamide, and N,N-dimethylacetamide. A preferable solvent herein is tetrahydrofuran or N,N-dimethylacetamide.

The reaction temperature in each step includes room temperature to 80°C. A preferable reaction temperature herein is room temperature.

[0163]

[Preparation method 1Z]: Preparation of Compound [15] or a salt thereof used in Preparation method 1.

Compound [15] or a salt thereof may be a commercially available product such as ethyl 6-isocyanato-hexanoate, methyl 2-isocyanato-2-methyl-propionate, methyl 3-isocyanato-propionate, ethyl 3-isocyanato-propionate, methyl 4-isocyanato-cyclohexanecarboxylate, and ethyl 4-isocyanatobenzoate, or may also be obtained by, for example, Preparation method 1Z shown as follows.



In the scheme, R³ is defined as defined above.

[0164]

Compound [15] or a salt thereof may be prepared by azidation reaction of Compound [Z1] such as a commercially

available product such as 3-(methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid, 3-(methoxycarbonyl)bicyclo[2.1.1]pentane-1-carboxylic acid, 1-(2-methoxy-2-oxoethyl)-5-oxopyrrolidine-3-carboxylic acid, and 3-[1-(ethoxycarbonyl)cyclopropyl]propanoic acid, or a salt thereof, in the presence of a base in a solvent, followed by Curtius rearrangement.

Such a solvent includes, for example, benzene, toluene, xylene, tetrahydrofuran, and a mixed solvent of any of them. A preferable solvent herein is toluene.

The azidation agent herein includes, for example, diphenylphosphoryl azide.

Such a base includes, for example, triethylamine and diisopropylethylamine. A preferable base herein is triethylamine.


The reaction temperature herein ranges, for example, from 0°C to 140°C, preferably from 100°C to 120°C.

When R^3 is $-Y^3-COO-R^{30}$ and R^{30} is C_{1-4} alkyl in Compound [15] or a salt thereof obtained from Preparation method 12 illustrated herein, Compound [I-1] wherein R^{30} is hydrogen may be obtained by preparation of Compound [I-1] wherein R^{30} is C_{1-4} alkyl from the aforementioned Compound [15] or a salt thereof in Preparation method 1, followed by hydrolysis under known methods.

[0165]

[Preparation method 2]: Preparation of Compound [I-2A] or a salt thereof

Compound [I] wherein X^2 is $=N-$, X^3 is $-C(R^5)(R^6)-$, R^3 is hydrogen, and R^5 is hydrogen (Compound [I-2A]), or a salt thereof, may be obtained by, for example, Preparation method 2 as follows.



In the scheme, R^1 , R^2 , R^6 , X^1 , X^4 , X^5 , and L^1 are defined as defined above.

10 [0166]

(Step 2-1)

Compound [22] or a salt thereof may be prepared by oxidation of Compound [21] or a salt thereof in a solvent.

Such a solvent includes, for example, ether solvents such as tetrahydrofuran; ester solvents such as ethyl acetate; hydrocarbon solvents such as toluene; halogenated

15

solvents such as dichloromethane; nitrile solvents such as acetonitrile; and a mixed solvent of any of them. A preferable solvent herein is dichloromethane.

An oxidizing agent used herein includes, for example,
5 sulfur trioxide-pyridine complex, dimethyl sulfoxide, pyridinium chlorochromate, and Dess-Martin reagent. A preferable oxidizing agent herein is sulfur trioxide-pyridine complex.

The reaction temperature herein ranges, for example,
10 from -60°C to 60°C, preferably from 0°C to room temperature.
[0167]

(Step 2-2)

Compound [24] or a salt thereof may be prepared by reaction of Compound [22] or a salt thereof with Compound
15 [23] or a salt thereof in a solvent in the presence of a base.

Such a solvent includes, for example, alcohol solvents such as ethanol; halogenated solvents such as chloroform; hydrocarbon solvents such as chlorobenzene; and a mixed
20 solvent of any of them. A preferable solvent herein is ethanol or water.

Such a base includes, for example, sodium hydrogen carbonate and triethylamine. A preferable base herein is sodium hydrogen carbonate.

25 The reaction temperature herein ranges, for example,

from -10°C to 100°C, preferably from 0°C to room temperature.

[0168]

(Step 2-3)

5 Compound [26] or a salt thereof may be prepared by reaction of Compound [24] or a salt thereof with Compound [11] or a salt thereof in a solvent.

 Such a solvent includes, for example, ether solvents such as tetrahydrofuran. A preferable solvent herein is
10 tetrahydrofuran or diethyl ether.

 A reagent used herein includes, for example, organometallic reagents such as n-butyllithium and Grignard reagents such as magnesium. A preferable reagent herein is n-butyllithium.

15 The reaction temperature herein ranges from -78°C to room temperature.

[0169]

(Step 2-4)

 Compound [I-2A] or a salt thereof may be prepared by
20 oxidation of Compound [26] or a salt thereof in a solvent.

 Such a solvent includes halogenated solvents such as dichloromethane. A preferable solvent herein is dichloromethane.

 An oxidizing agent used herein includes m-
25 chloroperoxybenzoic acid.

The reaction temperature herein is room temperature.

EXAMPLES

[0170]

5 The present invention is illustrated in more detail with Examples and Test Examples, but is not intended to be limited thereto.

¹H-NMR spectra were measured in CDCl₃, DMSO-D₆, or MeOH-D₄ with tetramethylsilane as an internal standard, and
10 all δ values are shown in ppm. Symbols in spectral data mean as follows.

s: singlet

d: doublet

t: triplet

15 q: quartet

dd: double doublet

ddd: double double doublet

brs: broad singlet

m: multiplet


20 J: coupling constant

[0171]

Example 1

(Step 1)

4-Bromo-2-chloro-1-(2,2-dimethyl-propoxy)benzene




4-Bromo-2-chloro-phenol (100 g) and cesium carbonate (126 g) were mixed in N,N-dimethylformamide (800 mL) under nitrogen gas, and thereto was added 1-iodo-2,2-dimethylpropanol (100 mL) at room temperature. The reaction solution was stirred under heating at 100°C for 2 days. The reaction solution was slowly cooled to room temperature, and then, thereto were added water (500 mL) and n-hexane (500 mL). The solution was separated. The organic layer was washed with 20w/w% aqueous sodium sulfite solution (100 mL), 2N aqueous sodium hydroxide solution (100 mL), water (100 mL), and saturated aqueous sodium chloride solution (100 mL), and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give the title compound (138 g).

¹H-NMR (400 MHz, CDCl₃) 1.07 (s, 9H), 3.61 (s, 2H), 6.76 (d, J=8.8Hz, 1H), 7.29 (dd, J=8.8, 2.3Hz, 1H), 7.49 (d, J=2.5Hz, 1H)

[0172]

(Step 2)

3-Methylsulfonyl-6,7-dihydro-5H-cyclopenta[1,2,4]triazine




Cyclopentane-1,2-dione (580 mg) was mixed in 1M aqueous sodium hydrogen carbonate solution (6.0 mL) and ethanol (6.0 mL) under nitrogen gas, and then, thereto was
 5 added S-methyl isothiosemicarbazide hydroiodide (1.38 g) at room temperature. The reaction solution was stirred at room temperature for 1 day. To the reaction solution were added ethyl acetate and water, and the reaction solution was separated. The organic layer was washed with saturated
 10 aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (20 g, ethyl acetate/chloroform = 1/5) to
 15 give the title compound (520 mg).

¹H-NMR (400 MHz, CDCl₃) 2.17-2.24 (m, 2H), 2.66 (s, 3H), 3.00 (t, J=7.7Hz, 2H), 3.17 (t, J=7.7Hz, 2H)

[0173]

(Step 3)

20 4a-[3-Chloro-4-(2,2-dihydro-propoxy)phenyl]-3-methylsulfanyl-4a,5,6,7-tetrahydro-2H-cyclopenta[1,2,4]triazine



Magnesium (30 mg) and diethyl ether (2.0 mL) were mixed under nitrogen gas, and then thereto was added one drop of a mixed solution of diethyl ether in iodine at room temperature. The reaction solution was stirred at room temperature for 30 minutes. To the reaction solution was added a mixed solution of 4-bromo-2-chloro-1-(2,2-dimethylpropoxy)benzene (420 mg) in diethyl ether (2.0 mL) at room temperature. The reaction solution was stirred under heating at 60°C for 4 hours. The reaction solution was slowly cooled to room temperature, and then, to the reaction solution was added a mixed solution of 3-methylsulfanyln-6,7-dihydro-5H-cyclopenta[1,2,4]triazine (100 mg) in tetrahydrofuran (2 mL). The reaction solution was stirred at room temperature for 1 day. To the reaction solution were added aqueous ammonium chloride solution and ethyl acetate under ice cooling, and the reaction solution was separated. The organic layer was washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. The resulting residue was purified by thin-layer silica gel column chromatography (ethyl acetate/chloroform


= 1/3) to give the title compound (17.4 mg).

¹H-NMR (400 MHz, CDCl₃) 1.06 (s, 9H), 1.68-1.78 (m, 2H),
2.13-2.20 (m, 1H), 2.39-2.43 (m, 1H), 2.44 (d, J=5.5Hz, 3H),
2.53-2.62 (m, 1H), 2.67-2.75 (m, 1H), 3.60 (s, 2H), 6.78 (d,
5 J=8.6Hz, 1H), 7.07 (dd, J=8.6, 2.3Hz, 1H), 7.30 (d, J=2.1Hz,
1H), 7.74 (br s, 1H)

[0174]

(Step 4)

4a-[3-Chloro-4-(2,2-dimethyl-propoxy)phenyl]-2,4,4a,5,6,7-
10 hexahydro-cyclopenta[1,2,4]triazin-3-one



4a-[3-Chloro-4-(2,2-dihydro-propoxy)phenyl]-3-
methylsulfanyl-4a,5,6,7-tetrahydro-2H-
cyclopenta[1,2,4]triazine (17.4 mg) and dichloromethane
15 (0.5 mL) were mixed under nitrogen gas, and thereto was
added meta-chloroperoxybenzoic acid (75 wt% of water
inclusive, 26 mg) under ice cooling. The reaction solution
was stirred at room temperature for 1 hour. To the
reaction solution were added 20w/w% aqueous sodium sulfite
20 solution (5 mL) and ethyl acetate (10 mL) under ice cooling.
The organic layer was washed with saturated aqueous sodium


hydrogen carbonate solution (5 mL) and saturated aqueous sodium chloride solution (5 mL) and dried over sodium sulfate. The resulting residue was purified by thin-layer silica gel column chromatography (methanol/chloroform = 5 1/15) to give the title compound (4.2 mg).

[0175]

Example 3

(Step 1)

Methyl 7,7-dimethyl-1,4-dioxaspiro[4.4]nonane-6-
10 carboxylate




2,2-Dimethyl-5-oxo-cyclopentanecarboxylic acid methyl ester (3.58 g) was mixed in ethylene glycol (1.76 mL) and toluene (40 mL) under nitrogen gas, and thereto was added
15 para-toluenesulfonic acid monohydrate (200 mg) at room temperature. The reaction solution was stirred under heating at 140°C for 3 hours to remove water. The resultant was slowly cooled to room temperature. Then, to the reaction solution were added 1M aqueous sodium
20 carbonate solution (1.1 mL) and ethyl acetate, and the solution was separated. The organic layer was washed with saturated aqueous sodium chloride solution and dried over

sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (4.24 g).

5 [0176]

(Step 2)

(7,7-Dimethyl-1,4-dioxa-spiro[4.4]non-6-yl)methanol



Lithium aluminum hydride (1.6 g) and tetrahydrofuran
 10 (30 mL) were mixed under nitrogen gas, and thereto was added a mixed solution of 7,7-dimethyl-1,4-dioxaspiro[4.4]none-6-carboxylic acid methyl ester (4.24 g) in tetrahydrofuran (10 mL) under ice cooling. The reaction solution was stirred at room temperature for 1 hour. Then,
 15 thereto were added sequentially water (1.6 mL), 2N aqueous sodium hydroxide solution (1.6 mL), and water (4.8 mL) under ice cooling. Celite (20 g) and magnesium sulfate (20 g) were added thereto, and the mixture was stirred at room temperature for 1 hour. The reaction solution was diluted
 20 with tetrahydrofuran (100 mL), and the resulting solid was removed through Celite. Then, the filtrate was concentrated under reduced pressure to give a crude product


of the title compound.

[0177]

(Step 3)

2-(7,7-Dimethyl-1,4-dioxa-spiro[4.4]non-6-

5 ylmethoxy)isoxazole-1,3-dione




(7,7-Dimethyl-1,4-dioxaspiro[4.4]non-6-yl)methanol,
 N-hydroxyphthalimide (5.14 g) and triphenylphosphine (8.26
 g) were mixed in tetrahydrofuran (50 mL) under nitrogen gas,
 10 and thereto was added di-2-methoxyethyl azodicarboxylate
 (7.37 g) under ice cooling. The reaction solution was
 stirred at room temperature for 3 hours. The reaction
 solution was concentrated under reduced pressure, and
 thereto were added ethyl acetate and water. The solution
 15 was separated. The organic layer was washed with saturated
 aqueous sodium chloride solution and dried over sodium
 sulfate. Sodium sulfate was removed with a filter, and
 then the filtrate was concentrated under reduced pressure.
 The resulting residue was purified by silica gel column
 20 chromatography (200 g, ethyl acetate/n-hexane = 1/3) to
 give the title compound (1.58 g).

¹H-NMR (400 MHz, CDCl₃) 1.02 (s, 3H), 1.22 (s, 3H), 1.49-

1.60 (m, 2H), 1.86-1.91 (m, 2H), 2.29 (t, J=6.6Hz, 1H),
 3.81 (dd, J=5.0, 4.5Hz, 1H), 3.91 (dt, J=6.9, 2.1Hz, 1H),
 4.03 (dq, J=16.9, 5.1Hz, 2H), 4.22 (dd, J=8.3, 6.7Hz, 1H),
 4.35 (dd, J=8.2, 6.6Hz, 1H), 7.73 (d, J=3.0Hz, 1H), 7.74 (d,
 5 J=3.0Hz, 1H), 7.82 (d, J=3.0Hz, 1H), 7.83 (d, J=3.0Hz, 1H)
 [0178]

(Step 4)

O-(7,7-Dimethyl-1,4-dioxaspiro[4.4]non-6-ylmethyl)-
 hydroxylamine



10


2-(7,7-dimethyl-1,4-dioxaspiro[4.4]non-6-
 ylmethoxy)isoxazole-1,3-dione (1.58 g) and chloroform (20
 mL) were mixed under nitrogen gas, and thereto was added N-
 methylhydrazine (0.3 mL) under ice cooling. The reaction
 15 solution was stirred at room temperature for 1 hour. The
 reaction solution was filtered through Celite, and then the
 filtrate was concentrated under reduced pressure to give a
 crude product of the title compound (1.61 g).

¹H-NMR (400 MHz, CDCl₃) 0.90 (s, 3H), 1.10 (s, 3H), 1.43-
 20 1.56 (m, 2H), 1.82-1.87 (m, 2H), 2.08 (dd, J=7.4, 6.5Hz,
 1H), 3.68-3.96 (m, 6H), 5.31 (br s, 2H)

[0179]

(Step 5)

4,4-Dimethyl-3a,4,5,6-tetrahydro-3H-cyclopenta[c]isoxazole



O-(7,7-Dimethyl-1,4-dioxaspiro[4.4]non-6-ylmethyl)-


5 hydroxylamine (1.61 g) and tetrahydrofuran (10 mL) were mixed under nitrogen gas, and thereto was added 3N hydrochloric acid (5 mL) at room temperature. The reaction solution was stirred at room temperature for 2 hours. To the reaction solution were added potassium carbonate (2.1 g) and diethyl ether, and the solution was separated. The organic layer was washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (20 g, diethyl ether/n-hexane = 1/1) to give the title compound (573 mg).

¹H-NMR (400 MHz, CDCl₃) 0.91 (s, 3H), 1.12 (s, 3H), 1.96 (ddd, J=13.0, 7.2, 3.6Hz, 1H), 2.05-2.13 (m, 1H), 2.43 - 2.49 (m, 2H), 3.53-3.59 (m, 1H), 3.89 (dd, J=11.9, 8.2Hz, 1H), 4.33 (dd, J=10.5, 8.2Hz, 1H)

[0180]

(Step 6)

6a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-4,4-dimethyl-hexahydro-cyclopenta[c]isoxazole




4-Bromo-2-chloro-1-(3,3-dimethyl-butyl)benzene (702
 5 mg) was mixed in toluene (7.0 mL) and tetrahydrofuran (2.8 mL) under argon gas, and thereto was added dropwise 1.6M n-butyllithium/n-hexane solution (1.55 mL) under cooling at -78°C. The reaction solution was stirred under cooling at -78°C for 1 hour (Reaction solution A). 4,4-Dimethyl-
 10 3a,4,5,6-tetrahydro-3H-cyclopenta[c]isoxazole (173 mg) and toluene (10 mL) were mixed, and thereto were added boron trifluoride-diethyl ether complex (0.314 mL) and Reaction solution A under cooling at -78°C. The reaction solution was stirred for 1 hour, and then thereto were added aqueous
 15 ammonium chloride solution (5 mL) and ethyl acetate at -78°C. The solution was separated. The organic layer was washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. The resulting residue was purified by silica gel column chromatography (10 g, diethyl
 20 ether/n-hexane = 1/1) to give the title compound (364 mg).

^1H -NMR (400 MHz, CDCl_3) 0.98 (s, 9H), 1.10 (s, 6H), 1.12-1.30 (m, 2H), 1.42-1.47 (m, 2H), 1.74-1.82 (m, 1H), 2.15-2.24 (m, 1H), 2.64-2.68 (m, 2H), 2.79 (dd, $J=6.7$, 5.5Hz, 1H), 3.80-4.20 (m, 2H), 4.98 (s, 1H), 7.17 (d, $J=7.9\text{Hz}$, 1H),
 5 7.30 (d, $J=7.9\text{Hz}$, 1H), 7.47 (s, 1H)

[0181]

(Step 7)

{2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-cyclopentyl}methanol



10

6a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-4,4-dimethyl-hexahydro-cyclopenta[c]isoxazole (320 mg) was mixed in acetic acid (4.5 mL), tetrahydrofuran (1.5 mL), and water (1.5 mL) under nitrogen gas, and thereto was
 15 added zinc powder (640 mg) in two parts under heating at 60°C. The reaction solution was stirred under heating at 60°C for 1.5 hours. The reaction solution was filtered through Celite at room temperature, and then the filtrate was concentrated under reduced pressure. The resulting
 20 residue was mixed with chloroform (30 mL) and 28 w/w%


ammonia water (7.5 mL) and separated. The organic layer was washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (383 mg).

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.11 (s, 3H), 1.16 (s, 3H), 1.41-1.47 (m, 2H), 1.70-1.88 (m, 3H), 2.25-2.34 (m, 4H), 2.63-2.68 (m, 2H), 3.77 (dd, J=7.3, 1.7Hz, 2H), 7.17 (d, J=8.1Hz, 1H), 7.33 (dd, J=8.1, 2.1Hz, 1H), 7.47 (d, J=2.1 Hz, 1H)

[0182]

(Step 8)

Ethyl 3-(3-{1-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-2-hydroxymethyl-3,3-dimethyl-cyclopentyl}ureido)propionate



{2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-cyclopentyl}methanol (190 mg) and tetrahydrofuran (2.0 mL) were mixed under nitrogen gas, and thereto was added ethyl 3-isocyanato-propionate (0.076 mL). The reaction solution was stirred at room temperature for 1


hour. The reaction solution was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (10 g, ethyl acetate/chloroform = 2/3) to give the title compound (163 mg).

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.12 (s, 3H), 1.13 (s, 3H), 1.27 (t, J=7.2Hz, 3H), 1.43-1.48 (m, 2H), 1.65-1.78 (m, 2H), 1.98 (dd, J=6.7, 4.2Hz, 1H), 2.26 (dq, J=27.3, 7.1Hz, 2H), 2.47 (td, J=5.9, 1.7Hz, 2H), 2.61 -2.66 (m, 2H), 3.40 (ddd, J=12.0, 6.0, 1.2Hz, 2H), 3.78 (ddd, J=9.9, 5.6, 3.3Hz, 2H), 4.14 (q, J=7.2Hz, 2H), 4.87 (t, J=6.1Hz, 1H), 5.93 (s, 1H), 7.13 (d, J=8.1Hz, 1H), 7.27 (dd, J=8.1, 2.1Hz, 1H), 7.38 (d, J=2.1Hz, 1H)

[0183]

(Step 9)

Ethyl 3-{7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}propionate



Ethyl 3-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-hydroxymethyl-3,3-dimethyl-cyclopentyl}ureido)propionate and dichloromethane (1.5 mL) were mixed under nitrogen gas,

and thereto were added (diacetoxyiodo)benzene (109 mg) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (2.7 mg) at room temperature. The reaction solution was stirred at room temperature for 16 hours, and then thereto was added
5 trifluoroacetic acid (0.1 mL) at room temperature. The mixture was stirred at room temperature for 4 hours. To the reaction solution were added saturated aqueous sodium hydrogen carbonate solution (2 mL) and ethyl acetate, and the solution was separated. Thereto was added aqueous
10 sodium carbonate solution (30 mL), and the mixture was separated. The organic layer was dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by thin-layer silica gel
15 column chromatography (ethyl acetate/chloroform = 1/10) to give a racemate of the title compound. Then, the racemate was purified by preparative chiral column chromatography (IA, isopropanol/n-hexane = 7/93, 15 ml/min) to give the title compound (55.1 mg, 94.5%ee).


20 ¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.10 (s, 3H), 1.24 (s, 3H), 1.26 (t, J=6.2Hz, 3H), 1.34-1.41 (m, 1H), 1.42-1.47 (m, 2H), 1.59 (ddd, J=9.6, 3.6, 2.8Hz, 1H), 2.12 (td, J=11.8, 7.2Hz, 1H), 2.38 (ddd, J=12.3, 6.2, 3.1Hz, 1H), 2.56 (t, J=6.7Hz, 2H), 2.62-2.67 (m, 2H), 3.47 (dt, J=14.1, 6.9Hz,
25 1H), 3.95 (dt, J=14.0, 6.4Hz, 1H), 4.15 (ddd, J=14.2, 7.1,

2.3Hz, 2H), 5.09 (s, 1H), 6.14 (s, 1H), 7.11 (dd, J=7.9, 1.8Hz, 1H), 7.15 (d, J=7.9Hz, 1H), 7.27 (s, 1H)

[0184]

(Step 10)

5 3-{7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}propionate



Ethyl 3-{7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-

10 5,5-dimethyl-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}propionate (55 mg) and methanol (0.5 mL) were mixed under nitrogen gas, and thereto was added 2N aqueous sodium hydroxide solution (0.24 mL) under ice cooling. The reaction solution was stirred at room temperature for 4 hours. To the reaction solution was added 2N hydrochloric acid (0.24 mL) under ice cooling, and the mixture was extracted with ethyl acetate. The organic layer was dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give the title compound (44.0 mg).


[0185]

Example 7

(Step 1)

Ethyl (1,1-dimethylallyloxy)acetate

5




2-Methyl-3-buten-2-ol (4.6 mL) and rhodium (II) acetate (97 mg) were mixed under nitrogen gas, and thereto was added dropwise ethyl diazoacetate (4.6 mL) for 2 hours under water cooling. The mixture was stirred for 1 hour and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate/n-hexane (ethyl acetate/n-hexane = 1/5 -> 1/2) to give the title compound (3.65 g).

¹H-NMR (400 MHz, CDCl₃) 1.27 (t, J=7.17Hz, 3H), 1.32 (s, 6H), 3.95 (s, 2H), 4.20 (q, J=7.17Hz, 2H), 5.20-5.15 (m, 2H), 5.83 (dd, J=17.34, 10.87Hz, 1H)

[0186]

(Step 2)

20 (1,1-Dimethylallyloxy)acetaldehyde oxime




Ethyl (1,1-dimethylallyloxy)acetate (1.65 g) and toluene (150 mL) were mixed under nitrogen gas, and thereto was added dropwise 1M diisobutylaluminum hydride/toluene solution (14 mL) under cooling at -78°C. The mixture was stirred for 30 minutes, and then the reaction solution was added to 1N hydrochloric acid (15 mL) under ice cooling. The mixture was stirred for 1 hour and then separated. Hydroxylamine hydrochloride (910 mg) was mixed in ethanol (10 mL) and water (2.5 mL), and thereto was added dropwise 4N aqueous solution of sodium hydroxide (3.35 mL) for 20 minutes under ice cooling. The resulting organic layer was added thereto, and the mixture was stirred at room temperature for 1 hour. The organic layer was washed sequentially with water and saturated aqueous sodium chloride solution and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate/n-hexane (ethyl acetate/n-hexane = 1/3 -> 2/3) to give a geometric isomer mixture (1:4) of the title compound (774 mg).

¹H-NMR (400 MHz, CDCl₃) 1.31 (s, 6H), 3.96 (d, J=5.55Hz, 1.6H), 4.23 (d, J=3.47Hz, 0.4H), 5.15-5.19 (m, 2H), 5.77-5.86 (m, 1H), 6.87 (t, J=3.47Hz, 0.2H), 7.42 (br s, 0.8H), 7.46 (t, J=5.55Hz, 0.8H), 7.69 (br s, 0.2H)

[0187]

(Step 3)

4,4-Dimethyl-3a,4-dihydro-3H,6H-furan[3,4-c]isoxazole




(1,1-Dimethylallyloxy)acetaldehyde oxime (774 mg) and dichloromethane (40 mL) were mixed, and thereto was added triethylamine (65 mL) under ice cooling. Then, thereto was added dropwise antiformin (10 mL) for 15 minutes, and then the mixture was stirred for 30 minutes. Then, thereto was added chloroform. The organic layer was washed sequentially with water and saturated aqueous sodium chloride solution and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate/n-hexane (ethyl acetate/n-hexane = 1/2) to give the title compound (771 mg).

¹H-NMR (400 MHz, CDCl₃) 1.20 (s, 3H), 1.41 (s, 3H), 4.01-3.92 (m, 2H), 4.38 (dd, J=13.99, 1.27Hz, 1H), 4.45-4.50 (m, 2H)

[0188]

(Step 4)

6a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-4,4-dimethyl-tetrahydro-furan[3,4-c]isoxazole




4-Bromo-2-chloro-1-(3,3-dimethyl-butyl)-benzene (1.23 g) was mixed in tetrahydrofuran (5 mL) and toluene (12 mL), and then thereto was added dropwise 1.6M n-butyllithium/n-hexane solution (2.66 mL) under cooling at -78°C . The reaction solution was stirred at -78°C for 1 hour (Reaction solution A). 4,4-Dimethyl-3a,4-dihydro-3H,6H-furan[3,4-c]isoxazole (300 mg) and toluene (18 mL) were mixed, and thereto was added boron trifluoride-diethyl ether complex (0.537 mL) under cooling at -78°C . The mixture was stirred for 10 minutes. Then, thereto was added dropwise Reaction solution A under cooling at -78°C . The reaction solution was stirred for 1 hour, and then thereto was added an aqueous solution of ammonium chloride (8 mL) under cooling at -78°C . The mixture was extracted with ethyl acetate, and the organic layer was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate/n-hexane (ethyl acetate/n-hexane = 1/5 $\rightarrow$ 1/3) to give the title compound (439 mg).

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.38 (s, 3H), 1.42 (s, 3H), 1.43-1.47 (m, 2H), 2.65-2.69 (m, 2H), 3.04 (dd, J=6.94, 4.16Hz, 1H), 3.86-4.22 (m, 4H), 5.20 (br s, 1H), 7.21 (d, J=8.55Hz, 1H), 7.33-7.28 (m, 1H), 7.52-7.46 (m, 1H)

5 [0189]

(Step 5)

{4-Amino-4-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2,2-dimethyl-tetrahydro-furan-3-yl}methanol



10 6a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-4,4-dimethyl-tetrahydro-furan[3,4-c]isoxazole (411 mg) was mixed in acetic acid (6 mL), tetrahydrofuran (2 mL), and water (2 mL), and thereto was added zinc powder (800 mg) under heating at 60°C. The mixture was stirred with
 15 heating at 60°C for 3 hours, and then ammonia water (10 mL) was added dropwise to the reaction solution at room temperature. The reaction solution was extracted with chloroform (15 mL, three times), and the organic layers were dried over sodium sulfate. Sodium sulfate was removed
 20 with a filter, and then the filtrate was concentrated under


reduced pressure to give a crude product of the title compound (229 mg).

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.36 (s, 3H), 1.41 (s, 3H), 1.43-1.47 (m, 2H), 2.19 (t, J=6.01Hz, 1H), 2.65-2.69 (m, 2H), 3.68 (d, J=9.25Hz, 1H), 3.83 (d, J=6.01Hz, 2H), 4.11 (d, J=9.25Hz, 1H), 7.20 (d, J=8.09Hz, 1H), 7.34 (dd, J=8.09, 2.08Hz, 1H), 7.49 (d, J=2.08Hz, 1H)

[0190]

(Step 6)

10 Ethyl 4-(3-{3-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-4-hydroxymethyl-5,5-dimethyl-tetrahydro-furan-3-yl}ureido)benzoate



15 {4-Amino-4-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2,2-dimethyl-tetrahydro-furan-3-yl}methanol (125 mg) and tetrahydrofuran (2.5 mL) were mixed, and thereto was added 4-ethoxycarbonylphenyl isocyanate (70 mg) under ice cooling. The mixture was stirred at room temperature for 51 minutes, and then the reaction solution was concentrated under


reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate/n-hexane (ethyl acetate/n-hexane = 1/10 -> 1/6 -> 1/2 -> 1/1) to give the title compound (142 mg).

5 ¹H-NMR (400 MHz, CDCl₃) 0.96 (s, 9H), 1.28 (s, 3H), 1.37 (t, J=7.00Hz, 3H), 1.40 (s, 3H), 1.41-1.44 (m, 2H), 2.61-2.65 (m, 2H), 2.85 (dd, J=9.66, 4.35Hz, 1H), 3.34 (br s, 1H), 3.75-3.87 (m, 2H), 3.90 (d, J=10.14Hz, 1H), 4.00 (d, J=10.14Hz, 1H), 4.34 (q, J=7.00Hz, 2H), 6.34 (br s, 1H),
10 7.08 (br s, 1H), 7.16 (d, J=8.21Hz, 1H), 7.32 (d, J=8.69Hz, 2H), 7.37 (dd, J=8.21, 2.17Hz, 1H), 7.52 (d, J=2.17Hz, 1H), 7.94 (d, J=8.69Hz, 2H)

[0191]

(Step 7)

15 Ethyl 4-{(S)-7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-2-oxo-1,2,7,7a-tetrahydro-5H-furan[3,4-d]pyrimidin-3-yl}benzoate



Ethyl 4-(3-{3-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-

20 4-hydroxymethyl-5,5-dimethyl-tetrahydro-furan-3-

yl}ureido)benzoate (124 mg) and chloroform (3 mL) were mixed, and thereto were added (diacetoxyiodo)benzene (75 mg) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (3.6 mg) at room temperature. The mixture was stirred at room temperature for 7 hours, and then thereto was added an aqueous solution of sodium thiosulfate under ice cooling. The mixture was extracted with chloroform, and the organic layer was dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was mixed in cyclopentyl methyl ether (5 mL), and thereto was added 4N hydrogen chloride/cyclopentyl methyl ether solution (0.137 mL) under ice cooling. The mixture was stirred under ice cooling for 2 hours and then stirred at room temperature for 1 hour. The mixture was stirred under heating at 40°C for 1 hour, and then the reaction solution was concentrated under reduced pressure. The resulting residue was purified by thin-layer silica gel column chromatography (ethyl acetate/n-hexane = 1/1) to give a racemate of the title compound (73.2 mg). The racemate was purified with a Recycling Preparative Liquid Chromatograph to give a single enantiomer of the title compound (30.8 mg).

Purification conditions for the preparative chromatography are shown as follows.

Preparative apparatus: Recycling preparative liquid

chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

Column: Daicel CHIRALPAK IA 2.0 cm ϕ x 25 cm

Mobile phase: n-hexane:2-propanol = 80:20

5 Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement of the resulting compound with a chiral column showed 6.5 minutes of the retention time for the resulting enantiomer with > 99%ee of optical purity. The
10 retention time for the opposite enantiomer on the phenyl group was 8.8 minutes.

Analytical conditions for the chiral column are shown as follows.

Measurement apparatus: HPLC system, Shimadzu Corporation,
15 High-Performance Liquid Chromatograph Prominence

Column: Daicel CHIRALPAK IA-3 0.46 cm ϕ x 15 cm

Column temperature: 30°C

Mobile phase: hexane:2-propanol = 80:20

Flow rate: 1.0 mL/min

20 Detection: UV (254 nm)

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.33 (s, 3H), 1.40 (t, J=7.12Hz, 3H), 1.42-1.46 (m, 2H), 1.50 (s, 3H), 2.65-2.69 (m, 2H), 4.09 (d, J=8.69Hz, 1H), 4.38 (q, J=7.12Hz, 2H), 4.51 (d, J=8.69Hz, 1H), 5.35 (s, 1H), 6.34 (s, 1H), 7.22 (d, J=7.97Hz, 1H), 7.30 (dd, J=7.97, 1.93Hz, 1H), 7.45 (d,


25

$J=9.02\text{Hz}$, 2H), 7.49 (d, $J=1.93\text{Hz}$, 1H), 8.07 (d, $J=9.02\text{Hz}$, 2H)

[0192]

(Step 8)

5 4-{(S)-7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-2-oxo-1,2,7,7a-tetrahydro-5H-furan[3,4-d]pyrimidin-3-yl}benzoic acid



Ethyl

4-{(S)-7a-[3-chloro-4-(3,3-dimethyl-


10 butyl)phenyl]-5,5-dimethyl-2-oxo-1,2,7,7a-tetrahydro-5H-furan[3,4-d]pyrimidin-3-yl}benzoate (28.0 mg) and ethanol (1 mL) were mixed, and thereto was added 2N aqueous solution of sodium hydroxide (0.110 mL) at room temperature. The reaction solution was stirred at room temperature for 4
15 hours 30 minutes, and then thereto was added 2N hydrochloric acid (0.136 mL). The precipitated solid was collected by filtration to give the title compound (22.1 mg).

[0193]

Example 10

(Step 1)

Ethyl 4-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-hydroxymethyl-3,3-dimethyl-cyclopentyl}ureido)benzoate



5

{2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-cyclopentyl}methanol (166 mg) was mixed with tetrahydrofuran (2 mL) under nitrogen gas, and thereto was added ethyl 4-isocyanatobenzoate (94 mg) under ice cooling.

10 The reaction solution was stirred at room temperature for 1 hour. The reaction solution was concentrated, and then purified by silica gel column chromatography (eluent: ethyl acetate/n-hexane (ethyl acetate/n-hexane = 25/1 -> 1/1) to give the title compound (144 mg).


15 ¹H-NMR (400 MHz, CDCl₃) 0.97 (s, 9H), 1.13 (s, 3H), 1.21 (s, 3H), 1.33-1.39 (m, 3H), 1.40-1.46 (m, 2H), 1.69-1.88 (m, 3H), 1.91-1.99 (m, 1H), 2.38-2.48 (m, 1H), 2.53-2.66 (m, 3H), 3.80-3.95 (m, 2H), 4.30-4.37 (m, 2H), 6.45-6.49 (m, 1H), 6.88-6.96 (m, 1H), 7.13-7.17 (m, 1H), 7.26-7.29 (m,

1H), 7.32-7.42 (m, 3H), 7.90-7.96 (m, 2H)

[0194]

(Step 2)

Ethyl 4-{7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}benzoate



Ethyl 4-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-hydroxymethyl-3,3-dimethyl-cyclopentyl}ureido)benzoate
 (135 mg) and dichloromethane (5 mL) were mixed under
 nitrogen gas, and thereto were added (diacetoxyiodo)benzene
 (82 mg) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (4
 mg) at room temperature. The reaction solution was stirred
 at room temperature for 3.5 hours, and then thereto was
 added trifluoroacetic acid (78 μ L) at room temperature.
 The mixture was stirred at room temperature overnight, and
 then thereto were added aqueous sodium thiosulfate solution
 and chloroform under ice cooling. The mixture was
 separated. The organic layer was dried over sodium sulfate.
 Sodium sulfate was removed with a filter, and then the

filtrate was concentrated under reduced pressure. The resulting residue was purified by thin-layer silica gel column chromatography (eluent: ethyl acetate/n-hexane = 1/1, R_f = 0.5) to give a racemate of the title compound (91 mg).

5 The racemate was purified with a Recycling Preparative Liquid Chromatograph to give a single enantiomer of the title compound (23 mg).

Purification conditions for the preparative chromatography are shown as follows.

10 Preparative apparatus: Recycling preparative liquid chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

Column: Daicel CHIRALPAK IA 2.0 cm ϕ x 25 cm

Mobile phase: n-hexane/2-propanol = 80/20

15 Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement of the resulting compound with a chiral column showed 12.8 minutes of the retention time for the resulting enantiomer with > 99%ee of optical purity. The
20 retention time for the opposite enantiomer was 10.6 minutes.

Analytical conditions for the chiral column are shown as follows.

Measurement apparatus: HPLC system, Shimadzu Corporation, High-Performance Liquid Chromatograph Prominence

25 Column: Daicel CHIRALPAK IA-3 0.46 cm ϕ x 15 cm

Column temperature: 40°C

Mobile phase: n-hexane/2-propanol = 90/10

Flow rate: 1.0 mL/min


Detection: UV (254 nm)

5 ¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.17 (s, 3H), 1.30 (s, 3H), 1.36-1.41 (m, 3H), 1.41-1.47 (m, 2H), 1.64-1.71 (m, 1H), 2.18-2.27 (m, 1H), 2.43-2.51 (m, 1H), 2.62-2.69 (m, 2H), 4.33-4.40 (m, 2H), 5.36 (br s, 1H), 6.36 (s, 1H), 7.17-7.24 (m, 2H), 7.37-7.39 (m, 1H), 7.41-7.47 (m, 2H),
10 8.01-8.06 (m, 2H)

[0195]

(Step 3)

4-{7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}benzoic acid
15



Ethyl 4-{7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-dimethyl-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}benzoate (21 mg) and ethanol (1


mL) were mixed under nitrogen gas, and thereto was added 2N aqueous sodium hydroxide solution (82 μ L) at room temperature. The reaction solution was stirred under heating at 60°C for 1 hour, and then concentrated under reduced pressure. To The resulting residue were added 2N hydrochloric acid and water at room temperature, and the resulting slurry was stirred at room temperature. The precipitated solid was collected by filtration to give the title compound (16 mg).

10 [0196]

Example 27

(Step 1)

Ethyl 6-(tert-butyl-dimethyl-silanyloxy)-3-vinyl-hexanoate



15 (E)-6-(tert-butyl-dimethyl-silanyloxy)hex-2-en-1-ol

(10.1 g) and triethyl orthoformate (102 mL) were mixed, and thereto was added propionate (51 mL). The reaction solution was stirred under heating at 150°C for 3.5 hours. The reaction solution was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, R_f = 0.33 (ethyl acetate/n-


hexane = 5/95)) to give the title compound (13.2 g).

¹H-NMR (400 MHz, CDCl₃) 0.04 (s, 6H), 0.89 (s, 9H), 1.24 (t, J=6.94Hz, 3H), 1.29-1.38 (m, 1H), 1.42-1.60 (m, 3H), 2.28 (dd, J=14.57, 8.55Hz, 1H), 2.36 (dd, J=14.57, 6.24Hz, 1H),
 5 2.48-2.57 (m, 1H), 3.59 (t, J=6.47Hz, 2H), 4.11 (q, J=7.17Hz, 2H), 4.99-5.06 (m, 2H), 5.62 (ddd, J=17.34, 10.40, 8.32Hz, 1H)

[0197]

(Step 2)

10 6-(tert-Butyl-dimethyl-silanyloxy)-3-vinyl-hexan-1-ol



Ethyl 6-(tert-butyl-dimethyl-silanyloxy)-3-vinyl-


hexanoate (11.5 g) and tetrahydrofuran (231 mL) were mixed under argon gas, and thereto was added dropwise 1M
 15 diisobutylaluminum hydride/toluene solution (92.2 mL) under cooling at -78°C. The reaction solution was stirred under ice cooling for 1 hour. To the reaction solution was added dropwise an aqueous solution of Rochelle salt (231 mL) under ice cooling. The reaction solution was stirred at
 20 room temperature for 16 hours. To the reaction solution was added ethyl acetate, and the solution was separated. The aqueous layer was extracted with ethyl acetate.

Combined organic layers were washed with brine, and then dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then concentrated under reduced pressure. The resulting residue was azeotroped with
 5 toluene to give a crude product of the title compound (11.3 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.04 (s, 6H), 0.89 (s, 9H), 1.24-1.34 (m, 2H), 1.40-1.61 (m, 3H), 1.64-1.72 (m, 1H), 2.10-2.19 (m, 1H), 3.59 (t, $J=6.47\text{Hz}$, 2H), 3.62-3.72 (m, 2H),
 10 4.99-5.04 (m, 2H), 5.57 (ddd, $J=18.73, 9.71, 9.02\text{Hz}$, 1H)
 [0198]

(Step 3)

[4-(2-Benzyloxy-ethyl)hex-5-enyloxy]-tert-butyl-dimethyl-silane



Sodium hydride (2.73 g) and dimethylformamide (125 mL) were mixed under argon gas, and thereto was added dropwise a mixed solution of 6-(tert-butyl-dimethyl-silanyloxy)-3-vinyl-hexan-1-ol (12.6 g) in dimethylformamide (60 mL)
 20 under ice cooling. The reaction solution was stirred under ice cooling for 1 hour. To the reaction solution was added dropwise a mixed solution of benzyl bromide (8.41 mL) in dimethylformamide (60 mL) under ice cooling. The reaction


solution was stirred at room temperature overnight. To the reaction solution were added aqueous ammonium chloride solution (125 mL), water (60 mL), and ethyl acetate/n-hexane = 1/3, and the solution was separated. The aqueous layer was extracted with ethyl acetate/n-hexane = 1/3. Combined organic layers were washed with water (three times) and brine, and then dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, R_f = 0.36 (ethyl acetate/n-hexane = 10/90)) to give the title compound (19.5 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.04 (s, 6H), 0.89 (s, 9H), 1.21-1.31 (m, 2H), 1.38-1.60 (m, 3H), 1.70-1.78 (m, 1H), 2.10-2.20 (m, 1H), 3.41-3.51 (m, 2H), 3.58 (t, $J=6.24\text{Hz}$, 2H), 4.46 (d, $J=11.79\text{Hz}$, 1H), 4.49 (d, $J=11.79\text{Hz}$, 1H), 4.93-4.99 (m, 2H), 5.52 (ddd, $J=17.11, 10.40, 9.02\text{Hz}$, 1H), 7.27-7.37 (m, 5H)

[0199]

(Step 4)

4-(2-Benzyloxy-ethyl)hex-5-en-1-ol




[4-(2-Benzyloxy-ethyl)hex-5-enyloxy]-tert-butyl-dimethyl-silane (12.88 g) and tetrahydrofuran (130 mL) were mixed under argon gas, and thereto was added dropwise 1M tetra-n-butylammonium fluoride/tetrahydrofuran solution
5 (55.4 mL) under ice cooling. The reaction solution was stirred at room temperature for 19.5 hours. To the reaction solution were added water (100 mL) and ethyl acetate, and the solution was separated. The aqueous layer was extracted with ethyl acetate (three times). Combined
10 organic layers were washed with brine, and then dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent:
15 ethyl acetate/n-hexane) to give the title compound (8.43 g).
¹H-NMR (400 MHz, CDCl₃) 1.35-1.21 (m, 2H), 1.42-1.67 (m, 3H), 1.70-1.79 (m, 1H), 2.13-2.22 (m, 1H), 3.42-3.52 (m, 2H), 3.60-3.64 (m, 2H), 4.46 (d, J=12.02Hz, 1H), 4.49 (d, J=12.02Hz, 1H), 4.94-5.01 (m, 2H), 5.52 (ddd, J=16.88,
20 10.17, 8.79Hz, 1H), 7.27-7.38 (m, 5H)

[0200]

(Step 5)

4-(2-Benzyloxy-ethyl)hex-5-enal




4-(2-Benzyloxy-ethyl)hex-5-en-1-ol (7.14 g) and (diacetoxyiodo)benzene (10.7 g) were mixed in chloroform (72 mL) under nitrogen gas, and thereto was added 2,2,6,6-tetramethylpiperidin-1-oxyl radical (0.477 g) at room temperature. The reaction solution was stirred at room temperature for 22 hours. To the reaction solution were added saturated aqueous sodium hydrogen carbonate solution (20 mL), water (10 mL), sodium thiosulfate (0.723 g), and ethyl acetate, and the solution was separated. The aqueous layer was extracted with ethyl acetate. Combined organic layers were washed with brine, and then dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, R_f = 0.50 (ethyl acetate/n-hexane = 20/80)) to give the title compound (7.24 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.50-1.60 (m, 2H), 1.71-1.81 (m, 2H), 2.13-2.23 (m, 1H), 2.35-2.51 (m, 2H), 3.42-3.52 (m, 2H), 4.46 (d, $J=11.79\text{Hz}$, 1H), 4.49 (d, $J=11.79\text{Hz}$, 1H), 4.96-5.05 (m, 2H), 5.47 (ddd, $J=17.11, 10.17, 9.02\text{Hz}$, 1H), 7.27-7.36 (m, 5H), 9.75 (t, $J=1.39\text{Hz}$, 1H)

[0201]

(Step 6)

4-(2-Benzyloxy-ethyl)hex-5-enal oxime



5 4-(2-Benzyloxy-ethyl)hex-5-enal (5.94 g), ethanol (51.2 mL), and water (25.6 mL) were mixed under nitrogen gas, and thereto were added sodium acetate (15.4 g) and hydroxylamine hydrochloride (7.78 g) at room temperature. The reaction solution was stirred under heating at 60°C for

10 21.5 hours. The reaction solution was concentrated under reduced pressure, and thereto was added toluene. The solution was separated. The aqueous layer was extracted with toluene (twice). Combined organic layers were washed with brine (twice), and then dried over sodium sulfate.

15 Sodium sulfate was removed with a filter, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, R_f = 0.27 (ethyl acetate/n-hexane = 20/80)) to give the

20 title compound (7.38 g).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.38-1.48 (m, 1H), 1.50-1.64 (m, 2H), 1.70-1.80 (m, 1H), 2.09-2.28 (m, 2H), 2.30-2.44 (m, 1H), 3.41-3.52 (m, 2H), 4.46 (d, $J=12.95\text{Hz}$, 1H), 4.49 (d,

$J=12.95\text{Hz}$, 1H), 4.97-5.05 (m, 2H), 5.56-5.45 (m, 1H), 6.70 (t, $J=5.55\text{Hz}$, 0.5H), 7.27-7.36 (m, 5H), 7.41 (t, $J=6.01\text{Hz}$, 0.5H)

[0202]

5 (Step 7)

4-(2-Benzyloxy-ethyl)-3a,4,5,6-tetrahydro-3H-cyclopenta[c]isoxazole



4-(2-Benzyloxy-ethyl)hex-5-enal oxime (5.90 g) and
 10 triethylamine (2.04 mL) were mixed in dichloromethane (118 mL), and then thereto was added a mixed aqueous solution of sodium hypochlorite 5-hydrate (3.61 g) in water (53.1 mL) under water cooling. The reaction solution was stirred at room temperature for 3.5 hours. To the reaction solution
 15 was added a mixed solution of sodium hypochlorite 5-hydrate (4.95 g) in water (53.1 mL) under water cooling. The reaction solution was stirred at room temperature for 17 hours. To the reaction solution was added ethyl acetate, and the solution was separated. The aqueous layer was
 20 extracted with ethyl acetate. Combined organic layers were washed with water (twice) and brine, and then dried over sodium sulfate. Sodium sulfate was removed with a filter, and then concentrated under reduced pressure. The


resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, R_f = 0.27 (ethyl acetate/n-hexane = 30/70)) and (Biotage flash purification systems, eluent: acetone/chloroform) to give the title compound (2.34 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.66-1.57 (m, 1H), 1.77-2.06 (m, 3H), 2.33-2.56 (m, 3H), 3.35-3.50 (m, 3H), 3.78 (dd, $J=12.26$, 8.07Hz, 1H), 4.41 (dd, $J=9.57$, 8.07Hz, 1H), 4.44 (d, $J=11.66\text{Hz}$, 1H), 4.48 (d, $J=11.66\text{Hz}$, 1H), 7.28-7.37 (m, 5H)

[0203]

(Step 8)

4-(2-Benzyloxy-ethyl)-6a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]hexahydro-cyclopenta[c]isoxazole




This step was performed according to Example 3 Step 6.

[0204]

(Step 9)

{2-Amino-5-(2-benzyloxy-ethyl)-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]cyclopentyl}methanol




This step was performed according to Example 3 Step 7.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.98 (s, 9H), 1.42-1.77 (m, 6H),
 1.88-1.96 (m, 1H), 2.02-2.12 (m, 1H), 2.25-2.37 (m, 2H),
 2.64-2.69 (m, 2H), 3.53-3.59 (m, 2H), 3.64 (dd, $J=11.91$,
 4.97Hz, 1H), 3.72 (dd, $J=11.91$, 3.12Hz, 1H), 3.77-3.82 (m,
 2H), 4.49 (d, $J=11.79\text{Hz}$, 1H), 4.53 (d, $J=11.79\text{Hz}$, 1H), 7.18
 (d, $J=8.09\text{Hz}$, 1H), 7.28-7.35 (m, 6H), 7.44 (d, $J=2.08\text{Hz}$,
 1H)

[0205]

(Step 10)

Ethyl 3-(3-{3-(2-benzyloxy-ethyl)-1-[3-chloro-4-(3,3-
 dimethyl-butyl)phenyl]-2-hydroxymethyl-
 cyclopentyl}ureido)propionate



This step was performed according to Example 3 Step 8.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.97 (s, 9H), 1.25 (t, $J=7.17\text{Hz}$,

3H), 1.42-1.68 (m, 6H), 1.77-1.85 (m, 1H), 2.08 -2.15 (m, 1H), 2.27 -2.41 (m, 2H), 2.45-2.50 (m, 2H), 2.60-2.65 (m, 2H), 3.34-3.42 (m, 2H), 3.52-3.68 (m, 4H), 4.12 (q, J=7.17Hz, 2H), 4.52 (s, 2H), 4.69 (t, J=5.53Hz, 1H), 6.52 (s, 1H), 7.13 (d, J=7.92Hz, 1H), 7.19 (dd, J=7.92, 1.94Hz, 1H), 7.29 -7.38 (m, 6H)

[0206]

(Step 11)

Ethyl 3-{5-(2-benzyloxy-ethyl)-7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}propionate




This step was performed according to Example 3 Step 9.

[0207]

(Step 12)

Ethyl 3-{7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5-(2-hydroxy-ethyl)-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}propionate



Ethyl 3-{5-(2-benzyloxy-ethyl)-7a-[3-chloro-4-(3,3-
 dimethyl-butyl)phenyl]-2-oxo-1,2,5,6,7,7a-hexahydro-
 cyclopentapyrimidin-3-yl}propionate (0.100 g) and
 5 dichloromethane (2 mL) were mixed under nitrogen gas, and
 thereto was added dropwise 1.01M boron
 tribromide/dichloromethane solution (0.42 mL) under cooling
 at -78°C. The reaction solution was stirred under cooling
 at -78°C for 1 hour. To the reaction solution was added a
 10 mixed solution of triethylamine (0.42 mL) in methanol (0.42
 mL) under cooling at -78°C. The reaction solution was
 stirred at room temperature for 10 minutes. To the
 reaction solution was added dropwise water (0.5 mL). To
 the reaction solution was added ethyl acetate, and the
 15 solution was separated. The aqueous layer was extracted
 with ethyl acetate. Combined organic layers were washed
 with brine, and then dried over sodium sulfate. Sodium
 sulfate was removed with a filter, and then concentrated
 under reduced pressure. The resulting residue was purified
 20 by silica gel column chromatography (Biotage flash
 purification systems, eluent: ethyl acetate/n-hexane, Rf =


0.07 (ethyl acetate/n-hexane = 2/3)) and (Biotage flash purification systems, eluent: acetone/chloroform, R_f = 0.43 (acetone/chloroform = 3/7), followed by purification by thin-layer silica gel column chromatography (acetone/chloroform = 1/4) to give the title compound (18.8 mg).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.98 (s, 9H), 1.14-1.08 (m, 1H), 1.18-1.33 (m, 5H), 1.42-1.46 (m, 2H), 1.65-1.73 (m, 1H), 1.97-2.06 (m, 2H), 2.45-2.49 (m, 1H), 2.53-2.61 (m, 2H), 2.62-2.67 (m, 2H), 2.91-2.99 (m, 1H), 3.45-3.53 (m, 1H), 3.64-3.73 (m, 2H), 3.89-3.96 (m, 1H), 4.08-4.19 (m, 2H), 5.11 (s, 1H), 6.25 (d, $J=1.85\text{Hz}$, 1H), 7.06-7.27 (m, 3H)

[0208]

(Step 13)

Ethyl 3-{7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5-(2-methoxy-ethyl)-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}propionate



Ethyl 3-{7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5-(2-hydroxy-ethyl)-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}propionate (52 mg) and


acetonitrile (0.75 mL) were mixed under nitrogen gas, and thereto were added methyl iodide (0.75 mL) and silver (I) oxide (38 mg) at room temperature. The reaction solution was stirred under heating at 80°C. The resulting solid was removed with Celite, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by thin-layer silica gel column chromatography (ethyl acetate/n-hexane = 1/1) and thin-layer silica gel column chromatography (acetone/chloroform = 7/93) to give the title compound (15.5 mg).

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.04-1.14 (m, 1H), 1.25 (t, J=7.17Hz, 3H), 1.41-1.46 (m, 2H), 1.50-1.59 (m, 2H), 1.66-1.74 (m, 1H), 1.94-2.06 (m, 2H), 2.42-2.47 (m, 1H), 2.55-2.59 (m, 2H), 2.62-2.66 (m, 2H), 2.88-2.95 (m, 1H), 3.34 (s, 3H), 3.35-3.53 (m, 3H), 3.89-3.95 (m, 1H), 4.12-4.19 (m, 2H), 5.20 (s, 1H), 6.21 (d, J=1.85Hz, 1H), 7.06-7.24 (m, 3H)

[0209]

(Step 14)

3-{7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5-(2-methoxyethyl)-2-oxo-1,2,5,6,7,7a-hexahydro-cyclopentapyrimidin-3-yl}propionate




This step was performed according to Example 3 Step 10.

[0210]

Example 34

5 (Step 1)

2,2,3-Trimethylpent-4-en-1-ol



Lithium aluminum hydride (4.0 g) and tetrahydrofuran (210 mL) were mixed under argon gas, and thereto was added a mixed solution of 2,2,3-trimethylpent-4-enoic acid (5.0 g) in tetrahydrofuran (70 mL) under ice cooling. The mixture was stirred under ice cooling for 20 minutes, and then the reaction solution was stirred with being heated to reflux for 2 hours 30 minutes. Then, thereto were added dropwise sequentially water (4 mL), 2N sodium hydroxide (4 mL), and water (12 mL) under ice cooling. The reaction solution was stirred at room temperature for 1 hour. To the reaction solution were added Celite (4 g) and magnesium sulfate (4 g), and Celite was removed with a filter, and


then the filtrate was concentrated under reduced pressure to give the title compound (4.2 g).

¹H-NMR (400MHz, CDCl₃) 0.84 (s, 3H), 0.87 (s, 3H), 0.97 (d, J=6.94Hz, 3H), 1.37 (br s, 1H), 2.13-2.21 (m, 1H), 3.37 (s, 2H), 4.97-5.05 (m, 2H), 5.82 (ddd, J=18.15, 9.25, 7.86Hz, 1H)

[02111]

(Step 2)

2,2,3-Trimethyl-pent-4-en-1-ol methanesulfonic acid ester



2,2,3-Trimethylpent-4-en-1-ol (4.2 g) and

triethylamine (7.2 mL) were mixed in chloroform (50 mL) under argon gas, and thereto was added dropwise a mixed solution of methanesulfonyl chloride (3.8 mL) in chloroform (15 mL) under ice cooling. The reaction solution was stirred under ice cooling for 5 minutes, and then warmed to room temperature. The reaction solution was stirred at room temperature for 3 hours, and then thereto was added water (13 mL). The solution was stirred at room temperature, and then thereto was added ethyl acetate (20 mL). The solution was separated. The aqueous layer was extracted with ethyl acetate (20 mL) once. The organic layer was washed with 1N hydrochloric acid (12 mL), saturated sodium hydrogen carbonate (12 mL), and saturated

15

20


aqueous sodium chloride solution (12 mL) and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give the title compound (7.1 g).

5 ^1H -NMR (400MHz, CDCl_3) 0.92 (s, 3H), 0.96 (s, 3H), 0.99 (d, $J=6.88\text{Hz}$, 3H), 2.16-2.24 (m, 1H), 3.00 (s, 3H), 3.94 (s, 2H), 5.01-5.08 (m, 2H), 5.67-5.78 (m, 1H)

[0212]

(Step 3)

10 5-Iodo-3,4,4-trimethylpent-1-ene



2,2,3-Trimethyl-pent-4-en-1-ol methanesulfonic acid ester (7.1 g) and N-methylpyridone (65 mL) were mixed under nitrogen gas, and thereto was added sodium iodide (24 g) at room temperature. The reaction solution was heated at 140°C for 4 hours 20 minutes. Then, thereto was added 20w/w% aqueous sodium thiosulfate solution (31 mL) at room temperature, and the mixture was stirred for 1 hour. The reaction solution was extracted with ethyl acetate/n-hexane = 1/2 (50 mL) twice. The organic layer was washed with water (21 mL) twice, 20w/w% aqueous sodium thiosulfate solution (21 mL), and saturated aqueous sodium chloride solution (21 mL) and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the


filtrate was concentrated under reduced pressure to give the title compound (5.4 g).

^1H -NMR (400MHz, CDCl_3) 0.95-0.99 (m, 9H), 2.20-2.28 (m, 1H), 3.16-3.24 (m, 2H), 5.00-5.11 (m, 2H), 5.70 (ddd, $J=18.03$, 9.25, 7.86Hz, 1H)

[0213]

(Step 4)

3,3,4-Trimethylhex-5-enenitrile



5-Iodo-3,4,4-trimethylpent-1-ene (3.8 g) and dimethyl sulfoxide (42 mL) were mixed under argon gas, and thereto was added tetraethylammonium cyanide (6.8 g) at room temperature. The reaction solution was stirred under heating at 80°C for 7 hours. To the reaction solution was added water (21 mL) at room temperature, and the aqueous layer was extracted with a mixed solvent of ethyl acetate/n-hexane = 1/2 (21 mL, twice). The organic layer was washed with water (12 mL, three times) and saturated aqueous sodium chloride solution (12 mL) and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was combined with a crude product (0.5 g) synthesized separately according to Example 34 Step 4 and purified by silica gel column


chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, 5/95 to 30/70) and then azeotroped with n-hexane to give the title compound (2.0 g).

¹H-NMR (400MHz, CDCl₃) 0.97-1.02 (m, 6H), 1.06 (s, 3H),
 5 2.13-2.22 (m, 1H), 2.24 (s, 2H), 5.04-5.12 (m, 2H), 5.69
 (ddd, J=18.15, 9.13, 7.98Hz, 1H)

[0214]

(Step 5)

3,3,4-Trimethylhex-5-enal



3,3,4-Trimethylhex-5-enenitrile (2.0 g) and dichloromethane (150 mL) were mixed under argon gas, and thereto was added dropwise 1.02M diisobutylaluminum hydride/n-hexane solution (22 mL) under cooling at -78°C.
 15 The reaction solution was stirred under cooling at -78°C for 15 minutes. The reaction solution was gradually warmed to 0°C and stirred under ice cooling for 2 hours. Then, thereto was added dropwise a saturated aqueous solution of Rochelle salt (150 mL) under ice cooling. The reaction
 20 solution was stirred at room temperature for 2 hours. The aqueous layer was extracted with ethyl acetate/n-hexane = 1/2 (20 mL, twice). The organic layer was washed with saturated aqueous sodium chloride solution (6 mL) and dried


over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give the title compound (1.8 g).

¹H-NMR (400MHz, CDCl₃) 0.99 (d, J=5.78Hz, 3H), 1.03 (s, 3H),
 5 1.05 (s, 3H), 2.06-2.15 (m, 1H), 2.27-2.30 (m, 2H), 4.98-
 5.05 (m, 2H), 5.74 (ddd, J=18.09, 9.19, 7.69Hz, 1H), 9.86
 (t, J=3.12 Hz, 1H)

[0215]

(Step 6)

10 3,3,4-Trimethylhex-5-enal oxime



3,3,4-Trimethylhex-5-enal (1.8 g) was mixed in ethanol (30 mL) and water (15 mL) under argon gas, and thereto were added sodium acetate (7.4 g) and hydroxylamine
 15 hydrochloride (3.1 g) at room temperature. The reaction solution was stirred under heating at 60°C for 1 day. The reaction solution was concentrated under reduced pressure, and thereto were added ethyl acetate and water. The solution was separated. The aqueous layer was extracted
 20 with ethyl acetate. The organic layer was washed with saturated aqueous sodium chloride solution (twice) and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under

reduced pressure. The resulting residue was mixed with ethyl acetate, and the resulting solid was filtered with Celite. Then, the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel

5 column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, 5/95 to 20/80) to give the title compound (1.7 g).


¹H-NMR (400MHz, CDCl₃) 0.90 (s, 1.5H), 0.91 (s, 1.5H), 0.93 (s, 1.5H), 0.94 (s, 1.5H), 0.97 (d, J=4.39Hz, 1.5H), 0.99

10 (d, J=4.39Hz, 1.5H), 1.98-2.09 (m, 1H), 2.12 (dd, J=6.82, 2.89Hz, 1H), 2.29 (dd, J=15.67, 5.55Hz, 0.5H), 2.38 (dd, J=15.67, 5.90Hz, 0.5H), 4.96 -5.03 (m, 2H), 5.72-5.82 (m, 1H), 6.83 (t, J=5.90Hz, 0.5H), 6.91 (br s, 1H), 7.48 (t, J=6.82 Hz, 0.5H)

15 [0216]

(Step 7)

4,5,5-Trimethyl-3a,4,5,6-tetrahydro-3H-cyclopenta[c]isoxazole



20 Heptan-6-al oxime (1.7 g) and methanol (34 mL) were mixed under nitrogen gas, and thereto was added trifluoroacetic acid (0.25 mL) under sodium chloride-ice cooling, followed by addition of (diacetoxyiodo)benzene


(5.1 g) over 40 minutes. The reaction solution was stirred under ice cooling for 20 minutes and at room temperature for 35 minutes. Then, thereto were added saturated aqueous sodium hydrogen carbonate solution (17 mL) and sodium sulfite (0.75 g) under ice cooling, and the mixture was stirred at room temperature for 30 minutes. The reaction solution was concentrated under reduced pressure. The aqueous layer was extracted with ethyl acetate (10 mL, twice). The organic layer was washed with saturated aqueous sodium chloride solution (5 mL, twice) and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane = 5/95 to 30/70), followed by further purification by silica gel column chromatography (Biotage flash purification systems, eluent: acetone/n-hexane = 5/95 to 20/80), and then azeotroped with n-hexane to give the title compound (0.87 g).

¹H-NMR (400MHz, CDCl₃) 0.96 (d, J=6.94Hz, 3H), 1.02 (s, 3H), 1.12 (s, 3H), 1.55-1.64 (m, 1H), 2.35-2.37 (m, 2H), 3.51-3.62 (m, 1H), 3.78 (dd, J=12.02, 7.86Hz, 1H), 4.48 (dd, J=9.48, 7.86Hz, 1H)

[0217]

(Step 8)

6a-(3-Chloro-4-(3,3-dimethylbutyl)phenyl)-4,5,5-trimethylhexahydro-1H-cyclopenta[c]isoxazole




5 This step was performed according to Example 3 Step 6.

¹H-NMR (400MHz, CDCl₃) 0.57-0.69 (m, 3H), 0.94-1.01 (m, 15H), 1.42-1.49 (m, 2H), 1.59-1.68 (m, 1H), 1.93-1.99 (m, 1H), 2.33-2.41 (m, 1H), 2.64-2.68 (m, 2H), 2.85-2.91 (m, 1H), 3.73-3.79 (m, 1H), 4.00-4.04 (m, 1H), 7.18 (d, 10 J=8.09Hz, 1H), 7.26-7.30 (m, 1H), 7.42-7.44 (m, 1H)

[0218]

(Step 9)

(2-Amino-2-(3-chloro-4-(3,3-dimethylbutyl)phenyl)-4,4,5-trimethylcyclopentyl)methanol




This step was performed according to Example 3 Step 7.

¹H-NMR (400MHz, CDCl₃) 0.85 (s, 3H), 0.94 (d, J=6.70Hz, 3H), 0.97 (s, 9H), 1.14 (s, 3H), 1.42-1.48 (m, 2H), 1.50-1.57 (m, 1H), 1.79-1.85 (m, 1H), 2.07-2.16 (m, 1H), 2.30 (d, J=14.10Hz, 1H), 2.63-2.69 (m, 2H), 3.28 (s, 3H), 3.66 (dd, J=12.02, 4.62Hz, 1H), 3.77-3.83 (m, 2H), 7.17 (d, J=7.86Hz, 1H), 7.25-7.28 (m, 1H), 7.42 (d, J=2.08Hz, 1H)

[0219]

(Step 10)

10 Methyl 3-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-(hydroxymethyl)-3,4,4-trimethylcyclopentyl}ureido)bicyclo[1.1.1]pentane-1-carboxylate



15 This step was performed according to Example 54 Step 8.


¹H-NMR (400MHz, CDCl₃) 0.85 (s, 3H), 0.92 (d, J=6.70 Hz, 3H), 0.97 (s, 10H), 0.99 (s, 3H), 1.14 (s, 3H), 1.41-1.46 (m, 3H), 1.57-1.63 (m, 1H), 2.13-2.18 (m, 1H), 2.22-2.26 (m, 1H), 2.33 (s, 6H), 2.57-2.65 (m, 3H), 3.66-3.69 (m, 5H),

4.63-4.67 (m, 1H), 6.67-6.70 (m, 1H), 7.10-7.18 (m, 2H)

[0220]

(Step 11)

Methyl 3-[(S)-7a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-
5 5,6,6-trimethyl-2-oxo-1,2,5,6,7,7a-hexahydro-3H-
cyclopenta[d]pyrimidin-3-yl]bicyclo[1.1.1]pentane-1-
carboxylate



This step was performed according to Example 54 Step 9.

10 Purification was performed with a chiral preparative
column. Purification conditions for the preparative column
are shown as follows.

Preparative apparatus: Recycling Preparative Liquid
Chromatograph LC-92XX NEXT SERIES, Japan Analytical
15 Industry Co., Ltd.

Column: Daicel CHIRALPAK IA 2.0 cmφ x 25 cmL

Mobile phase: n-hexane/2-propanol = 92/8

Flow rate: 10.0 mL/min

Detection: UV (254 nm)

20 Measurement with a chiral column showed 6.0 minutes of

the retention time for the resulting title compound (8.5 minutes of the retention time for the enantiomer of the title compound) with > 99% ee purity. Analytical conditions for the chiral column are shown as follows.

5 Measurement apparatus: HPLC system, Shimadzu Corporation, High-Performance Liquid Chromatograph Prominence

Column: Daicel CHIRALPAK IA-3 0.46 cm ϕ x 15cmL

Column temperature: 30°C

Mobile phase: n-hexane/2-propanol = 90/10

10 Flow rate: 1.0 mL/min


Detection: UV (254 nm)

¹H-NMR (400MHz, CDCl₃) 0.76 (s, 3H), 0.90 (d, J=7.34 Hz, 3H), 0.97 (s, 9H), 1.13 (s, 3H), 1.41-1.47 (m, 2H), 2.01 (d, J=13.33Hz, 1H), 2.30 (d, J=13.33Hz, 1H), 2.37-2.46 (m, 7H),
15 2.60-2.67 (m, 2H), 3.69 (s, 3H), 4.91 (s, 1H), 6.23 (d, J=0.73Hz, 1H), 7.05 (dd, J=7.83, 1.96Hz, 1H), 7.14 (d, J=7.83Hz, 1H), 7.20 (d, J=1.96Hz, 1H)

[0221]

(Step 12)

20 3-{(S)-7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5,6,6-trimethyl-2-oxo-1,2,5,6,7,7a-hexahydro-3H-cyclopenta[d]pyrimidin-3-yl}bicyclo[1.1.1]pentane-1-carboxylic acid




This step was performed according to Example 54 Step 10.

[0222]

5 Example 54

(Step 1)

Hept-6-enal



Hept-6-en-1-ol (150 g) and (diacetoxyiodo)benzene (508 g) were mixed in chloroform (1500 mL) under nitrogen gas, and thereto was added a mixed solution of 2,2,6,6-tetramethylpiperidin-1-oxyl radical (20.5 g) in chloroform (20.0 mL) at room temperature. The reaction solution was stirred at room temperature for 3 days. Then, thereto was added an aqueous solution (1500 mL) of sodium carbonate (278 g) and sodium thiosulfate (208 g) under water cooling, and the mixture was stirred at room temperature for 1 hour, and then separated. The aqueous layer was extracted with chloroform (1000 mL). Combined organic layers were washed


with saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was distilled under reduced pressure (85°C, 70 mmHg) to give a crude product of the title compound (214 g, 46.4w% of iodobenzene inclusive).

¹H-NMR (400 MHz, CDCl₃) 1.40-1.47 (m, 2H), 1.62-1.69 (m, 2H), 2.03-2.11 (m, 2H), 2.44 (td, J=7.40, 1.62Hz, 2H), 4.95-5.04 (m, 2H), 5.74-5.84 (m, 1H), 9.77 (t, J=1.62Hz, 1H).

[0223]

(Step 2)

Hept-6-enal oxime



Hept-6-enal (214 g, 46.4w% of iodobenzene inclusive) was mixed in water (1149 mL) and ethanol (2298 mL) under nitrogen gas, and then thereto were added sodium acetate (151 g) and hydroxylamine hydrochloride (107 g) at room temperature. The reaction solution was stirred at room temperature for 2 days. The reaction solution was concentrated under reduced pressure, and thereto were added ethyl acetate and water. The solution was separated. The aqueous layer was extracted with ethyl acetate (twice).


The organic layer was washed with saturated aqueous sodium chloride solution and concentrated under reduced pressure. The resulting residue was mixed with ethyl acetate, and the resulting solid was filtered with Celite. The filtrate was concentrated under reduced pressure to give a crude product of the title compound (203 g, 24.5w% of ethyl acetate, 2.5w% of ethanol, and 9.2w% of iodobenzene inclusive) (geometric isomer mixture).

¹H-NMR (400 MHz, CDCl₃) 1.40-1.56 (m, 4H), 2.05-2.10 (m, 2H), 2.18-2.24 (m, 1H), 2.38-2.43 (m, 1H), 4.94-5.04 (m, 2H), 5.74-5.85 (m, 1H), 6.72 (t, J=5.32Hz, 0.55H), 7.43 (t, J=6.01Hz, 0.45H)

[0224]

(Step 3)

3,3a,4,5,6,7-Hexahydro-benzo[c]isoxazole



Hept-6-enal oxime (203 g, 24.5w% of ethyl acetate, 2.5w% of ethanol, and 9.2w% of iodobenzene inclusive) and methanol (2333 mL) were mixed under nitrogen gas, and thereto was added trifluoroacetic acid (17.5 mL) under salt-ice cooling, followed by addition of (diacetoxyiodo)benzene (384 g) over 1 hour. The reaction solution was stirred at room temperature for 3 hours. Then, thereto was added an aqueous solution (1000 mL) of sodium

carbonate (194 g) and sodium sulfite (57.8 g) under ice cooling, and the solution was stirred at room temperature for 1 hour. The reaction solution was concentrated under reduced pressure, and thereto was added water (2500 mL).


5 The aqueous layer was extracted with ethyl acetate (1200 mL, twice). The organic layer was washed with saturated aqueous sodium chloride solution (twice) and concentrated under reduced pressure. The resulting residue was combined with a crude product (45.0 g) synthesized separately
10 according to Example 54 Steps 1 to 2 and purified by silica gel column chromatography (Yamazen Corporation, automated refining apparatus, eluent: ethyl acetate/n-hexane, R_f = 0.25 (ethyl acetate/n-hexane = 1/4)), and then azeotroped with n-hexane to give the title compound (92.2 g, 2.5w% of
15 n-hexane inclusive).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.24-1.45 (m, 3H), 1.84-1.90 (m, 1H), 1.96-2.20 (m, 3H), 2.77-2.82 (m, 1H), 3.10-3.20 (m, 1H), 3.77 (dd, $J=9.48$, 7.86Hz, 1H), 4.49 (dd, $J=9.48$, 7.86Hz, 1H)

20 [0225]

(Step 4)

7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]octahydro-benzo[c]isoxazole



4-Bromo-2-chloro-1-(3,3-dimethyl-butyl)benzene (119 g) and tetrahydrofuran (210 mL) were mixed under argon gas, and thereto was added 2.69M n-butyllithium/n-hexane solution (100 mL) under cooling at -78°C . The reaction solution was stirred under cooling at -78°C for 2.5 hours (Reaction solution A). 3,3a,4,5,6,7-Hexahydrobenzo[c]isoxazole (33.4 g) was mixed in toluene (900 mL) and tetrahydrofuran (180 mL), and thereto were added boron trifluoride-diethyl ether complex (37.7 mL) and Reaction solution A under cooling at -78°C . The reaction solution was stirred for 2 hours, and then thereto was added 2N aqueous sodium hydroxide solution (240 mL) under cooling at -78°C . The mixture was stirred at room temperature, and then thereto were added toluene (300 mL) and water (240 mL). The mixture was separated. The organic layer was washed with 13w/w% aqueous sodium chloride solution (400 mL) and saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (107


g).

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.43-1.86 (m, 7H),
1.86-1.98 (m, 3H), 2.64-2.77 (m, 3H), 3.37-4.20 (brm, 2H),
5.82 (brs, 1H), 7.18 (d, J=7.83Hz, 1H), 7.37 (d, J=6.60Hz,
5 1H), 7.52 (s, 1H)

[0226]

(Step 5)

{2-Amino-2-[3-chloro-4-(3,3-dimethyl-
butyl)phenyl]cyclohexyl}methanol



10

7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]octahydro-
benzo[c]isoxazole (107 g) was mixed in acetic acid (600 mL),
tetrahydrofuran (180 mL), and water (180 mL) under nitrogen
gas, and thereto was added zinc powder (24.4 g) in 5 parts
15 under heating at 80°C. The reaction solution was stirred
under heating at 80°C for 30 minutes. The reaction
solution was filtered with Celite at room temperature, and
then the filtrate was concentrated under reduced pressure.
The resulting residue was mixed in ethyl acetate (1000 mL)
20 and 28w/w% ammonia water (140 mL), and the mixture was


separated. The organic layer was washed with 20w/w% aqueous sodium carbonate solution (200 mL) and saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (99.6 g).

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.39-1.50 (m, 4H), 1.53-1.66 (m, 2H), 1.70-1.93 (m, 4H), 2.02 (td, J=13.20, 2.93Hz, 1H), 2.65-2.69 (m, 2H), 3.26 (dd, J=11.25, 2.93Hz, 1H), 3.44 (dd, J=11.25, 2.93Hz, 1H), 7.19 (d, J=8.07Hz, 1H), 7.33 (dd, J=8.07, 1.96Hz, 1H), 7.48 (d, J=1.96Hz, 1H)

[0227]

(Step 6)

{(R)-2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]cyclohexyl}methanol (2S,3S)-2,3-bis-benzoyloxy-succinate



{2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]cyclohexyl}methanol (99.6 g) and tetrahydrofuran (245 mL) were mixed under nitrogen gas, and


thereto was added a mixed solution of (2*S*,3*S*)-2,3-bis-benzoyloxy-succinic acid (94.0 g) in tetrahydrofuran (345 mL) under heating at 85°C. The reaction solution was stirred under heating at 85°C for 5.5 hours. The solution
 5 was stirred for 13.5 hours with being slowly cooled to room temperature, and then the precipitated solid was collected by filtration to give the title compound (62.2 g).

¹H-NMR (400 MHz, DMSO-D₆) 0.96 (s, 9H), 1.38 -1.65 (m, 4H), 1.78-1.70 (m, 8H), 1.86-1.94 (m, 1H), 1.99-2.06 (m, 1H),
 10 2.62-2.67 (m, 2H), 3.09 (dd, J=10.76, 5.62Hz, 1H), 3.19 (dd, J=10.76, 2.45Hz, 1H), 5.64 (s, 2H), 7.36-7.42 (m, 2H), 7.45-7.49 (m, 4H), 7.58-7.63 (m, 3H), 7.89-7.91 (m, 4H)

[0228]

(Step 7)

15 { (R)-2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]cyclohexyl}methanol



{ (R)-2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]cyclohexyl}methanol (2*S*,3*S*)-2,3-bis-benzoyloxy-succinate (62.2 g) and tetrahydrofuran (290 mL)
 20


were mixed under nitrogen gas. The reaction solution was stirred under heating at 85°C for 6 hours. The solution was stirred for 13.5 hours with being slowly cooled to room temperature, and then the precipitated solid was collected by filtration to give {2-amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]cyclohexyl}methanol (2S,3S)-2,3-bis-benzoyloxy-succinate (59.5 g). The resulting {2-amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]cyclohexyl}methanol (2S,3S)-2,3-bis-benzoyloxy-succinate was mixed in ethyl acetate (600 mL) and methanol (60 mL), and thereto was added 1N aqueous sodium hydroxide solution (150 mL) under ice cooling. The mixture was separated at room temperature. The organic layer was washed with 20w/w% aqueous sodium carbonate solution (135 mL for the first round and 50 mL for the second round), 13w/w% aqueous sodium chloride solution (130 mL), and saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give the title compound (25.5 g).

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.39-1.50 (m, 4H), 1.53-1.66 (m, 2H), 1.70-1.93 (m, 4H), 2.02 (td, J=13.20, 2.93Hz, 1H), 2.65-2.69 (m, 2H), 3.26 (dd, J=11.25, 2.93Hz, 1H), 3.44 (dd, J=11.25, 2.93Hz, 1H), 7.19 (d, J=8.07Hz, 1H), 7.33 (dd, J=8.07, 1.96Hz, 1H), 7.48 (d, J=1.96Hz, 1H)

[0229]

(Step 8)

Methyl 3-(3-{(R)-1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-
2-hydroxymethyl-cyclohexyl}ureido)bicyclo[1.1.1]pentane-1-
5 carboxylate



3-(Methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic
acid (14.7 g) and toluene (135 mL) were mixed under
nitrogen gas, and thereto were added diphenylphosphoryl
10 azide (20.3 mL) and triethylamine (13.1 mL) at room
temperature. The reaction solution was stirred under
heating at 100°C for 1 hour. The reaction solution was
added dropwise to a mixed solution of {(R)-2-amino-2-[3-
chloro-4-(3,3-dimethyl-butyl)phenyl]cyclohexyl}methanol
15 (25.5 g) in tetrahydrofuran (135 mL) over 35 minutes under
ice cooling. The reaction solution was stirred at room
temperature for 26 hours, and then the precipitated solid
was collected by filtration to give the title compound
(21.3 g). To the filtrate were added ethyl acetate (400
20 mL) and saturated aqueous sodium hydrogen carbonate

solution (150 mL), and the mixture was separated. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution (150 mL), water (100 mL), and saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was mixed with ethyl acetate (50 mL) and stirred under heating at 70°C for 1.5 hours. The reaction solution was stirred at room temperature for 2 hours, and then the precipitated solid was collected by filtration to give the title compound (13.9 g).

Measurement with a chiral column showed 9.4 minutes of the retention time for the resulting title compound (12.3 minutes of the retention time for the enantiomer of the title compound) with > 99% ee purity. Conditions for the chiral column are shown as follows.

Measurement apparatus: HPLC system, Shimadzu Corporation, High-Performance Liquid Chromatograph Prominence

Column: Daicel CHIRALPAK OZ-3R 0.46 cmφ x 15 cmL

Column temperature: 40°C

Mobile phase: 0.1v/v% aqueous formic acid solution/0.1v/v% formic acid-acetonitrile solution = 45/55

Flow rate: 1.0 mL/min


Detection: UV (254 nm)

^1H -NMR (400 MHz, CDCl_3) 0.97 (s, 9H), 1.42-1.48 (m, 3H),
 1.70-1.52 (m, 4H), 1.72-1.79 (m, 1H), 1.88-1.98 (m, 2H),
 2.37 (s, 6H), 2.61-2.66 (m, 2H), 2.98-3.04 (m, 1H), 3.37-
 3.41 (m, 1H), 3.45-3.49 (m, 1H), 3.69 (s, 3H), 4.71 (s, 1H),
 5 6.27 (s, 1H), 7.15-7.18 (m, 1H), 7.26-7.27 (m, 2H)

[0230]

(Step 9)

Methyl 3-[(S)-8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-
 oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-
 10 yl]bicyclo[1.1.1]pentane-1-carboxylate



Methyl 3-[(S)-8a-[3-chloro-4-(3,3-dimethyl-
 butyl)phenyl]-2-hydroxymethyl-
 cyclohexyl]ureido)bicyclo[1.1.1]pentane-1-carboxylate (16.7
 15 g) and chloroform (495 mL) were mixed under nitrogen gas,
 and thereto were added at room temperature
 (diacetoxyiodo)benzene (11.9 g) and 2,2,6,6-
 tetramethylpiperidin-1-oxyl radical (0.527 g). The
 reaction solution was stirred at room temperature for 21.5
 20 hours, and then thereto was added 2,2,6,6-


tetramethylpiperidin-1-oxyl radical (0.527 g) at room temperature. The mixture was stirred at room temperature for 4 hours, and then thereto were added (diacetoxyiodo)benzene (1.20 g) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (0.527 g) at room temperature. The mixture was stirred at room temperature for 17 hours, and then thereto were added 20w/w% aqueous sodium sulfite solution (200 mL) and 20w/w% aqueous sodium carbonate solution (30 mL) at room temperature. The mixture was separated. The organic layer was dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was mixed with toluene (300 mL), and thereto was added pentafluoroaniline trifluoromethanesulfonate (0.561 g) at room temperature. The reaction solution was stirred for 2 hours with being heated to 100°C, and then thereto was added trifluoroacetic acid (10 mL) at room temperature. The reaction solution was stirred at room temperature for 15 hours, and then concentrated under reduced pressure. The resulting residue was combined with a crude product (48.3 g) synthesized separately according to Example 54 Steps 1 to 8 and purified by silica gel column chromatography (ethyl acetate/n-hexane = 1/4 -> 1/2 -> 1/1) to give the title compound (35.3 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.98 (s, 9H), 1.26-1.42 (m, 2H),
 1.43-1.47 (m, 2H), 1.68-1.73 (m, 2H), 1.88-1.97 (m, 2H),
 2.13-2.17 (m, 1H), 2.44 (s, 6H), 2.48-2.53 (m, 1H), 2.64-
 2.69 (m, 2H), 3.70 (s, 3H), 4.54 (s, 1H), 5.96 (d, $J=1.47\text{Hz}$,
 5 1H), 7.15 (dd, $J=8.07$, 1.96Hz , 1H), 7.21 (d, $J=8.07\text{Hz}$, 1H),
 7.28 (d, $J=1.96\text{Hz}$, 1H)

[0231]

(Step 10)

3-{(S)-8a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-2-oxo-
 10 1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-
 yl}bicyclo[1.1.1]pentane-1-carboxylic acid



Methyl 3-{(S)-8a-[3-chloro-4-(3,3-dimethyl-
 butyl)phenyl]-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-
 15 yl}bicyclo[1.1.1]pentane-1-carboxylate (35.3 g) was mixed
 in tetrahydrofuran (150 mL) and methanol (600 mL) under
 nitrogen gas, and thereto was added 2N aqueous sodium
 hydroxide solution (72 mL) under ice cooling. The reaction
 solution was stirred at room temperature for 17.5 hours,


and then thereto was added water (300 mL). The reaction solution was concentrated under reduced pressure. To The resulting residue was added 2N hydrochloric acid (90 mL) under ice cooling, and the mixture was extracted with ethyl acetate/methanol = 100/1 (1010 mL). The organic layer was washed with 13w/w% aqueous sodium chloride solution (200 mL, twice) and saturated aqueous sodium chloride solution (twice) and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was mixed in ethyl acetate (55 mL) and n-hexane (220 mL) and stirred under heating at 55°C for 9 hours. The reaction solution was stirred for 9 hours with being slowly cooled to room temperature, and then the precipitated solid was collected by filtration to give the title compound (22.8 g).

Absolute configuration of the asymmetric carbon of the title compound was determined by single-crystal X-ray structural analysis.

[0232]

(Intermediate Step 1)

4-Bromo-2-chloro-1-(3,3-dimethyl-but-1-ynyl)benzene




4-Bromo-2-chloro-1-iodo-benzene (325 g), 3,3-dimethyl-
 but-1-yne (97 g), bis(triphenylphosphino)palladium (II)
 dichloride (14.3 g), copper (I) iodide (7.80 g),
 5 triphenylphosphine (10.7 g), and diisopropylamine (2.17 L)
 were mixed under argon gas and stirred under heating at
 100°C for 15 hours. Then, thereto was added n-hexane (2.20
 L) at room temperature, and the precipitated solid was
 filtered with Celite. Then, the filtrate was concentrated
 10 under reduced pressure. The resulting residue was mixed
 with n-hexane (2.00 L) and silica gel (327 g) and stirred
 at room temperature for 17 hours. The resulting solid was
 removed with a filter, and then the filtrate was
 concentrated under reduced pressure to give a crude product
 15 of the title compound (316 g).

¹H-NMR (400 MHz, CDCl₃) 1.33 (s, 9H), 7.26 (d, J=8.31Hz,
 1H), 7.30 (dd, J=8.31, 1.71Hz, 1H), 7.54 (d, J=1.96Hz, 1H)
 [0233]

(Intermediate Step 2)

20 4-Bromo-2-chloro-1-(3,3-dimethyl-butyl)benzene



4-Bromo-2-chloro-1-(3,3-dimethyl-but-1-ynyl)benzene

(66.0 g) was mixed with ethanol (330 mL), and thereto was added 5w/w% platinum/activated carbon (13.2 g). The reaction solution was stirred under ordinary pressure and hydrogen gas for 6 hours, and then the gas in the reaction vessel was replaced with nitrogen gas. Celite was added to the reaction solution and removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was mixed with ethanol (330 mL), and thereto was added 5w/w% platinum/activated carbon (13.6 g). The reaction solution was stirred under 0.1 MPa hydrogen gas for 55 hours, and then the gas in the reaction vessel was replaced with nitrogen gas. Celite was added to the reaction solution and removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was mixed with n-hexane (200 mL), and the mixture was separated. The organic layer was washed with water. All aqueous layers were extracted with n-hexane (20 mL). The organic layer was washed with water and saturated aqueous sodium chloride solution and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and

then the filtrate was concentrated under reduced pressure to give Crude product A (61.0 g).

4-Bromo-2-chloro-1-(3,3-dimethyl-but-1-ynyl)benzene (10.0 g) was mixed with ethanol (50.0 mL), and thereto was added 5w/w% platinum/activated carbon (2.03 g). The reaction solution was stirred under ordinary pressure and hydrogen gas for 18 hours, and then the gas in the reaction vessel was replaced with nitrogen gas. Celite was added to the reaction solution and removed with a filter, and then the filtrate was concentrated under reduced pressure to give Crude product B (9.81 g).

4-Bromo-2-chloro-1-(3,3-dimethyl-but-1-ynyl)benzene (82.1 g) was mixed with ethanol (411 mL), and thereto was added 5w/w% platinum/activated carbon (16.7 g). The reaction solution was stirred under ordinary pressure and hydrogen gas for 26 hours, and then the gas in the reaction vessel was replaced with nitrogen gas. Celite was added to the reaction solution and removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was mixed with n-hexane (250 mL) and water (20 mL) and separated. The organic layer was washed with water. All aqueous layers were extracted with n-hexane (50 mL). The organic layer was washed with water and saturated aqueous sodium chloride solution and dried over magnesium sulfate. Magnesium sulfate was removed with

a filter, and then the filtrate was concentrated under reduced pressure to give Crude product C (78.4 g).

4-Bromo-2-chloro-1-(3,3-dimethyl-but-1-ynyl)benzene (82.1 g) was mixed with ethanol (411 mL), and thereto was
5 added 5w/w% platinum/activated carbon (16.7 g). The reaction solution was stirred under ordinary pressure and hydrogen gas for 26 hours, and then the gas in the reaction vessel was replaced with nitrogen gas. Celite was added to the reaction solution and removed with a filter, and then
10 the filtrate was concentrated under reduced pressure. The resulting residue was mixed with n-hexane (250 mL) and water (20 mL), and the mixture was separated. The organic layer was washed with water. All aqueous layers were extracted with n-hexane (50 mL). The organic layer was
15 washed with water and saturated aqueous sodium chloride solution and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give Crude product D (74.9 g).

20 4-Bromo-2-chloro-1-(3,3-dimethyl-but-1-ynyl)benzene (82.1 g) was mixed with ethanol (410 mL), and thereto was added 5w/w% platinum/activated carbon (16.4 g). The reaction solution was stirred under ordinary pressure and hydrogen gas for 27 hours, and then the gas in the reaction
25 vessel was replaced with nitrogen gas. Celite was added to

the reaction solution and removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was mixed with n-hexane (250 mL) and water (20 mL), and the mixture was separated. The organic layer was washed with water. All aqueous layers were extracted with n-hexane (50 mL). The organic layers were washed with water and saturated aqueous sodium chloride solution and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give Crude product E (74.4 g).

The title compound (298.51 g) of Crude products A, B, C, D, and E was mixed with n-hexane (2.00 L) and silica gel (150 g) and stirred at room temperature for 3 hours. The resulting solid was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was azeotroped with toluene (300 mL) to give a crude product of the title compound (304 g).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.97 (s, 9H), 1.40-1.45 (m, 2H), 2.61-2.66 (m, 2H), 7.08 (d, $J=8.07\text{Hz}$, 1H), 7.29 (dd, $J=8.07$, 1.96Hz , 1H), 7.48 (d, $J=1.96\text{Hz}$, 1H)

[0234]

Example 63

(Step 1)

Ethyl 8,8-difluoro-1,4-dioxo-spiro[4.5]decane-6-carboxylate



Ethyl 5,5-difluoro-2-oxo-cyclohexanecarboxylate (4.45 g) and toluene (53 mL) were mixed under nitrogen gas, and thereto were added ethylene glycol (1.45 mL) and p-toluenesulfonic acid monohydrate (205 mg) at room temperature. The reaction solution was stirred under heating at 140°C with dehydration for 1 day. Then, thereto was added a mixed aqueous solution of sodium carbonate (114 mg) in water (18 mL) under ice cooling, and the mixture was separated. The aqueous layer was extracted with ethyl acetate (twice). The organic layer was washed with water (three times) and saturated aqueous sodium chloride solution and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, $R_f = 0.38$ (ethyl acetate/n-hexane = 1/4)) to give a crude product of the title compound (3.46 g, 15w% of ethyl acetate inclusive).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.28 (t, $J=7.1\text{Hz}$, 3H), 1.78–1.91 (m, 2H), 2.06–2.16 (m, 2H), 2.23–2.31 (m, 1H), 2.38–2.53 (m, 1H), 2.99 (ddd, $J=12.4, 4.6, 1.6\text{Hz}$, 1H), 3.90–4.01 (m, 4H),

4.17 (q, $J=7.1\text{Hz}$, 2H)

[0235]

(Step 2)

(8,8-Difluoro-1,4-dioxa-spiro[4.5]dec-6-yl)methanol



5

Lithium aluminum hydride (886 mg) and tetrahydrofuran (12 mL) were mixed under argon gas, and thereto was added dropwise a mixed solution in tetrahydrofuran (18 mL) of ethyl 8,8-difluoro-1,4-dioxaspiro[4.5]decane-6-carboxylate (2.92 g) azeotroped with toluene over 1 hour under ice cooling. The reaction solution was stirred under ice cooling for 20 minutes and stirred at room temperature for 4 hours. To the reaction solution were added slowly water (0.886 mL), 2N aqueous solution of sodium hydroxide (0.886 mL), and water (2.66 mL) under ice cooling. The reaction solution was stirred at room temperature for 30 minutes. To the reaction solution were added Celite (1 g), magnesium sulfate (1 g), and ethyl acetate (30 mL), and the mixture was stirred for 30 minutes. Solids were removed with Celite, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (3.06 g, 22w% of ethyl acetate inclusive).

15

20


$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.67-1.75 (m, 1H), 1.83-1.90 (m,

1H), 1.91-2.18 (m, 4H), 2.18-2.26 (m, 1H), 2.56 (dd, J=6.5, 5.3Hz, 1H), 3.62-3.76 (m, 2H), 3.98-4.08 (m, 4H)

[0236]

(Step 3)

5 2-(8,8-Difluoro-1,4-dioxaspiro[4.5]dec-6-ylmethoxy)-
isoindole-1,3-dione



(8,8-Difluoro-1,4-dioxaspiro[4.5]dec-6-yl)methanol

(2.38 g) and tetrahydrofuran (36 mL) were mixed, and
10 thereto were added N-hydroxyphthalimide (2.79 g) and
triphenylphosphine (4.49 g) under ice cooling. To the
mixed solution was added bis(2-methoxyethyl)azodicarboxylate
(4.01 g) in 8 parts over 40 minutes under ice cooling. The
reaction solution was stirred at room temperature for 15
15 hours. The reaction solution was concentrated under
reduced pressure, and the resulting residue was purified by
silica gel column chromatography (Biotage flash
purification systems, eluent: ethyl acetate/n-hexane, R_f =
0.46 (ethyl acetate/n-hexane = 2/3)) to give a crude
20 product of the title compound (3.44 g).


¹H-NMR (400 MHz, CDCl₃) 1.71-1.85 (m, 2H), 1.96-2.20 (m, 3H), 2.51-2.59 (m, 1H), 2.65-2.75 (m, 1H), 3.93-4.10 (m,

5H), 4.41 (ddd, J=9.1, 4.0, 2.0Hz, 1H), 7.74 (d, J=3.0Hz, 1H), 7.76 (d, J=3.0Hz, 1H), 7.83 (d, J=3.0Hz, 1H), 7.84 (d, J=3.0Hz, 1H)

[0237]

5 (Step 4)

O-((8,8-Difluoro-1,4-dioxaspiro[4.5]decan-6-yl)methyl)hydroxylamine




2-((8,8-Difluoro-1,4-dioxaspiro[4.5]dec-6-yl)methoxy)-
 10 isoindole-1,3-dione (3.44 g) and chloroform (34 mL) were
 mixed, and thereto was added a mixed solution of
 methylhydrazine (0.619 mL) in chloroform (2 mL) over 5
 minutes under ice cooling. The reaction solution was
 stirred at room temperature for 2 hours. Solids were
 15 removed with Celite, and then the filtrate was concentrated
 under reduced pressure to give a crude product of the title
 compound (3.49 g, 38w% of chloroform inclusive).

¹H-NMR (400 MHz, CDCl₃) 1.71-2.10 (m, 5H), 2.23-2.32 (m,
 1H), 2.39-2.47 (m, 1H), 3.52 (dd, J=9.9, 8.3Hz, 1H), 3.88
 20 (ddd, J=10.1, 4.3, 2.2Hz, 1H), 3.91-4.02 (m, 4H), 5.40 (s,
 2H)

[0238]

(Step 5)

5,5-Difluoro-3,3a,4,5,6,7-hexahydrobenzo[c]isoxazole



O-(8,8-Difluoro-1,4-dioxaspiro[4.5]dec-6-ylmethyl)-

5 hydroxylamine (2.17 g, 38w% of chloroform inclusive) and tetrahydrofuran (19.5 mL) were mixed, and then thereto was added slowly 6N hydrochloric acid (4.87 mL) under ice cooling. The reaction solution was stirred at room temperature for 1 hour, and then thereto was added
 10 tetrahydrofuran (20 mL). The mixture was stirred further for 4 hours. To the reaction solution was added water (5 mL), and the mixture was stirred under heating at 60°C for 6 hours. Then, thereto was added potassium carbonate (4.04 g) under ice cooling, and the mixture was concentrated
 15 under reduced pressure and then extracted with ethyl acetate (twice). The organic layers were washed with saturated aqueous sodium chloride solution (twice) and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under
 20 reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, R_f = 0.19 (ethyl acetate/n-hexane = 2/3)) to give the title


compound (1.11 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.76-2.02 (m, 2H), 2.33-2.42 (m, 1H), 2.44-2.60 (m, 2H), 2.89 (ddt, $J=14.6, 5.8, 2.1\text{Hz}$, 1H), 3.49-3.59 (m, 1H), 3.91 (dd, $J=9.8, 8.4\text{Hz}$, 1H), 4.57 (ddd, $J=10.4, 8.3, 1.5\text{Hz}$, 1H)

[0239]

(Step 6)

7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-difluoro-octahydro-benzo[c]isoxazole



10

4-Bromo-2-chloro-1-(3,3-dimethyl-butyl)-benzene (1.26 g) and tetrahydrofuran (5 mL) were mixed under argon gas, and thereto was added dropwise 1.54M n-butyllithium/n-hexane solution (2.42 mL) under cooling at -78°C . The reaction solution was stirred for 1 hour under cooling at -78°C (Reaction solution A). 5,5-Difluoro-3,3a,4,5,6,7-hexahydro-benzo[c]isoxazole (500 mg) and toluene (12.5 mL) were mixed, and thereto were added boron trifluoride-diethyl ether complex (0.464 mL) and Reaction solution A under cooling at -78°C . The reaction solution was stirred at -78°C for 3 hours, and then thereto was added saturated

20


aqueous sodium chloride solution (16 mL). The aqueous layer was extracted with ethyl acetate (twice) at room temperature. The organic layers were washed with saturated aqueous sodium chloride solution (twice) and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, $R_f = 0.43$ (ethyl acetate/n-hexane = 1/4)) to give a crude product of the title compound (1.09 g, 8w% of ethyl acetate inclusive).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.98 (s, 9H), 1.43-1.47 (m, 2H), 1.88-2.32 (m, 6H), 2.65-2.70 (m, 2H), 3.02 (qd, $J=7.0$, 3.8Hz, 1H), 3.79-3.88 (m, 2H), 5.67 (s, 1H), 7.21 (d, $J=7.9\text{Hz}$, 1H), 7.34 (dd, $J=8.0$, 2.0Hz, 1H), 7.52 (d, $J=1.4\text{Hz}$, 1H)

[0240]

(Step 7)

{2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-difluoro-cyclohexyl}methanol




7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-difluoro-octahydro-benzo[c]isoxazole (1.00 g), acetic acid (15 mL), tetrahydrofuran (10 mL), and water (5 mL) were mixed under nitrogen gas, and thereto was added zinc powder (1.83 g) in 5 parts under heating at 60°C. The reaction solution was stirred under heating at 60°C for 4 hours. Then, thereto was added 28w/w% ammonia water (25 mL) under ice cooling, and the mixture was extracted with cyclopentyl methyl ether (three times). The organic layers were washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (1.41 g, 31w% of cyclopentyl methyl ether inclusive).

¹H-NMR (400 MHz, CDCl₃) 0.99 (s, 9H), 1.43-1.48 (m, 2H), 1.50-1.75 (m, 7H), 2.04-2.44 (m, 3H), 2.66-2.70 (m, 2H), 3.30 (dd, J=11.7, 3.4Hz, 1H), 3.44 (dt, J=11.6, 2.9Hz, 1H), 7.22 (d, J=8.1Hz, 1H), 7.29 (dd, J=8.1, 2.1Hz, 1H), 7.47 (d, J=2.1Hz, 1H)

[0241]

(Step 8)

Ethyl 3-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-4,4-difluoro-2-hydroxymethyl-cyclohexyl}ureido)propionate



{2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-
 5,5-difluoro-cyclohexyl}methanol (300 mg, 31w% of
 cyclopentyl methyl ether inclusive) and tetrahydrofuran (3
 5 mL) were mixed under argon gas, and thereto was added ethyl
 3-isocyanato-propionate (0.122 mL) under ice cooling. The
 reaction solution was stirred at room temperature for 24
 hours. Then, thereto was added N,N,N'-
 trimethylethylenediamine, and the mixture was stirred at
 10 room temperature for 10 minutes. The organic layer was
 concentrated under reduced pressure. Then, thereto was
 added ethyl acetate, and the mixture was washed with 10w/w%
 aqueous citric acid solution, 10w/w% aqueous sodium
 chloride solution, and saturated aqueous sodium chloride
 15 solution. The organic layer was dried over magnesium
 sulfate. Magnesium sulfate was removed with a filter, and
 then the filtrate was concentrated under reduced pressure.
 The resulting residue was purified by silica gel column
 chromatography (Biotage flash purification systems, eluent:
 20 ethyl acetate/n-hexane, R_f = 0.41 (ethyl acetate/n-hexane =


1/2)) to give a crude product of the title compound (496 mg, 21w% of ethyl acetate inclusive).

¹H-NMR (400 MHz, CDCl₃) 0.97 (s, 9H), 1.29 (t, J=7.2Hz, 3H), 1.41-1.46 (m, 2H), 1.81-1.89 (m, 2H), 1.96-2.11 (m, 5H), 2.54 (dd, J=6.7, 5.2Hz, 2H), 2.61-2.67 (m, 2H), 3.20 (d, J=9.3Hz, 1H), 3.38-3.52 (m, 4H), 4.18 (q, J=7.2Hz, 2H), 4.91 (t, J=6.1Hz, 1H), 6.44 (s, 1H), 7.12 (dd, J=8.1, 1.8Hz, 1H), 7.17 (d, J=7.8Hz, 1H), 7.26 (d, J=1.8Hz, 1H)

[0242]

10 (Step 9)

Ethyl 3-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-6,6-difluoro-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}propionate



15 Ethyl 3-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-4,4-difluoro-2-hydroxymethyl-cyclohexyl}ureido)propionate (366 mg) and chloroform (3.7 mL) were mixed, and thereto were added (diacetoxyiodo)benzene (266 mg) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (11 mg) at room temperature. The reaction solution was stirred at room

20

temperature for 24 hours, and then thereto were added 20w/w% aqueous sodium sulfite solution and saturated aqueous sodium hydrogen carbonate solution at room temperature. The mixture was stirred at room temperature
5 for 30 minutes. The aqueous layer was extracted with ethyl acetate (twice) and washed with saturated aqueous sodium chloride solution. The organic layers were dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under
10 reduced pressure. The resulting residue and toluene (7.3 mL) were mixed, and thereto was added pentafluoroaniline trifluoromethanesulfonate (12.1 mg) at room temperature. The reaction solution was stirred under heating at 120°C for 6 hours, and then the reaction solution was
15 concentrated under reduced pressure. To the resulting residue was added saturated aqueous sodium hydrogen carbonate solution at room temperature, and the aqueous layer was extracted with ethyl acetate (twice). The organic layers were washed with saturated aqueous sodium
20 chloride solution and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent:
25 ethyl acetate/n-hexane, $R_f = 0.37$ (ethyl acetate/n-hexane =


2/3)) to give a crude product of the title compound (330 mg, 18w% of ethyl acetate inclusive).

¹H-NMR (400 MHz, CDCl₃) 0.98 (d, J=4.9Hz, 9H), 1.28 (t, J=7.1Hz, 3H), 1.42-1.47 (m, 2H), 1.61-1.77 (m, 1H), 2.13 (s, 1H), 2.21-2.38 (m, 2H), 2.53-2.73 (m, 6H), 3.58-3.65 (m, 1H), 3.83 (dt, J=14.1, 6.1Hz, 1H), 4.19 (q, J=7.2Hz, 2H), 4.78 (s, 1H), 6.22 (d, J=1.8Hz, 1H), 7.16 (dd, J=8.0, 2.0Hz, 1H), 7.24 (d, J=8.1Hz, 1H), 7.30 (d, J=2.1Hz, 1H)

[0243]

(Step 10)

3-{8a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-6,6-difluoro-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}propionic acid



Ethyl 3-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-6,6-difluoro-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}propionate (25 mg) was mixed in tetrahydrofuran (0.25 mL) and methanol (0.25 mL), and thereto was added 2N aqueous solution of sodium hydroxide (0.518 mL) at room temperature. The reaction solution was stirred at room


temperature for 16 hours, and then the reaction solution was concentrated under reduced pressure. To the resulting residue were added water and 2N hydrochloric acid (0.518 mL) under ice cooling, and the precipitated solid was collected by filtration to give the title compound (23.6 mg).

[0244]

Example 68

(Step 1)

7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5-methoxy-octahydro-benzo[c]isoxazole



4-Bromo-2-chloro-1-(3,3-dimethyl-butyl)-benzene (1.3 g) was mixed in toluene (11 mL) and tetrahydrofuran (4.4 mL) under argon gas, and thereto was added dropwise 2.66M n-butyllithium/n-hexane solution (1.45 mL) under cooling at -78°C. The reaction solution was stirred under cooling at -78°C for 1 hour (Reaction solution A). 5-methoxy-3,3a,4,5,6,7-hexahydrobenzo[c]isoxazole (0.5 g) and toluene (32 mL) were mixed, and thereto were added boron trifluoride-diethyl ether complex (0.49 mL) and Reaction


solution A under cooling at -78°C . The reaction solution was stirred under cooling at -78°C for 2 hours, and then thereto was added a saturated aqueous solution of ammonium chloride. The mixture was stirred at room temperature, and then thereto were added ethyl acetate and water. The mixture was separated. The aqueous layer was extracted with chloroform. The organic layer was concentrated under reduced pressure, and the crude product was purified by silica gel column chromatography (ethyl acetate/n-hexane = 1/9 $\rightarrow$ 2/3) to give the title compound (0.68 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.97 (s, 9H), 1.40-1.48 (m, 2H), 1.49-1.61 (m, 4H), 1.79-2.19 (m, 4H), 2.63-2.69 (m, 2H), 2.86-2.98 (m, 1H), 3.39 (s, 3H), 3.49-3.61 (m, 1H), 3.65-3.81 (m, 1H), 7.13-7.20 (m, 1H), 7.34-7.42 (m, 1H), 7.52-7.58 (m, 1H)

[0245]

(Step 2)

{2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5-methoxy-cyclohexyl}methanol



7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5-methoxy-octahydro-benzo[c]isoxazole (0.66 g) was mixed in acetic acid (10 mL), tetrahydrofuran (3.3 mL), and water (3.3 mL) under nitrogen gas, and thereto was added zinc powder (1.24 g) in parts at room temperature. The reaction solution was stirred under heating at 60°C for 1 hour 30 minutes. Then, thereto was added 28w/w% ammonia water (15 mL) under ice cooling, and the mixture was stirred at room temperature. To the reaction solution were added chloroform, methanol, and water, and the mixture was separated. The organic layer was dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The crude product was purified by silica gel column chromatography (methanol/ethyl acetate = 0/1 -> 1/99) to give the title compound (0.75 g).


¹H-NMR (400 MHz, CDCl₃) 0.96 (s, 9H), 1.11-1.28 (m, 1H), 1.40-1.48 (m, 2H), 1.48-1.65 (m, 3H), 1.67-1.78 (m, 1H), 1.89-2.05 (m, 5H), 2.29-2.43 (m, 1H), 2.62-2.69 (m, 2H), 3.18-3.25 (m, 1H), 3.36 (s, 3H), 3.43-3.48 (m, 1H), 3.64-3.69 (m, 1H), 7.15-7.19 (m, 1H), 7.31-7.35 (m, 1H), 7.47-7.50 (m, 1H)

[0246]

(Step 3)

Ethyl 3-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-

hydroxymethyl-4-methoxy-cyclohexyl}ureido)propionate




{2-Amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5-methoxy-cyclohexyl}methanol (0.18 g) and tetrahydrofuran (3
 5 mL) were mixed, and thereto was added ethyl 3-isocyanato-propionate (74 μ L) under ice cooling. The reaction solution was stirred at room temperature for 40 minutes and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography
 10 (methanol/ethyl acetate = 0/1 $\rightarrow$ 1/99) to give the title compound (0.26 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.97 (s, 9H), 1.22-1.33 (m, 3H), 1.40-1.49 (m, 2H), 1.64-1.72 (m, 2H), 1.72-1.86 (m, 2H), 1.86-2.00 (m, 2H), 2.05-2.21 (m, 2H), 2.49-2.57 (m, 2H),
 15 2.59-2.68 (m, 2H), 2.68-2.81 (m, 1H), 3.29-3.40 (m, 3H), 3.40-3.53 (m, 3H), 3.62-3.66 (m, 1H), 4.09-4.22 (m, 2H), 4.80 (br s, 1H), 6.35 (br s, 1H), 7.12-7.21 (m, 2H), 7.29-7.34 (m, 1H)

[0247]

20 (Step 4)

Ethyl 3-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-6-methoxy-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}propionate



5 Ethyl 3-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-hydroxymethyl-4-methoxy-cyclohexyl}ureido)propionate
 (0.257 g) and dichloromethane (3 mL) were mixed under nitrogen gas, and thereto were added (diacetoxyiodo)benzene (0.18 g) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (8
 10 mg) under ice cooling. The reaction solution was stirred at room temperature for 3 hours, and then thereto were added ethyl acetate, an aqueous solution of sodium thiosulfate, and an aqueous solution of sodium hydrogen carbonate at room temperature. The mixture was separated.
 15 The organic layer was washed with water and then dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue and toluene (4 mL) were mixed, and thereto was added pentafluoroaniline trifluoromethanesulfonate (8 mg) at room temperature. The
 20

reaction solution was stirred under heating at 120°C for 1 hour 30 minutes and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (ethyl acetate/n-hexane = 1/4 -> 1/2
5 -> 1/1) to give a diastereomeric mixture of the title compound (0.19 g). The diastereomeric mixture was purified with a Recycling Preparative Liquid Chromatograph to give a single enantiomer of the title compound (40 mg).

Purification conditions for the preparative
10 chromatography are shown as follows.

Preparative apparatus: Recycling preparative liquid chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

Column: Daicel CHIRALPAK IA 2.0 cmφ x 25 cm

15 Mobile phase: n-hexane:2-propanol = 85:15

Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement of the resulting compound with a chiral column showed 7.8 minutes of the retention time for the
20 resulting enantiomer with > 99%ee of optical purity. The retention time for the opposite enantiomer was 5.2 minutes.

Analytical conditions for the chiral column are shown as follows.

Measurement apparatus: HPLC system, Shimadzu Corporation,
25 High-Performance Liquid Chromatograph Prominence

Column: Daicel CHIRALPAK IA-3 0.46 cmφ x 15 cm

Column temperature: 40°C

Mobile phase: n-hexane:2-propanol = 85:15

Flow rate: 1.0 mL/min


5 Detection: UV (254 nm)

¹H-NMR (400 MHz, CDCl₃) 0.96 (s, 9H), 1.15-1.23 (m, 1H),
1.23-1.29 (m, 3H), 1.37-1.45 (m, 2H), 1.74-1.84 (m, 1H),
1.87-1.98 (m, 1H), 1.98-2.06 (m, 1H), 2.47-2.56 (m, 2H),
2.57-2.70 (m, 4H), 3.17-3.27 (m, 1H), 3.29 (s, 3H), 3.53-
10 3.63 (m, 1H), 3.75-3.85 (m, 1H), 4.13-4.21 (m, 2H), 4.65
(br s, 1H), 6.09-6.13 (m, 1H), 7.09-7.15 (m, 1H), 7.15-7.21
(m, 1H), 7.25-7.28 (m, 1H)

[0248]

(Step 5)

15 3-{(S)-8a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-6-
methoxy-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-
yl}propionic acid



Ethyl 3-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-6-
20 methoxy-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-

yl}propionate (36 mg) and ethanol (1 mL) were mixed, and thereto was added 2N aqueous solution of sodium hydroxide (0.11 mL) at room temperature. The reaction solution was stirred at room temperature for 1 hour 30 minutes and then concentrated under reduced pressure. To the resulting residue were added at room temperature 1N hydrochloric acid (0.23 mL) and water, and then the precipitated solid was collected by filtration to give the title compound (27.4 mg).


10 [0249]

Example 77

(Step 1)

Methyl 3-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-hydroxymethyl-4-methoxy-

15 cyclohexyl}ureido)bicyclo[1.1.1]pentane-1-carboxylate



3-(Methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid (96 mg) and toluene (2 mL) were mixed under argon gas, and thereto were added diphenylphosphoryl azide (0.13 mL) and triethylamine (0.087 mL) at room temperature. The

20


reaction solution was stirred under heating at 120°C for 1 hour. The reaction solution was added dropwise to a mixed solution of {2-amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5-methoxy-cyclohexyl}methanol (200 mg) in 5 tetrahydrofuran (3 mL) under ice cooling. The reaction solution was stirred at room temperature for 2 hours and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (methanol/ethyl acetate = 0/1 -> 1/99) to give the title 10 compound (0.29 g).

¹H-NMR (400 MHz, CDCl₃) 0.97 (s, 9H), 1.40-1.47 (m, 2H), 1.61-1.72 (m, 1H), 1.75-1.84 (m, 1H), 1.88-1.99 (m, 2H), 2.05-2.17 (m, 2H), 2.36 (s, 6H), 2.60-2.67 (m, 2H), 2.76-2.88 (m, 1H), 3.31-3.37 (m, 1H), 3.38 (s, 3H), 3.45-3.53 (m, 15 1H), 3.61-3.67 (m, 1H), 3.69 (s, 4H), 4.80-4.87 (m, 1H), 6.41 (br s, 1H), 7.13-7.16 (br m, 2H), 7.29 (br s, 1H)

[0250]

(Step 2)

Methyl 3-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-6-methoxy-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}bicyclo[1.1.1]pentane-1-carboxylate 20



Methyl

3-(3-{1-[3-chloro-4-(3,3-dimethyl-

butyl)phenyl]-2-hydroxymethyl-4-methoxy-

cyclohexyl}ureido)bicyclo[1.1.1]pentane-1-carboxylate

5 (0.287 g) and dichloromethane (7 mL) were mixed, and thereto were added (diacetoxyiodo)benzene (0.195 g) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (9 mg) under ice cooling. The reaction solution was stirred at room temperature for 14 hours 30 minutes, and then thereto were

10 added ethyl acetate, an aqueous solution of sodium thiosulfate, and saturated aqueous sodium hydrogen carbonate solution at room temperature. The mixture was separated and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate

15 was concentrated under reduced pressure. The resulting residue and toluene (3 mL) were mixed, and then thereto was added pentafluoroaniline trifluoromethanesulfonate (9 mg) at room temperature. The reaction solution was stirred under heating at 120°C for 1 hour and then concentrated

20 under reduced pressure. The resulting residue was purified

by silica gel column chromatography (ethyl acetate/n-hexane = 1/4 -> 1/2 -> 1/1) to give a diastereomeric mixture of the title compound (0.20 g). The diastereomeric mixture was purified with a Recycling Preparative Liquid Chromatograph to give a single enantiomer of the title compound (78 mg).

Purification conditions for the preparative chromatography are shown as follows.

Preparative apparatus: Recycling preparative liquid chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

Column: Daicel CHIRALPAK IA 2.0 cmφ x 25 cm

Mobile phase: n-hexane:2-propanol = 85:15

Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement of the resulting compound with a chiral column showed 6.5 minutes of the retention time for the resulting enantiomer with > 99%ee of optical purity. The retention time for the opposite enantiomer was 4.0 minutes.

Analytical conditions for the chiral column are shown as follows.

Measurement apparatus: HPLC system, Shimadzu Corporation, High-Performance Liquid Chromatograph Prominence

Column: Daicel CHIRALPAK IA-3 0.46 cmφ x 15 cm

Column temperature: 40°C

Mobile phase: n-hexane:2-propanol = 85:15

Flow rate: 1.0 mL/min


Detection: UV (254 nm)

¹H-NMR (400 MHz, CDCl₃) 0.98 (s, 9H), 1.16-1.28 (m, 1H),
 1.39-1.47 (m, 2H), 1.80-1.89 (m, 1H), 1.89-2.00 (m, 1H),
 2.00-2.08 (m, 1H), 2.45 (s, 6H), 2.51-2.59 (m, 2H), 2.62-
 2.69 (m, 2H), 3.20-3.29 (m, 1H), 3.31 (s, 3H), 3.70 (s, 3H),
 4.67-4.70 (m, 1H), 6.01-6.04 (m, 1H), 7.10-7.16 (m, 1H),
 7.16-7.23 (m, 1H), 7.24-7.29 (m, 1H)

[0251]

(Step 3)

3-{(S)-8a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-6-methoxy-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}bicyclo[1.1.1]pentane-1-carboxylic acid



Methyl

3-{(S)-8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-6-methoxy-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}bicyclo[1.1.1]pentane-1-carboxylate (69 mg) and methanol (2 mL) were mixed, and thereto was added 2N


aqueous solution of sodium hydroxide (0.21 mL) at room temperature. The reaction solution was stirred at room temperature for 2 hours and then concentrated under reduced pressure. To the resulting residue were added 2N
 5 hydrochloric acid (0.3 mL), chloroform, and methanol at room temperature, and the mixture was separated. The organic layer was dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give the title
 10 compound (60 mg).

[0252]

Example 87

(Step 1)

Methyl 3-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-4,4-
 15 difluoro-2-hydroxymethyl-
 cyclohexyl}ureido)bicyclo[1.1.1]pentane-1-carboxylate



3-(Methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic
 acid (319 mg) and toluene (4 mL) were mixed under argon gas,
 20 and thereto were added diphenylphosphoryl azide (0.445 mL)


and triethylamine (0.288 mL) at room temperature. The reaction solution was stirred under heating at 120°C for 1 hour 30 minutes. To the reaction solution was added tetrahydrofuran (2 mL), and the mixture was added dropwise to a mixed solution of {2-amino-2-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,5-difluoro-cyclohexyl}methanol (979 mg, 31w% of cyclopentyl methyl ether inclusive) in tetrahydrofuran (6.5 mL) over 10 minutes under ice cooling. The reaction solution was stirred at room temperature for 1 hour 30 minutes and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, Rf = 0.62 (ethyl acetate/n-hexane = 1/2)) to give the title compound (493 mg).

¹H-NMR (400 MHz, CDCl₃) 0.97 (s, 9H), 1.42-1.46 (m, 2H), 1.88-1.93 (m, 1H), 2.06-2.51 (m, 12H), 2.62-2.67 (m, 2H), 3.21-3.25 (m, 1H), 3.42 (d, J=10.4Hz, 1H), 3.50 (d, J=10.9Hz, 1H), 3.69 (s, 3H), 4.84 (d, J=2.8Hz, 1H), 6.45 (s, 1H), 7.11 (dd, J=8.2, 2.0Hz, 1H), 7.18 (d, J=8.2Hz, 1H), 7.25 (d, J=2.0Hz, 1H)

[0253]

(Step 2)

Methyl 3-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-6,6-difluoro-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}bicyclo[1.1.1]pentane-1-carboxylate



Methyl

3-(3-{1-[3-chloro-4-(3,3-dimethyl-

butyl)phenyl]-4,4-difluoro-2-hydroxymethyl-
cyclohexyl}ureido)bicyclo[1.1.1]pentane-1-carboxylate (493

5 mg) and chloroform (16 mL) were mixed, and thereto were added (diacetoxyiodo)benzene (332 mg) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (15 mg) at room temperature. The reaction solution was stirred at room temperature for 16 hours, and then thereto were added an
10 aqueous solution of sodium thiosulfate and saturated aqueous sodium hydrogen carbonate solution at room temperature. The aqueous layer was extracted with chloroform (twice). The organic layers were dried over magnesium sulfate. Magnesium sulfate was removed with a
15 filter, and then the filtrate was concentrated under reduced pressure. The resulting residue and toluene (20 mL) were mixed, and thereto was added pentafluoroaniline trifluoromethanesulfonate (16 mg) at room temperature. The reaction solution was stirred under heating at 120°C for 2
20 hours and then concentrated under reduced pressure. The

resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, $R_f = 0.46$ (ethyl acetate/n-hexane = 1/2)) to give a crude product of a racemate of the title compound (316 mg). The racemate was purified with a Recycling Preparative Liquid Chromatograph to give a single enantiomer of the title compound (112 mg).

Purification conditions for the preparative chromatography are shown as follows.

10 Preparative apparatus: Recycling preparative liquid chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

Column: Daicel CHIRALPAK IA 2.0 cmφ x 25 cm

Mobile phase: n-hexane:2-propanol = 85:15

15 Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement of the resulting compound with a chiral column showed 5.2 minutes of the retention time for the resulting enantiomer with > 99%ee of optical purity. The retention time for the opposite enantiomer was 3.1 minutes.

20 Analytical conditions for the chiral column are shown as follows.

Measurement apparatus: HPLC system, Shimadzu Corporation, High-Performance Liquid Chromatograph Prominence

25 Column: Daicel CHIRALPAK IA-3 0.46 cmφ x 15 cm

Column temperature: 40°C

Mobile phase: n-hexane:2-propanol = 80:20

Flow rate: 1.0 mL/min


Detection: UV (254 nm)

5 ¹H-NMR (400 MHz, CDCl₃) (s, 9H), 1.43-1.47 (m, 2H), 1.61-1.77 (m, 1H), 2.09-2.15 (m, 1H), 2.21-2.42 (m, 2H), 2.45 (s, 6H), 2.53-2.64 (m, 2H), 2.66-2.70 (m, 2H), 3.71 (s, 3H), 4.84 (s, 1H), 6.10 (d, J=1.8Hz, 1H), 7.15 (dd, J=7.9, 1.9Hz, 1H), 7.25 (d, J=8.1Hz, 1H), 7.29 (d, J=2.1Hz, 1H)

10 [0254]

(Step 3)

3-{(S)-8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-6,6-difluoro-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}bicyclo[1.1.1]pentane-1-carboxylic acid



15

Methyl

3-{(S)-8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-6,6-difluoro-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}bicyclo[1.1.1]pentane-1-carboxylate (94 mg) and methanol (2 mL) were mixed, and thereto was added 2N


aqueous solution of sodium hydroxide (0.28 mL) at room temperature. The reaction solution was stirred at room temperature for 26 hours and then concentrated under reduced pressure. To the resulting residue were added 2N
 5 hydrochloric acid (0.28 mL) and water under ice cooling, and the mixture was extracted with a mixed solution of chloroform and methanol. The organic layer was dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced
 10 pressure to give the title compound (73.0 mg).

[0255]

Example 109

(Step 1)

4-(2,2-Diethoxy-ethoxy)-3,4-dimethyl-pent-1-ene



15

60w% Sodium hydride (5.17 g) and tetrahydrofuran (66 mL) were mixed under argon gas, and thereto was added slowly a mixed solution of 2,3-dimethyl-pent-4-en-2-ol (5.0 g) in tetrahydrofuran (11 mL) under ice cooling. The
 20 reaction solution was stirred under ice cooling for 1 hour, and then thereto was added bromoacetaldehyde diethyl acetal (20.4 mL) under ice cooling. The reaction solution was


stirred under heating at 80°C for 18 hours. To the reaction solution was added water (20 mL) under ice cooling, and the mixture was stirred at room temperature for 1 hour. Then, the aqueous layer was extracted with ethyl acetate (twice). The organic layers were washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/nhexane, Rf = 0.24 (ethyl acetate/n-hexane = 1/19)). Evaporation of bromoacetaldehyde diethyl acetal included in the resulting crude product under reduced pressure gave a crude product of the title compound (2.80 g, 50w% of bromoacetaldehyde diethyl acetal inclusive).

¹H-NMR (400 MHz, CDCl₃) 1.00 (d, J=6.9Hz, 3H), 1.09 (s, 3H), 1.11 (s, 3H), 1.22 (t, J=7.1Hz, 6H), 2.33 (dt, J=14.6, 6.9Hz, 1H), 3.40 (dd, J=5.2, 2.7Hz, 2H), 3.54-3.62 (m, 2H), 3.67-3.75 (m, 2H), 4.54 (t, J=5.3Hz, 1H), 4.97-5.03 (m, 2H), 5.83 (ddd, J=17.7, 9.8, 7.3Hz, 1H)

[0256]

(Step 2)

(1,1,2-Trimethyl-but-3-enyloxy)acetaldehyde




4-(2,2-Diethoxy-ethoxy)-3,4-dimethyl-pent-1-ene (2.80 g, 50w% of bromoacetaldehyde diethyl acetal inclusive) and tetrahydrofuran (12.2 mL) were mixed, and thereto was added
 5 2N hydrochloric acid (6.1 mL) under ice cooling. The reaction solution was stirred under heating at 60°C for 4 hours. To the reaction solution was added potassium carbonate (0.84 g) under ice cooling. The reaction solution was concentrated under reduced pressure, and
 10 thereto was added saturated aqueous sodium chloride solution (10 mL). The aqueous layer was extracted with diethyl ether (twice). The organic layer was washed with saturated aqueous sodium chloride solution and dried over magnesium sulfate. Magnesium sulfate was removed with a
 15 filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (2.03 g, 44w% of tetrahydrofuran inclusive).

¹H-NMR (400 MHz, CDCl₃) 1.04 (d, J=6.9Hz, 3H), 1.14 (s, 3H), 1.15 (s, 3H), 2.36 (t, J=7.4Hz, 1H), 3.96 (t, J=1.3Hz, 2H),
 20 5.02-5.08 (m, 2H), 5.84 (ddd, J=17.9, 9.9, 7.4Hz, 1H), 9.72 (t, J=1.3Hz, 1H)

[0257]

(Step 3)

(1,1,2-Trimethyl-but-3-enyloxy)acetaldehyde oxime



(1,1,2-Trimethyl-but-3-enyloxy)acetaldehyde (2.03 g,
 5 44w% of tetrahydrofuran inclusive) was mixed in ethanol
 (16.1 mL) and water (8.1 mL) under argon gas, and thereto
 were added sodium acetate (4.62 g) and hydroxylamine
 hydrochloride (2.24 g) at room temperature. The reaction
 solution was stirred under heating at 60°C for 20 hours.
 10 The reaction solution was concentrated under reduced
 pressure, and thereto was added saturated aqueous sodium
 chloride solution (10 mL). The aqueous layer was extracted
 with ethyl acetate (twice). The organic layers were washed
 with saturated aqueous sodium chloride solution (twice) and
 15 dried over sodium sulfate. Sodium sulfate was removed with
 a filter, and then the filtrate was concentrated under
 reduced pressure. The resulting residue was purified by
 silica gel column chromatography (Biotage flash
 purification systems, eluent: ethyl acetate/n-hexane, R_f =
 20 0.38 (ethyl acetate/n-hexane = 1/4)) to give a crude
 product of the title compound (1.00 g, 32w% of ethyl
 acetate inclusive).


¹H-NMR (400 MHz, CDCl₃) 1.00 (d, J=6.7Hz, 1.38H), 1.01 (d,

J=6.7Hz, 1.62H), 1.12 (s, 3H), 1.15 (s, 1.62H), 1.16 (s, 1.38H), 2.33-2.40 (m, 1H), 2.95 (s, 1H), 4.03 (dd, J=5.5, 2.1Hz, 0.92H), 4.29 (d, J=3.2Hz, 1.08H), 5.00-5.06 (m, 2H), 5.77-5.86 (m, 1H), 6.87 (t, J=3.6Hz, 0.54H), 7.47 (t, J=5.4Hz, 0.46H)

[0258]

(Step 4)

4,5,5-Trimethyl-3,3a,4,5-tetrahydro-7H-pyrano[3,4-c]isoxazole



(1,1,2-Trimethylbut-3-en-1-yl)oxyacetaldehyde oxime (1.00 g, 32w% of ethyl acetate inclusive) and methanol (13.5 mL) were mixed, and thereto was added slowly trifluoroacetic acid (0.101 mL) under ice cooling. To the reaction solution was added (diacetoxyiodo)benzene (1.70 g) over 1 hour under ice cooling. The reaction solution was stirred under ice cooling for 20 minutes and stirred at room temperature for 1 hour. To the reaction solution were added saturated aqueous sodium hydrogen carbonate solution (7 mL) and sodium sulfite (248 mg), and then the mixture was stirred at room temperature for 30 minutes. The reaction solution was concentrated under reduced pressure, and the residue was extracted with ethyl acetate (twice).

The organic layers were washed with saturated aqueous sodium chloride solution (twice) and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure.


5 The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, $R_f = 0.24$ (ethyl acetate/n-hexane = 1/4)) to give the title compound (495 mg).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.94 (d, $J=6.9\text{Hz}$, 1.95H), 0.94 (d, $J=7.2\text{Hz}$, 1.05H), 1.22 (s, 1.95H), 1.23 (s, 1.05H), 1.26 (s, 1.95H), 1.42 (s, 1.05H), 1.64-1.72 (m, 0.65H), 1.84 (dt, $J=13.7, 6.1\text{Hz}$, 0.35H), 3.16 (q, $J=11.0\text{Hz}$, 0.65H), 3.77 (dd, $J=11.5, 7.9\text{Hz}$, 0.65H), 3.86-3.93 (m, 0.35H), 4.14 (t, $J=8.5\text{Hz}$, 0.35H), 4.34 (ddd, $J=15.3, 13.8, 1.1\text{Hz}$, 1H), 4.38 (dd, $J=7.6, 4.0\text{Hz}$, 0.35H), 4.49 (d, $J=9.6\text{Hz}$, 0.65H), 4.53 (d, $J=9.3\text{Hz}$, 0.35H), 4.60 (dd, $J=10.2, 7.8\text{Hz}$, 0.65H)

[0259]

(Step 5)

7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-4,5,5-trimethyl-
20 hexahydro-pyrano[3,4-c]isoxazole



4-Bromo-2-chloro-1-(3,3-dimethyl-butyl)-benzene (0.553 g) was mixed in tetrahydrofuran (2.2 mL) and toluene (5.53 mL) under argon gas, and then thereto was added dropwise

5 1.54M n-butyllithium/n-hexane solution (1.25 mL) under cooling at -78°C . The reaction solution was stirred under cooling at -78°C for 1 hour (Reaction solution A). 4,5,5-Trimethyl-3,3a,4,5-tetrahydro-7H-pyrano[3,4-c]isoxazole (250 mg) and toluene (15 mL) were mixed, and thereto was

10 added boron trifluoride-diethyl ether complex (0.241 mL) under cooling at -78°C . The mixture was stirred for 10 minutes. Then, thereto was added dropwise slowly Reaction solution A over 25 minutes under cooling at -78°C . The reaction solution was stirred for 3 hours, and then thereto

15 was added a saturated aqueous solution of ammonium chloride (8 mL) under cooling at -78°C . The mixture was stirred at room temperature for 30 minutes. The aqueous layer was extracted with ethyl acetate (twice). The organic layers were washed with saturated aqueous sodium chloride solution

20 (twice) and dried over sodium sulfate. Sodium sulfate was


removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, R_f = 0.48 (ethyl acetate/n-hexane = 1/4)) to give a crude product of the title compound (511 mg, 23w% of ethyl acetate inclusive).

¹H-NMR (400 MHz, CDCl₃) 0.96 (d, J=6.7Hz, 1.95H), 0.98 (s, 9H), 1.05 (d, J=7.4Hz, 1.05H), 1.28 (s, 1.05H), 1.29 (s, 1.95H), 1.30 (s, 1.05H), 1.32 (s, 1.95H), 1.41-1.47 (m, 2H), 1.63 (dt, J=18.0, 6.9Hz, 0.65H), 2.19 (t, J=7.1Hz, 0.35H), 2.43 (dd, J=11.0, 5.0Hz, 0.65H), 2.64-2.68 (m, 2H), 2.91 (td, J=9.2, 5.6Hz, 0.35H), 3.55 (dd, J=7.4, 5.1Hz, 0.65H), 3.64 (d, J=13.2Hz, 0.65H), 3.74 (d, J=13.2Hz, 0.35H), 3.82 (d, J=7.6Hz, 0.35H), 3.85 (d, J=13.2Hz, 1H), 3.89-4.20 (m, 1H), 5.73 (s, 0.35H), 6.24 (s, 0.65H), 7.18 (d, J=8.3Hz, 0.35H), 7.19 (d, J=8.1Hz, 0.65H), 7.36 (dd, J=8.2, 2.0Hz, 0.35H), 7.42 (dd, J=8.0, 2.2Hz, 0.65H), 7.52 (d, J=1.8Hz, 0.35H), 7.57 (d, J=1.8Hz, 0.65H)

[0260]

(Step 6)

{5-Amino-5-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2,2,3-trimethyl-tetrahydro-pyran-4-yl}methanol



7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-4,5,5-trimethyl-hexahydro-pyrano[3,4-c]isoxazole (392 mg) was mixed in acetic acid (5.9 mL), tetrahydrofuran (2 mL), and water (2 mL), and thereto was added zinc powder (700 mg) in 5 parts over 25 minutes under heating at 60°C. The reaction solution was stirred under heating at 60°C for 3 hours. To the reaction solution was added 28w/w% ammonia water (10 mL) under ice cooling, and the aqueous layer was extracted with cyclopentyl methyl ether (three times). The organic layers were washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (467 mg, 21w% of cyclopentyl methyl ether inclusive).


¹H-NMR (400 MHz, CDCl₃) 0.97 (d, J=6.7Hz, 1.95H), 0.99 (s, 9H), 1.14 (d, J=8.6Hz, 1.05H), 1.20 (s, 1.05H), 1.28 (s, 1.95H), 1.32 (s, 1.05H), 1.38 (s, 1.95H), 1.44-1.49 (m, 2H), 1.50-1.76 (m, 4H), 2.13-2.21 (m, 0.65H), 2.26-2.30 (m,

0.35H), 2.66-2.71 (m, 2H), 3.08 (d, J=12.5Hz, 0.65H), 3.41 (dd, J=12.5, 1.4Hz, 0.35H), 3.42 (dd, J=12.0, 1.4Hz, 0.65H), 3.53 (dd, J=11.9, 3.1Hz, 0.65H), 3.72 (d, J=5.1Hz, 0.70H), 4.04 (d, J=12.0Hz, 0.35H), 4.10 (d, J=12.0Hz, 0.65H), 7.21 (d, J=7.9Hz, 0.35H), 7.23 (d, J=7.6Hz, 0.65H), 7.27 (dd, J=8.0, 2.2Hz, 0.65H), 7.40 (d, J=2.1Hz, 0.65H), 7.42 (dd, J=7.7, 2.0Hz, 0.35H), 7.57 (d, J=1.8Hz, 0.35H)

[0261]

(Step 7)

10 Methyl 3-(3-{3-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-4-hydroxymethyl-5,6,6-trimethyl-tetrahydro-pyran-3-yl}ureido)bicyclo[1.1.1]pentane-1-carboxylate



3-(Methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic
 15 acid (197 mg) and toluene (4 mL) were mixed under argon gas, and thereto were added diphenylphosphoryl azide (0.270 mL) and triethylamine (0.174 mL) at room temperature. The reaction solution was stirred under heating at 120°C for 1 hour. To the reaction solution was added tetrahydrofuran


(4 mL), and the mixture was added dropwise to a mixed solution of {5-amino-5-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2,2,3-trimethyl-tetrahydro-pyran-4-yl}methanol (467 mg, 21w% of cyclopentyl methyl ether inclusive) in tetrahydrofuran (4 mL) over 10 minutes under ice cooling. The reaction solution was stirred at room temperature for 13 hours, and then thereto was added N,N,N'-trimethylethylenediamine (0.0195 mL). The reaction solution was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: acetone/n-hexane, Rf = 0.50 (acetone/n-hexane = 2/3)) to give a crude product of the title compound (511 mg, 5w% of ethyl acetate inclusive).

¹H-NMR (400 MHz, CDCl₃) 0.90-0.99 (m, 3H), 0.97 (s, 9H), 1.10 (s, 1.05H), 1.21 (s, 1.95H), 1.32 (s, 1.95H), 1.42-1.49 (m, 2H), 1.43 (s, 1.05H), 1.50-1.73 (m, 1H), 1.98-2.41 (m, 2H), 2.29 (s, 2.1H), 2.38 (s, 3.9H), 2.61-2.67 (m, 2H), 3.58-3.76 (m, 2H), 3.67 (s, 1.05H), 3.69 (s, 1.95H), 3.79-3.87 (m, 1.30H), 4.01-4.09 (m, 0.70H), 4.68 (s, 0.35H), 4.97 (s, 0.65H), 5.94 (s, 0.65H), 6.21 (s, 0.35H), 7.12-7.17 (m, 1.65H), 7.25-7.27 (m, 0.65H), 7.45 (d, J=8.1Hz, 0.35H), 7.51 (d, J=1.8Hz, 0.35H)

[0262]

(Step 8)

Methyl 3-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,6,6-trimethyl-2-oxo-1,2,5,6,8,8a-hexahydro-pyrano[3,4-d]pyrimidin-3-yl}bicyclo[1.1.1]pentane-1-carboxylate



5 Methyl 3-(3-{3-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-4-hydroxymethyl-5,6,6-trimethyl-tetrahydro-pyran-3-yl}ureido)bicyclo[1.1.1]pentane-1-carboxylate (511 mg, 5w% of ethyl acetate inclusive) and chloroform (4.8 mL) were mixed, and thereto were added (diacetoxyiodo)benzene

10 (330 mg) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (14.1 mg) at room temperature. The reaction solution was stirred at room temperature for 19 hours, and then thereto were added 20w/w% aqueous sodium sulfite solution and saturated aqueous sodium hydrogen carbonate solution at

15 room temperature. The mixture was stirred at room temperature for 30 minutes. The aqueous layer was extracted with ethyl acetate (twice). The organic layers were washed with saturated aqueous sodium chloride solution and dried over magnesium sulfate. Magnesium sulfate was

removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue and toluene (9.7 mL) were mixed, and thereto was added pentafluoroaniline trifluoromethanesulfonate (15 mg) at
5 room temperature. The reaction solution was stirred under heating at 120°C for 1 hour 30 minutes and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane,
10 $R_f = 0.40$ (ethyl acetate/n-hexane = 2/3)) to give a diastereomeric mixture of the title compound (434 mg). The diastereomeric mixture was purified with a Recycling Preparative Liquid Chromatograph to give a single enantiomer of the title compound (39.8 mg).

15 Purification conditions for the preparative chromatography are shown as follows.

Preparative apparatus: Recycling preparative liquid chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

20 Column: Daicel CHIRALPAK IA 2.0 cmφ x 25 cm

Mobile phase: n-hexane:2-propanol = 85:15

Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement of the resulting compound with a chiral
25 column showed 7.6 minutes of the retention time for the

resulting enantiomer with > 99%ee of optical purity. The retention time for a diastereomer on the methyl group was 9.8 minutes, and the retention time for a diastereomer on the phenyl group as well as the opposite enantiomer was 3.9 minutes.

Analytical conditions for the chiral column are shown as follows.

Measurement apparatus: HPLC system, Shimadzu Corporation, High-Performance Liquid Chromatograph Prominence

10 Column: Daicel CHIRALPAK IA-3 0.46 cmφ x 15 cm

Column temperature: 30°C

Mobile phase: n-hexane:2-propanol = 85:15

Flow rate: 1.0 mL/min


Detection: UV (254 nm)

15 ¹H-NMR (400 MHz, CDCl₃) 0.88 (d, J=6.9Hz, 3H), 0.98 (s, 9H), 1.14 (s, 3H), 1.21 (s, 3H), 1.42-1.47 (m, 2H), 2.09 (ddd, J=13.8, 7.1, 2.4Hz, 1H), 2.49 (s, 6H), 2.64-2.69 (m, 2H), 3.71 (s, 3H), 3.90 (d, J=11.6Hz, 1H), 4.08 (d, J=11.6Hz, 1H), 4.45 (s, 1H), 5.74 (d, J=1.8Hz, 1H), 7.19 (d, J=8.1Hz, 1H), 7.33 (dd, J=8.0, 2.0Hz, 1H), 7.50 (d, J=2.1Hz, 1H)

[0263]

(Step 9)

3-[(S)-8a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5,6,6-trimethyl-2-oxo-1,2,5,6,8,8a-hexahydro-pyrano[3,4-d]pyrimidin-3-yl}bicyclo[1.1.1]pentane-1-carboxylic acid



Methyl

3-[(S)-8a-[3-chloro-4-(3,3-dimethyl-

butyl)phenyl]-5,6,6-trimethyl-2-oxo-1,2,5,6,8,8a-hexahydro-
pyrano[3,4-d]pyrimidin-3-yl]bicyclo[1.1.1]pentane-1-

5 carboxylate (39.8 mg) was mixed in tetrahydrofuran (0.398 mL) and methanol (0.398 mL) under nitrogen gas, and thereto was added 2N aqueous solution of sodium hydroxide (0.0791 mL) at room temperature. The reaction solution was stirred at room temperature for 16 hours and then concentrated
10 under reduced pressure. To the resulting residue was added water (1 mL), and then thereto was added 2N hydrochloric acid (0.0791 mL) under ice cooling. The precipitated solid was collected by filtration to give the title compound (19.7 mg).


15 [0264]

Example 114

(Step 1)

Methyl 3-(3-[(3R,4S)-1-[4-(3-tert-butyl-2,2-difluoro-
cyclopropyl)-3-chloro-phenyl]-2-hydroxymethyl-3,4-dimethyl-

cyclohexyl}ureido)bicyclo[1.1.1]pentane-1-carboxylate



3-(Methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid (97.9 mg) was mixed in toluene (2 mL) and tetrahydrofuran (2 mL) under argon gas, and thereto were added triethylamine (0.866 mL) and diphenylphosphoryl azide (0.134 mL) at room temperature. The reaction solution was stirred under heating at 120°C for 1 hour. The reaction solution was added dropwise to a mixed solution of { (5S,6R)-2-amino-2-[4-(3-tert-butyl-2,2-difluorocyclopropyl)-3-chlorophenyl]-5,6-dimethylcyclohexyl}methanol (200 mg) in tetrahydrofuran (4 mL) under ice cooling. The reaction solution was stirred at room temperature for 13 hours, and then thereto was added trimethylethylenediamine (0.129 mL). The reaction solution was concentrated under reduced pressure, and then the resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: acetone/n-hexane) to give the title compound (328 mg).


¹H-NMR (400 MHz, CDCl₃) 1.02-1.05 (m, 6H), 1.11-1.11 (m,

9H), 1.30-1.34 (m, 1H), 1.37-1.42 (m, 1H), 1.46-1.48 (m, 1H), 1.61-1.73 (m, 2H), 1.78-1.85 (m, 1H), 2.39-2.35 (m, 7H), 2.73-2.80 (m, 1H), 3.04 (d, J=13.64Hz, 1H), 3.32 (d, J=11.79Hz, 1H), 3.69 (s, 3H), 3.74 (d, J=11.33Hz, 1H), 4.73 (s, 1H), 6.37 (s, 1H), 7.13-7.15 (m, 2H), 7.32 (d, J=14.57Hz, 1H)

[0265]

(Step 2)

Methyl 3-((5R,6S)-8a-[4-(3-tert-butyl-2,2-difluorocyclopropyl)-3-chloro-phenyl]-5,6-dimethyl-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl)bicyclo[1.1.1]pentane-1-carboxylate



Methyl 3-((3-((3R,4S)-1-[4-(3-tert-butyl-2,2-difluorocyclopropyl)-3-chloro-phenyl]-2-hydroxymethyl-3,4-dimethylcyclohexyl)ureido)bicyclo[1.1.1]pentane-1-carboxylate (216 mg) and chloroform (2.2 mL) were mixed, and thereto were added (diacetoxyiodo)benzene (139 mg) and 2,2,6,6-tetramethylpiperidin-1-oxyl radical (6.0 mg) at room temperature. The reaction solution was stirred at room

temperature for 14 hours, and then thereto were added 20w/w% aqueous sodium sulfite solution and 5w/w% aqueous solution of sodium hydrogen carbonate at room temperature. The mixture was stirred at room temperature for 30 minutes and then extracted with ethyl acetate (twice). The organic layer was washed with saturated aqueous sodium chloride solution and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue and toluene (4.3 mL) were mixed, and then thereto was added pentafluoroaniline trifluoromethanesulfonate (6.3 mg) at room temperature. The reaction solution was stirred under heating at 120°C for 3 hours 30 minutes, and then stirred under microwave radiation (100W, 120°C) for 30 minutes. The reaction solution was concentrated under reduced pressure, and then the resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane) to give a diastereomeric mixture of the title compound (61.2 mg). The diastereomeric mixture was purified with a Recycling Preparative Liquid Chromatograph to give a single enantiomer of the title compound (18.0 mg).

Purification conditions for the preparative chromatography are shown as follows.

Preparative apparatus: Recycling preparative liquid

chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

Column: Daicel CHIRALPAK IA 2.0 cm ϕ x 25 cm

Mobile phase: n-hexane:2-propanol = 88:12

5 Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement of the resulting compound with a chiral column showed 8.0 minutes of the retention time for the resulting enantiomer with > 99%ee of optical purity. The
10 retention time for the opposite enantiomer on the phenyl group was 4.8 minutes.

Analytical conditions for the chiral column are shown as follows.

Measurement apparatus: HPLC system, Shimadzu Corporation,
15 High-Performance Liquid Chromatograph Prominence

Column: Daicel CHIRALPAK IA-3 0.46 cm ϕ x 15 cm

Column temperature: 30°C

Mobile phase: n-hexane:2-propanol = 85:15

Flow rate: 1.0 mL/min

20 Detection: UV (254 nm)


^1H -NMR (400 MHz, CDCl_3) 0.65 (d, $J=7.40\text{Hz}$, 3H), 1.03 (d, $J=6.94\text{Hz}$, 3H), 1.12 (s, 9H), 1.36-1.42 (m, 1H), 1.62-1.74 (m, 2H), 1.77-1.86 (m, 1H), 2.05-2.09 (m, 1H), 2.11-2.16 (m, 1H), 2.31-2.25 (m, 1H), 2.48 (s, 6H), 2.79 (dd, $J=14.33$, 9.48Hz, 1H), 3.71 (s, 3H), 4.44 (s, 1H), 6.01 (s, 1H), 7.17
25

(d, $J=8.09\text{Hz}$, 1H), 7.24 (dd, $J=8.32$, 2.08Hz , 1H), 7.39 (d, $J=2.08\text{Hz}$, 1H)

[0266]

(Step 3)

5 3-{(5R,6S)-8a-[4-(3-tert-Butyl-2,2-difluoro-cyclopropyl)-3-chloro-phenyl]-5,6-dimethyl-2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}bicyclo[1.1.1]pentane-1-carboxylic acid



Methyl 3-{(5R,6S)-8a-[4-(3-tert-butyl-2,2-difluoro-cyclopropyl)-3-chloro-phenyl]-5,6-dimethyl-2-oxo-
 10 1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-yl}bicyclo[1.1.1]pentane-1-carboxylate (18.0 mg) was mixed
 in tetrahydrofuran (0.18 mL) and methanol (0.18 mL), and
 then thereto was added 2N aqueous solution of sodium
 15 hydroxide (0.0329 mL) at room temperature. The reaction
 solution was stirred at room temperature for 7 hours 30
 minutes and then concentrated under reduced pressure. To
 the resulting residue was added water (1 mL), and then
 thereto was added 2N hydrochloric acid (0.0329 mL) under
 20 ice cooling. The precipitated solid was collected by


filtration to give the title compound (14.7 mg).

[0267]

Example 115

(Step 1)

5 (2R,3S)-2,3-Dimethyl-pent-4-enoic acid methoxy-methyl-amide



(2R,3S)-2,3-Dimethyl-pent-4-enoic acid (3.5 g), N,O-dimethylhydroxylamine hydrochloride (4.0 g) and 1-hydroxybenzotriazole monohydrate (6.3 g) were mixed in
 10 dimethylformamide (20 mL) under nitrogen gas, and thereto were added triethylamine (7.6 mL) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (7.9 g) under ice cooling. The reaction solution was stirred at room temperature overnight. Then, thereto was added water
 15 (175 mL), and the aqueous layer was extracted with ethyl acetate/n-hexane = 1/1. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution, 1N hydrochloric acid, and saturated aqueous sodium chloride solution and dried over magnesium sulfate. Magnesium
 20 sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl


acetate/n-hexane, $R_f = 0.25$ (ethyl acetate/n-hexane = 1/4))
to give the title compound (4.4 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.99-1.03 (m, 3H), 1.06-1.11 (m, 3H), 2.40-2.49 (m, 1H), 2.74-2.87 (m, 1H), 3.15 (s, 3H),
5 3.66 (s, 3H), 4.89-5.03 (m, 2H), 5.69-5.83 (m, 1H)

[0268]

(Step 2)

(2R,3S)-2,3-Dimethyl-pent-4-enal



10 (2R,3S)-2,3-Dimethyl-pent-4-enoic acid methoxy-methyl-
amide (4.4 g) and tetrahydrofuran (44 mL) were mixed under
nitrogen gas, and thereto was added dropwise 1.02M
diisobutylaluminum hydride/n-hexane solution (30 mL) under
cooling at -78°C . The mixture was stirred under cooling at
15 -78°C for 2.5 hours, and then thereto was added dropwise
1.5M aqueous sulfuric acid solution (41 mL). The mixture
was stirred for 2 hours under ice cooling, and then the
aqueous layer was extracted with methyl tert-butyl ether.
The organic layer was washed with 1.0M sulfuric acid and
20 saturated aqueous sodium chloride solution and dried over
magnesium sulfate. Magnesium sulfate was removed with a
filter, and then the filtrate was concentrated under
reduced pressure to give a crude product of the title


compound (3.7 g, 39w% of tetrahydrofuran inclusive).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.99-1.05 (m, 6H), 2.30-2.41 (m, 1H), 2.55-2.66 (m, 1H), 4.97-5.09 (m, 2H), 5.72-5.85 (m, 1H), 9.63-9.68 (m, 1H)

5 [0269]

(Step 3)

Ethyl (E)-(4S,5S)-4,5-dimethyl-hepta-2,6-dienoate




Sodium hydride (960 mg) and tetrahydrofuran (34 mL)
 10 were mixed under nitrogen gas, and thereto was added
 dropwise ethyl diphenylphosphonoacetate (5.2 mL) under ice
 cooling. The mixture was stirred at room temperature for
 30 minutes, and then thereto was added dropwise a mixed
 solution of (2R,3S)-2,3-dimethyl-pent-4-enal (2.24 g) in
 15 tetrahydrofuran (22 mL) under ice cooling. The reaction
 solution was stirred at room temperature overnight. Then,
 thereto were added saturated aqueous ammonium chloride
 solution and water at room temperature, and the organic
 solvent was evaporated under reduced pressure. The aqueous
 20 layer was extracted with ethyl acetate, and the organic
 layer was washed with saturated aqueous sodium chloride
 solution and dried over magnesium sulfate. Magnesium

sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane) to give the title compound (3.2 g).

[0270]

(Step 4)

Ethyl (4S,5S)-4,5-dimethyl-hept-6-enoate



(E)-(4S,5S)-4,5-Dimethyl-hepta-2,6-dienoic acid ethyl ester (3.2 g) was mixed in tetrahydrofuran (45 mL) and methanol (26 mL) under nitrogen gas, and thereto was added copper (I) chloride. Then, thereto was added sodium borohydride (4.7 g) under ice cooling. The reaction solution was stirred for 5 hours under ice cooling. To the reaction solution was added 1N hydrochloric acid (33 mL). Insoluble substances were removed with a filter, and the filtrate was concentrated under reduced pressure. To the resulting residue was added ethyl acetate, and the mixture was separated. The organic layer was washed with 1N hydrochloric acid and saturated aqueous sodium chloride solution. The resultant was dried over magnesium sulfate,


and then magnesium sulfate was removed with a filter. The filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate/n-hexane = 1/1), and then azeotroped with n-hexane to give the title compound (2.8 g, 6w% of ethyl (E)-(4S,5S)-4,5-dimethyl-hepta-2,6-dienoate inclusive).

¹H-NMR (400 MHz, CDCl₃) 0.79-0.84 (m, 3H), 0.92-0.96 (m, 3H), 1.21-1.28 (m, 3H), 1.34-1.47 (m, 2H), 1.70-1.80 (m, 1H), 2.02-2.12 (m, 1H), 2.15-2.39 (m, 2H), 4.05-4.15 (m, 2H), 4.90 -4.98 (m, 2H), 5.64-5.79 (m, 1H)

[0271]

(Step 5)

(4S,5S)-4,5-Dimethyl-hept-6-enal



Ethyl (4S,5S)-4,5-dimethyl-hept-6-enoate (2.8 g) and dichloromethane (150 mL) were mixed under nitrogen gas, and thereto was added dropwise 1.02M diisobutylaluminum hydride/n-hexane solution (19 mL) under cooling at -78°C. The mixture was stirred under cooling at -78°C for 30 minutes, and then thereto was added dropwise methanol (7.6 mL). Then, thereto was added a saturated aqueous solution


of Rochelle salt (76 mL) under ice cooling. Then, the mixture was stirred at room temperature for 2 hours, and thereto was added diethyl ether. The mixture was separated, and the organic layer was washed with saturated aqueous sodium chloride solution. The solution was dried over magnesium sulfate, and then magnesium sulfate was removed with a filter. The filtrate was concentrated under reduced pressure to give a crude product of the title compound (2.6 g).

¹H-NMR (400 MHz, CDCl₃) 0.81-0.85 (m, 3H), 0.93-0.97 (m, 3H), 1.34-1.45 (m, 2H), 1.70-1.80 (m, 1H), 2.04-2.11 (m, 1H), 2.30-2.50 (m, 2H), 4.90-4.97 (m, 2H), 5.66-5.77 (m, 1H), 9.74-9.76 (m, 1H)

[0272]

(Step 6)

(4S,5S)-4,5-Dimethyl-hept-6-enal-oxime



(4S,5S)-4,5-Dimethyl-hept-6-enal (2.1 g) was mixed in water (15 mL) and ethanol (30 mL) under nitrogen gas, and thereto were added sodium acetate (8.7 g) and hydroxylamine hydrochloride (4.2 g) at room temperature. The reaction solution was stirred under heating at 60°C overnight. The reaction solution was concentrated under reduced pressure,


and thereto were added ethyl acetate and water. The solution was separated. The aqueous layer was extracted with ethyl acetate (twice). The organic layer was washed with saturated aqueous sodium chloride solution and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate/n-hexane, $R_f = 0.1$ (ethyl acetate/n-hexane = 1/10)) to give the title compound (2.0 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.83-0.90 (m, 3H), 0.92-0.97 (m, 3H), 1.21-1.67 (m, 3H), 2.04-2.44 (m, 3H), 4.93-5.01 (m, 2H), 5.66-5.80 (m, 1H), 6.67-6.72 (m, 0.5H), 7.38-7.46 (m, 0.5H)

[0273]

(Step 7)

(4R,5S)-4,5-Dimethyl-3,3a,4,5,6,7-hexahydro-benzo[c]isoxazole



(4S,5S)-4,5-Dimethyl-hept-6-enal-oxime (2.0 g) and methanol (40 mL) were mixed under nitrogen gas, and thereto were added trifluoroacetic acid (0.3 mL) and (diacetoxyiodo)benzene (5.5 g) under sodium chloride-ice cooling. The reaction solution was stirred under ice cooling for 20 minutes and stirred at room temperature for


1 hour. Then, thereto were added saturated aqueous sodium hydrogen carbonate solution (20 ml) and sodium sulfite (0.8 g) under ice cooling, and the mixture was stirred at room temperature for 30 minutes. The reaction solution was concentrated under reduced pressure, and the aqueous layer was extracted with ethyl acetate (twice). The organic layer was washed with saturated aqueous sodium chloride solution (twice) and dried over sodium sulfate. Then, sodium sulfate was removed with a filter, and the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, $R_f = 0.1$ (ethyl acetate/n-hexane = 1/20)) to give the title compound (1.4 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.93-0.98 (m, 6H), 1.08-1.34 (m, 2H), 1.77-1.99 (m, 2H), 2.09-2.21 (m, 1H), 2.69-2.78 (m, 2H), 2.78-2.89 (m, 1H), 3.76-3.85 (m, 1H), 4.45-4.52 (m, 1H)

[0274]

(Step 8)

(4R,5S)-7a-[4-(3-tert-Butyl-2,2-difluoro-cyclopropyl)-3-chloro-phenyl]-4,5-dimethyl-octahydro-benzo[c]isoxazole



4-Bromo-1-(3-tert-butyl-2,2-difluoro-cyclopropyl)-2-chloro-benzene (648 mg) and tetrahydrofuran (2.6 mL) were mixed under argon gas, and thereto was added dropwise 1.5M

5 n-butyllithium/n-hexane solution (13 mL) under cooling at -78°C. The reaction solution was stirred under cooling at -78°C for 1 hour (Reaction solution A). To a mixed solution of (4R,5S)-4,5-dimethyl-3,3a,4,5,6,7-hexahydrobenzo[c]isoxazole (245 mg) in toluene (6.5 mL) were added

10 boron trifluoride-diethyl ether complex and Reaction solution A under cooling at -78°C. The reaction solution was stirred for 2 hours, and then thereto was added saturated aqueous ammonium chloride solution (8 mL) under cooling at -78°C. The mixture was stirred at room

15 temperature, and then thereto was added ethyl acetate. The mixture was separated. The organic layer was washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced

20 pressure. The resulting residue was purified by silica gel


column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, $R_f = 0.30$ (ethyl acetate/n-hexane = 1/10)) to give the title compound (357 mg, 13w% of ethyl acetate inclusive).

5 $^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.97-1.01 (m, 6H), 1.05-1.20 (m, 11H), 1.27-1.37 (m, 1H), 1.59-1.78 (m, 2H), 1.83-1.99 (m, 2H), 2.16-2.28 (m, 1H), 2.71-2.83 (m, 1H), 3.43-3.51 (m, 1H), 3.76-3.82 (m, 1H), 5.83 (br s, 1H), 7.11-7.17 (m, 1H), 7.42 -7.49 (m, 1H), 7.58-7.67 (m, 1H)

10 [0275]

(Step 9)

{ (5S,6R)-2-Amino-2-[4-(3-tert-butyl-2,2-difluoro-cyclopropyl)-3-chloro-phenyl]-5,6-dimethyl-cyclohexyl}methanol



15

(4R,5S)-7a-[4-(3-tert-Butyl-2,2-difluoro-cyclopropyl)-3-chloro-phenyl]-4,5-dimethyl-octahydro-benzo[c]isoxazole (311 mg) was mixed in acetic acid (4.7 mL), tetrahydrofuran (1.6 mL), and water (1.6 mL) under nitrogen gas, and thereto was added zinc powder (510 mg) in several parts

20


under heating at 60°C. The reaction solution was stirred under heating at 60°C for 2 hours 20 minutes. To the reaction solution was added dropwise ammonia water (8 ml) under ice cooling. The aqueous layer was extracted with cyclopentyl methyl ether (three times), and then washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (492 mg).

¹H-NMR (400 MHz, CDCl₃) 1.03-1.06 (m, 6H), 1.11 (s, 9H), 1.18-1.33 (m, 2H), 1.34-1.42 (m, 1H), 1.44-1.77 (m, 5H), 2.01-2.14 (m, 1H), 2.73-2.83 (m, 1H), 3.21-3.25 (m, 1H), 3.50-3.58 (m, 1H), 3.75-3.82 (m, 2H), 7.13-7.21 (m, 1H), 7.30-7.39 (m, 1H), 7.44-7.56 (m, 1H)

[0276]

(Step 10)

Methyl 4-(3-((3R,4S)-1-[4-(3-tert-butyl-2,2-difluorocyclopropyl)-3-chloro-phenyl]-2-hydroxymethyl-3,4-dimethylcyclohexyl)ureido)bicyclo[2.1.1]hexane-1-carboxylate



3-(Methoxycarbonyl)bicyclo[2.1.1]pentane-1-carboxylic acid (59 mg) and toluene (1.2 mL) were mixed under nitrogen gas, and thereto were added diphenylphosphoryl azide (75
 5 μ L) and triethylamine (49 μ L) at room temperature. The reaction solution was stirred under heating at 120°C for 1 hour. The reaction solution was added dropwise to a mixed solution of {(5S,6R)-2-amino-2-[4-(3-tert-butyl-2,2-difluoro-cyclopropyl)-3-chloro-phenyl]-5,6-dimethyl-
 10 cyclohexyl}methanol (112 mg) in tetrahydrofuran (2.3 mL) under ice cooling. The reaction solution was stirred at room temperature for 13 hours, and then thereto was added N,N,N'-trimethyl-ethane-1,2-diamine (7.2 μ L). The mixture was concentrated under reduced pressure. The resulting
 15 residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: acetone/n-hexane, R_f = 0.40 (acetone/n-hexane = 1/4)) to give the title compound (201 mg, 31w% of ethyl acetate inclusive).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) 0.93-1.06 (m, 8H), 1.10 (s, 9H),
 20 1.54-1.83 (m, 5H), 1.89-1.93 (m, 5H), 1.98-2.02 (m, 1H),

2.12-2.29 (m, 4H), 2.70-2.81 (m, 1H), 2.99-3.09 (m, 1H),
 3.29-3.35 (m, 1H), 3.67 (s, 3H), 3.70-3.75 (m, 1H), 4.81
 (br s, 1H), 6.28 (br s, 1H), 7.09-7.23 (m, 2H), 7.26-7.36
 (m, 1H)

5 [0277]

(Step 11)

Methyl 4-{(5R,6S)-8a-[4-(3-tert-butyl-2,2-difluoro-
 cyclopropyl)-3-chloro-phenyl]-5,6-dimethyl-2-oxo-
 1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-
 10 yl}bicyclo[2.1.1]hexane-1-carboxylate



Methyl 4-(3-{(3R,4S)-1-[4-(3-tert-butyl-2,2-difluoro-
 cyclopropyl)-3-chloro-phenyl]-2-hydroxymethyl-3,4-dimethyl-
 cyclohexyl}ureido)bicyclo[2.1.1]hexane-1-carboxylate (140
 15 mg) and chloroform (1.4 mL) were mixed under nitrogen gas,
 and thereto were added (diacetoxyiodo)benzene (88 mg) and
 2,2,6,6-tetramethylpiperidin-1-oxyl radical (3.8 mg) at
 room temperature. The reaction solution was stirred at
 room temperature for 14 hours, and then thereto were added
 20 saturated aqueous sodium hydrogen carbonate solution and

sodium sulfite at room temperature. The mixture was stirred at room temperature for 30 minutes, and then extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium chloride solution, and then
5 dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was mixed with toluene (2.8 mL), and thereto was added pentafluoroaniline trifluoromethanesulfonate (4 mg) at room
10 temperature. The reaction solution was stirred for 7 hours under heating at 120°C, and then the reaction solution was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane,
15 $R_f = 0.25$ (ethyl acetate/n-hexane = 3/7)) to give the title compound (36 mg).

This racemate was purified with Recycling Preparative Liquid Chromatograph to give a single enantiomer of the title compound (9 mg).

20 Purification conditions for the preparative chromatography are shown as follows.

Preparative apparatus: Recycling Preparative Liquid Chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

25 Column: Daicel CHIRALPAK IA 2.0 cmφ x 25 cm

Mobile phase: n-hexane/2-propanol = 85/15

Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement of the resulting compound with a chiral
 5 column showed 6.9 minutes of the retention time for the
 resulting enantiomer with > 99%ee of optical purity. The
 retention time for the opposite enantiomer was 10.3 minutes.

Analytical conditions for the chiral column are shown
 as follows.

10 Measurement apparatus: HPLC system, Shimadzu Corporation,
 High-Performance Liquid Chromatograph Prominence

Column: Daicel CHIRALPAK IA-3 0.46 cm ϕ x 15 cm

Column temperature: 40°C

Mobile phase: n-hexane/2-propanol = 85/15

15 Flow rate: 1.0 mL/min

Detection: UV (254 nm)


¹H-NMR (400 MHz, CDCl₃) 0.62-0.67 (m, 3H), 0.99-1.05 (m,
 5H), 1.12 (s, 9H), 1.33-1.47 (m, 2H), 1.56-1.84 (m, 3H),
 1.93-2.13 (m, 9H), 2.22-2.31 (m, 1H), 2.70-2.83 (m, 1H),
 20 3.70 (s, 3H), 4.44 (br s, 1H), 5.96 (br s, 1H), 7.13-7.20
 (m, 1H), 7.21-7.23 (m, 1H), 7.38-7.40 (m, 1H)

[0278]

(Step 12)

4-[(5R,6S)-8a-[4-(3-tert-Butyl-2,2-difluoro-cyclopropyl)-3-
 25 chloro-phenyl]-5,6-dimethyl-2-oxo-1,5,6,7,8,8a-hexahydro-

2H-quinazolin-3-yl}bicyclo[2.1.1]hexane-1-carboxylic acid




Methyl 3-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-
2-oxo-1,5,6,7,8,8a-hexahydro-2H-quinazolin-3-

5 yl}bicyclo[1.1.1]pentane-1-carboxylate (8.7 mg) was mixed
in tetrahydrofuran (87 μ L) and methanol (136 μ L) under
nitrogen gas, and thereto was added 2N aqueous sodium
hydroxide solution (16 μ L) at room temperature. The
reaction solution was stirred at room temperature for 8
10 hours, and then thereto was added water (1 mL), followed by
addition of 2N hydrochloric acid under ice cooling. Then,
the mixture was stirred under ice cooling, and the
precipitated solid was collected by filtration to give the
title compound (7 mg).

15 [0279]

(Intermediate Step 1)

1-(4-Bromo-2-chloro-phenyl)-3,3-dimethyl-butan-1-ol




4-Bromo-2-chloro-1-iodo-benzene (10 g) and tetrahydrofuran (100 mL) were mixed under argon gas, and thereto was added dropwise 2.0M isopropylmagnesium chloride/tetrahydrofuran solution (17 mL) under cooling at -40°C. The mixture was stirred under cooling at -40°C for 1.5 hours, and then thereto was added dropwise 3,3-dimethylbutyraldehyde (4.8 mL). The mixture was stirred under cooling at -40°C for 1 hour 20 minutes, and then thereto was added 2N hydrochloric acid (17 mL) under ice cooling. Then, thereto was added ethyl acetate, and the mixture was separated. The organic layer was washed with saturated aqueous sodium chloride solution and dried over magnesium sulfate. Then, magnesium sulfate was removed with a filter, and the filtrate was concentrated under reduced pressure and azeotroped with toluene to give the title compound (9.8 g, 9.0w% of toluene inclusive).

¹H-NMR (400 MHz, CDCl₃) 1.03 (s, 9H), 1.53-1.55 (m, 2H), 1.75-1.77 (m, 1H), 5.19-5.24 (m, 1H), 7.37-7.48 (m, 3H)

[0280]

(Intermediate Step 2)

4-Bromo-2-chloro-1-((E)-3,3-dimethyl-but-1-enyl)benzene



1-(4-Bromo-2-chloro-phenyl)-3,3-dimethyl-butan-1-ol


(800 mg) and toluene (12 mL) were mixed, and thereto was added pentafluoroanilinium trifluoromethanesulfonate (494 mg) at room temperature. The mixture was stirred under heating at 120°C for 3 hours, and then the reaction solution was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate/n-hexane, R_f = 0.8 (ethyl acetate/n-hexane = 5/95)) to give the title compound (9.5 g, 16w% of ethyl acetate inclusive).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.12 (s, 9H), 6.16-6.24 (m, 1H), 6.53-6.62 (m, 1H), 7.27-7.38 (m, 2H), 7.47-7.50 (m, 1H)

[0281]

(Intermediate Step 3)

4-Bromo-1-(3-tert-butyl-2,2-difluoro-cyclopropyl)-2-chloro-benzene



To 4-bromo-2-chloro-1-((E)-3,3-dimethyl-but-1-enyl)benzene (800 mg) were added tetra-n-butylammonium bromide (48 mg) and (bromodifluoromethyl)trimethylsilane (0.69 mL) at room temperature under argon gas. The mixture
 5 was stirred under heating at 110°C overnight, and then thereto were added tetra-n-butylammonium bromide (48 mg) and (bromodifluoromethyl)trimethylsilane (0.69 mL). The mixture was stirred under heating at 120°C for 8 hours. Then, thereto was added water (12 mL) and ethyl acetate at
 10 room temperature, and the mixture was separated. The organic layer was washed with saturated aqueous sodium chloride solution and dried over magnesium sulfate. Then, magnesium sulfate was removed with a filter, and The resulting residue was purified by silica gel column
 15 chromatography (eluent: ethyl acetate/n-hexane, R_f = 0.75 (ethyl acetate/n-hexane = 5/95)) to give the title compound (697 mg, 7w% of toluene inclusive).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.10 (s, 9H), 1.60-1.73 (m, 1H), 2.66-2.78 (m, 1H), 7.01-7.08 (m, 1H), 7.31-7.37 (m, 1H),
 20 7.54-7.58 (m, 1H)

[0282]

Example 118

(Step 1)

Ethyl (E)-4,4,5-trimethylhepta-2,6-dienoate




60wt% Sodium hydride (1.1 g) and tetrahydrofuran (28 mL) were mixed under ice cooling and argon gas, and thereto was added dropwise triethyl phosphonoacetate (6.4 g). The reaction solution was stirred at room temperature for 30 minutes. Then, thereto was added dropwise a mixed solution of 2,2,3-trimethylpent-4-enal in tetrahydrofuran (28 mL) under ice cooling. The reaction solution was stirred under ice cooling for 10 minutes, and then stirred at room temperature for 12.5 hours. To the reaction solution were added saturated aqueous ammonium chloride solution (22 mL) and water (6 mL), and the mixture was concentrated under reduced pressure. The resultant aqueous layer was extracted with ethyl acetate (20 mL, twice). The organic layer was washed with saturated aqueous sodium chloride solution (10 mL, twice) and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (3.2 g).

¹H-NMR (400MHz, CDCl₃) 0.94 (d, J = 6.94 Hz, 3H), 1.01 (s, 3H), 1.02 (s, 3H), 1.30 (t, J=7.17Hz, 3H), 2.05-2.14 (m, 1H), 4.19 (q, J=7.17Hz, 2H), 4.97-5.03 (m, 2H), 5.65-5.75 (m, 2H), 6.95 (d, J=15.95Hz, 1H)

[0283]

(Step 2)

Ethyl 4,4,5-trimethylhept-6-enoate



5 Ethyl (E)-4,4,5-trimethylhepta-2,6-dienoate (3.2 g) and copper (I) chloride (1.6 g) were mixed in tetrahydrofuran (44 mL) and methanol (25 mL) under argon gas, and thereto was added sodium borohydride over 1 hour under ice cooling. The reaction solution was stirred under

10 ice cooling for 2 hours, and then thereto was added 1N hydrochloric acid (32 mL). Solid substances were removed with a filter, and then the resultant was washed with ethyl acetate and water. The filtrate was concentrated under reduced pressure. The resulting aqueous layer was

15 extracted with ethyl acetate (10 mL) twice. The organic layer was washed with 1N hydrochloric acid (9 mL) and saturated aqueous sodium chloride solution (9 mL) and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under

20 reduced pressure. To The resulting residue were added silica gel (15 g), ethyl acetate (16 mL), and n-hexane (16 mL), and the mixture was stirred for 30 minutes. Silica gel was removed with a filter, and then the filtrate was


washed with a mixed solvent of ethyl acetate/n-hexane = 1/1 (64 mL). The filtrate was concentrated under reduced pressure to give a crude product of the title compound (3.4 g).

5 $^1\text{H-NMR}$ (400MHz, CDCl_3) 0.83 (s, 3H), 0.83 (s, 3H), 0.95 (d, $J=6.94\text{Hz}$, 3H), 1.25 (t, $J=7.13\text{Hz}$, 3H), 1.54-1.61 (m, 2H), 1.94-2.02 (m, 1H), 2.22-2.29 (m, 2H), 4.12 (q, $J=7.13\text{Hz}$, 2H), 4.94-5.02 (m, 3H), 5.68-5.82 (m, 2H)

[0284]

10 (Step 3)

4,4,5-Trimethylhept-6-enal



Ethyl 4,4,5-trimethylhept-6-enoate (3.4 g) and dichloromethane (170 mL) were mixed under argon gas, and thereto was added dropwise 1.02M diisobutylaluminum hydride/n-hexane solution (25 mL) under cooling at -78°C . The reaction solution was stirred under cooling at -78°C for 2 hours. To the reaction solution was added methanol (10 mL) under cooling at -78°C . The reaction solution was warmed to 0°C , and thereto was added a saturated aqueous solution of Rochelle salt (85 mL). The reaction solution was stirred at room temperature for 3 hours. The aqueous layer was extracted with diethyl ether (40 mL, twice). The


organic layer was washed with saturated aqueous sodium chloride solution (10 mL, twice) and dried over magnesium sulfate. Magnesium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure to give a crude product of the title compound (2.5 g).

$^1\text{H-NMR}$ (400MHz, CDCl_3) 0.82-0.84 (m, 6H), 0.95-0.97 (m, 3H), 1.54-1.58 (m, 2H), 1.96-2.04 (m, 1H), 2.35-2.43 (m, 2H), 4.93-4.99 (m, 3H), 5.71-5.81 (m, 1H), 9.77 (t, $J=1.97\text{Hz}$, 1H)

[0285]

(Step 4)

4,4,5-Trimethylhept-6-enal oxime



4,4,5-Trimethylhept-6-enal (1.9 g) was mixed in ethanol (25 mL) and water (12 mL) under argon gas, and thereto were added sodium acetate (7.0 g) and hydroxylamine hydrochloride (3.4 g) at room temperature. The reaction solution was stirred under heating at 60°C for 14.5 hours. The reaction solution was concentrated under reduced pressure, and thereto were added ethyl acetate and water. The mixture was separated. The aqueous layer was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium chloride solution (twice) and


dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash
 5 purification systems, eluent: ethyl acetate/n-hexane, 5/95 to 20/80) to give the title compound (1.8 g).

$^1\text{H-NMR}$ (400MHz, CDCl_3) 0.83-0.98 (m, 9H), 1.37-1.43 (m, 2H), 1.97-2.08 (m, 1H), 2.13-2.20 (m, 1H), 2.31-2.37 (m, 1H), 4.93-5.00 (m, 3H), 5.51-5.81 (m, 2H), 6.67-6.73 (m, 1H),
 10 7.00 (t, $J=1.27\text{Hz}$, 0.5H), 7.41 (t, $J=6.13\text{Hz}$, 0.5H)

[0286]

(Step 5)

4,5,5-Trimethyl-3,3a,4,5,6,7-hexahydrobenzo[c]isoxazole



15 4,4,5-Trimethylhept-6-enal oxime (1.8 g) and methanol (36 mL) were mixed under nitrogen gas, and thereto was added trifluoroacetic acid (0.27 mL) under sodium chloride-ice cooling, followed by addition of (diacetoxyiodo)benzene (4.5 g) over 1 hour. The reaction solution was stirred
 20 under ice cooling for 20 minutes and at room temperature for 1.5 hours. Then, thereto were added saturated aqueous sodium hydrogen carbonate solution (18 mL) and sodium sulfite (0.66 g) under ice cooling, and the mixture was


stirred at room temperature for 30 minutes. The reaction solution was concentrated under reduced pressure. The aqueous layer was extracted with ethyl acetate (10 mL, twice). The organic layer was washed with saturated aqueous sodium chloride solution (5 mL, twice) and dried over sodium sulfate. Sodium sulfate was removed with a filter, and then the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane = 5/95 to 30/70), and then purified further by silica gel column chromatography (Biotage flash purification systems, eluent: acetone/n-hexane = 5/95 to 20/80) and azeotroped with n-hexane to give the title compound (0.42 g).

¹H-NMR (400MHz, CDCl₃) 0.80-0.98 (m, 9H), 1.25-1.45 (m, 2H), 1.63-1.70 (m, 1H), 3.76 (dd, J=10.87, 7.86Hz, 1H), 4.10-4.15 (m, 1H), 4.51 (dd, J=10.40, 7.86Hz, 1H)

[0287]

(Step 6)

7a-(3-Chloro-4-(3,3-dimethylbutyl)phenyl)-4,5,5-trimethyloctahydrobenzo[c]isoxazole




This step was performed according to Example 3 Step 6.

¹H-NMR (400MHz, CDCl₃) 0.93 (d, J=6.70Hz, 3H), 0.98 (s, 9H),
 0.99 (s, 3H), 1.01 (s, 3H), 1.27-1.34 (m, 1H), 1.36-1.41 (m,
 2H), 1.42-1.48 (m, 2H), 1.79-1.86 (m, 1H), 1.93-2.00 (m,
 1H), 2.28 -2.35 (m, 1H), 2.62-2.69 (m, 2H), 3.52 (dd,
 J=7.40, 5.55Hz, 1H), 3.81 (d, J=7.40Hz, 1H), 5.88 (s, 1H),
 7.16 (d, J=8.04Hz, 1H), 7.41 (dd, J=8.04, 1.91Hz, 1H), 7.56
 (d, J=1.91Hz, 1H)

[0288]

(Step 7)

(2-Amino-2-(3-chloro-4-(3,3-dimethylbutyl)phenyl)-5,5,6-
 trimethylcyclohexyl)methanol



15


This step was performed according to Example 3 Step 7.

¹H-NMR (400MHz, CDCl₃) 0.94 (s, 3H), 0.95-0.99 (m, 12H),
 1.05 (s, 3H), 1.18-1.23 (m, 1H), 1.36-1.87 (m, 4H), 2.22-
 2.31 (m, 1H), 2.65-2.70 (m, 2H), 3.28-3.32 (m, 1H), 3.45-
 3.49 (m, 1H), 3.77-3.83 (m, 1H), 7.19 (d, J=8.09Hz, 1H),
 5 7.30 (dd, J=8.09, 2.08Hz, 1H), 7.45 (d, J=2.08 Hz, 1H)

[0289]

(Step 8)

Methyl 3-(3-{1-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2-
 (hydroxymethyl)-3,4,4-
 10 trimethylcyclohexyl}ureido)bicyclo[1.1.1]pentane-1-
 carboxylate




This step was performed according to Example 54 Step 8.

¹H-NMR (400MHz, CDCl₃) 0.94 (s, 6H), 0.97 (s, 9H), 1.03 (s,
 15 3H), 1.29-1.68 (m, 5H), 1.86-2.00 (m, 2H), 2.36 (s, 6H),
 2.59-2.67 (m, 2H), 2.86-2.92 (m, 1H), 3.34-3.40 (m, 1H),
 3.62-3.69 (m, 4H), 4.71 (br s, 1H), 6.42 (s, 1H), 7.10-7.17
 (m, 2H), 7.22-7.27 (m, 3H)

[0290]

(Step 9)

Methyl 3-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,6,6-trimethyl-2-oxo-1,5,6,7,8,8a-hexahydroquinazolin-3(2H)-yl}bicyclo[1.1.1]pentane-1-carboxylate



This step was performed according to Example 54 Step 9. The product was purified with a chiral preparative column. Purification conditions for the preparative column are shown as follows.

10 Preparative apparatus: Recycling Preparative Liquid Chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

Column: Daicel CHIRALPAK IA 2.0cmφ x 25 cmL

Mobile phase: n-hexane/2-propanol = 90/10

15 Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement with a chiral column showed 11.2 minutes of the retention time for the resulting title compound (4.2 minutes of the retention time for the enantiomer of the title compound) with > 99% ee purity. Analytical conditions

20

for the chiral column are shown as follows.

Measurement apparatus: HPLC system, Shimadzu Corporation,
High-Performance Liquid Chromatograph Prominence

Column: Daicel CHIRALPAK IA-3 0.46cmφ x 15 cmL

5 Column temperature: 30°C

Mobile phase: n-hexane/2-propanol = 90/10

Flow rate: 1.0 mL/min


Detection: UV (254 nm)

¹H-NMR (400MHz, CDCl₃) 0.47 (d, J=7.40Hz, 3H), 0.82 (s, 3H),
10 0.98 (s, 9H), 1.04 (s, 3H), 1.18-1.32 (m, 2H), 1.40-1.48 (m,
2H), 1.96-2.11 (m, 2H), 2.35-2.41 (m, 1H), 2.48 (s, 6H),
2.63-2.68 (m, 2H), 3.71 (s, 3H), 4.46 (br s, 1H), 5.96-5.98
(m, 1H), 7.16-7.18 (m, 2H), 7.30-7.32 (m, 1H)

[0291]

15 (Step 10)

3-{7a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5,6,6-
trimethyl-2-oxo-1,2,5,6,7,7a-hexahydro-3H-
cyclopenta[d]pyrimidin-3-yl}bicyclo[1.1.1]pentane-1-
carboxylic acid




This step was performed according to Example 54 Step 10.

[0292]

Example 120

5 (Step 1)

Methyl 4-(3-{3-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-4-hydroxymethyl-5,6,6-trimethyl-tetrahydro-pyran-3-yl}ureido)bicyclo[2.1.1]hexane-1-carboxylate



10 Methyl bicyclo[2.1.1]hexane-1,4-dicarboxylate (144 mg)
and toluene (2.89 mL) were mixed under argon gas, and
thereto were added diphenylphosphoryl azide (0.183 mL) and
triethylamine (0.118 mL) at room temperature. The reaction
solution was stirred under heating at 120°C for 1 hour. To
15 the reaction solution was added tetrahydrofuran (2.89 mL),
and the mixture was added dropwise to a mixed solution of
{5-amino-5-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-2,2,3-
trimethyl-tetrahydro-pyran-4-yl}methanol (251 mg, 21w% of
cyclopentyl methyl ether inclusive) in tetrahydrofuran


(5.02 mL) over 10 minutes under ice cooling. The reaction solution was stirred at room temperature for 13 hours, and then thereto was added N,N,N'-trimethylethylenediamine (0.0176 mL). The reaction solution was concentrated under
 5 reduced pressure, and the resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: acetone/n-hexane, R_f = 0.29 (acetone/n-hexane = 1/2)) to give a crude product of the title compound (906 mg, 67w% of ethyl acetate inclusive).

10 ¹H-NMR (400 MHz, CDCl₃) 0.94 (d, J=7.2Hz, 3H), 0.96 (s, 9H), 1.08 (s, 1.05H), 1.20 (s, 1.95H), 1.31 (s, 1.95H), 1.40-1.48 (m, 2H), 1.42 (s, 1.05H), 1.66-2.34 (m, 7H), 1.91 (s, 1.40H), 2.03 (s, 2.60H), 2.60-2.66 (m, 2H), 3.55-3.69 (m, 2H), 3.66 (s, 1.95H), 3.69 (s, 1.05H), 3.81-3.87 (m, 1.30H),
 15 4.02-4.08 (m, 0.70H), 4.69 (s, 0.35H), 5.03 (s, 0.65H), 5.80 (s, 0.65H), 6.19 (s, 0.35H), 7.12-7.17 (m, 1.65H), 7.25 (d, J=1.8Hz, 0.65H), 7.46 (d, J=8.3Hz, 0.35H), 7.53 (d, J=1.8Hz, 0.35H)

[0293]

20 (Step 2)

Methyl 4-{8a-[3-chloro-4-(3,3-dimethyl-butyl)phenyl]-5,6,6-trimethyl-2-oxo-1,2,5,6,8,8a-hexahydro-pyrano[3,4-d]pyrimidin-3-yl}bicyclo[2.1.1]hexane-1-carboxylate



Methyl

4-(3-{3-[3-chloro-4-(3,3-dimethyl-

butyl)phenyl]-4-hydroxymethyl-5,6,6-trimethyl-tetrahydro-
pyran-3-yl}ureido)bicyclo[2.1.1]hexane-1-carboxylate (906

5 mg, 67w% of ethyl acetate inclusive) and chloroform (3.0
mL) were mixed, and then thereto were added

(diacetoxyiodo)benzene (202 mg) and 2,2,6,6-
tetramethylpiperidin-1-oxyl radical (8.6 mg) at room
temperature. The reaction solution was stirred at room

10 temperature for 14 hours, and then thereto were added
20w/w% aqueous sodium sulfite solution (2 mL) and saturated
aqueous sodium hydrogen carbonate solution (2 mL) at room
temperature. The mixture was stirred at room temperature

for 30 minutes. The aqueous layer was extracted with ethyl
15 acetate (twice). The organic layers were washed with

saturated aqueous sodium chloride solution and dried over
magnesium sulfate. Magnesium sulfate was removed with a
filter, and then the filtrate was concentrated under
reduced pressure. The resulting residue and toluene (6.1

20 mL) were mixed, and then thereto was added

pentafluoroaniline trifluoromethanesulfonate (9.1 mg) at room temperature. The reaction solution was stirred under heating at 120°C for 2 hours and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (Biotage flash purification systems, eluent: ethyl acetate/n-hexane, R_f = 0.42 (ethyl acetate/n-hexane = 2/3)) to give a diastereomeric mixture of the title compound (245 mg). The diastereomeric mixture was purified with a Recycling Preparative Liquid Chromatograph to give a single enantiomer of the title compound (50.3 mg).

Purification conditions for the preparative chromatography are shown as follows.

Preparative apparatus: Recycling preparative liquid chromatograph LC-92XX NEXT SERIES, Japan Analytical Industry Co., Ltd.

Column: Daicel CHIRALPAK IA 2.0 cmφ x 25 cm

Mobile phase: n-hexane:2-propanol = 85:15

Flow rate: 10.0 mL/min

Detection: UV (254 nm)

Measurement of the resulting compound with a chiral column showed 12.9 minutes of the retention time for the resulting enantiomer with > 99% ee of optical purity. The retention time for a diastereomer on the methyl group was 10.6 minutes, and the retention time for a diastereomer on

the phenyl group as well as the opposite enantiomer was 6.6 or 7.5 minutes.

Analytical conditions for the chiral column are shown as follows.

5 Measurement apparatus: HPLC system, Shimadzu Corporation,
High-Performance Liquid Chromatograph Prominence

Column: Daicel CHIRALPAK IA-3 0.46 cm ϕ x 15 cm

Column temperature: 30°C

Mobile phase: n-hexane:2-propanol = 85:15

10 Flow rate: 1.0 mL/min


Detection: UV (254 nm)

¹H-NMR (400 MHz, CDCl₃) 0.54 (d, J=7.2Hz, 3H), 0.98 (s, 9H),
1.12 (s, 3H), 1.34 (s, 3H), 1.41-1.46 (m, 2H), 2.00-2.10 (m,
9H), 2.64-2.69 (m, 2H), 3.73 (s, 3H), 3.90 (d, J=12.0Hz,
15 1H), 4.22 (d, J=11.8Hz, 1H), 4.38 (s, 1H), 5.91 (s, 1H),
7.17 (d, J=8.1Hz, 1H), 7.31 (dd, J=8.0, 2.0Hz, 1H), 7.47 (d,
J=2.1Hz, 1H)

[0294]

(Step 3)

20 4-{(S)-8a-[3-Chloro-4-(3,3-dimethyl-butyl)phenyl]-5,6,6-
trimethyl-2-oxo-1,2,5,6,8,8a-hexahydro-pyrano[3,4-
d]pyrimidin-3-yl}bicyclo[2.1.1]hexane-1-carboxylic acid



Methyl

4-[(S)-8a-[3-chloro-4-(3,3-dimethyl-

butyl)phenyl]-5,6,6-trimethyl-2-oxo-1,2,5,6,8,8a-hexahydro-
pyrano[3,4-d]pyrimidin-3-yl]bicyclo[2.1.1]hexane-1-

5 carboxylate (50.3 mg) was mixed in tetrahydrofuran (0.553 mL) and methanol (0.553 mL), and then thereto was added 2N aqueous solution of sodium hydroxide (0.0951 mL) at room temperature. The reaction solution was stirred at room temperature for 16 hours and then concentrated under
10 reduced pressure. To the resulting residue was added water (1 mL), and thereto was added 2N hydrochloric acid (0.0951 mL) under ice cooling. The precipitated solid was collected by filtration to give the title compound (42.4 mg).

15 [0295]

The following tables show structures, structural information, and physical property data for the above Example compounds and Example compounds prepared according to any of the above methods of Examples.

20 In the tables, "chiral column IA-3" means CHIRALPAK

IA-3 analytical column (Product code: 80524; internal diameter: 4.6 mm, length: 150 mm, particle size: 3 μ m; Daicel Corporation).

5 "Chiral column AS-3R" means CHIRALPAK AS-3R analytical column (Product code: 20824; internal diameter: 4.6 mm, length: 150 mm, particle size: 3 μ m; Daicel Corporation).

Each absolute configuration of the enantiomer shown in the structure of each example compound was estimated according to the following regularities:






- 10 1) a certain regularity in the retention time of chiral column for a methyl or ethyl ester derivative of example compound; and
- 2) a certain regularity in the degree of inhibitory activity against ROR γ transcription of each enantiomer of
- 15 example compound (see Test Example 1);





provided that the absolute configuration of Example 54 compound was determined by single-crystal X-ray structural analysis.




In the following table, the term "methyl ester derivative" or "ethyl ester derivative" described in the





20 structural information means a methyl-ester derivative or ethyl-ester derivative corresponding to each example compound, respectively.




[0296]




Example	Structure	Structural Information
1		Racemate
2		Racemate
3		Enantiomer of Example 4 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 93/7, flow 1 ml/min Retention time: 7.4 min
4		Enantiomer of Example 3 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 93/7, flow 1 ml/min Retention time: 8.7 min
5		Enantiomer of Example 6 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 6.6 min





6		Enantiomer of Example 5 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 6.9 min
7		Enantiomer of Example 8 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 80/20, flow 1 ml/min Retention time: 6.5 min
8		Enantiomer of Example 7 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 80/20, flow 1 ml/min Retention time: 8.8 min
9		Enantiomer of Example 10 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 10.6 min





10		Enantiomer of Example 9 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 12.8 min
11		Optically active. Optical purity was not determined. (Absolute configuration was not determined.) Single diastereomer. Configuration of methyl group on the cyclopentane ring was estimated to be R derived from a starting material. Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 5.3 min
12		Optically active. Optical purity was not determined. (Absolute configuration was not determined.) Single diastereomer. Configuration of methyl group on the cyclopentane ring was estimated to be R derived from a starting material. Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min





		Retention time: 6.2 min
13		Racemate. Estimated structure.
14		Racemate. Estimated structure.
15		Enantiomer of Example 16 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hex./IPA = 85/15, flow rate = 1.0 ml/min, Retention time: 6.4 min Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and benzyloxymethyl was estimated to be trans
16		Enantiomer of Example 15 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hex./IPA = 85/15, flow rate = 1.0 ml/min, Retention time: 8.8 min Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and benzyloxymethyl was





17		estimated to be trans Enantiomer of Example 18 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hex./IPA = 80/20, flow rate = 1.0 ml/min, Retention time: 6.3 min Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and hydroxymethyl was estimated to be trans
18		Enantiomer of Example 17 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hex./IPA = 80/20, flow rate = 1.0 ml/min, Retention time: 7.6 min Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and hydroxymethyl was estimated to be trans
19		Enantiomer of Example 20 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hex./IPA = 90/10, flow rate = 1.0 ml/min, Retention time: 8.7 min Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and methoxymethyl was estimated to be trans





20		Enantiomer of Example 19 (Absolute configuration was not determined, >98%ee) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hex./IPA = 90/10, flow rate = 1.0 ml/min, Retention time: 9.8 min Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and methoxymethyl was estimated to be trans
21		Enantiomer of Example 22 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hex./IPA = 90/10, flow rate = 1.0 ml/min, Retention time: 7.5 min, Optical purity was not determined Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and butoxymethyl was estimated to be trans
22		Enantiomer of Example 21 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hex./IPA = 90/10, flow rate = 1.0 ml/min, Retention time: 8.3 min, Optical purity was not determined Relative configuration





		of 3-chloro-4-(3,3-dimethylbutyl)phenyl and butoxymethyl was estimated to be trans
23		Racemate
24		Racemate
25		Racemate Configuration of hydroxyethyl and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans Single diastereomer
26		Racemate Configuration of benzyloxyethyl and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans Single diastereomer





27		Racemate Configuration of methoxyethyl and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans Single diastereomer
28	 Na⁺	Racemate The number of sodium salt was estimated. Configuration of dimethylaminoethyl and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans Single diastereomer
29		Racemate Configuration of hydroxymethyl and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans Single diastereomer
30		Racemate Configuration of methylacetamide and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans from the synthetic method Single diastereomer





31		Racemate Configuration of nitrilemethyl and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans
32		Enantiomer of Example 33 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 5.9 min
33		Enantiomer of Example 32 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 6.9 min
34		Enantiomer of Example 35 (Absolute configuration was not determined.) (>99%ee for corresponding methyl ester derivative) chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 6.0 min) Single stereoisomer, Relative configuration of methyl and 3-chloro-4-(3,3-dimethylbutyl)phenyl was uncertain





35		Enantiomer of Example 34 (Absolute configuration was not determined.) (>99%ee for corresponding methyl ester derivative chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 8.5 min) Single stereoisomer, Relative configuration of three methyl groups on the cyclopentane ring and 3-chloro-4-(3,3-dimethylbutyl)phenyl was uncertain
36		Racemate
37		Enantiomer of Example 38 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 4.9 min
38		Enantiomer of Example 37 (Absolute configuration was not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min





		Retention time: 8.5 min
39		Racemate
40		Optically active (Absolute configuration was not determined.) (>99%ee, IA-3 IPA/Hex. = 7%, flow rate = 1 ml/min, R.T. = 7.3 min) Enantiomer of Example 41
41		Optically active (Absolute configuration was not determined.) (>99%ee, IA-3 IPA/Hex. = 7%, flow rate = 1 ml/min, R.T. = 12.4 min) Enantiomer of Example 40
42		Estimated structure Racemate





43		Racemate Configuration of benzyloxyethyl and 3- chloro-4-(3,3- dimethylbutyl)phenyl was estimated to be trans
44		Racemate Configuration of hydroxyethyl and 3- chloro-4-(3,3- dimethylbutyl)phenyl was estimated to be trans
45		Racemate Configuration of carboxymethyl and 3- chloro-4-(3,3- dimethylbutyl)phenyl was estimated to be trans
46		Racemate Configuration of methoxyethyl and 3- chloro-4-(3,3- dimethylbutyl)phenyl was estimated to be trans





47	 HCl	Racemate Configuration of aminomethyl and 3- chloro-4-(3,3- dimethylbutyl)phenyl was estimated to be trans
48	 HCl	Racemate Configuration of mesyloxyethyl and 3- chloro-4-(3,3- dimethylbutyl)phenyl was estimated to be trans
49	 HCl	Racemate Configuration of methylaminoethyl and 3- chloro-4-(3,3- dimethylbutyl)phenyl was estimated to be trans The number of hydrochloride salt was estimated
50	 HCl	Racemate Configuration of dimethylaminoethyl and 3-chloro-4-(3,3- dimethylbutyl)phenyl was estimated to be trans The number of hydrochloride salt was estimated





51		Racemate Configuration of carboxamidemethyl and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans
52		Racemate Configuration of nitrilemethyl and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans
53		Optically active Enantiomer of Example 54 (>99%ee for corresponding methyl ester derivative chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 5.3 min)
54		Optically active Enantiomer of Example 53 (>99%ee for corresponding methyl ester derivative, the product of Example 54 Step 9) chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 8.5 min)




59	 <chem>CC(C)CCOC1=CC=C(Cl)C=C1C2=CC(=C(C=C2)C3=CC(=CC=C3)C(=O)NCCC(=O)O)C4=CC(=CC=C4)C5=CC(=CC=C5)C(=O)NCCC(=O)O</chem>	Racemate
60	 <chem>CC(C)(C)C(C)COC1=CC=C(Cl)C=C1C2=CC(=C(C=C2)C3=CC(=CC=C3)C(=O)NCCC(=O)O)C4=CC(=CC=C4)C5=CC(=CC=C5)C(=O)NCCC(=O)O</chem>	Racemate
61	 <chem>CC(C)(C)C(C)COC1=CC=C(Cl)C=C1C2=CC(=C(C=C2)C3=CC(=CC=C3)C(=O)NCCC(=O)O)C4=CC(=CC=C4)C5=CC(=CC=C5)C(=O)NCCC(=O)O</chem>	Racemate
62	 <chem>C1CCC1CCOC2=CC=C(Cl)C=C2C3=CC(=C(C=C3)C4=CC(=CC=C4)C(=O)NCCC(=O)O)C5=CC(=CC=C5)C(=O)NCCC(=O)O</chem>	Racemate

63		Racemate
64		Racemate. Estimated structure.
65		Racemate
66		Racemate Single diastereomer Relative configuration of 3-chloro-4-(3,3- dimethylbutyl)phenyl and methyl group on the cyclohexane ring was uncertain





67		Enantiomer of Example 68 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 5.2 min
68		Enantiomer of Example 67 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 7.8 min
69		Racemate Single diastereomer Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and methyl group on the cyclohexane ring was uncertain Diastereomer of Example 66
70		Optically active (Absolute configuration was not determined.) (Optical purity was not determined) Single diastereomer Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and methyl group on the tetrahydropyran ring was uncertain Enantiomer of Example 71





71		Optically active (Absolute configuration was not determined.) (Optical purity was not determined) Single diastereomer Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and methyl group on the tetrahydropyran ring was uncertain Enantiomer of Example 70
72		Optically active (Absolute configuration was not determined.) (Optical purity was not determined) Single diastereomer Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and methyl group on the tetrahydropyran was uncertain
73		Racemate Configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and methyl group on the cyclohexane ring was estimated to be cis
74		Racemate





75		Estimated to be optically active. (Absolute and relative configurations were not determined.) Estimated structure.
76		Enantiomer of Example 77 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 4.0 min
77		Enantiomer of Example 76 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 6.5 min
78		Optically active compound of Example 73 (Absolute and relative configurations were not determined.) Enantiomer of Example 79 (>99% ee for corresponding methyl ester derivative) chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min





		Retention time: 4.2 min)
79		Optically active compound of Example 73 (Absolute and relative configurations were not determined.) Enantiomer of Example 78 (>99%ee for corresponding methyl ester derivative) chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 7.2 min)
80		Optically active compound of Example 69 (Absolute configuration was not determined.) Enantiomer of Example 81 (>99%ee for corresponding ethyl ester derivative) chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 5.8 min) Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and methyl group on the cyclohexane ring was uncertain
81		Optically active compound of Example 69 (Absolute configuration was not determined.) Enantiomer of Example 80 (>99%ee for corresponding ethyl ester derivative) chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 8.6 min)

		Relative configuration of 3-chloro-4-(3,3-dimethylbutyl)phenyl and methyl group on the cyclohexane ring was uncertain
82		Optically active compound of Example 74 (Absolute configuration was not determined.) Enantiomer of Example 83 (>99%ee for corresponding ethyl ester derivative chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 5.2 min)
83		Optically active compound of Example 74 (Absolute configuration was not determined.) Enantiomer of Example 82 (>99%ee for corresponding ethyl ester derivative chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 8.9 min)
84		Enantiomer of Example 85 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 3.9 min




85		Enantiomer of Example 84 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 6.3 min
86		Enantiomer of Example 87 (Absolute configuration was not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 80/20, flow 1 ml/min Retention time: 3.1 min
87		Enantiomer of Example 86 (Absolute configuration was not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 80/20, flow 1 ml/min Retention time: 5.2 min
88		Racemate 1:1 diastereomeric mixture regarding the substituents on cyclopropyl




89		Enantiomer of Example 90 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 4.6 min
90		Enantiomer of Example 89 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 8.4 min
91		Enantiomer of Example 92 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 4.1 min
92		Enantiomer of Example 91 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min





		Retention time: 7.8 min
93		Enantiomer of Example 94 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 4.1 min
94		Enantiomer of Example 93 (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 7.3 min
95		Racemate
96		Racemate





97		Single diastereomer. Estimated structure. Estimated to be racemate.
98		Optically active (Absolute and relative configurations were not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, IPA/Hex. = 10/90, flow 1 ml/min Retention time: 3.7 min
99		Optically active (Absolute and relative configurations were not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, IPA/Hex. = 10/90, flow 1 ml/min Retention time: 6.5 min
100		Racemate




101		Racemate
102		Enantiomer of Example 103 (Absolute configuration was not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 4.3 min
103		Enantiomer of Example 102 (Absolute configuration was not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 9.7 min
104		Optically active (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column AS-3R, MeCN/H ₂ O = 80/20, flow 1 ml/min Retention time: 6.8 min Cyclopentane ring moiety was induced from methyl (1R,3S)-3-

		aminocyclopentanecarboxylate hydrochloride.
105		Optically active (Absolute and relative configurations were not determined.) Analytical conditions for corresponding methyl ester derivative: chiral column AS-3R, MeCN/H₂O = 80/20, flow 1 ml/min Retention time: 7.5 min Cyclopentane ring moiety was induced from methyl (1R,3S)-3-aminocyclopentanecarboxylate hydrochloride.
106		Optically active. (Absolute configuration was not determined.) Optical purity was not determined. Single stereoisomer. Configuration of two methyl groups on the cyclohexane ring was estimated to be cis. Configuration of the two methyl groups and 3-chloro-4-(3,3-dimethylbutyl)phenyl was estimated to be trans from NOE.
107		Optically active. (Absolute configuration was not determined.) Optical purity was not determined. Single stereoisomer. Estimated structure. Configuration of two methyl groups on the cyclohexane ring was estimated to be trans. Analytical conditions for corresponding methyl ester derivative: chiral column IA-3,

		Hexane/IPA = 90/10, flow 1 ml/min Retention time: 9.7 min
108		Diastereomeric mixture (Absolute configuration was not determined.) Enantiomeric mixture of Example 109 and Example 110 (>99%ee for corresponding methyl ester derivative chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 3.9 min)
109		Optically active (Absolute and relative configurations were not determined.) Enantiomer of Example 108 Diastereomer of Example 110 (>99%ee for corresponding methyl ester derivative chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 7.6 min)
110		Optically active (Absolute and relative configurations were not determined.) Enantiomer of Example 108 Diastereomer of Example 109 (>99%ee for corresponding methyl ester derivative chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 9.8 min)

111		Optically active (Absolute and relative configurations were not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, IPA/Hex. = 10/90, flow 1 ml/min Retention time: 6.4 min
112		Optically active (Absolute and relative configurations were not determined.) Analytical conditions for corresponding ethyl ester derivative: chiral column IA-3, IPA/Hex. = 10/90, flow 1 ml/min Retention time: 13.3 min
113		Optically active (Absolute and relative configurations were not determined.) Optical purity was not determined Enantiomer of Example 114
114		Optically active (Absolute and relative configurations were not determined.) Optical purity was not determined Enantiomer of Example 113

115		Optically active (Absolute and relative configurations were not determined.) Optical purity was not determined Enantiomer of Example 116
116		Optically active (Absolute and relative configurations were not determined.) Optical purity was not determined Enantiomer of Example 115
117		Optically active (Absolute and relative configurations were not determined.) Single stereoisomer Enantiomer of Example 118 (>99%ee for corresponding ethyl ester derivative chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min Retention time: 4.2 min)
118		Optically active (Absolute and relative configurations were not determined.) Single stereoisomer Enantiomer of Example 117 (>99%ee for corresponding ethyl ester derivative chiral column IA-3, Hexane/IPA = 90/10, flow 1 ml/min)

		Retention time: 11.2 min)
119		Diastereomeric mixture (Absolute configuration was not determined.) Enantiomeric mixture of Example 120 and Example 121 (>99%ee for corresponding ethyl ester derivative chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 6.6, 7.5 min)
120		Optically active (Absolute and relative configurations were not determined.) Single stereoisomer Enantiomer of Example 119 Diastereomer of Example 121 (>99%ee for corresponding ethyl ester derivative chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 12.9 min)
121		Optically active (Absolute and relative configurations were not determined.) Single stereoisomer Enantiomer of Example 119 Diastereomer of Example 120 (>99%ee for corresponding ethyl ester derivative chiral column IA-3, Hexane/IPA = 85/15, flow 1 ml/min Retention time: 10.5 min)

[0297]

Example	¹ H-NMR	MS	
		M+H	M-H
1	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 1.02 (s, 9H), 1.47-1.60 (m, 1H), 1.78-1.88 (m, 1H), 1.99-2.07 (m, 1H), 2.13-2.19 (m, 1H), 2.53-2.64 (m, 2H), 3.69 (s, 2H), 7.09 (d, J = 8.6 Hz, 1H), 7.12 (dd, J = 8.6, 2.1 Hz, 1H), 7.36 (d, J = 2.1 Hz, 1H), 8.09 (d, J = 1.8 Hz, 1H), 9.63 (d, J = 1.8 Hz, 1H).	336	334
2	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.93 (s, 9H), 1.08 (s, 3H), 1.20 (s, 3H), 1.21-1.26 (m, 2H), 1.43 (ddd, J = 12.0, 7.5, 3.0 Hz, 2H), 1.46-1.51 (m, 1H), 1.53-1.63 (m, 2H), 2.00 (t, J = 8.1 Hz, 2H), 2.06 (td, J = 11.2, 5.7 Hz, 1H), 2.23 (dq, J = 12.8, 3.2 Hz, 1H), 2.45-2.49 (m, 1H), 2.99-3.06 (m, 1H), 3.55 (dt, J = 14.0, 6.6 Hz, 1H), 6.27 (s, 1H), 7.12 (d, J = 8.3 Hz, 2H), 7.30 (d, J = 8.3 Hz, 2H), 7.42 (s, 1H), 12.14 (br s, 1H).	413	411
3	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.06 (s, 3H), 1.19 (s, 3H), 1.20-1.24 (m, 1H), 1.37-1.41 (m, 2H), 1.45-1.53 (m, 1H), 2.05 (dt, J = 18.3, 5.9 Hz, 1H), 2.19-2.25 (m, 1H), 2.33 (t, J = 6.8 Hz, 2H), 2.59-2.63 (m, 2H), 3.28-3.35 (m, 1H), 3.67 (dt, J = 14.2, 6.8 Hz, 1H), 6.38 (s, 1H), 7.23 (dd, J = 8.0, 1.7 Hz, 1H), 7.27 (d, J = 7.9 Hz, 1H), 7.42 (d, J = 1.6 Hz, 1H), 7.57 (s, 1H), 12.23 (br s, 1H).	433	431
4	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.06 (s, 3H), 1.19 (s, 3H), 1.20-1.24 (m, 1H), 1.37-1.41 (m, 2H), 1.45-1.53 (m, 1H), 2.05 (dt, J = 18.3, 5.9 Hz, 1H), 2.19-2.25 (m, 1H), 2.33 (t, J = 6.8 Hz, 2H), 2.59-2.63 (m, 2H), 3.28-3.35 (m, 1H), 3.67 (dt, J = 14.2, 6.8 Hz, 1H), 6.38 (s, 1H), 7.23 (dd, J	433	431

	= 8.0, 1.7 Hz, 1H), 7.27 (d, J = 7.9 Hz, 1H), 7.42 (d, J = 1.6 Hz, 1H), 7.57 (s, 1H), 12.23 (br s, 1H).		
5	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.18 (s, 3H), 1.36 (s, 3H), 1.37-1.41 (m, 2H), 2.40 (dt, J = 10.5, 3.5 Hz, 2H), 2.62 (ddd, J = 9.3, 4.2, 3.3 Hz, 2H), 3.37 (dt, J = 14.3, 6.8 Hz, 1H), 3.69-3.76 (m, 1H), 3.83 (d, J = 8.6 Hz, 1H), 4.25 (d, J = 8.6 Hz, 1H), 6.52 (s, 1H), 7.27 (dd, J = 8.0, 1.7 Hz, 1H), 7.30 (d, J = 7.9 Hz, 1H), 7.44 (d, J = 1.6 Hz, 1H), 7.59 (s, 1H), 12.28 (br s, 1H).	435	433
6	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.18 (s, 3H), 1.36 (s, 3H), 1.37-1.41 (m, 2H), 2.40 (dt, J = 10.5, 3.5 Hz, 2H), 2.62 (ddd, J = 9.3, 4.2, 3.3 Hz, 2H), 3.37 (dt, J = 14.3, 6.8 Hz, 1H), 3.69-3.76 (m, 1H), 3.83 (d, J = 8.6 Hz, 1H), 4.25 (d, J = 8.6 Hz, 1H), 6.52 (s, 1H), 7.27 (dd, J = 8.0, 1.7 Hz, 1H), 7.30 (d, J = 7.9 Hz, 1H), 7.44 (d, J = 1.6 Hz, 1H), 7.59 (s, 1H), 12.28 (br s, 1H).	435	433
7	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.93 (s, 9H), 1.27 (s, 3H), 1.33-1.41 (m, 2H), 1.44 (s, 3H), 2.57-2.66 (m, 2H), 3.90-3.97 (m, 2H), 4.28-4.35 (m, 2H), 6.86 (br s, 1H), 7.32-7.42 (m, 2H), 7.43-7.56 (m, 3H), 7.89-7.96 (m, 2H), 8.11 (br s, 1H), 12.90 (br s, 1H).	483	481
8	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.93 (s, 9H), 1.28 (s, 3H), 1.33-1.41 (m, 2H), 2.63 (s, 3H), 3.90-3.98 (m, 2H), 4.28-4.36 (m, 2H), 7.30-7.44 (m, 2H), 7.44-7.56 (m, 3H), 7.87-7.96 (m, 2H), 8.11 (br s, 1H), 12.91 (br s, 1H).	483	481
9	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.92 (s, 9H), 1.15 (s, 3H), 1.22-1.33 (m, 4H), 1.33-1.41 (m, 2H), 1.51-1.60 (m, 1H), 2.09-2.21 (m, 1H), 2.21-2.30 (m, 1H), 2.57-2.63 (m, 2H), 6.67 (s, 1H), 7.29-7.37 (m, 2H), 7.42-7.47 (m, 2H), 7.50-7.53 (m, 1H), 7.86-7.91 (m, 2H), 8.12	481	479

	(br s, 1H), 12.85 (br s, 1H).		
10	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.93 (s, 9H), 1.15 (s, 3H), 1.20-1.34 (m, 4H), 1.33-1.42 (m, 2H), 1.50-1.63 (m, 1H), 2.10-2.22 (m, 1H), 2.22-2.33 (m, 1H), 2.55-2.64 (m, 2H), 6.67 (s, 1H), 7.28-7.37 (m, 2H), 7.41-7.48 (m, 2H), 7.50-7.55 (m, 1H), 7.85-7.93 (m, 2H), 8.12 (s, 1H), 12.85 (br s, 1H).	481	479
11	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.97 (s, 9H), 1.11 (d, J = 19.08 Hz, 3H), 1.40-1.48 (m, 2H), 1.86-2.07 (m, 2H), 2.33-2.41 (m, 1H), 2.51-2.59 (m, 2H), 2.60-2.68 (m, 2H), 2.80-2.92 (m, 1H), 3.49-3.59 (m, 1H), 3.76-3.87 (m, 1H), 6.17-6.20 (m, 1H), 6.51 (br s, 1H), 7.14-7.16 (m, 2H), 7.30 (br s, 1H).	419	417
12	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.97 (s, 9H), 1.11 (d, J = 19.08 Hz, 3H), 1.40-1.48 (m, 2H), 1.86-2.07 (m, 2H), 2.33-2.41 (m, 1H), 2.51-2.59 (m, 2H), 2.60-2.68 (m, 2H), 2.80-2.92 (m, 1H), 3.49-3.59 (m, 1H), 3.76-3.87 (m, 1H), 6.17-6.20 (m, 1H), 6.51 (br s, 1H), 7.14-7.16 (m, 2H), 7.30 (br s, 1H).	419	417
13	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.96 (s, 9H), 1.19-1.29 (m, 5H), 1.37-1.50 (m, 11H), 2.59-2.69 (m, 2H), 3.52-3.61 (m, 2H), 3.86-4.22 (m, 6H), 5.39 (br s, 1H), 6.25 (br s, 1H), 7.07-7.14 (m, 1H), 7.17-7.22 (m, 1H), 7.23-7.25 (m, 1H).	534	532
14	¹ H-NMR (400 MHz, MeOH-d ₄) δ: 1.01 (s, 9H), 1.41-1.49 (m, 2H), 2.05 (s, 3H), 2.55-2.62 (m, 2H), 2.68-2.76 (m, 2H), 3.48-3.64 (m, 2H), 3.80-3.95 (m, 2H), 4.05-4.55 (m, 3H), 6.55 (br s, 1H), 7.21-7.33 (m, 2H), 7.37-7.41 (m, 1H).	448	446
15	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.94 (s, 9H), 1.33-1.38 (m, 2H), 2.35-2.40 (m, 2H), 2.57-2.61 (m, 2H), 3.35-3.41 (m, 3H), 3.65-3.72 (m, 1H), 3.77 (d, J = 8.55 Hz, 1H), 4.33 (d, J = 8.55 Hz, 1H), 4.42 (d, J = 12.95 Hz, 1H), 4.47 (d, J = 12.95 Hz, 1H), 4.74-4.78 (m, 1H), 6.56 (d, J = 1.62 Hz, 1H),	527	525

	7.20-7.35 (m, 7H), 7.41 (s, 1H), 7.65 (s, 1H), 12.18 (br s, 1H).		
16	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.94 (s, 9H), 1.33-1.38 (m, 2H), 2.35-2.40 (m, 2H), 2.57-2.61 (m, 2H), 3.35-3.41 (m, 3H), 3.65-3.72 (m, 1H), 3.77 (d, J = 8.55 Hz, 1H), 4.33 (d, J = 8.55 Hz, 1H), 4.42 (d, J = 12.95 Hz, 1H), 4.47 (d, J = 12.95 Hz, 1H), 4.74-4.78 (m, 1H), 6.56 (d, J = 1.62 Hz, 1H), 7.20-7.35 (m, 7H), 7.41 (s, 1H), 7.65 (s, 1H), 12.18 (br s, 1H).	527	525
17	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.94 (s, 9H), 1.35-1.39 (m, 2H), 2.39 (dd, J = 7.17, 2.77 Hz, 2H), 2.58-2.63 (m, 2H), 3.25-3.33 (m, 1H), 3.35-3.42 (m, 1H), 3.64-3.71 (m, 1H), 3.76 (d, J = 8.32 Hz, 1H), 4.29 (d, J = 8.55 Hz, 1H), 4.55 (dd, J = 7.17, 6.94 Hz, 1H), 4.87 (dd, J = 6.94, 5.78 Hz, 1H), 6.50 (d, J = 1.62 Hz, 1H), 7.26 (d, J = 8.09 Hz, 1H), 7.28 (d, J = 8.09 Hz, 1H), 7.43 (s, 1H), 7.64 (s, 1H), 12.21 (s, 1H).	437	435
18	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.94 (s, 9H), 1.35-1.39 (m, 2H), 2.39 (dd, J = 7.17, 2.77 Hz, 2H), 2.58-2.63 (m, 2H), 3.25-3.33 (m, 1H), 3.35-3.42 (m, 1H), 3.64-3.71 (m, 1H), 3.76 (d, J = 8.32 Hz, 1H), 4.29 (d, J = 8.55 Hz, 1H), 4.55 (dd, J = 7.17, 6.94 Hz, 1H), 4.87 (dd, J = 6.94, 5.78 Hz, 1H), 6.50 (d, J = 1.62 Hz, 1H), 7.26 (d, J = 8.09 Hz, 1H), 7.28 (d, J = 8.09 Hz, 1H), 7.43 (s, 1H), 7.64 (s, 1H), 12.21 (s, 1H).	437	435
19	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.94 (s, 9H), 1.36-1.40 (m, 2H), 2.31-2.45 (m, 2H), 2.59-2.63 (m, 2H), 3.18-3.25 (m, 1H), 3.21 (s, 3H), 3.27-3.33 (m, 1H), 3.35-3.42 (m, 1H), 3.64-3.71 (m, 1H), 3.76 (d, J = 8.55 Hz, 1H), 4.31 (d, J = 8.55 Hz, 1H), 4.71 (dd, J = 7.17, 4.16 Hz, 1H), 6.55 (d, J = 1.62 Hz, 1H), 7.24 (dd, J = 7.86, 1.62 Hz, 1H), 7.29 (d, J = 7.86 Hz, 1H), 7.42 (d, J = 1.62 Hz, 1H), 7.67	451	449

	(s, 1H), 12.22 (s, 1H).		
20	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.94 (s, 9H), 1.36-1.40 (m, 2H), 2.31-2.45 (m, 2H), 2.59-2.63 (m, 2H), 3.18-3.25 (m, 1H), 3.21 (s, 3H), 3.27-3.33 (m, 1H), 3.35-3.42 (m, 1H), 3.64-3.71 (m, 1H), 3.76 (d, J = 8.55 Hz, 1H), 4.31 (d, J = 8.55 Hz, 1H), 4.71 (dd, J = 7.17, 4.16 Hz, 1H), 6.55 (d, J = 1.62 Hz, 1H), 7.24 (dd, J = 7.86, 1.62 Hz, 1H), 7.29 (d, J = 7.86 Hz, 1H), 7.42 (d, J = 1.62 Hz, 1H), 7.67 (s, 1H), 12.22 (s, 1H).	451	449
21	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.09 (s, 9H), 1.36-1.41 (m, 2H), 2.35-2.44 (m, 2H), 2.60-2.64 (m, 2H), 3.20 (dd, J = 8.32, 5.78 Hz, 1H), 3.25-3.29 (m, 1H), 3.36-3.43 (m, 1H), 3.67-3.74 (m, 1H), 3.76 (t, J = 6.36 Hz, 1H), 4.31 (d, J = 8.32 Hz, 1H), 4.58 (dd, J = 6.24, 5.78 Hz, 1H), 6.52 (d, J = 1.85 Hz, 1H), 7.28 (d, J = 7.17 Hz, 1H), 7.30 (d, J = 7.17 Hz, 1H), 7.44 (s, 1H), 7.67 (s, 1H).	493	491
22	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.09 (s, 9H), 1.36-1.41 (m, 2H), 2.35-2.44 (m, 2H), 2.60-2.64 (m, 2H), 3.20 (dd, J = 8.32, 5.78 Hz, 1H), 3.25-3.29 (m, 1H), 3.36-3.43 (m, 1H), 3.67-3.74 (m, 1H), 3.76 (t, J = 6.36 Hz, 1H), 4.31 (d, J = 8.32 Hz, 1H), 4.58 (dd, J = 6.24, 5.78 Hz, 1H), 6.52 (d, J = 1.85 Hz, 1H), 7.28 (d, J = 7.17 Hz, 1H), 7.30 (d, J = 7.17 Hz, 1H), 7.44 (s, 1H), 7.67 (s, 1H).	493	491
23	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.79-0.90 (m, 1H), 0.97 (s, 9H), 1.42-1.46 (m, 2H), 1.78-1.88 (m, 1H), 2.03-2.11 (m, 1H), 2.32-2.42 (m, 2H), 2.46-2.54 (m, 1H), 2.63-2.67 (m, 2H), 5.42 (s, 1H), 6.03 (s, 1H), 6.10 (s, 1H), 7.15 (dd, J = 8.06, 1.61 Hz, 1H), 7.18 (d, J = 8.06 Hz, 1H), 7.27 (d, J = 1.61 Hz, 1H).	333	-
24	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.93	405	403

	(s, 9H), 1.35-1.39 (m, 3H), 1.65-1.73 (m, 1H), 1.89-1.97 (m, 1H), 2.12-2.17 (m, 1H), 2.22-2.34 (m, 3H), 2.40-2.45 (m, 1H), 2.57-2.61 (m, 2H), 3.36-3.42 (m, 1H), 3.52-3.59 (m, 1H), 6.29 (s, 1H), 7.19 (dd, J = 7.86, 1.85 Hz, 1H), 7.25 (d, J = 7.86 Hz, 1H), 7.37 (d, J = 1.85 Hz, 1H), 7.61 (br s, 1H).		
25	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.91-0.82 (m, 1H), 0.93 (s, 9H), 1.32-1.40 (m, 3H), 1.56-1.64 (m, 1H), 1.82-1.90 (m, 2H), 2.25-2.35 (m, 3H), 2.57-2.61 (m, 2H), 2.75-2.83 (m, 1H), 3.39-3.46 (m, 3H), 3.60-3.67 (m, 1H), 4.39 (t, J = 5.09 Hz, 1H), 6.34 (d, J = 1.85 Hz, 1H), 7.22 (dd, J = 8.09, 1.39 Hz, 1H), 7.25 (d, J = 8.09 Hz, 1H), 7.41 (d, J = 1.39 Hz, 1H), 7.56 (s, 1H), 12.20 (br s, 1H).	449	447
26	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.95 (s, 9H), 1.01-1.09 (m, 1H), 1.39-1.43 (m, 2H), 1.54-1.63 (m, 1H), 1.64-1.73 (m, 1H), 1.88-2.03 (m, 2H), 2.42 (dd, J = 11.21, 6.13 Hz, 1H), 2.49-2.53 (m, 2H), 2.59-2.63 (m, 2H), 2.89-2.97 (m, 1H), 3.39-3.54 (m, 3H), 3.75-3.82 (m, 1H), 4.46 (d, J = 11.79 Hz, 1H), 4.50 (d, J = 11.79 Hz, 1H), 6.04 (d, J = 1.62 Hz, 1H), 6.14 (s, 1H), 7.10 (dd, J = 8.32, 1.39 Hz, 1H), 7.12 (d, J = 8.32 Hz, 1H), 7.26-7.34 (m, 6H).	539	537
27	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.91-0.84 (m, 1H), 0.93 (s, 9H), 1.34-1.47 (m, 3H), 1.60-1.68 (m, 1H), 1.81-1.90 (m, 2H), 2.26-2.35 (m, 3H), 2.57-2.61 (m, 2H), 2.73-2.80 (m, 1H), 3.21 (s, 3H), 3.31-3.38 (m, 3H), 3.61-3.68 (m, 1H), 6.34 (d, J = 1.62 Hz, 1H), 7.22 (dd, J = 7.86, 1.39 Hz, 1H), 7.25 (d, J = 7.86 Hz, 1H), 7.41 (d, J = 1.39 Hz, 1H), 7.56 (s, 1H), 12.20 (s, 1H).	463	461
28	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.91-0.84 (m, 1H), 0.95 (s, 9H), 1.29-1.42 (m, 4H), 1.54-1.61 (m, 1H), 1.83-1.94 (m, 4H), 2.10 (s, 6H),	476	474

	2.17-2.28 (m, 3H), 2.59-2.63 (m, 2H), 3.20-3.27 (m, 1H), 3.48-3.55 (m, 1H), 6.41 (d, J = 1.49 Hz, 1H), 7.26 (d, J = 8.07 Hz, 1H), 7.28 (d, J = 8.07 Hz, 1H), 7.35 (s, 1H), 7.42 (s, 1H).		
29	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.92-0.86 (m, 1H), 0.93 (s, 9H), 1.35-1.40 (m, 2H), 1.77-1.92 (m, 2H), 2.27-2.35 (m, 3H), 2.57-2.61 (m, 2H), 2.79-2.87 (m, 1H), 3.17-3.37 (m, 3H), 3.57-3.64 (m, 1H), 4.65 (br s, 1H), 6.39 (d, J = 1.85 Hz, 1H), 7.24-7.26 (m, 2H), 7.42 (s, 1H), 7.57 (s, 1H).	435	433
30	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.92-0.85 (m, 1H), 0.94 (s, 9H), 1.36-1.40 (m, 2H), 1.79 (s, 3H), 1.81-1.90 (m, 2H), 2.26-2.38 (m, 3H), 2.58-2.62 (m, 2H), 2.83-2.92 (m, 2H), 3.03-3.11 (m, 1H), 3.31-3.37 (m, 1H), 3.59-3.66 (m, 1H), 6.39 (s, 1H), 7.24-7.28 (m, 2H), 7.44 (s, 1H), 7.57 (s, 1H), 7.85-7.88 (m, 1H), 12.23 (br s, 1H).	476	474
31	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.93 (s, 9H), 0.97-1.06 (m, 1H), 1.36-1.40 (m, 2H), 1.88-2.00 (m, 2H), 2.30-2.36 (m, 3H), 2.51-2.62 (m, 4H), 3.01-3.08 (m, 1H), 3.32-3.38 (m, 1H), 3.60-3.67 (m, 1H), 6.58 (d, J = 1.85 Hz, 1H), 7.21 (dd, J = 7.86, 1.85 Hz, 1H), 7.26 (d, J = 7.86 Hz, 1H), 7.43 (d, J = 1.85 Hz, 1H), 7.69 (s, 1H), 12.21 (br s, 1H).	444	442
32	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.95 (s, 9H), 0.99 (s, 3H), 1.10 (s, 3H), 1.38-1.45 (m, 2H), 1.96-2.02 (m, 1H), 2.13-2.23 (m, 2H), 2.26-2.33 (m, 1H), 2.54-2.66 (m, 4H), 3.45-3.56 (m, 1H), 3.79-3.92 (m, 1H), 5.70-5.76 (m, 1H), 6.14 (br s, 1H), 7.04-7.09 (m, 1H), 7.11-7.16 (m, 1H), 7.18-7.21 (m, 1H).	433	431
33	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.95 (s, 9H), 0.99 (s, 3H), 1.10 (s, 3H), 1.38-1.45 (m, 2H), 1.96-2.02 (m, 1H), 2.13-2.23 (m, 2H), 2.26-2.33 (m, 1H), 2.54-2.66 (m, 4H), 3.45-3.56 (m, 1H), 3.79-3.92 (m,	433	431

	1H), 5.70-5.76 (m, 1H), 6.14 (br s, 1H), 7.04-7.09 (m, 1H), 7.11-7.16 (m, 1H), 7.18-7.21 (m, 1H).		
34	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.65 (s, 3H), 0.84 (d, J = 6.47 Hz, 3H), 0.93 (s, 9H), 1.02 (s, 3H), 1.36-1.40 (m, 2H), 2.05 (s, 2H), 2.19 (s, 6H), 2.40 (q, J = 6.47 Hz, 1H), 2.57-2.61 (m, 2H), 6.45 (d, J = 0.92 Hz, 1H), 7.17 (dd, J = 8.09, 1.85 Hz, 1H), 7.25 (d, J = 8.09 Hz, 1H), 7.34 (d, J = 1.85 Hz, 1H), 7.52 (s, 1H), 12.43 (s, 1H).	485	483
35	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.65 (s, 3H), 0.84 (d, J = 6.47 Hz, 3H), 0.93 (s, 9H), 1.02 (s, 3H), 1.36-1.40 (m, 2H), 2.05 (s, 2H), 2.19 (s, 6H), 2.40 (q, J = 6.47 Hz, 1H), 2.57-2.61 (m, 2H), 6.45 (d, J = 0.92 Hz, 1H), 7.17 (dd, J = 8.09, 1.85 Hz, 1H), 7.25 (d, J = 8.09 Hz, 1H), 7.34 (d, J = 1.85 Hz, 1H), 7.52 (s, 1H), 12.43 (s, 1H).	485	483
36	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 1.02 (s, 9H), 1.15-1.08 (m, 1H), 1.40-1.48 (m, 1H), 1.61-1.65 (m, 1H), 1.73-1.81 (m, 2H), 2.08-2.18 (m, 1H), 2.41-2.55 (m, 2H), 3.71 (s, 2H), 7.11 (dd, J = 8.60, 1.76 Hz, 1H), 7.14 (d, J = 8.60 Hz, 1H), 7.30 (d, J = 1.76 Hz, 1H), 7.58 (d, J = 1.98 Hz, 1H), 9.64 (d, J = 1.98 Hz, 1H).	350	348
37	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.58 (s, 3H), 0.97 (s, 9H), 1.14 (s, 3H), 1.34-1.49 (m, 4H), 1.51-1.73 (m, 3H), 1.82-1.93 (m, 1H), 2.54-2.76 (m, 4H), 3.60-3.72 (m, 1H), 3.79-3.92 (m, 1H), 5.08-5.17 (br m, 1H), 6.12 (br s, 1H), 7.14-7.22 (m, 2H), 7.30-7.36 (m, 1H).	447	445
38	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.58 (s, 3H), 0.97 (s, 9H), 1.14 (s, 3H), 1.34-1.49 (m, 4H), 1.51-1.73 (m, 3H), 1.82-1.93 (m, 1H), 2.54-2.76 (m, 4H), 3.60-3.72 (m, 1H), 3.79-3.92 (m, 1H), 5.08-5.17 (br m, 1H), 6.12 (br s, 1H), 7.14-7.22 (m, 2H), 7.30-7.36 (m, 1H).	447	445

39	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 0.98-1.06 (m, 1H), 1.23-1.30 (m, 1H), 1.38-1.42 (m, 2H), 1.59-1.81 (m, 4H), 2.16-2.22 (m, 1H), 2.42 (t, J = 7.05 Hz, 2H), 2.52-2.57 (m, 1H), 2.61-2.65 (m, 2H), 3.35-3.42 (m, 1H), 3.58-3.65 (m, 1H), 6.23 (d, J = 1.62 Hz, 1H), 7.07 (s, 1H), 7.24 (dd, J = 8.09, 1.85 Hz, 1H), 7.33 (d, J = 8.09 Hz, 1H), 7.35 (d, J = 1.85 Hz, 1H), 12.24 (s, 1H).	419	417
40	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 0.98-1.06 (m, 1H), 1.23-1.30 (m, 1H), 1.38-1.42 (m, 2H), 1.59-1.81 (m, 4H), 2.16-2.22 (m, 1H), 2.42 (t, J = 7.05 Hz, 2H), 2.52-2.57 (m, 1H), 2.61-2.65 (m, 2H), 3.35-3.42 (m, 1H), 3.58-3.65 (m, 1H), 6.23 (d, J = 1.62 Hz, 1H), 7.07 (s, 1H), 7.24 (dd, J = 8.09, 1.85 Hz, 1H), 7.33 (d, J = 8.09 Hz, 1H), 7.35 (d, J = 1.85 Hz, 1H), 12.24 (s, 1H).	419	417
41	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 0.98-1.06 (m, 1H), 1.23-1.30 (m, 1H), 1.38-1.42 (m, 2H), 1.59-1.81 (m, 4H), 2.16-2.22 (m, 1H), 2.42 (t, J = 7.05 Hz, 2H), 2.52-2.57 (m, 1H), 2.61-2.65 (m, 2H), 3.35-3.42 (m, 1H), 3.58-3.65 (m, 1H), 6.23 (d, J = 1.62 Hz, 1H), 7.07 (s, 1H), 7.24 (dd, J = 8.09, 1.85 Hz, 1H), 7.33 (d, J = 8.09 Hz, 1H), 7.35 (d, J = 1.85 Hz, 1H), 12.24 (s, 1H).	419	417
42	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.00-1.06 (m, 1H), 1.23-1.31 (m, 1H), 1.37-1.42 (m, 2H), 1.55-1.74 (m, 4H), 2.00 (s, 3H), 2.15-2.23 (m, 1H), 2.32-2.44 (m, 3H), 2.59-2.64 (m, 2H), 3.52-3.59 (m, 1H), 3.77-3.84 (m, 1H), 7.15-7.18 (m, 2H), 7.28-7.31 (m, 2H), 12.19 (s, 1H).	433	431
43	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.92 (s, 9H), 0.94-1.09 (m, 2H), 1.32-1.42 (m, 3H), 1.53-1.66 (m, 5H), 2.40-2.48 (m, 3H), 2.58-2.68 (m, 3H), 3.16-3.22 (m, 1H), 3.33-3.40 (m, 1H), 3.67-3.74 (m, 1H), 4.30	553	551

	(d, $J = 11.79$ Hz, 1H), 4.34 (d, $J = 11.79$ Hz, 1H), 6.14 (s, 1H), 7.00 (s, 1H), 7.25–7.28 (m, 5H), 7.31–7.35 (m, 3H), 12.31 (s, 1H).		
44	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ : 0.94–0.88 (m, 2H), 0.95 (s, 9H), 1.33–1.41 (m, 3H), 1.52–1.66 (m, 4H), 2.46–2.50 (m, 3H), 2.60–2.67 (m, 3H), 3.12–3.17 (m, 1H), 3.23–3.29 (m, 1H), 3.36–3.43 (m, 1H), 3.68–3.75 (m, 1H), 4.19 (s, 1H), 6.23 (s, 1H), 6.97 (s, 1H), 7.24–7.31 (m, 3H), 12.30 (s, 1H).	463	461
45	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ : 0.95 (s, 9H), 1.20 (t, $J = 7.09$ Hz, 3H), 1.26–1.34 (m, 1H), 1.38–1.42 (m, 2H), 1.50–1.81 (m, 6H), 2.53 (t, $J = 6.94$ Hz, 2H), 2.62–2.67 (m, 3H), 2.80–2.85 (m, 1H), 3.39–3.46 (m, 1H), 3.70–3.77 (m, 1H), 4.10 (q, $J = 7.09$ Hz, 2H), 6.34 (s, 1H), 7.05 (s, 1H), 7.23 (dd, $J = 8.09, 2.08$ Hz, 1H), 7.30 (d, $J = 8.09$ Hz, 1H), 7.31 (d, $J = 2.08$ Hz, 1H), 11.97 (s, 1H).	505	503
46	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ : 0.90–1.02 (m, 2H), 0.95 (s, 9H), 1.35–1.40 (m, 3H), 1.53–1.67 (m, 4H), 2.37–2.42 (m, 1H), 2.46–2.52 (m, 2H), 2.60–2.67 (m, 3H), 3.02–3.07 (m, 1H), 3.11 (s, 3H), 3.14–3.21 (m, 1H), 3.35–3.42 (m, 1H), 3.70–3.77 (m, 1H), 6.18 (s, 1H), 7.00 (s, 1H), 7.25–7.32 (m, 3H), 12.29 (s, 1H).	477	475
47	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ : 0.93–0.98 (m, 1H), 0.96 (s, 9H), 1.21 (t, $J = 7.09$ Hz, 3H), 1.37–1.41 (m, 2H), 1.45–1.74 (m, 4H), 2.09–2.14 (m, 1H), 2.18–2.26 (m, 1H), 2.57–2.69 (m, 6H), 3.45–3.52 (m, 1H), 3.70–3.76 (m, 1H), 4.11 (q, $J = 7.09$ Hz, 2H), 6.36 (s, 1H), 7.16 (s, 1H), 7.25–7.34 (m, 3H), 7.51 (s, 3H).	476	–
48	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ : 0.95 (s, 9H), 1.03–1.10 (m, 1H), 1.18–1.26 (m, 1H), 1.32–1.41 (m, 3H), 1.57–1.69 (m, 4H), 2.44–2.54 (m, 3H), 2.61–2.68 (m, 3H), 3.14 (s, 3H), 3.35–3.42 (m, 1H), 3.69–3.76	541	539

	(m, 1H), 3.90-3.95 (m, 1H), 4.02-4.08 (m, 1H), 6.28 (s, 1H), 7.05 (s, 1H), 7.26-7.32 (m, 3H), 12.30 (s, 1H).		
49	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.96 (s, 9H), 1.03-1.12 (m, 2H), 1.22 (t, J = 7.17 Hz, 3H), 1.33-1.40 (m, 3H), 1.54-1.68 (m, 5H), 2.36-2.42 (m, 1H), 2.37 (s, 3H), 2.56-2.65 (m, 5H), 2.75-2.82 (m, 1H), 3.38-3.45 (m, 1H), 3.74-3.81 (m, 1H), 4.12 (q, J = 7.17 Hz, 2H), 6.33 (s, 1H), 7.10 (s, 1H), 7.24 (dd, J = 8.09, 1.85 Hz, 1H), 7.30 (d, J = 1.85 Hz, 1H), 7.31 (d, J = 8.09 Hz, 1H), 8.09 (s, 2H).	504	-
50	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.00-1.20 (m, 3H), 1.22 (t, J = 7.17 Hz, 3H), 1.30-1.70 (m, 8H), 2.33-2.68 (m, 80H), 3.40-3.47 (m, 1H), 3.72-3.80 (m, 1H), 4.12 (q, J = 7.17 Hz, 2H), 6.29 (s, 1H), 7.12 (s, 1H), 7.25-7.35 (m, 3H), 9.24 (br s, 1H).	518	-
51	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.31-1.70 (m, 6H), 2.41-2.52 (m, 5H), 2.61-2.67 (m, 3H), 2.84-2.91 (m, 1H), 3.35-3.44 (m, 1H), 3.65-3.71 (m, 1H), 6.36 (s, 1H), 6.64 (s, 1H), 7.00 (s, 1H), 7.21-7.38 (m, 3H), 12.29 (br s, 1H).	476	474
52	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.21-1.32 (m, 1H), 1.37-1.44 (m, 2H), 1.52-1.70 (m, 4H), 1.83 (dd, J = 16.76, 9.13 Hz, 1H), 2.13 (dd, J = 16.76, 6.82 Hz, 1H), 2.45-2.53 (m, 2H), 2.62-2.67 (m, 3H), 2.70-2.78 (m, 1H), 3.41-3.48 (m, 1H), 3.67-3.74 (m, 1H), 6.48 (s, 1H), 7.15 (s, 1H), 7.27-7.38 (m, 3H), 12.31 (br s, 1H).	458	456
53	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 0.96-1.05 (m, 1H), 1.21-1.33 (m, 1H), 1.37-1.42 (m, 2H), 1.56-1.73 (m, 3H), 1.76-1.85 (m, 1H), 2.25-2.29 (m, 1H), 2.25 (s, 6H), 2.48-2.57 (m, 1H), 2.61-2.65 (m, 2H), 6.20 (d, J = 1.47 Hz, 1H), 7.16 (s, 1H), 7.22 (dd, J = 8.07, 1.96 Hz, 1H), 7.34-7.36 (m,	457	455

	2H), 12.48 (s, 1H).		
54	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 0.96-1.05 (m, 1H), 1.21-1.33 (m, 1H), 1.37-1.42 (m, 2H), 1.56-1.73 (m, 3H), 1.76-1.85 (m, 1H), 2.25-2.29 (m, 1H), 2.25 (s, 6H), 2.48-2.57 (m, 1H), 2.61-2.65 (m, 2H), 6.20 (d, J = 1.47 Hz, 1H), 7.16 (s, 1H), 7.22 (dd, J = 8.07, 1.96 Hz, 1H), 7.34-7.36 (m, 2H), 12.48 (s, 1H).	457	455
55	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.17-1.26 (m, 1H), 1.30-1.42 (m, 3H), 1.44-1.52 (m, 1H), 1.58-1.68 (m, 1H), 1.76-1.83 (m, 1H), 2.41-2.50 (m, 2H), 2.57-2.69 (m, 4H), 2.80-2.87 (m, 1H), 3.35-3.44 (m, 1H), 3.62-3.73 (m, 1H), 4.35-4.40 (m, 1H), 6.31 (s, 1H), 7.02 (s, 1H), 7.22-7.34 (m, 3H), 12.30 (br s, 1H).	449	447
56	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.11-1.23 (m, 1H), 1.36-1.69 (m, 6H), 2.41-2.47 (m, 2H), 2.50-2.73 (m, 8H), 3.01 (s, 3H), 3.39-3.47 (m, 1H), 3.61-3.67 (m, 1H), 6.31 (s, 1H), 7.05 (s, 1H), 7.24-7.40 (m, 3H), 12.29 (br s, 1H).	463	461
57	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.60 (d, J = 7.63 Hz, 3H), 0.95 (s, 9H), 1.30-1.43 (m, 4H), 1.48-1.67 (m, 3H), 2.43-2.68 (m, 6H), 3.33-3.43 (m, 1H), 3.64-3.73 (m, 1H), 6.31 (s, 1H), 7.00 (s, 1H), 7.24-7.36 (m, 3H), 12.31 (br s, 1H).	433	431
58	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.96 (s, 9H), 1.35-1.42 (m, 2H), 1.97-2.08 (m, 2H), 2.43-2.50 (m, 2H), 2.58-2.67 (m, 2H), 3.34-3.46 (m, 3H), 3.65-3.72 (m, 1H), 3.88-3.95 (m, 1H), 4.32 (d, J = 11.10 Hz, 1H), 6.24 (s, 1H), 7.09 (s, 1H), 7.30-7.46 (m, 3H), 12.24 (br s, 1H).	421	419
59	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.94 (d, J = 5.50 Hz, 6H), 0.98-1.07 (m, 1H), 1.23-1.29 (m, 1H), 1.57-1.85 (m, 7H), 2.13-2.20 (m, 1H), 2.40 (t, J = 7.05 Hz, 2H), 2.49-2.56 (m, 1H), 3.34-3.42 (m, 1H),	421	419

	3.57-3.64 (m, 1H), 4.07 (t, J = 6.47 Hz, 2H), 6.22 (s, 1H), 7.04 (s, 1H), 7.11-7.38 (m, 3H), 12.28 (s, 1H).		
60	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.99-1.06 (m, 1H), 1.03 (s, 9H), 1.20-1.29 (m, 1H), 1.56-1.83 (m, 4H), 2.14-2.20 (m, 1H), 2.40 (t, J = 6.94 Hz, 2H), 2.47-2.56 (m, 1H), 3.34-3.41 (m, 1H), 3.57-3.65 (m, 1H), 3.70 (s, 2H), 6.22 (s, 1H), 7.04 (s, 1H), 7.09 (d, J = 8.55 Hz, 1H), 7.23 (dd, J = 8.55, 2.31 Hz, 1H), 7.37 (d, J = 2.31 Hz, 1H), 12.26 (br s, 1H).	421	419
61	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95-1.08 (m, 1H), 0.97 (s, 9H), 1.20-1.35 (m, 2H), 1.56-1.82 (m, 6H), 2.13-2.21 (m, 1H), 2.42 (t, J = 6.90 Hz, 2H), 2.48-2.56 (m, 1H), 3.33-3.42 (m, 1H), 3.58-3.66 (m, 1H), 4.10 (t, J = 6.90 Hz, 2H), 6.22 (d, J = 1.39 Hz, 1H), 7.05 (s, 1H), 7.15 (d, J = 8.73 Hz, 1H), 7.24 (dd, J = 8.73, 2.14 Hz, 1H), 7.36 (d, J = 2.14 Hz, 1H), 12.20 (br s, 1H).	435	433
62	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.96-1.07 (m, 1H), 1.20-1.31 (m, 1H), 1.55-1.87 (m, 10H), 2.01-2.09 (m, 2H), 2.13-2.20 (m, 1H), 2.39 (t, J = 7.05 Hz, 2H), 2.42-2.47 (m, 1H), 2.49-2.55 (m, 1H), 3.34-3.41 (m, 1H), 3.56-3.65 (m, 1H), 3.98 (t, J = 6.36 Hz, 2H), 6.22 (d, J = 1.62 Hz, 1H), 7.03 (s, 1H), 7.10 (d, J = 8.55 Hz, 1H), 7.23 (dd, J = 8.55, 2.31 Hz, 1H), 7.36 (d, J = 2.31 Hz, 1H), 12.29 (br s, 1H).	433	431
63	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.38-1.59 (m, 3H), 1.88-1.97 (m, 1H), 2.00-2.34 (m, 2H), 2.41-2.46 (m, 2H), 2.59-2.74 (m, 4H), 3.37-3.45 (m, 1H), 3.62-3.70 (m, 1H), 6.46 (d, J = 1.85 Hz, 1H), 7.25-7.41 (m, 4H), 12.28 (br s, 1H).	455	453
64	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.38 (s, 3H), 0.91 (s, 3H), 0.96 (s, 9H), 1.23-1.40 (m, 2H), 1.40-1.47 (m, 2H), 1.80-1.89 (m, 1H), 2.15-	447	445

	2.23 (m, 2H), 2.32-2.40 (m, 1H), 2.55-2.69 (m, 4H), 3.53-3.65 (m, 1H), 3.65-3.76 (m, 1H), 5.57 (br s, 1H), 6.12 (br s, 1H), 7.10-7.19 (m, 2H), 7.24-7.26 (m, 1H).		
65	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.97-1.13 (m, 5H), 1.21-1.29 (m, 1H), 1.56-1.82 (m, 4H), 2.14-2.20 (m, 1H), 2.39-2.43 (m, 2H), 2.49-2.59 (m, 1H), 3.34-3.42 (m, 1H), 3.58-3.65 (m, 1H), 4.19 (s, 2H), 6.22 (d, J = 1.62 Hz, 1H), 7.06 (br s, 1H), 7.12 (d, J = 8.79 Hz, 1H), 7.24 (dd, J = 8.67, 2.20 Hz, 1H), 7.38 (d, J = 2.31 Hz, 1H), 12.25 (br s, 1H).	473	471
66	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.90-0.99 (m, 12H), 1.28-1.43 (m, 4H), 1.81-2.05 (m, 4H), 2.22-2.34 (m, 1H), 2.39-2.45 (m, 2H), 2.58-2.67 (m, 2H), 3.34-3.43 (m, 1H), 3.59-3.67 (m, 1H), 6.19 (s, 1H), 7.07 (br s, 1H), 7.22-7.37 (m, 3H), 12.21 (br s, 1H).	433	431
67	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.95 (s, 9H), 1.07-1.27 (m, 1H), 1.38-1.47 (m, 2H), 1.75-2.06 (m, 2H), 2.45-2.61 (m, 2H), 2.60-2.69 (m, 4H), 3.18-3.27 (m, 1H), 3.30 (s, 3H), 3.57-3.68 (m, 1H), 3.71-3.82 (m, 1H), 5.40 (br s, 1H), 6.07 (br s, 1H), 7.08-7.21 (m, 2H), 7.27-7.31 (m, 1H).	449	447
68	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.95 (s, 9H), 1.07-1.27 (m, 1H), 1.38-1.47 (m, 2H), 1.75-2.06 (m, 2H), 2.45-2.61 (m, 2H), 2.60-2.69 (m, 4H), 3.18-3.27 (m, 1H), 3.30 (s, 3H), 3.57-3.68 (m, 1H), 3.71-3.82 (m, 1H), 5.40 (br s, 1H), 6.07 (br s, 1H), 7.08-7.21 (m, 2H), 7.27-7.31 (m, 1H).	449	447
69	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.65-0.78 (m, 4H), 0.77 (d, J = 7.51 Hz, 3H), 0.96 (s, 9H), 1.36-1.60 (m, 5H), 1.70-1.80 (m, 1H), 2.14-2.23 (m, 1H), 2.38-2.44 (m, 2H), 2.52-2.59 (m, 1H), 2.59-2.67 (m, 2H), 3.35-3.43 (m, 1H), 3.57-3.64 (m, 1H), 6.24 (s, 1H), 7.11 (s, 1H), 7.20-7.36 (m, 3H), 12.27 (br	433	431

	s, 1H).		
70	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.18 (d, J = 6.80 Hz, 3H), 1.36-1.42 (m, 2H), 1.83 (d, J = 13.64 Hz, 1H), 2.15-2.22 (m, 1H), 2.46 (t, J = 6.94 Hz, 2H), 2.60-2.66 (m, 2H), 3.39-3.47 (m, 1H), 3.65-3.78 (m, 2H), 3.96 (d, J = 11.79 Hz, 1H), 4.03-4.12 (m, 1H), 6.20 (d, J = 1.62 Hz, 1H), 7.10 (s, 1H), 7.31-7.33 (m, 2H), 7.41-7.43 (m, 1H), 12.27 (br s, 1H).	435	433
71	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.18 (d, J = 6.80 Hz, 3H), 1.36-1.42 (m, 2H), 1.83 (d, J = 13.64 Hz, 1H), 2.15-2.22 (m, 1H), 2.46 (t, J = 6.94 Hz, 2H), 2.60-2.66 (m, 2H), 3.39-3.47 (m, 1H), 3.65-3.78 (m, 2H), 3.96 (d, J = 11.79 Hz, 1H), 4.03-4.12 (m, 1H), 6.20 (d, J = 1.62 Hz, 1H), 7.10 (s, 1H), 7.31-7.33 (m, 2H), 7.41-7.43 (m, 1H), 12.27 (br s, 1H).	435	433
72	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.95 (s, 9H), 1.18 (d, J = 6.47 Hz, 3H), 1.36-1.41 (m, 2H), 1.92 (d, J = 13.18 Hz, 1H), 2.18-2.27 (m, 1H), 2.28 (s, 6H), 2.60-2.66 (m, 2H), 3.73 (d, J = 11.79 Hz, 1H), 3.96 (d, J = 11.79 Hz, 1H), 4.05-4.14 (m, 1H), 6.19 (d, J = 1.85 Hz, 1H), 7.17 (s, 1H), 7.26-7.42 (m, 3H), 12.46 (br s, 1H).	473	471
73	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.58 (d, J = 7.40 Hz, 3H), 0.95 (s, 9H), 1.34-1.41 (m, 5H), 1.49-1.66 (m, 3H), 2.23 (s, 6H), 2.59-2.68 (m, 3H), 6.25 (s, 1H), 7.02 (s, 1H), 7.21-7.34 (m, 3H).	471	469
74	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.79 (s, 3H), 0.90 (s, 3H), 0.95 (s, 9H), 1.25-1.32 (m, 1H), 1.36-1.43 (m, 2H), 1.63 (d, J = 13.76 Hz, 1H), 1.78 (dd, J = 13.76, 1.50 Hz, 1H), 1.83-1.93 (m, 1H), 2.37-2.51 (m, 2H), 2.40-2.44 (m, 2H), 2.59-2.66 (m, 2H), 3.36-3.44 (m, 1H), 3.59-3.67 (m, 1H), 6.19 (d, J = 1.39 Hz, 1H), 7.10 (s, 1H), 7.21-	447	445

	7.36 (m, 3H), 12.21 (br s, 1H).		
75	¹ H-NMR (400 MHz, CDCl ₃) δ : 0.97 (s, 9H), 1.39-1.48 (m, 2H), 1.53-1.65 (m, 1H), 1.90-2.00 (m, 1H), 2.15-2.35 (m, 5H), 2.41 (br s, 6H), 2.61-2.69 (m, 2H), 5.60-5.76 (br m, 1H), 6.03 (br s, 1H), 7.11-7.16 (m, 1H), 7.19-7.26 (m, 2H).	525	523
76	¹ H-NMR (400 MHz, DMSO-d ₆) δ : 0.94 (s, 9H), 1.34-1.41 (m, 2H), 1.53-1.64 (m, 1H), 1.65-1.77 (m, 1H), 1.84-1.94 (m, 1H), 2.24 (br s, 6H), 2.51-2.73 (m, 5H), 3.11-3.22 (m, 4H), 6.27-6.31 (m, 1H), 7.17-7.27 (m, 2H), 7.30-7.37 (m, 2H), 12.45 (br s, 1H).	487	485
77	¹ H-NMR (400 MHz, DMSO-d ₆) δ : 0.94 (s, 9H), 1.34-1.41 (m, 2H), 1.53-1.64 (m, 1H), 1.65-1.77 (m, 1H), 1.84-1.94 (m, 1H), 2.24 (br s, 6H), 2.51-2.73 (m, 5H), 3.11-3.22 (m, 4H), 6.27-6.31 (m, 1H), 7.17-7.27 (m, 2H), 7.30-7.37 (m, 2H), 12.45 (br s, 1H).	487	485
78	¹ H-NMR (400 MHz, DMSO-d ₆) δ : 0.58 (d, J = 7.40 Hz, 3H), 0.95 (s, 9H), 1.34-1.41 (m, 5H), 1.49-1.66 (m, 3H), 2.23 (s, 6H), 2.59-2.68 (m, 3H), 6.25 (s, 1H), 7.02 (s, 1H), 7.21-7.34 (m, 3H).	471	469
79	¹ H-NMR (400 MHz, DMSO-d ₆) δ : 0.58 (d, J = 7.40 Hz, 3H), 0.95 (s, 9H), 1.34-1.41 (m, 5H), 1.49-1.66 (m, 3H), 2.23 (s, 6H), 2.59-2.68 (m, 3H), 6.25 (s, 1H), 7.02 (s, 1H), 7.21-7.34 (m, 3H).	471	469
80	¹ H-NMR (400 MHz, DMSO-d ₆) δ : 0.65-0.78 (m, 4H), 0.77 (d, J = 7.51 Hz, 3H), 0.96 (s, 9H), 1.36-1.60 (m, 5H), 1.70-1.80 (m, 1H), 2.14-2.23 (m, 1H), 2.38-2.44 (m, 2H), 2.52-2.59 (m, 1H), 2.59-2.67 (m, 2H), 3.35-3.43 (m, 1H), 3.57-3.64 (m, 1H), 6.24 (s, 1H), 7.11 (s, 1H), 7.20-7.36 (m, 3H), 12.27 (br s, 1H).	433	431
81	¹ H-NMR (400 MHz, DMSO-d ₆) δ : 0.65-0.78 (m, 4H), 0.77 (d, J = 7.51 Hz, 3H), 0.96 (s, 9H), 1.36-1.60 (m, 5H), 1.70-1.80 (m, 1H), 2.14-2.23 (m, 1H), 2.38-2.44 (m, 2H),	433	431

	2.52-2.59 (m, 1H), 2.59-2.67 (m, 2H), 3.35-3.43 (m, 1H), 3.57-3.64 (m, 1H), 6.24 (s, 1H), 7.11 (s, 1H), 7.20-7.36 (m, 3H), 12.27 (br s, 1H).		
82	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.79 (s, 3H), 0.90 (s, 3H), 0.95 (s, 9H), 1.25-1.32 (m, 1H), 1.36-1.43 (m, 2H), 1.63 (d, J = 13.76 Hz, 1H), 1.78 (dd, J = 13.76, 1.50 Hz, 1H), 1.83-1.93 (m, 1H), 2.37-2.51 (m, 2H), 2.40-2.44 (m, 2H), 2.59-2.66 (m, 2H), 3.36-3.44 (m, 1H), 3.59-3.67 (m, 1H), 6.19 (d, J = 1.39 Hz, 1H), 7.10 (s, 1H), 7.21-7.36 (m, 3H), 12.21 (br s, 1H).	447	445
83	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.79 (s, 3H), 0.90 (s, 3H), 0.95 (s, 9H), 1.25-1.32 (m, 1H), 1.36-1.43 (m, 2H), 1.63 (d, J = 13.76 Hz, 1H), 1.78 (dd, J = 13.76, 1.50 Hz, 1H), 1.83-1.93 (m, 1H), 2.37-2.51 (m, 2H), 2.40-2.44 (m, 2H), 2.59-2.66 (m, 2H), 3.36-3.44 (m, 1H), 3.59-3.67 (m, 1H), 6.19 (d, J = 1.39 Hz, 1H), 7.10 (s, 1H), 7.21-7.36 (m, 3H), 12.21 (br s, 1H).	447	445
84	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.66-0.80 (m, 1H), 0.94 (s, 9H), 1.33-1.43 (m, 2H), 1.44-1.80 (m, 4H), 2.24 (s, 6H), 2.29-2.37 (m, 1H), 2.53-2.69 (m, 3H), 2.97-3.10 (m, 2H), 3.15 (s, 3H), 6.23 (s, 1H), 7.14-7.23 (m, 2H), 7.29-7.37 (m, 2H), 12.41 (br s, 1H).	501	499
85	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.66-0.80 (m, 1H), 0.94 (s, 9H), 1.33-1.43 (m, 2H), 1.44-1.80 (m, 4H), 2.24 (s, 6H), 2.29-2.37 (m, 1H), 2.53-2.69 (m, 3H), 2.97-3.10 (m, 2H), 3.15 (s, 3H), 6.23 (s, 1H), 7.14-7.23 (m, 2H), 7.29-7.37 (m, 2H), 12.41 (br s, 1H).	501	499
86	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.93 (s, 9H), 1.34-1.43 (m, 2H), 1.43-1.56 (m, 1H), 1.84-1.95 (m, 1H), 1.98-2.09 (m, 1H), 2.15-2.30 (m, 7H), 2.59-2.68 (m, 3H), 2.74-2.86 (m, 1H), 6.42-6.45 (m, 1H), 7.22-7.27 (m, 1H), 7.34-7.43 (m, 3H).	493	491
87	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.93	493	491

	(s, 9H), 1.34-1.43 (m, 2H), 1.43-1.56 (m, 1H), 1.84-1.95 (m, 1H), 1.98-2.09 (m, 1H), 2.15-2.30 (m, 7H), 2.59-2.68 (m, 3H), 2.74-2.86 (m, 1H), 6.42-6.45 (m, 1H), 7.22-7.27 (m, 1H), 7.34-7.43 (m, 3H).		
88	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.31 (t, J = 4.74 Hz, 1H), 0.56 (dd, J = 8.67, 4.28 Hz, 1H), 0.99-1.09 (m, 2H), 1.07 (s, 3H), 1.09 (s, 3H), 1.23-1.31 (m, 1H), 1.57-1.83 (m, 4H), 2.14-2.19 (m, 1H), 2.41 (t, J = 7.05 Hz, 2H), 2.49-2.55 (m, 1H), 3.35-3.42 (m, 1H), 3.57-3.65 (m, 1H), 3.88-3.95 (m, 1H), 4.17-4.24 (m, 1H), 6.22 (d, J = 1.62 Hz, 1H), 7.04 (s, 1H), 7.13 (d, J = 8.55 Hz, 1H), 7.21-7.25 (m, 1H), 7.35-7.37 (m, 1H), 12.25 (br s, 1H).	433	431
89	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.61-0.74 (m, 1H), 0.74-0.79 (m, 2H), 0.94 (s, 9H), 1.35-1.41 (m, 2H), 1.42-1.51 (m, 2H), 1.50-1.59 (m, 1H), 1.66-1.78 (m, 1H), 2.19-2.28 (m, 7H), 2.51-2.58 (m, 1H), 2.58-2.64 (m, 2H), 6.19 (br s, 1H), 7.16 (br s, 1H), 7.18-7.22 (m, 1H), 7.30-7.35 (m, 2H), 12.46 (br s, 1H).	471	469
90	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.61-0.74 (m, 1H), 0.74-0.79 (m, 2H), 0.94 (s, 9H), 1.35-1.41 (m, 2H), 1.42-1.51 (m, 2H), 1.50-1.59 (m, 1H), 1.66-1.78 (m, 1H), 2.19-2.28 (m, 7H), 2.51-2.58 (m, 1H), 2.58-2.64 (m, 2H), 6.19 (br s, 1H), 7.16 (br s, 1H), 7.18-7.22 (m, 1H), 7.30-7.35 (m, 2H), 12.46 (br s, 1H).	471	469
91	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.89-0.97 (m, 12H), 1.24-1.34 (m, 2H), 1.34-1.42 (m, 2H), 1.81-1.97 (m, 3H), 1.98-2.09 (m, 1H), 2.19-2.33 (m, 7H), 2.56-2.64 (m, 2H), 6.12-6.16 (m, 1H), 7.11 (s, 1H), 7.18-7.24 (m, 1H), 7.30-7.35 (m, 2H), 12.44 (br s, 1H).	471	469
92	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.89-0.97 (m, 12H), 1.24-1.34 (m, 2H), 1.34-1.42 (m, 2H), 1.81-1.97 (m,	471	469

	3H), 1.98-2.09 (m, 1H), 2.19-2.33 (m, 7H), 2.56-2.64 (m, 2H), 6.12-6.16 (m, 1H), 7.11 (s, 1H), 7.18-7.24 (m, 1H), 7.30-7.35 (m, 2H), 12.44 (br s, 1H).		
93	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.58-0.71 (m, 1H), 0.75-0.81 (m, 3H), 0.94 (s, 9H), 1.02-1.12 (m, 2H), 1.18-1.32 (m, 1H), 1.34-1.42 (m, 2H), 1.56-1.65 (m, 1H), 1.65-1.77 (m, 1H), 2.23 (s, 6H), 2.27-2.36 (m, 1H), 2.52-2.65 (m, 3H), 6.21 (br s, 1H), 7.16 (br s, 1H), 7.18-7.22 (m, 1H), 7.28-7.35 (m, 2H), 12.45 (br s, 1H).	485	483
94	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.58-0.71 (m, 1H), 0.75-0.81 (m, 3H), 0.94 (s, 9H), 1.02-1.12 (m, 2H), 1.18-1.32 (m, 1H), 1.34-1.42 (m, 2H), 1.56-1.65 (m, 1H), 1.65-1.77 (m, 1H), 2.23 (s, 6H), 2.27-2.36 (m, 1H), 2.52-2.65 (m, 3H), 6.21 (br s, 1H), 7.16 (br s, 1H), 7.18-7.22 (m, 1H), 7.28-7.35 (m, 2H), 12.45 (br s, 1H).	485	483
95	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.80-0.87 (m, 1H), 1.24-1.30 (m, 1H), 1.33-1.42 (m, 2H), 1.53-1.97 (m, 12H), 2.11-2.16 (m, 1H), 2.47-2.51 (m, 1H), 2.59-2.72 (m, 2H), 3.59-3.66 (m, 1H), 3.70-3.78 (m, 1H), 4.26-4.32 (m, 1H), 5.10 (s, 1H), 6.00 (d, J = 1.47 Hz, 1H), 6.93 (d, J = 8.80 Hz, 1H), 7.17 (dd, J = 8.80, 2.45 Hz, 1H), 7.33 (d, J = 2.45 Hz, 1H).	433	431
96	¹ H-NMR (400 MHz, CDCl ₃) δ: 1.43-1.21 (m, 2H), 1.38 (d, J = 6.11 Hz, 6H), 1.63-1.74 (m, 2H), 1.87-1.97 (m, 2H), 2.11-2.15 (m, 1H), 2.47-2.53 (m, 1H), 2.56-2.67 (m, 2H), 3.61 (dt, J = 14.43, 6.36 Hz, 1H), 3.73 (dt, J = 13.94, 6.36 Hz, 1H), 4.50-4.59 (m, 1H), 5.81 (s, 1H), 6.01 (d, J = 1.47 Hz, 1H), 6.92 (d, J = 8.56 Hz, 1H), 7.19 (dd, J = 8.56, 2.45 Hz, 1H), 7.34 (d, J = 2.45 Hz, 1H).	393	391
97	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.96 (s, 9H), 1.13-1.29 (m, 1H), 1.37-1.45 (m, 2H), 1.71-1.81 (m, 1H),	482	480

	1.94-2.05 (m, 2H), 2.22 (s, 6H), 2.55-2.74 (m, 4H), 2.78-2.89 (m, 1H), 6.35-6.38 (m, 1H), 7.19-7.23 (m, 1H), 7.30 (s, 1H), 7.33-7.40 (m, 2H).		
98	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.81 (dd, J = 6.60, 2.45 Hz, 6H), 0.97-0.93 (m, 1H), 0.98 (s, 9H), 1.19-1.34 (m, 2H), 1.44-1.48 (m, 2H), 1.58-1.65 (m, 1H), 1.71-1.76 (m, 1H), 1.90-1.98 (m, 1H), 2.18-2.23 (m, 1H), 2.47 (s, 6H), 2.54-2.59 (m, 1H), 2.64-2.69 (m, 2H), 5.15 (s, 1H), 5.97 (d, J = 1.47 Hz, 1H), 7.14 (dd, J = 7.83, 2.20 Hz, 1H), 7.20 (d, J = 7.83 Hz, 1H), 7.27 (d, J = 2.20 Hz, 1H).	499	497
99	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.81 (dd, J = 6.60, 2.45 Hz, 6H), 0.97-0.93 (m, 1H), 0.98 (s, 9H), 1.19-1.34 (m, 2H), 1.44-1.48 (m, 2H), 1.58-1.65 (m, 1H), 1.71-1.76 (m, 1H), 1.90-1.98 (m, 1H), 2.18-2.23 (m, 1H), 2.47 (s, 6H), 2.54-2.59 (m, 1H), 2.64-2.69 (m, 2H), 5.15 (s, 1H), 5.97 (d, J = 1.47 Hz, 1H), 7.14 (dd, J = 7.83, 2.20 Hz, 1H), 7.20 (d, J = 7.83 Hz, 1H), 7.27 (d, J = 2.20 Hz, 1H).	499	497
100	¹ H-NMR (400 MHz, CDCl ₃) δ: 1.10 (s, 3H), 1.18 (s, 3H), 1.24-1.39 (m, 2H), 1.78-1.68 (m, 4H), 1.87-1.98 (m, 4H), 2.10-2.16 (m, 1H), 2.48-2.54 (m, 1H), 2.58-2.76 (m, 3H), 3.63 (dt, J = 14.35, 7.34 Hz, 1H), 3.75 (dt, J = 14.35, 6.24 Hz, 1H), 3.95 (d, J = 6.11 Hz, 2H), 5.48 (s, 1H), 6.01 (d, J = 1.71 Hz, 1H), 6.88 (d, J = 8.56 Hz, 1H), 7.19 (dd, J = 8.56, 2.20 Hz, 1H), 7.35 (d, J = 2.20 Hz, 1H).	447	445
101	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.99 (t, J = 7.34 Hz, 3H), 1.24-1.30 (m, 1H), 1.32-1.37 (m, 1H), 1.48-1.58 (m, 2H), 1.67-1.73 (m, 2H), 1.78-1.85 (m, 2H), 1.89-1.99 (m, 2H), 2.12-2.16 (m, 1H), 2.47-2.53 (m, 1H), 2.64-2.73 (m, 2H), 3.61-3.68 (m, 1H), 3.73-3.80 (m, 1H), 4.03 (t, J = 6.48 Hz, 2H), 4.95 (s, 1H), 6.01 (d, J = 1.71 Hz,	407	405

	1H), 6.90 (d, J = 8.56 Hz, 1H), 7.19 (dd, J = 8.56, 2.45 Hz, 1H), 7.34 (d, J = 2.45 Hz, 1H).		
102	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.78 (s, 3H), 0.88 (s, 3H), 0.90-1.06 (m, 10H), 1.20-1.32 (m, 1H), 1.35-1.43 (m, 2H), 1.61-1.70 (m, 1H), 1.79-1.92 (m, 2H), 2.24 (s, 6H), 2.33-2.42 (m, 1H), 2.58-2.65 (m, 2H), 6.15 (br s, 1H), 7.15 (s, 1H), 7.17-7.22 (m, 1H), 7.30-7.35 (m, 2H), 12.41 (br s, 1H).	485	483
103	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.78 (s, 3H), 0.88 (s, 3H), 0.90-1.06 (m, 10H), 1.20-1.32 (m, 1H), 1.35-1.43 (m, 2H), 1.61-1.70 (m, 1H), 1.79-1.92 (m, 2H), 2.24 (s, 6H), 2.33-2.42 (m, 1H), 2.58-2.65 (m, 2H), 6.15 (br s, 1H), 7.15 (s, 1H), 7.17-7.22 (m, 1H), 7.30-7.35 (m, 2H), 12.41 (br s, 1H).	485	483
104	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.67-0.76 (m, 1H), 0.78 (dd, J = 6.60, 1.71 Hz, 6H), 0.95 (s, 9H), 1.13-1.30 (m, 2H), 1.37-1.42 (m, 2H), 1.48-1.55 (m, 1H), 1.59-1.99 (m, 8H), 2.35-2.39 (m, 1H), 2.58-2.64 (m, 3H), 2.68-2.76 (m, 1H), 4.57-4.66 (m, 1H), 6.35 (d, J = 1.22 Hz, 1H), 7.11 (s, 1H), 7.21 (dd, J = 7.95, 1.83 Hz, 1H), 7.32-7.34 (m, 2H), 12.18 (br s, 1H).	501	499
105	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.67-0.76 (m, 1H), 0.78 (dd, J = 6.60, 1.71 Hz, 6H), 0.95 (s, 9H), 1.13-1.30 (m, 2H), 1.37-1.42 (m, 2H), 1.48-1.55 (m, 1H), 1.59-1.99 (m, 8H), 2.35-2.39 (m, 1H), 2.58-2.64 (m, 3H), 2.68-2.76 (m, 1H), 4.57-4.66 (m, 1H), 6.35 (d, J = 1.22 Hz, 1H), 7.11 (s, 1H), 7.21 (dd, J = 7.95, 1.83 Hz, 1H), 7.32-7.34 (m, 2H), 12.18 (br s, 1H).	501	499
106	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.79 (d, J = 6.80 Hz, 3H), 0.95-0.98 (m, 3H), 0.95 (s, 9H), 1.36-1.45 (m, 4H), 1.70-1.74 (m, 1H), 1.85-1.93 (m, 1H), 2.09-2.15 (m, 1H), 2.28 (s, 6H), 2.30-2.37 (m, 1H), 2.59-2.65 (m, 2H), 5.89 (d, J = 1.62 Hz, 1H), 7.12 (s, 1H), 7.20	485	483

	(dd, $J = 8.09, 1.85$ Hz, 1H), 7.31-7.35 (m, 2H), 12.44 (br s, 1H).		
107	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ : 0.57-0.62 (m, 3H), 0.91-0.97 (m, 12H), 1.21-1.30 (m, 1H), 1.34-1.41 (m, 2H), 1.46-1.66 (m, 2H), 1.79-1.91 (m, 1H), 2.12-2.20 (m, 1H), 2.27 (s, 6H), 2.29-2.37 (m, 1H), 2.58-2.64 (m, 2H), 6.21 (s, 1H), 7.01 (s, 1H), 7.18-7.25 (m, 1H), 7.28-7.32 (m, 2H), 12.41 (br s, 1H).	485	483
108	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ : 0.44 (d, $J = 7.29$ Hz, 1.89H), 0.88 (d, $J = 6.94$ Hz, 1.11H), 0.95 (s, 9H), 1.02 (s, 1.89H), 1.04 (s, 1.11H), 1.11 (s, 1.11H), 1.25 (s, 1.89H), 1.34-1.41 (m, 2H), 1.94-1.99 (m, 0.37H), 2.13 (q, $J = 7.29$ Hz, 0.63H), 2.30 (s, 2.22H), 2.31 (s, 3.88H), 2.59-2.66 (m, 2H), 3.66 (d, $J = 12.02$ Hz, 0.63H), 3.69 (d, $J = 11.90$ Hz, 0.37H), 4.06 (d, $J = 11.90$ Hz, 0.37H), 4.23 (d, $J = 12.25$ Hz, 0.63H), 5.88 (d, $J = 1.85$ Hz, 0.37H), 6.21 (s, 0.63H), 7.06 (s, 0.63H), 7.16 (s, 0.37H), 7.26-7.42 (m, 3H), 12.50 (br s, 1H).	501	499
109	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ : 0.88 (d, $J = 6.94$ Hz, 3H), 0.95 (s, 9H), 1.04 (s, 3H), 1.11 (s, 3H), 1.37-1.41 (m, 2H), 1.94-1.99 (m, 1H), 2.30 (s, 6H), 2.60-2.65 (m, 2H), 3.69 (d, $J = 11.90$ Hz, 1H), 4.06 (d, $J = 11.90$ Hz, 1H), 5.88 (d, $J = 1.85$ Hz, 1H), 7.16 (s, 1H), 7.28-7.42 (m, 3H), 12.50 (br s, 1H).	501	499
110	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ : 0.44 (d, $J = 7.29$ Hz, 3H), 0.95 (s, 9H), 1.02 (s, 3H), 1.25 (s, 3H), 1.34-1.41 (m, 2H), 2.13 (q, $J = 7.29$ Hz, 1H), 2.31 (s, 6H), 2.59-2.66 (m, 2H), 3.66 (d, $J = 12.02$ Hz, 1H), 4.23 (d, $J = 12.25$ Hz, 1H), 6.21 (s, 1H), 7.06 (s, 1H), 7.26-7.38 (m, 3H), 12.49 (br s, 1H).	501	499
111	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 0.83 (d, $J = 5.38$ Hz, 3H), 0.98-0.92 (m, 1H), 0.99 (s, 9H), 1.45-1.49	485	483

	(m, 2H), 1.54-1.67 (m, 3H), 1.95-2.07 (m, 9H), 2.11-2.14 (m, 1H), 2.52-2.56 (m, 1H), 2.65-2.69 (m, 2H), 5.38 (s, 1H), 5.92 (s, 1H), 7.15 (dd, J = 8.07, 1.96 Hz, 1H), 7.20 (d, J = 8.07 Hz, 1H), 7.29 (d, J = 1.96 Hz, 1H).		
112	¹ H-NMR (400 MHz, CDCl ₃) δ: 0.83 (d, J = 5.38 Hz, 3H), 0.98-0.92 (m, 1H), 0.99 (s, 9H), 1.45-1.49 (m, 2H), 1.54-1.67 (m, 3H), 1.95-2.07 (m, 9H), 2.11-2.14 (m, 1H), 2.52-2.56 (m, 1H), 2.65-2.69 (m, 2H), 5.38 (s, 1H), 5.92 (s, 1H), 7.15 (dd, J = 8.07, 1.96 Hz, 1H), 7.20 (d, J = 8.07 Hz, 1H), 7.29 (d, J = 1.96 Hz, 1H).	485	483
113	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.62 (d, J = 7.40 Hz, 3H), 0.96 (d, J = 6.80 Hz, 3H), 1.09 (s, 9H), 1.24-1.31 (m, 1H), 1.49-1.65 (m, 2H), 1.84-1.92 (m, 1H), 2.06-2.22 (m, 2H), 2.26-2.37 (m, 1H), 2.29 (s, 6H), 2.89 (dd, J = 14.33, 9.25 Hz, 1H), 6.25 (s, 1H), 7.10 (s, 1H), 7.29-7.35 (m, 2H), 7.40-7.43 (m, 1H), 12.44 (br s, 1H).	533	531
114	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.62 (d, J = 7.40 Hz, 3H), 0.96 (d, J = 6.80 Hz, 3H), 1.09 (s, 9H), 1.24-1.31 (m, 1H), 1.49-1.65 (m, 2H), 1.84-1.92 (m, 1H), 2.06-2.22 (m, 2H), 2.26-2.37 (m, 1H), 2.29 (s, 6H), 2.89 (dd, J = 14.33, 9.25 Hz, 1H), 6.25 (s, 1H), 7.10 (s, 1H), 7.29-7.35 (m, 2H), 7.40-7.43 (m, 1H), 12.44 (br s, 1H).	533	531
115	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.63 (d, J = 7.60 Hz, 3H), 0.96 (d, J = 7.20 Hz, 3H), 1.10 (s, 9H), 1.24-1.31 (m, 1H), 1.48-1.64 (m, 2H), 1.80-1.96 (m, 9H), 2.06-2.22 (m, 2H), 2.87-2.93 (m, 1H), 6.19 (s, 1H), 6.98 (s, 1H), 7.32-7.35 (m, 2H), 7.43-7.46 (m, 1H), 12.25 (br s, 1H).	547	545
116	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.63 (d, J = 7.60 Hz, 3H), 0.96 (d, J = 7.20 Hz, 3H), 1.10 (s, 9H), 1.24-1.31 (m, 1H), 1.48-1.64 (m, 2H), 1.80-1.96 (m, 9H), 2.06-2.22 (m,	547	545

	2H), 2.87-2.93 (m, 1H), 6.19 (s, 1H), 6.98 (s, 1H), 7.32-7.35 (m, 2H), 7.43-7.46 (m, 1H), 12.25 (br s, 1H).		
117	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.39 (d, J = 7.28 Hz, 3H), 0.75 (s, 3H), 0.95 (s, 9H), 0.97 (s, 3H), 1.10-1.26 (m, 2H), 1.35-1.46 (m, 3H), 1.77-1.85 (m, 1H), 1.99-2.05 (m, 1H), 2.29 (s, 6H), 2.59-2.66 (m, 2H), 6.18 (s, 1H), 7.04 (s, 1H), 7.20-7.35 (m, 3H).	499	497
118	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.39 (d, J = 7.28 Hz, 3H), 0.75 (s, 3H), 0.95 (s, 9H), 0.97 (s, 3H), 1.10-1.26 (m, 2H), 1.35-1.46 (m, 3H), 1.77-1.85 (m, 1H), 1.99-2.05 (m, 1H), 2.29 (s, 6H), 2.59-2.66 (m, 2H), 6.18 (s, 1H), 7.04 (s, 1H), 7.20-7.35 (m, 3H).	499	497
119	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.45 (d, J = 7.40 Hz, 1.77H), 0.87 (d, J = 6.70 Hz, 1.23H), 0.95 (s, 9H), 1.01 (s, 1.77H), 1.05 (s, 1.23H), 1.11 (s, 1.23H), 1.25 (s, 1.77H), 1.36-1.41 (m, 2H), 1.80-1.98 (m, 8.41H), 2.09-2.14 (m, 0.59H), 2.59-2.66 (m, 2H), 3.67 (d, J = 12.25 Hz, 0.59H), 3.70 (d, J = 12.02 Hz, 0.41H), 4.06 (d, J = 11.79 Hz, 0.41H), 4.24 (d, J = 12.25 Hz, 0.59H), 5.82 (d, J = 1.85 Hz, 0.41H), 6.15 (s, 0.59H), 6.95 (s, 0.59H), 7.05 (s, 0.41H), 7.31-7.44 (m, 3H), 12.30 (br s, 1H).	545	513
120	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.45 (d, J = 7.40 Hz, 3H), 0.95 (s, 9H), 1.01 (s, 3H), 1.25 (s, 3H), 1.36-1.41 (m, 2H), 1.80-1.97 (m, 8H), 2.09-2.14 (m, 1H), 2.59-2.66 (m, 2H), 3.67 (d, J = 12.25 Hz, 1H), 4.24 (d, J = 12.25 Hz, 1H), 6.15 (s, 1H), 6.95 (s, 1H), 7.31-7.41 (m, 3H), 12.31 (s, 1H).	545	513
121	¹ H-NMR (400 MHz, DMSO-d ₆) δ: 0.87 (d, J = 6.70 Hz, 3H), 0.95 (s, 9H), 1.05 (s, 3H), 1.11 (s, 3H), 1.37-1.41 (m, 2H), 1.80-1.98 (m, 9H), 2.59-2.66 (m, 2H), 3.70 (d, J = 12.02 Hz, 1H), 4.06 (d, J =	545	513

	11.79 Hz, 1H), 5.82 (d, J = 1.85 Hz, 1H), 7.05 (s, 1H), 7.31-7.35 (m, 2H), 7.41-7.44 (m, 1H), 12.30 (br s, 1H).		
--	---	--	--

[0298]

[Test Example 1]

In-vitro assay for inhibitory activity against ROR γ
5 transcription

Inhibitory activity against ROR γ transcription was assessed for test compounds by either of the following Test Method A or B of reporter gene assay.

[0299]

10 (Test Method A)

cDNAs encoding human and mouse ROR γ ligand binding domains (LBD) were obtained based on the sequences of human ROR γ (Genbank registered number NM_005060.3) and mouse ROR γ (Genbank registered number NM_011281.2) (LBD sequences:
15 from Ser253 to Lys518 for human ROR γ ; from Ile251 to Lys516 for mouse ROR γ).

LBD cDNAs of human and mouse ROR γ were inserted into pFA-CMV vector (Stratagene), which expresses GAL4-DNA binding domain fusion protein. The resulting plasmids are
20 hereinafter referred to as GAL4-hROR γ plasmid and GAL4-mROR γ plasmid, respectively.

GAL4-hROR γ plasmid or GAL4-mROR γ plasmid was transiently co-transfected into Chinese hamster ovary cells

(CHO cells) with pG5-Luc (Promega), reporter plasmid expressing firefly luciferase in a GAL4-dependent manner.

TransIT (Registered trademark) CHO Transfection Kit (Mirus) was used to co-transfect CHO cells with the
5 plasmids. One day prior to the assay, CHO cells were suspended in HAM F-12 Nutrient medium containing 10%(v/v) fetal bovine serum and seeded at 6×10^6 cells per 175 cm² cell culture flask each. 54 μ L of TransIT (Registered trademark) CHO reagent was added into a 15 mL tube
10 containing 1.16 mL of HAM F-12 Nutrient medium without fetal bovine serum, and then mixed and incubated at room temperature for 10 minutes. 36 μ L of a plasmid solution containing 400 ng of GAL4-hROR γ plasmid, 9000 ng of pG5-Luc plasmid, and 8600 ng of pcDNA3 plasmid were added into the
15 tube and mixed gently. In the case of mouse assay, a plasmid solution containing 250 ng of GAL4-mROR γ plasmid, 9000 ng of pG5-Luc plasmid, and 8750 ng of pcDNA3 plasmid was added, instead. The mixture was incubated at room temperature for 10 minutes. 9 μ L each of CHO Mojo Reagent
20 were then added into each tube and mixed gently. The mixture was incubated at room temperature for 10 minutes. The resulting transfection reagent was applied to the cell culture. After incubation at 37°C, 5% CO₂ for 4 hours, the plasmid-transfected CHO cells were harvested by trypsin
25 treatment. The collected cells were resuspended in culture

medium and plated into a 384-well-white plate at 8,000 cells/50 μ L/well. The plate was let stand at room temperature for 1 hour and then further incubated at 37°C, 5% CO₂ for 3 hours. A test compound was dissolved in DMSO to obtain a concentration of 10 mM. Each solution was serially diluted with DMSO and further diluted with culture medium just before use. The test compound solutions were added to the cells in the plate at 8 different concentrations. The final concentration of DMSO was 0.2%(v/v). After the addition of test compound, the cells were incubated at 37°C, 5% CO₂ for 2 days.

Cell viability was tested by fluorescence method with Resazurin (Invitrogen). Two days after the addition of test compound, Resazurin was diluted to 20 μ M with a medium. 10 μ L each of the diluted solution of Resazurin was added into the 384-well-plate. Just after the addition, fluorescence at 615 nm was measured with a 570-nm excitation wavelength (0-hour value). After incubation at 37°C, 5% CO₂ for 2 hours, fluorescence at 615 nm was measured with the 570-nm excitation wavelength (2-hour value). A fluorescence count (2h-0h) was calculated by deducting the 0-hour value from the 2-hour value. The fluorescence count (2h-0h) in the cells treated with 0.2% DMSO alone was defined as 100%, and the cell viability after treatment with the test compound was calculated as a

percentage, %-of-control, based on the value of 0.2% DMSO alone. When the cell viability was 70% or less, a test compound was assessed to have cytotoxicity.

RORy transcriptional activity was detected as the
5 intracellular luciferase activity using SteadyLite HTS Reporter Gene Assay System (Perkin Elmer). SteadyLite Reagent was diluted five-fold with Extension buffer (10 mM Tricine, 0.2%(w/v) bovine serum albumin, 0.02%(v/v) Tween-20) to obtain a luciferase substrate solution. After the
10 measurement of the cell viability with Resazurin, the medium was removed in the plate and the luciferase substrate solution was added thereto. After the incubation at room temperature for 10 minutes, luminescence of each well was measured with a microplate reader. The luciferase
15 activity derived from the luminescence count in a vehicle-control well treated with 0.2% DMSO alone was defined as 100%, and the luciferase activity after treatment with the test compound was calculated as a percentage, %-of-control, based on the value of the vehicle-control. An EC₅₀ value
20 of the test compound was calculated by curve fitting with GraphPad Prism. The luminescence counts at the concentration of test compound where cytotoxicity was observed were excluded from data analysis.

[0300]

25 (Test Method B)

cDNAs encoding human and mouse ROR γ ligand binding domains (LBD) were obtained based on the sequences of human ROR γ (Genbank registered number NM_005060.3) and mouse ROR γ (Genbank registered number NM_011281.2) (LBD sequences: 5 from Ser253 to Lys518 for human ROR γ ; from Ile251 to Lys516 for mouse ROR γ).

LBD cDNAs of human and mouse ROR γ were inserted into pFA-CMV vector (Agilent Technologies, Inc.), which expresses GAL4-DNA binding domain fusion protein. The 10 resulting plasmids are hereinafter referred to as pFA/hROR γ plasmid and pFA/mROR γ plasmid, respectively.

pFA/hROR γ plasmid or pFA/mROR γ plasmid was transiently co-transfected into Chinese hamster ovary cells (CHO cells) with pG5-Luc (Promega), reporter plasmid expressing firefly 15 luciferase in a GAL4-dependent manner.

TransIT (Registered trademark) CHO Transfection Kit (Mirus) was used to co-transfect CHO cells with the plasmids. One day prior to the assay, CHO cells were suspended in HAM F-12 Nutrient medium containing 10%(v/v) 20 fetal bovine serum and seeded at 5.5×10^6 cells per 225 cm² cell culture flask each. 72 μ L of TransIT (Registered trademark) CHO reagent was added into a 2 mL tube containing 1.55 mL of Opti-MEM, and then mixed and incubated at room temperature for 10 minutes. 50.4 μ L of a 25 plasmid solution containing 300 ng of pFA/hROR γ plasmid,

12000 ng of pG5-Luc plasmid, and 11700 ng of pcDNA3.1 plasmid were added into the tube and mixed gently. In the case of mouse assay, a plasmid solution containing 300 ng of pFA/mRORγ plasmid, 12000 ng of pG5-Luc plasmid, and 5 11700 ng of pcDNA3.1 plasmid was added, instead. The mixture was incubated at room temperature for 10 minutes. 12 μL each of CHO Mojo Reagent were then added into each tube and mixed gently. The mixture was incubated at room temperature for 10 minutes. The resulting transfection 10 reagent was applied to the cell culture. After incubation at 37°C, 5% CO₂ for 4 hours, the plasmid-transfected CHO cells were harvested by trypsin treatment. The collected cells were resuspended in culture medium and plated into a 384-well-white plate at 8,000 cells/35 μL/well. The plate 15 was let stand at room temperature for 1 hour and then further incubated at 37°C, 5% CO₂ for 3 hours. A test compound was dissolved in dimethyl sulfoxide (DMSO) to obtain a concentration of 10 mM. Each solution was serially diluted with DMSO and further diluted with culture 20 medium just before use. The test compound solutions were added to the cells in the plate at 6 different concentrations. The final concentration of DMSO was 0.2%(v/v). After the addition of test compound, the cells were incubated at 37°C, 5% CO₂ for 2 days.

25 Cell viability was tested by luminescence method with

CellTiter-Glo (Promega). Two days after the addition of test compound, 40 μ L each of CellTiter-Glo was added into the 384-well-plate. Ten minutes after the addition, luminescence was measured for each well with a microplate reader. The luminescence count in the cells treated with 0.2% DMSO alone was defined as 100%, and the cell viability after treatment with the test compound was calculated as a percentage, %-of-control, based on the value of 0.2% DMSO alone. When the cell viability was 70% or less, a test compound was assessed to have cytotoxicity.

RORy transcriptional activity was detected as the intracellular luciferase activity using SteadyLite HTS Reporter Gene Assay System (Perkin Elmer). SteadyLite Reagent was diluted 2.5-fold with Extension buffer (10 mM Tricine, 0.2%(w/v) bovine serum albumin, 0.02%(v/v) Tween-20) to obtain a luciferase substrate solution. Two days after the addition of test compound, 40 μ L each of the luciferase substrate solution was added into the 384-well-plate. After the incubation at room temperature for 10 minutes, luminescence of each well was measured with a microplate reader. The luciferase activity derived from the luminescence count in a vehicle-control well treated with 0.2% DMSO alone was defined as 100%, and the luciferase activity after treatment of the test compound was calculated as a percentage, %-of-control, based on the

value of the vehicle-control. An EC₅₀ value of the test compound was calculated by curve fitting. The luminescence counts at the concentration of test compound where cytotoxicity was observed were excluded from data analysis.

5 [0301]

The results are shown in the table below.

The value with (%) is the activity after treatment with a test compound which was calculated as a %-of-control value based on 100% of the vehicle-control treated with
10 0.2% DMSO alone.

Example	LUC EC ₅₀ (μM)		Test Method
	hRORγ	mRORγ	
1	1.693	0.536	A
2	0.030	0.022	A
3	0.008	0.013	A
4	1.022	1.014	A
5	0.007	0.009	A
6	0.554	0.714	A
7	0.009	0.011	A
8	0.427	0.381	A
9	0.819	1.466	A
10	0.016	0.030	A
11	0.021	0.022	A
12	2.584	3.429	A
13	5.701	5.374	A
14	2.878	3.805	A
15	>8 (71%)	>8 (69%)	A
16	0.755	0.774	A

17	>20 (65%)	>20 (78%)	A
18	0.773	0.452	A
19	17.130	15.590	A
20	0.025	0.025	A
21	2.263	1.270	A
22	16.170	>8 (74%)	A
23	0.224	0.106	A
24	0.332	0.126	A
25	0.092	0.105	A
26	0.351	0.243	A
27	0.027	0.039	A
28	>20 (64%)	>20 (73%)	A
29	0.173	0.103	A
30	>20 (60%)	>20 (54%)	A
31	0.048	0.048	A
32	0.021	0.018	A
33	2.964	>3.2 (52%)	A
34	0.087	0.169	B
35	4.193	4.974	B
36	0.435	0.071	A
37	3.870	5.508	A
38	0.011	0.021	A
39	0.032	0.021	A
40	4.638	5.328	A
41	0.017	0.013	A
42	0.290	0.139	A
43	0.043	0.159	A
44	1.274	0.482	A
45	1.627	0.840	A
46	0.302	0.211	A

47	>8 (51%)	8.828	A
48	1.990	1.867	A
49	>3.2 (85%)	>3.2 (87%)	A
50	>8 (51%)	7.479	A
51	4.971	1.796	A
52	0.018	0.029	A
53	>3.2 (69%)	2.732	A
54	0.064	0.035	A
55	0.093	0.058	A
56	0.020	0.029	A
57	0.034	0.021	A
58	0.033	0.020	A
59	0.279	0.149	A
60	0.118	0.072	A
61	0.106	0.068	A
62	0.151	0.099	A
63	0.022	0.022	A
64	0.034	0.052	A
65	1.159	0.160	A
66	0.024	0.030	A
67	4.719	8.341	A
68	0.015	0.016	A
69	0.015	0.027	A
70	0.541	0.516	A
71	0.010	0.007	A
72	0.028	0.015	A
73	0.014	0.027	A
74	0.016	0.024	A
75	0.061	0.038	A
76	2.406	3.031	A

77	0.024	0.031	A
78	2.203	>3.2 (51%)	A
79	0.018	0.022	A
80	>10 (64%)	>10 (61%)	A
81	0.013	0.019	A
82	9.866	9.792	A
83	0.014	0.018	A
84	>10 (61%)	>10 (74%)	A
85	0.033	0.073	A
86	4.942	4.366	B
87	0.038	0.029	B
88	0.751	0.169	B
89	3.707	4.191	B
90	0.022	0.023	B
91	0.111	0.185	B
92	0.042	0.019	B
93	3.968	4.231	B
94	0.046	0.082	B
95	2.344	0.455	B
96	>10 (84%)	>10 (84%)	B
97	0.110	0.156	B
98	4.359	5.107	B
99	0.053	0.146	B
100	0.206	0.055	B
101	7.284	1.390	B
102	3.954	4.572	B
103	0.026	0.046	B
104	5.261	5.278	B
105	0.102	0.113	B
106	4.398	4.778	B

107	0.027	0.032	B
108	4.689	5.053	B
109	0.033	0.036	B
110	0.010	0.012	B
111	3.449	3.660	B
112	0.035	0.038	B
113	0.033	0.037	B
114	0.027	0.050	B
115	0.031	0.049	B
116	0.044	0.101	B
117	4.679	5.296	B
118	0.033	0.082	B
119	>10	>10	B
120	0.013	0.020	B
121	0.035	0.040	B

[0302]

[Formulation Examples]

Formulation Examples in the present invention include,
 5 for example, the following formulations. The present
 invention, however, is not intended to be limited to these
 Formulation Examples.

[0303]

Formulation Example 1 (Preparation of a capsule)

- | | | | |
|----|----|----------------------------|-------|
| 10 | 1) | Example 1 Compound | 30 mg |
| | 2) | Microcrystalline cellulose | 10 mg |
| | 3) | Lactose | 19 mg |
| | 4) | Magnesium stearate | 1 mg |

Ingredients 1), 2), 3), and 4) are mixed to be filled in a gelatin capsule.

Formulation Example 2 (Preparation of a tablet)

	1)	Example 1 Compound	10 g
5	2)	Lactose	50 g
	3)	Cornstarch	15 g
	4)	Carmellose calcium	44 g
	5)	Magnesium stearate	1 g

The total amount of Ingredients 1), 2), and 3) and 30
10 g of Ingredient 4) are combined with water, dried in vacuo,
and then granulated. The resulted granules are mixed with
14 g of Ingredient 4) and 1 g of Ingredient 5), and
tableted with a tableting machine. In this manner, 1000
tablets comprising 10 mg of Example 1 Compound per each are
15 obtained.


INDUSTRIAL APPLICABILITY

[0304]

A compound of Formula [I] or a pharmaceutically
20 acceptable salt thereof is expected to be useful for
treating or preventing autoimmune diseases, allergic
diseases, dry eye, fibrosis, cancers, metabolic disease,
ischemia, cardiomyopathy, hypertension, and periodontal
disease.

CLAIMS

1. A compound of Formula [I]:



5 wherein R^1 is

- (1) C_{1-8} alkyl,
- (2) halo- C_{1-8} alkyl,
- (3) C_{3-8} cycloalkyl optionally substituted with the same or different 1 to 3 substituents selected from Group A^1 , or
- 10 (4) C_{3-8} cycloalkyl- C_{1-4} alkyl wherein the C_{3-8} cycloalkyl moiety may be optionally substituted with the same or different 1 to 3 substituents selected from Group A^1 ,

Group A^1 is

- (1) halogen,
- 15 (2) C_{1-4} alkyl, and
- (3) halo- C_{1-4} alkyl,

X^1 is

- (1) a bond, or
- (2) -O-,
 R^2 is
- (1) hydrogen, or
- 5 (2) halogen,
 R^3 is
- (1) hydrogen, or
- (2) $-Y^3-COO-R^{30}$,
 Y^3 is
- 10 (1) C_{1-8} alkylene,
 (2) C_{3-8} cycloalkylene,
 (3) bridged C_{5-8} cycloalkylene, or
 (4) C_{6-14} arylene,
 R^{30} is
- 15 (1) hydrogen, or
 (2) C_{1-4} alkyl,
 X^2 is
- (1) $=C(R^4)-$, or
- (2) $=N-$,
 R^4 is
- 20 (1) hydrogen, or
 (2) C_{1-4} alkyl,
 X^3 is
- (1) $-C(R^5)(R^6)-$,
 X^4 is
- 25

- (1) a bond, or
- (2) $-C(R^7)(R^8)-$,
 X^5 is
 - (1) $-C(R^9)(R^{10})-$,
 - 5 (2) $-N(R^{11})-$, or
 - (3) $-O-$,
 R^5 and R^6 are each independently
 - (1) hydrogen,
 - (2) C_{1-4} alkyl,
 - 10 (3) halo- C_{1-4} alkyl,
 - (4) cyano- C_{1-4} alkyl, or
 - (5) C_{1-4} alkyl substituted with one substituent selected
 from the group consisting of $-O-R^{51}$, $-CO-R^{61}$, $-COO-R^{52}$, $-$
 $N(R^{71})(R^{72})$, $-CO-N(R^{73})(R^{74})$, $-N(R^{75})-CO-R^{62}$, $-N(R^{76})-COO-R^{53}$,
 15 and $-O-S(O)_2-R^{63}$,
 R^7 , R^8 , R^9 , and R^{10} are each independently
 - (1) hydrogen,
 - (2) halogen,
 - (3) cyano,
 - 20 (4) hydroxy,
 - (5) C_{1-4} alkyl,
 - (6) halo- C_{1-4} alkyl,
 - (7) cyano- C_{1-4} alkyl,
 - (8) C_{1-4} alkoxy, or
 - 25 (9) C_{1-4} alkyl substituted with one substituent selected

from the group consisting of $-O-R^{51}$, $-CO-R^{61}$, $-COO-R^{52}$, $-N(R^{71})(R^{72})$, $-CO-N(R^{73})(R^{74})$, $-N(R^{75})-CO-R^{62}$, $-N(R^{76})-COO-R^{53}$, and $-O-S(O)_2-R^{63}$,

R^{51} , R^{52} , and R^{53} are each independently

- 5 (1) hydrogen,
- (2) C_{1-4} alkyl, or
- (3) C_{6-14} aryl- C_{1-4} alkyl,

R^{61} , R^{62} , and R^{63} are each independently

- (1) C_{1-4} alkyl,

10 R^{71} , R^{72} , R^{73} , R^{74} , R^{75} , and R^{76} are each independently

- (1) hydrogen, or
- (2) C_{1-4} alkyl,

R^{11} is

- (1) $-CO-R^{111}$, or

15 (2) $-COO-R^{112}$,

R^{111} is


- (1) C_{1-4} alkyl,

R^{112} is

- (1) C_{1-4} alkyl, or a pharmaceutically acceptable salt

20 thereof.

2. The compound according to claim 1, having a structure of Formula [II]:



wherein each variable is defined as defined in claim 1, or a pharmaceutically acceptable salt thereof.

5 3. The compound according to claim 1 or 2, wherein X^2 is =N-, or a pharmaceutically acceptable salt thereof.

4. The compound according to claim 1 or 2, wherein X^2 is =C(R^4)- and R^4 is hydrogen, or a pharmaceutically
10 acceptable salt thereof.

5. The compound according to any one of claims 1 to 4, wherein R^3 is hydrogen, or a pharmaceutically acceptable salt thereof.

15

6. The compound according to any one of claims 1 to 4, wherein R^3 is $-Y^3-COO-R^{30}$,

Y^3 is

- (1) C_{1-8} alkylene,
- (2) C_{3-8} cycloalkylene, or
- (3) bridged C_{5-8} cycloalkylene,

5 R^{30} is hydrogen or C_{1-4} alkyl, or a pharmaceutically acceptable salt thereof.

7. The compound according to any one of claims 1 to 6, wherein R^2 is halogen, or a pharmaceutically acceptable
10 salt thereof.

8. The compound according to any one of claims 1 to 7, wherein R^1 is C_{1-8} alkyl and X^1 is a bond, or a pharmaceutically acceptable salt thereof.

15







9. The compound according to any one of claims 1 to 8, wherein R^5 and R^6 are each independently hydrogen or C_{1-4} alkyl, or a pharmaceutically acceptable salt thereof.









20 10. The compound according to any one of claims 1 to 9, wherein X^4 is a bond or $-C(R^7)(R^8)-$ and both of R^7 and R^8 are hydrogen, or a pharmaceutically acceptable salt thereof.


11. The compound according to any one of claims 1 to 10,
25 wherein X^5 is $-C(R^9)(R^{10})-$ or $-O-$ and both of R^9 and R^{10} are

hydrogen, or a pharmaceutically acceptable salt thereof.

12. The compound according to claim 1, selected from the following compound group:







or a pharmaceutically acceptable salt thereof.

13. A pharmaceutical composition comprising a compound according to any one of claims 1 to 12 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

14. An ROR γ antagonist comprising a compound according to any one of claims 1 to 12 or a pharmaceutically acceptable salt thereof.

15. A therapeutic or preventive agent for a disease selected from the group consisting of autoimmune diseases, allergic diseases, dry eye, fibrosis, cancers, metabolic disease, ischemia, cardiomyopathy, hypertension, and periodontal disease, comprising a compound according to any one of claims 1 to 12 or a pharmaceutically acceptable salt thereof.

16. A method of antagonizing ROR γ , comprising administering a therapeutically effective amount of a compound according to any one of claims 1 to 12 or a pharmaceutically acceptable salt thereof to a mammal.

5

17. A method of treating or preventing a disease selected from the group consisting of autoimmune diseases, allergic diseases, dry eye, fibrosis, cancers, metabolic disease, ischemia, cardiomyopathy, hypertension, and periodontal
10 disease, comprising administering a therapeutically effective amount of a compound according to any one of claims 1 to 12 or a pharmaceutically acceptable salt thereof to a mammal.

15 18. Use of a compound according to any one of claims 1 to 12 or a pharmaceutically acceptable salt thereof in the manufacture of an ROR γ antagonist.

19. Use of a compound according to any one of claims 1 to
20 12 or a pharmaceutically acceptable salt thereof in the manufacture of a therapeutic or preventive agent for a disease selected from the group consisting of autoimmune diseases, allergic diseases, dry eye, fibrosis, cancers, metabolic disease, ischemia, cardiomyopathy, hypertension,
25 and periodontal disease.

20. A compound according to any one of claims 1 to 12 or a pharmaceutically acceptable salt thereof for use in an ROR γ antagonist.

5

21. A compound according to any one of claims 1 to 12 or a pharmaceutically acceptable salt thereof for use in treating or preventing a disease selected from the group consisting of autoimmune diseases, allergic diseases, dry
10 eye, fibrosis, cancers, metabolic disease, ischemia, cardiomyopathy, hypertension, and periodontal disease.