

WHAT IS CLAIMED IS:

1. An antibody or antigen-binding fragment thereof that specifically binds to the same CD37 epitope as an antibody comprising the polypeptide of SEQ ID NO:9 and the polypeptide of SEQ ID NO:10.
2. An antibody or antigen-binding fragment thereof that specifically binds to CD37, wherein said antibody or fragment thereof competitively inhibits binding to CD37 of an antibody comprising the polypeptide of SEQ ID NO:9 and the polypeptide of SEQ ID NO:10.
3. The antibody or antigen-binding fragment thereof of claim 1 or 2, wherein the antibody or fragment thereof comprises the heavy chain variable region (VH) CDR1, CDR2, and CDR3 sequences of SEQ ID NOs: 3-5, respectively, and the light chain variable region (VL) CDR1, CDR2, and CDR3 sequences of SEQ ID NOs: 6-8, respectively.
4. An antibody or antigen-binding fragment thereof that specifically binds to CD37, wherein said antibody or fragment thereof comprises the VH CDR1, CDR2, and CDR3 sequences of SEQ ID NOs: 3-5 and the VL CDR1, CDR2, and CDR3 sequences of SEQ ID NOs: 6-8, respectively.
5. The antibody or antigen-binding fragment thereof of any one of claims 1-4, wherein the antibody or fragment thereof comprises the polypeptide sequences of SEQ ID NOs:9 and/or 10.
6. The antibody or antigen-binding fragment thereof of any of claims 1-5, wherein said antibody or antigen-binding fragment thereof is murine, humanized, chimeric, or human.
7. The antibody or antigen-binding fragment thereof of any of claims 1-6, which is a full length antibody.
8. The antibody or antigen-binding fragment thereof of any of claims 1-6, which is an antigen-binding fragment.
9. The antibody or antigen-binding fragment thereof of claim 8, wherein said antibody or antigen-binding fragment thereof comprises a Fab, Fab', F(ab')₂, single chain Fv or scFv, disulfide linked Fv, intrabody, IgGΔCH₂, minibody, F(ab')₃, tetrabody, triabody, diabody, DVD-Ig, mAb₂, (scFv)₂, or scFv-Fc.
10. The antibody or antigen-binding fragment thereof of any one of claims 1-9, which binds to a human CD37 with a K_d of about 1.0 nM or better.

11. The antibody or antigen-binding fragment thereof of any one of claims 1-10, wherein the antibody or antigen-binding fragment thereof is detectably labeled.
12. A method of detecting CD37 expression in a sample comprising contacting said sample with the antibody or antigen-binding fragment thereof of any of claims 1-11.
13. The method of claim 12, wherein CD37 expression is determined by immunohistochemical (IHC) assay.
14. A therapeutically active agent comprising an anti-CD37 antibody or antigen-binding fragment thereof of claims 1-11 for use in treating in a subject, wherein increased expression of CD37 has been detected in a cancerous sample from said subject using the antibody or antigen-binding fragment thereof of any one of claims 1-11.
15. The therapeutically active agent of claim 14, wherein the detecting is by immunohistochemistry (IHC).
16. The therapeutically active agent of claim 14 or 15, wherein the cancer is a leukemia or a lymphoma.
17. The therapeutically active agent of claim 14 or 15, wherein said cancer is selected from the group consisting of wherein said cancer is selected from the group consisting of B cell lymphomas, NHL, precursor B cell lymphoblastic leukemia/lymphoma and mature B cell neoplasms, B cell chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL), B cell prolymphocytic leukemia, lymphoplasmacytic lymphoma, mantle cell lymphoma (MCL), follicular lymphoma (FL), low grade, intermediate-grade and high-grade (FL), cutaneous follicle center lymphoma, marginal zone B cell lymphoma. MALT type marginal zone B cell lymphoma, nodal marginal zone B cell lymphoma, splenic type marginal zone B cell lymphoma, hairy cell leukemia, diffuse large B cell lymphoma, Burkitt's lymphoma, plasmacytoma, plasma cell myeloma, post-transplant lymphoproliferative disorder, Waldenstrom's macroglobulinemia, and anaplastic large-cell lymphoma (ALCL).
18. The therapeutically active agent of claim 17, wherein said cancer is diffuse large B cell lymphoma.
19. The therapeutically active agent of any of claims 14-18, wherein the therapeutically active agent is an antibody maytansinoid conjugate that comprises a CD37 antibody comprising a

heavy chain variable region comprising SEQ ID NO:28 and a light chain variable region comprising SEQ ID NO:29.

20. The therapeutically active agent of claim 19, wherein the therapeutically active agent comprises the maytansinoid DM1, the non-cleavable SMCC linker, and wherein the CD37 antibody comprises a full length heavy chain comprising SEQ ID NO:30 and a full length light chain comprising SEQ ID NO:31.