

NOVEL COMPOSITIONS AND METHODS

FIELD OF THE INVENTION

5 The present invention relates to polypeptides and polynucleotides for use in the treatment or prevention of tuberculosis, in particular for use in the treatment or prevention of latent tuberculosis and in the prevention or delay of reactivation of tuberculosis (and also to related methods). The present invention further relates to pharmaceutical and immunogenic compositions comprising said polypeptides and polynucleotides, and to methods for the
10 diagnosis of tuberculosis (in particular latent tuberculosis).

BACKGROUND OF THE INVENTION

Tuberculosis (TB) is a chronic infectious disease caused by infection with
15 *Mycobacterium tuberculosis* and other *Mycobacterium* species. It is a major disease in developing countries, as well as an increasing problem in developed areas of the world. More than 2 billion people are believed to be infected with TB bacilli, with about 9.2 million new cases of TB and 1.7 million deaths each year. 10% of those infected with TB bacilli will develop active TB, each person with active TB infecting an average of 10 to 15 others per
20 year. While annual incidence rates have peaked globally, the number of deaths and cases is still rising due to population growth (World Health Organisation *Tuberculosis Facts* 2008).

Mycobacterium tuberculosis infects individuals through the respiratory route. Alveolar macrophages engulf the bacterium, but it is able to survive and proliferate by inhibiting
25 phagosome fusion with acidic lysosomes. A complex immune response involving CD4+ and CD8+ T cells ensues, ultimately resulting in the formation of a granuloma. Central to the success of *Mycobacterium tuberculosis* as a pathogen is the fact that the isolated, but not eradicated, bacterium may persist for long periods, leaving an individual vulnerable to the later development of active TB.

30 Fewer than 5% of infected individuals develop active TB in the first years after infection. The granuloma can persist for decades and is believed to contain live *Mycobacterium tuberculosis* in a state of dormancy, deprived of oxygen and nutrients. However, recently it has been suggested that the majority of the bacteria in the dormancy state are located in non-
35 macrophage cell types spread throughout the body (Locht et al, *Expert Opin. Biol. Ther.* 2007 7(11):1665-1677). The development of active TB occurs when the balance between the host's natural immunity and the pathogen changes, for example as a result of an immunosuppressive

event (Anderson P *Trends in Microbiology* 2007 15(1):7-13; Ehlers S *Infection* 2009 37(2):87-95).

A dynamic hypothesis describing the balance between latent TB and active TB has also been proposed (Cardana P-J *Inflammation & Allergy – Drug Targets* 2006 6:27-39; Cardana P-J *Infection* 2009 37(2):80-86).

Although an infection may be asymptomatic for a considerable period of time, the active disease is most commonly manifested as an acute inflammation of the lungs, resulting in tiredness, weight loss, fever and a persistent cough. If untreated, serious complications and death typically result.

Tuberculosis can generally be controlled using extended antibiotic therapy, although such treatment is not sufficient to prevent the spread of the disease. Infected individuals may be asymptomatic, but contagious, for some time. In addition, although compliance with the treatment regimen is critical, patient behaviour is difficult to monitor. Some patients do not complete the course of treatment, which can lead to ineffective treatment and the development of drug resistance.

Multidrug-resistant TB (MDR-TB) is a form which fails to respond to first line medications. 5% of all TB cases are MDR-TB, with an estimated 490,000 new MDR-TB cases occurring each year. Extensively drug-resistant TB (XDR-TB) occurs when resistance to second line medications develops on top of MDR-TB. It is estimated that 40,000 new cases of the virtually untreatable XDR-TB arise annually (World Health Organisation *Tuberculosis Facts* 2008).

Even if a full course of antibiotic treatment is completed, infection with *M. tuberculosis* may not be eradicated from the infected individual and may remain as a latent infection that can be reactivated.

In order to control the spread of tuberculosis, effective vaccination and accurate early diagnosis of the disease are of utmost importance.

Diagnosis of latent TB infection is commonly achieved using the tuberculin skin test, which involves intradermal exposure to tuberculin protein-purified derivative (PPD). Antigen-specific T cell responses result in measurable induration at the injection site by 48-72 hours after injection, which indicates exposure to mycobacterial antigens. Sensitivity and specificity have, however, been a problem with this test, and individuals vaccinated with BCG cannot always be

easily distinguished from infected individuals (this is particularly important in light of the fact that BCG does not protect against latent infection). In general, individuals who have received BCG but are not infected by *M. tuberculosis* show a PPD reaction below 10 mm in diameter whereas people who have a PPD reaction above 10 mm in diameter are considered to have
 5 been infected by *M. tuberculosis*. However, this rule is not applicable to individuals with immunosuppression due to HIV infection, which may result in a PPD reaction below 10 mm in diameter); or in endemic countries, where people infected by non-tuberculosis mycobacteria can show a PPD reaction above 10 mm in diameter.

- 10 Progress over recent years has seen the development of *in vitro* T cell based assays, based on interferon-gamma release and using antigens which are more specific to *M. tuberculosis* than PPD, namely ESAT-6 and CFP-10. These high specificity tests appear to be at least as sensitive as the tuberculin skin test and also demonstrate less cross-reactivity due to BCG vaccination. See Pai M et al *Expert Rev. Mol. Diagn.* 2006 6(3):413-422 for a recent review of
 15 latent TB diagnosis. However, since ESAT-6/CFP-10 are early stage antigens, assays based on ESAT-6/CFP-10 may only perform optimally in recently infected people. Consequently, the identification of new antigens specifically associated with latent tuberculosis may aid the development of more sensitive assays that could detect longer-term latent infections.
- 20 There remains a need for effective strategies for the treatment and prevention of tuberculosis, in particular the treatment and prevention of latent TB and the prevention of reactivation of TB.

BRIEF SUMMARY OF THE INVENTION

- 25 The present invention relates generally to the identification of Rv3616c as a latent TB antigen and to related methods and uses in the prevention and treatment of latent TB, and the prevention or delay of TB reactivation.

The present invention provides a polypeptide which comprises:

- 30 (i) an Rv3616c protein sequence;
 (ii) a variant of an Rv3616c protein sequence; or
 (iii) an immunogenic fragment of an Rv3616c protein sequence;
 for use as a medicament in the treatment or prevention of latent TB.

- 35 A further aspect of the invention relates to a method for the treatment or prevention of latent TB comprising the administration of a safe and effective amount of a polypeptide comprising:
 (i) an Rv3616c protein sequence;

- (ii) a variant of an Rv3616c protein sequence; or
- (iii) an immunogenic fragment of an Rv3616c protein sequence;

to a subject in need thereof, wherein said polypeptide induces an immune response, in particular an immune response against *Mycobacterium tuberculosis*.

5

The use of a polypeptide comprising:

- (i) an Rv3616c protein sequence;
- (ii) a variant of an Rv3616c protein sequence; or
- (iii) an immunogenic fragment of an Rv3616c protein sequence;

10 in the manufacture of a medicament for the treatment or prevention of latent TB, represents another aspect of the invention.

Also provided is a polynucleotide comprising a nucleic acid sequence encoding a polypeptide which comprises:

- 15
- (i) an Rv3616c protein sequence;
 - (ii) a variant of an Rv3616c protein sequence; or
 - (iii) an immunogenic fragment of an Rv3616c protein sequence;

for use as a medicament in the treatment or prevention of latent TB.

20 A further aspect of the invention relates to a method for the treatment or prevention of latent TB comprising the administration of a safe and effective amount of a polynucleotide comprising a nucleic acid sequence encoding a polypeptide comprising:

- (i) an Rv3616c protein sequence;
- (ii) a variant of an Rv3616c protein sequence; or
- 25 (iii) an immunogenic fragment of an Rv3616c protein sequence;

to a subject in need thereof, wherein said polynucleotide induces an immune response, in particular an immune response against *Mycobacterium tuberculosis*.

Use of a polynucleotide comprising a nucleic acid sequence encoding a polypeptide comprising:

- 30
- (i) an Rv3616c protein sequence;
 - (ii) a variant of an Rv3616c protein sequence; or
 - (iii) an immunogenic fragment of an Rv3616c protein sequence;

in the manufacture of a medicament for the treatment or prevention of latent TB, represents
35 another aspect of the invention.

Additionally, there is provided a pharmaceutical composition comprising:

(a) a polypeptide which comprises:
 (i) an Rv3616c protein sequence;
 (ii) a variant of an Rv3616c protein sequence; or
 (iii) an immunogenic fragment of an Rv3616c protein sequence; or
5 (b) a polynucleotide comprising a nucleic acid sequence encoding the polypeptide of
 (a);
 and
 (c) a pharmaceutically acceptable carrier or excipient,
for use as a medicament in the treatment or prevention of latent TB.

10

Further, there is provided an immunogenic composition comprising:

(a) a polypeptide which comprises:
 (i) an Rv3616c protein sequence;
 (ii) a variant of an Rv3616c protein sequence; or
15 (iii) an immunogenic fragment of an Rv3616c protein sequence; or
(b) a polynucleotide comprising a nucleic acid sequence encoding the polypeptide of
 (a);
 and
 (c) a non-specific immune response enhancer,
20 for use as a medicament in the treatment or prevention of latent TB.

Additionally provided is an antibody or fragment thereof which specifically binds to a polypeptide comprising:

(i) an Rv3616c protein sequence;
25 (ii) a variant of an Rv3616c protein sequence; or
 (iii) an immunogenic fragment of an Rv3616c protein sequence,
for use in diagnosis of latent TB (such as methods for the diagnosis of latent tuberculosis comprising determining the presence of an antibody or fragment thereof which specifically binds to the polypeptides of the invention in a biological sample from a test subject).

30

There is also provided a method for the treatment of latent TB comprising the steps:

(i) Identifying a subject as having a latent TB infection (e.g. by PPD or T cell based assays); and
(ii) administering to said subject a safe and effective amount of a polypeptide or
35 polynucleotide as described herein (such as in the form of a pharmaceutical composition or immunogenic composition).

Also provided is the use of a polypeptide of the present invention in the manufacture of a diagnostic kit for the identification of latent TB in a test subject.

- 5 In one embodiment the subject receiving a polypeptide, polynucleotide or composition may have active tuberculosis (e.g. active infection by *M. tuberculosis*). In a second embodiment the subject may have latent tuberculosis (e.g. dormant infection by *M. tuberculosis*). In a third embodiment the subject may be free from tuberculosis (e.g. free from infection by *M. tuberculosis*).

10

A subject receiving a polypeptide, polynucleotide or composition may have previously been vaccinated for tuberculosis (e.g. vaccinated against infection by *M. tuberculosis*), such as having been vaccinated with *Bacillus Calmette-Guerin* (BCG). Alternatively, a subject receiving a polypeptide, polynucleotide or composition of the invention may not have

15 previously been vaccinated for tuberculosis (e.g. not vaccinated against infection by *M. tuberculosis*), such as not having been vaccinated with *Bacillus Calmette-Guerin* (BCG).

DESCRIPTION OF THE FIGURES

- 20 Figure 1: Rv3616c peptide alignment with full length sequence.
 Figure 2: PBMC responses to Rv3616c peptides.
 Figure 3: Percentage of CD4 and CD8 cells from immunised CB6F1 mice expressing IFN-gamma and/or IL-2 and/or TNF-alpha cytokines at day 21 (i.e. 7 days post second immunisation).
- 25 Figure 4: Cytokine profile at day 21 (i.e. 7 days post second immunisation) of the antigen specific CD4 response in immunised CB6F1 mice.
 Figure 5: Cytokine profile at day 21 (i.e. 7 days post second immunisation) of the antigen specific CD8 response in immunised CB6F1 mice.
 Figure 6: Percentage of CD4 and CD8 cells from immunised CB6F1 mice expressing IFN-gamma and/or IL-2 and/or TNF-alpha cytokines at day
- 30 35 (i.e. 7 days post third immunisation).
 Figure 7: Cytokine profile at day 35 (i.e. 7 days post third immunisation) of the antigen specific CD4 response in immunised CB6F1 mice.
 Figure 8: Cytokine profile at day 35 (i.e. 7 days post third immunisation) of the antigen specific CD8 response in immunised CB6F1 mice.
- 35

- Figure 9: Percentage of CD4 and CD8 cells from immunised C57BL/6 mice expressing IFN-gamma and/or IL-2 and/or TNF-alpha cytokines at day 21 (i.e. 7 days post second immunisation).
- 5 Figure 10: Cytokine profile at day 21 (i.e. 7 days post second immunisation) of the antigen specific CD4 response in immunised C57BL/6 mice.
- Figure 11: Percentage of CD4 and CD8 cells from immunised C57BL/6 mice expressing IFN-gamma and/or IL-2 and/or TNF-alpha cytokines at day 35 (i.e. 7 days post third immunisation).
- 10 Figure 12: Cytokine profile at day 35 (i.e. 7 days post third immunisation) of the antigen specific CD4 response in immunised C57BL/6 mice.
- Figure 13: Cytokine profile at day 35 (i.e. 7 days post third immunisation) of the antigen specific CD8 response in immunised C57BL/6 mice.
- Figure 14: Antigen-specific CD4 T cell responses in naive and latently infected humans.

15

DESCRIPTION OF THE LISTED SEQUENCES

- SEQ ID No: 1: polypeptide sequence of Rv3616c from *M. tuberculosis* H37Rv strain.
- SEQ ID No: 2: polynucleotide sequence of Rv3616c from *M. tuberculosis* H37Rv strain.
- 20 SEQ ID No: 3: polypeptide sequence of Rv3616c from *M. tuberculosis* CDC1551 strain.
- SEQ ID No: 4: polypeptide sequence of Rv3616c from *M. tuberculosis* F11 strain.
- SEQ ID No: 5: polypeptide sequence of Rv3616c from *M. tuberculosis* Haarlem A strain.
- SEQ ID No: 6: polypeptide sequence of Rv3616c from *M. tuberculosis* C strain.
- 25 SEQ ID No: 7: polypeptide sequence of Rv3616c from BCG.
- SEQ ID No: 8: polypeptide sequence of Mtb8.4.
- SEQ ID No: 9: polypeptide sequence of Mtb9.8.
- SEQ ID No: 10: polypeptide sequence of Mtb9.9.
- SEQ ID No: 11: polypeptide sequence of Ra12.
- 30 SEQ ID No: 12: polypeptide sequence of Ra35.
- SEQ ID No: 13: polypeptide sequence of TbH9.
- SEQ ID No: 14: polypeptide sequence of Mtb41.
- SEQ ID No: 15: polypeptide sequence of ESAT-6.
- SEQ ID No: 16: polypeptide sequence of Ag85A.
- 35 SEQ ID No: 17: polypeptide sequence of Ag85B.
- SEQ ID No: 18: polypeptide sequence of alpha-crystallin.
- SEQ ID No: 19: polypeptide sequence of MPT64.

	SEQ ID No: 20:	polypeptide sequence of Mtb32A.
	SEQ ID No: 21:	polypeptide sequence of Ser/Ala mutated mature Mtb32A.
	SEQ ID No: 22:	polypeptide sequence of TB10.4.
	SEQ ID No: 23:	polypeptide sequence of Mtb72f.
5	SEQ ID No: 24:	polypeptide sequence of M72.
	SEQ ID No: 25:	polypeptide sequence of Mtb71f.
	SEQ ID No: 26:	polypeptide sequence of M92 fusion.
	SEQ ID No: 27:	polypeptide sequence of M103 fusion.
	SEQ ID No: 28:	polypeptide sequence of M114 fusion.
10	SEQ ID No: 29:	putative human CD4 cell epitope 1.
	SEQ ID No: 30:	putative human CD4 cell epitope 2.
	SEQ ID No: 31:	putative human CD4 cell epitope 3.
	SEQ ID No: 32:	putative human CD4 cell epitope 4.
	SEQ ID No: 33:	putative human CD4 cell epitope 5.
15	SEQ ID No: 34:	putative human CD4 cell epitope 6.
	SEQ ID No: 35:	putative human CD4 cell epitope 7.
	SEQ ID No: 36:	putative human CD4 cell epitope 8.
	SEQ ID No: 37:	putative human CD4 cell epitope 9.
	SEQ ID No: 38:	putative human CD4 cell epitope 10.
20	SEQ ID No: 39:	putative human CD4 cell epitope 11.
	SEQ ID No: 40:	putative human CD4 cell epitope 12.
	SEQ ID No: 41:	putative human CD4 cell epitope 13.
	SEQ ID No: 42:	putative human CD4 cell epitope 14.
	SEQ ID No: 43:	putative human CD4 cell epitope 15.
25	SEQ ID No: 44:	putative human CD4 cell epitope 16.
	SEQ ID No: 45:	putative human CD4 cell epitope 17.
	SEQ ID No: 46:	putative human CD4 cell epitope 18.
	SEQ ID No: 47:	putative human CD4 cell epitope 19.
	SEQ ID No: 48:	putative human CD8 cell epitope 1.
30	SEQ ID No: 49:	putative human CD8 cell epitope 2.
	SEQ ID No: 50:	putative human CD8 cell epitope 3.
	SEQ ID No: 51:	putative human CD8 cell epitope 4.
	SEQ ID No: 52:	putative human CD8 cell epitope 5.
	SEQ ID No: 53:	putative human CD8 cell epitope 6.
35	SEQ ID No: 54:	putative human CD8 cell epitope 7.
	SEQ ID No: 55:	putative human CD8 cell epitope 8.
	SEQ ID No: 56:	putative human CD8 cell epitope 9.

	SEQ ID No: 57:	putative human CD8 cell epitope 10.
	SEQ ID No: 58:	putative human CD8 cell epitope 11.
	SEQ ID No: 59:	putative human CD8 cell epitope 12.
	SEQ ID No: 60:	putative human CD8 cell epitope 13.
5	SEQ ID No: 61:	putative human CD8 cell epitope 14.
	SEQ ID No: 62:	putative human CD8 cell epitope 15.
	SEQ ID No: 63:	putative human CD8 cell epitope 16.
	SEQ ID No: 64:	putative human CD8 cell epitope 17.
	SEQ ID No: 65:	putative human CD8 cell epitope 18.
10	SEQ ID No: 66:	putative human CD8 cell epitope 19.
	SEQ ID No: 67:	putative human CD8 cell epitope 20.
	SEQ ID No: 68:	putative human CD8 cell epitope 21.
	SEQ ID No: 69:	putative human CD8 cell epitope 22.
	SEQ ID No: 70:	putative human CD8 cell epitope 23.
15	SEQ ID No: 71:	putative human CD8 cell epitope 24.
	SEQ ID No: 72:	putative human CD8 cell epitope 25.
	SEQ ID No: 73:	putative human CD8 cell epitope 26.
	SEQ ID No: 74:	putative human CD8 cell epitope 27.
	SEQ ID No: 75:	putative human CD8 cell epitope 28.
20	SEQ ID No: 76:	putative human CD8 cell epitope 29.
	SEQ ID No: 77:	putative human CD8 cell epitope 30.
	SEQ ID No: 78:	putative human CD8 cell epitope 31.
	SEQ ID No: 79:	putative human CD8 cell epitope 32.
	SEQ ID No: 80:	putative human CD8 cell epitope 33.
25	SEQ ID No: 81:	putative human CD8 cell epitope 34.
	SEQ ID No: 82:	putative human CD8 cell epitope 35.
	SEQ ID No: 83:	putative human CD8 cell epitope 36.
	SEQ ID No: 84:	putative human CD8 cell epitope 37.
	SEQ ID No: 85:	putative human CD8 cell epitope 38.
30	SEQ ID No: 86:	putative human CD8 cell epitope 39.
	SEQ ID No: 87:	putative human CD8 cell epitope 40.
	SEQ ID No: 88:	putative human CD8 cell epitope 41.
	SEQ ID No: 89:	putative human CD8 cell epitope 42.
	SEQ ID No: 90:	putative human CD8 cell epitope 43.
35	SEQ ID No: 91:	putative human CD8 cell epitope 44.
	SEQ ID No: 92:	putative human CD8 cell epitope 45.
	SEQ ID No: 93:	putative human CD8 cell epitope 46.

	SEQ ID No: 94:	putative human CD8 cell epitope 47.
	SEQ ID No: 95:	putative human CD8 cell epitope 48.
	SEQ ID No: 96:	putative human CD8 cell epitope 49.
	SEQ ID No: 97:	putative human CD8 cell epitope 50.
5	SEQ ID No: 98:	putative human CD8 cell epitope 51.
	SEQ ID No: 99:	putative human CD8 cell epitope 52.
	SEQ ID No: 100:	putative human CD8 cell epitope 53.
	SEQ ID No: 101:	putative human CD8 cell epitope 54.
	SEQ ID No: 102:	putative human CD8 cell epitope 55.
10	SEQ ID No: 103:	putative human CD8 cell epitope 56.
	SEQ ID No: 104:	putative human CD8 cell epitope 57.
	SEQ ID No: 105:	putative human CD8 cell epitope 58.
	SEQ ID No: 106:	putative human CD8 cell epitope 59.
	SEQ ID No: 107:	putative human CD8 cell epitope 60.
15	SEQ ID No: 108:	putative human CD8 cell epitope 61.
	SEQ ID No: 109:	putative human CD8 cell epitope 62.
	SEQ ID No: 110:	putative human CD8 cell epitope 63.
	SEQ ID No: 111:	putative human CD8 cell epitope 64.
	SEQ ID No: 112:	putative human CD8 cell epitope 65.
20	SEQ ID No: 113:	putative human CD8 cell epitope 66.
	SEQ ID No: 114:	putative human CD8 cell epitope 67.
	SEQ ID No: 115:	putative human CD8 cell epitope 68.
	SEQ ID No: 116:	putative human CD8 cell epitope 69.
	SEQ ID No: 117:	putative human CD8 cell epitope 70.
25	SEQ ID No: 118:	putative human CD8 cell epitope 71.
	SEQ ID No: 119:	putative human CD8 cell epitope 72.
	SEQ ID No: 120:	putative human CD8 cell epitope 73.
	SEQ ID No: 121:	putative human CD8 cell epitope 74.
	SEQ ID No: 122:	putative human CD8 cell epitope 75.
30	SEQ ID No: 123:	putative human CD8 cell epitope 76.
	SEQ ID No: 124:	putative human CD8 cell epitope 77.
	SEQ ID No: 125:	putative human CD8 cell epitope 78.
	SEQ ID No: 126:	putative human CD8 cell epitope 79.
	SEQ ID No: 127:	peptide 1.
35	SEQ ID No: 128:	peptide 2.
	SEQ ID No: 129:	peptide 3.
	SEQ ID No: 130:	peptide 4.

	SEQ ID No: 131:	peptide 5.
	SEQ ID No: 132:	peptide 6.
	SEQ ID No: 133:	peptide 7.
	SEQ ID No: 134:	peptide 8.
5	SEQ ID No: 135:	peptide 9.
	SEQ ID No: 136:	peptide 10.
	SEQ ID No: 137:	peptide 11.
	SEQ ID No: 138:	peptide 12.
	SEQ ID No: 139:	peptide 13.
10	SEQ ID No: 140:	peptide 14.
	SEQ ID No: 141:	peptide 15.
	SEQ ID No: 142:	peptide 16.
	SEQ ID No: 143:	peptide 17.
	SEQ ID No: 144:	peptide 18.
15	SEQ ID No: 145:	peptide 19.
	SEQ ID No: 146:	peptide 20.
	SEQ ID No: 147:	peptide 21.
	SEQ ID No: 148:	peptide 22.
	SEQ ID No: 149:	peptide 23.
20	SEQ ID No: 150:	peptide 24.
	SEQ ID No: 151:	peptide 25.
	SEQ ID No: 152:	peptide 26.
	SEQ ID No: 153:	peptide 27.
	SEQ ID No: 154:	peptide 28.
25	SEQ ID No: 155:	peptide 29.
	SEQ ID No: 156:	peptide 30.
	SEQ ID No: 157:	polypeptide sequence of Rv1753c from <i>M. tuberculosis</i> H37Rv strain.
	SEQ ID No: 158:	polypeptide sequence of Rv2386c from <i>M. tuberculosis</i> H37Rv strain.
	SEQ ID No: 159:	polypeptide sequence of Rv2707c from <i>M. tuberculosis</i> H37Rv strain.
30		

DETAILED DESCRIPTION

Currently, vaccination with live bacteria is the most efficient method for inducing protective immunity. The most common *Mycobacterium* employed for this purpose is *Bacillus Calmette-Guerin* (BCG), an avirulent strain of *M. bovis* which was developed over 60 years ago. However, the safety and efficacy of BCG is a source of controversy - while protecting against severe disease manifestation in children, BCG does not prevent the establishment of latent TB

or reactivation of pulmonary disease in adult life. Additionally, some countries, such as the United States, do not vaccinate the general public with this agent.

Almost all new generation TB vaccines which are currently in clinical development have been designed as pre-exposure vaccines. These include subunit vaccines, which have been particularly effective in boosting immunity induced by prior BCG vaccination, and advanced live mycobacterial vaccines which aim to replace BCG with more efficient and/or safer strains. Although these vaccines aim to improve resistance to infection, they are likely to be less effective as post-exposure or therapeutic vaccines in latent TB cases (Lin MY et al *Endocrine, Metabolic & Immune Disorders – Drug Targets* 2008 8:15-29).

Several of the proteins which are strongly expressed during the early stages of *Mycobacterium* infection have been shown to provide strong protective efficacy in animal vaccination models. However, vaccination with antigens which are highly expressed during the early stages of infection may not provide an optimal immune response for dealing with later stages of infection. Adequate control during the later stages of infection may require T cells which are specific for the particular antigens which are expressed at that time.

Post-exposure vaccines which directly target the dormant persistent bacteria may aid in protecting against TB reactivation, thereby enhancing TB control, or even enabling clearance of the infection. A vaccine targeting latent TB could therefore significantly and economically reduce global TB infection rates.

Subunit vaccines based on late stage antigens could also be utilised in combination with early stage antigens to provide a multiphase vaccine. Alternatively, late stage antigens could be used to complement and improve BCG vaccination (either by boosting the BCG response or through the development of advanced recombinant BCG strains).

Rv3616c, also known as Mtb40 or HTCC1, has previously been implicated in the immune responses associated with tuberculosis (see, for example, WO98/53075). Al-Attayah et al. *Clin. Exp. Immunol.* 2004 138:139-144 have shown that Rv3616c is well recognised (through PMBC proliferation and IFN-gamma production) by pulmonary tuberculosis patients. Mustafa et al. *Infect. Immun.* 2006 74(8):4566-4572 have investigated the recognition of Rv3616c by *M. bovis* infected and BCG vaccinated cattle.

Recently, a range of *M. tuberculosis* vaccine candidates have been proposed based on a bioinformatics analysis of the whole genome *M. tuberculosis* genome (Zvi et al. *BMC Medical*

Genetics 2008 1:18) and on the testing of differentially expressed proteins in actively and latently infected individuals (Schuck SD et al. *PLoS ONE* 2009 4(5):e5590).

While macrophages have been shown to act as the principal effectors of *Mycobacterium*

immunity, T cells are the predominant inducers of such immunity. The essential role of T cells in protection against tuberculosis is illustrated by the increased rates of TB reactivation in human immunodeficiency virus infected individuals, due to the associated depletion of CD4+ T cells. Furthermore, adoptive transfer of CD4+ T cells taken at the height of the primary immune response to *M. tuberculosis* has been shown to confer protection against *M. tuberculosis* in T cell deficient mice (Orme et al *J. Exp. Med.* 1983 158:74-83).

Mycobacterium-reactive CD4+ T cells have been shown to be potent producers of γ -interferon (IFN- γ), which, in turn, has been shown to trigger the anti-mycobacterial effects of macrophages in mice (Flynn et al. *J. Exp. Med.* 1993 178:2249-2254). While the role of IFN- γ in humans is less clear, studies have shown that 1,25-dihydroxy-vitamin D3, either alone or in combination with IFN- γ or tumor necrosis factor- α , activates human macrophages to inhibit *M. tuberculosis* infection. Furthermore, it is known that IFN- γ stimulates human macrophages to make 1,25-dihydroxy-vitamin D3. Similarly, interleukin-12 (IL-12) has been shown to play a role in stimulating resistance to *M. tuberculosis* infection. For a review of the immunology of *M. tuberculosis* infection, see Chan & Kaufmann, *Tuberculosis: Pathogenesis, Protection and Control* (Bloom ed., 1994), *Tuberculosis* (2nd ed., Rom and Garay, eds., 2003), and *Harrison's Principles of Internal Medicine*, Chapter 150, pp. 953-966 (16th ed., Braunwald, et al., eds., 2005).

The present invention relates generally to the identification of Rv3616c as a TB antigen which is associated with latent TB and to related methods and uses in the prevention and treatment of latent TB, and the prevention or delay of TB reactivation.

The invention therefore provides an Rv3616c protein, variant thereof or immunogenic fragment thereof, or a polynucleotide encoding said protein, variant or fragment, for use in the treatment or prevention of latent TB. Suitably, the use may be specifically in the treatment of latent TB. Alternatively, the use may be in the prevention or delay of TB reactivation (especially the delay of TB reactivation, for example by a period of months, years or even indefinitely).

The term "*Mycobacterium* species of the tuberculosis complex" includes those species traditionally considered as causing the disease tuberculosis, as well as *Mycobacterium* environmental and opportunistic species that cause tuberculosis and lung disease in immune

compromised patients, such as patients with AIDS, e.g., *M. tuberculosis*, *M. bovis*, or *M. africanum*, BCG, *M. avium*, *M. intracellulare*, *M. celatum*, *M. genavense*, *M. haemophilum*, *M. kansasii*, *M. simiae*, *M. vaccae*, *M. fortuitum*, and *M. scrofulaceum* (see, e.g., *Harrison's Principles of Internal Medicine*, Chapter 150, pp. 953-966 (16th ed., Braunwald, et al., eds., 2005)). The present invention is particularly directed to infection with *M. tuberculosis*.

The term "active infection" refers to an infection (e.g. infection by *M. tuberculosis*) with manifested disease symptoms and/or lesions (suitably with manifested disease symptoms).

The terms "inactive infection", "dormant infection" or "latent infection" refer to an infection (e.g. infection by *M. tuberculosis*) without manifested disease symptoms and/or lesions (suitably without manifested disease symptoms).

The term "primary tuberculosis" refers to clinical illness (manifestation of disease symptoms) directly following infection (e.g. infection by *M. tuberculosis*). See, *Harrison's Principles of Internal Medicine*, Chapter 150, pp. 953-966 (16th ed., Braunwald, et al., eds., 2005).

The terms "secondary tuberculosis" or "postprimary tuberculosis" refer to the reactivation of a dormant, inactive or latent infection (e.g. infection by *M. tuberculosis*). See, *Harrison's Principles of Internal Medicine*, Chapter 150, pp. 953-966 (16th ed., Braunwald, et al., eds., 2005).

The term "tuberculosis reactivation" refers to the later manifestation of disease symptoms in an individual that tests positive for infection (e.g. in a tuberculin skin test, suitably in an *in vitro* T cell based assay) test but does not have apparent disease symptoms. The positive diagnostic test indicates that the individual is infected, however, the individual may or may not have previously manifested active disease symptoms that had been treated sufficiently to bring the tuberculosis into an inactive or latent state. It will be recognised that methods for the prevention, delay or treatment of tuberculosis reactivation can be initiated in an individual manifesting active symptoms of disease.

The term "drug resistant" tuberculosis refers to an infection (e.g. infection by *M. tuberculosis*) wherein the infecting strain is not held static or killed (i.e. is resistant to) one or more of so-called "front-line" chemotherapeutic agents effective in treating tuberculosis (e.g., isoniazid, rifampin, ethambutol, streptomycin and pyrazinamide).

The term "multi-drug resistant" tuberculosis refers to an infection (e.g. infection by *M. tuberculosis*) wherein the infecting strain is resistant to two or more of "front-line" chemotherapeutic agents effective in treating tuberculosis.

- 5 A "chemotherapeutic agent" refers to a pharmacological agent known and used in the art to treat tuberculosis (e.g. infection by *M. tuberculosis*). Exemplified pharmacological agents used to treat tuberculosis include, but are not limited to amikacin, aminosalicylic acid, capreomycin, cycloserine, ethambutol, ethionamide, isoniazid, kanamycin, pyrazinamide, rifamycins (*i.e.*, rifampin, rifapentine and rifabutin), streptomycin, ofloxacin, ciprofloxacin, clarithromycin, azithromycin and fluoroquinolones. "First-line" or "Front-line" chemotherapeutic agents used to treat tuberculosis that is not drug resistant include isoniazid, rifampin, ethambutol, streptomycin and pyrazinamide. "Second-line" chemotherapeutic agents used to treat tuberculosis that has demonstrated drug resistance to one or more "first-line" drugs include ofloxacin, ciprofloxacin, ethionamide, aminosalicylic acid, cycloserine, amikacin, kanamycin and capreomycin. Such pharmacological agents are reviewed in Chapter 48 of *Goodman and Gilman's The Pharmacological Basis of Therapeutics*, Hardman and Limbird eds., 2001.

- 20 The terms "polypeptide", "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms also apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer. Suitably a polypeptide according to the present invention will consist only of naturally occurring amino acid residues, especially those amino acids encoded by the genetic code.

- 25 The term "amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. Amino acid analogues refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, *i.e.*, an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, e.g., homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (e.g., norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid.

Suitably an amino acid is a naturally occurring amino acid or an amino acid analogue, especially a naturally occurring amino acid and in particular those amino acids encoded by the genetic code.

- 5 "Nucleic acid" refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. The term encompasses nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, which have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides.
- 10 Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs). Suitably the term "nucleic acid" refers to naturally occurring deoxyribonucleotides or ribonucleotides and polymers thereof.
- 15 Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated (suitably it refers to the sequence explicitly indicated). Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is
- 20 substituted with mixed-base and/or deoxyinosine residues (Batzer *et al.*, *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka *et al.*, *J. Biol. Chem.* 260:2605-2608 (1985); Rossolini *et al.*, *Mol. Cell. Probes* 8:91-98 (1994)). The term nucleic acid is used interchangeably with gene, cDNA, mRNA, oligonucleotide, and polynucleotide.
- 25 Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.
- 30 By the term 'Rv3616c protein sequence' as used herein is meant the polypeptide sequence provided in SEQ ID No: 1 or a homologue thereof from a *Mycobacterium* species of the tuberculosis complex, e.g., a species such as *M. tuberculosis*, *M. bovis*, or *M. africanum*, or a *Mycobacterium* species that is environmental or opportunistic and that causes opportunistic infections such as lung infections in immune compromised hosts (e.g., patients with AIDS),
- 35 e.g., *BCG*, *M. avium*, *M. intracellulare*, *M. celatum*, *M. genavense*, *M. haemophilum*, *M. kansasii*, *M. simiae*, *M. vaccae*, *M. fortuitum*, and *M. scrofulaceum* (see, e.g., Harrison's

Principles of Internal Medicine, Chapter 150, pp. 953-966, 16th ed., Braunwald, *et al.*, eds., 2005).

To ensure a high efficacy rate among vaccinated hosts, the components of a vaccine should be well conserved among the strains of clinical significance. Suitably, the Rv3616c protein is derived from *M. tuberculosis* H37Rv (i.e. the polypeptide sequence provided in SEQ ID No: 1) or a homologue thereof from another *M. tuberculosis* strain (such as CDC1551, F11, Haarlem A and C strains). Strains of *M. tuberculosis* which are associated with drug resistance (e.g. MDR or especially XDR) are a particularly valuable basis for the Rv3616c protein sequence.

Strains of interest include:

CDC1551 - transmissible and virulent strain

Haarlem family (such as Haarlem A) - Drug resistant strains found in crowded human populations. Members of the Haarlem family of *M. tuberculosis* strains have been found in many parts of the world. The first representative of the family was discovered in Haarlem, The Netherlands.

KZN4207 - Drug sensitive isolate from patients in KwaZulu-Natal, South Africa

KZN1435 - Multiple drug resistant (MDR) isolate from patients in KwaZulu-Natal, South Africa

KZN605 - Extensively drug resistant (XDR) isolate from patients in KwaZulu-Natal, South Africa

C - Highly transmitted in New York City. In one study this strain was found to be more common among injection drug users and resistant to reactive nitrogen intermediates (Friedman *et al.* *J. Infect. Dis.* 1997 176(2):478-84)

94_M4241A - Isolated in San Francisco in 1994 from a patient born in China. This strain was previously analysed by genomic deletion analysis (Gagneux *et al.*, *PNAS* 2006 103(8):2869-2873).

02_1987 - Isolated in San Francisco in 2002 from a patient born in South Korea. This strain was previously analyzed by genomic deletion analysis (Gagneux *et al.*, *PNAS* 2006 103(8):2869-2873).

T92 - Isolated in San Francisco in 1999 from a patient born in The Philippines. This strain was published in Hirsh et al. *PNAS* 2004 101:4871–4876).

5 T85 - Isolated in San Francisco in 1998 from a patient born in China. This strain was published in Hirsh et al. *PNAS* 2004 101:4871–4876).

EAS054 - Isolated in San Francisco in 1993 from a patient born in India. This strain was previously analyzed by genomic deletion analysis (Gagneux et al., *PNAS* 2006 103(8):2869-2873).

Gagneux et al., *PNAS* 2006 103(8):2869-2873 and Herbert et al. *Infect. Immun.* 2007 75(12):5798-5805 provide valuable background on the range of *M. tuberculosis* strains which are known to exist.

15 Most suitably, the Rv3616c protein is selected from the polypeptide sequences provided in SEQ ID Nos: 1 and 3-7, in particular SEQ ID Nos: 1 and 3-6, such as SEQ ID No: 1.

20 Polynucleotides of particular interest are those comprising (such as consisting of) a sequence encoding:

- (i) an Rv3616c protein sequence;
- (ii) a variant of an Rv3616c protein sequence; or
- (iii) an immunogenic fragment of an Rv3616c protein sequence.

25 Polynucleotides will suitably comprise (such as consisting of) a variant of SEQ ID NO: 2 or fragment of SEQ ID NO: 2 which encodes an immunogenic fragment of an Rv3616c protein.

COMBINATIONS

30 The Rv3616c related polypeptides of use in the present invention can further comprise other components designed to enhance their immunogenicity or to improve these antigens in other respects. For example, improved isolation of the polypeptide antigens may be facilitated through the addition of a stretch of histidine residues (commonly known as a his-tag) towards one end of the antigen.

35 The term "his-tag" refers to a string of histidine residues, typically six residues, that are inserted within the reference sequence. To minimise disruption of the activity associated with the reference sequence, a his-tag is typically inserted at the N-terminus, usually immediately

after the initiating methionine residue, or else at the C-terminus. They are usually heterologous to the native sequence but are incorporated since they facilitate isolation by improving the protein binding to immobilised metal affinity chromatography resins (IMAC). Generally speaking the presence or absence of a his-tag is not of significance from the point of view of eliciting a desirable immune response against the reference protein. However, to avoid the risk of an adverse reaction against the his-tag itself, it is considered best to minimise the length of the his-tag e.g. to four or fewer residues, in particular two residues (or to exclude the use of a his-tag entirely).

- 5
- 10 To improve the magnitude and/or breadth of the elicited immune response the compositions, polypeptides and nucleic acids can comprise multiple copies of the inventive antigen and/or additional heterologous polypeptides (or polynucleotides encoding them) from *Mycobacterium* species (in particular *M. tuberculosis*).
- 15 One skilled in the art will recognise that when a number of components are utilised in combination, the precise presentation can be varied. For example, a Rv3616c component and an additional copy of the antigen or an additional heterologous antigen component could be presented:
 - 20 (1) as two individual polypeptide components;
 - (2) as a fusion protein comprising both polypeptide components;
 - (3) as one polypeptide and one polynucleotide component;
 - (4) as two individual polynucleotide components;
 - (5) as a single polynucleotide encoding two individual polypeptide components; or
 - 25 (6) as a single polynucleotide encoding a fusion protein comprising both polypeptide components.

This flexibility applies equally to situations where three or more components are used in combination. However, for convenience, it is often desirable that when a number of

- 30 components are present they are contained within a single fusion protein or a polynucleotide encoding a single fusion protein. In one embodiment of the invention all antigen components are provided as polypeptides (e.g. within a single fusion protein). In an alternative embodiment of the invention all antigen components are provided as polynucleotides (e.g. a single polynucleotide, such as one encoding a single fusion protein).

35

The term "heterologous" when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not found in the same

relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, e.g., a promoter from one source and a coding region from another source. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature (e.g., a fusion protein).

"Fusion polypeptide" or "fusion protein" refers to a protein having at least two heterologous polypeptides (e.g. at least two *Mycobacterium* sp. polypeptides) covalently linked, either directly or via an amino acid linker. The polypeptides forming the fusion protein are typically linked C-terminus to N-terminus, although they can also be linked C-terminus to C-terminus, N-terminus to N-terminus, or N-terminus to C-terminus. The polypeptides of the fusion protein can be in any order. This term also refers to conservatively modified variants, polymorphic variants, alleles, mutants, immunogenic fragments, and interspecies homologs of the antigens that make up the fusion protein. *Mycobacterium tuberculosis* antigens are described in Cole *et al.*, *Nature* 393:537 (1998), which discloses the entire *Mycobacterium tuberculosis* genome. Antigens from other *Mycobacterium* species that correspond to *M. tuberculosis* antigens can be identified, e.g., using sequence comparison algorithms, as described herein, or other methods known to those of skill in the art, e.g., hybridisation assays and antibody binding assays.

The term "fused" refers to the covalent linkage between two polypeptides in a fusion protein. The polypeptides are typically joined via a peptide bond, either directly to each other or via an amino acid linker. Optionally, the peptides can be joined via non-peptide covalent linkages known to those of skill in the art.

Exemplary *M. tuberculosis* antigens which may be combined with Rv3616c include one or more of (e.g. 1 to 5, such as 1 to 3, in particular 1) the following (such as one or more of (i) to (xi)):

- (i) Mtb8.4 (also known as DPV and Rv1174c), the polypeptide sequence of which is described in SEQ ID No: 102 of WO97/09428 (cDNA in SEQ ID No: 101) and in Coler *et al Journal of Immunology* 1998 161:2356-2364. Of particular interest is the mature Mtb8.4 sequence which is absent the leading signal peptide (i.e. amino acid residues 15-96 from SEQ ID No: 102 of WO97/09428). The full-length polypeptide sequence of Mtb8.4 is shown in SEQ ID No: 8;

(ii) Mtb9.8 (also known as MSL and Rv0287), the polypeptide sequence of which is described in SEQ ID No: 109 of WO98/53075 (fragments of MSL are disclosed in SEQ ID Nos: 110-124 of WO98/53075, SEQ ID Nos: 119 and 120 being of particular interest) and also in Coler et al *Vaccine* 2009 27:223-233 (in particular the reactive fragments shown in Figure 2 therein). The full-length polypeptide sequence for Mtb9.8 is shown in SEQ ID No: 9;

(iii) Mtb9.9 (also known as Mtb9.9A, MTI, MTI-A and Rv1793) the polypeptide sequence of which is described in SEQ ID No: 19 of WO98/53075 and in Alderson et al *Journal of Experimental Medicine* 2000 7:551-559 (fragments of MTI are disclosed in SEQ ID Nos: 17 and 51-66 of WO98/53075, SEQ ID Nos: 17, 51, 52, 53, 56 and 62-65 being of particular interest). A number of polypeptide variants of MTI are described in SEQ ID Nos: 21, 23, 25, 27, 29 and 31 of WO98/53075 and in Alderson et al *Journal of Experimental Medicine* 2000 7:551-559. The full-length polypeptide sequence for Mtb9.9 is shown in SEQ ID No: 10;

(iv) Ra12 (also known as Mtb32A C-terminal antigen) the polypeptide sequence of which is described in SEQ ID No: 10 of WO01/98460 and in Skeiky et al *Journal of Immunology* 2004 172:7618-7682. The full-length polypeptide sequence for Ra12 is shown in SEQ ID No: 11;

(v) Ra35 (also known as Mtb32A N-terminal antigen) the polypeptide sequence of which is described in SEQ ID No: 8 of WO01/98460 and in Skeiky et al *Journal of Immunology* 2004 172:7618-7682. The full-length polypeptide sequence for Ra35 is shown in SEQ ID No: 12;

(vi) TbH9 (also known as Mtb39, Mtb39A, TbH9FL and Rv1196) the polypeptide sequence of which is described in SEQ ID No: 107 of WO97/09428, and also in Dillon et al *Infection and Immunity* 1999 67(6):2941-2950 and Skeiky et al *Journal of Immunology* 2004 172:7618-7682. The full-length polypeptide sequence for TbH9 is shown in SEQ ID No: 13;

(vii) Mtb41 (also known as MTCC2 and Rv0915c) the polypeptide sequence of which is described in SEQ ID No: 142 of WO98/53075 (cDNA in SEQ ID No: 140) and in Skeiky et al *Journal of Immunology* 2000 165:7140-7149. The full-length polypeptide sequence for Mtb41 is shown in SEQ ID No: 14;

(viii) ESAT-6 (also known as esxA and Rv3875) the polypeptide sequence of which is described in SEQ ID No: 103 of WO97/09428 (cDNA in SEQ ID No: 104) and in Sorensen et al *Infection and Immunity* 1995 63(5):1710-1717. The full-length polypeptide sequence for ESAT-6 is shown in SEQ ID No: 15;

5

(ix) Ag85 complex antigens (e.g. Ag85A, also known as fbpA and Rv3804c; or Ag85B, also known as fbpB and Rv1886c) which are discussed, for example, in Content et al *Infection and Immunity* 1991 59:3205-3212 and in Huygen et al *Nature Medicine* 1996 2(8):893-898. The full-length polypeptide sequence for Ag85A is shown in SEQ ID No: 16 (the mature protein of residues 43-338, i.e. lacking the signal peptide, being of particular interest). The full-length polypeptide sequence for Ag85B is shown in SEQ ID No: 17 (the mature protein of residues 41-325, i.e. lacking the signal peptide, being of particular interest);

10

15

(x) Alpha-crystallin (also known as hspX and Rv2031c) which is described in Verbon et al *Journal of Bacteriology* 1992 174:1352-1359 and Friscia et al *Clinical and Experimental Immunology* 1995 102:53-57 (of particular interest are the fragments corresponding to residues 71-91, 21-40, 91-110 and 111-130)). The full-length polypeptide sequence for alpha-crystallin is shown in SEQ ID No: 18;

20

(xi) Mpt64 (also known as Rv1980c) which is described in Roche et al *Scandinavian Journal of Immunology* 1996 43:662-670. The full-length polypeptide sequence for MPT64 is shown in SEQ ID No: 19 (the mature protein of residues 24-228, i.e. lacking the signal peptide, being of particular interest);

25

(xii) Mtb32A, the polypeptide sequence of which is described in SEQ ID No: 2 (full-length) and residues 8-330 of SEQ ID No: 4 (mature) of WO01/98460, especially variants having at least one of the catalytic triad mutated (e.g. the catalytic serine residue, which may for example be mutated to alanine). The full-length polypeptide sequence for Mtb32A is shown in SEQ ID No: 20. The mature form of Mtb32A having a Ser/Ala mutation is shown in SEQ ID No: 21;

30

(xiii) TB10.4, the full-length polypeptide sequence for TB10.4 is shown in SEQ ID No: 22;

35

(xiv) Rv1753c, the full-length polypeptide sequence for Rv1753c from *Mycobacterium tuberculosis* H37Rv is shown in SEQ ID No: 157;

(xv) Rv2386c, the full-length polypeptide sequence for Rv2386c from *Mycobacterium tuberculosis* H37Rv is shown in SEQ ID No: 158; and/or

5 (xvi) Rv2707c, the full-length polypeptide sequence for Rv2707c from *Mycobacterium tuberculosis* H37Rv is shown in SEQ ID No: 159.

or combinations thereof, such as (for example combinations such as (a) to (g)):

10 (a) a combination of Ra12, TbH9 and Ra35 components, for example in the form of a fusion protein, such as Mtb72f. The polypeptide sequence of Mtb72f is described in SEQ ID No: 6 of WO2006/117240 (cDNA in SEQ ID No: 5) and in Skeiky et al *Journal of Immunology* 2004 172:7618-7682 (where it incorporates an optional His-tag to aid purification, when utilised in the present invention suitably Mtb72f is absent the optional
15 histidine residues). The polypeptide sequence for Mtb72f is shown in SEQ ID No: 23;

(b) a combination of Ra12, TbH9 and Ser/Ala mutated Ra35 (i.e. where the catalytic serine residue has been replaced with alanine) components, for example in the form of a fusion protein, such as M72. The polypeptide sequence of M72 is described in SEQ
20 ID No: 4 of WO2006/117240 (cDNA in SEQ ID No: 3) where it incorporates an optional double histidine to aid manufacture, when utilised in the present invention M72 may also incorporate a double histidine though suitably M72 is absent the optional double histidine (i.e. residues 4-725 from SEQ ID No: 4 of WO2006/117240 are of particular interest). The polypeptide sequence for M72 is shown in SEQ ID No: 24;

25 (c) a combination of Mtb8.4, Mtb9.8, Mtb9.9 and Mtb41 components, for example in the form of a fusion protein, such as Mtb71f. The polypeptide sequence of Mtb71f is described in SEQ ID No: 16 of WO99/051748 (cDNA in SEQ ID No: 15), where it incorporates an optional His-tag to aid purification, when utilised in the present
30 invention suitably Mtb71f corresponds to amino acid residues 9-710 of SEQ ID NO: 16 from WO99/051748. The polypeptide sequence for Mtb71f is shown in SEQ ID No: 25;

(d) a combination of Mtb72f or M72 (suitably without optional histidine residues to aid expression) with Mtb9.8 and Mtb9.9, for example in a fusion protein. The polypeptide
35 sequence for an M72-Mtb9.9-Mtb9.8 fusion is shown in SEQ ID No: 26 (M92 fusion), when used in the present invention, the M72-Mtb9.9-Mtb9.8 fusion may optionally

incorporate a double histidine following the initiating methionine residue to aid manufacture;

(e) a combination of Mtb72f or M72 (suitably without optional histidine residues to aid expression) with Ag85B, for example in a fusion protein, such Mtb103f. The polypeptide sequence of Mtb103f is described in SEQ ID No: 18 of WO03/070187 (cDNA in SEQ ID No: 10), where it incorporates an optional His-tag to aid purification, when utilised in the present invention suitably Mtb103f corresponds to amino acid residues 8-1016 of SEQ ID No: 18 from WO03/070187. Also of particular interest is M103, i.e. Mtb103f incorporating a Ser/Ala mutation in the Ra35 component, when utilised in the present invention suitably M103 corresponds to amino acid residues 8-1016 of SEQ ID No: 18 from WO03/070187 wherein the Ser residue at position 710 has been replaced with Ala. The polypeptide sequence for M103 is shown in SEQ ID No: 27, when used in the present invention, the M72-Mtb9.9-Mtb9.8 fusion may optionally incorporate a double histidine following the initiating methionine residue to aid manufacture;

(f) a combination of Mtb72f or M72 (suitably without optional histidine residues to aid expression) with Mtb41, for example in a fusion protein, such Mtb114f. The polypeptide sequence of Mtb114f is described in SEQ ID No: 16 of WO03/070187 (cDNA in SEQ ID No: 9), where it incorporates an optional His-tag to aid purification, when utilised in the present invention suitably Mtb114f corresponds to amino acid residues 8-1154 of SEQ ID No: 16 from WO03/070187. Also of particular interest is M114, i.e. Mtb114f incorporating a Ser/Ala mutation in the Ra35 component, when utilised in the present invention suitably M114 corresponds to amino acid residues 8-1154 of SEQ ID No: 16 from WO03/070187 wherein the Ser residue at position 710 has been replaced with Ala. The polypeptide sequence for M114 is shown in SEQ ID No: 28, when used in the present invention, the M72-Mtb9.9-Mtb9.8 fusion may optionally incorporate a double histidine following the initiating methionine residue to aid manufacture;

(g) a combination of Ag85B and ESAT-6 components, such as in a fusion described in Doherty et al *Journal of Infectious Diseases* 2004 190:2146-2153; and/or

(h) a combination of Ag85B and TB10.4 components, such as in a fusion described in Dietrich et al *Journal of Immunology* 2005 174(10):6332-6339 190:2146-2153.

Combinations of an Rv3616c component and an Rv1753c component are of particular interest. Obviously such combinations could optionally contain other additional antigen components (e.g. an M72 component).

- 5 Another combination of interest comprises an Rv3616c component and an M72 component.

A further combination of interest comprises an Rv3616c component and an Rv2386c component.

- 10 Other combinations of interest include those comprising an Rv3616c component and an Rv2707c component.

An additional combination of interest comprises an Rv3616c component and an alpha-crystallin component.

15

The skilled person will recognise that combinations need not rely upon the specific sequences described in above in (i)-(xvi) and (a)-(h), and that conservatively modified variants (e.g. having at least 70% identity, such as at least 80% identity, in particular at least 90% identity and especially at least 95% identity) or immunogenic fragments (e.g. at least 20% of the full length antigen, such as at least 50% of the antigen, in particular at least 70% and especially at least 80%) of the described sequences can be used to achieve the same practical effect.

20

Each of the above individual antigen sequences is also disclosed in Cole et al *Nature* 1998 393:537-544 and Camus *Microbiology* 2002 148:2967-2973. The genome of *M. tuberculosis* H37Rv is publicly available, for example at the Wellcome Trust Sanger Institute website (www.sanger.ac.uk/Projects/M_tuberculosis/) and elsewhere.

25

Many of the above antigens are also disclosed in U.S. patent application numbers 08/523,435, 08/523,436, 08/658,800, 08/659,683, 08/818,111, 08/818,112, 08/942,341, 08/942,578, 08/858,998, 08/859,381, 09/056,556, 09/072,596, 09/072,967, 09/073,009, 09/073,010, 09/223,040, 09/287,849 and in PCT patent applications PCT/US98/10407, PCT/US98/10514, PCT/US99/03265, PCT/US99/03268, PCT/US99/07717, WO97/09428 and WO97/09429, WO98/16645, WO98/16646, each of which is herein incorporated by reference.

30

- 35 The compositions, polypeptides, and nucleic acids of the invention can also comprise additional polypeptides from other sources. For example, the compositions and fusion proteins of the invention can include polypeptides or nucleic acids encoding polypeptides, wherein the

polypeptide enhances expression of the antigen, e.g., NS1, an influenza virus protein (see, e.g. WO99/40188 and WO93/04175). The nucleic acids of the invention can be engineered based on codon preference in a species of choice, e.g., humans (in the case of *in vivo* expression) or a particular bacterium (in the case of polypeptide production).

5

The Rv3616c component may also be administered with one or more chemotherapeutic agents effective against tuberculosis (e.g. *M. tuberculosis* infection). Examples of such chemotherapeutic agents include, but are not limited to, amikacin, aminosalicylic acid, capreomycin, cycloserine, ethambutol, ethionamide, isoniazid, kanamycin, pyrazinamide, rifamycins (*i.e.*, rifampin, rifapentine and rifabutin), streptomycin, ofloxacin, ciprofloxacin, clarithromycin, azithromycin and fluoroquinolones. Such chemotherapy is determined by the judgment of the treating physician using preferred drug combinations. "First-line" chemotherapeutic agents used to treat tuberculosis (e.g. *M. tuberculosis* infection) that is not drug resistant include isoniazid, rifampin, ethambutol, streptomycin and pyrazinamide.

10

"Second-line" chemotherapeutic agents used to treat tuberculosis (e.g. *M. tuberculosis* infection) that has demonstrated drug resistance to one or more "first-line" drugs include ofloxacin, ciprofloxacin, ethionamide, aminosalicylic acid, cycloserine, amikacin, kanamycin and capreomycin.

15

Conventional chemotherapeutic agents are generally administered over a relatively long period (ca. 9 months). Combination of conventional chemotherapeutic agents with the administration of a Rv3616c component according to the present invention may enable the chemotherapeutic treatment period to be reduced (e.g. to 8 months, 7 months, 6 months, 5 months, 4 months, 3 months or less) without a decrease in efficacy.

20

25

Of particular interest is the use of an Rv3616c component in conjunction with *Bacillus Calmette-Guerin* (BCG). For example, in the form of a modified BCG which recombinantly expresses Rv3616c (or a variant or fragment thereof as described herein). Alternatively, the Rv3616c component may be used to enhance the response of a subject to BCG vaccination, either by co-administration or by boosting a previous BCG vaccination. When used to enhance the response of a subject to BCG vaccination, the Rv3616c component may obviously be provided in the form of a polypeptide or a polynucleotide (optionally in conjunction with additional antigenic components as described above).

30

The skilled person will recognise that combinations of components need not be administered together and may be applied: separately or in combination; at the same time, sequentially or within a short period; though the same or through different routes. Nevertheless, for

35

convenience it is generally desirable (where administration regimes are compatible) to administer a combination of components as a single composition.

The polypeptides, polynucleotides and compositions of the present invention will usually be administered to humans, but are effective in other mammals including domestic mammals (e.g., dogs, cats, rabbits, rats, mice, guinea pigs, hamsters, chinchillas) and agricultural mammals (e.g., cows, pigs, sheep, goats, horses).

IMMUNOGENIC FRAGMENTS

T cell epitopes are short contiguous stretches of amino acids which are recognised by T cells (e.g. CD4+ or CD8+ T cells). Identification of T cell epitopes may be achieved through epitope mapping experiments which are well known to the person skilled in the art (see, for example, Paul, *Fundamental Immunology*, 3rd ed., 243-247 (1993); Beißbarth et al *Bioinformatics* 2005 21(Suppl. 1):i29-i37).

Alternatively, epitopes may be predicted using the approaches discussed in the Examples.

As a result of the crucial involvement of the T cell response in tuberculosis, it is readily apparent that fragments of the full length Rv3616c polypeptide which contain at least one T cell epitope will be immunogenic and may contribute to immunoprotection. Such fragments are referred to herein as Immunogenic fragments.

Immunogenic fragments according to the present invention will typically comprise at least 9 contiguous amino acids from the full length polypeptide sequence (e.g. at least 10), such as at least 12 contiguous amino acids (e.g. at least 15 or at least 20 contiguous amino acids), in particular at least 50 contiguous amino acids, such as at least 100 contiguous amino acids (for example at least 200 contiguous amino acids). Suitably the immunogenic fragments will be at least 20%, such as at least 50%, at least 70% or at least 80% of the length of the full length polypeptide sequence.

It will be understood that in a diverse out-bred population, such as humans, different HLA types mean that specific epitopes may not be recognised by all members of the population. Consequently, to maximise the level of recognition and scale of immune response to a polypeptide, it is generally desirable that an immunogenic fragment contains a plurality of the epitopes from the full length sequence (suitably all epitopes).

Particular fragments of the Rv3616c protein which may be of use include those containing at least one CD4+ epitope, suitably at least two CD4+ epitopes and especially all CD4+ epitopes (such as those epitopes described in the Examples and in SEQ ID Nos: 29-47, particularly those associated with a plurality of HLA alleles, e.g. those associated with 2, 3, 4, 5 or more alleles).

Other fragments of the Rv3616c protein which may be of use include those containing at least one CD8 epitope, suitably at least two CD8 epitopes and especially all CD8 epitopes (such as those epitopes described in the Examples and in SEQ ID Nos: 48-126, particularly those associated with a plurality of HLA alleles, e.g. those associated with 2, 3, 4, 5 or more alleles).

Where an individual fragment of the full length polypeptide is used, such a fragment is considered to be immunogenic where it elicits a response which is at least 20%, suitably at least 50% and especially at least 75% (such as at least 90%) of the activity of the reference sequence in an *in vitro* restimulation assay of PBMC or whole blood with specific antigens (e.g. restimulation for a period of between several hours to up to two weeks, such as up to one day, 1 day to 1 week or 1 to 2 weeks) that measures the activation of the cells via lymphoproliferation, production of cytokines in the supernatant of culture (measured by ELISA, CBA etc) or characterisation of T and B cell responses by intra and extracellular staining (e.g. using antibodies specific to immune markers, such as CD3, CD4, CD8, IL2, TNFa, IFNg, CD40L, CD69 etc) followed by analysis with a flowcytometer. Suitably, a fragment is considered to be immunogenic where it elicits a response which is at least 20%, suitably at least 50% and especially at least 75% (such as at least 90%) of the activity of the reference sequence in a T cell proliferation and/or IFN-gamma production assay.

In some circumstances a plurality of fragments of the full length polypeptide (which may or may not be overlapping and may or may not cover the entirety of the full length sequence) may be used to obtain an equivalent biological response to the full length sequence itself. For example, at least two immunogenic fragments (such as three, four or five) as described above, which in combination provide at least 50%, suitably at least 75% and especially at least 90% activity of the reference sequence in an *in vitro* restimulation assay of PBMC or whole blood (e.g. a T cell proliferation and/or IFN-gamma production assay).

Peptides of SEQ ID Nos: 127-156 are fragments of particular interest (especially those in SEQ ID Nos: 127-133 and 143-156).

VARIANTS

"Variants" or "conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified

- 5 variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences.

- Due to the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all
10 encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations lead to "silent" or "degenerate" variants, which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the
15 nucleic acid. One of skill will recognise that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid that encodes a polypeptide is implicit in each described sequence.

- A polynucleotide of the invention may contain a number of silent variations (for example, 1-50,
20 such as 1-25, in particular 1-5, and especially 1 codon(s) may be altered) when compared to the reference sequence. A polynucleotide of the invention may contain a number of non-silent conservative variations (for example, 1-50, such as 1-25, in particular 1-5, and especially 1 codon(s) may be altered) when compared to the reference sequence. Non-silent variations are those which result in a change in the encoded amino acid sequence (either through the
25 substitution, deletion or addition of amino acid residues). Those skilled in the art will recognise that a particular polynucleotide sequence may contain both silent and non-silent conservative variations.

- In respect of variants of a protein sequence, the skilled person will recognise that individual substitutions, deletions or additions to polypeptide, which alters, adds or deletes a single
30 amino acid or a small percentage of amino acids is a "conservatively modified variant" where the alteration(s) results in the substitution of an amino acid with a functionally similar amino acid or the substitution/deletion/addition of residues which do not substantially impact the biological function of the variant.

Conservative substitution tables providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles of the invention.

- 5 A polypeptide of the invention may contain a number of conservative substitutions (for example, 1-50, such as 1-25, in particular 1-10, and especially 1 amino acid residue(s) may be altered) when compared to the reference sequence. In general, such conservative substitutions will fall within one of the amino-acid groupings specified below, though in some circumstances other substitutions may be possible without substantially affecting the
- 10 immunogenic properties of the antigen. The following eight groups each contain amino acids that are typically conservative substitutions for one another:

- 1) Alanine (A), Glycine (G);
- 2) Aspartic acid (D), Glutamic acid (E);
- 15 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V);
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W);
- 7) Serine (S), Threonine (T); and
- 20 8) Cysteine (C), Methionine (M)
- (see, e.g., Creighton, *Proteins* 1984).

Suitably such substitutions do not occur in the region of an epitope, and do not therefore have a significant impact on the immunogenic properties of the antigen.

- 25 Protein variants may also include those wherein additional amino acids are inserted compared to the reference sequence, for example, such insertions may occur at 1-10 locations (such as 1-5 locations, suitably 1 or 2 locations, in particular 1 location) and may, for example, involve the addition of 50 or fewer amino acids at each location (such as 20 or fewer, in particular 10
- 30 or fewer, especially 5 or fewer). Suitably such insertions do not occur in the region of an epitope, and do not therefore have a significant impact on the immunogenic properties of the antigen. One example of insertions includes a short stretch of histidine residues (e.g. 2-6 residues) to aid expression and/or purification of the antigen in question.

- 35 Protein variants include those wherein amino acids have been deleted compared to the reference sequence, for example, such deletions may occur at 1-10 locations (such as 1-5 locations, suitably 1 or 2 locations, in particular 1 location) and may, for example, involve the

deletion of 50 or fewer amino acids at each location (such as 20 or fewer, in particular 10 or fewer, especially 5 or fewer). Suitably such deletions do not occur in the region of an epitope, and do not therefore have a significant impact on the immunogenic properties of the antigen.

- 5 The skilled person will recognise that a particular protein variant may comprise substitutions, deletions and additions (or any combination thereof).

Methods of determining the epitope regions of an antigen are described and exemplified in the Examples.

10

Variants preferably exhibit at least about 70% identity, more preferably at least about 80% identity and most preferably at least about 90% identity (such as at least about 95%, at least about 98% or at least about 99%) to the associated reference sequence.

- 15 The terms "identical" or percent "identity," in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or sub-sequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (i.e., 70% identity, optionally 75%, 80%, 85%, 90%, 95%, 98% or 99% identity over a specified region), when compared and aligned for maximum correspondence over a comparison
20 window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Such sequences are then said to be "substantially identical." This definition also refers to the complement of a test sequence. Optionally, the identity exists over a region that is at least about 25 to about 50 amino acids or nucleotides in length, or optionally over a region that is 75-100 amino acids or nucleotides in
25 length. Suitably, the comparison is performed over a window corresponding to the entire length of the reference sequence.

- For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference
30 sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

35

A "comparison window", as used herein, references to a segment in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two

sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (see, e.g., *Current Protocols in Molecular Biology* (Ausubel et al., eds. 1995 supplement)).

One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J. Mol. Evol.* 35:351-360 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151-153 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. Using PILEUP, a reference sequence is compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps. PILEUP can be obtained from the GCG sequence analysis software package, e.g., version 7.0 (Devereaux et al., *Nuc. Acids Res.* 12:387-395 (1984)).

Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., *Nuc. Acids Res.* 25:3389-3402 (1997) and Altschul et al., *J. Mol. Biol.* 215:403-410 (1990), respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (website at www.ncbi.nlm.nih.gov/). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued

threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

The present invention also extends to polynucleotides comprising a first nucleotide sequence which selectively hybridises under moderately stringent conditions (such as under highly stringent conditions) to the complement of a second nucleotide sequence which encodes a polypeptide comprising:

- (i) an Rv3616c protein sequence;
- (ii) a variant of an Rv3616c protein sequence; or
- (iii) an immunogenic fragment of an Rv3616c protein sequence,

for the treatment or prevention of latent TB.

The phrase "highly stringent hybridisation conditions" refers to conditions under which a probe will hybridise to its target subsequence, typically in a complex mixture of nucleic acid, but to no other sequences. Highly stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridise specifically at higher temperatures. An

5 extensive guide to the hybridisation of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Probes*, "Overview of principles of hybridisation and the strategy of nucleic acid assays" (1993). Generally, highly stringent conditions are selected to be about 5–10°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength pH. The T_m is the temperature (under defined

10 ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridise to the target sequence at equilibrium (as the target sequences are present in excess, at T_m , 50% of the probes are occupied at equilibrium). Highly stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the

15 temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Highly stringent conditions may also be achieved with the addition of destabilising agents such as formamide. For selective or specific hybridisation, a positive signal is at least two times background, optionally 10 times background hybridisation.

20 Exemplary highly stringent hybridisation conditions can be as following: 50% formamide, 5x SSC, and 1% SDS, incubating at 42°C, or, 5x SSC, 1% SDS, incubating at 65°C, with wash in 0.2x SSC, and 0.1% SDS at 65°C.

25 Nucleic acids that do not hybridise to each other under highly stringent conditions are still functionally equivalent if the polypeptides which they encode are substantially identical. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridise under moderately stringent hybridisation conditions.

30 Exemplary "moderately stringent hybridisation conditions" include a hybridisation in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 1X SSC at 45°C. A positive hybridisation is at least twice background. Those of ordinary skill will readily recognise that alternative hybridisation and wash conditions can be utilised to provide conditions of similar

35 stringency.

The phrase "selectively (or specifically) hybridises to" refers to the binding, duplexing, or hybridising of a molecule only to a particular nucleotide sequence under stringent hybridisation conditions when that sequence is present in a complex mixture (e.g., total cellular or library DNA or RNA).

5

In any event, variants of a polypeptide sequence will have essentially the same activity as the reference sequence (in the case of polynucleotides, variant polynucleotide sequences will encode a polypeptide which has essentially the same activity as the reference sequence). By essentially the same activity is meant at least 50%, suitably at least 75% and especially at least 90% activity of the reference sequence in an *in vitro* restimulation assay of PBMC or whole blood with specific antigens (e.g. restimulation for a period of between several hours to up to two weeks, such as up to one day, 1 day to 1 week or 1 to 2 weeks) that measures the activation of the cells via lymphoproliferation, production of cytokines in the supernatant of culture (measured by ELISA, CBA etc) or characterisation of T and B cell responses by intra and extracellular staining (e.g. using antibodies specific to immune markers, such as CD3, CD4, CD8, IL2, TNFa, IFNg, CD40L, CD69 etc) followed by analysis with a flowcytometer. Suitably, by essentially the same activity is meant at least 50%, suitably at least 75% and especially at least 90% activity of the reference sequence in a T cell proliferation and/or IFN-gamma production assay.

10

15

20

POLYNUCLEOTIDE COMPOSITIONS

As used herein, the term "polynucleotide" refers to a molecule that has been isolated free of total genomic DNA of a particular species. Therefore, a polynucleotide encoding a polypeptide refers to a polynucleotide segment that contains one or more coding sequences yet is substantially isolated away from, or purified free from, total genomic DNA of the species from which the polynucleotide is obtained.

25

30

As will be understood by those skilled in the art, the polynucleotides of use in this invention can include genomic sequences, extra-genomic and plasmid-encoded sequences and smaller engineered gene segments that express, or may be adapted to express, proteins, polypeptides, peptides and the like. Such segments may be naturally isolated, or modified synthetically by the hand of man.

35

"Isolated," as used herein, means that a polynucleotide is substantially away from other coding sequences, and that the polynucleotide does not contain large portions of unrelated coding DNA, such as large chromosomal fragments or other functional genes or polypeptide coding

regions. An isolated nucleic acid is separated from other open reading frames that flank the gene and encode proteins other than the gene. Of course, this refers to the DNA segment as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

5

As will be recognised by the skilled artisan, polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules include HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain
10 introns. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

15

Polynucleotides may comprise a native sequence (i.e., an endogenous sequence that encodes a *Mycobacterium* antigen or a portion thereof) or may comprise a variant, or a biological or functional equivalent of such a sequence. Polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions, as further described below, preferably such that the immunogenicity of the encoded polypeptide is not diminished, relative to the reference protein. The effect on the immunogenicity of the encoded polypeptide may generally
20 be assessed as described herein.

25

In additional embodiments, the present invention provides isolated polynucleotides and polypeptides comprising various lengths of contiguous stretches of sequence identical to or complementary to one or more of the sequences disclosed herein. For example,
polynucleotides are provided by this invention that comprise at least about 30, 40, 50, 75, 100, 150, 200, 300, 400, 500 or 1000 or more contiguous nucleotides of the reference sequence disclosed herein as well as all intermediate lengths there between. It will be readily understood that "intermediate lengths", in this context, means any length between the quoted values, such as 30, 31, 32, etc.; 50, 51, 52, 53, etc.; 100, 101, 102, 103, etc.; 150, 151, 152,
30 153, etc.; including all integers through 200-500; 500-1,000, and the like.

35

Moreover, it will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear relatively low identity to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention, for example polynucleotides that are optimized for human and/or primate codon selection. Further, alleles

of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridisation, amplification and/or database sequence comparison).

POLYNUCLEOTIDE IDENTIFICATION AND CHARACTERISATION

Polynucleotides may be identified, prepared and/or manipulated using any of a variety of well established techniques. For example, a polynucleotide may be identified, as described in more detail below, by screening a microarray of cDNAs. Such screens may be performed, for example, using a Synteni microarray (Palo Alto, CA) according to the manufacturer's instructions (and essentially as described by Schena *et al.*, *Proc. Natl. Acad. Sci. USA* 93:10614-10619 (1996) and Heller *et al.*, *Proc. Natl. Acad. Sci. USA* 94:2150-2155 (1997)). Alternatively, polynucleotides may be amplified from cDNA prepared from cells expressing the proteins described herein, such as *M. tuberculosis* cells. Such polynucleotides may be amplified via polymerase chain reaction (PCR). For this approach, sequence-specific primers may be designed based on the sequences provided herein, and may be purchased or synthesised.

An amplified portion of a polynucleotide may be used to isolate a full length gene from a suitable library (e.g., a *M. tuberculosis* cDNA library) using well known techniques. Within such techniques, a library (cDNA or genomic) is screened using one or more polynucleotide probes or primers suitable for amplification. Preferably, a library is size-selected to include larger molecules. Random primed libraries may also be preferred for identifying 5' and upstream regions of genes. Genomic libraries are preferred for obtaining introns and extending 5' sequences.

For hybridisation techniques, a partial sequence may be labeled (e.g., by nick-translation or end-labeling with ³²P) using well known techniques. A bacterial or bacteriophage library is then generally screened by hybridising filters containing denatured bacterial colonies (or lawns containing phage plaques) with the labeled probe (see Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (2000)). Hybridising colonies or plaques are selected and expanded, and the DNA is isolated for further analysis. cDNA clones may be analyzed to determine the amount of additional sequence by, for example, PCR using a primer from the partial sequence and a primer from the vector. Restriction maps and partial sequences may be generated to

identify one or more overlapping clones. The complete sequence may then be determined using standard techniques, which may involve generating a series of deletion clones. The resulting overlapping sequences can then be assembled into a single contiguous sequence. A full length cDNA molecule can be generated by ligating suitable fragments, using well known techniques.

Alternatively, there are numerous amplification techniques for obtaining a full length coding sequence from a partial cDNA sequence. Within such techniques, amplification is generally performed via PCR. Any of a variety of commercially available kits may be used to perform the amplification step. Primers may be designed using, for example, software well known in the art. Primers are preferably 22-30 nucleotides in length, have a GC content of at least 50% and anneal to the target sequence at temperatures of about 68°C to 72°C. The amplified region may be sequenced as described above, and overlapping sequences assembled into a contiguous sequence.

One such amplification technique is inverse PCR (see Triglia *et al.*, *Nucl. Acids Res.* 16:8186 (1988)), which uses restriction enzymes to generate a fragment in the known region of the gene. The fragment is then circularised by intramolecular ligation and used as a template for PCR with divergent primers derived from the known region. Within an alternative approach, sequences adjacent to a partial sequence may be retrieved by amplification with a primer to a linker sequence and a primer specific to a known region. The amplified sequences are typically subjected to a second round of amplification with the same linker primer and a second primer specific to the known region. A variation on this procedure, which employs two primers that initiate extension in opposite directions from the known sequence, is described in WO 96/38591. Another such technique is known as "rapid amplification of cDNA ends" or RACE. This technique involves the use of an internal primer and an external primer, which hybridises to a polyA region or vector sequence, to identify sequences that are 5' and 3' of a known sequence. Additional techniques include capture PCR (Lagerstrom *et al.*, *PCR Methods Applic.* 1:111-19 (1991)) and walking PCR (Parker *et al.*, *Nucl. Acids. Res.* 19:3055-60 (1991)). Other methods employing amplification may also be employed to obtain a full length cDNA sequence.

In certain instances, it is possible to obtain a full length cDNA sequence by analysis of sequences provided in an expressed sequence tag (EST) database, such as that available from GenBank. Searches for overlapping ESTs may generally be performed using well known programs (e.g., NCBI BLAST searches), and such ESTs may be used to generate a

contiguous full length sequence. Full length DNA sequences may also be obtained by analysis of genomic fragments.

POLYNUCLEOTIDE EXPRESSION IN HOST CELLS

5

Polynucleotide sequences or fragments thereof which encode polypeptides, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of a polypeptide in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences that encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express a given polypeptide.

10

As will be understood by those of skill in the art, it may be advantageous in some instances to produce polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

15

Moreover, the polynucleotide sequences can be engineered using methods generally known in the art in order to alter polypeptide encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. For example, DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences.

20

In addition, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

25

Natural, modified, or recombinant nucleic acid sequences may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of polypeptide activity, it may be useful to encode a chimeric protein that can be recognised by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between the polypeptide-encoding sequence and the heterologous protein sequence, so that the polypeptide may be cleaved and purified away from the heterologous moiety.

30

35

Sequences encoding a desired polypeptide may be synthesised, in whole or in part, using chemical methods well known in the art (see Caruthers, M. H. *et al.*, *Nucl. Acids Res. Symp. Ser.* pp. 215-223 (1980), Hom *et al.*, *Nucl. Acids Res. Symp. Ser.* pp. 225-232 (1980)).

Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of a polypeptide, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge *et al.*, *Science* 269:202-204 (1995)) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer, Palo Alto, CA).

- 10 A newly synthesised peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, *Proteins, Structures and Molecular Principles* (1983)) or other comparable techniques available in the art. The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure). Additionally, the amino acid sequence of a polypeptide, or any part thereof, may
15 be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

- In order to express a desired polypeptide, the nucleotide sequences encoding the polypeptide, or functional equivalents, may be inserted into an appropriate expression vector, i.e., a vector
20 which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. Such
25 techniques are described in Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (2000), and Ausubel *et al.*, *Current Protocols in Molecular Biology* (updated annually).

- A variety of expression vector/host systems may be utilised to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as
30 bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

- 35 The "control elements" or "regulatory sequences" present in an expression vector are those non-translated regions of the vector—enhancers, promoters, 5' and 3' untranslated regions—

which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilised, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the PBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or PSFORT1 plasmid (Gibco BRL, Gaithersburg, MD) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be advantageously used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, for example for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke & Schuster, *J. Biol. Chem.* 264:5503-5509 (1989)); and the like. pGEX Vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. Other vectors containing constitutive or inducible promoters include GAP, PGK, GAL and ADH. For reviews, see Ausubel *et al.* (*supra*) and Grant *et al.*, *Methods Enzymol.* 153:516-544 (1987) and Romas *et al.* *Yeast* 8 423-88 (1992).

In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, *EMBO J.* 6:307-311 (1987)). Alternatively,

plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi et al., *EMBO J.* 3:1671-1680 (1984); Broglie et al., *Science* 224:838-843 (1984); and Winter et al., *Results Probl. Cell Differ.* 17:85-105 (1991)). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection.

- 5 Such techniques are described in a number of generally available reviews (see, e.g., Hobbs in *McGraw Hill Yearbook of Science and Technology* pp. 191-196 (1992)).

An insect system may also be used to express a polypeptide of interest. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to
 10 express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia larvae*. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect,
 15 for example, *S. frugiperda* cells or *Trichoplusia larvae* in which the polypeptide of interest may be expressed (Engelhard et al., *Proc. Natl. Acad. Sci. U.S.A.* 91 :3224-3227 (1994)).

In mammalian host cells, a number of viral-based expression systems are generally available. For example, in cases where an adenovirus is used as an expression vector, sequences
 20 encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells (Logan & Shenk, *Proc. Natl. Acad. Sci. U.S.A.* 81:3655-3659 (1984)). In addition, transcription enhancers, such as the Rous
 25 sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. Methods and protocols for working with adenovirus vectors are reviewed in Wold, *Adenovirus Methods and Protocols*, 1998. Additional references regarding use of adenovirus vectors can be found in *Adenovirus: A Medical Dictionary, Bibliography, and Annotated Research Guide to Internet References*, 2004.

30 Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional
 35 transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon

should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf. *et al.*, *Results Probl. Cell Differ.* 20:125-162 (1994)).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler *et al.*, *Cell* 11:223-32 (1977)) and adenine phosphoribosyltransferase (Lowy *et al.*, *Cell* 22:817-23 (1990)) genes which can be employed in tk.sup.- or aprt.sup.- cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 77:3567-70 (1980)); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin *et al.*, *J. Mol. Biol.* 150:1-14 (1981)); and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*). Additional selectable genes have been described, for example, trpB, which allows cells to utilise indole in

place of tryptophan, or hisD, which allows cells to utilise histinol in place of histidine (Hartman & Mulligan, *Proc. Natl. Acad. Sci. U.S.A.* 85:8047-51 (1988)). Recently, the use of visible markers has gained popularity with such markers as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes *et al.*, *Methods Mol. Biol.* 55:121-131 (1995)).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding a polypeptide is inserted within a marker gene sequence, recombinant cells containing sequences can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a polypeptide-encoding sequence under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells which contain and express a desired polynucleotide sequence may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridisations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using either polyclonal or monoclonal antibodies specific for the product are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilising monoclonal antibodies reactive to two non-interfering epitopes on a given polypeptide may be preferred for some applications, but a competitive binding assay may also be employed. These and other assays are described, among other places, in Hampton *et al.*, *Serological Methods, a Laboratory Manual* (1990) and Maddox *et al.*, *J. Exp. Med.* 158:1211-1216 (1983).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labelled hybridisation or PCR probes for detecting sequences related to polynucleotides include oligolabeling, nick translation, end-labelling or PCR amplification using a labelled nucleotide. Alternatively, the sequences, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available,

and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a polynucleotide sequence of interest may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides may be designed to contain signal sequences which direct secretion of the encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilised immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and the encoded polypeptide may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a polypeptide of interest and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilised metal ion affinity chromatography) as described in Porath *et al.*, *Prot. Exp. Purif.* 3:263-281 (1992) while the enterokinase cleavage site provides a means for purifying the desired polypeptide from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll *et al.*, *DNA Cell Biol.* 12:441-453 (1993).

IN VIVO POLYNUCLEOTIDE DELIVERY TECHNIQUES

In additional embodiments, genetic constructs comprising one or more of the polynucleotides of the invention are introduced into cells *in vivo*. This may be achieved using any of a variety of well known approaches, several of which are outlined below for the purpose of illustration.

1. ADENOVIRUS

One of the preferred methods for *in vivo* delivery of one or more nucleic acid sequences involves the use of an adenovirus expression vector. "Adenovirus expression vector" is meant to include those constructs containing adenovirus sequences sufficient to (a) support packaging of the construct and (b) to express a polynucleotide that has been cloned therein in a sense or antisense orientation. Of course, in the context of an antisense construct, expression does not require that the gene product be synthesised.

The expression vector comprises a genetically engineered form of an adenovirus. Knowledge of the genetic organisation of adenovirus, a 36 kb, linear, double-stranded DNA virus, allows substitution of large pieces of adenoviral DNA with foreign sequences up to 7 kb (Grunhaus & Horwitz, 1992). In contrast to retrovirus, the adenoviral infection of host cells does not result in chromosomal integration because adenoviral DNA can replicate in an episomal manner without potential genotoxicity. Also, adenoviruses are structurally stable, and no genome rearrangement has been detected after extensive amplification. Adenovirus can infect virtually all epithelial cells regardless of their cell cycle stage. So far, adenoviral infection appears to be linked only to mild disease such as acute respiratory disease in humans.

Adenovirus is particularly suitable for use as a gene transfer vector because of its mid-sized genome, ease of manipulation, high titre, wide target-cell range and high infectivity. Both ends of the viral genome contain 100-200 base pair inverted repeats (ITRs), which are *cis* elements necessary for viral DNA replication and packaging. The early (E) and late (L) regions of the genome contain different transcription units that are divided by the onset of viral DNA replication. The E1 region (E1A and E1B) encodes proteins responsible for the regulation of transcription of the viral genome and a few cellular genes. The expression of the E2 region (E2A and E2B) results in the synthesis of the proteins for viral DNA replication. These proteins are involved in DNA replication, late gene expression and host cell shut-off (Renan, 1990).

The products of the late genes, including the majority of the viral capsid proteins, are expressed only after significant processing of a single primary transcript issued by the major late promoter (MLP). The MLP, (located at 16.8 m.u.) is particularly efficient during the late phase of infection, and all the mRNA's issued from this promoter possess a 5'-tripartite leader (TPL) sequence which makes them preferred mRNA's for translation.

In a current system, recombinant adenovirus is generated from homologous recombination between shuttle vector and provirus vector. Due to the possible recombination between two

proviral vectors, wild-type adenovirus may be generated from this process. Therefore, it is critical to isolate a single clone of virus from an individual plaque and examine its genomic structure.

- 5 Generation and propagation of the current adenovirus vectors, which are replication deficient, depend on a unique helper cell line, designated 293, which was transformed from human embryonic kidney cells by Ad5 DNA fragments and constitutively expresses E1 proteins (Graham *et al.*, 1977). Since the E3 region is dispensable from the adenovirus genome (Jones & Shenk, 1978), the current adenovirus vectors, with the help of 293 cells, carry foreign DNA in
10 either the E1, the D3 or both regions (Graham & Prevec, 1991). In nature, adenovirus can package approximately 105% of the wild-type genome (Ghosh-Choudhury *et al.*, 1987), providing capacity for about 2 extra kB of DNA. Combined with the approximately 5.5 kB of DNA that is replaceable in the E1 and E3 regions, the maximum capacity of the current adenovirus vector is under 7.5 kB, or about 15% of the total length of the vector. More than
15 80% of the adenovirus viral genome remains in the vector backbone and is the source of vector-borne cytotoxicity. Also, the replication deficiency of the E1-deleted virus is incomplete. For example, leakage of viral gene expression has been observed with the currently available vectors at high multiplicities of infection (MOI) (Mulligan, 1993).
- 20 Helper cell lines may be derived from human cells such as human embryonic kidney cells, muscle cells, hematopoietic cells or other human embryonic mesenchymal or epithelial cells. Alternatively, the helper cells may be derived from the cells of other mammalian species that are permissive for human adenovirus. Such cells include, e.g., Vero cells or other monkey embryonic mesenchymal or epithelial cells. As stated above, the currently preferred helper cell
25 line is 293.

- 30 Racher *et al.* (1995) have disclosed improved methods for culturing 293 cells and propagating adenovirus. In one format, natural cell aggregates are grown by inoculating individual cells into 1 litre siliconised spinner flasks (Technique, Cambridge, UK) containing 100-200 ml of
35 medium. Following stirring at 40 rpm, the cell viability is estimated with trypan blue. In another format, Fibra-Cel microcarriers (Bibby Sterlin, Stone, UK) (5 g/l) is employed as follows. A cell inoculum, resuspended in 5 ml of medium, is added to the carrier (50 ml) in a 250 ml Erlenmeyer flask and left stationary, with occasional agitation, for 1 to 4 h. The medium is then replaced with 50 ml of fresh medium and shaking initiated. For virus production, cells are
40 allowed to grow to about 80% confluence, after which time the medium is replaced (to 25% of the final volume) and adenovirus added at an MOI of 0.05. Cultures are left stationary

overnight, following which the volume is increased to 100% and shaking commenced for another 72 h.

Other than the requirement that the adenovirus vector be replication defective, or at least conditionally defective, the nature of the adenovirus vector is not believed to be crucial to the successful practice of the invention. The adenovirus may be of any of the 42 different known serotypes or subgroups A-F. Adenovirus type 5 of subgroup C is the preferred starting material in order to obtain a conditional replication-defective adenovirus vector for use in the present invention, since Adenovirus type 5 is a human adenovirus about which a great deal of biochemical and genetic information is known, and it has historically been used for most constructions employing adenovirus as a vector.

As stated above, the typical vector according to the present invention is replication defective and will not have an adenovirus E1 region. Thus, it will be most convenient to introduce the polynucleotide encoding the gene of interest at the position from which the E1-coding sequences have been removed. However, the position of insertion of the construct within the adenovirus sequences is not critical to the invention. The polynucleotide encoding the gene of interest may also be inserted in lieu of the deleted E3 region in E3 replacement vectors as described by Karlsson *et al.* (1986) or in the E4 region where a helper cell line or helper virus complements the E4 defect.

Adenovirus is easy to grow and manipulate and exhibits broad host range *in vitro* and *in vivo*. This group of viruses can be obtained in high titres, e.g., 10^9 - 10^{11} plaque-forming units per ml, and they are highly infective. The life cycle of adenovirus does not require integration into the host cell genome. The foreign genes delivered by adenovirus vectors are episomal and, therefore, have low genotoxicity to host cells. No side effects have been reported in studies of vaccination with wild-type adenovirus (Couch *et al.*, 1963; Top *et al.*, 1971), demonstrating their safety and therapeutic potential as *in vivo* gene transfer vectors.

Adenovirus vectors have been used in eukaryotic gene expression (Levrero *et al.*, 1991; Gomez-Foix *et al.*, 1992) and vaccine development (Grunhaus & Horwitz, 1992; Graham & Prevec, 1992). Recently, animal studies suggested that recombinant adenovirus could be used for gene therapy (Stratford-Perricaudet & Perricaudet, 1991; Stratford-Perricaudet *et al.*, 1990; Rich *et al.*, 1993). Studies in administering recombinant adenovirus to different tissues include trachea instillation (Rosenfeld *et al.*, 1991; Rosenfeld *et al.*, 1992), muscle injection (Ragot *et al.*, 1993), peripheral intravenous injections (Herz & Gerard, 1993) and stereotactic inoculation into the brain (Le Gal La Salle *et al.*, 1993).

Adenovirus vectors may originate from human adenovirus. Alternatively they may originate from adenovirus of other species e.g. chimpanzee which may have the advantage that the viral vectors are not neutralised by antibodies against human adenovirus circulating in many human subjects (see e.g.: Tatsis N et al *Gene Therapy* 2006 13:421-429).

5

Adenovirus type 35, which is relatively uncommon and therefore there are low levels of pre-existing immunity to the vector itself, has been used as a delivery system in certain tuberculosis vaccines which are being developed (see for example, Radosevic et al *Infection and Immunity* 2007 75(8):4105-4115). Adenovirus type 35 may also be of particular value in the present invention as a delivery vector.

10

2. RETROVIRUSES

The retroviruses are a group of single-stranded RNA viruses characterised by an ability to convert their RNA to double-stranded DNA in infected cells by a process of reverse-transcription (Coffin, 1990). The resulting DNA then stably integrates into cellular chromosomes as a provirus and directs synthesis of viral proteins. The integration results in the retention of the viral gene sequences in the recipient cell and its descendants. The retroviral genome contains three genes, gag, pol, and env that code for capsid proteins, polymerase enzyme, and envelope components, respectively. A sequence found upstream from the gag gene contains a signal for packaging of the genome into virions. Two long terminal repeat (LTR) sequences are present at the 5' and 3' ends of the viral genome. These contain strong promoter and enhancer sequences and are also required for integration in the host cell genome (Coffin, 1990).

25

In order to construct a retroviral vector, a nucleic acid encoding one or more oligonucleotide or polynucleotide sequences of interest is inserted into the viral genome in the place of certain viral sequences to produce a virus that is replication-defective. In order to produce virions, a packaging cell line containing the gag, pol, and env genes but without the LTR and packaging components is constructed (Mann *et al.*, 1983). When a recombinant plasmid containing a cDNA, together with the retroviral LTR and packaging sequences is introduced into this cell line (by calcium phosphate precipitation for example), the packaging sequence allows the RNA transcript of the recombinant plasmid to be packaged into viral particles, which are then secreted into the culture media (Nicolas & Rubenstein, 1988; Temin, 1986; Mann *et al.*, 1983).

30

The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad variety of cell types.

35

However, integration and stable expression require the division of host cells (Paskind *et al.*, 1975).

A novel approach designed to allow specific targeting of retrovirus vectors was recently developed based on the chemical modification of a retrovirus by the chemical addition of lactose residues to the viral envelope. This modification could permit the specific infection of hepatocytes via sialoglycoprotein receptors.

A different approach to targeting of recombinant retroviruses was designed in which biotinylated antibodies against a retroviral envelope protein and against a specific cell receptor were used. The antibodies were coupled via the biotin components by using streptavidin (Roux *et al.*, 1989). Using antibodies against major histocompatibility complex class I and class II antigens, they demonstrated the infection of a variety of human cells that bore those surface antigens with an ecotropic virus *in vitro* (Roux *et al.*, 1989).

3. ADENO-ASSOCIATED VIRUSES

AAV (Ridgeway, 1988; Hermonat & Muzyczka, 1984) is a parovirus, discovered as a contamination of adenoviral stocks. It is a ubiquitous virus (antibodies are present in 85% of the US human population) that has not been linked to any disease. It is also classified as a dependovirus, because its replication is dependent on the presence of a helper virus, such as adenovirus. Five serotypes have been isolated, of which AAV-2 is the best characterised. AAV has a single-stranded linear DNA that is encapsidated into capsid proteins VP1, VP2 and VP3 to form an icosahedral virion of 20 to 24 nm in diameter (Muzyczka & McLaughlin, 1988).

The AAV DNA is approximately 4.7 kilobases long. It contains two open reading frames and is flanked by two ITRs. There are two major genes in the AAV genome: *rep* and *cap*. The *rep* gene codes for proteins responsible for viral replications, whereas *cap* codes for capsid protein VP1-3. Each ITR forms a T-shaped hairpin structure. These terminal repeats are the only essential *cis* components of the AAV for chromosomal integration. Therefore, the AAV can be used as a vector with all viral coding sequences removed and replaced by the cassette of genes for delivery. Three viral promoters have been identified and named p5, p19, and p40, according to their map position. Transcription from p5 and p19 results in production of *rep* proteins, and transcription from p40 produces the capsid proteins (Hermonat & Muzyczka, 1984).

There are several factors that prompted researchers to study the possibility of using rAAV as an expression vector. One is that the requirements for delivering a gene to integrate into the host chromosome are surprisingly few. It is necessary to have the 145-bp ITRs, which are only 6% of the AAV genome. This leaves room in the vector to assemble a 4.5-kb DNA
 5 insertion. While this carrying capacity may prevent the AAV from delivering large genes, it is amply suited for delivering antisense constructs.

AAV is also a good choice of delivery vehicles due to its safety. There is a relatively complicated rescue mechanism; not only wild type adenovirus but also AAV genes are
 10 required to mobilise rAAV. Likewise, AAV is not pathogenic and not associated with any disease. The removal of viral coding sequences minimises immune reactions to viral gene expression, and therefore, rAAV does not evoke an inflammatory response.

4. OTHER VIRAL VECTORS AS EXPRESSION CONSTRUCTS

Other viral vectors may be employed as expression constructs in the present invention for the delivery of oligonucleotide or polynucleotide sequences to a host cell. Vectors derived from viruses such as vaccinia virus (Ridgeway, 1988; Coupar *et al.*, 1988), lentiviruses, polio
 15 viruses and herpes viruses may be employed. Other poxvirus derived vectors, such as fowl-pox derived vectors, may also be expected to be of use. They offer several attractive features for various mammalian cells (Friedmann, 1989; Ridgeway, 1988; Coupar *et al.*, 1988; Horwich
 20 *et al.*, 1990).

With the recent recognition of defective hepatitis B viruses, new insight was gained into the
 25 structure-function relationship of different viral sequences. *In vitro* studies showed that the virus could retain the ability for helper-dependent packaging and reverse transcription despite the deletion of up to 80% of its genome (Horwich *et al.*, 1990). This suggested that large portions of the genome could be replaced with foreign genetic material. The hepatotropism and persistence (integration) were particularly attractive properties for liver-directed gene
 30 transfer. Chang *et al.* (1991) introduced the chloramphenicol acetyltransferase (CAT) gene into duck hepatitis B virus genome in the place of the polymerase, surface, and pre-surface coding sequences. It was cotransfected with wild-type virus into an avian hepatoma cell line. Culture media containing high titres of the recombinant virus were used to infect primary
 35 duckling hepatocytes. Stable CAT gene expression was detected for at least 24 days after transfection (Chang *et al.*, 1991).

Additional 'viral' vectors include virus like particles (VLPs) and phages.

5. NON-VIRAL VECTORS

In order to effect expression of the oligonucleotide or polynucleotide sequences of the present invention, the expression construct must be delivered into a cell. This delivery may be accomplished *in vitro*, as in laboratory procedures for transforming cells lines, or *in vivo* or *ex vivo*, as in the treatment of certain disease states. As described above, one preferred mechanism for delivery is *via* viral infection where the expression construct is encapsulated in an infectious viral particle.

Once the expression construct has been delivered into the cell the nucleic acid encoding the desired oligonucleotide or polynucleotide sequences may be positioned and expressed at different sites. In certain embodiments, the nucleic acid encoding the construct may be stably integrated into the genome of the cell. This integration may be in the specific location and orientation *via* homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the nucleic acid may be stably maintained in the cell as a separate, episomal segment of DNA. Such nucleic acid segments or "episomes" encode sequences sufficient to permit maintenance and replication independent of or in synchronisation with the host cell cycle. How the expression construct is delivered to a cell and where in the cell the nucleic acid remains is dependent on the type of expression construct employed.

In certain embodiments of the invention, the expression construct comprising one or more oligonucleotide or polynucleotide sequences may simply consist of naked recombinant DNA or plasmids. Transfer of the construct may be performed, for example, by any method which physically or chemically permeabilises the cell membrane. This is particularly applicable for transfer *in vitro* but it may be applied to *in vivo* use as well. Dubensky *et al.* (1984) successfully injected polyomavirus DNA in the form of calcium phosphate precipitates into liver and spleen of adult and newborn mice demonstrating active viral replication and acute infection. Benvenisty & Reshef (1986) also demonstrated that direct intraperitoneal injection of calcium phosphate-precipitated plasmids results in expression of the transfected genes. It is envisioned that DNA encoding a gene of interest may also be transferred in a similar manner *in vivo* and express the gene product.

Another embodiment of the invention for transferring a naked DNA expression construct into cells may involve particle bombardment. This method depends on the ability to accelerate DNA-coated microprojectiles to a high velocity allowing them to pierce cell membranes and

enter cells without killing them (Klein *et al.*, 1987). Several devices for accelerating small particles have been developed. One such device relies on a high voltage discharge to generate an electrical current, which in turn provides the motive force (Yang *et al.*, 1990). The microprojectiles used have consisted of biologically inert substances such as tungsten or gold beads.

Selected organs including the liver, skin, and muscle tissue of rats and mice have been bombarded *in vivo* (Yang *et al.*, 1990; Zelenin *et al.*, 1991). This may require surgical exposure of the tissue or cells, to eliminate any intervening tissue between the gun and the target organ, i.e., *ex vivo* treatment. Again, DNA encoding a particular gene may be delivered via this method and still be incorporated.

Bacteria may also be utilised as a delivery method (e.g. listeria, see WO2004/11048) and in particular BCG.

POLYPEPTIDE COMPOSITIONS

Generally, a polypeptide of use in the invention will be an isolated polypeptide (i.e. separated from those components with which it may usually be found in nature).

For example, a naturally-occurring protein is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. A polynucleotide is considered to be isolated if, for example, it is cloned into a vector that is not a part of the natural environment.

Polypeptides may be prepared using any of a variety of well known techniques. Recombinant polypeptides encoded by DNA sequences as described above may be readily prepared from the DNA sequences using any of a variety of expression vectors known to those of ordinary skill in the art. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast, and higher eukaryotic cells, such as mammalian cells and plant cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line such as COS or CHO. Supernatants from suitable host/vector systems which secrete recombinant protein or polypeptide into culture media may be first concentrated using a commercially available filter. Following concentration, the concentrate may be applied to a suitable purification matrix such as an affinity matrix or an ion

exchange resin. Finally, one or more reverse phase HPLC steps can be employed to further purify a recombinant polypeptide.

Polypeptides of the invention, immunogenic fragments thereof, and other variants having less than about 100 amino acids, and generally less than about 50 amino acids, may also be generated by synthetic means, using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesised using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, *J. Am. Chem. Soc.* 85:2149-2146 (1963). Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

Within certain specific embodiments, a polypeptide may be a fusion protein that comprises multiple polypeptides as described herein, or that comprises at least one polypeptide as described herein and an unrelated sequence, examples of such proteins include tetanus, tuberculosis and hepatitis proteins (see, e.g., Stoute *et al.*, *New Engl. J. Med.* 336:86-91 (1997)). A fusion partner may, for example, assist in providing T helper epitopes (an immunological fusion partner), preferably T helper epitopes recognised by humans, or may assist in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Certain preferred fusion partners are both immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the protein or to enable the protein to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags, which facilitate purification of the protein.

Fusion proteins may generally be prepared using standard techniques, including chemical conjugation. Preferably, a fusion protein is expressed as a recombinant protein, allowing the production of increased levels, relative to a non-fused protein, in an expression system.

Briefly, DNA sequences encoding the polypeptide components may be assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase. This permits translation into a single fusion protein that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea *et al.*, *Gene* 40:39-46 (1985); Murphy *et al.*, *Proc. Natl. Acad. Sci. USA* 83:8258-8262 (1986); U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in length. Linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

Within preferred embodiments, an immunological fusion partner is derived from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenza* B (WO 91/18926). Preferably, a protein D derivative comprises approximately the first third of the protein (e.g., the first N-terminal 100-110 amino acids), and a protein D derivative may be lipidated. Within certain preferred embodiments, the first 109 residues of a lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus functioning as an expression enhancer). The lipid tail ensures optimal presentation of the antigen to antigen presenting cells. Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *Lyta* gene; *Gene* 43:265-292 (1986)). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the

C-LYTA fragment at the amino terminus has been described (see *Biotechnology* 10:795-798 (1992)). Within a preferred embodiment, a repeat portion of LYTA may be incorporated into a fusion protein. A repeat portion is found in the C-terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

5

T CELLS

Immunotherapeutic compositions may also, or alternatively, comprise T cells specific for a *Mycobacterium* antigen. Such cells may generally be prepared *in vitro* or *ex vivo*, using
 10 standard procedures. For example, T cells may be isolated from bone marrow, peripheral blood, or a fraction of bone marrow or peripheral blood of a patient, using a commercially available cell separation system, such as the Isolex™ System, available from Nexell Therapeutics, Inc. (Irvine, CA; see also U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). Alternatively, T cells may be derived from
 15 related or unrelated humans, non-human mammals, cell lines or cultures.

T cells may be stimulated with a polypeptide of the invention, polynucleotide encoding such a polypeptide, and/or an antigen presenting cell (APC) that expresses such a polypeptide. Such stimulation is performed under conditions and for a time sufficient to permit the generation of T
 20 cells that are specific for the polypeptide. Preferably, the polypeptide or polynucleotide is present within a delivery vehicle, such as a microsphere, to facilitate the generation of specific T cells.

T cells are considered to be specific for a polypeptide of the invention if the T cells specifically
 25 proliferate, secrete cytokines or kill target cells coated with the polypeptide or expressing a gene encoding the polypeptide. T cell specificity may be evaluated using any of a variety of standard techniques. For example, within a chromium release assay or proliferation assay, a stimulation index of more than two fold increase in lysis and/or proliferation, compared to negative controls, indicates T cell specificity. Such assays may be performed, for example, as
 30 described in Chen *et al.*, *Cancer Res.* 54:1065-1070 (1994)). Alternatively, detection of the proliferation of T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring an increased rate of DNA synthesis (e.g., by pulse-labelling cultures of T cells with tritiated thymidine and measuring the amount of tritiated thymidine incorporated into DNA). Contact with a polypeptide of the invention (100 ng/ml - 100
 35 µg/ml, preferably 200 ng/ml - 25 µg/ml) for 3 - 7 days should result in at least a two fold increase in proliferation of the T cells. Contact as described above for 2-3 hours should result in activation of the T cells, as measured using standard cytokine assays in which a two fold

increase in the level of cytokine release (e.g., TNF or IFN- γ) is indicative of T cell activation (see Coligan *et al.*, *Current Protocols in Immunology*, vol. 1 (1998)). T cells that have been activated in response to a polypeptide, polynucleotide or polypeptide-expressing APC may be CD4⁺ and/or CD8⁺. Protein-specific T cells may be expanded using standard techniques.

- 5 Within preferred embodiments, the T cells are derived from a patient, a related donor or an unrelated donor, and are administered to the patient following stimulation and expansion.

- For therapeutic purposes, CD4⁺ or CD8⁺ T cells that proliferate in response to a polypeptide, polynucleotide or APC can be expanded in number either *in vitro* or *in vivo*. Proliferation of
 10 such T cells *in vitro* may be accomplished in a variety of ways. For example, the T cells can be re-exposed to a polypeptide, or a short peptide corresponding to an immunogenic portion of such a polypeptide, with or without the addition of T cell growth factors, such as interleukin-2, and/or stimulator cells that synthesise a polypeptide. Alternatively, one or more T cells that proliferate in the presence of the protein can be expanded in number by cloning. Methods for
 15 cloning cells are well known in the art, and include limiting dilution.

PHARMACEUTICAL COMPOSITIONS

- In additional embodiments, the polynucleotide, polypeptide, T-cell and/or antibody
 20 compositions disclosed herein will be formulated in pharmaceutically-acceptable or physiologically-acceptable solutions for administration to a cell or an animal, either alone, or in combination with one or more other modalities of therapy.

- It will also be understood that, if desired, the nucleic acid segment (e.g., RNA or DNA) that
 25 expresses a polypeptide as disclosed herein may be administered in combination with other agents as well, such as, e.g., other proteins or polypeptides or various pharmaceutically-active agents, including chemotherapeutic agents effective against a *M. tuberculosis* infection. In fact, there is virtually no limit to other components that may also be included, given that the additional agents do not cause a significant adverse effect upon contact with the target cells or
 30 host tissues. The compositions may thus be delivered along with various other agents as required in the particular instance. Such compositions may be purified from host cells or other biological sources, or alternatively may be chemically synthesised as described herein. Likewise, such compositions may further comprise substituted or derivatised RNA or DNA compositions.

Formulation of pharmaceutically-acceptable excipients and carrier solutions is well-known to those of skill in the art, as is the development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens, including e.g., oral, parenteral, intravenous, intranasal, and intramuscular administration and
 5 formulation. Other routes of administration include via the mucosal surfaces.

Typically, formulations comprising a therapeutically effective amount deliver about 0.1 ug to about 1000 ug of polypeptide per administration, more typically about 2.5 ug to about 100 ug of polypeptide per administration. In respect of polynucleotide compositions, these typically deliver about 10 ug to about 20 mg of the inventive polynucleotide per administration, more
 10 typically about 0.1 mg to about 10 mg of the inventive polynucleotide per administration

Naturally, the amount of active compound(s) in each therapeutically useful composition may be prepared in such a way that a suitable dosage will be obtained in any given unit dose of the compound. Factors such as solubility, bioavailability, biological half-life, route of
 15 administration, product shelf life, as well as other pharmacological considerations will be contemplated by one skilled in the art of preparing such pharmaceutical formulations, and as such, a variety of dosages and treatment regimens may be desirable.

1. ORAL DELIVERY

20 In certain applications, the pharmaceutical compositions disclosed herein may be delivered via oral administration to an animal. As such, these compositions may be formulated with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard- or soft-shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly
 25 with the food of the diet.

The active compounds may even be incorporated with excipients and used in the form of ingestible tablets, buccal tables, troches, capsules, elixirs, suspensions, syrups, wafers, and the like (Mathiowitz *et al.*, 1997; Hwang *et al.*, 1998; U. S. Patent 5,641,515; U. S. Patent
 30 5,580,579 and U. S. Patent 5,792,451, each specifically incorporated herein by reference in its entirety). The tablets, troches, pills, capsules and the like may also contain the following: a binder, as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or
 35 saccharin may be added or a flavouring agent, such as peppermint, oil of wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain, in addition to

materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar, or both. A syrup or elixir may contain the active component, sucrose as a sweetening agent methyl and propylparabens as

5 preservatives, a dye and flavouring, such as cherry or orange flavour. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active components may be incorporated into sustained-release preparation and formulations.

10 For oral administration the compositions of the present invention may alternatively be incorporated with one or more excipients in the form of a mouthwash, dentifrice, buccal tablet, oral spray, or sublingual orally-administered formulation. For example, a mouthwash may be prepared incorporating the active ingredient in the required amount in an appropriate solvent, such as a sodium borate solution (Dobell's Solution). Alternatively, the active ingredient may
15 be incorporated into an oral solution such as one containing sodium borate, glycerin and potassium bicarbonate, or dispersed in a dentifrice, or added in a therapeutically-effective amount to a composition that may include water, binders, abrasives, flavoring agents, foaming agents, and humectants. Alternatively the compositions may be fashioned into a tablet or solution form that may be placed under the tongue or otherwise dissolved in the mouth.

20

2. INJECTABLE DELIVERY

In certain circumstances it will be desirable to deliver the pharmaceutical compositions disclosed herein parenterally, intravenously, intramuscularly, intradermally, or even

25 intraperitoneally as described in U. S. Patent 5,543,158; U. S. Patent 5,641,515 and U. S. Patent 5,399,363 (each specifically incorporated herein by reference in its entirety). Solutions of the active compounds as free base or pharmacologically acceptable salts may be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils.

30 Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable

35 solutions or dispersions (U. S. Patent 5,466,468, specifically incorporated herein by reference in its entirety). In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and

must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils. Proper fluidity may be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be facilitated by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion (see, e.g., *Remington's Pharmaceutical Sciences*, 15th Edition, pp. 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, and the general safety and purity standards as required by FDA Office of Biologics standards.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

The compositions disclosed herein may be formulated in a neutral or salt form.

Pharmaceutically-acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms such as injectable solutions, drug-release capsules, and the like.

As used herein, "carrier" includes any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

The phrase "pharmaceutically-acceptable" refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a human. The preparation of an aqueous composition that contains a protein as an active ingredient is well understood in the art. Typically, such compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection can also be prepared. The preparation can also be emulsified.

3. NASAL AND BUCCAL DELIVERY

In certain embodiments, the pharmaceutical compositions may be delivered by intranasal sprays, buccal sprays, inhalation, and/or other aerosol delivery vehicles. Methods for delivering genes, nucleic acids, and peptide compositions directly to the lungs eg *via* nasal and buccal aerosol sprays has been described e.g., in U. S. Patent 5,756,353 and U. S. Patent 5,804,212 (each specifically incorporated herein by reference in its entirety). Likewise, the delivery of drugs using intranasal microparticle resins (Takenaga *et al.*, 1998) and lysophosphatidyl-glycerol compounds (U. S. Patent 5,725,871, specifically incorporated herein by reference in its entirety) are also well-known in the pharmaceutical arts. Likewise, transmucosal drug delivery in the form of a polytetrafluoroethylene support matrix is

described in U. S. Patent 5,780,045 (specifically incorporated herein by reference in its entirety).

4. LIPOSOME-, NANOCAPSULE-, AND MICROPARTICLE-MEDIATED DELIVERY

In certain embodiments, the inventors contemplate the use of liposomes, nanocapsules, microparticles, microspheres, lipid particles, vesicles, and the like, for the introduction of the compositions of the present invention into suitable host cells. In particular, the compositions of the present invention may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like.

Such formulations may be preferred for the introduction of pharmaceutically-acceptable formulations of the nucleic acids or constructs disclosed herein. The formation and use of liposomes is generally known to those of skill in the art (see for example, Couvreur *et al.*, 1977; Couvreur, 1988; Lasic, 1998; which describes the use of liposomes and nanocapsules in the targeted antibiotic therapy for intracellular bacterial infections and diseases). Recently, liposomes were developed with improved serum stability and circulation half-times (Gabizon & Papahadjopoulos, 1988; Allen and Choun, 1987; U. S. Patent 5,741,516, specifically incorporated herein by reference in its entirety). Further, various methods of liposome and liposome like preparations as potential drug carriers have been reviewed (Takakura, 1998; Chandran *et al.*, 1997; Margalit, 1995; U. S. Patent 5,567,434; U. S. Patent 5,552,157; U. S. Patent 5,565,213; U. S. Patent 5,738,868 and U. S. Patent 5,795,587, each specifically incorporated herein by reference in its entirety).

Liposomes have been used successfully with a number of cell types that are normally resistant to transfection by other procedures including T cell suspensions, primary hepatocyte cultures and PC 12 cells (Renneisen *et al.*, 1990; Muller *et al.*, 1990). In addition, liposomes are free of the DNA length constraints that are typical of viral-based delivery systems. Liposomes have been used effectively to introduce genes, drugs (Heath & Martin, 1986; Heath *et al.*, 1986; Balazsovits *et al.*, 1989; Fresta & Puglisi, 1996), radiotherapeutic agents (Pikul *et al.*, 1987), enzymes (Imaizumi *et al.*, 1990a; Imaizumi *et al.*, 1990b), viruses (Faller & Baltimore, 1984), transcription factors and allosteric effectors (Nicolau & Gersonde, 1979) into a variety of cultured cell lines and animals. In addition, several successful clinical trials examining the effectiveness of liposome-mediated drug delivery have been completed (Lopez-Berestein *et al.*, 1985a; 1985b; Coune, 1988; Sculier *et al.*, 1988). Furthermore, several studies suggest that the use of liposomes is not associated with autoimmune responses, toxicity or gonadal localization after systemic delivery (Mori & Fukatsu, 1992).

Liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric bilayer vesicles (also termed multilamellar vesicles (MLVs). MLVs generally have diameters of from 25 nm to 4 μ m. Sonication of MLVs
5 results in the formation of small unilamellar vesicles (SUVs) with diameters in the range of 200 to 500 Å, containing an aqueous solution in the core.

Liposomes bear resemblance to cellular membranes and are contemplated for use in connection with the present invention as carriers for the peptide compositions. They are
10 widely suitable as both water- and lipid-soluble substances can be entrapped, i.e. in the aqueous spaces and within the bilayer itself, respectively. It is possible that the drug-bearing liposomes may even be employed for site-specific delivery of active agents by selectively modifying the liposomal formulation.

15 In addition to the teachings of Couvreur *et al.* (1977; 1988), the following information may be utilized in generating liposomal formulations. Phospholipids can form a variety of structures other than liposomes when dispersed in water, depending on the molar ratio of lipid to water. At low ratios the liposome is the preferred structure. The physical characteristics of liposomes depend on pH, ionic strength and the presence of divalent cations. Liposomes can show low
20 permeability to ionic and polar substances, but at elevated temperatures undergo a phase transition which markedly alters their permeability. The phase transition involves a change from a closely packed, ordered structure, known as the gel state, to a loosely packed, less-ordered structure, known as the fluid state. This occurs at a characteristic phase-transition temperature and results in an increase in permeability to ions, sugars and drugs.

25 In addition to temperature, exposure to proteins can alter the permeability of liposomes. Certain soluble proteins, such as cytochrome c, bind, deform and penetrate the bilayer, thereby causing changes in permeability. Cholesterol inhibits this penetration of proteins, apparently by packing the phospholipids more tightly. It is contemplated that the most useful
30 liposome formations for antibiotic and inhibitor delivery will contain cholesterol.

The ability to trap solutes varies between different types of liposomes. For example, MLVs are moderately efficient at trapping solutes, but SUVs are extremely inefficient. SUVs offer the advantage of homogeneity and reproducibility in size distribution, however, and a compromise
35 between size and trapping efficiency is offered by large unilamellar vesicles (LUVs). These are prepared by ether evaporation and are three to four times more efficient at solute entrapment than MLVs.

In addition to liposome characteristics, an important determinant in entrapping compounds is the physicochemical properties of the compound itself. Polar compounds are trapped in the aqueous spaces and nonpolar compounds bind to the lipid bilayer of the vesicle. Polar compounds are released through permeation or when the bilayer is broken, but nonpolar compounds remain affiliated with the bilayer unless it is disrupted by temperature or exposure to lipoproteins. Both types show maximum efflux rates at the phase transition temperature.

Liposomes interact with cells *via* four different mechanisms: endocytosis by phagocytic cells of the reticuloendothelial system such as macrophages and neutrophils; adsorption to the cell surface, either by nonspecific weak hydrophobic or electrostatic forces, or by specific interactions with cell-surface components; fusion with the plasma cell membrane by insertion of the lipid bilayer of the liposome into the plasma membrane, with simultaneous release of liposomal contents into the cytoplasm; and by transfer of liposomal lipids to cellular or subcellular membranes, or vice versa, without any association of the liposome contents. It often is difficult to determine which mechanism is operative and more than one may operate at the same time.

The fate and disposition of intravenously injected liposomes depend on their physical properties, such as size, fluidity, and surface charge. They may persist in tissues for h or days, depending on their composition, and half lives in the blood range from min to several h. Larger liposomes, such as MLVs and LUVs, are taken up rapidly by phagocytic cells of the reticuloendothelial system, but physiology of the circulatory system restrains the exit of such large species at most sites. They can exit only in places where large openings or pores exist in the capillary endothelium, such as the sinusoids of the liver or spleen. Thus, these organs are the predominate site of uptake. On the other hand, SUVs show a broader tissue distribution but still are sequestered highly in the liver and spleen. In general, this *in vivo* behavior limits the potential targeting of liposomes to only those organs and tissues accessible to their large size. These include the blood, liver, spleen, bone marrow, and lymphoid organs.

Targeting is generally not a limitation in terms of the present invention. However, should specific targeting be desired, methods are available for this to be accomplished. Antibodies may be used to bind to the liposome surface and to direct the antibody and its drug contents to specific antigenic receptors located on a particular cell-type surface. Carbohydrate determinants (glycoprotein or glycolipid cell-surface components that play a role in cell-cell recognition, interaction and adhesion) may also be used as recognition sites as they have potential in directing liposomes to particular cell types. Mostly, it is contemplated that

intravenous injection of liposomal preparations would be used, but other routes of administration are also conceivable.

Alternatively, the invention provides for pharmaceutically-acceptable nanocapsule formulations of the compositions of the present invention. Nanocapsules can generally entrap compounds in a stable and reproducible way (Henry-Michelland *et al.*, 1987; Quintanar-Guerrero *et al.*, 1998; Douglas *et al.*, 1987). To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1 μm) should be designed using polymers able to be degraded *in vivo*. Biodegradable polyalkyl-cyanoacrylate nanoparticles that meet these requirements are contemplated for use in the present invention. Such particles may be easily made, as described (Couvreux *et al.*, 1980; 1988; zur Muhlen *et al.*, 1998; Zambaux *et al.* 1998; Pinto-Alphandry *et al.*, 1995 and U. S. Patent 5,145,684, specifically incorporated herein by reference in its entirety).

Skin patches may also be utilised for transcutaneous delivery.

IMMUNOGENIC COMPOSITIONS

In certain preferred embodiments of the present invention, immunogenic compositions are provided. The immunogenic compositions will generally comprise one or more polypeptides or polynucleotides, such as those discussed above, in combination with an immunostimulant. An immunostimulant may be any substance that enhances or potentiates an immune response (antibody and/or cell-mediated) to an exogenous antigen. Examples of immunostimulants include adjuvants, biodegradable microspheres (e.g., polylactic galactide) and liposomes (into which the compound is incorporated; see, e.g., Fullerton, U.S. Patent No. 4,235,877).

Preparation of immunogenic compositions is generally described in, for example, Powell & Newman, eds., *Vaccine Design* (the subunit and adjuvant approach) (1995). Pharmaceutical compositions and immunogenic compositions within the scope of the present invention may also contain other compounds, which may be biologically active or inactive. For example, one or more immunogenic portions of other *M. tuberculosis* antigens may be present, either incorporated into a fusion polypeptide or as a separate compound, within the pharmaceutical or immunogenic composition.

Illustrative immunogenic compositions may contain a polynucleotide (e.g. DNA) encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in situ* (thereby eliciting an immune response). As noted above, the DNA may be present within any

of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Numerous gene delivery techniques are well known in the art, such as those described by Rolland, *Crit. Rev. Therap. Drug Carrier Systems* 15:143-198 (1998), and references cited therein. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve the administration of a bacterium host cell (for example, a *Mycobacterium*, *Bacillus* or *Lactobacillus* strain, including *Bacillus-Calmette-Guerrin* or *Lactococcus lactis*) that expresses the polypeptide (e.g. on its cell surface or secretes the polypeptide) (see, for example, Ferreira, et al., *An Acad Bras Cienc* (2005) 77:113-124; and Raha, et al., *Appl Microbiol Biotechnol* (2005) PubMedID 15635459). In a preferred embodiment, the DNA may be introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are disclosed, for example, in Fisher-Hoch et al., *Proc. Natl. Acad. Sci. USA* 86:317-321 (1989); Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103 (1989); Flexner et al., *Vaccine* 8:17-21 (1990); U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627 (1988); Rosenfeld et al., *Science* 252:431-434 (1991); Kolls et al., *Proc. Natl. Acad. Sci. USA* 91:215-219 (1994); Kass-Eisler et al., *Proc. Natl. Acad. Sci. USA* 90:11498-11502 (1993); Guzman et al., *Circulation* 88:2838-2848 (1993); and Guzman et al., *Cir. Res.* 73:1202-1207 (1993). Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in Ulmer et al., *Science* 259:1745-1749 (1993) and reviewed by Cohen, *Science* 259:1691-1692 (1993). The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells. It will be apparent that a immunogenic composition may comprise both a polynucleotide and a polypeptide component. Such immunogenic composition may provide for an enhanced immune response.

It will be apparent that an immunogenic composition may contain pharmaceutically acceptable salts of the polynucleotides and polypeptides provided herein. Such salts may be prepared from pharmaceutically acceptable non-toxic bases, including organic bases (e.g., salts of primary, secondary and tertiary amines and basic amino acids) and inorganic bases (e.g., sodium, potassium, lithium, ammonium, calcium and magnesium salts).

While any suitable carrier known to those of ordinary skill in the art may be employed in the immunogenic compositions of this invention, the type of carrier will vary depending on the mode of administration. Compositions of the present invention may be formulated for any

appropriate manner of administration, including for example, topical, oral, nasal, intravenous, intracranial, intraperitoneal, subcutaneous or intramuscular administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (e.g., polylactate polyglycolate) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268; 5,075,109; 5,928,647; 5,811,128; 5,820,883; 5,853,763; 5,814,344 and 5,942,252. One may also employ a carrier comprising the particulate-protein complexes described in U.S. Patent No. 5,928,647, which are capable of inducing a class I-restricted cytotoxic T lymphocyte responses in a host.

Such compositions may also comprise buffers (e.g., neutral buffered saline or phosphate buffered saline), carbohydrates (e.g., glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, bacteriostats, chelating agents such as EDTA or glutathione, adjuvants (e.g., aluminum hydroxide), solutes that render the formulation isotonic, hypotonic or weakly hypertonic with the blood of a recipient, suspending agents, thickening agents and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate. Compounds may also be encapsulated within liposomes using well known technology.

Any of a variety of immunostimulants may be employed in the immunogenic compositions of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminium hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium* species or *Mycobacterium* derived proteins. For example, delipidated, deglycolipidated *M. vaccae* ("pVac") can be used. Suitable adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI); Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ); AS01B, AS02A, AS15, AS-2 and derivatives thereof (GlaxoSmithKline, Philadelphia, PA); CWS (cell wall skeleton from a tubercule bacillus), TDM (trehalose dicorynomycolate), Leif (Leishmania elongation initiation factor), aluminium salts such as aluminium hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatized polysaccharides; polyphosphazenes; biodegradable microspheres; monophosphoryl lipid A (MPL®); and quil A

(e.g. QS-21). Cytokines, such as GM-CSF or interleukin-2, -7, or -12, may also be used as adjuvants.

An adjuvant refers to the components in a vaccine or therapeutic composition that increase the specific immune response to the antigen (see, e.g., Edelman, *AIDS Res. Hum Retroviruses* 8:1409-1411 (1992)). Adjuvants induce immune responses of the Th1-type and Th-2 type response. Th1-type cytokines (e.g., IFN- γ , IL-2, and IL-12) tend to favour the induction of cell-mediated immune response to an administered antigen, while Th-2 type cytokines (e.g., IL-4, IL-5, IL-6, IL-10) tend to favour the induction of humoral immune responses. Adjuvants capable of preferential stimulation of a Th-1 cell-mediated immune response are described in WO 94/00153 and WO 95/17209.

Within the immunogenic compositions provided herein, the adjuvant composition is preferably designed to induce an immune response predominantly of the Th1 type. Following application of a immunogenic composition as provided herein, a patient will typically support an immune response that includes Th1- and Th2-type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may be readily assessed using standard assays. For a review of the families of cytokines, see Janeway, *et al.*, *Immunobiology*, 5th Edition, 2001.

The Rv3616c compositions usually comprise one or more adjuvants, e.g., AS01B (3-de-O-acylated monophosphoryl lipid A (3D-MPL®) and QS21 in a liposome formulation; see, U.S. Patent Publication No. 2003/0143240); AS02A (3D-MPL® and QS21 and an oil in water emulsion; see, Bojang, *et al.*, *Lancet* (2001) 358:1927); , ENHANZYN® (Detox); 3D-MPL®; saponins including Quil A and its components e.g. QS21 and saponin mimetics; CWS (cell wall skeleton from a tubercule bacillus); TDM (trehalose dicorynomycolate); aminoalkyl glucosaminide 4-phosphates (AGPs); immunostimulatory oligonucleotides e.g. CPG; Leif (Leishmania elongation initiation factor); and derivatives thereof. In a preferred embodiment, an Rv3616c polypeptide is administered with one or more adjuvants selected from the group consisting of 3D-MPL® and QS21 in a liposome formulation e.g. AS01B and 3D-MPL® and QS21 and an oil in water emulsion (e.g. AS02A). Adjuvant systems AS01B and AS02A are further described in Pichyangkul, *et al.*, *Vaccine* (2004) 22:3831-40.

When delivering the Rv3616c antigen as a nucleic acid, it can be delivered, for example, in a viral vector (*i.e.*, an adenovirus vector), or in a mutant bacterium host cell (*i.e.*, a mutant,

avirulent *Mycobacterium*, *Lactobacillus* or *Bacillus* host cell including *Bacillus Calmette-Guerin* (BCG) and *Lactococcus lactis*).

Preferred adjuvants for use in eliciting a predominantly Th1-type response include, for
 5 example, a combination of monophosphoryl lipid A (MPL®) preferably 3-O-deacylated
 monophosphoryl lipid A (3D-MPL®), optionally with an aluminium salt (see, for example, Ribi,
et al., 1986, Immunology and Immunopharmacology of Bacterial Endotoxins, Plenum Publ.
 Corp., NY, pp. 407-419; GB 2122204B; GB 2220211; and US 4,912,094). A preferred form of
 3D-MPL® is in the form of an emulsion having a small particle size less than 0.2mm in
 10 diameter, and its method of manufacture is disclosed in WO 94/21292. Aqueous formulations
 comprising monophosphoryl lipid A and a surfactant have been described in WO 98/43670.
 Exemplified preferred adjuvants include AS01B (MPL® and QS21 in a liposome formulation),
 3D-MPL® and QS21 in a liposome formulation, AS02A (MPL® and QS21 and an oil in water
 emulsion), 3D-MPL® and QS21 and an oil in water emulsion, and AS15, available from
 15 GlaxoSmithKline. MPL® adjuvants are available from GlaxoSmithKline (see US Patent Nos.
 4,436,727; 4,877,611; 4,866,034 and 4,912,094).

CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce
 a predominantly Th1 response. CpG is an abbreviation for cytosine-guanosine dinucleotide
 20 motifs present in DNA. Such oligonucleotides are well known and are described, for example,
 in WO 96/02555, WO 99/33488 and U.S. Patent Nos. 6,008,200 and 5,856,462.

Immunostimulatory DNA sequences are also described, for example, by Sato *et al.*, *Science*
 273:352 (1996). CpG when formulated into immunogenic compositions, is generally
 administered in free solution together with free antigen (WO 96/02555; McCluskie and Davis,
 25 supra) or covalently conjugated to an antigen (WO 98/16247), or formulated with a carrier such
 as aluminium hydroxide ((Hepatitis surface antigen) Davis *et al.* supra; Brazolot-Millan *et al.*,
Proc.Natl.Acad.Sci., USA, 1998, 95(26), 15553-8). CpG is known in the art as being an
 adjuvant that can be administered by both systemic and mucosal routes (WO 96/02555, EP
 468520, Davis *et al.*, *J.Immunol.*, 1998, 160(2):870-876; McCluskie and Davis, *J.Immunol.*,
 30 1998, 161(9):4463-6).

Another preferred adjuvant is a saponin or saponin mimetics or derivatives, such as Quil A,
 preferably QS21 (Aquila Biopharmaceuticals Inc., Framingham, MA), which may be used alone
 or in combination with other adjuvants. For example, an enhanced system involves the
 35 combination of a monophosphoryl lipid A (MPL®) and saponin derivative, such as the
 combination of QS21 and 3D-MPL® as described in WO 94/00153, or a less reactogenic

composition where the QS21 is quenched with cholesterol, as described in WO 96/33739.

Other preferred formulations comprise an oil-in-water emulsion and tocopherol. A particularly potent adjuvant formulation involving QS21, 3D-MPL® and tocopherol in an oil-in-water emulsion is described in WO 95/17210. Additional saponin adjuvants of use in the present

5 invention include QS7 (described in WO 96/33739 and WO 96/11711) and QS17 (described in U.S. Patent No. 5,057,540 and EP 0 362 279 B1).

Alternatively the saponin formulations may be combined with vaccine vehicles composed of chitosan or other polycationic polymers, polylactide and polylactide-co-glycolide particles, poly-
 10 N-acetyl glucosamine-based polymer matrix, particles composed of polysaccharides or chemically modified polysaccharides, liposomes and lipid-based particles, particles composed of glycerol monoesters, etc. The saponins may also be formulated in the presence of cholesterol to form particulate structures such as liposomes or ISCOM®s. Furthermore, the saponins may be formulated together with a polyoxyethylene ether or ester, in either a non-
 15 particulate solution or suspension, or in a particulate structure such as a paucilamellar liposome or ISCOM®. The saponins may also be formulated with excipients such as CARBOPOL® to increase viscosity, or may be formulated in a dry powder form with a powder excipient such as lactose.

20 In one embodiment, the adjuvant system includes the combination of a monophosphoryl lipid A and a saponin derivative, such as the combination of QS21 and 3D-MPL® adjuvant, as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol containing liposomes, as described in WO 96/33739. Other suitable formulations comprise an oil-in-water emulsion and tocopherol. Another suitable adjuvant
 25 formulation employing QS21, 3D-MPL® adjuvant and tocopherol in an oil-in-water emulsion is described in WO 95/17210.

Another enhanced adjuvant system involves the combination of a CpG-containing oligonucleotide and a saponin derivative particularly the combination of CpG and QS21 as
 30 disclosed in WO 00/09159. Suitably the formulation additionally comprises an oil in water emulsion and tocopherol.

Other suitable adjuvants include MONTANIDE® ISA 720 (Seppic, France), SAF (Chiron, California, United States), ISCOMS® (CSL), MF-59 (Chiron), the SBAS series of adjuvants
 35 (SmithKline Beecham, Rixensart, Belgium), Detox (Corixa), RC-529 (Corixa) and other aminoalkyl glucosaminide 4-phosphates (AGPs), such as those described in pending U.S.

Patent Application Serial Nos. 08/853,826 and 09/074,720, the disclosures of which are incorporated herein by reference in their entireties, and polyoxyethylene ether adjuvants such as those described in WO 99/52549A1. SmithKline Beecham and Corixa Corporation are now part of GlaxoSmithKline.

5

Other suitable adjuvants include adjuvant molecules of the general formula (I):



wherein, n is 1-50, A is a bond or $-\text{C}(\text{O})-$, R is C_{1-50} alkyl or Phenyl C_{1-50} alkyl.

- 10 A further adjuvant of interest is shiga toxin b chain, used for example as described in WO2005/112991.

One embodiment of the present invention consists of an immunogenic composition comprising a polyoxyethylene ether of general formula (I), wherein n is between 1 and 50, preferably 4-24,
 15 most preferably 9; the R component is C_{1-50} , preferably $\text{C}_4\text{-C}_{20}$ alkyl and most preferably C_{12} alkyl, and A is a bond. The concentration of the polyoxyethylene ethers should be in the range 0.1-20%, preferably from 0.1-10%, and most preferably in the range 0.1-1%. Preferred polyoxyethylene ethers are selected from the following group: polyoxyethylene-9-lauryl ether, polyoxyethylene-9-stearyl ether, polyoxyethylene-8-stearyl ether, polyoxyethylene-4-lauryl
 20 ether, polyoxyethylene-35-lauryl ether, and polyoxyethylene-23-lauryl ether. Polyoxyethylene ethers such as polyoxyethylene lauryl ether are described in the Merck index (12th edition: entry 7717). These adjuvant molecules are described in WO 99/52549.

Any immunogenic composition provided herein may be prepared using well known methods
 25 that result in a combination of antigen, immune response enhancer and a suitable carrier or excipient. The compositions described herein may be administered as part of a sustained release formulation (i.e., a formulation such as a capsule, sponge or gel (composed of polysaccharides, for example) that effects a slow release of compound following administration). Such formulations may generally be prepared using well known technology
 30 (see, e.g., Coombes *et al.*, *Vaccine* 14:1429-1438 (1996)) and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations may contain a polypeptide, polynucleotide or antibody dispersed in a carrier matrix and/or contained within a reservoir surrounded by a rate controlling membrane.

35

Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release.

Such carriers include microparticles of poly(lactide-co-glycolide), polyacrylate, latex, starch, cellulose, dextran and the like. Other delayed-release carriers include supramolecular biovectors, which comprise a non-liquid hydrophilic core (e.g., a cross-linked polysaccharide or oligosaccharide) and, optionally, an external layer comprising an amphiphilic compound, such as a phospholipid (see, e.g., U.S. Patent No. 5,151,254 and PCT applications WO 94/20078, WO/94/23701 and WO 96/06638). The amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release.

Any of a variety of delivery vehicles may be employed within pharmaceutical compositions and immunogenic compositions to facilitate production of an antigen-specific immune response. Delivery vehicles include antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen, to improve activation and/or maintenance of the T cell response and/or to be immunologically compatible with the receiver (i.e., matched HLA haplotype). APCs may generally be isolated from any of a variety of biological fluids and organs, and may be autologous, allogeneic, syngeneic or xenogeneic cells.

Certain preferred embodiments of the present invention use dendritic cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau & Steinman, *Nature* 392:245-251 (1998)) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic immunity (see Timmerman & Levy, *Ann. Rev. Med.* 50:507-529 (1999)). In general, dendritic cells may be identified based on their typical shape (stellate *in situ*, with marked cytoplasmic processes (dendrites) visible *in vitro*), their ability to take up, process and present antigens with high efficiency and their ability to activate naïve T cell responses. Dendritic cells may, of course, be engineered to express specific cell-surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells are contemplated by the present invention. As an alternative to dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within an immunogenic composition (see Zitvogel *et al.*, *Nature Med.* 4:594-600 (1998)).

Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, lymph nodes, spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF α to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells harvested from peripheral blood, umbilical cord blood or bone

marrow may be differentiated into dendritic cells by adding to the culture medium combinations of GM-CSF, IL-3, TNF α , CD40 ligand, LPS, flt3 ligand and/or other compound(s) that induce differentiation, maturation and proliferation of dendritic cells.

- 5 Dendritic cells are conveniently categorised as "immature" and "mature" cells, which allows a simple way to discriminate between two well characterised phenotypes. However, this nomenclature should not be construed to exclude all possible intermediate stages of differentiation. Immature dendritic cells are characterised as APC with a high capacity for antigen uptake and processing, which correlates with the high expression of Fc γ receptor and
10 mannose receptor. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and class II MHC, adhesion molecules (e.g., CD54 and CD11) and costimulatory molecules (e.g., CD40, CD80, CD86 and 4-1BB).
- 15 APCs may generally be transfected with a polynucleotide encoding a protein (or portion or other variant thereof) such that the polypeptide, is expressed on the cell surface. Such transfection may take place *ex vivo*, and a pharmaceutical composition or immunogenic composition comprising such transfected cells may then be used, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell
20 may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi *et al.*, *Immunology and Cell Biology* 75:456-460 (1997). Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with
25 the polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant bacterium or viruses (e.g., vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (e.g., a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.
30
- Immunogenic compositions and pharmaceutical compositions may be presented in unit-dose or multi-dose containers, such as sealed ampoules or vials. Such containers are preferably hermetically sealed to preserve sterility of the formulation until use. In general, formulations may be stored as suspensions, solutions or emulsions in oily or aqueous vehicles.
- 35 Alternatively, an immunogenic composition or pharmaceutical composition may be stored in a

freeze-dried condition requiring only the addition of a sterile liquid carrier immediately prior to use.

5 In some embodiments, a "priming" or first administration of an Rv3616c polypeptide (including variants, immunogenic fragments or fusion proteins), or polynucleotide encoding said polypeptide, is followed by one or more "boosting" or subsequent administrations of an Rv3616c polypeptide (including variants, immunogenic fragments or fusion proteins) or polynucleotide encoding said polypeptide ("prime and boost" method). For instance, a first administration with an Rv3616c polypeptide (including variants, immunogenic fragments or fusion proteins) or polynucleotide encoding said polypeptide is followed by one or more subsequent administrations of an Rv3616c polypeptide (including variants, immunogenic fragments or fusion proteins) or polynucleotide encoding said polypeptide.

15 In one embodiment, a first administration with an Rv3616c polypeptide or polynucleotide is followed by one or more subsequent administrations of an Rv3616c polypeptide. In one embodiment, a first administration with an Rv3616c polypeptide or polynucleotide is followed by one or more subsequent administrations of an Rv3616c polynucleotide. Usually the first or "priming" administration and the second or "boosting" administration are given about 2-12 weeks apart, or up to 4-6 months apart. Subsequent "booster" administrations are given about 20 6 months apart, or as long as 1, 2, 3, 4 or 5 years apart. Conventional booster treatment (e.g., a protein priming administration followed by a protein boosting administration) may also be useful in preventing or treating tuberculosis (e.g. preventing or treating latent tuberculosis, in particular preventing or delay tuberculosis reactivation).

25 **ANTIBODIES**

"Antibody" refers to a polypeptide comprising a framework region from an immunoglobulin gene or fragments thereof that specifically binds and recognises an antigen. The recognised immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

35 An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The N-terminus of each chain defines a

variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (V_L) and variable heavy chain (V_H) refer to these light and heavy chains respectively.

5 Antibodies exist, e.g., as intact immunoglobulins or as a number of well-characterised fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce $F(ab)'_2$, a dimer of Fab which itself is a light chain joined to V_H-C_H1 by a disulfide bond. The $F(ab)'_2$ may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the
10 $F(ab)'_2$ dimer into an Fab' monomer. The Fab' monomer is essentially Fab with part of the hinge region (see *Fundamental Immunology* (Paul ed., 3d ed. 1993). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized *de novo* either chemically or by using recombinant DNA methodology. Thus, the term antibody, as used herein, also includes antibody fragments
15 either produced by the modification of whole antibodies, or those synthesised *de novo* using recombinant DNA methodologies (e.g., single chain Fv) or those identified using phage display libraries (see, e.g., McCafferty *et al.*, *Nature* 348:552-554 (1990)).

For preparation of monoclonal or polyclonal antibodies, any technique known in the art can be
20 used (see, e.g., Kohler & Milstein, *Nature* 256:495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pp. 77-96 in *Monoclonal Antibodies and Cancer Therapy* (1985)). Techniques for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanised antibodies.
25 Alternatively, phage display technology can be used to identify antibodies and heteromeric Fab fragments that specifically bind to selected antigens (see, e.g., McCafferty *et al.*, *Nature* 348:552-554 (1990); Marks *et al.*, *Biotechnology* 10:779-783 (1992)).

The phrase "specifically (or selectively) binds" to an antibody or "specifically (or selectively)
30 immunoreactive with," when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and do not substantially bind in a significant amount to other proteins present in the sample. Specific binding to an antibody
35 under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, polyclonal antibodies raised to fusion proteins can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with fusion protein and

not with individual components of the fusion proteins. This selection may be achieved by subtracting out antibodies that cross-react with the individual antigens. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select antibodies specifically immunoreactive with a protein (see, e.g., Harlow & Lane, *Antibodies, A Laboratory Manual* (1988) and *Using Antibodies: A Laboratory Manual* (1998), for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity). Typically a specific or selective reaction will be at least twice background signal or noise and more typically more than 10, 20 or 100 times background (e.g. binding to other *Mycobacterium* proteins, such as other *Mycobacterium tuberculosis* proteins).

DIAGNOSTICS

In another aspect, this invention provides methods for using one or more of the polypeptides described above to diagnose latent tuberculosis (for example using T cell response based assays or antibody based assays of conventional format).

For example, there is provided a method for determining latent *M. tuberculosis* infection in an individual comprising:

- (a) obtaining a sample from the individual;
- (b) contacting said sample with an isolated polypeptide which comprises:
 - (i) an Rv3616c protein sequence;
 - (ii) a variant of an Rv3616c protein sequence; or
 - (iii) an immunogenic fragment of an Rv3616c protein sequence;
- (c) quantifying the sample response.

The sample may for example be whole blood or purified cells. Suitably the sample will contain peripheral blood mononucleated cells (PBMC). In one embodiment of the invention the individual will be seropositive. In a second embodiment of the invention the individual will be seronegative.

Suitably the individual will not previously have been vaccinated against *M. tuberculosis* infection (e.g. suitably the individual will not previously have been vaccinated with BCG).

The sample response may be quantified by a range of means known to those skilled in the art, including the monitoring of lymphocyte proliferation or the production of specific cytokines or antibodies. For example, T-cell ELISPOT may be used to monitor cytokines such as interferon

gamma (IFN γ), interleukin 2 (IL2) and interleukin 5 (IL5). B-cell ELLISPOT may be used to monitor the stimulation of *M. tuberculosis* specific antigens. The cellular response may also be characterised by the use of by intra- and extra-cellular staining and analysis by a flow cytometer.

5

Methods of quantifying a sample proliferation response include:

- (i) pulsing cultured cells with a radiolabel (e.g. tritiated thymidine) and monitoring tritium uptake (e.g. gas scintillation);
- (ii) carboxyfluorescein diacetate succinimidyl ester (CFSE) labelling and

10

fluorescence monitoring of cell division using flow cytometry.

Quantifying a sample cytokine response includes in particular the monitoring of interferon gamma production.

- 15 When using such quantification methods, a positive response to an antigen may be defined by a signal to noise ratio (S/N ratio) of at least 2:1 (for example, at least 3:1 or at least 5:1).

In a further aspect of the present invention methods are provided to diagnose latent *M. tuberculosis* infection using a skin test. As used herein, a "skin test" is any assay performed directly on a patient in which a delayed-type hypersensitivity (DTH) reaction (such as swelling, reddening or dermatitis) is measured following intradermal injection of an Rv153c polypeptide as described above (or variant, immunogenic fragments thereof or nucleotides encoding them). Such injection may be achieved using any suitable device sufficient to contact the antigen combinations with dermal cells of the patient, such as a tuberculin syringe or 1 mL

20 syringe. The reaction is measured after a period of time, for example at least 48 hours after injection, especially 48-72 hours.

25

The DTH reaction is a cell-mediated immune response, which is greater in patients that have been exposed previously to the test antigen. The response may be measured visually, using a ruler. In general, a response that is greater than about 0.5 cm in diameter, especially greater than about 1.0 cm in diameter, is a positive response, indicative of prior *M. tuberculosis* infection, which may or may not be manifested as an active disease.

30

For use in a skin test, the Rv3616c component is suitably formulated as a pharmaceutical composition containing a physiologically acceptable carrier. Suitably, the carrier employed in such pharmaceutical compositions is a saline solution with appropriate preservatives, such as phenol and/or Tween 80™.

35

The present invention further provides kits for use within any of the above diagnostic methods. Such kits typically comprise two or more components necessary for performing a diagnostic assay. Components may be compounds, reagents, containers and/or equipment.

5

For example, one container within a kit may contain a monoclonal antibody or fragment thereof that specifically binds to a protein. Such antibodies or fragments may be provided attached to a support material, as described above. One or more additional containers may enclose elements, such as reagents or buffers, to be used in the assay. Such kits may also, or
10 alternatively, contain a detection reagent as described above that contains a reporter group suitable for direct or indirect detection of antibody binding.

Alternatively, a kit may be designed to detect the level of mRNA encoding a protein in a biological sample. Such kits generally comprise at least one oligonucleotide probe or primer,
15 as described above, that hybridises to a polynucleotide encoding a protein. Such an oligonucleotide may be used, for example, within a PCR or hybridization assay. Additional components that may be present within such kits include a second oligonucleotide and/or a diagnostic reagent or container to facilitate the detection of a polynucleotide encoding a protein of the invention.

20

Other diagnostics kits include those designed for the detection of cell mediated responses (which may, for example, be of use in the diagnostic methods of the present invention). Such kits will typically comprise:

- (i) apparatus for obtaining an appropriate cell sample from a subject;
- 25 (ii) means for stimulating said cell sample with an Rv3616c polypeptide (or variant thereof, immunogenic fragments thereof, or DNA encoding such polypeptides);
- (iii) means for detecting or quantifying the cellular response to stimulation.

Suitable means for quantifying the cellular response include a B-cell ELISPOT kit or
30 alternatively a T-cell ELISPOT kit, which are known to those skilled in the art.

One possible kit comprises:

- (a) a polypeptide of the invention; and
- (b) a detection reagent suitable for direct or indirect detection of antibody binding.

35

Of particular interest are diagnostic kits tailored for quantifying T cell responses:

A diagnostic kit comprising:

- (a) a polypeptide of the invention; and
- (b) apparatus sufficient to contact said polypeptide with the dermal cells of an individual.

5

A diagnostic kit comprising:

- (a) a polypeptide of the invention;
- (b) apparatus sufficient to contact said polypeptide with a sample (e.g. whole blood or more suitably PBMC) from an individual; and
- 10 (c) means to quantify the T cell response (e.g. proliferation or IFN-gamma production).

EXAMPLES

- 15 The following examples are provided by way of illustration only and not by way of limitation. Those of skill in the art will readily recognize a variety of noncritical parameters that could be changed or modified to yield essentially similar results.

EXAMPLE 1 – IDENTIFICATION OF Rv3616c AS A LATENT TB VACCINE TARGET

20

The gene Rv3616c encodes for a conserved hypothetical alanine and glycine rich protein.

25

Rv3616c was selected based on a genome-wide analysis of *Mycobacterium tuberculosis* genes associated with dormancy phase maintenance and infectivity as in Murphy and Brown *BMC.Infect. Dis.* 2007 7:84-99. Potential dormancy phase gene targets in *Mycobacterium tuberculosis* were prioritised through a bioinformatics meta-analysis of published genome-wide DNA microarray datasets of bacterial gene expression under simulated dormancy conditions. Subcellular localisation of *M. tuberculosis* proteins encoded by genes, was subsequently carried out on the entire genome to identify vaccine targets.

30

Briefly, experimental conditions in the dormancy models were quite varied so a zero to five scoring system was developed to normalise these data based upon two criteria: 1) the relevance of the experimental conditions to the dormant state and 2) the rank order of expression. The maximum score for a particular experimental dataset was adjusted based on potential relevance to the clinical occurrence of dormancy phase *M. tuberculosis* infections. Table 1 shows the data sets collected for Step 1 together with the adjusted maximum scores for each dataset. Additional datasets on gene essentiality for growth were obtained from

35

published studies using transposon-based knockout experiments (TraSH). Genes which had no effect on growth received a score of zero.

Table 1 – Sources, experimental models, and scoring criteria for *M. tuberculosis* DNA microarray gene expression and genome-wide gene knock-out (growth phase essentiality).

Reference	Experimental model	Timepoint: Maximum score ^a
Betts JC et al. <i>Mol. Microbiol.</i> 2002 43:717-731	Starvation under controlled O ₂	96h: 3 24h: 2 4h: 1
Hampshire T et al. <i>Tuberculosis.(Edinb.)</i> 2004 84:228-238	Nutrient depletion under controlled O ₂	62 and 75d: 5 49d: 4 18d: 2
Muttucumaru DG et al. <i>Tuberculosis.(Edinb.)</i> 2004 84:239-246	Wayne model of hypoxia [#]	14d (NRP-2): 4 7d (NRP-1): 2
Voskuil MI et al. <i>Tuberculosis.(Edinb.)</i> 2004 84:218-227	Wayne model of hypoxia [#]	30 and 80d: 5 14 and 20d: 4 10 and 12d: 3 6 and 8d: 2
Schnappinger D et al. <i>J. Exp. Med.</i> 2003 198:693-704	Infection of mouse macrophages, +/- γ-INF	24 and 48h: 5
Karakousis PC et al. <i>J. Exp. Med.</i> 2004 200:647-657	Hollow fiber subcutaneous implant in mice	10d: 3
Talaat AM et al. <i>Proc. Natl. Acad. Sci. U.S.A</i> 2004, 101:4602-4607	Infection of mice. MTB harvested from lung ^b	28d: 3
Sasseti CM et al. <i>Mol. Microbiol.</i> 2003 48:77-84	TraSH mutated libraries grown on solid media	14d:5
Rengarajan J et al. <i>Proc.Natl.Acad.Sci.U.S.A</i> 2005, 102:8327-8332	Infection of mouse macrophages, +/- γ-INF with TraSH mutated libraries of <i>M. tuberculosis</i>	7d:5
Sasseti CM et al. <i>Proc.Natl.Acad.Sci.U.S.A</i> 2003 100:12989-12994	C57BL/6J mice infected with TraSH mutated libraries of <i>M. tuberculosis</i>	7, 14, 28 and 56d:5

^aMaximum score based on relevance as a dormancy model; h = hour ; d = day..

^bRatio of *M. tuberculosis* from Balb/c lung to MTB in aerated culture for 28d.

Wayne LG and Hayes LG *Infect. Immun.* 1996 64:2062-2069

5 Step 2 - In applying the second criterion, the rank order of gene expression, gene scores from each dataset were ordered from highest to lowest based on expression ratio (fold expression in the experimental condition versus cells in log-phase liquid culture). The highest scoring gene received the maximum score for that particular dataset (listed in column 3 of Table 1. (e.g. 5, 4 ... , 1 point)). The score was decreased by 0.005 points for each gene in order until zero, or the
10 end of the data set was reached. Thus when the maximum score was 4 points, the 100th ranked gene would receive a score of 3.500. For a maximum score of 5 points, 1000 genes or 25% of the *M. tuberculosis* genome received a score. For experiments where data from multiple time points were collected, the maximum score across all time points was used as the final score.

15

In Step 3 scores for each gene in each of the experimental conditions were collected into a Microsoft Access database. Reference fields were added to facilitate prioritisation, such as the Refseq ID, Genbank function, Genbank note, Tuberculist classification, and KEGG and Sanger Center links. By combining the data from different studies and sources, a consensus view was
20 reached about the particular genes and pathways most critical for survival in the dormant state.

In Step 4, a prioritised list of therapeutic targets was derived utilising the top 400 scoring genes (~10% of the genome) supplemented by expert computational and manual analysis of biochemical pathways, enzymology, drug tractability, homology to human genes and other
25 prior knowledge. The great majority of the high scoring genes come from the subset where two or three of the groups intersect.

In Step 5, the identification of subcellular localisation of *M. tuberculosis* proteins encoded by genes, was carried out on the entire genome. The heuristic used for membrane protein
30 prediction is described in Chalker et al. *J. Bacteriol.* 2001 183:1259-1268. Average hydropathy profiles (H) (von Heijne G *J. Mol. Biol.* 1992 225:487-494) were generated using GES hydropathy values (Engelman DM et al. *Annu. Rev. Biophys. Biophys. Chem.* 1986 15:321-353) weighted using a trapezoid window. Using a process similar to the initial steps of the TopPred II algorithm (Claros MG et al. *Comput. Appl. Biosci.* 1994 10:685-686), helical
35 transmembrane segments (TMS) were predicted for each peptide sequence by selecting 19 amino acids centered on the highest H value (MaxH), masking these from further consideration, and repeating the process until no peaks with a H of >0.5 remained. Subcellular

locations were assigned based on the peak MaxH value, number of segments with a H of >1.0, and distribution and peak H values of the putative TMS. A MaxH cutoff of 1.15 was chosen to maximize the discrimination between two SwissProtein release 34 test datasets containing transmembrane and cytoplasmic proteins, respectively (Boyd D et al. *Protein Sci.* 1998 7:201-205). Proteins with a MaxH of <1.15 were classified as cytoplasmic, while those with a MaxH of >1.15 and at least three possible TMS were classified as membrane proteins. Anchored proteins were defined as having exactly two TMS, one starting before amino acid (aa) 35 and one having a H of >1.15 with the other having a H not lower than 0.5. SignalP with Gram positive settings was specifically used for *M. bacterium* to identify secreted proteins amongst those classified as either cytoplasmic or "unknown" in the heuristic analysis (Nielsen H et al. *Protein Eng.* 1997 10:1-6).

Rv3616c ranked very high as a vaccine antigen according to several criteria:

- (i) Rv3616c is consistently up-regulated across all models of dormancy. Among the entire suite of 3999 genes scored in the meta-analysis, Rv3616c was ranked in the top quartile of overexpressed genes across all dormancy models. The up-regulated score for Rv3616c was 6.52 which favourably compared with the top gene score of 22.28.
- (ii) Rv3616c ranked as being highly essential for survival in the mouse spleen infection model (scoring 4.945, out of a possible scoring of 5).
- (iii) Subcellular localisation predicted that Rv3616c protein is a membrane bound protein and thus has significant extracellular exposure, indicating suitability as a vaccine target.
- (iv) Rv3616c can elicit a protective response against initial tuberculosis challenge.
- (v) Rv3616c is broadly recognised as an antigen.

EXAMPLE 2 – Rv3616c EPITOPE PREDICTION

Method

T cell epitope prediction was based on the following approaches:

Prediction	Name	URL/References
CD4 and CD8	Multipred	website: antigen.i2r.a-star.edu.sg/multipred/ Zhang, G.L., Khan, A.M., Srinivasan, K.N., August, J.T. and Brusic, V. (2005) "MULTIPRED: a computational system for prediction of promiscuous HLA binding peptides" <i>Nucleic Acids Res.</i> 33, W172 - W179.
	SVMHC	website: www-bs.informatik.uni-tuebingen.de/SVMHC "Prediction of MHC class I binding peptides, using SVMHC." Pierre Dönnes and Arne Elofsson in: <i>BMC Bioinformatics</i> 2002 3: 25
CD4	ProPred	website: www.imtech.res.in/raghava/propred/ Singh, H. and Raghava, G.P.S. (2001) "ProPred: Prediction of HLA-DR binding sites." <i>Bioinformatics</i>, 17(12), 1236-37.
	Tepitope2	In house program based on : H. Bian, J. Hammer (2004) "Discovery of promiscuous HLA-II-restricted T cell epitopes with TEPITOPE." <i>Methods</i> 34 : 468-75
CD8	nHLA	website: www.imtech.res.in/raghava/nhlapred/ Bhasin M. and Raghava G P S (2006) "A hybrid approach for predicting promiscuous MHC class I restricted T cell epitopes"; <i>J. Biosci.</i> 32:31-42
	NetCTL	website: www.cbs.dtu.dk/services/NetCTL/ "An integrative approach to CTL epitope prediction. A combined algorithm integrating MHC-I binding, TAP transport efficiency, and proteasomal cleavage predictions." Larsen M.V., Lundegaard C., Kasper Lamberth, Buus S., Brunak S., Lund O., and Nielsen M. <i>European Journal of Immunology.</i> 35(8): 2295-303. 2005

Prediction	Name	URL/References
	EpiJen	website: www.jenner.ac.uk/EpiJen/ Doytchinova, I. A., P. Guan, D. R. Flower. "EpiJen: a server for multi-step T cell epitope prediction." <i>BMC Bioinformatics</i>, 2006, 7, 131.
	Syfpeithi	website: www.syfpeithi.de/Scripts/MHCServer.dll/EpitopePrediction.htm Hans-Georg Rammensee, Jutta Bachmann, Niels Nikolaus Emmerich, Oskar Alexander Bachor, Stefan Stevanovic: "SYFPEITHI: database for MHC ligands and peptide motifs." <i>Immunogenetics</i> (1999) 50: 213-219
	PredTAP	website: antigen.i2r.a-star.edu.sg/predTAP/ Zhang, G.L., Petrovsky, N., Kwok, C.K., August, J.T. and Brusic, V. (2006) "PRED^{TAP}: a system for prediction of peptide binding to the human transporter associated with antigen processing." <i>Immunome Res.</i> 2(1), 3.
	PAPROC	http://www.paproc2.de/paproc1/paproc1.html C. Kuttler, A.K. Nussbaum, T.P. Dick, H.-G. Rammensee, H. Schild, K.P. Haderl, "An algorithm for the prediction of proteasomal cleavages", <i>J. Mol. Biol.</i> 298 (2000), 417-429 A.K. Nussbaum, C. Kuttler, K.P. Haderl, H.-G. Rammensee, H. Schild, "PAPROC: A Prediction Algorithm for Proteasomal Cleavages available on the WWW", <i>Immunogenetics</i> 53 (2001), 87-94

Results

5

Table 2 – Putative Rv3616c human CD4+ T cell epitopes

Putative CD4 epitope number	Amino acid position	Epitope sequence	SEQ ID No:	HLA allele
-----------------------------	---------------------	------------------	------------	------------

Putative CD4 epitope number	Amino acid position	Epitope sequence	SEQ ID No:	HLA allele
1	5	FIIDPTISA	SEQ ID No: 29	DRB1_0301, DRB1_0401, DRB1_1101
2	31	ILYSSLEYF	SEQ ID No: 30	DRB1_0301
3	36	LEYFEKALE	SEQ ID No: 31	DRB1_1301
4	63	YAGKNRNVH	SEQ ID No: 32	DRB1_0801
5	87	LIHDQANAV	SEQ ID No: 33	DRB1_0301, DRB1_0401
6	111	FVRPVAVDL	SEQ ID No: 34	DRB1_0101
7	119	LTYIPVVGH	SEQ ID No: 35	DRB1_0401
8	121	YIPVVGHAL	SEQ ID No: 36	DRB1_0101
9	151	YLVVKTLIN	SEQ ID No: 37	DRB1_0401
10	152	LVVVKTLINA	SEQ ID No: 38	DRB1_1301
11	154	VKTLINATQ	SEQ ID No: 39	DRB1_0401
12	164	LKLLAKLAE	SEQ ID No: 40	DRB1_0301, DRB1_0801, DRB1_1101, DRB1_1301
13	173	LVAAAIADI	SEQ ID No: 41	DRB1_0301, DRB1_1101, DRB1_1301
14	181	IISDVADII	SEQ ID No: 42	DRB1_0301
15	197	WEFITNALN	SEQ ID No: 43	DRB1_0401
16	252	LFGAAGLSA	SEQ ID No: 44	DRB1_1501
17	264	LAHADSLAS	SEQ ID No: 45	DRB1_0401
18	270	LASSASLPA	SEQ ID No: 46	DRB1_0401
19	288	FGGLPSLAQ	SEQ ID No: 47	DRB1_0401

Table 3 – Putative Rv3616c human CD8+ T cell epitopes

Putative CD8 epitope number	Amino acid position	Epitope sequence	SEQ ID No:	HLA allele
1	5	FIIDPTISA	SEQ ID No: 48	A2
2	6	IIDPTISAI	SEQ ID No: 49	A_0101, A2
3	9	PTISAIIDGL	SEQ ID No: 50	A2, A_0201, B7, B8
4	10	TISAIIDGLY	SEQ ID No: 51	A1, A_0101, A3, A_0301
5	12	SAIDGLYDL	SEQ ID No: 52	A2, B_3501
6	13	AIDGLYDLL	SEQ ID No: 53	A_0101, A_0201, B44
7	17	LYDLLGIGI	SEQ ID No: 54	A24
8	25	IPNQGGILY	SEQ ID No: 55	B7, A_0101, B_3501, B51
9	30	GILYSSLEY	SEQ ID No: 56	A1, A_0101, A3, A_0301
10	33	YSSLEYFEK	SEQ ID No: 57	A1, A_0301
11	35	SLEYFEKAL	SEQ ID No: 58	A_0201, B7, Cw_0401, Cw_0602
12	38	YFEKALEEL	SEQ ID No: 59	A24, A_2402, B8, Cw_0401, Cw_0602
13	39	FEKALEELA	SEQ ID No: 60	B44, B_4403
14	69	NHVNFFQEL	SEQ ID No: 61	A24, Cw_0602
15	76	ELADLDRQL	SEQ ID No: 62	A_0201
16	77	LADLDRQLI	SEQ ID No: 63	A_0101, B51
17	79	DLDRQLISL	SEQ ID No: 64	A_0101, A_0201
18	80	LDRQLISLI	SEQ ID No: 65	A24, B7, B51

Putative CD8 epitope number	Amino acid position	Epitope sequence	SEQ ID No:	HLA allele
19	94	AVQTTRDIL	SEQ ID No: 66	B7
20	103	EGAKKGLEF	SEQ ID No: 67	A24, B7
21	107	KGLEFVRPV	SEQ ID No: 68	A_0201, B51
22	108	GLEFVRPVA	SEQ ID No: 69	A_0101, A_0301
23	109	LEFVRPVAV	SEQ ID No: 70	B44
24	111	FVRPVAVDL	SEQ ID No: 71	B7, B8, B_3501
25	113	RPVAVDLTY	SEQ ID No: 72	B7, A_0101, B_3501, B51
26	116	AVDLTYIPV	SEQ ID No: 73	A2, A_0201
27	120	TYIPVVGHA	SEQ ID No: 74	A24
28	121	YIPVVGHAL	SEQ ID No: 75	A_0101, A2, A_0201, B7, B8
29	129	LSAAFQAPF	SEQ ID No: 76	A1, B7, B_3501
30	130	SAAFQAPFC	SEQ ID No: 77	A_0201
31	131	AAFQAPFCA	SEQ ID No: 78	A_0301, B_3501
32	133	FQAPFCAGA	SEQ ID No: 79	A2, A_0201
33	135	APFCAGAMA	SEQ ID No: 80	B7, B_3501
34	136	PFCAGAMAV	SEQ ID No: 81	A3
35	141	AMAVVGGAL	SEQ ID No: 82	A2, A_0201, A24, B7
36	143	AVVGGALAY	SEQ ID No: 83	A1, A3, A_0301, B7
37	147	GALAYLVVK	SEQ ID No: 84	A3, A_0301
38	149	LAYLVVKTL	SEQ ID No: 85	B8, B44, B51
39	150	AYLVVKTLI	SEQ ID No: 86	A24
40	155	KTINATQL	SEQ ID No: 87	A_0201, A2, A_0301, A24
41	156	TLINATQLL	SEQ ID No: 88	A2, A_0201, A3, A_0101, Cw_0401
42	158	INATQLLKL	SEQ ID No: 89	B7, B8, Cw_0602
43	159	NATQLLKL	SEQ ID No: 90	A_2402, B7, B_3501, B44, Cw_0401, Cw_0602
44	162	QLLKLLAKL	SEQ ID No: 91	A2, A_0201, A_0301, A_2402, B8, Cw_0401, Cw_0602
45	165	KLLAKLAEL	SEQ ID No: 92	A2, A_0201, A_0301, B7, B8, Cw_0602
46	166	LLAKLAELV	SEQ ID No: 93	A2, A_0201, A_0101, B8
47	169	KLAELVAAA	SEQ ID No: 94	A2
48	170	LAELVAAAI	SEQ ID No: 95	A1, A24, B51
49	173	LVAAAIADI	SEQ ID No: 96	B7, B51
50	177	AIADIISDV	SEQ ID No: 97	A2, A_0201, Cw_0602
51	178	IADIISDVA	SEQ ID No: 98	A_0101, B_3501
52	182	ISDVADIHK	SEQ ID No: 99	A1, A_0301
53	192	TLGEVWEFI	SEQ ID No: 100	A2, A_0201
54	199	FITNALNGL	SEQ ID No: 101	A2
55	202	NALNGLKEL	SEQ ID No: 102	B51, A_2402, B_3501, Cw_0602
56	213	KLTGWVTGL	SEQ ID No: 103	A2, A_0201
57	214	LTGWVTGLF	SEQ ID No: 104	A1, A_0101, A24
58	225	GWSNLESFF	SEQ ID No: 105	A24
59	228	NLESFFAGV	SEQ ID No: 106	A2, A_0201
60	231	SFFAGVPGL	SEQ ID No: 107	A2, A_0201, A24, Cw_0401
61	238	GLTGATSGL	SEQ ID No: 108	A2, A_0201
62	246	LSQVTGLFG	SEQ ID No: 109	A1, B8
63	247	SQVTGLFGA	SEQ ID No: 110	A2
64	258	LSASSGLAH	SEQ ID No: 111	A1, A3, B7, B8
65	260	ASSGLAHAD	SEQ ID No: 112	A1, A3, A_0301

Putative CD8 epitope number	Amino acid position	Epitope sequence	SEQ ID No:	HLA allele
66	262	SGLAHADSL	SEQ ID No: 113	A_0201
67	263	GLAHADSLA	SEQ ID No: 114	A_0101, A_0201, A_0301
68	269	SLASSASLP	SEQ ID No: 115	A_0201, A_0301
69	271	ASSASLPAL	SEQ ID No: 116	B7
70	286	SGFGGLPSL	SEQ ID No: 117	A2, A_0201, B51
71	291	LPSLAQVHA	SEQ ID No: 118	B7, B_3501, B51
72	298	HAASTRQAL	SEQ ID No: 119	B7, B8, B_3501
73	301	STRQALRPR	SEQ ID No: 120	A3, A_0301
74	307	RPRADGPVG	SEQ ID No: 121	B7, B_0702, B8, B51
75	319	EQVGGQSQL	SEQ ID No: 122	B7, B44
76	350	GASKGTTTK	SEQ ID No: 123	A3, A_0301
77	351	ASKGTTTKK	SEQ ID No: 124	A3, A_0301
78	353	KGTTTKKYS	SEQ ID No: 125	A_0301, B8
79	368	TEDAERAPV	SEQ ID No: 126	B44

As can be seen from Tables 2 and 3, Rv3616c contains a number of predicted CD4+ and CD8 T cell epitopes. Furthermore, this information suggests that the protein carries epitopes that can be recognised by HLAs which occur worldwide (that is HLAs from Caucasian, African,
5 Asian or Latin-American individuals – see website at www.allele frequencies.net).

EXAMPLE 3 – Rv3616c EPIOTOPE IDENTIFICATION

A range of 30 overlapping peptides covering the full length of Rv3616c were prepared (see
10 Figure 1 for details and SEQ ID Nos: 127-156) and tested for their ability to stimulate PBMC from four PPD+ donors.

The data, shown in Figure 2, reveals that peptides 1-7 and 17-30 were immunogenic for these individuals.

15

It should be noted that peptides 8-16 (amino acid residues 92-215) may be immunogenic in other individuals of differing HLA type.

EXAMPLE 4 – H37Rv HOMOLOGUES

20

Rv3616c sequences from a number of *M. tuberculosis* strains and BCG were identified using BLASTP searches of GenBank (H37Rv reference sequence accession number NP_218133.1):

Strain	Accession Number	% identity
CDC1551	NP_338263.1	99

F11	YP_001289574.1	99
Haarlem	ZP_02248979.1	99
C	ZP_00877472.1	99
BCG	YP_979759.1	99

Alignment of the homologue sequences indicates a high level of identity.

BIOLOGICAL ASSAYS

5

Quantification of T cell responses to Rv3616c

Polypeptides may be screened for their ability to activate T-cells (induction of proliferation and/or production of cytokines) in peripheral blood mononuclear cell (PBMC) or in whole blood preparations from infected (such as latently infected) individuals.

10

Latently infected individuals are usually identified by a skin test that has a diameter above 10mm and without symptoms, with no Mtb positive culture, with a negative sputum negative and with no lesion (as detected by chest X-Ray).

A range of *in vitro* assays can be used based on PBMC samples or whole blood: after restimulation in presence of the antigen (or variant/immunogenic fragment thereof as appropriate) the proliferation of the cells may be determined (as measured by CFSE/flow cytometry) or the production of cytokines quantified (present in the supernatant of cultured cells and measured by ELISA, or, after intracellular staining of CD4 and CD8 T cells and analysis by flow cytometry).

15

For example, PBMC samples may be obtained from heparinised whole blood by Ficoll-Hypaque density gradient centrifugation following standard procedures. The cells may then be washed and cryopreserved in liquid nitrogen until testing (for further details see Lalvani A et al. *J. Infect. Dis.* 1999 180:1656-1664).

20

T cell proliferation

25

The specific immune response may be characterised by performing lymphocyte proliferation analysis using the tritiated thymidine. This technique assesses the cellular expansion upon *in vitro* stimulation to an antigen. In practice, cell proliferation is determined by estimating

incorporation of tritiated-thymidine into DNA, a process closely related to underlying changes in cell number.

More suitably, lymphocyte proliferation may be performed using the succinimidyl ester of carboxyfluorescein diacetate (CFSE). CFSE spontaneously and irreversibly couples to both intracellular and cell surface proteins by reaction with lysine side chains and other available amine groups. When lymphocyte cells divide, CFSE labelling is distributed equally between the daughter cells, which are therefore half as fluorescent as the parents. As a result, halving of cellular fluorescence intensity marks each successive generation in a population of proliferating cells and is readily followed by flow cytometry (for further details see Hodgkins, PD et al *J. Exp. Med.* 1996 184:277-281).

Practically, after thawing, PMBC may be washed and stained with CFSE before being cultivated (2×10^5 cells) for 72 hrs with 10 ug/ml of antigen in culture media (RPMI-1640 supplemented with glutamine, non essential amino acid, pyruvate and heat inactivated human AB serum). Cells may then then harvested and their phenotype characterised using surface staining to identify memory CD8 and CD4+ T-Cells. Subsequently, flow cytometry analysis can be used to indicate the extent of lymphocyte proliferation in response to each antigen (proportion of cells with decreased CFSE intensity upon *in vitro* stimulation).

Cytokine production

IFN- γ production (or the production of other cytokines such as e.g. IL2, TNF-alpha, IL5, IL12 etc) may be measured using an enzyme-linked immunosorbent assay (ELISA). ELISA plates may be coated with a mouse monoclonal antibody directed to human IFN- γ (PharMingen, San Diego, CA) in PBS for four hours at room temperature. Wells are then blocked with PBS containing 5% (W/V) non-fat dried milk for 1 hour at room temperature. The plates are then washed, for example, six times in PBS/0.2% TWEEN-20 and samples diluted 1:2 in culture medium in the ELISA plates are incubated overnight at room temperature. The plates are again washed and a polyclonal rabbit anti-human IFN- γ serum, for example, diluted 1:3000 in PBS/10% normal goat serum may be added to each well. The plates are then incubated for two hours at room temperature, washed and horseradish peroxidase-coupled anti-rabbit IgG (Sigma Chemical So., St. Louis, MO) may be added, for example, at a 1:2000 dilution in PBS/5% non-fat dried milk. After a further two hour incubation at room temperature, the plates are washed and TMB substrate added. The reaction may be stopped after 20 min with 1 N sulfuric acid. Optical density can then be determined at 450 nm using 570 nm as a reference

wavelength. Typically, fractions that result in both replicates giving an OD two fold greater than the mean OD from cells cultured in medium alone may be considered positive.

EXAMPLE 5 – IMMUNOGENICITY IN CB6F1 MICE

- 5 The immunogenicity of the antigen was evaluated in CB6F1 mice (first generation cross of BALB/c and C57BL/6 mice).

- CB6F1 mice were immunised intramuscularly three times (on day 0, day 14 and day 28) with 0.5 ug of protein antigen in combination with the Adjuvant System AS01E (a liposomal
10 adjuvant formulation comprising 3D-MPL and QS21).

The experimental design was as follows:

Group	Day 0	Day 14	Day 28
1	0.5 ug Rv3616c/AS01E	0.5 ug Rv3616c/AS01E	0.5 ug Rv3616c/AS01E

- 15 A total of 24 mice were used in the protocol group.

- Peripheral blood lymphocytes (PBL) were collected and pooled on day 21 (i.e. 7 days post second immunisation) and day 35 (i.e. 7 days post third immunisation) and the antigen-specific CD4 & CD8 T cell responses (as determined by CD4 or CD8 T cells producing IL-2 and/or IFN-
20 gamma and/or TNF-alpha) were measured by flow cytometry after overnight *in vitro* restimulation with pools of 15mer peptides covering the sequences of interest.
The detection of mouse T cells that express IL-2 and/or IFN-gamma and/or TNF-alpha was done by using short-term antigen-driven *in vitro* amplification of cytokine expression.

- 25 Briefly, PharmLyse solution (BD-Pharmingen) was added to heparinised mouse peripheral blood in order to lyse the red blood cells. The PBLs (Peripheral Blood Lymphocytes) obtained were washed and then incubated in the presence of a pool of 15-mer peptides - overlapping by 11 amino acids - covering the sequence of the antigen of interest and of 1 ug/ml of antibodies to CD28 and CD49d (BD-Pharmingen). Each 15-mer peptide was used at a final
30 concentration of 1 ug/ml. Medium controls were also stimulated with antibodies to CD28 and CD49d.

The cytokine secretion blocking compound brefeldin-A (BD-Pharmingen) was added 2 h after the onset of the cultures at 37°C, 5% CO₂ and the cells maintained at 37°C, 5% CO₂ for 4 additional hours followed by overnight incubation at +4°C.

- 5 Cells were then harvested and stained with Pacific Blue-coupled anti-CD4 (BD – clone RM4-5, BD-Pharmingen) and peridinin chlorophyll A protein (PerCp) cyanin5.5 (Cy5.5)-coupled anti-CD8 alpha (clone 53-6.7, BD-Pharmingen) antibodies.

10 Cells were then washed, fixed, permeabilised (Cytotfix-cytoperm kit, BD-Pharmingen) and stained with allophycocyanin-coupled anti-IFN- γ antibodies (clone XMG1.2, BDPharmingen), fluorescein isothiocyanate (FITC)-coupled anti-IL-2 antibodies (clone JES 6-5H4, Beckman Coulter) and phycoerythrin (PE)-coupled anti-TNF alpha antibodies (clone MP6-XT22, BDPharmingen). After final washes, stained cells were analysed on a LSR II flow cytometer (Beckton-Dickinson). A minimum number of 10,000 cells were acquired in the CD8⁺ subset.

15 For further background see Walzer T et al *Cell Immunol.* 2000 206(1):16-25 and Maecker HT et al *J. Immunol. Methods* 2001 255(1-2):27-40.

20 As negative controls, some cells were also cultured overnight *in vitro* in culture medium (unstimulated). The antigen-specific responses were calculated by subtracting the average cytokine response produced by unstimulated cells from the average cytokine response produced by the peptide-stimulated cells.

25 At each timepoint and for each group, the data was collected from 4 pools of 6 mice each. The data below is presented as the % of CD4 or CD8 T cells producing IL-2 and/or IFN- γ and/or TNF- α . Each individual pool of mice is plotted (triangles) as well as the average value of the group (bar).

30 Figure 3 shows that on day 21 (i.e. 7 days post second immunisation), Rv3616c-specific CD4 and CD8 T cell responses are detected in mice immunised with 0.5 μ g of Rv3616c/AS01E.

Figure 4 shows the cytokine profile of CD4 T cell response from the Rv3616c peptide pool-stimulated PBL (not medium removed) on day 21 (i.e. 7 days post second immunisation).

35 Figure 5 shows the cytokine profile of CD8 T cell response from the Rv3616c peptide pool-stimulated PBL (not medium removed) on day 21 (i.e. 7 days post second immunisation).

Figure 6 shows that on day 35 (i.e. 7 days post third immunisation), Rv3616c-specific CD4 and CD8 T cell responses are detected in mice immunised with 0.5 ug of Rv3616c/AS01E. The third dose increases the CD4 T cell response but not the CD8 T cell response. Due to technical difficulties, data was only available for a single pool.

5

Figure 7 shows the cytokine profile of CD4 T cell response from the Rv3616c peptide pool-stimulated PBL (not medium removed) on day 35 (i.e. 7 days post third immunisation). Due to technical difficulties, data was only available for a single pool.

10 Figure 8 shows the cytokine profile of CD8 T cell response from the Rv3616c peptide pool-stimulated PBL (not medium removed) on day 35 (i.e. 7 days post third immunisation). Due to technical difficulties, data was only available for a single pool.

EXAMPLE 6 – IMMUNOGENICITY IN C57BL/6 MICE

15

The immunogenicity of the antigen was also evaluated in C57BL/6 mice.

C57BL/6 mice were immunised intramuscularly three times (on day 0, day 14 and day 28) with 1 ug protein antigen in combination with a the Adjuvant System AS01E (a liposomal adjuvant formulation comprising 3D-MPL and QS21)..

20

The experimental design was the following:

Group	Day 0	Day 14	Day 28
1	1 ug Rv3616c/AS01E	1 ug Rv3616c/AS01E	1 ug Rv3616c/AS01E

25 Peripheral blood lymphocytes (PBL) were collected and pooled on day 21 (i.e. 7 days post second immunisation) and day 35 (i.e. 7 days post third immunisation) and the antigen-specific CD4 & CD8 T cell responses (as determined by CD4 or CD8 T cells producing IL-2 and/or IFN-gamma and/or TNF-alpha) were measured by flow cytometry after overnight *in vitro* restimulation with pools of 15mer peptides covering the sequences of interest. The procedure followed was as described previously.

30

As negative controls, some cells were also cultured overnight *in vitro* in culture medium (unstimulated). The antigen-specific responses were calculated by subtracting the average

cytokine response produced by unstimulated cells from the average cytokine response produced by the peptide-stimulated cells.

At each timepoint and for each group, the data was collected from 4 pools of 6 mice each. The data below is presented as the % of CD4 or CD8 T cells producing IL-2 and/or IFN-gamma and/or TNF-alpha. Each individual pool of mice is plotted (triangles) as well as the average value of the group (bar).

Figure 9 shows that on day 21 (i.e. 7 days post second immunisation), Rv3616c-specific CD4 and CD8 T cell responses are detected in mice immunised with 1 ug of Rv3616c/AS01E, although the antigen-specific CD8 T cell response is very low (cytokine profile data is therefore not shown).

Figure 10 shows the cytokine profile of CD4 T cell response from the Rv3616c peptide pool-stimulated PBL (not medium removed) on day 21 (i.e. 7 days post second immunisation).

Figure 11 shows that on day 35 (i.e. 7 days post third immunisation), Rv3616c-specific CD4 and CD8 T cell responses are detected in mice immunised with 1 ug of Rv3616c/AS01E. A third immunisation dose increases the CD4 T cell responses but only slightly the CD8 T cell response.

Figure 12 shows the cytokine profile of CD4 T cell response from the Rv3616c peptide pool-stimulated PBL (not medium removed) on day 35 (i.e. 7 days post third immunisation).

Figure 13 shows the cytokine profile of CD8 T cell response from the Rv3616c peptide pool-stimulated PBL (not medium removed) on day 35 (i.e. 7 days post third immunisation).

EXAMPLE 7 – IN VITRO RECOGNITION BY PBMC FROM HUMANS WITH LATENT TB

Experiments were performed in order to assess the peripheral T cell response specific to the inventive antigen in 4 TB naïve healthy adults (PPD skin test = 0 mm) and 8 TB latently Infected healthy adults (PPD skin test = 15 mm or above) from South Africa

PPD Skin Test Data	
Individual ID	Induration
Number	diameter

	(mm)
4	0
5	0
33	0
38	0
36	15
46	15
13	15
7	16
58	25
74	26
8	53
60	55

The cell-mediated immune (CMI) response was assessed by measuring cytokines on isolated peripheral blood mononuclear cells (PBMCs) by intracellular cytokine staining (ICS) assay.

- 5 ICS carried out was an adaptation of previously described methodology (see Von Eschen et al, *Hum. Vaccin.* 2009 5(7)). PBMCs were stimulated *in vitro* by one pool of 15-mer peptides - overlapping by 11 amino acids - covering the entire sequence of the antigen of interest. Cells were stimulated with peptides for 2 hours, further cultured overnight in the presence of Brefeldin A, processed for ICS and analysed using flow cytometry. The frequencies of the
- 10 antigen-specific CD3+CD4+ or CD3+CD8+ T cells expressing IFN-gamma and/or TNF-alpha and/or IL-17 were measured. Medium-stimulated cell responses were subtracted from the responses obtained in peptide pools stimulated cells.

ICS: antibodies

- 15 Anti-CD3 PO (Invitrogen – cat CD0330)
 Anti-CD4 PB (BD - cat 558116)
 Anti-CD8 APC-H7 (BD – cat 641400)
 Anti-IFNγ AF700 (BD-Pharmingen – cat 557995)
 Anti-TNF PE-Cy7 (BD-Pharmingen – cat 557647)
- 20 Anti-IL17 AF647 (BD-Pharmingen – cat 51-7178-71)

The results are presented as number of antigen-specific CD3+CD4+ T cells expressing TNF-alpha and IFN-gamma, per million of CD3+CD4+ T cells since these cells represent the main population of the antigen-specific CD4 T cells (the background response level due to the

medium is removed). No antigen-specific CD3+CD8+ T cells were detected. Figure 14 shows that an antigen-specific CD4 T cell response is measured in 6 out of 8 latently infected individuals (not in individuals number 7 and 74) when compared to the non-specific CD4 T cell response measured in the naïve individuals.

5

In conclusion it may be noted that the Rv3616c antigen is capable of eliciting an immune response in both CB6F1 and C57BL/6 mice. Furthermore, the profile of cytokine production indicates that a large proportion of antigen-specific T-cells express a plurality of Th1 associated cytokines (i.e. a polyfunctional T-cell response is elicited). Importantly both CD4 and CD8 antigen-specific T-cells are present after immunisation, CD8 cells may be particularly important in a latent TB scenario. The relevance of Rv3616c to human infection is confirmed by the high level of recognition in latently infected individuals from South Africa and the absence of responses in naïve subjects. Rv3616c may therefore be expected to be of substantial value in the prevention, treatment and diagnosis of latent tuberculosis infection.

10

15

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to one of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

20

All references referred to in this application, including patents and patent applications, are incorporated herein by reference to the fullest extent possible as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

25

Throughout the specification and the claims which follow, unless the context requires otherwise, the word 'comprise', and variations such as 'comprises' and 'comprising', will be understood to imply the inclusion of a stated integer, step, group of integers or group of steps but not to the exclusion of any other integer, step, group of integers or group of steps.

30

WHAT IS CLAIMED IS:

1. An isolated polypeptide which comprises:
 - (i) an Rv3616c protein sequence;
 - (ii) a variant of an Rv3616c protein sequence; or
 - (iii) an immunogenic fragment of an Rv3616c protein sequence,for use in the treatment or prevention of latent TB.
2. A polypeptide according to claim 1, which comprises an Rv3616c protein sequence.
3. A polypeptide according to claim 2, which consists of an Rv3616c protein sequence.
4. A polypeptide according to claim 1, which comprises a variant of an Rv3616c protein sequence.
5. A polypeptide according to claim 4, which comprises a variant of an Rv3616c protein sequence having at least 70% identity thereto.
6. A polypeptide according to claim 5, which comprises a variant of an Rv3616c protein sequence having at least 90% identity thereto.
7. A polypeptide according to claim 6, which comprises a variant of an Rv3616c protein sequence having at least 95% identity thereto.
8. A polypeptide according to claim 4, which consists of a variant of an Rv3616c protein sequence.
9. A polypeptide according to claim 8, which consists of a variant of an Rv3616c protein sequence having at least 70% identity thereto.
10. A polypeptide according to claim 9, which consists of a variant of an Rv3616c protein sequence having at least 90% identity thereto.
11. A polypeptide according to claim 10, which consists of a variant of an Rv3616c protein sequence having at least 95% identity thereto.

12. A polypeptide according to claim 1, which comprises an immunogenic fragment of an Rv3616c protein sequence.
13. A polypeptide according to claim 12, which comprises an immunogenic fragment of
5 least 10 amino acid residues from an Rv3616c protein sequence.
14. A polypeptide according to claim 13, which comprises an immunogenic fragment of
least 20 amino acid residues from an Rv3616c protein sequence.
- 10 15. A polypeptide according to claim 14, which comprises an immunogenic fragment of
least 50 amino acid residues from an Rv3616c protein sequence.
16. A polypeptide according to claim 12, which consists of an immunogenic fragment of an
Rv3616c protein sequence.
- 15 17. A polypeptide according to claim 16, which consists of an immunogenic fragment of
least 10 amino acid residues from an Rv3616c protein sequence.
18. A polypeptide according to claim 17, which consists of an immunogenic fragment of
20 least 20 amino acid residues from an Rv3616c protein sequence.
19. A polypeptide according to claim 18, which consists of an immunogenic fragment of
least 50 amino acid residues from an Rv3616c protein sequence.
- 25 20. A polypeptide according to any one of claims 1 to 19, wherein the Rv3616c protein
sequence is from *Mycobacterium tuberculosis*.
21. A polypeptide according to claim 20, wherein the Rv3616c protein has the sequence of
SEQ ID NO: 1.
- 30 22. A polypeptide according to any one of claims 1 to 21, for use in the treatment of latent
tuberculosis.
23. A polypeptide according to any one of claims 1 to 21, for use in the prevention of latent
35 tuberculosis.

24. A polypeptide according to any one of claims 1 to 21, for use in the prevention of reactivation of tuberculosis.
- 5 25. A polypeptide according to any one of claims 1 to 21, for use in the delay of reactivation of tuberculosis.
26. A polypeptide according to any one of claims 22 to 25, wherein the tuberculosis is associated with a *Mycobacterium tuberculosis* infection
- 10 27. An isolated polynucleotide comprising a nucleic acid sequence encoding a polypeptide which comprises:
 - (i) an Rv3616c protein sequence;
 - (ii) a variant of an Rv3616c protein sequence; or
 - (iii) an immunogenic fragment of an Rv3616c protein sequence,15 for use in the treatment or prevention of latent TB.
28. A polynucleotide according to claim 27, comprising a nucleic acid sequence encoding a polypeptide which comprises an Rv3616c protein sequence.
- 20 29. A polynucleotide according to claim 28, comprising a nucleic acid sequence encoding a polypeptide which comprises a variant of an Rv3616c protein sequence.
30. A polynucleotide according to claim 29, comprising a nucleic acid sequence encoding a polypeptide which comprises a variant of an Rv3616c protein sequence having at least 25 70% identity thereto.
31. A polynucleotide according to claim 27, comprising a nucleic acid sequence encoding a polypeptide which comprises an immunogenic fragment of an Rv3616c protein sequence.
- 30 32. A polynucleotide according to claim 31, comprising a nucleic acid sequence encoding a polypeptide which comprises an immunogenic fragment of at least 10 amino acid residues from an Rv3616c protein sequence.
- 35 33. A polynucleotide according to any one of claims 27 to 32, comprising a nucleic acid sequence encoding a polypeptide which consists of:
 - (i) an Rv3616c protein sequence;

- (ii) a variant of an Rv3616c protein sequence; or
- (iii) an immunogenic fragment of an Rv3616c protein sequence

- 5 34. A polynucleotide according to claim 33, consisting of a nucleic acid sequence encoding a polypeptide which consists of:
- (i) an Rv3616c protein sequence;
 - (ii) a variant of an Rv3616c protein sequence; or
 - (iii) an immunogenic fragment of an Rv3616c protein sequence.
- 10 35. A polynucleotide according to any one of claims 27 to 34, wherein the Rv3616c protein sequence is from *Mycobacterium tuberculosis*.
- 15 36. A polynucleotide according to claim 35, wherein the Rv3616c protein has the sequence of SEQ ID NO: 1.
37. A polynucleotide according to claim 36, which comprises the sequence of SEQ ID NO: 2, or a fragment thereof.
- 20 38. A polynucleotide according to any one of claims 28 to 37, for use in the treatment of latent tuberculosis.
39. A polynucleotide according to any one of claims 28 to 37, for use in the prevention of latent tuberculosis.
- 25 40. A polynucleotide according to any one of claims 28 to 37, for use in the prevention of reactivation of tuberculosis.
41. A polynucleotide according to any one of claims 28 to 37, for use in the delay of reactivation of tuberculosis.
- 30 42. A polynucleotide according to any one of claims 38 to 41, wherein the tuberculosis is associated with a *Mycobacterium tuberculosis* infection
- 35 43. A pharmaceutical composition comprising:
- (a) a polypeptide which comprises:
 - (i) an Rv3616c protein sequence;
 - (ii) a variant of an Rv3616c protein sequence; or

- (iii) an immunogenic fragment of an Rv3616c protein sequence; or
- (b) a polynucleotide comprising a nucleic acid sequence encoding the polypeptide of (a);
- and
- 5 (c) a pharmaceutically acceptable carrier or excipient, for use in the treatment or prevention of latent TB.

44. An immunogenic composition comprising:

- (a) a polypeptide which comprises:

- 10 (i) an Rv3616c protein sequence;
- (ii) a variant of an Rv3616c protein sequence; or
- (iii) an immunogenic fragment of an Rv3616c protein sequence; or
- (b) a polynucleotide comprising a nucleic acid sequence encoding the polypeptide of (a);
- 15 and
- (c) a non-specific immune response enhancer, for use in the treatment or prevention of latent TB.

45. An immunogenic composition according to claim 44, wherein the non-specific immune response enhancer is an adjuvant.

46. A method for the treatment or prevention of latent tuberculosis comprising the administration of a safe and effective amount of a polypeptide comprising:

- 25 (i) an Rv3616c protein sequence;
 - (ii) a variant of an Rv3616c protein sequence; or
 - (iii) an immunogenic fragment of an Rv3616c protein sequence;
- to a subject in need thereof, wherein said polypeptide induces an immune response.

47. A method for the treatment or prevention of latent tuberculosis comprising the administration of a safe and effective amount of a polynucleotide comprising a nucleic acid sequence encoding a polypeptide comprising:

- 30 (i) an Rv3616c protein sequence;
 - (ii) a variant of an Rv3616c protein sequence; or
 - (iii) an immunogenic fragment of an Rv3616c protein sequence;
- 35 to a subject in need thereof, wherein said polynucleotide induces an immune response.

48. The method according to either claim 46 or 47, wherein the polypeptide or polynucleotide is administered as a pharmaceutical composition.
- 5 49. The method according to either claim 46 or 47, wherein the polypeptide or polynucleotide is administered as an immunogenic composition.
50. The method according to any one of claims 46 to 49, wherein the subject is a mammal.
- 10 51. The method according to claim 50, wherein the subject is a human.
52. The method according to any one of claims 46 to 51, wherein the subject has active tuberculosis.
- 15 53. The method according to any one of claims 46 to 51, wherein the subject has latent tuberculosis.
54. The method according to any one of claims 46 to 51, wherein the subject does not have tuberculosis.
- 20 55. The method according to any one of claims 46 to 54, wherein the subject was previously immunised with *Bacillus Calmette-Guerin* (BCG).
56. The method according to any one of claims 46 to 55, wherein the wherein the Rv3616c protein sequence is from *Mycobacterium tuberculosis*.
- 25 57. The method according to claim 56, wherein the Rv3616c protein has the sequence of SEQ ID NO: 1.
- 30 58. The method according to any one of claims 46 to 57, further comprising the administration of one or more chemotherapeutic agents effective in treating tuberculosis.
59. The method according to claim 58, wherein the one or more chemotherapeutic agents are selected from isoniazid and rifampin.
- 35 60. The method according to any one of claims 46 to 59, further comprising the administration of at least one additional *Mycobacterium tuberculosis* antigen.

61. The method according to claim 60, wherein the additional *Mycobacterium tuberculosis* antigen is provided in the form of a polypeptide.

5 62. The method according to claim 61, wherein the polypeptide comprising:
(i) an Rv3616c protein sequence;
(ii) a variant of an Rv3616c protein sequence; or
(iii) an immunogenic fragment of an Rv3616c protein sequence;
and the additional *Mycobacterium tuberculosis* antigen are provided in the form of a
10 fusion protein.

63. The method according to claim 60, wherein the additional *Mycobacterium tuberculosis* antigen is provided in the form of a polynucleotide.

15 64. The method according to claim 63, wherein the polynucleotide comprising a nucleic acid sequence encoding a polypeptide comprising:
(i) an Rv3616c protein sequence;
(ii) a variant of an Rv3616c protein sequence; or
(iii) an immunogenic fragment of an Rv3616c protein sequence;
20 and the additional *Mycobacterium tuberculosis* antigen are provided in the form of a single polynucleotide sequence.

65. The method according to claim 64, wherein said single polynucleotide sequence encodes a fusion protein containing the Rv3616c antigen and the additional
25 *Mycobacterium tuberculosis* antigen.

66. The method according to claim 60, wherein the additional *Mycobacterium tuberculosis* antigen is provided in the form of a *Bacillus Calmette-Guerin* (BCG).

30 67. The method according to any one of claims 46 to 66, wherein the treatment or prevention of latent tuberculosis refers to the treatment of latent tuberculosis.

68. The method according to any one of claims 46 to 66, wherein the treatment or prevention of latent tuberculosis refers to the prevention of latent tuberculosis.
35

69. The method according to any one of claims 46 to 66, wherein the treatment or prevention of latent tuberculosis refers to the prevention of reactivation of tuberculosis.

70. The method according to any one of claims 46 to 66, wherein the treatment or prevention of latent tuberculosis refers to the delay of reactivation of tuberculosis.
- 5 71. The method according to any one of claims 46 to 70, wherein the tuberculosis is associated with a *Mycobacterium tuberculosis* infection.
72. The method according to claim 71, wherein the *Mycobacterium tuberculosis* infection is associated with a multi-drug resistant strain of *Mycobacterium tuberculosis*.
- 10 73. The method according to any one of claims 46 to 72, which induces an immune response against *M. tuberculosis* infection.
74. Use of a polypeptide comprising:
- 15 (i) an Rv3616c protein sequence;
 (ii) a variant of an Rv3616c protein sequence; or
 (iii) an immunogenic fragment of an Rv3616c protein sequence;
 in the manufacture of a medicament for the treatment or prevention of latent tuberculosis.
- 20 75. Use of a polynucleotide comprising a nucleic acid sequence encoding a polypeptide comprising:
- (i) an Rv3616c protein sequence;
 (ii) a variant of an Rv3616c protein sequence; or
 25 (iii) an immunogenic fragment of an Rv3616c protein sequence;
 in the manufacture of a medicament for the treatment or prevention of latent tuberculosis.
76. A fusion protein comprising a polypeptide comprising:
- 30 (i) an Rv3616c protein sequence;
 (ii) a variant of an Rv3616c protein sequence; or
 (iii) an immunogenic fragment of an Rv3616c protein sequence,
 for use in the treatment or prevention of latent TB.
- 35 77. A polynucleotide comprising a nucleotide sequence encoding a fusion protein comprising a polypeptide comprising:
- (i) an Rv3616c protein sequence;

(ii) a variant of an Rv3616c protein sequence; or
 (iii) an immunogenic fragment of an Rv3616c protein sequence,
 for use in the treatment or prevention of latent TB.

5 78. A polynucleotide comprising a first nucleotide sequence which selectively hybridises under moderately stringent conditions to the complement of a second nucleotide sequence which encodes a polypeptide comprising:

(i) an Rv3616c protein sequence;
 (ii) a variant of an Rv3616c protein sequence; or
 10 (iii) an immunogenic fragment of an Rv3616c protein sequence,
 for use in the treatment or prevention of latent TB.

15 79. A polynucleotide according to claim 78, comprising a first nucleotide sequence which selectively hybridises under highly stringent conditions to the complement of a second nucleotide sequence which encodes a polypeptide comprising:

(i) an Rv3616c protein sequence;
 (ii) a variant of an Rv3616c protein sequence; or
 (iii) an immunogenic fragment of an Rv3616c protein sequence,
 for use in the treatment or prevention of latent TB.

20 80. A polynucleotide according to either claim 78 or 79, comprising a first nucleotide sequence which selectively hybridises to the complement of SEQ ID NO: 2.

25 81. A diagnostic kit comprising:
 (a) a polypeptide according to claim 1;
 (b) apparatus sufficient to contact said polypeptide with a sample (e.g. whole blood or more suitably PBMC) from an individual; and
 (c) means to quantify the T cell response of the sample.

30 82. A diagnostic kit according to claim 81, comprising:
 (a) a polypeptide according to claim 1;
 (b) apparatus sufficient to contact said polypeptide with a sample (e.g. whole blood or more suitably PBMC) from an individual; and
 (c) means to quantify and characterise the T cell response of the sample.

35 83. A diagnostic kit comprising:
 (a) a polypeptide according to claim 1;

- (b) apparatus sufficient to contact said polypeptide with a sample (e.g. whole blood or more suitably PBMC) from an individual; and
 - (c) means to quantify the B cell response of the sample.
- 5 84. A diagnostic kit comprising:
 - (a) a polypeptide according to claim 1; and
 - (b) apparatus sufficient to contact said polypeptide with the dermal cells of a patient.
- 10 85. A diagnostic kit comprising:
 - (a) a polypeptide according to claim 1; and
 - (c) means to quantify the T cell response in a sample from an individual (e.g. whole blood or more suitably PBMC).
- 15 86. A diagnostic kit comprising:
 - (a) a polypeptide according to claim 1; and
 - (c) means to quantify the B cell response in a sample from an individual (e.g. whole blood or more suitably PBMC).

eolf-seq1

SEQUENCE LISTING

<110> GlaxoSmithKline Biologicals s.a.
 Glaxo Group Limited
 Mettens, Pascal
 Brown, James
 Murphy, Dennis

<120> NOVEL COMPOSITIONS AND METHODS

<130> VB63090PCT

<150> US61/083720

<151> 2008-07-25

<160> 159

<170> PatentIn version 3.5

<210> 1

<211> 392

<212> PRT

<213> Mycobacterium tuberculosis

<220>

<221> MISC_FEATURE

<223> Strain H37Rv

<400> 1

Met Ser Arg Ala Phe Ile Ile Asp Pro Thr Ile Ser Ala Ile Asp Gly
 1 5 10 15

Leu Tyr Asp Leu Leu Gly Ile Gly Ile Pro Asn Gln Gly Gly Ile Leu
 20 25 30

Tyr Ser Ser Leu Glu Tyr Phe Glu Lys Ala Leu Glu Glu Leu Ala Ala
 35 40 45

Ala Phe Pro Gly Asp Gly Trp Leu Gly Ser Ala Ala Asp Lys Tyr Ala
 50 55 60

Gly Lys Asn Arg Asn His Val Asn Phe Phe Gln Glu Leu Ala Asp Leu
 65 70 75 80

Asp Arg Gln Leu Ile Ser Leu Ile His Asp Gln Ala Asn Ala Val Gln
 85 90 95

Thr Thr Arg Asp Ile Leu Glu Gly Ala Lys Lys Gly Leu Glu Phe Val
 100 105 110

Arg Pro Val Ala Val Asp Leu Thr Tyr Ile Pro Val Val Gly His Ala
 115 120 125

Leu Ser Ala Ala Phe Gln Ala Pro Phe Cys Ala Gly Ala Met Ala Val
 130 135 140

Val Gly Gly Ala Leu Ala Tyr Leu Val Val Lys Thr Leu Ile Asn Ala

eol f-seq1															
145				150				155				160			
Thr	Gln	Leu	Leu	Lys 165	Leu	Leu	Ala	Lys	Leu 170	Ala	Glu	Leu	Val	Ala 175	Ala
Ala	Ile	Ala	Asp 180	Ile	Ile	Ser	Asp	Val 185	Ala	Asp	Ile	Ile	Lys 190	Gly	Thr
Leu	Gly	Glu 195	Val	Trp	Glu	Phe	Ile 200	Thr	Asn	Ala	Leu	Asn 205	Gly	Leu	Lys
Glu	Leu 210	Trp	Asp	Lys	Leu	Thr 215	Gly	Trp	Val	Thr	Gly 220	Leu	Phe	Ser	Arg
Gly 225	Trp	Ser	Asn	Leu	Glu 230	Ser	Phe	Phe	Ala	Gly 235	Val	Pro	Gly	Leu	Thr 240
Gly	Ala	Thr	Ser	Gly 245	Leu	Ser	Gln	Val	Thr 250	Gly	Leu	Phe	Gly	Ala 255	Ala
Gly	Leu	Ser	Ala 260	Ser	Ser	Gly	Leu	Ala 265	His	Ala	Asp	Ser	Leu 270	Ala	Ser
Ser	Ala	Ser 275	Leu	Pro	Ala	Leu	Ala 280	Gly	Ile	Gly	Gly	Gly 285	Ser	Gly	Phe
Gly	Gly 290	Leu	Pro	Ser	Leu	Ala 295	Gln	Val	His	Ala	Ala 300	Ser	Thr	Arg	Gln
Ala 305	Leu	Arg	Pro	Arg	Ala 310	Asp	Gly	Pro	Val	Gly 315	Ala	Ala	Ala	Glu	Gln 320
Val	Gly	Gly	Gln	Ser 325	Gln	Leu	Val	Ser	Ala 330	Gln	Gly	Ser	Gln	Gly 335	Met
Gly	Gly	Pro	Val 340	Gly	Met	Gly	Gly	Met 345	His	Pro	Ser	Ser	Gly 350	Ala	Ser
Lys	Gly	Thr 355	Thr	Thr	Lys	Lys	Tyr 360	Ser	Glu	Gly	Ala	Ala 365	Ala	Gly	Thr
Glu	Asp 370	Ala	Glu	Arg	Ala	Pro 375	Val	Glu	Ala	Asp	Ala 380	Gly	Gly	Gly	Gln
Lys 385	Val	Leu	Val	Arg	Asn 390	Val	Val								

eo1f-seq1

<220>
 <221> misc_feature
 <223> Strain H37Rv

<400> 2
 atgagcagag cgttcatcat cgatccaacg atcagtgcc a ttgacggctt gtacgacctt 60
 ctggggattg gaatacccaa ccaagggggg atcctttact cctcactaga gtacttcgaa 120
 aaagccctgg aggagctggc agcagcgttt ccgggtgatg gctgggtagg ttcggccgcg 180
 gacaaatacg ccggcaaaaa ccgcaaccac gtgaattttt tccaggaact ggcagacctc 240
 gatcgtcagc tcatcagcct gatccacgac caggccaacg cgggtccagac gacccgcgac 300
 atcctggagg gcgccaagaa aggtctcgag ttcgtgcgcc cgggtggctgt ggacctgacc 360
 tacatcccgg tcgtcgggca cgccctatcg gccgcccttc aggcgccgtt ttgcgcgggc 420
 gcgatggccg tagtggggcg cgcgcttgcc tacttggtcg tgaaaacgct gatcaacgcg 480
 actcaactcc tcaaattgct tgccaaattg gcggagttgg tcgcggccgc cattgcggac 540
 atcatttcgg atgtggcgga catcatcaag ggcaccctcg gagaagtgtg ggagttcatc 600
 acaaacgcgc tcaacggcct gaaagagctt tgggacaagc tcacgggggtg ggtgaccgga 660
 ctgtttcttc gaggggtggtc gaacctggag tccttctttg cgggcgtccc cggcttgacc 720
 ggcgcgacca gcggcttgtc gcaagtgact ggcttggtcg gtgcggcccg tctgtccgca 780
 tcgtcgggct tggctcacgc ggatagcctg gcgagctcag ccagcttgcc cgccctggcc 840
 ggcattgggg gcgggtcccg ttttgggggc ttgccgagcc tggtcaggt ccatgccgcc 900
 tcaactcggc aggcgctacg gccccgagct gatggcccg tggcgccgc tgccgagcag 960
 gtcggcgggc agtcgcagct ggtctccgcg cagggttccc aaggtatggg cggacccgta 1020
 ggcattgggc gcatgcaccc ctcttcgggg gcgtcgaaag ggacgacgac gaagaagtac 1080
 tcggaaggcg cggcggcggg cactgaagac gccgagcgcg cgccagtcga agctgacgcg 1140
 ggcgggtggc aaaagggtgct ggtacgaaac gtcgtctaa 1179

<210> 3
 <211> 392
 <212> PRT
 <213> Mycobacterium tuberculosis

<220>
 <221> MISC_FEATURE
 <223> Strain CDC1551

<400> 3
 Met Ser Arg Ala Phe Ile Ile Asp Pro Thr Ile Ser Ala Ile Asp Gly
 1 5 10 15
 Leu Tyr Asp Leu Leu Gly Ile Gly Ile Pro Asn Gln Gly Gly Ile Leu
 20 25 30
 Tyr Ser Ser Leu Glu Tyr Phe Glu Lys Ala Leu Glu Glu Leu Ala Ala

35

40

45

Ala Phe Pro Gly Asp Gly Trp Leu Gly Ser Ala Ala Asp Lys Tyr Ala
 50 55 60

Gly Lys Asn Arg Asn His Val Asn Phe Phe Gln Glu Leu Ala Asp Leu
 65 70 75 80

Asp Arg Gln Leu Ile Ser Leu Ile His Asp Gln Ala Asn Ala Val Gln
 85 90 95

Thr Thr Arg Asp Ile Leu Glu Gly Ala Lys Lys Gly Leu Glu Phe Val
 100 105 110

Arg Pro Val Ala Val Asp Leu Thr Tyr Ile Pro Val Val Gly His Ala
 115 120 125

Leu Ser Ala Ala Phe Gln Ala Pro Phe Cys Ala Gly Ala Met Ala Val
 130 135 140

Val Gly Gly Ala Leu Ala Tyr Leu Val Val Lys Thr Leu Ile Asn Ala
 145 150 155 160

Thr Gln Leu Leu Lys Leu Leu Ala Lys Leu Ala Glu Leu Val Ala Ala
 165 170 175

Ala Ile Ala Asp Ile Ile Ser Asp Val Ala Asp Ile Ile Lys Gly Ile
 180 185 190

Leu Gly Glu Val Trp Glu Phe Ile Thr Asn Ala Leu Asn Gly Leu Lys
 195 200 205

Glu Leu Trp Asp Lys Leu Thr Gly Trp Val Thr Gly Leu Phe Ser Arg
 210 215 220

Gly Trp Ser Asn Leu Glu Ser Phe Phe Ala Gly Val Pro Gly Leu Thr
 225 230 235 240

Gly Ala Thr Ser Gly Leu Ser Gln Val Thr Gly Leu Phe Gly Ala Ala
 245 250 255

Gly Leu Ser Ala Ser Ser Gly Leu Ala His Ala Asp Ser Leu Ala Ser
 260 265 270

Ser Ala Ser Leu Pro Ala Leu Ala Gly Ile Gly Gly Gly Ser Gly Phe
 275 280 285

Gly Gly Leu Pro Ser Leu Ala Gln Val His Ala Ala Ser Thr Arg Gln
 290 295 300

Ala Leu Arg Pro Arg Ala Asp Gly Pro Val Gly Ala Ala Ala Glu Gln

130

135

140

Val Gly Gly Ala Leu Ala Tyr Leu Val Val Lys Thr Leu Ile Asn Ala
145 150 155 160

Thr Gln Leu Leu Lys Leu Leu Ala Lys Leu Ala Glu Leu Val Ala Ala
165 170 175

Ala Ile Ala Asp Ile Ile Ser Asp Val Ala Asp Ile Ile Lys Gly Ile
180 185 190

Leu Gly Glu Val Trp Glu Phe Ile Thr Asn Ala Leu Asn Gly Leu Lys
195 200 205

Glu Leu Trp Asp Lys Leu Thr Gly Trp Val Thr Gly Leu Phe Ser Arg
210 215 220

Gly Trp Ser Asn Leu Glu Ser Phe Phe Ala Gly Val Pro Gly Leu Thr
225 230 235 240

Gly Ala Thr Ser Gly Leu Ser Gln Val Thr Gly Leu Phe Gly Ala Ala
245 250 255

Gly Leu Ser Ala Ser Ser Gly Leu Ala His Ala Asp Ser Leu Ala Ser
260 265 270

Ser Ala Ser Leu Pro Ala Leu Ala Gly Ile Gly Gly Gly Ser Gly Phe
275 280 285

Gly Gly Leu Pro Ser Leu Ala Gln Val His Ala Ala Ser Thr Arg Gln
290 295 300

Ala Leu Arg Pro Arg Ala Asp Gly Pro Val Gly Ala Ala Ala Glu Gln
305 310 315 320

Val Gly Gly Gln Ser Gln Leu Val Ser Ala Gln Gly Ser Gln Gly Met
325 330 335

Gly Gly Pro Val Gly Met Gly Gly Met His Pro Ser Ser Gly Ala Ser
340 345 350

Lys Gly Thr Thr Thr Lys Lys Tyr Ser Glu Gly Ala Ala Ala Gly Thr
355 360 365

Glu Asp Ala Glu Arg Ala Pro Val Glu Ala Asp Ala Gly Gly Gly Gln
370 375 380

Lys Val Leu Val Arg Asn Val Val
385 390

<210> 5

eof-seq1

<211> 392
 <212> PRT
 <213> Mycobacterium tuberculosis

<220>
 <221> MISC_FEATURE
 <223> Strain Haarlem A

<400> 5

Met Ser Arg Ala Phe Ile Ile Asp Pro Thr Ile Ser Ala Ile Asp Gly
 1 5 10 15

Leu Tyr Asp Leu Leu Gly Ile Gly Ile Pro Asn Gln Gly Gly Ile Leu
 20 25 30

Tyr Ser Ser Leu Glu Tyr Phe Glu Lys Ala Leu Glu Glu Leu Ala Ala
 35 40 45

Ala Phe Pro Gly Asp Gly Trp Leu Gly Ser Ala Ala Asp Lys Tyr Ala
 50 55 60

Gly Lys Asn Arg Asn His Val Asn Phe Phe Gln Glu Leu Ala Asp Leu
 65 70 75 80

Asp Arg Gln Leu Ile Ser Leu Ile His Asp Gln Ala Asn Ala Val Gln
 85 90 95

Thr Thr Arg Asp Ile Leu Glu Gly Ala Lys Lys Gly Leu Glu Phe Val
 100 105 110

Arg Pro Val Ala Val Asp Leu Thr Tyr Ile Pro Val Val Gly His Ala
 115 120 125

Leu Ser Ala Ala Phe Gln Ala Pro Phe Cys Ala Gly Ala Met Ala Val
 130 135 140

Val Gly Gly Ala Leu Ala Tyr Leu Val Val Lys Thr Leu Ile Asn Ala
 145 150 155 160

Thr Gln Leu Leu Lys Leu Leu Ala Lys Leu Ala Glu Leu Val Ala Ala
 165 170 175

Ala Ile Ala Asp Ile Ile Ser Asp Val Ala Asp Ile Ile Lys Gly Ile
 180 185 190

Leu Gly Glu Val Trp Glu Phe Ile Thr Asn Ala Leu Asn Gly Leu Lys
 195 200 205

Glu Leu Trp Asp Lys Leu Thr Gly Trp Val Thr Gly Leu Phe Ser Arg
 210 215 220

Gly Trp Ser Asn Leu Glu Ser Phe Phe Ala Gly Val Pro Gly Leu Thr
 Page 7

ecdf-seq1
60

50

55

Gly Lys Asn Arg Asn His Val Asn Phe Phe Gln Glu Leu Ala Asp Leu
65 70 75 80

Asp Arg Gln Leu Ile Ser Leu Ile His Asp Gln Ala Asn Ala Val Gln
85 90 95

Thr Thr Arg Asp Ile Leu Glu Gly Ala Lys Lys Gly Leu Glu Phe Val
100 105 110

Arg Pro Val Ala Val Asp Leu Thr Tyr Ile Pro Val Val Gly His Ala
115 120 125

Leu Ser Ala Ala Phe Gln Ala Pro Phe Cys Ala Gly Ala Met Ala Val
130 135 140

Val Gly Gly Ala Leu Ala Tyr Leu Val Val Lys Thr Leu Ile Asn Ala
145 150 155 160

Thr Gln Leu Leu Lys Leu Leu Ala Lys Leu Ala Glu Leu Val Ala Ala
165 170 175

Ala Ile Ala Asp Ile Ile Ser Asp Val Ala Asp Ile Ile Lys Gly Ile
180 185 190

Leu Gly Glu Val Trp Glu Phe Ile Thr Asn Ala Leu Asn Gly Leu Lys
195 200 205

Glu Leu Trp Asp Lys Leu Thr Gly Trp Val Thr Gly Leu Phe Ser Arg
210 215 220

Gly Trp Ser Asn Leu Glu Ser Phe Phe Ala Gly Val Pro Gly Leu Thr
225 230 235 240

Gly Ala Thr Ser Gly Leu Ser Gln Val Thr Gly Leu Phe Gly Ala Ala
245 250 255

Gly Leu Ser Ala Ser Ser Gly Leu Ala His Ala Asp Ser Leu Ala Ser
260 265 270

Ser Ala Ser Leu Pro Ala Leu Ala Gly Ile Gly Gly Gly Ser Gly Phe
275 280 285

Gly Gly Leu Pro Ser Leu Ala Gln Val His Ala Ala Ser Thr Arg Gln
290 295 300

Ala Leu Arg Pro Arg Ala Asp Gly Pro Val Gly Ala Ala Ala Glu Gln
305 310 315 320

Val Gly Gly Gln Ser Gln Leu Val Ser Ala Gln Gly Ser Gln Gly Met

325 eolf-seq1 335
330

Gly Gly Pro Val Gly Met Gly Gly Met His Pro Ser Ser Gly Ala Ser
340 345 350

Lys Gly Thr Thr Thr Lys Lys Tyr Ser Glu Gly Ala Ala Ala Gly Thr
355 360 365

Glu Asp Ala Glu Arg Ala Pro Val Glu Ala Asp Ala Gly Gly Gly Gln
370 375 380

Lys Val Leu Val Arg Asn Val Val
385 390

<210> 7
<211> 392
<212> PRT
<213> Mycobacterium bovis

<220>
<221> MISC_FEATURE
<223> Strain BCG

<400> 7

Met Ser Arg Val Phe Ile Ile Asp Pro Thr Ile Ser Ala Ile Asp Gly
1 5 10 15

Leu Tyr Asp Leu Leu Gly Ile Gly Ile Pro Asn Gln Gly Gly Ile Leu
20 25 30

Tyr Ser Ser Leu Glu Tyr Phe Glu Lys Ala Leu Glu Glu Leu Ala Ala
35 40 45

Ala Phe Pro Gly Asp Gly Trp Leu Gly Ser Ala Ala Asp Lys Tyr Ala
50 55 60

Gly Lys Asn Arg Asn His Val Asn Phe Phe Gln Glu Leu Ala Asp Leu
65 70 75 80

Asp Arg Gln Leu Ile Ser Leu Ile His Asp Gln Ala Asn Ala Val Gln
85 90 95

Thr Thr Arg Asp Ile Leu Glu Gly Ala Lys Lys Gly Leu Glu Phe Val
100 105 110

Arg Pro Val Ala Val Asp Leu Thr Tyr Ile Pro Val Val Gly His Ala
115 120 125

Leu Ser Ala Ala Phe Gln Ala Pro Phe Cys Ala Gly Ala Met Ala Val
130 135 140

Val Gly Gly Ala Leu Ala Tyr Leu Val Val Lys Thr Leu Ile Asn Ala
Page 10

145	150	eolf-seq1				160
		155				
Thr Gln Leu Leu Lys	Leu Leu Ala Lys	Leu Ala Glu Leu Val	Ala Ala			
	165	170	175			
Ala Ile Ala Asp Ile Ile Ser Asp	Val Ala Asp Ile Ile	Lys Gly Ile				
	180	185	190			
Leu Gly Glu Val Trp Glu Phe Ile Thr Asn Ala Leu Asn Gly Leu Lys						
	195	200	205			
Glu Leu Trp Asp Lys Leu Thr Gly Trp Val Thr Gly Leu Phe Ser Arg						
	210	215	220			
Gly Trp Ser Asn Leu Glu Ser Phe Phe Ala Gly Val Pro Gly Leu Thr						
	225	230	235			240
Gly Ala Thr Ser Gly Leu Ser Gln Val Thr Gly Leu Phe Gly Ala Ala						
	245	250				255
Gly Leu Ser Ala Ser Ser Gly Leu Ala His Ala Asp Ser Leu Ala Ser						
	260	265				270
Ser Ala Ser Leu Pro Ala Leu Ala Gly Ile Gly Gly Gly Ser Gly Phe						
	275	280				285
Gly Gly Leu Pro Ser Leu Ala Gln Val His Ala Ala Ser Thr Arg Gln						
	290	295				300
Ala Leu Arg Pro Arg Ala Asp Gly Pro Val Gly Ala Ala Ala Glu Gln						
	305	310				315
Val Gly Gly Gln Ser Gln Leu Val Ser Ala Gln Gly Ser Gln Gly Met						
	325	330				335
Gly Gly Pro Val Gly Met Gly Gly Met His Pro Ser Ser Gly Ala Ser						
	340	345				350
Lys Gly Thr Thr Thr Lys Lys Tyr Ser Glu Gly Ala Ala Ala Gly Thr						
	355	360				365
Glu Asp Ala Glu Arg Ala Pro Val Glu Ala Asp Ala Gly Gly Gly Gln						
	370	375				380
Lys Val Leu Val Arg Asn Val Val						
	385	390				

<210> 8
 <211> 110
 <212> PRT
 <213> Mycobacterium tuberculosis

eof-seq1

<220>

<221> mat_peptide

<222> (29)..(110)

<400> 8

Met Arg Leu Ser Leu Thr Ala Leu Ser Ala Gly Val Gly Ala Val Ala
-25 -20 -15

Met Ser Leu Thr Val Gly Ala Gly Val Ala Ser Ala Asp Pro Val Asp
-10 -5 -1 1

Ala Val Ile Asn Thr Thr Cys Asn Tyr Gly Gln Val Val Ala Ala Leu
5 10 15 20

Asn Ala Thr Asp Pro Gly Ala Ala Ala Gln Phe Asn Ala Ser Pro Val
25 30 35

Ala Gln Ser Tyr Leu Arg Asn Phe Leu Ala Ala Pro Pro Pro Gln Arg
40 45 50

Ala Ala Met Ala Ala Gln Leu Gln Ala Val Pro Gly Ala Ala Gln Tyr
55 60 65

Ile Gly Leu Val Glu Ser Val Ala Gly Ser Cys Asn Asn Tyr
70 75 80

<210> 9

<211> 97

<212> PRT

<213> Mycobacterium tuberculosis

<400> 9

Met Ser Leu Leu Asp Ala His Ile Pro Gln Leu Val Ala Ser Gln Ser
1 5 10 15

Ala Phe Ala Ala Lys Ala Gly Leu Met Arg His Thr Ile Gly Gln Ala
20 25 30

Glu Gln Ala Ala Met Ser Ala Gln Ala Phe His Gln Gly Glu Ser Ser
35 40 45

Ala Ala Phe Gln Ala Ala His Ala Arg Phe Val Ala Ala Ala Ala Lys
50 55 60

Val Asn Thr Leu Leu Asp Val Ala Gln Ala Asn Leu Gly Glu Ala Ala
65 70 75 80

Gly Thr Tyr Val Ala Ala Asp Ala Ala Ala Ala Ser Thr Tyr Thr Gly
85 90 95

Phe

eof-seq1

<210> 10
 <211> 94
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 10

Met Thr Ile Asn Tyr Gln Phe Gly Asp Val Asp Ala His Gly Ala Met
 1 5 10 15

Ile Arg Ala Gln Ala Ala Ser Leu Glu Ala Glu His Gln Ala Ile Val
 20 25 30

Arg Asp Val Leu Ala Ala Gly Asp Phe Trp Gly Gly Ala Gly Ser Val
 35 40 45

Ala Cys Gln Glu Phe Ile Thr Gln Leu Gly Arg Asn Phe Gln Val Ile
 50 55 60

Tyr Glu Gln Ala Asn Ala His Gly Gln Lys Val Gln Ala Ala Gly Asn
 65 70 75 80

Asn Met Ala Gln Thr Asp Ser Ala Val Gly Ser Ser Trp Ala
 85 90

<210> 11
 <211> 132
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 11

Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly Phe
 1 5 10 15

Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg Ser
 20 25 30

Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu Gly
 35 40 45

Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg Val
 50 55 60

Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp Val
 65 70 75 80

Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met Ala
 85 90 95

Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr Trp
 100 105 110

Gln Thr Lys Ser Gly Gly Thr Arg Thr ^{eolf-seq1} Gly Asn Val Thr Leu Ala Glu
 115 120 125

Gly Pro Pro Ala
 130

<210> 12
 <211> 195
 <212> PRT
 <213> Mycobacterium tuberculosis
 <400> 12

Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
 1 5 10 15

Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
 20 25 30

Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
 35 40 45

Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
 50 55 60

Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
 65 70 75 80

Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
 85 90 95

Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
 100 105 110

Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
 115 120 125

Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
 130 135 140

Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
 145 150 155 160

Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ser
 165 170 175

Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
 180 185 190

Ala Ala Ser
 195

<210> 13

eof-seq1

<211> 391
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 13

Met Val Asp Phe Gly Ala Leu Pro Pro Glu Ile Asn Ser Ala Arg Met
 1 5 10 15

Tyr Ala Gly Pro Gly Ser Ala Ser Leu Val Ala Ala Ala Gln Met Trp
 20 25 30

Asp Ser Val Ala Ser Asp Leu Phe Ser Ala Ala Ser Ala Phe Gln Ser
 35 40 45

Val Val Trp Gly Leu Thr Val Gly Ser Trp Ile Gly Ser Ser Ala Gly
 50 55 60

Leu Met Val Ala Ala Ala Ser Pro Tyr Val Ala Trp Met Ser Val Thr
 65 70 75 80

Ala Gly Gln Ala Glu Leu Thr Ala Ala Gln Val Arg Val Ala Ala Ala
 85 90 95

Ala Tyr Glu Thr Ala Tyr Gly Leu Thr Val Pro Pro Pro Val Ile Ala
 100 105 110

Glu Asn Arg Ala Glu Leu Met Ile Leu Ile Ala Thr Asn Leu Leu Gly
 115 120 125

Gln Asn Thr Pro Ala Ile Ala Val Asn Glu Ala Glu Tyr Gly Glu Met
 130 135 140

Trp Ala Gln Asp Ala Ala Ala Met Phe Gly Tyr Ala Ala Ala Thr Ala
 145 150 155 160

Thr Ala Thr Ala Thr Leu Leu Pro Phe Glu Glu Ala Pro Glu Met Thr
 165 170 175

Ser Ala Gly Gly Leu Leu Glu Gln Ala Ala Ala Val Glu Glu Ala Ser
 180 185 190

Asp Thr Ala Ala Ala Asn Gln Leu Met Asn Asn Val Pro Gln Ala Leu
 195 200 205

Gln Gln Leu Ala Gln Pro Thr Gln Gly Thr Thr Pro Ser Ser Lys Leu
 210 215 220

Gly Gly Leu Trp Lys Thr Val Ser Pro His Arg Ser Pro Ile Ser Asn
 225 230 235 240

Met Val Ser Met Ala Asn Asn His Met Ser Met Thr Asn Ser Gly Val
 245 250 255

ecolf-seq1

Ser Met Thr Asn Thr Leu Ser Ser Met Leu Lys Gly Phe Ala Pro Ala
260 265 270

Ala Ala Ala Gln Ala Val Gln Thr Ala Ala Gln Asn Gly Val Arg Ala
275 280 285

Met Ser Ser Leu Gly Ser Ser Leu Gly Ser Ser Gly Leu Gly Gly Gly
290 295 300

Val Ala Ala Asn Leu Gly Arg Ala Ala Ser Val Gly Ser Leu Ser Val
305 310 315 320

Pro Gln Ala Trp Ala Ala Ala Asn Gln Ala Val Thr Pro Ala Ala Arg
325 330 335

Ala Leu Pro Leu Thr Ser Leu Thr Ser Ala Ala Glu Arg Gly Pro Gly
340 345 350

Gln Met Leu Gly Gly Leu Pro Val Gly Gln Met Gly Ala Arg Ala Gly
355 360 365

Gly Gly Leu Ser Gly Val Leu Arg Val Pro Pro Arg Pro Tyr Val Met
370 375 380

Pro His Ser Pro Ala Ala Gly
385 390

<210> 14
<211> 423
<212> PRT
<213> Mycobacterium tuberculosis
<400> 14

Met Asp Phe Gly Leu Leu Pro Pro Glu Val Asn Ser Ser Arg Met Tyr
1 5 10 15

Ser Gly Pro Gly Pro Glu Ser Met Leu Ala Ala Ala Ala Ala Trp Asp
20 25 30

Gly Val Ala Ala Glu Leu Thr Ser Ala Ala Val Ser Tyr Gly Ser Val
35 40 45

Val Ser Thr Leu Ile Val Glu Pro Trp Met Gly Pro Ala Ala Ala Ala
50 55 60

Met Ala Ala Ala Ala Thr Pro Tyr Val Gly Trp Leu Ala Ala Thr Ala
65 70 75 80

Ala Leu Ala Lys Glu Thr Ala Thr Gln Ala Arg Ala Ala Ala Glu Ala
85 90 95

eo1F-seq1

Phe Gly Thr Ala Phe Ala Met Thr Val Pro Pro Ser Leu Val Ala Ala
 100 105 110
 Asn Arg Ser Arg Leu Met Ser Leu Val Ala Ala Asn Ile Leu Gly Gln
 115 120 125
 Asn Ser Ala Ala Ile Ala Ala Thr Gln Ala Glu Tyr Ala Glu Met Trp
 130 135 140
 Ala Gln Asp Ala Ala Val Met Tyr Ser Tyr Glu Gly Ala Ser Ala Ala
 145 150 155 160
 Ala Ser Ala Leu Pro Pro Phe Thr Pro Pro Val Gln Gly Thr Gly Pro
 165 170 175
 Ala Gly Pro Ala Ala Ala Ala Ala Thr Gln Ala Ala Gly Ala Gly
 180 185 190
 Ala Val Ala Asp Ala Gln Ala Thr Leu Ala Gln Leu Pro Pro Gly Ile
 195 200 205
 Leu Ser Asp Ile Leu Ser Ala Leu Ala Ala Asn Ala Asp Pro Leu Thr
 210 215 220
 Ser Gly Leu Leu Gly Ile Ala Ser Thr Leu Asn Pro Gln Val Gly Ser
 225 230 235 240
 Ala Gln Pro Ile Val Ile Pro Thr Pro Ile Gly Glu Leu Asp Val Ile
 245 250 255
 Ala Leu Tyr Ile Ala Ser Ile Ala Thr Gly Ser Ile Ala Leu Ala Ile
 260 265 270
 Thr Asn Thr Ala Arg Pro Trp His Ile Gly Leu Tyr Gly Asn Ala Gly
 275 280 285
 Gly Leu Gly Pro Thr Gln Gly His Pro Leu Ser Ser Ala Thr Asp Glu
 290 295 300
 Pro Glu Pro His Trp Gly Pro Phe Gly Gly Ala Ala Pro Val Ser Ala
 305 310 315 320
 Gly Val Gly His Ala Ala Leu Val Gly Ala Leu Ser Val Pro His Ser
 325 330 335
 Trp Thr Thr Ala Ala Pro Glu Ile Gln Leu Ala Val Gln Ala Thr Pro
 340 345 350
 Thr Phe Ser Ser Ser Ala Gly Ala Asp Pro Thr Ala Leu Asn Gly Met
 355 360 365

eof-seq1

Pro Ala Gly Leu Leu Ser Gly Met Ala Leu Ala Ser Leu Ala Ala Arg
370 375 380

Gly Thr Thr Gly Gly Gly Gly Thr Arg Ser Gly Thr Ser Thr Asp Gly
385 390 395 400

Gln Glu Asp Gly Arg Lys Pro Pro Val Val Val Ile Arg Glu Gln Pro
405 410 415

Pro Pro Gly Asn Pro Pro Arg
420

<210> 15
<211> 95
<212> PRT
<213> Mycobacterium tuberculosis

<220>
<221> INIT_MET
<222> (1)..(1)

<220>
<221> mat_peptide
<222> (2)..(95)

<400> 15

Met Thr Glu Gln Gln Trp Asn Phe Ala Gly Ile Glu Ala Ala Ala Ser
-1 1 5 10 15

Ala Ile Gln Gly Asn Val Thr Ser Ile His Ser Leu Leu Asp Glu Gly
20 25 30

Lys Gln Ser Leu Thr Lys Leu Ala Ala Ala Trp Gly Gly Ser Gly Ser
35 40 45

Glu Ala Tyr Gln Gly Val Gln Gln Lys Trp Asp Ala Thr Ala Thr Glu
50 55 60

Leu Asn Asn Ala Leu Gln Asn Leu Ala Arg Thr Ile Ser Glu Ala Gly
65 70 75

Gln Ala Met Ala Ser Thr Glu Gly Asn Val Thr Gly Met Phe Ala
80 85 90

<210> 16
<211> 338
<212> PRT
<213> Mycobacterium tuberculosis

<220>
<221> mat_peptide
<222> (43)..(338)

<400> 16

eo1f-seq1

Met Gln Leu Val Asp Arg Val Arg Gly Ala Val Thr Gly Met Ser Arg
-40 -35 -30

Arg Leu Val Val Gly Ala Val Gly Ala Ala Leu Val Ser Gly Leu Val
-25 -20 -15

Gly Ala Val Gly Gly Thr Ala Thr Ala Gly Ala Phe Ser Arg Pro Gly
-10 -5 -1 1 5

Leu Pro Val Glu Tyr Leu Gln Val Pro Ser Pro Ser Met Gly Arg Asp
10 15 20

Ile Lys Val Gln Phe Gln Ser Gly Gly Ala Asn Ser Pro Ala Leu Tyr
25 30 35

Leu Leu Asp Gly Leu Arg Ala Gln Asp Asp Phe Ser Gly Trp Asp Ile
40 45 50

Asn Thr Pro Ala Phe Glu Trp Tyr Asp Gln Ser Gly Leu Ser Val Val
55 60 65 70

Met Pro Val Gly Gly Gln Ser Ser Phe Tyr Ser Asp Trp Tyr Gln Pro
75 80 85

Ala Cys Gly Lys Ala Gly Cys Gln Thr Tyr Lys Trp Glu Thr Phe Leu
90 95 100

Thr Ser Glu Leu Pro Gly Trp Leu Gln Ala Asn Arg His Val Lys Pro
105 110 115

Thr Gly Ser Ala Val Val Gly Leu Ser Met Ala Ala Ser Ser Ala Leu
120 125 130

Thr Leu Ala Ile Tyr His Pro Gln Gln Phe Val Tyr Ala Gly Ala Met
135 140 145 150

Ser Gly Leu Leu Asp Pro Ser Gln Ala Met Gly Pro Thr Leu Ile Gly
155 160 165

Leu Ala Met Gly Asp Ala Gly Gly Tyr Lys Ala Ser Asp Met Trp Gly
170 175 180

Pro Lys Glu Asp Pro Ala Trp Gln Arg Asn Asp Pro Leu Leu Asn Val
185 190 195

Gly Lys Leu Ile Ala Asn Asn Thr Arg Val Trp Val Tyr Cys Gly Asn
200 205 210

Gly Lys Pro Ser Asp Leu Gly Gly Asn Asn Leu Pro Ala Lys Phe Leu
215 220 225 230

ecdf-seq1

Glu Gly Phe Val Arg Thr Ser Asn Ile Lys Phe Gln Asp Ala Tyr Asn
235 240 245

Ala Gly Gly Gly His Asn Gly Val Phe Asp Phe Pro Asp Ser Gly Thr
250 255 260

His Ser Trp Glu Tyr Trp Gly Ala Gln Leu Asn Ala Met Lys Pro Asp
265 270 275

Leu Gln Arg Ala Leu Gly Ala Thr Pro Asn Thr Gly Pro Ala Pro Gln
280 285 290

Gly Ala
295

<210> 17
<211> 325
<212> PRT
<213> Mycobacterium tuberculosis

<220>
<221> mat_peptide
<222> (41)..(325)

<400> 17

Met Thr Asp Val Ser Arg Lys Ile Arg Ala Trp Gly Arg Arg Leu Met
-40 -35 -30 -25

Ile Gly Thr Ala Ala Ala Val Val Leu Pro Gly Leu Val Gly Leu Ala
-20 -15 -10

Gly Gly Ala Ala Thr Ala Gly Ala Phe Ser Arg Pro Gly Leu Pro Val
-5 -1 1 5

Glu Tyr Leu Gln Val Pro Ser Pro Ser Met Gly Arg Asp Ile Lys Val
10 15 20

Gln Phe Gln Ser Gly Gly Asn Asn Ser Pro Ala Val Tyr Leu Leu Asp
25 30 35 40

Gly Leu Arg Ala Gln Asp Asp Tyr Asn Gly Trp Asp Ile Asn Thr Pro
45 50 55

Ala Phe Glu Trp Tyr Tyr Gln Ser Gly Leu Ser Ile Val Met Pro Val
60 65 70

Gly Gly Gln Ser Ser Phe Tyr Ser Asp Trp Tyr Ser Pro Ala Cys Gly
75 80 85

Lys Ala Gly Cys Gln Thr Tyr Lys Trp Glu Thr Phe Leu Thr Ser Glu
90 95 100

eof-seq1

Leu Pro Gln Trp Leu Ser Ala Asn Arg Ala Val Lys Pro Thr Gly Ser
105 110 115 120

Ala Ala Ile Gly Leu Ser Met Ala Gly Ser Ser Ala Met Ile Leu Ala
125 130 135

Ala Tyr His Pro Gln Gln Phe Ile Tyr Ala Gly Ser Leu Ser Ala Leu
140 145 150

Leu Asp Pro Ser Gln Gly Met Gly Pro Ser Leu Ile Gly Leu Ala Met
155 160 165

Gly Asp Ala Gly Gly Tyr Lys Ala Ala Asp Met Trp Gly Pro Ser Ser
170 175 180

Asp Pro Ala Trp Glu Arg Asn Asp Pro Thr Gln Gln Ile Pro Lys Leu
185 190 195 200

Val Ala Asn Asn Thr Arg Leu Trp Val Tyr Cys Gly Asn Gly Thr Pro
205 210 215

Asn Glu Leu Gly Gly Ala Asn Ile Pro Ala Glu Phe Leu Glu Asn Phe
220 225 230

Val Arg Ser Ser Asn Leu Lys Phe Gln Asp Ala Tyr Asn Ala Ala Gly
235 240 245

Gly His Asn Ala Val Phe Asn Phe Pro Pro Asn Gly Thr His Ser Trp
250 255 260

Glu Tyr Trp Gly Ala Gln Leu Asn Ala Met Lys Gly Asp Leu Gln Ser
265 270 275 280

Ser Leu Gly Ala Gly
285

<210> 18
<211> 144
<212> PRT
<213> Mycobacterium tuberculosis

<220>
<221> INIT_MET
<222> (1)..(1)

<220>
<221> mat_peptide
<222> (2)..(144)

<400> 18

Met Ala Thr Thr Leu Pro Val Gln Arg His Pro Arg Ser Leu Phe Pro
-1 1 5 10 15

eo1f-seq1

Glu Phe Ser Glu Leu Phe Ala Ala Phe Pro Ser Phe Ala Gly Leu Arg
20 25 30

Pro Thr Phe Asp Thr Arg Leu Met Arg Leu Glu Asp Glu Met Lys Glu
35 40 45

Gly Arg Tyr Glu Val Arg Ala Glu Leu Pro Gly Val Asp Pro Asp Lys
50 55 60

Asp Val Asp Ile Met Val Arg Asp Gly Gln Leu Thr Ile Lys Ala Glu
65 70 75

Arg Thr Glu Gln Lys Asp Phe Asp Gly Arg Ser Glu Phe Ala Tyr Gly
80 85 90 95

Ser Phe Val Arg Thr Val Ser Leu Pro Val Gly Ala Asp Glu Asp Asp
100 105 110

Ile Lys Ala Thr Tyr Asp Lys Gly Ile Leu Thr Val Ser Val Ala Val
115 120 125

Ser Glu Gly Lys Pro Thr Glu Lys His Ile Gln Ile Arg Ser Thr Asn
130 135 140

<210> 19
<211> 228
<212> PRT
<213> Mycobacterium tuberculosis

<220>
<221> mat_peptide
<222> (24)..(228)

<400> 19

Met Arg Ile Lys Ile Phe Met Leu Val Thr Ala Val Val Leu Leu Cys
-20 -15 -10

Cys Ser Gly Val Ala Thr Ala Ala Pro Lys Thr Tyr Cys Glu Glu Leu
-5 -1 1 5

Lys Gly Thr Asp Thr Gly Gln Ala Cys Gln Ile Gln Met Ser Asp Pro
10 15 20 25

Ala Tyr Asn Ile Asn Ile Ser Leu Pro Ser Tyr Tyr Pro Asp Gln Lys
30 35 40

Ser Leu Glu Asn Tyr Ile Ala Gln Thr Arg Asp Lys Phe Leu Ser Ala
45 50 55

Ala Thr Ser Ser Thr Pro Arg Glu Ala Pro Tyr Glu Leu Asn Ile Thr
60 65 70

eo1f-seq1

Ser Ala Thr Tyr Gln Ser Ala Ile Pro Pro Arg Gly Thr Gln Ala Val
75 80 85

Val Leu Lys Val Tyr Gln Asn Ala Gly Gly Thr His Pro Thr Thr Thr
90 95 100 105

Tyr Lys Ala Phe Asp Trp Asp Gln Ala Tyr Arg Lys Pro Ile Thr Tyr
110 115 120

Asp Thr Leu Trp Gln Ala Asp Thr Asp Pro Leu Pro Val Val Phe Pro
125 130 135

Ile Val Gln Gly Glu Leu Ser Lys Gln Thr Gly Gln Gln Val Ser Ile
140 145 150

Ala Pro Asn Ala Gly Leu Asp Pro Val Asn Tyr Gln Asn Phe Ala Val
155 160 165

Thr Asn Asp Gly Val Ile Phe Phe Phe Asn Pro Gly Glu Leu Leu Pro
170 175 180 185

Glu Ala Ala Gly Pro Thr Gln Val Leu Val Pro Arg Ser Ala Ile Asp
190 195 200

Ser Met Leu Ala
205

<210> 20
<211> 355
<212> PRT
<213> Mycobacterium tuberculosis

<220>
<221> mat_peptide
<222> (33)..(355)

<400> 20

Met Ser Asn Ser Arg Arg Arg Ser Leu Arg Trp Ser Trp Leu Leu Ser
-30 -25 -20

Val Leu Ala Ala Val Gly Leu Gly Leu Ala Thr Ala Pro Ala Gln Ala
-15 -10 -5 -1

Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
1 5 10 15

Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
20 25 30

Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
35 40 45

eof- seq1

Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
50 55 60

Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
65 70 75 80

Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
85 90 95

Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
100 105 110

Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
115 120 125

Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
130 135 140

Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
145 150 155 160

Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ser
165 170 175

Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
180 185 190

Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly Phe Ala
195 200 205

Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg Ser Gly
210 215 220

Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu Gly Leu
225 230 235 240

Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg Val Val
245 250 255

Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp Val Ile
260 265 270

Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met Ala Asp
275 280 285

Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr Trp Gln
290 295 300

Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala Glu Gly
305 310 315 320

Pro Pro Ala

<210> 21
 <211> 323
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Ser/Ala mutant of mature Mtb32A

<400> 21

Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
 1 5 10 15

Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
 20 25 30

Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
 35 40 45

Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
 50 55 60

Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
 65 70 75 80

Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
 85 90 95

Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
 100 105 110

Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
 115 120 125

Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
 130 135 140

Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
 145 150 155 160

Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ala
 165 170 175

Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
 180 185 190

Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly Phe Ala
 195 200 205

Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg Ser Gly
 Page 25

[illegible]

```
<210> 22
<211> 96
<212> PRT
<213> Mycobacterium tuberculosis
<400> 22
```

Met Ser Gln Ile Met Tyr Asn Tyr Pro Ala Met Leu Gly His Ala Gly
1 5 10 15

Asp Met Ala Gly Tyr Ala Gly Thr Leu Gln Ser Leu Gly Ala Glu Ile
20 25 30

Ala Val Gln Gln Ala Ala Leu Gln Ser Ala Trp Gln Gly Asp Thr Gly
35 40 45

Ile Thr Tyr Gln Ala Trp Gln Ala Gln Trp Asn Gln Ala Met Glu Asp
50 55 60

Leu Val Arg Ala Tyr His Ala Met Ser Ser Thr His Glu Ala Asn Thr
65 70 75 80

Met Ala Met Met Ala Arg Asp Thr Ala Glu Ala Ala Lys Trp Gly Gly
85 90 95

```
<210> 23
<211> 723
<212> PRT
<213> Artificial sequence
<220>
```

<223> Mtb72f

<400> 23

Met Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly
 1 5 10 15

Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg
 20 25 30

Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu
 35 40 45

Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg
 50 55 60

Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp
 65 70 75 80

Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met
 85 90 95

Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr
 100 105 110

Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala
 115 120 125

Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu Pro Pro
 130 135 140

Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly Pro Gly Ser Ala Ser Leu
 145 150 155 160

Val Ala Ala Ala Gln Met Trp Asp Ser Val Ala Ser Asp Leu Phe Ser
 165 170 175

Ala Ala Ser Ala Phe Gln Ser Val Val Trp Gly Leu Thr Val Gly Ser
 180 185 190

Trp Ile Gly Ser Ser Ala Gly Leu Met Val Ala Ala Ala Ser Pro Tyr
 195 200 205

Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr Ala Ala
 210 215 220

Gln Val Arg Val Ala Ala Ala Ala Tyr Glu Thr Ala Tyr Gly Leu Thr
 225 230 235 240

Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met Ile Leu
 245 250 255

eolf-seq1

Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala Val Asn
260 265 270

Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Ala Met Phe
275 280 285

Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu Pro Phe
290 295 300

Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu Gln Ala
305 310 315 320

Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln Leu Met
325 330 335

Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr Gln Gly
340 345 350

Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val Ser Pro
355 360 365

His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn His Met
370 375 380

Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser Ser Met
385 390 395 400

Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln Thr Ala
405 410 415

Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
420 425 430

Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala
435 440 445

Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln
450 455 460

Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser
465 470 475 480

Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro Val Gly
485 490 495

Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu Arg Val
500 505 510

Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly Asp Ile
515 520 525

colf-seq1

Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
530 535 540

Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
545 550 555 560

Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
565 570 575

Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
580 585 590

Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
595 600 605

Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
610 615 620

Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
625 630 635 640

Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
645 650 655

Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
660 665 670

Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
675 680 685

Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ser
690 695 700

Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
705 710 715 720

Ala Ala Ser

<210> 24
<211> 723
<212> PRT
<213> Artificial Sequence

<220>
<223> M72

<400> 24

Met Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly
1 5 10 15

Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg
20 25 30

colf-seq1

Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu
35 40 45

Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg
50 55 60

Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp
65 70 75 80

Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met
85 90 95

Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr
100 105 110

Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala
115 120 125

Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu Pro Pro
130 135 140

Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly Pro Gly Ser Ala Ser Leu
145 150 155 160

Val Ala Ala Ala Gln Met Trp Asp Ser Val Ala Ser Asp Leu Phe Ser
165 170 175

Ala Ala Ser Ala Phe Gln Ser Val Val Trp Gly Leu Thr Val Gly Ser
180 185 190

Trp Ile Gly Ser Ser Ala Gly Leu Met Val Ala Ala Ala Ser Pro Tyr
195 200 205

Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr Ala Ala
210 215 220

Gln Val Arg Val Ala Ala Ala Ala Tyr Glu Thr Ala Tyr Gly Leu Thr
225 230 235 240

Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met Ile Leu
245 250 255

Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala Val Asn
260 265 270

Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Ala Met Phe
275 280 285

Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu Pro Phe
290 295 300

eof-seq1

Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu Gln Ala
 305 310 315 320
 Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln Leu Met
 325 330 335
 Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr Gln Gly
 340 345 350
 Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val Ser Pro
 355 360 365
 His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn His Met
 370 375 380
 Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser Ser Met
 385 390 395 400
 Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln Thr Ala
 405 410 415
 Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
 420 425 430
 Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala
 435 440 445
 Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln
 450 455 460
 Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser
 465 470 475 480
 Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro Val Gly
 485 490 495
 Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu Arg Val
 500 505 510
 Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly Asp Ile
 515 520 525
 Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
 530 535 540
 Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
 545 550 555 560
 Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
 565 570 575

ecdf-seq1

Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
580 585 590

Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
595 600 605

Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
610 615 620

Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
625 630 635 640

Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
645 650 655

Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
660 665 670

Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
675 680 685

Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ala
690 695 700

Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
705 710 715 720

Ala Ala Ser

<210> 25
<211> 702
<212> PRT
<213> Artificial Sequence

<220>
<223> Mtb71f

<400> 25

Asp Pro Val Asp Ala Val Ile Asn Thr Thr Cys Asn Tyr Gly Gln Val
1 5 10 15

Val Ala Ala Leu Asn Ala Thr Asp Pro Gly Ala Ala Ala Gln Phe Asn
20 25 30

Ala Ser Pro Val Ala Gln Ser Tyr Leu Arg Asn Phe Leu Ala Ala Pro
35 40 45

Pro Pro Gln Arg Ala Ala Met Ala Ala Gln Leu Gln Ala Val Pro Gly
50 55 60

eolf-seq1

Ala Ala Gln Tyr Ile Gly Leu Val Glu Ser Val Ala Gly Ser Cys Asn
65 70 75 80

Asn Tyr Glu Leu Met Thr Ile Asn Tyr Gln Phe Gly Asp Val Asp Ala
85 90 95

His Gly Ala Met Ile Arg Ala Gln Ala Ala Ser Leu Glu Ala Glu His
100 105 110

Gln Ala Ile Val Arg Asp Val Leu Ala Ala Gly Asp Phe Trp Gly Gly
115 120 125

Ala Gly Ser Val Ala Cys Gln Glu Phe Ile Thr Gln Leu Gly Arg Asn
130 135 140

Phe Gln Val Ile Tyr Glu Gln Ala Asn Ala His Gly Gln Lys Val Gln
145 150 155 160

Ala Ala Gly Asn Asn Met Ala Gln Thr Asp Ser Ala Val Gly Ser Ser
165 170 175

Trp Ala Thr Ser Met Ser Leu Leu Asp Ala His Ile Pro Gln Leu Val
180 185 190

Ala Ser Gln Ser Ala Phe Ala Ala Lys Ala Gly Leu Met Arg His Thr
195 200 205

Ile Gly Gln Ala Glu Gln Ala Ala Met Ser Ala Gln Ala Phe His Gln
210 215 220

Gly Glu Ser Ser Ala Ala Phe Gln Ala Ala His Ala Arg Phe Val Ala
225 230 235 240

Ala Ala Ala Lys Val Asn Thr Leu Leu Asp Val Ala Gln Ala Asn Leu
245 250 255

Gly Glu Ala Ala Gly Thr Tyr Val Ala Ala Asp Ala Ala Ala Ala Ser
260 265 270

Thr Tyr Thr Gly Phe Asp Ile Met Asp Phe Gly Leu Leu Pro Pro Glu
275 280 285

Val Asn Ser Ser Arg Met Tyr Ser Gly Pro Gly Pro Glu Ser Met Leu
290 295 300

Ala Ala Ala Ala Ala Trp Asp Gly Val Ala Ala Glu Leu Thr Ser Ala
305 310 315 320

Ala Val Ser Tyr Gly Ser Val Val Ser Thr Leu Ile Val Glu Pro Trp
325 330 335

eolr-seq1

Met Gly Pro Ala Ala Ala Ala Met Ala Ala Ala Ala Thr Pro Tyr Val
340 345 350

Gly Trp Leu Ala Ala Thr Ala Ala Leu Ala Lys Glu Thr Ala Thr Gln
355 360 365

Ala Arg Ala Ala Ala Glu Ala Phe Gly Thr Ala Phe Ala Met Thr Val
370 375 380

Pro Pro Ser Leu Val Ala Ala Asn Arg Ser Arg Leu Met Ser Leu Val
385 390 395 400

Ala Ala Asn Ile Leu Gly Gln Asn Ser Ala Ala Ile Ala Ala Thr Gln
405 410 415

Ala Glu Tyr Ala Glu Met Trp Ala Gln Asp Ala Ala Val Met Tyr Ser
420 425 430

Tyr Glu Gly Ala Ser Ala Ala Ala Ser Ala Leu Pro Pro Phe Thr Pro
435 440 445

Pro Val Gln Gly Thr Gly Pro Ala Gly Pro Ala Ala Ala Ala Ala Ala
450 455 460

Thr Gln Ala Ala Gly Ala Gly Ala Val Ala Asp Ala Gln Ala Thr Leu
465 470 475 480

Ala Gln Leu Pro Pro Gly Ile Leu Ser Asp Ile Leu Ser Ala Leu Ala
485 490 495

Ala Asn Ala Asp Pro Leu Thr Ser Gly Leu Leu Gly Ile Ala Ser Thr
500 505 510

Leu Asn Pro Gln Val Gly Ser Ala Gln Pro Ile Val Ile Pro Thr Pro
515 520 525

Ile Gly Glu Leu Asp Val Ile Ala Leu Tyr Ile Ala Ser Ile Ala Thr
530 535 540

Gly Ser Ile Ala Leu Ala Ile Thr Asn Thr Ala Arg Pro Trp His Ile
545 550 555 560

Gly Leu Tyr Gly Asn Ala Gly Gly Leu Gly Pro Thr Gln Gly His Pro
565 570 575

Leu Ser Ser Ala Thr Asp Glu Pro Glu Pro His Trp Gly Pro Phe Gly
580 585 590

Gly Ala Ala Pro Val Ser Ala Gly Val Gly His Ala Ala Leu Val Gly
595 600 605

Ala Leu Ser Val Pro His Ser Trp Thr Thr Ala Ala Pro Glu Ile Gln
610 615 620

Leu Ala Val Gln Ala Thr Pro Thr Phe Ser Ser Ser Ala Gly Ala Asp
625 630 635 640

Pro Thr Ala Leu Asn Gly Met Pro Ala Gly Leu Leu Ser Gly Met Ala
645 650 655

Leu Ala Ser Leu Ala Ala Arg Gly Thr Thr Gly Gly Gly Gly Thr Arg
660 665 670

Ser Gly Thr Ser Thr Asp Gly Gln Glu Asp Gly Arg Lys Pro Pro Val
675 680 685

Val Val Ile Arg Glu Gln Pro Pro Pro Gly Asn Pro Pro Arg
690 695 700

<210> 26
<211> 920
<212> PRT
<213> Artificial sequence

<220>
<223> M72-Mtb9.9-Mtb9.8

<400> 26

Met Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly
1 5 10 15

Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg
20 25 30

Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu
35 40 45

Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg
50 55 60

Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp
65 70 75 80

Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met
85 90 95

Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr
100 105 110

Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala
115 120 125

Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu Pro Pro
130 135 140

eo1f-seq1

Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly Pro Gly Ser Ala Ser Leu
 145 150 155 160
 Val Ala Ala Ala Gln Met Trp Asp Ser Val Ala Ser Asp Leu Phe Ser
 165 170 175
 Ala Ala Ser Ala Phe Gln Ser Val Val Trp Gly Leu Thr Val Gly Ser
 180 185 190
 Trp Ile Gly Ser Ser Ala Gly Leu Met Val Ala Ala Ala Ser Pro Tyr
 195 200 205
 Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr Ala Ala
 210 215 220
 Gln Val Arg Val Ala Ala Ala Tyr Glu Thr Ala Tyr Gly Leu Thr
 225 230 235 240
 Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met Ile Leu
 245 250 255
 Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala Val Asn
 260 265 270
 Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Ala Met Phe
 275 280 285
 Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu Pro Phe
 290 295 300
 Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu Gln Ala
 305 310 315 320
 Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln Leu Met
 325 330 335
 Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr Gln Gly
 340 345 350
 Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val Ser Pro
 355 360 365
 His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn His Met
 370 375 380
 Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser Ser Met
 385 390 395 400
 Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln Thr Ala
 405 410 415

eo1f-seq1

Ala Gln Asn Gly val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
420 425 430

Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala
435 440 445

Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln
450 455 460

Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser
465 470 475 480

Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro Val Gly
485 490 495

Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu Arg Val
500 505 510

Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly Asp Ile
515 520 525

Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
530 535 540

Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
545 550 555 560

Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
565 570 575

Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
580 585 590

Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
595 600 605

Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
610 615 620

Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
625 630 635 640

Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
645 650 655

Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
660 665 670

Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
675 680 685

eof-seq1

Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ala
690 695 700

Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
705 710 715 720

Ala Ala Ser Ser Thr Met Thr Ile Asn Tyr Gln Phe Gly Asp Val Asp
725 730 735

Ala His Gly Ala Met Ile Arg Ala Gln Ala Ala Ser Leu Glu Ala Glu
740 745 750

His Gln Ala Ile Val Arg Asp Val Leu Ala Ala Gly Asp Phe Trp Gly
755 760 765

Gly Ala Gly Ser Val Ala Cys Gln Glu Phe Ile Thr Gln Leu Gly Arg
770 775 780

Asn Phe Gln Val Ile Tyr Glu Gln Ala Asn Ala His Gly Gln Lys Val
785 790 795 800

Gln Ala Ala Gly Asn Asn Met Ala Gln Thr Asp Ser Ala Val Gly Ser
805 810 815

Ser Trp Ala Thr Ser Met Ser Leu Leu Asp Ala His Ile Pro Gln Leu
820 825 830

Val Ala Ser Gln Ser Ala Phe Ala Ala Lys Ala Gly Leu Met Arg His
835 840 845

Thr Ile Gly Gln Ala Glu Gln Ala Ala Met Ser Ala Gln Ala Phe His
850 855 860

Gln Gly Glu Ser Ser Ala Ala Phe Gln Ala Ala His Ala Arg Phe Val
865 870 875 880

Ala Ala Ala Ala Lys Val Asn Thr Leu Leu Asp Val Ala Gln Ala Asn
885 890 895

Leu Gly Glu Ala Ala Gly Thr Tyr Val Ala Ala Asp Ala Ala Ala Ala
900 905 910

Ser Thr Tyr Thr Gly Phe Pro Trp
915 920

<210> 27
<211> 1010
<212> PRT
<213> Artificial Sequence
<220>

<223> M103

<400> 27

Met Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly
 1 5 10 15

Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg
 20 25 30

Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu
 35 40 45

Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg
 50 55 60

Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp
 65 70 75 80

Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met
 85 90 95

Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr
 100 105 110

Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala
 115 120 125

Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu Pro Pro
 130 135 140

Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly Pro Gly Ser Ala Ser Leu
 145 150 155 160

Val Ala Ala Ala Gln Met Trp Asp Ser Val Ala Ser Asp Leu Phe Ser
 165 170 175

Ala Ala Ser Ala Phe Gln Ser Val Val Trp Gly Leu Thr Val Gly Ser
 180 185 190

Trp Ile Gly Ser Ser Ala Gly Leu Met Val Ala Ala Ala Ser Pro Tyr
 195 200 205

Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr Ala Ala
 210 215 220

Gln Val Arg Val Ala Ala Ala Tyr Glu Thr Ala Tyr Gly Leu Thr
 225 230 235 240

Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met Ile Leu
 245 250 255

eolf-seql

Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala Val Asn
260 265 270

Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Ala Met Phe
275 280 285

Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu Pro Phe
290 295 300

Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu Gln Ala
305 310 315 320

Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln Leu Met
325 330 335

Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr Gln Gly
340 345 350

Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val Ser Pro
355 360 365

His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn His Met
370 375 380

Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser Ser Met
385 390 395 400

Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln Thr Ala
405 410 415

Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
420 425 430

Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala
435 440 445

Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln
450 455 460

Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser
465 470 475 480

Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro Val Gly
485 490 495

Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu Arg Val
500 505 510

Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly Asp Ile
515 520 525

eolf-seq1

Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
530 535 540

Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
545 550 555 560

Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
565 570 575

Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
580 585 590

Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
595 600 605

Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
610 615 620

Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
625 630 635 640

Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
645 650 655

Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
660 665 670

Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
675 680 685

Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ala
690 695 700

Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
705 710 715 720

Ala Ala Ser Ser Gly Phe Ser Arg Pro Gly Leu Pro Val Glu Tyr Leu
725 730 735

Gln Val Pro Ser Pro Ser Met Gly Arg Asp Ile Lys Val Gln Phe Gln
740 745 750

Ser Gly Gly Asn Asn Ser Pro Ala Val Tyr Leu Leu Asp Gly Leu Arg
755 760 765

Ala Gln Asp Asp Tyr Asn Gly Trp Asp Ile Asn Thr Pro Ala Phe Glu
770 775 780

Trp Tyr Tyr Gln Ser Gly Leu Ser Ile Val Met Pro Val Gly Gly Gln
785 790 795 800

eof-seq1

Ser Ser Phe Tyr Ser Asp Trp Tyr Ser Pro Ala Cys Gly Lys Ala Gly
805 810 815

Cys Gln Thr Tyr Lys Trp Glu Thr Phe Leu Thr Ser Glu Leu Pro Gln
820 825 830

Trp Leu Ser Ala Asn Arg Ala Val Lys Pro Thr Gly Ser Ala Ala Ile
835 840 845

Gly Leu Ser Met Ala Gly Ser Ser Ala Met Ile Leu Ala Ala Tyr His
850 855 860

Pro Gln Gln Phe Ile Tyr Ala Gly Ser Leu Ser Ala Leu Leu Asp Pro
865 870 875 880

Ser Gln Gly Met Gly Pro Ser Leu Ile Gly Leu Ala Met Gly Asp Ala
885 890 895

Gly Gly Tyr Lys Ala Ala Asp Met Trp Gly Pro Ser Ser Asp Pro Ala
900 905 910

Trp Glu Arg Asn Asp Pro Thr Gln Gln Ile Pro Lys Leu Val Ala Asn
915 920 925

Asn Thr Arg Leu Trp Val Tyr Cys Gly Asn Gly Thr Pro Asn Glu Leu
930 935 940

Gly Gly Ala Asn Ile Pro Ala Glu Phe Leu Glu Asn Phe Val Arg Ser
945 950 955 960

Ser Asn Leu Lys Phe Gln Asp Ala Tyr Asn Ala Ala Gly Gly His Asn
965 970 975

Ala Val Phe Asn Phe Pro Pro Asn Gly Thr His Ser Trp Glu Tyr Trp
980 985 990

Gly Ala Gln Leu Asn Ala Met Lys Gly Asp Leu Gln Ser Ser Leu Gly
995 1000 1005

Ala Gly
1010

<210> 28
<211> 1148
<212> PRT
<213> Artificial sequence

<220>
<223> M114

<400> 28

Met Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly
1 5 10 15

co1f seq1

Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg
 20 25 30
 Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu
 35 40 45
 Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg
 50 55 60
 Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp
 65 70 75 80
 Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met
 85 90 95
 Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr
 100 105 110
 Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala
 115 120 125
 Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu Pro Pro
 130 135 140
 Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly Pro Gly Ser Ala Ser Leu
 145 150 155 160
 Val Ala Ala Ala Gln Met Trp Asp Ser Val Ala Ser Asp Leu Phe Ser
 165 170 175
 Ala Ala Ser Ala Phe Gln Ser Val Val Trp Gly Leu Thr Val Gly Ser
 180 185 190
 Trp Ile Gly Ser Ser Ala Gly Leu Met Val Ala Ala Ala Ser Pro Tyr
 195 200 205
 Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr Ala Ala
 210 215 220
 Gln Val Arg Val Ala Ala Ala Ala Tyr Glu Thr Ala Tyr Gly Leu Thr
 225 230 235 240
 Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met Ile Leu
 245 250 255
 Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala Val Asn
 260 265 270
 Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Ala Met Phe
 275 280 285

eo1f-seq1

Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Leu Leu Pro Phe
290 295 300

Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu Gln Ala
305 310 315 320

Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln Leu Met
325 330 335

Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr Gln Gly
340 345 350

Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val Ser Pro
355 360 365

His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn His Met
370 375 380

Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser Ser Met
385 390 395 400

Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln Thr Ala
405 410 415

Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
420 425 430

Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala
435 440 445

Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln
450 455 460

Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser
465 470 475 480

Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro Val Gly
485 490 495

Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu Arg Val
500 505 510

Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly Asp Ile
515 520 525

Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
530 535 540

Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
545 550 555 560

colf-seq1

Asn Ile Asn Thr Lys 565 Leu Gly Tyr Asn Asn 570 Ala Val Gly Ala Gly Thr
 Gly Ile Val Ile 580 Asp Pro Asn Gly Val 585 Val Leu Thr Asn Asn 590 His Val
 Ile Ala Gly 595 Ala Thr Asp Ile Asn 600 Ala Phe Ser Val Gly 605 Ser Gly Gln
 Thr Tyr Gly Val Asp Val Val 615 Gly Tyr Asp Arg Thr 620 Gln Asp Val Ala
 Val 625 Leu Gln Leu Arg Gly 630 Ala Gly Gly Leu Pro 635 Ser Ala Ala Ile Gly 640
 Gly Gly Val Ala Val 645 Gly Glu Pro Val Val 650 Ala Met Gly Asn Ser Gly 655
 Gly Gln Gly Gly Thr Pro Arg Ala Val 665 Pro Gly Arg Val Val 670 Ala Leu
 Gly Gln Thr Val 675 Gln Ala Ser Asp 680 Ser Leu Thr Gly Ala 685 Glu Glu Thr
 Leu Asn Gly Leu Ile Gln Phe 695 Asp Ala Ala Ile Gln 700 Pro Gly Asp Ala
 Gly Gly Pro Val Val 710 Asn Gly Leu Gly Gln Val 715 Val Gly Met Asn Thr 720
 Ala Ala Ser Ser Thr 725 Met Asp Phe Gly Leu 730 Leu Pro Pro Glu Val 735 Asn
 Ser Ser Arg Met 740 Tyr Ser Gly Pro Gly 745 Pro Glu Ser Met Leu 750 Ala Ala
 Ala Ala Ala Trp Asp Gly Val 760 Ala Ala Glu Leu Thr Ser 765 Ala Ala Val
 Ser Tyr Gly Ser Val Val 775 Ser Thr Leu Ile Val Glu 780 Pro Trp Met Gly
 Pro Ala Ala Ala Ala Met 790 Ala Ala Ala Ala Thr 795 Pro Tyr Val Gly Trp 800
 Leu Ala Ala Thr 805 Ala Ala Leu Ala Lys Glu 810 Thr Ala Thr Gln Ala Arg 815
 Ala Ala Ala Glu 820 Ala Phe Gly Thr Ala 825 Phe Ala Met Thr Val 830 Pro Pro

eo1f-seq1

Ser Leu Val Ala Ala Asn Arg Ser Arg Leu Met Ser Leu Val Ala Ala
835 840 845

Asn Ile Leu Gly Gln Asn Ser Ala Ala Ile Ala Ala Thr Gln Ala Glu
850 855 860

Tyr Ala Glu Met Trp Ala Gln Asp Ala Ala Val Met Tyr Ser Tyr Glu
865 870 875 880

Gly Ala Ser Ala Ala Ala Ser Ala Leu Pro Pro Phe Thr Pro Pro Val
885 890 895

Gln Gly Thr Gly Pro Ala Gly Pro Ala Ala Ala Ala Ala Thr Gln
900 905 910

Ala Ala Gly Ala Gly Ala Val Ala Asp Ala Gln Ala Thr Leu Ala Gln
915 920 925

Leu Pro Pro Gly Ile Leu Ser Asp Ile Leu Ser Ala Leu Ala Ala Asn
930 935 940

Ala Asp Pro Leu Thr Ser Gly Leu Leu Gly Ile Ala Ser Thr Leu Asn
945 950 955 960

Pro Gln Val Gly Ser Ala Gln Pro Ile Val Ile Pro Thr Pro Ile Gly
965 970 975

Glu Leu Asp Val Ile Ala Leu Tyr Ile Ala Ser Ile Ala Thr Gly Ser
980 985 990

Ile Ala Leu Ala Ile Thr Asn Thr Ala Arg Pro Trp His Ile Gly Leu
995 1000 1005

Tyr Gly Asn Ala Gly Gly Leu Gly Pro Thr Gln Gly His Pro Leu
1010 1015 1020

Ser Ser Ala Thr Asp Glu Pro Glu Pro His Trp Gly Pro Phe Gly
1025 1030 1035

Gly Ala Ala Pro Val Ser Ala Gly Val Gly His Ala Ala Leu Val
1040 1045 1050

Gly Ala Leu Ser Val Pro His Ser Trp Thr Thr Ala Ala Pro Glu
1055 1060 1065

Ile Gln Leu Ala Val Gln Ala Thr Pro Thr Phe Ser Ser Ser Ala
1070 1075 1080

Gly Ala Asp Pro Thr Ala Leu Asn Gly Met Pro Ala Gly Leu Leu
1085 1090 1095

eof seq

Ser Gly Met Ala Leu Ala Ser Leu Ala Ala Arg Gly Thr Thr Gly
1100 1105 1110

Gly Gly Gly Thr Arg Ser Gly Thr Ser Thr Asp Gly Gln Glu Asp
1115 1120 1125

Gly Arg Lys Pro Pro Val Val Val Ile Arg Glu Gln Pro Pro Pro
1130 1135 1140

Gly Asn Pro Pro Arg
1145

<210> 29
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 29

Phe Ile Ile Asp Pro Thr Ile Ser Ala
1 5

<210> 30
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 30

Ile Leu Tyr Ser Ser Leu Glu Tyr Phe
1 5

<210> 31
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 31

Leu Glu Tyr Phe Glu Lys Ala Leu Glu
1 5

<210> 32
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 32

Tyr Ala Gly Lys Asn Arg Asn His Val
1 5

<210> 33
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 33

Leu Ile His Asp Gln Ala Asn Ala Val
1 5

<210> 34
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis
<400> 34

Phe Val Arg Pro Val Ala Val Asp Leu
1 5

<210> 35
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis
<400> 35

Leu Thr Tyr Ile Pro Val Val Gly His
1 5

<210> 36
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis
<400> 36

Tyr Ile Pro Val Val Gly His Ala Leu
1 5

<210> 37
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis
<400> 37

Tyr Leu Val Val Lys Thr Leu Ile Asn
1 5

<210> 38
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis
<400> 38

Leu Val Val Lys Thr Leu Ile Asn Ala
1 5

<210> 39
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis
<400> 39

Val Lys Thr Leu Ile Asn Ala Thr Gln

1

5

<210> 40
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 40

Leu Lys Leu Leu Ala Lys Leu Ala Glu
 1 5

<210> 41
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 41

Leu Val Ala Ala Ala Ile Ala Asp Ile
 1 5

<210> 42
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 42

Ile Ile Ser Asp Val Ala Asp Ile Ile
 1 5

<210> 43
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 43

Trp Glu Phe Ile Thr Asn Ala Leu Asn
 1 5

<210> 44
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 44

Leu Phe Gly Ala Ala Gly Leu Ser Ala
 1 5

<210> 45
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 45

Leu Ala His Ala Asp Ser Leu Ala Ser
 1 5

<210> 46
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 46

Leu Ala Ser Ser Ala Ser Leu Pro Ala
 1 5

<210> 47
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 47

Phe Gly Gly Leu Pro Ser Leu Ala Gln
 1 5

<210> 48
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 48

Phe Ile Ile Asp Pro Thr Ile Ser Ala
 1 5

<210> 49
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 49

Ile Ile Asp Pro Thr Ile Ser Ala Ile
 1 5

<210> 50
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 50

Pro Thr Ile Ser Ala Ile Asp Gly Leu
 1 5

<210> 51
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 51

Thr Ile Ser Ala Ile Asp Gly Leu Tyr
 1 5

<210> 52

<211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 52

Ser Ala Ile Asp Gly Leu Tyr Asp Leu
 1 5

<210> 53
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 53

Ala Ile Asp Gly Leu Tyr Asp Leu Leu
 1 5

<210> 54
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 54

Leu Tyr Asp Leu Leu Gly Ile Gly Ile
 1 5

<210> 55
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 55

Ile Pro Asn Gln Gly Gly Ile Leu Tyr
 1 5

<210> 56
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 56

Gly Ile Leu Tyr Ser Ser Leu Glu Tyr
 1 5

<210> 57
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 57

Tyr Ser Ser Leu Glu Tyr Phe Glu Lys
 1 5

<210> 58
 <211> 9
 <212> PRT

<213> Mycobacterium tuberculosis

<400> 58

Ser Leu Glu Tyr Phe Glu Lys Ala Leu
1 5

<210> 59

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 59

Tyr Phe Glu Lys Ala Leu Glu Glu Leu
1 5

<210> 60

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 60

Phe Glu Lys Ala Leu Glu Glu Leu Ala
1 5

<210> 61

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 61

Asn His Val Asn Phe Phe Gln Glu Leu
1 5

<210> 62

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 62

Glu Leu Ala Asp Leu Asp Arg Gln Leu
1 5

<210> 63

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 63

Leu Ala Asp Leu Asp Arg Gln Leu Ile
1 5

<210> 64

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 64

Asp Leu Asp Arg Gln Leu Ile Ser Leu
 1 5

<210> 65

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 65

Leu Asp Arg Gln Leu Ile Ser Leu Ile
 1 5

<210> 66

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 66

Ala Val Gln Thr Thr Arg Asp Ile Leu
 1 5

<210> 67

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 67

Glu Gly Ala Lys Lys Gly Leu Glu Phe
 1 5

<210> 68

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 68

Lys Gly Leu Glu Phe Val Arg Pro Val
 1 5

<210> 69

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 69

Gly Leu Glu Phe Val Arg Pro Val Ala
 1 5

<210> 70

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 70

Leu Glu Phe Val Arg Pro Val Ala Val
1 5

<210> 71
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 71

Phe Val Arg Pro Val Ala Val Asp Leu
1 5

<210> 72
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 72

Arg Pro Val Ala Val Asp Leu Thr Tyr
1 5

<210> 73
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 73

Ala Val Asp Leu Thr Tyr Ile Pro Val
1 5

<210> 74
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 74

Thr Tyr Ile Pro Val Val Gly His Ala
1 5

<210> 75
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 75

Tyr Ile Pro Val Val Gly His Ala Leu
1 5

<210> 76
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 76

Leu Ser Ala Ala Phe Gln Ala Pro Phe
1 5

<210> 77
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 77

Ser Ala Ala Phe Gln Ala Pro Phe Cys
 1 5

<210> 78
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 78

Ala Ala Phe Gln Ala Pro Phe Cys Ala
 1 5

<210> 79
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 79

Phe Gln Ala Pro Phe Cys Ala Gly Ala
 1 5

<210> 80
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 80

Ala Pro Phe Cys Ala Gly Ala Met Ala
 1 5

<210> 81
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 81

Pro Phe Cys Ala Gly Ala Met Ala Val
 1 5

<210> 82
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 82

Ala Met Ala Val Val Gly Gly Ala Leu
 1 5

<210> 83
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 83

Ala Val Val Gly Gly Ala Leu Ala Tyr
 1 5

<210> 84
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 84

Gly Ala Leu Ala Tyr Leu Val Val Lys
 1 5

<210> 85
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 85

Leu Ala Tyr Leu Val Val Lys Thr Leu
 1 5

<210> 86
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 86

Ala Tyr Leu Val Val Lys Thr Leu Ile
 1 5

<210> 87
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 87

Lys Thr Leu Ile Asn Ala Thr Gln Leu
 1 5

<210> 88
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 88

Thr Leu Ile Asn Ala Thr Gln Leu Leu
 1 5

<210> 89
 <211> 9

<212> PRT
 <213> Mycobacterium tuberculosis

<400> 89

Ile Asn Ala Thr Gln Leu Leu Lys Leu
 1 5

<210> 90
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 90

Asn Ala Thr Gln Leu Leu Lys Leu Leu
 1 5

<210> 91
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 91

Gln Leu Leu Lys Leu Leu Ala Lys Leu
 1 5

<210> 92
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 92

Lys Leu Leu Ala Lys Leu Ala Glu Leu
 1 5

<210> 93
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 93

Leu Leu Ala Lys Leu Ala Glu Leu Val
 1 5

<210> 94
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 94

Lys Leu Ala Glu Leu Val Ala Ala Ala
 1 5

<210> 95
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 95

Leu Ala Glu Leu Val Ala Ala Ala Ile
 1 5

<210> 96

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 96

Leu Val Ala Ala Ala Ile Ala Asp Ile
 1 5

<210> 97

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 97

Ala Ile Ala Asp Ile Ile Ser Asp Val
 1 5

<210> 98

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 98

Ile Ala Asp Ile Ile Ser Asp Val Ala
 1 5

<210> 99

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 99

Ile Ser Asp Val Ala Asp Ile Ile Lys
 1 5

<210> 100

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 100

Thr Leu Gly Glu Val Trp Glu Phe Ile
 1 5

<210> 101

<211> 9

<212> PRT

<213> Mycobacterium tuberculosis

<400> 101

Phe Ile Thr Asn Ala Leu Asn Gly Leu
1 5

<210> 102
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 102

Asn Ala Leu Asn Gly Leu Lys Glu Leu
1 5

<210> 103
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 103

Lys Leu Thr Gly Trp Val Thr Gly Leu
1 5

<210> 104
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 104

Leu Thr Gly Trp Val Thr Gly Leu Phe
1 5

<210> 105
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 105

Gly Trp Ser Asn Leu Glu Ser Phe Phe
1 5

<210> 106
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 106

Asn Leu Glu Ser Phe Phe Ala Gly Val
1 5

<210> 107
<211> 9
<212> PRT
<213> Mycobacterium tuberculosis

<400> 107

Ser Phe Phe Ala Gly Val Pro Gly Leu

1

5

<210> 108
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 108

Gly Leu Thr Gly Ala Thr Ser Gly Leu
 1 5

<210> 109
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 109

Leu Ser Gln Val Thr Gly Leu Phe Gly
 1 5

<210> 110
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 110

Ser Gln Val Thr Gly Leu Phe Gly Ala
 1 5

<210> 111
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 111

Leu Ser Ala Ser Ser Gly Leu Ala His
 1 5

<210> 112
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 112

Ala Ser Ser Gly Leu Ala His Ala Asp
 1 5

<210> 113
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 113

Ser Gly Leu Ala His Ala Asp Ser Leu
 1 5

<210> 114
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 114

Gly Leu Ala His Ala Asp Ser Leu Ala
 1 5

<210> 115
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 115

Ser Leu Ala Ser Ser Ala Ser Leu Pro
 1 5

<210> 116
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 116

Ala Ser Ser Ala Ser Leu Pro Ala Leu
 1 5

<210> 117
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 117

Ser Gly Phe Gly Gly Leu Pro Ser Leu
 1 5

<210> 118
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 118

Leu Pro Ser Leu Ala Gln Val His Ala
 1 5

<210> 119
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 119

His Ala Ala Ser Thr Arg Gln Ala Leu
 1 5

<210> 120

<211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 120

Ser Thr Arg Gln Ala Leu Arg Pro Arg
 1 5

<210> 121
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 121

Arg Pro Arg Ala Asp Gly Pro Val Gly
 1 5

<210> 122
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 122

Glu Gln Val Gly Gly Gln Ser Gln Leu
 1 5

<210> 123
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 123

Gly Ala Ser Lys Gly Thr Thr Thr Lys
 1 5

<210> 124
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 124

Ala Ser Lys Gly Thr Thr Thr Lys Lys
 1 5

<210> 125
 <211> 9
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 125

Lys Gly Thr Thr Thr Lys Lys Tyr Ser
 1 5

<210> 126
 <211> 9
 <212> PRT

<213> Mycobacterium tuberculosis

<400> 126

Thr Glu Asp Ala Glu Arg Ala Pro Val
1 5

<210> 127

<211> 20

<212> PRT

<213> Mycobacterium tuberculosis

<400> 127

Met Ser Arg Ala Phe Ile Ile Asp Pro Thr Ile Ser Ala Ile Asp Gly
1 5 10 15

Leu Tyr Asp Leu
20

<210> 128

<211> 20

<212> PRT

<213> Mycobacterium tuberculosis

<400> 128

Ile Asp Gly Leu Tyr Asp Leu Leu Gly Ile Gly Ile Pro Asn Gln Gly
1 5 10 15

Gly Ile Leu Tyr
20

<210> 129

<211> 20

<212> PRT

<213> Mycobacterium tuberculosis

<400> 129

Asn Gln Gly Gly Ile Leu Tyr Ser Ser Leu Glu Tyr Phe Glu Lys Ala
1 5 10 15

Leu Glu Glu Leu
20

<210> 130

<211> 20

<212> PRT

<213> Mycobacterium tuberculosis

<400> 130

Glu Lys Ala Leu Glu Glu Leu Ala Ala Ala Phe Pro Gly Asp Gly Trp
1 5 10 15

Leu Gly Ser Ala
20

eolf-seq1

<210> 131
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 131

Asp Gly Trp Leu Gly Ser Ala Ala Asp Lys Tyr Ala Gly Lys Asn Arg
 1 5 10 15

Asn His Val Asn
 20

<210> 132
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 132

Lys Asn Arg Asn His Val Asn Phe Phe Gln Glu Leu Ala Asp Leu Asp
 1 5 10 15

Arg Gln Leu Ile
 20

<210> 133
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 133

Asp Leu Asp Arg Gln Leu Ile Ser Leu Ile His Asp Gln Ala Asn Ala
 1 5 10 15

Val Gln Thr Thr
 20

<210> 134
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 134

Ala Asn Ala Val Gln Thr Thr Arg Asp Ile Leu Glu Gly Ala Lys Lys
 1 5 10 15

Gly Leu Glu Phe
 20

<210> 135
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 135

Ala Lys Lys Gly Leu Glu Phe Val Arg Pro Val Ala Val Asp Leu Thr

1 5 eol f-seq1 15
 10
 Tyr Ile Pro Val
 20
 <210> 136
 <211> 27
 <212> PRT
 <213> Mycobacterium tuberculosis
 <400> 136
 Phe Val Arg Pro Val Ala Val Asp Leu Thr Tyr Ile Pro Val Val Gly
 1 5 10 15
 His Ala Leu Ser Ala Ala Phe Gln Ala Pro Phe
 20 25
 <210> 137
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis
 <400> 137
 Ala Ala Phe Gln Ala Pro Phe Cys Ala Gly Ala Met Ala Val Val Gly
 1 5 10 15
 Gly Ala Leu Ala
 20
 <210> 138
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis
 <400> 138
 Val Val Gly Gly Ala Leu Ala Tyr Leu Val Val Lys Thr Leu Ile Asn
 1 5 10 15
 Ala Thr Gln Leu
 20
 <210> 139
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis
 <400> 139
 Leu Ile Asn Ala Thr Gln Leu Leu Lys Leu Leu Ala Lys Leu Ala Glu
 1 5 10 15
 Leu Val Ala Ala
 20
 <210> 140

eo1f-seq1

<211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 140

Leu Ala Glu Leu Val Ala Ala Ala Ile Ala Asp Ile Ile Ser Asp Val
 1 5 10 15

Ala Asp Ile Ile
 20

<210> 141
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 141

Ser Asp Val Ala Asp Ile Ile Lys Gly Thr Leu Gly Glu Val Trp Glu
 1 5 10 15

Phe Ile Thr Asn
 20

<210> 142
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 142

Val Trp Glu Phe Ile Thr Asn Ala Leu Asn Gly Leu Lys Glu Leu Trp
 1 5 10 15

Asp Lys Leu Thr
 20

<210> 143
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 143

Glu Leu Trp Asp Lys Leu Thr Gly Trp Val Thr Gly Leu Phe Ser Arg
 1 5 10 15

Gly Trp Ser Asn
 20

<210> 144
 <211> 20
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 144

Phe Ser Arg Gly Trp Ser Asn Leu Glu Ser Phe Phe Ala Gly Val Pro
 1 5 10 15

eof-seq1

Gly Leu Thr Gly
20

<210> 145
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis

<400> 145

Gly Val Pro Gly Leu Thr Gly Ala Thr Ser Gly Leu Ser Gln Val Thr
1 5 10 15

Gly Leu Phe Gly
20

<210> 146
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis

<400> 146

Gln Val Thr Gly Leu Phe Gly Ala Ala Gly Leu Ser Ala Ser Ser Gly
1 5 10 15

Leu Ala His Ala
20

<210> 147
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis

<400> 147

Ser Ser Gly Leu Ala His Ala Asp Ser Leu Ala Ser Ser Ala Ser Leu
1 5 10 15

Pro Ala Leu Ala
20

<210> 148
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis

<400> 148

Ala Ser Leu Pro Ala Leu Ala Gly Ile Gly Gly Gly Ser Gly Phe Gly
1 5 10 15

Gly Leu Pro Ser
20

<210> 149
<211> 20

eolf-seq1

<212> PRT
<213> Mycobacterium tuberculosis

<400> 149

Gly Phe Gly Gly Leu Pro Ser Leu Ala Gln Val His Ala Ala Ser Thr
1 5 10 15

Arg Gln Ala Leu
20

<210> 150
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis

<400> 150

Ala Ser Thr Arg Gln Ala Leu Arg Pro Arg Ala Asp Gly Pro Val Gly
1 5 10 15

Ala Ala Ala Glu
20

<210> 151
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis

<400> 151

Pro Val Gly Ala Ala Ala Glu Gln Val Gly Gly Gln Ser Gln Leu Val
1 5 10 15

Ser Ala Gln Gly
20

<210> 152
<211> 19
<212> PRT
<213> Mycobacterium tuberculosis

<400> 152

Gln Leu Val Ser Ala Gln Gly Ser Gln Gly Met Gly Gly Pro Val Gly
1 5 10 15

Met Gly Gly

<210> 153
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis

<400> 153

Pro Val Gly Met Gly Gly Met His Pro Ser Ser Gly Ala Ser Lys Gly
1 5 10 15

colf-seq1

Thr Thr Thr Lys
20

<210> 154
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis
<400> 154

Ser Lys Gly Thr Thr Thr Lys Lys Tyr Ser Glu Gly Ala Ala Ala Gly
1 5 10 15

Thr Glu Asp Ala
20

<210> 155
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis
<400> 155

Ala Ala Gly Thr Glu Asp Ala Glu Arg Ala Pro Val Glu Ala Asp Ala
1 5 10 15

Gly Gly Gly Gln
20

<210> 156
<211> 20
<212> PRT
<213> Mycobacterium tuberculosis
<400> 156

Arg Ala Pro Val Glu Ala Asp Ala Gly Gly Gly Gln Lys Val Leu Val
1 5 10 15

Arg Asn Val Val
20

<210> 157
<211> 1053
<212> PRT
<213> Mycobacterium tuberculosis
<400> 157

Met Asn Phe Ser Val Leu Pro Pro Glu Ile Asn Ser Ala Leu Ile Phe
1 5 10 15

Ala Gly Ala Gly Pro Glu Pro Met Ala Ala Ala Ala Thr Ala Trp Asp
20 25 30

Gly Leu Ala Met Glu Leu Ala Ser Ala Ala Ala Ser Phe Gly Ser Val
35 40 45

eof-seq1

```

rhr Ser Gly Leu Val Gly Gly Ala Trp Gln Gly Ala Ser Ser Ser Ala
 50          55          60

Met Ala Ala Ala Ala Ala Pro Tyr Ala Ala Trp Leu Ala Ala Ala Ala
65          70          75          80

Val Gln Ala Glu Gln Thr Ala Ala Gln Ala Ala Ala Met Ile Ala Glu
      85          90          95

Phe Glu Ala Val Lys Thr Ala Val Val Gln Pro Met Leu Val Ala Ala
      100        105        110

Asn Arg Ala Asp Leu Val Ser Leu Val Met Ser Asn Leu Phe Gly Gln
      115        120        125

Asn Ala Pro Ala Ile Ala Ala Ile Glu Ala Thr Tyr Glu Gln Met Trp
      130        135        140

Ala Ala Asp Val Ser Ala Met Ser Ala Tyr His Ala Gly Ala Ser Ala
      145        150        155        160

Ile Ala Ser Ala Leu Ser Pro Phe Ser Lys Pro Leu Gln Asn Leu Ala
      165        170        175

Gly Leu Pro Ala Trp Leu Ala Ser Gly Ala Pro Ala Ala Ala Met Thr
      180        185        190

Ala Ala Ala Gly Ile Pro Ala Leu Ala Gly Gly Pro Thr Ala Ile Asn
      195        200        205

Leu Gly Ile Ala Asn Val Gly Gly Gly Asn Val Gly Asn Ala Asn Asn
      210        215        220

Gly Leu Ala Asn Ile Gly Asn Ala Asn Leu Gly Asn Tyr Asn Phe Gly
      225        230        235        240

Ser Gly Asn Phe Gly Asn Ser Asn Ile Gly Ser Ala Ser Leu Gly Asn
      245        250        255

Asn Asn Ile Gly Phe Gly Asn Leu Gly Ser Asn Asn Val Gly Val Gly
      260        265        270

Asn Leu Gly Asn Leu Asn Thr Gly Phe Ala Asn Thr Gly Leu Gly Asn
      275        280        285

Phe Gly Phe Gly Asn Thr Gly Asn Asn Asn Ile Gly Ile Gly Leu Thr
      290        295        300

Gly Asn Asn Gln Ile Gly Ile Gly Gly Leu Asn Ser Gly Thr Gly Asn
      305        310        315        320

```

eof-seq1

Phe Gly Leu Phe Asn Ser Gly Ser Gly Asn Val Gly Phe Phe Asn Ser
 325 330 335
 Gly Asn Gly Asn Phe Gly Ile Gly Asn Ser Gly Asn Phe Asn Thr Gly
 340 345 350
 Gly Trp Asn Ser Gly His Gly Asn Thr Gly Phe Phe Asn Ala Gly Ser
 355 360 365
 Phe Asn Thr Gly Met Leu Asp Val Gly Asn Ala Asn Thr Gly Ser Leu
 370 375 380
 Asn Thr Gly Ser Tyr Asn Met Gly Asp Phe Asn Pro Gly Ser Ser Asn
 385 390 395 400
 Thr Gly Thr Phe Asn Thr Gly Asn Ala Asn Thr Gly Phe Leu Asn Ala
 405 410 415
 Gly Asn Ile Asn Thr Gly Val Phe Asn Ile Gly His Met Asn Asn Gly
 420 425 430
 Leu Phe Asn Thr Gly Asp Met Asn Asn Gly Val Phe Tyr Arg Gly Val
 435 440 445
 Gly Gln Gly Ser Leu Gln Phe Ser Ile Thr Thr Pro Asp Leu Thr Leu
 450 455 460
 Pro Pro Leu Gln Ile Pro Gly Ile Ser Val Pro Ala Phe Ser Leu Pro
 465 470 475 480
 Ala Ile Thr Leu Pro Ser Leu Asn Ile Pro Ala Ala Thr Thr Pro Ala
 485 490 495
 Asn Ile Thr Val Gly Ala Phe Ser Leu Pro Gly Leu Thr Leu Pro Ser
 500 505 510
 Leu Asn Ile Pro Ala Ala Thr Thr Pro Ala Asn Ile Thr Val Gly Ala
 515 520 525
 Phe Ser Leu Pro Gly Leu Thr Leu Pro Ser Leu Asn Ile Pro Ala Ala
 530 535 540
 Thr Thr Pro Ala Asn Ile Thr Val Gly Ala Phe Ser Leu Pro Gly Leu
 545 550 555 560
 Thr Leu Pro Ser Leu Asn Ile Pro Ala Ala Thr Thr Pro Ala Asn Ile
 565 570 575
 Thr Val Gly Ala Phe Ser Leu Pro Gly Leu Thr Leu Pro Ser Leu Asn
 580 585 590

eof-seq1

Ile Pro Ala Ala Thr Thr Pro Ala Asn Ile Thr Val Gly Ala Phe Ser
595 600 605

Leu Pro Gly Leu Thr Leu Pro Ser Leu Asn Ile Pro Ala Ala Thr Thr
610 615 620

Pro Ala Asn Ile Thr Val Ser Gly Phe Gln Leu Pro Pro Leu Ser Ile
625 630 635 640

Pro Ser Val Ala Ile Pro Pro Val Thr Val Pro Pro Ile Thr Val Gly
645 650 655

Ala Phe Asn Leu Pro Pro Leu Gln Ile Pro Glu Val Thr Ile Pro Gln
660 665 670

Leu Thr Ile Pro Ala Gly Ile Thr Ile Gly Gly Phe Ser Leu Pro Ala
675 680 685

Ile His Thr Gln Pro Ile Thr Val Gly Gln Ile Gly Val Gly Gln Phe
690 695 700

Gly Leu Pro Ser Ile Gly Trp Asp Val Phe Leu Ser Thr Pro Arg Ile
705 710 715 720

Thr Val Pro Ala Phe Gly Ile Pro Phe Thr Leu Gln Phe Gln Thr Asn
725 730 735

Val Pro Ala Leu Gln Pro Pro Gly Gly Gly Leu Ser Thr Phe Thr Asn
740 745 750

Gly Ala Leu Ile Phe Gly Glu Phe Asp Leu Pro Gln Leu Val Val His
755 760 765

Pro Tyr Thr Leu Thr Gly Pro Ile Val Ile Gly Ser Phe Phe Leu Pro
770 775 780

Ala Phe Asn Ile Pro Gly Ile Asp Val Pro Ala Ile Asn Val Asp Gly
785 790 795 800

Phe Thr Leu Pro Gln Ile Thr Thr Pro Ala Ile Thr Thr Pro Glu Phe
805 810 815

Ala Ile Pro Pro Ile Gly Val Gly Gly Phe Thr Leu Pro Gln Ile Thr
820 825 830

Thr Gln Glu Ile Ile Thr Pro Glu Leu Thr Ile Asn Ser Ile Gly Val
835 840 845

Gly Gly Phe Thr Leu Pro Gln Ile Thr Thr Pro Pro Ile Thr Thr Pro
850 855 860

eof-seq1

Pro Leu Thr Ile Asp Pro Ile Asn Leu Thr Gly Phe Thr Leu Pro Gln
865 870 875 880

Ile Thr Thr Pro Pro Ile Thr Thr Pro Pro Leu Thr Ile Asp Pro Ile
885 890 895

Asn Leu Thr Gly Phe Thr Leu Pro Gln Ile Thr Thr Pro Pro Ile Thr
900 905 910

Thr Pro Pro Leu Thr Ile Glu Pro Ile Gly Val Gly Gly Phe Thr Thr
915 920 925

Pro Pro Leu Thr Val Pro Gly Ile His Leu Pro Ser Thr Thr Ile Gly
930 935 940

Ala Phe Ala Ile Pro Gly Gly Pro Gly Tyr Phe Asn Ser Ser Thr Ala
945 950 955 960

Pro Ser Ser Gly Phe Phe Asn Ser Gly Ala Gly Gly Asn Ser Gly Phe
965 970 975

Gly Asn Asn Gly Ser Gly Leu Ser Gly Trp Phe Asn Thr Asn Pro Ala
980 985 990

Gly Leu Leu Gly Gly Ser Gly Tyr Gln Asn Phe Gly Gly Leu Ser Ser
995 1000 1005

Gly Phe Ser Asn Leu Gly Ser Gly Val Ser Gly Phe Ala Asn Arg
1010 1015 1020

Gly Ile Leu Pro Phe Ser Val Ala Ser Val Val Ser Gly Phe Ala
1025 1030 1035

Asn Ile Gly Thr Asn Leu Ala Gly Phe Phe Gln Gly Thr Thr Ser
1040 1045 1050

<210> 158
<211> 450
<212> PRT
<213> Mycobacterium tuberculosis

<400> 158

Met Ser Glu Leu Ser Val Ala Thr Gly Ala Val Ser Thr Ala Ser Ser
1 5 10 15

Ser Ile Pro Met Pro Ala Gly Val Asn Pro Ala Asp Leu Ala Ala Glu
20 25 30

Leu Ala Ala Val Val Thr Glu Ser Val Asp Glu Asp Tyr Leu Leu Tyr
35 40 45

colf-seq1

Glu Cys Asp Gly Gln Trp Val Leu Ala Ala Gly Val Gln Ala Met Val
50 55 60

Glu Leu Asp Ser Asp Glu Leu Arg Val Ile Arg Asp Gly Val Thr Arg
65 70 75 80

Arg Gln Gln Trp Ser Gly Arg Pro Gly Ala Ala Leu Gly Glu Ala Val
85 90 95

Asp Arg Leu Leu Leu Glu Thr Asp Gln Ala Phe Gly Trp Val Ala Phe
100 105 110

Glu Phe Gly Val His Arg Tyr Gly Leu Gln Gln Arg Leu Ala Pro His
115 120 125

Thr Pro Leu Ala Arg Val Phe Ser Pro Arg Thr Arg Ile Met Val Ser
130 135 140

Glu Lys Glu Ile Arg Leu Phe Asp Ala Gly Ile Arg His Arg Glu Ala
145 150 155 160

Ile Asp Arg Leu Leu Ala Thr Gly Val Arg Glu Val Pro Gln Ser Arg
165 170 175

Ser Val Asp Val Ser Asp Asp Pro Ser Gly Phe Arg Arg Arg Val Ala
180 185 190

Val Ala Val Asp Glu Ile Ala Ala Gly Arg Tyr His Lys Val Ile Leu
195 200 205

Ser Arg Cys Val Glu Val Pro Phe Ala Ile Asp Phe Pro Leu Thr Tyr
210 215 220

Arg Leu Gly Arg Arg His Asn Thr Pro Val Arg Ser Phe Leu Leu Gln
225 230 235 240

Leu Gly Gly Ile Arg Ala Leu Gly Tyr Ser Pro Glu Leu Val Thr Ala
245 250 255

Val Arg Ala Asp Gly Val Val Ile Thr Glu Pro Leu Ala Gly Thr Arg
260 265 270

Ala Leu Gly Arg Gly Pro Ala Ile Asp Arg Leu Ala Arg Asp Asp Leu
275 280 285

Glu Ser Asn Ser Lys Glu Ile Val Glu His Ala Ile Ser Val Arg Ser
290 295 300

Ser Leu Glu Glu Ile Thr Asp Ile Ala Glu Pro Gly Ser Ala Ala Val
305 310 315 320

Ile Asp Phe Met Thr Val Arg Glu Arg Gly Ser Val Gln His Leu Gly
 325 330 335
 Ser Thr Ile Arg Ala Arg Leu Asp Pro Ser Ser Asp Arg Met Ala Ala
 340 345 350
 Leu Glu Ala Leu Phe Pro Ala Val Thr Ala Ser Gly Ile Pro Lys Ala
 355 360 365
 Ala Gly Val Glu Ala Ile Phe Arg Leu Asp Glu Cys Pro Arg Gly Leu
 370 375 380
 Tyr Ser Gly Ala Val Val Met Leu Ser Ala Asp Gly Gly Leu Asp Ala
 385 390 395 400
 Ala Leu Thr Leu Arg Ala Ala Tyr Gln Val Gly Gly Arg Thr Trp Leu
 405 410 415
 Arg Ala Gly Ala Gly Ile Ile Glu Glu Ser Glu Pro Glu Arg Glu Phe
 420 425 430
 Glu Glu Thr Cys Glu Lys Leu Ser Thr Leu Thr Pro Tyr Leu Val Ala
 435 440 445
 Arg Gln
 450

<210> 159
 <211> 324
 <212> PRT
 <213> Mycobacterium tuberculosis
 <400> 159

Met Ser Asp Gln Val Pro Lys Pro His Arg His His Ile Trp Arg Ile
 1 5 10 15
 Thr Arg Arg Thr Leu Ser Lys Ser Trp Asp Asp Ser Ile Phe Ser Glu
 20 25 30
 Ser Ala Gln Ala Ala Phe Trp Ser Ala Leu Ser Leu Pro Pro Leu Leu
 35 40 45
 Leu Gly Met Leu Gly Ser Leu Ala Tyr Val Ala Pro Leu Phe Gly Pro
 50 55 60
 Asp Thr Leu Pro Ala Ile Glu Lys Ser Ala Leu Ser Thr Ala His Ser
 65 70 75 80
 Phe Phe Ser Pro Ser Val Val Asn Glu Ile Ile Glu Pro Thr Ile Gly
 85 90 95
 Asp Ile Thr Asn Asn Ala Arg Gly Glu Val Ala Ser Leu Gly Phe Leu

eolf-seq1

100	105	110
Ile Ser Leu Trp Ala Gly Ser Ser Ala Ile Ser Ala Phe val Asp Ala	115	120
val val Glu Ala His Asp Gln Thr Pro Leu Arg His Pro val Arg Gln	130	140
Arg Phe Phe Ala Leu Phe Leu Tyr val val Met Leu val Phe Leu val	145	155
Ala Thr Ala Pro val Met val val Gly Pro Arg Lys val Ser Glu His	165	170
Ile Pro Glu Ser Leu Ala Asn Leu Leu Arg Tyr Gly Tyr Tyr Pro Ala	180	185
Leu Ile Leu Gly Leu Thr val Gly val Ile Leu Leu Tyr Arg val Ala	195	200
Leu Pro val Pro Leu Pro Thr His Arg Leu val Leu Gly Ala val Leu	210	215
Ala Ile Ala val Phe Leu Ile Ala Thr Leu Gly Leu Arg val Tyr Leu	225	230
Ala Trp Ile Thr Arg Thr Gly Tyr Thr Tyr Gly Ala Leu Ala Thr Pro	245	250
Ile Ala Phe Leu Leu Phe Ala Phe Phe Gly Gly Phe Ala Ile Met Leu	260	265
Gly Ala Glu Leu Asn Ala Ala val Gln Glu Glu Trp Pro Ala Pro Ala	275	280
Thr His Ala His Arg Leu Gly Asn Trp Leu Lys Ala Arg Ile Gly val	290	295
Gly Thr Thr Thr Tyr Ser Ser Thr Ala Gln His Ser Ala val Ala Ala	305	310
Glu Pro Pro Ser		