

CARBAMATE DERIVATIVES AND USES THEREOF

FIELD OF THE DISCLOSURE

[001] The present disclosure relates to carbamate derivatives, prodrugs, and pharmaceutically acceptable salts thereof, which may possess inflammasome inhibitory activity and are accordingly useful in methods of treatment of the human or animal body. The present disclosure also relates to processes for the preparation of these compounds, to pharmaceutical compositions comprising them and to their use in the treatment of disorders in which inflammasome activity is implicated, such as inflammatory, autoinflammatory, autoimmune and oncological diseases.

BACKGROUND

[002] Autoimmune diseases are associated with the overproduction of pro-inflammatory factors. One of them is interleukin-1 (IL-1), produced by activated macrophages, monocytes, fibroblasts, and other components of the innate immune system like dendritic cells. IL-1 is involved in a variety of cellular activities, including cell proliferation, differentiation and apoptosis (Seth L. al. *Rev. Immunol.* 2009. 27:621–68).

[003] In humans, 22 NLR proteins are divided into four NLR subfamilies according to their N-terminal domains. NLRA contains a CARD-AT domain, NLRB (NAIP) contains a BIR domain, NLRC (including NOD1 and NOD2) contains a CARD domain, and NLRP contains a pyrin domain. Multiple NLR family members are associated with inflammasome formation.


[004] Although inflammasome activation appears to have evolved as an important component of host immunity to pathogens, the NLRP3 inflammasome is unique in its ability to activate in response to endogenous sterile danger signals. Many such sterile signals have been elucidated, and their formation is associated with specific disease states. For example, uric acid crystals found in gout patients are effective triggers of NLRP3 activation. Similarly, cholesterol crystals found in atherosclerotic patients can also promote NLRP3 activation. Recognition of the role of sterile danger signals as NLRP3 activators led to IL-1 and IL-18 being implicated in a diverse range of pathophysiological indications including metabolic, physiologic, inflammatory, hematologic and immunologic disorders.

[005] The disclosure arises from a need to provide further compounds for the specific modulation of NLRP3-dependent cellular processes. In particular, compounds with improved

physicochemical, pharmacological and pharmaceutical properties to existing compounds are desirable.

SUMMARY

[006] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (I):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

X is O or NR₅;

R₁ is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with

one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R_{3S}**;

each **R_{3S}** is independently selected from -OC(=O)**R_{3Sa}** or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)**R_{3Sb}**;

R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;


R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₅ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₆ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈

heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[007] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (I'):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

R₁ is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or R_{3S};

each **R_{3S}** is independently selected from -OC(=O)R_{3Sa} or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;


R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)OR_{3Sb};

R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₆ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[008] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (II):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

X is O or NR₅;

R₁ is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R_{3S}**;

each **R_{3S}** is independently selected from -OC(=O)**R_{3Sa}** or 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;


R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)OR_{3Sb};

R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₅ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[009] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (II'):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

R₁ is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R**_{1SS} is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R**_{3S};

each **R**_{3S} is independently selected from -OC(=O)**R**_{3Sa} or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)**R**_{3Sb};

R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl,

C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[010] In some aspects, the present disclosure provides a compound obtainable by, or obtained by, a method for preparing a compound as described herein (e.g., a method comprising one or more steps described in Scheme 1 or Scheme 2).

[011] In some aspects, the present disclosure provides a pharmaceutical composition comprising a compound of the present disclosure, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable diluent or carrier.

[012] In some aspects, the present disclosure provides an intermediate as described herein, being suitable for use in a method for preparing a compound as described herein (e.g., the intermediate is selected from the intermediates described in Examples 1-32).

[013] In some aspects, the present disclosure provides a method of inhibiting inflammasome (e.g., the NLRP3 inflammasome) activity (e.g., *in vitro* or *in vivo*), comprising contacting a cell with an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof.

[014] In some aspects, the present disclosure provides a method of treating or preventing a disease or disorder disclosed herein in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[015] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in inhibiting inflammasome (e.g., the NLRP3 inflammasome) activity (e.g., *in vitro* or *in vivo*).

[016] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating or preventing a disease or disorder disclosed herein.

[017] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for inhibiting inflammasome (e.g., the NLRP3 inflammasome) activity (e.g., *in vitro* or *in vivo*).

In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing a disease or disorder disclosed herein.

[018] In some aspects, the present disclosure provides a method of preparing a compound of the present disclosure.

[019] In some aspects, the present disclosure provides a method of preparing a compound, comprising one or more steps described herein.

[020] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. In the specification, the singular forms also include the plural unless the context clearly dictates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. All publications, patent applications, patents and other references mentioned herein are incorporated by reference. The references cited herein are not admitted to be prior art to the claimed invention. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods and examples are illustrative only and are not intended to be limiting. In the case of conflict between the chemical structures and names of the compounds disclosed herein, the chemical structures will control.

[021] Other features and advantages of the disclosure will be apparent from the following detailed description and claims.

DETAILED DESCRIPTION

[022] Autoimmune diseases are associated with the overproduction of proinflammatory factors. One of them is interleukin-1 (IL-1), produced by activated macrophages, monocytes, fibroblasts, and other components of the innate immune system like dendritic cells, involved in a variety of cellular activities, including cell proliferation, differentiation and apoptosis (Seth L. al. Rev. Immunol. 2009. 27:621–68).

[023] Cytokines from the IL-1 family are highly active and, as important mediators of inflammation, primarily associated with acute and chronic inflammation (Sims J. et al. Nature Reviews Immunology 10, 89-102 (February 2010)). The overproduction of IL-1 is considered to be a mediator of some autoimmune and autoinflammatory diseases. Autoinflammatory diseases

are characterised by recurrent and unprovoked inflammation in the absence of autoantibodies, infection, or antigen-specific T lymphocytes.

[024] Proinflammatory cytokines of the IL-1 superfamily include IL-1 α , IL-1 β , IL-18, and IL-36 α , β , λ and are produced in response to pathogens and other cellular stressors as part of a host innate immune response. Unlike many other secreted cytokines, which are processed and released via the standard cellular secretory apparatus consisting of the endoplasmic reticulum and Golgi apparatus, IL-1 family members lack leader sequences required for endoplasmic reticulum entry and thus are retained intracellularly following translation. In addition, IL-1 β , IL-18, and IL-36 α , β , λ are synthesised as procytokines that require proteolytic activation to become optimal ligands for binding to their cognate receptors on target cells.

[025] In the case of IL-1 α , IL-1 β and IL-18, it is now appreciated that a multimeric protein complex known as an inflammasome is responsible for activating the proforms of IL-1 β and IL-18 and for release of these cytokines extracellularly. An inflammasome complex typically consists of a sensor molecule, such as an NLR (Nucleotide-Oligomerisation Domain (NOD)-like receptor), an adaptor molecule ASC (Apoptosis-associated speck-like protein containing a CARD (Caspase Recruitment Domain)) and procaspase-1. In response to a variety of “danger signals”, including pathogen-associated molecule patterns (PAMPs) and danger associated molecular patterns (DAMPs), subunits of an inflammasome oligomerise to form a supramolecular structure within the cell. PAMPs include molecules such as peptidoglycan, viral DNA or RNA and bacterial DNA or RNA. DAMPs, on the other hand, consist of a wide range of endogenous or exogenous sterile triggers including monosodium urate crystals, silica, alum, asbestos, fatty acids, ceramides, cholesterol crystals and aggregates of beta-amyloid peptide. Assembly of an inflammasome platform facilitates autocatalysis of procaspase-1 yielding a highly active cysteine protease responsible for activation and release of pro-IL-1 β and pro-IL-18. Thus, release of these highly inflammatory cytokines is achieved only in response to inflammasome sensors detecting and responding to specific molecular danger signals.

[026] In humans, 22 NLR proteins are divided into four NLR subfamilies according to their N-terminal domains. NLRA contains a CARD-AT domain, NLRB (NAIP) contains a BIR domain, NLRC (including NOD1 and NOD2) contains a CARD domain, and NLRP contains a pyrin domain. Multiple NLR family members are associated with inflammasome formation including NLRP1, NLRP3, NLRP6, NLRP7, NLRP12 and NLRC4 (IPAF).

[027] Two other structurally distinct inflammasome structures containing a PYHIN domain (pyrin and HIN domain containing protein) namely Absent in Melanoma 2 (AIM2) and IFN λ -inducible protein 16 (IFI16) (Latz et al., Nat Rev Immunol 2013 13(6) 397-311) serve as intracellular DNA sensors. Pyrin (encoded by the MEFV gene) represents another type of inflammasome platform associated with proIL-1 β activation (Chae et al., Immunity 34, 755-768, 2011).

[028] Requiring assembly of an inflammasome platform to achieve activation and release of IL-1 β and IL-18 from monocytes and macrophages ensures their production is carefully orchestrated via a 2-step process. First, the cell must encounter a priming ligand (such as the TLR4 receptor ligand LPS, or an inflammatory cytokine such as TNF α) which leads to NF κ B dependent transcription of NLRP3, pro-IL-1 β and pro-IL-18. The newly translated procytokines remain intracellular and inactive unless producing cells encounter a second signal leading to activation of an inflammasome scaffold and maturation of procaspase-1.

[029] In addition to proteolytic activation of pro-IL-1 β and pro-IL-18, active caspase-1 also triggers a form of inflammatory cell death known as pyroptosis through cleavage of gasdermin-D. Pyroptosis allows the mature forms of IL-1 β and IL-18 to be externalised along with release of alarmin molecules (compounds that promote inflammation and activate innate and adaptive immunity) such as high mobility group box 1 protein (HMGB1), IL-33, and IL-1 α .

[030] Although inflammasome activation appears to have evolved as an important component of host immunity to pathogens, the NLRP3 inflammasome is unique in its ability activate in response to endogenous and exogenous sterile danger signals. Many such sterile signals have been elucidated, and their formation is associated with specific disease states. For example, uric acid crystals found in gout patients are effective triggers of NLRP3 activation. Similarly, cholesterol crystals found in atherosclerotic patients can also promote NLRP3 activation. Recognition of the role of sterile danger signals as NLRP3 activators led to IL-1 β and IL-18 being implicated in a diverse range of pathophysiological indications including metabolic, physiologic, inflammatory, hematologic and immunologic disorders.

[031] A link to human disease is best exemplified by discovery that mutations in the NLRP3 gene which lead to gain-of-function confer a range of autoinflammatory conditions collectively known as cryopyrin-associated periodic syndromes (CAPS) including familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS) and Neonatal onset

multisystem inflammatory disease (NOMID) (Hoffman et al., Nat Genet. 29(3) (2001) 301-305). Likewise, sterile mediator-induced activation of NLRP3 has been implicated in a wide range of disorders including joint degeneration (gout, rheumatoid arthritis, osteoarthritis), cardiometabolic (type 2 diabetes, atherosclerosis, hypertension), Central Nervous System (Alzheimer's Disease, Parkinson's disease, multiple sclerosis), gastrointestinal (Crohn's disease, ulcerative colitis), lung (chronic obstructive pulmonary disease (COPD), asthma, idiopathic pulmonary fibrosis) and liver (fibrosis, non-alcoholic fatty liver disease, non-alcoholic steatohepatitis (NASH)). It is further believed that NLRP3 activation promotes kidney inflammation and thus contributes to chronic kidney disease (CKD).

[032] Current treatment options for diseases where IL-1 is implicated as a contributor to pathogenesis include the IL-1 receptor antagonist anakinra, an Fc-containing fusion construct of the extracellular domains of the IL-1 receptor and IL-1 receptor accessory protein (riloncept) and the anti-IL-1 β monoclonal antibody canakinumab. For example, canakinumab is licensed for CAPS, Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS), Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD), Familial Mediterranean Fever (FMF) and gout.

[033] Some small molecules have been reported to inhibit function of the NLRP3 inflammasome. Glyburide, for example, is a specific inhibitor of NLRP3 activation, albeit at micromolar concentrations which are unlikely attainable *in vivo*. Non-specific agents such as parthenolide, Bay 11-7082, and 3,4-methylenedioxy- β -nitrostyrene are reported to impair NLRP3 activation but are expected to possess limited therapeutic utility due to their sharing of a common structural feature consisting of an olefin activated by substitution with an electron withdrawing group; this can lead to undesirable formation of covalent adducts with protein-bearing thiol groups. A number of natural products, for example β -hydroxybutyrate, sulforaphane, quercetin, and salvianolic acid, also are reported to suppress NLRP3 activation. Likewise, numerous effectors/modulators of other molecular targets have been reported to impair NLRP3 activation including agonists of the G-protein coupled receptor TGR5, an inhibitor of sodium-glucose co-transport epigliflozin, the dopamine receptor antagonist A-68930, the serotonin reuptake inhibitor fluoxetine, fenamate non-steroidal anti-inflammatory drugs, and the β -adrenergic receptor blocker nebivolol. Utility of these molecules as therapeutics for the chronic treatment of NLRP3-dependent inflammatory disorders remains to be established. A

series of sulphonylurea-containing molecules was previously identified as potent and selective inhibitors of post-translational processing of pro-IL-1 β (Perregaux et al., J Pharmacol. Exp. Ther. 299, 187-197, 2001). The exemplar molecule CP-456,773 from this work was recently characterised as a specific inhibitor of NLRP3 activation (Coll et al., Nat Med 21.3 (2015): 248-255.).

[034] The disclosure relates to compounds useful for the specific modulation of NLRP3-dependent cellular processes. In particular, compounds with improved physicochemical, pharmacological and pharmaceutical properties to existing NLRP3-modulating compounds are desired.

Definitions

[035] Unless otherwise stated, the following terms used in the specification and claims have the following meanings set out below.

[036] As used herein, “alkyl”, “C₁, C₂, C₃, C₄, C₅ or C₆ alkyl” or “C₁-C₆ alkyl” is intended to include C₁, C₂, C₃, C₄, C₅ or C₆ straight chain (linear) saturated aliphatic hydrocarbon groups and C₃, C₄, C₅ or C₆ branched saturated aliphatic hydrocarbon groups. For example, C₁-C₆ alkyl is intended to include C₁, C₂, C₃, C₄, C₅ and C₆ alkyl groups. Examples of alkyl include, moieties having from one to six carbon atoms, such as, but not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, or n-hexyl. In some embodiments, a straight chain or branched alkyl has six or fewer carbon atoms (*e.g.*, C₁-C₆ for straight chain, C₃-C₆ for branched chain), and in another embodiment, a straight chain or branched alkyl has four or fewer carbon atoms.

[037] As used herein, the term “optionally substituted alkyl” refers to unsubstituted alkyl or alkyl having designated substituents replacing one or more hydrogen atoms on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulphhydryl, alkylthio, arylthio, thiocarboxylate, sulphates,

alkylsulphinyl, sulphonato, sulphamoyl, sulphonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[038] As used herein, the term “alkenyl” includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double bond. For example, the term “alkenyl” includes straight chain alkenyl groups (*e.g.*, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl), and branched alkenyl groups. In certain embodiments, a straight chain or branched alkenyl group has six or fewer carbon atoms in its backbone (*e.g.*, C₂-C₆ for straight chain, C₃-C₆ for branched chain). The term “C₂-C₆” includes alkenyl groups containing two to six carbon atoms. The term “C₃-C₆” includes alkenyl groups containing three to six carbon atoms.

[039] As used herein, the term “optionally substituted alkenyl” refers to unsubstituted alkenyl or alkenyl having designated substituents replacing one or more hydrogen atoms on one or more hydrocarbon backbone carbon atoms. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonate, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulphhydryl, alkylthio, arylthio, thiocarboxylate, sulphates, alkylsulphinyl, sulphonato, sulphamoyl, sulphonamido, nitro, trifluoromethyl, cyano, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[040] As used herein, the term “alkynyl” includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but which contain at least one triple bond. For example, “alkynyl” includes straight chain alkynyl groups (*e.g.*, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl), and branched alkynyl groups. In certain embodiments, a straight chain or branched alkynyl group has six or fewer carbon atoms in its backbone (*e.g.*, C₂-C₆ for straight chain, C₃-C₆ for branched chain). The term “C₂-C₆” includes alkynyl groups containing two to six carbon atoms. The term “C₃-C₆” includes alkynyl groups containing three to six carbon atoms. As used herein, “C₂-C₆ alkenylene linker” or “C₂-C₆ alkynylene linker” is intended to include C₂, C₃, C₄, C₅ or C₆ chain

(linear or branched) divalent unsaturated aliphatic hydrocarbon groups. For example, C₂-C₆ alkenylene linker is intended to include C₂, C₃, C₄, C₅ and C₆ alkenylene linker groups.

[041] As used herein, the term “optionally substituted alkynyl” refers to unsubstituted alkynyl or alkynyl having designated substituents replacing one or more hydrogen atoms on one or more hydrocarbon backbone carbon atoms. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulphhydryl, alkylthio, arylthio, thiocarboxylate, sulphates, alkylsulphinyl, sulphonato, sulphamoyl, sulphonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[042] Other optionally substituted moieties (such as optionally substituted cycloalkyl, heterocycloalkyl, aryl, or heteroaryl) include both the unsubstituted moieties and the moieties having one or more of the designated substituents. For example, substituted heterocycloalkyl includes those substituted with one or more alkyl groups, such as 2,2,6,6-tetramethyl-piperidinyl and 2,2,6,6-tetramethyl-1,2,3,6-tetrahydropyridinyl.

[043] As used herein, the term “cycloalkyl” refers to a saturated or partially unsaturated hydrocarbon monocyclic or polycyclic (e.g., fused, bridged, or spiro rings) system having 3 to 30 carbon atoms (e.g., C₃-C₁₂, C₃-C₁₀, or C₃-C₈). Examples of cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, 1,2,3,4-tetrahydronaphthalenyl, and adamantyl. In the case of polycyclic cycloalkyl, only one of the rings in the cycloalkyl needs to be non-aromatic.

[044] As used herein, the term “heterocycloalkyl” refers to a saturated or partially unsaturated 3-8 membered monocyclic, 7-12 membered bicyclic (fused, bridged, or spiro rings), or 11-14 membered tricyclic ring system (fused, bridged, or spiro rings) having one or more heteroatoms (such as O, N, S, P, or Se), e.g., 1 or 1-2 or 1-3 or 1-4 or 1-5 or 1-6 heteroatoms, or *e.g.*, 1, 2, 3, 4, 5, or 6 heteroatoms, independently selected from the group consisting of nitrogen, oxygen and sulphur, unless specified otherwise. Examples of heterocycloalkyl groups include, but are not

limited to, piperidinyl, piperazinyl, pyrrolidinyl, dioxanyl, tetrahydrofuranyl, isoindolinyl, indolinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, oxiranyl, azetidiny, oxetanyl, thietanyl, 1,2,3,6-tetrahydropyridinyl, tetrahydropyranyl, dihydropyranyl, pyranyl, morpholinyl, tetrahydrothiopyranyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, 2-oxa-6-azaspiro[3.3]heptanyl, 2,6-diazaspiro[3.3]heptanyl, 1,4-dioxo-8-azaspiro[4.5]decanyl, 1,4-dioxaspiro[4.5]decanyl, 1-oxaspiro[4.5]decanyl, 1-azaspiro[4.5]decanyl, 3'H-spiro[cyclohexane-1,1'-isobenzofuran]-yl, 7'H-spiro[cyclohexane-1,5'-furo[3,4-b]pyridin]-yl, 3'H-spiro[cyclohexane-1,1'-furo[3,4-c]pyridin]-yl, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[3.1.0]hexan-3-yl, 1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazolyl, 3,4,5,6,7,8-hexahydropyrido[4,3-d]pyrimidinyl, 4,5,6,7-tetrahydro-1H-pyrazolo[3,4-c]pyridinyl, 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidinyl, 2-azaspiro[3.3]heptanyl, 2-methyl-2-azaspiro[3.3]heptanyl, 2-azaspiro[3.5]nonanyl, 2-methyl-2-azaspiro[3.5]nonanyl, 2-azaspiro[4.5]decanyl, 2-methyl-2-azaspiro[4.5]decanyl, 2-oxa-azaspiro[3.4]octanyl, 2-oxa-azaspiro[3.4]octan-6-yl, and the like. In the case of multicyclic heterocycloalkyl, only one of the rings in the heterocycloalkyl needs to be non-aromatic (*e.g.*, 4,5,6,7-tetrahydrobenzo[c]isoxazolyl).

[045] As used herein, the term “aryl” includes groups with aromaticity, including “conjugated,” or multicyclic systems with one or more aromatic rings and do not contain any heteroatom in the ring structure. The term aryl includes both monovalent species and divalent species. Examples of aryl groups include, but are not limited to, phenyl, biphenyl, naphthyl and the like. Conveniently, an aryl is phenyl.

[046] As used herein, the term “heteroaryl” is intended to include a stable 5-, 6-, or 7-membered monocyclic or 7-, 8-, 9-, 10-, 11- or 12-membered bicyclic aromatic heterocyclic ring which consists of carbon atoms and one or more heteroatoms, *e.g.*, 1 or 1-2 or 1-3 or 1-4 or 1-5 or 1-6 heteroatoms, or *e.g.*, 1, 2, 3, 4, 5, or 6 heteroatoms, independently selected from the group consisting of nitrogen, oxygen and sulphur. The nitrogen atom may be substituted or unsubstituted (*i.e.*, N or NR wherein R is H or other substituents, as defined). The nitrogen and sulphur heteroatoms may optionally be oxidised (*i.e.*, N→O and S(O)_p, where p = 1 or 2). It is to be noted that total number of S and O atoms in the aromatic heterocycle is not more than 1. Examples of heteroaryl groups include pyrrole, furan, thiophene, thiazole, isothiazole, imidazole, triazole, tetrazole, pyrazole, oxazole, isoxazole, pyridine, pyrazine, pyridazine, pyrimidine, and

the like. Heteroaryl groups can also be fused or bridged with alicyclic or heterocyclic rings, which are not aromatic so as to form a multicyclic system (*e.g.*, 4,5,6,7-tetrahydrobenzo[c]isoxazolyl).

[047] Furthermore, the terms “aryl” and “heteroaryl” include multicyclic aryl and heteroaryl groups, *e.g.*, tricyclic, bicyclic, *e.g.*, naphthalene, benzoxazole, benzodioxazole, benzothiazole, benzoimidazole, benzothiophene, quinoline, isoquinoline, naphthrydine, indole, benzofuran, purine, benzofuran, deazapurine, indolizine.

[048] The cycloalkyl, heterocycloalkyl, aryl, or heteroaryl ring can be substituted at one or more ring positions (*e.g.*, the ring-forming carbon or heteroatom such as N) with such substituents as described above, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkylaminocarbonyl, aralkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aralkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonate, phosphinate, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulphhydryl, alkylthio, arylthio, thiocarboxylate, sulphates, alkylsulphinyl, sulphonate, sulphamoyl, sulphonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Aryl and heteroaryl groups can also be fused or bridged with alicyclic or heterocyclic rings, which are not aromatic so as to form a multicyclic system (*e.g.*, tetralin, methylenedioxyphenyl such as benzo[d][1,3]dioxole-5-yl).

[049] As used herein, the term “substituted,” means that any one or more hydrogen atoms on the designated atom is replaced with a selection from the indicated groups, provided that the designated atom’s normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is oxo or keto (*i.e.*, =O), then 2 hydrogen atoms on the atom are replaced. Keto substituents are not present on aromatic moieties. Ring double bonds, as used herein, are double bonds that are formed between two adjacent ring atoms (*e.g.*, C=C, C=N or N=N). “Stable compound” and “stable structure” are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

[050] When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom in the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such formula.

Combinations of substituents and/or variables are permissible, but only if such combinations result in stable compounds.

[051] When any variable (*e.g.*, R) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R moieties, then the group may optionally be substituted with up to two R moieties and R at each occurrence is selected independently from the definition of R. Also, combinations of substituents and/or variables are permissible, but only if such combinations result in stable compounds.

[052] As used herein, the term “hydroxy” or “hydroxyl” includes groups with an -OH or -O[•].

[053] As used herein, the term “halo” or “halogen” refers to fluoro, chloro, bromo and iodo.

[054] The term “haloalkyl” or “haloalkoxyl” refers to an alkyl or alkoxyl substituted with one or more halogen atoms.

[055] As used herein, the term “optionally substituted haloalkyl” refers to unsubstituted haloalkyl having designated substituents replacing one or more hydrogen atoms on one or more hydrocarbon backbone carbon atoms. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulphhydryl, alkylthio, arylthio, thiocarboxylate, sulphates, alkylsulphinyl, sulphonato, sulphamoyl, sulphonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[056] As used herein, the term “alkoxy” or “alkoxyl” includes substituted and unsubstituted alkyl, alkenyl and alkynyl groups covalently linked to an oxygen atom. Examples of alkoxy groups or alkoxyl radicals include, but are not limited to, methoxy, ethoxy, isopropoxy, propoxy, butoxy and pentoxy groups. Examples of substituted alkoxy groups include

halogenated alkoxy groups. The alkoxy groups can be substituted with groups such as alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonato, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulphhydryl, alkylthio, arylthio, thiocarboxylate, sulphates, alkylsulphinyl, sulphonato, sulphonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moieties. Examples of halogen substituted alkoxy groups include, but are not limited to, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chloromethoxy, dichloromethoxy and trichloromethoxy.

[057] As used herein, the expressions “one or more of A, B, or C,” “one or more A, B, or C,” “one or more of A, B, and C,” “one or more A, B, and C,” “selected from the group consisting of A, B, and C,” “selected from A, B, and C”, and the like are used interchangeably and all refer to a selection from a group consisting of A, B, and/or C, i.e., one or more As, one or more Bs, one or more Cs, or any combination thereof, unless indicated otherwise.

[058] It is to be understood that the present disclosure provides methods for the synthesis of the compounds of any of the Formulae described herein. The present disclosure also provides detailed methods for the synthesis of various disclosed compounds of the present disclosure according to the following schemes as well as those shown in the Examples.

[059] It is to be understood that, throughout the description, where compositions are described as having, including, or comprising specific components, it is contemplated that compositions also consist essentially of, or consist of, the recited components. Similarly, where methods or processes are described as having, including, or comprising specific process steps, the processes also consist essentially of, or consist of, the recited processing steps. Further, it should be understood that the order of steps or order for performing certain actions is immaterial so long as the invention remains operable. Moreover, two or more steps or actions can be conducted simultaneously.

[060] It is to be understood that the synthetic processes of the disclosure can tolerate a wide variety of functional groups, therefore various substituted starting materials can be used. The processes generally provide the desired final compound at or near the end of the overall process,

although it may be desirable in certain instances to further convert the compound to a pharmaceutically acceptable salt thereof.

[061] It is to be understood that compounds of the present disclosure can be prepared in a variety of ways using commercially available starting materials, compounds known in the literature, or from readily prepared intermediates, by employing standard synthetic methods and procedures either known to those skilled in the art, or which will be apparent to the skilled artisan in light of the teachings herein. Standard synthetic methods and procedures for the preparation of organic molecules and functional group transformations and manipulations can be obtained from the relevant scientific literature or from standard textbooks in the field. Although not limited to any one or several sources, classic texts such as Smith, M. B., March, J., *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 5th edition, John Wiley & Sons: New York, 2001; Greene, T.W., Wuts, P.G. M., *Protective Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons: New York, 1999; R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); and L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995), incorporated by reference herein, are useful and recognised reference textbooks of organic synthesis known to those in the art

[062] One of ordinary skill in the art will note that, during the reaction sequences and synthetic schemes described herein, the order of certain steps may be changed, such as the introduction and removal of protecting groups. One of ordinary skill in the art will recognise that certain groups may require protection from the reaction conditions via the use of protecting groups. Protecting groups may also be used to differentiate similar functional groups in molecules. A list of protecting groups and how to introduce and remove these groups can be found in Greene, T.W., Wuts, P.G. M., *Protective Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons: New York, 1999.

[063] It is to be understood that, unless otherwise stated, any description of a method of treatment includes use of the compounds to provide such treatment or prophylaxis as is described herein, as well as use of the compounds to prepare a medicament to treat or prevent such condition. The treatment includes treatment of human or non-human animals including rodents and other disease models.

[064] As used herein, the term “subject” is interchangeable with the term “subject in need thereof”, both of which refer to a subject having a disease or having an increased risk of developing the disease. A “subject” includes a mammal. The mammal can be *e.g.*, a human or appropriate non-human mammal, such as primate, mouse, rat, dog, cat, cow, horse, goat, camel, sheep or a pig. The subject can also be a bird or fowl. In one embodiment, the mammal is a human. A subject in need thereof can be one who has been previously diagnosed or identified as having a disease or disorder disclosed herein. A subject in need thereof can also be one who is suffering from a disease or disorder disclosed herein. Alternatively, a subject in need thereof can be one who has an increased risk of developing such disease or disorder relative to the population at large (i.e., a subject who is predisposed to developing such disorder relative to the population at large). A subject in need thereof can have a refractory or resistant a disease or disorder disclosed herein (i.e., a disease or disorder disclosed herein that does not respond or has not yet responded to treatment). The subject may be resistant at start of treatment or may become resistant during treatment. In some embodiments, the subject in need thereof received and failed all known effective therapies for a disease or disorder disclosed herein. In some embodiments, the subject in need thereof received at least one prior therapy.

[065] As used herein, the term “treating” or “treat” describes the management and care of a patient for the purpose of combating a disease, condition, or disorder and includes the administration of a compound of the present disclosure, or a pharmaceutically acceptable salt, polymorph or solvate thereof, to alleviate the symptoms or complications of a disease, condition or disorder, or to eliminate the disease, condition or disorder. The term “treat” can also include treatment of a cell *in vitro* or an animal model. It is to be appreciated that references to “treating” or “treatment” include the alleviation of established symptoms of a condition. “Treating” or “treatment” of a state, disorder or condition therefore includes: (1) preventing or delaying the appearance of clinical symptoms of the state, disorder or condition developing in a human that may be afflicted with the state, disorder or condition but does not yet experience or display clinical or subclinical symptoms of the state, disorder or condition, (2) inhibiting the state, disorder or condition, i.e., arresting, reducing or delaying the development of the disease or a relapse thereof (in case of maintenance treatment) or at least one clinical or subclinical symptom thereof, or (3) relieving or attenuating the disease, i.e., causing regression of the state, disorder or condition or at least one of its clinical or subclinical symptoms.

[066] It is to be understood that a compound of the present disclosure, or a pharmaceutically acceptable salt, polymorph or solvate thereof, can or may also be used to prevent a relevant disease, condition or disorder, or used to identify suitable candidates for such purposes.

[067] As used herein, the term “preventing,” “prevent,” or “protecting against” describes reducing or eliminating the onset of the symptoms or complications of such disease, condition or disorder.

[068] It is to be understood that one skilled in the art may refer to general reference texts for detailed descriptions of known techniques discussed herein or equivalent techniques. These texts include Ausubel *et al.*, *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc. (2005); Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (3rd edition), Cold Spring Harbor Press, Cold Spring Harbor, New York (2000); Coligan *et al.*, *Current Protocols in Immunology*, John Wiley & Sons, N.Y.; Enna *et al.*, *Current Protocols in Pharmacology*, John Wiley & Sons, N.Y.; Fingl *et al.*, *The Pharmacological Basis of Therapeutics* (1975), *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 18th edition (1990). These texts can, of course, also be referred to in making or using an aspect of the disclosure.

[069] It is to be understood that the present disclosure also provides pharmaceutical compositions comprising any compound described herein in combination with at least one pharmaceutically acceptable excipient or carrier.

[070] As used herein, the term “pharmaceutical composition” is a formulation containing the compounds of the present disclosure in a form suitable for administration to a subject. In one embodiment, the pharmaceutical composition is in bulk or in unit dosage form. The unit dosage form is any of a variety of forms, including, for example, a capsule, an IV bag, a tablet, a single pump on an aerosol inhaler or a vial. The quantity of active ingredient (*e.g.*, a formulation of the disclosed compound or salt, hydrate, solvate or isomer thereof) in a unit dose of composition is an effective amount and is varied according to the particular treatment involved. One skilled in the art will appreciate that it is sometimes necessary to make routine variations to the dosage depending on the age and condition of the patient. The dosage will also depend on the route of administration. A variety of routes are contemplated, including oral, pulmonary, rectal, parenteral, transdermal, subcutaneous, intravenous, intramuscular, intraperitoneal, inhalational, buccal, sublingual, intrapleural, intrathecal, intranasal, and the like. Dosage forms for the topical or transdermal administration of a compound of this disclosure include powders, sprays,

ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. In one embodiment, the active compound is mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants that are required.

[071] As used herein, the term “pharmaceutically acceptable” refers to those compounds, anions, cations, materials, compositions, carriers, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[072] As used herein, the term “pharmaceutically acceptable excipient” means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes excipient that is acceptable for veterinary use as well as human pharmaceutical use. A “pharmaceutically acceptable excipient” as used in the specification and claims includes both one and more than one such excipient.

[073] It is to be understood that a pharmaceutical composition of the disclosure is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, ingestion), inhalation, transdermal (topical), and transmucosal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulphite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[074] It is to be understood that a compound or pharmaceutical composition of the disclosure can be administered to a subject in many of the well-known methods currently used for chemotherapeutic treatment. For example, a compound of the disclosure may be injected into the blood stream or body cavities or taken orally or applied through the skin with patches. The dose chosen should be sufficient to constitute effective treatment but not so high as to cause

unacceptable side effects. The state of the disease condition (*e.g.*, a disease or disorder disclosed herein) and the health of the patient should preferably be closely monitored during and for a reasonable period after treatment.

[075] A “therapeutically effective amount” means the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect such treatment for the disease. The “therapeutically effective amount” will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated. Therapeutically effective amounts for a given situation can be determined by routine experimentation that is within the skill and judgment of the clinician.

[076] As used herein, the term “effective amount”, refers to an amount of a pharmaceutical agent to treat, ameliorate, or prevent an identified disease or condition, or to exhibit a detectable therapeutic or inhibitory effect. The effect can be detected by any assay method known in the art. The precise effective amount for a subject will depend upon the subject’s body weight, size, and health; the nature and extent of the condition; and the therapeutic or combination of therapeutics selected for administration. Effective amounts for a given situation can be determined by routine experimentation that is within the skill and judgment of the clinician.

[077] It is to be understood that, for any compound, the therapeutically effective amount can be estimated initially either in cell culture assays, *e.g.*, of neoplastic cells, or in animal models, usually rats, mice, rabbits, dogs, or pigs. It is to be understood that, for any compound, the effective amount can be estimated initially either in cell culture assays, *e.g.*, of neoplastic cells, or in animal models, usually rats, mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans. Therapeutic/prophylactic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD₅₀/ED₅₀. Pharmaceutical compositions that exhibit large therapeutic indices are preferred. The dosage may vary within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

[078] Dosage and administration are adjusted to provide sufficient levels of the active agent(s) or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

[079] The pharmaceutical compositions containing active compounds of the present disclosure may be manufactured in a manner that is generally known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilising processes. Pharmaceutical compositions may be formulated in a conventional manner using one or more pharmaceutically acceptable carriers comprising excipients and/or auxiliaries that facilitate processing of the active compounds into preparations that can be used pharmaceutically. Of course, the appropriate formulation is dependent upon the route of administration chosen.

[080] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol and sorbitol, and sodium chloride in the composition. Prolonged absorption of the

injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

[081] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilisation. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[082] Oral compositions generally include an inert diluent or an edible pharmaceutically acceptable carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[083] For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser, which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebuliser.

[084] Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or

suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[085] The active compounds can be prepared with pharmaceutically acceptable carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

[086] It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the disclosure are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved.

[087] In therapeutic applications, the dosages of the pharmaceutical compositions used in accordance with the disclosure vary depending on the agent, the age, weight, and clinical condition of the recipient patient, and the experience and judgment of the clinician or practitioner administering the therapy, among other factors affecting the selected dosage. Generally, the dose should be sufficient to result in slowing, and preferably regressing, the symptoms of the disease or disorder disclosed herein and also preferably causing complete regression of the disease or disorder. Dosages can range from about 0.01 mg/kg per day to about 5000 mg/kg per day. In preferred aspects, dosages can range from about 1 mg/kg per day to about 1000 mg/kg per day. In an aspect, the dose will be in the range of about 0.1 mg/day to about 50 g/day; about 0.1 mg/day to about 25 g/day; about 0.1 mg/day to about 10 g/day; about 0.1 mg to about 3 g/day; or about 0.1 mg to about 1 g/day, in single, divided, or continuous doses (which dose may be adjusted for the patient's weight in kg, body surface area in m², and age in years). An effective

amount of a pharmaceutical agent is that which provides an objectively identifiable improvement as noted by the clinician or other qualified observer. Improvement in survival and growth indicates regression. As used herein, the term “dosage effective manner” refers to amount of an active compound to produce the desired biological effect in a subject or cell.

[088] It is to be understood that the pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

[089] It is to be understood that, for the compounds of the present disclosure being capable of further forming salts, all of these forms are also contemplated within the scope of the claimed disclosure.

[090] As used herein, the term “pharmaceutically acceptable salts” refer to derivatives of the compounds of the present disclosure wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines, alkali or organic salts of acidic residues such as carboxylic acids, and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include, but are not limited to, those derived from inorganic and organic acids selected from 2-acetoxybenzoic, 2-hydroxyethane sulphonic, acetic, ascorbic, benzene sulphonic, benzoic, bicarbonic, carbonic, citric, edetic, ethane disulphonic, 1,2-ethane sulphonic, fumaric, glucoheptonic, gluconic, glutamic, glycolic, glycollyarsanilic, hexylresorcinic, hydrabamic, hydrobromic, hydrochloric, hydroiodic, hydroxymaleic, hydroxynaphthoic, isethionic, lactic, lactobionic, lauryl sulphonic, maleic, malic, mandelic, methane sulphonic, napsylic, nitric, oxalic, pamoic, pantothenic, phenylacetic, phosphoric, polygalacturonic, propionic, salicylic, stearic, subacetic, succinic, sulphamic, sulphanilic, sulphuric, tannic, tartaric, toluene sulphonic, and the commonly occurring amine acids, *e.g.*, glycine, alanine, phenylalanine, arginine, etc.

[091] In some embodiments, the pharmaceutically acceptable salt is a sodium salt, a potassium salt, a calcium salt, a magnesium salt, a diethylamine salt, a choline salt, a meglumine salt, a benzathine salt, a tromethamine salt, an ammonia salt, an arginine salt, or a lysine salt.

[092] Other examples of pharmaceutically acceptable salts include hexanoic acid, cyclopentane propionic acid, pyruvic acid, malonic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, 4-chlorobenzenesulphonic acid, 2-naphthalenesulphonic acid, 4-toluenesulphonic

acid, camphorsulphonic acid, 4-methylbicyclo-[2.2.2]-oct-2-ene-1-carboxylic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, muconic acid, and the like. The present disclosure also encompasses salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, *e.g.*, an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine, and the like. In the salt form, it is understood that the ratio of the compound to the cation or anion of the salt can be 1:1, or any ratio other than 1:1, *e.g.*, 3:1, 2:1, 1:2, or 1:3.

[093] It is to be understood that all references to pharmaceutically acceptable salts include solvent addition forms (solvates) or crystal forms (polymorphs) as defined herein, of the same salt.

[094] The compounds, or pharmaceutically acceptable salts thereof, are administered orally, nasally, transdermally, pulmonary, inhalationally, buccally, sublingually, intraperitoneally, subcutaneously, intramuscularly, intravenously, rectally, intrapleurally, intrathecally and parenterally. In one embodiment, the compound is administered orally. One skilled in the art will recognise the advantages of certain routes of administration.

[095] The dosage regimen utilising the compounds is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the condition.

[096] Techniques for formulation and administration of the disclosed compounds of the disclosure can be found in *Remington: the Science and Practice of Pharmacy*, 19th edition, Mack Publishing Co., Easton, PA (1995). In an embodiment, the compounds described herein, and the pharmaceutically acceptable salts thereof, are used in pharmaceutical preparations in combination with a pharmaceutically acceptable carrier or diluent. Suitable pharmaceutically acceptable carriers include inert solid fillers or diluents and sterile aqueous or organic solutions. The compounds will be present in such pharmaceutical compositions in amounts sufficient to provide the desired dosage amount in the range described herein.

[097] All percentages and ratios used herein, unless otherwise indicated, are by weight. Other features and advantages of the present disclosure are apparent from the different examples. The provided examples illustrate different components and methodology useful in practicing the present disclosure. The examples do not limit the claimed disclosure. Based on the present disclosure the skilled artisan can identify and employ other components and methodology useful for practicing the present disclosure.


[098] In the synthetic schemes described herein, compounds may be drawn with one particular configuration for simplicity. Such particular configurations are not to be construed as limiting the disclosure to one or another isomer, tautomer, regioisomer or stereoisomer, nor does it exclude mixtures of isomers, tautomers, regioisomers or stereoisomers; however, it will be understood that a given isomer, tautomer, regioisomer or stereoisomer may have a higher level of activity than another isomer, tautomer, regioisomer or stereoisomer.

[099] All publications and patent documents cited herein are incorporated herein by reference as if each such publication or document was specifically and individually indicated to be incorporated herein by reference. Citation of publications and patent documents is not intended as an admission that any is pertinent prior art, nor does it constitute any admission as to the contents or date of the same. The invention having now been described by way of written description, those of skill in the art will recognize that the invention can be practiced in a variety of embodiments and that the foregoing description and examples below are for purposes of illustration and not limitation of the claims that follow.

[0100] As use herein, the phrase “compound of the disclosure” refers to those compounds which are disclosed herein, both generically and specifically.

Compounds of the Present Disclosure

[0101] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (I):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

X is O or NR₅;

R₁ is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R_{3S}**;

each **R_{3S}** is independently selected from -OC(=O)**R_{3Sa}** or 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)OR_{3Sb};


R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₅ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₆ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0102] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (I'):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

R₁ is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R_{3S}**;

each **R_{3S}** is independently selected from -OC(=O)**R_{3Sa}** or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;


R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)**R_{3Sb}**;

R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₆ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0103] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (II):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

X is **O** or **NR₅**;

R₁ is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R**_{1S} is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R**_{1SS};

each **R**_{1SS} is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-

(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or R_{3S};

each **R_{3S}** is independently selected from -OC(=O)R_{3Sa} or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)OR_{3Sb};


R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈

heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₅ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0104] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (II'):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

R₁ is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-

C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R_{3S}**;


each **R_{3S}** is independently selected from -OC(=O)**R_{3Sa}** or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)**R_{3Sb}**;

R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0105] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (III):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

X is O or NR₅;

R₁ is C₃-C₁₆ cycloalkyl, 5- or 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl,


C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₅ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₆ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0106] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (III'):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

R₁ is C₃-C₁₆ cycloalkyl, 5- or 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-

membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R**_{1SS};

each **R**_{1SS} is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;


Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0107] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (IV):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

X is O or NR₅;

R₁ is 5- to 10-membered heteroaryl or C₅-C₁₀ aryl, wherein the 5- or 6-membered heteroaryl or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H or C₁-C₆ alkyl optionally substituted with one or more C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R_{3S}**;

each **R_{3S}** is independently selected from -OC(=O)**R_{3Sa}** or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)OR_{3Sb};


R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₅ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₆ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0108] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (IV'):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

R₁ is 5- to 10-membered heteroaryl or C₅-C₁₀ aryl, wherein the 5- or 6-membered heteroaryl or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R**_{1SS} is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H or C₁-C₆ alkyl optionally substituted with one or more C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R**_{3S};


each **R**_{3S} is independently selected from -OC(=O)**R**_{3Sa} or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)**R**_{3Sb};

R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₄ is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0109] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (V):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

X is O or NR₅;

R₁ is 5- to 10-membered heteroaryl or C₅-C₁₀ aryl, wherein the 5- or 6-membered heteroaryl or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;


R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₅ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0110] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (V'):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

R₁ is 5- to 10-membered heteroaryl or C₅-C₁₀ aryl, wherein the 5- or 6-membered heteroaryl or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;


R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-

membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₄ is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0111] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (VI):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

X is O or NR₅;

R₁ is 5- to 10-membered heteroaryl or C₅-C₁₀ aryl, wherein the 5- or 6-membered heteroaryl or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;


R₃ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₅ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₆ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0112] In some aspects, the present disclosure provides, *inter alia*, a compound of Formula (VI):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

R₁ is 5- to 10-membered heteroaryl or C₅-C₁₀ aryl, wherein the 5- or 6-membered heteroaryl or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN;

wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R**_{1SS};

each **R**_{1SS} is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₆ is H or C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0113] It is understood that, for a compound of Formula (I), **X**, **R**₁, **R**_{1S}, **R**_{1SS}, **Q**, **R**₂, **R**₃, **R**_{3Sa}, **R**_{3Sb}, **R**₄, **R**₅, and **R**₆ can each be, where applicable, selected from the groups described herein, and any group described herein for any of **X**, **R**₁, **R**_{1S}, **R**_{1SS}, **Q**, **R**₂, **R**₃, **R**_{3Sa}, **R**_{3Sb}, **R**₄, **R**₅, and **R**₆ can be combined, where applicable, with any group described herein for one or more of the remainder of **X**, **R**₁, **R**_{1S}, **R**_{1SS}, **Q**, **R**₂, **R**₃, **R**₄, **R**₅, and **R**₆.

[0114] In some embodiments, **X** is O or NR₅. In some embodiments, **X** is O. In some embodiments, **X** is NR₅.

[0115] In some embodiments, **R₁** is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**.

[0116] In some embodiments, **R₁** is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl or C₅-C₆ aryl is optionally substituted with one or more **R_{1S}**.

[0117] In some embodiments, **R₁** is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the 5- to 10-membered heteroaryl or C₅-C₆ aryl is optionally substituted with one or more **R_{1S}**.

[0118] In some embodiments, **R₁** is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl or 5- to 10-membered heteroaryl is optionally substituted with one or more **R_{1S}**.

[0119] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₆ aryl is optionally substituted with one or more **R_{1S}**.

[0120] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₈ monocyclic cycloalkyl, or C₈-C₁₆ polycyclic cycloalkyl is optionally substituted with one or more **R_{1S}**.

[0121] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, or C₁₂-C₁₆ tricyclic cycloalkyl is optionally substituted with one or more **R_{1S}**.

[0122] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, or C₅-C₆ aryl is optionally substituted with one or more **R_{1S}**.

[0123] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, is optionally substituted with one or more **R_{1S}**.

[0124] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, or C₅-C₁₀ aryl, is optionally substituted with one or more **R_{1S}**.

[0125] In some embodiments, **R₁** is C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**.

[0126] In some embodiments, **R₁** is C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl or C₅-C₆ aryl is optionally substituted with one or more **R_{1S}**.

[0127] In some embodiments, **R₁** is C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the 5- or 6-membered heteroaryl or C₅-C₆ aryl is optionally substituted with one or more **R_{1S}**.

[0128] In some embodiments, **R₁** is C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl or 5- or 6-membered heteroaryl is optionally substituted with one or more **R_{1S}**.

[0129] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₆ aryl is optionally substituted with one or more **R_{1S}**.

[0130] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₈ monocyclic cycloalkyl, or C₈-C₁₆ polycyclic cycloalkyl is optionally substituted with one or more **R_{1S}**.

[0131] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, or C₁₂-C₁₆ tricyclic cycloalkyl is optionally substituted with one or more **R_{1S}**.

[0132] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇ monocyclic cycloalkyl, C₈-C₁₆ polycyclic cycloalkyl, or C₅-C₆ aryl is optionally substituted with one or more **R_{1S}**.

[0133] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇

monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, is optionally substituted with one or more **R_{1S}**.

[0134] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₇ monocyclic cycloalkyl, C₉-C₁₀ bicyclic cycloalkyl, C₁₂-C₁₆ tricyclic cycloalkyl, or C₅-C₁₀ aryl, is optionally substituted with one or more **R_{1S}**.

[0135] In some embodiments, **R₁** is C₃-C₁₆ cycloalkyl optionally substituted with one or more **R_{1S}**.

[0136] In some embodiments, **R₁** is C₃-C₇ monocyclic cycloalkyl optionally substituted with one or more **R_{1S}**.

[0137] In some embodiments, **R₁** is C₃-C₇ monocyclic saturated cycloalkyl optionally substituted with one or more **R_{1S}**.

[0138] In some embodiments, **R₁** is C₃-C₇ monocyclic partially saturated cycloalkyl optionally substituted with one or more **R_{1S}**.

[0139] In some embodiments, **R₁** is C₉-C₁₀ bicyclic cycloalkyl optionally substituted with one or more **R_{1S}**.

[0140] In some embodiments, **R₁** is C₉-C₁₀ bicyclic saturated cycloalkyl optionally substituted with one or more **R_{1S}**.

[0141] In some embodiments, **R₁** is C₉-C₁₀ bicyclic partially saturated cycloalkyl optionally substituted with one or more **R_{1S}**.

[0142] In some embodiments, **R₁** is C₁₂-C₁₆ tricyclic cycloalkyl optionally substituted with one or more **R_{1S}**.

[0143] In some embodiments, **R₁** is C₁₂-C₁₆ tricyclic saturated cycloalkyl optionally substituted with one or more **R_{1S}**.

[0144] In some embodiments, **R₁** is C₁₂-C₁₆ tricyclic partially unsaturated cycloalkyl optionally substituted with one or more **R_{1S}**.

[0145] In some embodiments, **R₁** is C₈-C₁₆ polycyclic cycloalkyl substituted with one or more **R_{1S}**.

[0146] In some embodiments, **R₁** is 2,3-dihydro-1*H*-indenyl optionally substituted with one or more **R_{1S}**.

[0147] In some embodiments, **R₁** is hexahydroindacenyl optionally substituted with one or more **R_{1S}**.

[0148] In some embodiments, **R₁** is 5- to 10-membered heteroaryl optionally substituted with one or more **R_{1S}**.

[0149] In some embodiments, **R₁** is 5- to 10-membered heteroaryl. In some embodiments, **R₁** is 5- to 9-membered heteroaryl optionally substituted with one or more **R_{1S}**.

[0150] In some embodiments, **R₁** is 5- to 9-membered heteroaryl.

[0151] In some embodiments, **R₁** is 9-membered heteroaryl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is 9-membered heteroaryl.

[0152] In some embodiments, **R₁** is quinoliny.

[0153] In some embodiments, **R₁** is 5- to 8-membered heteroaryl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is 5- to 8-membered heteroaryl.

[0154] In some embodiments, **R₁** is 5- to 7-membered heteroaryl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is 5- to 7-membered heteroaryl.







[0155] In some embodiments, **R₁** is 5- or 6-membered heteroaryl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is 5- or 6-membered heteroaryl.

[0156] In some embodiments, **R₁** is 5-membered heteroaryl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is 5-membered heteroaryl.

[0157] In some embodiments, **R₁** is pyrrolyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is pyrazolyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is imidazolyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is triazolyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is tetrazolyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is isoxazolyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is furanyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is oxazolyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is isothiazolyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is thiazolyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is thiadiazolyl optionally substituted with one or more **R_{1S}**.

[0158] In some embodiments, **R₁** is 6-membered heteroaryl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is 6-membered heteroaryl.

[0159] In some embodiments, **R₁** is pyridinyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is diazinyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is pyridazinyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is pyrimidinyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is pyrazinyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is triazinyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is tetrazinyl optionally substituted with one **R_{1S}**. In some embodiments, **R₁** is pentazinyl.

[0160] In some embodiments, **R₁** is . In some embodiments, **R₁** is . In some embodiments, **R₁** is . In some embodiments, **R₁** is . In some embodiments, **R₁** is . In some embodiments, **R₁** is .

[0161] In some embodiments, **R₁** is C₅-C₁₀ aryl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is C₅-C₉ aryl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is C₅-C₈ aryl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is C₅-C₇ aryl optionally substituted with one or more **R_{1S}**.

[0162] In some embodiments, **R₁** is C₅-C₁₀ aryl substituted with one or more **R_{1S}**. In some embodiments, **R₁** is C₅-C₉ aryl substituted with one or more **R_{1S}**. In some embodiments, **R₁** is C₅-C₈ aryl substituted with one or more **R_{1S}**. In some embodiments, **R₁** is C₅-C₇ aryl substituted with one or more **R_{1S}**.


[0163] In some embodiments, **R₁** is C₅-C₆ monocyclic aryl optionally substituted with one or more **R_{1S}**.

[0164] In some embodiments, **R₁** is C₅-C₆ monocyclic aryl substituted with one or more **R_{1S}**.



[0165] In some embodiments, **R₁** is C₅-C₆ monocyclic aryl.

[0166] In some embodiments, **R₁** is phenyl. In some embodiments, **R₁** is phenyl optionally substituted with one or more **R_{1S}**. In some embodiments, **R₁** is phenyl substituted with one or more **R_{1S}**.




[0167] In some embodiments, **R₁** is phenyl substituted with one **R_{1S}**.



[0169] In some embodiments, **R₁** is phenyl substituted with two **R_{1S}**.



[0171] In some embodiments, **R₁** is phenyl substituted with three **R_{1S}**.



[0174] In some embodiments, **R₁** is phenyl substituted with four **R_{1S}**.

[0175] In some embodiments, **R₁** is phenyl substituted with five **R_{1S}**.

[0176] In some embodiments, **R₁** is phenyl substituted with one or more substituents independently selected from C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**.

[0177] In some embodiments, **R₁** is phenyl substituted with one or more substituents independently selected from C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN.

[0178] In some embodiments, **R₁** is phenyl substituted with one or more substituents independently selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, C₁-C₆ alkoxy, halo, or CN.

[0179] In some embodiments, **R₁** is phenyl substituted with one or more substituents independently selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, or C₁-C₆ alkoxy.

[0180] In some embodiments, **R₁** is phenyl substituted with one or more substituents independently selected from halo or CN.

[0181] In some embodiments, **R₁** is phenyl substituted with one or more halo. In some embodiments, **R₁** is phenyl substituted with one or more F, Cl, Br, or I. In some embodiments, **R₁** is phenyl substituted with one or more F or Cl. In some embodiments, **R₁** is phenyl substituted with one or more F. In some embodiments, **R₁** is phenyl substituted with one or more Cl.

[0182] In some embodiments, **R₁** is phenyl substituted with one or more CN.

[0183] In some embodiments, **R₁** is phenyl substituted with one or more substituents independently selected from methyl, ethyl, isopropyl, tert-butyl, ethoxy, cyclopropyl, chloro, or –CN.

[0184] In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl.

[0185] In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl optionally substituted with one or more **R_{iss}**.

[0186] In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0187] In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or

more C₃-C₈ cycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more C₅-C₆ aryl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -O-(C₃-C₈ cycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -O-(C₅-C₆ aryl). In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -O-(5- or 6-membered heteroaryl). In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -O-(C₃-C₈ heterocycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more halo. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -CN. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -OH. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -NH₂. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more oxo.

[0188] In some embodiments, **R₁** is phenyl substituted with C₁-C₆ alkyl substituted with one or more Cl, F, Br, or I.

[0189] In some embodiments, **R₁** is phenyl substituted with C₁-C₆ alkyl substituted with one or more F.

[0190] In some embodiments, **R₁** is phenyl substituted with C₁-C₆ alkyl substituted with one or three F.

[0191] In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl.

[0192] In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl optionally substituted with one or more **R_{1SS}**.

[0193] In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0194] In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more C₅-C₆ aryl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more -O-(C₃-C₈ cycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more -O-(C₅-C₆ aryl). In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more -O-(5- or 6-membered heteroaryl). In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more -O-(C₃-C₈ heterocycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more halo. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more -CN. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more -OH. In some

embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more -NH₂. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl substituted with one or more or oxo.

[0195] In some embodiments, **R₁** is phenyl substituted with C₂-C₆ alkenyl substituted with one or more Cl, F, Br, or I.

[0196] In some embodiments, **R₁** is phenyl substituted with C₂-C₆ alkenyl substituted with one or more F.

[0197] In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl.

[0198] In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl optionally substituted with one or more **R_{1SS}**.

[0199] In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0200] In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more C₅-C₆ aryl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more -O-(C₃-C₈ cycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl substituted with

one or more -O-(C₅-C₆ aryl). In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more -O-(5- or 6-membered heteroaryl). In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more -O-(C₃-C₈ heterocycloalkyl). In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more halo. In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more -CN. In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more -OH. In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more -NH₂. In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R**₁ is phenyl substituted with one or more C₂-C₆ alkynyl substituted with one or more or oxo.

[0201] In some embodiments, **R**₁ is phenyl substituted with C₂-C₆ alkynyl substituted with one or more Cl, F, Br, or I.

[0202] In some embodiments, **R**₁ is phenyl substituted with C₂-C₆ alkynyl substituted with one or more F.

[0203] In some embodiments, **R**₁ is phenyl substituted with one or more C₃-C₈ cycloalkyl.

[0204] In some embodiments, **R**₁ is phenyl substituted with one or more C₃-C₈ cycloalkyl optionally substituted with one or more **R**_{1SS}.

[0205] In some embodiments, **R**₁ is phenyl substituted with one or more C₃-C₈ cycloalkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0206] In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more C₅-C₆ aryl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more -O-(C₃-C₈ cycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more -O-(C₅-C₆ aryl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more -O-(5- or 6-membered heteroaryl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more -O-(C₃-C₈ heterocycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more halo. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more -CN. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more -OH. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more -NH₂. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl substituted with one or more or oxo.

[0207] In some embodiments, **R₁** is phenyl substituted with C₃-C₈ cycloalkyl substituted with one or more Cl, F, Br, or I.

[0208] In some embodiments, **R₁** is phenyl substituted with C₃-C₈ cycloalkyl substituted with one or more F.

[0209] In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl.

[0210] In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl optionally substituted with one or more **R_{1SS}**.

[0211] In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0212] In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₁** is phenyl substituted with one or C₅-C₆ aryl substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more C₅-C₆ aryl. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more -O-(C₃-C₈ cycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more -O-(C₅-C₆ aryl). In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more -O-(5- or 6-membered heteroaryl). In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more -O-(C₃-C₈ heterocycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more halo. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more

–CN. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more –OH. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more –NH₂. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more –NH(C₁-C₆ alkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more –N(C₁-C₆ alkyl)₂. In some embodiments, **R₁** is phenyl substituted with one or more C₅-C₆ aryl substituted with one or more or oxo.

[0213] In some embodiments, **R₁** is phenyl substituted with C₅-C₆ aryl substituted with one or more Cl, F, Br, or I.

[0214] In some embodiments, **R₁** is phenyl substituted with C₅-C₆ aryl substituted with one or more F.

[0215] In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl.

[0216] In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl optionally substituted with one or more **R_{1SS}**.

[0217] In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, –O-(C₃-C₈ cycloalkyl), –O-(C₅-C₆ aryl), –O-(5- or 6-membered heteroaryl), –O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, –CN, –OH, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0218] In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more C₅-C₆ aryl. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-

membered heteroaryl substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more -O-(C₃-C₈ cycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more -O-(C₅-C₆ aryl). In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more -O-(5- or 6-membered heteroaryl). In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more -O-(C₃-C₈ heterocycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more halo. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more -CN. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more -OH. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more -NH₂. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl substituted with one or more oxo.

[0219] In some embodiments, **R₁** is phenyl substituted with 5- or 6-membered heteroaryl substituted with one or more Cl, F, Br, or I.

[0220] In some embodiments, **R₁** is phenyl substituted with 5- or 6-membered heteroaryl substituted with one or more F.

[0221] In some embodiments, **R₁** is phenyl substituted with one or C₃-C₈ heterocycloalkyl.

[0222] In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl optionally substituted with one or more **R_{1SS}**.

[0223] In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈

cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0224] In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more C₅-C₆ aryl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more -O-(C₃-C₈ cycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more -O-(C₅-C₆ aryl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more -O-(5- or 6-membered heteroaryl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more -O-(C₃-C₈ heterocycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more halo. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more -CN. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more -OH. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more -NH₂. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with

one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl substituted with one or more or oxo.

[0225] In some embodiments, **R₁** is phenyl substituted with C₃-C₈ heterocycloalkyl substituted with one or more Cl, F, Br, or I.

[0226] In some embodiments, **R₁** is phenyl substituted with C₃-C₈ heterocycloalkyl substituted with one or more F.

[0227] In some embodiments, **R₁** is phenyl substituted with one or C₁-C₆ alkoxy.

[0228] In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy optionally substituted with one or more **R_{1SS}**.


[0229] In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.


[0230] In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₅-C₆ aryl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more -O-(C₃-C₈ cycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more -O-(C₅-C₆ aryl). In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more -O-(5- or 6-membered heteroaryl). In some

embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more -O-(C₃-C₈ heterocycloalkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more halo. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more -CN. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more -OH. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more -NH₂. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more oxo.

[0231] In some embodiments, **R₁** is phenyl substituted with C₁-C₆ alkoxy substituted with one or more Cl, F, Br, or I.

[0232] In some embodiments, **R₁** is phenyl substituted with C₁-C₆ alkoxy substituted with one or more F.





[0234] In some embodiments, at least one **R_{1S}** is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**.

[0235] In some embodiments, at least one **R_{1S}** is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN.

[0236] In some embodiments, at least one **R_{1S}** is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, or C₁-C₆ alkoxy.

[0237] In some embodiments, at least one **R_{1S}** is C₁-C₆ alkyl (*e.g.* linear or branched).

[0238] In some embodiments, at least one **R_{1S}** is C₁-C₆ alkyl (*e.g.* linear or branched) optionally substituted with one or more **R_{1SS}**.

[0239] In some embodiments, at least one **R_{1S}** is methyl. In some embodiments, at least one **R_{1S}** is ethyl. In some embodiments, at least one **R_{1S}** is propyl. In some embodiments, at least one **R_{1S}** is butyl. In some embodiments, at least one **R_{1S}** is pentyl. In some embodiments, at least one **R_{1S}** is hexyl. In some embodiments, at least one **R_{1S}** is isopropyl. In some embodiments, at least one **R_{1S}** is isobutyl. In some embodiments, at least one **R_{1S}** is isopentyl. In some embodiments, at least one **R_{1S}** is isohexyl. In some embodiments, at least one **R_{1S}** is secbutyl. In some embodiments, at least one **R_{1S}** is secpentyl. In some embodiments, at least one **R_{1S}** is sechexyl. In some embodiments, at least one **R_{1S}** is tertbutyl.

[0240] In some embodiments, at least one **R_{1S}** is C₂-C₆ alkenyl. In some embodiments, at least one **R_{1S}** is C₂-C₆ alkynyl.

[0241] In some embodiments, at least one **R_{1S}** is C₂-C₆ alkenyl optionally substituted with one or more **R_{1SS}**. In some embodiments, at least one **R_{1S}** is C₂-C₆ alkynyl optionally substituted with one or more **R_{1SS}**.

[0242] In some embodiments, at least one **R_{1S}** is C₁-C₆ alkoxy (*e.g.* methyl, ethyl, propyl, butyl, pentyl, or hexyl substituted with one or more halogen).

[0243] In some embodiments, at least one **R_{1S}** is methoxy. In some embodiments, at least one **R_{1S}** is ethoxy. In some embodiments, at least one **R_{1S}** is propoxy. In some embodiments, at least one **R_{1S}** is butoxy. In some embodiments, at least one **R_{1S}** is pentoxy. In some embodiments, at least one **R_{1S}** is hexoxy.

[0244] In some embodiments, at least one **R_{1S}** is C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0245] In some embodiments, at least one **R_{1S}** is C₃-C₈ cycloalkyl.

[0246] In some embodiments, at least one **R_{1S}** is C₃-C₈ cycloalkyl optionally substituted with one or more **R_{1SS}**.

[0247] In some embodiments, at least one **R_{1S}** is cyclopropyl. In some embodiments, at least one **R_{1S}** is cyclobutyl. In some embodiments, at least one **R_{1S}** is cyclopentyl. In some embodiments, at least one **R_{1S}** is cyclohexyl. In some embodiments, at least one **R_{1S}** is cycloheptyl. In some embodiments, at least one **R_{1S}** is cyclooctyl.

[0248] In some embodiments, at least one **R_{1S}** is C₅-C₆ aryl.

[0249] In some embodiments, at least one **R_{1S}** is C₅-C₆ aryl optionally substituted with one or more **R_{1SS}**. In some embodiments, at least one **R_{1S}** is C₅ aryl. In some embodiments, at least one **R_{1S}** is C₆ aryl.

[0250] In some embodiments, at least one **R_{1S}** is C₆ phenyl.

[0251] In some embodiments, at least one **R_{1S}** is 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0252] In some embodiments, at least one **R_{1S}** is 5- or 6-membered heteroaryl.

[0253] In some embodiments, at least one **R_{1S}** is 5- or 6-membered heteroaryl optionally substituted with one or more **R_{1SS}**. In some embodiments, at least one **R_{1S}** is 5-membered heteroaryl. In some embodiments, at least one **R_{1S}** is 6-membered heteroaryl.

[0254] In some embodiments, at least one **R_{1S}** is C₃-C₈ heterocycloalkyl.

[0255] In some embodiments, at least one **R_{1S}** is C₃-C₈ heterocycloalkyl optionally substituted with one or more **R_{1SS}**.

[0256] In some embodiments, at least one **R_{1S}** is C₃ heterocycloalkyl. In some embodiments, at least one **R_{1S}** is C₄ heterocycloalkyl. In some embodiments, at least one **R_{1S}** is C₅ heterocycloalkyl. In some embodiments, at least one **R_{1S}** is C₆ heterocycloalkyl. In some embodiments, at least one **R_{1S}** is C₇ heterocycloalkyl. In some embodiments, at least one **R_{1S}** is C₈ heterocycloalkyl.

[0257] In some embodiments, at least one **R_{1S}** is halo or CN.

[0258] In some embodiments, at least one **R_{1S}** is halo. In some embodiments, at least one **R_{1S}** is F, Cl, Br, or I. In some embodiments, at least one **R_{1S}** is F or Cl. In some embodiments, at least one **R_{1S}** is F. In some embodiments, at least one **R_{1S}** is Cl.

[0259] In some embodiments, at least one **R_{1S}** is CN.

[0260] In some embodiments, at least one **R_{1S}** is Cl, F, CN, methyl, methoxy, ethoxy, or cyclopropyl.

[0261] In some embodiments, at least one **R_{1SS}** is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0262] In some embodiments, at least one **R_{1SS}** is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0263] In some embodiments, when **R_{1S}** is ethenyl or ethynyl, then each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂.

[0264] In some embodiments, at least one **R_{1SS}** is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl.

[0265] In some embodiments, at least one **R_{1SS}** is C₁-C₆ alkyl (*e.g.* linear or branched).

[0266] In some embodiments, at least one **R**_{1SS} is methyl. In some embodiments, at least one **R**_{1SS} is ethyl. In some embodiments, at least one **R**_{1SS} is propyl. In some embodiments, at least one **R**_{1SS} is butyl. In some embodiments, at least one **R**_{1SS} is pentyl. In some embodiments, at least one **R**_{1SS} is hexyl. In some embodiments, at least one **R**_{1SS} is isopropyl. In some embodiments, at least one **R**_{1SS} is isobutyl. In some embodiments, at least one **R**_{1SS} is isopentyl. In some embodiments, at least one **R**_{1SS} is isohexyl. In some embodiments, at least one **R**_{1SS} is secbutyl. In some embodiments, at least one **R**_{1SS} is secpentyl. In some embodiments, at least one **R**_{1SS} is sechexyl. In some embodiments, at least one **R**_{1SS} is tertbutyl.

[0267] In some embodiments, at least one **R**_{1SS} is C₂-C₆ alkenyl. In some embodiments, at least one **R**_{1SS} is C₂-C₆ alkynyl.

[0268] In some embodiments, at least one **R**_{1SS} is C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0269] In some embodiments, at least one **R**_{1SS} is C₃-C₈ cycloalkyl.

[0270] In some embodiments, at least one **R**_{1SS} is cyclopropyl. In some embodiments, at least one **R**_{1SS} is cyclobutyl. In some embodiments, at least one **R**_{1SS} is cyclopentyl. In some embodiments, at least one **R**_{1SS} is cyclohexyl. In some embodiments, at least one **R**_{1SS} is cycloheptyl. In some embodiments, at least one **R**_{1SS} is cyclooctyl.

[0271] In some embodiments, at least one **R**_{1SS} is C₅-C₆ aryl. In some embodiments, at least one **R**_{1SS} is C₅ aryl. In some embodiments, at least one **R**_{1SS} is C₆ aryl.

[0272] In some embodiments, at least one **R**_{1SS} is phenyl.

[0273] In some embodiments, at least one **R**_{1SS} 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0274] In some embodiments, at least one **R**_{1SS} 5- or 6-membered heteroaryl. In some embodiments, at least one **R**_{1SS} 5-membered heteroaryl. In some embodiments, at least one **R**_{1SS} 6-membered heteroaryl.

[0275] In some embodiments, at least one **R**_{1SS} C₃-C₈ heterocycloalkyl.

[0276] In some embodiments, at least one **R**_{1SS} is C₃ heterocycloalkyl. In some embodiments, at least one **R**_{1SS} is C₄ heterocycloalkyl. In some embodiments, at least one **R**_{1SS} is C₅ heterocycloalkyl. In some embodiments, at least one **R**_{1SS} is C₆ heterocycloalkyl. In some embodiments, at least one **R**_{1SS} is C₇ heterocycloalkyl. In some embodiments, at least one **R**_{1SS} is C₈ heterocycloalkyl.

- [0277] In some embodiments, at least one **R**_{1SS} is -OH, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), or -O-(C₃-C₈ heterocycloalkyl).
- [0278] In some embodiments, at least one **R**_{1SS} is -OH.
- [0279] In some embodiments, at least one **R**_{1SS} is -O-(C₃-C₈ cycloalkyl) or -O-(C₅-C₆ aryl).
- [0280] In some embodiments, at least one **R**_{1SS} is -O-(C₃-C₈ cycloalkyl).
- [0281] In some embodiments, at least one **R**_{1SS} is -O-(cyclopropyl). In some embodiments, at least one **R**_{1SS} is -O-(cyclobutyl). In some embodiments, at least one **R**_{1SS} is -O-(cyclopentyl). In some embodiments, at least one **R**_{1SS} is -O-(cyclohexyl). In some embodiments, at least one **R**_{1SS} is -O-(cycloheptyl). In some embodiments, at least one **R**_{1SS} is -O-(cyclooctyl).
- [0282] In some embodiments, at least one **R**_{1SS} is -O-(C₅-C₆ aryl).
- [0283] In some embodiments, at least one **R**_{1SS} is -O-(C₅ aryl). In some embodiments, at least one **R**_{1SS} is -O-(C₆ aryl).
- [0284] In some embodiments, at least one **R**_{1SS} is -O-(phenyl).
- [0285] In some embodiments, at least one **R**_{1SS} is -O-(5- or 6-membered heteroaryl) or -O-(C₃-C₈ heterocycloalkyl).
- [0286] In some embodiments, at least one **R**_{1SS} is -O-(5- or 6-membered heteroaryl).
- [0287] In some embodiments, at least one **R**_{1SS} is -O-(5-membered heteroaryl). In some embodiments, at least one **R**_{1SS} is -O-(6-membered heteroaryl).
- [0288] In some embodiments, at least one **R**_{1SS} is -O-(C₃-C₈ heterocycloalkyl).
- [0289] In some embodiments, at least one **R**_{1SS} is -O-(C₃ heterocycloalkyl). In some embodiments, at least one **R**_{1SS} is -O-(C₄ heterocycloalkyl). In some embodiments, at least one **R**_{1SS} is -O-(C₅ heterocycloalkyl). In some embodiments, at least one **R**_{1SS} is -O-(C₆ heterocycloalkyl). In some embodiments, at least one **R**_{1SS} is -O-(C₇ heterocycloalkyl). In some embodiments, at least one **R**_{1SS} is -O-(C₈ heterocycloalkyl).
- [0290] In some embodiments, at least one **R**_{1SS} is C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.
- [0291] In some embodiments, at least one **R**_{1SS} is C₁-C₆ haloalkyl.
- [0292] In some embodiments, at least one **R**_{1SS} is halomethyl. In some embodiments, at least one **R**_{1SS} is haloethyl. In some embodiments, at least one **R**_{1SS} is halopropyl. In some embodiments, at least one **R**_{1SS} is halobutyl. In some embodiments, at least one **R**_{1SS} is halopentyl. In some embodiments, at least one **R**_{1SS} is haloheptyl.

- [0293] In some embodiments, at least one **R**_{1SS} is C₁-C₆ alkoxy.
- [0294] In some embodiments, at least one **R**_{1SS} is methoxy. In some embodiments, at least one **R**_{1SS} is ethoxy. In some embodiments, at least one **R**_{1SS} is propoxy. In some embodiments, at least one **R**_{1SS} is butoxy. In some embodiments, at least one **R**_{1SS} is pentoxy. In some embodiments, at least one **R**_{1SS} is hexoxy.
- [0295] In some embodiments, at least one **R**_{1SS} is C₁-C₆ haloalkoxy.
- [0296] In some embodiments, at least one **R**_{1SS} is halomethoxy. In some embodiments, at least one **R**_{1SS} is haloethoxy. In some embodiments, at least one **R**_{1SS} is halopropoxy. In some embodiments, at least one **R**_{1SS} is halobutoxy. In some embodiments, at least one **R**_{1SS} is halopentoxy. In some embodiments, at least one **R**_{1SS} is haloethoxy.
- [0297] In some embodiments, at least one **R**_{1SS} is halo, -CN, or oxo.
- [0298] In some embodiments, at least one **R**_{1SS} is halo. In some embodiments, at least one **R**_{1SS} is F, Cl, Br, or I. In some embodiments, at least one **R**_{1SS} is F or Cl. In some embodiments, at least one **R**_{1SS} is F. In some embodiments, at least one **R**_{1SS} is Cl.
- [0299] In some embodiments, at least one **R**_{1SS} is -CN.
- [0300] In some embodiments, at least one **R**_{1SS} is oxo.
- [0301] In some embodiments, at least one **R**_{1SS} is -NH₂, -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
- [0302] In some embodiments, at least one **R**_{1SS} is -NH₂. In some embodiments, at least one **R**_{1SS} is -NH(C₁-C₆ alkyl).
- [0303] In some embodiments, at least one **R**_{1SS} is -NH(methyl). In some embodiments, at least one **R**_{1SS} is -NH(ethyl). In some embodiments, at least one **R**_{1SS} is -NH(propyl). In some embodiments, at least one **R**_{1SS} is -NH(butyl). In some embodiments, at least one **R**_{1SS} is -NH(pentyl). In some embodiments, at least one **R**_{1SS} is -NH(hexyl).
- [0304] In some embodiments, at least one **R**_{1SS} is -N(C₁-C₆ alkyl)₂.
- [0305] In some embodiments, at least one **R**_{1SS} is F, methyl, phenyl, or -O-(phenyl).
- [0306] In some embodiments, **Q** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
- [0307] In some embodiments, **Q** is C₁-C₆ alkyl.

[0308] In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0309] In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more halo. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more -CN. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more -OH. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more -NH₂. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **Q** is C₁-C₆ alkyl optionally substituted with one or more oxo.

[0310] In some embodiments, **Q** is methyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0311] In some embodiments, **Q** is methyl.

[0312] In some embodiments, **Q** is methyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **Q** is methyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **Q** is methyl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **Q** is methyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **Q** is methyl optionally substituted with one or more C₅-C₆ aryl. In some

embodiments, **Q** is methyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **Q** is methyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **Q** is methyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **Q** is methyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **Q** is methyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **Q** is methyl optionally substituted with one or more halo. In some embodiments, **Q** is methyl optionally substituted with one or more -CN. In some embodiments, **Q** is methyl optionally substituted with one or more -OH. In some embodiments, **Q** is methyl optionally substituted with one or more -NH₂. In some embodiments, **Q** is methyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **Q** is methyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **Q** is methyl optionally substituted with one or more oxo.

[0313] In some embodiments, **Q** is C₂-C₆ alkenyl.

[0314] In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0315] In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more halo. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more -CN. In some embodiments, **Q** is C₂-C₆ alkenyl optionally substituted with one or more -OH. In some embodiments, **Q** is C₂-C₆ alkenyl

optionally substituted with one or more -NH_2 . In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkenyl optionally substituted with one or more $\text{-NH}(\text{C}_1\text{-C}_6 \text{ alkyl})$. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkenyl optionally substituted with one or more $\text{-N}(\text{C}_1\text{-C}_6 \text{ alkyl})_2$. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkenyl optionally substituted with one or more oxo.

[0316] In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl.

[0317] In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_5\text{-C}_6$ aryl, 5- or 6-membered heteroaryl, $\text{C}_3\text{-C}_8$ heterocycloalkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ haloalkoxy, halo, -CN , -OH , -NH_2 , $\text{-NH}(\text{C}_1\text{-C}_6 \text{ alkyl})$, $\text{-N}(\text{C}_1\text{-C}_6 \text{ alkyl})_2$, or oxo.

[0318] In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_1\text{-C}_6$ alkyl. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_2\text{-C}_6$ alkenyl. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_2\text{-C}_6$ alkynyl. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_3\text{-C}_8$ cycloalkyl. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_5\text{-C}_6$ aryl. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_3\text{-C}_8$ heterocycloalkyl. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_1\text{-C}_6$ haloalkyl. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_1\text{-C}_6$ alkoxy. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{C}_1\text{-C}_6$ haloalkoxy. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more halo. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more -CN . In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more -OH . In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more -NH_2 . In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{-NH}(\text{C}_1\text{-C}_6 \text{ alkyl})$. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more $\text{-N}(\text{C}_1\text{-C}_6 \text{ alkyl})_2$. In some embodiments, **Q** is $\text{C}_2\text{-C}_6$ alkynyl optionally substituted with one or more oxo.

[0319] In some embodiments, **R**₂ is 5- to 10-membered heteroaryl or 3- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_5\text{-C}_6$ aryl, 5- or 6-membered heteroaryl, $\text{C}_3\text{-C}_8$ heterocycloalkyl, $\text{C}_1\text{-}$

C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0320] In some embodiments, **R₂** is 5- to 10-membered heteroaryl.

[0321] In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0322] In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more halo. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more -CN. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more -OH. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more -NH₂. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₂** is 5- to 10-membered heteroaryl optionally substituted with one or more oxo.

[0323] In some embodiments, **R₂** is 5- to 9-membered heteroaryl.

[0324] In some embodiments, **R₂** is 5- to 9-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-

membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0325] In some embodiments, **R₂** is 5- to 8-membered heteroaryl.

[0326] In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0327] In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more halo. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more -CN. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more -OH. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more -NH₂. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₂** is 5- to 8-membered heteroaryl optionally substituted with one or more oxo.

[0328] In some embodiments, **R₂** is 5- to 7-membered heteroaryl.

[0329] In some embodiments, **R₂** is 5- to 7-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-

membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0330] In some embodiments, **R₂** is 5- or 6-membered heteroaryl.

[0331] In some embodiments, **R₂** is 5- or 6-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0332] In some embodiments, **R₂** is 5-membered heteroaryl.

[0333] In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0334] In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more halo. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more -CN. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more -OH. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more -NH₂. In some embodiments, **R₂** is 5-membered heteroaryl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₂** is 5-membered heteroaryl optionally

substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R**₂ is 5-membered heteroaryl optionally substituted with one or more oxo.

[0335] In some embodiments, **R**₂ is 6-membered heteroaryl.

[0336] In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0337] In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more halo. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more -CN. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more -OH. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more -NH₂. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R**₂ is 6-membered heteroaryl optionally substituted with one or more oxo.

[0338] In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl.

[0339] In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or

6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0340] In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more halo. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -CN. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -OH. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -NH₂. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R**₂ is 4- to 8-membered heterocycloalkyl optionally substituted with one or more oxo.

[0341] In some embodiments, **R**₂ is 4-membered heterocycloalkyl.

[0342] In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0343] In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally

substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more halo. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more -CN. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more -OH. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more -NH₂. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R**₂ is 4-membered heterocycloalkyl optionally substituted with one or more oxo.

[0344] In some embodiments, **R**₂ is 5-membered heterocycloalkyl.

[0345] In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0346] In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted

with one or more 5- or 6-membered heteroaryl. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more halo. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more -CN. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more -OH. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more -NH₂. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R**₂ is 5-membered heterocycloalkyl optionally substituted with one or more oxo.

[0347] In some embodiments, **R**₂ is 6-membered heterocycloalkyl.

[0348] In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0349] In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally

substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more halo. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more -CN. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more -OH. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more -NH₂. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R**₂ is 6-membered heterocycloalkyl optionally substituted with one or more oxo.

[0350] In some embodiments, **R**₂ is 7-membered heterocycloalkyl.

[0351] In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0352] In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more halo. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more -CN. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more -OH. In some embodiments, **R**₂ is 7-membered heterocycloalkyl optionally substituted with one or more -NH₂.

In some embodiments, **R₂** is 7-membered heterocycloalkyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₂** is 7-membered heterocycloalkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₂** is 7-membered heterocycloalkyl optionally substituted with one or more oxo.

[0353] In some embodiments, **R₂** is 8-membered heterocycloalkyl.

[0354] In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0355] In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₂-C₆ alkynyl. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more halo. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more -CN. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more -OH. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more -NH₂. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂. In some embodiments, **R₂** is 8-membered heterocycloalkyl optionally substituted with one or more oxo.

[0356] In some embodiments, **R₂** is pyrazolyl.

[0357] In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0358] In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more methyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more ethyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more propyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more butyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more pentyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more hexyl.

[0359] In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₂-C₆ alkynyl.

[0360] In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more cyclopropyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more cyclobutyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more cyclopentyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more cyclohexyl.

[0361] In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₅ aryl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₆ aryl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more phenyl.

[0362] In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more 5-membered heteroaryl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more 6-membered heteroaryl.

[0363] In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or more C₃ heterocycloalkyl. In some embodiments, **R₂** is pyrazolyl optionally substituted with one or

more C₄ heterocycloalkyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more C₅ heterocycloalkyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more C₆ heterocycloalkyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more C₇ heterocycloalkyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more C₈ heterocycloalkyl.

[0364] In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more halomethyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more haloethyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more halopropyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more halobutyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more halopentyl. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more haloethyl.

[0365] In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more methoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more ethoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more propoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more butoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more pentoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more hexoxy.

[0366] In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more halomethoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more haloethoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more halopropoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more halobutoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more halopentoxy. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more haloethoxy.

[0367] In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more halo. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more F or Cl. In some

embodiments, **R**₂ is pyrazolyl optionally substituted with one or more F. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more Cl.

[0368] In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –CN. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –OH. In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –NH₂.

[0369] In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –NH(C₁–C₆ alkyl). In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –NH(methyl). In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –NH(ethyl). In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –NH(propyl). In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –NH(butyl). In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –NH(pentyl). In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –NH(hexyl).

[0370] In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more –N(C₁–C₆ alkyl)₂.

[0371] In some embodiments, **R**₂ is pyrazolyl optionally substituted with one or more oxo.

[0372] In some embodiments, **R**₂ is .

[0373] In some embodiments, **R**₂ is triazolyl.

[0374] In some embodiments, **R**₂ is triazolyl optionally substituted with one or more C₁–C₆ alkyl, C₂–C₆ alkenyl, C₂–C₆ alkynyl, C₃–C₈ cycloalkyl, C₅–C₆ aryl, 5- or 6-membered heteroaryl, C₃–C₈ heterocycloalkyl, C₁–C₆ haloalkyl, C₁–C₆ alkoxy, C₁–C₆ haloalkoxy, halo, –CN, –OH, –NH₂, –NH(C₁–C₆ alkyl), –N(C₁–C₆ alkyl)₂, or oxo.

[0375] In some embodiments, **R**₂ is triazolyl optionally substituted with one or more C₁–C₆ alkyl. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more methyl. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more ethyl. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more propyl. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more butyl. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more pentyl. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more hexyl.

[0376] In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₂-C₆ alkynyl.

[0377] In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more cyclopropyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more cyclobutyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more cyclopentyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more cyclohexyl.

[0378] In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₅ aryl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₆ aryl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more phenyl.

[0379] In some embodiments, **R₂** is triazolyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more 5-membered heteroaryl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more 6-membered heteroaryl.

[0380] In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₃ heterocycloalkyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₄ heterocycloalkyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₅ heterocycloalkyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₆ heterocycloalkyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₇ heterocycloalkyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₈ heterocycloalkyl.

[0381] In some embodiments, **R₂** is triazolyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more halomethyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more haloethyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more halopropyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more halobutyl. In some embodiments, **R₂** is triazolyl optionally substituted with one or more

halopentyl. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more haloheptyl.

[0382] In some embodiments, **R**₂ is triazolyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more methoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more ethoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more propoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more butoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more pentoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more hexoxy.

[0383] In some embodiments, **R**₂ is triazolyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more halomethoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more haloethoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more halopropoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more halobutoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more halopentoxy. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more haloheptyl.

[0384] In some embodiments, **R**₂ is triazolyl optionally substituted with one or more halo. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more F or Cl. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more F. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more Cl.

[0385] In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -CN. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -OH. In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -NH₂.

[0386] In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -NH(methyl). In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -NH(ethyl). In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -NH(propyl). In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -NH(butyl). In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -

NH(pentyl). In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -NH(hexyl).

[0387] In some embodiments, **R**₂ is triazolyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0388] In some embodiments, **R**₂ is triazolyl optionally substituted with one or more oxo.

[0389] In some embodiments, **R**₂ is .

[0390] In some embodiments, **R**₂ is pyrimidinyl.

[0391] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0392] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more C₁-C₆ alkyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more methyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more ethyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more propyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more butyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more pentyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more hexyl.

[0393] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more C₂-C₆ alkenyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more C₂-C₆ alkynyl.

[0394] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more C₃-C₈ cycloalkyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more cyclopropyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more cyclobutyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more cyclopentyl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more cyclohexyl.

[0395] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more C₅-C₆ aryl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more C₅ aryl. In

some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₆ aryl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more phenyl.

[0396] In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more 5- or 6-membered heteroaryl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more 5-membered heteroaryl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more 6-membered heteroaryl.

[0397] In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₃-C₈ heterocycloalkyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₃ heterocycloalkyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₄ heterocycloalkyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₅ heterocycloalkyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₆ heterocycloalkyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₇ heterocycloalkyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₈ heterocycloalkyl.

[0398] In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₁-C₆ haloalkyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more halomethyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more haloethyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more halopropyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more halobutyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more halopentyl. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more haloethyl.

[0399] In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more C₁-C₆ alkoxy. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more methoxy. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more ethoxy. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more propoxy. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more butoxy. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more pentoxy. In some embodiments, **R₂** is pyrimidinyl optionally substituted with one or more hexoxy.

[0400] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more C₁-C₆ haloalkoxy. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more halomethoxy. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more haloethoxy. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more halopropoxy. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more halobutoxy. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more halopentoxy. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more halohexoxy.

[0401] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more halo. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more F or Cl. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more F. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more Cl.

[0402] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -CN. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -OH. In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -NH₂.




[0403] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -NH(C₁-C₆ alkyl). In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -NH(methyl). In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -NH(ethyl). In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -NH(propyl). In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -NH(butyl). In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -NH(pentyl). In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -NH(hexyl).




[0404] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.


[0405] In some embodiments, **R**₂ is pyrimidinyl optionally substituted with one or more oxo.

[0406] In some embodiments, **R**₂ is .

[0407] In some embodiments, **R**₂ is , , or .

[0408] In some embodiments, **Q** and **R₂** together forms , , or .

[0409] In some embodiments, **Q** and **R₂** together forms , , ,



[0410] In some embodiments, **R₃** is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0411] In some embodiments, **R₃** is H.

[0412] In some embodiments, **R₃** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0413] In some embodiments, **R₃** is C₂-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₂-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0414] In some embodiments, **R₃** is C₁-C₆ alkyl, C₃-C₆ alkenyl, or C₃-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₃-C₆ alkenyl, or C₃-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0415] In some embodiments, **R₃** is C₂-C₆ alkyl, C₃-C₆ alkenyl, or C₃-C₆ alkynyl; wherein the C₂-C₆ alkyl, C₃-C₆ alkenyl, or C₃-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0416] In some embodiments, **R₃** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0417] In some embodiments, **R₃** is C₁-C₆ alkyl. In some embodiments, **R₃** is methyl. In some embodiments, **R₃** is ethyl. In some embodiments, **R₃** is propyl. In some embodiments, **R₃** is butyl. In some embodiments, **R₃** is pentyl. In some embodiments, **R₃** is hexyl. In some embodiments, **R₃** is isopropyl. In some embodiments, at least one **R₃** is isobutyl. In some embodiments, at least one **R₃** is isopentyl. In some embodiments, at least one **R₃** is isohexyl. In some embodiments, at least one **R₃** is secbutyl. In some embodiments, at least one **R₃** is secpentyl. In some embodiments, at least one **R₃** is sechexyl. In some embodiments, at least one **R₃** is tertbutyl.

[0418] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R_{3s}**.

[0419] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R_{3s}**.

[0420] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0421] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, or **R_{3s}**.

[0422] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, or halo.

[0423] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0424] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0425] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl.

[0426] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkenyl.

[0427] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkynyl.

[0428] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0429] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0430] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₅-C₆ aryl.

[0431] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0432] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0433] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0434] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0435] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkyl.

- [0436] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkoxy.
- [0437] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkoxy.
- [0438] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.
- [0439] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.
- [0440] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more halo or -CN.
- [0441] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.
- [0442] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more halo.
- [0443] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more F, Cl, Br, or I.
- [0444] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more F or Cl.
- [0445] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more F.
- [0446] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more Cl.
- [0447] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -CN.
- [0448] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -OH.
- [0449] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -NH₂.
- [0450] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more oxo.
- [0451] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
- [0452] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl).
- [0453] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.
- [0454] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more -OH.
- [0455] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more -NH₂.
- [0456] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more oxo.

- [0457] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
- [0458] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl).
- [0459] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.
- [0460] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more **R_{3S}**.
- [0461] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more **R_{3S}**.
- [0462] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -OC(=O)**R_{3Sa}** or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
- [0463] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more -OC(=O)**R_{3Sa}** or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
- [0464] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -OC(=O)**R_{3Sa}**.
- [0465] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl.
- [0466] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)**R_{3Sb}**.
- [0467] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more -OH.
- [0468] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more halo.
- [0469] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more -CN.
- [0470] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more -OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more -NH₂.

[0471] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more –NH(C₁-C₆ alkyl).

[0472] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more –N(C₁-C₆ alkyl)₂.

[0473] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more oxo.

[0474] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more –C(=O)**R_{3Sb}**.

[0475] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**.

[0476] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein **R_{3Sb}** is 4- to 8-membered heterocycloalkyl or –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more –OH, halo, –CN, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0477] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted one or more –OH, halo, –CN, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0478] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted one or more –OH.

[0479] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein **R_{3Sb}** is 4- to 8-membered heterocycloalkyl.

[0480] In some embodiments, **R₃** is C₁-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein

R_{3Sb} is $-(C_1-C_6 \text{ alkyl})-(4\text{- to } 8\text{-membered heterocycloalkyl})$ optionally substituted with one or more $-OH$, halo, $-CN$, $-NH_2$, $-NH(C_1-C_6 \text{ alkyl})$, $-N(C_1-C_6 \text{ alkyl})_2$, or oxo.

[0481] In some embodiments, **R₃** is C_1-C_6 alkyl optionally substituted with one or more $-OC(=O)R_{3Sa}$; wherein **R_{3Sa}** is C_1-C_6 alkyl substituted with one or more $-C(=O)OR_{3Sb}$; wherein **R_{3Sb}** is $-(C_1-C_6 \text{ alkyl})-(4\text{- to } 8\text{-membered heterocycloalkyl})$ optionally substituted with one or more $-OH$.

[0482] In some embodiments, **R₃** is C_1-C_6 alkyl optionally substituted with one or more $-OC(=O)R_{3Sa}$; wherein **R_{3Sa}** is C_1-C_6 alkyl substituted with one or more $-C(=O)OR_{3Sb}$; wherein **R_{3Sb}** is $-(C_1-C_6 \text{ alkyl})-(4\text{- to } 8\text{-membered heterocycloalkyl})$.

[0483] In some embodiments, **R₃** is C_1-C_6 alkyl optionally substituted with one or more 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more $-OH$, halo, $-CN$, $-NH_2$, $-NH(C_1-C_6 \text{ alkyl})$, $-N(C_1-C_6 \text{ alkyl})_2$, or oxo.

[0484] In some embodiments, **R₃** is C_1-C_6 alkyl optionally substituted with one or more 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more $-OH$.

[0485] In some embodiments, **R₃** is C_1-C_6 alkyl optionally substituted with one or more 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one, two, or three $-OH$.

[0486] In some embodiments, **R₃** is C_1-C_6 alkyl optionally substituted with one or more 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with three $-OH$.

[0487] In some embodiments, **R₃** is C_2-C_6 alkyl optionally substituted with one or more $-OC(=O)R_{3Sa}$.

[0488] In some embodiments, **R₃** is C_2-C_6 alkyl optionally substituted with one or more $-OC(=O)R_{3Sa}$; wherein **R_{3Sa}** is C_1-C_6 alkyl optionally substituted with one or more $-OH$, halo, $-CN$, $-NH_2$, $-NH(C_1-C_6 \text{ alkyl})$, $-N(C_1-C_6 \text{ alkyl})_2$, oxo, or $-C(=O)OR_{3Sb}$.

[0489] In some embodiments, **R₃** is C_2-C_6 alkyl optionally substituted with one or more $-OC(=O)R_{3Sa}$; wherein **R_{3Sa}** is C_1-C_6 alkyl.

[0490] In some embodiments, **R₃** is C_2-C_6 alkyl optionally substituted with one or more $-OC(=O)R_{3Sa}$; wherein **R_{3Sa}** is C_1-C_6 alkyl optionally substituted with one or more $-C(=O)OR_{3Sb}$.

[0491] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**.

[0492] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein **R_{3Sb}** is 4- to 8-membered heterocycloalkyl or –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more –OH, halo, –CN, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0493] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more –OH, halo, –CN, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0494] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more –OH.

[0495] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein **R_{3Sb}** is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more –OH, halo, –CN, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0496] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more –OC(=O)**R_{3Sa}**; wherein **R_{3Sa}** is C₁-C₆ alkyl substituted with one or more –C(=O)**R_{3Sb}**; wherein **R_{3Sb}** is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more –OH.

[0497] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more –OH, halo, –CN, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0498] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more –OH.

[0499] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one, two, or three -OH.

[0500] In some embodiments, **R₃** is C₂-C₆ alkyl optionally substituted with one or more 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with three -OH.

[0501] In some embodiments, **R₃** is ethyl.

[0502] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0503] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0504] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₁-C₆ alkyl.

[0505] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₂-C₆ alkenyl.

[0506] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₂-C₆ alkynyl.

[0507] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0508] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0509] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₅-C₆ aryl.

[0510] In some embodiments, **R₃** is ethyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0511] In some embodiments, **R₃** is ethyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0512] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0513] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0514] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0515] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₁-C₆ alkoxy.

[0516] In some embodiments, **R₃** is ethyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0517] In some embodiments, **R₃** is ethyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0518] In some embodiments, **R₃** is ethyl optionally substituted with one or more halo. In some embodiments, **R₃** is ethyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R₃** is ethyl optionally substituted with one or more F or Cl. In some embodiments, **R₃** is ethyl optionally substituted with one or more F. In some embodiments, **R₃** is ethyl optionally substituted with one or more Cl.

[0519] In some embodiments, **R₃** is ethyl optionally substituted with one or more -CN.

[0520] In some embodiments, **R₃** is ethyl optionally substituted with one or more -OH.

[0521] In some embodiments, **R₃** is ethyl optionally substituted with one or more -NH₂.

[0522] In some embodiments, **R₃** is ethyl optionally substituted with one or more oxo.

[0523] In some embodiments, **R₃** is ethyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0524] In some embodiments, **R₃** is ethyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0525] In some embodiments, **R₃** is ethyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0526] In some embodiments, **R₃** is isopropyl. In some embodiments, **R₃** is isopropyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0527] In some embodiments, **R₃** is isopropyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0528] In some embodiments, **R₃** is isopropyl optionally substituted with one or more C₁-C₆ alkyl.

[0529] In some embodiments, **R₃** is isopropyl optionally substituted with one or more C₂-C₆ alkenyl.

[0530] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more C₂-C₈ alkynyl.

[0531] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0532] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0533] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more C₅-C₆ aryl.

[0534] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0535] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0536] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0537] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0538] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0539] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more C₁-C₆ alkoxy.

[0540] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0541] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0542] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more halo. In some embodiments, **R**₃ is isopropyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R**₃ is isopropyl optionally substituted with one or more F or Cl. In some embodiments, **R**₃ is isopropyl optionally substituted with one or more F. In some embodiments, **R**₃ is isopropyl optionally substituted with one or more Cl.

[0543] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more -CN.

[0544] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more -OH.

[0545] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more -NH₂.

[0546] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more oxo.

[0547] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0548] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0549] In some embodiments, **R**₃ is isopropyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0550] In some embodiments, **R**₃ is C₂-C₆ alkenyl.

[0551] In some embodiments, **R**₃ is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0552] In some embodiments, **R**₃ is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, or -N(C₁-C₆ alkyl)₂.

[0553] In some embodiments, **R**₃ is C₃-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0554] In some embodiments, **R**₃ is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0555] In some embodiments, **R**₃ is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl.

[0556] In some embodiments, **R**₃ is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkenyl.

[0557] In some embodiments, **R**₃ is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkynyl.

[0558] In some embodiments, **R**₃ is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0559] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0560] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more C₅-C₆ aryl.

[0561] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0562] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0563] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0564] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0565] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0566] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkoxy.

[0567] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0568] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0569] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more halo or -CN.

[0570] In some embodiments, **R₃** is C₃-C₆ alkenyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0571] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more halo.

[0572] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more F, Cl, Br, or I.

[0573] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more F or Cl.

[0574] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more F.

[0575] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more Cl.

- [0576] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more -CN.
- [0577] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more -OH.
- [0578] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more -NH₂.
- [0579] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more oxo.
- [0580] In some embodiments, **R₃** is C₃-C₆ alkenyl optionally substituted with one or more -OH.
- [0581] In some embodiments, **R₃** is C₃-C₆ alkenyl optionally substituted with one or more -NH₂.
- [0582] In some embodiments, **R₃** is C₃-C₆ alkenyl optionally substituted with one or more oxo.
- [0583] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
- [0584] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl).
- [0585] In some embodiments, **R₃** is C₂-C₆ alkenyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.
- [0586] In some embodiments, **R₃** is C₃-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
- [0587] In some embodiments, **R₃** is C₃-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl).
- [0588] In some embodiments, **R₃** is C₃-C₆ alkenyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.
- [0589] In some embodiments, **R₃** is C₂-C₆ alkynyl.
- [0590] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
- [0591] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, or -N(C₁-C₆ alkyl)₂.
- [0592] In some embodiments, **R₃** is C₃-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered

heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0593] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0594] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl.

[0595] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₂-C₆ alkenyl.

[0596] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₂-C₆ alkynyl.

[0597] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0598] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0599] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₅-C₆ aryl.

[0600] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0601] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0602] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0603] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0604] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0605] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkoxy.

[0606] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0607] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0608] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more halo or -CN.

[0609] In some embodiments, **R₃** is C₃-C₆ alkynyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0610] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more halo.

[0611] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more F, Cl, Br, or I.

[0612] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more F or Cl.

[0613] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more F.

[0614] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more Cl.

[0615] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more -CN.

[0616] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more -OH.

[0617] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more -NH₂.

[0618] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more oxo.

[0619] In some embodiments, **R₃** is C₃-C₆ alkynyl optionally substituted with one or more -OH.

[0620] In some embodiments, **R₃** is C₃-C₆ alkynyl optionally substituted with one or more -NH₂.

[0621] In some embodiments, **R₃** is C₃-C₆ alkynyl optionally substituted with one or more oxo.

[0622] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0623] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0624] In some embodiments, **R₃** is C₂-C₆ alkynyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0625] In some embodiments, **R₃** is C₃-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0626] In some embodiments, **R₃** is C₃-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0627] In some embodiments, **R₃** is C₃-C₆ alkynyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0628] In some embodiments, **R₃** is ethyl or isopropyl.

[0629] In some embodiments, **R_{3S}** is independently selected from -OC(=O)**R_{3Sa}** or 5- to 6-membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0630] In some embodiments, **R_{3S}** is -OC(=O)**R_{3Sa}** or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

[0631] In some embodiments, **R_{3S}** is -OC(=O)**R_{3Sa}**.

[0632] In some embodiments, **R_{3S}** is 5- to 6- membered heterocycloalkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0633] In some embodiments, **R_{3S}** is 5- to 6- membered heterocycloalkyl substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0634] In some embodiments, **R_{3S}** is 5- to 6- membered heterocycloalkyl substituted with one or more -OH, halo, -CN, or oxo.

[0635] In some embodiments, **R_{3S}** is 5- to 6- membered heterocycloalkyl substituted with one or more -OH.

[0636] In some embodiments, **R_{3S}** is 5- to 6- membered heterocycloalkyl substituted with one, two or three -OH.

[0637] In some embodiments, **R_{3S}** is 5- to 6- membered heterocycloalkyl substituted with three -OH.

[0638] In some embodiments, **R_{3S}** is 5- to 6- membered heterocycloalkyl substituted with one or more halo.

[0639] In some embodiments, **R_{3S}** is 5- to 6- membered heterocycloalkyl substituted with one or more -CN.

[0640] In some embodiments, **R_{3S}** is 5- to 6- membered heterocycloalkyl substituted with one or more oxo.

[0641] In some embodiments, R_{3S} is 5- to 6- membered heterocycloalkyl substituted with one or more $-NH_2$, $-NH(C_1-C_6 \text{ alkyl})$, or $-N(C_1-C_6 \text{ alkyl})_2$.

[0642] In some embodiments, R_{3S} is 5- to 6- membered heterocycloalkyl substituted with one or more $-NH_2$.

[0643] In some embodiments, R_{3S} is 5- to 6- membered heterocycloalkyl substituted with one or more $-NH(C_1-C_6 \text{ alkyl})$.

[0644] In some embodiments, R_{3S} is 5- to 6- membered heterocycloalkyl substituted with one or more $-N(C_1-C_6 \text{ alkyl})_2$.

[0645] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-OH$, halo, $-CN$, $-NH_2$, $-NH(C_1-C_6 \text{ alkyl})$, $-N(C_1-C_6 \text{ alkyl})_2$, oxo, or $-C(=O)OR_{3Sb}$.

[0646] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-OH$, halo, $-CN$, $-NH_2$, $-NH(C_1-C_6 \text{ alkyl})$, $-N(C_1-C_6 \text{ alkyl})_2$, or oxo.

[0647] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-OH$, halo, or $-CN$.

[0648] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-OH$.

[0649] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more halo.

In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more F, Cl, Br, or I.

In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more F or Cl. In

some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more F. In some

embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more Cl.

[0650] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-CN$.

[0651] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-NH_2$, $-NH(C_1-C_6 \text{ alkyl})$, $-N(C_1-C_6 \text{ alkyl})_2$, or oxo.

[0652] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more oxo.

[0653] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-NH_2$, $-NH(C_1-C_6 \text{ alkyl})$, or $-N(C_1-C_6 \text{ alkyl})_2$.

[0654] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-NH_2$.

[0655] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-NH(C_1-C_6 \text{ alkyl})$.

[0656] In some embodiments, R_{3Sa} is C_1-C_6 alkyl optionally substituted with one or more $-N(C_1-C_6 \text{ alkyl})_2$.

[0657] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}.

[0658] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl or –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more –OH, halo, –CN, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0659] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl optionally substituted with one or more –OH, halo, –CN, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0660] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl optionally substituted with one or more –OH.

[0661] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl optionally substituted with one or more halo.

[0662] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl optionally substituted with one or more –CN.

[0663] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl optionally substituted with one or more –NH₂.

[0664] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl optionally substituted –NH(C₁-C₆ alkyl).

[0665] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl optionally substituted with one or more –N(C₁-C₆ alkyl)₂.

[0666] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl optionally substituted with one or more oxo.

[0667] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is 4- to 8-membered heterocycloalkyl.

[0668] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more –OH, halo, –CN, –NH₂, –NH(C₁-C₆ alkyl), –N(C₁-C₆ alkyl)₂, or oxo.

[0669] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more –OH.

[0670] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more halo.

[0671] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more –CN.

[0672] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more –NH₂.

[0673] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more –NH(C₁-C₆ alkyl).

[0674] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more –N(C₁-C₆ alkyl)₂.

[0675] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more oxo.

[0676] In some embodiments, R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more –C(=O)OR_{3Sb}; wherein, R_{3Sb} is –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl).

[0677] In some embodiments, R_{3Sb} is 4- to 8-membered heterocycloalkyl or –(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or –(C₁-C₆

alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0678] In some embodiments, **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0679] In some embodiments, **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -OH.

[0680] In some embodiments, **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more halo.

[0681] In some embodiments, **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -CN.

[0682] In some embodiments, **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -NH₂.

[0683] In some embodiments, **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0684] In some embodiments, **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0685] In some embodiments, **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more oxo.

[0686] In some embodiments, **R_{3Sb}** is -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

[0687] In some embodiments, **R_{3Sb}** is -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more 4- to 8- membered heterocycloalkyl optionally substituted with one or more -OH.

[0688] In some embodiments, **R_{3Sb}** is -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more -OH.

[0689] In some embodiments, **R_{3Sb}** is -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more halo.





[0690] In some embodiments, **R_{3Sb}** is -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more -CN.


[0691] In some embodiments, R_{3Sb} is $-(C_1-C_6 \text{ alkyl})-(4- \text{ to } 8\text{-membered heterocycloalkyl})$ optionally substituted with one or more $-NH_2$.

[0692] In some embodiments, R_{3Sb} is $-(C_1-C_6 \text{ alkyl})-(4- \text{ to } 8\text{-membered heterocycloalkyl})$ optionally substituted with one or more $-NH(C_1-C_6 \text{ alkyl})$.


[0693] In some embodiments, R_{3Sb} is $-(C_1-C_6 \text{ alkyl})-(4- \text{ to } 8\text{-membered heterocycloalkyl})$ optionally substituted with one or more $-N(C_1-C_6 \text{ alkyl})_2$.

[0694] In some embodiments, R_{3Sb} is $-(C_1-C_6 \text{ alkyl})-(4- \text{ to } 8\text{-membered heterocycloalkyl})$ optionally substituted with one or more oxo.





[0701] In some embodiments, **R₃** is



[0702] In some embodiments, **R₃** is

[0703] In some embodiments, **R₄** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0704] In some embodiments, **R₄** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0705] In some embodiments, **R₄** is C₁-C₆ alkyl. In some embodiments, **R₄** is methyl. In some embodiments, **R₄** is ethyl. In some embodiments, **R₄** is propyl. In some embodiments, **R₄** is butyl. In some embodiments, **R₄** is pentyl. In some embodiments, **R₄** is hexyl. In some embodiments, at least one **R₄** is isopropyl. In some embodiments, at least one **R₄** is isobutyl. In some embodiments, at least one **R₄** is isopentyl. In some embodiments, at least one **R₄** is isohexyl. In some embodiments, at least one **R₄** is secbutyl. In some embodiments, at least one **R₄** is secpentyl. In some embodiments, at least one **R₄** is sechexyl. In some embodiments, at least one **R₄** is tertbutyl.

[0706] In some embodiments, **R₄** is  or .

[0707] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0708] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0709] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl.

[0710] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkenyl.

[0711] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkynyl.

[0712] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0713] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0714] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₅-C₆ aryl.

[0715] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0716] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0717] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0718] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0719] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0720] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkoxy.

[0721] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0722] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0723] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more halo. In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more F or Cl. In some

embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more F. In some

embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more Cl.

[0724] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more -CN.

[0725] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more -OH.

[0726] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more -NH₂.

[0727] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more oxo.

[0728] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0729] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0730] In some embodiments, **R₄** is C₁-C₆ alkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0731] In some embodiments, **R₄** is methyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0732] In some embodiments, **R₄** is methyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0733] In some embodiments, **R₄** is methyl optionally substituted with one or more C₁-C₆ alkyl.

[0734] In some embodiments, **R₄** is methyl optionally substituted with one or more C₂-C₆ alkenyl.

[0735] In some embodiments, **R₄** is methyl optionally substituted with one or more C₂-C₆ alkynyl.

[0736] In some embodiments, **R₄** is methyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0737] In some embodiments, **R₄** is methyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0738] In some embodiments, **R₄** is methyl optionally substituted with one or more C₅-C₆ aryl.

[0739] In some embodiments, **R₄** is methyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

- [0740] In some embodiments, **R₄** is methyl optionally substituted with one or more 5- or 6-membered heteroaryl.
- [0741] In some embodiments, **R₄** is methyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.
- [0742] In some embodiments, **R₄** is methyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.
- [0743] In some embodiments, **R₄** is methyl optionally substituted with one or more C₁-C₆ haloalkyl.
- [0744] In some embodiments, **R₄** is methyl optionally substituted with one or more C₁-C₆ alkoxy.
- [0745] In some embodiments, **R₄** is methyl optionally substituted with one or more C₁-C₆ haloalkoxy.
- [0746] In some embodiments, **R₄** is methyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.
- [0747] In some embodiments, **R₄** is methyl optionally substituted with one or more halo. In some embodiments, **R₄** is methyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R₄** is methyl optionally substituted with one or more F or Cl. In some embodiments, **R₄** is methyl optionally substituted with one or more F. In some embodiments, **R₄** is methyl optionally substituted with one or more Cl.
- [0748] In some embodiments, **R₄** is methyl optionally substituted with one or more -CN.
- [0749] In some embodiments, **R₄** is methyl optionally substituted with one or more -OH.
- [0750] In some embodiments, **R₄** is methyl optionally substituted with one or more -NH₂.
- [0751] In some embodiments, **R₄** is methyl optionally substituted with one or more oxo.
- [0752] In some embodiments, **R₄** is methyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
- [0753] In some embodiments, **R₄** is methyl optionally substituted with one or more -NH(C₁-C₆ alkyl).
- [0754] In some embodiments, **R₄** is methyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.
- [0755] In some embodiments, **R₄** is C₂-C₆ alkenyl.

[0756] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0757] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0758] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl.

[0759] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkenyl.

[0760] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkynyl.

[0761] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0762] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0763] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₅-C₆ aryl.

[0764] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0765] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0766] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0767] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0768] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0769] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkoxy.

[0770] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0771] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0772] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more halo. In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more F or Cl. In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more F. In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more Cl.

[0773] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more -CN.

[0774] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more -OH.

[0775] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more -NH₂.

[0776] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more oxo.

[0777] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0778] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0779] In some embodiments, **R₄** is C₂-C₆ alkenyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0780] In some embodiments, **R₄** is C₂-C₆ alkynyl.

[0781] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0782] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0783] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl.

[0784] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₂-C₆ alkenyl.

[0785] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₂-C₆ alkynyl.

[0786] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0787] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0788] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₅-C₆ aryl.

[0789] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0790] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0791] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0792] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0793] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0794] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkoxy.

[0795] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0796] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0797] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more halo.

[0798] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more F, Cl, Br, or I.

[0799] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more F or Cl.

[0800] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more F.

[0801] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more Cl.

- [0802] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more -CN.
- [0803] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more -OH.
- [0804] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more -NH₂.
- [0805] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more oxo.
- [0806] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
- [0807] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl).
- [0808] In some embodiments, **R₄** is C₂-C₆ alkynyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.
- [0809] In some embodiments, **R₅** is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
- [0810] In some embodiments, **R₅** is H.
- [0811] In some embodiments, **R₅** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
- [0812] In some embodiments, **R₅** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.
- [0813] In some embodiments, **R₅** is C₁-C₆ alkyl. In some embodiments, **R₅** is methyl. In some embodiments, **R₅** is ethyl. In some embodiments, **R₅** is propyl. In some embodiments, **R₅** is butyl. In some embodiments, **R₅** is pentyl. In some embodiments, **R₅** is hexyl. In some embodiments, at least one **R₅** is isopropyl. In some embodiments, at least one **R₅** is isobutyl. In some embodiments, at least one **R₅** is isopentyl. In some embodiments, at least one **R₅** is isohexyl. In some embodiments, at least one **R₅** is secbutyl. In some embodiments, at least one **R₅** is secpentyl. In some embodiments, at least one **R₅** is sechexyl. In some embodiments, at least one **R₅** is tertbutyl.

[0814] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0815] In some embodiments, **R₅** is C₂-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0816] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0817] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl.

[0818] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkenyl.

[0819] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkynyl.

[0820] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0821] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0822] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₅-C₆ aryl.

[0823] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0824] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0825] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0826] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0827] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0828] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkoxy.

[0829] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0830] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0831] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more halo. In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more F or Cl. In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more F. In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more Cl.

[0832] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more -CN.

[0833] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more -OH.

[0834] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more -NH₂.

[0835] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more oxo.

[0836] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0837] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0838] In some embodiments, **R₅** is C₁-C₆ alkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0839] In some embodiments, **R₅** is C₂-C₆ alkenyl. In some embodiments, **R₅** is C₃-C₆ alkenyl.

[0840] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0841] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl,

C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0842] In some embodiments, **R₅** is C₃-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0843] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0844] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl.

[0845] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkenyl.

[0846] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkynyl.

[0847] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0848] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0849] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₅-C₆ aryl.

[0850] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0851] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0852] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0853] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0854] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkyl.

- [0855] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkoxy.
- [0856] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkoxy.
- [0857] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.
- [0858] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more halo or -CN.
- [0859] In some embodiments, **R₅** is C₃-C₆ alkenyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.
- [0860] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more halo.
- [0861] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more F, Cl, Br, or I.
- [0862] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more F or Cl.
- [0863] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more F.
- [0864] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more Cl.
- [0865] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more -CN.
- [0866] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more -OH.
- [0867] In some embodiments, **R₅** is C₃-C₆ alkenyl optionally substituted with one or more -OH.
- [0868] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more -NH₂.
- [0869] In some embodiments, **R₅** is C₃-C₆ alkenyl optionally substituted with one or more -NH₂.
- [0870] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more oxo.
- [0871] In some embodiments, **R₅** is C₃-C₆ alkenyl optionally substituted with one or more oxo.
- [0872] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
- [0873] In some embodiments, **R₅** is C₃-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
- [0874] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0875] In some embodiments, **R₅** is C₃-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0876] In some embodiments, **R₅** is C₂-C₆ alkenyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0877] In some embodiments, **R₅** is C₃-C₆ alkenyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0878] In some embodiments, **R₅** is C₂-C₆ alkynyl.

[0879] In some embodiments, **R₅** is C₃-C₆ alkynyl.

[0880] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0881] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0882] In some embodiments, **R₅** is C₃-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0883] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0884] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl.

[0885] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more C₂-C₆ alkenyl.

[0886] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more C₂-C₆ alkynyl.

[0887] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

- [0888] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ cycloalkyl.
- [0889] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more C₅-C₆ aryl.
- [0890] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.
- [0891] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more 5- or 6-membered heteroaryl.
- [0892] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.
- [0893] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.
- [0894] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkyl.
- [0895] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkoxy.
- [0896] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkoxy.
- [0897] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.
- [0898] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more halo.
- [0899] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more F, Cl, Br, or I.
- [0900] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more F or Cl.
- [0901] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more F.
- [0902] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more Cl.
- [0903] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more -CN.
- [0904] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more -OH.
- [0905] In some embodiments, **R**₅ is C₂-C₆ alkynyl optionally substituted with one or more -NH₂.

[0906] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more oxo.

[0907] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0908] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0909] In some embodiments, **R₅** is C₂-C₆ alkynyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0910] In some embodiments, **R₆** is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0911] In some embodiments, **R₆** is H.

[0912] In some embodiments, **R₆** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0913] In some embodiments, **R₆** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0914] In some embodiments, **R₆** is C₁-C₆ alkyl. In some embodiments, **R₆** is methyl. In some embodiments, **R₆** is ethyl. In some embodiments, **R₆** is propyl. In some embodiments, **R₆** is butyl. In some embodiments, **R₆** is pentyl. In some embodiments, **R₆** is hexyl. In some embodiments, at least one **R₆** is isopropyl. In some embodiments, at least one **R₆** is isobutyl. In some embodiments, at least one **R₆** is isopentyl. In some embodiments, at least one **R₆** is isohexyl. In some embodiments, at least one **R₆** is secbutyl. In some embodiments, at least one **R₆** is secpentyl. In some embodiments, at least one **R₆** is sechexyl. In some embodiments, at least one **R₆** is tertbutyl.

[0915] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0916] In some embodiments, **R₆** is C₂-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0917] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0918] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkyl.

[0919] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkenyl.

[0920] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₂-C₆ alkynyl.

[0921] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0922] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0923] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₅-C₆ aryl.

[0924] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0925] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0926] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0927] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0928] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0929] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ alkoxy.

[0930] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0931] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0932] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more halo. In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more F or Cl. In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more F. In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more Cl.

[0933] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more -CN.

[0934] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more -OH.

[0935] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more -NH₂.

[0936] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more oxo.

[0937] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0938] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0939] In some embodiments, **R₆** is C₁-C₆ alkyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0940] In some embodiments, **R₆** is C₂-C₆ alkenyl. In some embodiments, **R₆** is C₃-C₆ alkenyl.

[0941] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0942] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0943] In some embodiments, **R₆** is C₃-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl,

C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0944] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0945] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkyl.

[0946] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkenyl.

[0947] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₂-C₆ alkynyl.

[0948] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0949] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0950] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₅-C₆ aryl.

[0951] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0952] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0953] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0954] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0955] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0956] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ alkoxy.

[0957] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0958] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0959] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more halo or -CN.

[0960] In some embodiments, **R₆** is C₃-C₆ alkenyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0961] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more halo. In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more F or Cl. In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more F. In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more Cl.

[0962] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more -CN.

[0963] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more -OH.

[0964] In some embodiments, **R₆** is C₃-C₆ alkenyl optionally substituted with one or more -OH.

[0965] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more -NH₂.

[0966] In some embodiments, **R₆** is C₃-C₆ alkenyl optionally substituted with one or more -NH₂.

[0967] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more oxo.

[0968] In some embodiments, **R₆** is C₃-C₆ alkenyl optionally substituted with one or more oxo.

[0969] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0970] In some embodiments, **R₆** is C₃-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0971] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0972] In some embodiments, **R₆** is C₃-C₆ alkenyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[0973] In some embodiments, **R₆** is C₂-C₆ alkenyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0974] In some embodiments, **R₆** is C₃-C₆ alkenyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[0975] In some embodiments, **R₆** is C₂-C₆ alkynyl. In some embodiments, **R₆** is C₃-C₆ alkynyl.

[0976] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0977] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[0978] In some embodiments, **R₆** is C₃-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

[0979] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

[0980] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkyl.

[0981] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more C₂-C₆ alkenyl.

[0982] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more C₂-C₆ alkynyl.

[0983] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ cycloalkyl or C₅-C₆ aryl.

[0984] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ cycloalkyl.

[0985] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more C₅-C₆ aryl.

[0986] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more 5- or 6-membered heteroaryl or C₃-C₈ heterocycloalkyl.

[0987] In some embodiments, **R₆** is C₂-C₆ alkynyl optionally substituted with one or more 5- or 6-membered heteroaryl.

[0988] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more C₃-C₈ heterocycloalkyl.

[0989] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.

[0990] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkyl.

[0991] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ alkoxy.

[0992] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more C₁-C₆ haloalkoxy.

[0993] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more halo, -CN, -OH, -NH₂, or oxo.

[0994] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more halo. In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more F, Cl, Br, or I. In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more F or Cl. In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more F. In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more Cl.

[0995] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more -CN.

[0996] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more -OH.

[0997] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more -NH₂.



[0998] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more oxo.

[0999] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.

[01000] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more -NH(C₁-C₆ alkyl).

[01001] In some embodiments, **R**₆ is C₂-C₆ alkynyl optionally substituted with one or more -N(C₁-C₆ alkyl)₂.

[01002] In some embodiments, the compound is of Formula (I-a) or (I-b):





or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01003] In some embodiments, the compound is of Formula (I-a) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01004] In some embodiments, the compound is of Formula (I-b) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01005] In some embodiments, the compound is of Formula (I-a1), (I-a2), (I-a3), or (I-a4):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01006] In some embodiments, the compound is of Formula (I-a1) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01007] In some embodiments, the compound is of Formula (I-a2) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01008] In some embodiments, the compound is of Formula (I-a3) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01009] In some embodiments, the compound is of Formula (I-a4) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01010] In some embodiments, the compound is of Formula (I-a5):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01011] In some embodiments, the compound is of Formula (I-a6), (I-a7), (I-a8), or (I-a9):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01012] In some embodiments, the compound is of Formula (I-a6) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01013] In some embodiments, the compound is of Formula (I-a7) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01014] In some embodiments, the compound is of Formula (I-a8) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01015] In some embodiments, the compound is of Formula (I-a9) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01016] In some embodiments, the compound is of Formula (I-b1), (I-b2), (I-b3), or (I-b4):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01017] In some embodiments, the compound is of Formula (I-b1) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01018] In some embodiments, the compound is of Formula (I-b2) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01019] In some embodiments, the compound is of Formula (I-b3) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01020] In some embodiments, the compound is of Formula (I-b4) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01021] In some embodiments, the compound is of Formula (I-b5):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01022] In some embodiments, the compound is of Formula (I-b6), (I-b7), (I-b8), or (I-b9):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01023] In some embodiments, the compound is of Formula (I-b6) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01024] In some embodiments, the compound is of Formula (I-b7) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01025] In some embodiments, the compound is of Formula (I-b8) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01026] In some embodiments, the compound is of Formula (I-b9) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01027] In some embodiments, the compound is of Formula (I-c), (I-d), or (I-e):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01028] In some embodiments, the compound is of Formula (I-c) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01029] In some embodiments, the compound is of Formula (I-d) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01030] In some embodiments, the compound is of Formula (I-e) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01031] In some embodiments, the compound is of Formula (I-f), (I-g), or (I-h):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01032] In some embodiments, the compound is of Formula (I-f) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01033] In some embodiments, the compound is of Formula (I-g) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01034] In some embodiments, the compound is of Formula (I-h) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01035] In some embodiments, the compound is of Formula (I-i), (I-j), or (I-k):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01036] In some embodiments, the compound is of Formula (I-i) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01037] In some embodiments, the compound is of Formula (I-j) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01038] In some embodiments, the compound is of Formula (I-k) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01039] In some embodiments, the compound is of Formula (I-l) or (I-m):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01040] In some embodiments, the compound is of Formula (I-l) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01041] In some embodiments, the compound is of Formula (I-m) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01042] In some embodiments, the compound is of Formula (I-n), (I-o), or (I-p):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01043] In some embodiments, the compound is of Formula (I-n) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01044] In some embodiments, the compound is of Formula (I-o) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01045] In some embodiments, the compound is of Formula (I-p) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01046] In some embodiments, the compound is of Formula (Ia-a):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01047] In some embodiments, the compound is of Formula (Ia-a) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01048] In some embodiments, the compound is of Formula (Ia-b), (Ia-c) or (Ia-d):






or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01049] In some embodiments, the compound is of Formula (Ia-b) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01050] In some embodiments, the compound is of Formula (Ia-c) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01051] In some embodiments, the compound is of Formula (Ia-d) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01052] In some embodiments, the compound is of Formula (Ia-e), (Ia-f), or (Ia-g):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01053] In some embodiments, the compound is of Formula (Ia-e) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01054] In some embodiments, the compound is of Formula (Ia-f) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01055] In some embodiments, the compound is of Formula (Ia-g) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01056] In some embodiments, the compound is of Formula (Ia-h), or (Ia-i):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01057] In some embodiments, the compound is of Formula (Ia-h) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01058] In some embodiments, the compound is of Formula (Ia-i) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01059] In some embodiments, the compound is of Formula (Ia-j), (Ia-k), or (Ia-l):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01060] In some embodiments, the compound is of Formula (Ia-j) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01061] In some embodiments, the compound is of Formula (Ia-k) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01062] In some embodiments, the compound is of Formula (Ia-l) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01063] In some embodiments, the compound is of Formula (Ia-m), (Ia-n), (Ia-o), or (Ia-p):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01064] In some embodiments, the compound is of Formula (Ia-m) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01065] In some embodiments, the compound is of Formula (Ia-n) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01066] In some embodiments, the compound is of Formula (Ia-o) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01067] In some embodiments, the compound is of Formula (Ia-p) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01068] In some embodiments, the compound is of Formula (Ia-q), (Ia-r), (Ia-s), or (Ia-t):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01069] In some embodiments, the compound is of Formula (Ia-q) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01070] In some embodiments, the compound is of Formula (Ia-r) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01071] In some embodiments, the compound is of Formula (Ia-s) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01072] In some embodiments, the compound is of Formula (Ia-t) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01073] In some embodiments, the compound is of Formula (Ia-u), (Ia-v), (Ia-w), (Ia-x), (Ia-y), or (Ia-z):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01074] In some embodiments, the compound is of Formula (Ia-u) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01075] In some embodiments, the compound is of Formula (Ia-v) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01076] In some embodiments, the compound is of Formula (Ia-w) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01077] In some embodiments, the compound is of Formula (Ia-x) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01078] In some embodiments, the compound is of Formula (Ia-y) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01079] In some embodiments, the compound is of Formula (Ia-z) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.



[01080] In some embodiments, the compound is of Formula (Ib-a):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01081] In some embodiments, the compound is of Formula (Ib-a) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01082] In some embodiments, the compound is of Formula (Ib-b), (Ib-c) or (Ib-d):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01083] In some embodiments, the compound is of Formula (Ib-b) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01084] In some embodiments, the compound is of Formula (Ib-c) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01085] In some embodiments, the compound is of Formula (Ib-d) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01086] In some embodiments, the compound is of Formula (Ib-e), (Ib-f), or (Ib-g):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01087] In some embodiments, the compound is of Formula (Ib-e) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01088] In some embodiments, the compound is of Formula (Ib-f) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01089] In some embodiments, the compound is of Formula (Ib-g) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01090] In some embodiments, the compound is of Formula (Ib-h), or (Ib-i):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01091] In some embodiments, the compound is of Formula (Ib-h) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01092] In some embodiments, the compound is of Formula (Ib-i) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01093] In some embodiments, the compound is of Formula (Ib-j), (Ib-k), or (Ib-l):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01094] In some embodiments, the compound is of Formula (Ib-j) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01095] In some embodiments, the compound is of Formula (Ib-k) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01096] In some embodiments, the compound is of Formula (Ib-l) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01097] In some embodiments, the compound is of Formula (Ib-m), (Ib-n), (Ib-o), or (Ib-p):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01098] In some embodiments, the compound is of Formula (Ib-m) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01099] In some embodiments, the compound is of Formula (Ib-n) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01100] In some embodiments, the compound is of Formula (Ib-o) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01101] In some embodiments, the compound is of Formula (Ib-p) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01102] In some embodiments, the compound is of Formula (Ib-q), (Ib-r), (Ib-s), or (Ib-t):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01103] In some embodiments, the compound is of Formula (Ib-q) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01104] In some embodiments, the compound is of Formula (Ib-r) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01105] In some embodiments, the compound is of Formula (Ib-s) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01106] In some embodiments, the compound is of Formula (Ib-t) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01107] In some embodiments, the compound is of Formula (Ib-u), (Ib-v), (Ib-w) (Ib-x), (Ib-y), or (Ib-z):





or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01108] In some embodiments, the compound is of Formula (Ib-u) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01109] In some embodiments, the compound is of Formula (Ib-v) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01110] In some embodiments, the compound is of Formula (Ib-w) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.



[01111] In some embodiments, the compound is of Formula (Ib-x) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01112] In some embodiments, the compound is of Formula (Ib-y) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01113] In some embodiments, the compound is of Formula (Ib-z) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01114] In some embodiments, the compound is of Formula (Ic-a), (Ic-b), (Ic-c), (Ic-d), or (Ic-e):





or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01115] In some embodiments, the compound is of Formula (Ic-a) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


[01116] In some embodiments, the compound is of Formula (Ic-b) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.



[01117] In some embodiments, the compound is of Formula (Ic-c) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01118] In some embodiments, the compound is of Formula (Ic-d) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01119] In some embodiments, the compound is of Formula (Ic-e) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01120] In some embodiments, the compound is of Formula (Id-a), (Id-b), (Id-c), (Id-d), or (Id-e):





or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01121] In some embodiments, the compound is of Formula (Id-a) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01122] In some embodiments, the compound is of Formula (Id-b) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01123] In some embodiments, the compound is of Formula (Id-c) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01124] In some embodiments, the compound is of Formula (Id-d) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01125] In some embodiments, the compound is of Formula (Id-e) or a prodrug, solvate, or pharmaceutically acceptable salt thereof.

[01126] It is understood that, for a compound of any one of the formulae described herein, **X**, **R₁**, **R_{1S}**, **R_{1SS}**, **Q**, **R₂**, **R₃**, **R_{3Sa}**, **R_{3Sb}**, **R₄**, and **R₅** can each be, where applicable, selected from the groups described herein, and any group described herein for any of **X**, **R₁**, **R_{1S}**, **R_{1SS}**, **Q**, **R₂**, **R₃**, **R_{3Sa}**, **R_{3Sb}**, **R₄**, and **R₅** can be combined, where applicable, with any group described herein for one or more of the remainder of **X**, **R₁**, **R_{1S}**, **R_{1SS}**, **Q**, **R₂**, **R₃**, **R_{3Sa}**, **R_{3Sb}**, **R₄**, and **R₅**.










[01127] In some embodiments, the compound is selected from the compounds described in Table 1 and prodrugs and pharmaceutically acceptable salts thereof.










[01128] In some embodiments, the compound is selected from the compounds described in Table 1 and pharmaceutically acceptable salts thereof.










[01129] In some embodiments, the compound is selected from the prodrugs of compounds described in Table 1 and pharmaceutically acceptable salts thereof.






[01130] In some embodiments, the compound is selected from the compounds described in Table 1.

Table 1


Compound No.	Structure
1	
2	
3	
4	
5	
6	
7	
8	
9	


10	
11	
12	
13	
14	
15	
16	
17	
18	

19	
20	
21	
22	
23	
24	
25	
26	
27	


28	
29	
30	
31	
32	

[01131] In some embodiments, the compound is of Formula (Ic-a), (Ic-b), (Ic-c), (Ic-d), or (Ic-e):







(Ic-d);




(Ic-e);




(Id-a);




(Id-b);



(Id-c);



(Id-d); or






(Id-e),

or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₄**, and **-Q-R₂** are each independently selected from Table 2 below.

Table 2

Variable	Options
----------	---------

R₁	
R₄	
-Q-R₂	

[01132] In some aspects, the present disclosure provides a compound being an isotopic derivative (e.g., isotopically labeled compound) of any one of the compounds of the Formulae disclosed herein.

[01133] In some embodiments, the compound is an isotopic derivative of any one of the compounds described in Table 1 and prodrugs and pharmaceutically acceptable salts thereof.

[01134] In some embodiments, the compound is an isotopic derivative of any one of the compounds described in Table 1 and pharmaceutically acceptable salts thereof.

[01135] In some embodiments, the compound is an isotopic derivative of any one of prodrugs of the compounds described in Table 1 and pharmaceutically acceptable salts thereof.

[01136] In some embodiments, the compound is an isotopic derivative of any one of the compounds described in Table 1.

[01137] It is understood that the isotopic derivative can be prepared using any of a variety of art-recognised techniques. For example, the isotopic derivative can generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples described herein, by substituting an isotopically labeled reagent for a non-isotopically labeled reagent.

[01138] In some embodiments, the isotopic derivative is a deuterium labeled compound.

[01139] In some embodiments, the isotopic derivative is a deuterium labeled compound of any one of the compounds of the Formulae disclosed herein.

[01140] In some embodiments, the compound is a deuterium labeled compound of any one of the compounds described in Table 1 and prodrugs and pharmaceutically acceptable salts thereof.

[01141] In some embodiments, the compound is a deuterium labeled compound of any one of the compounds described in Table 1 and pharmaceutically acceptable salts thereof.

[01142] In some embodiments, the compound is a deuterium labeled compound of any one of the prodrugs of the compounds described in Table 1 and pharmaceutically acceptable salts thereof.

[01143] In some embodiments, the compound is a deuterium labeled compound of any one of the compounds described in Table 1.

[01144] It is understood that the deuterium labeled compound comprises a deuterium atom having an abundance of deuterium that is substantially greater than the natural abundance of deuterium, which is 0.015%.

[01145] In some embodiments, the deuterium labeled compound has a deuterium enrichment factor for each deuterium atom of at least 3500 (52.5% deuterium incorporation at each deuterium atom), at least 4000 (60% deuterium incorporation), at least 4500 (67.5% deuterium incorporation), at least 5000 (75% deuterium), at least 5500 (82.5% deuterium incorporation), at least 6000 (90% deuterium incorporation), at least 6333.3 (95% deuterium incorporation), at least 6466.7 (97% deuterium incorporation), at least 6600 (99% deuterium incorporation), or at least 6633.3 (99.5% deuterium incorporation). As used herein, the term “deuterium enrichment factor” means the ratio between the deuterium abundance and the natural abundance of a deuterium.

[01146] It is understood that the deuterium labeled compound can be prepared using any of a variety of art-recognised techniques. For example, the deuterium labeled compound can

generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples described herein, by substituting a deuterium labeled reagent for a non-deuterium labeled reagent.

[01147] A compound of the invention or a pharmaceutically acceptable salt or solvate thereof that contains the aforementioned deuterium atom(s) is within the scope of the invention. Further, substitution with deuterium (*i.e.*, ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability, e.g., increased *in vivo* half-life or reduced dosage requirements.

[01148] In some embodiments, the compound is a ^{18}F labeled compound.

[01149] In some embodiments, the compound is a ^{123}I labeled compound, a ^{124}I labeled compound, a ^{125}I labeled compound, a ^{129}I labeled compound, a ^{131}I labeled compound, a ^{135}I labeled compound, or any combination thereof.

[01150] In some embodiments, the compound is a ^{33}S labeled compound, a ^{34}S labeled compound, a ^{35}S labeled compound, a ^{36}S labeled compound, or any combination thereof.

[01151] It is understood that the ^{18}F , ^{123}I , ^{124}I , ^{125}I , ^{129}I , ^{131}I , ^{135}I , ^3S , ^{34}S , ^{35}S , and/or ^{36}S labeled compound, can be prepared using any of a variety of art-recognised techniques. For example, the deuterium labeled compound can generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples described herein, by substituting a ^{18}F , ^{123}I , ^{124}I , ^{125}I , ^{129}I , ^{131}I , ^{135}I , ^3S , ^{34}S , ^{35}S , and/or ^{36}S labeled reagent for a non-isotope labeled reagent.

[01152] A compound of the invention or a pharmaceutically acceptable salt or solvate thereof that contains one or more of the aforementioned ^{18}F , ^{123}I , ^{124}I , ^{125}I , ^{129}I , ^{131}I , ^{135}I , ^3S , ^{34}S , ^{35}S , and ^{36}S atom(s) is within the scope of the invention. Further, substitution with isotope (*e.g.*, ^{18}F , ^{123}I , ^{124}I , ^{125}I , ^{129}I , ^{131}I , ^{135}I , ^3S , ^{34}S , ^{35}S , and/or ^{36}S) may afford certain therapeutic advantages resulting from greater metabolic stability, e.g., increased *in vivo* half-life or reduced dosage requirements.

[01153] For the avoidance of doubt it is to be understood that, where in this specification a group is qualified by “described herein”, the said group encompasses the first occurring and broadest definition as well as each and all of the particular definitions for that group.

[01154] The various functional groups and substituents making up the compounds of the Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') are typically chosen such that the molecular weight of the compound does not exceed 1000 daltons. More usually, the molecular weight of the compound will be less than 900, for example less than 800, or less than

750, or less than 700, or less than 650 daltons. More conveniently, the molecular weight is less than 600 and, for example, is 550 daltons or less.

[01155] A suitable pharmaceutically acceptable salt of a compound of the disclosure is, for example, an acid-addition salt of a compound of the disclosure which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, formic, citric methane sulphonate or maleic acid. In addition, a suitable pharmaceutically acceptable salt of a compound of the disclosure which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a pharmaceutically acceptable cation, for example a salt with methylamine, dimethylamine, diethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

[01156] It will be understood that the compounds of any one of the Formulae disclosed herein and any pharmaceutically acceptable salts thereof, comprise stereoisomers, mixtures of stereoisomers, polymorphs of all isomeric forms of said compounds.

[01157] As used herein, the term “isomerism” means compounds that have identical molecular formulae but differ in the sequence of bonding of their atoms or in the arrangement of their atoms in space. Isomers that differ in the arrangement of their atoms in space are termed “stereoisomers.” Stereoisomers that are not mirror images of one another are termed “diastereoisomers,” and stereoisomers that are non-superimposable mirror images of each other are termed “enantiomers” or sometimes optical isomers. A mixture containing equal amounts of individual enantiomeric forms of opposite chirality is termed a “racemic mixture.”

[01158] As used herein, the term “chiral centre” refers to a carbon atom bonded to four nonidentical substituents.

[01159] As used herein, the term “chiral isomer” means a compound with at least one chiral centre. Compounds with more than one chiral centre may exist either as an individual diastereomer or as a mixture of diastereomers, termed “diastereomeric mixture.” When one chiral centre is present, a stereoisomer may be characterised by the absolute configuration (R or S) of that chiral centre. Absolute configuration refers to the arrangement in space of the substituents attached to the chiral centre. The substituents attached to the chiral centre under consideration are ranked in accordance with the *Sequence Rule* of Cahn, Ingold and Prelog.

(Cahn *et al.*, *Angew. Chem. Inter. Edit.* 1966, 5, 385; errata 511; Cahn *et al.*, *Angew. Chem.* 1966, 78, 413; Cahn and Ingold, *J. Chem. Soc.* 1951 (London), 612; Cahn *et al.*, *Experientia* 1956, 12, 81; Cahn, *J. Chem. Educ.* 1964, 41, 116).

[01160] As used herein, the term “geometric isomer” means the diastereomers that owe their existence to hindered rotation about double bonds or a cycloalkyl linker (e.g., 1,3-cyclobutyl). These configurations are differentiated in their names by the prefixes *cis* and *trans*, or *Z* and *E*, which indicate that the groups are on the same or opposite side of the double bond in the molecule according to the Cahn-Ingold-Prelog rules.

[01161] It is to be understood that the compounds of the present disclosure may be depicted as different chiral isomers or geometric isomers. It is also to be understood that when compounds have chiral isomeric or geometric isomeric forms, all isomeric forms are intended to be included in the scope of the present disclosure, and the naming of the compounds does not exclude any isomeric forms, it being understood that not all isomers may have the same level of activity.

[01162] It is to be understood that the structures and other compounds discussed in this disclosure include all atropic isomers thereof. It is also to be understood that not all atropic isomers may have the same level of activity.

[01163] As used herein, the term “atropic isomers” are a type of stereoisomer in which the atoms of two isomers are arranged differently in space. Atropic isomers owe their existence to a restricted rotation caused by hindrance of rotation of large groups about a central bond. Such atropic isomers typically exist as a mixture, however as a result of recent advances in chromatography techniques, it has been possible to separate mixtures of two atropic isomers in select cases.

[01164] As used herein, the term “tautomer” is one of two or more structural isomers that exist in equilibrium and is readily converted from one isomeric form to another. This conversion results in the formal migration of a hydrogen atom accompanied by a switch of adjacent conjugated double bonds. Tautomers exist as a mixture of a tautomeric set in solution. In solutions where tautomerisation is possible, a chemical equilibrium of the tautomers will be reached. The exact ratio of the tautomers depends on several factors, including temperature, solvent and pH. The concept of tautomers that are interconvertible by tautomerisations is called tautomerism. Of the various types of tautomerism that are possible, two are commonly observed. In keto-enol tautomerism a simultaneous shift of electrons and a hydrogen atom occurs. Ring-chain

tautomerism arises as a result of the aldehyde group (-CHO) in a sugar chain molecule reacting with one of the hydroxy groups (-OH) in the same molecule to give it a cyclic (ring-shaped) form as exhibited by glucose.

[01165] It is to be understood that the compounds of the present disclosure may be depicted as different tautomers. It should also be understood that when compounds have tautomeric forms, all tautomeric forms are intended to be included in the scope of the present disclosure, and the naming of the compounds does not exclude any tautomer form. It will be understood that certain tautomers may have a higher level of activity than others.

[01166] Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers". Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers" and those that are non-superimposable mirror images of each other are termed "enantiomers". When a compound has an asymmetric centre, for example, it is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterised by the absolute configuration of its asymmetric centre and is described by the R- and S-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarised light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture".

[01167] The compounds of this disclosure may possess one or more asymmetric centres; such compounds can therefore be produced as individual (R)- or (S)-stereoisomers or as mixtures thereof. Unless indicated otherwise, the description or naming of a particular compound in the specification and claims is intended to include both individual enantiomers and mixtures, racemic or otherwise, thereof. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art (see discussion in Chapter 4 of "Advanced Organic Chemistry", 4th edition J. March, John Wiley and Sons, New York, 2001), for example by synthesis from optically active starting materials or by resolution of a racemic form. Some of the compounds of the disclosure may have geometric isomeric centres (E- and Z- isomers). It is to be understood that the present disclosure encompasses all optical, diastereoisomers and geometric isomers and mixtures thereof that possess inflammasome inhibitory activity.

[01168] The present disclosure also encompasses compounds of the disclosure as defined herein which comprise one or more isotopic substitutions.

[01169] It is to be understood that the compounds of any Formula described herein include the compounds themselves, as well as their salts, and their solvates, if applicable. A salt, for example, can be formed between an anion and a positively charged group (e.g., amino) on a substituted compound disclosed herein. Suitable anions include chloride, bromide, iodide, sulphate, bisulphate, sulphamate, nitrate, phosphate, citrate, methanesulphonate, trifluoroacetate, glutamate, glucuronate, glutarate, malate, maleate, succinate, fumarate, tartrate, tosylate, salicylate, lactate, naphthalenesulphonate, and acetate (e.g., trifluoroacetate).

[01170] As used herein, the term “pharmaceutically acceptable anion” refers to an anion suitable for forming a pharmaceutically acceptable salt. Likewise, a salt can also be formed between a cation and a negatively charged group (e.g., carboxylate) on a substituted compound disclosed herein. Suitable cations include sodium ion, potassium ion, magnesium ion, calcium ion, and an ammonium cation such as tetramethylammonium ion or diethylamine ion. The substituted compounds disclosed herein also include those salts containing quaternary nitrogen atoms.

[01171] It is to be understood that the compounds of the present disclosure, for example, the salts of the compounds, can exist in either hydrated or unhydrated (the anhydrous) form or as solvates with other solvent molecules. Nonlimiting examples of hydrates include monohydrates, dihydrates, etc. Nonlimiting examples of solvates include ethanol solvates, acetone solvates, etc.

[01172] As used herein, the term “solvate” means solvent addition forms that contain either stoichiometric or non-stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent is water the solvate formed is a hydrate; and if the solvent is alcohol, the solvate formed is an alcoholate. Hydrates are formed by the combination of one or more molecules of water with one molecule of the substance in which the water retains its molecular state as H₂O.

[01173] As used herein, the term “analog” refers to a chemical compound that is structurally similar to another but differs slightly in composition (as in the replacement of one atom by an atom of a different element or in the presence of a particular functional group, or the replacement of one functional group by another functional group). Thus, an analog is a compound that is similar or comparable in function and appearance, but not in structure or origin to the reference compound.


[01174] As used herein, the term “derivative” refers to compounds that have a common core structure and are substituted with various groups as described herein.

[01175] As used herein, the term “bioisostere” refers to a compound resulting from the exchange of an atom or of a group of atoms with another, broadly similar, atom or group of atoms. The objective of a bioisosteric replacement is to create a new compound with similar biological properties to the parent compound. The bioisosteric replacement may be physicochemically or topologically based. Examples of carboxylic acid bioisosteres include, but are not limited to, acyl sulphonamides, tetrazoles, sulphonates and phosphonates. See, *e.g.*, Patani and LaVoie, *Chem. Rev.* 96, 3147-3176, 1996.

[01176] It is also to be understood that certain compounds of any one of the Formulae disclosed herein may exist in solvated as well as unsolvated forms such as, for example, hydrated forms. A suitable pharmaceutically acceptable solvate is, for example, a hydrate such as hemi-hydrate, a mono-hydrate, a di-hydrate or a tri-hydrate. It is to be understood that the disclosure encompasses all such solvated forms that possess inflammasome inhibitory activity.

[01177] It is also to be understood that certain compounds of any one of the Formulae disclosed herein may exhibit polymorphism, and that the disclosure encompasses all such forms, or mixtures thereof, which possess inflammasome inhibitory activity. It is generally known that crystalline materials may be analysed using conventional techniques such as X-Ray Powder Diffraction analysis, Differential Scanning Calorimetry, Thermal Gravimetric Analysis, Diffuse Reflectance Infrared Fourier Transform (DRIFT) spectroscopy, Near Infrared (NIR) spectroscopy, solution and/or solid state nuclear magnetic resonance spectroscopy. The water content of such crystalline materials may be determined by Karl Fischer analysis.

[01178] Compounds of any one of the Formulae disclosed herein may exist in a number of different tautomeric forms and references to compounds of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') include all such forms. For the avoidance of doubt, where a compound can exist in one of several tautomeric forms, and only one is specifically described or shown, all others are nevertheless embraced by Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI'). Examples of tautomeric forms include keto-, enol-, and enolate-forms, as in, for example, the following tautomeric pairs: keto/enol (illustrated below), imine/enamine, amide/imino alcohol, amidine/amidine, nitroso/oxime, thioketone/enethiol, and nitro/aci-nitro.



[01179] Compounds of any one of the Formulae disclosed herein containing an amine function may also form N-oxides. A reference herein to a compound of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') that contains an amine function also includes the N-oxide. Where a compound contains several amine functions, one or more than one nitrogen atom may be oxidised to form an N-oxide. Particular examples of N-oxides are the N-oxides of a tertiary amine or a nitrogen atom of a nitrogen-containing heterocycle. N-oxides can be formed by treatment of the corresponding amine with an oxidising agent such as hydrogen peroxide or a peracid (e.g. a peroxy-carboxylic acid), see for example Advanced Organic Chemistry, by Jerry March, 4th Edition, Wiley Interscience, pages. More particularly, N-oxides can be made by the procedure of L. W. Deady (Syn. Comm. 1977, 7, 509-514) in which the amine compound is reacted with meta-chloroperoxybenzoic acid (mCPBA), for example, in an inert solvent such as dichloromethane.

[01180] The compounds of any one of the Formulae disclosed herein may be administered in the form of a prodrug which is broken down in the human or animal body to release a compound of the disclosure. A prodrug may be used to alter the physical properties and/or the pharmacokinetic properties of a compound of the disclosure. A prodrug can be formed when the compound of the disclosure contains a suitable group or substituent to which a property-modifying group can be attached. Examples of prodrugs include derivatives containing *in vivo* cleavable alkyl or acyl substituents at the ester or amide group in any one of the Formulae disclosed herein.

[01181] Accordingly, the present disclosure includes those compounds of any one of the Formulae disclosed herein as defined hereinbefore when made available by organic synthesis and when made available within the human or animal body by way of cleavage of a prodrug thereof. Accordingly, the present disclosure includes those compounds of any one of the Formulae disclosed herein that are produced by organic synthetic means and also such compounds that are produced in the human or animal body by way of metabolism of a precursor compound, that is a compound of any one of the Formulae disclosed herein may be a synthetically-produced compound or a metabolically-produced compound.

[01182] A suitable pharmaceutically acceptable prodrug of a compound of any one of the Formulae disclosed herein is one that is based on reasonable medical judgment as being suitable for administration to the human or animal body without undesirable pharmacological activities and without undue toxicity. Various forms of prodrug have been described, for example in the following documents: a) Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, et al. (Academic Press, 1985); b) Design of Pro-drugs, edited by H. Bundgaard, (Elsevier, 1985); c) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Pro-drugs", by H. Bundgaard p. 113-191 (1991); d) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992); e) H. Bundgaard, et al., Journal of Pharmaceutical Sciences, 77, 285 (1988); f) N. Kakeya, et al., Chem. Pharm. Bull., 32, 692 (1984); g) T. Higuchi and V. Stella, "Pro-Drugs as Novel Delivery Systems", A.C.S. Symposium Series, Volume 14; and h) E. Roche (editor), "Bioreversible Carriers in Drug Design", Pergamon Press, 1987.

[01183] A suitable pharmaceutically acceptable prodrug of a compound of any one of the Formulae disclosed herein that possesses a hydroxy group is, for example, an *in vivo* cleavable ester or ether thereof. An *in vivo* cleavable ester or ether of a compound of any one of the Formulae disclosed herein containing a hydroxy group is, for example, a pharmaceutically acceptable ester or ether which is cleaved in the human or animal body to produce the parent hydroxy compound. Suitable pharmaceutically acceptable ester forming groups for a hydroxy group include inorganic esters such as phosphate esters (including phosphoramidic cyclic esters). Further suitable pharmaceutically acceptable ester forming groups for a hydroxy group include C₁-C₁₀ alkanoyl groups such as acetyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl groups, C₁-C₁₀ alkoxycarbonyl groups such as ethoxycarbonyl, N,N-(C₁-C₆ alkyl)₂carbamoyl, 2-dialkylaminoacetyl and 2-carboxyacetyl groups. Examples of ring substituents on the phenylacetyl and benzoyl groups include aminomethyl, N-alkylaminomethyl, N,N-dialkylaminomethyl, morpholinomethyl, piperazin-1-ylmethyl and 4-(C₁-C₄ alkyl)piperazin-1-ylmethyl. Suitable pharmaceutically acceptable ether forming groups for a hydroxy group include α -acyloxyalkyl groups such as acetoxymethyl and pivaloyloxymethyl groups.

[01184] A suitable pharmaceutically acceptable prodrug of a compound of any one of the Formulae disclosed herein that possesses a carboxy group is, for example, an *in vivo* cleavable

amide thereof, for example an amide formed with an amine such as ammonia, a C₁₋₄alkylamine such as methylamine, a (C_{1-C4} alkyl)₂amine such as dimethylamine, N-ethyl-N-methylamine or diethylamine, a C_{1-C4} alkoxy-C_{2-C4} alkylamine such as 2-methoxyethylamine, a phenyl-C_{1-C4} alkylamine such as benzylamine and amino acids such as glycine or an ester thereof.

[01185] It is understood that a compound of any one of the Formulae disclosed herein, wherein **R₃** is not H, may be used as a prodrug of the corresponding compound, wherein **R₃** is H. For example, a compound of any one of the Formulae disclosed herein, wherein **R₃** is C_{1-C6} alkyl, C_{2-C6} alkenyl, or C_{2-C6} alkynyl; wherein the C_{1-C6} alkyl, C_{2-C6} alkenyl, or C_{2-C6} alkynyl is optionally substituted with one or more C_{1-C6} alkyl, C_{2-C6} alkenyl, C_{2-C6} alkynyl, C_{3-C8} cycloalkyl, C_{5-C6} aryl, 5- or 6-membered heteroaryl, C_{3-C8} heterocycloalkyl, C_{1-C6} haloalkyl, C_{1-C6} alkoxy, C_{1-C6} haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C_{1-C6} alkyl), -N(C_{1-C6} alkyl)₂, oxo, or **R_{3s}**, may be used as a prodrug of the corresponding compound, wherein **R₃** is H.

[01186] A suitable pharmaceutically acceptable prodrug of a compound of any one of the Formulae disclosed herein that possesses an amino group is, for example, an *in vivo* cleavable amide derivative thereof. Suitable pharmaceutically acceptable amides from an amino group include, for example an amide formed with C_{1-C10} alkanoyl groups such as an acetyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl groups. Examples of ring substituents on the phenylacetyl and benzoyl groups include aminomethyl, N-alkylaminomethyl, N,N-dialkylaminomethyl, morpholinomethyl, piperazin-1-ylmethyl and 4-(C_{1-C4} alkyl)piperazin-1-ylmethyl.

[01187] The *in vivo* effects of a compound of any one of the Formulae disclosed herein may be exerted in part by one or more metabolites that are formed within the human or animal body after administration of a compound of any one of the Formulae disclosed herein. As stated hereinbefore, the *in vivo* effects of a compound of any one of the Formulae disclosed herein may also be exerted by way of metabolism of a precursor compound (a prodrug).

[01188] Suitably, the present disclosure excludes any individual compounds not possessing the biological activity defined herein.

Methods of Synthesis

[01189] In some aspects, the present disclosure provides a method of preparing a compound of the present disclosure.

[01190] In some aspects, the present disclosure provides a method of a compound, comprising one or more steps as described herein.

[01191] In some aspects, the present disclosure provides a compound obtainable by, or obtained by, or directly obtained by a method for preparing a compound as described herein.

[01192] In some aspects, the present disclosure provides an intermediate as described herein, being suitable for use in a method for preparing a compound as described herein.

[01193] The compounds of the present disclosure can be prepared by any suitable technique known in the art. Particular processes for the preparation of these compounds are described further in the accompanying examples.

[01194] In the description of the synthetic methods described herein and in any referenced synthetic methods that are used to prepare the starting materials, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of the experiment and workup procedures, can be selected by a person skilled in the art.

[01195] It is understood by one skilled in the art of organic synthesis that the functionality present on various portions of the molecule must be compatible with the reagents and reaction conditions utilised.

[01196] It will be appreciated that during the synthesis of the compounds of the disclosure in the processes defined herein, or during the synthesis of certain starting materials, it may be desirable to protect certain substituent groups to prevent their undesired reaction. The skilled chemist will appreciate when such protection is required, and how such protecting groups may be put in place, and later removed. For examples of protecting groups see one of the many general texts on the subject, for example, 'Protective Groups in Organic Synthesis' by Theodora Green (publisher: John Wiley & Sons). Protecting groups may be removed by any convenient method described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with the minimum disturbance of groups elsewhere in the molecule. Thus, if reactants include, for example, groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

[01197] By way of example, a suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group,

for example a methoxycarbonyl, ethoxycarbonyl, or t-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxy carbonyl group or an aroyl group may be removed by, for example, hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a tert-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium on carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

[01198] A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium, sodium hydroxide or ammonia. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium on carbon.

[01199] A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a tert-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium on carbon.

[01200] Once a compound of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') has been synthesised by any one of the processes defined herein, the processes may then further comprise the additional steps of: (i) removing any protecting groups present; (ii) converting the compound Formula (I) into another compound of Formula (I), (I'), (II), (II'), (III),

(III'), (IV), (IV'), (V), (V'), (VI), or (VI'); (iii) forming a pharmaceutically acceptable salt, hydrate or solvate thereof; and/or (iv) forming a prodrug thereof.

[01201] The resultant compounds of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') can be isolated and purified using techniques well known in the art.

[01202] Conveniently, the reaction of the compounds is carried out in the presence of a suitable solvent, which is preferably inert under the respective reaction conditions. Examples of suitable solvents comprise but are not limited to hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichlorethylene, 1,2-dichloroethane, tetrachloromethane, chloroform or dichloromethane; alcohols, such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol; ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF), 2-methyltetrahydrofuran, cyclopentylmethyl ether (CPME), methyl tert-butyl ether (MTBE) or dioxane; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether or ethylene glycol dimethyl ether (diglyme); ketones, such as acetone, methylisobutylketone (MIBK) or butanone; amides, such as acetamide, dimethylacetamide, dimethylformamide (DMF) or N-methylpyrrolidinone (NMP); nitriles, such as acetonitrile; sulfoxides, such as dimethyl sulfoxide (DMSO); nitro compounds, such as nitromethane or nitrobenzene; esters, such as ethyl acetate or methyl acetate, or mixtures of the said solvents or mixtures with water.

[01203] The reaction temperature is suitably between about -100 °C and 300 °C, depending on the reaction step and the conditions used.

[01204] Reaction times are generally in the range between a fraction of a minute and several days, depending on the reactivity of the respective compounds and the respective reaction conditions. Suitable reaction times are readily determinable by methods known in the art, for example reaction monitoring. Based on the reaction temperatures given above, suitable reaction times generally lie in the range between 10 minutes and 48 hours.


[01205] Moreover, by utilising the procedures described herein, in conjunction with ordinary skills in the art, additional compounds of the present disclosure can be readily prepared. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds.

[01206] As will be understood by the person skilled in the art of organic synthesis, compounds of the present disclosure are readily accessible by various synthetic routes, some of which are

exemplified in the accompanying examples. The skilled person will easily recognise which kind of reagents and reactions conditions are to be used and how they are to be applied and adapted in any particular instance – wherever necessary or useful – in order to obtain the compounds of the present disclosure. Furthermore, some of the compounds of the present disclosure can readily be synthesised by reacting other compounds of the present disclosure under suitable conditions, for instance, by converting one particular functional group being present in a compound of the present disclosure, or a suitable precursor molecule thereof, into another one by applying standard synthetic methods, like reduction, oxidation, addition or substitution reactions; those methods are well known to the skilled person. Likewise, the skilled person will apply – whenever necessary or useful – synthetic protecting (or protective) groups; suitable protecting groups as well as methods for introducing and removing them are well-known to the person skilled in the art of chemical synthesis and are described, in more detail, in, e.g., P.G.M. Wuts, T.W. Greene, “Greene’s Protective Groups in Organic Synthesis”, 4th edition (2006) (John Wiley & Sons).

[01207] General routes for the preparation of a compound of the application are described in Scheme 1 and Scheme 2 herein.

Scheme 1




[01208] Reaction (i') may be performed by reacting primary amine **3** with $\text{R}_6\text{-X'}$ (wherein X' is a suitable leaving group) in a suitable solvent and, optionally, at an elevated temperature (e.g., 70° C or 80° C), optionally in the presence of a base, to yield the intermediate **10**.

[01209] Reaction (ii') may be performed by reacting amine **10** with a suitable reagent (e.g., diphosgene) in a suitable solvent and, optionally, at an elevated temperature (e.g., 70° C or 80° C) to yield the intermediate **11**.

[01210] Reaction (iii') may be performed by reacting intermediate **11** with intermediate **9** in an inert environment in a suitable solvent (e.g. tetrahydrofuran). In some embodiments, reaction (iii') may be performed at a cooled temperature (e.g. 0° C), optionally, in the presence of an

additive (e.g. CuCl), yielding a compound of **Formula (I)**. The compound of **Formula (I)** may be isolated by purification (e.g., by flash column chromatography or by preparative TLC or HPLC).

Scheme 2



[01211] Reaction (i) may be performed by reacting ketone **1** with *(S)*-2-methylpropane-2-sulphinamide and titanium isopropoxide in a suitable solvent (e.g., THF) and, optionally, heating to an elevated temperature (e.g., 70° C or 80° C) until the intermediate sulphinamide has formed. The reduction of the intermediate may be carried out by cooling and diluting the reaction mixture with a suitable solvent (e.g. MeOH) and adding a suitable reducing agent (e.g. LiBH₄), yielding Intermediate **2**.

[01212] Reaction (ii) may be performed by treating sulphinamide **2** or **4** with a suitable acidic solution (e.g. 4M HCl in methanol), yielding intermediate **3** or **5**.

[01213] Reaction (iii) may be performed by reacting ketone **1** with *(R)*-2-methylpropane-2-sulphinamide and titanium isopropoxide in a suitable solvent (e.g. THF) and, optionally, heating to an elevated temperature (e.g., 70° C or 80° C) until the intermediate sulphinamide has formed. The reduction of the intermediate may be carried out by cooling and diluting the reaction mixture with a suitable solvent (e.g. MeOH) and adding a suitable reducing agent (e.g. LiBH₄), yielding Intermediate **4**.

[01214] Reaction (iv) may be performed by reacting ketone **1** with hydroxylamine hydrochloride and sodium acetate in a suitable solvent mixture (e.g. ethanol and water) and, optionally, heating to an elevated temperature (e.g., 70° C or 80° C), yielding Intermediate **6**.

[01215] Reaction (v) may be performed by reacting intermediate **6** with Raney Nickel in a suitable solvent (e.g. methanol) to yield intermediate **7**.

[01216] Reaction (vi) may be performed by reacting primary amine **3**, **5** or **7** with a suitable reagent (e.g. trichloromethyl chloroformate) in a suitable solvent (e.g. dioxane) and, optionally, at an elevated temperature (e.g., 70° C or 80° C) to yield the intermediate **8**.

[01217] Reaction (vii) may be performed by reacting intermediate **8** with intermediate **9** in an inert environment in a suitable solvent (e.g. tetrahydrofuran). In some embodiments, reaction (vii) may be performed at a cooled temperature (e.g. 0° C), optionally, in the presence of an additive (e.g. CuCl), yielding a compound of **Formula (II)**. The compound of **Formula (II)** may be isolated by purification (e.g., by flash column chromatography or by preparative TLC or HPLC).

[01218] It should be understood that in the description and formulae shown above, the various groups are as defined herein, except where otherwise indicated. Furthermore, for synthetic purposes, the compounds in the Schemes are mere representatives with elected substituents to illustrate the general synthetic methodology of a compound disclosed herein.

[01219] It is understood that a neutral compound of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') may be converted to a salt (e.g., sodium salt) using routine techniques in the art (e.g., pH adjustment and, optionally, extraction (e.g., into an organic phase)). Further, a salt (e.g., sodium salt) of a compound of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') may be converted to a neutral compound using routine techniques in the art (e.g., pH adjustment and, optionally, extraction (e.g., into an aqueous phase)).

Biological Assays

[01220] Compounds designed, selected and/or optimised by methods described above, once produced, can be characterised using a variety of assays known to those skilled in the art to determine whether the compounds have biological activity. For example, the molecules can be

characterised by conventional assays, including but not limited to those assays described below, to determine whether they have a predicted activity, binding activity and/or binding specificity. [01221] Furthermore, high-throughput screening can be used to speed up analysis using such assays. As a result, it can be possible to rapidly screen the molecules described herein for activity, using techniques known in the art. General methodologies for performing high-throughput screening are described, for example, in Devlin (1998) *High Throughput Screening*, Marcel Dekker; and U.S. Patent No. 5,763,263. High-throughput assays can use one or more different assay techniques including, but not limited to, those described below.

[01222] Various *in vitro* or *in vivo* biological assays may be suitable for detecting the effect of the compounds of the present disclosure. These *in vitro* or *in vivo* biological assays can include, but are not limited to, enzymatic activity assays, electrophoretic mobility shift assays, reporter gene assays, *in vitro* cell viability assays, and the assays described herein.

[01223] In some embodiments, the compounds of the present disclosure may be tested for their inhibitory activity in various cell lines (*e.g.*, peripheral blood mononuclear cells). In some embodiments, the compounds of the present disclosure may be tested for their inhibitory activity in peripheral blood mononuclear cells. In some embodiments, the compounds of the present disclosure may be tested for their inhibitory activity against IL-1 β release upon NLRP3 activation.

[01224] In some embodiments, a PBMC IC₅₀ determination assay may be used to characterize the compounds of the present disclosure.

[01225] PBMC may be isolated, seeded into the wells of a plate, and incubated with a saccharide. Following medium exchange, the compounds of the present disclosure may be added to a well and incubated. The cells may be stimulated and the cell culture media collected for analysis.

[01226] PBMC may be isolated by density gradient centrifugation, seeded into the wells of a plate, and incubated with a saccharide. The compounds of the present disclosure may be added to a well and incubated. The cells may be stimulated and the cell culture media collected for analysis.

[01227] In some embodiments, release of IL-1 β may be determined by a quantitative detection. In some embodiments, release of IL-1 β may be determined by a quantitative detection of IL-1 β using an IL-1 β enzyme-linked immunosorbent assay (ELISA). A microplate spectrophotometer may be used to detect signals (*e.g.*, at 450 nm).

[01228] In some embodiments, release of IL-1 β may be determined by quantitative detection of IL-1 β using Homogenous Time-Resolved Fluorescence (HTRF®). A microplate spectrophotometer may be used to detect signals (e.g., at 655 nm and 620 nm).

[01229] In some embodiments, the biological assay is described in the Examples herein.

Pharmaceutical Compositions

[01230] In some aspects, the present disclosure provides a pharmaceutical composition comprising a compound of the present disclosure as an active ingredient. In some embodiments, the present disclosure provides a pharmaceutical composition comprising at least one compound of each of the formulae described herein, or a pharmaceutically acceptable salt or solvate thereof, and one or more pharmaceutically acceptable carriers or excipients. In some embodiments, the present disclosure provides a pharmaceutical composition comprising at least one compound selected from Tables 1 and 2.

[01231] As used herein, the term “composition” is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

[01232] The compounds of present disclosure can be formulated for oral administration in forms such as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups and emulsions. The compounds of present disclosure can also be formulated for intravenous (bolus or infusion), intraperitoneal, topical, subcutaneous, intramuscular or transdermal (e.g., patch) administration, all using forms well known to those of ordinary skill in the pharmaceutical arts.

[01233] The formulation of the present disclosure may be in the form of an aqueous solution comprising an aqueous vehicle. The aqueous vehicle component may comprise water and at least one pharmaceutically acceptable excipient. Suitable acceptable excipients include those selected from the group consisting of a solubility enhancing agent, chelating agent, preservative, tonicity agent, viscosity/suspending agent, buffer, and pH modifying agent, and a mixture thereof.

[01234] Any suitable solubility enhancing agent can be used. Examples of a solubility enhancing agent include cyclodextrin, such as those selected from the group consisting of hydroxypropyl- β -cyclodextrin, methyl- β -cyclodextrin, randomly methylated- β -cyclodextrin, ethylated- β -

cyclodextrin, triacetyl- β -cyclodextrin, peracetylated- β -cyclodextrin, carboxymethyl- β -cyclodextrin, hydroxyethyl- β -cyclodextrin, 2-hydroxy-3-(trimethylammonio)propyl- β -cyclodextrin, glucosyl- β -cyclodextrin, sulphated β -cyclodextrin (S- β -CD), maltosyl- β -cyclodextrin, β -cyclodextrin sulphobutyl ether, branched- β -cyclodextrin, hydroxypropyl- γ -cyclodextrin, randomly methylated- γ -cyclodextrin, and trimethyl- γ -cyclodextrin, and mixtures thereof.

[01235] Any suitable chelating agent can be used. Examples of a suitable chelating agent include those selected from the group consisting of ethylenediaminetetraacetic acid and metal salts thereof, disodium edetate, trisodium edetate, and tetrasodium edetate, and mixtures thereof.

[01236] Any suitable preservative can be used. Examples of a preservative include those selected from the group consisting of quaternary ammonium salts such as benzalkonium halides (preferably benzalkonium chloride), chlorhexidine gluconate, benzethonium chloride, cetyl pyridinium chloride, benzyl bromide, phenylmercury nitrate, phenylmercury acetate, phenylmercury neodecanoate, merthiolate, methylparaben, propylparaben, sorbic acid, potassium sorbate, sodium benzoate, sodium propionate, ethyl p-hydroxybenzoate, propylaminopropyl biguanide, and butyl-p-hydroxybenzoate, and sorbic acid, and mixtures thereof.

[01237] The aqueous vehicle may also include a tonicity agent to adjust the tonicity (osmotic pressure). The tonicity agent can be selected from the group consisting of a glycol (such as propylene glycol, diethylene glycol, triethylene glycol), glycerol, dextrose, glycerin, mannitol, potassium chloride, and sodium chloride, and a mixture thereof.

[01238] The aqueous vehicle may also contain a viscosity/suspending agent. Suitable viscosity/suspending agents include those selected from the group consisting of cellulose derivatives, such as methyl cellulose, ethyl cellulose, hydroxyethylcellulose, polyethylene glycols (such as polyethylene glycol 300, polyethylene glycol 400), carboxymethyl cellulose, hydroxypropylmethyl cellulose, and cross-linked acrylic acid polymers (carbomers), such as polymers of acrylic acid cross-linked with polyalkenyl ethers or divinyl glycol (Carbopols - such as Carbopol 934, Carbopol 934P, Carbopol 971, Carbopol 974 and Carbopol 974P), and a mixture thereof.

[01239] In order to adjust the formulation to an acceptable pH (typically a pH range of about 5.0 to about 9.0, more preferably about 5.5 to about 8.5, particularly about 6.0 to about 8.5, about 7.0 to about 8.5, about 7.2 to about 7.7, about 7.1 to about 7.9, or about 7.5 to about 8.0), the

formulation may contain a pH modifying agent. The pH modifying agent is typically a mineral acid or metal hydroxide base, selected from the group of potassium hydroxide, sodium hydroxide, and hydrochloric acid, and mixtures thereof, and preferably sodium hydroxide and/or hydrochloric acid. These acidic and/or basic pH modifying agents are added to adjust the formulation to the target acceptable pH range. Hence it may not be necessary to use both acid and base - depending on the formulation, the addition of one of the acid or base may be sufficient to bring the mixture to the desired pH range.

[01240] The aqueous vehicle may also contain a buffering agent to stabilise the pH. When used, the buffer is selected from the group consisting of a phosphate buffer (such as sodium dihydrogen phosphate and disodium hydrogen phosphate), a borate buffer (such as boric acid, or salts thereof including disodium tetraborate), a citrate buffer (such as citric acid, or salts thereof including sodium citrate), and ϵ -aminocaproic acid, and mixtures thereof.

[01241] The formulation may further comprise a wetting agent. Suitable classes of wetting agents include those selected from the group consisting of polyoxypropylene-polyoxyethylene block copolymers (poloxamers), polyethoxylated ethers of castor oils, polyoxyethylenated sorbitan esters (polysorbates), polymers of oxyethylated octyl phenol (Tyloxapol), polyoxyl 40 stearate, fatty acid glycol esters, fatty acid glyceryl esters, sucrose fatty esters, and polyoxyethylene fatty esters, and mixtures thereof.

[01242] Oral compositions generally include an inert diluent or an edible pharmaceutically acceptable carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavouring agent such as peppermint, methyl salicylate, or orange flavoring.

[01243] According to a further aspect of the disclosure there is provided a pharmaceutical composition which comprises a compound of the disclosure as defined hereinbefore, or a pharmaceutically acceptable salt, hydrate or solvate thereof, in association with a pharmaceutically acceptable diluent or carrier.

[01244] The compositions of the disclosure may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular, intraperitoneal or intramuscular dosing or as a suppository for rectal dosing).

[01245] The compositions of the disclosure may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

[01246] An effective amount of a compound of the present disclosure for use in therapy is an amount sufficient to treat or prevent an inflammasome related condition referred to herein, slow its progression and/or reduce the symptoms associated with the condition.

[01247] The size of the dose for therapeutic or prophylactic purposes of a compound of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well-known principles of medicine.

Methods of Use

[01248] In some aspects, the present disclosure provides a method of inhibiting inflammasome (e.g., the NLRP3 inflammasome) activity (e.g., *in vitro* or *in vivo*), comprising contacting a cell with an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof.

[01249] In some aspects, the present disclosure provides a method of treating or preventing a disease or disorder disclosed herein in a subject in need thereof, comprising administering to the

subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01250] In some aspects, the present disclosure provides a method of treating or preventing a disease or disorder disclosed herein in a subject in need thereof, comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01251] In some aspects, the present disclosure provides a method of treating a disease or disorder disclosed herein in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01252] In some aspects, the present disclosure provides a method of treating a disease or disorder disclosed herein in a subject in need thereof, comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01253] In some embodiments, the disease or disorder is associated with an implicated inflammasome activity. In some embodiments, the disease or disorder is a disease or disorder in which inflammasome activity is implicated.

[01254] In some embodiments, the disease or disorder is an inflammatory disorder, autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease, or cancer.

[01255] In some embodiments, the disease or disorder is an inflammatory disorder, autoinflammatory disorder and/or an autoimmune disorder.

[01256] In some embodiments, the disease or disorder is selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological disease (e.g. acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases).

[01257] In some embodiments, the disease or disorder is a neurodegenerative disease.

[01258] In some embodiments, the disease or disorder is Parkinson's disease or Alzheimer's disease.

[01259] In some embodiments, the disease or disorder is a dermatological disease.

[01260] In some embodiments, the dermatological disease is acne.

[01261] In some embodiments, the disease or disorder is cancer.

[01262] In some embodiments, the cancer is metastasising cancer, gastrointestinal cancer, skin cancer, non-small-cell lung carcinoma, brain cancer (e.g. glioblastoma) or colorectal adenocarcinoma.

[01263] In some embodiments, the subject is predisposed to the state, disorder, or condition (e.g., presence of a genetic variant).

[01264] In some aspects, the present disclosure provides a method of treating or preventing an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease or cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01265] In some aspects, the present disclosure provides a method of treating or preventing an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease or cancer in a subject in need thereof, comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01266] In some aspects, the present disclosure provides a method of treating an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease or cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01267] In some aspects, the present disclosure provides a method of treating an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease or cancer in a subject in need thereof, comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01268] In some aspects, the present disclosure provides a method of treating or preventing an inflammatory disorder, autoinflammatory disorder and/or an autoimmune disorder selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological disease (e.g. acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases) in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01269] In some aspects, the present disclosure provides a method of treating or preventing an inflammatory disorder, autoinflammatory disorder and/or an autoimmune disorder selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological disease (e.g. acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases) in a subject in need thereof, comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01270] In some aspects, the present disclosure provides a method of treating an inflammatory disorder, autoinflammatory disorder and/or an autoimmune disorder selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial

Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological disease (e.g. acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases) in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01271] In some aspects, the present disclosure provides a method of treating an inflammatory disorder, autoinflammatory disorder and/or an autoimmune disorder selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological disease (e.g. acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases) in a subject in need thereof, comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01272] In some aspects, the present disclosure provides a method of treating or preventing a neurodegenerative disease (e.g., Parkinson's disease or Alzheimer's disease) in a subject in need thereof, said method comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01273] In some aspects, the present disclosure provides a method of treating or preventing a neurodegenerative disease (e.g., Parkinson's disease or Alzheimer's disease) in a subject in need thereof, said method comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01274] In some aspects, the present disclosure provides a method of treating a neurodegenerative disease (e.g., Parkinson's disease or Alzheimer's disease) in a subject in need thereof, said method comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01275] In some aspects, the present disclosure provides a method of treating a neurodegenerative disease (e.g., Parkinson's disease or Alzheimer's disease) in a subject in need thereof, said method comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01276] In some aspects, the present disclosure provides a method of treating or preventing cancer in a subject in need thereof, said method comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01277] In some aspects, the present disclosure provides a method of treating or preventing cancer in a subject in need thereof, said method comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01278] In some aspects, the present disclosure provides a method of treating cancer in a subject in need thereof, said method comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01279] In some aspects, the present disclosure provides a method of treating cancer in a subject in need thereof, said method comprising administering to the subject an effective amount of a compound of the present disclosure or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the present disclosure.

[01280] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in inhibiting inflammasome (e.g., the NLRP3 inflammasome) activity (e.g., *in vitro* or *in vivo*).

[01281] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating or preventing a disease or disorder disclosed herein.

[01282] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating a disease or disorder disclosed herein.

[01283] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating or preventing an inflammatory disorder, an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease or cancer in a subject in need thereof.

[01284] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating an inflammatory disorder, an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease or cancer in a subject in need thereof.

[01285] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating or preventing an inflammatory disorder, an autoinflammatory disorder and/or an autoimmune disorder selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases) in a subject in need thereof.

[01286] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating an inflammatory disorder, an autoinflammatory disorder and/or an autoimmune disorder selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial

Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases) in a subject in need thereof.

[01287] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating or preventing a neurodegenerative disease (e.g., Parkinson's disease or Alzheimer's disease) in a subject in need thereof.

[01288] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating a neurodegenerative disease (e.g., Parkinson's disease or Alzheimer's disease) in a subject in need thereof.

[01289] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating or preventing cancer in a subject in need thereof.

[01290] In some aspects, the present disclosure provides a compound of the present disclosure or a pharmaceutically acceptable salt thereof for use in treating cancer in a subject in need thereof.

[01291] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for inhibiting inflammasome (e.g., the NLRP3 inflammasome) activity (e.g., *in vitro* or *in vivo*).

[01292] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing a disease or disorder disclosed herein.

[01293] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating a disease or disorder disclosed herein.

[01294] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing an inflammatory disorder, an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease or cancer in a subject in need thereof.

[01295] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for

treating an inflammatory disorder, an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease or cancer in a subject in need thereof.

[01296] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing an inflammatory disorder, an autoinflammatory disorder and/or an autoimmune disorder selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological disorders (e.g., acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases) in a subject in need thereof.

[01297] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating an inflammatory disorder, an autoinflammatory disorder and/or an autoimmune disorder selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological disorders (e.g., acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases) in a subject in need thereof.

[01298] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing a neurodegenerative disease (e.g., Parkinson's disease or Alzheimer's disease) in a subject in need thereof.

[01299] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating a neurodegenerative disease (e.g., Parkinson's disease or Alzheimer's disease) in a subject in need thereof.

[01300] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing cancer in a subject in need thereof.

[01301] In some aspects, the present disclosure provides use of a compound of the present disclosure or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating cancer in a subject in need thereof.

[01302] The present disclosure provides compounds that function as inhibitors of inflammasome activity. The present disclosure therefore provides a method of inhibiting inflammasome activity *in vitro* or *in vivo*, said method comprising contacting a cell with an effective amount of a compound, or a pharmaceutically acceptable salt thereof, as defined herein.

[01303] Effectiveness of compounds of the disclosure can be determined by industry-accepted assays/ disease models according to standard practices of elucidating the same as described in the art and are found in the current general knowledge.

[01304] The present disclosure also provides a method of treating a disease or disorder in which inflammasome activity is implicated in a patient in need of such treatment, said method comprising administering to said patient a therapeutically effective amount of a compound, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition as defined herein.

[01305] The present disclosure also provides a method of treating a disease or disorder in which inflammasome activity is implicated in a patient in need of such treatment, said method comprising administering to said patient an effective amount of a compound, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition as defined herein.

[01306] On a general level, the compounds of the present disclosure, which inhibit the maturation of cytokines of the IL-1 family, are effective in all therapeutic indications that are mediated or associated with elevated levels of active forms of cytokines belonging to IL-1 family of cytokines (Sims J. et al. Nature Reviews Immunology 10, 89-102 (February 2010)).

[01307] Exemplary diseases and the corresponding references will be given in the following: inflammatory, autoinflammatory and autoimmune diseases like CAPS (Dinarello CA. Immunity.

2004 Mar;20(3):243-4; Hoffman HM. al. *Reumatología* 2005; 21(3)), gout, rheumatoid arthritis (Gabay C et al. *Arthritis Research & Therapy* 2009, 11:230; Schett G. et al. *Nat Rev Rheumatol.* 2016 Jan;12(1):14-24.), Crohn's disease (Jung Mogg Kim *Korean J Gastroenterol* Vol. 58 No. 6, 300-310), COPD (Mortaz E. et al. *Tanaffos.* 2011; 10(2): 9–14.), fibrosis (Gasse P. et al. *Am J Respir Crit Care Med.* 2009 May 15;179(10):903-13), obesity, type 2 diabetes ((Dinarello CA. et al. *Curr Opin Endocrinol Diabetes Obes.* 2010 Aug;17(4):314-21)) multiple sclerosis (see EAE-model in Coll RC. et al. *Nat Med.* 2015 Mar;21(3):248-55) and many others (Martinon F. et al. *Immunol.* 2009. 27:229–65) like Parkinson's disease or Alzheimer's disease (Michael T. et al. *Nature* 493, 674–678 (31 January 2013); Halle A. et al., *Nat Immunol.* 2008 Aug;9(8):857-65; Saresella M. et al. *Mol Neurodegener.* 2016 Mar 3;11:23) and some oncological disorders.

[01308] Suitably, the compounds according to the present disclosure can be used for the treatment of a disease selected from the group consisting of an inflammatory disease, an autoinflammatory disease, an autoimmune disease, a neurodegenerative disease and cancer. Said inflammatory, autoinflammatory and autoimmune disease is suitably selected from the group consisting of a cryopyrin-associated autoinflammatory syndrome (CAPS, such as for example familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), chronic kidney disease (CKD), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, COPD, fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological diseases (e.g., acne) and neuroinflammation occurring in protein misfolding diseases, such as Prion diseases. Said neurodegenerative disease includes, but is not limited, to Parkinson's disease and Alzheimer's disease.

[01309] Accordingly, the compounds of the present disclosure can be used for the treatment of a disease selected from the group consisting of cryopyrin-associated autoinflammatory syndrome (CAPS, such as for example familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), chronic kidney disease (CKD), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, COPD, fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological diseases (e.g., acne)

neuroinflammation occurring in protein misfolding diseases, such as Prion diseases, neurodegenerative diseases (e.g., Parkinson's disease, Alzheimer's disease) and oncological disorders.

Treatment in Cancer; links with inflammasome

[01310] Chronic inflammation responses have long been observed to be associated with various types of cancer. During malignant transformation or cancer therapy inflammasomes may become activated in response to danger signals and this activation may be both beneficial and detrimental in cancer.

[01311] IL-1 β expression is elevated in a variety of cancers (including breast, prostate, colon, lung, head and neck cancers and melanomas) and patients with IL-1 β producing tumours generally have a worse prognosis (Lewis, Anne M., et al. "Interleukin-1 and cancer progression: the emerging role of interleukin-1 receptor antagonist as a novel therapeutic agent in cancer treatment." *Journal of translational medicine* 4.1 (2006): 48).

[01312] Cancers derived from epithelial cells (carcinoma) or epithelium in glands (adenocarcinoma) are heterogeneous; consisting of many different cell types. This may include fibroblasts, immune cells, adipocytes, endothelial cells and pericytes amongst others, all of which may be cytokine/ chemokine secreting (Grivennikov, Sergei I., Florian R. Greten, and Michael Karin. "Immunity, inflammation, and cancer." *Cell* 140.6 (2010): 883-899). This can lead to cancer-associated inflammation through the immune cell infiltration. The presence of leukocytes in tumours is known but it has only recently become evident that an inflammatory microenvironment is an essential component of all tumours. Most tumours (>90%) are the result of somatic mutations or environmental factors rather than germline mutations and many environmental causes of cancer are associated with chronic inflammation (20% of cancers are related to chronic infection, 30% to smoking/ inhaled pollutants and 35% to dietary factors (20% of all cancers are linked to obesity) (Aggarwal, Bharat B., R. V. Vijayalekshmi, and Bokyung Sung. "Targeting inflammatory pathways for prevention and therapy of cancer: short-term friend, long-term foe." *Clinical Cancer Research* 15.2 (2009): 425-430).

GI cancer

[01313] Cancers of the gastrointestinal (GI) tract are frequently associated with chronic inflammation. For example, *H. pylori* infection is associated with gastric cancer (Amieva, Manuel, and Richard M. Peek. "Pathobiology of *Helicobacter pylori*-Induced Gastric Cancer." *Gastroenterology* 150.1 (2016): 64-78). Colorectal cancer is associated with inflammatory bowel disease (Bernstein, Charles N., et al. "Cancer risk in patients with inflammatory bowel disease." *Cancer* 91.4 (2001): 854-862). Chronic inflammation in stomach leads to the upregulation of IL-1 and other cytokines (Basso D, et al., (1996) *Helicobacter pylori* infection enhances mucosal interleukin-1 beta, interleukin-6, and the soluble receptor of interleukin-2. *Int J Clin Lab Res* 26:207–210) and polymorphisms in IL-1 β gene can increase risk of gastric cancer (Wang P, et al., (2007) Association of interleukin-1 gene polymorphisms with gastric cancer: a meta-analysis. *Int J Cancer* 120:552–562).

[01314] In 19% of gastric cancer cases, caspase-1 expression is decreased which correlates with stage, lymph node metastasis and survival (Jee et al., 2005). *Mycoplasma hyorhinis* is associated with the development of gastric cancer its activation of the NLRP3 inflammasome may be associated with its promotion of gastric cancer metastasis (Xu et al., 2013).

Skin cancers

[01315] Ultraviolet radiation is the greatest environmental risk for skin cancer which is promoted by causing DNA damage, immunosuppression and inflammation. The most malignant skin cancer, melanoma, is characterised by the upregulation of inflammatory cytokines, all of which can be regulated by IL-1 β (Lázár-Molnár, Eszter, et al. "Autocrine and paracrine regulation by cytokines and growth factors in melanoma." *Cytokine* 12.6 (2000): 547-554). Systemic inflammation induces an enhancement of melanoma cell metastasis and growth by IL-1-dependent mechanisms in vivo. Using thymoquinone inhibition of metastasis in a B16F10 mouse melanoma model was shown to be dependent on inhibition of the NLRP3 inflammasome (Ahmad, Israr, et al. "Thymoquinone suppresses metastasis of melanoma cells by inhibition of NLRP3 inflammasome." *Toxicology and applied pharmacology* 270.1 (2013): 70-76).

Glioblastoma

[01316] NLRP3 contributes to radiotherapy resistance in glioma. Ionising radiation can induce NLRP3 expression whereas NLRP3 inhibition reduced tumour growth and prolonged mouse

survival following radiation therapy. NLRP3 inflammasome inhibition can therefore provide a therapeutic strategy for radiation-resistant glioma (Li, Lianling, and Yuguang Liu. "Aging-related gene signature regulated by Nlrp3 predicts glioma progression." *American journal of cancer research* 5.1 (2015): 442).

Metastasis

[01317] More widely, NLRP3 is considered by the applicants to be involved in the promotion of metastasis and consequently modulation of NLRP3 should plausibly block this. IL-1 is involved in tumour genesis, tumour invasiveness, metastasis, tumour host interactions (Apte, Ron N., et al. "The involvement of IL-1 in tumorigenesis, tumour invasiveness, metastasis and tumour-host interactions." *Cancer and Metastasis Reviews* 25.3 (2006): 387-408) and angiogenesis (Voronov, Elena, et al. "IL-1 is required for tumor invasiveness and angiogenesis." *Proceedings of the National Academy of Sciences* 100.5 (2003): 2645-2650).

[01318] The IL-1 gene is frequently expressed in metastases from patients with several types of human cancers. For example, IL-1 mRNA was highly expressed in more than half of all tested metastatic human tumour specimens including specifically non-small-cell lung carcinoma, colorectal adenocarcinoma, and melanoma tumour samples (Elaraj, Dina M., et al. "The role of interleukin 1 in growth and metastasis of human cancer xenografts." *Clinical Cancer Research* 12.4 (2006): 1088-1096) and IL-1RA inhibits xenograft growth in IL-1 producing tumours but without anti-proliferative effects in vitro.

[01319] Further, IL-1 signalling is a biomarker for predicting breast cancer patients at increased risk for developing bone metastasis. In mouse models IL-1 β and its receptor are upregulated in breast cancer cells that metastasise to bone compared with cells that do not. In a mouse model the IL-1 receptor antagonist anakinra reduced proliferation and angiogenesis in addition to exerting significant effects on the tumour environment reducing bone turnover markers, IL-1 β and TNF alpha (Holen, Ingunn, et al. "IL-1 drives breast cancer growth and bone metastasis in vivo." *Oncotarget* (2016).

[01320] IL-18 induced the production of MMP-9 in the human leukaemia cell line HL-60, thus favouring degradation of the extracellular matrix and the migration and invasiveness of cancer cells (Zhang, Bin, et al. "IL-18 increases invasiveness of HL-60 myeloid leukemia cells: up-regulation of matrix metalloproteinases-9 (MMP-9) expression." *Leukemia research* 28.1 (2004):

91-95). Additionally IL-18 can support the development of tumour metastasis in the liver by inducing expression of VCAM-1 on hepatic sinusoidal endothelium (Carrascal, Maria Teresa, et al. "Interleukin-18 binding protein reduces b16 melanoma hepatic metastasis by neutralizing adhesiveness and growth factors of sinusoidal endothelium." *Cancer Research* 63.2 (2003): 491-497).

CD36

[01321] The fatty acid scavenger receptor CD36 serves a dual role in priming gene transcription of pro-IL-1 β and inducing assembly of the NLRP3 inflammasome complex. CD36 and the TLR4-TLR6 heterodimer recognise oxLDL, which initiates a signalling pathway leading to transcriptional upregulation of NLRP3 and pro-IL-1 β (signal 1). CD36 also mediates the internalisation of oxLDL into the lysosomal compartment, where crystals are formed that induce lysosomal rupture and activation of the NLRP3 inflammasome (signal 2) (Kagan, J. and Horng T., "NLRP3 inflammasome activation: CD36 serves double duty." *Nature immunology* 14.8 (2013): 772-774).

[01322] A subpopulation of human oral carcinoma cells express high levels of the fatty acid scavenger receptor CD36 and are unique in their ability to initiate metastasis. Palmitic acid or a high fat diet boosted the metastatic potential of the CD36⁺ cells. Neutralising anti-CD36 antibodies blocked metastasis in orthotopic mouse models of human oral cancer. The presence of CD36⁺ metastasis-initiating cells correlates with a poor prognosis for numerous types of carcinomas. It is suggested that dietary lipids may promote metastasis (Pasqual, G, Avgustinova, A., Mejetta, S, Martin, M, Castellanos, A, Attolini, CS-O, Berenguer, A., Prats, N, Toll, A, Hueto, JA, Bescos, C, Di Croce, L, and Benitah, SA. 2017 "Targeting metastasis-initiating cells through the fatty acid receptor CD36" *Nature* 541:41-45).

[01323] In hepatocellular carcinoma exogenous palmitic acid activated an epithelial-mesenchymal transition (EMT)-like program and induced migration that was decreased by the CD36 inhibitor, sulpho-N-succinimidyl oleate (Nath, Aritro, et al. "Elevated free fatty acid uptake via CD36 promotes epithelial-mesenchymal transition in hepatocellular carcinoma." *Scientific reports* 5 (2015). Body mass index was not associated with the degree of EMT highlighting that it is actually CD36 and free fatty acids that are important.

[01324] Cancer stems cells (CSCs) use CD36 to promote their maintenance. Oxidised phospholipids, ligands of CD36, were present in glioblastoma and the proliferation of CSCs but not non-CSCs increased with exposure to oxidised LDL. CD36 also correlated with patient prognosis.

Chemotherapy resistance

[01325] In addition to direct cytotoxic effects, chemotherapeutic agents harness the host immune system which contributes to anti-tumour activity. However, gemcitabine and 5-FU were shown to activate NLRP3 in myeloid-derived suppressor cells leading to production of IL-1 β which curtails anti-tumour efficacy. Mechanistically these agents destabilised the lysosome to release cathepsin B to activate NLRP3. IL-1 β drove the production of IL-17 from CD4⁺ T cells which in turn blunted the efficacy of the chemotherapy. Higher anti-tumoral effects for both gemcitabine and 5-FU were observed when tumours were established in NLRP3^{-/-} or Caps1^{-/-} mice, or WT mice treated with IL-1RA. Myeloid-derived suppressor cell NLRP3 activation therefore limits the anti-tumour efficacy of gemcitabine and 5-FU (Bruchard, Mélanie, et al. "Chemotherapy-triggered cathepsin B release in myeloid-derived suppressor cells activates the Nlrp3 inflammasome and promotes tumour growth." *Nature medicine* 19.1 (2013): 57-64.).

Compounds of the present disclosure may therefore be useful in chemotherapy to treat a range of cancers.

[01326] Compounds of the present disclosure, or pharmaceutically acceptable salts thereof, may be administered alone as a sole therapy or can be administered in addition with one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment.

[01327] For example, therapeutic effectiveness may be enhanced by administration of an adjuvant (i.e. by itself the adjuvant may only have minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the individual is enhanced). Alternatively, by way of example only, the benefit experienced by an individual may be increased by administering the compound of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') with another therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit.

[01328] In the instances where the compound of the present disclosure is administered in combination with other therapeutic agents, the compound of the disclosure need not be administered via the same route as other therapeutic agents, and may, because of different physical and chemical characteristics, be administered by a different route. For example, the compound of the disclosure may be administered orally to generate and maintain good blood levels thereof, while the other therapeutic agent may be administered intravenously. The initial administration may be made according to established protocols known in the art, and then, based upon the observed effects, the dosage, modes of administration and times of administration can be modified by the skilled clinician.

[01329] The particular choice of other therapeutic agent will depend upon the diagnosis of the attending physicians and their judgment of the condition of the individual and the appropriate treatment protocol. According to this aspect of the disclosure there is provided a combination for use in the treatment of a disease in which inflammasome activity is implicated comprising a compound of the disclosure as defined hereinbefore, or a pharmaceutically acceptable salt thereof, and another suitable agent.

[01330] According to a further aspect of the disclosure there is provided a pharmaceutical composition which comprises a compound of the disclosure, or a pharmaceutically acceptable salt thereof, in combination with a suitable, in association with a pharmaceutically acceptable diluent or carrier.

[01331] In addition to its use in therapeutic medicine, compounds of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') and pharmaceutically acceptable salts thereof are also useful as pharmacological tools in the development and standardisation of in vitro and in vivo test systems for the evaluation of the effects of inhibitors of inflammasome in laboratory animals such as dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

[01332] In any of the above-mentioned pharmaceutical composition, process, method, use, medicament, and manufacturing features of the instant disclosure, any of the alternate embodiments of macromolecules of the present disclosure described herein also apply.

Routes of Administration

[01333] The compounds of the disclosure or pharmaceutical compositions comprising these compounds may be administered to a subject by any convenient route of administration, whether systemically/ peripherally or topically (i.e., at the site of desired action).

[01334] Routes of administration include, but are not limited to, oral (e.g. by ingestion); buccal; sublingual; transdermal (including, e.g., by a patch, plaster, etc.); transmucosal (including, e.g., by a patch, plaster, etc.); intranasal (e.g., by nasal spray); ocular (e.g., by eye drops); pulmonary (e.g., by inhalation or insufflation therapy using, e.g., via an aerosol, e.g., through the mouth or nose); rectal (e.g., by suppository or enema); vaginal (e.g., by pessary); parenteral, for example, by injection, including subcutaneous, intradermal, intramuscular, intravenous, intra-arterial, intracardiac, intrathecal, intraspinal, intracapsular, subcapsular, intraorbital, intraperitoneal, intratracheal, subcuticular, intraarticular, subarachnoid, and intrasternal; by implant of a depot or reservoir, for example, subcutaneously or intramuscularly.

EXAMPLES

[01335] For exemplary purpose, neutral compounds of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') are synthesized and tested in the examples. It is understood that the neutral compounds of Formula (I), (I'), (II), (II'), (III), (III'), (IV), (IV'), (V), (V'), (VI), or (VI') may be converted to the corresponding pharmaceutically acceptable salts of the compounds using routine techniques in the art (e.g., by saponification of an ester to the carboxylic acid salt, or by hydrolyzing an amide to form a corresponding carboxylic acid and then converting the carboxylic acid to a carboxylic acid salt).

[01336] Nuclear magnetic resonance (NMR) spectra were recorded at 400 MHz or 300 MHz as stated and at 300.3 K unless otherwise stated; the chemical shifts (δ) are reported in parts per million (ppm). Spectra were recorded using a Bruker or Varian instrument with 8, 16 or 32 scans.

[01337] LC-MS chromatograms and spectra were recorded using an Agilent 1200 or Shimadzu LC-20 AD&MS 2020 instrument using a C-18 column such as a Luna-C18 2.0x30 mm or Xbridge Shield RPC18 2.1x50 mm. Injection volumes were 0.7 – 8.0 μ l and the flow rates were typically 0.8 or 1.2 ml/min. Detection methods were diode array (DAD) or evaporative light scattering (ELSD) as well as positive ion electrospray ionisation. MS range was 100 - 1000 Da.


Solvents were gradients of water and acetonitrile both containing a modifier (typically 0.01 – 0.04 %) such as trifluoroacetic acid or ammonium carbonate.

[01338] Abbreviations:

ACN	Acetonitrile
ACN- <i>d</i> ₃	Trideuteroacetonitrile
BH ₃ .THF	Borane tetrahydrofuran complex solution
CDCl ₃	Chloroform- <i>d</i>
DCC	N,N'-Dicyclohexylcarbodiimide
DCM	Dichloromethane
DIPEA	N,N-Diisopropylethylamine
DMAP	4-Dimethylaminopyridine
DMF	N,N-dimethylformamide
DMSO	Dimethylsulphoxide
DMSO- <i>d</i> ₆	Hexadeuterodimethylsulphoxide
dppf	1,1'-Bis(diphenylphosphino)ferrocene
eq.	Equivalents
ESI	Electrospray ionisation
EtOAc	Ethyl acetate
FCC	Flash column chromatography
H	Hour(s)
HATU	1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate
¹ H NMR	Proton nuclear magnetic resonance spectroscopy
HPLC	High performance liquid chromatography
LC-MS	Liquid chromatography-mass spectrometry
MeOD	Methanol- <i>d</i> ₄
MeOH	Methanol
min	Minute(s)
NaOAc	Sodium acetate
Pd(PPh ₃) ₄	Tetrakis(triphenylphosphine)palladium(0)
pet. ether	Petroleum ether


ppm	Parts per million
RM	Reaction mixture
TEA	Triethylamine
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran
TLC	Thin layer chromatography
Y	Yield

General Procedure A:




[01339] To a solution of the ketone (1 eq.) in THF (0.6 M) was added (*S*)-2-methylpropane-2-sulphinamide (1.5 eq.) and titanium isopropoxide (3 eq.). The solution was stirred for 12 h at 80° C under N₂. The RM was cooled to 0° C and MeOH (1.5 eq.) and LiBH₄ (1.5 eq.) were added. The RM was stirred at 25° C for 1 h then diluted with water and extracted twice with EtOAc. The combined organic phases were concentrated to give the desired product.

General procedure B:




[01340] The sulphinamide from General procedure A (1 eq.) was dissolved in 4M HCl in MeOH (0.1 M) and the solution was stirred at 25° C for 1 h. The solution was concentrated *in vacuo* to give the desired product.

General procedure C:




[01341] To a solution of amine hydrochloride from General procedure B (1 eq.) in dioxane (0.2 M) was added trichloromethyl chloroformate (1.1 eq.) and the mixture heated at 70° C for 1 h or until complete. The solution was concentrated *in vacuo* to give the desired product.

General procedure D:



[01342] To a mixture of the alcohol (1 eq.) and CuCl (1.1 eq.) in THF (0.2 M) was added the isocyanate from general procedure C (1 eq.) in one portion at 0° C under N₂. The mixture was stirred at 25° C for 10 h. The RM was diluted with H₂O and extracted three times with EtOAc. The combined organic layers were washed (brine), dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by prep-TLC or prep HPLC to give the title compound.

Example 1. Ethyl 2-(((1S)-1-(4-chlorophenyl)ethyl)carbamoyl)oxy-3-(1H-pyrazol-1-yl)propanoate




[01343] **Step 1: 1-chloro-4-[(1S)-1-isocyanatoethyl]benzene.** To a stirred solution of (1S)-1-(4-chlorophenyl)ethan-1-amine (0.1 g, 0.64 mmol, 1 eq.) in THF (1 ml) was added 20 % phosgene in toluene (0.4 ml, 0.77 mmol, 1.2 eq.) at 0° C under N₂. The RM was heated at 80° C for 3 h and concentrated *in vacuo* to obtain the 1-chloro-4-[(1S)-1-isocyanatoethyl]benzene, which was used crude in the next reaction (Y = 86 %).

[01344] **Step-2: Ethyl 2-(((S)-1-(4-chlorophenyl)ethyl)carbamoyl)oxy-3-(1H-pyrazol-1-yl)propanoate.** To a mixture of ethyl 2-hydroxy-3-(1H-pyrazol-1-yl)propanoate (synthesized in WO 2019025467 A1) (0.1 g, 0.54 mmol, 1 eq.) and 1-chloro-4-[(1S)-1-isocyanatoethyl]benzene (0.1 g, 0.54 mmol, 1 eq.) in DMF (2 ml) was added CuCl (0.06 g, 0.59 mmol, 1.1 eq.) at room

temperature under N₂. The RM was stirred at RT for 15 min then poured into water (20 ml) and extracted with EtOAc (2 x 10 ml). The organic phases were combined, washed with cold water (5 x 10 ml), dried (Na₂SO₄) and concentrated *in vacuo*. Purification by FCC twice (SiO₂, 25 % EtOAc in hexane) gave the ethyl 2-((((S)-1-(4-chlorophenyl)ethyl)carbamoyl)oxy)-3-(1H-pyrazol-1-yl)propanoate as a colourless viscous liquid (Y = 8 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.09 (d, *J* = 8 Hz, 1H), 7.73 – 7.71 (m, 1H), 7.46 – 7.43 (m, 1H), 7.38 – 7.26 (m, 5H), 6.26 – 6.25 (m, 1H), 5.19 – 5.13 (m, 1H), 4.67 – 4.58 (m, 1H), 4.55 – 4.52 (m, 2H), 4.12 – 4.01 (m, 2H), 1.33 – 1.24 (m, 5H), 1.19 – 1.13 (m, 2H), 1.09 – 1.01 (m, 2H). LC-MS (ESI): *m/z*: [M+H]⁺ = 366.1


Example 2. Propan-2-yl 2-((((1S)-1-(3,4-dichlorophenyl)ethyl)carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate



[01345] **Step 1: 1,2-Dichloro-4-[(1S)-1-isocyanatoethyl]benzene.** To a solution of triphosgene (390 mg, 1.32 mmol, 1 eq.) in DCM (3 ml) was added (1S)-1-(3,4-dichlorophenyl)ethanamine (250 mg, 1.32 mmol, 1 eq.) and TEA (384 µl, 2.76 mmol, 2.1 eq.) at 20° C. After 1 h, TLC (pet. ether: EtOAc, 2:1) indicated the reaction was complete. The solvent was removed *in vacuo* to afford the crude product as a yellow solid.

[01346] **Step 2: Propan-2-yl 2-((((1S)-1-(3,4-dichlorophenyl)ethyl)carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate.** General procedure D was followed using isopropyl 2-hydroxy-3-pyrimidin-2-ylpropanoate (synthesized in WO 2019025467 A1) and 1,2-dichloro-4-[(1S)-1-isocyanatoethyl]benzene. Prep-TLC (SiO₂, pet. ether : EtOAc, 1:2) gave the propan-2-yl 2-((((1S)-1-(3,4-dichlorophenyl)ethyl)carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate as a yellow gum (Y = 39 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.69 - 8.65 (m, 2H), 7.44 - 7.30 (m, 2H), 7.17 - 7.11 (m, 2H), 5.67 - 5.61 (m, 1H), 5.15 - 5.05 (m, 1H), 4.80 - 4.70 (m, 1H), 3.56 - 3.40 (m, 2H), 1.44 - 1.41 (m, 3H), 1.26 - 1.13 (m, 6H). LC-MS (ESI): *m/z*: [M+H]⁺ = 426.0


Example 3. Propan-2-yl 2-({[(1S)-1-(4-chlorophenyl)ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate



[01347] **Step 1: 1-Chloro-4-[(1S)-1-isocyanatoethyl]benzene.** To a solution of triphosgene (477 mg, 1.61 mmol, 1 eq.) in DCM (3 ml) was added (1S)-1-(4-chlorophenyl)ethanamine (250 mg, 1.61 mmol, 1 eq.) and TEA (470 μ l, 3.37 mmol, 2.1 eq.) at 20 °C. After 1 h, TLC (pet. ether : EtOAc, 2:1) indicated that the reaction was complete. The solvent was removed *in vacuo* to give 1-chloro-4-[(1S)-1-isocyanatoethyl]benzene as a yellow solid.

[01348] **Step 2: Propan-2-yl 2-({[(1S)-1-(4-chlorophenyl)ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate.** General procedure D was followed using isopropyl 2-hydroxy-3-pyrimidin-2-ylpropanoate and 1-chloro-4-[(1S)-1-isocyanatoethyl]benzene. The crude product was purified by prep-TLC (SiO₂, pet. ether: EtOAc, 1:2) to give propan-2-yl 2-({[(1S)-1-(4-chlorophenyl)ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate as a yellow gum (Y = 47 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.69 (d, *J* = 3 Hz, 1H), 8.64 (d, *J* = 5 Hz, 1H) 7.35 - 7.25 (m, 1H), 7.20 - 7.15 (m, 4H), 5.66 - 5.61 (m, 1H), 5.16 - 5.01 (m, 2H), 4.79 - 4.76 (m, 1H), 3.53 - 3.46 (m, 2H), 1.48 - 1.40 (m, 3H), 1.32 - 1.11 (m, 6H). LC-MS (ESI): *m/z*: [M+H] = 392.1

Example 4. Propan-2-yl 2-({[(1S)-1-(4-cyanophenyl)ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate




[01349] **Step 1: 4-[(1S)-1-Isocyanatoethyl]benzonitrile.** To a solution of triphosgene (406 mg, 1.37 mmol, 1 eq.) in DCM (5 ml) was added 4-[(1S)-1-aminoethyl]benzonitrile (250 mg, 1.37 mmol, 1 eq. HCl salt) and TEA (400 μ l, 2.87 mmol, 2.1 eq.) at 20° C. After 1 h, TLC (pet. ether:

EtOAc, 1:1) indicated that the reaction was complete. The mixture was concentrated *in vacuo* to give 4-[(1S)-1-isocyanatoethyl]benzonitrile as a yellow solid, which was used immediately.

[01350] **Step 2: Propan-2-yl 2-({[(1S)-1-(4-cyanophenyl)ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate.** General procedure D was followed, using isopropyl 2-hydroxy-3-pyrimidin-2-yl-propanoate and 4-[(1S)-1-isocyanatoethyl]benzonitrile. The crude product was purified by prep-TLC (SiO₂, pet. ether: EtOAc, 1:2) to give propan-2-yl 2-({[(1S)-1-(4-cyanophenyl)ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate as a colourless gum (Y = 14 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.77 (d, *J* = 5 Hz, 1H), 8.74 (d, *J* = 5 Hz, 1H), 8.07 (t, *J* = 8 Hz, 1H), 7.80 - 7.77 (m, 2H), 7.49 (d, *J* = 8 Hz, 1H), 7.43 - 7.39 (m, 2H), 5.41 - 5.37 (m, 1H), 4.97 - 4.88 (m, 1H), 4.70 - 4.65 (m, 1H), 3.40 - 3.36 (m, 2H), 1.31 - 1.27 (m, 3H), 1.15 (d, *J* = 6 Hz, 1.5H), 1.12 - 1.10 (m, 3H), 0.95 (d, *J* = 6 Hz, 1.5H). LC-MS (ESI): *m/z*: [M+H] = 383.1

Example 5. Propan-2-yl 3-(pyrimidin-2-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoate




[01351] **Step 1: 1-[(1S)-1-Isocyanatoethyl]-4-(trifluoromethyl)benzene.** To a solution of triphosgene (314 mg, 1.06 mmol, 1 eq.) in DCM (5 ml) was added (1S)-1-[4-(trifluoromethyl)phenyl]ethanamine (200 mg, 1.06 mmol, 1 eq.) and TEA (309 µl, 2.22 mmol, 2.1 eq.) at 0° C. After 1 h, TLC (pet. ether: EtOAc, 1:1) indicated that the reaction was complete. The mixture was concentrated *in vacuo* to give the crude product as a yellow solid. This product was used directly in the following reaction.

[01352] **Step 2: Propan-2-yl 3-(pyrimidin-2-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoate.** General procedure D was followed using isopropyl 2-hydroxy-3-pyrimidin-2-yl-propanoate and 1-[(1S)-1-isocyanatoethyl]-4-(trifluoromethyl)benzene. The crude product was purified by prep-TLC (SiO₂, pet. ether: EtOAc = 1:2) to give propan-2-yl 3-(pyrimidin-2-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoate as a white solid (Y = 9 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.77 - 8.72 (m, 2H), 8.07 (t, *J* = 8 Hz, 1H), 7.68 - 7.65 (m, 2H),

7.51 (d, $J = 8$ Hz, 1H), 7.46 - 7.37 (m, 2H), 5.41 - 5.35 (m, 1H), 4.95 - 4.78 (m, 1H), 4.68 - 4.62 (m, 1H), 3.39 - 3.34 (m, 2H), 1.33 - 1.29 (m, 3H), 1.15 - 0.90 (m, 6H). LC-MS (ESI): m/z : $[M+H] = 426.1$

Example 6. Propan-2-yl (2R)-2-([(1S)-1-(4-tert-butylphenyl)ethyl]carbamoyl)oxy-3-(1H-1,2,4-triazol-1-yl)propanoate



[01353] Step 1: (S)-N-[(1S)-1-(4-Tert-butylphenyl)ethyl]-2-methylpropane-2-sulphinamide.

General procedure A was followed using 1-(4-tert-butylphenyl)ethan-1-one. The crude product was purified by column chromatography (SiO_2 , pet. ether: EtOAc, 3:1) to give (S)-N-[(1S)-1-(4-tert-butylphenyl)ethyl]-2-methylpropane-2-sulphinamide as a white solid ($Y = 43\%$). ^1H NMR (400 MHz, CDCl_3) δ ppm 7.36 - 7.33 (m, 2H), 7.26 - 7.24 (m, 2H), 4.52 - 4.50 (m, 1H), 3.37 (s, 1H), 1.48 (d, $J = 6$ Hz, 3H), 1.30 (s, 9H), 1.21 (s, 9H).


[01354] Step 2: (1S)-1-(4-Tert-butylphenyl)ethan-1-aminium chloride. General procedure B was followed using (S)-N-[(1S)-1-(4-(tert-butyl)phenyl)ethyl]-2-methylpropane-2-sulphinamide to give (1S)-1-(4-tert-butylphenyl)ethan-1-aminium chloride as a white solid ($Y = 75\%$). ^1H NMR (400 MHz, CDCl_3) δ ppm 8.45 - 8.25 (br. s, 3H), 7.44 - 7.38 (m, 4H), 4.42 - 4.35 (m, 1H), 1.67 (d, $J = 7$ Hz, 3H), 1.27 (s, 9H).

[01355] Step 3: 1-Tert-butyl-4-[(1S)-1-isocyanatoethyl]benzene. General procedure C was followed using (1S)-1-(4-tert-butylphenyl)ethanamine to give 1-tert-butyl-4-[(1S)-1-isocyanatoethyl]benzene as yellow oil ($Y = \text{quantitative}$). LC-MS (ESI): m/z : $[M+\text{MeOH}+H]^+ = 236.2$

[01356] Step 4: Propan-2-yl (2R)-2-([(1S)-1-(4-tert-butylphenyl)ethyl]carbamoyl)oxy-3-(1H-1,2,4-triazol-1-yl)propanoate. General procedure D was followed using isopropyl (R)-

isopropyl 2-hydroxy-3-(1H-1,2,4-triazol-1-yl)propanoate (synthesised in WO 2019025467 A1) and (S)-1-(tert-butyl)-4-(1-isocyanatoethyl)benzene. The crude product was purified by prep-HPLC (column: Xbridge, 10 μ m, 100 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 40 - 100 %, 10 min) to give propan-2-yl (2R)-2-({[(1S)-1-(4-tert-butylphenyl)ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate as a white solid (Y = 6 %). ^1H NMR (400 MHz, CDCl_3) δ ppm 8.37 (s, 1H), 8.05 (s, 1H), 7.36 (d, J = 8 Hz, 2H), 7.24 (d, J = 8 Hz, 2H), 5.37 (t, J = 5 Hz, 1H), 5.30 (d, J = 8 Hz, 1H), 5.07 - 5.04 (m, 1H), 4.78 (t, J = 7 Hz, 1H), 4.69 (d, J = 4 Hz, 2H), 1.49 (d, J = 7 Hz, 3H), 1.31 (s, 9H), 1.21 - 1.16 (m, 6H). LC-MS (ESI): m/z : $[\text{M}+\text{H}] = 403.2$

Example 7. Propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-({[(1S)-1-[4-(2,2,2-trifluoroethoxy)phenyl]ethyl]carbamoyl}oxy)propanoate



[01357] **Step 1: 1-[4-(2,2,2-Trifluoroethoxy)phenyl]ethan-1-one.** To a solution of NaH (705 mg, 17.63 mmol, 60 % in mineral oil, 1.2 eq.) in DMF (20 ml) was added 1-(4-hydroxyphenyl)ethanone (2.0 g, 14.69 mmol, 1 eq.) dropwise at 25° C and the mixture was stirred for 0.5 h. A solution of 2,2,2-trifluoroethyl 4-methylbenzenesulphonate (4.48 g, 17.63 mmol, 1.2 eq.) in DMF (10 ml) was added dropwise at 25° C and the mixture stirred at 130° C for 12 h. The RM was concentrated *in vacuo* and the residue purified by prep-HPLC (column: Phenomenex Luna C18, 10 μ m, 250 x 50 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 40 - 70 %, 20 min) to give 1-[4-(2,2,2-trifluoroethoxy)phenyl]ethan-1-one as a white solid (Y = 62 %). ^1H NMR (400 MHz, CDCl_3) δ ppm 7.98 - 7.96 (m, 2H), 7.00 - 6.98 (m, 2H), 4.45 - 4.39 (m, 2H), 2.58 (s, 3H).

[01358] **Step 2: (S)-2-Methyl-N-[(1S)-1-[4-(2,2,2-trifluoroethoxy)phenyl]ethyl]propane-2-sulphinamide.** General procedure A was followed using 1-[4-(2,2,2-

trifluoroethoxy)phenyl]ethan-1-one. The crude product was purified by column chromatography (SiO₂, pet. ether: EtOAc, 1:0 to 1:1) to give (S)-2-methyl-N-[(1S)-1-[4-(2,2,2-trifluoroethoxy)phenyl]ethyl]propane-2-sulphinamide as a white solid (Y = quantitative).


¹H NMR (400 MHz, CDCl₃) δ ppm 7.32 - 7.27 (m, 2H), 6.93 - 6.89 (m, 2H), 4.53 - 4.49 (m, 1H), 4.37 - 4.31 (m, 2H), 3.40 (d, *J* = 2 Hz, 1H), 1.48 (d, *J* = 7 Hz, 3H), 1.22 (s, 9H).

[01359] **Step 3: (1S)-1-[4-(2,2,2-Trifluoroethoxy)phenyl]ethan-1-aminium chloride.** General procedure B was followed using (S)-2-methyl-N-((S)-1-(4-(2,2,2-trifluoroethoxy)phenyl)ethyl)propane-2-sulphinamide to give (1S)-1-[4-(2,2,2-trifluoroethoxy)phenyl]ethan-1-aminium chloride as a white solid (Y = 25 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.43 (s, 3H), 7.47 (d, *J* = 8 Hz, 2H), 6.93 (d, *J* = 8 Hz, 2H), 4.38 - 4.30 (m, 3H), 1.67 (d, *J* = 6 Hz, 3H).

[01360] **Step 4: 1-[(1S)-1-Isocyanatoethyl]-4-(2,2,2-trifluoroethoxy)benzene.** General procedure C was followed using (1S)-1-(4-(2,2,2-trifluoroethoxy)phenyl)ethan-1-aminium chloride to give the title compound (Y = 95 %). LC-MS (ESI): *m/z*: [M - NCO] = 203.1

[01361] **Step 5: Propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-({[(1S)-1-[4-(2,2,2-trifluoroethoxy)phenyl]ethyl]carbamoyl}oxy)propanoate.** General procedure D was followed using 1-[(1S)-1-(1-isocyanatoethyl)-4-(2,2,2-trifluoroethoxy)benzene and isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate. The crude product was purified by prep-TLC to give 1-[(1S)-1-isocyanatoethyl]-4-(2,2,2-trifluoroethoxy)benzene as a white solid (Y = 5 %). ¹H NMR (400 MHz, ACN-*d*₃) δ ppm 8.23 (s, 1H), 7.85 (s, 1H), 7.26 (d, *J* = 9 Hz, 2H), 6.94 (d, *J* = 9 Hz, 2H), 6.26 (d, *J* = 8 Hz, 1H), 5.14 (t, *J* = 10 Hz, 1H), 4.95 - 4.89 (m, 1H), 4.68 - 4.61 (m, 1H), 4.59 (d, *J* = 5 Hz, 2H), 4.53 - 4.47 (m, 2H), 1.37 (d, *J* = 7 Hz, 3H), 1.14 (d, *J* = 6 Hz, 3H), 1.10 (d, *J* = 6 Hz, 3H). LC-MS (ESI): *m/z*: [M+H]⁺ = 445.2

Example 8. Propan-2-yl (2R)-2-({[(1S)-1-[4-(phenoxyethyl)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate



[01362] **Step 1: 1-[4-(Phenoxyethyl)phenyl]ethan-1-one.** To a solution of 1-[4-(bromomethyl)phenyl]ethan-1-one (2.0 g, 9.39 mmol, 1 eq.) in ACN (20 ml) was added phenol (883 mg, 9.39 mmol, 826 μ l, 1 eq.) and K_2CO_3 (3.24 g, 23.47 mmol, 2.5 eq.) at 25° C. The RM was stirred at 80° C for 12 h, when LC-MS indicated the product had formed. The mixture was concentrated *in vacuo* and the residue partitioned between EtOAc (20 ml) and H_2O (20 ml). The organic phase was separated, washed (brine), dried (Na_2SO_4) and concentrated *in vacuo* to give 1-[4-(phenoxyethyl)phenyl]ethan-1-one as a white solid (Y = 94 %). 1H NMR (400 MHz, $DMSO-d_6$) δ ppm 7.98 - 7.96 (m, 2H), 7.58 (d, J = 8 Hz, 2H), 7.33 - 7.24 (m, 2H), 7.01 (d, J = 8 Hz, 2H), 6.94 (t, J = 7 Hz, 1H), 5.20 (s, 2H), 2.57 (s, 3H).

[01363] **Step 2: (S)-2-Methyl-N-[(1S)-1-[4-(phenoxyethyl)phenyl]ethyl]propane-2-sulphinamide.** General procedure A was followed using 1-[4-(phenoxyethyl)phenyl]ethan-1-one. The crude product was purified by column chromatography (SiO_2 , pet. ether: EtOAc, 1:0 to 1:1 to give (S)-2-methyl-N-[(1S)-1-[4-(phenoxyethyl)phenyl]ethyl]propane-2-sulphinamide as a white solid (Y = 51 %). 1H NMR (400 MHz, $CDCl_3$) δ ppm 7.44 (d, J = 8 Hz, 2H), 7.38 (d, J = 8 Hz, 2H), 7.33 - 7.27 (m, 2H), 7.00 - 6.96 (m, 3H), 5.01 (s, 2H), 4.55 - 4.60 (m, 1H), 3.44 (s, 1H), 1.53 (d, J = 6 Hz, 3H), 1.25 (s, 9H).

[01364] **Step 3: (1S)-1-[4-(Phenoxyethyl)phenyl]ethan-1-aminium chloride.** General procedure B was followed using (S)-2-methyl-N-[(S)-1-[4-(phenoxyethyl)phenyl]ethyl]propane-2-sulphinamide to give (1S)-1-[4-(phenoxyethyl)phenyl]ethan-1-aminium chloride as a white solid (Y = 50 %). 1H NMR (400 MHz, $CDCl_3$) δ ppm 8.80 - 8.60 (br. s, 3H), 7.53 - 7.46 (m, 4H), 7.31 - 7.27 (m, 2H), 7.00 - 6.92 (m, 3H), 5.02 (s, 2H), 4.45 - 4.27 (m, 1H), 1.75 - 1.60 (m, 3H).

[01365] **Step 4: 1-[(1S)-1-Isocyanatoethyl]-4-(phenoxyethyl)benzene.** General procedure C was followed using (1S)-1-[4-(phenoxyethyl)phenyl]ethan-1-aminium chloride to give 1-[(1S)-


1-isocyanatoethyl]-4-(phenoxyethyl)benzene as a yellow oil (Y = quantitative). LC-MS (ESI): m/z : $[M+MeOH+H]^+ = 286.1$

[01366] **Step 5: Propan-2-yl (2R)-2-({[(1S)-1-[4-**

(phenoxyethyl)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate. General procedure D was followed using isopropyl (2R)-2-carbamoyloxy-3-(1,2,4-triazol-1-yl)propanoate and 1-(1S)-1-(1-isocyanatoethyl)-4-(phenoxyethyl)benzene. The crude material was purified by prep-HPLC (column: Xbridge, 10 μ m, 150 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 35 - 65 %, 10 min) to give propan-2-yl (2R)-2-({[(1S)-1-[4-(phenoxyethyl)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate as a white solid (Y = 40 %). 1H NMR (400 MHz, $ACN-d_3$) δ ppm 8.28 (s, 1H), 7.88 (s, 1H), 7.40 (d, $J = 8$ Hz, 2H), 7.32 - 7.27 (m, 4H), 7.02 - 6.91 (m, 3H), 6.31 (d, $J = 7$ Hz, 1H), 5.16 (t, $J = 5$ Hz, 1H), 5.07 (s, 2H), 4.94 - 4.88 (m, 1H), 4.72 - 4.65 (m, 1H), 4.60 (d, $J = 5$ Hz, 2H), 1.39 (d, $J = 7$ Hz, 3H), 1.15 - 1.09 (m, 6H). LC-MS (ESI): m/z : $[M+H]^+ = 453.2$.

Example 9. Propan-2-yl (2R)-2-({[(1S)-1-[3-fluoro-4-

(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate



[01367] **Step 1: (S)-N-[(1S)-1-[3-Fluoro-4-(trifluoromethyl)phenyl]ethyl]-2-methylpropane-2-sulphinamide.** General procedure A was followed using 1-[3-fluoro-4-(trifluoromethyl)phenyl]ethan-1-one. The crude product was purified by column chromatography (SiO_2 , pet. ether: EtoAc, 1:1) to give (S)-N-[(1S)-1-[3-fluoro-4-(trifluoromethyl)phenyl]ethyl]-2-methylpropane-2-sulphinamide as a yellow oil (Y = 99 %). 1H NMR (400 MHz, $CDCl_3$) δ ppm

7.62 - 7.57 (m, 1H), 7.26 - 7.19 (m, 2H), 4.61 - 4.56 (m, 1H), 3.47 (d, $J = 3$ Hz, 1H), 1.55 - 1.52 (m, 3H), 1.25 (s, 9H).

[01368] Step 2: (1S)-1-[3-Fluoro-4-(trifluoromethyl)phenyl]ethan-1-aminium chloride.


General procedure B was followed using (S)-N-[(1S)-1-[3-fluoro-4-(trifluoromethyl)phenyl]ethyl]-2-methylpropane-2-sulphinamide to give (1S)-1-[3-fluoro-4-(trifluoromethyl)phenyl]ethan-1-aminium chloride as a white solid (Y = quantitative). ^1H NMR (400 MHz, CDCl_3) δ ppm 8.79 (s, 3H), 7.66 - 7.63 (m, 1H), 7.47 (d, $J = 11$ Hz, 1H), 7.39 (d, $J = 8$ Hz, 1H), 4.51 - 4.40 (m, 1H), 1.75 - 1.65 (m, 3H).

[01369] Step 3: 2-Fluoro-4-[(1S)-1-isocyanatoethyl]-1-(trifluoromethyl)benzene. General procedure C was followed using (1S)-1-[3-fluoro-4-(trifluoromethyl)phenyl]ethan-1-aminium chloride to give 2-fluoro-4-[(1S)-1-isocyanatoethyl]-1-(trifluoromethyl)benzene as a white solid (Y = quantitative).

[01370] Step 4: Propan-2-yl (2R)-2-({[(1S)-1-[3-fluoro-4-

(trifluoromethyl)phenyl]ethyl}carbonyl}oxy)-3-(1H-1,2,4-triazol-1-yl). General procedure D was followed using isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate and 2-fluoro-4-[(1S)-1-isocyanatoethyl]-1-(trifluoromethyl)benzene. The crude product was purified by prep-HPLC (column: Phenomenex Luna C18, 10 μm , 100 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 35 - 60 %, 12 min) to give propan-2-yl (2R)-2-({[(1S)-1-[3-fluoro-4-(trifluoromethyl)phenyl]ethyl}carbonyl}oxy)-3-(1H-1,2,4-triazol-1-yl) as a white solid (Y = 8 %). ^1H NMR (400 MHz, CDCl_3) δ ppm 8.34 (s, 1H), 8.04 (s, 1H), 7.58 (t, $J = 7$ Hz, 1H), 7.23 - 7.10 (m, 2H), 5.38 (s, 2H), 5.13 - 5.01 (m, 1H), 4.81 (s, 1H), 4.69 (s, 2H), 1.50 (d, $J = 6$ Hz, 3H), 1.30 - 1.15 (m, 6H). ^1H NMR (400 MHz, MeOD) δ ppm 8.53 (s, 1H), 7.99 (s, 1H), 7.62 (t, $J = 8$ Hz, 1H), 7.36 - 7.23 (m, 2H), 5.27 (t, $J = 5$ Hz, 1H), 4.98 - 4.95 (m, 1H), 4.76 - 4.72 (m, 2H), 1.44 (d, $J = 7$ Hz, 3H), 1.26 - 1.21 (m, 6H). LC-MS (ESI): m/z : $[\text{M}+\text{H}]^+ = 433.1$.

Example 10. Propan-2-yl (2R)-2-({[(1S)-1-(4-cyclopropylphenyl)ethyl]carbonyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate



[01371] **Step 1: 1-(4-Cyclopropylphenyl)ethan-1-one.** A mixture of 1-(4-bromophenyl)ethan-1-one (2.5 g, 12.56 mmol, 1 eq.), cyclopropylboronic acid (1.62 g, 18.84 mmol, 1.5 eq.), K_3PO_4 (8.00 g, 37.68 mmol, 3 eq.) and $Pd(dppf)Cl_2$ (919 mg, 1.26 mmol, 0.1 eq.) in DMSO (25 ml) and H_2O (8 ml) was heated at 100° C under N_2 for 4 h. The reaction was filtered and the filtrate concentrated *in vacuo*. Prep-HPLC (column: Agela Durashell, 10 μm , 250 x 50 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 20 - 50 %, 20 min) gave 1-(4-cyclopropylphenyl)ethan-1-one as a brown solid (Y = 25 %). 1H NMR (400 MHz, $CDCl_3$) δ ppm 7.85 (d, J = 8 Hz, 2H), 7.12 (d, J = 8 Hz, 2H), 2.57 (s, 3H), 2.01 - 1.89 (m, 1H), 1.10 - 1.03 (m, 2H), 0.82 - 0.75 (m, 2H).


[01372] **Step 2: (S)-N-[(1S)-1-(4-Cyclopropylphenyl)ethyl]-2-methylpropane-2-sulphinamide.** General procedure A was followed using 1-(4-cyclopropylphenyl)ethan-1-one. The crude product was purified by FCC (SiO_2 , pet. ether: EtOAc, 5:1 to 1:1) to give (S)-N-[(1S)-1-(4-cyclopropylphenyl)ethyl]-2-methylpropane-2-sulphinamide as a white solid (Y = 31 %). 1H NMR (400 MHz, $CDCl_3$) δ ppm 7.24 (d, J = 8 Hz, 2H), 7.05 (d, J = 8 Hz, 2H), 4.54 - 4.49 (m, 1H), 3.38 (s, 1H), 1.90 - 1.88 (m, 1H), 1.49 (d, J = 6 Hz, 3H), 1.23 (s, 9H), 0.99 - 0.94 (m, 2H), 0.70 - 0.68 (m, 2H).

[01373] **Step 3: (1S)-1-(4-Cyclopropylphenyl)ethan-1-aminium chloride.** General procedure B was followed using (S)-N-[(1S)-1-(4-cyclopropylphenyl)ethyl]-2-methylpropane-2-sulphinamide to give (1S)-1-(4-cyclopropylphenyl)ethan-1-aminium chloride as a white solid (Y = 43 %). 1H NMR (400 MHz, $CDCl_3$) δ ppm 8.60 (s, 3H), 7.35 (d, J = 8 Hz, 2H), 7.06 (d, J = 7 Hz, 2H), 4.32 (s, 1H), 1.89 - 1.85 (m, 1H), 1.63 (d, J = 5 Hz, 3H), 1.00 - 0.96 (m, 2H), 0.69 - 0.67 (m, 2H).

[01374] **Step 4: 1-Cyclopropyl-4-[(1S)-1-isocyanatoethyl]benzene.** General procedure C was followed using (1S)-1-(4-cyclopropylphenyl)ethan-1-aminium chloride to give 1-cyclopropyl-4-[(1S)-1-isocyanatoethyl]benzene (Y = quantitative).

[01375] **Step 5: Propan-2-yl (2R)-2-({[(1S)-1-(4-cyclopropylphenyl)ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.** General procedure D was followed using isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate and 1-cyclopropyl-4-[(1S)-1-isocyanatoethyl]benzene. The crude product was purified by prep-TLC (SiO₂, 100 % EtOAc) to give propan-2-yl (2R)-2-({[(1S)-1-(4-cyclopropylphenyl)ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate as a white solid (Y = 14 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.10 (s, 1H), 7.95 (s, 1H), 7.18 (d, *J* = 8 Hz, 2H), 7.04 (d, *J* = 8 Hz, 2H), 5.36 (t, *J* = 5 Hz, 1H), 5.21 (d, *J* = 7 Hz, 1H), 5.10 - 5.03 (m, 1H), 4.80 - 4.72 (m, 1H), 4.64 (d, *J* = 4 Hz, 2H), 1.90 - 1.85 (m, 1H), 1.46 (d, *J* = 7 Hz, 3H), 1.23 - 1.18 (m, 6H), 0.95 (d, *J* = 7 Hz, 2H), 0.67 (d, *J* = 4 Hz, 2H). LC-MS (ESI): *m/z*: [M+H]⁺ = 387.1

Example 11. Propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoate



[01376] **Step 1: (S)-2-Methyl-N-[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]propane-2-sulphinamide.** General procedure A was followed using 1-[4-(trifluoromethyl)phenyl]ethan-1-one to give (S)-2-methyl-N-[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]propane-2-sulphinamide as a white solid (Y = 65 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.62 (d, *J* = 8 Hz, 2H), 7.48 (d, *J* = 8 Hz, 2H), 4.65 - 4.57 (m, 1H), 3.44 (d, *J* = 2 Hz, 1H), 1.53 (d, *J* = 6 Hz, 3H), 1.25 - 1.24 (m, 9H).


[01377] **Step 2: (1S)-1-[4-(Trifluoromethyl)phenyl]ethan-1-aminium chloride.** General procedure B was following using (S)-2-methyl-N-[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]propane-2-sulphinamide to give (1S)-1-[4-

(trifluoromethyl)phenyl]ethan-1-aminium chloride as a yellow solid (Y = 78 %). ^1H NMR (400 MHz, CDCl_3) δ ppm 8.75 (s, 3H), 7.66 - 7.63 (m, 4H), 4.44 (s, 1H), 1.68 (d, J = 6 Hz, 3H).

[01378] **Step 3: 1-(1-Isocyanatoethyl)-4-(trifluoromethyl)benzene.** General procedure C was followed using (1S)-1-[4-(trifluoromethyl)phenyl]ethan-1-aminium chloride give 1-(1-isocyanatoethyl)-4-(trifluoromethyl)benzene as yellow oil (Y = quantitative). LC-MS (ESI): m/z : $[\text{M}+\text{H}]^+ = 248.0$

[01379] **Step 4: Propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoate.** General procedure D was followed using isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate and (S)-1-(1-isocyanatoethyl)-4-(trifluoromethyl)benzene. Prep-HPLC (column: Phenomenex Synergi C18, 4 μm , 100 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 25 - 55 %, 10 min) gave propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoate as a white solid (Y = 3 %). ^1H NMR (400 MHz, CDCl_3) δ ppm 8.11 (s, 1H), 7.96 (s, 1H), 7.60 (d, J = 8 Hz, 2H), 7.42 (d, J = 8 Hz, 2H), 5.36 (t, J = 5 Hz, 1H), 5.29 (d, J = 7 Hz, 1H), 5.08 - 5.02 (m, 1H), 4.88 - 4.80 (m, 1H), 4.65 (d, J = 5 Hz, 2H), 1.49 (d, J = 7 Hz, 3H), 1.19 (dd, J = 6, 17 Hz, 6H). LC-MS (ESI): m/z : $[\text{M}+\text{H}]^+ = 415.2$.

Example 12. Propan-2-yl (2R)-2-({[(1S)-1-(4-ethylphenyl)ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate



[01380] **Step 1: (S)-N-[(1S)-1-(4-Ethylphenyl)ethyl]-2-methylpropane-2-sulphinamide.**

General procedure A was followed using 1-(4-ethylphenyl)ethan-1-one. Purification by FCC (SiO_2 , pet. ether: EtOAc, 1:0 to 1:1) gave (S)-N-[(1S)-1-(4-ethylphenyl)ethyl]-2-methylpropane-2-sulphinamide as a white solid (Y = 57 %). ^1H NMR (400 MHz, CDCl_3) δ ppm 7.26 (d, J = 8


Hz, 2H), 7.18 (d, $J = 8$ Hz, 2H), 4.53 - 4.51 (m, 1H), 3.39 (s, 1H), 2.67 - 2.61 (m, 2H), 1.50 (d, $J = 6$ Hz, 3H), 1.25 - 1.21 (m, 12H).

[01381] **Step 2: (1S)-1-(4-Ethylphenyl)ethan-1-aminium chloride.** General procedure B was followed using (S)-N-[(1S)-1-(4-ethylphenyl)ethyl]-2-methylpropane-2-sulphinamide to give (1S)-1-(4-ethylphenyl)ethan-1-aminium chloride as a white solid (Y = 76 %). ^1H NMR (400 MHz, CDCl_3) δ ppm 8.55 (s, 3H), 7.38 (d, $J = 8$ Hz, 2H), 7.18 (d, $J = 8$ Hz, 2H), 4.33 - 4.30 (m, 1H), 2.65 - 2.59 (m, 2H), 1.62 (d, $J = 2$ Hz, 3H), 1.20 - 1.19 (m, 3H).

[01382] **Step 3: 1-Ethyl-4-[(1S)-1-isocyanatoethyl]benzene.** General procedure C was followed using (1S)-1-(4-ethylphenyl)ethan-1-aminium chloride to give 1-ethyl-4-[(1S)-1-isocyanatoethyl]benzene as a yellow oil (Y = quantitative). LC-MS (ESI): m/z : $[\text{M}+\text{MeOH}+\text{H}] = 208.1$

[01383] **Step 4: Propan-2-yl (2R)-2-([(1S)-1-(4-ethylphenyl)ethyl]carbamoyl)oxy-3-(1H-1,2,4-triazol-1-yl)propanoate.** General procedure D was followed using 1-ethyl-4-[(1S)-1-isocyanatoethyl]benzene and propan-2-yl (2R)-2-hydroxy-3-(1H-1,2,4-triazol-1-yl)propanoate. Prep-HPLC (column: Phenomenex Synergi C18, 4 μm , 100 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 30 - 60 %, 10 min) to give propan-2-yl (2R)-2-([(1S)-1-(4-ethylphenyl)ethyl]carbamoyl)oxy-3-(1H-1,2,4-triazol-1-yl)propanoate as a white solid (Y = 4 %). ^1H NMR (400 MHz, CDCl_3) δ ppm 8.09 (s, 1H), 7.94 (s, 1H), 7.22 - 7.16 (m, 4H), 5.37 - 5.34 (m, 1H), 5.30 (d, $J = 7$ Hz, 1H), 5.07 - 5.04 (m, 1H), 4.80 - 4.73 (m, 1H), 4.64 (d, $J = 5$ Hz, 2H), 2.65 - 2.59 (m, 2H), 1.48 (d, $J = 7$ Hz, 3H), 1.22 (m, 9H). LC-MS (ESI): m/z : $[\text{M}+\text{H}]^+ = 375.2$

Example 13. Propan-2-yl (2R)-2-([(1S)-1-[4-(propan-2-yl)phenyl]ethyl]carbamoyl)oxy-3-(1H-1,2,4-triazol-1-yl)propanoate




[01384] **Step 1: (S)-2-Methyl-N-[(1S)-1-[4-(propan-2-yl)phenyl]ethyl]propane-2-sulphinamide.** General procedure A was followed using 1-[4-(propan-2-yl)phenyl]ethan-1-one to give (S)-2-methyl-N-[(1S)-1-[4-(propan-2-yl)phenyl]ethyl]propane-2-sulphinamide (Y = 53 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.28 (d, *J* = 8 Hz, 2H), 7.21 (d, *J* = 8 Hz, 2H), 4.57 - 4.50 (m, 1H), 3.43 - 3.40 (m, 1H), 2.95 - 2.80 (m, 1H), 1.49 (d, *J* = 7 Hz, 3H), 1.31 - 1.19 (m, 15H).

[01385] **Step 2: (1S)-1-[4-(Propan-2-yl)phenyl]ethan-1-aminium chloride.** General procedure B was followed using (S)-N-((S)-1-(4-isopropylphenyl)ethyl)-2-methylpropane-2-sulphinamide to give (1S)-1-[4-(propan-2-yl)phenyl]ethan-1-aminium chloride as a white solid (Y = quantitative).

[01386] **Step 3: 1-[(1S)-1-Isocyanatoethyl]-4-(propan-2-yl)benzene.** General procedure C was followed using (1S)-1-(4-isopropylphenyl)ethanamine to give 1-[(1S)-1-isocyanatoethyl]-4-(propan-2-yl)benzene as a white solid (Y = 86 %). LC-MS (ESI): *m/z*: [M+CH₃OH+H]⁺ = 222.2

[01387] **Step 4: Propan-2-yl (2R)-2-([(1S)-1-[4-(propan-2-yl)phenyl]ethyl]carbamoyl)oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.** General procedure D was followed using propan-2-yl (2R)-2-hydroxy-3-(1H-1,2,4-triazol-1-yl)propanoate and 1-[(1S)-1-isocyanatoethyl]-4-(propan-2-yl)benzene. Prep-TLC (EtOAc) gave propan-2-yl (2R)-2-([(1S)-1-[4-(propan-2-yl)phenyl]ethyl]carbamoyl)oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate as a white solid (Y = 14 %). ¹H NMR (400 MHz, MeOD) δ ppm 8.52 (s, 1H), 7.99 (s, 1H), 7.25 (d, *J* = 8 Hz, 2H), 7.19 (d, *J* = 8 Hz, 2H), 5.26 (t, *J* = 5 Hz, 1H), 4.99 - 4.91 (m, 1H), 4.74 - 4.70 (m, 2H), 4.70 - 4.63 (m, 1H), 2.97 - 2.80 (m, 1H), 1.42 (d, *J* = 7 Hz, 3H), 1.24 (d, *J* = 7 Hz, 6H), 1.15 (d, *J* = 6 Hz, 3H), 1.07 (d, *J* = 6 Hz, 3H). LC-MS (ESI): *m/z*: [M+H]⁺ = 389.2

Example 14. Propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-([(1S)-1-[6-(trifluoromethyl)pyridin-3-yl]ethyl]carbamoyl)oxy)propanoate




[01388] **Step 1: (S)-2-Methyl-N-[(1S)-1-[6-(trifluoromethyl)pyridin-3-yl]ethyl]propane-2-sulphinamide.** General procedure A was followed using 1-[6-(trifluoromethyl)pyridin-3-yl]ethan-1-one. Purification by FCC (SiO₂, 0 – 100 % EtOAc in pet. ether) gave (S)-2-methyl-N-[(1S)-1-[6-(trifluoromethyl)pyridin-3-yl]ethyl]propane-2-sulphinamide as a white solid (Y = 39 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.73 (s, 1H), 7.90 (d, *J* = 8 Hz, 1H), 7.70 (d, *J* = 8 Hz, 1H), 4.75 - 4.63 (m, 1H), 3.51 - 3.45 (m, 1H), 1.60 (d, *J* = 7 Hz, 1H), 1.26 (s, 9H).

[01389] **Step 2: (1S)-1-[6-(Trifluoromethyl)pyridin-3-yl]ethan-1-aminium chloride.** General procedure B was followed using (S)-2-methyl-N-[(1S)-1-[6-(trifluoromethyl)pyridin-3-yl]ethyl]propane-2-sulphinamide to give (1S)-1-[6-(trifluoromethyl)pyridin-3-yl]ethan-1-aminium chloride as a white solid (Y = quantitative).

[01390] **Step 3: 5-[(1S)-1-Isocyanatoethyl]-2-(trifluoromethyl)pyridine.** General procedure C was followed using (1S)-1-[6-(trifluoromethyl)pyridin-3-yl]ethan-1-aminium chloride to give 5-[(1S)-1-isocyanatoethyl]-2-(trifluoromethyl)pyridine as a white solid (Y = quantitative). LC-MS in MeOH (ESI): *m/z*: [M+MeOH+H]⁺ = 249.2

[01391] **Step 4: (R)-Isopropyl 3-(1H-1,2,4-triazol-1-yl)-2-(((S)-1-(6-(trifluoromethyl)pyridin-3-yl)ethyl)carbamoyl)oxy)propanoate.** General procedure D was followed using propan-2-yl (2R)-2-hydroxy-3-(1H-1,2,4-triazol-1-yl)propanoate and 5-[(1S)-1-isocyanatoethyl]-2-(trifluoromethyl)pyridine. Prep-TLC (SiO₂, 100 % EtOAc) gave (R)-isopropyl 3-(1H-1,2,4-triazol-1-yl)-2-(((S)-1-(6-(trifluoromethyl)pyridin-3-yl)ethyl)carbamoyl)oxy)propanoate as a white solid (Y = 7 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.70 (s, 1H), 8.53 (s, 1H), 8.25 (d, *J* = 8 Hz, 1H), 8.02 - 7.98 (br. s, 1H), 7.98 (s, 1H), 7.88 (d, *J* = 8 Hz, 1H), 5.19 (t, *J* = 5 Hz, 1H), 4.86 - 4.73 (m, 2H), 4.67 - 4.62 (m, 2H), 1.40 (d, *J* = 7 Hz, 3H), 1.06 (d, *J* = 6 Hz, 3H), 0.94 (d, *J* = 6 Hz, 3H). LC-MS (ESI): *m/z*: [M+H]⁺ = 416.2.

Example 15. Propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-([(1R)-1-[6-(trifluoromethyl)pyridin-3-yl]ethyl]carbamoyl}oxy)propanoate




[01392] **Step 1: (R)-2-Methyl-N-([(1R)-1-[6-[trifluoromethyl]pyridin-3-yl]ethyl]propane-2-sulphinamide.** General procedure A was followed using 1-[6-(trifluoromethyl)pyridin-3-yl]ethan-1-one and (R)-2-methylpropane-2-sulphinamide. Purification by FCC (SiO₂, 0 – 100 % EtOAc in pet. ether) gave (R)-2-methyl-N-([(1R)-1-[6-[trifluoromethyl]pyridin-3-yl]ethyl]propane-2-sulphinamide as a white solid (Y = 64 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.74 (d, *J* = 2 Hz, 1H), 7.90 (dd, *J*₁ = 2 Hz, *J*₂ = 8 Hz, 1H), 7.69 (d, *J* = 8 Hz, 1H), 4.72 - 4.65 (m, 1H), 3.50 - 3.44 (br. s, 1H), 1.60 (d, *J* = 8 Hz, 3H), 1.26 (s, 9H).

[01393] **Step 2: (1R)-1-[6-(Trifluoromethyl)pyridin-3-yl]ethan-1-aminium chloride.** General procedure B was followed using (R)-2-methyl-N-([(1R)-1-[6-[trifluoromethyl]pyridin-3-yl]ethyl]propane-2-sulphinamide to give (1R)-1-[6-(trifluoromethyl)pyridin-3-yl]ethan-1-aminium chloride as a white solid (Y = quantitative). ¹H NMR (400 MHz, D₂O) δ ppm 8.66 (d, *J* = 2 Hz, 1H), 8.10 (dd, *J*₁ = 2 Hz, *J*₂ = 8 Hz, 1H), 7.87 (d, *J* = 8 Hz, 1H), 4.69 - 4.61 (m, 1H), 1.60 (d, *J* = 8 Hz, 3H).

[01394] **Step 3: 5-[(1R)-1-Isocyanatoethyl]-2-(trifluoromethyl)pyridine.** General procedure C was followed using (1R)-1-[6-(trifluoromethyl)pyridin-3-yl]ethan-1-aminium chloride to give 5-[(1R)-1-isocyanatoethyl]-2-(trifluoromethyl)pyridine as a white solid (Y = quantitative). LC-MS in MeOH (ESI): *m/z*: [M+MeOH+H]⁺ = 249.1.

[01395] **Step 4: (R)-Isopropyl 3-(1H-1,2,4-triazol-1-yl)-2-(((R)-1-(6-(trifluoromethyl)pyridin-3-yl)ethyl)carbamoyl)oxy)propanoate.** General procedure D was followed using isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate and 5-[(1R)-1-isocyanatoethyl]-2-(trifluoromethyl)pyridine. Prep-TLC (SiO₂, EtOAc) gave (R)-isopropyl 3-(1H-1,2,4-triazol-1-yl)-2-(((R)-1-(6-(trifluoromethyl)pyridin-3-yl)ethyl)carbamoyl)oxy)propanoate as a white solid (Y = 17%). ¹H NMR (400 MHz, ACN-*d*₃) δ ppm 8.67 (s, 1H), 8.23 (s, 1H), 7.87 (d, *J* = 8 Hz, 1H), 7.83 (s, 1H), 7.75 (d, *J* = 8 Hz, 1H), 6.47 (d, *J* = 7 Hz, 1H), 5.15 (t, *J* = 5 Hz, 1H), 5.04 - 4.96 (m, 1H), 4.84 - 4.76 (m, 1H), 4.60 (d, *J* = 5 Hz, 2H), 1.46 (d, *J* = 4 Hz, 3H), 1.25 - 1.17 (m, 6H). LC-MS (ESI): *m/z*: [M+H]⁺ = 416.2.

Example 16. Isopropyl 2-[1-[2-chloro-4-(trifluoromethyl)phenyl]ethylcarbamoyloxy]-3-pyrimidin-2-yl-propanoate



Step 1: 1-[2-Chloro-4-(trifluoromethyl) phenyl] ethylidene}hydroxylamine.

[01396] To a solution of 1-[2-chloro-4-(trifluoromethyl)phenyl]ethan-1-one (500 mg, 2.25 mmol) in EtOH (5 ml) and H₂O (1 ml) were added NH₂OH.HCl (234 mg, 3.37 mmol) and NaOAc (295 mg, 3.59 mmol) at 25° C. The RM was stirred at 80° C for 2 h. The solvent was removed *in vacuo*. The residue was dissolved in water and extracted three times with EtOAc. The organic phase was dried (Na₂SO₄) and concentrated *in vacuo* to give 1-[2-chloro-4-(trifluoromethyl) phenyl] ethylidene}hydroxylamine as a yellow solid (Y = 94 %). ¹H NMR (400 MHz, DMSO-*d*₆) (Mixture of cis and trans) δ ppm 11.55 (s, 1H), 10.82 (s, 1H), 7.93 (d, *J* = 8 Hz, 2H), 7.76 (d, *J* = 8 Hz, 2H), 7.60 (d, *J* = 8 Hz, 1H), 7.49 (d, *J* = 8 Hz, 1H), 2.13 (s, 3H), 2.09 (s, 3H).


Step 2: 1-[2-Chloro-4-(trifluoromethyl) phenyl] ethanamine.

[01397] To a solution of 1-[2-chloro-4-(trifluoromethyl)phenyl] ethylidene}hydroxylamine (200 mg of cis and trans mixture, 841 μ mol) in MeOH (10 ml) was added Raney nickel (200 mg, in water) at 25° C. The RM was stirred at 25° C for 12 h under H₂ (50 psi). The solution was filtered and the filtrate concentrated *in vacuo* to give 1-[2-chloro-4-(trifluoromethyl) phenyl] ethanamine as a white solid (Y = 80 %). LC-MS (ESI): m/z: [M+H]⁺ = 224.0.

[01398] **Step 3: 2-Chloro-1-(1-isocyanatoethyl)-4-(trifluoromethyl) benzene.** General procedure C was followed using 1-[2-chloro-4-(trifluoromethyl)phenyl]ethanamine to give 2-chloro-1-(1-isocyanatoethyl)-4-(trifluoromethyl) benzene as yellow oil (Y = quantitative). LC-MS (ESI): m/z: [M+H]⁺ = 282.1

[01399] **Step 4: Isopropyl 2-[1-[2-chloro-4-(trifluoromethyl)phenyl]ethylcarbamoyloxy]-3-pyrimidin-2-yl-propanoate.** General procedure D was followed using isopropyl 2-hydroxy-3-pyrimidin-2-yl-propanoate and 2-chloro-1-(1-isocyanatoethyl)-4-(trifluoromethyl)benzene. Prep-HPLC (column: Welch Ultimate AQ-C18, 5 μ m, 150 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 45 - 75 %, 12 min). gave isopropyl 2-[1-[2-chloro-4-(trifluoromethyl)phenyl]ethylcarbamoyloxy]-3-pyrimidin-2-yl-propanoate as a white solid (Y = 6 %). ¹H NMR (400 MHz, MeOD) δ ppm 8.78 – 8.72 (m, 2H), 7.71 - 7.67 (m, 1H), 7.64 - 7.58 (m, 1H), 7.53 - 7.49 (m, 1H), 7.43 - 7.36 (m, 1H), 5.54 - 5.47 (m, 1H), 5.17 - 4.94 (m, 2H), 3.51 - 3.39 (m, 2H), 1.44 - 1.36 (m, 3H), 1.27 - 1.18 (m, 4H), 1.08 (d, *J* = 8 Hz, 2H). LC-MS (ESI): m/z: [M+H]⁺ = 460.1

Example 17. Propan-2-yl (2R)-2-({[(1S)-1-[4-(benzyloxy)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate



[01400] **Step 1: (S)-N-[(1S)-1-[4-(Benzyloxy)phenyl]ethyl]-2-methylpropane-2-sulfonamide.** General procedure A was followed using 1-(4-benzyloxyphenyl)ethenone. Purification by FCC


(SiO₂, pet. ether: EtOAc, 5:1 to 1:1) gave (S)-N-[(1S)-1-[4-(benzyloxy)phenyl]ethyl]-2-methylpropane-2-sulfinamide as a white solid (Y = 39 %). ¹H NMR (400 MHz, MeOD) δ ppm 7.43 - 7.40 (m, 2H), 7.37 - 7.33 (m, 3H), 7.31 - 7.28 (m, 2H), 6.98 - 6.94 (m, 2H), 5.06 (s, 2H), 4.44 - 4.39 (m, 1H), 1.48 (d, *J* = 7 Hz, 3H), 1.21 (s, 9H).

[01401] **Step 2: [4-(Benzyloxy)phenyl]ethan-1-aminium chloride.** General procedure B was followed using (S)-N-((S)-1-(4-(benzyloxy)phenyl)ethyl)-2-methylpropane-2-sulfinamide to give [4-(benzyloxy)phenyl]ethan-1-aminium chloride as a light yellow solid. Y = 94 %. ¹H NMR (400 MHz, MeOD) δ ppm 7.43 - 7.42 (m, 2H), 7.40 - 7.34 (m, 4H), 7.32 - 7.30 (m, 1H), 7.09 - 7.05 (m, 2H), 5.12 (s, 2H), 4.43 - 4.38 (m, 1H), 1.61 (d, *J* = 7 Hz, 3H).

[01402] **Step 3: 1-(Benzyloxy)-4-[(1S)-1-isocyanatoethyl]benzene.** General procedure C was followed using (1S)-1-(4-benzyloxyphenyl)ethanamine to give 1-(benzyloxy)-4-[(1S)-1-isocyanatoethyl]benzene as a yellow oil. Y = quantitative.

[01403] **Step 4: Propan-2-yl (2R)-2-({[(1S)-1-[4-(benzyloxy)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.** General procedure D was followed using isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate and 1-benzyloxy-4-[(1S)-1-isocyanatoethyl]benzene. Purification by prep-TLC (SiO₂, pet. ether: EtOAc, 1:1) gave propan-2-yl (2R)-2-({[(1S)-1-[4-(benzyloxy)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate as a white solid (Y = 12 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.09 (s, 1H), 7.95 (s, 1H), 7.44 - 7.31 (m, 5H), 7.23 (d, *J* = 9 Hz, 2H), 6.94 (d, *J* = 9 Hz, 2H), 5.36 (t, *J* = 6 Hz, 1H), 5.17 (d, *J* = 8 Hz, 1H), 5.11 - 5.02 (m, 3H), 4.76 - 4.74 (m, 1H), 4.64 (d, *J* = 5 Hz, 2H), 1.47 (d, *J* = 7 Hz, 3H), 1.25 - 1.16 (m, 6H). LCMS (ESI): *m/z*: [M+H]⁺ = 453.2.

Example 18. Propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-[(1-[5-(trifluoromethyl)pyrimidin-2-yl]ethyl]carbamoyl]oxy]propanoate



[01404] **Step 1: 2-(1-Ethoxyethenyl)-5-(trifluoromethyl)pyrimidine.** To a degassed solution of 2-chloro-5-(trifluoromethyl)pyrimidine (660 mg, 3.62 mmol) and tributyl(1-ethoxyvinyl)stannane (1.57 g, 4.34 mmol, 1.46 ml) in DMF (6 ml) under N₂ was added PdCl₂(PPh₃)₂ (254 mg, 361 μmol). The reaction was stirred at 100° C for 3 h. The solution was cooled, quenched with water (6 ml) and extracted into EtOAc (3 x 3 ml). The organic phase was concentrated *in vacuo* to give 2-(1-ethoxyethenyl)-5-(trifluoromethyl)pyrimidine as a red oil (700 mg, quantitative). LC-MS (ESI): m/z: [M+H]⁺ = 219.0.

[01405] **Step 2: 1-[5-(Trifluoromethyl)pyrimidin-2-yl]ethan-1-one.** A solution of 2-(1-ethoxyvinyl)-5-(trifluoromethyl)pyrimidine (700 mg, 3.21 mmol) in 2 M HCl (3 ml) and THF (3 ml) was stirred at 25° C for 1 hr. The solution was filtered, quenched with water (3 ml) and extracted into DCM (3 x 2 ml). The yellow solution of 1-[5-(trifluoromethyl)pyrimidin-2-yl]ethan-1-one in DCM was used directly in the next step. LC-MS (ESI): m/z: [M+H]⁺ = 191.0.

[01406] **Step 3: (Diphenylmethyl)({1-[5-(trifluoromethyl)pyrimidin-2-yl]ethyl})amine.** To a solution of 1-[5-(trifluoromethyl)pyrimidin-2-yl]ethan-1-one (300 mg, 1.58 mmol) in DCM (3 ml) at 0° C was added acetic acid (0.1 ml) and diphenylmethanamine (289 mg, 1.58 mmol, 273 μl). The solution was stirred at 25° C for 30 min. NaBH(OAc)₃ (502 mg, 2.37 mmol) was added and the solution stirred at 25° C for 1 h. The solution was filtered and concentrated *in vacuo*. Purification by prep-HPLC (column: Waters Xbridge Prep OBD C18, 10 μm, 150 x 30 mm; mobile phase: [water (10 mM NH₄HCO₃) - ACN]; B: 50 - 80 %, 11 min) gave (diphenylmethyl)({1-[5-(trifluoromethyl)pyrimidin-2-yl]ethyl})amine as a white solid (Y = 37


%). ^1H NMR (400 MHz, MeOD- d_4) δ 9.07 (s, 2H), 7.38 (d, J = 7 Hz, 2H), 7.32 - 7.26 (m, 4H), 7.25 - 7.18 (m, 3H), 7.17 - 7.11 (m, 1H), 4.70 (s, 1H), 4.03-4.01 (m, 1H), 1.45 (d, J = 7 Hz, 3H).

[01407] **Step 4: 1-[5-(Trifluoromethyl)pyrimidin-2-yl]ethan-1-amine.** A solution of (diphenylmethyl)({1-[5-(trifluoromethyl)pyrimidin-2-yl]ethyl})amine (180 mg, 503 μmol) in TFA (0.2 ml) and 12 M HCl (0.2 ml) was stirred at 80° C for 1 h. The solution was filtered and concentrated *in vacuo* to give 1-[5-(trifluoromethyl)pyrimidin-2-yl]ethan-1-amine as a black oil. (110 mg, quantitative). LC-MS (ESI): m/z : $[\text{M}+\text{H}]^+ = 192.1$.

[01408] **Step 5a: Propan-2-yl (2R)-2-[(4-nitrophenoxy)carbonyl]oxy-3-(1H-1,2,4-triazol-1-yl)propanoate.** To a solution of propan-2-yl (2R)-2-hydroxy-3-(1H-1,2,4-triazol-1-yl)propanoate (500 mg, 2.51 mmol) in THF (5 ml) at 0° C was added TEA (419 μl , 3.01 mmol) and (4-nitrophenyl) carbonochloridate (506 mg, 2.51 mmol). The reaction mixture was stirred at 25° C for 1 h. The reaction mixture was concentrated *in vacuo* to give propan-2-yl (2R)-2-[(4-nitrophenoxy)carbonyl]oxy-3-(1H-1,2,4-triazol-1-yl)propanoate as a yellow oil. (700 mg, quantitative). LC-MS (ESI): m/z : $[\text{M}+\text{H}]^+ = 364.9$.

[01409] **Step 5: Propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-[(1-[5-(trifluoromethyl)pyrimidin-2-yl]ethyl)carbamoyl]oxy]propanoate.** To a solution of propan-2-yl (2R)-2-[(4-nitrophenoxy)carbonyl]oxy-3-(1H-1,2,4-triazol-1-yl)propanoate (366 mg, 1.00 mmol) in DCM (1 ml) at 0° C was added TEA (84 μl , 603 μmol) and 1-[5-(trifluoromethyl)pyrimidin-2-yl]ethan-1-amine (96 mg, 502 μmol). The reaction mixture was stirred at 25° C for 1 h. The solution was filtered and concentrated *in vacuo*. Purification by prep-HPLC (column: Welch Ultimate AQ-C18, 10 μm , 150 x 30 mm; mobile phase: [water (0.1 % TFA) -ACN]; B: 25 – 55 %, 12 min) gave propan-2-yl (2R)-3-(1H-1,2,4-triazol-1-yl)-2-[(1-[5-(trifluoromethyl)pyrimidin-2-yl]ethyl)carbamoyl]oxy]propanoate as a colourless oil. (Y = 27 %). ^1H NMR (400 MHz, MeOD) δ ppm 9.05 (s, 2H), 8.60 (d, J = 2 Hz, 1H), 8.05 (d, J = 8 Hz, 1H), 5.29 - 5.23 (m, 1H), 5.06 - 4.91 (m, 2H), 4.79 - 4.71 (m, 2H), 1.57 - 1.49 (m, 3H), 1.28 - 1.13 (m, 6H). LCMS (ESI): m/z : $[\text{M}+\text{H}]^+ = 417.1$.

Example 19. Propan-2-yl 2-[(1-[2-methoxy-4-(trifluoromethyl)phenyl]ethyl)carbamoyl]oxy-3-(pyrimidin-2-yl)propanoate



[01410] **Step 1: 1-[2-Methoxy-4-(trifluoromethyl)phenyl]ethan-1-ol.** 2-Methoxy-4-(trifluoromethyl)benzaldehyde (700 mg, 3.43 mmol) and 3 M bromo(methyl)magnesium (1.37 ml) were stirred in THF (10 ml) at 0° C for 2 h under N₂. The RM was quenched with water (15 ml) and extracted into EtOAc (15 ml x 3). The organic phase was rinsed (brine, 15 ml), dried (Na₂SO₄) and concentrated *in vacuo*. Purification by FCC (SiO₂, pet. ether: EtOAc, 10:1 - 5:1) gave 1-[2-methoxy-4-(trifluoromethyl)phenyl]ethan-1-ol as a white solid (Y = 79 %). ¹H NMR (400MHz, CDCl₃) δ ppm 7.51 (d, *J* = 8 Hz, 1H), 7.25 (d, *J* = 8 Hz, 1H), 7.08 (s, 1H), 5.17 - 5.15 (m, 1H), 3.92 (s, 3H), 2.40 (s, 1H), 1.51 (d, *J* = 6 Hz, 3H).

[01411] **Step 2: 1-[2-Methoxy-4-(trifluoromethyl)phenyl]ethan-1-one.** To a solution of 1-[2-methoxy-4-(trifluoromethyl)phenyl]ethan-1-ol (600 mg, 2.72 mmol) in DCM (15 ml) was added Dess-Martin periodinane (1.69 ml, 2.31 g, 5.45 mmol) and stirred at 25° C for 2 h. The RM was quenched with 2 M NaHCO₃ (5 ml) and filtered. The filtrate was diluted with water (15 ml) and extracted with DCM (3 x 15 ml). The organic phase was washed (brine, 15 ml), dried (Na₂SO₄) and concentrated *in vacuo*. Purification by FCC (SiO₂, pet. ether: EtOAc, 10:1 - 5:1) gave 1-[2-methoxy-4-(trifluoromethyl)phenyl]ethan-1-one as a white solid (Y = 84 %). ¹H NMR (400MHz, CDCl₃) δ ppm 7.87 (d, *J* = 8 Hz, 1H), 7.36 - 7.34 (m, 1H), 7.27 (s, 1H), 4.06 (s, 3H), 2.71 (s, 3H).


[01412] **Step 3: 1-[2-Methoxy-4-(trifluoromethyl)phenyl]ethan-1-amine.** To a solution of 1-[2-methoxy-4-(trifluoromethyl)phenyl]ethan-1-one (500 mg, 2.29 mmol) in MeOH (3 ml) and ACN (3 ml) was added ammonium acetate (1.77 g, 22.9 mmol) at 25° C. The RM was heated to 65° C under N₂. After 0.5 h at 65° C, NaBH₃CN (360 mg, 5.73 mmol) was added and the RM stirred at 65° C for 12 h under N₂. The RM was filtered and concentrated *in vacuo*. Prep HPLC (column: Phenomenex Luna C18, 10 μm, 250 x 50 mm; mobile phase: [water (0.1 % TFA) -

ACN]; B: 20 – 50 %, 20min) gave 1-[2-methoxy-4-(trifluoromethyl)phenyl]ethan-1-amine as a white solid (Y = 80%). ¹H NMR (400MHz, MeOD) δ ppm 7.55 (d, *J* = 8 Hz, 1H), 7.39 - 7.35 (m, 2H), 4.75 – 4.70 (m, 1H), 3.99 (s, 3H), 1.64 (d, *J* = 7 Hz, 3H). LCMS (ESI): *m/z*: [M+H]⁺ = 220.0.

[01413] **Step 4: 1-(1-Isocyanatoethyl)-2-methoxy-4-(trifluoromethyl)benzene.** General procedure C was followed using 1-[2-methoxy-4-(trifluoromethyl)phenyl]ethan-1-amine to give 1-(1-isocyanatoethyl)-2-methoxy-4-(trifluoromethyl)benzene (200 mg, crude) as a light yellow oil. LC-MS in MeOH (ESI): *m/z*: [M+MeOH+H]⁺ = 278.0.

[01414] **Step 5: Propan-2-yl 2-[(1-[2-methoxy-4-(trifluoromethyl)phenyl]ethyl)carbamoyloxy]-3-(pyrimidin-2-yl)propanoate.** General procedure D was followed using 1-(1-isocyanatoethyl)-2-methoxy-4-(trifluoromethyl)benzene and propan-2-yl 2-hydroxy-3-(pyrimidin-2-yl)propanoate. Purification by prep-HPLC (column: Nano-micro Kromasil C18, 10 μm, 100 x 40 mm; mobile phase: water (0.1 % TFA) - ACN]; B: 55 – 65 %, 10 min) gave propan-2-yl 2-[(1-[2-methoxy-4-(trifluoromethyl)phenyl]ethyl)carbamoyloxy]-3-(pyrimidin-2-yl)propanoate as a colourless gum (Y = 4%). ¹H NMR (400 MHz, ACN-*d*₃) δ ppm 8.80 - 8.66 (m, 2H), 7.41 - 7.09 (m, 4H), 6.15 - 6.13 (m, 1H), 5.44 - 5.39 (m, 1H), 5.02 - 4.88 (m, 2H), 3.92 - 3.81 (m, 3H), 3.43 - 3.29 (m, 2H), 1.33 - 1.29 (m, 3H), 1.44 – 1.21 (m, 5H), 1.16 - 1.04 (d, *J* = 6 Hz, 1H). LCMS (ESI): *m/z*: [M+H]⁺ = 456.2.

Example 20. Isopropyl 2-[1-[2-methyl-4-(trifluoromethyl)phenyl]ethylcarbamoyloxy]-3-pyrimidin-2-yl-propanoate.




[01415] **Step 1: 1-[2-Methyl-4-(trifluoromethyl)phenyl]ethan-1-one.** 1-Bromo-2-methyl-4-(trifluoromethyl)benzene (1.0 g, 4.18 mmol), tributyl(1-ethoxyvinyl)stannane (2.27 g, 6.28 mmol, 2.12 ml) and Pd(PPh₃)₄ (483 mg, 418 μ mol) in toluene (10 ml) were stirred at 120° C under N₂ for 12 h. The RM was cooled and partitioned between EtOAc (10 ml x 3) and water (10 ml). The organic phase was rinsed (brine, 10 ml), dried (Na₂SO₄) and concentrated *in vacuo*. Purification by FCC (SiO₂, 0- 100% EtOAc in pet. ether) gave 1-[2-methyl-4-(trifluoromethyl)phenyl]ethan-1-one as a yellow oil (600 mg). ¹H NMR (400MHz, CDCl₃) δ 7.75 (d, *J* = 8 Hz, 1H), 7.55 - 7.53 (m, 2H), 2.61 (s, 3H), 2.57 (s, 3H).

[01416] **Step 2: 1-[2-Methyl-4-(trifluoromethyl)phenyl]ethan-1-amine.** To a solution of 1-[2-methyl-4-(trifluoromethyl)phenyl]ethan-1-one (400 mg, 1.98 mmol) in MeOH (4 ml) and ACN (4 ml) was added ammonium acetate (1.53 g, 19.8 mmol) at 25°C. The mixture was stirred at 65° C for 0.5 h. NaBH₃CN (311 mg, 4.95 mmol) was added and the RM stirred at 65° C for 12 h under N₂. The RM was filtered and the filtrate concentrated *in vacuo*. Prep-HPLC (column: Welch Xtimate C18, 10 μ m, 250 x 50 mm; mobile phase: water (0.1 % TFA) - ACN]; B: 35 %, 10 min) gave 1-[2-methyl-4-(trifluoromethyl)phenyl]ethan-1-amine as a white solid (Y = 50 %). ¹H NMR (400MHz, CDCl₃) δ ppm 8.26 (s, 2H), 7.54 (d, *J* = 9 Hz, 1H), 7.47 - 7.45 (m, 2H), 4.59 - 4.57 (m, 1H), 2.37 (s, 3H), 1.53 (d, *J* = 7 Hz, 3H). LCMS (ESI): *m/z*: [M+H]⁺ = 204.1.

[01417] **Step 3: 1-(1-Isocyanatoethyl)-2-methyl-4-(trifluoromethyl)benzene.** General procedure C was followed using 1-[2-methyl-4-(trifluoromethyl)phenyl]ethan-1-amine to give 1-(1-isocyanatoethyl)-2-methyl-4-(trifluoromethyl)benzene as a light yellow oil (110 mg, crude). LCMS in MeOH (ESI): *m/z*: [M+MeOH+H]⁺ = 262.0.

[01418] **Step 4: Isopropyl 2-[1-[2-methyl-4-(trifluoromethyl)phenyl]ethylcarbamoyloxy]-3-pyrimidin-2-yl-propanoate.** General procedure D was followed using propan-2-yl 2-hydroxy-3-(pyrimidin-2-yl)propanoate and 1-(1-isocyanatoethyl)-2-methyl-4-(trifluoromethyl)benzene. Purification by prep-HPLC (column: Nano-micro Kromasil C18, 5 μ m, 100 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 48 - 57 %, 10 min) gave isopropyl 2-[1-[2-methyl-4-(trifluoromethyl)phenyl]ethylcarbamoyloxy]-3-pyrimidin-2-yl-propanoate as a colourless oil (Y = 15 %). ¹H NMR (400 MHz, ACN-*d*₃) δ ppm 8.71 (d, *J* = 5 Hz, 1H), 8.65 (d, *J* = 5 Hz, 1H), 7.51 - 7.46 (m, 2H), 7.33 - 7.22 (m, 2H), 6.20 - 6.17 (m, 1H), 5.43 - 5.38 (m, 1H), 5.02 - 4.88 (m, 2H), 3.42 - 3.28 (m, 2H), 2.45 - 2.38 (m, 3H), 1.34 - 1.12 (m, 8H), 1.04 - 1.00 (m, 1H). LCMS (ESI): *m/z*: [M+H]⁺ = 440.2.

Example 21. Propan-2-yl 3-(pyrimidin-2-yl)-2-[(1-[4-(trifluoromethyl)phenyl]butyl}carbamoyl)oxy]propanoate.



[01419] **Step 1: 1-[4-(Trifluoromethyl)phenyl]butan-1-ol.** To a solution of 4-(trifluoromethyl)benzaldehyde (2.0 g, 11.49 mmol, 1.54 ml) in THF (20 ml) was added 1 M bromo(propyl)magnesium (11.49 ml) under N₂ at 0° C and stirred at 15° C for 1 h. The RM was quenched (NH₄Cl, 20 ml), diluted (H₂O, 20 ml), extracted (EtOAc, 15 ml x 3), washed (brine, 20 ml), dried (Na₂SO₄) and concentrated *in vacuo*. Purification by FCC (SiO₂, pet. ether: EtOAc 3:1) gave 1-[4-(trifluoromethyl)phenyl]butan-1-ol as a yellow oil (Y = 48 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.61 (d, *J* = 8 Hz, 2H), 7.47 (d, *J* = 8 Hz, 2H), 4.78 – 4.75 (m, 1H), 1.80 – 1.69 (m, 2H), 1.50 – 1.28 (m, 2H), 0.97 – 0.93 (m, 3H).

[01420] **Step 2: 1-[4-(Trifluoromethyl)phenyl]butan-1-one.** To a solution of 1-[4-(trifluoromethyl)phenyl]butan-1-ol (1.2 g, 5.50 mmol) in DCM (12 ml) was added Dess-Martin periodinane (5.83 g, 13.75 mmol, 4.26 ml). The RM was stirred at 25° C for 3 h. The solution was washed sequentially with 10 % NaOH (10 ml) and Na₂SO₃ (10 ml), dried (Na₂SO₄) filtered and concentrated *in vacuo*. Purification by FCC (SiO₂, pet. ether: EtOAc, 3:1) gave 1-[4-(Trifluoromethyl)phenyl]butan-1-one as a yellow oil. Y = 67 %. ¹H NMR (400 MHz, CDCl₃) δ ppm 8.07 (d, *J* = 8 Hz, 2H), 7.73 (d, *J* = 8 Hz, 2H), 2.98 (t, *J* = 8 Hz, 2H), 1.84 – 1.75 (m, 2H), 1.03 (t, *J* = 7 Hz, 3H).


[01421] **Step 3: 1-[4-(Trifluoromethyl)phenyl]butan-1-amine.** To a solution of 1-[4-(trifluoromethyl)phenyl]butan-1-one (300 mg, 1.39 mmol) in MeOH (5 ml) and ACN (5 ml)

was added ammonium acetate (1.07 g, 13.88 mmol) at 25° C. The RM was heated to 65° C. After 0.5 h, NaBH₃CN (218.00 mg, 3.47 mmol) was added and the RM stirred at 65° C for 12 h. The solution was concentrated *in vacuo*. Purification by prep-HPLC (column: Nano-micro Kromasil C18, 5 µm, 100 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 20 – 35 %, 10 min) gave 1-[4-(trifluoromethyl)phenyl]butan-1-amine as a white solid (Y = 33 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.24 (s, 3H), 7.62 (d, *J* = 8 Hz, 2H), 7.43 (d, *J* = 8 Hz, 2H), 4.07 (t, *J* = 8 Hz, 1H), 1.91 - 1.81 (m, 2H), 1.26 - 1.10 (m, 2H), 0.88 (t, *J* = 6 Hz, 3H).

[01422] **Step 4: 1-(1-Isocyanatobutyl)-4-(trifluoromethyl)benzene.** General procedure C was followed using 1-[4-(trifluoromethyl)phenyl]butan-1-amine to give 1-(1-isocyanatobutyl)-4-(trifluoromethyl)benzene as a yellow solid (Y = 100%). LC-MS in MeOH (ESI): *m/z*: [M+MeOH+H]⁺ = 276.1

[01423] **Step 5: Propan-2-yl 3-(pyrimidin-2-yl)-2-[(1-[4-(trifluoromethyl)phenyl]butyl}carbamoyleoxy]propanoate.** General procedure D was followed using isopropyl 2-hydroxy-3-pyrimidin-2-yl-propanoate and 1-(1-isocyanatobutyl)-4-(trifluoromethyl)benzene. Purification by prep-HPLC (column: Welch Ultimate AQ-C18, 5 µm, 150 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 55 – 70 %, 10 min) gave propan-2-yl 3-(pyrimidin-2-yl)-2-[(1-[4-(trifluoromethyl)phenyl]butyl}carbamoyleoxy]propanoate as a colourless oil (Y = 18 %). ¹H NMR (400 MHz, ACN-*d*₃) δ ppm 8.71 (d, *J* = 8 Hz, 1H), 8.66 (d, *J* = 4 Hz, 1H), 7.75 - 7.60 (m, 2H), 7.46 (d, *J* = 8 Hz, 1H), 7.38 (d, *J* = 8 Hz, 1H), 7.31 - 7.25 (m, 1H), 6.14 (d, *J* = 8 Hz, 1H), 5.46 - 5.37 (m, 1H), 5.03 - 4.87 (m, 1H), 4.61 - 4.57 (m, 1H), 3.45 - 3.28 (m, 2H), 1.70 - 1.60 (m, 2H), 1.37 - 1.17 (m, 5H), 1.14 - 0.97 (m, 3H), 0.95-0.92 (m, 3H). LCMS (ESI): *m/z*: [M + H]⁺ = 454.1.

Example 22. Propan-2-yl 2-[(1-[2-ethyl-4-(trifluoromethyl)phenyl]ethyl}carbamoyleoxy]-3-(pyrimidin-2-yl)propanoate.



[01424] **Step 1: 2-Ethenyl-4-(trifluoromethyl)benzoic acid.** 2-Bromo-4-

(trifluoromethyl)benzoic acid (3 g, 11.15 mmol), potassium vinyltrifluoroborate (14.94 g, 111.52 mmol), PPh₃ (176 mg, 669 μ mol), PdCl₂ (198 mg, 1.12 mmol) and Cs₂CO₃ (10.90 g, 33.46 mmol) in THF (30 ml), H₂O (3 ml) were stirred at 85° C for 12 h under N₂. The RM was cooled, filtered and concentrated *in vacuo* to give a yellow solid (4 g, crude), which was used without purification.

[01425] **Step 2: 2-Ethenyl-N-methoxy-N-methyl-4-(trifluoromethyl)benzamide.** To a solution

of 2-ethenyl-4-(trifluoromethyl)benzoic acid (2.4 g, 11.10 mmol) in DMF (40 ml) was added HATU (6.33 g, 16.65 mmol) and stirred at 0° C for 0.5 h. N-methoxymethanamine hydrochloride (1.62 g, 16.65 mmol) and DIPEA (7.74 ml, 44.41 mmol) were added at 0° C and the RM stirred at 25° C for 3.5 h. The RM was filtered and concentrated *in vacuo*. The residue was diluted with water and extracted with EtOAc (3 x 50 ml). The organic phase was washed (brine, 30 ml), dried (Na₂SO₄) and concentrated *in vacuo*. Purification by FCC (SiO₂, pet. ether: EtOAc, 5:1 to 3:1) gave 2-ethenyl-N-methoxy-N-methyl-4-(trifluoromethyl)benzamide as a yellow oil. (2.4 g, crude). ¹H NMR (400MHz, CDCl₃) δ 7.86-7.85 (m, 1H), 7.70 - 7.62 (m, 1H), 7.46 - 7.39 (m, 1H), 6.83 - 6.81 (m, 1H), 5.85 (d, *J* = 18 Hz, 1H), 5.46 (d, *J* = 11 Hz, 1H), 3.58 - 3.33 (m, 6H).

[01426] **Step 3: 1-[2-Ethenyl-4-(trifluoromethyl)phenyl]ethan-1-one.** To a solution of 2-ethenyl-N-methoxy-N-methyl-4-(trifluoromethyl)benzamide (500 mg, 1.93 mmol) in THF (5 ml) was added 3 M bromo(methyl)magnesium (0.96 ml) at 0° C. The mixture was stirred at 0° C for 2 h under N₂. 2 M HCl (5 ml), and water (15 ml) were added. The mixture was extracted into EtOAc (3 x 15 ml) and the combined organic phases washed (brine), dried (Na₂SO₄) and concentrated *in vacuo*. Purification by FCC (SiO₂, pet. ether: EtOAc, 10:1 to 3:1) gave 1-[2-ethenyl-4-(trifluoromethyl)phenyl]ethan-1-one as a colourless oil (Y = 41 %). LCMS (ESI): m/z: [M+H]⁺ = 215.0.

[01427] **Step 4: 1-[2-Ethenyl-4-(trifluoromethyl)phenyl]ethan-1-amine.** To a solution of 1-[2-ethenyl-4-(trifluoromethyl)phenyl]ethan-1-one (1.1 g, 5.14 mmol) in ACN (8 ml) and MeOH (8 ml) was added ammonium acetate (3.96 g, 51.4 mmol) at 25° C. The RM was stirred at 65° C for 0.5 h. NaBH₃CN (807 mg, 12.84 mmol) was added and the RM stirred at 65° C for 12 h under N₂. Prep HPLC (column: Phenomenex Luna C18, 10 µm, 250 x 50 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 15 -35 %, 10 min) gave 1-[2-ethenyl-4-(trifluoromethyl)phenyl]ethan-1-amine as a white solid. Y = 41 %. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.55-8.35 (br s, 2H), 7.86 - 7.82 (m, 2H), 7.75 - 7.73 (m, 1H), 7.13 (dd, *J* = 17 Hz, 11 Hz, 1H), 5.92 (d, *J* = 17 Hz, 1H), 5.57 (d, *J* = 11 Hz, 1H), 4.80 (d, *J* = 6 Hz, 1H), 1.45 (d, *J* = 6 Hz, 3H). LCMS (ESI): m/z: [M+H]⁺ = 216.0.


[01428] **Step 5: 1-[2-Ethyl-4-(trifluoromethyl)phenyl]ethan-1-amine.** A mixture of 1-[2-ethenyl-4-(trifluoromethyl)phenyl]ethan-1-amine (250 mg, 1.16 mmol) and Raney-Ni (149 mg, 1.74 mmol) in MeOH (15 ml) was stirred at 25° C for 12 h under a balloon of hydrogen. The RM was filtered and the filtrate concentrated *in vacuo* to give 1-[2-ethyl-4-(trifluoromethyl)phenyl]ethan-1-amine as a white solid (Y = 40 %). ¹H NMR (400MHz, DMSO-*d*₆) δ ppm 8.22 (s, 2H), 7.75 - 7.72 (m, 2H), 7.62 (s, 1H), 4.67 - 4.64 (m, 1H), 2.90 - 2.60 (m, 2H), 1.48 (d, *J* = 6 Hz, 3H), 1.24 - 1.13 (m, 3H).

[01429] **Step 6: 2-Ethyl-1-(1-isocyanatoethyl)-4-(trifluoromethyl)benzene.** General procedure C was followed using 1-[2-ethyl-4-(trifluoromethyl)phenyl]ethan-1-amine to give 2-ethyl-1-(1-isocyanatoethyl)-4-(trifluoromethyl)benzene as a light yellow oil. (30 mg, crude). LC-MS in MeOH (ESI): m/z: [M+MeOH+H]⁺ = 276.1.

[01430] **Step 7: Propan-2-yl 2-[(1-[2-ethyl-4-(trifluoromethyl)phenyl]ethyl)carbamoyl]oxy]-3-(pyrimidin-2-yl)propanoate.** General

procedure D was followed using propan-2-yl 2-hydroxy-3-(pyrimidin-2-yl)propanoate and 2-ethyl-1-(1-isocyanatoethyl)-4-(trifluoromethyl)benzene. Purification by prep-HPLC (column: Welch Ultimate AQ-C18 5 μ m, 150 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 45 - 75 %, 12 min) gave propan-2-yl 2-[(1-[2-ethyl-4-(trifluoromethyl)phenyl]ethyl}carbamoxy]oxy]-3-(pyrimidin-2-yl)propanoate as a light yellow solid (Y = 18 %). ^1H NMR (400 MHz, ACN-d_3) δ ppm 8.74 - 8.66 (m, 2H), 7.49 (d, J = 9 Hz, 2H), 7.36 - 7.09 (m, 2H), 6.22 (s, 1H), 5.44 - 5.38 (m, 1H), 5.01 - 4.88 (m, 2H), 3.43 - 3.29 (m, 2H), 2.81 - 2.73 (m, 2H), 1.39 - 1.30 (m, 3H), 1.28 - 1.16 (m, 6H), 1.13 - 1.00 (m, 3H). LCMS (ESI): m/z: $[\text{M}+\text{H}]^+ = 454.2$.

Example 23. Propan-2-yl (2R)-3-(pyrimidin-2-yl)-2-([(1S)-1-[4-(trifluoromethyl)phenyl]ethyl}carbamoxy]oxy)propanoate.




[01431] **Step 1: (9H-Fluorenyl)methyl (2S)-2-([(2R)-1-oxo-1-(propan-2-yloxy)-3-(pyrimidin-2-yl)propan-2-yl]oxy}carbonyl)pyrrolidine-1-carboxylate.** To a solution of propan-2-yl 2-hydroxy-3-(pyrimidin-2-yl)propanoate (5.3 g, 25.21 mmol), (2S)-1-[(9H-fluorenyl)methoxy]carbonylpyrrolidine-2-carboxylic acid (8.51 g, 25.21 mmol) and DCC (5.20 g, 25.21 mmol, 5.10 ml) in DCM (100 ml) was added DMAP (246 mg, 2.02 mmol) at 0 °C. The RM was stirred at 25° C for 12 h, filtered and the resulting yellow solution concentrated *in vacuo* to afford a yellow solid. Purification by FCC (SiO_2 , 16 – 100 % EtOAc in DCM) gave (9H-fluorenyl)methyl (2S)-2-([(2R)-1-oxo-1-(propan-2-yloxy)-3-(pyrimidin-2-yl)propan-2-yl]oxy}carbonyl)pyrrolidine-1-carboxylate as a yellow oil (Y = 26 %).

[01432] **Step 2: (2R)-2-Hydroxy-3-(pyrimidin-2-yl)propanoic acid.** To solution of NaOH (1.52 g, 37.95 mmol) in H₂O (40 ml) and MeOH (40 ml) was added (9H-fluoren-9-yl)methyl (2S)-2-({[(2R)-1-oxo-1-(propan-2-yloxy)-3-(pyrimidin-2-yl)propan-2-yl]oxy}carbonyl)pyrrolidine-1-carboxylate (6.7 g, 12.65 mmol). The RM was stirred at 40° C for 3 h. The RM was cooled, filtered and the resulting yellow liquid concentrated *in vacuo* to afford a yellow oil, which was used without further purification.

[01433] **Step 3: Propan-2-yl (2R)-2-hydroxy-3-(pyrimidin-2-yl)propanoate.** To solution of (2R)-2-Hydroxy-3-(pyrimidin-2-yl)propanoic acid (100 mg, 595 µmol) in isopropanol (1 ml) was added SOCl₂ (43.1 µl 1 mg, 595 µmol). The RM was stirred at 40° C for 1 h. The solution was concentrated *in vacuo* to afford a yellow solid. Purification by prep-TLC (silica, 100 % EtOAc) gave propan-2-yl (2R)-2-hydroxy-3-(pyrimidin-2-yl)propanoate as a colourless gum (Y = 36 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.69 (d, *J* = 5 Hz 2H), 7.20 (t, *J* = 5 Hz, 1H), 5.11 – 5.07 (m, 1H), 4.73 - 4.69 (m, 1H), 4.25 (d, *J* = 6.4 Hz, 2H), 3.53 - 3.39 (m, 2H), 1.26 - 1.19 (m, 6H).


[01434] **Step 4: Propan-2-yl (2R)-3-(pyrimidin-2-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoate.** To a solution of 1-[(1S)-1-isocyanatoethyl]-4-(trifluoromethyl)benzene (47.7 mg, 222 µmol) in THF (1 ml) was added propan-2-yl (2R)-2-hydroxy-3-(pyrimidin-2-yl)propanoate (46.6 mg, 222 µmol) and CuCl (43.9 mg, 443 µmol). The RM was stirred at 0° C for 12 h. The solution was concentrated *in vacuo*. Purification by prep-HPLC (column: Nano-micro Kromasil C18, 3 µm, 80 x 25 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 35- 55 %, 10 min) gave propan-2-yl (2R)-3-(pyrimidin-2-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoate as a white solid (Y = 20 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.80 - 8.48 (m, 2H), 7.62 - 7.53 (m, 2H), 7.46 - 7.38 (m, 2H), 7.37 - 7.28 (m, 1H), 5.63 - 5.59 (m, 1H), 5.28 – 5.02 (m, 2H), 4.87 - 4.76 (m, 1H), 3.58 - 3.40 (m, 2H), 1.46 - 1.40 (m, 3H), 1.30 - 1.21 (m, 4H), 1.14 (d, *J* = 4 Hz, 2H). LCMS (ESI): *m/z*: [M + H]⁺ = 426.2.

Example 24. (2R)-3-(Pyrimidin-2-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoic acid.



[01435] To a solution of propan-2-yl (2R)-3-(pyrimidin-2-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoate (17 mg, 40 μ mol) in dioxane (1 ml) was added 4 M HCl (1 ml) at 25° C. The RM was heated at 40° C for 12 h. The reaction mixture was cooled, filtered and concentrated *in vacuo*. Purification by prep-HPLC (column: Ximate C18 3 μ m, 100 x 30 mm; mobile phase: [water (0.05 % HCl) - ACN]; B: 20 – 50 %, 12 min) gave (2R)-3-(pyrimidin-2-yl)-2-({[(1S)-1-[4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)propanoic acid as a yellow solid (Y= 20 %). ^1H NMR (400 MHz, ACN- d_3) δ ppm 8.97 - 8.55 (m, 2H), 7.65 - 7.25 (m, 5H), 6.70 - 6.62 (m, 1H), 5.46 - 5.43 (m, 1H), 4.75 - 4.68 (m, 1H), 3.68 - 3.47 (m, 2H), 1.32 - 1.28 (m, 3H). LCMS (ESI): m/z : $[\text{M} + \text{H}]^+ = 384.1$.

Example 25. Propan-2-yl (2R)-2-({[(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate.




[01436] **Step 1: 2-Chloro-1-[(1S)-1-isocyanatoethyl]-4-(trifluoromethyl)benzene.** General procedure C was followed using (1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethan-1-amine to give 2-chloro-1-[(1S)-1-isocyanatoethyl]-4-(trifluoromethyl)benzene as a yellow liquid (62 mg, crude). LC-MS in MeOH (ESI): m/z : $[\text{M} + \text{MeOH} + \text{H}]^+ = 282.6$.

[01437] **Step 2: Propan-2-yl (2R)-2-({[(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate.** General procedure D was followed using propan-2-yl (2R)-2-hydroxy-3-(pyrimidin-2-yl)propanoate (synthesised in example 23) and 2-chloro-1-[(1S)-1-isocyanatoethyl]-4-(trifluoromethyl)benzene. Prep-HPLC (column: Nano-micro Kromasil C18, 3 μ m, 80 x 25 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 45 -65 %, 10 min) gave propan-2-yl (2R)-2-({[(1S)-1-[2-chloro-4-


(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate as a white solid (Y = 69 %). ^1H NMR (400 MHz, $\text{ACN-}d_3$) δ ppm 8.80 - 8.36 (m, 2H), 7.71 (s, 1H), 7.67 - 7.44 (m, 2H), 7.35 - 7.30 (m, 1H), 6.38 - 6.25 (m, 1H), 5.47 - 5.34 (m, 1H), 5.08 - 5.00 (m, 1H), 4.97 - 4.82 (m, 1H), 3.47 - 3.34 (m, 2H), 1.34 (d, $J = 7$ Hz, 3H), 1.21 - 0.99 (m, 6H). LCMS (ESI): m/z : $[\text{M}+\text{H}]^+ = 460.1$.

Example 26. (2R)-2-({[(1S)-1-[2-Chloro-4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoic acid.



[01438] To a solution of propan-2-yl (2R)-2-({[(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate (40 mg, 87 μmol) in dioxane (1 ml) was added 4 M HCl (1 ml) and the resultant mixture stirred at 40° C for 12 h. The RM was concentrated *in vacuo* to afford a yellow liquid. Purification by prep-HPLC (column: Xtimate C18 3 μm , 100 x 30 mm; mobile phase: [water (0.05 % HCl) - ACN]; B: 25 – 55 %, 12 min) gave (2R)-2-({[(1S)-1-[2-Chloro-4-(trifluoromethyl)phenyl]ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoic acid as a white solid (Y = 27 %). ^1H NMR (400 MHz, $\text{ACN-}d_3$) δ ppm 8.98 (d, $J = 5$ Hz, 2H), 7.76 - 7.65 (m, 2H), 7.64 (s, 1H), 7.55 - 7.19 (m, 1H), 6.94 (d, $J = 6$ Hz, 1H), 5.53 - 5.38 (m, 1H), 5.10 - 4.99 (m, 1H), 3.73 - 3.67 (m, 1H), 3.60 - 3.54 (m, 1H), 1.37 (d, $J = 7$ Hz, 3H). LCMS (ESI): m/z : $[\text{M}+\text{H}]^+ = 418.1$.

Example 27. Propan-2-yl (2R)-2-({[(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl](methyl)carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate.



[01439] **Step 1: N-[(1S)-1-[2-Chloro-4-(trifluoromethyl)phenyl]ethyl]formamide.** To a solution of (1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethan-1-amine (300 mg, 1.34 mmol) in ethyl formate (3 ml) was added TEA (0.28 ml, 2.01 mmol). The RM was stirred at 50° C for 16 h. The reaction mixture was filtered to afford a yellow solution and the filtrate concentrated *in vacuo* to afford a yellow solid (330 mg, crude).


[01440] **Step 2: [(1S)-1-[2-Chloro-4-(trifluoromethyl)phenyl]ethyl](methyl)amine hydrochloride.** N-[(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl]formamide (300 mg, 1.19 mmol) was dissolved in 1 M BH₃.THF (5 ml) at 0° C and the stirred at 65° C for 12 h. The RM was quenched (water, 4 ml) and 3 M HCl (3 ml) was added. The solution was stirred at 60° C for 30 min, cooled and concentrated *in vacuo*. Purification by prep-HPLC (column: Phenomenex Luna C18, 5 µm, 100 x 30 mm; mobile phase: [water (0.04 % HCl) - ACN]; B: 15 - 30 %, 10 min) gave [(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl](methyl)amine hydrochloride as a white solid (Y = 53 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 10.64 (s, 1H), 10.12 (s, 1H), 8.26 (d, *J* = 8 Hz, 1H), 7.80 - 7.61 (m, 2H), 4.88 (m, 1H), 2.55 (s, 3H), 1.88 (d, *J* = 7 Hz, 3H). LCMS (ESI): *m/z*: [M+H]⁺ = 238.0.

[01441] **Step 3: N-[(1S)-1-[2-Chloro-4-(trifluoromethyl)phenyl]ethyl]-N-methylcarbamoyl chloride.** To a solution of [(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl](methyl)amine (50 mg, 210 µmol) and DIPEA (55 µl, 316 µmol) in THF (2 ml) was added triphosgene (94 mg, 316 µmol) and the resultant mixture stirred at 25° C for 2 h. The reaction mixture was concentrated *in vacuo* to afford a yellow residue. The crude product was suspended in water (2 ml) and extracted (EtOAc, 3 x 2 ml). The combined organic phases were washed (brine, 1 ml), dried (Na₂SO₄) and concentrated *in vacuo* to afford N-[(1S)-1-[2-chloro-4-

(trifluoromethyl)phenyl]ethyl]-N-methylcarbamoyl chloride as a yellow oil (63 mg, crude). LC-MS (ESI): m/z : $[M+H]^+ = 300.0/302.0$.

[01442] **Step 4: Propan-2-yl (2R)-2-([(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl](methyl)carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate.** To a solution of isopropyl (2R)-2-hydroxy-3-pyrimidin-2-yl-propanoate (37.8 mg, 180 μ mol) in dioxane (0.5 ml) at 25° C was added NaH (60 % in mineral oil, 13.6 mg, 340 μ mol). After 30 mins, N-[(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl]-N-methyl-carbamoyl chloride (60 mg, 200 μ mol) in dioxane (0.5 ml) was added and the RM was stirred at 25° C for 2 h. The RM was adjusted to pH 7 to 8 with 0.5 M HCl and the solution concentrated *in vacuo*. Purification by prep-TLC (silica, 100 % EtOAc) gave propan-2-yl (2R)-2-([(1S)-1-[2-chloro-4-(trifluoromethyl)phenyl]ethyl](methyl)carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate as a colourless gum (Y = 52%). ¹H NMR (400 MHz, ACN-*d*₃) δ ppm 8.76 - 8.50 (m, 1H), 7.80 - 7.42 (m, 3H), 7.33 - 7.09 (m, 1H), 5.58 - 5.24 (m, 2H), 5.05 - 4.95 (br. s, 1H), 3.46 - 3.20 (m, 2H), 2.76 - 2.43 (m, 3H), 2.23 - 2.06 (s, 6H), 1.46 (d, $J = 7$ Hz, 3H), 1.28 - 1.11 (m, 6H). LCMS (ESI): m/z : $[M+H]^+ = 474.1$.

Example 28. Propan-2-yl 3-(pyrimidin-2-yl)-2-([(1-[4-(trifluoromethyl)phenyl]propyl}carbamoyl)oxy]propanoate.




[01443] **Step 1: 1-[4-(Trifluoromethyl)phenyl]propan-1-amine.** To a solution of 1-[4-(trifluoromethyl)phenyl]propan-1-one (1 g, 4.95 mmol) in ACN (5 ml) and MeOH (5 ml) was added ammonium acetate (3.81 g, 49.5 mmol) at 65° C for 0.5 h. NaBH₃CN (777 mg, 12.4 mmol) was added and the RM stirred at 65° C for 12 h. The solution was concentrated *in vacuo*. Purification by prep-HPLC (column: Phenomenex Luna C18, 10 μ m, 250 x 80 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 15 - 45 %, 22 min) gave 1-[4-

(trifluoromethyl)phenyl]propan-1-amine as a white solid (Y = 60 %). HNMR (400 MHz, CDCl₃) δ ppm 8.33 (s, 2H), 7.63 (d, J = 8 Hz, 2H), 7.42 (d, J = 8 Hz, 2H), 4.10 - 3.95 (m, 1H), 2.02 - 1.80 (m, 2H), 0.81 (t, J = 7 Hz, 3H).

[01444] **Step 2: 1-(1-Isocyanatopropyl)-4-(trifluoromethyl)benzene.** General procedure C was followed using 1-[4-(trifluoromethyl)phenyl]propan-1-amine to give 1-(1-isocyanatopropyl)-4-(trifluoromethyl)benzene as a yellow solid (226 mg, crude). LCMS in MeOH (ESI): m/z : $[M+MeOH+H]^+ = 262.1$.

[01445] **Step 3: Propan-2-yl 3-(pyrimidin-2-yl)-2-[(1-[4-(trifluoromethyl)phenyl]propyl}carbamoyl)oxy]propanoate.** General procedure D was followed using propan-2-yl 2-hydroxy-3-(pyrimidin-2-yl)propanoate and 1-(1-isocyanatopropyl)-4-(trifluoromethyl)benzene. Purification by prep-HPLC (column: Welch Xtimate C18, 10 μ m, 250 x 50 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 50 - 70 %, 10 min) gave propan-2-yl 3-(pyrimidin-2-yl)-2-[(1-[4-(trifluoromethyl)phenyl]propyl}carbamoyl)oxy]propanoate as a white solid (Y = 12 %). ¹HNMR (400 MHz, ACN-*d*₃) δ ppm 8.80 - 8.44 (m, 2H), 7.65 - 7.55 (m, 2H), 7.48 - 7.35 (m, 2H), 7.31 - 7.28 (m, 1H), 6.23 - 6.15 (m, 1H), 5.46 - 5.35 (m, 1H), 5.08 - 4.82 (m, 1H), 4.52 - 4.45 (m, 1H), 3.45 - 3.28 (m, 2H), 1.76 - 1.61 (m, 2H), 1.25 - 1.16 (m, 3H), 1.12 - 1.01 (m, 3H), 0.91 - 0.74 (m, 3H). LCMS (ESI): m/z : $[M+H]^+ = 440.0$.

Example 29. Propan-2-yl (2R)-2-[(1S)-1-[4-(2-phenylethyl)phenyl]ethyl]carbamoyl]oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.



[01446] **Step 1: 1-[4-(2-Phenylethyl)phenyl]ethan-1-one.** 2-Phenylethylboronic acid (6.0 g, 40 mmol), 1-(4-bromophenyl)ethanone (7.96 g, 40 mmol), Pd(dppf)Cl₂.CH₂Cl₂ (3.27 g, 4.0 mmol) and Cs₂CO₃ (13.03 g, 40 mmol) in THF (60 ml) and H₂O (6 ml) was degassed and purged with N₂ three times. The reaction was stirred at 100° C for 12 h under N₂. The solvent was removed *in*

vacuo and the residue purified by column chromatography (SiO₂, pet. ether: EtOAc, 5:1 to 3:1) to give 1-[4-(2-phenylethyl)phenyl]ethan-1-one as a brown solid (Y = 64 %). ¹H NMR (400 MHz, MeOD) δ ppm 7.87 (d, *J* = 8 Hz, 2H), 7.28 - 7.21 (m, 4H), 7.16 - 7.13 (m, 3H), 3.00 - 2.90 (m, 4H), 2.55 (s, 3H).


[01447] **Step 2: (S)-2-Methyl-N-[(1S)-1-[4-(2-phenylethyl)phenyl]ethyl]propane-2-sulfinamide.** General procedure A was followed using 1-[4-(2-phenylethyl)phenyl]ethenone. The crude product was purified by column chromatography (SiO₂, pet. ether: EtOAc, 5:1 to 1:1) to give (S)-2-methyl-N-[(1S)-1-[4-(2-phenylethyl)phenyl]ethyl]propane-2-sulfinamide as a light-yellow solid (Y = 37 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.26 - 7.21 (m, 4H), 7.17 - 7.12 (m, 5H), 4.52 - 4.47 (m, 1H), 3.37 (s, 1H), 2.88 (s, 4H), 1.46 (d, *J* = 7 Hz, 3H), 1.20 (s, 9H).

[01448] **Step 3: (1S)-1-[4-(2-Phenylethyl)phenyl]ethan-1-aminium chloride.** General procedure B was followed using (S)-2-methyl-N-((S)-1-(4-phenethylphenyl)ethyl)propane-2-sulfinamide. Purification by FCC (SiO₂, pet. ether: EtOAc, 5:1 to 1:1) gave (1S)-1-[4-(2-phenylethyl)phenyl]ethan-1-aminium chloride as a white solid (Y = quantitative). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.61 (br. s, 2H), 7.41 (d, *J* = 6 Hz, 2H), 7.31 - 7.24 (m, 2H), 7.22 - 7.13 (m, 5H), 4.44 - 4.28 (m, 1H), 2.89 (s, 4H), 1.66 (s, 3H).

[01449] **Step 4: 1-[(1S)-1-Isocyanatoethyl]-4-(2-phenylethyl)benzene.** General procedure C was followed using (1S)-1-[4-(2-phenylethyl)phenyl]ethanamine to give 1-[(1S)-1-isocyanatoethyl]-4-(2-phenylethyl)benzene (Y = quantitative), which was used without purification.

[01450] **Step 5: Propan-2-yl (2R)-2-([(1S)-1-[4-(2-phenylethyl)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.** General procedure D was followed using isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate and 1-[(1S)-1-isocyanatoethyl]-4-(2-phenylethyl)benzene. Purification by prep-HPLC (column: Huapu C8 Extreme BDS, 5 μm, 150 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 50 - 70 %, 10 min) gave propan-2-yl (2R)-2-([(1S)-1-[4-(2-phenylethyl)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate as a colourless gum (Y = 14 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.51 (s, 1H), 7.95 (s, 1H), 7.26 - 7.21 (m, 4H), 7.18 - 7.14 (m, 5H), 5.15 (t, *J* = 5 Hz, 1H), 4.84 - 4.79 (m, 1H), 4.62 - 4.59 (m, 2H), 4.58 - 4.52 (m, 1H), 2.81 (s, 4H), 1.29 (d, *J* = 7 Hz, 3H), 1.07 (d, *J* = 6 Hz, 3H), 1.01 (d, *J* = 6 Hz, 3H). LCMS (ESI): *m/z*: [M+H]⁺ = 451.2.

Example 30. Propan-2-yl (2R)-2-([(1S)-1-{[1,1'-biphenyl]-4-yl}ethyl]carbamoyl)oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.



[01451] Step 1: (S)-N-[(1S)-1-{[1,1'-Biphenyl]-4-yl}ethyl]-2-methylpropane-2-sulfonamide.

General procedure A was followed using 1-(4-phenylphenyl)ethenone. Purification by FCC (SiO₂, pet. ether: EtOAc, 1:1) gave (S)-N-[(1S)-1-{[1,1'-Biphenyl]-4-yl}ethyl]-2-methylpropane-2-sulfonamide as a white solid. Y = 45 %. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.62 - 7.56 (m, 4H), 7.49 - 7.41 (m, 4H), 7.39 - 7.33 (m, 1H), 4.64 - 4.59 (m, 1H), 3.48 (d, *J* = 2 Hz, 1H), 1.56 (d, *J* = 7 Hz, 3H), 1.27 (s, 9H).

[01452] Step 2: (1S)-1-{[1,1'-Biphenyl]-4-yl}ethan-1-amine hydrochloride. General procedure B was followed using (S)-N-[(1S)-1-{[1,1'-biphenyl]-4-yl}ethyl]-2-methylpropane-2-sulfonamide to give (1S)-1-{[1,1'-Biphenyl]-4-yl}ethan-1-amine hydrochloride as a white solid. Y = 64 %. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.57 (s, 3H), 7.68 (dd, *J* = 8, 17 Hz, 4H), 7.59 (d, *J* = 8 Hz, 2H), 7.50 - 7.43 (m, 2H), 7.41 - 7.33 (m, 1H), 4.43 - 4.40 (m, 1H), 1.53 (d, *J* = 7 Hz, 3H).


[01453] Step 3: 4-[(1S)-1-isocyanatoethyl]-1,1'-biphenyl. General procedure C was followed using (1S)-1-(4-phenylphenyl)ethanamine to give 4-[(1S)-1-isocyanatoethyl]-1,1'-biphenyl as a light yellow oil (Y = 100 %).

[01454] Step 4: Propan-2-yl (2R)-2-([(1S)-1-{[1,1'-biphenyl]-4-yl}ethyl]carbamoyl)oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate. General procedure D was followed using isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate and 1-[(1S)-1-isocyanatoethyl]-4-phenyl-benzene. Purification by prep-TLC (SiO₂, EtOAc) gave propan-2-yl (2R)-2-([(1S)-1-{[1,1'-biphenyl]-4-yl}ethyl]carbamoyl)oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate as a white solid (Y = 31 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.11 (s, 1H), 7.96 (s, 1H), 7.57 (d, *J* = 8 Hz, 4H), 7.44 (t, *J* = 8 Hz, 2H), 7.41 - 7.34 (m, 3H), 5.38 (t, *J* = 5 Hz, 1H), 5.31 (d, *J* = 8 Hz, 1H), 5.09 - 5.03 (m, 1H),

4.89 - 4.82 (m, 1H), 4.65 (d, $J = 5$ Hz, 2H), 1.53 (d, $J = 7$ Hz, 3H), 1.20 (dd, $J = 6, 13$ Hz, 6H).

LCMS (ESI): m/z : $[M+H]^+$ 423.2.

Example 31. Propan-2-yl (2R)-2-({[(1S)-1-[4-(2-methylpropyl)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.



[01455] **Step 1: (S)-2-Methyl-N-[(1S)-1-[4-(2-methylpropyl)phenyl]ethyl]propane-2-sulfonamide.** General procedure A was followed using 1-(4-isobutylphenyl)ethanone. The crude product was purified by column chromatography (SiO_2 , pet. ether:EtOAc, 5:1 to 1:1) to give (S)-2-methyl-N-[(1S)-1-[4-(2-methylpropyl)phenyl]ethyl]propane-2-sulfonamide as a white solid (Y = 62 %). ^1H NMR (400 MHz, MeOD) δ ppm 7.29 (d, $J = 8$ Hz, 2H), 7.12 (d, $J = 8$ Hz, 2H), 4.47 - 4.42 (m, 1H), 2.46 (d, $J = 7$ Hz, 2H), 1.89 - 1.78 (m, 1H), 1.50 (d, $J = 7$ Hz, 3H), 1.24 (s, 9H), 0.90 (d, $J = 7$ Hz, 6H).


[01456] **Step 2: (1S)-1-[4-(2-Methylpropyl)phenyl]ethan-1-aminium chloride.** General procedure B was followed using (S)-N-((S)-1-(4-isobutylphenyl)ethyl)-2-methylpropane-2-sulfonamide to give (1S)-1-[4-(2-Methylpropyl)phenyl]ethan-1-aminium chloride as a white solid (Y = 97 %). ^1H NMR (400 MHz, MeOD) δ ppm 7.38 (d, $J = 8$ Hz, 2H), 7.24 (d, $J = 8$ Hz, 2H), 4.46 - 4.42 (m, 1H), 2.51 (d, $J = 7$ Hz, 2H), 1.90 - 1.83 (m, 1H), 1.17 (d, $J = 10$ Hz, 3H), 0.90 (d, $J = 7$ Hz, 6H).

[01457] **Step 3: 1-[(1S)-1-Isocyanatoethyl]-4-(2-methylpropyl)benzene.** General procedure C was followed using (1S)-1-(4-isobutylphenyl)ethanamine to give 1-[(1S)-1-isocyanatoethyl]-4-(2-methylpropyl)benzene (Y = 100 %).

[01458] **Step 4: Propan-2-yl (2R)-2-({[(1S)-1-[4-(2-methylpropyl)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.** General procedure was followed using isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate and (S)-1-isobutyl-4-(1-isocyanatoethyl)benzene. Purification by prep-HPLC (column: Luna C18, 5

μm , 100 x 30 mm; mobile phase: [water (0.1 % TFA) - MeOH]; B: 60 - 80 %, 12 min) gave Propan-2-yl (2R)-2-({[(1S)-1-[4-(2-methylpropyl)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate as a white solid (Y = 7 %). LCMS (ESI): m/z : $[M+H]^+ = 403.2$. ^1H NMR (400 MHz, CDCl_3) δ ppm 8.10 (s, 1H), 7.95 (s, 1H), 7.23 - 7.19 (m, 2H), 7.12 - 7.10 (m, 2H), 5.36 (t, $J = 5$ Hz, 1H), 5.23 (d, $J = 7$ Hz, 1H), 5.07 - 5.04 (m, 1H), 4.80 - 4.78 (m, 1H), 4.64 (d, $J = 5$ Hz, 2H), 2.47 - 2.44 (m, 2H), 1.87 - 1.81 (m, 1H), 1.48 (d, $J = 7$ Hz, 3H), 1.26 - 1.20 (m, 6H), 0.90 (d, $J = 6$ Hz, 6H).

Example 32. Propan-2-yl (2R)-2-({[(1S)-1-[4-(benzyloxy)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.



[01459] **Step 1: (S)-N-[(1S)-1-[4-(Benzyloxy)phenyl]ethyl]-2-methylpropane-2-sulfonamide.**


General procedure A was followed using 1-(4-benzyloxyphenyl)ethanone. Purification by FCC (SiO_2 , pet. ether:EtOAc, 5:1 to 1:1) gave (S)-N-[(1S)-1-[4-(benzyloxy)phenyl]ethyl]-2-methylpropane-2-sulfonamide as a white solid (Y = 39 %). ^1H NMR (400 MHz, MeOD) δ ppm 7.43 - 7.40 (m, 2H), 7.37 - 7.33 (m, 3H), 7.31 - 7.28 (m, 2H), 6.98 - 6.94 (m, 2H), 5.06 (s, 2H), 4.44 - 4.39 (m, 1H), 1.48 (d, $J = 7$ Hz, 3H), 1.21 (s, 9H).

[01460] **Step 2: [4-(Benzyloxy)phenyl]ethan-1-aminium chloride.** General procedure B was followed using (S)-N-[(S)-1-(4-(benzyloxy)phenyl)ethyl]-2-methylpropane-2-sulfonamide to give [4-(benzyloxy)phenyl]ethan-1-aminium chloride as a light yellow solid (Y = 94 %). ^1H NMR (400 MHz, MeOD) δ ppm 7.43 - 7.42 (m, 2H), 7.40 - 7.34 (m, 4H), 7.32 - 7.30 (m, 1H), 7.09 - 7.05 (m, 2H), 5.12 (s, 2H), 4.43 - 4.38 (m, 1H), 1.61 (d, $J = 7$ Hz, 3H).

[01461] **Step 3: 1-(Benzyloxy)-4-[(1S)-1-isocyanatoethyl]benzene.** General procedure C was followed using (1S)-1-(4-benzyloxyphenyl)ethanamine to give 1-(benzyloxy)-4-[(1S)-1-isocyanatoethyl]benzene (Y = quantitative).

[01462] **Step 4: Propan-2-yl (2R)-2-([(1S)-1-[4-(benzyloxy)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate.** General procedure D was followed using isopropyl (2R)-2-hydroxy-3-(1,2,4-triazol-1-yl)propanoate and 1-benzyloxy-4-[(1S)-1-isocyanatoethyl]benzene. Purification by prep-TLC (SiO₂, pet. ether: EtOAc, 1:1) gave propan-2-yl 2-([(1S)-1-[4-(benzyloxy)phenyl]ethyl]carbamoyl}oxy)-3-(1H-1,2,4-triazol-1-yl)propanoate as a white solid (Y = 12 %). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.09 (s, 1H), 7.95 (s, 1H), 7.44 - 7.31 (m, 5H), 7.23 (d, *J* = 9 Hz, 2H), 6.94 (d, *J* = 9 Hz, 2H), 5.36 (t, *J* = 6 Hz, 1H), 5.17 (d, *J* = 8 Hz, 1H), 5.11 - 5.02 (m, 3H), 4.76 - 4.74 (m, 1H), 4.64 (d, *J* = 5 Hz, 2H), 1.47 (d, *J* = 7 Hz, 3H), 1.25 - 1.16 (m, 6H). LCMS (ESI): *m/z*: [M+H]⁺ = 453.2.

Example 33. Propan-2-yl 2-([(1S)-1-[4-(2-methylpropyl)phenyl]ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate.



[01463] **Step 1: [(1S)-1-Isocyanatoethyl]benzene.** General procedure C was used with (S)-1-phenylethanamine to give [(1S)-1-isocyanatoethyl]benzene as a yellow liquid. Y = 98 %. LCMS in MeOH (ESI): *m/z*: [M+MeOH+H]⁺ = 179.9.

[01464] **Step 2: Propan-2-yl 2-([(1S)-1-[4-(2-methylpropyl)phenyl]ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate.** General procedure D was followed using [(1S)-1-isocyanatoethyl]benzene and isopropyl 2-hydroxy-3-(pyrimidin-2-yl)propanoate. Purification by prep-HPLC (column: Nano-micro Kromasil C18, 5 μm, 100 x 30 mm; mobile phase: [water (0.1 % TFA) - ACN]; B: 28 - 58 %, 10 min) gave propan-2-yl 2-([(1S)-1-[4-(2-methylpropyl)phenyl]ethyl]carbamoyl}oxy)-3-(pyrimidin-2-yl)propanoate as a colourless gum (Y = 34 %). ¹H NMR (400 MHz, ACN-*d*₃) δ ppm 8.72 - 8.63 (m, 2H), 7.34 - 7.20 (m, 6H), 6.09 - 6.05 (m, 1H), 5.45 - 5.39 (m, 1H), 5.01 - 4.92 (m, 1H), 4.67 - 4.64 (m, 1H), 3.40 - 3.28 (m, 2H), 1.37 - 1.33 (m, 3H), 1.21 - 1.15 (m, 5H), 1.08 - 1.04 (m, 1H). LCMS (ESI): *m/z*: [M+H]⁺ = 358.1.

Biological Activity of the Compounds of the Present Disclosure

[01465] The biological activity of the compounds of the present disclosure was determined utilising the assay described herein.

PBMC IC₅₀ determination assay

[01466] The compounds of the present disclosure were tested for their inhibitory activity against IL-1 β release upon NLRP3 activation in peripheral blood mononuclear cells (PBMC).

[01467] **Protocol A.** PBMC were isolated from buffy coats by density gradient centrifugation on Histopaque-1077 (Sigma, cat no. 10771). Isolated cells were seeded into the wells of a 96-well plate and incubated for 3 h with lipopolysaccharide (LPS). Following medium exchange, the compounds of the present disclosure were added (a single compound per well) and the cells were incubated for 30 min. Next, the cells were stimulated either with ATP (5 mM) or nigericin (10 μ M) for 1 h and the cell culture media from the wells were collected for further analysis.

[01468] The release of IL-1 β into the media was determined by a quantitative detection of IL-1 β in the media using an IL-1 β enzyme-linked immunosorbent assay (ELISA) Ready-SET-Go!, eBioscience cat. No. 88-7261-88. Briefly, in a first step, high affinity binding plates (Corning, Costar 9018 or NUNC Maxisorp Cat No. 44-2404) were coated overnight at 4 °C with specific capture antibody included in the kit (anti-human IL-1 β ref. 14-7018-68). Subsequently, plates were blocked with blocking buffer for 1 h at room temperature (rt) and after washing with a buffer (PBS with 0.05 % Tween-20) incubated with protein standard and culture media. After 2 h of incubation at rt, plates were washed and incubated with biotinylated detection antibody included in the kit (anti-human IL-1 β Biotin ref. 33-7110-68) for 1 h at rt. Plates were washed and incubated with HRP-streptavidin for 30 min at rt and washed again. The signal was developed after addition of 3,3',5,5'-tetramethylbenzidine-peroxidase (TMB) until colour appeared and the reaction was stopped by 2 M H₂SO₄. A microplate spectrophotometer (BioTek) was used to detect signals with 450 nm. The detection range of IL-1 β ELISA was 2-150 ng/ml.

[01469] **Protocol B.** PBMC were isolated from buffy coats by density gradient centrifugation on Histopaque-1077 (Sigma, cat no. 10771). Isolated cells were seeded into the wells (280,000 cells/well) of a 96-well plate and incubated for 3 h with lipopolysaccharide (LPS, 1 μ g/ml diluted 1000x from a 1 mg/ml stock solution). The compounds of the present disclosure were added (a single compound per well) and the cells were incubated for 30 min. Next, the cells were

stimulated with ATP (5 mM final concentration diluted 20x from a 100 mM stock solution) for 1 h and the cell culture media from the wells were collected for further analysis.

[01470] The release of IL-1 β into the media was determined by quantitative detection of IL-1 β in the media using HTRF®, CisBio cat. No. 62HIL1BPEH. Briefly, cell culture supernatant were dispensed directly into the assay plate containing antibodies labelled with the HTRF® donor and acceptor. A microplate spectrophotometer (BMG) was used to detect signals at 655 nm and 620 nm. The detection range of IL-1 β HTRF® was 39-6500 pg/ml.

[01471] The determination of the IC₅₀ values was preformed using the Graph Pad Prism software and the measured IC₅₀ values of compounds of the present disclosure are shown in Table A below (“++++” means <1 μ M; “+++” means ≥ 1 and <3 μ M; “++” means ≥ 3 and <10 μ M; “+” means ≥ 10 and <50 μ M). These results show that the compounds of the present disclosure are capable of inhibiting IL-1 β release upon inflammasome activation.

Table A

Compound No.	Average PBMC IC ₅₀ (μ M)
1	+
2	+
3	++++
4	+++
5	++++
6	++++
7	++++
8	+++
9	++++
10	++++
11	++++
12	++++
13	++++
14	+
15	+
16	++++
18	++
19	++++
20	++++
21	+++
22	++++
23	++++
24	++

Compound No.	Average PBMC IC ₅₀ (μM)
25	++++
26	++
27	+++


EQUIVALENTS

[01472] The details of one or more embodiments of the disclosure are set forth in the accompanying description above. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, the preferred methods and materials are now described. Other features, objects, and advantages of the disclosure will be apparent from the description and from the claims. In the specification and the appended claims, the singular forms include plural referents unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. All patents and publications cited in this specification are incorporated by reference.

[01473] The foregoing description has been presented only for the purposes of illustration and is not intended to limit the disclosure to the precise form disclosed, but by the claims appended hereto.

CLAIMS:

1. A compound of Formula (I'):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein:

R₁ is C₃-C₁₆ cycloalkyl, 5- to 10-membered heteroaryl, or C₅-C₁₀ aryl, wherein the C₃-C₁₆ cycloalkyl, 5- or 6-membered heteroaryl, or C₅-C₁₀ aryl is optionally substituted with one or more **R_{1S}**;

each **R_{1S}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ alkoxy, halo, or CN; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₁₀ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, or C₁-C₆ alkoxy is optionally substituted with one or more **R_{1SS}**;

each **R_{1SS}** is independently C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, -O-(C₃-C₈ cycloalkyl), -O-(C₅-C₆ aryl), -O-(5- or 6-membered heteroaryl), -O-(C₃-C₈ heterocycloalkyl), C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

Q is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₂ is 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl; wherein the 5- to 10-membered heteroaryl or 4- to 8-membered heterocycloalkyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₃ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or R_{3S};

each **R_{3S}** is independently selected from -OC(=O)R_{3Sa} or 5- to 6- membered heterocycloalkyl; wherein the 5- or 6-membered heterocycloalkyl is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;


R_{3Sa} is C₁-C₆ alkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or -C(=O)OR_{3Sb};

R_{3Sb} is 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl); wherein the 4- to 8-membered heterocycloalkyl or -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) is optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo;

R₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo; and

R₆ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

2. The compound of claim 1, wherein the compound is of Formula (II'):





or a prodrug, solvate, or pharmaceutically acceptable salt thereof.


3. The compound of any one of the preceding claims, wherein **X** is O.
4. The compound of any one of the preceding claims, wherein **R₁** is C₃-C₁₆ cycloalkyl optionally substituted with one or more **R_{1S}**.
5. The compound of any one of the preceding claims, wherein **R₁** is 5- to 10-membered heteroaryl optionally substituted with one or more **R_{1S}**.
6. The compound of any one of the preceding claims, wherein **R₁** is 5- or 6-membered heteroaryl optionally substituted with one or more **R_{1S}**.
7. The compound of any one of the preceding claims, wherein **R₁** is 6-membered heteroaryl optionally substituted with one or more **R_{1S}**.
8. The compound of any one of the preceding claims, wherein **R₁** is pyridinyl optionally substituted with one or more **R_{1S}**.
9. The compound of any one of the preceding claims, wherein **R₁** is pyrazinyl optionally substituted with one or more **R_{1S}**.


10. The compound of any one of the preceding claims, wherein **R₁** is .

11. The compound of any one of the preceding claims, wherein **R₁** is .

12. The compound of any one of the preceding claims, wherein **R₁** is .

13. The compound of any one of the preceding claims, wherein **R₁** is .

14. The compound of any one of the preceding claims, wherein **R₁** is .




15. The compound of any one of the preceding claims, wherein **R₁** is .

16. The compound of any one of the preceding claims, wherein **R₁** is C₅-C₁₀ aryl optionally substituted with one or more **R_{1S}**.

17. The compound of any one of the preceding claims, wherein **R₁** is C₅-C₆ monocyclic aryl optionally substituted with one or more **R_{1S}**.



18. The compound of any one of the preceding claims, wherein **R₁** is phenyl optionally substituted with one or more **R_{1S}**.

19. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one **R_{1S}**.


20. The compound of any one of the preceding claims, wherein **R₁** is , , or .

21. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with two **R_{1S}**.


22. The compound of any one of the preceding claims, wherein **R₁** is





23. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with three **R_{1S}**.



24. The compound of any one of the preceding claims, wherein **R₁** is



25. The compound of any one of the preceding claims, wherein **R₁** is



26. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more substituents independently selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, C₁-C₆ alkoxy, halo, or CN.


27. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more substituents independently selected from methyl, ethyl, isopropyl, tert-butyl, ethoxy, cyclopropyl, chloro, or -CN.
28. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₁-C₆ alkyl optionally substituted with one or more **R_{1SS}**.
29. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₁-C₆ alkyl.
30. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more C₁-C₆ alkyl.
31. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more -O-(C₅-C₆ aryl).
32. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one or more halo.
33. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₁-C₆ alkyl substituted with one, two, or three F.
34. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl optionally substituted with one or more **R_{1SS}**.
35. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₂-C₆ alkenyl.
36. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl optionally substituted with one or more **R_{1SS}**.

37. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₂-C₆ alkynyl.
38. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl optionally substituted with one or more **R_{1SS}**.
39. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₃-C₈ cycloalkyl.
40. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₅-C₆ aryl optionally substituted with one or more **R_{1SS}**.
41. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₅-C₆ aryl.
42. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl optionally substituted with one or more **R_{1SS}**.
43. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more 5- or 6-membered heteroaryl.
44. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₃-C₈ heterocycloalkyl optionally substituted with one or more **R_{1SS}**.
45. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or C₃-C₈ heterocycloalkyl.
46. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy optionally substituted with one or more **R_{1SS}**.

47. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or C₁-C₆ alkoxy.

48. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₅-C₆ aryl.

49. The compound of any one of the preceding claims, wherein **R₁** is phenyl substituted with one or more C₁-C₆ alkoxy substituted with one or more C₁-C₆ haloalkyl.



51. The compound of any one of the preceding claims, wherein at least one **R_{1S}** is C₁-C₆ alkyl.


52. The compound of any one of the preceding claims, wherein at least one **R_{1S}** is C₁-C₆ alkoxy.
53. The compound of any one of the preceding claims, wherein at least one **R_{1S}** is C₅-C₆ aryl.
54. The compound of any one of the preceding claims, wherein at least one **R_{1S}** is 5- or 6-membered heteroaryl.
55. The compound of any one of the preceding claims, wherein at least one **R_{1S}** is C₃-C₈ heterocycloalkyl.
56. The compound of any one of the preceding claims, wherein at least one **R_{1S}** is halo or CN.
57. The compound of any one of the preceding claims, wherein at least one **R_{1SS}** is C₁-C₆ alkyl.
58. The compound of any one of the preceding claims, wherein at least one **R_{1SS}** is C₅-C₆ aryl.
59. The compound of any one of the preceding claims, wherein at least one **R_{1SS}** is phenyl.
60. The compound of any one of the preceding claims, wherein at least one **R_{1SS}** is 5- or 6-membered heteroaryl.
61. The compound of any one of the preceding claims, wherein at least one **R_{1SS}** is C₃-C₈ heterocycloalkyl.
62. The compound of any one of the preceding claims, wherein at least one **R_{1SS}** is -O-(C₅-C₆ aryl).

63. The compound of any one of the preceding claims, wherein at least one **R**_{1SS} is -O-(phenyl).
64. The compound of any one of the preceding claims, wherein at least one **R**_{1SS} is -O-(5- or 6-membered heteroaryl).
65. The compound of any one of the preceding claims, wherein at least one **R**_{1SS} is -O-(C₃-C₈ heterocycloalkyl).
66. The compound of any one of the preceding claims, wherein at least one **R**_{1SS} is C₁-C₆ haloalkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkoxy.
67. The compound of any one of the preceding claims, wherein at least one **R**_{1SS} is halo, -CN, or oxo.
68. The compound of any one of the preceding claims, wherein at least one **R**_{1SS} is -NH₂, -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂.
69. The compound of any one of the preceding claims, wherein **Q** is methyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
70. The compound of any one of the preceding claims, wherein **Q** is methyl.
71. The compound of any one of the preceding claims, wherein **R**₂ is 5- to 10-membered heteroaryl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.


72. The compound of any one of the preceding claims, wherein **R**₂ is pyrazolyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.


73. The compound of any one of the preceding claims, wherein **R**₂ is .


74. The compound of any one of the preceding claims, wherein **R**₂ is triazolyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

75. The compound of any one of the preceding claims, wherein **R**₂ is .


76. The compound of any one of the preceding claims, wherein **R**₂ is pyrimidinyl optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

77. The compound of any one of the preceding claims, wherein **R**₂ is .


78. The compound of any one of the preceding claims, wherein **R**₂ is , , or



79. The compound of any one of the preceding claims, wherein **Q** and **R₂** together forms



80. The compound of any one of the preceding claims, wherein **Q** and **R₂** together forms



81. The compound of any one of the preceding claims, wherein **R₃** is H, C₁-C₆ alkyl, C₃-C₆ alkenyl, or C₃-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, oxo, or **R_{3s}**.


82. The compound of any one of the preceding claims, wherein **R₃** is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

83. The compound of any one of the preceding claims, wherein **R₃** is H.

84. The compound of any one of the preceding claims, wherein **R₃** is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.

85. The compound of any one of the preceding claims, wherein **R₃** is C₁-C₆ alkyl.

86. The compound of any one of the preceding claims, wherein **R₃** is ethyl.
87. The compound of any one of the preceding claims, wherein **R₃** is propyl.
88. The compound of any one of the preceding claims, wherein **R₃** is C₁-C₆ alkyl optionally substituted with one or more **R_{3s}**.
89. The compound of any one of the preceding claims, wherein **R_{3s}** is -OC(=O)**R_{3Sa}**.
90. The compound of any one of the preceding claims, wherein **R_{3s}** is 5- to 6- membered heterocycloalkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
91. The compound of any one of the preceding claims, wherein **R_{3s}** is 5- to 6- membered heterocycloalkyl optionally substituted with one or more -OH.
92. The compound of any one of the preceding claims, wherein **R_{3Sa}** is C₁-C₆ alkyl optionally substituted with one or more -C(=O)OR_{3Sb}.
93. The compound of any one of the preceding claims, wherein **R_{3Sb}** is 4- to 8-membered heterocycloalkyl optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
94. The compound of any one of the preceding claims, wherein **R_{3Sb}** is -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more -OH, halo, -CN, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.
95. The compound of any one of the preceding claims, wherein **R_{3Sb}** is -(C₁-C₆ alkyl)-(4- to 8-membered heterocycloalkyl) optionally substituted with one or more -OH.



97. The compound of any one of the preceding claims, wherein **R**₄ is C₁-C₆ alkyl, C₂-C₆ alkenyl or C₂-C₆ alkynyl.

98. The compound of any one of the preceding claims, wherein **R**₄ is C₁-C₆ alkyl.

99. The compound of any one of the preceding claims, wherein **R**₄ is methyl.


100. The compound of any one of the preceding claims, wherein **R**₄ is  or .

101. The compound of any one of the preceding claims, wherein **R**₅ is H.

102. The compound of any one of the preceding claims, wherein **R**₅ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl; wherein the C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl is optionally substituted with one or more C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₅-C₆ aryl, 5- or 6-membered heteroaryl, C₃-C₈ heterocycloalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, halo, -CN, -OH, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, or oxo.


103. The compound of any one of the preceding claims, wherein **R**₆ is C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl.

104. The compound of any one of the preceding claims, wherein the compound is of Formula (I-a):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₂**, **R₃**, **R₄**, and **Q** are as described herein.

105. The compound of any one of the preceding claims, wherein the compound is of Formula (I-a1), (I-a2), (I-a3), or (I-a4):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₂**, **R₃**, **R₄**, and **Q** are as described herein.

106. The compound of any one of the preceding claims, wherein the compound is of Formula (I-a5):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₂**, **R₄**, and **Q** are as described herein.

107. The compound of any one of the preceding claims, wherein the compound is of Formula (I-a6), (I-a7), (I-a8), or (I-a9):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₂**, **R₄**, and **Q** are as described herein.


108. The compound of any one of the preceding claims, wherein the compound is of Formula (I-c), (I-d), or (I-e):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R_{1S}**, **R₂**, **R₃**, **R₄**, **X**, and **Q** are as described herein.


109. The compound of any one of the preceding claims, wherein the compound is of Formula (I-f), (I-g), or (I-h):






or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₂**, **R₃**, **R₄**, and **X** are as described herein.

110. The compound of any one of the preceding claims, wherein the compound is of Formula (I-i), (I-j), or (I-k):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₃**, **R₄**, **X**, and **Q** are as described herein.

111. The compound of any one of the preceding claims, wherein the compound is of Formula (I-l) or (I-m):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₂**, **R₄**, **X**, and **Q** are as described herein.

112. The compound of any one of the preceding claims, wherein the compound is of Formula (I-n), (I-o), or (I-p):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₂**, **R₃**, **X**, and **Q** are as described herein.


113. The compound of any one of the preceding claims, wherein the compound is of Formula (Ia-a):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R_{1S}**, **R₂**, **R₃**, **R₄**, and **Q** are as described herein.


114. The compound of any one of the preceding claims, wherein the compound is of Formula (Ia-b), (Ia-c) or (Ia-d):






or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R_{1S}**, **R₂**, **R₃**, and **R₄** are as described herein.

115. The compound of any one of the preceding claims, wherein the compound is of Formula (Ia-e), (Ia-f), or (Ia-g):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R_{1S}**, **R₃**, **R₄**, and **Q** are as described herein.

116. The compound of any one of the preceding claims, wherein the compound is of Formula (Ia-h), or (Ia-i):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R_{1S}**, **R₂**, **R₄**, and **Q** are as described herein.

117. The compound of any one of the preceding claims, wherein the compound is of Formula (Ia-j), (Ia-k), or (Ia-l):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein R_{1S} , R_2 , R_3 , and Q are as described herein.

118. The compound of any one of the preceding claims, wherein the compound is of Formula (Ia-m), (Ia-n), (Ia-o), or (Ia-p):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein R_{1S} , R_3 , and R_4 are as described herein.

119. The compound of any one of the preceding claims, wherein the compound is of Formula (Ia-q), (Ia-r), (Ia-s), or (Ia-t):




or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R_{1s}**, **R₂**, and **R₄** are as described herein.

120. The compound of any one of the preceding claims, wherein the compound is of Formula (Ia-u), (Ia-v), (Ia-w), (Ia-x), (Ia-y), or (Ia-z):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R_{1s}**, **R₂**, and **R₃** are as described herein.

121. The compound of any of the preceding claims, being of Formula (Ic-a), (Ic-b), (Ic-c), (Ic-d), or (Ic-e):



or a prodrug, solvate, or pharmaceutically acceptable salt thereof, wherein **R₁**, **R₄**, **Q**, and **R₂** are as described herein.

122. The compound of any one of the claims 1-121, being selected from Compound Nos. 1-32 and prodrugs and pharmaceutically acceptable salts thereof.

123. The compound of any one of claims 1-121, being selected from Compound Nos. 1-32 and pharmaceutically acceptable salts thereof.

124. The compound of any one of claims 1-121, being selected from Compound Nos. 1-32.

125. A compound being an isotopic derivative of the compound of any one of claims 1-124.

126. The compound of claim 125, being a deuterium labeled compound of any one of Compound Nos. 1-32 and prodrugs and pharmaceutically acceptable salts thereof.

127. The compound of claim 125, being a deuterium labeled compound of any one of Compound Nos. 1-32.
128. A compound obtainable by, or obtained by, a method described herein; optionally, the method comprises one or more steps described in Scheme 1 or Scheme 2.
129. A pharmaceutical composition comprising the compound of any one of claims 1-128 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable diluent or carrier.
130. The pharmaceutical composition of claim 129, wherein the compound is selected from Compound Nos. 1-32.
131. A method of inhibiting inflammasome activity, comprising contacting a cell with an effective amount of the compound of any one of claims 1-128 or a pharmaceutically acceptable salt thereof; optionally, the inflammasome is NLRP3 inflammasome, and the activity is *in vitro* or *in vivo*.
132. A method of treating or preventing a disease or disorder in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of the compound of any one of claims 1-128 or a pharmaceutically acceptable salt thereof, or the pharmaceutical composition of any one of claims 129 or 130.
133. The method of claim 132, wherein the disease or disorder is associated with an implicated inflammasome activity; optionally, the disease or disorder is a disease or disorder in which inflammasome activity is implicated.
134. The method of claim 132 or claim 133, wherein the disease or disorder is an inflammatory disorder, an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease, or cancer.

135. The method of any one of claims 132-134, wherein the disease or disorder is an inflammatory disorder, an autoinflammatory disorder or an autoimmune disorder; optionally, the disease or disorder is selected from cryopyrin-associated auto-inflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological disease (e.g., acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases).

136. The method of any one of claims 132-134, wherein disease or disorder is a neurodegenerative disease; optionally, the disease or disorder is Parkinson's disease or Alzheimer's disease.

137. The method of any one of claims 132-134, wherein the disease or disorder is cancer; optionally, the cancer is metastasising cancer, brain cancer, gastrointestinal cancer, skin cancer, non-small-cell lung carcinoma, head and neck squamous cell carcinoma or colorectal adenocarcinoma.

138. The compound of any one of claims 1-128 or the pharmaceutical composition of any one of claims 129 or 130, for use in inhibiting inflammasome activity; optionally, the inflammasome is NLRP3 inflammasome, and the activity is *in vitro* or *in vivo*.

139. The compound of any one of claims 1-128, or the pharmaceutical composition of any one of claims 129 or 130, for use in treating or preventing a disease or disorder.

140. The compound or pharmaceutical composition of claim 139, wherein the disease or disorder is associated with an implicated inflammasome activity; optionally, the disease or disorder is a disease or disorder in which inflammasome activity is implicated.

141. The compound or pharmaceutical composition of claim 139 or claim 140, wherein the disease or disorder is an inflammatory disorder, an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease, or cancer.

142. The compound or pharmaceutical composition of any one of claims 139-141, wherein the disease or disorder is an inflammatory disorder, an autoinflammatory disorder or an autoimmune disorder; optionally, the disease or disorder is selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological diseases (e.g., acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases).

143. The compound or pharmaceutical composition of any one of claims 139-141, wherein disease or disorder is a neurodegenerative disease; optionally, the disease or disorder is Parkinson's disease or Alzheimer's disease.

144. The compound or pharmaceutical composition of any one of claims 139-141, wherein the disease or disorder is cancer; optionally, the cancer is metastasising cancer, gastrointestinal cancer, brain cancer, skin cancer, non-small-cell lung carcinoma, head and neck squamous cell carcinoma or colorectal adenocarcinoma.

145. Use of the compound of any one of claims 1-128 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for inhibiting inflammasome activity; optionally, the inflammasome is NLRP3 inflammasome, and the activity is *in vitro* or *in vivo*.

146. Use of the compound of any one of claims 1-128 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing a disease or disorder.

147. The use of claim 146, wherein the disease or disorder is associated with an implicated inflammasome activity; optionally, the disease or disorder is a disease or disorder in which inflammasome activity is implicated.

148. The use of claim 146 or claim 147, wherein the disease or disorder is an inflammatory disorder, an autoinflammatory disorder, an autoimmune disorder, a neurodegenerative disease, or cancer.

149. The use of any one of claims 146-148, wherein the disease or disorder is an inflammatory disorder, an autoinflammatory disorder or an autoimmune disorder; optionally, the disease or disorder is selected from cryopyrin-associated autoinflammatory syndrome (CAPS; e.g., familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome/ neonatal-onset multisystem inflammatory disease (NOMID)), familial Mediterranean fever (FMF), nonalcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), gout, rheumatoid arthritis, osteoarthritis, Crohn's disease, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), fibrosis, obesity, type 2 diabetes, multiple sclerosis, dermatological diseases (e.g., acne) and neuroinflammation occurring in protein misfolding diseases (e.g., Prion diseases).

150. The use of any one of claims 146-148, wherein the disease or disorder is a neurodegenerative disease; optionally, the disease or disorder is Parkinson's disease or Alzheimer's disease.

151. The use of any one of claims 146-148 wherein the disease or disorder is a dermatological disease; optionally, the disease or disorder is acne.

152. The use of any one of claims 146-148, wherein the disease or disorder is cancer; optionally, the cancer is metastasising cancer, brain cancer, gastrointestinal cancer, skin cancer,

non-small-cell lung carcinoma, head and neck squamous cell carcinoma or colorectal adenocarcinoma.