

Substituted 5-phenyl-3,6-dihydro-2-oxo-6H-1,3,4-thiadiazines

5 BACKGROUND OF THE INVENTION

The invention had the object of finding novel compounds having valuable properties, in particular those which can be used for the preparation of medicaments.

10 The present invention relates to compounds and to the use of compounds in which the inhibition, regulation and/or modulation of signal transduction by kinases, in particular tyrosine kinases and/or serine/threonine kinases, plays a role, furthermore to pharmaceutical compositions which comprise
15 these compounds, and to the use of the compounds for the treatment of kinase-induced diseases.

In particular, the present invention relates to compounds and to the use of compounds in which the inhibition, regulation and/or modulation of signal
20 transduction by Met kinase plays a role.

One of the principal mechanisms by which cellular regulation is effected is through the transduction of extracellular signals across the membrane that in turn modulate biochemical pathways within the cell. Protein phosphoryl-
25 ation represents one course by which intracellular signals are propagated from molecule to molecule resulting finally in a cellular response. These signal transduction cascades are highly regulated and often overlap, as is evident from the existence of many protein kinases as well as phosphata-
30 ses. Phosphorylation of proteins occurs predominantly at serine, threonine or tyrosine residues, and protein kinases have therefore been classified by their specificity of phosphorylation site, i.e. serine/threonine kinases and tyrosine kinases. Since phosphorylation is such a ubiquitous process
35 within cells and since cellular phenotypes are largely influenced by the activity of these pathways, it is currently believed that a number of disease

states and/or diseases are attributable to either aberrant activation or functional mutations in the molecular components of kinase cascades. Consequently, considerable attention has been devoted to the characterisation of these proteins and compounds that are able to modulate their activity (for a review see: Weinstein-Oppenheim et al. *Pharma. & Therap.*, 2000, 88, 229-279).

The role of the receptor tyrosine kinase Met in human oncogenesis and the possibility of inhibition of HGF (hepatocyte growth factor) dependent Met activation are described by S. Berthou et al. in *Oncogene*, Vol. 23, No. 31, pages 5387-5393 (2004). The inhibitor SU11274 described therein, a pyrrole-indoline compound, is potentially suitable for combating cancer.

Another Met kinase inhibitor for cancer therapy is described by J.G. Christensen et al. in *Cancer Res.* 2003, 63(21), 7345-55.

A further tyrosine kinase inhibitor for combating cancer is reported by H. Hov et al. in *Clinical Cancer Research* Vol. 10, 6686-6694 (2004). The compound PHA-665752, an indole derivative, is directed against the HGF receptor c-Met. It is furthermore reported therein that HGF and Met make a considerable contribution to the malignant process of various forms of cancer, such as, for example, multiple myeloma.

The synthesis of small compounds which specifically inhibit, regulate and/or modulate signal transduction by tyrosine kinases and/or serine/-threonine kinases, in particular Met kinase, is therefore desirable and an aim of the present invention.

It has been found that the compounds according to the invention and salts thereof have very valuable pharmacological properties while being well tolerated.

The present invention specifically relates to compounds of the formula I which inhibit, regulate and/or modulate signal transduction by Met kinase,

to compositions which comprise these compounds, and to processes for the use thereof for the treatment of Met kinase-induced diseases and complaints, such as angiogenesis, cancer, tumour formation, growth and propagation, arteriosclerosis, ocular diseases, such as age-induced macular degeneration, choroidal neovascularisation and diabetic retinopathy, inflammatory diseases, arthritis, thrombosis, fibrosis, glomerulonephritis, neurodegeneration, psoriasis, restenosis, wound healing, transplant rejection, metabolic diseases and diseases of the immune system, also autoimmune diseases, cirrhosis, diabetes and diseases of the blood vessels, also instability and permeability and the like in mammals.

Solid tumours, in particular fast-growing tumours, can be treated with Met kinase inhibitors. These solid tumours include monocytic leukaemia, brain, urogenital, lymphatic system, stomach, laryngeal and lung carcinoma, including lung adenocarcinoma and small-cell lung carcinoma.

The present invention is directed to processes for the regulation, modulation or inhibition of Met kinase for the prevention and/or treatment of diseases in connection with unregulated or disturbed Met kinase activity. In particular, the compounds of the formula I can also be employed in the treatment of certain forms of cancer. The compounds of the formula I can furthermore be used to provide additive or synergistic effects in certain existing cancer chemotherapies, and/or can be used to restore the efficacy of certain existing cancer chemotherapies and radiotherapies.

The compounds of the formula I can furthermore be used for the isolation and investigation of the activity or expression of Met kinase. In addition, they are particularly suitable for use in diagnostic methods for diseases in connection with unregulated or disturbed Met kinase activity.

It can be shown that the compounds according to the invention have an antiproliferative action in vivo in a xenotransplant tumour model. The com-

pounds according to the invention are administered to a patient having a hyperproliferative disease, for example to inhibit tumour growth, to reduce inflammation associated with a lymphoproliferative disease, to inhibit transplant rejection or neurological damage due to tissue repair, etc. The present compounds are suitable for prophylactic or therapeutic purposes. As used herein, the term "treatment" is used to refer to both prevention of diseases and treatment of pre-existing conditions. The prevention of proliferation is achieved by administration of the compounds according to the invention prior to the development of overt disease, for example to prevent the growth of tumours, prevent metastatic growth, diminish restenosis associated with cardiovascular surgery, etc. Alternatively, the compounds are used for the treatment of ongoing diseases by stabilising or improving the clinical symptoms of the patient.

The host or patient can belong to any mammalian species, for example a primate species, particularly humans; rodents, including mice, rats and hamsters; rabbits; horses, cows, dogs, cats, etc. Animal models are of interest for experimental investigations, providing a model for treatment of human disease.

The susceptibility of a particular cell to treatment with the compounds according to the invention can be determined by in vitro tests. Typically, a culture of the cell is combined with a compound according to the invention at various concentrations for a period of time which is sufficient to allow the active agents to induce cell death or to inhibit migration, usually between about one hour and one week. In vitro testing can be carried out using cultivated cells from a biopsy sample. The viable cells remaining after the treatment are then counted.

The dose varies depending on the specific compound used, the specific disease, the patient status, etc. A therapeutic dose is typically sufficient considerably to reduce the undesired cell population in the target tissue while the viability of the patient is maintained. The treatment is generally

continued until a considerable reduction has occurred, for example an at least about 50% reduction in the cell burden, and may be continued until essentially no more undesired cells are detected in the body.

5

For identification of a signal transduction pathway and for detection of interactions between various signal transduction pathways, various scientists have developed suitable models or model systems, for example cell culture models (for example Khwaja et al., EMBO, 1997, 16, 2783-93) and models of transgenic animals (for example White et al., Oncogene, 2001, 20, 7064-7072). For the determination of certain stages in the signal transduction cascade, interacting compounds can be utilised in order to modulate the signal (for example Stephens et al., Biochemical J., 2000, 351, 95-105). The compounds according to the invention can also be used as reagents for testing kinase-dependent signal transduction pathways in animals and/or cell culture models or in the clinical diseases mentioned in this application.

20

Measurement of the kinase activity is a technique which is well known to the person skilled in the art. Generic test systems for the determination of the kinase activity using substrates, for example histone (for example Alessi et al., FEBS Lett. 1996, 399, 3, pages 333-338) or the basic myelin protein, are described in the literature (for example Campos-González, R. and Glenney, Jr., J.R. 1992, J. Biol. Chem. 267, page 14535).

25

For the identification of kinase inhibitors, various assay systems are available. In scintillation proximity assay (Sorg et al., J. of Biomolecular Screening, 2002, 7, 11-19) and flashplate assay, the radioactive phosphorylation of a protein or peptide as substrate with γ ATP is measured. In the presence of an inhibitory compound, a decreased radioactive signal, or none at all, is detectable. Furthermore, homogeneous time-resolved fluorescence resonance energy transfer (HTR-FRET) and fluorescence polari-

30

35

sation (FP) technologies are suitable as assay methods (Sills et al., J. of Biomolecular Screening, 2002, 191-214).

Other non-radioactive ELISA assay methods use specific phospho-anti-
bodies (phospho-ABs). The phospho-AB binds only the phosphorylated
substrate. This binding can be detected by chemiluminescence using a
second peroxidase-conjugated anti-sheep antibody (Ross et al., 2002,
Biochem. J.).

There are many diseases associated with deregulation of cellular proliferation and cell death (apoptosis). The conditions of interest include, but are not limited to, the following. The compounds according to the invention are suitable for the treatment of various conditions where there is proliferation and/or migration of smooth muscle cells and/or inflammatory cells into the intimal layer of a vessel, resulting in restricted blood flow through that vessel, for example in the case of neointimal occlusive lesions. Occlusive graft vascular diseases of interest include atherosclerosis, coronary vascular disease after grafting, vein graft stenosis, peri-anastomatic prosthetic restenosis, restenosis after angioplasty or stent placement, and the like.

PRIOR ART

Other thiadiazinones are described in WO 03/037349.


4,5-dihydropyrazoles for combating cancer are described in WO 03/079973 A2.

Quinoline derivatives are disclosed as Met kinase inhibitors in EP 1 411 046 A1.

Pyrrole-indoline derivatives are known as Met kinase inhibitors from WO 02/096361 A2.

SUMMARY OF THE INVENTION

The invention relates to compounds of the formula I



in which

R^1 denotes H, A, Hal, OH, OA, SH, SA, SOA, SO_2A , NO_2 , NH_2 , NHA, NAA' , SO_2NH_2 , SO_2NHA , SO_2NAA' , $CONH_2$, CONHA, $CONAA'$, $NACOA'$, $NASO_2A'$, COOH, COOA or CN,

R^2 denotes H, A, Hal, SO_2NH_2 , SO_2NHA , SO_2NAA' , $CONH_2$, CONHA, $CONAA'$, $NACOA'$, $NASO_2A'$, COOH, COOA or CN,

R^1 and R^2 together also denote methylenedioxy,

B denotes $NHCOO(CH_2)_nR^3$ or $NHCO(CH_2)_nR^3$,

Q is absent or denotes linear or branched alkylene having 1-4 C atoms,

R^3 denotes R^1 or Het,

alkyl having 1-6 C atoms or cycloalkyl having 3-8 C atoms, each of which is unsubstituted or mono-, di-, tri- or tetrasubstituted by R^4 ,

R^4 denotes A, Hal, OH, OA, SH, SA, SOA, SO_2A , NO_2 , NH_2 , NHA, NAA' , SO_2NH_2 , SO_2NHA , SO_2NAA' , $CONH_2$, CONHA, $CONAA'$, $NACOA'$, $NASO_2A'$, COOH, COOA or CN,

Het denotes a mono- or bicyclic saturated or aromatic heterocycle having 1 to 4 N, O and/or S atoms, which may be unsubstituted or mono-, di- or trisubstituted by R^4 , CHO, COA, =S, =NH, =NA and/or =O (carbonyl oxygen),

- A, A' each, independently of one another, denote unbranched or branched alkyl having 1-10 C atoms,
in which 1-7 H atoms may be replaced by F, Cl and/or Br,
cycloalkyl having 3-8 C atoms or
cycloalkylalkylene having 4-10 C atoms,
Hal denotes F, Cl, Br or I,
m denotes 0 or 1,
n denotes 0, 1, 2 or 3,
and pharmaceutically usable derivatives, solvates, salts, tautomers and stereoisomers thereof, including mixtures thereof in all ratios.
- The invention also relates to the optically active forms (stereoisomers), the enantiomers, the racemates, the diastereomers and the hydrates and solvates of these compounds. The term solvates of the compounds is taken to mean adductions of inert solvent molecules onto the compounds which form owing to their mutual attractive force. solvates are, for example, mono- or dihydrates or alkoxides.
- The term pharmaceutically usable derivatives is taken to mean, for example, the salts of the compounds according to the invention and also so-called prodrug compounds.
- The term prodrug derivatives is taken to mean compounds of the formula I which have been modified by means of, for example, alkyl or acyl groups, sugars or oligopeptides and which are rapidly cleaved in the organism to form the effective compounds according to the invention.
- These also include biodegradable polymer derivatives of the compounds according to the invention, as described, for example, in Int. J. Pharm. 115, 61-67 (1995).
- The expression "effective amount" denotes the amount of a medicament or of a pharmaceutical active ingredient which causes in a tissue, system,

animal or human a biological or medical response which is sought or desired, for example, by a researcher or physician.

In addition, the expression "therapeutically effective amount" denotes an amount which, compared with a corresponding subject who has not received this amount, has the following consequence:


improved treatment, healing, prevention or elimination of a disease, syndrome, condition, complaint, disorder or side-effects or also the reduction in the advance of a disease, complaint or disorder.

The expression "therapeutically effective amount" also encompasses the amounts which are effective for increasing normal physiological function.

The invention also relates to the use of mixtures of the compounds of the formula I, for example mixtures of two diastereomers, for example in the ratio 1:1, 1:2, 1:3, 1:4, 1:5, 1:10, 1:100 or 1:1000.

These are particularly preferably mixtures of stereoisomeric compounds.

The invention relates to the compounds of the formula I and salts thereof and to a process for the preparation of compounds of the formula I according to Claims 1-11 and pharmaceutically usable derivatives, salts, solvates and stereoisomers thereof, characterised in that a compound of the formula Ia



in which

B denotes NH₂,

and R¹, R², m and Q have the meanings indicated in Claim 1,

is converted into a compound of the formula I in which

B denotes $\text{NHCOO}(\text{CH}_2)_n\text{R}^3$,

5 by reacting a compound of the formula Ia with a coupling reagent selected from the group

- a) isopropylidene chloroformate,
- b) p-nitrophenyl chloroformate,
- c) diphosgene,
- 10 d) triphosgene,

and a compound of the formula II

15 $\text{HO}(\text{CH}_2)_n\text{R}^3$ II

in which n and R^3 have the meanings indicated in Claim 1,

20 and/or

a base or acid of the formula I is converted into one of its salts.

Above and below, the radicals R^1 , R^2 , Q and B have the meanings indicated for the formula I, unless expressly stated otherwise.

25

A or A' denotes alkyl, is unbranched (linear) or branched, and has 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 C atoms. A preferably denotes methyl, furthermore ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl, furthermore
30 also pentyl, 1-, 2- or 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-, 2-, 3- or 4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-, 2,3- or 3,3-dimethylbutyl, 1- or 2-ethylbutyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1,1,2- or 1,2,2-trimethylpropyl, furthermore preferably, for
35 example, trifluoromethyl.

A very particularly preferably denotes alkyl having 1, 2, 3, 4, 5 or 6 C atoms, preferably methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl,

tert-butyl, pentyl, hexyl, trifluoromethyl, pentafluoroethyl or 1,1,1-trifluoroethyl.

cycloalkyl preferably denotes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

Cycloalkylalkylene preferably denotes cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl or cycloheptylmethyl.

Irrespective of further substitutions, Het preferably denotes, for example, 2- or 3-furyl, 2- or 3-thienyl, 1-, 2- or 3-pyrrolyl, 1-, 2, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, furthermore preferably 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or 5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 3- or 4-pyridazinyl, pyrazinyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 4- or 5-isoindolyl, indazolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5-, 6- or 7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 4-, 5-, 6- or 7-benz-2,1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl, 3-, 4-, 5-, 6-, 7- or 8-cinnolinyl, 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, 5- or 6-quinoxaliny, 2-, 3-, 5-, 6-, 7- or 8-2H-benzo-1,4-oxazinyl, further preferably 1,3-benzodioxol-5-yl, 1,4-benzodioxan-6-yl, 2,1,3-benzothiadiazol-4-, -5-yl or 2,1,3-benzoxadiazol-5-yl or dibenzofuranyl.

The heterocyclic radicals may also be partially or fully hydrogenated.

Irrespective of further substitutions, Het can thus also denote, for example, 2,3-dihydro-2-, -3-, -4- or -5-furyl, 2,5-dihydro-2-, -3-, -4- or 5-furyl, tetrahydro-2- or -3-furyl, 1,3-dioxolan-4-yl, tetrahydro-2- or -3-thienyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 2,5-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 1-, 2- or 3-pyrrolidinyl, tetrahydro-1-, -2- or -4-imidazolyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrazolyl, tetrahydro-1-, -3- or -4-pyrazolyl, 1,4-dihydro-1-, -2-, -3- or -4-pyridyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1-,

2-, 3- or 4-piperidiny, 2-, 3- or 4-morpholinyl, tetrahydro-2-, -3- or -4-pyranyl, 1,4-dioxanyl, 1,3-dioxan-2-, -4- or -5-yl, hexahydro-1-, -3- or -4-pyridaziny, hexahydro-1-, -2-, -4- or -5-pyrimidiny, 1-, 2- or 3-piperaziny, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-quinolyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-isoquinolyl, 2-, 3-, 5-, 6-, 7- or 8- 3,4-dihydro-2H-benzo-1,4-oxazinyl, furthermore preferably 2,3-methylenedioxyphenyl, 3,4-methylenedioxyphenyl, 2,3-ethylenedioxyphenyl, 3,4-ethylenedioxyphenyl, 3,4-(difluoromethylenedioxy)phenyl, 2,3-dihydrobenzofuran-5- or 6-yl, 2,3-(2-oxomethylenedioxy)phenyl or also 3,4-dihydro-2H-1,5-benzodioxepin-6- or -7-yl, furthermore preferably 2,3-dihydrobenzofuranyl, 2,3-dihydro-2-oxofuranyl, 3,4-dihydro-2-oxo-1H-quinazoliny, 2,3-dihydrobenzoxazolyl, 2-oxo-2,3-dihydrobenzoxazolyl, 2,3-dihydrobenzimidazolyl, 1,3-dihydroindole, 2-oxo-1,3-dihydroindole or 2-oxo-2,3-dihydrobenzimidazolyl.

Het preferably denotes a monocyclic saturated or aromatic heterocycle having 1 to 2 N and/or O atoms, which may be mono- or disubstituted by A and/or =O (carbonyl oxygen).

Het particularly preferably denotes piperidiny, pyrrolidiny, morpholin-4-yl, piperaziny, 1,3-oxazolidin-3-yl, imidazolidiny, pyridyl, pyrimidiny, furanyl, thienyl, thiazolyl, indolyl or indazolyl, where the radicals may also be mono- or disubstituted by A.

R¹ preferably denotes H, Hal, A, OH, OA, SO₂A, CN, NO₂, COOA or COOH, in particular Cl, very particularly preferably 4-Cl.

R² preferably denotes H or Hal.

B preferably denotes NHCOO(CH₂)_nR³, furthermore NHCO(CH₂)_nR³.

The radical B is preferably in the meta-position to Q.

Q preferably denotes CH₂, CH(CH₃) or CH(CH₂CH₃).

R³ preferably denotes Het or alkyl having 1-6 C atoms or cycloalkyl having 3-8 C atoms, each of which is unsubstituted or mono-, di-, tri- or tetrasubstituted by R⁴.

R^3 particularly preferably denotes 2-hydroxyethyl, pyrrolidinyl, N-methyl-pyrrolidinyl-, morpholin-4-yl, 3-(N,N-diethylamino)propyl, 3-(N,N-dimethyl-amino)propyl, methyl, ethyl, propyl, pyridyl, 1-methylpiperazin-4-yl, piperazin-4-yl,

R^4 preferably denotes OH, amino, methylamino, dimethylamino, ethyl-amino, diethylamino, ter-butylamino, isopropylamino

Hal preferably denotes F, Cl or Br, but also I, particularly preferably F or Cl.

Throughout the invention, all radicals which occur more than once may be identical or different, i.e. are independent of one another.

The compounds of the formula I may have one or more chiral centres and can therefore occur in various stereoisomeric forms. The formula I encompasses all these forms.

Accordingly, the invention relates, in particular, to the compounds of the formula I in which at least one of the said radicals has one of the preferred meanings indicated above. Some preferred groups of compounds may be expressed by the following sub-formulae Ia to II, which conform to the formula I and in which the radicals not designated in greater detail have the meaning indicated for the formula I, but in which

in Ia R^1 denotes H, Hal, A, OH, OA, SO_2A , CN, NO_2 , COOA or COOH;

in Ib R^2 denotes H or Hal;

in Ic Het denotes a monocyclic saturated or aromatic heterocycle having 1 to 2 N and/or O atoms, which may be unsubstituted or mono- or disubstituted by A;

in Id Q denotes CH_2 , $CH(CH_3)$ or $CH(CH_2CH_3)$;

5	in le	R^3	denotes Het or alkyl having 1-6 C atoms or cycloalkyl having 3-8 C atoms, each of which is unsubstituted or mono-, di-, tri- or tetrasubstituted by R^4 ;
	in lf	R^4	denotes OH, NH_2 , NHA or NAA';
10	in lg	A, A'	each, independently of one another, denote unbranched or branched alkyl having 1-6 C atoms, in which 1-5 H atoms may be replaced by F and/or chlorine;
15	in lh	Het	denotes piperidinyl, pyrrolidinyl, morpholin-4-yl, piperazinyl, 1,3-oxazolidin-3-yl, imidazolidinyl, pyridyl, pyrimidinyl, furanyl, thienyl, thiazolyl, indolyl or indazolyl, where the radicals may also be mono- or disubstituted by A;
20	in li	R^1	denotes H, Hal, A, OH, OA, SO_2A , CN, NO_2 , COOA or COOH,
		R^2	denotes H or Hal,
		R^1 and R^2	together also denote methylenedioxy,
25		B	denotes $NHCOO(CH_2)_nR^3$ or $NHCO(CH_2)_nR^3$,
		Q	denotes CH_2 , $CH(CH_3)$ or $CH(CH_2CH_3)$,
		R^3	denotes Het or alkyl having 1-6 C atoms or cycloalkyl having 3-8 C atoms, each of which is unsubstituted or mono-, di-, tri- or tetrasubstituted by R^4 ,
30		R^4	denotes OH, NH_2 , NHA or NAA',
		A, A'	each, independently of one another, denote unbranched or branched alkyl having 1-6 C atoms, in which 1-5 H atoms may be replaced by F and/or chlorine,
35		Het	denotes a monocyclic saturated or aromatic heterocycle having 1 to 2 N and/or O atoms, which may be unsubstituted or mono- or disubstituted by A,

Hal denotes F, Cl, Br or I,

m denotes 0 or 1,

n denotes 0, 1, 2 or 3;

5 and pharmaceutically usable derivatives, salts, solvates, tautomers and stereoisomers thereof, including mixtures thereof in all ratios.

10 The compounds of the formula I and also the starting materials for their preparation are, in addition, prepared by methods known per se, as described in the literature (for example in the standard works, such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart), to be precise under reaction conditions which are known and suitable for the said reactions. Use can
15 also be made here of variants known per se which are not mentioned here in greater detail.

20 The starting compounds of the formulae Ia and II are generally known. If they are novel, however, they can be prepared by methods known per se.

Compounds of the formula I can preferably be obtained by reacting a compound of the formula Ia with a coupling reagent selected from the group
25

- a) isopropylidene chloroformate,
- b) p-nitrophenyl chloroformate,
- c) diphosgene,
- d) triphosgene,

30 and a compound of the formula II. The reaction is preferably carried out as a one-pot reaction.

The reaction is generally carried out in an inert solvent.
35 Depending on the conditions used, the reaction time is between a few minutes and 14 days, the reaction temperature is between about -15° and

150°, normally between -5° and 90°, particularly preferably between 20° and 60°C.

5 Examples of suitable inert solvents are hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichloroethylene, 1,2-dichloroethane, carbon tetrachloride, chloroform or dichloromethane; alcohols, such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol; ethers, such as diethyl ether,
10 diisopropyl ether, tetrahydrofuran (THF) or dioxane; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether, ethylene glycol dimethyl ether (diglyme); ketones, such as acetone or butanone; amides, such as acetamide, dimethylacetamide or dimethylformamide (DMF); nitriles, such
15 as acetonitrile; sulfoxides, such as dimethyl sulfoxide (DMSO); carbon disulfide; carboxylic acids, such as formic acid or acetic acid; nitro compounds, such as nitromethane or nitrobenzene; esters, such as ethyl acetate, or mixtures of the said solvents.

20 Particular preference is given to THF, dichloromethane and/or DMF.

The reaction is generally carried out in the presence of an acid-binding agent, preferably an organic base, such as DIPEA, triethylamine, dimethylaniline, pyridine or quinoline.
25 The addition of an alkali or alkaline-earth metal hydroxide, carbonate or bicarbonate or of another salt of a weak acid of the alkali or alkaline-earth metals, preferably of potassium, sodium, calcium or caesium, may also be favourable.

30 Pharmaceutical salts and other forms

The said compounds according to the invention can be used in their final non-salt form. On the other hand, the present invention also encompasses
35 the use of these compounds in the form of their pharmaceutically acceptable salts, which can be derived from various organic and inorganic acids and bases by procedures known in the art. Pharmaceutically acceptable

salt forms of the compounds of the formula I are for the most part prepared by conventional methods. If the compound of the formula I contains a carboxyl group, one of its suitable salts can be formed by reacting the compound with a suitable base to give the corresponding base-addition salt. Such bases are, for example, alkali metal hydroxides, including potassium hydroxide, sodium hydroxide and lithium hydroxide; alkaline-earth metal hydroxides, such as barium hydroxide and calcium hydroxide; alkali metal alkoxides, for example potassium ethoxide and sodium propoxide; and various organic bases, such as piperidine, diethanolamine and N-methylglutamine. The aluminium salts of the compounds of the formula I are likewise included. In the case of certain compounds of the formula I, acid-addition salts can be formed by treating these compounds with pharmaceutically acceptable organic and inorganic acids, for example hydrogen halides, such as hydrogen chloride, hydrogen bromide or hydrogen iodide, other mineral acids and corresponding salts thereof, such as sulfate, nitrate or phosphate and the like, and alkyl- and monoarylsulfonates, such as ethanesulfonate, toluenesulfonate and benzenesulfonate, and other organic acids and corresponding salts thereof, such as acetate, trifluoroacetate, tartrate, maleate, succinate, citrate, benzoate, salicylate, ascorbate and the like. Accordingly, pharmaceutically acceptable acid-addition salts of the compounds of the formula I include the following: acetate, adipate, alginate, arginate, aspartate, benzoate, benzenesulfonate (besylate), bisulfate, bisulfite, bromide, butyrate, camphorate, camphorsulfonate, caprylate, chloride, chlorobenzoate, citrate, cyclopentanepropionate, digluconate, dihydrogenphosphate, dinitrobenzoate, dodecylsulfate, ethanesulfonate, fumarate, galacterate (from mucic acid), galacturonate, glucoheptanoate, gluconate, glutamate, glycerophosphate, hemisuccinate, hemisulfate, heptanoate, hexanoate, hippurate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, iodide, isethionate, isobutyrate, lactate, lactobionate, malate, maleate, malonate, mandelate, metaphosphate, methanesulfonate, methylbenzoate, monohydrogenphosphate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, oleate, palmo-

ate, pectinate, persulfate, phenylacetate, 3-phenylpropionate, phosphate, phosphonate, phthalate, but this does not represent a restriction.

5 Furthermore, the base salts of the compounds according to the invention include aluminium, ammonium, calcium, copper, iron(III), iron(II), lithium, magnesium, manganese(III), manganese(II), potassium, sodium and zinc salts, but this is not intended to represent a restriction. Of the above-men-
10 tioned salts, preference is given to ammonium; the alkali metal salts sodium and potassium, and the alkaline-earth metal salts calcium and magnesium. Salts of the compounds of the formula I which are derived from pharmaceutically acceptable organic non-toxic bases include salts of
15 primary, secondary and tertiary amines, substituted amines, also including naturally occurring substituted amines, cyclic amines, and basic ion ex- changer resins, for example arginine, betaine, caffeine, chlorprocaine, choline, N,N'-dibenzylethylenediamine (benzathine), dicyclohexylamine, diethanolamine, diethylamine, 2-diethylaminoethanol, 2-dimethylamino-
20 ethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpipe- ridine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lidocaine, lysine, meglumine, N-methyl-D-glucamine, morpholine, piper- azine, piperidine, polyamine resins, procaine, purines, theobromine, tri- ethanolamine, triethylamine, trimethylamine, tripropylamine and tris-
25 (hydroxymethyl)methylamine (tromethamine), but this is not intended to represent a restriction.

30 Compounds of the present invention which contain basic nitrogen-contain- ing groups can be quaternised using agents such as (C₁-C₄)alkyl halides, for example methyl, ethyl, isopropyl and tert-butyl chloride, bromide and iodide; di(C₁-C₄)alkyl sulfates, for example dimethyl, diethyl and diamyl sulfate; (C₁₀-C₁₈)alkyl halides, for example decyl, dodecyl, lauryl, myristyl
35 and stearyl chloride, bromide and iodide; and aryl(C₁-C₄)alkyl halides, for example benzyl chloride and phenethyl bromide. Both water- and oil-solu-

ble compounds according to the invention can be prepared using such salts.

5 The above-mentioned pharmaceutical salts which are preferred include acetate, trifluoroacetate, besylate, citrate, fumarate, gluconate, hemisuccinate, hippurate, hydrochloride, hydrobromide, isethionate, mandelate, meglumine, nitrate, oleate, phosphonate, pivalate, sodium phosphate, stearate, sulfate, sulfosalicylate, tartrate, thiomalate, tosylate and trometh-
10 amine, but this is not intended to represent a restriction.

The acid-addition salts of basic compounds of the formula I are prepared by bringing the free base form into contact with a sufficient amount of the
15 desired acid, causing the formation of the salt in a conventional manner. The free base can be regenerated by bringing the salt form into contact with a base and isolating the free base in a conventional manner. The free base forms differ in a certain respect from the corresponding salt forms thereof with respect to certain physical properties, such as solubility in
20 polar solvents; for the purposes of the invention, however, the salts otherwise correspond to the respective free base forms thereof.

25 As mentioned, the pharmaceutically acceptable base-addition salts of the compounds of the formula I are formed with metals or amines, such as alkali metals and alkaline-earth metals or organic amines. Preferred metals are sodium, potassium, magnesium and calcium. Preferred organic amines are N,N'-dibenzylethylenediamine, chloroprocaine, choline, di-
30 ethanolamine, ethylenediamine, N-methyl-D-glucamine and procaine.

The base-addition salts of acidic compounds according to the invention are prepared by bringing the free acid form into contact with a sufficient
35 amount of the desired base, causing the formation of the salt in a conventional manner. The free acid can be regenerated by bringing the salt form into contact with an acid and isolating the free acid in a conventional man-

ner. The free acid forms differ in a certain respect from the corresponding salt forms thereof with respect to certain physical properties, such as solubility in polar solvents; for the purposes of the invention, however, the salts otherwise correspond to the respective free acid forms thereof.

If a compound according to the invention contains more than one group which is capable of forming pharmaceutically acceptable salts of this type, the invention also encompasses multiple salts. Typical multiple salt forms include, for example, bitartrate, diacetate, difumarate, dimeglumine, diphosphate, disodium and trihydrochloride, but this is not intended to represent a restriction.

With regard to that stated above, it can be seen that the expression "pharmaceutically acceptable salt" in the present connection is taken to mean an active ingredient which comprises a compound of the formula I in the form of one of its salts, in particular if this salt form imparts improved pharmacokinetic properties on the active ingredient compared with the free form of the active ingredient or any other salt form of the active ingredient used earlier. The pharmaceutically acceptable salt form of the active ingredient can also provide this active ingredient for the first time with a desired pharmacokinetic property which it did not have earlier and can even have a positive influence on the pharmacodynamics of this active ingredient with respect to its therapeutic efficacy in the body.

The invention furthermore relates to medicaments comprising at least one compound of the formula I and/or pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios, and optionally excipients and/or adjuvants.

Pharmaceutical formulations can be administered in the form of dosage units which comprise a predetermined amount of active ingredient per dosage unit. Such a unit can comprise, for example, 0.5 mg to 1 g, prefer-

ably 1 mg to 700 mg, particularly preferably 5 mg to 100 mg, of a compound according to the invention, depending on the condition treated, the method of administration and the age, weight and condition of the patient, or pharmaceutical formulations can be administered in the form of dosage units which comprise a predetermined amount of active ingredient per dosage unit. Preferred dosage unit formulations are those which comprise a daily dose or part-dose, as indicated above, or a corresponding fraction thereof of an active ingredient. Furthermore, pharmaceutical formulations of this type can be prepared using a process which is generally known in the pharmaceutical art.

Pharmaceutical formulations can be adapted for administration via any desired suitable method, for example by oral (including buccal or sublingual), rectal, nasal, topical (including buccal, sublingual or transdermal), vaginal or parenteral (including subcutaneous, intramuscular, intravenous or intradermal) methods. Such formulations can be prepared using all processes known in the pharmaceutical art by, for example, combining the active ingredient with the excipient(s) or adjuvant(s).

Pharmaceutical formulations adapted for oral administration can be administered as separate units, such as, for example, capsules or tablets; powders or granules; solutions or suspensions in aqueous or non-aqueous liquids; edible foams or foam foods; or oil-in-water liquid emulsions or water-in-oil liquid emulsions.

Thus, for example, in the case of oral administration in the form of a tablet or capsule, the active-ingredient component can be combined with an oral, non-toxic and pharmaceutically acceptable inert excipient, such as, for example, ethanol, glycerol, water and the like. Powders are prepared by comminuting the compound to a suitable fine size and mixing it with a pharmaceutical excipient comminuted in a similar manner, such as, for

example, an edible carbohydrate, such as, for example, starch or mannitol. A flavour, preservative, dispersant and dye may likewise be present.

5 Capsules are produced by preparing a powder mixture as described above and filling shaped gelatine shells therewith. Glidants and lubricants, such as, for example, highly disperse silicic acid, talc, magnesium stearate, calcium stearate or polyethylene glycol in solid form, can be added to the powder mixture before the filling operation. A disintegrant or solubiliser,
10 such as, for example, agar-agar, calcium carbonate or sodium carbonate, may likewise be added in order to improve the availability of the medication after the capsule has been taken.

15 In addition, if desired or necessary, suitable binders, lubricants and disintegrants as well as dyes can likewise be incorporated into the mixture. Suitable binders include starch, gelatine, natural sugars, such as, for example, glucose or beta-lactose, sweeteners made from maize, natural
20 and synthetic rubber, such as, for example, acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. The lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. The disintegrants include, without being restricted
25 thereto, starch, methylcellulose, agar, bentonite, xanthan gum and the like. The tablets are formulated by, for example, preparing a powder mixture, granulating or dry-pressing the mixture, adding a lubricant and a disintegrant and pressing the entire mixture to give tablets. A powder mixture is
30 prepared by mixing the compound comminuted in a suitable manner with a diluent or a base, as described above, and optionally with a binder, such as, for example, carboxymethylcellulose, an alginate, gelatine or polyvinylpyrrolidone, a dissolution retardant, such as, for example, paraffin, an absorption accelerator, such as, for example, a quaternary salt, and/or an
35 absorbant, such as, for example, bentonite, kaolin or dicalcium phosphate. The powder mixture can be granulated by wetting it with a binder, such as,

for example, syrup, starch paste, acacia mucilage or solutions of cellulose or polymer materials and pressing it through a sieve. As an alternative to granulation, the powder mixture can be run through a tableting machine, giving lumps of non-uniform shape, which are broken up to form granules. The granules can be lubricated by addition of stearic acid, a stearate salt, talc or mineral oil in order to prevent sticking to the tablet casting moulds. The lubricated mixture is then pressed to give tablets. The compounds according to the invention can also be combined with a free-flowing inert excipient and then pressed directly to give tablets without carrying out the granulation or dry-pressing steps. A transparent or opaque protective layer consisting of a shellac sealing layer, a layer of sugar or polymer material and a gloss layer of wax may be present. Dyes can be added to these coatings in order to be able to differentiate between different dosage units.

Oral liquids, such as, for example, solution, syrups and elixirs, can be prepared in the form of dosage units so that a given quantity comprises a pre-specified amount of the compound. Syrups can be prepared by dissolving the compound in an aqueous solution with a suitable flavour, while elixirs are prepared using a non-toxic alcoholic vehicle. Suspensions can be formulated by dispersion of the compound in a non-toxic vehicle. Solubilisers and emulsifiers, such as, for example, ethoxylated isostearyl alcohols and polyoxyethylene sorbitol ethers, preservatives, flavour additives, such as, for example, peppermint oil or natural sweeteners or saccharin, or other artificial sweeteners and the like, can likewise be added.

The dosage unit formulations for oral administration can, if desired, be encapsulated in microcapsules. The formulation can also be prepared in such a way that the release is extended or retarded, such as, for example, by coating or embedding of particulate material in polymers, wax and the like.

The compounds of the formula I and salts, solvates and physiologically functional derivatives thereof can also be administered in the form of liposome delivery systems, such as, for example, small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from various phospholipids, such as, for example, cholesterol, stearylamine or phosphatidylcholines.

The compounds of the formula I and the salts, solvates and physiologically functional derivatives thereof can also be delivered using monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds can also be coupled to soluble polymers as targeted medicament carriers. Such polymers may encompass polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidophenol, polyhydroxyethylaspartamidophenol or polyethylene oxide polylysine, substituted by palmitoyl radicals. The compounds may furthermore be coupled to a class of biodegradable polymers which are suitable for achieving controlled release of a medicament, for example polylactic acid, poly-epsilon-caprolactone, polyhydroxybutyric acid, polyorthoesters, polyacetals, polydihydroxypyranes, polycyanoacrylates and crosslinked or amphipathic block copolymers of hydrogels.

Pharmaceutical formulations adapted for transdermal administration can be administered as independent plasters for extended, close contact with the epidermis of the recipient. Thus, for example, the active ingredient can be delivered from the plaster by iontophoresis, as described in general terms in *Pharmaceutical Research*, 3(6), 318 (1986).

Pharmaceutical compounds adapted for topical administration can be formulated as ointments, creams, suspensions, lotions, powders, solutions, pastes, gels, sprays, aerosols or oils.

For the treatment of the eye or other external tissue, for example mouth and skin, the formulations are preferably applied as topical ointment or cream. In the case of formulation to give an ointment, the active ingredient can be employed either with a paraffinic or a water-miscible cream base. Alternatively, the active ingredient can be formulated to give a cream with an oil-in-water cream base or a water-in-oil base.

Pharmaceutical formulations adapted for topical application to the eye include eye drops, in which the active ingredient is dissolved or suspended in a suitable carrier, in particular an aqueous solvent.

Pharmaceutical formulations adapted for topical application in the mouth encompass lozenges, pastilles and mouthwashes.

Pharmaceutical formulations adapted for rectal administration can be administered in the form of suppositories or enemas.

Pharmaceutical formulations adapted for nasal administration in which the carrier substance is a solid comprise a coarse powder having a particle size, for example, in the range 20-500 microns, which is administered in the manner in which snuff is taken, i.e. by rapid inhalation via the nasal passages from a container containing the powder held close to the nose. Suitable formulations for administration as nasal spray or nose drops with a liquid as carrier substance encompass active-ingredient solutions in water or oil.

Pharmaceutical formulations adapted for administration by inhalation encompass finely particulate dusts or mists, which can be generated by various types of pressurised dispensers with aerosols, nebulisers or insufflators.

Pharmaceutical formulations adapted for vaginal administration can be administered as pessaries, tampons, creams, gels, pastes, foams or spray formulations.

5

Pharmaceutical formulations adapted for parenteral administration include aqueous and non-aqueous sterile injection solutions comprising antioxidants, buffers, bacteriostatics and solutes, by means of which the formulation is rendered isotonic with the blood of the recipient to be treated; and
10 aqueous and non-aqueous sterile suspensions, which may comprise suspension media and thickeners. The formulations can be administered in single-dose or multidose containers, for example sealed ampoules and vials, and stored in freeze-dried (lyophilised) state, so that only the addition
15 of the sterile carrier liquid, for example water for injection purposes, immediately before use is necessary. Injection solutions and suspensions prepared in accordance with the recipe can be prepared from sterile powders, granules and tablets.

20

It goes without saying that, in addition to the above particularly mentioned constituents, the formulations may also comprise other agents usual in the art with respect to the particular type of formulation; thus, for example, formulations which are suitable for oral administration may comprise flavours.
25

A therapeutically effective amount of a compound of the formula I depends on a number of factors, including, for example, the age and weight of the animal, the precise condition that requires treatment, and its severity, the
30 nature of the formulation and the method of administration, and is ultimately determined by the treating doctor or vet. However, an effective amount of a compound according to the invention for the treatment of neoplastic growth, for example colon or breast carcinoma, is generally in the range
35 from 0.1 to 100 mg/kg of body weight of the recipient (mammal) per day and particularly typically in the range from 1 to 10 mg/kg of body weight per day. Thus, the actual amount per day for an adult mammal weighing

70 kg is usually between 70 and 700 mg, where this amount can be administered as a single dose per day or usually in a series of part-doses (such as, for example, two, three, four, five or six) per day, so that the total daily dose is the same. An effective amount of a salt or solvate or of a physiologically functional derivative thereof can be determined as the fraction of the effective amount of the compound according to the invention *per se*. It can be assumed that similar doses are suitable for the treatment of other conditions mentioned above.

The invention furthermore relates to medicaments comprising at least one compound of the formula I and/or pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios, and at least one further medicament active ingredient.

The invention also relates to a set (kit) consisting of separate packs of

- (a) an effective amount of a compound of the formula I and/or pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios, and
- (b) an effective amount of a further medicament active ingredient.

The set comprises suitable containers, such as boxes, individual bottles, bags or ampoules. The set may, for example, comprise separate ampoules, each containing an effective amount of a compound of the formula I and/or pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios, and an effective amount of a further medicament active ingredient in dissolved or lyophilised form.

USE

5 The present compounds are suitable as pharmaceutical active ingredients for mammals, especially for humans, in the treatment of tyrosine kinase-induced diseases. These diseases include the proliferation of tumour cells, pathological neovascularisation (or angiogenesis) which promotes the growth of solid tumours, ocular neovascularisation (diabetic retinopathy, age-induced macular degeneration and the like) and inflammation (psoriasis, rheumatoid arthritis and the like).

15 The present invention encompasses the use of the compounds of the formula I and/or physiologically acceptable salts and solvates thereof for the preparation of a medicament for the treatment or prevention of cancer. Preferred carcinomas for the treatment originate from the group cerebral carcinoma, urogenital tract carcinoma, carcinoma of the lymphatic system, stomach carcinoma, laryngeal carcinoma and lung carcinoma. A further group of preferred forms of cancer are monocytic leukaemia, lung adenocarcinoma, small-cell lung carcinomas, pancreatic cancer, glioblastomas and breast carcinoma.

20 Also encompassed is the use of the compounds according to Claim 1 according to the invention and/or physiologically acceptable salts and solvates thereof for the preparation of a medicament for the treatment or prevention of a disease in which angiogenesis is implicated.

30 Such a disease in which angiogenesis is implicated is an ocular disease, such as retinal vascularisation, diabetic retinopathy, age-induced macular degeneration and the like.

35 The use of compounds of the formula I and/or physiologically acceptable salts and solvates thereof for the preparation of a medicament for the treatment or prevention of inflammatory diseases also falls within the scope of the present invention. Examples of such inflammatory diseases include rheumatoid arthritis, psoriasis, contact dermatitis, delayed hypersensitivity reaction and the like.

Also encompassed is the use of the compounds of the formula I and/or physiologically acceptable salts and solvates thereof for the preparation of a medicament for the treatment or prevention of a tyrosine kinase-induced disease or a tyrosine kinase-induced condition in a mammal, in which to this method a therapeutically effective amount of a compound according to the invention is administered to a sick mammal in need of such treatment. The therapeutic amount varies according to the specific disease and can be determined by the person skilled in the art without undue effort.

The present invention also encompasses the use compounds of the formula I and/or physiologically acceptable salts and solvates thereof for the preparation of a medicament for the treatment or prevention of retinal vascularisation.

Methods for the treatment or prevention of ocular diseases, such as diabetic retinopathy and age-induced macular degeneration, are likewise part of the invention. The use for the treatment or prevention of inflammatory diseases, such as rheumatoid arthritis, psoriasis, contact dermatitis and delayed hypersensitivity reaction, as well as the treatment or prevention of bone pathologies from the group osteosarcoma, osteoarthritis and rickets, likewise falls within the scope of the present invention.

The expression "tyrosine kinase-induced diseases or conditions" refers to pathological conditions that depend on the activity of one or more tyrosine kinases. Tyrosine kinases either directly or indirectly participate in the signal transduction pathways of a variety of cellular activities, including proliferation, adhesion and migration and differentiation. Diseases associated with tyrosine kinase activity include proliferation of tumour cells, pathological neovascularisation that promotes the growth of solid tumours, ocular neovascularisation (diabetic retinopathy, age-induced macular degeneration and the like) and inflammation (psoriasis, rheumatoid arthritis and the like).

The compounds of the formula I can be administered to patients for the treatment of cancer, in particular fast-growing tumours.

5 The invention thus relates to the use of compounds of the formula I, and pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios, for the preparation of a medicament for the treatment of diseases in which the inhibition, regulation and/or modulation of kinase signal transduction plays a role.

10 Preference is given here to Met kinase.

Preference is given to the use of compounds of the formula I, and pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios,
15 for the preparation of a medicament for the treatment of diseases which are influenced by inhibition of tyrosine kinases by the compounds according to Claim 1.

20 Particular preference is given to the use for the preparation of a medicament for the treatment of diseases which are influenced by inhibition of Met kinase by the compounds according to Claim 1.

Especial preference is given to the use for the treatment of a disease where the disease is a solid tumour.
25

The solid tumour is preferably selected from the group of tumours of the lung, squamous epithelium, the bladder, the stomach, the kidneys, of head and neck, the oesophagus, the cervix, the thyroid, the intestine, the liver,
30 the brain, the prostate, the urogenital tract, the lymphatic system, the stomach and/or the larynx.

The solid tumour is furthermore preferably selected from the group lung adenocarcinoma, small-cell lung carcinomas, pancreatic cancer, glioblastomas, colon carcinoma and breast carcinoma.
35

5 Preference is furthermore given to the use for the treatment of a tumour of the blood and immune system, preferably for the treatment of a tumour selected from the group of acute myeloid leukaemia, chronic myeloid leukaemia, acute lymphatic leukaemia and/or chronic lymphatic leukaemia.

10 The disclosed compounds of the formula I can be administered in combination with other known therapeutic agents, including anticancer agents. As used here, the term "anticancer agent" relates to any agent which is administered to a patient with cancer for the purposes of treating the cancer.

15 The anti-cancer treatment defined herein may be applied as a sole therapy or may involve, in addition to the compound of the invention, conventional surgery or radiotherapy or chemotherapy. Such chemotherapy may include one or more of the following categories of anti- tumour agents:

- 20 (i) antiproliferative/antineoplastic/DNA-damaging agents and combinations thereof, as used in medical oncology, such as alkylating agents (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chloroambucil, busulphan and nitrosoureas); antimetabolites (for example antifolates such as fluoropyrimidines like 5-fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside, hydroxyurea and gemcitabine); antitumour antibiotics (for example anthracyclines, such as adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin) ; antimitotic agents (for example vinca alkaloids, like vincristine, vinblastine, vindesine and vinorelbine, and taxoids, like taxol and taxotere) ; topoisomerase inhibitors (for example epipodophyllotoxins, like etoposide and teniposide, amsacrine, topotecan, irinotecan and camptothecin) and cell-differentiating agents (for example all-trans-retinoic acid, 13-cis-retinoic acid and fenretinide);
- 35 (ii) cytostatic agents, such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene and idoxifene), oestrogen receptor downregulators (for example fulvestrant), antiandrogens (for example bi-

calutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example goserelin, leuprorelin and buserelin), progesterones (for example megestrol acetate), aromatase inhibitors (for example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 α -reductase, such as finasteride;

(iii) agents which inhibit cancer cell invasion (for example metalloproteinase inhibitors, like marimastat, and inhibitors of urokinase plasminogen activator receptor function);

(iv) inhibitors of growth factor function, for example such inhibitors include growth factor antibodies, growth factor receptor antibodies (for example the anti-erbB2 antibody trastuzumab [HerceptinTM] and the anti-erbB1 antibody cetuximab [C225]), farnesyl transferase inhibitors, tyrosine kinase inhibitors and serine/threonine kinase inhibitors, for example inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors, such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)-quinazolin-4-amine (CI 1033)), for example inhibitors of the platelet-derived growth factor family and for example inhibitors of the hepatocyte growth factor family;

(v) antiangiogenic agents, such as those which inhibit the effects of vascular endothelial growth factor, (for example the anti-vascular endothelial cell growth factor antibody bevacizumab [AvastinTM], compounds such as those disclosed in published international patent applications

WO 97/22596, WO 97/30035, WO 97/32856 and WO 98/13354) and compounds that work by other mechanisms (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function and angiostatin);

(vi) vessel-damaging agents, such as combretastatin A4 and compounds disclosed in international patent applications WO 99/02166,

WO 00/40529, WO 00/41669, WO 01/92224, WO 02/04434 and
WO 02/08213;

- (vii) antisense therapies, for example those which are directed to the
5 targets listed above, such as ISIS 2503, an anti-Ras antisense;
- (viii) gene therapy approaches, including, for example, approaches for
replacement of aberrant genes, such as aberrant p53 or aberrant BRCA1
or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches,
10 such as those using cytosine deaminase, thymidine kinase or a bacterial
nitroreductase enzyme, and approaches for increasing patient tolerance to
chemotherapy or radiotherapy, such as multi-drug resistance gene ther-
apy; and
- (ix) immunotherapy approaches, including, for example, ex-vivo and
15 in-vivo approaches for increasing the immunogenicity of patient tumour
cells, such as transfection with cytokines, such as interleukin 2, interleukin
4 or granulocyte-macrophage colony stimulating factor, approaches for
decreasing T-cell anergy, approaches using transfected immune cells,
20 such as cytokine-transfected dendritic cells, approaches using cytokine-
transfected tumour cell lines, and approaches using anti-idiotypic anti-
bodies.

25 The medicaments from Table 1 below are preferably, but not exclusively,
combined with the compounds of the formula I.

	Table 1.		
30	alkylating agents	cyclophosphamide Busulfan Ifosfamide Melfhalan Hexamethylmelamine Thiotepa chloroambucil Dacarbazine Carmustine	Lomustine Procarbazine Altretamine Estramustine phosphate Mechloroethamine Streptozocin Temozolomide Semustine
35	platinum agents	Cisplatin	Carboplatin

5		Oxaliplatin Spiroplatin Carboxyphthalatoplatinum Tetraplatin Ormiplatin Iproplatin	ZD-0473 (AnorMED) Lobaplatin (Aetema) Satraplatin (Johnson Matthey) BBR-3464 (Hoffmann-La Roche) SM-11355 (Sumitomo) AP-5280 (Access)
10	Antimetabolites	Azacytidine Gemcitabine Capecitabine 5-fluorouracil Floxuridine 2-chlorodesoxyadenosine 6-Mercaptopurine 6-Thioguanine Cytarabine 2-fluorodesoxycytidine Methotrexate Idatrexate	Tomudex Trimetrexate Deoxycoformycin Fludarabine Pentostatin Raltitrexed Hydroxyurea Decitabine (SuperGen) Clofarabine (Bioenvision) Irofulven (MGI Pharma) DMDC (Hoffmann-La Roche) Ethinylcytidine (Taiho)
20	Topoisomerase inhibitors	Amsacrine Epirubicin Etoposide Teniposide or mitoxantrone Irinotecan (CPT-11) 7-Ethyl-10- hydroxycamptothecin Topotecan Dexrazoxanet (TopoTarget) Pixantrone (Novuspharma) Rebeccamycin analogue (Exelixis) BBR-3576 (Novuspharma)	Rubitecan (SuperGen) Exatecan mesylate (Daiichi) Quinamed (ChemGenex) Gimatecan (Sigma- Tau) Diflomotecan (Beaufour- Ipsen) TAS-103 (Taiho) Elsamitrucin (Spectrum) J-107088 (Merck & Co) BNP-1350 (BioNumerik) CKD-602 (Chong Kun Dang) KW-2170 (Kyowa Hakko)
35	Antitumour antibiotics	Dactinomycin (Actinomycin D) Doxorubicin (Adriamycin) Deoxyrubicin Valrubicin Daunorubicin (Daunomycin) Epirubicin	Amonafide Azonafide Anthrapyrazole Oxantrazole Losoantrone Bleomycin sulfate (Blenoxan) Bleomycinic acid

5		Therarubicin Idarubicin Rubidazon Plicamycinp Porfiromycin Cyanomorpholinodoxo- rubicin Mitoxantron (Novantron)	Bleomycin A Bleomycin B Mitomycin C MEN-10755 (Menarini) GPX-100 (Gem Pharmaceuticals)
10	Antimitotic agents	Paclitaxel Docetaxel Colchicine Vinblastine Vincristine Vinorelbine Vindesine Dolastatin 10 (NCI) Rhizoxin (Fujisawa) Mivobulin (Warner- Lambert) Cemadotin (BASF) RPR 109881A (Aventis) TXD 258 (Aventis) Epothilone B (Novartis) T 900607 (Tularik) T 138067 (Tularik) Cryptophycin 52 (Eli Lilly) Vinflunine (Fabre) Auristatin PE (Teikoku Hormone) BMS 247550 (BMS) BMS 184476 (BMS) BMS 188797 (BMS) Taxoprexin (Protarga)	SB 408075 (GlaxoSmithKline) E7010 (Abbott) PG-TXL (Cell Therapeutics) IDN 5109 (Bayer) A 105972 (Abbott) A 204197 (Abbott) LU 223651 (BASF) D 24851 (ASTA Medica) ER-86526 (Eisai) Combretastatin A4 (BMS) Isohomohalichondrin-B (PharmaMar) ZD 6126 (AstraZeneca) PEG-Paclitaxel (Enzon) AZ10992 (Asahi) IDN-5109 (Indena) AVLB (Prescient NeuroPharma) Azaepothilon B (BMS) BNP- 7787 (BioNumerik) CA-4-Prodrug (OXIGENE) Dolastatin-10 (NrH) CA-4 (OXIGENE)
30	Aromatase inhibitors	Aminoglutethimide Letrozole Anastrozole Formestan	Exemestan Atamestan (BioMedicines) YM-511 (Yamanouchi)
35	Thymidylate synthase inhibitors	Pemetrexed (Eli Lilly) ZD-9331 (BTG)	Nolatrexed (Eximias) CoFactor™ (BioKeys)
	DNA antagonists	Trabectedin (PharmaMar) Glufosfamide (Baxter International)	Mafosfamide (Baxter International) Apaziquone (Spectrum)

	Albumin + ³² P (Isotope Solutions) Thymectacin (NewBiotics) Edotreotid (Novartis)	Pharmaceuticals) O6-Benzylguanine (Paligent)
5	Farnesyl transferase inhibitors Arglabin (NuOncology Labs) lonafarnib (Schering-Plough) BAY-43-9006 (Bayer)	Tipifarnib (Johnson & Johnson) Perillyl alcohol (DOR BioPharma)
10	Pump inhibitors CBT-1 (CBA Pharma) Tariquidar (Xenova) MS-209 (Schering AG)	Zosuquidar trihydrochloride (Eli Lilly) Biricodar dicitrate (Vertex)
15	Histone acetyl transferase inhibitors Tacedinaline (Pfizer) SAHA (Aton Pharma) MS-275 (Schering AG)	Pivaloyloxymethyl butyrate (Titan) Depsipeptide (Fujisawa)
20	Metalloproteinase inhibitors Ribonucleoside reductase inhibitors Neovastat (Aeterna Laboratories) Marimastat (British Biotech) Gallium maltolate (Titan) Triapin (Vion)	CMT -3 (CollaGenex) BMS-275291 (Celltech) Tezacitabine (Aventis) Didox (Molecules for Health)
	TNF-alpha agonists/antagonists Virulizin (Lorus Therapeutics) CDC-394 (Celgene)	Revimid (Celgene)
25	Endothelin-A receptor antagonists Atrasentan (Abbot) ZD-4054 (AstraZeneca)	YM-598 (Yamanouchi)
	Retinoic acid receptor agonists Fenretinide (Johnson & Johnson) LGD-1550 (Ligand)	Alitretinoin (Ligand)
30	Immunomodulators Interferon Oncophage (Antigenics) GMK (Progenics) Adenocarcinoma vaccine (Biomira) CTP-37 (AVI BioPharma) JRX-2 (Immuno-Rx) PEP-005 (Peplin Biotech) Synchrovax vaccines (CTL Immuno)	Dexosome therapy (Anosys) Pentrix (Australian Cancer Technology) JSF-154 (Tragen) Cancer vaccine (Intercell) Norelin (Biostar) BLP-25 (Biomira) MGV (Progenics) !3-Alethin (Dovetail)
35		

	Melanoma vaccine (CTL Immuno) p21-RAS vaccine (Gem-Vax)	CLL-Thera (Vasogen)
5	Hormonal and antihormonal agents	Oestrogens Conjugated oestrogens Ethinylloestradiol chlorotrianisene Idenestrol Hydroxyprogesterone caproate Medroxyprogesterone Testosterone Testosterone propionate Fluoxymesterone Methyltestosterone Diethylstilbestrol Megestrol Tamoxifen Toremofin Dexamethasone
10		Prednisone Methylprednisolone Prednisolone Aminoglutethimide Leuprolide Goserelin Leuporelin Bicalutamide Flutamide Octreotide Nilutamide Mitotan P-04 (Novogen) 2-Methoxyoestradiol (EntreMed) Arzoxifen (Eli Lilly)
15		
20	Photodynamic agents	Talaporfin (Light Sciences) Theralux (Theratechnologies) Motexafin-Gadolinium (Pharmacyclics) Pd-Bacteriopheophorbid (Yeda) Lutetium-Texaphyrin (Pharmacyclics) Hypericin
25	Tyrosine kinase inhibitors	Imatinib (Novartis) Leflunomide (Sugen/Pharmacia) ZD1839 (AstraZeneca) Erlotinib (Oncogene Science) Canertinib (Pfizer) Squalamine (Genaera) SU5416 (Pharmacia) SU6668 (Pharmacia) ZD4190 (AstraZeneca) ZD6474 (AstraZeneca) Vatalanib (Novartis) PKI166 (Novartis) GW2016 (GlaxoSmith-Kline) EKB-509 (Wyeth) EKB-569 (Wyeth)
30		Kahalide F (PharmaMar) CEP-701 (Cephalon) CEP-751 (Cephalon) MLN518 (Millenium) PKC412 (Novartis) Phenoxodiol O Trastuzumab (Genentech) C225 (ImClone) rhu-Mab (Genentech) MDX-H210 (Medarex) 2C4 (Genentech) MDX-447 (Medarex) ABX-EGF (Abgenix) IMC-1C11 (ImClone)
35		

5	Various agents	SR-27897 (CCK-A inhibitor, Sanofi-Synthelabo) Tocladesine (cyclic AMP agonist, Ribapharm) Alvocidib (CDK inhibitor, Aventis) CV-247 (COX-2 inhibitor, Ivy Medical) P54 (COX-2 inhibitor, Phytopharm) CapCell™ (CYP450 stimulant, Bavarian Nordic) GCS-100 (gal3 antagonist, GlycoGenesys) G17DT immunogen (gastrin inhibitor, Aphton) Efaproxiral (oxygenator, Allos Therapeutics) PI-88 (heparanase inhibitor, Progen) Tesmilifen (histamine antagonist, YM BioSciences) Histamine (histamine H2 receptor agonist, Maxim) Tiazofurin (IMPDH inhibitor, Ribapharm) Cilengitide (integrin antagonist, Merck KGaA) SR-31747 (IL-1 antagonist, Sanofi-Synthelabo) CCI-779 (mTOR kinase inhibitor, Wyeth) Exisulind (PDE-V inhibitor, Cell Pathways) CP-461 (PDE-V inhibitor, Cell Pathways) AG-2037 (GART inhibitor, Pfizer) WX-UK1 (plasminogen activator inhibitor, Willex) PBI-1402 (PMN stimulant, ProMetic LifeSciences) Bortezomib (proteasome inhibitor, Millennium) SRL-172 (T-cell stimulant, SR Pharma) TLK-286 (glutathione-S	BCX-1777 (PNP inhibitor, BioCryst) Ranpirnase (ribonuclease stimulant, Alfacell) Galarubicin (RNA synthesis inhibitor, Dong-A) Tirapazamine (reducing agent, SRI International) N-Acetylcysteine (reducing agent, Zambon) R-Flurbiprofen (NF-kappaB inhibitor, Encore) 3CPA (NF-kappaB inhibitor, Active Biotech) Seocalcitol (vitamin D receptor agonist, Leo) 131-I-TM-601 (DNA antagonist, TransMolecular) Eflornithin (ODC inhibitor, ILEX Oncology) Minodronic acid (osteoclast inhibitor, Yamanouchi) Indisulam (p53 stimulant, Eisai) Aplidin (PPT inhibitor, PharmaMar) Rituximab (CD20 antibody, Genentech) Gemtuzumab (CD33 antibody, Wyeth Ayerst) PG2 (haematopoiesis promoter, Pharmagenesis) Immunol™ (triclosan mouthwash, Endo) Triacetyluridine (uridine prodrug, Wellstat) SN-4071 (sarcoma agent, Signature BioScience) TransMID-107™ (immunotoxin, KS Biomedix) PCK-3145 (apoptosis promoter, Procyon) Doranidazole (apoptosis promoter, Pola)
10			
15			
20			
25			
30			
35			

5	transferase inhibitor, Telik) PT-100 (growth factor agonist, Point Therapeu- tics) Midostaurin (PKC inhibitor, Novartis) Bryostatin-1 (PKC stimu- lant, GPC Biotech) CDA-II (apoptosis pro- moter, Everlife) SDX-101 (apoptosis pro- moter, Salmedix) Ceflatonin (apoptosis pro- moter, ChemGenex)	CHS-828 (cytotoxic agent, Leo) Trans-retinic acid (differentiator, NIH) MX6 (apoptosis promoter, MAXIA) Apomine (apoptosis promoter, ILEX Oncology) Urocidin (apoptosis promoter, Bioniche) Ro-31-7453 (apoptosis promoter, La Roche) Brostallicin (apoptosis promoter, Pharmacia)
15	alkylating agents cyclophosphamide Busulfan Ifosfamide Melphalan Hexamethylmelamine Thiotepa chloroambucil Dacarbazine Carmustine	Lomustine Procarbazine Altretamine Estramustine phosphate Mechloroethamine Streptozocin Temozolomide Semustine
20	Platinum agents Cisplatin Oxaliplatin Spiroplatin Carboxyphthalatoplatinum Tetraplatin Ormiplatin Iproplatin	Carboplatin ZD-0473 (AnorMED) Lobaplatin (Aetema) Satraplatin (Johnson Matthey) BBR-3464 (Hoffmann-La Roche) SM-11355 (Sumitomo) AP-5280 (Access)
30	Antimetabolites Azacytidine Gemcitabine Capecitabine 5-fluorouracil Floxuridine 2-chlorodesoxyadenosine 6-Mercaptopurine 6-Thioguanine Cytarabine 2-fluorodesoxycytidine Methotrexate Idatrexate	Tomudex Trimetrexate Deoxycoformycin Fludarabine Pentostatin Raltitrexed Hydroxyurea Decitabine (SuperGen) Clofarabine (Bioenvision) Irofulven (MGI Pharma) DMDC (Hoffmann-La Roche)
35		

	Ethinylcytidine (Taiho)	
5	Topoisomerase inhibitors	Amsacrine Epirubicin Etoposide Teniposide or mitoxantrone Irinotecan (CPT-11) 7-Ethyl-10-hydroxycamptothecin Topotecan Dexrazoxanet (TopoTarget) Pixantrone (Novuspharma) Rebeccamycin analogue (Exelixis) BBR-3576 (Novuspharma)
10		Rubitecan (SuperGen) Exatecan mesylate (Daiichi) Quinamed (ChemGenex) Gimatecan (Sigma- Tau) Diflomotecan (Beaufour-Ipsen) TAS-103 (Taiho) Elsamitrucin (Spectrum) J-107088 (Merck & Co) BNP-1350 (BioNumerik) CKD-602 (Chong Kun Dang) KW-2170 (Kyowa Hakko)
15	Antitumour antibiotics	Dactinomycin (Actinomycin D) Doxorubicin (Adriamycin) Deoxyrubicin Valrubicin Daunorubicin (Daunomycin) Epirubicin Therarubicin Idarubicin Rubidazon Plicamycin Porfiromycin Cyanomorpholinodoxo-rubicin Mitoxantron (Novantron)
20		Amonafide Azonafide Anthrapyrazole Oxantrazole Losoxantrone Bleomycin sulfate (Blenoxan) Bleomycinic acid Bleomycin A Bleomycin B Mitomycin C MEN-10755 (Menarini) GPX-100 (Gem Pharmaceuticals)
25		

30

35

5

10

15

20

25

30

35

Antimitotic agents	Paclitaxel	SB 408075
	Docetaxel	(GlaxoSmithKline)
	Colchicine	E7010 (Abbott)
	Vinblastine	PG-TXL (Cell
	Vincristine	Therapeutics)
	Vinorelbine	IDN 5109 (Bayer)
	Vindesine	A 105972 (Abbott)
	Dolastatin 10 (NCI)	A 204197 (Abbott)
	Rhizoxin (Fujisawa)	LU 223651 (BASF)
	Mivobulin (Warner-	D 24851 (ASTA Medica)
	Lambert)	ER-86526 (Eisai)
	Cemadotin (BASF)	Combretastatin A4 (BMS)
	RPR 109881A (Aventis)	Isohomohalichondrin-B
	TXD 258 (Aventis)	(PharmaMar)
	Epothilone B (Novartis)	ZD 6126 (AstraZeneca)
	T 900607 (Tularik)	PEG-Paclitaxel (Enzon)
	T 138067 (Tularik)	AZ10992 (Asahi)
	Cryptophycin 52 (Eli Lilly)	IDN-5109 (Indena)
	Vinflunine (Fabre)	AVLB (Prescient
	Auristatin PE (Teikoku	NeuroPharma)
	Hormone)	Azaepothilon B (BMS)
	BMS 247550 (BMS)	BNP- 7787 (BioNumerik)
	BMS 184476 (BMS)	CA-4-Prodrug (OXiGENE)
	BMS 188797 (BMS)	Dolastatin-10 (NrH)
	Taxoprexin (Protarga)	CA-4 (OXiGENE)
Aromatase inhibitors	Aminoglutethimide	Exemestan
	Letrozole	Atamestan (BioMedicines)
	Anastrozole	YM-511 (Yamanouchi)
	Formestan	
Thymidylate synthase inhibitors	Pemetrexed (Eli Lilly)	Nolatrexed (Eximias)
	ZD-9331 (BTG)	CoFactor™ (BioKeys)
DNA antagonists	Trabectedin (PharmaMar)	Mafosfamide (Baxter
	Glufosfamide (Baxter	International)
	International)	Apaziquone (Spectrum
	Albumin + 32P (Isotope	Pharmaceuticals)
	Solutions)	O6-Benzylguanine
	Thymectacin (NewBiotics)	(Paligent)
	Edotreotid (Novartis)	

5

10

15

20

25

30

35

Farnesyl transferase inhibitors	Arglabin (NuOncology Labs) Ionafernib (Schering-Plough) BAY-43-9006 (Bayer)	Tipifarnib (Johnson & Johnson) Perillyl alcohol (DOR BioPharma)
Pump inhibitors	CBT-1 (CBA Pharma) Tariquidar (Xenova) MS-209 (Schering AG)	Zosuquidar trihydrochloride (Eli Lilly) Biricodar dicitrate (Vertex)
Histone acetyl transferase inhibitors	Tacedinaline (Pfizer) SAHA (Aton Pharma) MS-275 (Schering AG)	Pivaloyloxymethyl butyrate (Titan) Depsipeptide (Fujisawa)
Metalloproteinase inhibitors Ribonucleoside reductase inhibitors	Neovastat (Aeterna Laboratories) Marimastat (British Biotech) Gallium maltolate (Titan) Triapin (Vion)	CMT -3 (CollaGenex) BMS-275291 (Celltech) Tezacitabine (Aventis) Didox (Molecules for Health)
TNF-alpha agonists/ antagonists	Virulizin (Lorus Therapeutics) CDC-394 (Celgene)	Revimid (Celgene)
Endothelin-A receptor antagonists	Atrasentan (Abbot) ZD-4054 (AstraZeneca)	YM-598 (Yamanouchi)
Retinoic acid receptor agonists	Fenretinide (Johnson & Johnson) LGD-1550 (Ligand)	Alitretinoin (Ligand)
Immuno-modulators	Interferon Oncophage (Antigenics) GMK (Progenics) Adenocarcinoma vaccine (Biomira) CTP-37 (AVI BioPharma) JRX-2 (Immuno-Rx) PEP-005 (Peplin Biotech) Synchrovax vaccines (CTL Immuno) Melanoma vaccine (CTL Immuno) p21-RAS vaccine (GemVax)	Dexosome therapy (Anosys) Pentrix (Australian Cancer Technology) JSF-154 (Tragen) Cancer vaccine (Intercell) Norelin (Biostar) BLP-25 (Biomira) MGV (Progenics) I3-Alethin (Dovetail) CLL-Thera (Vasogen)

5	Hormonal and antihormonal agents	Oestrogens Conjugated oestrogens Ethinylloestradiol chlorotrianisene Idenestrol Hydroxyprogesterone caproate Medroxyprogesterone Testosterone Testosterone propionate Fluoxymesterone Methyltestosterone Diethylstilbestrol Megestrol Tamoxifen Toremofin Dexamethasone	Prednisone Methylprednisolone Prednisolone Aminoglutethimide Leuprolide Goserelin Leuporelin Bicalutamide Flutamide Octreotide Nilutamide Mitotan P-04 (Novogen) 2-Methoxyoestradiol (EntreMed) Arzoxifen (Eli Lilly)
15	Photodynamic agents	Talaporfin (Light Sciences) Theralux (Theratechnologies) Motexafin-Gadolinium (Pharmacyclics)	Pd-Bacteriopheophorbid (Yeda) Lutetium-Texaphyrin (Pharmacyclics) Hypericin
20	Tyrosine kinase inhibitors	Imatinib (Novartis) Leflunomide (Sugen/Pharmacia) ZD1839 (AstraZeneca) Erlotinib (Oncogene Science) Canertinib (Pfizer) Squalamine (Genaera) SU5416 (Pharmacia) SU6668 (Pharmacia) ZD4190 (AstraZeneca) ZD6474 (AstraZeneca) Vatalanib (Novartis) PKI166 (Novartis) GW2016 (GlaxoSmithKline) EKB-509 (Wyeth) EKB-569 (Wyeth)	Kahalide F (PharmaMar) CEP-701 (Cephalon) CEP-751 (Cephalon) MLN518 (Millenium) PKC412 (Novartis) Phenoxodiol O Trastuzumab (Genentech) C225 (ImClone) rhu-Mab (Genentech) MDX-H210 (Medarex) 2C4 (Genentech) MDX-447 (Medarex) ABX-EGF (Abgenix) IMC-1C11 (ImClone)
35	Various agents	SR-27897 (CCK-A inhibitor, Sanofi-Synthelabo)	BCX-1777 (PNP inhibitor, BioCryst) Ranpirnase (ribonuclease)

5	Tocladesine (cyclic AMP agonist, Ribapharm)	stimulant, Alfacell)
	Alvocidib (CDK inhibitor, Aventis)	Galarubicin (RNA synthesis inhibitor, Dong-A)
	CV-247 (COX-2 inhibitor, Ivy Medical)	Tirapazamine (reducing agent, SRI International)
	P54 (COX-2 inhibitor, Phytopharm)	N-Acetylcysteine (reducing agent, Zambon)
	CapCell™ (CYP450 stimulant, Bavarian Nordic)	R-Flurbiprofen (NF-kappaB inhibitor, Encore)
10	GCS-100 (gal3 antagonist, GlycoGenesys)	3CPA (NF-kappaB inhibitor, Active Biotech)
	G17DT immunogen (gastrin inhibitor, Aphton)	Seocalcitol (vitamin D receptor agonist, Leo)
	Efaproxiral (oxygenator, Allos Therapeutics)	131-I-TM-601 (DNA antagonist, TransMolecular)
15	PI-88 (heparanase inhibitor, Progen)	Eflornithin (ODC inhibitor, ILEX Oncology)
	Tesmilifen (histamine antagonist, YM BioSciences)	Minodronic acid (osteoclast inhibitor, Yamanouchi)
	Histamine (histamine H2 receptor agonist, Maxim)	Indisulam (p53 stimulant, Eisai)
20	Tiazofurin (IMPDH inhibitor, Ribapharm)	Aplidin (PPT inhibitor, PharmaMar)
	Cilengitide (integrin antagonist, Merck KGaA)	Rituximab (CD20 antibody, Genentech)
	SR-31747 (IL-1 antagonist, Sanofi-Synthelabo)	Gemtuzumab (CD33 antibody, Wyeth Ayerst)
25	CCI-779 (mTOR kinase inhibitor, Wyeth)	PG2 (haematopoiesis promoter, Pharmagenesis)
	Exisulind (PDE-V inhibitor, Cell Pathways)	Immunol™ (triclosan mouthwash, Endo)
	CP-461 (PDE-V inhibitor, Cell Pathways)	Triacetyluridine (uridine prodrug, Wellstat)
	AG-2037 (GART inhibitor, Pfizer)	SN-4071 (sarcoma agent, Signature BioScience)
30	WX-UK1 (plasminogen activator inhibitor, Willex)	TransMID-107™ (immunotoxin, KS Biomedix)
	PBI-1402 (PMN stimulant, ProMetic LifeSciences)	PCK-3145 (apoptosis promoter, Procyon)
	Bortezomib (proteasome inhibitor, Millennium)	Doranidazole (apoptosis promoter, Pola)
35	SRL-172 (T-cell stimulant, SR Pharma)	CHS-828 (cytotoxic agent, Leo)
	TLK-286 (glutathione-S transferase inhibitor, Telik)	

5	PT-100 (growth factor agonist, Point Therapeutics)	Trans-retinic acid (differentiator, NIH)
	Midostaurin (PKC inhibitor, Novartis)	MX6 (apoptosis promoter, MAXIA)
	Bryostat-1 (PKC stimulant, GPC Biotech)	Apomine (apoptosis promoter, ILEX Oncology)
	CDA-II (apoptosis promoter, Everlife)	Urocidin (apoptosis promoter, Bioniche)
10	SDX-101 (apoptosis promoter, Salmedix)	Ro-31-7453 (apoptosis promoter, La Roche)
	Ceflatonin (apoptosis promoter, ChemGenex)	Brostallicin (apoptosis promoter, Pharmacia)

15 A combined treatment of this type can be achieved with the aid of simultaneous, consecutive or separate dispensing of the individual components of the treatment. Combination products of this type employ the compounds according to the invention.

20 ASSAYS

The compounds of the formula I described in the examples were tested by the assays described below and were found to have kinase inhibitory activity. Other assays are known from the literature and could readily be performed by the person skilled in the art (see, for example, Dhanabal et al., *Cancer Res.* 59:189-197; Xin et al., *J. Biol. Chem.* 274:9116-9121; Sheu et al., *Anticancer Res.* 18:4435-4441; Ausprunk et al., *Dev. Biol.* 38:237-248; Gimbrone et al., *J. Natl. Cancer Inst.* 52:413-427; Nicosia et al., *In Vitro* 18:538- 549).

Measurement of Met kinase activity

35 According to the manufacturer's data (Met, active, upstate, catalogue No. 14-526), Met kinase is expressed for the purposes of protein production in insect cells (Sf21; *S. frugiperda*) and subsequent affinity-chromatographic

purification as "N-terminal 6His-tagged" recombinant human protein in a baculovirus expression vector.

5 The kinase activity can be measured using various available measurement systems. In the scintillation proximity method (Sorg et al., J. of Biomolecular Screening, 2002, 7, 11-19), the flashplate method or the filter binding test, the radioactive phosphorylation of a protein or peptide as substrate is
10 measured using radioactively labelled ATP (^{32}P -ATP, ^{33}P -ATP). In the case of the presence of an inhibitory compound, a reduced radioactive signal, or none at all, can be detected. Furthermore, homogeneous time-resolved fluorescence resonance energy transfer (HTR-FRET) and fluorescence polarisation (FP) technologies can be used as assay methods
15 (Sills et al., J. of Biomolecular Screening, 2002, 191-214).

Other non-radioactive ELISA assay methods use specific phospho-antibodies (phospho-ABs). The phospho-antibody only binds the phosphorylated substrate. This binding can be detected by chemiluminescence
20 using a second peroxidase-conjugated antibody (Ross et al., 2002, Biochem. J.).

Flashplate method (Met kinase)

25 The test plates used are 96-well Flashplate^R microtitre plates from Perkin Elmer (Cat. No. SMP200). The components of the kinase reaction described below are pipetted into the assay plate. The Met kinase and the substrate poly Ala-Glu-Lys-Tyr, (pAGLT, 6:2:5:1), are incubated for 3 hrs at
30 room temperature with radioactively labelled ^{33}P -ATP in the presence and absence of test substances in a total volume of 100 μl . The reaction is terminated using 150 μl of a 60 mM EDTA solution. After incubation for a further 30 min at room temperature, the supernatants are filtered off with
35 suction, and the wells are washed three times with 200 μl of 0.9% NaCl solution each time. The measurement of the bound radioactivity is carried

out by means of a scintillation measuring instrument (Topcount NXT, Perkin-Elmer).

5 The full value used is the inhibitor-free kinase reaction. This should be approximately in the range 6000-9000 cpm. The pharmacological zero value used is staurosporin in a final concentration of 0.1 mM. The inhibitory values (IC₅₀) are determined using the RS1_MTS program.

10 Kinase reaction conditions per well:

- 30 µl of assay buffer
- 10 µl of substance to be tested in assay buffer with 10% of DMSO
- 10 µl of ATP (final concentration 1 µM cold, 0.35 µCi of ³³P-ATP)
- 50 µl of Met kinase/substrate mixture in assay buffer;
- 15 (10 ng of enzyme/well, 50 ng of pAGLT/well)

Solutions used:

- Assay buffer:
 - 20 50 mM HEPES
 - 3 mM magnesium chloride
 - 3 µM sodium orthovanadate
 - 3 mM manganese(II) chloride
 - 1 mM dithiothreitol (DTT)
 - pH = 7.5 (to be set using sodium hydroxide)
- Stop solution:
 - 25 60 mM Titriplex III (EDTA)
- ³³P-ATP: Perkin-Elmer;
- 30 - Met kinase: Upstate, Cat. No. 14-526, Stock 1 µg/10 µl; spec. activity 954 U/mg;
- Poly-Ala-Glu-Lys-Tyr, 6 : 2 : 5 : 1 : Sigma Cat. No. P1152

Above and below, all temperatures are indicated in °C. In the following examples, "conventional work-up" means: water is added if necessary, the pH is adjusted, if necessary, to values between 2 and 10, depending on the constitution of the end product, the mixture is extracted with ethyl acetate or dichloromethane, the phases are separated, the organic phase is dried over sodium sulfate and evaporated, and the residue is purified by chromatography on silica gel and/or by crystallisation. Rf values on silica gel; eluent: ethyl acetate/methanol 9:1.

Mass spectrometry (MS): EI (electron impact ionisation) M^+
FAB (fast atom bombardment) $(M+H)^+$
ESI (electrospray ionisation) $(M+H)^+$
APCI-MS (atmospheric pressure chemical ionisation - mass spectrometry) $(M+H)^+$.

Retention time RT [min]: determination by HPLC

Column: ChromolithPerformance RP-18e (Merck KGaA, Cat. 1.02129.0001)

Eluents:

Eluent A: 0.1 M aqueous NaH_2PO_4

Eluent B: acetonitrile + 10% of water

Flow rate: 4 ml/min

Gradient:

0 min 1% of B

1 min 1% of B


7 min 99% of B

8 min 99% of B

Wavelength (detection): 220 nm

Example 1

The preparation of 3-diethylaminopropyl {3-[5-(4-chlorophenyl)-2-oxo-6H-1,3,4-thiadiazin-3-ylmethyl]phenyl}carbamate ("A1") is carried out analogously to the following scheme



25

a)

If the requisite haloacetophenones are not commercially available, they can be prepared analogously to the following synthetic procedure:

30

5.57 g of 3,4-dimethoxyacetophenone are dissolved in 60 ml of diethyl ether and 30 ml of 1,4-dioxane in a 250 ml three-necked flask provided with magnetic stirrer, condenser, thermometer, dropping funnel with pressure equalisation and drying tube, and 1.54 ml of bromine are added dropwise with stirring at RT, during which a precipitate forms after only a short time. The mixture is stirred at RT for a further 1 h, during which the precipitate re-dissolves, the temperature is raised by about 3°C, and a

35

pale-yellow, clear solution forms. This is poured onto ice, stirred well, and the precipitate formed between the phases is filtered off with suction. It is washed with water and then with a little MTB ether and dried (=K1). The mother liquor is extracted with MTB ether, dried, filtered and evaporated to dryness. The residue is triturated with a little MTB ether, filtered off with suction and dried (=K2). K1 and K2 are combined, giving 2'-bromo-4-chloroacetophenone, m.p. 91-92°; yield: 5.88 g (76%).

10

b)

15

8.09 ml of hydrazinium hydroxide are slowly added dropwise to a solution of 25.65 g of potassium O-ethyldithiocarbonate in 24 ml of water with stirring, and the mixture is stirred at room temperature for a further 6 h. The mixture is left to stand at room temperature for 16 h, then 12 ml of water are added, and the mixture is extracted with ether. The combined ether phases are dried, filtered and evaporated to dryness, giving 16.4 g of ethyl hydrazinecarbothionate.

20

c)

25

5.17 g of ethyl hydrazinecarbothionate (43 mmol) are added to a solution of 10.04 g of 2'-bromo-4-chloroacetophenone (43 mmol) in 40 ml of acetonitrile, and the mixture is stirred at RT for 3 h, during which a precipitate forms little by little. The reaction mixture is filtered with suction, washed with a little acetonitrile and then with ether and dried, giving 6.59 g (68%) of 5-(4-chlorophenyl)-3,6-dihydro-1,3,4-thiadiazin-2-one.

30

d)

35

4.19 g of 3-nitrobenzyl bromide and 9.95 g of potassium carbonate are added to a solution of 4.00 g of 5-(4-chlorophenyl)-3,6-dihydro-1,3,4-thiadiazin-2-one in 80 ml of acetonitrile, and the mixture is stirred at 80° for a further 2 h. The mixture is poured into water, extracted 2x with diethyl ether, dried, filtered and evaporated to dryness. A little diethyl ether is

added to the residue, which is crystallised and dried in a vacuum drying cabinet at 50°C, giving 5.5 g (86%) of 5-(4-chlorophenyl)-3-(3-nitrobenzyl)-3,6-dihydro-1,3,4-thiadiazin-2-one.

5

e)

10

5.47 g of 5-(4-chlorophenyl)-3-(3-nitrobenzyl)-3,6-dihydro-1,3,4-thiadiazin-2-one is dissolved in 100 ml of THF, and 1.3 g of Raney nickel is subsequently added. Hydrogen is subsequently passed in until starting material is no longer detectable. For work-up, the catalyst is filtered off, washed with THF, and the filtrate is evaporated to dryness and recrystallised from dichloromethane/diethyl ether, giving 4.6 g (94%) of 3-(3-aminobenzyl)-5-(4-chlorophenyl)-3,6-dihydro-1,3,4-thiadiazin-2-one.

15

f)

20

25






199 mg (0.6 mmol) of 3-(3-aminobenzyl)-5-(4-chlorophenyl)-3,6-dihydro-1,3,4-thiadiazin-2-one are dissolved in 4 ml of THF in an 8 ml multi-stirrer vessel, 107 µl of diisopropylethylamine are added, the mixture is cooled in an ice/water bath, 78 µl of isopropylidene chloroformate are added dropwise, and the mixture is subsequently stirred at 70°C for 1 h. The mixture is subsequently cooled to room temperature, and 107 µl (0.72 mmol) of 3-diethylaminopropan-1-ol is added, and the mixture is refluxed for 96 h. For work-up, the reaction mixture is diluted with 30 ml of dichloromethane, shaken with 20 ml of 1 N HCl and 20 ml of 2 N NaOH, dried over sodium sulfate, evaporated to dryness and purified by chromatography (dichloromethane/methanol: 9:1). The combined fractions containing the product are evaporated to dryness and crystallised from diethyl ether/petroleum ether 40-60, giving 128 mg of 3-diethylaminopropyl {3-[5-(4-chlorophenyl)-3,6-dihydro-2-oxo-6H-1,3,4-thiadiazin-3-ylmethyl]phenyl}carbamate ("A1"), m.p. 115-117°;







30









35









HPLC (100%): RT : 4.67 min.








The following compounds are obtained analogously








No.	Structure	M.p. [°C]; RT [min]
5 "A2"		157-158°
10	¹ H NMR (250 MHz, DMSO-d ₆) δ 9.680 (S, 1H), 7.849 (D, 2H), 7.546 (D, 2H), 7.475 (S, 1H), 7.394 (D, 1H), 7.235 (T, 1H), 6.960 (D, 1H), 4.993 (S, 2H) 4.803 (T, 1H), 4.331 (S, 2H), 4.093 (T, 2H), 3.614 (D, 2H)	
15 "A3"		70-105 (decomposition; evolution of gas); 4.56
20 "A4"		112-117 (decomposition; evolution of gas); 4.53
25	¹ H NMR (250 MHz, DMSO-d ₆) δ 9.719 (S, 1H), 7.856 (D, 2H), 7.558 (D, 2H), 7.472 (S, 1H), 7.400 (D, 1H), 7.253 (T, 1H), 6.982 (D, 1H), 5.003 (S, 2H), 4.347 (S, 2H), 4.046 (M, 2H), 2.985-2.621 (M, 7H), 2.556 (S, 3H), 2.038 (M, 1H), 1.639 (M, 1H)	
30 "A5"		106-107; 4.59
35 "A6"		96-97; 4.59

"A7"		155-156; 5.84
"A8"		59-61; 4.59
	¹ H NMR (250 MHz, DMSO-d ₆) δ 9.623 (S, 1H), 7.819 (Ml, 2H), 7.522 (M, 3H), 7.385 (DB, 1H), 7.240 (T, 1H), 6.967 (D, 1H), 5.005 (S, 2H), 4.346 (S, 2H), 4.108 (T, 2H), 2.553 (T, 2H), 2.325 (S, 6H), 1.830 (M, 2H)	
"A9"		66-68; 4.27
	¹ H NMR (250 MHz, DMSO-d ₆) δ 9.583 (S, 1H), 7.888 (M+, 2H), 7.473(SB, 1H), 7.300 (Ml, 4H), 6.963 (D, 1H), 4.986 (S, 2H), 4.324 (S, 2H), 4.093 (T, 2H), 2.335 (T, 2H), 2.160 (S, 6H), 1.767 (M, 2H)	
"A10"		
"A11"		61-64
"A12"		

5	"A13"		65-71
10	"A14"		167-168
15	"A15"		78-83
20	"A16"		59-65
25	"A17"		58-64
30	"A18"		58-63
35	"A19"		56-62
	"A20"		94-95

5	"A29"		
	"A30"		
10	"A31"		
15	"A32"		123-124
20	"A33"		93-94
25	"A34"		
30	"A35"		
35	"A36"		




"A37"		
"A38"		
"A39"		
"A40"		
"A41"		116-117
"A42"		153-154
"A43"		116-117

"A44"		
"A45"		
"A46"		
"A47"		
"A48"		
"A49"		
"A50"		

5

10

15

"A51"		
"A52"		
"A53"		
	¹ H NMR (250 MHz, DMSO-d ₆) δ [ppm] 9.636 (s, 1H), 7.784 (m, 2H), 7.461 (sb, 1H), 7.424 (d, 1H), 7.371 (m, 2H), 7.281 (m, 1H), 6.970 (d, 1H), 5.100 (s, 2H), 4.090 (t, 2H), 3.3 (m, 2H), 2.356 (m, 8H), 2.157 (s, 3H), 1.759 (dd, 2H)	

20


Example 2a

25

The preparation of N-{3-[5-(5-chloro-2-hydroxyphenyl)-2-oxo-6H-1,3,4-thiadiazin-3-ylmethyl]phenyl}-2-methylaminoacetamide ("B1") is carried out analogously to the following scheme

30

35



2a.1 87.6 mg of Boc-N-methylglycine are added to a solution of 260 mg of EDCI HCl [N-ethyl-N,N'-(dimethylaminopropyl)carbodiimide] and 169 mg of DMAP (dimethylaminopyridine) in 20 ml of dichloromethane and 3 drops of DMF, and the mixture is stirred for 30 minutes. A solution of 200 mg of "2be" (from Example 2b) in dichloromethane is then added, and the mixture is stirred at room temperature under an N₂ atmosphere for 16 hours. Conventional work-up gives "2aa".


2a.2 10 ml of 20% ethanolic KOH solution are added to a solution of 150 mg of "2aa" in 10 ml of ethanol at 0°. The mixture is stirred at room temperature for 10 minutes, subjected to conventional work-up, giving "2ab". This is dissolved in dichloromethane, and 2 ml of TFA is added at 0°, and the mixture is stirred at room temperature for 20 minutes. Conventional work-up gives "B1".

The following compounds are obtained analogously to the preparation of "B1"


No.	Structure	M.p. [°C]; RT [min]
"B7"		
"B8"		
"B9"		

Example 2b


The preparation of N-{3-[5-(5-chloro-2-hydroxyphenyl)-2-oxo-6H-1,3,4-thiadiazin-3-ylmethyl]phenyl}acetamide ("B2") is carried out analogously to the following scheme



2b.1 17.8 g of triethylamine are added dropwise to a solution of 15 g of 4-chloro-2-hydroxyacetophenone in 300 ml of dichloromethane, and the mixture is cooled to 0° . 12.72 g of pivalyl chloride are added, and the mixture is stirred under an N_2 atmosphere for 16 hours. Conventional work-up gives "2ba"




2b.2 A solution of 15.68 g of bromine in 200 ml of 1,4-dioxane is added dropwise to a solution of 25 g of "2ba" in 600 ml of 1,4-dioxane at 0° . The mixture is stirred for a further 3 hours and subjected to conventional work-up, giving "2bb"




"2bb" .

2b.3 A solution of 16.21 g of O-ethyl hydrazinecarbothioate




in 100 ml of tert-butanol is added to a solution of 45 g of "2bb" in 400 ml of tert-butanol. The mixture is boiled at 80° under an N₂ atmosphere for 16 hours. The mixture is cooled, subjected to conventional work-up, giving




"2bc" .


2b.4 4 g of "2bc" are added to a mixture of 5.075 g of potassium carbonate in 40 ml of DMF, and the mixture is stirred for 30 minutes. A solution of 2.64 g of m-nitrobenzyl bromide in DMF is then added, and the mixture is stirred at room temperature under an N₂ atmosphere for 16 hours. The mixture is subjected to conventional work-up, purified by column chromatography (hexane:ethyl acetate = 95:5), giving 1.2 g of "2bd"



2b.5 A solution of 1.87 g of NH₄Cl in 10 ml of water is added to a solution of 1.8 g of "2bd" in 20 ml of methanol. 1.3 g of iron powder are added, and the mixture is boiled under reflux for 3 hours. The mixture is cooled, filtered, subjected to conventional work-up, giving 900 mg of "2be"







2b.6 27.2 mg of acetyl chloride and a solution of 150 mg of "2be" in THF are added to a suspension of 16.6 mg of NaH (60% in mineral oil) in THF at 0°. The mixture is stirred at room temperature for 16 hours, subjected to conventional work-up, giving 120 mg of "2bf"



2b.7 10 ml of 20% ethanolic KOH solution are added to a solution of 120 mg of "2bf" in 10 ml of ethanol at 0°. The mixture is stirred at room

temperature for 10 minutes, subjected to conventional work-up, giving 60 mg of "B2".

5 The following compounds are obtained analogously to the preparation of "B2"

No.	Structure	M.p. [°C]; RT [min]
10 15 "B3"		
20 "B4"		
25 30 "B5"		
35 "B6"		

Pharmacological data

Met kinase inhibition

Table 1

5

10

15

20

Compound No.	IC ₅₀ (enzyme)
"A6"	A
"A7"	A
"A11"	A
"A12"	A
"A17"	A
"A20"	A
"A29"	A
"A30"	A
"A31"	A
"A33"	A
"B1"	A
"B2"	A
"B4"	A
"B6"	A

25

IC₅₀: 10 nM - 1 μ M = A1 μ M - 10 μ M = B> 10 μ M = C

30

35

The following examples relate to medicaments:

5 **Example A: Injection vials**

A solution of 100 g of an active ingredient of the formula I and 5 g of disodium hydrogenphosphate in 3 l of bidistilled water is adjusted to pH 6.5 using 2 N hydrochloric acid, sterile filtered, transferred into injection vials, lyophilised under sterile conditions and sealed under sterile conditions.
10 Each injection vial contains 5 mg of active ingredient.

Example B: Suppositories

A mixture of 20 g of an active ingredient of the formula I with 100 g of soya lecithin and 1400 g of cocoa butter is melted, poured into moulds and
15 allowed to cool. Each suppository contains 20 mg of active ingredient.

Example C: Solution

A solution is prepared from 1 g of an active ingredient of the formula I, 9.38 g of $\text{NaH}_2\text{PO}_4 \cdot 2 \text{H}_2\text{O}$, 28.48 g of $\text{Na}_2\text{HPO}_4 \cdot 12 \text{H}_2\text{O}$ and 0.1 g of benzalkonium chloride in 940 ml of bidistilled water. The pH is adjusted to 6.8, and the solution is made up to 1 l and sterilised by irradiation. This
20 solution can be used in the form of eye drops.
25

Example D: Ointment

500 mg of an active ingredient of the formula I are mixed with 99.5 g of Vaseline under aseptic conditions.
30

Example E: Tablets

A mixture of 1 kg of active ingredient of the formula I, 4 kg of lactose, 1.2 kg of potato starch, 0.2 kg of talc and 0.1 kg of magnesium stearate is pressed in a conventional manner to give tablets in such a way that each
35 tablet contains 10 mg of active ingredient.

Example F: Dragees

5 Tablets are pressed analogously to Example E and subsequently coated in a conventional manner with a coating of sucrose, potato starch, talc, tragacanth and dye.

Example G: Capsules

10 2 kg of active ingredient of the formula I are introduced into hard gelatine capsules in a conventional manner in such a way that each capsule contains 20 mg of the active ingredient.

Example H: Ampoules

15 A solution of 1 kg of active ingredient of the formula I in 60 l of bidistilled water is sterile filtered, transferred into ampoules, lyophilised under sterile conditions and sealed under sterile conditions. Each ampoule contains 10 mg of active ingredient.

20

25

30


35

Patent Claims

1. Compounds of the formula I

5

10



in which

15

R^1 denotes H, A, Hal, OH, OA, SH, SA, SOA, SO_2A , NO_2 , NH_2 , NHA, NAA' , SO_2NH_2 , SO_2NHA , SO_2NAA' , $CONH_2$, $CONHA$, $CONAA'$, $NACOA'$, $NASO_2A'$, $COOH$, $COOA$ or CN ,

20

R^2 denotes H, A, SO_2NH_2 , SO_2NHA , SO_2NAA' , $CONH_2$, $CONHA$, $CONAA'$, $NACOA'$, $NASO_2A'$, $COOH$, $COOA$ or CN ,

25

B denotes $NHCOO(CH_2)_nR^3$ or $NHCO(CH_2)_nR^3$,
 Q is absent or denotes linear or branched alkylene having 1-4 C atoms,

30

R^3 denotes R^1 , Het, alkyl having 1-6 C atoms or cycloalkyl having 3-8 C atoms, each of which is unsubstituted or mono-, di-, tri- or tetra-substituted by R^4 ,

35

R^4 denotes A, Hal, OH, OA, SH, SA, SOA, SO_2A , NO_2 , NH_2 , NHA, NAA' , SO_2NH_2 , SO_2NHA , SO_2NAA' , $CONH_2$, $CONHA$, $CONAA'$, $NACOA'$, $NASO_2A'$, $COOH$, $COOA$ or CN ,

- 5 Het denotes a mono- or bicyclic saturated or aromatic hetero-
cycle having 1 to 4 N, O and/or S atoms, which may be
unsubstituted or mono-, di- or trisubstituted by R⁴, CHO,
COA, =S, =NH, =NA and/or =O (carbonyl oxygen),
A, A' each, independently of one another, denote unbranched or
branched alkyl having 1-10 C atoms,
in which 1-7 H atoms may be replaced by F, Cl and/or
Br,
10 cycloalkyl having 3-8 C atoms or
cycloalkylalkylene having 4-10 C atoms,
Hal denotes F, Cl, Br or I,
m denotes 0 or 1,
15 n denotes 0, 1, 2 or 3,

and pharmaceutically usable solvates, salts, tautomers
and stereoisomers thereof, including mixtures thereof in all ratios.

- 20 2. Compounds according to Claim 1 in which
R¹ denotes H, Hal, A, OH, OA, SO₂A, CN, NO₂, COOA or
COOH,
and pharmaceutically usable solvates, salts, tautomers
25 and stereoisomers thereof, including mixtures thereof in all ratios.
3. Compounds according to Claim 1 or 2 in which
R² denotes H
30 and pharmaceutically usable solvates, salts, tautomers
and stereoisomers thereof, including mixtures thereof in all ratios.
4. Compounds according to one or more of Claims 1-3 in which
35 Het denotes a monocyclic saturated or aromatic heterocycle
having 1 to 2 N and/or O atoms, which may be unsubsti-
tuted or mono- or disubstituted by A,

and pharmaceutically usable solvates, salts, tautomers
and stereoisomers thereof, including mixtures thereof in all ratios.

5 5. Compounds according to one or more of Claims 1-4 in which
Q denotes CH_2 , $\text{CH}(\text{CH}_3)$ or $\text{CH}(\text{CH}_2\text{CH}_3)$,
and pharmaceutically usable solvates, salts, tautomers
and stereoisomers thereof, including mixtures thereof in all ratios.

10 6. Compounds according to one or more of Claims 1-5 in which
 R^3 denotes Het or alkyl having 1-6 C atoms or cycloalkyl hav-
ing 3-8 C atoms, each of which is unsubstituted or mono-,
di-, tri- or tetrasubstituted by R^4 ,
15 and pharmaceutically usable solvates, salts, tautomers
and stereoisomers thereof, including mixtures thereof in all ratios.

20 7. Compounds according to one or more of Claims 1-6 in which
 R^4 denotes OH, NH_2 , NHA or NAA' ,
and pharmaceutically usable solvates, salts, tautomers
and stereoisomers thereof, including mixtures thereof in all ratios.

25 8. Compounds according to one or more of Claims 1-7 in which
A, A' each, independently of one another, denote unbranched
or branched alkyl having 1-6 C atoms, in which 1-5 H
atoms may be replaced by F and/or chlorine,
and pharmaceutically usable solvates, salts, tautomers
30 and stereoisomers thereof, including mixtures thereof in all ratios.

35 9. Compounds according to one or more of Claims 1-8 in which
Het denotes piperidinyl, pyrrolidinyl, morpholin-4-yl, piperazinyl,
1,3-oxazolidin-3-yl, imidazolidinyl, pyridyl, pyrimidinyl,
furanyl, thienyl, thiazolyl, indolyl or indazolyl, where the
radicals may also be mono- or disubstituted by A,







and pharmaceutically usable solvates, salts, tautomers
and stereoisomers thereof, including mixtures thereof in all ratios.






- 5 10. Compounds according to one or more of Claims 1-9 in which
- R^1 denotes H, Hal, A, OH, OA, SO_2A , CN, NO_2 , COOA or
COOH,
- R^2 denotes
- 10 B denotes $NHCOO(CH_2)_nR^3$ or $NHCO(CH_2)_nR^3$,
Q denotes CH_2 , $CH(CH_3)$ or $CH(CH_2CH_3)$,
 R^3 denotes Het or alkyl having 1-6 C atoms or cycloalkyl hav-
ing 3-8 C atoms, each of which is unsubstituted or mono-,
15 di-, tri- or tetrasubstituted by R^4 ,
 R^4 denotes OH, NH_2 , NHA or NAA',
A, A' each, independently of one another, denote unbranched or
branched alkyl having 1-6 C atoms, in which 1-5 H atoms
20 may be replaced by F and/or chlorine,
Het denotes a monocyclic saturated or aromatic heterocycle
having 1 to 2 N and/or O atoms, which may be unsubstitu-
ted or mono- or disubstituted by A,
Hal denotes F, Cl, Br or I,
25 m denotes 0 or 1,
n denotes 0, 1, 2 or 3,
and pharmaceutically usable solvates, salts, tautomers
and stereoisomers thereof, including mixtures thereof in all ratios.

30


11. Compounds according to Claim 1, selected from the group

35

5	"A9"	
10	"A10"	
15	"A11"	
20	"A12"	
25	"A14"	
30		
35	"A16"	


5	"A25"	
10	"A26"	
15	"A27"	
20	"A28"	
25		
30		
35	"A32"	

5




10


15




20




25



30



35



5

10


15

20

25

30


35



5	"B7"	
10	"B8"	
15	"B9"	

and pharmaceutically usable solvates, salts, tautomers
and stereoisomers thereof, including mixtures thereof in all ratios.

12. Process for the preparation of compounds of the formula I according
to Claims 1-11 and pharmaceutically usable salts, sol-
vates, tautomers and stereoisomers thereof, characterised in that a
compound of the formula Ia



in which

B denotes NH₂,

and R¹, R² and Q have the meanings indicated in Claim 1,

is converted into a compound of the formula I in which

B denotes $\text{NHCOO}(\text{CH}_2)_n\text{R}^3$,

5

by reacting a compound of the formula Ia with a coupling reagent selected from the group

a) isopropylidene chloroformate,

b) p-nitrophenyl chloroformate,


10

c) diphosgene,

d) triphosgene,

and a compound of the formula II

15



in which n and R^3 have the meanings indicated in Claim 1,

20

and/or

a base or acid of the formula I is converted into one of its salts.

25

13. Medicaments comprising at least one compound of the formula I according to Claim 1-11 and/or pharmaceutically usable salts, solvates, tautomers and stereoisomers thereof, including mixtures thereof in all ratios, and optionally excipients and/or adjuvants.

30

35

5

10

15

14. Use of compounds according to Claim 1-11 and pharmaceutically usable salts, solvates, tautomers and stereoisomers thereof, including mixtures thereof in all ratios, for the preparation of a medicament for the treatment of diseases where the disease to be treated is a solid tumor.

20

15. Use according to Claim 14, where the solid tumour originates from the group of tumours of the squamous epithelium, the bladder, the stomach, the kidneys, of head and neck, the oesophagus, the cervix, the thyroid, the intestine, the liver, the brain, the prostate, the urogenital tract, the lymphatic system, the stomach, the larynx and/or the lung.

25

16. Use according to Claim 14, where the solid tumour originates from the group monocytic leukaemia, lung adenocarcinoma, small-cell lung carcinomas, pancreatic cancer, glioblastomas and breast carcinoma.

30

17. Use according to Claim 15, where the solid tumour originates from the group of lung adenocarcinoma, small-cell lung carcinomas, pancreatic cancer, glioblastomas, colon carcinoma and breast carcinoma.

35

18. Use of compounds according to Claim 1-11 and pharmaceutically usable salts, solvates, tautomers and stereoisomers thereof, including mixtures thereof in all ratios, for the preparation of a medicament for the treatment of diseases, where the disease to be treated is a tumor of the blood and immune system.

5 19. Use according to Claim 18, where the tumour originates from the group of acute myeloid leukaemia, chronic myeloid leukaemia, acute lymphatic leukaemia and/or chronic lymphatic leukaemia.

10 20. Medicaments comprising at least one compound of the formula I according to one or more of Claims 1 to 11, and/or pharmaceutically usable solvates and stereoisomers thereof, including mixtures thereof in all ratios, and at least one further medicament active ingredient.

15 21. Set (kit) consisting of separate packs of
(a) an effective amount of a compound of the formula I according to one or more of Claims 1 to 11, and/or pharmaceutically usable solvates, salts and stereoisomers thereof, including mix-
20 tures thereof in all ratios,
and
(b) an effective amount of a further medicament active ingredi-
25 ent.