

NOVEL CURCUMIN ANALOGUES AND USES THEREOF

Related Applications

This application is a continuation-in-part of PCT International Application No.
5 PCT/US2003/009350, filed March 27, 2003, titled *Novel Curcumin Analogues and
Uses Thereof*, published in English on October 30, 2003, which claims priority from
United States Patent Application Serial No. 10/124,642, filed April 17, 2002, titled
Novel Curcumin Analogues and Uses Thereof, now issued as United States Patent No.
6,790,979, the disclosures of which are incorporated by reference herein in their
10 entirety.

Statement of Federal Support

This invention was made with Government support under Grant No. CA-
17625 and Grant No. CA-55639. The government has certain rights in this invention.
15

Field of the Invention

The present invention relates to compounds capable of acting as androgen
receptor antagonists, pharmaceutical formulations containing the same, and methods
of use thereof.
20

Background of the Invention

The androgen receptor (AR) is a member of a large family of ligand-
dependent transcriptional factors known as the steroid receptor superfamily. Chang et
al., *Proc. Natl. Acad. Sci. USA*, 85, 7211-7215 (1988). Beato, M., *Cell*, 56, 335-344
25 (1989). Androgens and the AR play an important role in the growth of the normal
prostate and prostate cancer. Prostate cancer represents the most common male
malignancy in the United States. Landis et al., *Cancer J. Clin.*, 48, 6-29 (1998).
Recently, antiandrogens such as hydroxyflutamide (HF) in combination with surgical
or medical castration have been widely used for the treatment of prostate cancer.
30 Crawford et al., *New Engl. J. Med.*, 321, 419-424 (1989). Several compounds,
including cyprosterone, HF, and bicalutamide (shown below), have been used
clinically in the treatment of prostate cancer.




Chart 1. Structures of cyproterone, hydroxyflutamide, and bicalutamide.

The synthetic steroidal antiandrogen cyproterone is one of the first antiandrogens used clinically in Europe, McLeod, D., G., *Cancer*, 71, 1046-1049 (1993) but it has many side effects. Neumann et al., *J. Clin. Oncol.*, 1, 41-65 (1982). HF and bicalutamide are both nonsteroidal antiandrogens. Bicalutamide is a newer nonsteroidal antiandrogen originally thought to have a pure antiandrogen activity without agonist activity. It has a longer half-life (6 days) and a higher binding affinity to the AR than HF. Verhelst et al., *Clin. Endocrinol.*, 41, 525-530 (1994). (a) Kelly et al., *J. Urol.* (1993), 149, 607-609; (b) Scher et al., *Prostate Cancer. J. Clin. Oncol.*, 11, 1566-1572 (1993).

Although antiandrogen hormone therapy has been widely used for the treatment for prostate cancer, some antiandrogens may act as AR agonists which may result in “antiandrogen withdrawal syndrome.” Miyamoto et al., *Proc. Natl. Acad. Sci. USA*, 95, 7379-7384 (1998). A currently accepted hypothesis postulates that mutations in androgen receptors may account for why HF, the active metabolite of flutamide, can activate androgen receptor target genes and stimulate prostate cancer growth. Miyamoto et al., *Proc. Natl. Acad. Sci. USA*, 95, 7379-7384 (1998). The same mechanism is used to explain the “flutamide withdrawal syndrome,” in which patients who experience an increase in prostate-specific antigen (PSA) while taking flutamide, have a decrease in PSA after withdrawal of treatment. Indeed, HF can activate androgen receptor target genes, such as PSA and MMTV-LTR (a reporter gene which expressed androgen-response element), in the presence of ARA70, the first identified androgen receptor co-activator. Yeh et al., *The Lancet*, 349, 852-853 (1997). Because this syndrome often leads to the failure of androgen-ablative therapy, it is desirable to develop better antiandrogens without agonist activity.



The phenolic diarylheptanoid curcumin (1) is the major pigment in turmeric. Curcumin and its analogs show potent anti-oxidant activity, anti-inflammatory activity, Nurfin et al., *Eur. J. Med. Chem.*, 32, 321-328 (1997) cytotoxicity against tumor cells, Syu et al., *J. Nat. Prod.*, 61, 1531-1534 (1998), antitumor-promoting activities, Sugiyama et al., *Biochem. Pharmacol.*, 52, 519-525 (1996). Ruby et al., *Cancer Lett.*, 94, 79-83 (1995) and antiangiogenesis activity (J.L. Arbiser et al. *Mol. Med.*, 4: 376 (1998)).

Two cyclic diarylheptanoids, 13-oxomyricanol and myricanone, exhibiting potent antitumor promoting effects on DMBA-initiated and TPA-induced mouse skin carcinogenesis have been reported. Ishida et al., *Cancer Lett.*, 159, 135-140 (2000). In the present study, a number of novel curcumin analogues have been prepared and evaluated for antagonistic activity against the AR in the presence of androgen receptor coactivator, ARA70, using two human prostate cancer cell lines, PC-3 and DU-145. PC-3 cells are androgen-independent tumor cells that do not express functional AR. DU-145 cells are androgen-independent tumor cells that also do not express functional AR.

Summary of the Invention

According to embodiments of the present invention, the present invention relates to a compound according to formula I:

A first aspect of the present invention is a compound according to formula I or Ia:



wherein:

R' and R" are each independently selected from the group consisting of H, alkoxy, and halo;

R_a and R_b are each independently selected from the group consisting of H, alkyl and halo;

5 R₁ and R₂ are each independently selected from the group consisting of H, hydroxy, alkoxy, nitro, amino, and dialkylamino;

R₃ and R₄ are each independently selected from the group consisting of H, hydroxy, halo, alkoxy, and -OR₇C(O)R₈, wherein R₇ is lower alkylene and R₈ is alkoxy;

10 or R₁ and R₃ together are alkylenedioxy;

or R₂ and R₄ together are alkylenedioxy;

R₅ and R₆ are each independently selected from the group consisting of H, halogen, nitro and alkoxy;

X₁ is N, or X₁ is C bonded to a H, alkoxy or nitro; and


15 X₂ is N, or X₂ is C bonded to a H, alkoxy or nitro;

subject to the proviso that curcumin is excluded therefrom;

or a pharmaceutically acceptable salt thereof.

A further aspect of the present invention is a compound according to formula

II:



20

wherein:

R₁₁ and R₁₂ are each independently selected from the group consisting of alkoxy, nitro, amino, and dialkylamino;

25 R₁₃ and R₁₄ are each independently selected from the group consisting of hydroxy, alkoxy, -OR₁₈C(O)R₁₉ wherein R₁₈ is lower alkylene or alkenylene and R₁₉ is alkoxy, and tetrahydropyranyl;

or R₁₁ and R₁₃ together are alkylenedioxy;

or R₁₂ and R₁₄ together are alkylenedioxy;

R_{15} and R_{16} are each independently selected from the group consisting of H, halogen, and nitro;

R_{17} is $-R_{20}C(O)OR_{21}$, wherein R_{20} is alkylene or alkenylene and R_{21} is H or alkyl;


5 X_3 is N, or X_3 is C bonded to a H, alkoxy or nitro; and

X_4 is N, or X_4 is C bonded to a H, alkoxy or nitro;

or a pharmaceutically acceptable salt thereof.

According to yet other embodiments of the present invention, the present invention relates to compounds according to the formula III:

10



(III)

15

wherein:

R_{25} and R_{26} are each independently H or lower alkyl;


R_{27} , R_{28} , R_{29} and R_{30} are each alkoxy;

R_{31} is H or lower alkyl;

or a pharmaceutically acceptable salt thereof.

20

A further aspect of the present invention is a compound of Formula IV:



(IV)


wherein:

R_{41} and R_{42} are each independently hydroxy or alkoxy; and

R_{43} and R_{44} are each independently hydroxy or alkoxy;

25 or a pharmaceutically acceptable salt thereof.

A further aspect of the invention is a compound of the Formula:



or a pharmaceutically acceptable salt thereof.

According to still other embodiments of the present invention, the present invention relates to a method of treating cancer, comprising administering to a subject in need thereof a treatment effective amount of a compound according to the formulas above or curcumin. Examples of cancers that may be treated include, but are not limited to, skin cancer, small cell lung cancer, testicular cancer, lymphoma, leukemia, esophageal cancer, stomach cancer, colon cancer, breast cancer, endometrial cancer, ovarian cancer, central nervous system cancer, liver cancer and prostate cancer.

According to yet other embodiments of the present invention, the present invention relates to a method of inducing androgen receptor antagonist activity, the method comprising contacting a cancer cell with an androgen receptor antagonist effective amount of a compound according to the formulas above or curcumin.

According to other embodiments of the present invention, the present invention relates to a method of inducing androgen receptor antagonist activity in a subject afflicted with an androgen-related affliction by administering a compound according to the formulas above or curcumin. Examples of androgen-related afflictions include, but are not limited to, baldness, hirsutism, behavioral disorders, acne, and uninhibited spermatogenesis wherein inhibition of spermatogenesis is so desired.

Brief Description of Drawings

Figure 1 illustrates structures of curcumin analogues (1-20);

Figure 2 illustrates structures of curcumin analogues (21-44);

Figure 3A illustrates suppression of DHT-mediated MMTV transcription AR activity by hydroxyflutamide (HF) and selected compounds;

Figure 3B illustrates suppression of DHT-mediated MMTV transcription AR activity by hydroxyflutamide (HF) and selected compounds; and

Figure 3C illustrates suppression of DHT-mediated MMTV transcription AR activity by hydroxyflutamide (HF) and selected compounds.

Detailed Description of the Invention


5 The present invention will now be described more fully hereinafter with reference to the accompanying figures, which further illustrate the invention described herein. This invention may, however, be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these
10 embodiments are provided so that this disclosure will be thorough and complete, and will fully convey the scope of the invention to those skilled in the art.

 The terminology used in the description of the invention herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention. As used in the description of the invention and the appended claims, the singular forms "a", "an" and "the" are intended to include the plural forms as well,
15 unless the context clearly indicates otherwise.

 Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. All publications, patent applications, patents and other references mentioned herein are incorporated by reference in their entirety.

20 The term "alkyl" or "loweralkyl" as used herein refers to C1 to C4, C6 or C8 alkyl, which may be linear or branched and saturated or unsaturated.

 "Curcumin" as used herein refers to a compound of the Formula:



Curcumin, or a pharmaceutically acceptable salt thereof, may be used to carry out the
25 methods described herein.

 "Cycloalkyl" is specified as such herein, and is typically C3, C4 or C5 to C6 or C8 cycloalkyl.

 "Alkenyl" or "loweralkenyl" as used herein likewise refers to C1 to C4 alkenyl, and alkoxy or loweralkoxy as used herein likewise refers to C1 to C4 alkoxy.

"Alkoxy" as used herein refers to linear or branched, saturated or unsaturated oxo-hydrocarbon chains, including for example methoxy, ethoxy, propoxy, isopropoxy, butoxy, and *t*-butoxy.


The term "aryl" as used herein refers to C3 to C10 cyclic aromatic groups such as phenyl, naphthyl, and the like, and includes substituted aryl groups such as tolyl.

"Halo" as used herein refers to any halogen group, such as chloro, fluoro, bromo, or iodo. In some embodiments halo is preferably fluoro.

The term "hydroxyalkyl" as used herein refers to C1 to C4 linear or branched hydroxy-substituted alkyl, i.e., -CH₂OH, -(CH₂)₂OH, etc.

The term "aminoalkyl" as used herein refers to C1 to C4 linear or branched amino-substituted alkyl, wherein the term "amino" refers to the group NR'R'', wherein R' and R'' are independently selected from H or lower alkyl as defined above, i.e., -NH₂, -NHCH₃, -N(CH₃)₂, etc.

The term "tetrahydropyranyl" refers to a group of the formula:



The term "oxyalkyl" as used herein refers to C1 to C4 oxygen-substituted alkyl, i.e., -OCH₃, and the term "oxyaryl" as used herein refers to C3 to C10 oxygen-substituted cyclic aromatic groups.

The term "alkylenedioxy" refers to a group of the general formula -ORNO-, -ORNORN-, or -RNORNORN- where each RN is independently alkyl.

"Treat" or "treating" as used herein refers to any type of treatment that imparts a benefit to a patient afflicted with a disease, including improvement in the condition of the patient (*e.g.*, in one or more symptoms), delay in the progression of the disease, prevention or delay of the onset of the disease, etc.

"Pharmaceutically acceptable" as used herein means that the compound or composition is suitable for administration to a subject to achieve the treatments described herein, without unduly deleterious side effects in light of the severity of the disease and necessity of the treatment.

"Inhibit" as used herein means that a potential effect is partially or completely eliminated.

"Androgen" as used herein refers to sex hormones generally known to those skilled in the art and include, but are not limited to, testosterone, dihydrotestosterone, and androstenedione and compounds known to act in mechanisms similar to androgens such as androgen receptor agonists. "Androgen" relates to a hormone or
5 compound, or a combination thereof.

"Antiandrogen withdrawal syndrome" as used herein refers to a phenomenon characterized by either no change or an increase in serum prostate-specific antigen (PSA) concentration upon administration of antiandrogen therapy, and a subsequent decreased PSA concentration observed after withdrawal of antiandrogen therapy.



10 "Androgen receptor antagonist" as used herein refers to a compound that partially or completely inhibits the activity of an androgen receptor agonist.

"Androgen-related affliction" as used herein refers to conditions wherein an androgen or combination of androgens play a role in the condition observed.

The present invention is concerned primarily with the treatment of human
15 subjects, but may also be employed for the treatment of other animal subjects (i.e., mammals, avians) for veterinary purposes. Mammals are preferred, with humans being particularly preferred.

As noted above, a first aspect of the present invention is a compound according to Formulas I or Ia:

20



wherein:

R' and R'' are each independently selected from the group consisting of H,
25 alkoxy, and halo;

R_a and R_b are each independently selected from the group consisting of H, alkyl and halo;

R_1 and R_2 are each independently selected from the group consisting of H, hydroxy, alkoxy, nitro, amino, and dialkylamino;

R_3 and R_4 are each independently selected from the group consisting of H, hydroxy, halo, alkoxy, and $-OR_7C(O)R_8$, wherein R_7 is lower alkylene and R_8 is

5 alkoxy;

or R_1 and R_3 together are alkylenedioxy;

or R_2 and R_4 together are alkylenedioxy;

R_5 and R_6 are each independently selected from the group consisting of H, halogen, nitro and alkoxy;

10 X_1 is N, or X_1 is C bonded to a H, alkoxy or nitro; and

X_2 is N, or X_2 is C bonded to a H, alkoxy or nitro;

subject to the proviso that curcumin is excluded therefrom;

or a pharmaceutically acceptable salt thereof.

In some particular embodiments of compounds of Formulas I and Ia, at least
15 one of R' and R'' are alkoxy or halo.

In some particular embodiments of compounds of Formulas I and Ia, R_a is halo and R_b is H, alkyl or halo, preferably alkyl.


In some particular embodiments of compounds of Formulas I and Ia, R_1 and R_2 are each independently selected from the group consisting of alkoxy, nitro, amino,
20 and dimethylamino.

In some particular embodiments of compounds of Formulas I and Ia, R_1 and R_3 together are methylenedioxy or ethylenedioxy.

In some particular embodiments of compounds of Formulas I and Ia, R_2 and R_4 together are methylenedioxy or ethylenedioxy.

25 In some particular embodiments of compounds of Formulas I and Ia, R_5 and R_6 are each independently selected from the group consisting of H, F, and nitro.

A further aspect of the present invention is a compound according to Formula II:



wherein:

R₁₁ and R₁₂ are each independently selected from the group consisting of alkoxy, nitro, amino, and dialkylamino;

5 R₁₃ and R₁₄ are each independently selected from the group consisting of hydroxy, alkoxy, -OR₁₈C(O)R₁₉ wherein R₁₈ is lower alkylene or alkenylene and R₁₉ is alkoxy, and tetrahydropyranyl;

or R₁₁ and R₁₃ together are alkylenedioxy;

or R₁₂ and R₁₄ together are alkylenedioxy;

10 R₁₅ and R₁₆ are each independently selected from the group consisting of H, halogen, and nitro;

R₁₇ is -R₂₀C(O)OR₂₁, wherein R₂₀ is alkylene or alkenylene and R₂₁ is H or alkyl;

X₃ is N, or X₃ is C bonded to a H, alkoxy or nitro; and

15 X₄ is N, or X₄ is C bonded to a H, alkoxy or nitro;

or a pharmaceutically acceptable salt thereof.

In some particular embodiments of compounds of Formula II, R₂₀ is alkenylene.

20 In some particular embodiments of compounds of Formula II, R₁₃ and R₁₄ are tetrahydropyranyl.

In some particular embodiments of compounds of Formula II, R₁₁ and R₁₂ are each independently selected from the group consisting of alkoxy, nitro, amino, and dimethylamino.


25 In some particular embodiments of compounds of Formula II, R₁₁ and R₁₃ together are methylenedioxy or ethylenedioxy.

In some particular embodiments of compounds of Formula II, R₁₂ and R₁₄ together are methylenedioxy or ethylenedioxy.

In some particular embodiments of compounds of Formula II, R₁₅ and R₁₆ are each independently selected from the group consisting of H, F, and nitro.

30 According to yet other embodiments of the present invention, the present invention relates to compounds according to the formula III:

(III)



wherein:


R₂₅ and R₂₆ are each independently H or lower alkyl;

5 R₂₇, R₂₈, R₂₉ and R₃₀ are each alkoxy;

R₃₁ is H or lower alkyl;

or a pharmaceutically acceptable salt thereof.

A further aspect of the present invention is a compound of Formula IV:




10 wherein:

R₄₁ and R₄₂ are each independently hydroxy or alkoxy; and

R₄₃ and R₄₄ are each independently hydroxy or alkoxy;

or a pharmaceutically acceptable salt thereof.

A further aspect of the invention is a compound of the Formula:




15

or a pharmaceutically acceptable salt thereof.


A. Specific compounds

20 Specific compounds within the scope of the present invention include, but are not limited to:




S-1

S-2




5

S-3




S-4



10

S-5



Additional examples of compounds of the invention are set forth below.

15

B. Synthesis of Compounds

Variations on the following general synthetic methods will be readily apparent to those skilled in the art and are deemed to be within the scope of the present invention.

20

Figures 1 and 2 show the structures of curcumin analogues and 1,3-diaryl-1,3-diketopropane derivatives. Curcumin (1), demethoxycurcumin (2) and bisdemethoxycurcumin (3) were obtained by column chromatography (silica gel, CHCl₃-MeOH) of commercially available curcumin (Aldrich), which contained 2 and 3 as minor components. Treatment of 1 with diazomethane gave dimethylated curcumin (4) and monomethylated curcumin (9). Methylation of 1 with methyl iodide and K₂CO₃ furnished the trimethylated derivative 10, in which a methyl group was

25

also introduced at the C-4 position. Compounds 5–8 were synthesized by heating 1–4 with histidine hydrazide, AcOH and *p*-TsOH overnight. Hydrogenation of 1 with 10% Pd-C gave a mixture of 11–13. Similarly, compounds 14–16 and 17–18 were obtained by hydrogenation of 4, and 10, respectively. Heating 1 with methyl chloroacetate, NaI and K₂CO₃ in acetone furnished a mixture of monomethoxycarbonylmethyl ether 18 and bis-methoxycarbonylmethyl ether 19, which were separated by preparative TLC (PLC). Compounds 21–23 were prepared from benzene or vanillin and ethyl 4-acetyl-5-oxohexanoate by a method known in the art. Pedersen et al., *Liebigs Ann. Chem.*, 1557-1569 (1985). Compounds 21–23 constitute an unseparable mixture of keto-enol tautomeric isomers. The syntheses of 24–38 were described in our previous paper. Ishida et al., *Cancer Lett.*, 159. 135-140 (2000). Ishida et al., Synthesis and Evaluation of Curcumin Analogues as Cytotoxic Agents. Unpublished data. Compounds 39–44 were purchased from Aldrich, Inc (Milwaukee, WI).

15

C. Pharmaceutically acceptable salts

The term “active agent” as used herein, includes the pharmaceutically acceptable salts of the compound. Pharmaceutically acceptable salts are salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects. Examples of such salts are (a) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; and salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, *p*-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (b) salts formed from elemental anions such as chlorine, bromine, and iodine.

Active agents used to prepare compositions for the present invention may alternatively be in the form of a pharmaceutically acceptable free base of active agent. Because the free base of the compound is less soluble than the salt, free base compositions are employed to provide more sustained release of active agent to the target area. Active agent present in the target area which has not gone into solution is

not available to induce a physiological response, but serves as a depot of bioavailable drug which gradually goes into solution.

D. Pharmaceutical Formulations

5 Curcumin and the curcumin analogues of the present invention are useful as pharmaceutically active agents and may be utilized in bulk form. More preferably, however, these compounds are formulated into pharmaceutical formulations for administration. Any of a number of suitable pharmaceutical formulations may be utilized as a vehicle for the administration of the compounds of the present invention.

10 The compounds of the present invention may be formulated for administration for the treatment of a variety of conditions. In the manufacture of a pharmaceutical formulation according to the invention, the compounds of the present invention and the physiologically acceptable salts thereof, or the acid derivatives of either (hereinafter referred to as the "active compound") are typically admixed with, *inter*
15 *alia*, an acceptable carrier. The carrier must, of course, be acceptable in the sense of being compatible with any other ingredients in the formulation and must not be deleterious to the patient. The carrier may be a solid or a liquid, or both, and is preferably formulated with the compound as a unit-dose formulation, for example, a tablet, which may contain from 0.5% to 95% by weight of the active compound. One
20 or more of each of the active compounds may be incorporated in the formulations of the invention, which may be prepared by any of the well-known techniques of pharmacy consisting essentially of admixing the components, optionally including one or more accessory ingredients.

 The formulations of the invention include those suitable for oral, rectal,
25 topical, buccal (e.g., sub-lingual), parenteral (e.g., subcutaneous, intramuscular, intradermal, or intravenous) and transdermal administration, although the most suitable route in any given case will depend on the nature and severity of the condition being treated and on the nature of the particular active compound which is being used.

30 Formulations suitable for oral administration may be presented in discrete units, such as capsules, cachets, lozenges, or tablets, each containing a predetermined amount of the active compound; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water or water-in-oil

emulsion. Such formulations may be prepared by any suitable method of pharmacy which includes the step of bringing into association the active compound and a suitable carrier (which may contain one or more accessory ingredients as noted above).

5 In general, the formulations of the invention are prepared by uniformly and intimately admixing the active compound with a liquid or finely divided solid carrier, or both, and then, if necessary, shaping the resulting mixture. For example, a tablet may be prepared by compressing or molding a powder or granules containing the active compound, optionally with one or more accessory ingredients. Compressed
10 tablets may be prepared by compressing, in a suitable machine, the compound in a free-flowing form, such as a powder or granules optionally mixed with a binder, lubricant, inert diluent, and/or surface active/dispersing agent(s). Molded tablets may be made by molding, in a suitable machine, the powdered compound moistened with an inert liquid binder.

15 Formulations suitable for buccal (sub-lingual) administration include lozenges comprising the active compound in a flavoured base, usually sucrose and acacia or tragacanth; and pastilles comprising the compound in an inert base such as gelatin and glycerin or sucrose and acacia.

 Formulations of the present invention suitable for parenteral administration
20 conveniently comprise sterile aqueous preparations of the active compound, which preparations are preferably isotonic with the blood of the intended recipient. These preparations may be administered by means of subcutaneous, intravenous, intramuscular, or intradermal injection. Such preparations may conveniently be prepared by admixing the compound with water or a glycine buffer and rendering the
25 resulting solution sterile and isotonic with the blood.

 Formulations suitable for rectal administration are preferably presented as unit dose suppositories. These may be prepared by admixing the active compound with one or more conventional solid carriers, for example, cocoa butter, and then shaping the resulting mixture.

30 Formulations suitable for topical application to the skin preferably take the form of an ointment, cream, lotion, paste, gel, spray, aerosol, or oil. Carriers which may be used include vaseline, lanoline, polyethylene glycols, alcohols, transdermal enhancers, and combinations of two or more thereof.

Formulations suitable for transdermal administration may be presented as discrete patches adapted to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. Formulations suitable for transdermal administration may also be delivered by iontophoresis (*see, for example,*
5 *Pharmaceutical Research* 3(6):318 (1986)) and typically take the form of an optionally buffered aqueous solution of the active compound. Suitable formulations comprise citrate or bis/tris buffer (pH 6) or ethanol/water and contain from 0.01 to 0.2M active ingredient.

10 E. Methods of Use

In addition to the compounds of the formulas described herein, the present invention also provides useful therapeutic methods. For example, the present invention provides a method of inducing cytotoxicity against tumor cells, antitumor-promoting activities, anti-inflammatory activity. More specifically, the present
15 invention provides a method of inducing androgen receptor antagonist activity. The androgen receptor antagonist activity is a useful means of inhibiting androgen related tumor or cancer cell growth.

Cancer cells which may be inhibited include cells from skin cancer, small cell lung cancer, testicular cancer, lymphoma, leukemia, esophageal cancer, stomach
20 cancer, colon cancer, breast cancer, endometrial cancer, ovarian cancer, central nervous system cancer, liver cancer and prostate cancer.

The present invention also provides a method of treating cancer in a subject afflicted with cancer. These subjects also include subjects afflicted with antiandrogen withdrawal syndrome. The method includes administering to the subject in an
25 effective cancer-treating amount a compound of the formulas of the present invention. The method is useful for the treatment of a variety of cancer cells which include but are not limited to skin cancer, small cell lung cancer, testicular cancer, lymphoma, leukemia, esophageal cancer, stomach cancer, colon cancer, breast cancer, ovarian cancer, central nervous system cancer, liver cancer and prostate cancer.

30 Compounds with anti-androgen activity also have the potential to be therapeutically useful for treatment of androgen-potentiated hair disorders such as baldness and hirsutism. Anti-androgenic compounds may also be therapeutically useful as a form of male contraception where it is generally known and understood by

those skilled in the art that androgens are required to maintain spermatogenesis. Additionally, compounds with anti-androgenic activity may be useful for the treatment of behavioral disorders which include, but are not limited to, aggressiveness, violent behavior and sexual aggression. Anti-androgenic compounds may also be therapeutically useful for the treatment of acne due to the altered levels of hormones, including androgens, associated with acne disorders.

Subjects which may be treated using the methods of the present invention are typically human subjects although the methods of the present invention may be useful for veterinary purposes with other subjects, particularly mammalian subjects including, but not limited to, horses, cows, dogs, rabbits, fowl, sheep, and the like. As noted above, the present invention provides pharmaceutical formulations comprising the compounds of formulae described herein, or pharmaceutically acceptable salts thereof, in pharmaceutically acceptable carriers for any suitable route of administration, including but not limited to oral, rectal, topical, buccal, parenteral, intramuscular, intradermal, intravenous, and transdermal administration.

The therapeutically effective dosage of any specific compound will vary somewhat from compound to compound, patient to patient, and will depend upon the condition of the patient and the route of delivery. As a general proposition, a dosage from about 0.1 to about 50 mg/kg will have therapeutic efficacy, with still higher dosages potentially being employed for oral and/or aerosol administration. Toxicity concerns at the higher level may restrict intravenous dosages to a lower level such as up to about 10 mg/kg, all weights being calculated based upon the weight of the active base, including the cases where a salt is employed. Typically a dosage from about 0.5 mg/kg to about 5 mg/kg will be employed for intravenous or intramuscular administration. A dosage from about 10 mg/kg to about 50 mg/kg may be employed for oral administration.

The present invention is explained in greater detail in the following non-limiting examples.

EXAMPLE 1

A. Materials and Methods

Compounds 1–3 were obtained by column chromatography (silica gel, CHCl₃-MeOH) of commercially available curcumin (Aldrich), which contained 2 and 3 as

minor components. Compounds 39 –44 were purchased from Aldrich, Inc (Milwaukee, WI).

Dimethylcurcumin (4). Curcumin (1) in Et₂O and MeOH was treated with excess of diazomethane in ether for 24 h. The solvents were removed *in vacuo* and the residue was purified by silica gel column chromatography and PLC to yield yellow needles of 4 (yield 19.8%); mp 129-130°C (MeOH) (Roughley et al., *J. Chem. Soc. Perkin I*, 2379-2388 (1973))(128-130°C); ¹H NMR (300 MHz, CDCl₃): δ 3.93 (12H, s, OCH₃ x 4), 5.82 (1H, s, 1-H), 6.48 (2H, d, 16Hz), 6.88 (2H, d, *J* = 8Hz), 7.08 (2H, bs), 7.15 (2H, bd), 7.61 (2H, *J* = 16Hz); ¹³CNMR (300 MHz, CDCl₃): δ 55.9, 56.0, 101.3, 109.8, 111.1, 122.0, 122.6, 128.1, 140.4, 149.2, 151.0, 183.2.

Preparation of pyrazol derivative (8). To a solution of 1–4 in butanol and ethanol were added histidine hydrazide (1 equiv.), acetic acid and *p*-TsOH. The solution was refluxed for 24h, and then the solvent was removed *in vacuo*. The residue was purified by silica gel column chromatography and PLC.

Compound 8. Yellow powder (yield 17.5%), mp 166-168°C (MeOH); ¹H NMR (300 MHz, CDCl₃): δ 3.92 (6H, s, OCH₃ x 2), 3.94 (6H, s, OCH₃ x 2), 6.62 (1H, s, 1-H), 6.86 (2H, d, *J* = 8Hz), 6.93 (2H, d, *J* = 16Hz), 7.04 (2H, dd, *J* = 8, 2Hz), 7.06 (2H, bs), 7.05 (2H, d, *J* = 16Hz); ¹³CNMR (300 MHz, CDCl₃): δ 55.8, 55.9, 99.6, 108.6, 111.2, 115.8, 120.1, 129.7, 130.6, 149.1, 149.3; Anal. calcd. for C₂₃H₂₄N₂O₄·1.1/4H₂O: Theory: C, 66.57; H, 6.44; N, 6.75. Found C, 66.44; H, 6.19; N, 6.27.

Monomethylcurcumin (9). Curcumin (1) in MeOH was treated with excess diazomethane in Et₂O for 24 h. After removal of solvents, the residue was purified by silica gel column chromatography and PLC to yield a yellow amorphous solid (yield 20%); mp 89-91°C, [α]_D –3.6 (*c*=1.14, CHCl₃); ¹H NMR (300 MHz, CDCl₃): δ 3.93 (9H, s, OCH₃ x 3), 5.81 (1H, s, 1-H), 5.94 (1H, bs, OH), 6.49 (2H, bd, *J* = 15Hz), 6.93 (1H, d, *J* = 8Hz), 6.97 (1H, d, *J* = 8Hz), 7.10 (4H, m), 7.60 (2H, bd, *J* = 15Hz); EIMS *m/z* 382 (M⁺), HRFABMS 382.1396 (M+H⁺) (calcd for C₂₂H₂₂O₆: 382.1416).

Hydrogenation of 1, 4 and 10 (11–18). A solution of starting material in EtOAc was shaken with 10% Pd-C under H₂ (45psi) overnight using a Parr's apparatus. The solution was filtered and concentrated *in vacuo* to give a residue, which was purified by silica gel column chromatography and PLC.

5 **Tetrahydrocurcumin (11).** White powder, mp 92-93°C (Roughley et al., *J. Chem. Soc. Perkin I*, 2379-2388 (1973), 95-96°C), ¹H NMR (300 MHz, CDCl₃): δ 2.53-2.58 (3H, m), 2.78-2.88 (5H, m), 3.87 (6H, s, OCH₃ x 2), 5.43 (1H, s, 1-H), 5.50 (2H, s, ArOH), 6.65 (2H, d, *J* = 8Hz), 6.69 (2H, s), 6.83 (2H, d, *J* = 8Hz); ¹³CNMR (300 MHz, CDCl₃): δ 31.3, 40.4, 55.8, 99.8, 111.0, 114.3, 120.8, 132.6, 144.0, 146.4
10 and 193.2.

Hexahydrocurcumin (12). White powder, mp 87-88°C (Roughley, P.J. et al., *J. Chem. Soc. Perkin I*, 2379-2388 (1973), 78-80°C), ¹H NMR (300 MHz, CDCl₃): δ 1.60-1.81 (2H, m), 2.53-2.97 (8H, m), 3.85 (6H, s, OCH₃ x 2), 4.06 (1H, m, 2-H), 6.70 (4H, m), 6.80 (2H, d, *J* = 8Hz); ¹³CNMR (300 MHz, CDCl₃): δ 29.7, 31.7,
15 38.8, 45.8, 49.8, 56.3, 67.4, 111.5, 111.6, 114.8, 114.9, 121.2, 121.4, 133.0, 134.2, 144.2, 144.5, 146.9, 147.9 and 211.9.

Octahydrocurcumin (13). Colorless oil, ¹H NMR (300 MHz, CDCl₃): δ 1.61 (2H, m), 1.75 (4H, m), 2.53-2.70 (4H, m), 3.80 (6H, s, OCH₃ x 2), 3.91 (2H, brs), 6.13 (2H, s, ArOH), 6.65 (2H, d, *J* = 8Hz), 6.69 (2H, bs) 6.82 (2H, bd, *J* = 8Hz),
20 ¹³CNMR (300 MHz, acetone-*d*₆): δ 31.1, 39.8, 42.6, 35.6, 72.0, 111.0, 114.3, 120.6, 133.6, 143.6 and 146.4.

Compound 14. White powder (yield 26.0%), mp 60-61°C, ¹H NMR (300 MHz, CDCl₃): δ 2.56 (3H, m), 2.86 (5H, m), 3.85 (12H, s, OCH₃ x 4), 5.44 (1H, s, 1-H), 6.71 (4H, m), 6.78 (2H, bd); Anal. calcd. for C₂₃H₂₈O₆·1/4H₂O: Theory: C, 68.21; H, 7.09. Found C, 68.25; H, 7.06.
25

Compound 15. White powder (yield 20.0%), mp 94-95°C, ¹H NMR (300 MHz, CDCl₃): δ 1.65-1.80 (2H, m), 2.53-2.84 (8H, m), 3.85 (12H, s, OCH₃ x 4), 4.05(1H, bs, 2'-H), 6.68-7.23 (4H, m), 6.79 (2H, bd), Anal. calcd. for C₂₃H₃₀O₆·1/4H₂O: Theory: C, 67.88; H, 7.55. Found C, 67.73; H, 7.49.

Compound 16. Colorless oil (yield 4.2%), mp 60-61°C, ^1H NMR (300 MHz, CDCl_3): δ 1.55-1.65 (4H, m), 1.73-1.82 (3H, m), 2.60-2.72 (3H, m), 3.86 (6H, s, $\text{OCH}_3 \times 2$), 3.87 (8H, bs, $\text{OCH}_3 \times 2$, 2,2'-H), 6.72-6.78 (4H, m), 6.79 (2H, bd), 7.27 (2H, s, OH $\times 2$), EIMS m/z : 404 (M^+), HRFAB-MS m/z 404.219070 ($\text{M}+\text{H}$) $^+$
5 (calcd for $\text{C}_{23}\text{H}_{32}\text{O}_6$: 404.2198891).

Compound 17. Colorless oil (yield 5.9%), ^1H NMR (300 MHz, CDCl_3): δ 1.10 (3H, d), 1.80 (1H, m), 2.43-2.82 (8H, m), 3.86 (6H, s, $\text{OCH}_3 \times 2$), 3.87 (6H, s, $\text{OCH}_3 \times 2$), 3.94 (1H, bs, 2'-H), 6.70-6.78 (6H, m), EIMS m/z 416 (M^+).

Compound 18. Colorless oil (yield 6.95%), ^1H NMR (300 MHz, CDCl_3): δ
10 0.95 (3H, d, 1- CH_3), 1.52 (1H, m), 1.84 (2H, m), 2.67 (6H, m), 3.83 (14H, bs, $\text{OCH}_3 \times 4$, 2, 2'-H), 6.78 (6H, m); EIMS m/z : 418 (M^+), HRFAB-MS m/z 418.236618 ($\text{M}+\text{H}$) $^+$ (calcd for $\text{C}_{24}\text{H}_{34}\text{O}_6$: 418.2355392).

Preparation of 19 and 20. A mixture of curcumin (1, 100 mg, 0.81 mmol) in acetone (20 mL) with methylchloroacetate (2 mL) and NaI (20 mg) was refluxed with
15 anhydrous potassium carbonate (176 mg) for 24 h with stirring. After filtration and removal of solvent, the residue was purified by silica gel column chromatography to yield the corresponding methyl acetates 19 and 20.

Compound 19: Yellow powder (yield 20.0%), mp 60-61°C, mp 66-67°C, $[\alpha]_D$ -2.4 ($c = 2.08$, CHCl_3); ^1H NMR (300 MHz, acetone- d_6): δ 3.73 (3H, s, -
20 COOCH_3), 3.86 (6H, s, $\text{OCH}_3 \times 2$), 4.79 (2H, s, $\text{O-CH}_2\text{-COO}$), 5.99 (1H, s, 1-H), 6.70 and 6.73 (both 1H, d, $J = 15.3\text{Hz}$), 6.88 (1H, d, $J = 8\text{Hz}$), 6.94 (1H, d, $J = 8\text{Hz}$), 7.17 (2H, m), 7.33 (2H, m), 7.59 and 7.61 (both 1H, d, $J = 15.3\text{Hz}$), ^{13}C NMR (300 MHz, CDCl_3): δ 51.8, 55.9, 55.9, 65.9, 101.4, 111.2, 111.6, 114.3, 115.9, 121.8, 122.6, 123.0, 123.5, 127.6, 128.7, 129.8, 140.3, 141.3, 148.4, 149.8, 150.0, 150.4,
25 169.4, 183.4, 184.6; Anal. calcd. for $\text{C}_{24}\text{H}_{24}\text{O}_8 \cdot 3/4\text{H}_2\text{O}$: Theory: C, 63.50; H, 5.66. Found C, 63.53; H, 5.65.

Compound 20: Yellow powder (yield 20.0%), mp 141-142°C (MeOH), $[\alpha]_D$ -0.29 ($c = 5.86$, CHCl_3); ^1H NMR (300 MHz, CDCl_3): δ 3.80 (6H, s), 3.93 (6H, s), 4.73 (4H, s, $\text{O-CH}_2\text{-COO} \times 2$), 5.82 (1H, s, 1-H), 6.50 (2H, d, $J = 16\text{Hz}$), 6.79 (2H, d,

$J = 8\text{Hz}$), 7.09 (4H, bs), 7.58 (2H, d, $J = 16\text{Hz}$), ^{13}C NMR (300 MHz, CDCl_3): δ 52.3, 56.0, 66.0, 101.4, 110.7, 113.6, 122.0, 122.7, 129.5, 140.1, 149.0, 149.7, 169.0, 183.1; Anal. calcd. for $\text{C}_{27}\text{H}_{28}\text{O}_{10} \cdot 1/2\text{H}_2\text{O}$: Theory: C, 62.18; H, 5.60. Found C, 62.31; H, 5.57.

5 **Compound 21:** Yellow amorphous solid (yield 3.0%), ^1H NMR (300 MHz, CDCl_3): δ 2.58 (2H, m), 2.95 (2H, m), 7.12 (2H, d, $J=15\text{Hz}$), 7.40 (6H, m), 7.60 (4H, m), 7.81 (2H, d, $J = 15\text{Hz}$), 12.65 (1H, bs); Anal. calcd. for $\text{C}_{22}\text{H}_{20}\text{O}_4$: Theory: C, 75.84, H, 5.79. Found C, 75.56, H, 5.74.

10 **Compound 22:** Yellow amorphous solid (yield 25.0%), Anal. calcd. for $\text{C}_{26}\text{H}_{28}\text{O}_8$: Theory: C, 66.66, H, 6.02. Found C, 66.38, H, 6.16.

Compound 23: Yellow powder (yield 45.0%), mp 144-146°C (MeOH) (Pedersen et al., *Liebigs Ann. Chem.* 1557-1569 (1985), 71-73°C (CH_2Cl_2)) Anal. calcd. for $\text{C}_{24}\text{H}_{26}\text{O}_8 \cdot 2 \cdot 1/2\text{H}_2\text{O}$: Theory: C, 59.87; H, 5.23. Found C, 59.94; H, 5.11.

15 The structures of 1-4, 10-13, 22, and 23 were confirmed by comparison of their physical spectral data with those reported in the literature. Pedersen et al., *Liebigs Ann. Chem.* 1557-1569 (1985), Roughley et al., *J. Chem. Soc. Perkin I*, 2379-2388 (1973).

B. Suppression of DHT-Mediated Transcription Activity.

20 **Cell Culture and Transfections.** Human prostate cancer DU145 and PC-3 cells were maintained in Dulbecco's minimum essential medium (DMEM) containing penicillin (25 units/mL), streptomycin (25 $\mu\text{g/mL}$), and 10% fetal calf serum (FCS). For AR transactivation assay, PC-3 cells were transfected with an AR expression plasmid and reporter gene. Because of a low content of endogenous AR coactivators, 25 DU-145 cells were transfected with expression plasmids for AR and ARA70, and reporter gene. The conditions were followed as previously described in Miyamoto et al., *Proc. Natl. Acad. Sci. USA*, 95, 7379-7384 (1998), with minor modifications.

30 Transfections were performed using the SuperFect kit according to manufacturer's procedures (Qiagen, Chatsworth, CA). Briefly, 1×10^5 cells were plated on 35-mm dishes 24 h before transfection, and then a reporter plasmid, MMTV-Luciferase, which contains MMTV-LTR promoter and AR-binding element,

was co-transfected with an AR expression plasmid (wild type or mutant), or pSG5ARA70. PRL-TK was used as an internal control for transfection efficiency. The total amount of DNA was adjusted to 3.0 g with pSG5 in all transcriptional activation assays. After a 2 h transfection, the medium was changed to DMEM-10% charcoal stripped serum medium, and 14–16 h later, the cells were treated with DHT, antiandrogen, or test compounds. After another 14–16 h, the cells were harvested and tested for luciferase activity in luciferase assays (Promega, Dual Luciferase Assay System, Madison, WI). Data were expressed in relative luciferase activity as compared to an internal luciferase positive control.

10

C. Results and Discussion

The aim of this work was to investigate novel curcumin analogues for antiandrogen receptor antagonist activity. The synthesis and evaluation of novel curcumin analogues as antiandrogen receptor antagonists and antitumor agents are reported herein.

Forty-seven curcumin derivatives (1-47) were tested for antagonistic activity against the AR using two different human prostate cancer cells, PC-3 and DU-145 (Figures 3A–C). The parental compound, curcumin (1), was inactive in all cases. However, dimethylated curcumin (4) showed significant antagonistic activity (reducing 70% of DHT-induced AR activity) when assayed in PC-3 cells transfected with wild-type AR and was more potent than HF (which reduced 16% of DHT-induced AR activity, Figure 3A). Compound 4 also showed the highest antagonist activity when assayed in DU-145 cells transfected with a mutant LNCaP AR and ARA70 (showing a 45% reduction in DHT-induced AR activity, Figure 3B), indicating that compound 4 is an effective antagonist for both normal and mutant AR.

To determine the structural requirements for AR antagonist activity in this series of compounds, a structure-activity relationship (SAR) study was conducted in a PC-3 cell assay system. Compared with 4, monomethylated curcumin (9) lacks one *O*-methyl groups at the *p*-position on one benzene ring, and was significantly less active than 4 (Figure 3B). Thus, the bis(3,4-dimethoxyphenyl) groups of 4 are important to the activity. Introducing a methyl group at C-4 of 4 (10) resulted in decreased activity (Figure 3B). Compounds 14 and 15, which were obtained by hydrogenation of 4, were as potent as HF with an 18% reduction in DHT-induced AR

activity, but were considerably less active compared to **4** (Figure 3A). Converting the β -diketone moiety of **4** to the corresponding pyrazol derivative **8** greatly reduced the activity. Furthermore, 1,3-bis(3,4-dimethoxyphenyl)-1,3-propanedione (**39**), which contains the bis-aryl groups found in **4** but lacks the conjugated double bonds, was less active than **4** (Figure 3A and 3B), indicating that the conjugated double bonds also contribute to the activity of **4**. These observations suggested that the bis(3,4-dimethoxyphenyl) groups and the conjugated β -diketone moiety are crucial for the activity.

Data in Figure 3C show a somewhat different cell assay system where antiandrogen activity was assayed in DU-145 cells transfected with wild-type AR and ARA70. Compounds **4**, **20**, **22**, **23**, and **39** showed comparable or more potent antiandrogen activity than HF in this assay system. Compounds **20** and **22** were almost equipotent (54% and 53.8% reduction, respectively) and were slightly more active than **4** (49.9%). Because curcumin (**1**) itself was not active, introducing either methoxycarbonylmethyl groups at the phenolic hydroxyls (**20**) or an ethoxycarbonyl ethyl group at C-4 (**22**) greatly contributed to the anti-AR activity in DU-145 cells in the presence of wild-type AR and ARA70.

In this study, we also examined the antiandrogen activity of fluorodiarylheptanoids **24–29** and cyclic diarylheptanoids **30–38**. Compounds **24–29** have fluorine or trifluoromethyl substituents on both benzene rings, but showed weak activity or were inactive. Among the cyclic diarylheptanoids **30–38**, compound **30** was the most active and was almost as active as HF (Figure 3A and 3C). The remaining cyclic diarylheptanoids showed weak antagonistic activity.

In conclusion, we have prepared a number of curcumin analogues and evaluated their potential antiandrogen activity in three different assay conditions using human prostate cancer cell lines. Compounds **4** showed promising antiandrogen activities in all assays. Compounds **4**, **20**, **22**, **23** and **39** have been identified as a new class of antiandrogen agents. The SAR study revealed that bis(3,4-dimethoxyphenyl) moieties, a conjugated β -diketone, and an ethoxycarbonyl ethyl group at the C-4 position play important roles in the antagonistic activity.






EXAMPLE 2







Synthesis and Characterization of Additional Curcumin Analogs






Additional curcumin analogs are shown in Table 1 below.

5

Table 1. Curcumin analogues.

Compound	Structure
LL-7	
LL-10	
LL-17	
LL-18	
LL-27-B	


Compound	Structure
LL-32-B	 <chem>COc1cc(OC)c(OC)cc1/C=C/C(O)/C=C/C(=O)/C=C/c1ccc(OC)c(OC)c1</chem>
LL-35	 <chem>COc1cc(OC)c(OC)cc1/C=C/C(O)/C=C/C(=O)/C=C/c1ccc(OC)c(OC)c1</chem>
LL-36	 <chem>COc1cc(OC)c(OC)cc1/C=C/C(O)/C=C/C(=O)/C=C/c1ccc(OC)c(OC)c1</chem>
LL-41	 <chem>COc1cc(OC)c(OC)cc1/C=C/C(=O)/C=C/c1ccc(OC)c(OC)c1</chem>
LL-46	 <chem>COc1cc(OC)c(OC)cc1/C=C/C(=O)C(C(=O)OCC)C(=O)/C=C/c1ccc(OC)c(OC)c1</chem>
LL-55	 <chem>COc1cc(OC)c(OC)cc1/C=C/C(O)CC(=O)/C=C/c1ccc(OC)c(OC)c1</chem>

Compound	Structure
LL-61	 <chem>COc1cc(OC)c(C=C/C(O)/C(=O)C(=O)/C=C/c2cc(OC)c(OC)c2)cc1CCOC(=O)CC</chem>
LL-62	 <chem>COc1cc(OC)c(C=C/C(=O)C(F)C(=O)/C=C/c2cc(OC)c(OC)c2)cc1</chem>
LL-65	 <chem>COc1cc(OC)c(C=C/C(O)/C(=O)C(=O)/C=C/c2cc(OC)c(OC)c2)cc1CCOC(=O)CC</chem>
LL-66	 <chem>COc1cc(OC)c(C=C/C(O)/C(=O)C(=O)/C=C/c2cc(OC)c(OC)c2)cc1CCOC(=O)CC</chem>
LL-80	 <chem>COc1cc(OC)c(C=C/C(O)/C(=O)C(=O)/C=C/c2cc(OC)c(OC)c2)cc1CCOC(=O)CC</chem>

Compound	Structure
HOJ-7	
HOJ-9	
HOJ-10	

The foregoing compounds are synthesized as follows:


5



LL-7: The synthetic method was modified from the strategy developed by Pedersen et al. Acetylacetone (0.2 ml, 2 mmol) and boric anhydride (100 mg, 1.4 mmol) were dissolved in 15 ml of ethyl acetate. The solution was stirred at 70°C for 0.5 hour. 5-hydroxymethylfuran (506 mg, 4 mmol) and tributyl borate (1.08 ml, 4 mmol) were added. The mixture was stirred for 30 min. Then butylamine (0.3 ml, 3 mmol) dissolved in 4 ml of EtOAc was added dropwise during 15 min. The stirring continued for 5 hours at 85 °C. The mixture was then hydrolyzed by adding 8 ml of 1N HCl and stirring for 0.5 hour at 60 °C. The organic layer was separated. The aqueous layer was extracted with ethyl acetate. The combined organic layers were washed until neutral, dried with anhydrous sodium sulphate. The solven was removed


in vacu. The crude products were purified by CombiFlash® column chromatography eluting with hexane-EtOAc. 68 mg red powder was obtained in 12% yield. ^1H NMR (300 MHz, CDCl_3): δ 4.67 (4H, s), 5.74 (H, s), 6.40 (2H, d, $J = 3.3$ Hz), 6.53 (2H, d, $J = 15.3$ Hz), 6.58 (2H, d, $J = 3.3$ Hz), 7.43 (2H, d, $J = 15.6$ Hz).

5



LL-10: vanillin (1.52 g, 10 mmol) and tributyl borate (2.7 ml, 10 mmol) were dissolved in 8 ml ethyl acetate. To the stirring solution, acetone (0.37 ml, 5 mmol) was added. The solution was stirring for 10 min. Butylamine (1 ml, 10 mmol) in 5 ml EA was added dropwise. The solution was stirring for hours at 50 °C. 7 ml of 1N HCl was added. The mixture was stirring for 0.5 hour at 50 °C. ^1H NMR (300 MHz, CDCl_3): δ 3.94 (6H, s), 6.90 (2H, d, $J = 8.4$ Hz), 6.95 (2H, d, $J = 15.6$ Hz), 7.11 (2H, d, $J = 1.8$ Hz), 7.20 (2H, dd, $J = 1.8$ Hz, $J = 8.4$ Hz), 7.68 (2H, d, $J = 15.6$ Hz).


15




LL-1: Acetylacetone (3.1 ml, 30 mmol) and boric anhydride (1.5 g, 21 mmol) were dissolved in 20 ml of ethyl acetate. The solution was stirred at 70°C for 1 hour. To the solution vanillin (1.52 g, 10 mmol) and tributyl borate (2.7 ml, 10 mmol) were added. The mixture was stirred for 30 min. At 85 °C, butylamine (1 ml, 10 mmol) dissolved in 7 ml of EtOAc was added dropwise during 15 min. The stirring continued for 1 hour at 100 °C. The mixture was then hydrolyzed by adding 20 ml of 1N HCl at 50 °C and stirring for 0.5 hour at 50 °C. The organic layer was separated. The aqueous layer was extracted with ethyl acetate. The combined organic layers were washed until neutral, dried with anhydrous sodium sulphate. The solven was

removed in vacu. The crude products were purified by flash column chromatography eluting with hexane-EtOAc. Yellow powder was obtained in 49% yield. ^1H NMR (300 MHz, CDCl_3): δ 2.16 (3H, s), 3.94 (3H, s), 5.63 (H, s), 6.33 (H, d, $J=15.9$ Hz), 6.92 (H, d, $J=8.1$ Hz), 7.01 (H, d, $J=1.8$ Hz), 7.10 (H, dd, $J=8.1$ Hz, $J=1.8$ Hz), 7.53 (H, d, $J=15.9$ Hz).

LL-17: At 80 °C, **LL-1** (468, 2 mmol) and boric anhydride (100 mg, 1.4 mmol) were dissolved in 10 ml of EtOAc. The solution was stirring for 1 hour. To the mixture, 10 ml of EtOAc solution containing 3,4-dimethoxybenzaldehyde (365 mg, 2.2 mmol) and tributyl borate (0.54 ml, 2 mmol) was added. After stirring for 0.5 hour, piperidine (0.08 ml) was added to the mixture. After stirring for 2 hours, 4 ml of 0.4 N HCl was added at 50 °C. The mixture was vigorously stirred at 50 °C for 0.5 hour. The combined organic layers were washed until neutral, dried with anhydrous sodium sulphate. The solven was removed in vacu. The crude products were purified to offer 67.2 mg **LL-17** in 8.8% yield. ESI-MS m/z 381.1 ($\text{M}-1$) $^+$ ^1H NMR (300 MHz, CDCl_3): δ 3.94 (9H, s), 5.81 (H, s), 6.48 (2H, d, $J=12.6$ Hz), 6.87- 7.14 (6H, m), 7.60 (2H, d, $J=12.6$ Hz).




LL-18: the same procedure as the preparation of **LL-17**. 50% yield (started with **LL-1** and 4-hydroxybenzaldehyde). Orange powder. ESI-MS m/z 337.1 ($\text{M}-1$) $^+$; ^1H NMR (300 MHz, CD_3COCD_3): δ 3.93 (3H, s), 5.98 (H, s), 6.70 (2H, d, $J=15.9$ Hz), 6.87-6.92 (3H, m), 7.19 (H, dd, $J=8.4$ Hz, $J=1.8$ Hz), 7.35 (H, d, $J=1.8$ Hz), 7.56-7.64 (4H, m).




LL-27-B: Acetylacetone (3.1 ml, 30 mmol) and boric anhydride (1.5 g, 21 mmol) were dissolved in 20 ml of ethyl acetate. The solution was stirred at 70°C for 0.5 hour. 3-hydroxy-4-methoxybenzaldehyde (1.52 g, 10 mmol) and tributyl borate (2.7 ml, 10 mmol) were added. The mixture was stirred for 30 min at 70 °C. Then butylamine (1 ml, 10 mmol) dissolved in 7 ml of EtOAc was added dropwise during 15 min at 85 °C. The stirring continued for 1 hour at 100 °C. The mixture was then hydrolyzed by adding 20 ml of 1N HCl and stirring for 0.5 hour at 50 °C. The organic layer was separated. The aqueous layer was extracted with ethyl acetate. The combined organic layers were washed until neutral, dried with anhydrous sodium sulphate. The solvent was removed in vacu. The crude products were purified. 1.04 g **LL-27-A** and 248 mg **LL-27-B** were obtained. ESI-MS m/z 367.1 ($M-1$)⁺; ¹H NMR (300 MHz, CD₃COCD₃): δ 4.01 (6H, s), 6.01 (H, s), 6.66 (2H, d, J = 15.6 Hz), 7.03 (2H, d, J = 8.1 Hz), 7.20 (2H, dd, J = 2.1 Hz, J = 8.1 Hz), 7.26 (2H, d, J =2.1 Hz), 7.65 (2H, d, J = 15.3 Hz).


15




LL-32-B: Acetylacetone (3.1 ml, 30 mmol) and boric anhydride (1.5 g, 21 mmol) were dissolved in 20 ml of ethyl acetate. The solution was stirred at 70°C for 0.5 hour. 2,3,4-trimethoxybenzaldehyde (1.96 g, 10 mmol) and tributyl borate (2.7 ml, 10 mmol) were added. The mixture was stirred for 30 min at 70 °C. Then butylamine (1 ml, 10 mmol) dissolved in 7 ml of EtOAc was added dropwise during 15 min at 85 °C. The stirring continued for 1 hour at 100 °C. The mixture was then hydrolyzed by adding 20 ml of 1N HCl and stirring for 0.5 hour at 50 °C. The organic layer was separated. The aqueous layer was extracted with ethyl acetate. The combined organic layers were washed until neutral, dried with anhydrous sodium sulphate. The solvent was removed in vacu. The crude products were purified. 1.25 g **LL-32-A** and 150 mg **LL-32-B** were obtained. ¹H NMR (300 MHz, CDCl₃): δ 3.89 (6H, s), 3.90 (6H, s), 3.94 (6H, s), 5.83 (H, s), 6.64 (2H, d, J = 15.9 Hz), 6.71 (2H, d, J = 9 Hz), 7.31 (2H, d, J = 8.7 Hz), 7.85 (2H, d, J =16.2 Hz).



LL-35: The same procedure as the preparation of **LL-17**. 55% yield (started with **LL-32-A** and 3,4-dimethoxybenzaldehyde). ESI-MS m/z 427.2 ($M+H$) $^+$, 449.3 ($M+Na$) $^+$; 1H NMR (300 MHz, $CDCl_3$): δ 3.89(3H, s), 3.91 (3H, s), 3.93 (3H, s), 3.94 (6H, s), 5.83 (H, s), 6.51 (H, d, $J=15.9$ Hz), 6.63 (H, d, $J=15.9$ Hz), 6.71 (H, d, $J=8.7$ Hz), 6.89 (H, d, $J=8.4$ Hz), 7.09 (H, d, $J=2.1$ Hz), 7.15 (H, dd, $J=1.8$ Hz, $J=8.7$ Hz), 7.31 (H, d, $J=8.7$ Hz), 7.61 (H, d, $J=15.9$ Hz), 7.85 (H, d, $J=16.2$ Hz).




LL-36: The same procedure as the preparation of **LL-17**. 48% yield (started with **LL-33-A** and 3,4-dimethoxybenzaldehyde). ESI-MS m/z 427.2 ($M+H$) $^+$, 449.3 ($M+Na$) $^+$; 1H NMR (300 MHz, $CDCl_3$): δ 3.89(3H, s), 3.90 (3H, s), 3.93 (3H, s), 3.94 (3H, s), 3.95 (3H, s), 5.84 (H, s), 6.50 (H, d, $J=15.6$ Hz), 6.51(H, s), 6.57 (H, d, $J=16.2$ Hz), 6.88 (H, d, $J=8.1$ Hz), 7.06 (H, s), 7.08 (H, d, $J=2.1$ Hz), 7.14 (H, dd, $J=2.1$ Hz, $J=8.1$ Hz), 7.60 (H, d, $J=15.9$ Hz), 7.96 (H, d, $J=15.9$ Hz).



LL-41: To a solution of acetone (0.36 ml, 5 mmol) and 3,4-dimethoxybenzaldehyde (1.66 g, 10 ml) in 50 ml of a 0.25 M solution of aqueous NaOH was added 1.5 ml of a 25% w/w aqueous solution of cetyltrimethylammonium


bromide. The mixture was allowed to stir vigorously at room temperature for 20 hours, diluted with brine and extracted with ethyl acetate. The ethyl acetate solution was concentrated and then underwent column chromatography. 1.34 g bright yellow powder was obtained. 75% yield. ESI-MS m/z 355.2 ($M+H$)⁺; ¹H NMR (300 MHz, CD₃COCD₃): δ 3.94 (6H, s), 3.96 (6H, s), 6.90 (2H, d, J = 8.4 Hz), 6.95 (2H, d, J = 15.9 Hz), 7.15 (2H, d, J = 1.8 Hz), 7.20 (2H, dd, J = 1.8 Hz, J = 8.4 Hz), 7.69 (2H, d, J = 15.9 Hz).



10

LL-46: The same procedure as the preparation of LL-17. 38% yield (started with LL-40 and 3,4-dimethoxybenzaldehyde). ESI-MS m/z 483.4 ($M+H$)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.25 (3H, t, J = 7.2 Hz), 2.32-2.37 (2H, m), 2.55 (2H, t, J = 7.8 Hz), 2.95 (0.5H, t, J = 7.8 Hz), 3.91-3.97 (9H, m), 4.14 (2H, q, J = 7.2 Hz), 6.70 (H, dd, J = 4.2 Hz, J = 15.6 Hz), 6.84-7.19 (7H, m), 7.62-7.76 (2H, m);


15




LL-55: To a stirred solution of LDA (0.29 ml, 2M hex/THF, 0.58 mmol) in 3 ml of THF at -78 °C was added a solution of 3,4-dimethoxycinnamone (100 mg, 0.48 mmol) in THF (3 ml). After 15 min, 3,4-dimethoxycinnamaldehyde (85 mg, 0.44 mmol) in THF (3 ml) was added. The mixture was stirred for 20 min at -78 °C. And then the mixture was quenched with saturated NH₄Cl. The solution was allowed to warm to ambient temperature and extracted with ethyl acetate. 22 mg LL-55 was obtained by column chromatography. 13% yield. ¹H NMR (300 MHz, CD₃COCD₃): δ 2.93 (2H, d), 3.80 (6H, s), 3.85 (6H, s), 4.13 (H, d), 6.25 (H, dd, J = 6 Hz, J = 15.9 Hz),

25

6.58 (H, d, $J=15.9$ Hz), 6.80 (H, d, $J=16.2$ Hz), 6.88 (H, d, $J=8.7$ Hz), 6.90 (H, dd, $J=0.9$ Hz, $J=8.7$ Hz), 7.00 (H, d, $J=8.4$ Hz), 7.04 (H, d, $J=0.9$ Hz), 7.25 (H, dd, $J=0.9$ Hz, $J=8.7$ Hz), 7.34 (H, d, $J=0.9$ Hz), 7.61 (1H, d, $J=16.2$ Hz).




LL-61: 4-thoxycarbonyl ethyl curcumin (153.1 mg, 0.33 mmol) and dihydropyran (0.73 mL, 7.32 mmol) were dissolved in 3 mL of dry dichloromethane containing PPTS (8.3 mg, 0.033 mmol). The solution was stirred at room temperature for 48 hours. The solution was then washed by water. The solvent was removed in vacu. The crude products was purified by CombiFlash® chromatography eluting with hexane-EtOAc to give **LL-61** (134 mg), 59% yield, yellow powder, mp 60- 61 °C; ESI MS m/z 635.2 ($M-1$)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.25 (3H, t), 1.57-2.17 (12H, m), 2.96 (0.57H, t), 3.62 (4H, t), 3.91 (6H, s), 4.13 (2H, q), 5.47 (2H, t), 6.72 (2H, d, $J=15.6$ Hz), 6.90-7.18 (6H, m), 7.44 (2H, d, $J=15.6$ Hz);



LL-62: A DMF solution (2 mL) of 1,7-bis-(3,4-dimethoxyphenyl)-4-methyl-1,6-heptadiene-3,5-dione (50.3mg, 0.12 mmol) was added to an oil-free suspension of NaH (10 mg of 60%, 6 mg, 0.4 mmol) in DMF (2 mL) under nitrogen at 0 °C. The solution was stirred at 0 °C for 30 min and then at room temperature for 2 h. To the sodium salt solution SelectfluorTM (86.7 mg, 0.25 mmol) in DMF (2 mL) was added. After stirring for 2 h, the solution was extracted with EtOAc and washed with 5% H₂SO₄ (10 mL) and subsequently saturated NaHCO₃ solution (10 mL). Then the solvent was evaporated in vacuo. The crude products were purified by flash column chromatography eluting with hexane-EtOAc to afford **LL-62** (25 mg) in 49% yield. yellow powder. ¹H NMR (300 MHz, CDCl₃): δ 1.97 (3H, s), 3.90 (6H, s), 3.95 (6H,

s), 6.83 (2H, d, $J=7.8$ Hz), 6.87 (2H, dd, $J=3$ Hz, $J=15.6$ Hz), 6.90 (2H, d, $J=1.8$ Hz), 6.95 (2H, dd, $J=1.8$ Hz, $J=7.8$ Hz), 7.75 (2H, d, $J=15.6$ Hz);



5

LL-64B: Recrystallized curcumin (1.08 g, 2.94 mmol) and dihydropyran (2 mL, 20 mmol) were dissolved in 30 mL of dry dichloromethane containing PPTS (74 mg, 0.29 mmol). The solution was stirred at room temperature for 24 hours. The solution was then washed by water. The solvent was removed in vacuo. The crude products was purified by CombiFlash® chromatography eluting with hexane-EtOAc to give **LL-64B** (1.12 g), 66.8% yield, yellow powder; ESI MS m/z 535.0 ($M-1$)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.57-2.17 (12H, m), 3.62 (4H, t), 3.91 (6H, s), 5.47 (2H, t), 5.83 (1H, s), 6.50 (2H, d, $J=15.9$ Hz), 7.09-7.16 (6H, m), 7.60 (2H, d, $J=15.9$ Hz); Anal. Calcd (theoretical) for C₃₁H₃₆O₈·1/4H₂O: C, 68.81; H, 6.80. Found: C, 68.86; H, 6.83.


15

LL-65: A THF solution (3 mL) of **LL-64B** (55mg, 0.10 mmol) was added to an oil-free suspension of NaH (7 mg of 60%, 4.2 mg, 0.17 mmol) in THF (2 mL) under nitrogen at 0 °C. The solution was stirred at 0 °C for 30 min and then at room temperature for 2 h. To the sodium salt solution ethyl propiolate (0.02 mL, mmol) was added. After stirring for 2 h, the solution was extracted with EtOAc and washed with 5% H₂SO₄ (10 mL) and subsequently saturated NaHCO₃ solution (10 mL). Then the solvent was evaporated in vacuo. The crude products were purified by CombiFlash® chromatography eluting with hexane-EtOAc to afford **LL-65** (39.4 mg) 62% yield, orange powder; mp 72- 73 °C; ESI MS m/z 634.7 M⁺; ¹H NMR (300 MHz, CDCl₃): δ

20

1.34 (3H, t), 1.5-2.2 (12H, m), 3.62 (4H, t), 3.92 (6H, s), 4.28 (2H, q), 5.49 (2H, t) 5.96 (H, d, $J=15.6$ Hz), 7.00 (2H, d, $J=15.6$ Hz), 7.08-7.16 (6H, m), 7.76 (2H, d, $J=15.3$ Hz), 7.83 (H, d, $J=15.9$ Hz); Anal. Calcd (theoretical) for $C_{36}H_{42}O_{10}$: C, 68.12; H, 6.67. Found: C, 67.82; H, 6.73.

- 5 **LL-66**: The EtOH (2.5 mL) solution of **LL-65** (14.8 mg, 0.023 mmol) and PPTS (5 mg, 0.02 mmol) was stirred at room temperature for 3 h. The solution was evaporated in vacuo. **LL-66** (10 mg) was obtained by CombiFlash® chromatography eluting with hexane-EtOAc. 93% yield, orange powder, mp 106- 106.5 °C; ESI MS m/z 465.2 ($M-1$)⁺; ¹H NMR (300 MHz, $CDCl_3$): δ 1.34 (3H, t), 3.95 (6H, s), 4.29 (2H, quart), 5.96 (2H, d, $J=15.6$ Hz), 6.95 (2H, d, $J=8.2$ Hz), 6.96 (1H, d, $J=15.6$ Hz), 10 7.05 (2H, d, $J=2.1$ Hz), 7.17 (2H, dd, $J=8.2$ Hz, $J=2.1$ Hz), 7.75 (2H, d, $J=15.3$ Hz), 7.90 (1H, d, $J=15.9$ Hz).



15

- LL-80**: The same synthetic procedure as the preparation of **LL-66**. 62% yield (started with 1,7-bis (3,4-dimethoxyphenyl)-1,6-pentadiene-3, 6-dione). Red powder. ¹H NMR (300 MHz, $CDCl_3$): δ 1.34 (3H, t), 3.95 (12H, s), 4.29 (2H, quart), 5.98 (2H, d, $J=15.6$ Hz), 6.95 (2H, d, $J=8.4$ Hz), 7.00 (1H, d, $J=15.6$ Hz), 7.08 (2H, d, $J=1.8$ Hz), 20 7.22 (2H, dd, $J=8.4$ Hz, $J=1.8$ Hz), 7.77 (2H, d, $J=15.3$ Hz), 7.91 (1H, d, $J=15.6$ Hz).

- HOJ-7**: 3% yield (started with 5 mmol of 2,4-difluorobenzaldehyde), yellow powder, mp 155.5–156 °C (*n*-hexane–EtOAc). ¹H NMR ($CDCl_3$) δ 5.86 (1H, s), 6.68 (2H, d, $J=16.0$ Hz), 6.84–6.96 (4H, m), 7.56 (2H, m), 7.71 (2H, d, $J=16.0$ Hz). ESI- 25 MS m/z 347 [$M-1$]⁺. Anal. calcd for $C_{19}H_{12}F_4O_2$: C, 65.52; H, 3.47. Found: C, 65.66; H, 3.57.

HOJ-9: 10% yield (started with 5 mmol of 2-fluoro-4-methoxybenzaldehyde), yellow powder, mp 163–165 °C (*n*-hexane–EtOAc). ¹H NMR ($CDCl_3$) δ 3.84 (6H, s), 5.82 (1H, s), 6.62 (2H, d, $J=16.0$ Hz), 6.70 (4H, m), 7.48 (2H, t, $J=8.7$ Hz), 7.71 (2H,

d, $J=16.0$ Hz). ESI-MS m/z 371 $[M-1]^+$. Anal. calcd for $C_{21}H_{18}F_2O_4$: C, 67.74; H, 4.87. Found: C, 67.58; H, 4.92.

HOJ-10: 9% yield (started with 5 mmol of 2-fluoro-6-methoxybenzaldehyde), yellow needles, mp 144–145 °C (*n*-hexane–EtOAc). 1H NMR ($CDCl_3$) δ 3.82 (6H, s),
5 5.90 (1H, s), 6.72 (2H, d, $J=16.0$ Hz), 6.88 (2H, m), 7.56 (2H, m), 7.00–7.07 (4H, m),
7.74 (2H, d, $J=16.0$ Hz). ESI-MS m/z 371 $[M-1]^+$. Anal. calcd for $C_{21}H_{18}F_2O_4$: C, 67.74; H, 4.87. Found: C, 67.60; H, 4.95.

Biological activity. The compounds shown above are screened as follows and found to have anticancer and anti-androgen receptor activity as shown in **Table 2**
10 below.

Cell culture and transfection—Human prostate cancer LNCaP and PC-3 cells were maintained in RPMI medium and Dulbecco's minimum essential medium (DMEM), respectively. Both media were strengthened with penicillin (25 units/mL), streptomycin (25 μ g/mL), and 10% fetal calf serum. Androgen receptor
15 transactivation assay, an androgen-dependent reporter gene transcription test, was employed as the primary screening for potential antiandrogen identification. This assay was first performed in LNCaP cells, which express a clinically relevant mutant AR. Once potential compounds were indicated by LNCaP AR transactivation assay, they were re-examined for their potential activity against wild type AR. Wild type
20 AR transactivation assay was performed in PC-3 host cells, which lack an endogenous, functional AR.

Our previously described conditions for cell transfection were followed (Ohtsu *et al*, *J Medicinal Chem* 45: 5037-5042 (2002); Ohtsu *et al.*, *Bioorganic & Medicinal Chemistry* 11:5083-5090 (2003)). In brief, cells were plated in 24- or 48-well tissue
25 culture dishes twenty-four (PC-3 cells) or forty-eight (LNCaP cells) hours prior to transfection. Subsequently, LNCaP cells were transfected with a reporter gene, MMTV-luciferase, which contains MMTV-LTR promoter and androgen receptor binding element, and PRL-SV40, which served as an internal control for transfection efficiency. PC-3 cells were transfected with a wild type AR expression plasmid, pSG5AR, in addition to the above-mentioned MMTV-luciferase reporter gene and
30 PRL-SV40 internal control. SuperFect (Qiagen, Chatsworth, CA) was employed as the transfection reagent following manufacturer's recommendations. At the end of a five-hour transfection, the medium was changed to DMEM or RPMI supplemented

with 10% charcoal dextran-stripped, androgen-depleted serum. Twenty-four hours later, the cells were treated with 1 nM of DHT and/or test compounds at the designated concentration for another twenty-four hours. The cells were harvested for luciferase activity assay using Dual Luciferase Assay System (Promega, Madison, WI). The derived data were expressed as relative luciferase activity normalized to the internal luciferase control. DHT incubation induced a marked expression of reporter gene. Test compounds capable of significantly suppressing this androgen-induced reporter gene expression were identified as potential antiandrogens.

LNCaP cell growth assay: As mentioned above, LNCaP cells contain a substantial amount of mutant AR, and thus these cells' growth can be significantly activated upon androgen incubation. LNCaP cell growth assay thus was used as an alternative to confirm potential antiandrogens identified by the above-mentioned AR transactivation assay. An MTT analysis, relying upon the conversion of a colorless substrate to reduced tetrazolium by the mitochondrial dehydrogenase, was used. The experimental conditions were detailed elsewhere (Ohtsu *et al.*, *Bioorganic & Medicinal Chemistry* 11:5083-5090 (2003)). Briefly, cells were plated in 96-well tissue culture plates and then incubated for 5 consecutive days in the presence of 1 nM of DHT and/or test compounds in 10% charcoal dextran-stripped serum-containing RPMI. At the end of incubation, the cells were given MTT (5 mg/ml in PBS) for three hours at 37°C. The resultant precipitate was dissolved in a lysis buffer and then quantitated at a wavelength of 595 nm using a microplate reader. To ensure the accuracy of data derived from the MTT analysis, cell count was performed using duplicate samples. Test compounds that displayed an adverse effect on the androgen-induced prostate tumor cell growth were identified as potential antiandrogens.

Table 2. The anti-AR activity of curcumin analogues.



Compound	wrAR/PC3	LNCaP
LL-7	0	+
LL-10	0	+
LL-17	+	0
LL-18	+	0
LL-27B	+	0

Compound	wrAR/PC3	LNCaP
LL-32B	+	0
LL-35	+	0
LL-36	+	0
HOJ-7	+	+
HOJ-9	+	0
HOJ-10	+	+
LL-41	+	+
LL-46	+	+
LL-55	0	+
LL-61A	+	+
LL-62	0	+
LL-65/LL-80	++	++
LL-66	0	+

The foregoing is illustrative of the present invention, and is not to be construed as limiting thereof. The invention is defined by the following claims, with equivalents of the claims to be included therein.

What is claimed is:

1. A compound according to formula I or Ia:



5

wherein:

R' and R'' are each independently selected from the group consisting of H, alkoxy, and halo;

10 R_a and R_b are each independently selected from the group consisting of H, alkyl and halo;

R₁ and R₂ are each independently selected from the group consisting of H, hydroxy, alkoxy, nitro, amino, and dialkylamino;

15 R₃ and R₄ are each independently selected from the group consisting of H, hydroxy, halo, alkoxy, and -OR₇C(O)R₈, wherein R₇ is lower alkylene and R₈ is alkoxy;

or R₁ and R₃ together are alkylenedioxy;

or R₂ and R₄ together are alkylenedioxy;

20 R₅ and R₆ are each independently selected from the group consisting of H, halogen, nitro and alkoxy;

X₁ is N, or X₁ is C bonded to a H, alkoxy or nitro; and

X₂ is N, or X₂ is C bonded to a H, alkoxy or nitro;

subject to the proviso that curcumin is excluded therefrom;

or a pharmaceutically acceptable salt thereof.

25

2. The compound according to claim 1, wherein R₁ and R₂ are each independently selected from the group consisting of alkoxy, nitro, amino, and dimethylamino.


3. The compound according to claim 1, wherein R₁ and R₃ together are methylenedioxy or ethylenedioxy.

5 4. The compound according to claim 1; wherein R₂ and R₄ together are methylenedioxy or ethylenedioxy.

5. The compound according to claim 1, wherein R₅ and R₆ are each independently selected from the group consisting of H, F, and nitro.

10

6. A compound according to formula II:



15

wherein:

R₁₁ and R₁₂ are each independently selected from the group consisting of alkoxy, nitro, amino, and dialkylamino;

R₁₃ and R₁₄ are each independently selected from the group consisting of hydroxy, alkoxy, -OR₁₈C(O)R₁₉ wherein R₁₈ is lower alkylene or alkenylene and R₁₉ is alkoxy, and tetrahydropyranyl;

20

or R₁₁ and R₁₃ together are alkylenedioxy;

or R₁₂ and R₁₄ together are alkylenedioxy;

R₁₅ and R₁₆ are each independently selected from the group consisting of H, halogen, and nitro;

25

R₁₇ is -R₂₀C(O)OR₂₁, wherein R₂₀ is alkylene or alkenylene and R₂₁ is H or alkyl;

X₃ is N, or X₃ is C bonded to a H, alkoxy or nitro; and

X_4 is N, or X_4 is C bonded to a H, alkoxy or nitro;
or a pharmaceutically acceptable salt thereof.

7. The compound according to claim 6, wherein R_{11} and R_{12} are each
5 independently selected from the group consisting of alkoxy, nitro, amino, and
dimethylamino.


8. The compound according to claim 6, wherein R_{11} and R_{13} together are
methylenedioxy or ethylenedioxy.

10

9. The compound according to claim 6, wherein R_{12} and R_{14} together are
methylenedioxy or ethylenedioxy.

10. The compound according to claim 6, wherein R_{15} and R_{16} are each
15 independently selected from the group consisting of H, F, and nitro.

11. A compound of Formula IV:




wherein:

20 R_{41} and R_{42} are each independently hydroxy or alkoxy; and
 R_{43} and R_{44} are each independently hydroxy or alkoxy;
or a pharmaceutically acceptable salt thereof.

12. A compound of the Formula:

25



or a pharmaceutically acceptable salt thereof.

13. A pharmaceutical formulation comprising a compound according to claim
5 1, 6, 11 or 12 in a pharmaceutically acceptable carrier.

14. The pharmaceutical formulation according to claim 13, wherein said
carrier is an aqueous carrier.

10 15. A method of treating a cancer, comprising administering to a subject in
need thereof a treatment effective amount of a compound according to claim 1, 6, 11,
12, or curcumin or a pharmaceutically acceptable salt thereof.

15 16. The method according to claim 15, wherein said cancer is selected from
the group consisting of skin cancer, small cell lung cancer, testicular cancer,
lymphoma, leukemia, esophageal cancer, stomach cancer, colon cancer, breast cancer,
endometrial cancer, ovarian cancer, central nervous system cancer, liver cancer and
prostate cancer.

20 17. The method according to claim 16, wherein said cancer is prostate cancer.

18. The method according to claim 16, wherein said cancer is colon cancer.

25 19. The method according to claim 16, wherein said subject is afflicted with
antiandrogen withdrawal syndrome.

20. A method of inducing androgen receptor antagonist activity, said method
comprising contacting a cell with an androgen receptor antagonist effective amount of
a compound according to claim 1, 6, 11, 12 or curcumin or a pharmaceutically
30 acceptable salt thereof.

21. The method according to claim 20, wherein said cell is a cancer cell.

22. The method according to claim 21, wherein said contacting step is carried out *in vivo*.

23. The method according to claim 21, wherein said contacting step is carried
5 out *in vitro*.

24. A method of inducing androgen receptor antagonist activity in a subject afflicted with an androgen-related affliction, said method comprising administering an androgen receptor antagonist effective amount of a compound according to claim 1, 6,
10 11, 12 or curcumin or a pharmaceutically acceptable salt thereof.

25. The method according to claim 24, wherein said subject is afflicted with baldness.

15 26. The method according to claim 24, wherein said subject is afflicted with hirsutism.

27. The method according to claim 24, wherein said subject is afflicted with a behavioral disorder.
20

28. The method according to claim 24, wherein said subject is afflicted with acne.

29. The method according to claim 24, wherein said subject is a male subject
25 and said androgen receptor antagonist effective amount of said inhibits spermatogenesis.