

Method for producing a solid with sufficiently low hygroscopicity which comprises glutamic acid-N,N-diacetic acid (GLDA) or a derivative thereof


- 5 Glutamic acid-N,N-diacetic acid (GLDA) and salts and/or derivatives thereof are of great interest as complexing agents for alkaline earth metal ions and heavy metal ions in a very wide variety of technical fields of application.
- 10 Complexing agents for alkaline earth metal ions and heavy metal ions, as are used for example in detergents and cleaners, are usually synthesized in aqueous solution. For certain application cases, they are required in solid form.
- 15 Customary methods for producing solids from solutions are in particular crystallization and spray-drying methods. It is known that crystalline solid, as is produced for example during evaporative or cooling crystallization methods, can comprise water of crystallization and, under ambient
- 20 conditions, is in most cases less hygroscopic and more storage-stable than amorphous solid. By means of spray-drying methods (e.g. in the spray tower or in the spray fluidized bed), on the other hand, the solid is obtained in amorphous form. In this form, the solid is often highly hygroscopic and,
- 25 upon open storage under ambient conditions, it loses within a short time the ability to be poured. Consequently, measures to increase the storage stability of spray powders are described in the literature, e.g. the compaction or post-treatment of builders for detergents with benzoic acid in US 3,932,316.

30

It was an object of the present invention to provide a solid which comprises glutamic acid-N,N-diacetic acid (GLDA) and/or

salts or derivatives thereof and which preferably has a sufficiently low hygroscopicity for processing and application.


- 5 The present invention relates to a solid which preferably has a sufficiently low hygroscopicity e.g. for processing and application which comprises the following constituents:
- a. a core which comprises glutamic acid-N,N-diacetic acid (GLDA) and/or a salt and/or a derivative thereof (in particular glutamic acid-N,N-diacetic acid (GLDA) and/or a salt thereof), and
 - 10 b. a coating which comprises a compound (or two or more different compounds) of the general formula I



(I)


in which

R is C₁- to C₃₀-alkyl or C₂- to C₃₀-alkenyl, which additionally carry, as substituents, up to 5 hydroxyl groups, formyl groups, C₁- to C₄-alkoxy groups, phenoxy groups or C₁- to C₄-alkoxycarbonyl groups and can be interrupted by up to 5 nonadjacent oxygen atoms, is alkoxylate groups of the formula



in which A¹ and A², independently of one another, are 1,2-alkylene groups having 2 to 4 carbon atoms, Y is hydrogen, C₁- to C₁₂-alkyl, phenyl or C₁- to C₄-alkoxycarbonyl, and k is the number 1, 2 or 3, and m

and n, in each case independently of one another, are numbers from 0 to 50, where the sum of m + n must be at least 4, is phenylalkyl groups having 1 to 20 carbon atoms in the alkyl, a five- or six-membered unsaturated or saturated heterocyclic ring having up to three heteroatoms from the group nitrogen, oxygen and sulfur, which can additionally be benzo-fused, where all of the phenyl rings and heterocyclic rings specified in the meanings for R can also additionally carry, as substituents, up to three C₁- to C₄-alkyl groups, hydroxyl groups, carboxyl groups, sulfo groups and/or C₁- to C₄-alkoxycarbonyl groups, or is a radical for the formula



where A is a C₁- to C₁₂-alkylene bridge or a chemical bond, and

each M, independently of the others, is hydrogen, alkali metal, alkaline earth metal, ammonium or substituted ammonium (e.g. organic amine salts) in the corresponding stoichiometric amounts (in particular each M, independently of the others, is hydrogen, alkali metal, ammonium or substituted ammonium (e.g. organic amine salts) in the corresponding stoichiometric amounts).

The present invention also relates to a method for producing the solid according to the invention, wherein the seed initially introduced is a solid which comprises (or consists of) glutamic acid-N,N-diacetic acid (GLDA) and/or a salt and/or a derivative thereof (in particular glutamic acid-


N,N-diacetic acid (GLDA) and/or a salt thereof), and a spray granulation or a coating (preferably in a fluidized-bed apparatus) with at least one compound of the formula I is carried out.

5

If the core material (which comprises glutamic acid-N,N-diacetic acid (GLDA) and/or a salt and/or a derivative thereof) is used as powder, then this powder, during the spray-granulation, is preferably firstly stuck together by the
 10 coating material (which comprises a compound of the general formula I) to give larger particles (granulation), before said particles are coated by the coating material. If the core material is initially introduced in the form of relatively large granules, then these granules, during the spray-
 15 granulation, are preferably coated directly with the coating material (coating).

Preferably, the glutamic acid-N,N-diacetic acid (GLDA) and/or the salt is a compound of the formula (II)

20



where M^1 is hydrogen, ammonium or an alkali metal (e.g. sodium, potassium) and/or an organic amine (organic amine salt).

25

Particular preference is given to the L form thereof.

Furthermore, preference is given to the sodium salt.

Particular preference is given to compounds of the formula I, as are described in DE 196 49 681.

30

For the method of the present invention, powders and/or granules of the core material are introduced as initial charge

(seeds) in a manner customary per se in a suitable apparatus known per se (preferably a fluidized-bed spray-granulation apparatus, e.g. of the AGT 400 type from Glatt) and fluidized. Here, particle sizes of 30 μm - 300 μm (in particular 50 μm - 200 μm) for the granulation and 200 μm - 1000 μm (in particular 300 μm - 1000 μm) for the coating are preferred.

Preferably, the core material (material comprising GLDA) is dry material, i.e. in particular the core material has a water content of less than 10% (in particular less than 5%).

Preferably, the core material (material comprising GLDA) is produced in a manner customary per se by crystallization, spray-drying or by fluidized-bed granulation.

15

The seeds are held in a fluidized bed in the suspended state (fluidization) and form the surface for a layer-wise drying of atomized droplets which comprise at least one compound of the formula I. The particle produced in this way can be continuously removed from the drying space during the production via a granulation through a classifying discharge in a flexible manner - e.g. with freely adjustable particle sizes - without interrupting the drying operation. In the case of production by means of pure coating, a batch operation is preferred.

25

As regards the spray-granulation method, see e.g.: H. Uhlemann, L. Mörl, "Wirbelschicht - Sprühgranulation [Fluidized-bed - spray-granulation]" Springer - Verlag 2000 (ISBN 3-540-66985-X).

30

The method of the present invention is one wherein a solid which comprises glutamic acid-N,N-diacetic acid (GLDA) and/or

a salt and/or a derivative thereof is initially introduced as seed and then, in a manner customary per se, a spray-granulation/coating with at least one compound of the formula I, which is preferably present in solution (in particular in aqueous solution, preferably ca. 35-43% strength, particularly preferably ca. 35-42% strength, especially preferably ca. 40% strength) is carried out.

Preferably, a spray-granulation/coating with the following parameters is carried out:

- preferred incoming air temperature: 90-160°C,
- preferred product temperature: 50-120°C,
- preferred outgoing air temperature: 40-110°C,
- preferred temperature of the feed: 40-100°C.


In the method according to the invention, for example liquid raw material (e.g. a 35 to 43% strength, preferably a 35-42% strength, in particular a 40% strength, aqueous solution of a compound of the formula I) is sprayed onto the seeds (comprising glutamic acid-N,N-diacetic acid (GLDA) and/or a salt and/or a derivative thereof) fluidizing in the stream of hot air, which dries as a result and the seeds are coated or granulated and then coated.

The fluidized-bed apparatus is preferably a fluidized-bed spray-granulator which is equipped for example with a cyclone and/or a filter and/or a wet scrubber.

Here, a solid is referred to as non-hygroscopic or of being of sufficiently low hygroscopicity when, upon open storage under normal ambient conditions, e.g. 25°C and relative atmospheric humidity of 76%, over a period of at least one day, preferably

one week, it keeps its consistency as (preferably free-flowing) powder or granules.

The core of the solid according to the invention comprises
 5 glutamic acid-N,N-diacetic acid (GLDA) and/or salt and/or a derivative thereof, preferably a compound of the general formula (II):



10

where M^1 is hydrogen, ammonium, an organic amine or an alkali metal (preferably sodium). Particularly preferably, the core comprises L-glutamic acid-N,N-diacetic acid tetrasodium salt ($C_9H_9NO_8Na_4$, e.g. CAS No.: 51981-21-6). Preferably, the core
 15 material comprises Dissolvine® GL from Akzo Nobel.


The core can optionally comprise one or more further complexing agents for alkaline earth metal ions and/or heavy metal ions, such as e.g. polymers such as polyacrylates or
 20 sulfonated polymers (for example sulfonated multipolymer such as Acusol 588 from Room and Haas (Dow)) or ethylenediaminetetraacetic acid (EDTA, for example Dissolvine NA from Akzo Nobel), in particular as tetrasodium salt.

25 The core particularly preferably comprises GLDA granules or co-granules (i.e. granules with auxiliaries which simplify the granulation of GLDA, e.g. polycarboxylates).

Within the context of the present invention, the expression
 30 GLDA preferably refers to glutamic acid-N,N-diacetic acid or one of the salts described herein (e.g. the tetrasodium salt).

The solid produced according to the invention comprises, as coating, preferably essentially compounds of the formula I, where small amounts of starting materials and/or by-products from the preparation of the compounds of the formula I may additionally be present. Depending on the synthesis method used, customary purities for the compounds I are 70 to 99.9% by weight, in particular 80 to 99.5% by weight, in each case based on the solids content.

- 10 For the present invention, those compounds of the formula I in which R is C₁- to C₂₀-alkyl, C₂- to C₂₀-alkenyl or a radical of the formula



- 15 are preferably suitable.

As compound of the formula I, particular preference is given to using α-alanine-N,N-diacetic acid (R = CH₃, MGDA) and its salts. Preference is given to using e.g. its alkali metal salts, ammonium salts and substituted ammonium salts.

Of suitability as such salts are primarily the sodium, potassium and ammonium salts, in particular the trisodium, tripotassium and triammonium salt, and also organic triamine salts with a tertiary nitrogen atom.

Suitable bases underlying organic amine salts are in particular tertiary amines, such as trialkylamines having 1 to 4 carbon atoms in each alkyl, such as trimethyl- and triethylamine, and trialkanolamines having 2 or 3 carbon atoms

in the alkanol radical, preferably triethanolamine, tri-n-propanolamine or triisopropanolamine.

The alkaline earth metal salts used are in particular the
5 calcium and magnesium salts.

Besides methyl, of suitability for the radical R as straight-chain or branched alk(en)yl radicals, are in particular C₂- to C₁₇-alkyl and -alkenyl, here in particular straight-chain
10 radicals derived from saturated or unsaturated fatty acids.

Examples of individual radicals R are: ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, tert-pentyl, neopentyl, n-hexyl, n-heptyl, 3-heptyl
15 (derived from 2-ethylhexanoic acid), n-octyl, isooctyl (derived from isononanoic acid), n-nonyl, n-decyl, n-undecyl, n-dodecyl, isododecyl (derived from isotridecanoic acid), n-tridecyl, n-tetradecyl, n-pentadecyl, n-hexadecyl, n-heptadecyl, n-octadecyl, n-nonadecyl, n-eicosyl and
20 n-heptadecenyl (derived from oleic acid). For R, it is also possible for mixtures to arise, in particular those which are derived from naturally occurring fatty acids and from synthetically produced technical-grade acids, for example by
25 oxo synthesis.

25

The C₁- to C₁₂-alkylene bridges A are in particular polymethylene groups of the formula (CH₂)_k, in which k is a number from 2 to 12, in particular from 2 to 8, i.e. 1,2-ethylene, 1,3-propylene, 1,4-butylene, pentamethylene,
30 hexamethylene, heptamethylene, octamethylene, nonamethylene, decamethylene, undecamethylene and dodecamethylene. Hexamethylene, octamethylene, 1,2-ethylene and 1,4-butylene

are particularly preferred here. In addition, however, branched C₃- to C₁₂-alkylene groups can also occur, e.g. -CH₂CH(CH₃)CH₂-, -CH₂C(CH₃)₂CH₂-, -CH₂CH(C₂H₅)- or CH₂CH(CH₃)-.

- 5 The C₁- to C₃₀-alkyl and C₂- to C₃₀-alkenyl groups can carry up to 5, in particular up to 3, additional substituents of the specified type and be interrupted by up to 5, in particular up to 3, nonadjacent oxygen atoms. Examples of such substituted alk(en)yl groups are -CH₂OH, -CH₂CH₂OH, -CH₂CH₂-O-CH₃,
 10 -CH₂CH₂-O-CH₂CH₂-O-CH₃, -CH₂-O-CH₂CH₃, -CH₂-O-CH₂CH₂-OH, -CH₂-CHO, -CH₂-OPh, -CH₂-COOCH₃ or -CH₂CH₂-COOCH₃.

Suitable alkoxy(ate) groups are in particular those in which m and n, in each case independently of one another, are
 15 numbers from 0 to 30, in particular from 0 to 15. A¹O and A²O are preferably groups derived from butylene oxide and in particular from propylene oxide and from ethylene oxide. Of particular interest are pure ethoxylates and pure propoxylates, but ethylene oxide-propylene oxide block
 20 structures can also arise.

Suitable five- or six-membered unsaturated or saturated heterocyclic rings having up to three heteroatoms from the group nitrogen, oxygen and sulfur, which can be additionally
 25 benzo-fused and substituted by the designated radicals, are e.g.:

tetrahydrofuran, furan, tetrahydrothiophene, thiophene, 2,5-dimethylthiophene, pyrrolidine, pyrroline, pyrrole, isoxazole, oxazole, thiazole, pyrazole, imidazoline,
 30 imidazole, 1,2,3-triazolidine, 1,2,3- and 1,2,4-triazole, 1,2,3-, 1,2,4- and 1,2,5-oxadiazole, tetrahydropyran, dihydropyran, 2H- and 4H-pyran, piperidine, 1,3- and

1,4-dioxane, morpholine, pyrazan, pyridine, α -, β - and γ -picoline, α - and γ -piperidone, pyrimidine, pyridazine, pyrazine, 1,2,5-oxathiazine, 1,3,5-, 1,2,3- and 1,2,4-triazine, benzofuran, thionaphthene, indoline, indole,
 5 isoindoline, benzoxazole, indazole, benzimidazole, chromane, isochromane, 2H- and 4H-chromene, quinoline, isoquinoline, 1,2,3,4-tetrahydroisoquinoline, cinnoline, quinazoline, quinoxaline, phthalazine and benzo-1,2,3-triazine.

10 N-H groups in the specified heterocyclic rings should be present as far as possible in derivatized form, for example as N-alkyl group (e.g. having 1 to 12 carbon atoms).

15 In the case of substitution on the phenyl rings or the heterocyclic rings, preferably two (identical or different) or in particular one individual substituent, arise.

Examples of optionally substituted phenylalkyl groups and alkyl groups carrying heterocyclic rings for R are benzyl,
 20 2-phenylethyl, 3-phenylpropyl, 4-phenylbutyl, o-, m- or p-hydroxybenzyl, o-, m- or p-carboxybenzyl, o-, m- or p-sulfobenzyl, o-, m- or p-methoxy or -ethoxycarbonylbenzyl, 2-furylmethyl, N-methylpiperidin-4-ylmethyl or 2-, 3- or 4-pyridinylmethyl.

25

For the substitution on phenyl rings and also on heterocyclic rings, preference is given to using water-solubilizing groups such as hydroxyl groups, carboxyl groups or sulfo groups.

30 Examples of the specified C₁- to C₄-, C₁- to C₁₂- and C₁- to C₂₀-alkyl groups are also to be understood as meaning the corresponding radicals listed above for R.

The solid according to the invention is suitable to a particular degree as a component for solid detergent and cleaner formulations. Subsequently, solid detergent and cleaner formulations which comprise the solid according to the invention as complexing agent for alkaline earth metal ions and heavy metal ions in the amounts customary for this purpose, as well as other customary constituents of such formulations, are also provided by the present invention. Compositions and customary constituents of such solid detergent and cleaner formulations are known to the person skilled in the art and therefore do not need to be explained in more detail here.

The example below is intended to illustrate the invention in more detail. The glycine-N,N-diacetic acid derivative I used was α -alanine-N,N-diacetic acid (methylglycine-N,N-diacetic acid, "MGDA") trisodium salt.

Examples

20

The following processes were carried out on a pilot fluidized-bed spray-granulation apparatus model AGT 400 from Glatt which was equipped with a cyclone, a filter and a wet scrubber.

25 Example 1

It was an aim of this experiment to coat granules of GLDA with Trilon M liquid (BASF). The target grain size was 200-1000 μm .
Experimental procedure and results

30

Trilon M liquid was heated to 90-95°C and sprayed onto the initial charge of GLDA granules. This gave granules in the

desired grain spectrum. The oversize fraction was sieved off over a 1000 μm sieve.

At the following processing parameters, the coating was able
5 to run in a stable manner:

incoming air temperature 140°C

incoming air amount: 1400 m³/h

product temperature 100-104°C

10 outgoing air temperature: 98-101°C

spraying pressure: 2.0 bar

spraying rate: ca. 350-450 g/min

storage container: 90°C

15 After establishing the processing parameter specified above, it was possible to achieve a stable process.

By means of subsequent protective sieving over 1000 μm ,
15.9 kg of granules were separated. The oversize fraction
20 amounted to 6.0 kg.

Example 2

The aim of this experiment was to coat co-granules of GLDA and
25 a polymer (homopolymer Sokalan PA 30 CL (BASF)) with Trilon M Liquid. Target grain size of the coated granules: 200-1000 μm .

10 kg of Trilon M liquid were heated to 90-95°C and sprayed on
to the initial charge of co-granules (15 kg). This gave
30 granules in the desired grain spectrum. The oversize fraction was sieved over a 1000 μm sieve.

At the following processing parameters, the coating is able to run in a stable manner:

incoming air temperature 130-140°C
5 incoming air amount: 1200 m³/h
product temperature 104-106°C
outgoing air temperature: 100-102°C
spraying pressure: 2.0 bar
spraying rate: ca. 100-150 g/min
10 storage container: 95°C

The following example amounts were prepared:

17.0 kg of granules (sieved over 1000 µm). The oversize
15 fraction was 5.1 kg.

After establishing the processing parameters given above, it was possible to achieve a stable process.

20 Example 3

The aim of this experiment was to coat co-granules of GLDA and a polymer (homopolymer Sokalan PA 30 CL (BASF)) with Trilon M Liquid. Target grain size of the coated granules: 200-1000 µm.

25 20 kg of Trilon M liquid were heated to 90-95°C and sprayed on to the initial charge of co-granules (15 kg). This gave granules in the desired grain spectrum. The oversize fraction was sieved over a 1000 µm sieve.

30

At the following processing parameters, the coating is able to run in a stable manner:

incoming air temperature 130-140°C

incoming air amount: 1200 m³/h

product temperature 104-106°C

5 outgoing air temperature: 100-102°C

spraying pressure: 2.0 bar

spraying rate: ca. 100-150 g/min

storage container: 95°C


10 The following example amounts were prepared:

20.3 kg of granules (sieved over 1000 µm). The oversize fraction was 5.2 kg.

15 After establishing the processing parameters given above, it was possible to achieve a stable process.

Claims


1. A solid which comprises the following constituents:
- 5 a. a core which comprises glutamic acid-N,N-diacetic acid and/or a salt and/or a derivative thereof, and
- b. a coating which comprises a compound of the general formula I



(I)


in which

R is C₁- to C₃₀-alkyl or C₂- to C₃₀-alkenyl, which additionally carries, as substituents, up to 5 hydroxyl groups, formyl groups, C₁- to C₄-alkoxy groups, phenoxy groups or C₁- to C₄-alkoxycarbonyl groups and can be interrupted by up to 5 nonadjacent oxygen atoms, is alkoxylate groups of the formula



in which A¹ and A², independently of one another, are 1,2-alkylene groups having 2 to 4 carbon atoms, Y is hydrogen, C₁- to C₁₂-alkyl, phenyl or C₁- to C₄-alkoxycarbonyl, and k is the number 1, 2 or 3, and m and n, in each case independently of one another, are numbers from 0 to 50, where the sum of m + n must be at least 4, is phenylalkyl groups having 1 to 20 carbon atoms in the alkyl, a five- or six-membered unsaturated or


saturated heterocyclic ring having up to three heteroatoms from the group nitrogen, oxygen and sulfur, which can additionally be benzo-fused, where all of the phenyl rings and heterocyclic rings specified in the meanings for R can also additionally carry, as substituents, up to three C₁- to C₄-alkyl groups, hydroxyl groups, carboxyl groups, sulfo groups and/or C₁- to C₄-alkoxycarbonyl groups, or is a radical for the formula



where A is a C₁- to C₁₂-alkylene bridge or a chemical bond, and


each M, independently of the others, is hydrogen, alkali metal, alkaline earth metal, ammonium or substituted ammonium (e.g. organic amine salts) in the corresponding stoichiometric amounts.

2. The solid according to claim 1, where R is C₁- to C₂₀-alkyl, C₂- to C₂₀-alkenyl or a radical of the formula



3. The solid according to claim 1 or 2, where the coating consists essentially of α-alanine-N,N-diacetic acid (MGDA) or its alkali metal, ammonium or substituted amine salts.

4. The solid according to any one of claims 1 to 3, where the glutamic acid-N,N-diacetic acid (GLDA) and/or the salt or the derivative is a compound of the formula (II)



where M^1 is hydrogen, ammonium and/or an alkali metal or an organic amine.

- 10 5. The method for producing a solid according to any one of claims 1 to 4, wherein the seed initially introduced is a solid which comprises glutamic acid-N,N-diacetic acid (GLDA) and/or a derivative and/or a salt thereof, and a spray-granulation with at least one compound of the
15 formula I is carried out.
6. The method according to claim 5, wherein an aqueous solution of a compound of the formula I is used for the spray-granulation.
- 20 7. A solid detergent or cleaner formulation comprising a solid according to any one of claims 1 to 4 as complexing agent for alkaline earth metal ions and heavy metal ions.