

DETECTION OF MISFOLDED TAU PROTEIN**CROSS-REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims priority from U.S. Provisional Patent Application No. 62/507,166, filed on May 16, 2017, the entire contents of which are incorporated herein by reference.

BACKGROUND

[0002] Tauopathies may include, for example, Alzheimer's disease (AD), Parkinson's Disease (PD), Progressive Supranuclear Palsy (PSP), Frontotemporal Dementia (FTD), Corticobasal degeneration (CBD), Mild cognitive impairment (MCI), Argyrophilic grain disease (AgD) Traumatic Brain Injury (TBI), Chronic Traumatic Encephalopathy (CTE), and Dementia Pugilistica (DP), and the like. Misfolded tau aggregates and fibrils may be formed and accumulate via nucleation and growth. The misfolded tau aggregates may induce cellular dysfunction and tissue damage, among other effects.

[0003] Real time quaking-induced conversion (RT-QuIC) has been shown to cause replication of 3-repeat (3R) tau isoforms from brain homogenate and cerebrospinal fluid samples drawn from Pick disease subjects, allowing sensitive detection of this rare disease and discrimination from other tauopathies. Surprisingly, however, for more common tauopathies of clinical importance that include misfolding of 4R tau, the efficacy of RT-QuIC was reduced by 3 to 5 orders of magnitude, rendering it ineffective and impractical for clinical and laboratory use. Such adverse results were obtained by seeding with brain samples containing predominant 4-repeat (4R) tau aggregates from cases of CBD, AgD, and FTDP-17, and PSP, as well as AD, a 4R + 3R tauopathy. Some AD and PSP samples gave signals above the detection limit, but the signals were outliers and much weaker compared to Pick disease brain samples. Additionally, the AD and PSP samples which generated weak responses were not analyzed for contamination. The RT-QuIC analyses of 4R or 4R + 3R tauopathies in general do not appear to be significantly different from controls using diseased subjects with no immunohistologically detected tau pathology. Such controls included diagnoses of senile change (SC), cerebrovascular disease (CVD), diffuse Lewy body disease (DLBD), frontotemporal dementia with TDP-43 (FTD-TDP), and amyotrophic lateral sclerosis (ALS). In sum, RT-QuIC analyses were shown to be generally ineffective and impractical for 4R tauopathies including 4R predominant and 4R + 3R mixed tauopathies.

[0004] The present application appreciates that detection of misfolded tau protein, e.g., for diagnosis of related diseases, may be a challenging endeavor.

SUMMARY

[0005] In one embodiment, a method is provided for determining a presence or absence in a sample of a first misfolded protein aggregate. The method may include performing a first protein misfolding cyclic amplification (PMCA) procedure. The first PMCA procedure may include forming a first incubation mixture by contacting a first portion of the sample with a first substrate protein. The first substrate protein may include 4R tau protein. The first PMCA procedure may include conducting an incubation cycle two or more times under conditions effective to form as first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the first incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by analyzing the first incubation mixture for the presence or absence of the first amplified, misfolded protein aggregate. The first misfolded protein aggregate may include the first substrate protein. The first amplified, misfolded protein aggregate may include the first substrate protein.

[0006] In another embodiment, a method is provided for determining a presence or absence in a subject of a tauopathy corresponding to a first misfolded protein aggregate. The method may include providing a sample from the subject. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting a first portion of the sample with a first substrate protein. The first substrate protein may include a tau isoform. The first substrate protein may be subject to pathological misfolding and/or aggregation in vivo to form the first misfolded protein aggregate. The first PMCA procedure may include conducting an incubation cycle two or more times under conditions effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the first incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by analyzing the first incubation mixture for the presence or absence of the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence of the tauopathy in the subject according the presence or absence of the first misfolded protein aggregate in the sample. The first misfolded protein aggregate may include the first substrate protein. The first amplified, misfolded

protein aggregate may include the first substrate protein. The method may provide that the tauopathy excludes Pick's disease when the first substrate protein consists of monomeric 3R tau.

[0007] In one embodiment, a method is provided using capturing for determining a presence or absence in a sample of a first misfolded protein aggregate. The method may include capturing the first misfolded protein aggregate from the sample to form a captured first misfolded protein aggregate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the captured first misfolded protein aggregate with a molar excess of a first substrate protein. The first substrate protein may be subject to pathological misfolding and/or aggregation in vivo to form the first misfolded protein aggregate. The molar excess may be greater than an amount of protein monomer included in the captured first misfolded protein aggregate. The method may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the captured first misfolded protein aggregate. Each incubation cycle may include disrupting the first incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence of the first misfolded protein aggregate in the sample by detecting the first amplified, misfolded protein aggregate. The first misfolded protein aggregate may include the first substrate protein. The first amplified, misfolded protein aggregate may include the first substrate protein.

[0008] In another embodiment, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including Alzheimer's disease (AD). The method may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of

the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of AD according to the presence or absence of the first misfolded protein aggregate in the sample.

[0009] In one embodiment, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including Parkinson's disease (PD). The method may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of PD according to the presence or absence of the first misfolded protein aggregate in the sample.

[0010] In another embodiment, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including Progressive Supranuclear Palsy (PSP). The method may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of

the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of PSP according to the presence or absence of the first misfolded protein aggregate in the sample.

[0011] In one embodiment, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including FrontoTemporal Dementia (FTD). The method may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of FTD according to the presence or absence of the first misfolded protein aggregate in the sample.

[0012] In another embodiment, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including Corticobasal degeneration (CBD). The method may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of

the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of CBD according to the presence or absence of the first misfolded protein aggregate in the sample.

[0013] In one embodiment, a kit is provided for determining a presence or absence in a sample of a first misfolded protein aggregate. The kit may include a first substrate protein that may include 4R tau. The kit may include an indicator of the first misfolded protein aggregate. The first misfolded protein aggregate may include the first substrate protein. The first misfolded protein aggregate may correspond to a tauopathy. The kit may include a buffer. The kit may include heparin. The kit may include a salt. The kit may include instructions. The instructions may direct a user to obtain the sample. The instructions may direct the user to perform at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting a first portion of the sample with the first substrate protein, the indicator of the first misfolded protein aggregate, the buffer, the heparin, and the salt. The first incubation mixture may be formed with a concentration of one or more of: the first substrate protein of less than about 20 μ M; the heparin of less than about 75 μ M; the salt as NaCl of less than about 190 mM; and the indicator of the first misfolded protein aggregate as Thioflavin T of less than about 9.5 μ M. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The instructions may direct the user to determine the presence or absence in the sample of the first misfolded protein aggregate by analyzing the first incubation mixture for the presence or absence of the first amplified, misfolded protein aggregate according to the indicator of the first misfolded protein aggregate.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014] The accompanying figures, which are incorporated in and constitute a part of the specification, illustrate example methods and results, and are used merely to illustrate example embodiments.

[0015] **FIG. 1A** shows electron micrographs taken at 0h, 5h, 10h, and 24h of incubation.

[0016] **FIG. 1B** is a western blot of soluble oligomeric A β protein aggregates.

[0017] **FIG. 2A** is a graph showing non-amplified amyloid formation measured by ThT fluorescence as a function of time seeded by various concentrations of synthetic soluble oligomeric A β protein of **EXAMPLE 1**.

[0018] **FIG. 2B** is a graph showing amplification cycle-accelerated amyloid formation measured by ThT fluorescence as a function of time seeded by various concentrations of synthetic soluble oligomeric A β protein of **EXAMPLE 1**.

[0019] **FIG. 3A** is a graph of amyloid formation versus time, measured as a function of ThT fluorescence labeling, showing the average kinetics of A β aggregation seeded by CSF from 5 representative samples from the AD, NND, and NAND groups.

[0020] **FIG. 3B** is a graph of the lag phase time in h for A β aggregation in the presence of samples from the AD, NND, and NAND groups.

[0021] **FIG. 3C** is a graph showing the extent of amyloid formation obtained after 180 A β -PMCA cycles, e.g. 90 h of incubation (P90) in the presence of CSF samples from AD, NND and NAND patients.

[0022] **FIGS. 4A-D** are plots of the true positive rate (sensitivity) as a function of the false positive rate (specificity) for different cut-off points using the lag phase values showed in **FIG. 3B** for AD vs NAND (**FIG. 4A**), AD vs NND (**FIG. 4B**) and AD vs All control samples (**FIG. 4C**). **FIG. 4D** estimates the most reliable cut-off point for the different set of group comparisons.

[0023] **FIG. 5, Table 1** shows estimations of the sensitivity, specificity and predictive value of the A β -PMCA test, calculated using the lag phase numbers.

[0024] **FIG. 6** is a graph of the lag phase time in h for samples obtained after 300 A β -PMCA cycles, e.g. 150 h of incubation (P90) in the presence of CSF samples from AD and control patients.

[0025] **FIG. 7A** is a western blot showing results of immunodepletion using synthetically prepared A β oligomers spiked into human CSF.

[0026] **FIG. 7B** is a graph showing the kinetics of A β aggregation seeded by control and immunodepleted CSF samples.

[0027] **FIG. 7C** is a graph showing the kinetics of A β aggregation seeded by control and immunodepleted CSF samples, depleted only with the A11 conformational antibody.

[0028] **FIG. 8A** is a schematic representation of an ELISA solid phase method employed to capture A β oligomers from complex biological samples.

[0029] **FIG. 8B** is a schematic representation of a magnetic bead solid phase method employed to capture A β oligomers from complex biological samples.

[0030] **FIG. 9, Table 2** shows the ability of specific antibodies to capture the A β oligomers.

[0031] **FIG. 10** is a graph of amyloid formation versus time showing the acceleration of A β aggregation by the presence of different quantities of synthetic oligomers spiked in human plasma.

[0032] **FIG. 11** is a graph showing time to reach 50% aggregation in an A β -PMCA assay in the presence of plasma samples from AD patients and controls.

[0033] **FIG. 12** is a western blot showing the results of amplification of A β aggregation by cycles of incubation/sonication in the presence of distinct quantities of synthetic A β oligomers monitored by Western blot after protease digestion.

[0034] **FIG. 13A** is a graph of Thioflavin T fluorescence versus time showing the detection of α S seeds by PD-PMCA.

[0035] **FIG. 13B** is a graph of time to reach 50% aggregation plotted as a function of the indicated amounts α S seeds.

[0036] **FIG. 14** shows detection of α S seeds in CSF samples from human PD patients by PD-PMCA, versus controls with Alzheimer's disease (AD) or a non-neurodegenerative disease (NND).

[0037] **FIG. 15, Table 3** demonstrates the ability of different sequence or conformational antibodies to capture α S oligomers.

[0038] **FIG. 16A** is a schematic representation of an ELISA solid phase method employed to capture α S oligomers.

[0039] **FIG. 16B** is a schematic representation of a magnetic bead solid phase method employed to capture α S oligomers.

[0040] **FIGS. 17A, 17B, and 17C** are a series of graphs that show the results of immunoprecipitation/aggregation of α -Synuclein oligomers from human blood plasma using three different α -Synuclein antibodies. **FIG. 17A** shows results with antibody N-19. **FIG. 17B** shows results with antibody 211. **FIG. 17C** shows results with antibody C-20.

[0041] **FIGS. 18A, 18B, and 18C** are a series of graphs that show the results of detection for α S seeds in CSF samples. **FIG. 18A** shows results in control samples. **FIG. 18B** shows results in PD patients. **FIG. 18C** shows results in patients with Multiple System Atrophy (MSA).

[0042] **FIG. 19** is a flow chart showing the preparation and purification of recombinant full-length 4R tau protein.

[0043] **FIG. 20A** is a graph of aggregation in % according to ThT fluorescence for various initial amounts of tau seeds and a control. The values in **FIG. 20A** are the mean of two replicates, with the error bars indicating standard deviation.

[0044] **FIG. 20B** is a graph of T_{50} , the time to 50% aggregation as measured by ThT fluorescence versus the log of the amount of oligomeric tau seeds in fmol.

[0045] **FIG. 20C** is a graph of aggregation followed over time by ThT fluorescence.

[0046] **FIG. 20D** is a graph of the relationship between the quantity of tau oligomers and the Tau-PMCA signal (time to reach 50% aggregation).

[0047] **FIGS. 20E-20L** are a series of graphs that display the aggregation results based on ThT fluorescence of 8 of the conditions tested, including 4 different time points (0, 7, 14 and 30 days) with samples subjected to freezing and thawing or not and in the presence of buffer or CSF.

[0048] **FIG. 20E** is a graph of aggregation based on ThT fluorescence of a first seed preparation at 0 days.

[0049] **FIG. 20F** is a graph of aggregation based on ThT fluorescence of a first seed preparation at 7 days.

[0050] **FIG. 20G** is a graph of aggregation based on ThT fluorescence of a first seed preparation at 14 days.

[0051] **FIG. 20H** is a graph of aggregation based on ThT fluorescence of a first seed preparation at 30 days.

[0052] **FIG. 20I** is a graph of aggregation based on ThT fluorescence of a second seed preparation at 0 days.

[0053] **FIG. 20J** is a graph of aggregation based on ThT fluorescence of a second seed preparation at 7 days.

[0054] **FIG. 20K** is a graph of aggregation based on ThT fluorescence of a second seed preparation at 14 days.

[0055] **FIG. 20L** is a graph of aggregation based on ThT fluorescence of a second seed preparation at 30 days.

[0056] **FIG. 20M** is a table of T_{50} values showing reproducibility across 16 different conditions.

[0057] **FIG. 20N** is a graph of ThT fluorescence vs time for the tau assay seeded with 1 pm of tau, A β 40, AB42, His α Syn, Hu α Syn, and a control with no seeds.

[0058] **FIG. 21A** is a graph showing ThT fluorescence at 447h of incubation for patients with AD, patients with MCI, patients with other tauopathies, positive controls using samples of healthy CSF spiked with synthetic Tau oligomers (12.5 fmol), negative controls of samples of healthy CSF without Tau seeds; and control patients with other neurological diseases.

[0059] **FIG. 21B** shows fluorescence signals for samples from patients with AD or other tauopathies for tau-PMCA comparable to that observed in samples containing recombinant tau oligomers.

[0060] **FIG. 22** is a graph showing aggregation % based on ThT versus time for patients affected by AD, FTD (frontotemporal dementia), CBD (corticobasal degeneration), and PSP (progressive supranuclear palsy), versus representative CSF samples from a control.

DETAILED DESCRIPTION

[0061] Methods and kits are provided for the detection or characterization of misfolded tau protein in a sample, including for the determination or diagnosis of tauopathies in a subject from which the sample is taken. Misfolded aggregates of tau proteins may be formed and accumulate. The misfolded aggregates may induce cellular dysfunction and tissue damage among other effects. For example, tauopathies may include those that predominantly regard misfolding of 4R, or misfolding of mixtures of 4R and 3R: Alzheimer's disease (AD), Parkinson's Disease (PD), Progressive Supranuclear Palsy (PSP), FrontoTemporal Dementia (FTD), Corticobasal degeneration (CBD), Mild cognitive impairment (MCI), Argyrophilic grain disease (AgD) Traumatic Brain Injury (TBI), Chronic Traumatic Encephalopathy (CTE), Dementia Pugilistica (DP), and the like.

[0062] In some embodiments, tauopathies herein may exclude Pick's disease. In some embodiments, the tauopathies described herein may exclude those that predominantly regard 3R tau misfolding, e.g., Pick's disease.

[0063] The methods may include protein misfolding cyclic amplification (PMCA), which may provide ultra-sensitive detection of misfolded protein aggregates such as tau through artificial acceleration and amplification of the misfolding and aggregation process *in vitro*. The basic concept of PMCA has been previously demonstrated experimentally for prions (Soto et al, WO 2002/04954; Estrada, et al., U.S. Pat. App. Pub. No. 20080118938, each of which is entirely incorporated herein by reference) and for other protein misfolding, such as of "A β " or "beta amyloid" in Alzheimer's disease and alpha synuclein in Parkinson's disease (Soto et al, WO 2016/040907, which is entirely incorporated herein by reference). However, prior to the filing date of the present document, no reference has described PCMA for the amplification and detection of misfolded tau protein corresponding to any tauopathy that predominantly regards misfolding of 4R, or that regards misfolding of 3R tau in the presence of 4R tau, or for any tauopathy other than Pick's disease. This document discloses specific examples and details which enable PMCA technology for the detecting the presence or absence of misfolded tau aggregates, and, in various embodiments, one or more additional PMCA procedures for the detection of other misfolded

proteins such as misfolded A β in Alzheimer's disease and alpha synuclein in Parkinson's disease. Such one or more additional PMCA procedures may provide discrimination among the various tauopathies, for example, to distinguish AD and PD from each other and from PSP, FTD, CBD, MCI, AgD, TBI, CTE, DP, and the like.

[0064] As used herein, "A β " or "beta amyloid" refers to a peptide formed via sequential cleavage of the amyloid precursor protein (APP). Various A β isoforms may include 38-43 amino acid residues. The A β protein may be formed when APP is processed by β - and/or γ -secretases in any combination. The A β may be a constituent of amyloid plaques in brains of individuals suffering from or suspected of having AD. Various A β isoforms may include and are not limited to Abeta40 and Abeta42. Various A β peptides may be associated with neuronal damage associated with AD.

[0065] As used herein, " α S" or "alpha-synuclein" refers to full-length, 140 amino acid α -synuclein protein, e.g., " α S-140." Other isoforms or fragments may include " α S-126," alpha-synuclein-126, which lacks residues 41-54, e.g., due to loss of exon 3; and " α S-112" alpha-synuclein-112, which lacks residue 103-130, e.g., due to loss of exon 5. The α S may be present in brains of individuals suffering from PD or suspected of having PD. Various α S isoforms may include and are not limited to α S-140, α S-126, and α S-112. Various α S peptides may be associated with neuronal damage associated with PD.

[0066] As used herein, "tau" refers to proteins are the product of alternative splicing from a single gene, e.g., MAPT (microtubule-associated protein tau) in humans. Tau proteins include up to full-length and truncated forms of any of tau's isoforms. Various isoforms include, but are not limited to, the six tau isoforms known to exist in human brain tissue, which correspond to alternative splicing in exons 2, 3, and 10 of the tau gene. Three isoforms have three binding domains and the other three have four binding domains. Misfolded tau may be present in brains of individuals suffering from AD or suspected of having AD, or other tauopathies that, like AD, regard misfolding in the presence of both 4R and 3R tau isoforms. Misfolded tau may also be present in diseases that regard misfolding of primarily 4R tau isoforms, such as progressive supranuclear palsy (PSP), tau-dependent frontotemporal dementia (FTD), corticobasal degeneration (CBD), mild cognitive impairment (MCI), argyrophilic grain disease (AgD), and the like.

[0067] As used herein, a "misfolded protein aggregate" is a protein that contains in part or in full a structural conformation of the protein that differs from the structural conformation that exists when involved in its typical, non-pathogenic normal function within a biological system. A misfolded protein may aggregate. A misfolded protein may localize in a protein aggregate. A

misfolded protein may be a non-functional protein. A misfolded protein may be a pathogenic conformer of the protein. Monomeric protein compositions may be provided in native, nonpathogenic conformations without the catalytic activity for misfolding, oligomerization, and aggregation associated with seeds (a misfolded protein oligomer capable of catalyzing misfolding under PMCA conditions). Monomeric protein compositions may be provided in seed-free form.

[0068] As used herein, “monomeric protein” refers to single protein molecules. “Soluble, aggregated misfolded protein” refers to oligomers or aggregations of monomeric protein that remain in solution. Examples of soluble, misfolded protein may include any number of protein monomers so long as the misfolded protein remains soluble. For example, soluble, misfolded protein may include monomers or aggregates of between 2 and about 50 units of monomeric protein.

[0069] Monomeric and/or soluble, misfolded protein may aggregate to form insoluble aggregates, higher oligomers, and/or tau fibrils. For example, aggregation of A β or tau protein may lead to protofibrils, fibrils, and eventually misfolded plaques or tangles that may be observed in AD or tauopathy subjects. “Seeds” or “nuclei” refer to misfolded protein or short fragmented fibrils, particularly soluble, misfolded protein with catalytic activity for further misfolding, oligomerization, and aggregation. Such nucleation-dependent aggregation may be characterized by a slow lag phase wherein aggregate nuclei may form, which may then catalyze rapid formation of further aggregates and larger oligomers and polymers. The lag phase may be minimized or removed by addition of pre-formed nuclei or seeds. Monomeric protein compositions may be provided without the catalytic activity for misfolding and aggregation associated with misfolded seeds. Monomeric protein compositions may be provided in seed-free form.

[0070] As used herein, “soluble” species may form a solution in biological fluids under physiological conditions, whereas “insoluble” species may be present as precipitates, fibrils, deposits, tangles, or other non-dissolved forms in such biological fluids under physiological conditions. Such biological fluids may include, for example, fluids, or fluids expressed from one or more of: amniotic fluid; bile; blood; cerebrospinal fluid; cerumen; skin; exudate; feces; gastric fluid; lymph; milk; mucus, e.g. nasal secretions; mucosal membrane, e.g., nasal mucosal membrane; peritoneal fluid; plasma; pleural fluid; pus; saliva; sebum; semen; sweat; synovial fluid; tears; urine; and the like. Insoluble species may include, for example, fibrils of A β , α S, 4R tau, 3R tau, combinations thereof such as 3R tau + 4R tau, and the like. A species that dissolves in a non-biological fluid but not one of the aforementioned biological fluids under physiological conditions may be considered insoluble. For example, fibrils of A β , α S, 4R tau, 3R tau, combinations thereof such as 3R tau + 4R tau, and the like may be dissolved in a solution of, e.g.,

a surfactant such as sodium dodecyl sulfate (SDS) in water, but may still be insoluble in one or more of the mentioned biological fluids under physiological conditions.

[0071] In some embodiments, the sample may exclude insoluble species of the misfolded proteins such as A β , α S, 4R tau, 3R tau, combinations thereof such as 3R tau + 4R tau and the like as a precipitate, fibril, deposit, tangle, plaque, or other form that may be insoluble in one or more of the described biological fluids under physiological conditions.

[0072] For example, in some embodiments, the sample may exclude tau in fibril form. The sample may exclude misfolded tau proteins in insoluble form, e.g., the sample may exclude the misfolded tau proteins as precipitates, fibrils, deposits, tangles, plaques, or other insoluble forms, e.g., in fibril form. The methods described herein may include preparing the sample by excluding the misfolded protein in insoluble form, e.g., by excluding from the sample the misfolded tau protein as precipitates, fibrils, deposits, tangles, plaques, or other insoluble forms, e.g., in fibril form. The kits described herein may include instructions directing a user to prepare the sample by excluding from the sample the misfolded tau protein as precipitates, fibrils, deposits, tangles, plaques, or other insoluble forms, e.g., in fibril form. The exclusion of such insoluble forms of the described misfolded proteins from the sample may be substantial or complete.

[0073] As used herein, aggregates of misfolded protein refer to non-covalent associations of protein including soluble, misfolded protein. Aggregates of misfolded protein may be “de-aggregated”, or disrupted to break up or release soluble, misfolded protein. The catalytic activity of a collection of soluble, misfolded protein seeds may scale, at least in part with the number of such seeds in a mixture. Accordingly, disruption of aggregates of misfolded protein in a mixture to release misfolded protein seeds may lead to an increase in catalytic activity for oligomerization or aggregation of monomeric protein.

[0074] In various embodiments, a method is provided for determining a presence or absence in a sample of a first misfolded protein aggregate. The method may include performing a first protein misfolding cyclic amplification (PMCA) procedure. The first PMCA procedure may include forming a first incubation mixture by contacting a first portion of the sample with a first substrate protein. The first substrate protein may include 4R tau protein. The first PMCA procedure may include conducting an incubation cycle two or more times under conditions effective to form as first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the first incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or

absence in the sample of the first misfolded protein aggregate by analyzing the first incubation mixture for the presence or absence of the first amplified, misfolded protein aggregate. The first misfolded protein aggregate may include the first substrate protein. The first amplified, misfolded protein aggregate may include the first substrate protein.

[0075] In various embodiments, a method is provided for determining a presence or absence in a subject of a tauopathy corresponding to a first misfolded protein aggregate. The method may include providing a sample from the subject. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting a first portion of the sample with a first substrate protein. The first substrate protein may include a tau isoform. The first substrate protein may be subject to pathological misfolding and/or aggregation in vivo to form the first misfolded protein aggregate. The first PMCA procedure may include conducting an incubation cycle two or more times under conditions effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the first incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by analyzing the first incubation mixture for the presence or absence of the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence of the tauopathy in the subject according the presence or absence of the first misfolded protein aggregate in the sample. The first misfolded protein aggregate may include the first substrate protein. The first amplified, misfolded protein aggregate may include the first substrate protein. The method may provide that the tauopathy excludes Pick's disease when the first substrate protein consists of monomeric 3R tau.

[0076] In various embodiments, a method is provided using capturing for determining a presence or absence in a sample of a first misfolded protein aggregate. The method may include capturing the first misfolded protein aggregate from the sample to form a captured first misfolded protein aggregate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the captured first misfolded protein aggregate with a molar excess of a first substrate protein. The first substrate protein may be subject to pathological misfolding and/or aggregation in vivo to form the first misfolded protein aggregate. The molar excess may be greater than an amount of protein monomer included in the captured first misfolded protein aggregate. The method may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein

aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the captured first misfolded protein aggregate. Each incubation cycle may include disrupting the first incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence of the first misfolded protein aggregate in the sample by detecting the first amplified, misfolded protein aggregate. The first misfolded protein aggregate may include the first substrate protein. The first amplified, misfolded protein aggregate may include the first substrate protein.

[0077] In various embodiments, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including Alzheimer's disease (AD). The method may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of AD according to the presence or absence of the first misfolded protein aggregate in the sample.

[0078] In some embodiments, determining the presence or absence in the subject of AD may include distinguishing AD from one or more additional tauopathies by determining a signature of AD tau protein aggregate. The signature AD tau protein aggregate may include one or more of: one or more AD-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a conformational epitope of AD tau protein aggregate; an indicator selective for AD tau protein aggregate; and a spectrum characteristic of AD tau protein aggregate.

[0079] In various embodiments, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including Parkinson's disease (PD). The method

may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of PD according to the presence or absence of the first misfolded protein aggregate in the sample.

[0080] In some embodiments, determining the presence or absence in the subject of PD may include distinguishing PD from one or more additional tauopathies by determining a signature of PD tau protein aggregate. The signature PD tau protein aggregate may include one or more of: one or more PD-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a conformational epitope of PD tau protein aggregate; an indicator selective for PD tau protein aggregate; and a spectrum characteristic of PD tau protein aggregate.

[0081] In various embodiments, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including Progressive Supranuclear Palsy (PSP). The method may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate.

The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of PSP according to the presence or absence of the first misfolded protein aggregate in the sample.

[0082] In some embodiments, determining the presence or absence in the subject of PSP may include distinguishing PSP from one or more additional tauopathies by determining a signature of PSP tau protein aggregate. The signature PSP tau protein aggregate may include one or more of: one or more PSP-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a conformational epitope of PSP tau protein aggregate; an indicator selective for PSP tau protein aggregate; and a spectrum characteristic of PSP tau protein aggregate.

[0083] In various embodiments, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including FrontoTemporal Dementia (FTD). The method may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of FTD according to the presence or absence of the first misfolded protein aggregate in the sample.

[0084] In some embodiments, determining the presence or absence in the subject of FTD may include distinguishing FTD from one or more additional tauopathies by determining a signature of FTD tau protein aggregate. The signature FTD tau protein aggregate may include one or more of: one or more FTD-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a

conformational epitope of FTD tau protein aggregate; an indicator selective for FTD tau protein aggregate; and a spectrum characteristic of FTD tau protein aggregate.

[0085] In various embodiments, a method is provided for determining a presence or absence of a tauopathy in a subject, the tauopathy including Corticobasal degeneration (CBD). The method may include providing the subject. The method may include obtaining a sample from the subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The method may include performing at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting the sample with a first substrate protein. The first substrate protein may include 4R tau. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The first PMCA procedure may include determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate. The method may include determining the presence or absence in the subject of CBD according to the presence or absence of the first misfolded protein aggregate in the sample.

[0086] In some embodiments, determining the presence or absence in the subject of CBD may include distinguishing CBD from one or more additional tauopathies by determining a signature of CBD tau protein aggregate. The signature CBD tau protein aggregate may include one or more of: one or more CBD-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a conformational epitope of CBD tau protein aggregate; an indicator selective for CBD tau protein aggregate; and a spectrum characteristic of CBD tau protein aggregate.

[0087] In further embodiments, each of the methods described herein above may incorporate one or more of the following features. In particular, each feature described with reference to any protein substrate, misfolded protein aggregate, amplified misfolded protein aggregate, incubation mixture, PMCA procedure, portion of the sample, and the like, should be understood to describe, independently selected in various other embodiments, any other protein substrate, misfolded protein aggregate, amplified misfolded protein aggregate, incubation mixture, PMCA procedure, portion of the sample, and the like. For example, features described for a “first” protein substrate may, in some embodiments, also be independently selected to describe a “second”

protein substrate; features described for a “first” misfolded protein aggregate may also be independently selected to describe a “second” misfolded protein aggregate; features described for a “first” incubation mixture may also be independently selected to describe a “second” incubation mixture; features described for a “first” PMCA procedure may also be independently selected to describe a “second” PMCA procedure; and the like. Further, for example, features described with reference to “each” protein substrate, misfolded protein aggregate, amplified misfolded protein aggregate, incubation mixture, PMCA procedure, portion of the sample, and the like, should be understood to describe, independently selected in various other embodiments, any other enumerated element, e.g., “first,” “second,” “third,” and the like, as applied to the protein substrate, misfolded protein aggregate, amplified misfolded protein aggregate, incubation mixture, PMCA procedure, portion of the sample, and the like. For example, a description with reference to “each substrate protein” may be independently selected to describe and support recitations of a “first substrate protein,” a “second substrate protein,” a “third substrate protein,” and the like.

[0088] In several embodiments, features described generally for enumerated or specified elements, e.g., “first,” “second,” “each,” and the like, may be independently selected to be the same or distinct. For example, in some embodiments, a first substrate protein may include a 4R tau and a second substrate protein may include A β ; a condition such as a temperature may be selected independently for a first and second PMCA procedure, and the like. In several embodiments, some features described generally for such first and second elements may be selected to be the same, or to overlap, while other features described generally for such first and second elements may be independently selected to be distinct. For example, in some embodiments, first and second portions of the sample may be the same or combined, and first and second incubation mixtures may be the same or combined, while corresponding first and second PMCA procedures may be conducted in parallel or in series in the combined incubation mixture using different first and second substrate proteins, e.g., 4R tau and A β .

[0089] In several embodiments, one, two, or more instances may be independently selected for each protein substrate, misfolded protein aggregate, amplified misfolded protein aggregate, incubation mixture, PMCA procedure, portion of the sample, and the like. For example, various method embodiments may include a first PMCA procedure using 4R tau as a first substrate protein, a second PMCA procedure using A β as a second substrate protein, a third PMCA procedure using alpha synuclein as a third substrate protein, a fourth PMCA procedure using 3R tau as a fourth substrate protein, and the like. Such multiple PMCA procedures may be performed for a sample, e.g., a laboratory sample, or a sample drawn from a subject, such as a subject having a tauopathy. Such multiple PMCA procedures may be performed in parallel for each protein substrate,

misfolded protein aggregate, amplified misfolded protein aggregate, incubation mixture, PMCA procedure, portion of the sample, and the like, for example as follows.

[0090] In various embodiments, the method may include determining the presence in the sample of the first misfolded protein aggregate. The method may include performing at least a second PMCA procedure to determine the presence or absence in the sample of at least a second misfolded protein aggregate. The second PMCA procedure may include forming a second incubation mixture by contacting a second portion of the sample with a second substrate protein. The second substrate protein may be subject to pathological misfolding and/or aggregation in vivo to form the second misfolded protein aggregate. The second PMCA procedure may include conducting an incubation cycle two or more times under conditions effective to form a second amplified, misfolded protein aggregate. Each incubation cycle may include incubating the second incubation mixture effective to cause misfolding and/or aggregation of the second substrate protein in the presence of the second misfolded protein aggregate. Each incubation cycle may include disrupting the second incubation mixture effective to form the second amplified, misfolded protein aggregate. The second PMCA procedure may include determining the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate. The second misfolded protein aggregate may include the second substrate protein. The second amplified, misfolded protein aggregate may include the second substrate protein.

[0091] In some embodiments, the method may include determining the presence in the sample of the first misfolded protein aggregate and the second misfolded protein aggregate. The method may include performing at least a third PMCA procedure to determine the presence or absence in the sample of at least a third misfolded protein aggregate. The third PMCA procedure may include forming a third incubation mixture by contacting a third portion of the sample with a third substrate protein. The third substrate protein may be subject to pathological misfolding and/or aggregation in vivo to form the third misfolded protein aggregate. The third PMCA procedure may include conducting an incubation cycle two or more times under conditions effective to form a third amplified, misfolded protein aggregate. Each incubation cycle may include incubating the third incubation mixture effective to cause misfolding and/or aggregation of the third substrate protein in the presence of the third misfolded protein aggregate. Each incubation cycle may include disrupting the third incubation mixture effective to form the third amplified, misfolded protein aggregate. The third PMCA procedure may include determining the presence or absence in the sample of the third misfolded protein aggregate by analyzing the third incubation mixture for the presence or absence of the third amplified, misfolded protein aggregate.

The third misfolded protein aggregate may include the third substrate protein. The third amplified, misfolded protein aggregate may include the third substrate protein.

[0092] In several embodiments, the method may include determining the presence in the sample of the first misfolded protein aggregate, the second misfolded protein aggregate, and the fourth misfolded protein aggregate. The method may include performing at least a fourth PMCA procedure to determine the presence or absence in the sample of a fourth misfolded protein aggregate. The fourth PMCA procedure may include forming a fourth incubation mixture by contacting a fourth portion of the sample with a fourth substrate protein. The fourth substrate protein may be subject to pathological misfolding and/or aggregation *in vivo* to form the fourth misfolded protein aggregate. The fourth PMCA procedure may include conducting an incubation cycle two or more times under conditions effective to form a fourth amplified, misfolded protein aggregate. Each incubation cycle may include incubating the fourth incubation mixture effective to cause misfolding and/or aggregation of the fourth substrate protein in the presence of the fourth misfolded protein aggregate. Each incubation cycle may include disrupting the fourth incubation mixture effective to form the fourth amplified, misfolded protein aggregate. The fourth PMCA procedure may include determining the presence or absence in the sample of the fourth misfolded protein aggregate by analyzing the fourth incubation mixture for the presence or absence of the fourth amplified, misfolded protein aggregate. The fourth misfolded protein aggregate may include the fourth substrate protein. The fourth amplified, misfolded protein aggregate may include the fourth substrate protein.

[0093] In various embodiments, the first substrate protein may independently include a tau isoform, e.g., 3R tau, 4R tau, and the like. In several embodiments, the first substrate protein may include 4R tau. The first substrate protein may include 3R tau. The first substrate protein may exclude 3R tau, for example, when the sample corresponds to Pick's disease or is drawn from a subject having Pick's disease. The first substrate protein may be soluble. The first substrate protein may be monomeric. The first substrate protein may be in a native *in vivo* conformation, e.g., folded. The first substrate protein may be distinct from each other substrate protein.

[0094] In various embodiments, the second substrate protein may independently include one of: a tau isoform, e.g., 3R tau, 4R tau, and the like; A β , alpha synuclein, and the like. The second substrate protein may include one of: 3R tau, A β , alpha synuclein, and the like. The second substrate protein may include 3R tau. The second substrate protein may include A β . The second substrate protein may include alpha synuclein. The second substrate protein may consist essentially of, or consist of, one of: 3R tau, 4R tau, A β , or alpha synuclein. The second substrate protein may be soluble. The second substrate protein may be monomeric. The second substrate

protein may be in a native in vivo conformation, e.g., folded. The second substrate protein may be distinct from each other substrate protein.

[0095] In various embodiments, the third substrate protein may independently include one of: a tau isoform, e.g., 3R tau, 4R tau, and the like; A β , alpha synuclein, and the like. The third substrate protein may include one of: 3R tau, A β , alpha synuclein, and the like. The third substrate protein may include 3R tau. The third substrate protein may include A β . The third substrate protein may include alpha synuclein. The third substrate protein may consist essentially of, or consist of, one of: 3R tau, 4R tau, A β , or alpha synuclein. The third substrate protein may be soluble. The third substrate protein may be monomeric. The third substrate protein may be in a native in vivo conformation, e.g., folded. The third substrate protein may be distinct from each other substrate protein.

[0096] In various embodiments, the fourth substrate protein may independently include one of: a tau isoform, e.g., 3R tau, 4R tau, and the like; A β , alpha synuclein, and the like. The fourth substrate protein may include one of: 3R tau, A β , alpha synuclein, and the like. The fourth substrate protein may include 3R tau. The fourth substrate protein may include A β . The fourth substrate protein may include alpha synuclein. The fourth substrate protein may consist essentially of, or consist of, one of: 3R tau, 4R tau, A β , or alpha synuclein. The fourth substrate protein may be soluble. The fourth substrate protein may be monomeric. The fourth substrate protein may be in a native in vivo conformation, e.g., folded. The fourth substrate protein may be distinct from each other substrate protein.

[0097] In some embodiments the sample may be taken from a subject. The method may include determining or diagnosing the presence or absence of a tauopathy in the subject according to the presence or absence of the first misfolded protein aggregate in the sample.

[0098] In several embodiments, the method may include performing at least a second PMCA procedure to determine the presence or absence in the sample of second misfolded protein aggregate, e.g., a misfolded protein aggregate that includes a second substrate protein. The second PMCA procedure may include forming a second incubation mixture by contacting a second portion of the sample with a second substrate protein. The second substrate protein may be subject to pathological misfolding and/or aggregation in vivo to form the second misfolded protein aggregate. The methods may include determining the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate. The second misfolded protein aggregate may include the second substrate protein. The second amplified, misfolded protein

aggregate may include the second substrate protein. The second substrate protein may include one of: amyloid-beta (A β), alpha synuclein, and 3R tau.

[0099] In some embodiments, the sample may be taken from a subject. The methods may include determining or diagnosing the presence or absence of a tauopathy in the subject according to the presence or absence of the first misfolded protein aggregate in the sample. The methods may include performing at least a second PMCA procedure to determine the presence or absence in the sample of a second misfolded protein aggregate. The second PMCA procedure may include forming a second incubation mixture by contacting a second portion of the sample with a second substrate protein. The second substrate protein may be subject to pathological misfolding and/or aggregation in vivo to form the second misfolded protein aggregate. The second PMCA procedure may include conducting an incubation cycle two or more times under conditions effective to form a second amplified, misfolded protein aggregate. Each incubation cycle may include incubating the second incubation mixture effective to cause misfolding and/or aggregation of the second substrate protein in the presence of the second misfolded protein aggregate. Each incubation cycle may include disrupting the second incubation mixture effective to form the second amplified, misfolded protein aggregate. The second PMCA procedure may include determining the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate. The second misfolded protein aggregate may include the second substrate protein. The second amplified, misfolded protein aggregate may include the second substrate protein.

[00100] In various embodiments, the subject may have the tauopathy. The methods may include characterizing an identity of the tauopathy in the subject according to: the presence in the sample of the first misfolded protein aggregate; and the presence or absence in the sample of the second misfolded protein aggregate. The second substrate protein may include one of: amyloid-beta (A β), alpha synuclein, and 3R tau. For example, the presence of misfolded 4R tau aggregate as the first misfolded protein aggregate and A β as the second misfolded protein aggregate may indicate the tauopathy in the subject is AD; the presence of misfolded 4R tau aggregate as the first misfolded protein aggregate and alpha synuclein as the second misfolded protein aggregate may indicate the tauopathy in the subject is PD; the presence of misfolded 4R tau aggregate as the first misfolded protein aggregate and the absence of the second misfolded protein aggregate including A β , alpha synuclein, or 3R may indicate a 4R tauopathy, such as PSP, CBD, AGD, and the like.

[00101] In various embodiments, the tauopathy may include a primary tauopathy or a secondary tauopathy. The tauopathy may be characterized at least in part by misfolding and/or aggregation of 4R tau protein. The tauopathy may be characterized at least in part by misfolding

and/or aggregation of 4R tau protein and 3R tau protein. The tauopathy may be characterized at least in part by misfolded and/or aggregated 4R tau protein, in a ratio to misfolded and/or aggregated 3R tau protein, of one of about: 1:99, 5:95, 10:90, 15:85, 20:80, 25:75, 30:70, 35:65, 40:60, 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5, and 99:1, or a range between any two of the preceding ratios, for example, between 1:99 and 99:1.

[00102] In several embodiments, the methods may include characterizing an identity of the tauopathy by analyzing the first amplified, misfolded protein aggregate or one or more corresponding PMCA kinetic parameters thereof for a signature of at least one of: Alzheimer's disease (AD), Parkinson's Disease (PD), Progressive Supranuclear Palsy (PSP), Frontotemporal Dementia (FTD), Corticobasal degeneration (CBD), Mild cognitive impairment (MCI), Argrophilic grain disease (AgD) Traumatic Brain Injury (TBI), Chronic Traumatic Encephalopathy (CTE), and Dementia Pugilistica (DP). For example, characterizing the identity of the tauopathy may include determining the one or more corresponding PMCA kinetic parameters, including one or more of: lag phase, T_{50} , amplification rate, and amplification extent. Characterizing the identity of the tauopathy may include comparing the one or more corresponding PMCA kinetic parameters to one or more corresponding predetermined corresponding PMCA kinetic parameters that are characteristic of the identity of the tauopathy to determine a similarity or difference effective to characterize the identity of the tauopathy.

[00103] In some embodiments, the methods may include characterizing the identity of the tauopathy using an antibody selective for a conformational epitope of a tauopathy-specific misfolded tau protein aggregate. The methods may include characterizing the identity of the tauopathy using an indicator selective for each tauopathy-specific misfolded tau protein aggregate. The indicator selective for each tauopathy-specific misfolded tau protein aggregate may include a small molecule, a peptide, or a DNA or RNA aptamer; and the like. The methods may include characterizing the identity of the tauopathy using a spectrum characteristic of each tauopathy-specific misfolded tau protein aggregate.

[00104] In some embodiments, the methods may include, for example, characterizing the identity of the tauopathy by analyzing the proteolytic resistance of each tauopathy-specific misfolded tau protein aggregate. For example, each tauopathy-specific misfolded tau protein aggregate may be contacted with a proteinase, e.g., proteinase K, trypsin, chymotrypsin, and the like, at a proteinase concentration of from 0.1 to 5000 $\mu\text{g/mL}$, at various temperatures from 20° C to 120° C and for various times, e.g., from 1 min to 4 h. The proteolytic resistance of each tauopathy-specific misfolded tau protein aggregate may be characterized and used to distinguish the various tauopathy-specific misfolded tau protein aggregates.

[00105] In several embodiments, the methods may include characterizing the identity of the tauopathy by analyzing the stability to denaturation of each tauopathy-specific misfolded tau protein aggregate. For example, each tauopathy-specific misfolded tau protein aggregate may be treated with guanidinium or urea at a sufficiently elevated temperature to induce protein denaturation of each tauopathy-specific misfolded tau protein aggregate. The concentration of guanidinium or urea may range from 0.1 M to 8 M. The temperature may range between 20° C to 120° C. The stability of each tauopathy-specific misfolded tau protein aggregate may be characterized and used to distinguish the various tauopathy-specific misfolded tau protein aggregates.

[00106] The methods may include sedimentation of each tauopathy-specific misfolded tau protein aggregate. The methods may include gel chromatography to characterize the size of each tauopathy-specific misfolded tau protein aggregate. The methods may include circular dichroism spectroscopy of each tauopathy-specific misfolded tau protein aggregate. The methods may include Fourier transform infrared spectroscopy to analyze secondary structure of each tauopathy-specific misfolded tau protein aggregate. The methods may include nuclear magnetic resonance spectroscopy to analyze structure of each tauopathy-specific misfolded tau protein aggregate. The methods may include mass spectrometry, e.g., fragmentation and collision induced dissociation to analyze secondary and tertiary structure of each tauopathy-specific misfolded tau protein aggregate. The methods may include microscopy, e.g., atomic force microscopy, cryo-electron microscopy, and the like to analyze morphology of each tauopathy-specific misfolded tau protein aggregate. Each of these methods may be coupled with substitution using atomic isotopes of different mass, magnetic properties, and/or isotopic stability to complement the methods; for example, nuclear magnetic resonance spectroscopy may be coupled with deuterium exchange in each tauopathy-specific misfolded tau protein aggregate to obtain structural information.

[00107] In various embodiments, the methods are provided such that the tauopathy specifically excludes Pick's disease. In various embodiments, the exclusion of Pick's disease does not encompass the remainder of Pick's complex of diseases.

[00108] In several embodiments, the methods may include determining or diagnosing the presence or absence of a tauopathy in the subject including comparing the presence or absence of the first misfolded protein aggregate in the sample to a control sample taken from a control subject. The detecting may include detecting an amount of the first misfolded protein aggregate in the sample. The sample may be taken from a subject. The methods may include determining or diagnosing the presence or absence of a tauopathy in the subject by comparing the amount of the first misfolded protein aggregate in the sample to a predetermined threshold amount. The sample

may be taken from a subject exhibiting no clinical signs of dementia according to cognitive testing. The methods may include determining or diagnosing the presence or absence of a tauopathy in the subject according to the presence or absence of the first misfolded protein aggregate in the sample. The sample may be taken from a subject exhibiting no cortex plaques or tangles according to contrast imaging. The methods may include determining or diagnosing the presence or absence of a tauopathy in the subject according to the presence or absence of the first misfolded protein aggregate in the sample. The sample may be taken from a subject exhibiting clinical signs of dementia according to cognitive testing. The methods may include determining or diagnosing the presence or absence of a tauopathy as a contributing factor to the clinical signs of dementia in the subject according to the presence or absence of the first misfolded protein aggregate in the sample. The sample may be taken from a subject exhibiting no clinical signs of dementia according to cognitive testing. The subject may exhibit a predisposition to dementia according to genetic testing. The genetic testing may indicate, for example, an increased risk of tauopathy according to one or two copies of the ApoE4 allele, variants of the brain derived neurotrophic factor (BDNF) gene, such as the val66met allele, in which valine at AA position 66 is replaced by methionine, and the like. The methods may include determining or diagnosing the presence or absence of a tauopathy in the subject according to the presence or absence of the first misfolded protein aggregate in the sample.

[00109] In some embodiments, the methods may include preparing the first incubation mixture characterized by at least one concentration of: the first substrate protein of less than about 20 μM ; heparin of less than about 75 μM ; NaCl of less than about 190 mM; and Thioflavin T of less than about 9.5 μM .

[00110] In various embodiments, the methods may include preparing the first incubation mixture including the first substrate protein at a concentration in μM of one or more of about: 0.001, 0.01, 0.1, 0.25, 0.5, 0.75, 1, 1.25, 1.5, 1.75, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5, 11, 11.5, 12, 12.5, 13, 13.5, 14, 14.5, 15, 15.5, 16, 16.5, 17, 17.5, 18, 18.5, 19, 19.5, 20, 25, 50, 70, 100, 150, 200, 250, 500, 750, 1000, 1500, or 2000, or a range between any two of the preceding values, for example, between about 0.001 μM and about 2000 μM . The methods may include preparing the first incubation mixture characterized by heparin at a concentration in μM of one or more of about: 0.001, 0.01, 0.1, 0.25, 0.5, 0.75, 1, 1.25, 1.5, 1.75, 2, 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9, 10, 11, 12, 12.5, 15, 17.5, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, and 75, or a range between any two of the preceding values, for example, between about 0.001 μM and about 75 μM . The methods may include preparing the first incubation mixture including a buffer composition of one or more of: Tris-HCL, PBS, MES, PIPES, MOPS, BES, TES, and

HEPES. The methods may include preparing the first incubation mixture including the buffer composition at a total concentration of one or more of about: 1 μ M, 10 μ M, 100 μ M, 250 μ M, 500 μ M, 750 μ M, 1 mM, 10 mM, 100 mM, 250 mM, 500 mM, 750 mM, and 1M, or a range between any two of the preceding values, for example, between about 1 μ M and about 1 M. The methods may include preparing the first incubation mixture including a salt composition at a total concentration of one or more of: 1 μ M, 10 μ M, 100 μ M, 250 μ M, 500 μ M, 750 μ M, 1 mM, 10 mM, 20 mM, 30 mM, 40 mM, 50 mM, 60 mM, 70 mM, 80 mM, 90 mM, 100 mM, 110 mM, 120 mM, 130 mM, 140 mM, 150 mM, 160 mM, 170 mM, 180 mM, 190 mM, 200 mM, 250 mM, 500 mM, 750 mM, and 1M, or a range between any two of the preceding values, for example, between about 1 μ M and about 1 M. The salt composition may include one or more of: NaCl and KCl.

[00111] In various embodiments, the methods may include preparing or maintaining the first incubation mixture at a pH of one or more of about: 5, 5.5, 6, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9, 8, 8.1, 8.2, 8.3, 8.4, 8.5, or 9, or a range between any two of the preceding values, e.g., from about pH 5 to about pH 9.

[00112] In some embodiments, the methods may include preparing the first incubation mixture including an indicator at a total concentration of one or more of: 1 nM, 10 nM, 100 nM, 250 nM, 500 nM, 750 nM, 1 μ M, 2 μ M, 3 μ M, 4 μ M, 5 μ M, 6 μ M, 7 μ M, 8 μ M, 9 μ M, 9.5 μ M, 10 μ M, 25 μ M, 50 μ M, 100 μ M, 250 μ M, 500 μ M, 750 μ M, 1 mM, or a range between any two of the preceding values, for example, between about 1 nM and about 1 mM.

[00113] In some embodiments of the methods, the incubating may include heating or maintaining the first incubation mixture at a temperature in $^{\circ}$ C of one of: 5, 10, 15, 20, 22.5, 25, 27.5, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 50, 55, 60, or a range between any two of the preceding values, for example, between about 5 $^{\circ}$ C and about 60 $^{\circ}$ C.

[00114] In several embodiments, the methods may include contacting an indicator of the first misfolded protein aggregate to the first incubation mixture. The indicator of the first misfolded protein aggregate may be characterized by an indicating state in the presence of the first misfolded protein aggregate and a non-indicating state in the absence of the first misfolded protein aggregate. Determining the presence of the first misfolded protein aggregate in the sample may include detecting the indicating state of the indicator of the first misfolded protein aggregate. The indicating state of the indicator and the non-indicating state of the indicator may be characterized by a difference in fluorescence. Determining the presence of the first misfolded protein aggregate in the sample may include detecting the difference in fluorescence. The methods may include contacting a molar excess of the indicator of the first misfolded protein aggregate to the first incubation mixture. The molar excess may be greater than a total molar amount of protein

monomer included in the first substrate protein and the first misfolded protein aggregate in the first incubation mixture. The indicator of the first misfolded protein aggregate may include one or more of: a thioflavin, e.g., thioflavin T or thioflavin S; Congo Red, m-I-Stilbene, Chrysamine G, PIB, BF-227, X-34, TZDM, FDDNP, MeO-X-04, IMPY, NIAD-4, luminescent conjugated polythiophenes, a fusion with a fluorescent protein such as green fluorescent protein and yellow fluorescent protein, derivatives thereof, and the like.

[00115] In various embodiments, the method may include determining an amount of the first misfolded protein aggregate in the sample. For example, known amounts of in vitro, synthetic misfolded protein aggregate seeds may be added to various portions of a biological fluid of a healthy patient, e.g., CSF. Subsequently, PMCA may be performed on the various portions. In each of the various portions, a fluorescent indicator of the misfolded protein aggregate may be added, and fluorescence may be measured as a function of, e.g., number of PMCA cycles, to determine various PMCA kinetics parameters, e.g., number of PMCA cycles to maximum fluorescence signal, number of PMCA cycles to 50% of maximum fluorescence signal, lag phase in increase of fluorescence signal, rate of increase in fluorescence signal versus PMCA cycles, and the like. A calibration curve showing the relationship between the concentration of synthetic seeds added and the PMCA kinetic parameters. The kinetic parameters may be measured for unknown samples and compared to the calibration curve to determine the expected amount of seeds present in a particular sample. Alternatively, the amount of the first misfolded protein aggregate in the sample may be determined by a series of known dilutions of the sample, and PMCA of each serial dilution to determine whether the first misfolded protein aggregate can be detected or not in a particular dilution. The amount of the first misfolded protein aggregate in the undiluted sample can be estimated based on the known dilution that results in no detection of the first misfolded protein aggregate by PMCA. In another example, the amount of the first misfolded protein aggregate in the sample may be determined by a series of known dilutions of the sample, and PMCA to determine a detection signal in each serial dilution. The collected detection signals in the serial dilutions can be fit, e.g., via least squares analysis, to determine whether the first misfolded protein aggregate can be detected or not in a particular dilution. In another example, the amount of the first misfolded protein aggregate in the sample may be determined by known amounts of antibodies to the first misfolded protein aggregate.

[00116] In some embodiments, the methods may include detecting the amount of the first misfolded protein aggregate in the sample at a sensitivity of at least about one or more of: 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, and 100%, e.g., at least about

70%. The methods may include detecting the amount of the first misfolded protein aggregate in the sample at less than about one or more of: 625, 62.5, 6.25, 630 μ g, 63 μ g, 6.3 μ g, 630 ng, 63 ng, 6.3 ng, 630 pg, 200 pg, 63 pg, 6.3 pg, 630 fg, 300 fg, 200 fg, 125 fg, 63 fg, 50 fg, 30 fg, 15 fg, 12.5 fg, 10 fg, 5 fg, or 2.5 fg. The methods may include detecting the amount of the first misfolded protein aggregate in the sample at less than about one or more of: 100 nmol, 10 nmol, 1 nmol, 100 pmol, 10 pmol, 1 pmol, 100 fmol, 10 fmol, 3 fmol, 1 fmol, 100 attomol, 10 attomol, 5 attomol, 2 attomol, 1 attomol, 0.75 attomol, 0.5 attomol, 0.25 attomol, 0.2 attomol, 0.15 attomol, 0.1 attomol, and 0.05 attomol, e.g., less than about 100 nmol. The methods may include detecting the amount of the first misfolded protein aggregate in the sample in a molar ratio to the first substrate protein included by the sample. The molar ratio may be less than about one or more of: 1: 100, 1: 10,000, 1: 100,000, and 1: 1,000,000, e.g., less than about 1: 100. The methods may include determining the amount of the first misfolded protein aggregate in the sample compared to a control sample.

[00117] In several embodiments, the methods may include detecting the first misfolded protein aggregate in the sample with a specificity of at least about one or more of: 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, and 100%, e.g. at least about 70%. The methods may include detecting the first misfolded protein aggregate including one or more of: a Western Blot assay, a dot blot assay, an enzyme-linked immunosorbent assay (ELISA), a fluorescent protein/peptide binding assay, a thioflavin binding assay, a Congo Red binding assay, a sedimentation assay, electron microscopy, atomic force microscopy, surface plasmon resonance, and spectroscopy. The ELISA may include a two-sided sandwich ELISA. The spectroscopy may include one or more of: quasi-light scattering spectroscopy, multispectral ultraviolet spectroscopy, confocal dual-color fluorescence correlation spectroscopy, Fourier-transform infrared spectroscopy, capillary electrophoresis with spectroscopic detection, electron spin resonance spectroscopy, nuclear magnetic resonance spectroscopy, and Fluorescence Resonance Energy Transfer (FRET) spectroscopy. Detecting the first misfolded protein aggregate may include contacting the first incubation mixture with a protease; and detecting the first misfolded protein aggregate using anti-misfolded protein antibodies or antibodies specific for a misfolded tau aggregate in one or more of: a Western Blot assay, a dot blot assay, and an ELISA.

[00118] In various embodiments, the methods may include providing the first substrate protein in labeled form. The first substrate protein in labeled form may include one or more of: a covalently incorporated radioactive amino acid, a covalently incorporated, isotopically labeled amino acid, and a covalently incorporated fluorophore. The methods may include detecting the

first substrate protein in labeled form as incorporated into the first amplified, misfolded protein aggregate.

[00119] In some embodiments, the sample may include one or more of a bio-fluid, e.g., blood, a biomaterial, e.g., cerumen, a homogenized tissue, and a cell lysate. The sample may include one or more of: amniotic fluid; bile; blood; cerebrospinal fluid; cerumen; skin; exudate; feces; gastric fluid; lymph; milk; mucus; mucosal membrane; peritoneal fluid; plasma; pleural fluid; pus; saliva; sebum; semen; sweat; synovial fluid; tears; and urine. The sample may be derived from cells or tissue of one or more of: skin, brain, heart, liver, pancreas, lung, kidney, gastro-intestine, nerve, mucous membrane, blood cell, gland, and muscle. The methods may include obtaining the sample from a subject, such as by drawing a bio-fluid or biomaterial, performing a tissue biopsy, and the like. The volume of each portion of the sample added to a particular PMCA reaction, e.g., in fluid or homogenized form, may be a volume in μL of one of about 5,000, 4,000, 3,000, 2,000, 1000, 900, 800, 750, 700, 650, 600, 550, 500, 450, 400, 350, 300, 250, 200, 150, 125, 100, 90, 80, 70, 60, 50, 40, 30, 25, 20, 15, 10, 5, or 1, or a range between any two of the preceding values, e.g., from about 1 μL to about 1000 μL . In some embodiments, when the sample is CSF, the amount of each portion added to a particular PMCA reaction may be a volume in μL of any of the preceding, for example, one of about 80, 70, 60, 50, 40, 30, 25, 20, 15, or 10, or a range between any two of the preceding values, e.g., e.g., from about 10 μL to about 80 μL , e.g., about 40 μL . In some embodiments, when the sample is plasma, the amount of each portion added to a particular PMCA reaction may be a volume in μL of any of the preceding, for example, one of about 750, 700, 650, 600, 550, 500, 450, 400, 350, 300, 250, or a range between any two of the preceding values, e.g., e.g., from about 250 μL to about 750 μL , e.g., about 500 μL . In some embodiments, when the sample is blood, the amount of each portion added to a particular PMCA reaction may be a volume in μL of any of the preceding, for example, one of about 5,000, 4,000, 3,000, 2,000, 1000, 900, 800, 750, 700, 650, 600, 550, 500, 450, 400, 350, 300, 250, or 200, or a range between any two of the preceding values, e.g., from about 200 μL to about 1000 μL .

[00120] In several embodiments, the subject may be one of a: human, mouse, rat, dog, cat, cattle, horse, deer, elk, sheep, goat, pig, and non-human primate. The subject may be one or more of: at risk of a tauopathy, having the tauopathy, and under treatment for the tauopathy. The methods may include determining a progression or homeostasis of a tauopathy in the subject by comparing the amount of the first misfolded protein aggregate in the sample to an amount of the first misfolded protein aggregate in a comparison sample taken from the subject at a different time compared to the sample. The subject may be treated with a tauopathy modulating therapy. The methods may include comparing the amount of the first misfolded protein aggregate in the sample

to an amount of the first misfolded protein aggregate in a comparison sample. The sample and the comparison sample may be taken from the subject at different times over a period of time under the tauopathy modulating therapy. The methods may include determining the subject is one of: responsive to the tauopathy modulating therapy according to a change in the first misfolded protein aggregate over the period of time, or non-responsive to the tauopathy modulating therapy according to homeostasis of the first misfolded protein aggregate over the period of time. The methods may include treating the subject determined to be responsive to the tauopathy modulating therapy with the tauopathy modulating therapy. The methods may include treating the subject with a tauopathy modulating therapy to inhibit production of the first substrate protein or to inhibit aggregation of the first misfolded protein aggregate.

[00121] In some embodiments, the subject may be treated with a protein misfolding disorder (PMD) modulating therapy. The method may include comparing the amount of the each misfolded protein aggregate in the sample to an amount of the each misfolded protein aggregate in a comparison sample. The sample and the comparison sample may be taken from the subject at different times over a period of time under the each misfolded protein aggregate modulating therapy. The method may include determining or diagnosing the subject is one of: responsive to the each misfolded protein aggregate modulating therapy according to a change in the each misfolded protein aggregate over the period of time, or non-responsive to the each misfolded protein aggregate modulating therapy according to homeostasis of the each misfolded protein aggregate over the period of time. The method may include treating the subject determined to be responsive to the each misfolded protein aggregate modulating therapy with the each misfolded protein aggregate modulating therapy. For AD, for example, the PMD modulating therapy may include administration of one or more of: an inhibitor of BACE1 (beta-secretase 1); an inhibitor of γ -secretase; and a modulator of A β homeostasis, e.g., an immunotherapeutic modulator of A β homeostasis. The A β modulating therapy may include administration of one or more of: E2609; MK-8931; LY2886721; AZD3293; semagacestat (LY-450139); avagacestat (BMS-708163); solanezumab; crenezumab; bapineuzumab; BIIB037; CAD106; 8F5 or 5598 or other antibodies raised against A β globulomers, e.g., as described by Barghorn et al, "Globular amyloid β -peptide₁₋₄₂ oligomer--a homogenous and stable neuropathological protein in Alzheimer's disease" *J. Neurochem.*, **2005**, 95, 834-847, the entire teachings of which are incorporated herein by reference; ACC-001; V950; Affitrope AD02; and the like.

[00122] For PD, for example, the PMD modulating therapy may include active immunization, such as PD01A+ or PD03A+, passive immunization such as PRX002, and the like. The PMD modulating therapy may also include treatment with GDNF (Glia cell-line derived

neurotrophic factor), inosine, Calcium-channel blockers, specifically Cav1.3 channel blockers such as isradipine, nicotine and nicotine-receptor agonists, GM-CSF, glutathione, PPAR-gamma agonists such as pioglitazone, and dopamine receptor agonists, including D2/D3 dopamine receptor agonists and LRRK2 (leucine-rich repeat kinase 2) inhibitors.

[00123] In several embodiments, the amount of misfolded protein may be measured in samples from patients using PMCA. Patients with elevated misfolded protein measurements may be treated with disease modifying therapies for a PMD. Patients with normal misfolded protein measurements may not be treated. A response of a patient to disease-modifying therapies may be followed. For example, misfolded protein levels may be measured in a patient sample at the beginning of a therapeutic intervention. Following treatment of the patient for a clinical meaningful period of time, another patient sample may be obtained and misfolded protein levels may be measured. Patients who show a change in misfolded protein levels following therapeutic intervention may be considered to respond to the treatment. Patients who show unchanged misfolded protein levels may be considered non-responding. The methods may include detection of misfolded protein aggregates in patient samples containing components that may interfere with the PMCA reaction.

[00124] In various embodiments, the methods may include selectively concentrating the first misfolded protein aggregate in one or more of the sample and the first incubation mixture. The selectively concentrating the first misfolded protein aggregate may include pre-treating the sample prior to forming the first incubation mixture. The selectively concentrating the first misfolded protein aggregate may include pre-treating the first incubation mixture prior to incubating the first incubation mixture. The selectively concentrating the first misfolded protein aggregate may include contacting one or more antibodies capable of binding the first misfolded protein aggregate to form a captured first misfolded protein aggregate. The one or more antibodies capable of binding the first misfolded protein aggregate may include one or more of: an antibody specific for an amino acid epitope sequence of the first misfolded protein aggregate, and an antibody specific for a conformation of the first misfolded protein aggregate. The antibody specific for a conformation of the first misfolded protein aggregate may be selective for a conformational epitope of a tauopathy-specific misfolded tau aggregate. The one or more one or more antibodies capable of binding the first misfolded protein aggregate may be coupled to a solid phase. The solid phase may include one or more of a magnetic bead and a multiwell plate.

[00125] For example, ELISA plates may be coated with the antibodies used to capture first misfolded protein aggregate from the patient sample. The antibody-coated ELISA plates may be incubated with a patient sample, unbound materials may be washed off, and the PMCA reaction

may be performed. Antibodies may also be coupled to beads. The beads may be incubated with the patient sample and used to separate first misfolded protein aggregate -antibody complexes from the remainder of the patient sample.

[00126] In some embodiments, contacting the sample with the first substrate protein to form the first incubation mixture may include contacting a molar excess of the first substrate protein to the sample including the captured first misfolded protein aggregate. The molar excess of the first substrate protein may be greater than a total molar amount of protein monomer included in the captured first misfolded protein aggregate. Incubating the first incubation mixture may be effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the captured first misfolded protein aggregate to form the first amplified, misfolded protein aggregate.

[00127] In several embodiments, disrupting the first incubation mixture may include physically disrupting and/or thermally disrupting. For example, the disrupting may include one or more of: sonication, stirring, shaking, freezing/thawing, laser irradiation, autoclave incubation, high pressure, and homogenization. Disrupting the first incubation mixture may include cyclic agitation. The cyclic agitation may be conducted for one or more of: between about 50 rotations per minute (RPM) and 10,000 RPM, between about 200 RPM and about 2000 RPM, and at about 500 RPM. Disrupting the first incubation mixture may be conducted in each incubation cycle for one or more of: between about 5 seconds and about 10 minutes, between about 30 sec and about 1 minute, between about 45 sec and about 1 minute, and about 1 minute. Incubating the first incubation mixture may be independently conducted, in each incubation cycle for one or more of: between about 1 minute and about 5 hours, between about 5 minutes and about 5 hours, between about 10 minutes and about 2 hours, between about 15 minutes and about 1 hour, and between about 25 minutes and about 45 minutes. Each incubation cycle may include independently incubating and disrupting the first incubation mixture for one or more of: incubating between about 1 minute and about 5 hours and disrupting between about 5 seconds and about 10 minutes; incubating between about 5 minutes and about 5 hours and disrupting between about 5 seconds and about 10 minutes; incubating between about 10 minutes and about 2 hours and disrupting between about 30 sec and about 1 minute; incubating between about 15 minutes and about 1 hour and disrupting between about 45 sec and about 1 minute; incubating between about 25 minutes and about 45 minutes and disrupting between about 45 sec and about 90 seconds; incubating for about 29 minutes and for about 1 minute; and incubating about 1 minute and disrupting about 1 minute. Conducting the incubation cycle may be repeated for one or more of: between about 2 times and about 1000 times, between about 5 times and about 500 times, between about 50 times and about 500 times, and between about 150 times and about 250 times.

[00128] In various embodiments, contacting the sample with the first substrate protein to form the first incubation mixture may be conducted under physiological conditions. the methods may include contacting the sample with a molar excess of the first substrate protein to form the first incubation mixture. The molar excess may be greater than a total molar amount of protein monomer included in the first misfolded protein aggregate in the sample.

[00129] In some embodiments, the methods may include contacting the sample with a thioflavin, e.g., thioflavin T or thioflavin S, and a molar excess of the first substrate protein to form the first incubation mixture. The molar excess may be greater than an amount of the first substrate protein included in the first misfolded protein aggregate in the sample. The method may include conducting the incubation cycle two or more times effective to form the first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least the portion of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include shaking the first incubation mixture effective to form the first amplified, misfolded protein aggregate. The methods may include determining the presence of the first misfolded protein aggregate in the sample by detecting a fluorescence of the thioflavin corresponding to the first misfolded protein aggregate.

[00130] In several embodiments, the first substrate protein may be produced by one of: chemical synthesis, recombinant production, and extraction from non-recombinant biological samples. The first misfolded protein aggregate may include one or more of a soluble first misfolded protein aggregate and an insoluble first misfolded protein aggregate. The first amplified, misfolded protein aggregate may include one or more of: a soluble portion and an insoluble portion. The first misfolded protein aggregate may be substantially be a soluble first misfolded protein aggregate. In some embodiments, the methods may provide that the sample excludes tau fibrils. For example, the sample may be filtered or centrifuged to remove tau fibrils.

[00131] In various embodiments, the second substrate protein may be distinct from the first substrate protein. The second substrate protein may include one of: amyloid-beta (A β), alpha synuclein, 3R tau, and 4R tau. The first substrate protein may include 4R tau.

[00132] In some embodiments, the methods may include performing at least a second PMCA procedure to determine the presence or absence in the sample of a second misfolded protein aggregate. The second PMCA procedure may include forming a second incubation mixture by contacting a second portion of the sample with a second substrate protein. The second substrate protein may be subject to pathological misfolding and/or aggregation in vivo to form the second misfolded protein aggregate. The second PMCA procedure may include conducting an incubation

cycle two or more times under conditions effective to form a second amplified, misfolded protein aggregate. Each incubation cycle may include incubating the second incubation mixture effective to cause misfolding and/or aggregation of the second substrate protein in the presence of the second misfolded protein aggregate. Each incubation cycle may include disrupting the second incubation mixture effective to form the second amplified, misfolded protein aggregate. The second PMCA procedure may include determining the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate. The second misfolded protein aggregate may include the second substrate protein. The second amplified, misfolded protein aggregate may include the second substrate protein.

[00133] In several embodiments, the tauopathy may be present in the subject. The methods may include characterizing an identity of the tauopathy in the subject according to the presence in the sample of the first misfolded protein aggregate. The methods may include characterizing an identity of the tauopathy in the subject according to the presence or absence in the sample of the second misfolded protein aggregate.

[00134] In some embodiments, the methods may provide that the tauopathy is not primarily characterized by misfolding and/or aggregation of 3R tau protein. For example, the tauopathy may be characterized at least in part by misfolded and/or aggregated 4R tau protein, in a ratio to misfolded and/or aggregated 3R tau protein, of one of about: 1:99, 5:95, 10:90, 15:85, 20:80, 25:75, 30:70, 35:65, 40:60, 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5, and 99:1, or a range between any two of the preceding ratios, for example, between 1:99 and 99:1.

[00135] In various embodiments, the methods may include contacting the sample with a thioflavin, e.g., thioflavin S or thioflavin T, and a molar excess of the first substrate protein to form the first incubation mixture. The molar excess may be greater than an amount of protein monomer included in the first misfolded protein aggregate in the sample. The methods may include conducting the incubation cycle two or more times effective to form the first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least the portion of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include shaking the first incubation mixture effective to form the first amplified, misfolded protein aggregate. The methods may include determining the presence or absence of the first misfolded protein aggregate in the sample by detecting a fluorescence of the thioflavin corresponding to the first misfolded protein aggregate.

[00136] In some embodiments of the methods, the capturing the first misfolded protein aggregate from the sample to form a captured first misfolded protein aggregate may be conducted using one or more antibodies specific for the first misfolded protein aggregate. The one or more antibodies specific for the first misfolded protein aggregate may include one or more of: an antibody specific for an amino acid epitope sequence of the first misfolded protein aggregate and an antibody specific for a conformation of the first misfolded protein aggregate. The antibody specific for a conformation of the first misfolded protein aggregate may be selective for a conformational epitope of a tauopathy-specific first misfolded protein aggregate. The antibody specific for the conformation of the first misfolded protein aggregate may correspond to one of: Alzheimer's disease (AD), Parkinson's Disease (PD), Progressive Supranuclear Palsy (PSP), Frontotemporal Dementia (FTD), Corticobasal degeneration (CBD), Mild cognitive impairment (MCI), Argrophilic grain disease (AgD) Traumatic Brain Injury (TBI), Chronic Traumatic Encephalopathy (CTE), and Dementia Pugilistica (DP). The one or more antibodies specific for the first misfolded protein aggregate may be coupled to a solid phase. The solid phase may include one or more of a magnetic bead and a multiwell plate. Contacting the sample with the first substrate protein to form the first incubation mixture may include contacting a molar excess of the first substrate protein to the sample. The molar excess of the first substrate protein may be greater than a total molar amount of protein monomer included in the captured first misfolded protein aggregate. Incubating the first incubation mixture may be effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the captured first misfolded protein aggregate to form the first amplified, misfolded protein aggregate. The first substrate protein may include 4R tau protein.

[00137] In various embodiments, the methods may include performing at least a second PMCA procedure to determine the presence or absence in the sample of a second misfolded protein aggregate. The second PMCA procedure may include forming a second incubation mixture by contacting a second portion of the sample with a second substrate protein, the second substrate protein may be subject to pathological misfolding and/or aggregation. The second PMCA procedure may include conducting an incubation cycle two or more times under conditions effective to form a second amplified, misfolded protein aggregate. Each incubation cycle may include incubating the second incubation mixture effective to cause misfolding and/or aggregation of the second substrate protein in the presence of the second misfolded protein aggregate. Each incubation cycle may include disrupting the second incubation mixture effective to form the second amplified, misfolded protein aggregate. The second PMCA procedure may include determining the presence or absence in the sample of the second misfolded protein aggregate by

analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate. The tauopathy may be present in the subject. The methods may include characterizing an identity of the tauopathy in the subject according to: the presence in the sample of the first misfolded protein aggregate; and the presence or absence in the sample of the second misfolded protein aggregate. The second misfolded protein aggregate may include the second substrate protein. The second amplified, misfolded protein aggregate may include the second substrate protein. The second substrate protein may be distinct from the first substrate protein. The second substrate protein may include one of: amyloid-beta ($A\beta$), alpha synuclein, and 3R tau.

[00138] In some embodiments, the methods may include distinguishing each tauopathy from one or more additional tauopathies by analyzing for at least one signature of one or more misfolded aggregates each corresponding to one of $A\beta$, alpha synuclein, and 3R tau. Each signature may correspond to one or more of: an assay using an antibody selective for a conformational epitope of any of the one or more misfolded aggregates; an assay using an antibody selective for a conformational epitope of any of the one or more misfolded aggregates; one or more PMCA kinetic parameters of the one or more misfolded aggregates, including one or more of: lag phase, T_{50} , amplification rate, and amplification extent; an indicator selective for any of the one or more misfolded aggregates; and a spectrum characteristic of any of the one or more misfolded aggregates.

[00139] Further, for example, specific antibodies may be employed for second misfolded protein aggregates. For example, for AD, amyloid antibodies may include one or more of: 6E10, 4G8, 82E1, A11, X-40/42, 16ADV; and the like. Such antibodies may be obtained as follows: 6E10 and 4G8 (Covance, Princeton, NJ); 82E1 (IBL America, Minneapolis, MN); A11 (Invitrogen, Carlsbad, CA); X-40/42 (MyBioSource, Inc., San Diego, CA); and 16ADV (Acumen Pharmaceuticals, Livermore, CA).

[00140] Further, for PD, for example, the one or more synuclein specific antibodies may include PD specific antibodies including one or more of: α/β -synuclein N-19; α -synuclein C-20-R; α -synuclein 211; α -synuclein Syn 204; α -synuclein 2B2D1; α -synuclein LB 509; α -synuclein SPM451; α -synuclein 3G282; α -synuclein 3H2897; α/β -synuclein Syn 202; α/β -synuclein 3B6; $\alpha/\beta/\gamma$ -synuclein FL-140; and the like. In some examples, the one or more specific antibodies may include one or more of: α/β -synuclein N-19; α -synuclein C-20-R; α -synuclein 211; α -synuclein Syn 204; and the like. Such antibodies may be obtained as follows: α/β -synuclein N-19 (cat. No. SC-7012, Santa Cruz Biotech, Dallas, TX); α -synuclein C-20-R (SC-7011-R); α -synuclein 211 (SC-12767); α -synuclein Syn 204 (SC-32280); α -synuclein 2B2D1 (SC-53955); α -synuclein LB

509 (SC-58480); α -synuclein SPM451 (SC-52979); α -synuclein 3G282 (SC-69978); α -synuclein 3H2897 (SC-69977); α/β -synuclein Syn 202 (SC-32281); α/β -synuclein 3B6 (SC-69699); or $\alpha/\beta/\gamma$ -synuclein FL-140 (SC-10717).

[00141] In various embodiments, a kit is provided for determining a presence or absence in a sample of a first misfolded protein aggregate. The kit may include a first substrate protein that may include 4R tau. The kit may include an indicator of the first misfolded protein aggregate. The first misfolded protein aggregate may include the first substrate protein. The first misfolded protein aggregate may correspond to a tauopathy. The kit may include a buffer. The kit may include heparin. The kit may include a salt. The kit may include instructions. The instructions may direct a user to obtain the sample. The instructions may direct the user to perform at least a first PMCA procedure. The first PMCA procedure may include forming a first incubation mixture by contacting a first portion of the sample with the first substrate protein, the indicator of the first misfolded protein aggregate, the buffer, the heparin, and the salt. The first incubation mixture may be formed with a concentration of one or more of: the first substrate protein of less than about 20 μ M; the heparin of less than about 75 μ M; the salt as NaCl of less than about 190 mM; and the indicator of the first misfolded protein aggregate as Thioflavin T of less than about 9.5 μ M. The first PMCA procedure may include conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate. Each incubation cycle may include incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate. The instructions may direct the user to determine the presence or absence in the sample of the first misfolded protein aggregate by analyzing the first incubation mixture for the presence or absence of the first amplified, misfolded protein aggregate according to the indicator of the first misfolded protein aggregate.

[00142] In several embodiments, the kit may include any element of the methods described herein. Moreover, the kit may include instructions directing the user to conduct any of the steps of the methods described herein.

[00143] In some embodiments, for example, the instructions may include directing the user to obtain the sample from a subject. The sample may include one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate. The instructions directing the user to determine or diagnose a tauopathy in the subject according to the presence or absence in the sample of the first misfolded protein aggregate.

[00144] In various embodiments, the kit may include a second substrate protein and an indicator of a second misfolded protein aggregate. The second misfolded protein aggregate may include the second substrate protein. The second substrate protein may be distinct from the first substrate protein. The second substrate protein may include one of: amyloid-beta ($A\beta$), alpha synuclein, 3R tau, and 4R tau. The instructions may direct the user to perform at least a second PMCA procedure. The second PMCA procedure may include forming a second incubation mixture by contacting a second portion of the sample with the second substrate protein and the indicator of the second misfolded protein aggregate. The second PMCA procedure may include conducting an incubation cycle two or more times effective to form a second amplified, misfolded protein aggregate. Each incubation cycle may include incubating the second incubation mixture effective to cause misfolding and/or aggregation of the second substrate protein in the presence of the second misfolded protein aggregate. Each incubation cycle may include disrupting the incubation mixture effective to form the second amplified, misfolded protein aggregate. The second PMCA procedure may include determining the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate. The instructions may also direct the user to characterize the sample for an identity of a tauopathy according to: the presence in the sample of the first misfolded protein aggregate; and the presence or absence in the sample of the second misfolded protein aggregate.

[00145] In some embodiments, the kit may include a PMCA apparatus. The PMCA apparatus may include one or more of: a multiwell microtitre plate; a microfluidic plate; a shaking apparatus; a spectrometer; and an incubator. The apparatus may be included either as one or more of the individual plates or apparatuses, as a combination device, and the like. For example, a shaking microplate reader may be used to perform cycles of incubation and shaking and automatically measure the ThT fluorescence emission during the course of an experiment (e.g., FLUOstar OPTIMA, BMG LABTECH Inc., Cary, NC).

[00146] The antibody specific for the conformation of the first misfolded protein aggregate may correspond to one of: Alzheimer's disease (AD), Parkinson's Disease (PD), Progressive Supranuclear Palsy (PSP), Frontotemporal Dementia (FTD), Corticobasal degeneration (CBD), Mild cognitive impairment (MCI), Argrophilic grain disease (AgD) Traumatic Brain Injury (TBI), Chronic Traumatic Encephalopathy (CTE), and Dementia Pugilistica (DP). The instructions may include determining, according to a binding assay using the antibody specific for the conformation of the first misfolded protein aggregate, the presence or absence in the subject of one of AD, PD, PSP, FTD, CBD, MCI, AgD, TBI, CTE, and DP.

EXAMPLES

EXAMPLE 1: PREPARATION OF SYNTHETIC A β OLIGOMERS

[00147] A β 1-42 was synthesized using solid-phase N-tert-butyloxycarbonyl chemistry at the W. Keck Facility at Yale University and purified by reverse-phase HPLC. The final product was lyophilized and characterized by amino acid analysis and mass spectrometry. To prepare stock solutions free of aggregated, misfolded A β protein, aggregates were dissolved high pH and filtration through 30 kDa cut-off filters to remove remaining aggregates. To prepare different types of aggregates, solutions of seed-free A β 1-42 (10 μ M) were incubated for different times at 25 °C in 0.1 M Tris-HCl, pH 7.4 with agitation. This preparation contained a mixture of A β monomers as well as fibrils, protofibrils and soluble, misfolded A β protein in distinct proportions depending on the incubation time. The degree of aggregation was characterized by ThT fluorescence emission, electron microscopy after negative staining, dot blot studies with the A11 conformational antibody and western blot after gel electrophoresis using the 4G8 anti-A β antibody.

[00148] A mixture of A β oligomers of different sizes were generated during the process of fibril formation. Specifically, soluble, misfolded A β protein was prepared by incubation of monomeric synthetic A β 1-42 (10 μ M) at 25 °C with stirring. After 5 h of incubation, an abundance of soluble, misfolded A β protein, globular in appearance, was observed by electron microscopy after negative staining, with only a small amount of protofibrils and fibrils observed. At 10 h there are mostly protofibrils and at 24 h, a large amount of long fibrils are observed. **FIG. 1A** shows electron micrographs taken at 0h, 5h, 10h, and 24h of incubation.

[00149] The soluble, misfolded A β protein aggregates tested positive using A11 anti-oligomer specific antibody according to the method of Kaye, et al. "Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis," *Science* **2003**, 300, 486-489. After further incubation at 10 h and 24 h, protofibrillar and fibrillar structures were observed. The size of the aggregates was determined by filtration through filters of defined pore size and western blotting after SDS-PAGE separation. Soluble, misfolded A β protein formed by incubation for 5 h was retained in filters of 30 kDa cut-off and passed through 1000 kDa cutoff filters. **FIG. 1B** is a western blot of soluble, misfolded A β protein aggregates. Electrophoretic separation of this soluble, misfolded A β protein showed that the majority of the material migrated as ~170 kDa SDS-resistant aggregates, with a minor band at 17 kDa.

EXAMPLE 2: A β -PMCA DETECTS SYNTHETIC A β OLIGOMERS

[00150] **EXAMPLE 2A.** Seeding of A β aggregation was studied by incubating a solution of seed-free A β 1-42 in the presence of Thioflavin T with or without different quantities of synthetic

soluble, misfolded A β protein (Control (no A β oligomer); or 3, 80, 300, and 8400 femtomolar in synthetic soluble, misfolded A β protein). A β -PMCA general procedure: Solutions of 2 μ M aggregate-free A β 1-42 in 0.1 M Tris-HCl pH 7.4 (200 μ L total volume) were placed in opaque 96-wells plates and incubated alone or in the presence of synthetic A β aggregates (prepared by incubation over 5h as described in **EXAMPLE 1**) or 40 μ L of CSF aliquots. Samples were incubated in the presence of 5 μ M Thioflavin T (ThT) and subjected to cyclic agitation (1 min at 500 rpm followed by 29 min without shaking) using an Eppendorf thermomixer, at a constant temperature of 22 °C. At various time points, ThT fluorescence was measured in the plates at 485 nm after excitation at 435 nm using a plate spectrofluorometer. **FIG. 2A** is a graph of amyloid formation (without cyclic amplification) versus time as measured by Thioflavin T fluorescence, using the indicated femtomolar concentration of synthetic soluble, misfolded A β protein seeds. The peptide concentration, temperature and pH of the buffer were monitored to control the extent of the lag phase and reproducibility among experiments. Under these conditions, no spontaneous A β aggregation was detected during the time in which the experiment was performed (125 h). Aggregation of monomeric A β 1-42 protein was observed in the presence of 0.3 to 8.4 fmol of the synthetic soluble, misfolded A β protein of **EXAMPLE 1**.

[00151] **EXAMPLE 2B:** Amplification cycles, combining phases of incubation and physical disruption were employed. The same samples as in **FIG. 2A** were incubated with cyclic agitation (1 min stirring at 500 rpm followed by 29 min without shaking). Aggregation was measured over time by the thioflavin T (ThT) binding to amyloid fibrils using a plate spectrofluorometer (excitation: 435; emission: 485 nm). Graphs show the mean and standard error of 3 replicates. The concentration of A β oligomers was estimated assuming an average molecular weight of 170 kDa. **FIG. 2B** is a graph showing amplification cycle-accelerated amyloid formation measured by ThT fluorescence as a function of time for various concentrations of the synthetic soluble, misfolded A β protein of **EXAMPLE 1**. Under these conditions, the aggregation of monomeric A β 1-42 protein induced by 8400, 300, 80 and 3 fmol of the synthetic soluble, misfolded A β protein was clearly faster and more easily distinguished from that observed in the absence of the synthetic soluble, misfolded A β protein. This result indicates the detection limit, under these conditions, is 3 fmol of soluble, misfolded A β protein or less in a given sample.

EXAMPLE 3: A β -PMCA DETECTS MISFOLDED A β IN THE CEREBROSPINAL FLUID OF AD PATIENTS

[00152] Aliquots of CSF were obtained from 50 AD patients, 39 cognitively normal individuals affected by non-degenerative neurological diseases (NND), and 37 patients affected by non-AD neurodegenerative diseases including other forms of dementia (NAND). Test CSF

samples were obtained from 50 patients with the diagnosis of probable AD as defined by the DSM-IV and the NINCDS-ADRA guidelines (McKhann et al., 1984) and determined using a variety of tests, including routine medical examination, neurological evaluation, neuropsychological assessment, magnetic resonance imaging and measurements of CSF levels of A β 1-42, total Tau and phospho-Tau. The mean age of AD patients at the time of sample collection was 71.0 ± 8.1 years (range 49-84). Control CSF samples were obtained from 39 patients affected by non-degenerative neurological diseases (NND), including 12 cases of normal pressure hydrocephalus, 7 patients with peripheral neuropathy, 7 with diverse forms of brain tumor, 2 with ICTUS, 1 with severe cephalgia, 3 with encephalitis, 1 with hypertension and 6 with unclear diagnosis. The mean age at which CSF samples were taken from this group of patients was 64.6 ± 14.7 years (range 31-83). Control CSF samples were also taken from 37 individuals affected by non-AD neurodegenerative diseases (NAND), including 10 cases of fronto-temporal dementia (5 behavioral and 5 language variants), 6 patients with Parkinson's disease (including 4 associated with dementia and 1 with motor neuron disease), 6 with progressive supranuclear palsy, 6 with spinocerebellar ataxia (1 associated with dementia), 4 with amyotrophic lateral sclerosis, 2 with Huntington's disease, 1 with MELAS, 1 with Lewy body dementia, and 1 with vascular dementia. The mean age at sample collection for this group was 63.8 ± 11.1 years (range 41-80). CSF samples were collected in polypropylene tubes following lumbar puncture at the L4/L5 or L3/L4 interspace with atraumatic needles after one night fasting. The samples were centrifuged at 3,000 g for 3 min at 4°C, aliquoted and stored at -80°C until analysis. CSF cell counts, glucose and protein concentration were determined. Albumin was measured by rate nephelometry. To evaluate the integrity of the blood brain barrier and the intrathecal IgG production, the albumin quotient (CSF albumin/serum albumin) $\times 10^3$ and the IgG index (CSF albumin/serum albumin)/(CSF IgG/serum IgG) were calculated. The study was conducted according to the provisions of the Helsinki Declaration and was approved by the Ethics Committee.

[00153] The experiments as well as the initial part of the analysis were conducted blind.

FIG. 3A is a graph of amyloid formation versus time, measured as a function of ThT fluorescence labeling, showing the average kinetics of A β aggregation of 5 representative samples from the AD, NND, and NAND groups.

[00154] The results indicate that CSF from AD patients significantly accelerates A β aggregation as compared to control CSF ($P < 0.001$). The significance of the differences in A β aggregation kinetics in the presence of human CSF samples was analyzed by one-way ANOVA, followed by the Tukey's multiple comparison post-test. The level of significance was set at

$P < 0.05$. The differences between AD and samples from the other two groups were highly significant with $P < 0.001$ (***).

[00155] **FIG. 3B** is a graph of the lag phase time in h for samples from the AD, NND, and NAND groups. To determine the effect of individual samples on A β aggregation, the lag phase was estimated, defined as the time to ThT fluorescence larger than 40 arbitrary units after subtraction of a control buffer sample. This value was selected considering that it corresponds to ~5 times the reading of the control buffer sample.

[00156] **FIG. 3C** is a graph showing the extent of amyloid formation obtained after 180 A β -PMCA cycles, e.g. 90 h of incubation (P90). Comparison of the lag phase and P90 among the experimental groups reveals a significant difference between AD and control samples from individuals with non-degenerative neurological diseases or with non-AD neurodegenerative diseases. Further, no correlation was detected between the aggregation parameters and the age of the AD patients, which indicates that the levels of the marker corresponds to aggregated A β protein in patient CSF, and not patient age.

[00157] **FIG. 5, Table 1** shows estimations of the sensitivity, specificity and predictive value of the A β -PMCA test, calculated using the lag phase numbers.

[00158] To study reproducibility, an experiment similar to the one shown in **FIGS. 3A-C** was independently done with different samples, reagents and a new batch of A β peptide as substrate for A β -PMCA. The extent of amyloid formation obtained after 300 A β -PMCA cycles, e.g. 150 h of incubation (P150), was measured in each patient. The control group includes both people affected by other neurodegenerative diseases and non-neurologically sick patients. Data for each sample represent the average of duplicate tubes. Statistical differences were analyzed by student-t test. **FIG. 6** is a graph of the lag phase time in h for samples obtained after 300 A β -PMCA cycles, e.g. 150 h of incubation (P90).

[00159] During the course of the study an entire set of CSF samples coming from a fourth location did not aggregate at all, even after spiking with large concentrations of synthetic oligomers. It is expected that reagent contamination during sample collection interfered with the assay.

[00160] The differences in aggregation kinetics between different samples were evaluated by the estimation of various different kinetic parameters, including the lag phase, A50, and P90. Lag phase is defined as the time required to reach a ThT fluorescence higher than 5 times the background value of the buffer alone. The A50 corresponds to the time to reach 50% of maximum aggregation. P90 corresponds to the extent of aggregation (measured as ThT fluorescence) at 90 h. Sensitivity, specificity and predictive value were determined using this data, with cutoff

thresholds determined by Receiver Operating Characteristics (ROC) curve analysis, using MedCalc software (MedCalc Software, Ostend, Belgium).

EXAMPLE 4: DETERMINATION OF THRESHOLD VALUES OF MISFOLDED A β FOR A β -PMCA DETECTION OF AD IN CSF

[00161] In support of **FIG. 5, TABLE 1**, sensitivity, specificity and predictive value were determined using the lag phase data, with cutoff thresholds determined by Receiver Operating Characteristics (ROC) curve analysis, using the MedCalc software (version 12.2.1.0, MedCalc, Belgium). As shown in **FIG. 5, TABLE 1**, a 90.0% sensitivity and 84.2% specificity was estimated for the control group consisting of age-matched individuals with non-degenerative neurological diseases. By contrast, for the clinically more relevant differentiation of AD from other neurodegenerative diseases including other forms of dementia, 100% sensitivity and 94.6% specificity was estimated. This ability of A β -PMCA to distinguish AD from other forms of neurodegenerative diseases is clinically significant. The overall sensitivity and specificity considering all control individuals was 90% and 92%, respectively.

[00162] To evaluate the performance of the A β -PMCA test to distinguish AD patients from controls, the true positive rate (sensitivity) was plotted as a function of the false positive rate (specificity) for different cut-off points. For this analysis the lag phase values for AD vs NAND (**FIG. 4A**), AD vs NND (**FIG. 4B**) and AD vs All control samples (**FIG. 4C**) was used. The performance of the test, estimated as the area under the curve was 0.996 ± 0.0033 , 0.95 ± 0.020 and 0.97 ± 0.011 for the comparison of AD with NAND, NND and all controls, respectively. Statistical analysis was done using the MedCalc ROC curve analysis software (version 12.2.1.0) and the result indicated that the test can distinguish AD from the controls with a $P < 0.0001$. To estimate the most reliable cut-off point for the different set of group comparisons, sensitivity (blue line) and specificity (red line) were plotted for each cut-off value (**FIG. 4D**). The graph shows the curve and the 95% confidence intervals for the AD vs all control samples (including NAND and NND groups). These cut-off values were used to estimate sensitivity, specificity and predictive value in **FIG. 5, Table 1**.

EXAMPLE 5: A β -OLIGOMER IMMUNODEPLETION REMOVES A β SEEDS IN HUMAN CEREBROSPINAL FLUID AND CONFIRMS A β -PMCA DETECTS SOLUBLE MISFOLDED A β PROTEIN IN AD CSF

[00163] Immunodepletion experiments were performed to confirm that A β -PMCA detects a seeding activity associated to soluble, misfolded A β protein present in CSF. The methodology for efficient immunodepletion of soluble, misfolded A β protein was first optimized by using synthetically prepared soluble, misfolded A β protein. Immunodepletion was performed by

incubation with dynabeads conjugated with a mixture of antibodies recognizing specifically the sequence of A β (4G8) and conformational (A11) antibodies. **FIG. 7A** is a western blot showing results of immunodepletion using synthetically prepared A β oligomers spiked into human CSF. Soluble, misfolded A β protein was efficiently removed by this immunodepletion.

[00164] **FIGs. 7A and 7B** are graphs of amyloid formation versus time as measured by Thioflavin T fluorescence, demonstrating that seeding activity in AD CSF is removed by soluble, misfolded A β protein immuno-depletion. Samples of AD CSF before or after immunodepletion with 4G8 and A11 antibodies were used to seed A β aggregation in the A β -PMCA assay. Immunodepletion was applied to 3 AD CSF. **FIG. 7B** is a graph showing the kinetics of control and immunodepleted CSF samples. **FIG. 7B** shows that for immunodepleted AD CSF, the kinetics of A β aggregation in the A β -PMCA reaction was comparable to that observed in control CSF samples, and both were significantly different from the aggregation observed with AD CSF prior to immunodepletion. **FIG. 7C** is a graph showing the kinetics of control and immunodepleted CSF samples, depleted only with the A11 conformational antibody and aggregation monitored by A β -PMCA assay. **FIG. 7C** shows similar results, obtained using AD CSF immunodepleted using the A11 conformational antibody, which specifically recognizes, misfolded A β . These results confirm that A β -PMCA detects soluble, misfolded β protein in AD CSF.

EXAMPLE 6: SOLID PHASE IMMUNO CAPTURING

[00165] **FIGs. 8A and 8B** are schematic representations of two solid phase methods used to capture soluble, misfolded A β protein from complex samples such as blood plasma. Strategy 1 employed ELISA plates pre-coated with specific antibodies bound to a solid phase on the ELISA plate. After washing the plates, the A β -PMCA reaction was carried out in the same plates. Strategy 2 used magnetic beads as the solid phase coated with specific antibodies. This approach provided concentration of the samples.

EXAMPLE 7: SPECIFICITY OF IMMUNO CAPTURING

[00166] **FIG. 9** shows **Table 2**, demonstrating the ability of specific antibodies to capture the A β oligomers. The top panel shows a schematic representation of the epitope recognition site on the A β protein of the diverse sequence antibodies used in this study. **Table 2** in **FIG. 9** demonstrates the efficiency of different sequence or conformational antibodies to capture A β oligomers. The capacity to capture oligomers was measured by spiking synthetic A β oligomers in healthy human blood plasma and detection by A β -PMCA. The symbols indicate that the detection limits using the different antibodies were: <12 fmol (+++); between 10-100 fmol (++); >1pmol (+) and not significantly higher than without capturing reagent (-).

EXAMPLE 8: DETECTION OF A β OLIGOMERS SPIKED IN HUMAN PLASMA

[00167] **FIG. 10** is a graph of amyloid formation versus time as measured by Thioflavin T fluorescence showing detection of soluble, misfolded A β protein spiked in human plasma. ELISA plates pre-coated with protein G were coated with an anti-conformational antibody (16ADV from Acumen). Thereafter, plates were incubated with human blood plasma (100 μ l) as such (control) or spiked with different concentrations of synthetic soluble, misfolded A β protein. After incubation, plates were subjected to A β -PMCA (29 min incubation and 30 s shaking) in the presence of A β 40 monomer (2 μ M) and ThT (5 μ M). Amyloid formation was measured by Thioflavin fluorescence. **FIG. 10** is representative of several experiments done with 3 different antibodies which worked similarly.

EXAMPLE 9: CAPTURING OF SOLUBLE MISFOLDED A β FROM AD PATIENT SAMPLES VS CONTROLS

[00168] **FIG. 11** is a graph showing time to reach 50% aggregation in an A β -PMCA assay in plasma samples from AD patients and controls. Blood plasma samples from patients affected by AD, non-AD neurodegenerative diseases (NAD), and healthy controls were incubated with anti-A β antibody (82E1) coated beads. A β -PMCA was carried out as described in **EXAMPLE 2**. The time needed to reach 50% aggregation was recorded in individual patients in each group. Differences were analyzed by one-way ANOVA followed by the Tukey's post-hoc test. ROC analysis of this data showed a 82% sensitivity and 100% specificity for correctly identifying AD patients from controls.

EXAMPLE 10: SONICATION AND SHAKING ARE EFFECTIVE WITH VARIOUS DETECTION METHODS

[00169] **FIG. 12** is a western blot showing the results of amplification of A β aggregation by using sonication instead of shaking as a mean to fragment aggregates. The experiment was done in the presence of distinct quantities of synthetic A β oligomers. Samples of 10 μ g/ml of seed-free monomeric A β 1-42 were incubated alone (lane 1) or with 300 (lane 2), 30 (lane 3) and 3 (lane 4) fmols of, misfolded A β . Samples were either frozen without amplification (non-amplified) or subjected to 96 PMCA cycles (amplified), each including 30 min incubation followed by 20 sec sonication. Aggregated A β was detected by western blot using anti-A β antibody after treatment of the samples with proteinase K (PK). In our experiments, it was observed that detection using thioflavin T fluorescence was not compatible with sonication, but works very well with shaking as a physical disruption method. **FIG. 12** shows that using a different detection method for the A β aggregates, in this case Western Blotting, sonication works as well as shaking.

EXAMPLE 11: PRODUCTION OF MONOMERIC A β AS PMCA SUBSTRATE

[00170] Seed-free monomeric A β was obtained by size exclusion chromatography. Briefly, an aliquot of a 1 mg/mL peptide solution prepared in dimethylsulfoxide was fractionated using a Superdex 75 column eluted at 0.5 mL/min with 10 mM sodium phosphate at pH 7.5. Peaks will be detected by UV absorbance at 220 nm. The peak corresponding to 4-10 kDa molecular weight containing monomer/dimers of A β was collected and concentration determined by amino acid analysis. Samples were stored lyophilized at -80 °C.

EXAMPLE 12: PRODUCTION AND PURIFICATION OF A β

[00171] E. coli cells harboring pET28 GroES-Ub-A β 42 plasmid were grown in Luria broth (LB) at 37 °C, and expression was induced with 0.4 mM IPTG. After 4 h, cells were harvested and lysed in a lysis buffer (20 mM Tris-Cl, pH 8.0, 10 mM NaCl, 0.1 mM PMSF, 0.1 mM EDTA and 1 mM β -mercaptoethanol) and centrifuged at 18,000 rpm for 30 min. Inclusion bodies were re-suspended in a resuspension buffer (50 mM Tris-Cl, pH 8.0, 150 mM NaCl, and 1 mM DTT) containing 6 M urea. Insoluble protein was removed by centrifugation at 18,000 rpm for 30 min. The supernatant containing GroES-Ub-A β 42 fusion protein will be collected. To cleave off A β 42 from fusion protein, the fusion protein was diluted 2-fold with resuspension buffer and treated with recombinant de-ubiquinating enzyme (Usp2cc) 1: 100 enzyme to substrate molar ratio at 37 °C for 2 h. After that, samples was loaded on a C18 column (25 mm x 250 mm, Grace Vydac, USA). A β 42 was purified with a solvent system buffer 1 (30 mM ammonium acetate, pH 10, 2% acetonitrile) and buffer 2 (70% acetonitrile) at a flow rate 10 ml/min using a 20–40% linear gradient of buffer 2 over 35 min. Purified A β 42 was lyophilized and stored at -80 °C, until use.

EXAMPLE 13: DETECTION OF α S SEEDS BY PD-PMCA

[00172] **EXAMPLE 13A:** Seeding of α S aggregation was studied by incubating a solution of seed-free α S in the presence of Thioflavin T with or without different quantities of synthetic soluble oligomeric α S protein: Control (no α S oligomer); or 1 ng/mL, 10 ng/mL, 100 ng/mL, and 1 μ g/mL of the synthetic soluble oligomeric α S protein seed. α S-PMCA general procedure: Solutions of 100 μ g/mL α S seed-free α S in PBS, pH 7.4 (200 μ L total volume) were placed in opaque 96-wells plates and incubated alone or in the presence of the indicated concentrations of synthetic α S aggregates or 40 μ L of CSF aliquots. Samples were incubated in the presence of 5 μ M Thioflavin T (ThT) and subjected to cyclic agitation (1 min at 500 rpm followed by 29 min without shaking) using an Eppendorf thermomixer, at a constant temperature of 22 °C. At various time points, ThT fluorescence was measured in the plates at 485 nm after excitation at 435 nm using a plate spectrofluorometer. **FIG. 13A** is a graph of Thioflavin T fluorescence as a function of time, showing the detection of α S seeds by PD-PMCA, using the indicated concentration of synthetic soluble oligomeric α S protein seeds. The peptide concentration, temperature and pH of

the buffer were monitored to control the extent of the lag phase and reproducibility among experiments. Aggregation of monomeric α S protein was observed in the presence of 1 ng/mL, 10 ng/mL, 100 ng/mL, and 1 μ g/mL α S of the synthetic soluble oligomeric α S protein seed.

[00173] **EXAMPLE 13B:** The time to reach 50% aggregation as a function of amounts of α S seeds added was determined using the samples in **EXAMPLE 1A**. **FIG. 13B** is a graph showing time to reach 50% aggregation plotted as a function of amounts of α S seeds added.

EXAMPLE 14: α S-PMCA DETECTS OLIGOMERIC α S IN THE CEREBROSPINAL FLUID OF PD PATIENTS

[00174] Detection of seeding activity in human CSF samples from controls and PD patients was performed by PD-PMCA. Purified seed free alpha-synuclein (100 μ g/mL) in PBS, pH 7.4 was allowed to aggregate at 37 °C with shaking at 500 rpm in the presence of CSF from human patients with confirmed PD, AD or non-neurodegenerative neurological diseases (NND). The extend of aggregation was monitored by Thioflavin fluorescence at 485 nm after excitation at 435 nm using a plate spectrofluorometer.

[00175] Aliquots of CSF were obtained from PD patients, cognitively normal individuals affected by non-degenerative neurological diseases (NND), and patients affected by Alzheimer's disease (AD). Test CSF samples were obtained from patients with the diagnosis of probable PD as defined by the DSM-IV and determined using a variety of tests, including routine medical examination, neurological evaluation, neuropsychological assessment, and magnetic resonance imaging. CSF samples were collected in polypropylene tubes following lumbar puncture at the L4/L5 or L3/L4 interspace with atraumatic needles after one night fasting. The samples were centrifuged at 3,000 g for 3 min at 4°C, aliquoted and stored at -80°C until analysis. CSF cell counts, glucose and protein concentration were determined. Albumin was measured by rate nephelometry. To evaluate the integrity of the blood brain barrier and the intrathecal IgG production, the albumin quotient (CSF albumin/serum albumin) $\times 10^3$ and the IgG index (CSF albumin/serum albumin)/(CSF IgG/serum IgG) were calculated. The study was conducted according to the provisions of the Helsinki Declaration and was approved by the Ethics Committee.

[00176] The experiments as well as the initial part of the analysis were conducted blind. **FIG. 14** is a graph of α S oligomerization versus time, measured as a function of ThT fluorescence labeling, showing the average kinetics of α S aggregation of representative samples from the PD, AD, and NND groups.

[00177] The results indicate that CSF from PD patients significantly accelerates α S aggregation as compared to control CSF ($P < 0.001$). The significance of the differences in α S aggregation kinetics in the presence of human CSF samples was analyzed by one-way ANOVA,

followed by the Tukey's multiple comparison post-test. The level of significance was set at $P < 0.05$. The differences between PD and samples from the other two groups were highly significant with $P < 0.001$ (***).

EXAMPLE 15: SPECIFICITY OF IMMUNO CAPTURING

[00178] **FIG. 15** shows **Table 3**, demonstrating the ability of different sequence or conformational antibodies to capture α S oligomers. The capacity to capture oligomers was measured by spiking synthetic α S oligomers in healthy human blood plasma and detection by α S-PMCA. The first column shows various antibodies tested and corresponding commercial sources. The second column lists the epitope recognition site on the α S protein of the diverse sequence antibodies used in this study. The third column indicates the observed ability of specific antibodies to capture the α S oligomers. The symbols indicate that the detection limits using the different antibodies were: <12 fmol (+++); between 10-100 fmol (++); >1 pmol (+) and not significantly higher than without capturing reagent (-). Alpha/beta-synuclein antibody N-19 (N-terminal epitope) and alpha-synuclein antibody C-20-R (C-terminal epitope) showed the best results; and alpha-synuclein antibody 211 (epitope: amino acids 121-125) showed very good results; alpha-synuclein antibody 204 (epitope: fragment 1-130) showed good results; and 16 ADV Mouse IgG1 (conformational epitope) showed no result.

EXAMPLE 16: SOLID PHASE IMMUNO CAPTURING

[00179] **FIGs. 16A** and **16B** are schematic representations of two solid phase methods used to capture soluble, misfolded α S protein from complex samples such as blood plasma. Strategy 1 employed ELISA plates pre-coated with specific antibodies bound to a solid phase on the ELISA plate. After washing the plates, the α S-PMCA reaction was carried out in the same plates. Strategy 2 used magnetic beads as the solid phase coated with specific antibodies. This approach provided concentration of the samples.

EXAMPLE 17: α S-PMCA FOR THE DETECTION OF α -SYNUCLEIN OLIGOMERS SPIKED IN HUMAN BLOOD PLASMA

[00180] Immunoprecipitation of α -Synuclein oligomers from human blood plasma was performed by anti- α -Synuclein antibody-coated beads (Dynabeads) and a seeding aggregation assay using α -Synuclein monomers as seeding substrate along with thioflavin-T for detection. The anti- α -Synuclein coated beads (1×10^7 beads) were incubated with human blood plasma (500 μ L) with α -Synuclein seeds (+ 0.2 μ g Seed) and without α -Synuclein seeds (- Seed). After immunoprecipitation, the beads were re-suspended in 20 μ L of reaction buffer (1X PBS), and 10 μ L of beads were added to each well of a 96-well plate. The aggregation assay was performed by adding α -Synuclein monomers (200 μ g/mL) and thioflavin-T (5 μ M). The increase in fluorescence

was monitored by a fluorimeter using an excitation of 435 nm and emission of 485 nm. **FIG. 17A** illustrates immunoprecipitation/aggregation results with N-19 antibody in blood plasmas with and without seed. **FIG. 17B** illustrates immunoprecipitation/aggregation results with 211 antibody in blood plasmas with and without seed. **FIG. 17C** illustrates immunoprecipitation/aggregation results with C-20 antibody in blood plasmas with and without seed.

EXAMPLE 18: α S-PMCA DETECTS OLIGOMERIC α S IN THE CEREBROSPINAL FLUID OF PATIENTS AFFECTED BY PD AND MULTIPLE SYSTEM ATROPHY WITH HIGH SENSITIVITY AND SPECIFICITY.

[00181] To study the efficiency of α S-PMCA for biochemical diagnosis of PD and related α -synucleinopathies, such as multiple system atrophy (MSA), tests were performed on CSF from many patients affected by these diseases as well as controls affected by other diseases. **FIGS. 18A, 18B, and 18C** show detection of seeding activity in human CSF samples from controls and patients affected by PD and MSA, respectively, using α S-PMCA. Purified seed free alpha-synuclein (100 μ g/mL) in buffer MES, pH 6.0 was allowed to aggregate at 37 °C with shaking at 500 rpm in the presence of CSF from human patients and controls. The extent of aggregation was monitored by Thioflavin T fluorescence at 485 nm after excitation at 435 nm using a plate spectrofluorometer.

[00182] Test CSF samples were obtained from patients with the diagnosis of probable PD and MSA as defined by the DSM-IV and determined using a variety of tests, including routine medical examination, neurological evaluation, neuropsychological assessment, and magnetic resonance imaging. CSF samples were collected in polypropylene tubes following lumbar puncture at the L4/L5 or L3/L4 interspace with atraumatic needles after one night fasting. The samples were centrifuged at 3,000 g for 3 min at 4°C, aliquoted and stored at -80°C until analysis. CSF cell counts, glucose and protein concentration were determined. Albumin was measured by rate nephelometry. The study was conducted according to the provisions of the Helsinki Declaration and was approved by the Ethics Committee.

[00183] The experiments as well as the initial part of the analysis were conducted blind. **FIGS. 18A, 18B, and 18C** are graphs of α S aggregation versus time, measured as a function of ThT fluorescence labeling, showing the average kinetics of α S aggregation, respectively, for controls and two representative samples from the PD and MSA groups.

[00184] The results indicate that CSF from PD patients significantly accelerates α S aggregation as compared to control CSF ($P < 0.001$). The significance of the differences in α S aggregation kinetics in the presence of human CSF samples was analyzed by one-way ANOVA, followed by the Tukey's multiple comparison post-test. The level of significance was set at

$P < 0.05$. The differences between PD and samples from the other two groups were highly significant with $P < 0.001$ (***).

[00185] The outcome of the overall set of 29 PD or MSA samples and 41 controls was that 26 of the 29 PD or MSA samples were positive, whereas 3 of the 41 control samples were positive, which corresponded to a 90% sensitivity and 93% specificity.

EXAMPLE 19: SYNTHESIS OF FULL-LENGTH 4R TAU PROTEIN AND SEEDS

[00186] **FIG. 19** is a flow chart showing the preparation and purification of recombinant full-length 4R tau protein. A gene for hTau40 was transfected into *E. coli* and incubated under standard conditions to express the hTau40. After a period of growth, the *E. coli* cells were pelleted, lysed, heat treated, and precipitated with ammonium sulfate to produce a crude product. The crude product was subjected to cation exchange chromatography, dialysis, then concentrated and the buffer exchanged. Cut-off filtration at a mass of 100kDa was employed to further purify the hTau40. The yield of the purified hTau 40 was 20 mg/L of bacterial culture. Full-length 4R Tau seeds were then prepared by incubating hTau 40 monomer with 12.5 μ M heparin in 10 mM HEPES pH 7.4, 100 mM NaCl for 3 days at 37°C using cyclic agitation (1 min shaking at 500 rpm followed by 29 min without shaking).

EXAMPLE 20A: TAU PMCA

[00187] A Tau-PMCA assay was performed on 96 well plates using 12.5 μ M Tau monomer, 1.25 μ M heparin, 5 μ M Thioflavin T, using cyclic agitation (1 min shaking at 500 rpm followed by 29 min without shaking). Seeds were added to the wells in amounts of 12.5 pmol, 1.25 pmol, 125 fmol, 12.5 fmol, and 1.25 fmol. Controls were performed without seeds. Aggregation was followed over time by ThT fluorescence using a plate spectrofluorometer (excitation: 435; emission: 485). **FIG. 20A** is a graph of aggregation in % for the various initial amounts of seeds and the control. The values in **FIG. 20A** are the mean of two replicates, with the error bars indicating standard deviation. **FIG. 20B** is a graph of T_{50} , the time to 50% aggregation as measured by ThT fluorescence versus the log of the amount of oligomeric tau seeds in fmol.

EXAMPLE 20B: TAU PMCA

[00188] For optimization of the tau-PMCA assay, full-length human Tau40 that includes four imperfect tandem microtubule binding repeats (4R). Tau oligomers were generated by incubation of full-length Tau (50 μ M) in the presence of heparin (12.5 μ M) for 3 days at 37°C. Seeds were characterized by ability to seed tau aggregation, binding to thioflavin T, western blot and electron microscopy. The preformed aggregates were used to nucleate and induce the aggregation of Tau. For the assay, seed-free monomeric tau (15 μ M) in 10 mM HEPES pH 7.4, 100 mM NaCl containing 3 μ M of heparin in the absence or the presence of different quantities of

synthetic seeds was subjected to cycles of tau-PMCA by incubating at 20°C for 29.5 min followed by shaking for 30 sec at 500 rpm. Under these conditions, Tau was only observed to aggregate in the presence of preformed seeds and the kinetic of aggregation was dependent on the amount of seeds added. Importantly, the PMCA signal was observed to be directly proportional to the amount of seeds added to the reaction. This assay corresponded to a detection threshold of 0.125 pg of oligomeric tau. This detection threshold corresponds to -2 atto-mol, based on the molecular weight of the tau monomer, or 0.15 atto-mol, based on the molecular weight of a 12-mer oligomer as a proxy for average oligomer size.

EXAMPLE 20C: TAU PMCA

[00189] A further Tau-PMCA assay was performed using full-length Tau seeds prepared by incubating Tau monomer with 12.5 μ M heparin in 10 mM HEPES pH 7.4, 100 mM NaCl for 3 days at 37°C with shaking. The assay was performed on 96 well plates using 12.5 μ M Tau monomer, 1.25 μ M heparin, and 5 μ M Thioflavin T, using cyclic agitation (1 min shaking at 500 rpm followed by 29 min without shaking). **FIG. 20C** is a graph of aggregation followed over time by ThT fluorescence using a plate spectrofluorometer (excitation: 435; emission: 485). **FIG. 20C** shows the mean and SD of two replicates. **FIG. 20D** is a graph of the relationship between the quantity of tau oligomers and the Tau-PMCA signal (time to reach 50% aggregation).

[00190] EXAMPLE 20D: TAU PMCA IS REPRODUCIBLE

[00191] A large scale experiment was conducted to evaluate the robustness and reproducibility of the tau-PMCA assay to analyze the performance at four different times (0, 14, 28 and 30 days) with or without freezing/thawing. Two different set of synthetic seeds and five different concentrations of synthetic seeds (1250, 125, 12.5, 1.25 and 0.125 pg of seeds) were employed, spiked either in buffer or control CSF. Each sample was run in triplicate. The experiment encompassed several steps, including the large-scale expression and purification of tau in quantities needed to perform all experiments, quality control of the material produced, generation and characterization of synthetic tau oligomeric seeds and the tau-PMCA experiments to investigate assay precision and reproducibility in buffer and in the biological matrix (CSF). In total the experiment employed 32 different conditions (2 different seeds x 4 time points x: 2 manners of dilution (freezing or not freezing) x 2 different matrices (buffer or CSF)). Since all conditions were tested with five different concentrations of oligomeric seeds and each was done in triplicate, the entire experiment involved 480 wells. The protein concentration, buffer and PMCA conditions were the same as **EXAMPLE 20B**. From the 32 conditions tested only one gave results that were slightly significantly different from the others, indicating high precision and reproducibility. **FIGS. 20E-20L** are a series of graphs that display the aggregation results based

on ThT fluorescence of 8 of the conditions tested, including 4 different time points (0, 7, 14 and 30 days) with samples subjected to freezing and thawing or not and in the presence of buffer or CSF, and two different seed preparations (**FIGS. 20E-20H** and **FIGS. 20I-20L**). **FIGS. 20E-20L** demonstrate that the results obtained are very similar between the triplicates, different time points, and distinct seeds. Data correspond to average $\pm$ standard error of triplicate samples. Tau substrate in the absence of seeds was not observed to aggregate under any condition within the time in which experiments were done.

[00192] To analyze the reproducibility of the assay, T_{50} values (time to reach 50% aggregation) for the experiments in the presence of 1250 pg of Tau seeds. The T_{50} values for the experiments done at different days, with one of two seed preparations A or B and using fresh or frozen seeds did not show any statistically significant difference and an average T_{50} of 71.5 ± 1.8 h was obtained. Similar non-significant differences were observed for the studies done in the presence of all the other seed concentrations or for the experiments done in buffer. **FIG. 20M** is a table of T_{50} values showing reproducibility across 16 different conditions. All values were analyzed by one way ANOVA, followed by Tukey multiple comparison test.

[00193] **EXAMPLE 20E: TAU PMCA IS SPECIFIC**

[00194] The tau PMCA assay was investigated for specificity, particularly for the ability to detect aggregates composed of other amyloidogenic proteins. $A\beta$ and α Syn oligomeric species were prepared and used to cross-seed monomeric tau. **FIG. 20N** is a graph of ThT fluorescence vs time for the tau assay seeded with 1 pm of tau, $A\beta$ 40, AB42, His α Syn, Hu α Syn, and no seeds. **FIG. 20N** shows that no significant signal was detectable in the presence of $A\beta$ or α Syn seeds and no signal was detected before about 100 h, even when the concentration of these particles was relatively high (equivalent to 2 ng of tau seeds). These $A\beta$ or α Syn seeds are very efficient in inducing aggregation in the respective $A\beta$ - or α Syn-PMCA assays described in preceding EXAMPLES. These results indicate that under the conditions and concentrations used there is no cross-seeding between other protein aggregates and that tau-PMCA is specific for detecting tau oligomers.

EXAMPLE 21A: TAU PMCA DETECTS TAU IN HUMAN CSF

[00195] Human CSF samples from AD patients (7 cases), 5 other Tauopathies (1FTD, 2PSP, 2CBD), people affected by mild cognitive impairment (MCI) and controls affected by other neurological diseases (7 samples) were analyzed by Tau-PMCA. **FIG. 21A** is a graph showing ThT fluorescence at 447h of incubation, in which most samples have reached the maximum fluorescence. Positive controls used samples of healthy CSF spiked with synthetic Tau oligomers (12.5 fmol). Negative controls correspond to samples of healthy CSF without Tau seeds. **FIG.**

21A shows that patients with AD, other tauopathies, and MCI showed Tau aggregation significantly above the negative control and consistent with the positive control. 6 of the 7 control patients with other neurological diseases were consistent with the negative control. One control patients in the other neurological disease group showed maximum fluorescence consistent with a tauopathy. This may indicate an undiagnosed tauopathy in that patient, or alternatively, inadvertent contamination.

EXAMPLE 21B: TAU PMCA DETECTS TAU IN HUMAN CSF

[00196] Human CSF samples from 11 patients affected by AD, 11 from other tauopathies (4 PSP, 1 FTD, 5 CBD and 1 CTE) and 7 controls affected from unrelated neurological disorders were examined. Positive controls were prepared by spiking CSF with 20 ng of recombinant tau oligomers. Negative controls were healthy CSF without tau seeds. **FIG. 21B** is a graph of fluorescence signals for samples from patients with AD or other tauopathies for tau-PMCA comparable to that observed in samples containing recombinant tau oligomers. Consequently, samples from patients with AD or other tauopathies were able to accelerate tau aggregation. Conversely, 6 out of 7 of the controls produced a low signal in tau-PMCA, with values equivalent to those observed in CSF without seeds. Despite the small sample size, the differences between controls and patients were statistically significant. These positive results indicate that tau-PMCA is capable of detecting tau aggregates in CSF of patients. **FIG. 21B** shows the maximum ThT fluorescence, expressed as arbitrary units. Differences were analyzed by one-way ANOVA followed by the Tukey's multiple comparison post-test. ** $P < 0.01$.

EXAMPLE 22: TAU PMCA DETECTS TAU IN HUMAN CSF

[00197] The performance of Tau-PMCA assay was examined in the presence of representative CSF samples from a control, and patients affected by AD, FTD (frontotemporal dementia), CBD (corticobasal degeneration), and PSP (progressive supranuclear palsy). The Tau-PMCA assay was performed on 96 well plates using 12.5 μM Tau monomer, 1.25 μM heparin, 5 μM Thioflavin T, using cyclic agitation (1 min shaking at 500 rpm followed by 29 min without shaking). Aggregation was followed over time by ThT fluorescence using a plate spectrofluorometer (excitation: 435; emission: 485). **FIG. 22** is a graph showing aggregation % based on ThT versus time. The various tauopathies differed in T_{50} , amplification rate, and amplification extent. For example, the T_{50} of PSP and AD samples was about the same at 150 h, while the PSP sample appeared to have a shorter lag phase, a lower amplification rate, and a lower extent of amplification compared to AD. CBD appeared to have a T_{50} of about 175 h, with a lower amplification rate, and a lower extent of amplification compared to both AD and PSP. FTD appeared to have a T_{50} of about 210 h, with a higher amplification rate, and a lower extent of

amplification compared to each of AD, PSP, and CBD. The control CSF sample had a similar T_{50} and amplification rate compared to FTD, with a far lower extent of amplification. The amplification in the control CSF sample may be due to spontaneous (seed-free) amplification, inadvertent contamination, or an undiagnosed tauopathy with somewhat similar kinetics to FTD.

DISCUSSION

[00198] The present tau-PMCA may provide various advantages. For example, embodiments of the present invention may detect the tau misfolded protein directly, by contrast with known indirect measures such as non-pathogenic biomarkers, measurement of the total pool of tau of which only a small fraction forms the synapto-toxic oligomeric aggregates, or measurement of variously phosphorylated species of tau. In various embodiments, the present invention detects the misfolded tau oligomers that seed tau misfolding and are believed to contribute to spreading the damage in the brain during the disease.

[00199] To the extent that the term “includes” or “including” is used in the specification or the claims, it is intended to be inclusive in a manner similar to the term “comprising” as that term is interpreted when employed as a transitional word in a claim. Furthermore, to the extent that the term “or” is employed (e.g., A or B) it is intended to mean “A or B or both.” When the applicants intend to indicate “only A or B but not both” then the term “only A or B but not both” will be employed. Thus, use of the term “or” herein is the inclusive, and not the exclusive use. See Bryan A. Garner, *A Dictionary of Modern Legal Usage* 624 (2d. Ed. 1995). Also, to the extent that the terms “in” or “into” are used in the specification or the claims, it is intended to additionally mean “on” or “onto.” To the extent that the term “selectively” is used in the specification or the claims, it is intended to refer to a condition of a component wherein a user of the apparatus may activate or deactivate the feature or function of the component as is necessary or desired in use of the apparatus. To the extent that the term “operatively connected” is used in the specification or the claims, it is intended to mean that the identified components are connected in a way to perform a designated function. To the extent that the term “substantially” is used in the specification or the claims, it is intended to mean that the identified components have the relation or qualities indicated with degree of error as would be acceptable in the subject industry.

[00200] As used in the specification and the claims, the singular forms “a,” “an,” and “the” include the plural unless the singular is expressly specified. For example, reference to “a compound” may include a mixture of two or more compounds, as well as a single compound.

[00201] As used herein, the term “about” in conjunction with a number is intended to include $\pm 10\%$ of the number. In other words, “about 10” may mean from 9 to 11.

[00202] As used herein, the terms “optional” and “optionally” mean that the subsequently described circumstance may or may not occur, so that the description includes instances where the circumstance occurs and instances where it does not.

[00203] In addition, where features or aspects of the disclosure are described in terms of Markush groups, those skilled in the art will recognize that the disclosure is also thereby described in terms of any individual member or subgroup of members of the Markush group. As will be understood by one skilled in the art, for any and all purposes, such as in terms of providing a written description, all ranges disclosed herein also encompass any and all possible sub-ranges and combinations of sub-ranges thereof. Any listed range can be easily recognized as sufficiently describing and enabling the same range being broken down into at least equal halves, thirds, quarters, fifths, tenths, and the like. As a non-limiting example, each range discussed herein can be readily broken down into a lower third, middle third and upper third, and the like. As will also be understood by one skilled in the art all language such as “up to,” “at least,” “greater than,” “less than,” include the number recited and refer to ranges which can be subsequently broken down into sub-ranges as discussed above. Finally, as will be understood by one skilled in the art, a range includes each individual member. For example, a group having 1-3 cells refers to groups having 1, 2, or 3 cells. Similarly, a group having 1-5 cells refers to groups having 1, 2, 3, 4, or 5 cells, and so forth. While various aspects and embodiments have been disclosed herein, other aspects and embodiments will be apparent to those skilled in the art.

[00204] As stated above, while the present application has been illustrated by the description of embodiments thereof, and while the embodiments have been described in considerable detail, it is not the intention of the applicants to restrict or in any way limit the scope of the appended claims to such detail. Additional advantages and modifications will readily appear to those skilled in the art, having the benefit of the present application. Therefore, the application, in its broader aspects, is not limited to the specific details, illustrative examples shown, or any apparatus referred to. Departures may be made from such details, examples, and apparatuses without departing from the spirit or scope of the general inventive concept.

[00205] The various aspects and embodiments disclosed herein are for purposes of illustration and are not intended to be limiting, with the true scope and spirit being indicated by the following claims.

CLAIMS

1. A method for determining a presence or absence in a sample of a first misfolded protein aggregate, the method comprising:
 - performing a first protein misfolding cyclic amplification (PMCA) procedure, the first PMCA procedure comprising:
 - forming a first incubation mixture by contacting a first portion of the sample with a first substrate protein, the first substrate protein comprising 4R tau;
 - conducting an incubation cycle two or more times under conditions effective to form a first amplified, misfolded protein aggregate, each incubation cycle comprising:
 - incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the first misfolded protein aggregate;
 - disrupting the first incubation mixture effective to form the first amplified, misfolded protein aggregate; and
 - determining the presence or absence in the sample of the first misfolded protein aggregate by analyzing the first incubation mixture for the presence or absence of the first amplified, misfolded protein aggregate,
 - the first misfolded protein aggregate comprising the first substrate protein and the first amplified, misfolded protein aggregate comprising the first substrate protein.
2. The method of claim 1, comprising determining the presence in the sample of the first misfolded protein aggregate, the first misfolded protein aggregate being misfolded 4R tau aggregate.
3. The method of claim 1, further comprising determining the presence or absence in the sample of at least a second misfolded protein aggregate.
4. The method of claim 3, further comprising performing at least a second PMCA procedure to determine the presence or absence in the sample of at least the second misfolded protein aggregate, comprising:
 - forming a second incubation mixture by contacting a second portion of the sample with a second substrate protein, the second substrate protein being subject to pathological misfolding and/or aggregation in vivo to form the second misfolded protein aggregate;
 - conducting an incubation cycle two or more times under conditions effective to form a second amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the second incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the second substrate protein in the presence of the second misfolded protein aggregate;

disrupting the second incubation mixture effective to form the second amplified, misfolded protein aggregate; and

determining the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate,

the second misfolded protein aggregate comprising the second substrate protein and the second amplified, misfolded protein aggregate comprising the second substrate protein.

5. The method of claim 4, the second substrate protein comprising one of: amyloid-beta (A β), alpha synuclein, and 3R tau.

6. The method of claim 4, the second substrate protein comprising 3R tau, the method further comprising determining a ratio of the 4R tau and the 3R tau in the sample of between about 1:99 and about 99:1.

7. The method of claim 3, comprising determining the presence in the sample of the first misfolded protein aggregate and the second misfolded protein aggregate.

8. The method of claim 3, further comprising determining the presence or absence in the sample of at least a third misfolded protein aggregate.

9. The method of claim 8, further comprising performing at least a third PMCA procedure to determine the presence or absence in the sample of at least the third misfolded protein aggregate, comprising:

forming a third incubation mixture by contacting a third portion of the sample with a third substrate protein, the third substrate protein being subject to pathological misfolding and/or aggregation in vivo to form the third misfolded protein aggregate;

conducting an incubation cycle two or more times under conditions effective to form a third amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the third incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the third substrate protein in the presence of the third misfolded protein aggregate;

disrupting the third incubation mixture effective to form the third amplified, misfolded protein aggregate; and

determining the presence or absence in the sample of the third misfolded protein aggregate by analyzing the third incubation mixture for the presence or absence of the third amplified, misfolded protein aggregate,

the third misfolded protein aggregate comprising the third substrate protein and the third amplified, misfolded protein aggregate comprising the third substrate protein.

10. The method of claim 9, the first substrate protein comprising 4R tau, the second substrate protein comprising A β , and the third substrate protein comprising alpha synuclein.

11. The method of claim 8, comprising determining the presence in the sample of the first misfolded protein aggregate, the second misfolded protein aggregate, and the third misfolded protein aggregate.

12. The method of claim 8, further comprising determining the presence or absence in the sample of at least a third misfolded protein aggregate.

13. The method of claim 12, further comprising performing at least a fourth PMCA procedure to determine the presence or absence in the sample of at least the fourth misfolded protein aggregate, comprising:

forming a fourth incubation mixture by contacting a fourth portion of the sample with a fourth substrate protein, the fourth substrate protein being subject to pathological misfolding and/or aggregation in vivo to form the fourth misfolded protein aggregate;

conducting an incubation cycle two or more times under conditions effective to form a fourth amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the fourth incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the fourth substrate protein in the presence of the fourth misfolded protein aggregate;

disrupting the fourth incubation mixture effective to form the fourth amplified, misfolded protein aggregate; and

determining the presence or absence in the sample of the fourth misfolded protein aggregate by analyzing the fourth incubation mixture for the presence or absence of the fourth amplified, misfolded protein aggregate,

the fourth misfolded protein aggregate comprising the fourth substrate protein and the fourth amplified, misfolded protein aggregate comprising the fourth substrate protein.

14. The method of claim 13, the first substrate protein comprising 4R tau, the second substrate protein comprising A β , the third substrate protein comprising alpha synuclein, and the fourth substrate protein comprising 3R tau.
15. The method of claim 1, the sample being taken from a subject, further comprising determining or diagnosing the presence or absence of a tauopathy in the subject according to the presence or absence of the first misfolded protein aggregate in the sample.
16. The method of claim 15, the method further comprising:
performing at least a second PMCA procedure to determine the presence or absence in the sample of a misfolded protein aggregate comprising a second substrate protein, comprising:
forming a second incubation mixture by contacting a second portion of the sample with a second substrate protein, the second substrate protein being subject to pathological misfolding and/or aggregation in vivo to form the second misfolded protein aggregate;
conducting an incubation cycle two or more times under conditions effective to form a second amplified, misfolded protein aggregate, each incubation cycle comprising:
incubating the second incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the second substrate protein in the presence of the second misfolded protein aggregate;
disrupting the second incubation mixture effective to form the second amplified, misfolded protein aggregate; and
determining the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate,
the second misfolded protein aggregate comprising the second substrate protein and the second amplified, misfolded protein aggregate comprising the second substrate protein.
17. The method of claim 16, the subject comprising the tauopathy, further comprising characterizing an identity of the tauopathy in the subject according to: the presence in the sample of the first misfolded protein aggregate; and the presence or absence in the sample of the second misfolded protein aggregate.
18. The method of claim 16, the second substrate protein comprising one of: amyloid-beta (A β), alpha synuclein, and 3R tau.
19. The method of claim 16, the method further comprising:

performing at least a third PMCA procedure to determine the presence or absence in the sample of a misfolded protein aggregate comprising a third substrate protein, comprising:

forming a third incubation mixture by contacting a third portion of the sample with a third substrate protein, the third substrate protein being subject to pathological misfolding and/or aggregation in vivo to form the third misfolded protein aggregate;

conducting an incubation cycle two or more times under conditions effective to form a third amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the third incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the third substrate protein in the presence of the third misfolded protein aggregate;

disrupting the third incubation mixture effective to form the third amplified, misfolded protein aggregate; and

determining the presence or absence in the sample of the third misfolded protein aggregate by analyzing the third incubation mixture for the presence or absence of the third amplified, misfolded protein aggregate,

the third misfolded protein aggregate comprising the third substrate protein and the third amplified, misfolded protein aggregate comprising the third substrate protein.

20. The method of claim 19, the method further comprising:

performing at least a fourth PMCA procedure to determine the presence or absence in the sample of a misfolded protein aggregate comprising a fourth substrate protein, comprising:

forming a fourth incubation mixture by contacting a fourth portion of the sample with a fourth substrate protein, the fourth substrate protein being subject to pathological misfolding and/or aggregation in vivo to form the fourth misfolded protein aggregate;

conducting an incubation cycle two or more times under conditions effective to form a fourth amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the fourth incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the fourth substrate protein in the presence of the fourth misfolded protein aggregate;

disrupting the fourth incubation mixture effective to form the fourth amplified, misfolded protein aggregate; and

determining the presence or absence in the sample of the fourth misfolded protein aggregate by analyzing the fourth incubation mixture for the presence or absence of the fourth amplified, misfolded protein aggregate,

the fourth misfolded protein aggregate comprising the fourth substrate protein and the fourth amplified, misfolded protein aggregate comprising the fourth substrate protein.

21. The method of claim 15, the tauopathy comprising a primary tauopathy or a secondary tauopathy.
22. The method of claim 15, the tauopathy characterized at least in part by misfolding and/or aggregation of 4R tau protein.
23. The method of claim 15, the second substrate protein comprising 3R tau, the method further comprising characterizing the tauopathy by a ratio between the first misfolded protein aggregate and the second misfolded protein aggregate, and determining the tauopathy corresponds to a known tauopathy according to a known ratio of misfolding and/or aggregation of 4R tau protein to 3R tau protein that is characteristic of the known tauopathy.
24. The method of claim 15, further comprising characterizing an identity of the tauopathy by analyzing the first amplified, misfolded protein aggregate or one or more corresponding PMCA kinetic parameters thereof for a signature of at least one of: Alzheimer's disease (AD), Parkinson's Disease (PD), Progressive Supranuclear Palsy (PSP), Frontotemporal Dementia (FTD), Corticobasal degeneration (CBD), Mild cognitive impairment (MCI), Argrophilic grain disease (AgD) Traumatic Brain Injury (TBI), Chronic Traumatic Encephalopathy (CTE), and Dementia Pugilistica (DP).
25. The method of claim 24, characterizing the identity of the tauopathy comprising:
determining the one or more corresponding PMCA kinetic parameters, comprising one or more of: lag phase, T_{50} , amplification rate, and amplification extent;
comparing the one or more corresponding PMCA kinetic parameters to one or more corresponding predetermined corresponding PMCA kinetic parameters that are characteristic of the identity of the tauopathy to determine a similarity or difference effective to characterize the identity of the tauopathy.
26. The method of claim 24, characterizing the identity of the tauopathy comprising using one or more of: an antibody selective for a conformational epitope of a tauopathy-specific misfolded tau protein aggregate; an indicator selective for the tauopathy-specific misfolded tau protein aggregate; a spectrum characteristic of the tauopathy-specific misfolded tau protein aggregate; a proteolytic resistance of the tauopathy-specific misfolded tau protein aggregate; and a stability to denaturation of the tauopathy-specific misfolded tau protein aggregate.

27. The method of claim 15, provided that the tauopathy excludes Pick's disease.
28. The method of claim 15, determining or diagnosing the presence or absence of a tauopathy in the subject comprising comparing the presence or absence of the first misfolded protein aggregate in the sample to a control sample taken from a control subject.
29. The method of claim 1, the detecting comprising detecting an amount of the first misfolded protein aggregate in the sample, the sample being taken from a subject, further comprising determining or diagnosing the presence or absence of a tauopathy in the subject by comparing the amount of the first misfolded protein aggregate in the sample to a predetermined threshold amount.
30. The method of claim 1, the sample being taken from a subject exhibiting no clinical signs of dementia according to cognitive testing, further comprising determining or diagnosing the presence or absence of a tauopathy in the subject according to the presence or absence of the first misfolded protein aggregate in the sample.
31. The method of claim 1, the sample being taken from a subject exhibiting no cortex plaques or tangles according to contrast imaging, further comprising determining or diagnosing the presence or absence of a tauopathy in the subject according to the presence or absence of the first misfolded protein aggregate in the sample.
32. The method of claim 1, the sample being taken from a subject exhibiting clinical signs of dementia according to cognitive testing, further comprising determining or diagnosing the presence or absence of a tauopathy as a contributing factor to the clinical signs of dementia in the subject according to the presence or absence of the first misfolded protein aggregate in the sample.
33. The method of claim 1, the sample being taken from a subject exhibiting no clinical signs of dementia according to cognitive testing and the subject exhibiting a predisposition to dementia according to genetic testing, further comprising determining or diagnosing the presence or absence of a tauopathy in the subject according to the presence or absence of the first misfolded protein aggregate in the sample.
34. The method of claim 1, comprising preparing the first incubation mixture characterized by at least one concentration of:
the first substrate protein of less than about 20 μ M;

heparin of less than about 75 μ M;
NaCl of less than about 190 mM; and
Thioflavin T of less than about 9.5 μ M.

35. The method of claim 1, comprising preparing the first incubation mixture comprising the first substrate protein at a concentration between about 0.001 μ M and about 2000 μ M.
36. The method of claim 1, comprising preparing the first incubation mixture characterized by heparin at a concentration between about 0.001 μ M and about 75 μ M.
37. The method of claim 1, comprising preparing the first incubation mixture comprising a buffer composition of one or more of: Tris-HCL, PBS, MES, PIPES, MOPS, BES, TES, and HEPES.
38. The method of claim 31, comprising preparing the first incubation mixture comprising the buffer composition at a total concentration of between about 1 μ M and about 1 M.
39. The method of claim 1, comprising preparing the first incubation mixture comprising a salt composition at a total concentration of between about 1 μ M and about 1 M.
40. The method of claim 39, the salt composition comprising one or more of: NaCl and KCl.
41. The method of claim 1, comprising preparing or maintaining the first incubation mixture at a pH of between about 5 and about 9.
42. The method of claim 1, comprising preparing the first incubation mixture comprising an indicator at a total concentration of between about 1 nM and about 1 mM.
43. The method of claim 1, the incubating comprising heating or maintaining the first incubation mixture at a temperature between about 5°C and about 60°C.
44. The method of claim 1:
further comprising contacting an indicator of the first misfolded protein aggregate to the first incubation mixture, the indicator of the first misfolded protein aggregate being characterized by an indicating state in the presence of the first misfolded protein aggregate and a non-indicating state in the absence of the first misfolded protein aggregate; and
wherein the determining the presence of the first misfolded protein aggregate in the sample comprises detecting the indicating state of the indicator of the first misfolded protein aggregate.

45. The method of claim 44, the indicating state of the indicator and the non-indicating state of the indicator being characterized by a difference in fluorescence, the determining the presence of the first misfolded protein aggregate in the sample comprising detecting the difference in fluorescence.
46. The method of claim 44, comprising contacting a molar excess of the indicator of the first misfolded protein aggregate to the first incubation mixture, the molar excess being greater than a total molar amount of protein monomer included in the first substrate protein and the first misfolded protein aggregate in the first incubation mixture.
47. The method of claim 44, the indicator of the first misfolded protein aggregate comprising one or more of: a thioflavin, Congo Red, m-I-Stilbene, Chrysamine G, PIB, BF-227, X-34, TZDM, FDDNP, MeO-XO4, IMPY, NIAD-4, luminescent conjugated polythiophenes, a fluorescent protein, and derivatives thereof.
48. The method of claim 1, the determining the presence of the first misfolded protein aggregate in the sample comprising determining an amount of the first misfolded protein aggregate in the sample.
49. The method of claim 47, comprising detecting the amount of the first misfolded protein aggregate in the sample at a sensitivity of at least about 70%.
50. The method of claim 47, comprising detecting the amount of the first misfolded protein aggregate in the sample at less than about 100 nmol.
51. The method of claim 47, comprising detecting the amount of the first misfolded protein aggregate in the sample in a molar ratio to the first substrate protein comprised by the sample, the molar ratio being less than about 1:100.
52. The method of claim 47, comprising determining the amount of the first misfolded protein aggregate in the sample compared to a control sample.
53. The method of claim 1, comprising detecting the first misfolded protein aggregate in the sample with a specificity of at least about 70%.
54. The method of claim 1, the detecting the first misfolded protein aggregate comprising one or more of: a Western Blot assay, a dot blot assay, an enzyme-linked immunosorbent assay (ELISA), a fluorescent protein/peptide binding assay, a thioflavin binding assay, a Congo Red

binding assay, a sedimentation assay, electron microscopy, atomic force microscopy, surface plasmon resonance, and spectroscopy.

55. The method of claim 54, the ELISA comprising a two-sided sandwich ELISA.

56. The method of claim 54, the spectroscopy comprising one or more of: quasi-light scattering spectroscopy, multispectral ultraviolet spectroscopy, confocal dual-color fluorescence correlation spectroscopy, Fourier-transform infrared spectroscopy, capillary electrophoresis with spectroscopic detection, electron spin resonance spectroscopy, nuclear magnetic resonance spectroscopy, and Fluorescence Resonance Energy Transfer (FRET) spectroscopy.

57. The method of claim 1, the detecting the first misfolded protein aggregate comprising:
contacting the first incubation mixture with a protease; and
detecting the first misfolded protein aggregate using anti-misfolded protein antibodies or antibodies specific for a misfolded tau aggregate in one or more of: a Western Blot assay, a dot blot assay, and an ELISA.

58. The method of claim 1, further comprising providing the first substrate protein in labeled form.

59. The method of claim 58, the first substrate protein in labeled form comprising one or more of: a covalently incorporated radioactive amino acid, a covalently incorporated, isotopically labeled amino acid, and a covalently incorporated fluorophore.

60. The method of claim 59, the detecting the first misfolded protein aggregate comprising detecting the first substrate protein in labeled form as incorporated into the first amplified, misfolded protein aggregate.

61. The method of claim 1, the sample comprising one or more of: amniotic fluid; bile; blood; cerebrospinal fluid; cerumen; skin; exudate; feces; gastric fluid; lymph; milk; mucus; mucosal membrane; peritoneal fluid; plasma; pleural fluid; pus; saliva; sebum; semen; sweat; synovial fluid; tears; and urine.

62. The method of claim 1, the sample derived from cells or tissue of one or more of: skin, brain, heart, liver, pancreas, lung, kidney, gastro-intestine, nerve, mucous membrane, blood cell, gland, and muscle.

63. The method of claim 1, each portion of the sample characterized by a volume from about 1 μ L to about 1000 μ L.
64. The method of claim 1, each portion of the sample comprising CSF characterized by a volume from about 10 μ L to about 80 μ L.
65. The method of claim 1, each portion of the sample comprising one of:
- plasma characterized by a volume from about 250 μ L to about 750 μ L; and
- blood characterized by a volume from about 200 μ L to about 1000 μ L.
66. The method of claim 1, further comprising obtaining the sample from a subject.
67. The method of claim 66, the subject being one of a: human, mouse, rat, dog, cat, cattle, horse, deer, elk, sheep, goat, pig, and non-human primate.
68. The method of claim 66, the subject being one or more of: at risk of a tauopathy, having the tauopathy, and under treatment for the tauopathy.
69. The method of claim 1, the sample being taken from a subject, further comprising determining a progression or homeostasis of a tauopathy in the subject by comparing the amount of the first misfolded protein aggregate in the sample to an amount of the first misfolded protein aggregate in a comparison sample taken from the subject at a different time compared to the sample.
70. The method of claim 1, the subject being treated with a tauopathy modulating therapy, further comprising:
- comparing the amount of the first misfolded protein aggregate in the sample to an amount of the first misfolded protein aggregate in a comparison sample, the sample and the comparison sample being taken from the subject at different times over a period of time under the tauopathy modulating therapy; and
- determining the subject is one of: responsive to the tauopathy modulating therapy according to a change in the first misfolded protein aggregate over the period of time, or non-responsive to the tauopathy modulating therapy according to homeostasis of the first misfolded protein aggregate over the period of time.
71. The method of claim 70, further comprising treating the subject determined to be responsive to the tauopathy modulating therapy with the tauopathy modulating therapy.

72. The method of claim 1, the sample being taken from a subject, the subject being treated with a tauopathy modulating therapy to inhibit production of the first substrate protein or to inhibit aggregation of the first misfolded protein aggregate.
73. The method of claim 1, further comprising selectively concentrating the first misfolded protein aggregate in one or more of the sample and the first incubation mixture.
74. The method of claim 73, the selectively concentrating the first misfolded protein aggregate comprising pre-treating the sample prior to forming the first incubation mixture.
75. The method of claim 70, the selectively concentrating the first misfolded protein aggregate comprising pre-treating the first incubation mixture prior to incubating the first incubation mixture.
76. The method of claim 73, the selectively concentrating the first misfolded protein aggregate comprising contacting one or more antibodies capable of binding the first misfolded protein aggregate to form a captured first misfolded protein aggregate.
77. The method of claim 76, the one or more antibodies capable of binding the first misfolded protein aggregate comprising one or more of: an antibody specific for an amino acid epitope sequence of the first misfolded protein aggregate, and an antibody specific for a conformation of the first misfolded protein aggregate.
78. The method of claim 77, the antibody specific for a conformation of the first misfolded protein aggregate being selective for a conformational epitope of a tauopathy-specific misfolded tau aggregate.
79. The method of claim 77, the one or more one or more antibodies capable of binding the first misfolded protein aggregate being coupled to a solid phase.
80. The method of claim 79, the solid phase comprising one or more of a magnetic bead and a multiwell plate.
81. The method of claim 76, the contacting the sample with the first substrate protein to form the first incubation mixture comprising contacting a molar excess of the first substrate protein to the sample comprising the captured first misfolded protein aggregate, the molar excess of the first substrate protein being greater than a total molar amount of protein monomer included in the captured first misfolded protein aggregate.

82. The method of claim 81, the incubating the first incubation mixture being effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the captured first misfolded protein aggregate to form the first amplified, misfolded protein aggregate.

83. The method of claim 1, the disrupting the first incubation mixture comprising one or more of: sonication, stirring, shaking, freezing/thawing, laser irradiation, autoclave incubation, high pressure, and homogenization.

84. The method of claim 1, the disrupting the first incubation mixture comprising cyclic agitation.

85. The method of claim 84, the cyclic agitation being conducted for one or more of: between about 50 rotations per minute (RPM) and 10,000 RPM, between about 200 RPM and about 2000 RPM, and at about 500 RPM.

86. The method of claim 1, the disrupting the first incubation mixture being conducted in each incubation cycle for one or more of: between about 5 seconds and about 10 minutes, between about 30 sec and about 1 minute, between about 45 sec and about 1 minute, and about 1 minute.

87. The method of claim 1, the incubating the first incubation mixture being independently conducted, in each incubation cycle for one or more of: between about 1 minute and about 5 hours, between about 5 minutes and about 5 hours, between about 10 minutes and about 2 hours, between about 15 minutes and about 1 hour, and between about 25 minutes and about 45 minutes.

88. The method of claim 1, each incubation cycle comprising independently incubating and disrupting the first incubation mixture for one or more of: incubating between about 1 minute and about 5 hours and disrupting between about 5 seconds and about 10 minutes; incubating between about 5 minutes and about 5 hours and disrupting between about 5 seconds and about 10 minutes; incubating between about 10 minutes and about 2 hours and disrupting between about 30 sec and about 1 minute; incubating between about 15 minutes and about 1 hour and disrupting between about 45 sec and about 1 minute; incubating between about 25 minutes and about 45 minutes and disrupting between about 45 sec and about 90 seconds; incubating for about 29 minutes and for about 1 minute; and incubating about 1 minute and disrupting about 1 minute.

89. The method of claim 1, the conducting the incubation cycle being repeated for one or more of: between about 2 times and about 1000 times, between about 5 times and about 500 times, between about 50 times and about 500 times, and between about 150 times and about 250 times.
90. The method of claim 1, the contacting the sample with the first substrate protein to form the first incubation mixture being conducted under physiological conditions.
91. The method of claim 1, comprising contacting the sample with a molar excess of the first substrate protein to form the first incubation mixture, the molar excess being greater than a total molar amount of protein monomer included in the first misfolded protein aggregate in the sample.
92. The method of claim 1, comprising:
contacting the sample with a thioflavin and a molar excess of the first substrate protein to form the first incubation mixture, the molar excess being greater than an amount of the first substrate protein included in the first misfolded protein aggregate in the sample;
conducting the incubation cycle two or more times effective to form the first amplified, misfolded protein aggregate, each incubation cycle comprising:
incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate;
shaking the first incubation mixture effective to form the first amplified, misfolded protein aggregate; and
determining the presence of the first misfolded protein aggregate in the sample by detecting a fluorescence of the thioflavin corresponding to the first misfolded protein aggregate.
93. The method of claim 1, the first substrate protein being produced by one of: chemical synthesis, recombinant production, and extraction from non-recombinant biological samples.
94. The method of claim 1,
the first misfolded protein aggregate comprising one or more of a soluble first misfolded protein aggregate and an insoluble first misfolded protein aggregate; and
the first amplified, misfolded protein aggregate comprising one or more of: a soluble portion and an insoluble portion,

95. The method of claim 94, the first misfolded protein aggregate substantially being the soluble first misfolded protein aggregate.
96. The method of claim 1, provided the sample excludes tau fibrils.
97. A method for determining a presence or absence in a subject of a tauopathy corresponding to a first misfolded protein aggregate, comprising:
- providing a sample from the subject;
 - performing at least a first protein misfolding cyclic amplification (PMCA) procedure, the first PMCA procedure comprising:
 - forming a first incubation mixture by contacting a first portion of the sample with a first substrate protein, first substrate protein the first substrate protein comprising at a tau isoform, the first substrate protein being subject to pathological misfolding and/or aggregation in vivo to form the first misfolded protein aggregate;
 - conducting an incubation cycle two or more times under conditions effective to form the first amplified, misfolded protein aggregate, each incubation cycle comprising:
 - incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the first misfolded protein aggregate;
 - disrupting the first incubation mixture effective to form the first amplified, misfolded protein aggregate; and
 - determining the presence or absence in the sample of the first misfolded protein aggregate by analyzing the first incubation mixture for the presence or absence of the first amplified, misfolded protein aggregate; and
 - determining the presence or absence of the tauopathy in the subject according the presence or absence of the first misfolded protein aggregate in the sample,
 - the first misfolded protein aggregate comprising the first substrate protein and the first amplified, misfolded protein aggregate comprising the first substrate protein. and
 - provided that the tauopathy excludes Pick's disease when the first substrate protein consists of monomeric 3R tau.
98. The method of claim 97, the method further comprising:
- performing at least a second PMCA procedure to determine the presence or absence in the sample of at least a second misfolded protein aggregate, comprising:

forming a second incubation mixture by contacting a second portion of the sample with a second substrate protein, the second substrate protein being subject to pathological misfolding and/or aggregation in vivo to form the second misfolded protein aggregate;

conducting an incubation cycle two or more times under conditions effective to form a second amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the second incubation mixture effective to cause misfolding and/or aggregation of the second substrate protein in the presence of the second misfolded protein aggregate;

disrupting the second incubation mixture effective to form the second amplified, misfolded protein aggregate; and

determining the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate,

the second misfolded protein aggregate comprising the second substrate protein and the second amplified, misfolded protein aggregate comprising the second substrate protein.

99. The method of claim 98, the second substrate protein being distinct from the first substrate protein and the second substrate protein comprising one of: amyloid-beta (A β), alpha synuclein, 3R tau, and 4R tau.

100. The method of claim 98, the first substrate protein comprising 4R tau.

101. The method of claim 98, the method further comprising:

performing at least a third PMCA procedure to determine the presence or absence in the sample of at least a third misfolded protein aggregate, comprising:

forming a third incubation mixture by contacting a third portion of the sample with a third substrate protein, the third substrate protein being subject to pathological misfolding and/or aggregation in vivo to form the third misfolded protein aggregate;

conducting an incubation cycle two or more times under conditions effective to form a third amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the third incubation mixture effective to cause misfolding and/or aggregation of the third substrate protein in the presence of the third misfolded protein aggregate;

disrupting the third incubation mixture effective to form the third amplified, misfolded protein aggregate; and

determining the presence or absence in the sample of the third misfolded protein aggregate by analyzing the third incubation mixture for the presence or absence of the third amplified, misfolded protein aggregate,

the third misfolded protein aggregate comprising the third substrate protein and the third amplified, misfolded protein aggregate comprising the third substrate protein.

102. The method of claim 101, the tauopathy being present in the subject, further comprising characterizing an identity of the tauopathy in the subject according to: the presence in the sample of the first misfolded protein aggregate; and the presence or absence in the sample of the second misfolded protein aggregate.

103. The method of claim 97, provided that the tauopathy is not primarily characterized by misfolding and/or aggregation of 3R tau protein.

104. The method of claim 97, comprising:

contacting the sample with a thioflavin and a molar excess of the first substrate protein to form the first incubation mixture, the molar excess being greater than an amount of protein monomer included in the first misfolded protein aggregate in the sample;

conducting the incubation cycle two or more times effective to form the first amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least the portion of the first substrate protein in the presence of the first misfolded protein aggregate;

shaking the first incubation mixture effective to form the first amplified, misfolded protein aggregate; and

determining the presence or absence of the first misfolded protein aggregate in the sample by detecting a fluorescence of the thioflavin corresponding to the first misfolded protein aggregate.

105. A method for determining a presence or absence in a sample of a first misfolded protein aggregate using capturing, comprising:

capturing the first misfolded protein aggregate from the sample to form a captured first misfolded protein aggregate;

performing at least a first protein misfolding cyclic amplification (PMCA) procedure, the first PMCA procedure comprising:

forming a first incubation mixture by contacting the captured first misfolded protein aggregate with a molar excess of a first substrate protein, the first substrate protein being

subject to pathological misfolding and/or aggregation in vivo to form the first misfolded protein aggregate;

the molar excess being greater than an amount of protein monomer included in the captured first misfolded protein aggregate;

conducting an incubation cycle two or more times effective to form the first amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the captured first misfolded protein aggregate;

disrupting the first incubation mixture effective to form the first amplified, misfolded protein aggregate; and

determining the presence or absence of the first misfolded protein aggregate in the sample by detecting at least a portion of the first amplified, misfolded protein aggregate, the first misfolded protein aggregate comprising the first substrate protein and the first amplified, misfolded protein aggregate comprising the first substrate protein.

106. The method of claim 105, the capturing the first misfolded protein aggregate from the sample to form a captured first misfolded protein aggregate being conducted using one or more antibodies specific for the first misfolded protein aggregate, comprising one or more of: an antibody specific for an amino acid epitope sequence of the first misfolded protein aggregate and an antibody specific for a conformation of the first misfolded protein aggregate.

107. The method of claim 106, the antibody specific for a conformation of the first misfolded protein aggregate being selective for a conformational epitope of a tauopathy-specific first misfolded protein aggregate.

108. The method of claim 106, the antibody specific for the conformation of the first misfolded protein aggregate corresponding to one of: Alzheimer's disease (AD), Parkinson's Disease (PD), Progressive Supranuclear Palsy (PSP), FrontoTemporal Dementia (FTD), Corticobasal degeneration (CBD), Mild cognitive impairment (MCI), Argrophilic grain disease (AgD) Traumatic Brain Injury (TBI), Chronic Traumatic Encephalopathy (CTE), and Dementia Pugilistica (DP).

109. The method of claim 105, the one or more antibodies specific for the first misfolded protein aggregate being coupled to a solid phase.

110. The method of claim 109, the solid phase comprising one or more of a magnetic bead and a multiwell plate.
111. The method of claim 109, the contacting the sample with the first substrate protein to form the first incubation mixture comprising contacting a molar excess of the first substrate protein to the sample, the molar excess of the first substrate protein being greater than a total molar amount of protein monomer included in the captured first misfolded protein aggregate.
112. The method of claim 109, the incubating the first incubation mixture being effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the captured first misfolded protein aggregate to form the first amplified, misfolded protein aggregate.
113. The method of claim 105, the first substrate protein comprising 4R tau protein.
114. A method for determining a presence or absence of a tauopathy in a subject, the tauopathy comprising Alzheimer's disease (AD), the method comprising:
- providing the subject;
 - obtaining a sample from the subject, the sample comprising one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate;
 - performing at least a first protein misfolding cyclic amplification (PMCA) procedure, the first PMCA procedure comprising:
 - forming a first incubation mixture by contacting the sample with a first substrate protein, the first substrate protein comprising 4R tau;
 - conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate, each incubation cycle comprising:
 - incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the first misfolded protein aggregate;
 - disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate; and
 - determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate; and
 - determining the presence or absence in the subject of AD according to the presence or absence of the first misfolded protein aggregate in the sample.

115. The method of claim 114, determining the presence or absence in the subject of AD further comprising distinguishing AD from one or more additional tauopathies by determining a signature of AD tau protein aggregate comprising one or more of: one or more AD-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a conformational epitope of AD tau protein aggregate; an indicator selective for AD tau protein aggregate; and a spectrum characteristic of AD tau protein aggregate.

116. A method for determining a presence or absence of a tauopathy in a subject, the tauopathy comprising Parkinson's Disease (PD), the method comprising:

- providing the subject;
- obtaining a sample from the subject, the sample comprising one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate;
- performing at least a first PMCA procedure comprising:
 - forming a first incubation mixture by contacting the sample with a first substrate protein, the first substrate protein comprising 4R tau;
 - conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate, each incubation cycle comprising:
 - incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the first misfolded protein aggregate;
 - disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate; and
 - determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate; and
 - determining the presence or absence in the subject of PD according to the presence or absence of the first misfolded protein aggregate in the sample.

117. The method of claim 116, determining the presence or absence in the subject of PD further comprising distinguishing PD from one or more additional tauopathies by determining a signature of PD tau protein aggregate comprising one or more of: one or more PD-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a conformational epitope of PD tau protein

aggregate; an indicator selective for PD tau protein aggregate; and a spectrum characteristic of PD tau protein aggregate.

118. A method for determining a presence or absence of a tauopathy in a subject, the tauopathy comprising Progressive Supranuclear Palsy (PSP), the method comprising:

- providing the subject;
- obtaining a sample from the subject, the sample comprising one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate;
- performing at least a first PMCA procedure comprising:
 - forming a first incubation mixture by contacting the sample with a first substrate protein, the first substrate protein comprising 4R tau;
 - conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate, each incubation cycle comprising:
 - incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the first misfolded protein aggregate;
 - disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate; and
 - determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate; and
 - determining the presence or absence in the subject of PSP according to the presence or absence of the first misfolded protein aggregate in the sample.

119. The method of claim 118, determining the presence or absence in the subject of PSP further comprising distinguishing PSP from one or more additional tauopathies by determining a signature of PSP tau protein aggregate comprising one or more of: one or more PSP-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a conformational epitope of PSP tau protein aggregate; an indicator selective for PSP tau protein aggregate; and a spectrum characteristic of PSP tau protein aggregate.

120. A method for determining a presence or absence of a tauopathy in a subject, the tauopathy comprising FrontoTemporal Dementia (FTD), the method comprising:

- providing the subject;

obtaining a sample from the subject, the sample comprising one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate;

performing at least a first PMCA procedure comprising:

- forming a first incubation mixture by contacting the sample with a first substrate protein, the first substrate protein comprising 4R tau;
- conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate, each incubation cycle comprising:
 - incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the first misfolded protein aggregate;
 - disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate; and
 - determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate; and
- determining the presence or absence in the subject of FTD according to the presence or absence of the first misfolded protein aggregate in the sample.

121. The method of claim 120, determining the presence or absence in the subject of FTD further comprising distinguishing FTD from one or more additional tauopathies by determining a signature of FTD tau protein aggregate comprising one or more of: one or more FTD-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a conformational epitope of FTD tau protein aggregate; an indicator selective for FTD tau protein aggregate; and a spectrum characteristic of FTD tau protein aggregate.

122. A method for determining a presence or absence of a tauopathy in a subject, the tauopathy comprising Corticobasal degeneration (CBD), the method comprising:

- providing the subject;
- obtaining a sample from the subject, the sample comprising one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate;
- performing at least a first protein misfolding cyclic amplification (PMCA) procedure, the first PMCA procedure comprising:
 - forming a first incubation mixture by contacting the sample with a first substrate protein, the first substrate protein comprising 4R tau;

conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the first incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the first substrate protein in the presence of the first misfolded protein aggregate;

disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate; and

determining the presence or absence in the sample of the first misfolded protein aggregate by detecting in the first incubation mixture the presence or absence of the first amplified, misfolded protein aggregate; and

determining the presence or absence in the subject of CBD according to the presence or absence of the first misfolded protein aggregate in the sample.

123. The method of claim 122, determining the presence or absence in the subject of CBD further comprising distinguishing CBD from one or more additional tauopathies by determining a signature of CBD tau protein aggregate comprising one or more of: one or more CBD-specific corresponding PMCA kinetic parameters of: lag phase, T_{50} , amplification rate, and amplification extent; an assay using an antibody selective for a conformational epitope of CBD tau protein aggregate; an indicator selective for CBD tau protein aggregate; and a spectrum characteristic of CBD tau protein aggregate.

124. The method of any one of claims 114, 116, 118, 120, and 120, further comprising:

performing at least a second PMCA procedure to determine the presence or absence in the sample of at least a second misfolded protein aggregate, comprising:

forming a second incubation mixture by contacting a second portion of the sample with a second substrate protein, the second substrate protein being subject to pathological misfolding and/or aggregation;

conducting an incubation cycle two or more times under conditions effective to form a second amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the second incubation mixture effective to cause misfolding and/or aggregation of at least a portion of the second substrate protein in the presence of the second misfolded protein aggregate;

disrupting the second incubation mixture effective to form the second amplified, misfolded protein aggregate; and

determining the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate; and

the tauopathy being present in the subject, further comprising characterizing an identity of the tauopathy in the subject according to: the presence in the sample of the first misfolded protein aggregate; and the presence or absence in the sample of the second misfolded protein aggregate.

the second misfolded protein aggregate comprising the second substrate protein, the second amplified, misfolded protein aggregate comprising the second substrate protein, and

the second substrate protein being distinct from the first substrate protein and the second substrate protein comprising one of: amyloid-beta (A β), alpha synuclein, and 3R tau.

125. The method of claim 124, determining the presence or absence in the subject of each tauopathy further comprising distinguishing each tauopathy from one or more additional tauopathies by analyzing for at least one signature of one or more misfolded aggregates each corresponding to one of A β , alpha synuclein, and 3R tau, each signature corresponding to one or more of: an assay using an antibody selective for a conformational epitope of any of the one or more misfolded aggregates; an assay using an antibody selective for a conformational epitope of any of the one or more misfolded aggregates; one or more PMCA kinetic parameters of the one or more misfolded aggregates, comprising one or more of: lag phase, T₅₀, amplification rate, and amplification extent; an indicator selective for any of the one or more misfolded aggregates; and a spectrum characteristic of any of the one or more misfolded aggregates.

126. A kit for determining a presence or absence in a sample of a first misfolded protein aggregate, comprising:

a first substrate protein comprising 4R tau;

an indicator of the first misfolded protein aggregate, the first misfolded protein aggregate comprising the first substrate protein and the first misfolded protein aggregate corresponding to a tauopathy;

a buffer;

heparin; and

a salt;

instructions, the instructions directing a user to:

obtain the sample;

perform at least a first PMCA procedure, comprising:

forming a first incubation mixture by contacting a first portion of the sample with the first substrate protein, the indicator of the first misfolded protein aggregate, the buffer, the heparin, and the salt, the first incubation mixture being formed with a concentration of one or more of: the first substrate protein of less than about 20 μ M; the heparin of less than about 75 μ M; the salt as NaCl of less than about 190 mM; and the indicator of the first misfolded protein aggregate as Thioflavin T of less than about 9.5 μ M;

conducting an incubation cycle two or more times effective to form a first amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the first incubation mixture effective to cause misfolding and/or aggregation of the first substrate protein in the presence of the first misfolded protein aggregate;

disrupting the incubation mixture effective to form the first amplified, misfolded protein aggregate; and

determine the presence or absence in the sample of the first misfolded protein aggregate by analyzing the first incubation mixture for the presence or absence of the first amplified, misfolded protein aggregate according to the indicator of the first misfolded protein aggregate.

127. The kit of claim 126, the instructions directing the user to obtain the sample from a subject, the sample comprising one or more of: a bio-fluid, a biomaterial, a homogenized tissue, and a cell lysate, and the instructions directing the user to determine or diagnose a tauopathy in the subject according to the presence or absence in the sample of the first misfolded protein aggregate.

128. The kit of claim 126:

further comprising a second substrate protein and an indicator of a second misfolded protein aggregate, the second misfolded protein aggregate comprising the second substrate protein, the second substrate protein being distinct from the first substrate protein and the second substrate protein comprising one of: amyloid-beta ($A\beta$), alpha synuclein, 3R tau, and 4R tau;

the instructions further directing the user to:

perform at least a second PMCA procedure, comprising:

forming a second incubation mixture by contacting a second portion of the sample with the second substrate protein and the indicator of the second misfolded protein aggregate;

conducting an incubation cycle two or more times effective to form a second amplified, misfolded protein aggregate, each incubation cycle comprising:

incubating the second incubation mixture effective to cause misfolding and/or aggregation of the second substrate protein in the presence of the second misfolded protein aggregate;

disrupting the incubation mixture effective to form the second amplified, misfolded protein aggregate;

determine the presence or absence in the sample of the second misfolded protein aggregate by analyzing the second incubation mixture for the presence or absence of the second amplified, misfolded protein aggregate,

characterize the sample for an identity of a tauopathy according to: the presence in the sample of the first misfolded protein aggregate; and the presence or absence in the sample of the second misfolded protein aggregate.

129. The kit of claim 126, further comprising a PMCA apparatus, the PMCA apparatus comprising one or more of: a multiwall microtitre plate; a microfluidic plate; a shaking apparatus; a spectrometer; and an incubator.

130. The kit of claim 126, the antibody specific for the conformation of the first misfolded protein aggregate corresponding to one of: Alzheimer's disease (AD), Parkinson's Disease (PD), Progressive Supranuclear Palsy (PSP), Frontotemporal Dementia (FTD), Corticobasal degeneration (CBD), Mild cognitive impairment (MCI), Argrophilic grain disease (AgD) Traumatic Brain Injury (TBI), Chronic Traumatic Encephalopathy (CTE), and Dementia Pugilistica (DP), the instructions comprising determining, according to a binding assay using the antibody specific for the conformation of the first misfolded protein aggregate, the presence or absence in the subject of one of AD, PD, PSP, FTD, CBD, MCI, AgD, TBI, CTE, and DP.