

2- (1H-INDOLYLSULFANYL) -ARYL AMINE DERIVATIVES FOR USE IN THE TREATMENT OF AFFECTIVE DISORDERS, PAIN, ADHD AND STRESS URINARY INCONTINENCE

FIELD OF THE INVENTION

The present invention relates to compounds, which are serotonin reuptake inhibitors and preferably also norepinephrine reuptake inhibitors and/or dopamine reuptake inhibitors and the medical use of such compounds, e.g., in the treatment of affective disorders, pain disorders, attention deficit hyperactivity disorder (ADHD) and stress urinary incontinence.

BACKGROUND OF THE INVENTION

10 All currently available antidepressants can be classified in 3 classes:

- 1) monoamine oxidase inhibitors (MAOIs),
- 2) biogenic amine neurotransmitter [serotonin (5-HT), norepinephrine (NE) and dopamine (DA)] transporter reuptake blockers, and
- 3) modulators, especially blockers of one or more of the 5-HT and/or NE receptors.

15

Since depression is associated with a relative deficiency of the biogenic amines, the use of 5-HT and/or NE-receptor blockers (i.e. 5-HT and or NE-antagonist's) have not proven very successful in the treatment of depression and anxiety and the preferred and currently most efficient treatments are based on the enhancement of 5-HT and/or NE neurotransmission by blocking their reuptake back from the synaptic cleft (Slattery, D.A. et al., "The evolution of antidepressant mechanisms", *fundamental and Clinical pharmacology*, **2004**, 18, 1-21; Schloss, P. et al, "new insights into the mechanism of antidepressant therapy", *Pharmacology and therapeutics*, **2004**, 102, 47-60).

20

25 Selective serotonin reuptake inhibitors (hereinafter referred to as SSRIs) have become first choice therapeutics in the treatment of depression, certain forms of anxiety and social phobias, because they generally are effective, well tolerated and have a favourable safety profile compared to the classic tricyclic antidepressants. Drugs claimed to be SSRIs are for example fluoxetine, sertraline and paroxetine.

30

However, clinical studies on depression indicate that non-response to the known SSRIs is substantial, up to 30%. Another, often neglected, factor in antidepressant treatment is the fact that there is generally a delay in therapeutic effect of the SSRIs. Sometimes symptoms

even worsen during the first weeks of treatment. Furthermore, sexual dysfunction is generally a side effect common to SSRIs. Accordingly, there is a desire for the development of compounds capable of improving the treatment of depression and other serotonin related diseases.

5

A newer strategy has been the development of dual re-uptake inhibitors, e.g., the combined effect of 5-HT reuptake inhibition and NE (norepinephrine is also named noradrenaline, NA) reuptake inhibition on depression is explored in clinical studies of compounds such as Duloxetine (Wong, "Duloxetine (LY-248686): an inhibitor of serotonin and noradrenaline uptake and an antidepressant drug candidate", *Expert Opinion on Investigational Drugs*, 10 **1998**, 7, 10, 1691-1699) and Venlafaxine (Khan-A et al, 30 "Venlafaxine in depressed outpatients", *Psychopharmacology Bulletin*, **1991**, 27, 141-144). Compounds having such dual effect are also named SNRIs, "serotonin and noradrenaline reuptake inhibitors", or NSRIs, "noradrenaline and serotonin reuptake inhibitors".

15

Since treatment with the selective NE reuptake inhibitor reboxetine has been shown to stimulate 5-HT neurons and mediate the release of 5-HT in the brain (Svensson, T. et al, *J. Neural. Transmission*, **2004**, 111, 127) there might be a synergistic advantage using SNRI's in the treatment of depression or anxiety.

20

The use of SNRI's have been shown in clinical studies to have a beneficial effect on pain (e.g. Fibromyalgia syndrome, overall pain, back pain, shoulder pain, headache, pain while awake and during daily activities) and especially pain associated with depression (Berk, M. *Expert Rev. Neurotherapeutics* **2003**, 3, 47-451; Fishbain, D.A., et al. "Evidence-based data from animal and human experimental studies on pain relief with antidepressants: A structured review" *Pain Medicine* 2000 1:310-316).

25

SNRI's have also been shown in clinical studies to have a beneficial effect in attention deficit hyperactivity disorder (ADHD) (N. M. Mukaddes; Venlafaxine in attention deficit hyperactivity disorder, *European Neuropsychopharmacology*, Volume 12, Supplement 3, 30 October 2002, Page 421).

Furthermore, SNRI's have been shown to be effective for the treatment of stress urinary incontinence (Dmochowski R.R. et al. "Duloxetine versus placebo for the treatment of North American women with stress urinary incontinence", *Journal of Urology* 2003, 170:4, 1259-1263.)

5

Naranjo, C. *et al.* "The role of the brain reward system in depression" *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* **2001**, 25, 781– 823 discloses clinical and preclinical findings of links between lack of extra cellular dopamine in the mesocorticolimbic system and anhedonia, which is one of the main symptoms of depression.

10

Furthermore, Axford L. et al. describe the development of triple 5-HT, NE and DA re-uptake inhibitors for treatment of depression. (2003, *Bioorganic & Medical Chemistry Letters*, 13, 3277-3280: "Bicyclo[2.2.1.]heptanes as novel triple re-uptake inhibitors for the treatment of depression"). Wellbutrin (bupropion) which has DA re-uptake activity in vitro and in vivo, show antidepressant efficacy. Other combination studies have indicated that addition of some affinity at the DA uptake site may have some clinical benefit (Nelson, J. C. J. *Clin. Psychiatry* **1998**, 59, 65; Masand, P. S. et al. *Depression Anxiety* **1998**, 7, 89; Bodkin, J. A et al. *J. Clin. Psychiatry* **1997**, 58, 137).

15


20 The combination of an SSRI and a norepinephrine and dopamine reuptake inhibitor, has been shown to have better efficacy in SSRI-non-responders (Lam R. W. *et al.* "Citalopram and Bupropion-SR: Combining Versus Switching in Patients With Treatment-Resistant Depression." *J. Clin. Psychiatry* **2004**, 65, 337-340).

25

There is clinical evidence suggesting that the combination of an SSRI and a norepinephrine and dopamine reuptake inhibitor induces less sexual dysfunction, than treatment with SSRI's alone (Kennedy S. H. *et al.* "Combining Bupropion SR With Venlafaxine, Paroxetine, or Duloxetine: A Preliminary Report on Pharmacokinetic, Therapeutic, and Sexual Dysfunction Effects" *J. Clin. Psychiatry* **2002**, 63, 181-186).

30


Diphenyl sulphides of Formula II and variations thereof have been disclosed as serotonin re-uptake inhibitors and have been suggested for use in treatment of depression, cf. e.g. WO03029232(A1).



(II)

- 5 The above-mentioned reference does not disclose compounds comprising an indole group like the indolyl-sulfanyl arylamines of the present invention.

Diphenyl sulphides of Formula (IIA) and variations thereof have been disclosed as serotonin re-uptake inhibitors and have been suggested for use in treatment of depression, cf. e.g. US
 10 5095039, US 4056632, EP 396827 A1 and WO 9312080. EP 402097 describes halogen substituted diphenylsulfides claimed to be selective serotonin inhibitors for treatment of depression. Likewise WO 9717325 disclose derivatives of N,N-dimethyl-
 2-(arylthio)benzylamine claimed to be selective serotonin transport inhibitors and suggest their use as antidepressants. J. Jilek et al., 1989, Collect. Czech Chem. Commun., 54,
 15 3294-3338 also discloses various derivatives of diphenyl sulphides, "phenyl-thio-benzylamines" as antidepressants. Furthermore, diphenyl sulphides are also disclosed in US 3803143 and claimed useful as antidepressant.




(IIA)

20

K. Sindelar et al., "Collection of Czechoslovak Chemical Communications, (1991), 56(2), 449-58, by K. Sindelar et al" disclose compounds of Formula (IIB) in which one of the rings is substituted with a thiophene ring with test for selectivity as 5-HT re-uptake inhibitor and NA re-uptake inhibitor, respectively, for use as antidepressants.

5




The present invention provides 2-(1H-indolylsulfanyl)-aryl amine derivatives of formula I which are serotonin reuptake inhibitors. A particular aspect of the invention provides
 5 compounds possessing the combined effect of serotonin reuptake inhibition and norepinephrine reuptake inhibition. Another particular aspect of the invention provides compounds possessing the combined effect of serotonin reuptake inhibition and dopamine reuptake inhibition. Furthermore, some of the compounds are also triple 5-HT, NE and DA re-uptake inhibitors.

10

SUMMARY OF THE INVENTION

One object of the invention is the provision of compounds, which are serotonin reuptake inhibitors. Another object of the invention is the provision of compounds, which are both serotonin reuptake inhibitors and noradrenaline reuptake inhibitors. Yet another object of
 15 the invention is the provision of compounds, which are both serotonin reuptake inhibitors and dopamine reuptake inhibitors. Yet another object of the invention is the provision of compounds, which are serotonin reuptake inhibitors, noradrenaline reuptake inhibitors and dopamine reuptake inhibitors.

20 The compounds of the invention are substituted indole derivatives of the general formula I as the free base or salts thereof



wherein Z, the dotted line and R¹-R¹⁰ are as defined below.

The invention provides a compound according to the above for use as a medicament.

The invention provides a pharmaceutical composition comprising a compound according to
5 the above and at least one pharmaceutically acceptable carrier or diluent.

The invention provides the use of a compound according to the above for the preparation of
a medicament for the treatment of affective disorders, pain disorders, ADHD and stress
urinary incontinence.

10

The invention furthermore concerns the use of a compound according to the above in a
method of treatment of affective disorders, pain disorders, ADHD and stress urinary
incontinence.

15

DEFINITION OF SUBSTITUENTS

The term heteroatom refers to a nitrogen, oxygen or sulphur atom.

Halo means halogen. Halogen means fluoro, chloro, bromo or iodo.

20

The expression "C₁₋₆-alk(en/yn)yl" means a C₁₋₆-alkyl, a C₂₋₆-alkenyl or a C₂₋₆-alkynyl
group. The expression "C₁₋₆-alk(en)yl" means a C₁₋₆-alkyl or a C₂₋₆-alkenyl group.

The term "C₁₋₆-alkyl" refers to a branched or unbranched alkyl group having from one to six
carbon atoms inclusive, including but not limited to methyl, ethyl, 1-propyl, 2-propyl,
25 1-butyl, 2-butyl, 2-methyl-2-propyl and 2-methyl-1-propyl.

The term "C₂₋₆-alkenyl" designates such groups having from two to six carbon atoms,
including one double bond, including but not limited to ethenyl, propenyl, and butenyl.
Such C₂₋₆-alkenyl group may be branched or unbranched.

The term "C₂₋₆-alkynyl" designates such groups having from two to six carbon atoms,
30 including one triple bond, including but not limited to ethynyl, propynyl and butynyl. Such
C₂₋₆-alkynyl group may be branched or unbranched.

The expression "C₃₋₈-cycloalk(en)yl" means a C₃₋₈-cycloalkyl or a C₃₋₈-cycloalkenyl group. The term "C₃₋₈-cycloalkyl" designates a monocyclic or bicyclic carbocycle having three to eight C-atoms, including but not limited to cyclopropyl, cyclopentyl, and cyclohexyl. The term "C₃₋₈-cycloalkenyl" designates a monocyclic or bicyclic carbocycle having three to
5 eight C-atoms and one double bond, including but not limited to cyclopropenyl, cyclopentenyl and cyclohexenyl.

In the expression "C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl", "C₁₋₆-alk(en/yn)ylamino", "di-(C₁₋₆-alk(en/yn)yl)amino", "C₁₋₆-alk(en/yn)ylcarbonyl",
10 "C₁₋₆-alk(en/yn)ylaminocarbonyl", "di-(C₁₋₆-alk(en/yn)yl)aminocarbonyl", "C₁₋₆-alk(en/yn)ylloxy", "C₁₋₆-alk(en/yn)ylsulfanyl", "halo-C₁₋₆-alk(en/yn)yl", "halo-C₁₋₆-alk(en/yn)ylsulfonyl", "halo-C₁₋₆-alk(en/yn)ylsulfanyl", "C₁₋₆-alk(en/yn)yl-O-CO-" and "C₁₋₆-alk(en/yn)ylsulfonyl" the terms "C₃₋₈-cycloalk(en)yl", "C₁₋₆-alk(en/yn)yl", "C₁₋₆-alk(en)yl" and "halo" are as defined above.


15

DESCRIPTION OF THE INVENTION

The present invention relates to 2-(1*H*-indolylsulfanyl)-aryl amine-derivatives which are serotonin reuptake inhibitors, some of the compounds are also noradrenaline reuptake
20 inhibitors. Furthermore, some compounds of the invention are serotonin reuptake inhibitors and dopamine reuptake inhibitors. Furthermore, some compounds of the invention are serotonin reuptake inhibitors, noradrenaline reuptake inhibitors and dopamine reuptake inhibitors. Accordingly, the compounds of the invention are considered to be useful in the treatment of affective disorders such as for example depression and anxiety.

25

The present invention relates to a compound represented by the general formula I as the free base or salts thereof



wherein

the dotted line represents an optional bond;

5 and

each of R^1 - R^4 is independently selected from the group consisting of hydrogen, halogen, cyano, nitro, C_{1-6} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl, C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, amino, C_{1-6} -alk(en/yn)ylamino, di-(C_{1-6} -alk(en/yn)yl)amino, C_{1-6} -alk(en/yn)ylcarbonyl, aminocarbonyl, C_{1-6} -alk(en/yn)ylaminocarbonyl, di-(C_{1-6} -alk(en)yl)aminocarbonyl,

10 hydroxy, C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)ylsulfonyl, halo- C_{1-6} -alk(en/yn)yl, halo- C_{1-6} -alk(en/yn)ylsulfonyl, halo- C_{1-6} -alk(en/yn)ylsulfonyl and C_{1-6} -alk(en/yn)ylsulfonyl;
and

R^5 is selected from the group consisting of hydrogen, halogen, cyano, C_{1-6} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl, C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, amino, C_{1-6} -alk(en/yn)ylamino, di-(C_{1-6} -alk(en/yn)yl)amino, C_{1-6} -alk(en/yn)ylcarbonyl, aminocarbonyl,

15 C_{1-6} -alk(en/yn)ylaminocarbonyl, di-(C_{1-6} -alk(en)yl)aminocarbonyl, hydroxy,

C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)ylsulfonyl, halo- C_{1-6} -alk(en/yn)yl, halo-

C_{1-6} -alk(en/yn)ylsulfonyl, halo- C_{1-6} -alk(en/yn)ylsulfonyl and C_{1-6} -alk(en/yn)ylsulfonyl;

and

20 each of R^6 - R^9 is independently selected from the group consisting of hydrogen, halogen, cyano, nitro, C_{1-6} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl, C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, amino, C_{1-6} -alk(en/yn)ylamino, di-(C_{1-6} -alk(en/yn)yl)amino, C_{1-6} -alk(en/yn)ylcarbonyl, aminocarbonyl, C_{1-6} -alk(en/yn)ylaminocarbonyl, di-(C_{1-6} -alk(en)yl)aminocarbonyl,

hydroxy, C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)ylsulfonyl, halo- C_{1-6} -alk(en/yn)yl, halo-

25 C_{1-6} -alk(en/yn)ylsulfonyl, halo- C_{1-6} -alk(en/yn)ylsulfonyl, C_{1-6} -alk(en/yn)ylsulfonyl and C_{1-6} -alk(en/yn)yl-O-CO-;

and

R¹⁰ is selected from the group consisting of hydrogen, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl;
and

Z is selected from the group consisting of N, CH and C, with the proviso that the dotted line
5 represents a bond when Z is C and with the proviso that the dotted line does not represent a bond when Z is N or CH.

In one embodiment of the compound of formula I, the dotted line ---- represents a bond. In another embodiment of the compound of formula I the dotted line ---- does not represent a
10 bond.

In a further embodiment of the compound of formula I, the dotted line ---- does not represent a bond and Z represents a nitrogen atom. In a further embodiment of the compound of formula I, the dotted line ---- does not represent a bond and Z represents CH.

15 In a further embodiment of the compound of formula I, the dotted line ---- represents a bond and Z represents a carbon atom.

In a further embodiment of the compound of formula I each of R¹⁻⁴ is independently selected from the group consisting of nitro, amino, C₁₋₆-alk(en/yn)ylamino,
20 di-(C₁₋₆-alk(en/yn)yl)amino, C₁₋₆-alk(en/yn)ylcarbonyl, aminocarbonyl, C₁₋₆-alk(en/yn)ylaminocarbonyl, di-(C₁₋₆-alk(en/yn)yl)aminocarbonyl, hydroxy, C₁₋₆-alk(en/yn)ylloxy, C₁₋₆-alk(en/yn)ylsulfanyl, halo-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)ylsulfonyl, halo-C₁₋₆-alk(en/yn)ylsulfanyl and C₁₋₆-alk(en/yn)ylsulfonyl; in another embodiment each of R¹⁻⁴ is independently selected from the group consisting of
25 hydrogen, halogen, cyano, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl. Typically, each of R¹⁻⁴ is independently selected from the group consisting of hydrogen, halogen and C₁₋₆-alk(en/yn)yl. To further illustrate without limiting the invention, an embodiment concerns such compounds wherein at least one of R¹⁻⁴ is hydrogen; in a further embodiment at least one of R¹⁻⁴ is halogen such as fluoro; in a further
30 embodiment at least one of R¹⁻⁴ is C₁₋₆-alk(en/yn)yl such as methyl.

In a further embodiment of the compound of formula I, R¹ is hydrogen.

In a further embodiment of the compound of formula I, R² is selected from the group consisting of hydrogen, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl. Typically, R² is selected from the group consisting of hydrogen and C₁₋₆-alk(en/yn)yl. To further illustrate without limiting the invention an embodiment of R² is
5 hydrogen; another embodiment of R² is C₁₋₆-alk(en/yn)yl typically C₁₋₆-alkyl such as methyl.

In a further embodiment of the compound of formula I, R³ is hydrogen.

10 In a further embodiment of the compound of formula I, R⁴ is selected from the group consisting of hydrogen, halogen and cyano. Typically, R⁴ is selected from the group consisting of hydrogen and halogen. To further illustrate without limiting the invention an embodiment of R⁴ is hydrogen; another embodiment of R⁴ is halogen such as fluoro.

15 In a further embodiment of the compound of formula I R⁵ is selected from the group consisting of halogen, cyano, amino, C₁₋₆-alk(en/yn)ylamino, di-(C₁₋₆-alk(en/yn)yl)amino, C₁₋₆-alk(en/yn)ylcarbonyl, aminocarbonyl, C₁₋₆-alk(en/yn)ylaminocarbonyl, di-(C₁₋₆-alk(en)yl)aminocarbonyl, hydroxy, C₁₋₆-alk(en/yn)ylloxy, C₁₋₆-alk(en/yn)ylsulfanyl, halo-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)ylsulfonyl, halo-C₁₋₆-alk(en/yn)ylsulfanyl and
20 C₁₋₆-alk(en/yn)ylsulfonyl; in another embodiment R⁵ is selected from the group consisting of hydrogen, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl. Typically, R⁵ is selected from the group consisting of hydrogen and C₁₋₆-alk(en/yn)yl. To further illustrate without limiting the invention an embodiment of R⁵ is hydrogen; another embodiment of R⁵ is C₁₋₆-alk(en/yn)yl typically C₁₋₆-alkyl such as
25 methyl.

In a further embodiment of the compound of formula I, R⁵ is attached to position 2 of the indole moiety. In a further embodiment of the compound of formula I, R⁵ is attached to position 3 of the indole moiety.

30

In a further embodiment of the compound of formula I each of R⁶⁻⁹ is independently selected from the group consisting of amino, C₁₋₆-alk(en/yn)ylamino, di-(C₁₋₆-alk(en/yn)yl)amino, C₁₋₆-alk(en/yn)ylcarbonyl, aminocarbonyl,

- C₁₋₆-alk(en/yn)ylaminocarbonyl, di-(C₁₋₆-alk(en/yn)yl)aminocarbonyl, hydroxy, C₁₋₆-alk(en/yn)ylsulfanyl, halo-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)ylsulfonyl, halo-C₁₋₆-alk(en/yn)ylsulfanyl and C₁₋₆-alk(en/yn)ylsulfonyl; in another embodiment each of R⁶⁻⁹ is independently selected from the group consisting of hydrogen, halogen, cyano, nitro,
- 5 C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)ylxy and C₁₋₆-alk(en/yn)yl-O-CO-. Typically, each of R⁶⁻⁹ is independently selected from the group consisting of hydrogen, halogen, nitro, C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)ylxy and C₁₋₆-alk(en/yn)yl-O-CO-. To further illustrate without limiting the invention, an embodiment concerns such compounds wherein at least one of R⁶⁻⁹ is
- 10 hydrogen; in a further embodiment at least one of R⁶⁻⁹ is halogen such as chloro or fluoro; in a further embodiment at least one of R⁶⁻⁹ is nitro; in a further embodiment at least one of R⁶⁻⁹ is C₁₋₆-alk(en/yn)yl typically C₁₋₆-alkyl such as methyl or ethyl; in a further embodiment at least one of R⁶⁻⁹ is C₁₋₆-alk(en/yn)ylxy typically C₁₋₆-alkylxy such as methoxy; in a further embodiment at least one of R⁶⁻⁹ is C₁₋₆-alk(en/yn)yl-O-CO- typically
- 15 C₁₋₆-alkyl-O-CO- such as CH₃-O-CO-.

- In a further embodiment of the compound of formula I, R⁶ is selected from the group consisting of hydrogen, halogen, cyano, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl and C₁₋₆-alk(en/yn)ylxy. Typically, R⁶ is selected
- 20 from the group consisting of hydrogen, halogen, C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)ylxy. To further illustrate without limiting the invention an embodiment of R⁶ is hydrogen, another embodiment of R⁶ is halogen such as chloro or fluoro; another embodiment of R⁶ is C₁₋₆-alk(en/yn)yl typically C₁₋₆-alkyl such as methyl; another embodiment of R⁶ is C₁₋₆-alk(en/yn)ylxy typically C₁₋₆-alkylxy such as methoxy.

25

- In a further embodiment of the compound of formula I, R⁷ is selected from the group consisting of hydrogen, halogen, cyano, nitro, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)ylxy and C₁₋₆-alk(en/yn)yl-O-CO-. Typically, R⁷ is selected from the group consisting of hydrogen, halogen, nitro,
- 30 C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)ylxy and C₁₋₆-alk(en/yn)yl-O-CO-. To further illustrate without limiting the invention an embodiment of R⁷ is hydrogen, another embodiment of R⁷ is halogen such as chloro or fluoro; another embodiment of R⁷ is nitro; another embodiment of R⁷ is C₁₋₆-alk(en/yn)yl typically C₁₋₆-alkyl such as methyl, another embodiment of R⁷ is

C₁₋₆-alk(en/yn)yl oxy typically C₁₋₆-alkyloxy such as methoxy; another embodiment of R⁷ is C₁₋₆-alk(en/yn)yl-O-CO- typically C₁₋₆-alkyl-O-CO- such as Me-O-CO-.

- In a further embodiment of the compound of formula I, R⁸ is selected from the group
- 5 consisting of hydrogen, halogen, cyano, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl and C₁₋₆-alk(en/yn)yl oxy. Typically, R⁸ is selected from the group consisting of hydrogen, halogen, C₁₋₆-alk(en/yn)yl and C₁₋₆-alk(en/yn)yl oxy. To further illustrate without limiting the invention an embodiment of R⁸ is hydrogen; another embodiment of R⁸ is halogen such as chloro or fluoro; another embodiment of R⁸ is
- 10 C₁₋₆-alk(en/yn)yl typically C₁₋₆-alkyl such as methyl; another embodiment of R⁸ is C₁₋₆-alk(en/yn)yl oxy typically C₁₋₆-alkyloxy such as methoxy.

- In a further embodiment of the compound of formula I, R⁹ is selected from the group consisting of hydrogen, halogen, cyano, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl,
- 15 C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yl oxy and C₁₋₆-alk(en/yn)yl-O-CO-. Typically, R⁹ is selected from the group consisting of hydrogen, halogen, C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yl oxy, C₁₋₆-alk(en/yn)yl-O-CO-. To further illustrate without limiting the invention an embodiment of R⁹ is hydrogen; another embodiment of R⁹ is halogen such as chloro or fluoro; another embodiment of R⁹ is C₁₋₆-alk(en/yn)yl typically C₁₋₆-alkyl such as
- 20 methyl or ethyl; another embodiment of R⁹ is C₁₋₆-alk(en/yn)yl oxy typically C₁₋₆-alkyloxy such as methoxy; another embodiment of R⁹ is C₁₋₆-alk(en/yn)yl-O-CO- typically C₁₋₆-alkyl-O-CO- such as Me-O-CO-.

- In a further embodiment of the compound of formula I R¹⁰ is selected from the group
- 25 consisting of hydrogen and C₁₋₆-alk(en/yn)yl. To further illustrate without limiting the invention an embodiment of R¹⁰ is hydrogen; another embodiment of R¹⁰ is C₁₋₆-alk(en/yn)yl typically C₁₋₆-alkyl such as methyl.

- To further illustrate without limiting the invention independent embodiments concerns such
- 30 compounds wherein

- halogen typically is fluoro or chloro.
- C₁₋₆-alk(en/yn)yl typically is C₁₋₆-alkyl, such as methyl.
- C₃₋₈-cycloalk(en)yl typically is C₃₋₈-cycloalkyl.

- C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl typically is C₃₋₈-cycloalkyl-C₁₋₆-alkyl.
- C₁₋₆-alk(en/yn)ylamino typically is C₁₋₆-alkylamino.
- di-(C₁₋₆-alk(en/yn)yl)amino typically is di-(C₁₋₆-alkyl)amino.
- C₁₋₆-alk(en/yn)ylcarbonyl typically is C₁₋₆-alkylcarbonyl.
- 5 • C₁₋₆-alk(en/yn)ylaminocarbonyl typically is C₁₋₆-alkylaminocarbonyl.
- di-(C₁₋₆-alk(en)yl)aminocarbonyl typically is di-(C₁₋₆-alkyl)aminocarbonyl.
- C₁₋₆-alk(en/yn)ylloxy typically is C₁₋₆-alkoxy, such as methoxy.
- C₁₋₆-alk(en/yn)ylsulfanyl typically is C₁₋₆-alkylsulfanyl, such as methylsulfanyl.
- halo-C₁₋₆-alk(en/yn)yl typically is halo-C₁₋₆-alkyl, such as trifluoromethyl.
- 10 • halo-C₁₋₆-alk(en/yn)ylsulfonyl typically is trifluoromethylsulfonyl.
- halo-C₁₋₆-alk(en/yn)ylsulfanyl typically is trifluoromethylsulfanyl
- C₁₋₆-alk(en/yn)ylsulfonyl typically is C₁₋₆-alkylsulfonyl.

In a further embodiment, the invention concerns such compounds wherein 0, 1, 2 or 3 of

15 R¹⁻¹⁰ are different from hydrogen, typically 1 or 2 of R¹⁻¹⁰ are different from hydrogen. In a further embodiment 0 or 1 of R¹⁻⁴ is different from hydrogen and 0, 1 or 2 of R⁶⁻⁹ is/are different from hydrogen. In a further embodiment, none of R¹⁻¹⁰ is different from hydrogen. In a further embodiment, one of R¹⁻¹⁰ is different from hydrogen. A further embodiment concerns such compounds wherein one of R¹⁻⁴ is different from hydrogen and none of R⁶⁻¹⁰

20 is hydrogen; in a further embodiment one of R⁶⁻¹⁰ is different from hydrogen and none of R¹⁻⁴ is hydrogen. In a further embodiment two of R¹⁻¹⁰ are different from hydrogen. To further illustrate without limiting the invention an embodiment concerns such compounds wherein zero of R¹⁻⁴ are different from hydrogen and two of R⁶⁻⁹ are different from hydrogen; in another embodiment one of R¹⁻⁴ is different from hydrogen and one of R⁶⁻⁹ is


25 different from hydrogen. In a further embodiment three of R¹⁻¹⁰ are different from hydrogen. To further illustrate without limiting the invention an embodiment concerns such compounds wherein one of R¹⁻⁴ is different from hydrogen and two of R⁶⁻⁹ are different from hydrogen. In each embodiment, as mentioned the remaining substituents are hydrogen.

30 To further illustrate without limiting the invention, an embodiment concerns such compounds wherein R¹ is different from hydrogen;

another embodiment concerns such compounds wherein R^2 is different from hydrogen;
 another embodiment concerns such compounds wherein R^3 is different from hydrogen;
 another embodiment concerns such compounds wherein R^4 is different from hydrogen;
 another embodiment concerns such compounds wherein R^5 is different from hydrogen;
 5 another embodiment concerns such compounds wherein R^6 is different from hydrogen;
 another embodiment concerns such compounds wherein R^7 is different from hydrogen;
 another embodiment concerns such compounds wherein R^8 is different from hydrogen;
 another embodiment concerns such compounds wherein R^9 is different from hydrogen;
 another embodiment concerns such compounds wherein R^{10} is different from hydrogen.


10

In a further embodiment of the compound of formula I, R^5 is attached to position 2 of the indole. Such compounds are embraced by the below formula IA:



wherein Z, the dotted line and R^1 - R^{10} are as defined under formula I. Any one of the
 15 embodiments related to formula I is also an embodiment of formula IA.

In a further embodiment of the compound of formula I, R^5 is attached to position 3 of the indole. Such compounds are embraced by the below formula IB:



20

wherein Z, the dotted line and R^1 - R^{10} are as defined under formula I. Any one of the
 embodiments related to formula I is also an embodiment of formula IB.

In a further embodiment of the compound of formula IB, each of R^{1-4} is independently selected from the group consisting of hydrogen and C_{1-6} -alk(en/yn)yl.


- 5 In a further embodiment of the compound of formula IB, R^5 is C_{1-6} -alk(en/yn)yl

In a further embodiment of the compound of formula IB, each of R^{6-9} are hydrogen.

In a further embodiment of the compound of formula IB, R^{10} is hydrogen.

10

In a further embodiment of the compound of formula I the dotted line does not represent a line and Z is N. Such compounds are embraced by the below formula IC:



- 15 wherein R^1 - R^{10} are as defined under formula I. Any one of the embodiments related to formula I is also an embodiment of formula IC.

In a further embodiment of the compound of formula IC, each of R^{1-4} is independently selected from the group consisting of hydrogen, halogen and C_{1-6} -alk(en/yn)yl. Typically,

- 20 R^1 is hydrogen; R^2 is hydrogen or C_{1-6} -alk(en/yn)yl; R^3 is hydrogen and R^4 is hydrogen or halogen.

In a further embodiment of the compound of formula IC, R^5 is hydrogen or C_{1-6} -alk(en/yn)yl.

25


In a further embodiment of the compound of formula IC, each of R^{6-9} is independently selected from the group consisting of hydrogen, halogen, C_{1-6} -alk(en/yn)yl,

- C₁₋₆-alk(en/yn)yloxy, C₁₋₆-alk(en/yn)yl-O-CO- and NO₂. Typically, R⁶ is hydrogen, halogen, C₁₋₆-alk(en/yn)yl or C₁₋₆-alk(en/yn)yloxy; R⁷ is hydrogen, halogen, C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yloxy or NO₂; R⁸ is hydrogen, halogen, C₁₋₆-alk(en/yn)yl or C₁₋₆-alk(en/yn)yloxy; and R⁹ is hydrogen, halogen, C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yloxy or C₁₋₆-alk(en/yn)yl-O-CO-.

In a further embodiment of the compound of formula IC, R¹⁰ is hydrogen or C₁₋₆-alk(en/yn)yl.

10

In a further embodiment of the compound of formula I the dotted line represents a line and Z is C. Such compounds are embraced by the below formula ID:




- wherein R¹-R¹⁰ are as defined under formula I. Any one of the embodiments related to formula I is also an embodiment of formula ID.

15

In a further embodiment of the compound of formula ID, each of R¹⁻¹⁰ are hydrogen atoms

In a further embodiment of the compound of formula I the dotted line does not represent a line and Z is CH. Such compounds are embraced by the below formula IE:



wherein R^1 - R^{10} are as defined under formula I. Any one of the embodiments related to
5 formula I is also an embodiment of formula IE.

In a further embodiment of the compound of formula IE, each of R^{1-4} are hydrogen.

In a further embodiment of the compound of formula IE, R^5 is hydrogen or
10 C_{1-6} -alk(en/yn)yl.

In a further embodiment of the compound of formula IE, each of R^{6-9} is hydrogen, halogen,
 C_{1-6} -alk(en/yn)yl or C_{1-6} -alk(en/yn)yl oxy. Typically, R^6 is hydrogen, halogen or
 C_{1-6} -alk(en/yn)yl; R^7 is hydrogen, halogen, C_{1-6} -alk(en/yn)yl or C_{1-6} -alk(en/yn)yl oxy; R^8 is
15 hydrogen or halogen; R^9 is hydrogen, halogen, C_{1-6} -alk(en/yn)yl or C_{1-6} -alk(en/yn)yl oxy.

In a further embodiment of the compound of formula IE, R^{10} is hydrogen or
 C_{1-6} -alk(en/yn)yl.

20 In a further embodiment of the compound of formula I said compound is selected from the following list of compounds

Compound no.	Name
1	3-(2-Piperazin-1-yl-phenylsulfanyl)-1H-indole
2	4-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
3	3-(2-Piperazin-1-yl-phenylsulfanyl)-1H-indole-7-carboxylic acid methyl ester

4	5-Nitro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
5	6-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
6	6-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
7	6-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
8	7-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
9	3-[2-(1,2,3,6-Tetrahydro-pyridin-4-yl)-phenylsulfanyl]-1H-indole
10	6-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
11	4-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
12	5-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
13	4-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
14	6-Fluoro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
15	4-Fluoro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
16	7-Fluoro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
17	5-Methoxy-4-methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
18	5-Methoxy-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
19	7-Methoxy-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
20	4-Methoxy-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
21	4-Chloro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
22	7-Chloro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
23	1-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
24	3-Methyl-2-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
25	2-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
26	5-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
27	4-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
28	7-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
29	7-Ethyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
30	5-Methoxy-4-methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
31	5-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
32	7-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
33	5-Fluoro-2-methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
34	5-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole

35	4-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
36	7-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
37	1-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
38	3-Methyl-2-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
39	6-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
40	2-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
41	5-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
42	7-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
44	3-(2-Piperidin-4-yl-phenylsulfanyl)-1H-indole
45	2-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
46	1-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
47	5-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
48	7-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
49	6-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
50	5-Methoxy-4-methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
51	4-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
52	7-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
53	4-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
54	4-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
55	6-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
56	5-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
57	5-Methoxy-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
58	5-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
59	7-Methoxy-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
60	7-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
61	7-Ethyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
62	3-Methyl-2-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
63	6-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
64	5-Fluoro-2-methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole

as the free base or a salt thereof. Each of these compounds is considered a specific embodiment and may be subjected to individual claims.

The present invention comprises the free base and salts of the compounds of the invention, typically, pharmaceutically acceptable salts. The salts of the invention include acid addition salts, metal salts, ammonium and alkylated ammonium salts.

The salts of the invention are preferably acid addition salts. The acid addition salts of the invention are preferably pharmaceutically acceptable salts of the compounds of the invention formed with non-toxic acids. Acid addition salts include salts of inorganic acids as well as organic acids. Examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, sulfamic, nitric acids and the like. Examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, itaconic, lactic, methanesulfonic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methane sulfonic, ethanesulfonic, tartaric, ascorbic, pamoic, bismethylene salicylic, ethanedisulfonic, gluconic, citraconic, aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids, theophylline acetic acids, as well as the 8-halothephyllines, for example 8-bromothephylline and the like. Further examples of pharmaceutical acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in J. Pharm. Sci. 1977,66,2, which is incorporated herein by reference.

Also intended as acid addition salts are the hydrates, which the present compounds, are able to form.

Examples of metal salts include lithium, sodium, potassium, magnesium salts and the like.

Examples of ammonium and alkylated ammonium salts include ammonium, methyl-, dimethyl-, trimethyl-, ethyl-, hydroxyethyl-, diethyl-, n-butyl-, sec-butyl-, tert-butyl-, tetramethylammonium salts and the like.

Further, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

5

The compounds of the present invention may have one or more asymmetric centre and it is intended that any optical isomers (i.e. enantiomers or diastereomers), as separated, pure or partially purified optical isomers and any mixtures thereof including racemic mixtures, i.e. a mixture of stereoisomeres, are included within the scope of the invention

10

Racemic forms can be resolved into the optical antipodes by known methods, for example, by separation of diastereomeric salts thereof with an optically active acid, and liberating the optically active amine compound by treatment with a base. Another method for resolving racemates into the optical antipodes is based upon chromatography on an optically active matrix. Racemic compounds of the present invention can also be resolved into their optical antipodes, e.g. by fractional crystallization. The compounds of the present invention may also be resolved by the formation of diastereomeric derivatives. Additional methods for the resolution of optical isomers, known to those skilled in the art, may be used. Such methods include those discussed by J. Jaques, A. Collet and S. Wilen in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981). Optically active compounds can also be prepared from optically active starting materials, or by stereoselective synthesis.

15

20

Furthermore, when a double bond or a fully or partially saturated ring system is present in the molecule geometric isomers may be formed. It is intended that any geometric isomers, as separated, pure or partially purified geometric isomers or mixtures thereof are included within the scope of the invention. Likewise, molecules having a bond with restricted rotation may form geometric isomers. These are also intended to be included within the scope of the present invention.

25

30

Furthermore, some of the compounds of the present invention may exist in different tautomeric forms and it is intended that any tautomeric forms that the compounds are able to form are included within the scope of the present invention.

The invention also encompasses prodrugs of the present compounds, which on administration undergo chemical conversion by metabolic processes before becoming pharmacologically active substances. In general, such prodrugs will be functional derivatives of the compounds of the general formula I, IA, IB, IC, ID or IE, which are readily convertible in vivo into the required compound of the formula I, IA, IB, IC, ID or IE. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

10 The invention also encompasses active metabolites of the present compounds.

Some compounds according to the invention inhibit the serotonin transporter and are thus serotonin reuptake inhibitors. Typically, the compounds have an in vitro uptake inhibition (IC₅₀) of 5 μ M or less, typically of 1 μ M or less, preferably less than 500nM or less than 100 nM or less than 50 nM, preferably as measured by the method described in Example 3 - Measurements of "[³H]-5-HT uptake into rat cortical synaptosomes".

Some compounds according to the invention inhibit the norepinephrine transporter and are thus norepinephrine reuptake inhibitors. The compounds typically have an in vitro uptake inhibition (IC₅₀) of 5 μ M or less, typically of 1 μ M or less, preferably less than 500nM, less than 100 nM or less than 50 nM, as measured by the method described in Example 3 - Measurements of "[³H]noradrenaline uptake into rat cortical synaptosomes".

Some compounds according to the invention inhibit the dopamine transporter and are thus dopamine reuptake inhibitors. Typically, such compounds have an in vitro uptake inhibition (IC₅₀) of 5 μ M or less, typically of 1 μ M or less, preferably less than 500nM, less than 100 nM or less than 50 nM, preferably as measured by the method described in Example 3 - "Measurements of [³H]dopamine uptake into rat cortical synaptosomes".

30 As already mentioned, the compounds according to the invention are serotonin reuptake inhibitors and they are thus considered to be applicable in the treatment of one or more of the following disorders: affective disorders, pain disorders, ADHD and stress urinary incontinence.

An embodiment concerns compounds of the invention having dual action being serotonin reuptake inhibitors and dopamine reuptake inhibitors at the same time. Typically, compounds according to the invention having dual action have an in vitro uptake inhibition for the serotonin transporter which is at least 5, preferably at least 10 or even more preferred at least 20 or 30 times higher than the in vitro uptake inhibition for the dopamine transporter as measured by the methods described in Example 3 – “Measurements of [³H]-5-HT uptake into rat cortical synaptosomes” and “Measurements of [³H]dopamine uptake into rat cortical synaptosomes”.

10

An embodiment concerns compounds of the invention having dual action being serotonin reuptake inhibitors and norepinephrine reuptake inhibitors at the same time. Typically, compounds according to the invention having dual action have an in vitro uptake inhibition for the serotonin transporter which is at least 5, preferably at least 10 or even more preferred at least 20 or 30 times higher than the in vitro uptake inhibition for the norepinephrine transporter as measured by the methods described in Example 3 – “Measurements of [³H]-5-HT uptake into rat cortical synaptosomes” and “Measurements of [³H]noradrenaline uptake into rat cortical synaptosomes”.

20 A further embodiment concerns compounds of the invention having triple action and thus being serotonin reuptake inhibitors, norepinephrine reuptake inhibitors and dopamine reuptake inhibitors.

In a further aspect the invention provides a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for use as a medicament.

25

An embodiment of the invention provides a pharmaceutical composition comprising a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof and at least one pharmaceutically acceptable carrier or diluent. The composition may comprise any one of the embodiments of formula I, IA, IB, IC, ID or IE described above.

30

A further embodiment of the invention relates to the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical

composition for the treatment of a disease or disorder wherein a serotonin reuptake inhibitor is beneficial. Such pharmaceutical composition may comprise any one of the embodiments of formula I, IA, IB, IC, ID or IE described above.

- 5 A further embodiment of the invention relates to the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of a disease or disorder, wherein a combined serotonin and norepinephrine reuptake inhibitor is beneficial. Such pharmaceutical composition may comprise any one of the embodiments of formula I, IA, IB, IC, ID or IE described above.

10

A further embodiment of the invention relates to the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of a disease or disorder, wherein a combined serotonin and dopamine reuptake inhibitor is beneficial. Such pharmaceutical composition may comprise
15 any one of the embodiments of formula I, IA, IB, IC, ID or IE described above.

- A further embodiment of the present invention relates to the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of a disease or disorder, wherein a combined
20 serotonin, norepinephrine and dopamine reuptake inhibitor is beneficial. Such pharmaceutical composition may comprise any one of the embodiments of formula I, IA, IB, IC, ID or IE described above.

- A further embodiment of the invention relates to the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of affective disorders, pain disorders, ADHD and stress
25 urinary incontinence.

- In a further embodiment the present invention relates to the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of affective disorders. To further illustrate without limiting
30 the invention, the affective disorder to be treated is selected from the group consisting of depressive disorders and anxiety disorders.

A further embodiment concerns the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of depressive disorders. Typically, the depressive disorder to be treated is selected
5 from the group consisting of major depressive disorder, postnatal depression, dysthymia and depression associated with bipolar disorder, alzheimers, psychosis or parkinsons. To further illustrate without limiting the invention, an embodiment of the invention concerns the treatment of major depressive disorder; another embodiment concerns the treatment of postnatal depression; another embodiment concerns the treatment of dysthymia; another
10 embodiment concerns the treatment of depression associated with bipolar disorder, alzheimers, psychosis or parkinsons. To further illustrate without limiting the invention, an embodiment of the invention concerns the treatment of depression associated with bipolar disorder; another embodiment concerns the treatment of depression associated with alzheimers; another embodiment concerns the treatment of depression associated with
15 psychosis; another embodiment concerns the treatment of depression associated with parkinsons.

A further embodiment concerns the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical composition for the
20 treatment of anxiety disorders. Typically, the anxiety disorders to be treated are selected from the group consisting of general anxiety disorder, social anxiety disorder, post traumatic stress disorder, obsessive compulsive disorder, panic disorder, panic attacks, specific phobias, social phobia and agoraphobia. To further illustrate without limiting the invention, an embodiment of the invention concerns the treatment of general anxiety disorder; another
25 embodiment concerns the treatment of social anxiety disorder; another embodiment concerns the treatment of post traumatic stress disorder; another embodiment concerns the treatment of obsessive compulsive disorder; another embodiment concerns the treatment of panic disorder; another embodiment concerns the treatment of panic attacks; another embodiment concerns the treatment of specific phobias; another embodiment concerns the
30 treatment of social phobia; another embodiment concerns the treatment of agoraphobia.

In a further embodiment the present invention relates to the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical

composition for the treatment of pain disorders. To further illustrate without limiting the invention, the pain disorder to be treated is selected from the group consisting of fibromyalgia syndrome (FMS), overall pain, back pain, shoulder pain, headache as well as pain while awake and during daily activities. To further illustrate without limiting the invention, an embodiment of the invention concerns the treatment of fibromyalgia syndrome; another embodiment concerns the treatment of overall pain; another embodiment concerns the treatment of back pain; another embodiment concerns the treatment of shoulder pain; another embodiment concerns the treatment of headache; another embodiment concerns the treatment of pain while awake and during daily activities.

10



In a further embodiment the present invention relates to the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of attention deficit hyperactivity disorder.

15 In a further embodiment the present invention relates to the use of a compound of formula I, IA, IB, IC, ID or IE as the free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of stress urinary incontinence.

The term "treatment" as used herein in connection with a disease or disorders includes also prevention as the case may be.

20

In a further aspect, the present invention relates to a method of preparing a compound of formula I, comprising deprotection or cleavage from a polymer support of a compound of formula VA or VB



25

wherein Z, the dotted line and R^1 - R^{10} are as previously described, and R' is a carbamate or a benzyl-derived protective group.

PHARMACEUTICAL COMPOSITIONS

The present invention also relates to a pharmaceutical composition. The compounds of the invention as the free base or the salt thereof may be administered alone or in combination
5 with pharmaceutically acceptable carriers or diluents, in either single or multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19 Edition, Gennaro, Ed., Mack
10 Publishing Co., Easton, PA, 1995.

The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as the oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including
15 subcutaneous, intramuscular, intrathecal, intravenous and intradermal) route, the oral route being preferred. It will be appreciated that the preferred route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient chosen.

20 The pharmaceutical compositions formed by combining the compound of the invention and the pharmaceutical acceptable carriers are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

25 The compounds of this invention are generally utilized as the free substance (base) or as a pharmaceutically acceptable salt thereof. One example is an acid addition salt of a compound having the utility of a free base. When a compound of the invention contains a free base such salts are prepared in a conventional manner by treating a solution or suspension of a free base of the invention with a chemical equivalent of a pharmaceutically
30 acceptable acid. Representative examples are mentioned above.

Pharmaceutical compositions for oral administration may be solid or liquid. Solid dosage forms for oral administration include e.g. capsules, tablets, dragees, pills, lozenges,

powders, granules and tablette e.g. placed in a hard gelatine capsule in powder or pellet form or e.g. in the form of a troche or lozenge. Where appropriate, pharmaceutical compositions for oral administration may be prepared with coatings such as enteric coatings or they can be formulated so as to provide controlled release of the active ingredient such as

5 sustained or prolonged release according to methods well known in the art. Liquid dosage forms for oral administration include e.g. solutions, emulsions, suspensions, syrups and elixirs.

Formulations of the present invention suitable for oral administration may be presented as

10 discrete units such as capsules or tablets, each containing a predetermined amount of the active ingredient, and which may include a suitable excipient. Furthermore, the orally available formulations may be in the form of a powder or granules, a solution or suspension in an aqueous or non-aqueous liquid, or an oil-in-water or water-in-oil liquid emulsion.

15 Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents. Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, gelatine, agar, pectin, acacia, magnesium stearate, stearic acid, lower alkyl ethers of cellulose, corn starch, potato starch, gums and the like. Examples of liquid carriers are syrup, peanut oil, olive oil, phospho lipids, fatty acids, fatty acid amines,

20 polyoxyethylene and water.

The carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax.

25 Any adjuvants or additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are compatible with the active ingredients.

The amount of solid carrier may vary but will usually be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft

30 gelatine capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

Tablets may be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine.

Pharmaceutical compositions for parenteral administration include sterile aqueous and
5 nonaqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Depot injectable formulations are also contemplated as being within the scope of the present invention.

10 For parenteral administration, solutions of the compound of the invention in sterile aqueous solution, aqueous propylene glycol, aqueous vitamin E or sesame or peanut oil may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal
15 administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

Solutions for injections may be prepared by dissolving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution
20 to the desired volume, sterilising the solution and filling it in suitable ampoules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

Other suitable administration forms include suppositories, sprays, ointments, cremes, gels,
25 inhalants, dermal patches, implants, etc.

A typical oral dosage is in the range of from about 0.001 to about 100 mg/kg body weight per day, preferably from about 0.01 to about 50 mg/kg body weight per day, and more preferred from about 0.05 to about 10 mg/kg body weight per day administered in one or
30 more dosages such as 1 to 3 dosages. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

The formulations may conveniently be presented in unit dosage form by methods known to those skilled in the art. A typical unit dosage form for oral administration one or more times per day such as 1 to 3 times per day may contain from 0.01 to about 1000 mg, such as about
 5 0.01 to 100 mg, preferably from about 0.05 to about 500 mg, and more preferred from about 0.5 mg to about 200 mg.

For parenteral routes such as intravenous, intrathecal, intramuscular and similar
 administration, typically doses are in the order of about half the dose employed for oral
 10 administration.

Typical examples of recipes for the formulation of the invention are as follows:

- 1) Tablets containing 5.0 mg of a compound of the invention calculated as the free
 15 base:

Compound of the invention	5.0 mg
Lactose	60 mg
Maize starch	30 mg
Hydroxypropylcellulose	2.4 mg
20 Microcrystalline cellulose	19.2 mg
Croscarmellose Sodium Type A	2.4 mg
Magnesium stearate	0.84 mg

- 2) Tablets containing 0.5 mg of a compound of the invention calculated as
 25 the free base:

Compound of the invention	0.5 mg
Lactose	46.9 mg
Maize starch	23.5 mg
Povidone	1.8 mg
30 Microcrystalline cellulose	14.4 mg
Croscarmellose Sodium Type A	1.8 mg
Magnesium stearate	0.63 mg

3) Syrup containing per millilitre:

	Compound of the invention	25 mg
	Sorbitol	500 mg
	Hydroxypropylcellulose	15 mg
5	Glycerol	50 mg
	Methyl-paraben	1 mg
	Propyl-paraben	0.1 mg
	Ethanol	0.005 mL
	Flavour	0.05 mg
10	Saccharin sodium	0.5 mg
	Water	ad 1 mL

4) Solution for injection containing per millilitre:

	Compound of the invention	0.5 mg
15	Sorbitol	5.1 mg
	Acetic Acid	0.05 mg
	Saccharin sodium	0.5 mg
	Water	ad 1mL

20

By the expression a compound of the invention is meant any one of the embodiments of formula I, IA, IB, IC, ID and IE as described herein.



In a further aspect the present invention relates to a method of preparing a compound of the invention as described in the following.

25

METHODS OF PREPARATION OF THE COMPOUNDS OF THE INVENTION

The compounds of the invention may be prepared as follows:

30 Deprotection or cleavage from a polymer support of a compound of formula VA or VB



wherein Z, the dotted line and R¹-R¹⁰ are as previously described, and R' is a carbamate (such as methyl-, ethyl-, tert-butyl-, allyl-, or benzyl-carbamate) or a benzyl-derived


- 5 protective group, wherein the protective groups may be linked to a solid support, like the Wang resin-based carbamate linker, whereupon the compound of formula I is isolated as the free base or an acid addition salt thereof.

The deprotection according to "Methods of preparation of the compounds of the invention" was performed by standard techniques, known to the persons skilled in the art and detailed in the textbook *Protective Groups in Organic Synthesis* Greene and Wuts, Wiley Interscience, (1991), ISBN 0471623016. The cleavage from a polymer support, such as from the Wang resin based carbamate linker, according to method A) was performed according to literature known procedures (Zaragoza *Tetrahedron Lett.* **1995**, 36, 8677-8678 and Conti *et al. Tetrahedron Lett.* **1997**, 38, 2915-2918).

15

Intermediates of formula VA and VB can be prepared by the following methods:

- Method A** Chemical transformation of a substituted thiophenol compound of formula VIA or of a substituted diphenyl disulfide compound of formula VIB to the corresponding benzenesulfonyl chloride compound and subsequent reaction with an indole of formula VIII or IX gives the protected intermediates VA or VB, respectively.
- 20




wherein Z is CH or N, and R¹-R¹⁰ are as previously described and R' is a nitrogen protecting group such as a carbamate group (such as methyl-, ethyl-, tert-butyl-, allyl-, or benzyl-carbamate) or a benzyl-derived protective group, wherein the protective groups may be linked to a solid support, like the Wang resin-based carbamate linker. Compounds of formula VIA (with Z being CH) can be prepared by demethylation of the corresponding 4-(2-methoxy-phenyl)-piperidine, protection of the nitrogen of the piperidine ring (with R' as described above), activation of the phenol functionality (with e.g. a triflyl- or nonafllyl group) and palladium catalysed exchange with trialkyl silanethiol (e.g. tri-iso-propyl silanethiol) using a catalyst such as *e.g.* tris(dibenzylideneacetone)dipalladium in combination with bis(2-diphenylphosphinophenyl)ether. Removal of the silane-protection group then gives compounds of formula VIA. Compounds of formula VIB (with Z being N) can be prepared by the reaction of mono-protected piperazine (with R' as described above) and 2-fluoro-1-nitrobenzenes, reduction of the nitro functionality, diazonium salt formation, followed by reaction with a xanthate salt (e.g. potassium ethylxanthate). Hydrolysis of the newly formed xanthate moiety followed by oxidation gives disulphides of formula VIB. Indoles of formula VIII and IX are either commercially available or can be prepared

according to methods described in standard works such as Houben-Weyl, Methoden der organischen Chemie (Methods of Organic Chemistry), Georg-Thieme-Verlag, Stuttgart; Organic Reactions, John Wiley & Sons, Inc. New York, namely under reaction conditions such as those which are known and suitable for such reactions.

5

Method B Chemical transformation of a thiophenol compound with the formula X to the corresponding benzenesulfonyl chloride compound and subsequent reaction with an indole of formula VIII or IX resulting in the intermediate XIIA or XIIB, which can be coupled with a suitable protected amine of formula XIII under palladium catalysis, such as *e.g.*


10 bis-(dibenzylideneacetone)palladium) in combination with 2-phosphino-2'-(N,N-dimethylamino) biphenyl, to give the compounds VA or VB.



15 wherein Z is nitrogen, R^1 - R^{10} are as previously described and R' is a carbamate (such as methyl-, ethyl-, tert-butyl-, allyl-, or benzyl-carbamate) or a benzyl-derived protective

group, wherein the protective groups may be linked to a solid support, like the Wang resin-based carbamate linker.

- Method C** Chemical transformation of a thiophenol compound with the formula X to the
- 5 corresponding benzenesulfonyl chloride compound and subsequent reaction with an indole of formula VIII or IX resulting in the intermediate XIIA or XIIB, which can be coupled with a suitable protected amine boronic acid ester of formula XIV under palladium catalysis to give the compounds VA or VB using a catalyst such as *e.g.*
- 1,1'-bis(diphenylphosphino)ferrocenedichloropalladium(II).



wherein Z is C, the dotted line represents a bond, R¹-R¹⁰ are as previously described and R'
 5 is a carbamate (such as methyl-, ethyl-, tert-butyl-, allyl-, or benzyl-carbamate) or a benzyl-
 derived protective group, wherein the protective groups may be linked to a solid support,
 like the Wang resin-based carbamate linker.

Method D Reduction of a tetrahydropyridine compound of formula VA (in which Z is C and the dotted line represents a bond) to the corresponding piperidine compound of formula VA (in which Z is CH), and reduction of a tetrahydropyridine compound of formula VB (in which Z is C and the dotted line represents a bond) to the corresponding piperidine

- 5 compound of formula VB (in which Z is CH). Subsequently the compound of formula I is isolated as the free base or an acid addition salt thereof.

The invention disclosed herein is further illustrated by the following non-limiting examples.

10 EXAMPLES

General Methods

Analytical LC-MS data were obtained on a PE Sciex API 150EX instrument equipped with photoionization (APPI) ion source and Shimadzu LC-8A/SLC-10A LC system. The LC conditions (Symmetry C18 column 4.6 × 30 mm with a particle size of 3.5 μm) were linear
15 gradient elution with eluents A (water containing 0.05% TFA) and B (acetonitrile containing 5% water and 0.035% TFA). Gradient (time[min]/%B): (0.00/10.0); (4.00/100.0); (4.10/10.0); (5.00/10.0) with 2 mL/min. Purity was determined by integration of the UV trace (254 nm) and ELS (SEDERE SEDEX 55 with Heto CBN 8-30 cooling bath). The retention times, R_t , are expressed in minutes.

20

Mass spectra were obtained by an alternating scan method to give molecular weight information. The molecular ion, MH^+ , was obtained at low orifice voltage (5V) and fragmentation at high orifice voltage (100V).

- 25 Preparative LC-MS-separation was performed on the same instrument. The LC conditions (Symmetry C18 column 10 × 50 mm) were linear gradient elution with eluents A (water containing 0.05% TFA) and B (acetonitrile containing 5% water and 0.035% TFA): (time[min]/%B): (0.00/20.0); (7.00/100.0); (7.10/20.0); (8.00/20.0) with 5.7 mL/min. Fraction collection was performed by split-flow MS detection.

30

For column chromatography silica gel of type Kieselgel 60, 230-400 mesh ASTM was used. For ion-exchange chromatography (SCX, 1 g, Varian Mega Bond Elut®, Chrompack cat.

No. 220776) was used. Prior use of the SCX-columns was pre-conditioned with 10% solution of acetic acid in methanol (3 mL).

Preperation of intermediates

5

4-(2-Methoxy-phenyl)-piperidine (Method A)

Purchased from Maybridge (Product Number BTB 13447)

4-(2-Hydroxy-phenyl)-piperidine-1-carboxylic acid tert-butyl ester (Method A)

- 10 BBr₃ (7.0 mL; 74 mmol) was added slowly to 4-(2-methoxy-phenyl)-piperidine (6.25 g; 32.7 mmol) in 100 dichloromethane at 0°C under argon. The reaction mixture was allowed to warm to room temperature and stirred 16 hours. The reaction mixture was quenched by carefully adding 100 mL water at 0°C. Another 300 mL water was added, followed by the addition of 50 mL 28% aqueous NaOH. The organic phase was discarded. Di-tert-butyl
- 15 dicarbonate (9.0 g; 41 mmol) was added to the aqueous phase. The reaction mixture was stirred 1 hour at room temperature, neutralized with 1 N HCl, extracted with ethyl acetate (2 x 200 mL). The organic phase was washed with brine (100 mL), dried with MgSO₄ and concentrated *in vacuo* to give 4-(2-hydroxy-phenyl)-piperidine-1-carboxylic acid tert-butyl ester.

20

4-[2-(Nonafluorobutane-1-sulfonyloxy)-phenyl]-piperidine-1-carboxylic acid tert-butyl ester (Method A)

- Nonafluoro-butane-1-sulfonyl fluoride (2.71 mL; 15.1 mmol) was added to a mixture of ethyl-diisopropyl-amine (2.87 mL; 16.5 mmol), 4-(*N,N*-dimethylamino)-pyridine (25 mg)
- 25 and 4-(2-hydroxy-phenyl)-piperidine-1-carboxylic acid tert-butyl ester (3.81 g; 13.7 mmol) in 50 ml 1,2-dichloro-ethane at 0°C under argon. The reaction mixture was allowed to warm to room temperature and stirred 16 hours. The reaction mixture was extracted with water (100 mL) and brine (100 mL). The combined aqueous phases were extracted with ethyl acetate (100 mL). The combined organic phases were washed with brine (100 mL), dried
- 30 with MgSO₄ and concentrated *in vacuo*. The residue was purified by flash chromatography (silica gel; ethyl acetate/heptane) to give 4.34 g 4-[2-(nonafluorobutane-1-sulfonyloxy)-phenyl]-piperidine-1-carboxylic acid tert-butyl ester (7.76 mmol; 57 %).

4-(2-Triisopropylsilanylsulfanyl-phenyl)-piperidine-1-carboxylic acid tert-butyl ester (Method A)

Bis[(2-diphenylphosphino) phenyl] ether (0.43 g; 0.80 mmol) and tris(dibenzylideneacetone)-dipalladium (0.37 mg; 0.40 mmol) were dissolved in 40 mL
5 toluene. 4-[2-(Nonafluorobutane-1-sulfonyloxy)-phenyl]-piperidine-1-carboxylic acid tert-butyl ester (4.34 g; 7.76 mmol), triisopropyl-silanethiol (1.83 mL; 8.53 mmol) and sodium *tert*-butoxide (0.90 g; 9.3 mmol) were added and the reaction mixture was stirred under argon for 3 hours at 100°C. The reaction mixture was cooled to room temperature filtered through a pad of silica and concentrated *in vacuo*. The residue was purified by flash
10 chromatography (silica gel; ethyl acetate/heptane) to give 1.58 g 4-(2-triisopropylsilanylsulfanyl-phenyl)-piperidine-1-carboxylic acid tert-butyl ester (3.51 mmol; 45 %).

4-(5-Methyl-2-nitro-phenyl)-piperazine-1-carboxylic acid benzyl ester (Method A)

15 Piperazine-1-carboxylic acid benzyl ester (8.00 g; 36.3 mmol), 2-fluoro-4-methyl-1-nitro-benzene (4.70 g; 30.3 mmol) and potassium carbonate (8.5 g; 61.5 mmol) were stirred 3 hours in 50 mL DMSO at 80°C. The reaction mixture was cooled to room temperature and 250 mL water was added. The mixture was extracted with diethyl ether (2 x 250 mL). The organic phase was washed with water (100 mL), 1 N HCl (2 x 100 mL), brine (2 x 100 mL),
20 dried with MgSO₄ and concentrated *in vacuo* to give 10.5 g (29.5 mmol; 97.6 %) 4-(5-methyl-2-nitro-phenyl)-piperazine-1-carboxylic acid benzyl ester.

4-(2-Amino-5-methyl-phenyl)-piperazine-1-carboxylic acid benzyl ester (Method A)

4-(5-Methyl-2-nitro-phenyl)-piperazine-1-carboxylic acid benzyl ester (5.60 g; 15.7 mmol)
25 and SnCl₂·H₂O (17.5 g; 77.6 mmol) were refluxed in 100 mL ethanol for 2½ hours. Ethanol was removed *in vacuo*. The residue was poured into 300 mL saturated NaHCO₃. The mixture was extracted with ethyl acetate (2 x 250 mL). The organic phase was washed with brine (2 x 100 mL), dried with MgSO₄, filtered through a pad of silica and concentrated *in vacuo* to give 4.71 g (14.5 mmol; 91.6 %) 4-(2-amino-5-methyl-phenyl)-piperazine-1-
30 carboxylic acid benzyl ester.

2-(4-(Benzyloxy-carbonyl)-piperazin-1-yl)-4-methyl-benzenedisulphide (Method A)

Sulphuric acid (2.37 g; 24.2 mmol) in 20 mL of water was added to 4-(2-amino-5-methyl-phenyl)-piperazine-1-carboxylic acid benzyl ester in 20 mL THF. THF was removed *in vacuo*. The mixture was cooled to 0°C and NaNO₂ (0.611 g; 8.86 mmol) in 10 mL water was added dropwise. The reaction mixture was stirred 15 minutes at room temperature and
5 added dropwise to potassium ethylxanthate (3.87 g; 24.2 mmol) in 30 mL water at 70°C (CAUTION!: gas evolution). The reaction mixture was stirred 2 hours at 70°C. 10 mL 28% NaOH and 30 mL ethanol were added and the reaction mixture was stirred 16 hours at 60°C in an open flask. The reaction mixture was cooled to room temperature and extracted with diethyl ether (2 x 250 mL). The organic phase was washed with brine (2 x 100 mL), dried
10 with MgSO₄ and concentrated *in vacuo*. The residue was purified by flash chromatography (silica gel; ethyl acetate/heptane) to give 1.19g 2-(4-(benzyloxy-carbonyl)-piperazin-1-yl)-4-methyl-benzenedisulphide (1.74 mmol; 43%).

2-(4-(Benzyloxy-carbonyl)-piperazin-1-yl)- benzenedisulphide (Method A) was prepared in
15 a similar way.

3-(2-Bromo-phenylsulfanyl)-1H-indole (Method B, C)

69.9 mmol of N-chlorosuccinimide was suspended in 200 mL dry CH₂Cl₂ and cooled in an ice bath to 0 °C. It was slowly treated with 66.6 mmol 2-bromothiophenol. After addition
20 was completed the solution cleared up and turned orange. Stirring was continued at room temperature for 1 hour.

The above solution was added slowly to a solution of 66.6 mmol indole in 180 mL dry DMF at 0 °C under argon. After addition was completed the solution was stirred for 2½ hour at 0 °C.

25 The reaction was quenched with 200 mL water, and 150 mL sat. NaHCO₃ was added. 250 mL ethyl acetate was added and the phases were separated. The water phase was back extracted with ethyl acetate. The combined organic phases were then washed with brine, dried over MgSO₄ and evaporated under vacuum. The title compound was recrystallized from acetonitrile.

30

3-(2-Bromo-phenylsulfanyl)-indole-1-carboxylic acid tert-butyl ester (Method B, C)

To a solution of 26.3 mmol 3-(2-bromo-phenylsulfanyl)-1H-indole in dry CH₂Cl₂ (120 mL) was added 1.3 mmol of 4-dimethyl aminopyridine and 28.9 of mmol triethylamine. 28.9 mmol of di-tert-butyl-dicarbonate were added and the solution was stirred at room temperature over night under argon. The solution was washed with 0.5 M aqueous HCl. The water phase was extracted with CH₂Cl₂. The combined organic phase was dried over K₂CO₃ and evaporated under vacuum. The compound was applied to a silica column in 1 / 4 ethyl acetate / heptane and filtered through using 1 / 4 ethyl acetate / heptane. The eluent was evaporated under vacuum to give the title compound.

10 **3-[2-(4-tert-Butoxycarbonyl-piperazin-1-yl)-phenylsulfanyl]-indole-1-carboxylic acid tert-butyl ester (Method B)**

0.0086 mmol of Pd(dba)₂ (bis-(dibenzylideneacetone)palladium) and 0.0172 mmol of 2-phosphino-2'-(N,N-dimethylamino) biphenyl were added to a dry flask and stored under argon.

15 0.043 mmol of 3-(2-bromo-phenylsulfanyl)-indole-1-carboxylic acid tert-butyl ester and 0.052 mmol of *N*-tert-butoxycarbonyl-piperazine were dissolved in 2.5 mL dry toluene and 0.061 mmol sodium-tert-butoxide in toluene were added. The solution was degassed and added to the flask with bis(dibenzylideneacetone)-palladium and of 2-phosphino-2'-(N,N-dimethylamino)biphenyl. The reaction was stirred for 18 hour at 100 °C. Ethyl acetate (3 mL) was added and the organic phase was washed with aqueous NH₄Cl and brine, dried over MgSO₄ and evaporated under vacuum to give the title compound.

4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (Method C)

25 Was made according to the procedure described in Tetrahedron Letters 41 (2000) 3705-3708.

Example 1 Compounds of the invention of Formula IA

Synthesis of 3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole

30 3 M HCl-gas in dry ethyl acetate (3 mL) was added to 3-[2-(4-tert-butoxycarbonyl-piperazin-1-yl)-phenylsulfanyl]-indole-1-carboxylic acid *tert*-butyl ester. The solution was stirred for 1 h at 0 °C and evaporated under vacuum. The crude product was purified by preparative LC-MS. The isolated product was submitted for testing as DMSO solution.

The following compounds were prepared in a similar way and analytical data are shown in table 1:

1. 3-(2-Piperazin-1-yl-phenylsulfanyl)-1H-indole
- 5 2. 4-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
3. 3-(2-Piperazin-1-yl-phenylsulfanyl)-1H-indole-7-carboxylic acid methyl ester
4. 5-Nitro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
5. 6-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
6. 6-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
- 10 7. 6-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
8. 7-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
9. 3-[2-(1,2,3,6-Tetrahydro-pyridin-4-yl)-phenylsulfanyl]-1H-indole
11. 4-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
12. 5-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole

15

Example 2 Compounds of Formula IA or IB

Synthesis of 3-(2-Piperidin-4-yl-phenylsulfanyl)-1H-indole

- 1.58 g 4-(2-Triisopropylsilanylsulfanyl-phenyl)-piperidine-1-carboxylic acid *tert*-butyl ester (3.51 mmol) in 10 mL THF was added to 1.21 g tetrabutylammonium fluoride dihydrate (3.85 mmol) in 5 mL THF at 0°C. The reaction mixture was stirred 1 hour at 0°C, filtered through a plug of silica (eluted with EtOAc/heptane 1:1) and concentrated *in vacuo*. The residue was redissolved in 14 mL THF. 2 mL of this solution (≈ 0.5 mmol) was added to 67 mg *N*-chloro succinimide (0.50 mmol) in 2 mL 1,2-dichloro-ethane at 0°C. The reaction mixture was stirred 30 minutes at room temperature. The resulting solution was added to 90 mg indole (0.75 mmol) in 2 mL THF at 0°C. The reaction mixture was stirred 2 hour at 0°C, poured into 20 mL saturated NaHCO₃ and extracted with ethyl acetate (2x50 mL). The combined organic phases were washed with brine (20 mL), dried with MgSO₄ and concentrated *in vacuo*. The residue was purified by flash chromatography (silica gel; ethyl acetate/heptane) to give 89 mg 4-[2-(1H-indol-3-ylsulfanyl)-phenyl]-piperidine-1-carboxylic acid *tert*-butyl ester (0.22 mmol, $\approx 44\%$). 2 mL diethyl ether saturated with HCl was added to 4-[2-(1H-indol-3-ylsulfanyl)-phenyl]-piperidine-1-carboxylic acid *tert*-butyl ester and stirred over night at room temperature. The reaction mixture was neutralized with saturated NaHCO₃ and extracted with ethyl acetate (2x50 mL). The combined organic

phases were washed with brine (20 mL), dried with MgSO_4 and concentrated *in vacuo*.
3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole was isolated after preparative HPLC.

The following compounds were prepared in a similar way and analytical data are

5 shown in Table 1:

- 44. 3-(2-Piperidin-4-yl-phenylsulfanyl)-1H-indole
- 45. 2-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 46. 1-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 47. 5-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 10 48. 7-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 49. 6-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 50. 5-Methoxy-4-methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 51. 4-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 52. 7-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 15 53. 4-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 54. 4-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 55. 6-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 56. 5-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 57. 5-Methoxy-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 20 58. 5-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 59. 7-Methoxy-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 60. 7-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 61. 7-Ethyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
- 62. 3-Methyl-2-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole

25

Synthesis of 3-(4-Methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole

SO_2Cl_2 (120 μL ; 1.50 mmol) was added to 2-(4-(benzyloxy-carbonyl)-piperazin-1-yl)-4-methyl-benzenedisulphide (0.956 g; 1.40 mmol) in 28 mL 1,2-dichloroethane and stirred 15 minutes at room temperature. 1 mL of this mixture was added to indole (23 mg; 0.20
30 mmol) in 1 mL THF at 0°C under argon. The reaction mixture was stirred 2 hours at 0°C . 2 mL saturated NaHCO_3 and 2 mL ethyl acetate was added. The organic phase was separated and concentrated *in vacuo*. The residue was dissolved in 1 mL DMSO. 0.5 mL 20% aqueous KOH was added. The reaction mixture was stirred 4 minutes at 150°C under microwave

irradiation. 3 mL saturated aqueous NH_4Cl was added. The mixture was extracted with ethyl acetate (2 x 5 mL). The organic phase was concentrated *in vacuo*. 3-(4-Methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole was isolated after preparative HPLC.

5 The following compounds were prepared in a similar way and analytical data are shown in Table 1:

10. 6-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
13. 4-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
14. 6-Fluoro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
- 10 15. 4-Fluoro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
16. 7-Fluoro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
17. 5-Methoxy-4-methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
18. 5-Methoxy-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
19. 7-Methoxy-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
- 15 20. 4-Methoxy-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
21. 4-Chloro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
22. 7-Chloro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
23. 1-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
24. 3-Methyl-2-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
- 20 25. 2-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
26. 5-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
27. 4-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
28. 7-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
29. 7-Ethyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
- 25 30. 5-Methoxy-4-methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
31. 5-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
32. 7-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
33. 5-Fluoro-2-methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
34. 5-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
- 30 35. 4-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
36. 7-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
37. 1-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
38. 3-Methyl-2-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole

39. 6-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
 40. 2-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
 41. 5-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
 42. 7-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole

5

Table 1. Measured molecular mass, measured HPLC-retention time (RT, min) and UV- and ELSD-purities (%).

compound	RT min.	UV-purity (%)	ELSD- purity (%)	M+H ⁺	Synthesis method
1	2.04	98.3	65.3	310	B
2	1.77	100	91.3	340	B
3	1.93	100	93.42	368	B
4	1.82	90.5	57.96	355	B
5	2.005	98.69	93.33	346	B
6	1.78	98.43	97.6	340	B
7	1.875	99.24	89.92	328	B
8	1.92	95.03	98.24	324	B
9	2	93.16	77.61	307.1	C
10	2.017	73.77	96.96	342.2	A
11	1.86	93.4	99.6	328	B
12	2.04	96.06	97.98	328	B
13	2.335	91.77	95.19	338.3	A
14	2.3	84.45	99.8	341.9	A
15	2.213	74.5	95.49	341.8	A
16	2.302	83.07	98.83	342.0	A
17	2.37	79.23	94.36	398.1	A
18	2.193	72.02	88.95	354.2	A
19	2.282	81.72	94.25	354.0	A
20	2.185	71.34	93.84	354.2	A
21	2.3	82.42	96.53	358.0	A
22	2.442	82.34	98.28	358.0	A
23	2.435	84.72	96.25	338.0	A

24	2.568	87.44	97.55	338.3	A
25	1.913	91.99	99.79	324.1	A
26	1.983	74.46	95.27	323.9	A
27	1.967	82.68	100	324.1	A
28	1.928	96.71	99.85	328.1	A
29	2.137	71.62	99.05	338.3	A
30	1.92	75.46	100	354.2	A
31	1.815	70.05	88.36	340.0	A
32	1.907	74.16	93.41	340.1	A
33	1.965	94.37	99.87	342.0	A
34	2.047	93.32	100	344.1	A
35	2.04	95.23	99.88	344.1	A
36	2.078	94.53	100	344.1	A
37	2.07	95.53	100	324.0	A
38	2.213	95.06	99.92	324.1	A
39	2.02	79.43	98.47	338.3	A
40	2.03	80.36	95.66	338.3	A
41	2.09	77.01	98.45	338.1	A
42	2.108	84.83	99.76	338.3	A
44	1.933	93.78	99.68	309.1	A
45	1.995	91.34	99.51	323.1	A
46	2.133	98.98	99.47	323	A
47	2.06	87.52	98.71	323.1	A
48	2.012	91.07	98.64	327.2	A
49	2.008	95.96	98.6	326.9	A
50	1.963	83.51	98.37	354.2	A
51	1.937	88.63	97.5	327.2	A
52	2.072	95.08	97.16	323.1	A
53	2.013	89.2	97.12	323.1	A
54	1.987	79.17	96.72	342.9	A
55	2.158	92	95.27	342.9	A
56	1.987	94.53	95.17	327	A

57	1.9	78.75	94.62	339.1	A
58	2.042	88.63	87.81	342.9	A
59	1.985	90.24	80.44	339.3	A
60	2.108	95.87	99.68	342.9	A
61	2.177	73.11	98.96	337.2	A
62	2.242	95.29	98.7	323	A
63	1,987	92.82	99.65	323,1	A
64	1,958	75.72	95.59	341,1	A

Example 3 **Transporter inhibition assay**

5 **Measurements of [³H]-5-HT uptake into rat cortical synaptosomes**

Whole brains from male Wistar rats (125-225 g), excluding cerebellum, are homogenized in 0.32 M sucrose supplemented with 1mM nialamid with a glass/teflon homogenizer. The homogenate is centrifuged at 600 x g for 10 min at 4 °C. The pellet is discarded and the supernatant is centrifuged at 20.000 x g for 55 min. The final pellet is homogenized (20 sec) in this assay buffer (0.5 mg original tissue/well). Test compounds (or buffer) and 10 nM [³H]-5-HT are added to 96 well plates and shaken briefly. Composition of assay buffer: 123 mM NaCl, 4.82 mM KCl, 0.973 mM CaCl₂, 1.12 mM MgSO₄, 12.66 mM Na₂HPO₄, 2.97 mM NaH₂PO₄, 0.162 mM EDTA, 10 mM glucose and 1 mM ascorbic acid. Buffer is oxygenated with 95% O₂/5% CO₂ for 10 min at 37 °C and pH is adjusted 7.4. The incubation is started by adding tissue to a final assay volume of 0.2 mL. After 15 min incubation with radioligand at 37 °C, samples are filtered directly on Unifilter GF/C glass fiber filters (soaked for 1 hour in 0.1% polyethylenimine) under vacuum and immediately washed with 3 x 0.2 ml assay buffer. Non-specific uptake is determined using citalopram (10 µM final concentration). Citalopram is included as reference in all experiments as dose-response curve.

20

Measurements of [³H]noradrenaline uptake into rat cortical synaptosomes

Fresh cortex from male Wistar rats (125-225 g) are homogenized in 0.4M sucrose with a glass/teflon homogenizer. The homogenate is centrifuged at 600 x g for 10 min at 4 °C. The pellet is discarded and the supernatant is centrifuged at 20.000 x g for 55 min. The final pellet is homogenized (20 sec) in this assay buffer (6 mg original tissue/mL = 4 mg/well).

25

Test compounds (or buffer) and 10 nM [^3H]-noradrenaline are added to deep 96 well plates and shaken briefly. Composition of assay buffer: 123 mM NaCl, 4.82 mM KCl, 0.973 mM CaCl_2 , 1.12 mM MgSO_4 , 12.66 mM Na_2HPO_4 , 2.97 mM NaH_2PO_4 , 0.162 mM EDTA, 10 mM glucose and 1 mM ascorbic acid. Buffer is oxygenated with 95% O_2 /5% CO_2 for 10 min at 37 °C and pH is adjusted 7.4. The incubation is started by adding tissue to a final assay volume of 1 ml. After 15 min incubation with radioligand at 37 °C, samples are filtered directly on Unifilter GF/C glass fiber filters (soaked for 1 hour in 0.1% polyethylenimine) under vacuum and immediately washed with 3 x 1 mL assay buffer. Non-specific uptake is determined using talsupram (10 μM final concentration). Duloxetine is included as reference in all experiments as dose-response curve.

Measurements of [^3H]dopamine uptake into rat cortical synaptosomes

Tissue preparation: male wistar rats (125-250 g) are sacrificed by decapitated and striatum quickly dissected out and placed in 40 vol (w/v) ice cold 0,40 M sucrose. The tissue is gently homogenised (glass teflon homogeniser) and the P2 fraction is obtained by centrifugation (1000 g, 10 minutes and 40000 g, 20 minutes, 4°C) and suspended in 560 volumes of a modified Krebs-Ringer-phosphate buffer, pH 7.4.

Tissue 0,25 mg/well(140 μl) (original tissue) is mixed with test suspension. After 5 minutes pre-incubation 12.5 nM 3H-dopamine is added and the mixture is incubated for 5 minutes at RT.

The incubation is terminated by filtering the samples under vacuum through Whatman GF/C filters with a wash of 1 ml buffer. The filters are dried and appropriate scintillation fluid (Optiphase Supermix) is added. After storage for 2 hours in the dark the content of radioactivity is determined by liquid scintillation counting. Uptake is obtained by subtracting the non-specific binding and passive transport measured in the presence of 100 μM of benztropin. For determination of the inhibition of uptake ten concentrations of drugs covering 6 decades are used.

^3H -DA = 3,4-(ring-2,5,6- ^3H)dopamine hydrochloride from New England Nuclear, specific activity 30-50 Ci/mmol.


Hyttel, Biochem. Pharmacol. 1978, 27, 1063-1068;

Hyttel, Prog. Neuro-Psychopharmacol. & bil. Psychiat. 1982, 6, 277-295;

Hyttel & Larsen, Acta Pharmacol. Tox. 1985, 56, suppl. 1, 146-153.

CLAIMS

1. A compound having the general formula I as the free base or salts thereof



5 wherein

the dotted line represents an optional bond;

and

each of R¹-R⁴ is independently selected from the group consisting of hydrogen, halogen, cyano, nitro, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-

10 C₁₋₆-alk(en/yn)yl, amino, C₁₋₆-alk(en/yn)ylamino, di-(C₁₋₆-alk(en/yn)yl)amino,

C₁₋₆-alk(en/yn)ylcarbonyl, aminocarbonyl, C₁₋₆-alk(en/yn)ylaminocarbonyl,

di-(C₁₋₆-alk(en)yl)aminocarbonyl, hydroxy, C₁₋₆-alk(en/yn)ylloxy,

C₁₋₆-alk(en/yn)ylsulfanyl, halo-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)ylsulfonyl, halo-

C₁₋₆-alk(en/yn)ylsulfanyl and C₁₋₆-alk(en/yn)ylsulfonyl;

15 and

R⁵ is selected from the group consisting of hydrogen, halogen, cyano, C₁₋₆-alk(en/yn)yl,

C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, amino,

C₁₋₆-alk(en/yn)ylamino, di-(C₁₋₆-alk(en/yn)yl)amino, C₁₋₆-alk(en/yn)ylcarbonyl,

aminocarbonyl, C₁₋₆-alk(en/yn)ylaminocarbonyl, di-(C₁₋₆-alk(en)yl)aminocarbonyl,

20 hydroxy, C₁₋₆-alk(en/yn)ylloxy, C₁₋₆-alk(en/yn)ylsulfanyl, halo-C₁₋₆-alk(en/yn)yl, halo-

C₁₋₆-alk(en/yn)ylsulfonyl, halo-C₁₋₆-alk(en/yn)ylsulfanyl and C₁₋₆-alk(en/yn)ylsulfonyl;

and

each of R⁶-R⁹ is independently selected from the group consisting of hydrogen, halogen, cyano, nitro, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-

25 C₁₋₆-alk(en/yn)yl, amino, C₁₋₆-alk(en/yn)ylamino, di-(C₁₋₆-alk(en/yn)yl)amino,

C₁₋₆-alk(en/yn)ylcarbonyl, aminocarbonyl, C₁₋₆-alk(en/yn)ylaminocarbonyl,

di-(C₁₋₆-alk(en)yl)aminocarbonyl, hydroxy, C₁₋₆-alk(en/yn)ylloxy,

C₁₋₆-alk(en/yn)ylsulfanyl, halo-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)ylsulfonyl, halo-C₁₋₆-alk(en/yn)ylsulfanyl, C₁₋₆-alk(en/yn)ylsulfonyl and C₁₋₆-alk(en/yn)yl-O-CO-;

and

R¹⁰ is selected from the group consisting of hydrogen, C₁₋₆-alk(en/yn)yl,

5 C₃₋₈-cycloalk(en)yl and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl;

and

Z is selected from the group consisting of N, CH and C, with the proviso that the dotted line represents a bond when Z is C and the dotted line does not represent a bond when Z is CH or N.

10

2. A compound according to claim 1 wherein each of R¹-R⁴ is independently selected from the group consisting of hydrogen, halogen and C₁₋₆-alk(en/yn)yl.

15 3. A compound according to any one of claims 1-2 wherein R⁵ is selected from the group consisting of hydrogen and C₁₋₆-alk(en/yn)yl.

4. A compound according to any one of claims 1-3 wherein each of R⁶-R⁹ is independently selected from the group consisting of hydrogen, halogen, nitro, C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)oxy and C₁₋₆-alk(en/yn)yl-O-CO-.

20

5. A compound according to any one of claims 1-4 wherein R¹⁰ is selected from the group consisting of hydrogen and C₁₋₆-alk(en/yn)yl.

25 6. A compound according to any one of claims 1-5 wherein the dotted line represents a bond and Z is a carbon atom

7. A compound according to any one of claims 1-5 wherein Z is a nitrogen atom and the dotted line does not represent a line.

30 8. A compound according to any one of claims 1-5 wherein Z represents CH and the dotted line does not represent a line.

9. A compound according to any one of claims 1-8, said compounds being selected from the following list of compounds

Compound no.	Name
1	3-(2-Piperazin-1-yl-phenylsulfanyl)-1H-indole
2	4-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
3	3-(2-Piperazin-1-yl-phenylsulfanyl)-1H-indole-7-carboxylic acid methyl ester
4	5-Nitro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
5	6-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
6	6-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
7	6-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
8	7-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
9	3-[2-(1,2,3,6-Tetrahydro-pyridin-4-yl)-phenylsulfanyl]-1H-indole
10	6-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
11	4-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
12	5-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
13	4-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
14	6-Fluoro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
15	4-Fluoro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
16	7-Fluoro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
17	5-Methoxy-4-methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)- 1H-indole
18	5-Methoxy-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
19	7-Methoxy-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
20	4-Methoxy-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
21	4-Chloro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
22	7-Chloro-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
23	1-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
24	3-Methyl-2-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
25	2-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
26	5-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole

27	4-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
28	7-Fluoro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
29	7-Ethyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
30	5-Methoxy-4-methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
31	5-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
32	7-Methoxy-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
33	5-Fluoro-2-methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
34	5-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
35	4-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
36	7-Chloro-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
37	1-Methyl-3-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
38	3-Methyl-2-(2-piperazin-1-yl-phenylsulfanyl)-1H-indole
39	6-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
40	2-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
41	5-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
42	7-Methyl-3-(4-methyl-2-piperazin-1-yl-phenylsulfanyl)-1H-indole
44	3-(2-Piperidin-4-yl-phenylsulfanyl)-1H-indole
45	2-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
46	1-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
47	5-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
48	7-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
49	6-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
50	5-Methoxy-4-methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
51	4-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
52	7-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
53	4-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
54	4-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
55	6-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
56	5-Fluoro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
57	5-Methoxy-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
58	5-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
59	7-Methoxy-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole

60	7-Chloro-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
61	7-Ethyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
62	3-Methyl-2-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
63	6-Methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole
64	5-Fluoro-2-methyl-3-(2-piperidin-4-yl-phenylsulfanyl)-1H-indole

as the free base or a salt thereof.

10. A pharmaceutical composition comprising a compound according to any one of claims 1-9 and at least one pharmaceutically acceptable carrier or diluent.
- 5 11. Use of a pharmaceutical composition according to claim 10 for the inhibition of the serotonin transporter.
12. Use according to claim 11 for the treatment of a disease or disorder wherein a serotonin reuptake inhibitor is beneficial.
- 10 13. Use of a pharmaceutical composition according to claim 10 for the inhibition of the serotonin and noradrenaline transporters.
- 15 14. Use according to claim 13 for the treatment of a disease or disorder, wherein a combined serotonin and norepinephrine reuptake inhibitor is beneficial.
15. Use of a pharmaceutical composition according to claim 10 for the inhibition of the serotonin, noradrenaline and dopamine transporters.
- 20 16. Use according to claim 15 for the treatment of a disease or disorder, wherein a combined serotonin, norepinephrine and dopamine reuptake inhibitor is beneficial.
17. Use of a pharmaceutical composition according to claim 10 for the inhibition of the serotonin and dopamine transporters
- 25 18. Use according to claim 17 for the treatment of a disease or disorder, wherein a combined serotonin and dopamine reuptake inhibitor is beneficial.

19. Use according to any one of claims 12, 14, 16 and 18, wherein said disorder or disease is an affective disorder.
- 5 20. Use according to claim 19, wherein the affective disorder is a depressive disorder such as major depressive disorder, postnatal depression, dysthymia or depression associated with bipolar disorder, alzheimers, psychosis or parkinsons.
- 10 21. Use according to claim 19, wherein the affective disorder is an anxiety disorder such as general anxiety disorder, social anxiety disorder, post traumatic stress disorder, obsessive compulsive disorder, panic disorder, panic attacks, specific phobias, social phobia and agoraphobia.
- 15 22. Use according to any one of claims 14 and 16, wherein said disorder or disease is a pain disorder such as fibromyalgia syndrome, overall pain, back pain, shoulder pain, headache as well as pain while awake and during daily activities.
- 20 23. Use according to any one of claims 14 and 16, wherein said disorder or disease is attention deficit hyperactivity disorder.
24. Use according to any one of claims 14 and 16, wherein said disorder or disease is stress urinary incontinence.