
[0007] The pituitary tumor to be treated (i.e., shrunk) may be a non-functional (non-secreting) tumor, a hormone-secreting tumor, a benign tumor, or a malignant tumor, with or without metastasis. The pituitary tumor is typically not a non-pituitary metastatic tumor. However, in example, the pituitary tumor may be an invasive tumor. Thus, the pituitary tumor to be treated may be an adenoma or carcinoma, a non-functional (non-secreting) tumor, a hormone-secreting tumor (e.g., a hormone-secreting neuroendocrine tumor), or other tumor. In example, the pituitary tumor to be treated is a pituitary adenoma. In example, the pituitary adenoma to be treated is a macroadenoma. In example, the pituitary adenoma is an adenoma selected from a corticotroph (including a corticotroph with significant amounts of Crooke's hyalinazation, a corticotroph with little or no Crooke's hyalinization, and a silent corticotroph); a somatotroph; a lactotroph; a thyrotroph; a gonadotroph; and an oncocytoma. In example, the pituitary tumor to be reduced in size is pituitary tumor selected from a nonfunctional pituitary tumor; a growth hormone-secreting tumor; a gonadotropin-secreting tumor (e.g., secreting LH or FSH); or other tumor.

[0007a] In one aspect, the present disclosure refers to a use of relacorilant in the manufacture of a medicament for reducing the size of a visible pituitary macroadenoma tumor measuring at least 10 millimeters (mm) in at least one dimension in a patient, the use comprising selecting a patient suffering from a pituitary tumor suitable for reduction in size, said selecting comprising imaging said pituitary tumor effective to determine that said tumor is a visible macroadenoma, and that an effective amount of relacorilant is to be administered to the selected patient, wherein relacorilant is ((R)-(1-(4-fluorophenyl)-6-((1-methyl-1H-pyrazol-4-yl)sulfonyl)-4,4a,5,6,7,8-hexahydro-1H-pyrazolo[3,4-g]isoquinolin-4a-yl)(4-(trifluoromethyl)pyridin-2-yl)methanone), which has the following structure:




,

effective to reduce the size of the pituitary tumor.

CLAIMS:

1. Use of relacorilant in the manufacture of a medicament for reducing the size of a visible pituitary macroadenoma tumor measuring at least 10 millimeters (mm) in at least one dimension in a patient, the use comprising selecting a patient suffering from a pituitary tumor suitable for reduction in size, said selecting comprising imaging said pituitary tumor effective to determine that said tumor is a visible macroadenoma, and that an effective amount of relacorilant is to be administered to the selected patient, wherein relacorilant is ((R)-(1-(4-fluorophenyl)-6-((1-methyl-1H-pyrazol-4-yl)sulfonyl)-4,4a,5,6,7,8-hexahydro-1H-pyrazolo[3,4-g]isoquinolin-4a-yl)(4-(trifluoromethyl)pyridin-2-yl)methanone), which has the following structure:



,

effective to reduce the size of the pituitary tumor.

2. The use of claim 1, wherein said selecting further comprises determining that said patient suffers from a symptom of Cushing's syndrome.

3. The use of claim 1, further comprising measuring said pituitary tumor.

4. The use of claim 1, further comprising measuring said pituitary tumor following relacorilant administration, wherein said measuring comprises a second imaging of the pituitary tumor.

5. The use of claim 1, wherein said relacorilant administration comprises that relacorilant is to be administered orally.

6. The use of claim 1, wherein said pituitary tumor is a non-secreting pituitary tumor.

7. The use of claim 1, wherein said pituitary tumor is a hormone-secreting tumor, and said treatment is effective to reduce the size of the pituitary tumor and to reduce hormone secretion from said pituitary tumor.

8. The use of claim 1, wherein said relacorilant administration comprises that between ~~about~~ 20 milligrams (mg) to ~~about~~ 800 mg of relacorilant is to be administered daily.

9. The use of claim 1, wherein said relacorilant administration comprises that relacorilant is to be administered daily in single unit dosage forms each containing an amount of relacorilant selected from 50 milligrams (mg), 100 mg, 150 mg, 200 mg, 250 mg, 300 mg, 350 mg, 400 mg, 450 mg, and 500 mg of relacorilant.

10. The use of claim 4, further comprising application of a further pituitary tumor targeting treatment following said second imaging of the pituitary tumor.

11. The use of claim 10, wherein said pituitary tumor targeting treatment comprises surgical treatment.

12. The use of claim 5, wherein said administration of relacorilant comprises that relacorilant is to be administered orally to a fasted patient.

13. The use of claim 12, wherein said fasted patient is a patient who has not eaten a meal or otherwise ingested food for at least two hours prior to the administration of relacorilant.

14. The use of claim 5, wherein said administration of relacorilant comprises that relacorilant is to be administered orally to a fed patient.

15. The use of claim 14, wherein said fed patient is a patient who has begun eating a meal about 30 minutes or less prior to the administration of relacorilant.

16. The use of claim 14, wherein a fed patient is a patient who has begun eating a meal about one hour or less prior to the administration of relacorilant.

17. The use of claim 1, wherein said administration of relacorilant comprises that relacorilant is to be administered prior to surgery for resection of the pituitary tumor.

18. The use of claim 1, wherein said administration of relacorilant comprises that relacorilant is to be administered following surgery for resection of the pituitary tumor.

19. The use of claim 18, wherein said relacorilant administration following surgery for resection of the tumor is effective to reduce the size of, or growth of, tumor tissue remaining in the patient following said surgical treatment.

20. The use of claim 19, further comprising that radiation treatment or cancer chemotherapy treatment, or both, is to be administered to said patient.

21. The use of claim 1, wherein said administration of relacorilant comprises that relacorilant is to be administered to a patient who has been administered another pharmaceutical or medical treatment prior to, or at the same time as, said relacorilant.

22. The use of claim 1, wherein said administration of relacorilant comprises that relacorilant is to be administered to a patient, followed by surgical treatment in that said pituitary tumor is to be resected, then followed by a further step that relacorilant is to be administered.

23. The use of claim 1, wherein said administration of relacorilant comprises that relacorilant is to be administered to a patient, followed by surgical treatment in that said pituitary tumor is to be resected, then followed by radiation treatment targeting the pituitary tumor and/or regions near the location previously occupied by said pituitary tumor.

24. The use of claim 5, wherein said administration of relacorilant comprises that relacorilant is to be administered to a patient, followed by surgical treatment in that said pituitary tumor is to be resected, then followed by a further step that relacorilant is to be administered.

25. The use of claim 5, wherein said administration of relacorilant comprises that relacorilant is to be administered to a patient, followed by surgical treatment in that said pituitary tumor is to be resected, then followed by radiation treatment targeting the pituitary tumor and/or regions near the location previously occupied by said pituitary tumor.

26. The use of claim 14, wherein said administration of relacorilant comprises that relacorilant is to be administered to a patient, followed by surgical treatment in that said pituitary tumor is to be resected, then followed by a further step that relacorilant is to be administered.

27. The use of claim 14, wherein said administration of relacorilant comprises that relacorilant is to be administered to a patient, followed by surgical treatment in that said pituitary tumor is to be resected, then followed by radiation treatment targeting the pituitary tumor and/or regions near the location previously occupied by said pituitary tumor.

28. The use of claim 1, wherein said pituitary tumor is hormone-secreting tumor, and said tumor secretes a hormone selected from adrenocorticotrophic hormone, prolactin, growth hormone, thyroid stimulating hormone, follicle stimulating hormone, luteinizing hormone, melanocyte stimulating hormone, β-lipotropin, oxytocin, and vasopressin.

29. The use of claim 5, wherein said pituitary tumor is hormone-secreting tumor, and said tumor secretes a hormone selected from adrenocorticotrophic hormone, prolactin, growth hormone, thyroid stimulating hormone, follicle stimulating hormone, luteinizing hormone, melanocyte stimulating hormone, β-lipotropin, oxytocin, and vasopressin.

30. The use of claim 7, wherein said pituitary tumor is hormone-secreting tumor, and said tumor secretes a hormone selected from adrenocorticotrophic hormone, prolactin, growth hormone, thyroid stimulating hormone, follicle stimulating hormone, luteinizing hormone, melanocyte stimulating hormone, β-lipotropin, oxytocin, and vasopressin.