

(51) International Patent Classification:
C12P 19/18 (2006.01)(21) International Application Number:
PCT/US2019/041957(22) International Filing Date:
16 July 2019 (16.07.2019)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/698,617 16 July 2018 (16.07.2018) US

(71) Applicant: MANUS BIO, INC. [US/US]; 1030 Massachusetts Avenue, Cambridge, MA 02138 (US).

(72) Inventors: KUMARAN, Ajikumar, Parayil; C/o Manus Bio, Inc., 1030 Massachusetts Avenue, Cambridge, MA 02138 (US). SANTOS, Christine, Nicole S.; C/o Manus Bio, Inc., 1030 Massachusetts Avenue, Cambridge, MA 02138 (US). DONALD, Jason, Eric; C/o Manus Bio,


Inc., 1030 Massachusetts Avenue, Cambridge, MA 02138 (US). FOWLER, Mary, Elizabeth; C/o Manus Bio, Inc., 1030 Massachusetts Avenue, Cambridge, MA 02138 (US). PHILIPPE, Ryan, N.; C/o Manus Bio, Inc., 1030 Massachusetts Avenue, Cambridge, MA 02138 (US). FREI, Christopher, Scott; C/o Manus Bio, Inc., 1030 Massachusetts Avenue, Cambridge, MA 02138 (US). LOVE, Aaron; C/o Manus Bio, Inc., 1030 Massachusetts Avenue, Cambridge, MA 02138 (US).

(74) Agent: HAYMAN, Mark, L. et al.; Morgan, Lewis & Bockius LLP, 1111 Pennsylvania Avenue, NW, Washington, DC 20004 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,

(54) Title: PRODUCTION OF STEVIOL GLYCOSIDES THROUGH WHOLE CELL BIOTRANSFORMATION

FIG. 5



(57) Abstract: In various aspects and embodiments, the invention provides microbial cells and methods for producing advanced glycosylation products from lower glycosylated intermediates. The microbial cell expresses one or more UDP-dependent glycosyl transferase enzymes in the cytoplasm, for glycosylation of the intermediates. When incubating the microbial strain with a plant extract or fraction thereof comprising the intermediates, these glycosylated intermediates are available for further glycosylation by the cell, and the advanced glycosylation products can be recovered from the media and/or microbial cells.

[Continued on next page]

OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*