

Cyclopropyl Amines as Modulators of the Histamine H₃ Receptor

FIELD OF THE INVENTION

5 The present invention relates to a series of cyclopropyl amines, their synthesis, and methods for using them to treat disorders and conditions in which the histamine H₃ receptor is involved. As a consequence of these activities, the compounds of the present invention will have therapeutic utility for the treatment of a variety of CNS related disorders including, but not limited
10 to, narcolepsy, sleep disorders, obesity, neurodegenerative disorders, cognitive disorders, and hyperactivity disorders.

BACKGROUND OF THE INVENTION

Histamine {2-(imidazol-4-yl)ethylamine} is a biologically-active molecule. Histamine exerts a physiological effect via multiple distinct G-protein coupled
15 receptors. The histamine H₃ receptor was first described as a presynaptic autoreceptor in the central nervous system (CNS) (Arrang, J.-M. et al., Nature 1983, 302, 832-837) controlling the synthesis and release of histamine. Evidence has emerged showing that H₃ receptors are also located presynaptically as heteroreceptors on serotonergic, noradrenergic,
20 dopaminergic, cholinergic, and GABAergic (gamma-aminobutyric acid containing) neurons. These H₃ receptors have also recently been identified in peripheral tissues such as vascular smooth muscle. Consequently, there are many potential therapeutic applications for histamine H₃ agonists, antagonists, and inverse agonists. (See: "The Histamine H₃ Receptor-A Target for New
25 Drugs", Leurs, R. and Timmerman, H., (Eds.), Elsevier, 1998; Morisset, S. et al., Nature 2000, 408, 860-864.)

Several indications for histamine H₃ antagonists and inverse agonists have similarly been proposed based on animal pharmacology and other experiments with known histamine H₃ antagonists (e.g. thioperamide). These
30 include dementia, Alzheimer's disease (Panula, P. et al., Soc. Neurosci. Abstr. 1995, 21, 1977), epilepsy (Yokoyama, H. et al., Eur. J. Pharmacol. 1993, 234, 129-133), narcolepsy, with or without associated cataplexy, cataplexy,

disorders of sleep/wake homeostasis, idiopathic somnolence, excessive daytime sleepiness (EDS), circadian rhythm disorders, sleep/fatigue disorders, fatigue, drowsiness associated with sleep apnea, sleep impairment due to perimenopausal hormonal shifts, jet lag, Parkinson's-related fatigue, multiple sclerosis (MS)-related fatigue, depression-related fatigue, chemotherapy-induced fatigue, eating disorders (Machidori, H. et al., *Brain Res.* 1992, 590, 180-186), motion sickness, vertigo, attention deficit hyperactivity disorders (ADHD), learning and memory (Barnes, J.C. et al., *Soc. Neurosci. Abstr.* 1993, 19, 1813), and schizophrenia (Schlicker, E. and Marr, I., *Naunyn-Schmiedeberg's Arch. Pharmacol.* 1996, 353, 290-294). (Also see: Stark, H. et al., *Drugs Future* 1996, 21(5), 507-520; and Leurs, R. et al., *Prog. Drug Res.* 1995, 45, 107-165 and references cited therein.) Histamine H₃ antagonists, alone or in combination with a histamine H₁ antagonist, are reported to be useful for the treatment of upper airway allergic responses (U.S. Patent Nos. 5,217,986; 5,352,707 and 5,869,479). A more recent review of this topic was presented by Tozer and Kalindjian (*Exp. Opin. Ther. Patents* 2000, 10, 1045). For additional reviews, see: Celanire, S. *Drug Discovery Today* 2005, 10(23/24), 1613-1627; Hancock, A.A. *Biochem. Pharmacol.* 2006, 71, 1103-1113.

The compounds of the present invention display potency at the human H₃ receptor as determined by receptor binding to the human histamine H₃ receptor (see Lovenberg, T.W. et al., *Mol. Pharmacol.* 1999, 55, 1101-1107). Screening using the human receptor is particularly important for the identification of new therapies for the treatment of human disease.

Conventional binding assays, for example, are determined using rat synaptosomes (Garbarg, M. et al., *J. Pharmacol. Exp. Ther.* 1992, 263(1), 304-310), rat cortical membranes (West, R.E. et al., *Mol. Pharmacol.* 1990, 38, 610-613), and guinea pig brain (Korte, A. et al., *Biochem. Biophys. Res. Commun.* 1990, 168(3), 979-986). Only limited studies have been performed previously using human tissue or the human receptor, but these indicate significant differences in the pharmacology of rodent and primate receptors (West, R.E. et al. *Eur. J. Pharmacol.* 1999, 377, 233-239; Ireland, D. et al. *Eur. J. Pharmacol.* 2001, 433, 141-150).

To achieve a desired pharmacological effect, a compound must display potency against the biological target, as well as a suitable pharmacokinetic profile. First, the compound must be able to travel to its site of action, whether in the CNS, requiring adequate permeation of the blood-brain barrier, or in the periphery. Absorption through various biological membranes is dependent on the physical properties of the drug (degree of ionization at physiological pH, partition coefficient, molecular size, among other factors). Once the desired pharmacological effect is produced, a drug must be eliminated from the organism at a suitable rate. Where an elimination process is too slow, an accumulation of the drug can occur, potentially causing undesirable side effects.

Various H₃-mediated diseases may require compounds with distinct and different pharmacokinetic profiles. In particular, administration of a compound with a short half-life provides greater control over exposure and duration of action of the drug, which may be advantageous in treating or preventing a particular disease or condition. A compound with such an optimized profile allows for the use of tailored formulations, dosing regimens, and/or delivery strategies to accomplish these results. For example, a compound with an attenuated pharmacokinetic profile may produce a shorter pharmacodynamic effect, which may be preferable in treating certain disease states. In contrast, a compound with a long half-life may be preferred for conditions in which constant occupancy of the target by the drug, with no or only very minor changes in drug concentration, are desirable.

Various piperazinyl benzamides were disclosed in U.S. Patent Appl. Publication No. US-2004-0110746-A1 (June 10, 2004), which is hereby incorporated by reference.

The features and advantages of the invention are apparent to one of ordinary skill in the art. Based on this disclosure, including the summary, detailed description, background, examples, and claims, one of ordinary skill in the art will be able to make modifications and adaptations to various conditions and usages. Publications described herein are incorporated by reference in their entirety.

Described herein is a series of N-cyclopropyl amine compounds with the ability to modulate the activity of the histamine receptor, specifically the H₃ receptor.

SUMMARY OF THE INVENTION

5 The invention features a compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-10 4-ylmethyl)-phenyl]-methanone, and enantiomers, hydrates, solvates, and pharmaceutically acceptable salts thereof.

In particular embodiments, the compound is (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone citrate salt dihydrochloride or (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone 15 citrate salt. In further embodiments, the compound is (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone citrate salt, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone citrate salt, or (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone citrate salt.

20 The present invention provides methods of treating or preventing diseases and conditions mediated by histamine H₃ receptor activity. The invention also features pharmaceutical compositions containing such compounds and methods of using such compositions in the treatment or prevention of diseases mediated by histamine H₃ receptor activity. The 25 present invention also contemplates a method of treating or preventing a disease or condition in which histamine is involved with a combination therapy of compounds of the present invention administered with any of the following: histamine H₁ antagonists, histamine H₂ antagonists, neurotransmitter re-uptake blockers, selective serotonin re-uptake inhibitors (SSRIs), serotonin- 30 norepinephrine reuptake inhibitors, noradrenergic reuptake inhibitors, non-selective serotonin-, dopamine- or norepinephrine- re-uptake inhibitors, modafinil, and topiramate.

The pK_a of a given compound affects the degree of ionization at physiological pH. As unionized forms are more lipophilic, they can penetrate membranes, including the blood-brain barrier, more readily. Although calculated pK_a values for the a cyclopropyl amine compound of the present invention (Example 1) were similar to that obtained for an isopropyl analog (Comparative Example 1), the measured pK_a 's differed greatly. Although it has been suggested that a cyclopropyl amine is approximately ten times less basic than aliphatic amines (Zaragoza, et al. J. Med. Chem. 2004, 47, 2833-2838), no direct experimental evidence has been reported (Love, et al. J. Am. Chem. Soc. 1968, 90(10), 2455-2462). Therefore, replacement of a cyclopropyl group for an aliphatic one may produce a higher fraction of non-protonated amine at physiological pH and, thus, serve to improve permeability of membranes (Zaragoza, et al. J. Med. Chem. 2005, 48, 306-311). These data indicate that cyclopropyl amines may tend toward a greater permeability and volume of distribution (V_d) than aliphatic amines of similar structure.

The present invention provides experimental evidence demonstrating a significant decrease in basicity for Example 1, a cyclopropyl amine, relative to its isopropyl and cyclobutyl amine analogs. However, cyclopropyl amines of the present invention actually showed a pharmacokinetic profile that contrasted with the results predicted in the literature. For example, Example 1B displayed a shorter half-life ($T_{1/2}$) and lower volume of distribution (V_d) than Comparative Examples 1B and 2B. Similarly, Example 2B displayed a shorter $T_{1/2}$ and lower V_d than Comparative Examples 3B and 4B; Example 3B displayed a shorter $T_{1/2}$ and lower V_d than Comparative Example 5B, and Example 4B displayed a shorter $T_{1/2}$ and lower V_d than Comparative Example 6B.

Additional features and advantages of the invention will become apparent from the detailed description and examples below, and the appended claims.

DETAILED DESCRIPTION OF THE INVENTION

The invention may be more fully appreciated by reference to the following description, including the following glossary of terms and the concluding examples. For the sake of brevity, the disclosures of the publications cited in this specification are herein incorporated by reference.

As used herein, the terms "including", "containing" and "comprising" are used herein in their open, non-limiting sense.

Isomeric forms of the compounds of the present invention, and of their pharmaceutically acceptable salts, are encompassed within the present invention, and reference herein to one of such isomeric forms is meant to refer to at least one of such isomeric forms. One of ordinary skill in the art will recognize that compounds according to this invention may exist, for example in a single isomeric form whereas other compounds may exist in the form of an isomeric mixture. For example, the present invention encompasses optical isomers of the compounds described herein, including enantiomers and mixtures thereof. In addition, certain compounds referred to herein can exist in solvated or hydrated forms as well as unsolvated forms. It is understood that this invention encompasses all such solvated and unsolvated forms of the compounds of this invention.


Compounds according to the present invention that have been modified to be detectable by some analytic technique are also within the scope of this invention. The compounds of the present invention may be labeled with radioactive elements such as ^{125}I , ^{18}F , ^{11}C , ^{64}Cu , ^3H , ^{14}C , and the like for use in imaging or for radioactive treatment of patients. An example of such compounds is an isotopically labeled compound, such as an ^{18}F isotopically labeled compound that may be used as a probe in detection and/or imaging techniques, such as positron emission tomography (PET) and single-photon emission computed tomography (SPECT). Preferably, compounds of the present invention labeled with ^{18}F or ^{11}C may be used as a positron emission tomography (PET) molecular probe for studying histamine-mediated disorders. Alternatively, compounds of the present invention labeled with ^{14}C may be used in metabolic studies. Another example of such compounds is an isotopically labeled compound, such as a deuterium and/or tritium labeled compound, that may be used in reaction kinetic studies. The compounds described herein may be reacted with an appropriate functionalized radioactive reagents using conventional chemistry to provide radiolabeled compounds.

The compounds as described above may be made according to processes within the skill of the art and/or that are described in the schemes

and examples that follow. To obtain the various compounds herein, starting materials may be employed that carry the ultimately desired substituents though the reaction scheme with or without protection as appropriate. This may be achieved by means of conventional protecting groups, such as those
5 described in "Protective Groups in Organic Chemistry", ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, "Protective Groups in Organic Synthesis", 3rd ed., John Wiley & Sons, 1999. The protecting groups may be removed at a convenient subsequent stage using methods known from the art. Alternatively, it may be necessary to employ, in the place of the
10 ultimately desired substituent, a suitable group that may be carried through the reaction scheme and replaced as appropriate with the desired substituent. Such compounds, precursors, or prodrugs are also within the scope of the invention. Reactions may be performed between the melting point and the reflux temperature of the solvent, and preferably between 0 °C and the reflux
15 temperature of the solvent.

The compounds as described above may be made according to Schemes A-C below. Persons skilled in the art will recognize that certain compounds are more advantageously produced by one scheme as compared to the other. In addition, synthetic sequences described in U.S. Pat. Appl. No.
20 10/690,115 are hereby incorporated by reference and may be applied to the preparation of compounds of the present invention. One skilled in the art will recognize that compounds of Formula (I) where where R¹R²N- is morpholinyl, 4-fluoropiperidinyl, thiomorpholinyl, or 2-hydroxymethyl-morpholin-4-yl are compounds of the present invention.


Scheme A



Referring to Scheme A, amines of formula (X), where PG is cyclopropyl or a suitable protecting group such as a benzyl or tert-butylcarbamoyl (Boc), may be coupled with benzoic acids of formula (XI), either through activation of the acid to the acid chloride or acid fluoride followed by reaction with the amine, or directly under peptide coupling conditions, such as 1,1'-carbonyldiimidazole (CDI) or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC)/1-hydroxybenzotriazole (HOBt). Preferably, reactions are run with EDC/HOBt in the presence of 4-(dimethylamino)pyridine (DMAP), in a solvent such as CH_2Cl_2 . Benzaldehydes of formula (XII) may then be reacted with suitable amines $\text{R}^1\text{NR}^2\text{H}$ (where $\text{R}^1\text{NR}^2\text{H}$ is morpholine, 4-fluoropiperidine, thiomorpholine, or morpholin-2-yl-methanol) under reductive amination conditions to provide benzyl amines (XIII). Suitable reducing agents include NaCNBH_3 or $\text{NaB(OAc)}_3\text{H}$ in a solvent such as methanol or dichloroethane. Preferred conditions include $\text{NaB(OAc)}_3\text{H}$ in methanol. The protecting group "PG" may then be removed under standard deprotection conditions to provide amines of formula (XIV). Where PG is Boc, deprotection may be effected using HCl in 1,4-dioxane or trifluoroacetic acid (TFA) in CH_2Cl_2 . Amines (XV) are converted to the corresponding cyclopropyl amines of Formula (I) through reaction with [(1-methoxycyclopropyl)oxy]trimethylsilane or [(1-ethoxycyclopropyl)oxy]-trimethylsilane (R is methyl or ethyl) under conditions


similar to those described in J. Med. Chem. 2004, 47(11), 2733-2738 and Tetrahedron Lett. 1995, 36(41), 7399-7402.

Scheme B



- 5 Alternatively, compounds of the present invention may be prepared as in Schemes B and C. To this end, protected heterocycle (X), where PG is as defined previously, may be reacted with [(1-methoxycyclopropyl)oxy]-trimethylsilane or [(1-ethoxycyclopropyl)oxy]trimethylsilane as described in J. Med. Chem. 2004, 47(11), 2733-2738 and Tetrahedron Lett. 1995, 36(41), 7399-7402. Preferably, PG is a Boc group. The group "PG" may then be removed under standard deprotection conditions to provide cyclopropyl amines of formula (XVI). Where PG is Boc, preferred conditions include a mixture of HCl in a solvent such as 1,4-dioxane.

Scheme C



- 15 Acids of formula (XI) may be coupled (through activation or directly as described in Scheme A) with amines (XVI) to form amides (XVII). Reductive amination with a suitable amine as described in Scheme A gives rise to compounds of Formula (I).

- 20 Compounds of the present invention may be converted to their corresponding salts using methods known to those skilled in the art. For example, free base forms of compounds of the present invention may be

treated with TFA, HCl, or citric acid in a solvent such as methanol (MeOH) or ethanol (EtOH) to provide the corresponding salt forms.

Compounds prepared according to the schemes described above may be obtained as single enantiomers or diastereomers, or as racemic mixtures or
5 mixtures of enantiomers or diastereomers. Where such mixtures are obtained, isomers may be separated using conventional methods such as chromatography or crystallization. Where racemic (1:1) and non-racemic (not 1:1) mixtures of enantiomers are obtained, single enantiomers may be isolated using conventional separation methods known to one skilled in the art.
10 Particularly useful separation methods may include chiral chromatography, recrystallization, resolution, diastereomeric salt formation, or derivatization into diastereomeric adducts followed by separation.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional
15 derivatives of the compounds that are readily convertible *in vivo* into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various disorders described with a compound of the present invention or with a compound that converts to a compound of the present invention *in vivo* after
20 administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985. In addition to salts, the invention provides the esters, amides, and other protected or derivatized forms of the described compounds.

25 For therapeutic use, salts of the compounds of the present invention are those that are pharmaceutically acceptable. However, salts of acids and bases that are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not, are included within the
30 ambit of the present invention.

Pharmaceutically acceptable salts of compounds according to the present invention refer to those salt forms of the compounds of the present invention which would be apparent to the pharmaceutical chemist, *i.e.*, those

that are non-toxic and that would favorably affect the pharmacokinetic properties of said compounds of the present invention, such as sufficient palatability, absorption, distribution, metabolism and excretion. Other factors, more practical in nature, which are also important in the selection, are cost of raw materials, ease of crystallization, yield, stability, hygroscopicity and flowability of the resulting bulk drug.

Examples of acids that may be used in the preparation of pharmaceutically acceptable salts include the following: acetic acid, 2,2-dichloroacetic acid, acylated amino acids, adipic acid, alginic acid, ascorbic acid, L-aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, boric acid, (+)-camphoric acid, camphorsulfonic acid, (+)-(1*S*)-camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, cinnamic acid, citric acid, cyclamic acid, cyclohexanesulfamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxy-ethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, D-gluconic acid, D-glucuronic acid, L-glutamic acid, α -oxo-glutaric acid, glycolic acid, hippuric acid, hydrobromic acid, hydrochloric acid, hydroiodic acid, (+)-L-lactic acid, ($\pm$)-DL-lactic acid, lactobionic acid, lauric acid, maleic acid, (-)-L-malic acid, malonic acid, ($\pm$)-DL-mandelic acid, methanesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, nitric acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, perchloric acid, phosphoric acid, L-pyroglutamic acid, saccharic acid, salicylic acid, 4-amino-salicylic acid, sebacic acid, stearic acid, succinic acid, sulfuric acid, tannic acid, (+)-L-tartaric acid, thiocyanic acid, *p*-toluenesulfonic acid, undecylenic acid, and valeric acid.

Compounds of the present invention containing acidic protons may be converted into their therapeutically active non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts; the alkali and earth alkaline metal salts (e.g. lithium, sodium, potassium, magnesium, calcium salts, which may be prepared by treatment with, for example, magnesium hydroxide, calcium hydroxide, potassium hydroxide, zinc hydroxide, or sodium hydroxide); and amine salts made with organic bases (e.g. primary, secondary

and tertiary aliphatic and aromatic amines such as L-arginine, benethamine, benzathine, choline, deanol, diethanolamine, diethylamine, dimethylamine, dipropylamine, diisopropylamine, 2-(diethylamino)-ethanol, ethanolamine, ethylamine, ethylenediamine, isopropylamine, N-methyl-glucamine, 5 hydrabamine, 1*H*-imidazole, L-lysine, morpholine, 4-(2-hydroxyethyl)-morpholine, methylamine, piperidine, piperazine, propylamine, pyrrolidine, 1-(2-hydroxyethyl)-pyrrolidine, pyridine, quinuclidine, quinoline, isoquinoline, secondary amines, triethanolamine, trimethylamine, triethylamine, *N*-methyl-D-glucamine, 2-amino-2-(hydroxymethyl)-1,3-propanediol, and tromethamine). 10 See, e.g., S.M. Berge, *et al.*, "Pharmaceutical Salts", J. Pharm. Sci., 1977, 66:1-19, and *Handbook of Pharmaceutical Salts, Properties, Selection, and Use*; Stahl, P.H., Wermuth, C.G., Eds.; Wiley-VCH and VHCA: Zurich, 2002, which are incorporated herein by reference.

The compounds of the present invention are modulators of the 15 histamine H₃ receptor, and as such, the compounds are useful in the treatment of disease states mediated by histamine H₃ receptor activity. Thus, the compounds of the present invention may be used in a method of treating a subject suffering from or diagnosed with a disease mediated by histamine H₃ receptor activity, comprising administering to a subject in need of such 20 treatment an effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof. Particularly, the compounds may be used in methods for treating or preventing neurologic or neuropsychiatric disorders including sleep/wake and arousal/vigilance disorders (e.g. insomnia and jet lag), attention deficit hyperactivity disorders (ADHD), learning and 25 memory disorders, cognitive dysfunction, migraine, neurogenic inflammation, dementia, mild cognitive impairment (pre-dementia), Alzheimer's disease, epilepsy, narcolepsy with or without associated cataplexy, cataplexy, disorders of sleep/wake homeostasis, idiopathic somnolence, excessive daytime sleepiness (EDS), circadian rhythm disorders, sleep/fatigue disorders, fatigue, 30 drowsiness associated with sleep apnea, sleep impairment due to perimenopausal hormonal shifts, Parkinson's-related fatigue, MS-related fatigue, depression-related fatigue, chemotherapy-induced fatigue, eating disorders, obesity, motion sickness, vertigo, schizophrenia, substance abuse,

bipolar disorders, manic disorders and depression, as well as other disorders in which the histamine H₃ receptor is involved, such as upper airway allergic response, asthma, itch, nasal congestion and allergic rhinitis in a subject in need thereof. For example, the invention features methods for preventing,
5 inhibiting the progression of, or treating upper airway allergic response, asthma, itch, nasal congestion and allergic rhinitis. Excessive daytime sleepiness (EDS) may occur with or without associated sleep apnea, shift work, fibromyalgia, MS, and the like.

The compounds of the present invention may be used in methods for
10 treating or preventing disease states selected from the group consisting of: cognitive disorders, sleep disorders, psychiatric disorders, and other disorders.

Cognitive disorders include, for example, dementia, Alzheimer's disease (Panula, P. et al., Soc. Neurosci. Abstr. 1995, 21, 1977), cognitive dysfunction, mild cognitive impairment (pre-dementia), attention deficit hyperactivity
15 disorders (ADHD), attention-deficit disorders, and learning and memory disorders (Barnes, J.C. et al., Soc. Neurosci. Abstr. 1993, 19, 1813). Learning and memory disorders include, for example, learning impairment, memory impairment, age-related cognitive decline, and memory loss. H₃ antagonists have been shown to improve memory in a variety of memory tests, including
20 the elevated plus maze in mice (Miyazaki, S. et al. *Life Sci.* 1995, 57(23), 2137-2144), a two-trial place recognition task (Orsetti, M. et al. *Behav. Brain Res.* 2001, 124(2), 235-242), the passive avoidance test in mice (Miyazaki, S. et al. *Meth. Find. Exp. Clin. Pharmacol.* 1995, 17(10), 653-658) and the radial maze in rats (Chen, Z. *Acta Pharmacol. Sin.* 2000, 21(10), 905-910). Also, in the
25 spontaneously hypertensive rat, an animal model for the learning impairments in attention-deficit disorders, H₃ antagonists were shown to improve memory (Fox, G.B. et al. *Behav. Brain Res.* 2002, 131(1-2), 151-161).

Sleep disorders include, for example, insomnia, disturbed sleep, narcolepsy (with or without associated cataplexy), cataplexy, disorders of
30 sleep/wake homeostasis, idiopathic somnolence, excessive daytime sleepiness (EDS), circadian rhythm disorders, fatigue, lethargy, REM-behavioral disorder, and jet lag. Fatigue and/or sleep impairment may be caused by or associated with various sources, such as, for example, sleep apnea, perimenopausal

hormonal shifts, Parkinson's disease, multiple sclerosis (MS), depression, chemotherapy, or shift work schedules.

Psychiatric disorders include, for example, schizophrenia (Schlicker, E. and Marr, I., Naunyn-Schmiedeberg's Arch. Pharmacol. 1996, 353, 290-294),
5 bipolar disorders, manic disorders, depression (Lamberti, C. et al. *Br. J. Pharmacol.* 1998, 123(7), 1331-1336; Perez-Garcia, C. et al. *Psychopharmacology* 1999, 142(2), 215-220) (Also see: Stark, H. et al., *Drugs Future* 1996, 21(5), 507-520; and Leurs, R. et al., *Prog. Drug Res.* 1995, 45, 107-165 and references cited therein.), obsessive-compulsive disorder, and
10 post-traumatic stress disorder.

Other disorders include, for example, motion sickness, vertigo (including vertigo and benign postural vertigo), tinnitus, epilepsy (Yokoyama, H. et al., *Eur. J. Pharmacol.* 1993, 234, 129-133), migraine, neurogenic inflammation, eating disorders (Machidori, H. et al., *Brain Res.* 1992, 590, 180-186), obesity,
15 substance abuse disorders, movement disorders (e.g. restless leg syndrome), and eye-related disorders (e.g. macular degeneration and retinitis pigmentosa).

Said methods of treating and preventing comprise the step of administering to a mammal suffering therefrom an effective amount of at least
20 one compound of the present invention.

The present invention also contemplates a method of treating or preventing a histamine-mediated disease or condition with a combination therapy for the treatment of allergic rhinitis, nasal congestion, and allergic congestion, comprising: a) administering an effective amount of at least one
25 compound of the present invention, and b) administering an effective amount of one or more histamine H₁ or H₂ antagonists. Suitable histamine H₁ antagonists include: loratidine (CLARITIN™), desloratidine (CLARINEX™), fexofenadine (ALLEGRA™) and cetirizine (ZYRTEC™).

The present invention also contemplates a method of treating or
30 preventing a histamine-mediated disease or condition with a combination therapy for the treatment of depression, mood disorders or schizophrenia, comprising: a) administering an effective amount of at least one compound of the present invention, and b) administering an effective amount of one or more

neurotransmitter re-uptake blockers. Suitable neurotransmitter re-uptake blockers include: selective serotonin re-uptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors, noradrenergic reuptake inhibitors, or non-selective serotonin-, dopamine- or norepinephrine re-uptake inhibitors.

- 5 Particular examples of neurotransmitter re-uptake blockers include fluoxetine (PROZAC™), sertraline (ZOLOFT™), paroxetine (PAXIL™), and amitriptyline.

The present invention also contemplates a method of treating or preventing a histamine-mediated disease or condition with a combination therapy for the treatment of narcolepsy, excessive daytime sleepiness (EDS),
10 Alzheimer's disease, depression, attention deficit disorders, MS-related fatigue, post-anesthesia grogginess, cognitive impairment, schizophrenia, spasticity associated with cerebral palsy, age-related memory decline, idiopathic somnolence, or jet-lag, comprising: a) administering an effective amount of at least one compound of the present invention, and b) administering an effective
15 amount of modafinil.

In another embodiment, the present invention contemplates a method of treating or preventing a histamine-mediated disease or condition with a combination therapy comprising: a) administering an effective amount of at least one compound of the present invention, and b) administering an effective
20 amount of topiramate (Topamax). In particular, such methods are useful for the treatment of obesity. Preferably, the combination method employs doses of topiramate in the range of about 20 to 300 mg per dose.

Compounds of the present invention may be administered in pharmaceutical compositions to treat patients (humans and other mammals)
25 with disorders mediated by the H₃ receptor. Thus, the invention features pharmaceutical compositions containing at least one compound of the present invention and a pharmaceutically acceptable carrier. A composition of the invention may further include at least one other therapeutic agent such as H₁ antagonists, SSRIs, topiramate, or modafinil (for example, a combination
30 formulation or combination of differently formulated active agents for use in a combination therapy method).

The present invention also features methods of using or preparing or formulating such pharmaceutical compositions. The pharmaceutical

compositions can be prepared using conventional pharmaceutical excipients and compounding techniques known to those skilled in the art of preparing dosage forms. It is anticipated that the compounds of the invention can be administered by oral, parenteral, rectal, topical, or ocular routes, or by
5 inhalation. Preparations may also be designed to give slow release of the active ingredient. The preparation may be in the form of tablets, capsules, sachets, vials, powders, granules, lozenges, powders for reconstitution, liquid preparations, or suppositories. Preferably, compounds may be administered by intravenous infusion or topical administration, but more preferably by oral
10 administration.

For oral administration, the compounds of the invention can be provided in the form of tablets or capsules, or as a solution, emulsion, or suspension. Tablets for oral use may include the active ingredient mixed with pharmaceutically acceptable excipients such as inert diluents, disintegrating
15 agents, binding agents, lubricating agents, sweetening agents, flavoring agents, coloring agents and preservatives. Suitable inert fillers include sodium and calcium carbonate, sodium and calcium phosphate, lactose, starch, sugar, glucose, methyl cellulose, magnesium stearate, mannitol, sorbitol, and the like; typical liquid oral excipients include ethanol, glycerol, water and the like.
20 Starch, polyvinyl-pyrrolidone, sodium starch glycolate, microcrystalline cellulose, and alginic acid are suitable disintegrating agents. Binding agents may include starch and gelatin. The lubricating agent, if present, will generally be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or glyceryl distearate to
25 delay absorption in the gastrointestinal tract, or may be coated with an enteric coating. Capsules for oral use include hard gelatin capsules in which the active ingredient is mixed with a solid, semi-solid, or liquid diluent, and soft gelatin capsules wherein the active ingredient is mixed with water, an oil such as peanut oil or olive oil, liquid paraffin, a mixture of mono and di-glycerides of
30 short chain fatty acids, polyethylene glycol 400, or propylene glycol.

Liquids for oral administration may be suspensions, solutions, emulsions or syrups or may be presented as a dry product for reconstitution with water or other suitable vehicles before use. Compositions of such liquid may contain

pharmaceutically-acceptable excipients such as suspending agents (for example, sorbitol, methyl cellulose, sodium alginate, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel and the like); non-aqueous vehicles, which include oils (for example, almond oil or fractionated coconut oil), propylene glycol, ethyl alcohol or water; preservatives (for example, methyl or propyl p-hydroxybenzoate or sorbic acid); wetting agents such as lecithin; and, if needed, flavoring or coloring agents.

The compounds of this invention may also be administered by non-oral routes. The compositions may be formulated for rectal administration as a suppository. For parenteral use, including intravenous, intramuscular, intraperitoneal, or subcutaneous routes, the compounds of the invention will generally be provided in sterile aqueous solutions or suspensions, buffered to an appropriate pH and isotonicity or in parenterally acceptable oil. Suitable aqueous vehicles include Ringer's solution and isotonic sodium chloride. Such forms will be presented in unit dose form such as ampules or disposable injection devices, in multi-dose forms such as vials from which the appropriate dose may be withdrawn, or in a solid form or pre-concentrate that can be used to prepare an injectable formulation. Another mode of administration of the compounds of the invention may utilize a patch formulation to affect transdermal delivery. The compounds of this invention may also be administered by inhalation, via the nasal or oral routes using a spray formulation consisting of the compound of the invention and a suitable carrier.

Methods are known in the art for determining effective doses for therapeutic (treatment) and prophylactic (preventative) purposes for the pharmaceutical compositions or the drug combinations of the present invention, whether or not formulated in the same composition. The specific dosage level required for any particular patient will depend on a number of factors, including severity of the condition being treated, the route of administration, metabolic rate, and the weight of the patient. For therapeutic purposes, "effective dose" or "effective amount" refers to that amount of each active compound or pharmaceutical agent, alone or in combination, that elicits the biological or medicinal response in a tissue system, animal, or human that is being sought by a researcher, veterinarian, medical doctor, or other clinician,

which includes alleviation of the symptoms of the disease or disorder being treated. For prophylactic purposes (i.e., preventing or inhibiting the onset or progression of a disorder), the term "effective dose" or "effective amount" refers to that amount of each active compound or pharmaceutical agent, alone or in
5 combination, that inhibits in a subject the onset or progression of a disorder as being sought by a researcher, veterinarian, medical doctor, or other clinician, the delaying of which disorder is mediated, at least in part, by the modulation of the histamine H₃ receptor. Methods of combination therapy include co-administration of a single formulation containing all active agents; essentially
10 contemporaneous administration of more than one formulation; and administration of two or more active agents separately formulated.

It is anticipated that the daily dose (whether administered as a single dose or as divided doses) will be in the range 0.01 to 1000 mg per day, more usually from 1 to 500 mg per day, and most usually from 10 to 200 mg per day.
15 Expressed as dosage per unit body weight, a typical dose will be expected to be between 0.0001 mg/kg and 15 mg/kg, especially between 0.01 mg/kg and 7 mg/kg, and most especially between 0.15 mg/kg and 2.5 mg/kg.

Preferably, oral doses range from about 0.05 to 200 mg/kg, daily, taken in 1 to 4 separate doses. Some compounds of the invention may be orally
20 dosed in the range of about 0.05 to about 50 mg/kg daily, others may be dosed at 0.05 to about 20 mg/kg daily, while still others may be dosed at 0.1 to about 10 mg/kg daily. Infusion doses can range from about 1 to 1000 µg/kg/min of inhibitor, admixed with a pharmaceutical carrier over a period ranging from several minutes to several days. For topical administration compounds of the
25 present invention may be mixed with a pharmaceutical carrier at a concentration of about 0.1% to about 10% of drug to vehicle.

Examples

In order to illustrate the invention, the following examples are included.
30 These examples do not limit the invention. They are only meant to suggest a method of practicing the invention. Those skilled in the art may find other methods of practicing the invention, which are obvious to them. However, those methods are deemed to be within the scope of this invention.


Chemical Examples and Data:

Mass spectra were obtained on an Agilent series 1100 MSD using electrospray ionization (ESI) in either positive or negative modes as indicated.


- 5 Nuclear magnetic resonance (NMR) spectra were obtained on a Bruker model DPX400 (400 MHz), DPX500 (500 MHz), or DPX600 (600 MHz) spectrometer. The format of the ^1H NMR data below is: chemical shift in ppm down field of the tetramethylsilane reference (multiplicity, coupling constant J in Hz, integration).

- 10 Reversed-phase high-pressure liquid chromatography (HPLC) was performed on a Hewlett Packard HPLC, Zorbax Eclipse XDB-C8, 5 μm , 4.6 x 150 mm column, with a gradient of 1 to 99% acetonitrile/water/0.05% TFA over 8 min.

- Chemical names were generated using ChemDraw Ultra 6.0.2
15 (CambridgeSoft Corp., Cambridge, MA).




Example 1; (4-Cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone.




- 20 Step A. 4-(4-Formyl-benzoyl)-piperazine-1-carboxylic acid tert-butyl ester. A suspension of 4-carboxybenzaldehyde (3.10 g) in CH_2Cl_2 was treated sequentially with piperazine-1-carboxylic acid tert-butyl ester (3.6 g), EDC (3.86 g), HOBT (2.68 g), and DMAP (~0.020 g). After 18 h, the mixture was extracted
25 with 1 N NaOH and then with 1 N HCl. The organic layer was dried (Na_2SO_4) and concentrated to give the title compound (5.11 g, 78%). MS (ESI): mass calcd. for $\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}_4$, 318.16; m/z found, 219.3 $[(M-100)+\text{H}]^+$. ^1H NMR

(CDCl₃): 10.04 (s, 1H), 7.93 (d, J = 8.2, 2H), 7.54 (d, J = 8.1, 2H), 3.82-3.67 (m, 2H), 3.58-3.30 (m, 6H), 1.46 (s, 9H).



Step B. 4-(4-Morpholin-4-ylmethyl-benzoyl)-piperazine-1-carboxylic acid tert-

- 5 butyl ester. A solution of 4-(4-formyl-benzoyl)-piperazine-1-carboxylic acid tert-butyl ester (2.06 g) in methanol (100 mL) was treated with morpholine (4 mL) and NaB(OAc)₃H (6.98 g, in portions over 1 h). After 3 h, the mixture was diluted with satd. aq. NaHCO₃ and extracted with CH₂Cl₂. The organic layer was dried (Na₂SO₄) and concentrated. The residue was purified by column
- 10 chromatography (SiO₂) to give the title compound (1.22 g, 48%). MS (ESI): mass calcd. for C₂₁H₃₁N₃O₄, 389.23; m/z found, 390.4 [M+H]⁺. ¹H NMR (CDCl₃): 7.39-7.33 (m, 4H), 3.75-3.66 (m, 6H), 3.50 (s, 2H), 3.51-3.33 (m, 6H), 2.45-2.41 (m, 4H), 1.46 (s, 9H).




- 15 Step C. (4-Morpholin-4-ylmethyl-phenyl)-piperazine-1-yl-methanone. A solution of 4-(4-morpholin-4-ylmethyl-benzoyl)-piperazine-1-carboxylic acid tert-butyl ester (1.163 g) in CH₂Cl₂ (10 mL) was treated with TFA (~4 mL). After 30 min, additional TFA (5 mL) was added, and the mixture was stirred for a further 2 h. The mixture was diluted with satd. aq. NaHCO₃ and extracted with
- 20 CH₂Cl₂. The organic layer was dried (Na₂SO₄) and concentrated. The residue was purified by column chromatography (SiO₂) to give the title compound (0.255 g, 30%). MS (ESI): mass calcd. for C₁₆H₂₃N₃O₂, 289.18; m/z found, 290.4 [M+H]⁺. ¹H NMR (CDCl₃): 7.41-7.35 (m, 4H), 3.95-3.70 (m, 6H), 3.52 (s, 2H), 3.09-2.80 (m, 6H), 2.49-2.42 (m, 4H).


- 25 Step D. A solution of (4-morpholin-4-ylmethyl-phenyl)-piperazine-1-yl-methanone (0.128 g) in methanol (7.5 mL) was treated with (1-ethoxycyclopropoxy)-trimethyl-silane (1.5 mL), acetic acid (0.2 mL), and NaBH₃CN (~400 mg). The mixture was heated at 60 °C for 18 h, and then was cooled to

rt and concentrated. The residue was diluted with 1 N NaOH and extracted with CH₂Cl₂. The organic layer was dried (Na₂SO₄) and concentrated. The residue was purified by column chromatography (SiO₂) to give the title compound (0.0548 g, 38%). MS (ESI): mass calcd. for C₁₉H₂₇N₃O₂, 329.21; m/z found, 330.4 [M+H]⁺. ¹H NMR (CDCl₃): 7.36 (s, 4H), 3.79-3.68 (m, 6H), 3.50 (s, 2H), 3.44-3.32 (m, 2H), 2.74-2.61 (m, 2H), 2.60-2.50 (s, 2H), 2.45-2.40 (m, 4H), 1.66-1.62 (m, 1H), 0.49-0.44 (m, 2H), 0.44-0.39 (m, 2H).

Alternative Preparation of Example 1.




Step A. *tert*-Butyl 4-cyclopropylpiperazine-1-carboxylate. A mixture of *tert*-butyl piperazine-1-carboxylate (75.0 g), tetrahydrofuran (THF) (500 mL), methanol (500 mL), [(1-ethoxycyclopropyl)oxy]trimethylsilane (161 mL), NaBH₃CN (38.0 g), and acetic acid (37 mL) was heated at 60 °C for 5 h. The mixture was cooled to rt, treated with water (30 mL) and stirred for 5 min. The mixture was then treated with 1 N NaOH (130 mL) and was further stirred for 15 min. The mixture was concentrated, and the remaining aqueous solution was extracted with CH₂Cl₂ (500 mL). The organic layer was washed with 1 N NaOH (500 mL). The combined aqueous layers were extracted with CH₂Cl₂ (150 mL). The combined organic layers were washed with brine (400 mL), dried (Na₂SO₄), and concentrated to give the title compound as a white solid (92 g, 100%). MS (ESI): mass calcd. for C₁₂H₂₂N₂O₂, 226.17; m/z found, 227.2 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): 3.39 (t, *J* = 5.0 Hz, 4H), 2.55 (t, *J* = 4.9 Hz, 4H), 1.60 (ddd, *J* = 10.3, 6.5, 3.8 Hz, 1H), 1.46 (s, 9H), 0.49-0.38 (m, 4H).



Step B. 1-Cyclopropylpiperazine dihydrochloride. A solution of *tert*-butyl 4-cyclopropylpiperazine-1-carboxylate (92 g) in 1,4-dioxane (200 mL) was treated with HCl (4 M in 1,4-dioxane, 500 mL) over 10 min while maintaining the temperature below 40 °C. After the addition was complete, the mixture was


heated at 45 °C for 9 h and then was cooled to rt. The thick suspension was diluted with hexanes (400 mL) and was cooled to 10 °C. The resulting solid was collected by filtration, washed with hexanes, and dried to give the title compound as a white solid (78 g, 96%). MS (ESI): mass calcd. for C₇H₁₄N₂, 126.12; m/z found, 127.0 [M+H⁺]. ¹H NMR (400 MHz, D₂O): 3.65 (br t, *J* = 4.7 Hz, 4H), 3.47 (br t, *J* = 5.5 Hz, 4H), 2.85 (br quintet, *J* = 5.8 Hz, 1H), 0.94 (br s, 2H), 0.92 (br s, 2H).



Step C. 4-(4-Cyclopropyl-piperazine-1-carbonyl)-benzaldehyde. A mixture of 4-formyl-benzoic acid (54.4 g), toluene (500 mL), N,N-dimethylformamide (DMF) (3.6 mL), and thionyl chloride (30.4 mL) was heated at 60 °C for 2 h and then was cooled to 5 °C. In a separate flask, a 5 °C mixture of NaOH (50.7 g), water (550 mL), and toluene (150 mL) was treated with 1-cyclopropyl-piperazine dihydrochloride (70.0 g) in portions while the temperature was maintained below 10 °C. After the addition was complete, the mixture was cooled to 5 °C and treated with the crude acyl chloride solution prepared as above at a rate such that the temperature did not exceed 10 °C. After the addition was complete, the mixture was allowed to warm to rt and was stirred overnight. The biphasic mixture was basified to pH ~10 with 1 N NaOH (300 mL). The layers were separated and the aqueous layer was extracted with toluene (100 mL x 2). The combined organic layers were washed with brine (200 mL), dried (Na₂SO₄), and concentrated to give the title compound as pale yellow viscous oil (56.0 g, 62%). HPLC: R_T = 5.19 min. MS (ESI): mass calcd. for C₁₅H₁₈N₂O₂, 258.14; m/z found, 258.9 [M+H⁺]. ¹H NMR (400 MHz, CDCl₃): 10.1 (s, 1H), 7.94 (pseudo d, *J* = 8.2 Hz, 2H), 7.56 (pseudo d, *J* = 8.1 Hz, 2H), 3.77 (br s, 2H), 3.33 (br s, 2H), 2.71 (br s, 2H), 2.55 (br s, 2H), 1.66 (ddd, *J* = 10.2, 6.6, 3.7 Hz, 1H), 0.52-0.46 (m, 2H), 0.45-0.40 (br s, 2H).

Step D. (4-Cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone. To a solution of 4-(4-cyclopropyl-piperazine-1-carbonyl)-benzaldehyde (56.0 g) in 1,2-dichloroethane (550 mL) was added morpholine

(37.8 mL) dropwise over 5 min. The mixture was cooled to 10 °C and was treated with NaB(OAc)₃H (64.3 g) in portions over 1 h. After a further 2 h, the mixture was warmed to rt, and a water bath was used to keep the temperature below 20 °C. After 18 h, water (60 mL) was added while the temperature was kept under 20 °C by the addition of small amounts of ice. After 20 min, the mixture was basified to pH ~10 with 1 N NaOH (450 mL) and the mixture was stirred for 10 min. The layers were separated, and the organic layer was washed with 1 N NaOH (150 mL). The combined aqueous layers were extracted with CH₂Cl₂ (200 mL). The combined organic layers were washed with brine (200 mL), dried (Na₂SO₄), and concentrated to give the title compound as pale yellow viscous oil (68.0 g, 95%). HPLC: R_T = 4.39 min. MS (ESI): mass calcd. for C₁₉H₂₇N₃O₂, 329.21; m/z found, 330.2 [M+H⁺]. ¹H NMR (400 MHz, CDCl₃): 7.35 (br s, 4H), 3.73 (br s, 2H), 3.69 (t, *J* = 4.6 Hz, 4H), 3.50 (s, 2H), 3.37 (br s, 2H), 2.67 (br s, 2H), 2.53 (br s, 2H), 2.43 (t, *J* = 4.2 Hz, 4H), 1.63 (ddd, *J* = 10.3, 6.7, 3.7 Hz, 1H), 0.49-0.43 (m, 2H), 0.42-0.39 (br s, 2H). ¹³C NMR (101 MHz, CDCl₃): 170.6, 140.0, 135.1, 129.5, 127.5, 67.4, 63.4, 54.0, 38.7, 6.3.



Example 1A; (4-Cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone dihydrochloride.


A solution of (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone (68.0 g) in ethanol (400 mL) was heated to 60 °C and treated with conc. HCl (37.8 mL) dropwise over 40 min. A precipitate started to form after ~20 mL of HCl had been added. After the addition was complete, the thick suspension was slowly cooled to 20 °C over 3 h. The solid was collected by filtration, washed with ethanol, and dried at 50 °C overnight in a vacuum oven to provide the title compound as a white solid (56.2 g, 68%). HPLC: R_T = 4.30 min. MS (ESI): mass calcd. for C₁₉H₂₇N₃O₂, 329.21; m/z found, 330.0 [M+H⁺]. ¹H NMR (400 MHz, D₂O): 7.64 (pseudo d, *J* = 8.3 Hz, 2H), 7.58 (pseudo d, *J* =

8.3 Hz, 2H), 4.44 (br s, 2H), 4.20-3.10 (m, 16H), 2.88 (ddd, $J = 11.2, 6.6, 4.8$ Hz, 1H), 1.03-0.98 (m, 4H). ^{13}C NMR (101 MHz, D_2O): 172.1, 135.3, 132.2, 130.9, 128.0, 64.0, 60.5, 52.6, 52.4, 51.7, 44.8, 39.7, 39.5, 3.9.

- 5 Example 1B; (4-Cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone citrate salt.

A mixture of (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone (0.476 g, 1.45 mmol) and citric acid (0.281 g, 1.46 mmol) was diluted with methanol (~10 mL). The mixture was heated until homogeneous
10 and then concentrated. The resulting oil was triturated with ethyl acetate and the solid material that formed was dried under vacuum to yield the citrate salt (0.760 g).


- The compounds in Examples 2-4 were prepared using methods
15 analogous to those described in the preceding examples, with the appropriate substituent changes.



Example 2; (4-Cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone.

- 20 MS (ESI): mass calcd. for $\text{C}_{20}\text{H}_{28}\text{FN}_3\text{O}$, 345.22; m/z found, 346.4 $[\text{M}+\text{H}^+]$. ^1H NMR (400 MHz, CDCl_3): 7.39-7.33 (m, 4H), 4.78-4.58 (m, 1H), 3.82-3.66 (m, 2H), 3.51 (s, 2H), 3.46-3.33 (m, 2H), 2.77-2.49 (m, 6H), 2.43-2.32 (m, 2H), 1.97-1.82 (m, 4H), 1.68-1.63 (m, 1H), 0.52-0.38 (m, 4H).


- 25 Example 2B; (4-Cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone citrate salt.



Example 3; (4-Cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone.

MS (ESI): mass calcd. for $C_{19}H_{27}N_3OS$, 345.19; m/z found, 346 $[M+H]^+$. 1H NMR (400 MHz, $CDCl_3$): 7.37-7.32 (m, 4H), 3.86-3.60 (bm, 2H), 3.53 (s, 2H),
 5 3.53-3.25 (bm, 2H), 2.75-2.61 (bm, 10H), 2.61-2.45 (bm, 2H), 1.66-1.60 (m, 1H), 0.51-0.44 (m, 2H), 0.44-0.38 (m, 6H).


Example 3B; (4-Cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone citrate salt.



Example 4; (4-Cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone.

MS (ESI): mass calcd. for $C_{20}H_{29}N_3O_3$, 359.22; m/z found, 360 $[M+H]^+$. 1H NMR (400 MHz, $CDCl_3$): 7.39-7.34 (m, 4H), 3.93-3.87 (m, 1H), 3.85-3.25 (m,
 15 10H), 2.75-2.45 (bm, 6H), 2.25-2.15 (m, 1H), 2.05-1.93 (m, 2H), 1.67-1.60 (m, 1H), 0.52-0.45 (m, 2H), 0.45-0.40 (m, 2H).

Example 4B; (4-Cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone citrate salt.




Comparative Example 1; (4-Isopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone.

Preparation and analytical data for the title compound was presented in U.S. Patent Appl. No. 10/690,115 (Oct. 21, 2003). The corresponding salt forms
 25 were prepared as described for Examples 1A and 1B.

Comparative Example 1A; (4-Isopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone dihydrochloride.

Comparative Example 1B; (4-Isopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone citrate salt.




5 Comparative Example 2; (4-Cyclobutyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone.

The title compound was prepared according to the methods described in Example 1. MS (ESI): mass calcd. for $C_{20}H_{29}N_3O_2$, 343.23; m/z found, 344.4 $[M+H]^+$. 1H NMR (400 MHz, $CDCl_3$): 7.39-7.32 (m, 4H), 3.87-3.65 (m, 6H),
 10 3.66-3.36 (m, 4H), 2.80-2.69 (m, 1H), 2.50-2.18 (m, 8H), 2.08-1.99 (m, 2H), 1.93-1.81 (m, 2H), 1.79-1.61 (m, 2H). The corresponding salt forms were prepared as described for Examples 1A and 1B.

Comparative Example 2A; (4-Cyclobutyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone dihydrochloride.

Comparative Example 2B; (4-Cyclobutyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone citrate salt.

The compounds in Comparative Examples 3-6 and their corresponding
 20 salt forms were prepared using methods analogous to those described in the preceding examples, with the appropriate substituent changes.




Comparative Example 3; [4-(4-Fluoro-piperidin-1-ylmethyl)-phenyl]-(4-isopropyl-piperazin-1-yl)-methanone.

25 MS (ESI): mass calcd. for $C_{20}H_{30}FN_3O$, 347.24; m/z found, 348.4 $[M+H]^+$. 1H NMR (400 MHz, $CDCl_3$): 7.36-7.33 (m, 4H), 4.75-4.59 (m, 1H), 3.84-3.71 (m, 2H), 3.50 (s, 2H), 3.49-3.38 (m, 2H), 2.71 (heptet, $J = 6.6$ Hz, 1H), 2.64-2.31 (m, 8H), 1.94-1.82 (m, 4H), 1.04 (d, $J = 6.3$ Hz, 6H).

Comparative Example 3B; [4-(4-Fluoro-piperidin-1-ylmethyl)-phenyl]-(4-isopropyl-piperazin-1-yl)-methanone citrate salt.

5



Comparative Example 4; (4-Cyclobutyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone.


MS (ESI): mass calcd. for $C_{21}H_{30}FN_3O$, 359.24; m/z found, 360.4 $[M+H]^+$. 1H

NMR (400 MHz, $CDCl_3$): 7.38-7.34 (m, 4H), 4.77-4.60 (m, 1H), 3.88-3.72 (m, 2H), 3.52 (s, 2H), 3.51-3.37 (m, 2H), 2.80-2.71 (m, 1H), 2.63-2.54 (m, 2H), 2.46-2.20 (m, 6H), 2.09-2.01 (m, 2H), 1.97-1.64 (m, 8H).

10

Comparative Example 4B; (4-Cyclobutyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone citrate salt.

15



Comparative Example 5; (4-Cyclobutyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone.


MS (ESI): mass calcd. for $C_{20}H_{29}N_3OS$, 359.20; m/z found, 360 $[M+H]^+$. 1H

NMR (400 MHz, $CDCl_3$): 7.37-7.32 (m, 4H), 3.88-3.65 (bm, 2H), 3.52 (s, 2H), 3.52-3.35 (bm, 2H), 2.80-2.63 (bm, 9H), 2.45-2.28 (bm, 4H), 2.10-1.98 (m, 2H), 1.95-1.79 (m, 2H), 1.78-1.65 (m, 2H).

20

Comparative Example 5B; (4-Cyclobutyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone citrate salt.

25



Comparative Example 6; [4-(2-Hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-(4-isopropyl-piperazin-1-yl)-methanone.

MS (ESI): mass calcd. for $C_{20}H_{31}N_3O_3$, 361.24; m/z found, 362 $[M+H]^+$. 1H

5 NMR (400 MHz, $CDCl_3$): 7.38-7.34 (m, 4H), 3.93-3.87 (m, 1H), 3.87-3.35 (m, 10H), 2.73 (heptet, $J = 6.6$ Hz, 1H), 2.70-2.62 (m, 2H), 2.62-2.36 (bm, 4H), 2.24-2.15 (m, 1H), 2.05-1.92 (bm, 2H), 1.05 (d, $J = 6.5$ Hz, 6H).

10 Comparative Example 6B; [4-(2-Hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-(4-isopropyl-piperazin-1-yl)-methanone citratae salt.

Physical Chemical Data

The measured pK_a values for Example 1, a cyclopropyl amine, was significantly lower than that predicted through calculation (using the Pallas software package from CompuDrug, Inc.), and also were significantly lower than that observed for Comparative Example 1 (isopropyl amine). Measured pK_a , log P, and log D values were determined by pION, Inc. (Woburn, Mass.).

Table 1. Physical Chemical Data

EX	pK_a (calc.)	pK_a (meas.)	log P	log D
Ex. 1	8.5, 6.2	6.5, 5.3	0.89	0.86
Comp. Ex. 1	8.5, 6.2	7.6, 6.2	0.55	0.13
Comp. Ex. 2	7.8, 6.2	7.4, 6.5	2.09	1.77

20 Biological Examples and Data

A. Transfection of Cells with Human Histamine Receptor

Cells were grown to about 70% to 80% confluence and removed from the plate with trypsin and pelleted in a clinical centrifuge. The pellet was then re-suspended in 400 μ L of complete media and transferred to an
 25 electroporation cuvette with a 0.4 cm gap between the electrodes (Bio-Rad #165-2088). One μ g supercoiled H_3 receptor cDNA was added to the cells and

mixed gently. The voltage for the electroporation was set at 0.25 kV and the capacitance was set at 960 μ F. After electroporation the cells were diluted with 10 mL of complete media and were plated onto four 10 cm dishes at the following ratios: 1:20, 1:10, 1:5, and 1:2. The cells were allowed to recover for 24 h before adding 600 μ g G-418. Colonies that survived selection were grown and tested. SK-N-MC cells were used because they give efficient coupling for inhibition of adenylate cyclase. The clones that gave the most robust inhibition of adenylate cyclase in response to histamine were used for further study.

10 B. [3 H]-N-Methylhistamine Binding

Cell pellets from histamine H_3 receptor-expressing SK-N-MC cells were homogenized in 50 mM TrisHCl/0.5 mM EDTA. Supernatants from an 800 g spin were collected and were recentrifuged at 30,000 g for 30 min. Pellets were re-homogenized in 50 mM Tris/5 mM EDTA (pH 7.4). Membranes were incubated with 0.8 nM [3 H]-N-methylhistamine plus/minus test compounds for 60 min at 25 $^{\circ}$ C and were harvested by rapid filtration over GF/C glass fiber filters (pretreated with 0.3% polyethylenimine) followed by four washes with buffer. Filters were added to 5 mL of scintillation cocktail, and the signal was then counted on a liquid scintillation counter. Non-specific binding was defined with 10 μ M histamine. pK_i values were calculated based on a K_D of 0.8 nM and a ligand concentration ($[L]$) of 0.8 nM according to the formula $K_i = (IC_{50}) / (1 + ([L] / (K_D)))$. Data for compounds tested in this assay are presented in Table 2 as an average of results obtained. Binding data for Comparative Example 1 was presented in U.S. Patent Appl. No. 10/690,115 (Oct. 21, 2003).

25

Table 2. Binding Activity at the Human H_3 Receptor.

Ex.	K_i (nM)	pK_i
Example 1	5.4	8.3
Example 1B	6.5	8.2
Example 2	2.0	8.7
Example 2B	2.0	8.7
Example 3	2.5	8.6

Example 4	6.0	8.2
Example 4B	23	7.6
Comparative Example 2	1.0	9.0
Comparative Example 2A	1.7	8.8
Comparative Example 2B	2.0	8.7
Comparative Example 3	2.0	8.7
Comparative Example 3B	1.0	9.0
Comparative Example 4	0.7	9.2
Comparative Example 5	1.0	9.0
Comparative Example 6	6.0	8.2
Comparative Example 6B	5.0	8.3

C. Cyclic AMP accumulation

Sublines of SK-N-MC cells were created that expressed a reporter construct and the human H₃ receptor. The pA₂ values were obtained as described by Barbier, A.J. et al. (Br. J. Pharmacol. 1994, 143(5), 649-661).
 5 Data for compounds tested in this assay are presented in Table 3, as an average of the results obtained.

Table 3. Functional Activity

Ex.	pA ₂
Example 1	8.4
Example 2B	9.0
Comparative Example 1	9.0
Comparative Example 1A	8.9
Comparative Example 2	9.5
Comparative Example 3B	9.6
Comparative Example 4	10.0
Comparative Example 6	8.2

10 D. Pharmacokinetics and bioanalysis

One group of six male Sprague Dawley Rats (approx. 300 g body weight; three animals per time point) was used. They were group-housed,

provided food and water ad libitum, and were maintained on a 12-h light and dark cycle. Animals were acclimatized for at least 7 days after receipt from the vendor prior to investigations.

For oral dosing, test compounds were formulated at 1 mg/mL in 0.5% hydroxypropyl methyl cellulose and delivered at a dose of 10 mg/kg. Citric acid salt forms (prepared as described for Example 1B) of the test compounds were used. Animals received a bolus dose of 10 mg/kg (10 mL/kg) for each compound via a 16 gauge intragastric gavage. For intravenous dosing, test compounds were formulated at 1 mg/mL in 5% dextrose in water and dosed at 1 mg/kg (1 mL/kg) with a bolus intravenous dose via a 24-gauge Terumo® Surflo® catheter in the lateral tail vein. All dosing solutions were prepared immediately prior to injection.

Blood samples (250 µL) were taken from the lateral tail vein into heparinized Natelson blood collection tubes and expelled into 1.5 mL microcentrifuge tubes. The blood samples were centrifuged for 5 min at 14,000 rpm in a micro-centrifuge. Plasma was retained and kept in a -20 °C freezer until analysis by LC-MS/MS.

Data analysis was performed using WinNonlin version 3.3 or 4.0.1. A non-compartmental model (#200 for Extravascular administration and #201 for i.v.) was used to determine the pharmacokinetic parameters shown in Tables 4 and 5 (NA = not applicable or not determined).

Table 4. Pharmacokinetic Profiles in the Rat

Ex.	Mode	T _{max} (h)	C _{max} (ng/mL)	AUC _{INF} (h-ng/mL)	T _{1/2} (h)	Cl (mL/min/kg)	V _d (L/kg)
Ex. 1B	oral	0.5	2787	8716	3.11	20.2	5.45
	i.v.	NA	935	777	0.92	21.6	1.72
Comp. Ex. 1B	oral	1.5	1050	9743	4.71	17.1	6.98
	i.v.	NA	4113	1560	3.2	12.8	3.55
Comp. Ex. 2B	oral	1.33	683	4104	2.89	43.2	10.33
	i.v.	NA	285	464	1.86	34.8	6.41

Table 5. Pharmacokinetic Profiles in the Rat

Ex.	Mode	T _{max} (h)	C _{max} (μmol/mL)	AUC _{INF} (h-μmol/mL)	T _{1/2} (h)	Cl (mL/min/kg)	V _d (L/kg)
Ex. 2B	oral	1.67	0.42	1.91	2.72	431	112
	i.v.	NA	3.78	1.37	1.11	41.4	3.94
Comp. Ex. 3B	oral	1.33	0.16	1.95	8.22	270	180
	i.v.	NA	1.49	1.16	7.60	44.2	29.2
Comp. Ex. 4B	oral	0.67	0.48	1.28	3.44	1135	397
	i.v.	NA	0.46	0.41	2.08	113	20.3
Ex. 3B	oral	0.25	0.19	0.18	0.41	NA	NA
	i.v.	NA	0.86	0.18	0.27	282	6.48
Comp. Ex. 5B	oral	0.38	0.07	0.12	0.95	NA	NA
	i.v.	NA	2.68	0.42	1.33	154	17.8
Ex. 4B	oral	0.83	8.73	33.2	3.58	NA	NA
	i.v.	NA	2.92	3.07	1.47	15.2	1.91
Comp. Ex. 6B	oral	0.50	2.96	19.99	4.35	NA	NA
	i.v.	NA	1.41	1.34	2.36	34.9	7.04

What is claimed is:

1. A compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, and enantiomers, hydrates, solvates, and pharmaceutically acceptable salts thereof.
2. A compound that is (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, or an enantiomer, hydrate, solvate, or pharmaceutically acceptable salt thereof.
3. A compound that is (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, or an enantiomer, hydrate, solvate, or pharmaceutically acceptable salt thereof.
4. A compound that is (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, or an enantiomer, hydrate, solvate, or pharmaceutically acceptable salt thereof.
5. A compound that is (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, or an enantiomer, hydrate, solvate, or pharmaceutically acceptable salt thereof.
6. A compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone and (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone dihydrochloride.
7. A compound that is (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone.

8. A compound that is (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone dihydrochloride.
9. A compound that is (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone citrate salt.
10. A compound that is (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone citrate salt, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone citrate salt, or (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone citrate salt.
11. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective amount of at least one compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, and enantiomers, hydrates, solvates and pharmaceutically acceptable salts thereof.
12. A method for the treatment or prevention of a CNS disorder selected from the group consisting of: neurologic disorders including sleep/wake and arousal/vigilance disorders (e.g. insomnia and jet lag), attention deficit hyperactivity disorders (ADHD), learning and memory disorders, cognitive dysfunction, migraine, neurogenic inflammation, dementia, mild cognitive impairment (pre-dementia), Alzheimer's disease, epilepsy, narcolepsy with or without associated cataplexy, cataplexy, disorders of sleep/wake homeostasis, idiopathic somnolence, excessive daytime sleepiness (EDS), circadian rhythm disorders, sleep/fatigue disorders, fatigue, drowsiness associated with sleep apnea, sleep impairment due to perimenopausal hormonal shifts, Parkinson's-related fatigue, MS-related fatigue, depression-related fatigue, chemotherapy-induced fatigue, eating disorders, obesity, motion sickness, vertigo,

schizophrenia, substance abuse, bipolar disorders, manic disorders and depression in mammals, comprising the step of administering to a mammal suffering there from an effective amount of at least one compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, and enantiomers, hydrates, solvates and pharmaceutically acceptable salts thereof.

10 13. A method for the treatment or prevention of upper airway allergic response, asthma, itch, nasal congestion, or allergic rhinitis in mammals, comprising the step of administering to a mammal suffering there from an effective amount of at least one compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, and enantiomers, hydrates, solvates and pharmaceutically acceptable salts thereof.

20 14. A method for the treatment or prevention of allergic rhinitis, nasal congestion, and allergic congestion, comprising: a) administering an effective amount of at least one compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, and enantiomers, hydrates, solvates, and pharmaceutically acceptable salts thereof, and b) administering an effective amount of one or more histamine H₁ or H₂ antagonists.

15. A method for the treatment or prevention of depression, mood disorders or schizophrenia, comprising: a) administering an effective amount of at least one compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, and enantiomers, hydrates, solvates, and pharmaceutically acceptable salts thereof, and b) administering an effective amount of one or more neurotransmitter re-uptake blockers.

16. A method for the treatment or prevention of narcolepsy, excessive daytime sleepiness (EDS), Alzheimer's disease, depression, attention deficit disorders, MS-related fatigue, post-anesthesia grogginess, cognitive impairment, schizophrenia, spasticity associated with cerebral palsy, age-related memory decline, idiopathic somnolence, or jet-lag, comprising: a) administering an effective amount of at least one compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, and enantiomers, hydrates, solvates, and pharmaceutically acceptable salts thereof, and b) administering an effective amount of modafinil.

17. A compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, isotopically-labelled to be detectable by PET or SPECT.

18. A method for studying histamine-mediated disorders comprising the step of using an ^{18}F -labeled or ^{11}C -labelled compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, as a positron emission tomography (PET) molecular probe.
19. A pharmaceutical composition according to claim 11, further comprising topiramate.
20. A method of treating a subject suffering from or diagnosed with a disease mediated by histamine H_3 receptor activity, comprising administering to a subject in need of such treatment an effective amount of a compound selected from the group consisting of: (4-cyclopropyl-piperazin-1-yl)-(4-morpholin-4-ylmethyl-phenyl)-methanone, (4-cyclopropyl-piperazin-1-yl)-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-methanone, (4-cyclopropyl-piperazin-1-yl)-(4-thiomorpholin-4-ylmethyl-phenyl)-methanone, and (4-cyclopropyl-piperazin-1-yl)-[4-(2-hydroxymethyl-morpholin-4-ylmethyl)-phenyl]-methanone, and pharmaceutically acceptable salts thereof.
21. The method according to claim 20, wherein the disease is selected from the group consisting of: cognitive disorders, sleep disorders, psychiatric disorders, and other disorders.
22. The method according to claim 20, wherein the disease is selected from the group consisting of: dementia, Alzheimer's disease, cognitive dysfunction, mild cognitive impairment, pre-dementia, attention deficit hyperactivity disorders, attention-deficit disorders, and learning and memory disorders.

23. The method according to claim 20, wherein the disease is selected from the group consisting of: learning impairment, memory impairment, and memory loss.
- 5 24. The method according to claim 20, wherein the disease is selected from the group consisting of: insomnia, disturbed sleep, narcolepsy with or without associated cataplexy, cataplexy, disorders of sleep/wake homeostasis, idiopathic somnolence, excessive daytime sleepiness, circadian rhythm disorders, fatigue, lethargy, and jet lag.
- 10 25. The method according to claim 20, wherein the disease is selected from the group consisting of: sleep apnea, perimenopausal hormonal shifts, Parkinson's disease, multiple sclerosis, depression, chemotherapy, and shift work schedules.
- 15 26. The method according to claim 20, wherein the disease is selected from the group consisting of: schizophrenia, bipolar disorders, manic disorders, depression, obsessive-compulsive disorder, and post-traumatic stress disorder.
- 20 27. The method according to claim 20, wherein the disease is selected from the group consisting of: motion sickness, vertigo, epilepsy, migraine, neurogenic inflammation, eating disorders, obesity, and substance abuse disorders.
- 25 28. The method according to claim 20, wherein the disease is selected from the group consisting of: depression, disturbed sleep, fatigue, lethargy, cognitive impairment, memory impairment, memory loss, learning impairment, attention-deficit disorders, and eating disorders.
- 30 29. The method according to claim 20, wherein the disease is selected from the group consisting of: age-related cognitive decline, REM-behavioral disorder, benign postural vertigo, tinnitus, movement disorders, restless leg

syndrome, eye-related disorders, macular degeneration, and retinitis pigmentosa.