

5

Compounds Which Modulate The CB2 Receptor

APPLICATION DATA

This application claims benefit to US provisional application serial no. 60/826,819 filed
10 September 25, 2006.

BACKGROUND OF THE INVENTION

1. TECHNICAL FIELD

The present invention relates to novel compounds which modulate the CB2 receptor and their
15 use as medicaments.

2. BACKGROUND INFORMATION

Cannabinoids are a group of about 60 distinct compounds found in *Cannabis sativa* (also
know as marijuana) with cannabinal, cannabidiol and Δ^9 -tetrahydrocannabinol (THC) being
20 the most representative molecules. The therapeutic usage of *Cannabis* can be dated back to
ancient dynasties of China and includes applications for various illnesses ranging from lack of
appetite, emesis, cramps, menstrual pain, spasticity to rheumatism. The long history of
Cannabis use has led to the development of several pharmaceutical drugs. For example,
Marinol and Cesamet which are based on THC and its analogous nabilone, respectively, are
25 used as anti-emetic and appetite stimulant. Despite of the clinical benefits, the therapeutic
usage of cannabis is limited by its psychoactive effects including hallucination, addiction and
dependence. Mechoulam R, ed. *Cannabinoids as Therapeutic Agents*, Boca Raton, FL; CRC
Press, 1986 provides a review of the medicinal use of cannabis.

30 The physiological effects of cannabinoids are mediated by at least two G-protein coupled
receptors, CB1 and CB2. Autoradiographic studies have demonstrated that CB1 receptors are
expressed primarily in the central nervous system, specifically in the cerebral cortex,

5 hippocampus, basal ganglia and cerebellum. They are also found to a lesser degree in the reproductive system and other peripheral tissues including that of the immune system. CB1 receptors regulate the release of neurotransmitters from the pre-synaptic neurons and are believed to mediate most of the euphoric and other central nervous system effects of cannabis, such as THC-induced ring-catalepsy, hypomobility, and hypothermia, which were found to be
10 completely absent in mice with a deletion of the CB1 gene (Zimmer et al., Increased mortality, hypoactivity, and hypoalgesia in cannabinoid CB1 receptor knockout mice. *Proc Natl Acad Sci U S A.* (1999) 96:5780-5785.)

CB2 receptors are almost exclusively found in the immune system, with the greatest density in the spleen. It is estimated that the expression level of CB2 in the immune cells is about 10 to
15 100 times higher than CB1. Within the immune system, CB2 is found in various cell types, including B cells, NK cells, monocytes, microglial cells, neutrophils, T cells, dendritic cells and mast cells, suggesting that a wide range of immune functions can be regulated through CB2 modulators (Klein et al., The cannabinoid system and immune system. *J Leukoc Biol* (2003) 74:486-496). This is supported by the finding that the immunomodulatory effect of
20 THC is absent in CB2 deficient mice (Bicklet et al., Immunomodulation by cannabinoid is absent in mice deficient for the cannabinoid CB2 receptor. *Eur J Pharmacol* (2000) 396:141-149). CB2 selective ligands have been developed and tested for their effects in various inflammatory settings. For example, in animal models of inflammation, CB2 selective agonists, inverse agonists and antagonists have been shown to be effective in suppressing
25 inflammation (Hanus et al., HU-308: a specific agonist for CB(2), a peripheral cannabinoid receptor. *Proc Natl Acad Sci U S A.* (1999) 96:14228-14233, Ueda et al., Involvement of cannabinoid CB(2) receptor-mediated response and efficacy of cannabinoid CB(2) receptor inverse agonist, JTE-907, in cutaneous inflammation in mice. *Eur J Pharmacol.* (2005) 520:164-171 and Smith et al., The anti-inflammatory activities of cannabinoid receptor ligands
30 in mouse peritonitis models *Eur J Pharmacol.* (2001) 432:107-119.). Furthermore, CB2 selective agonists inhibit disease severity and spasticity in animal models for multiple sclerosis (Baker et al., Cannabinoids control spasticity and tremor in a multiple sclerosis

5 model. Nature (2000) 404:84-87. Arevalo-Martin et al., Therapeutic action of cannabinoids in a murine model of multiple sclerosis J Neurosci. (2003) 23:2511-2516.). Taken together, these results support the notion that CB2 receptor modulators can be employed for the treatment of medical conditions having an inflammatory component.

In addition to inflammation, CB2 agonists have been shown to inhibit pain and emesis. For
10 instance, CB2 selective agonists blunt the pain response induced by thermal or other stimuli (Malan et al., CB2 cannabinoid receptor-mediated peripheral antinociception. Pain. (2001) 93:239-45 and Nackley et al., Selective activation of cannabinoid CB(2) receptors suppresses spinal fos protein expression and pain behavior in a rat model of inflammation. Neuroscience (2003) 119:747-57.) CB2 activation has also been demonstrated to inhibit neuropathic pain
15 response (Ibrahim et al., Activation of CB2 cannabinoid receptors by AM1241 inhibits experimental neuropathic pain: pain inhibition by receptors not present in the CNS. Proc Natl Acad Sci U S A. (2003) 100:10529-33.) Finally, in contrast to the earlier data which did not find CB2 in the brain, a recent article demonstrated the expression of CB2 in the brain, at about 1.5 % of the level in the spleen. CB2 activation is shown by this article to be
20 responsible for the anti-emetic effect of endocannabinoid (Van Sickle et al., Identification and functional characterization of brainstem cannabinoid CB2 receptors. Science. 2005 310:329-332.) The foregoing results confirm that CB2 agonists can be used for the treatment of inflammatory and neuropathic pain as well as emesis.


25

BRIEF SUMMARY OF THE INVENTION

The present invention provides novel compounds which bind to and modulate the CB2 receptor. The invention also provides a method and pharmaceutical compositions for treating inflammation by way of the administration of therapeutic amounts of these compounds.
30 Lastly, the invention provides a method and pharmaceutical compositions for treating pain by way of the administration of therapeutic amounts of the new compounds which are CB2 agonists.

5 DETAILED DESCRIPTION OF THE INVENTION

In its broadest generic aspect the invention provides compounds of the formula



10

wherein:

R¹ is C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, C₃₋₁₀ cycloalkyl, 3-10 membered saturated heterocyclic ring, benzyl or phenethyl each optionally independently substituted with 1-3 substituents
 15 chosen from C₁₋₁₀ alkyl optionally substituted with 1 to 3 halogen atoms, C₃₋₁₀ cycloalkyl, 3-10 membered saturated heterocyclic ring optionally substituted with acyl, oxo or methyl sulfone, acyl, cyano, phenyl, oxo, hydroxyl and halogen;

R² and R³ are independently hydrogen or C₁₋₆ alkyl; or **R² and R³** together with the
 20 carbon atom to which they are attached form a 3- to 6-membered cycloalkyl or heterocyclic ring;

R⁴ is hydrogen or methyl;

R⁵ is aryl or heteroaryl each optionally independently substituted with 1 to 3 substituents
 25 chosen from C₁₋₆ alkyl optionally substituted with 1 to 3 halogen atoms or with a heterocyclyl group, C₁₋₆ alkoxy optionally substituted with 1 to 3 halogen atoms, C₃₋₆ cycloalkyl, phenoxy, halogen, cyano, dimethylaminoC₁₋₄alkyl, phenyl optionally substituted with 1 to 2 halogen atoms or C₁₋₄ alkyl optionally substituted with halogen, thienyl
 30 optionally substituted with 1 to 2 halogen atoms or C₁₋₄ alkyl optionally substituted with

5 halogen and pyridinyl optionally substituted with 1 to 2 C₁-C₄ alkyl optionally substituted with halogen;

or a tautomer or pharmaceutically acceptable salt thereof.

10 In a first subgeneric aspect, the invention provides compounds of the formula I wherein,

R¹ is methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, tetrahydropyranyl, bicyclo(3.3.0)octanyl, bicyclo[4.3.0]nonyl, benzyl or phenethyl each optionally independently
15 substituted with 1 to 3 substituents chosen from methyl, ethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrolidinyl (optionally substituted with acyl, oxo or methylsulfonyl), piperidinyl (optionally substituted with acyl, oxo or methylsulfone, acyl, cyano, phenyl, oxo, fluoro, chloro, bromo and hydroxyl, ;

20 **R²** and **R³** are independently hydrogen, methyl, ethyl, n-propyl, isopropyl, isobutyl, tert-butyl, or **R²** and **R³** together with the carbon to which they are attached form a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or tetrahydropyranyl ring;

R⁴ is hydrogen;

25

R⁵ is phenyl, naphthyl, pyridinyl, quinolinyl, isoquinolinyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, indolyl, benzoxazolyl, benzopyrazolyl, benzoisoxazolyl, benzothiazolyl, benzoisothiazolyl, benzimidazolyl, tetrahydrobenzoxazolyl, tetrahydrobenzothiazolyl or
30 tetrahydrobenzimidazolyl each optionally independently substituted with 1 to 3 substituents chosen from C₁-C₆ alkyl optionally substituted with 1 to 3 halogen atoms, C₁-C₆ alkoxy optionally substituted with 1 to 3 halogen atoms, C₃-C₆ cycloalkyl, phenoxy, halogen, cyano,

5 dimethylaminoC₁-C₄alkyl, phenyl optionally substituted with 1 to 2 halogen atoms or C₁-C₄ alkyl optionally substituted with halogen, thienyl optionally substituted with 1 to 2 halogen atoms or C₁-C₄ alkyl optionally substituted with halogen and pyridinyl optionally substituted with 1 to 2 C₁-C₄ alkyl optionally substituted with halogen.

10 In a further subgeneric aspect, the invention provides compounds of the formula I wherein,

R¹ is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, tetrahydropyranyl, , bicyclo(3.3.0)octanyl, or bicyclo[4.3.0]nonyl
15 each optionally independently substituted with 1 to 3 substituents chosen from methyl, ethyl, pivaloyl, cyclopropyl, cyclobutyl, cyclohexyl, tetrahydropyranyl, tetrahydrofuranyl, pyrrolidinyl optionally substituted with methylsulfone, fluoro, chloro, bromo, oxo and hydroxy;

20 **R²** and **R³** are independently hydrogen, methyl, ethyl, n-propyl, isopropyl, isobutyl, tert-butyl, or **R²** and **R³** together with the carbon to which they are attached form a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or tetrahydropyranyl ring;

R⁴ is hydrogen;

25

R⁵ is phenyl, naphthyl, pyridinyl, quinolinyl, isoquinolinyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, indolyl, benzoxazolyl, benzopyrazolyl, benzoisoxazolyl, benzothiazolyl, benzoisothiazolyl, benzimidazolyl, tetrahydrobenzoxazolyl, tetrahydrobenzothiazolyl, or
30 tetrahydrobenzimidazolyl each optionally independently substituted with 1 to 3 substituents chosen from C₁-C₆ alkyl optionally substituted with 1 to 3 halogen atoms, C₁-C₆ alkoxy optionally substituted with 1 to 3 halogen atoms, C₃-C₆ cycloalkyl, phenoxy, halogen, cyano,

5 dimethylaminoC₁-C₄alkyl, phenyl optionally substituted with 1 to 2 halogen atoms or C₁-C₄ alkyl optionally substituted with halogen, thienyl optionally substituted with 1 to 2 halogen atoms or C₁-C₄ alkyl optionally substituted with halogen and pyridinyl optionally substituted with 1 to 2 C₁-C₄ alkyl optionally substituted with halogen.

10 In another subgeneric aspect, the invention provides compounds of the formula I wherein,

R¹ is benzyl or phenethyl each optionally independently substituted with 1 to 3 substituents chosen from methyl, fluoro, chloro and bromo.

15

In another subgeneric aspect, the invention provides compounds of the formula I wherein,

R¹ is methyl, ethyl, n-propyl, isopropyl, tert-butyl, n-butyl, cyclopropyl, cyclopentyl, 20 cyclohexyl, cycloheptyl, tetrahydropyranyl or benzyl each optionally substituted with methyl, pivaloyl, cyclopropyl, cyclobutyl, cyclohexyl tetrahydropyranyl, tetrahydrofuranyl, pyrrolidinyl optionally substituted with methylsulfonyl or chloro;

R² and R³ are independently hydrogen, methyl, isopropyl, tert-butyl, or **R² and R³** together 25 with the carbon to which they are attached form a cyclopropyl or cyclobutyl, ring;

R⁴ is hydrogen;

R⁵ is phenyl, pyridinyl, quinolinyl, isoquinolinyl, pyrazolyl, isoxazolyl, benzothiazolyl, 30 thiadiazolyl, or thiazolyl each optionally independently substituted with 1 to 2 substituents chosen from, methyl, ethyl, tert-butyl, neopentyl, cyclohexyl, trifluoromethyl or phenyl optionally substituted with a chlorine atom.

- 5 The term “alkyl”, or any substituent containing an alkyl group such as alkoxy or acylamino, shall be understood to be branched or unbranched alkyl groups, preferably C₁-C₆ and shall be understood to be optionally partially or fully halogenated.

10 The compounds of formula I may be made using the general synthetic methods described below, which also constitute part of the invention.


GENERAL SYNTHETIC METHODS

15 The invention also provides processes for making compounds of Formula (I). In all methods, unless specified otherwise, R¹, R², R³, R⁴ and R⁵ in the formulas below shall have the meaning of R¹, R², R³, R⁴ and R⁵ in Formula (I) of the invention described herein above.

Optimum reaction conditions and reaction times may vary depending on the particular reactants used. Unless otherwise specified, solvents, temperatures, pressures, and other reaction conditions may be readily selected by one of ordinary skill in the art. Specific
20 procedures are provided in the Synthetic Examples section. Typically, reaction progress may be monitored by thin layer chromatography (TLC), if desired, and intermediates and products may be purified by chromatography on silica gel and/or by recrystallization.

The examples which follow are illustrative and, as recognized by one skilled in the art, particular reagents or conditions could be modified as needed for individual compounds
25 without undue experimentation. Starting materials and intermediates used, in the methods below, are either commercially available or easily prepared from commercially available materials by those skilled in the art.

Compounds of Formula (I) may be synthesized by the method illustrated in Method A
30




Method A

- As illustrated in Method A, reaction of a thiol of formula II with a bromo ethyl ester of formula III, in a suitable solvent in the presence of a suitable base, provides a thioether of formula IV. Reacting the thioether of formula IV with a suitable oxidizing agent provides the corresponding sulfone of formula V. Hydrolysis of the ester group of sulfone of formula V in a suitable solvent in the presence of a suitable base such as lithium hydroxide, provides the corresponding acid of formula VI. Reacting the acid of formula VI with a reagent such as thionyl chloride or oxalyl chloride, provides the acid chloride which is then reacted with an amine of formula VII in a suitable solvent in the presence of a suitable base, to provide a compound of formula (I). Alternatively, the acid of formula VI may also be coupled with an amine of formula VII under standard coupling conditions, to provide a compound of formula (I). Standard peptide coupling reactions known in the art (see for example M. Bodanszky, 1984, The Practice of Peptide Synthesis, Springer-Verlag) may be employed in these

5 syntheses. An example of suitable coupling conditions is treatment of a solution of the carboxylic acid in a suitable solvent such as DMF with EDC, HOBT, and a base such as diisopropylethylamine, followed by the desired amine. Further modification of the initial product of formula (I) by methods known in the art and illustrated in the Examples below, may be used to prepare additional compounds of this invention.

10


Compounds of Formula (I) may also be synthesized by the method illustrated in Method B




Method B

As shown in Method B, sulfonyl chloride of formula VIII is converted to the corresponding sulfonic acid sodium salt of formula IX, using procedures reported in the literature. Reaction of the sulfone of formula IX with a bromo ethyl ester of formula III in a suitable solvent, provides a sulfone of formula V. The sulfone of formula V is subjected to the sequence of reactions as shown in Method A, to provide a compound of formula (I).

25 Compounds of Formula (I) may be synthesized by the method illustrated in Method C






5

Method C

As illustrated in Method C, reacting the acid of formula X with a reagent such as thionyl chloride or oxalyl chloride, provides the acid chloride which is then reacted with an amine of
 10 formula VII in a suitable solvent in the presence of a suitable base, to provide a compound of formula (XI). Alternatively, the acid of formula X may also be coupled with an amine of formula VII under standard coupling conditions, to provide a compound of formula (XI). Reaction of the amide of formula XI with a sulfinic acid sodium salt of formula IX, in a suitable solvent in the presence of a suitable base, provides a compound of formula (I).

15

Compounds of Formula (I) may be synthesized by the method illustrated in Method D




20

Method D

As illustrated in Method A, reaction of a thiol of formula II with a bromo amide of formula XI, in a suitable solvent in the presence of a suitable base, provides a thioether of formula XII.
 25 Reacting the thioether of formula XII with a suitable oxidizing agent in a suitable solvent, provides a compound of formula (I).


- 5 Compounds of Formula (I) may be synthesized by the method illustrated in Method E



Method E

- 10 Reaction of the starting bromide of formula XIII with a reagent such as potassium thioacetate, in a suitable solvent, provides a thioacetic acid ester of formula XIV. Reaction of the thioacetic acid ester XIV with the bromo ethyl ester of formula III, in a suitable solvent in the presence of a suitable base, provides the corresponding sulfanyl acid ethyl ester of formula IV which may be converted to a compound of formula (I) by the sequence of steps shown in
- 15 Method A.


Compounds of Formula (I) may be synthesized by the method illustrated in Method F



Method F

- 20 Reaction of a bromoamide of formula XI with a reagent such as potassium thioacetate, in a suitable solvent provides a thio compound of formula XV. Reaction of the thio compound of formula XV with the appropriate bromo compound, in a suitable solvent, in the presence of a
- 25 suitable base provides the thio amide of formula XII which may be converted to a compound of formula (I) as in Method D.

5 Compounds of Formula (I) may be synthesized by the method outlined in Method G.




Method G

Reaction of an acylated thio-compound of formula XV with bromomethyltetra-
 10 the presence of a suitable base, such as sodium methoxide, in a suitable solvent, provides an
 intermediate of formula XVI. Reaction of the intermediate of formula XVI with a suitable
 oxidizing agent such as hydrogen peroxide provides a compound of formula I.

Compounds of Formula (I) may be synthesized by the method shown in Method H.

15




Method H

As illustrated above, reaction of an acylated thio-compound of formula XV with N-Boc-
 bromomethylpiperidine, in the presence of a suitable base, such as sodium methoxide, in a
 20 suitable solvent, provides an intermediate of formula XVII. Removal of the protecting group
 in the presence of an acid, and oxidation under standard reaction conditions, provides a
 compound of formula I.

Compounds of Formula (I) may be synthesized by the method shown in Method I.


25



Method I

As outlined above, deprotection of a N-Boc protected piperidine of formula XVII, with a suitable acid followed by acylation with a reagent such as acetic anhydride, in the presence of a suitable base, provides a compound of formula XVIII. Reaction of the intermediate of
 10 formula XVIII with a suitable oxidizing agent such as m-chloroperbenzoic acid, provides a compound of formula I.


Compounds of Formula (I) may be synthesized by the method illustrated in Method J.



Method J

As outlined in the above method J, reaction of a compound of formula XIX with methane sulfonyl chloride, in the presence of a suitable base, in a suitable solvent, provides an intermediate of formula XX. Reaction of the intermediate of formula XX with a suitable
 20 oxidizing agent such as m-chloroperbenzoic acid, provides a compound of formula I.

Compounds of Formula (I) may be synthesized by the method illustrated in Method K.




Method K

A compound of formula I may be converted to another compound of formula I as shown in method K. Alkylation of a compound of formula 1 with a reagent such as methyl iodide, in the presence of a suitable base such as lithium diisopropylamide, provides a methylated compound of formula 1.

EXAMPLES


The manner in which the compounds of the invention can be made will be further understood by way of the following Examples.

Experimental Procedures for Compounds of Formula I



Method A

**Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
(Example 1 in Table 17)**



5

Step 1: Synthesis of 2-Cyclohexylsulfanyl-2-methyl-propionic acid ethyl ester

Prepared as described by adaptation of the following reference: Brown *et al. J. Med. Chem.*

10 **1999**, 42, 3785-3788

To a solution of 2.5 g (21.5 mmol) of cyclohexyl thiol in ethanol (75 mL) were added 1.2 g (21.5 mmol) of KOH pellets, followed by 4.2 g (21.5 mmol) of ethyl α -bromoisobutyrate. The reaction was heated to reflux for 18 h and then cooled to room temperature. The solid (KBr) was separated by filtration and rinsed with ethanol (20 mL). The filtrate was concentrated under reduced pressure and the residue dissolved in DCM (50 mL). The organic layer was washed with water (2 x 20 mL). The aqueous washes were back-extracted with DCM (10 mL). The combined organics were washed with brine, dried over Na₂SO₄. Filtration and concentration under reduced pressure afforded 4.15 g of 2-cyclohexylsulfanyl-2-methyl-propionic acid ethyl ester.









15



20

According to this procedure the following thioethers were synthesized:

25

Table 1

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	2-Cyclohexylsulfanyl-2-methyl-propionic acid ethyl ester	84	231
	2-Isopropylsulfanyl-2-methyl-propionic acid ethyl ester	83	191
	1-Cyclohexylsulfanyl-cyclobutanecarboxylic acid ethyl ester	68	243
	2-Cyclopentylsulfanyl-2-methyl-propionic acid ethyl ester	77	217
	2-(4-Chloro-benzylsulfanyl)-2-methyl-propionic acid	40*	245/247
	2-tert-Butylsulfanyl-2-methyl-propionic acid ethyl ester	88	205
	2-Isopropylsulfanyl-propionic acid ethyl ester	50	177
	2-Isopropylsulfanyl-3-methyl-butyric acid ethyl ester	45	structure confirmed by ¹ H NMR

	2-sec-Butylsulfanyl-2-methyl-propionic acid ethyl ester	55	205
	2-Ethylsulfanyl-2-methyl-propionic acid ethyl ester	44	177

5 *hydrolysis of the ethyl ester occurred during reaction conditions

Step 2: Synthesis of 2-Cyclohexanesulfonyl-2-methyl-propionic acid ethyl ester

Prepared as described by adaptation of the following reference:











10 Aranapakam, V. *et al. J. Med. Chem.*, **2004**, 47, 6255-6269.

To a solution of 4.15 g (18 mmol) of 2-cyclohexylsulfanyl-2-methyl-propionic acid ethyl ester in dioxane/water (5/1, 40 mL) were added in one portion 33.2 g (54 mmol) of potassium monopersulfate triple salt (OXONE[®]). The white suspension was stirred at room temperature
 15 for 18 h. The white solid was separated by filtration and washed with dioxane (10 mL). The filtrate was concentrated under reduced pressure to remove the organic solvent. The resulting aqueous solution was extracted with DCM (3 x 40 mL). The combined organic extracts were washed with brine, dried over Na₂SO₄ and filtered. The filtrate was concentrated under reduced pressure to afford 3.47 g of 2-cyclohexanesulfonyl-2-methyl-propionic acid ethyl
 20 ester.

According to this procedure the following sulfones were synthesized:

Table 2

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
--------------	------	-----------	-------------------------

	2-Cyclohexanesulfonyl-2-methyl-propionic acid ethyl ester	50	263
	2-Methyl-2-(propane-2-sulfonyl)-propionic acid ethyl ester	75	223
	1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid ethyl ester	76	275
	2-Cyclopentanesulfonyl-2-methyl-propionic acid ethyl ester	78	249
	2-(4-Chloro-phenylmethanesulfonyl)-2-methyl-propionic acid	81*	277/279, 294/296 [M+H ₂ O ⁺]
	2-Methyl-2-(2-methyl-propane-2-sulfonyl)-propionic acid ethyl ester	86	237
	2-(Propane-2-sulfonyl)-propionic acid ethyl ester	99	209
	3-Methyl-2-(propane-2-sulfonyl)-butyric acid ethyl ester	100	237
	2-(Butane-2-sulfonyl)-2-methyl-propionic acid ethyl ester	77	237
	2-Ethanesulfonyl-2-methyl-propionic acid ethyl ester	85	209

5 *2-(4-Chloro-benzylsulfonyl)-2-methyl-propionic acid used for oxidation step

5

Step 3: Synthesis of 2-Cyclohexanesulfonyl-2-methyl-propionic acid

Prepared as described by adaptation of the following reference:

Troeger, Uhde., *J. Prakt. Chem.* **1899**, 59, 320- 349








10

To a solution of 3.47g (13.2 mmol) of 2-cyclohexanesulfonyl-2-methyl-propionic acid ethyl ester in THF/water (4/1, 50 mL) were added 1.11 g (26.4 mmol) of lithium hydroxide monohydrate. The reaction was stirred at room temperature for 18 h. The reaction was further diluted with water (20 mL) and then washed with DCM (2 x 15 mL). The basic aqueous layer was cooled in an ice bath and then acidified with 1M aqueous HCl solution to pH 2. The acidic aqueous layer was extracted with DCM (3 x 20 mL). The combined organic extracts were washed with brine, dried over Na₂SO₄ and filtered. Concentration of the filtrate under reduced pressure afforded 2.96 g of 2-cyclohexanesulfonyl-2-methyl-propionic acid.

20 According to this procedure the following acids were synthesized:

Table 3

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	2-Cyclohexanesulfonyl-2-methyl-propionic acid	64	258 [M+Na ⁺ +H ⁺]
	2-Methyl-2-(propane-2-sulfonyl)-propionic acid	92	195, 212 [M+H ₂ O ⁺]

	1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid	89	247
	2-Cyclopentanesulfonyl-2-methyl-propionic acid	92	221
	2-Methyl-2-(2-methyl-propane-2-sulfonyl)-propionic acid	69	209, 226 [M+H ₂ O ⁺]
	2-(Propane-2-sulfonyl)-propionic acid	29	181
	3-Methyl-2-(propane-2-sulfonyl)-butyric acid	32	209
	2-(Butane-2-sulfonyl)-2-methyl-propionic acid	78	209, 226 [M+H ₂ O ⁺]
	2-Ethanesulfonyl-2-methyl-propionic acid	89	198 [M+H ₂ O ⁺]

5

Step 4: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide

Activation of 100 mg (0.4 mmol) of 2-cyclohexanesulfonyl-2-methyl-propionic acid as the corresponding acid chloride was achieved by treatment with thionyl chloride (1 mL) at 80 °C

10

5 for 2 h. The reaction was cooled to room temperature and excess thionyl chloride was removed under reduced pressure.

The crude acid chloride was dissolved in DCM (0.5 mL) and added to a solution of 67 mg (0.48 mmol) of 3-amino-5-tert-butylisoxazole and N,N-diisopropylethylamine (83 mL, 0.48 mmol) in DCM (1 mL). The reaction was stirred at room temperature for 18 h, before it was
 10 evaporated to dryness. The reaction mixture was diluted with DCM (4 mL) and washed with saturated aqueous NaHCO₃ solution (3 mL). The organic layers was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (silica, eluent: DCM, 0-5% ethyl acetate) to afford 14 mg of N-(5-tert-butyl-
 isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide.


15

Examples listed in Table 17, Method A were made according to this procedure.

Method B

Synthesis of N-(4-tert-Butyl-thiazol-2-yl)-2-methanesulfonyl-2-methyl-propionamide

20 **(Example 51 in Table 17)**



Step 1: Synthesis of 2-Methanesulfonyl-2-methyl-propionic acid ethyl ester

Prepared as described by adaptation of the following references:

25 Faucher, A.-M. *et al. J. Med. Chem.* **2004**, 47, 19-21.

Binsiti, C. *Eur. J. Med. Chem. Chim. Ther.* **2001**, 36, 809-828.

Field, L.; Clark, R.D. *Org. Synth.* **1958**, 38, 62-64.

Troeger; U., *J. Prakt. Chem.* **1899**, 59, 320-349.

5 To a suspension of 5 g (49 mmol) of sodium methane sulfite in DMF (50 mL) were added pyridine (6.3 mL) and ethyl α -bromoisobutyrate (2.9 mL). The reaction was stirred for 18 h at 50 °C under nitrogen. The reaction mixture was diluted with ethyl acetate (250 mL), washed with saturated aqueous NaHCO₃ (2 x 100 mL), 2M aqueous HCl solution (50 mL), brine (1 x 50 mL) and dried over Na₂SO₄. Filtration, concentration under reduced pressure afforded 3.04
10 g of 2-methanesulfonyl-2-methyl-propionic acid ethyl ester.
The compound did not ionise under LCMS conditions, thus the structure was confirmed by ¹H-NMR spectroscopy: ¹H NMR (360 MHz, *CHLOROFORM-d*) δ ppm 1.33 (3 H, t, $J=7.1$ Hz), 1.66 (6 H, s), 3.06 (3 H, s), 4.28 (2 H, q, $J=7.1$ Hz).

15 **Step 2: Synthesis of 2-Methanesulfonyl-2-methyl-propionic acid**

2-Methanesulfonyl-2-methyl-propionic acid was generally prepared as described in step 3, method A:

To a solution of 500 mg (2.6 mmol) of 2-methanesulfonyl-2-methyl-propionic acid ethyl ester
20 in THF/water (4/1, 5 mL) were added 270 mg (6.6 mmol) of lithium hydroxide monohydrate. The reaction was stirred at room temperature for 18 h. The reaction was further diluted with water (20 mL) and then washed with DCM (2 x 15 mL). The basic aqueous layer was cooled in an ice bath and acidified with 1M aqueous HCl solution to pH 2. The acidic aqueous layer was extracted with isopropanol/chloroform (1/1, 3 x 20 mL). The combined organic extracts
25 were washed with brine, dried over Na₂SO₄ and filtered. Concentration of the filtrate under reduced pressure afforded 413 mg of 2-methanesulfonyl-2-methyl-propionic acid. The compound did not ionise under LCMS conditions, thus the structure was confirmed by ¹H-NMR spectroscopy:

¹H NMR (400 MHz, *DMSO-d*₆) δ ppm 1.49 (6 H, s), 3.08 (3 H, s).

30

Step 3: Synthesis of N-(4-tert-butyl-thiazol-2-yl)-2-methanesulfonyl-2-methyl-propionamide

5


Activation of 97 mg (0.6 mmol) of 2-methanesulfonyl-2-methyl-propionic acid as the corresponding acid chloride was achieved by treatment with thionyl chloride (2 mL) at 80 °C for 2 h. The reaction was cooled to room temperature and excess thionyl chloride was removed under reduced pressure.

- 10 The crude acid chloride was dissolved in DCM (2 mL) and added to a solution of 91 mg (0.6 mmol) of 2-amino-4-tert-butylthiazole and N, N-diisopropylethylamine (0.13 mL) in DCM (1 mL). The reaction was stirred at room temperature for 18 h. The reaction mixture was diluted with DCM, washed with saturated aqueous NaHCO₃ solution (2 mL), brine (2 mL) and dried over Na₂SO₄. Filtration and concentration of the filtrate afforded the crude product. Further
- 15 purification by column chromatography (silica, eluent: heptanes, 0-20% ethyl acetate) yielded 116 mg of N-(4-tert-butyl-thiazol-2-yl)-2-methanesulfonyl-2-methyl-propionamide.

Examples listed in Table 17, Method B were made according to this procedure.

20 **Method C**

Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide (Example 10 in Table 17)



Step 1: Synthesis of 1-propanesulfinic acid sodium salt

25

Prepared as described by adaptation of the following references:

Faucher, A.-M. *et al. J. Med. Chem.* **2004**, 47, 19-21.

Binsiti, C. *Eur. J. Med. Chem. Chim. Ther.* **2001**, 36, 809-828.

5 Field, L.; Clark, R.D. *Org. Synth.* **1958**, 38, 62-64.

To a solution of 0.26 g (3.2 mmol) of NaHCO₃ and 0.4 g (3.2 mmol) of Na₂SO₃ in water (1.5 mL) were added 0.23 g (1.6 mmol) of 1-propanesulfonyl chloride. The reaction was heated at 80 °C for 3 h. The solvent was removed under reduced pressure. The crude material was
10 suspended in boiling ethanol (20 mL) and the inorganic solids were removed by filtration. The filtrate was concentrated under reduced pressure to afford 0.18 g of 1-propanesulfinic acid sodium salt.

Step 2: Synthesis of 2-Bromo-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide

15

Prepared as described by adaptation of the following references: Katoh A. *et al. Heterocycles* **1999**, 50, 299-308

Activation of 5 g (30 mmol) of 2-bromo-2-methylpropionic acid as the corresponding acid
20 chloride was achieved by treatment with thionyl chloride (10 mL) at 60 °C for 2 h. The reaction was cooled and excess thionyl chloride was removed under reduced pressure to afford the corresponding acid chloride.

To a solution of 4.2 g (30 mmol) of 3-amino-5-tert-butylisoxazole and 5.2 mL (30 mmol) of N,N-diisopropylethylamine in DCM (20 mL) was added dropwise the acid chloride, as
25 solution in DCM (15 mL). The reaction was stirred at room temperature for 18 h. The reaction mixture was washed with saturated aqueous NaHCO₃ solution (2 x 15 mL) and the layers were separated. The organic layer was further washed with brine, dried over MgSO₄, filtered and concentrated under reduced pressure. The crude was purified by dry flash column chromatography (silica, eluent DCM, 0-20% ethyl acetate) to yield 8.5 g of 2-bromo-N-(5-
30 tert-butyl-isoxazol-3-yl)-2-methyl-propionamide.

According to this procedure the following amides were synthesized:

5 **Table 4**

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	2-Bromo-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide	97	289/291
	2-Bromo-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	55	311/313
	2-Bromo-N-(5-tert-butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-propionamide	51	302/304
	2-Bromo-2-methyl-N-(6-trifluoromethyl-pyridin-2-yl)-propionamide	72	311/313
	2-Bromo-N-(3-tert-butyl-isoxazol-5-yl)-2-methyl-propionamide	68	289/291

Step 3: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide

10 Prepared as described by adaptation of the following references:

Faucher, A.-M. *et al. J. Med. Chem.* **2004**, 47, 19-21.

Troeger; Uhde; *J. Prakt. Chem.* **1899**, 59, 320-349.

Binsiti, C. *Eur. J. Med. Chem. Chim. Ther.* **2001**, 36, 809-828.

Field, L.; Clark, R.D. *Org. Synth.* **1958**, 38, 62-64.


- 5 To a solution of 160 mg (0.55 mmol) of 2-bromo-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide in DMF (3 mL) were added 87 mg (0.71 mmol) of 1-propanesulfinic acid sodium salt in one portion. Pyridine (48 μ L) was added and the reaction was stirred at 50 °C for 18 h. The mixture was acidified with 1M aqueous HCl solution (1 mL) and extracted with diethyl ether (3 x 3 mL). The combined organic layers were dried over Na₂SO₄ and filtered.
- 10 The filtrate was concentrated under reduced pressure. The crude material was purified by column chromatography (silica, eluent heptanes, 0-20% ethyl acetate) to afford 86 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide.

Examples listed in Table 17, Method C were made according to this procedure.

15

Method D

Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide (Example 52 in Table 17)





- 20 **Step 1: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-isobutylsulfanyl-2-methyl-propionamide**

- To a stirred solution of 82 μ L (1.31 mmol) of 2-methyl-1-propanethiol in ethanol (2 mL) were added 37 mg (0.65 mmol) of KOH pellets, followed by 190 mg (0.65 mmol) 2-bromo-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide (prepared as described in step 2, method B).
- 25 The reaction mixture was heated to reflux for 18 h. The reaction was cooled to room temperature. The solid (KBr) was separated by filtration and rinsed with ethanol (15 mL). The filtrate was concentrated under reduced pressure to yield 160 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-isobutylsulfanyl-2-methyl-propionamide.

5

According to this procedure the following amides were synthesized:

Table 5


MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	N-(5-tert-Butyl-isoxazol-3-yl)-2-isobutylsulfonyl-2-methyl-propionamide	83	299
	N-(3-tert-Butyl-isoxazol-5-yl)-2-(4-fluoro-benzylsulfonyl)-2-methyl-propionamide	63	351

10 **Step 2: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-isobutylsulfonyl-2-methyl-propionamide**

To a solution of 160 mg (0.54 mmol) of N-(5-tert-butyl-isoxazol-3-yl)-2-isobutylsulfonyl-2-methyl-propionamide in dioxane/water (5/1, 2.5 mL) were added in one portion 0.99 g (1.61 mmol) of potassium monopersulfate triple salt (OXONE[®]). The white suspension was stirred at room temperature for 18 h. The white solid was separated by filtration and washed with dioxane (5 mL). The filtrate was concentrated under reduced pressure to remove the organic solvent. The resulting aqueous solution was extracted with DCM (2 x 5 mL). The combined organic extracts were washed with saturated aqueous NaHCO₃ solution (5 mL), brine (5 mL), dried over Na₂SO₄ and filtered. The filtrate was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica, eluent DCM, 0-10 % ethyl acetate) to afford 98 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-isobutylsulfonyl-2-methyl-propionamide.

25 Examples listed in Table 17, Method D were made according to this procedure.

5

Method E**Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide (Example 56, Table 17)****Step 1: Synthesis of Thioacetic acid S-(tetrahydro-pyran-4-ylmethyl) ester**

Prepared as described by adaptation of the following literature reference:


Watson, R.J. *et al. Tetrahedron Lett.* **2002**, 43, 683-685.

To a solution of 0.97 g (5.4 mmol) of 4-bromomethyl tetrahydropyran in DMF (9.7 mL) were added 1.27 g (11.2 mmol) of potassium thioacetate. The reaction was stirred at room temperature for 3 h. Diethyl ether (100 mL) was added and the reaction mixture was washed with saturated aqueous NaHCO₃ solution (2 x 25 mL) and brine (25 mL). The organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica, eluent: heptanes, 10 % ethyl acetate) to afford 0.81 g of thioacetic acid S-(tetrahydro-pyran-4-ylmethyl) ester.

According to this procedure the following thioacetates were synthesized:

Table 6

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	Thioacetic acid S-(tetrahydro-pyran-4-ylmethyl) ester	86	175

	Thioacetic acid S-(tetrahydro-pyran-4-yl) ester	68	161
---	---	----	-----



5


Step 2: Synthesis of 2-Methyl-2-(tetrahydro-pyran-4-ylmethylsulfanyl)-propionic acid ethyl ester

A solution of 0.86 g (15.2 mmol) of potassium hydroxide in ethanol (78 mL, degassed and under nitrogen) was added to 0.81 g (4.66 mmol) of thioacetic acid S-(tetrahydro-pyran-4-ylmethyl) ester. The reaction was stirred at room temperature under nitrogen for 0.5 h. Then 2.1 mL (14.1 mmol) of ethyl α -bromoisobutyrate were added and the reaction stirred for 4 h. The resulting precipitate was removed by filtration and the filtrate concentrated under reduced pressure. The residue was dissolved in DCM (100 mL) and washed with saturated aqueous NaHCO_3 (50 mL), brine (50 mL) and dried over Na_2SO_4 . Filtration and concentration of the filtrate under reduced pressure, followed by column chromatography of the residue (silica, eluent: heptanes, 0-20 % ethyl acetate) afforded 0.76 g of 2-methyl-2-(tetrahydro-pyran-4-ylmethylsulfanyl)-propionic acid ethyl ester.

20 According to this procedure the following ethyl esters were synthesized:

Table 7

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	2-Methyl-2-(tetrahydro-pyran-4-ylmethylsulfanyl)-propionic acid ethyl ester	66	247
	2-Methyl-2-(tetrahydro-pyran-4-ylsulfanyl)-propionic acid ethyl ester	75	233

	1-(Tetrahydro-pyran-4-ylmethylsulfanyl)-cyclobutanecarboxylic acid ethyl ester	77	259
---	--	----	-----

5



Step 3: Synthesis of 2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionic acid ethyl ester


Prepared as described by adaptation of Method A, step 2.

- 10 To a solution of 0.76 g (3.1 mmol) of 2-methyl-2-(tetrahydro-pyran-4-ylmethylsulfanyl)-propionic acid ethyl ester in dioxane/water (1/1, 10 mL) was added in one portion 4.1 g (6.6 mmol) of potassium monopersulfate triple salt (OXONE[®]). The white suspension was stirred at room temperature for 1 h. The white solid was separated by filtration and washed with dioxane (5 mL). The filtrate was concentrated under reduced pressure to remove the organic
- 15 solvent. The resulting mixture was diluted with DCM (50 mL) and water (5 mL). The organic layer was separated and washed with a saturated aqueous NaHCO₃/brine (1/1) mixture (20 mL) and dried over Na₂SO₄. Filtration and concentration under reduced pressure afforded 0.77 g of 2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionic acid ethyl ester.

- 20 According to this procedure the following sulfones were synthesized:

Table 8

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionic acid ethyl ester	90	279
	2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionic acid ethyl ester	84	265

	1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid ethyl ester	60	291
---	---	----	-----

5

Step 4: Synthesis of 2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionic acid



Prepared as described by adaptation of Method A, step 3.


- 10 To a solution of 769 mg (2.77 mmol) of 2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionic acid ethyl ester in THF/water (1/1, 10 mL) were added 228 mg (5.43 mmol) of lithium hydroxide monohydrate. The reaction was stirred at room temperature for 18 h. The reaction was concentrated under reduced pressure. The residue was dissolved in water (10 mL) and washed with diethyl ether (2 x 25 mL). The aqueous layer was cooled in an ice bath
- 15 and then acidified with 1M aqueous HCl solution to pH 2. The acidic aqueous layer was extracted with ethyl acetate (2 x 50 mL) and with isopropanol/chloroform (3 x 50 mL). The combined organic extracts were dried over Na₂SO₄ and filtered. Concentration of the filtrate under reduced pressure afforded 734 mg of 2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionic acid.

20

According to this procedure the following acids were synthesized:

Table 9

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionic acid	100	251
	2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionic acid	93	237

	1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid	80	263
---	---	----	-----

5

Step 5: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide


- 10 Prepared as described by adaptation of Method A, step 4.
- Activation of 122 mg (0.49 mmol) of 2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionic acid as the corresponding acid chloride was achieved by treatment with thionyl chloride (0.4 mL) at 50 °C for 1 h. The reaction was cooled to room temperature and excess thionyl chloride was removed under reduced pressure.
- 15 The crude acid chloride was dissolved in DCM (2 mL) and added to a solution of 63 mg (0.49 mmol) of 3-amino-5-tert-butylisoxazole and N,N-diisopropylethylamine (145 mL, 0.88 mmol) in DCM (2 mL). The reaction was stirred at room temperature for 18 h. The reaction mixture was diluted with DCM (2 mL) and washed with a saturated aqueous NaHCO₃ solution/brine mixture (1/1, 2 mL). The organic layer was dried over Na₂SO₄, filtered and concentrated under
- 20 reduced pressure. The crude product was purified by column chromatography (silica, eluent: heptanes, 0-20 % ethyl acetate, then DCM, 0-20 % ethyl acetate) to afford 41 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide.

Examples listed in Table 17, Method E were made according to this procedure.

25

Method F

Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-4-ylmethanesulfonyl)-propionamide (Example 74, Table 17)



Step 1: Synthesis of Thioacetic acid S-[1-(5-tert-butyl-isoxazol-3-yl)carbamoyl]-1-methyl-ethyl ester

To a solution of 200 mg (0.69 mmol) of 2-bromo-N-(5-tert-butyl-isoxazol-3-yl)-2-methylpropionamide (prepared as described in Method B, step 2) in DMF (3 mL) were added 157 mg (1.4 mmol) of potassium thioacetate. The reaction was stirred at room temperature for 3 h. The reaction mixture was diluted with diethyl ether (5 mL) and washed with 2M aqueous HCl solution (5 mL). The organic layer was dried over Na₂SO₄. Filtration and concentration under reduced pressure afforded 151 mg of thioacetic acid S-[1-(5-tert-butyl-isoxazol-3-yl)carbamoyl]-1-methyl-ethyl ester.

According to this procedure the following thioacetates were synthesized:

Table 10

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	Thioacetic acid S-[1-(5-tert-butyl-isoxazol-3-yl)carbamoyl]-1-methyl-ethyl ester	76	285
	Thioacetic acid S-[1-(5-tert-butyl-2-methyl-2H-pyrazol-3-yl)carbamoyl]-1-methyl-ethyl ester	73	298

	Thioacetic acid S-[1-methyl-1-(5-trifluoromethyl-pyridin-2-ylcarbamoyl)-ethyl] ester	62	307
	Thioacetic acid S-[1-(3-tert-butyl-isoxazol-5-ylcarbamoyl)-1-methyl-ethyl] ester	62	285

5

Step 2: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethylsulfanyl)-propionamide

- 10 To a solution of 100 mg (0.35 mmol) of thioacetic acid S-[1-(5-tert-butyl-isoxazol-3-ylcarbamoyl)-1-methyl-ethyl] ester in ethanol (2 mL) were added 70 mg (0.42 mmol) of 2-(bromomethyl)tetrahydrofuran and 340 μ L (1.05 mmol) of sodium ethoxide solution (21% in ethanol) at room temperature. The reaction was heated to 50 °C for 18 h. The reaction mixture was concentrated under reduced pressure and the crude purified by column chromatography
- 15 (silica, eluent: heptanes, 0-20% ethyl acetate) to afford 82 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethylsulfanyl)-propionamide.

According to this procedure the following thioethers were synthesized:

Table 11

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethylsulfanyl)-propionamide	71	327

20

Step 3: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide

5

Prepared as described by adaptation of Method D, step 2.


To a solution of 82 mg (0.25 mmol) of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethylsulfanyl)-propionamide in dioxane/water (5/1, 4 mL) were added in one portion 460 mg (0.75 mmol) of potassium monopersulfate triple salt (OXONE[®]). The white
 10 suspension was stirred at room temperature for 3 h. The white solid was separated by filtration and washed with dioxane (5 mL). The filtrate was concentrated under reduced pressure to remove the organic solvent. The resulting aqueous solution was extracted with DCM (2 x 5 mL). The combined organic extracts were washed with saturated aqueous NaHCO₃ solution (5 mL), brine (5 mL), dried over Na₂SO₄ and filtered. The filtrate was concentrated under
 15 reduced pressure. The crude residue was purified by column chromatography (silica, eluent: DCM, 0-10 % ethyl acetate) to afford 5 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide.

Examples listed in Table 17, Method F were made according to this procedure.

20

Method G

Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-propionamide (Example 82, Table 17)



25




Step 1: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethylsulfanyl)-propionamide







To a solution of 100 mg (0.352 mmol) of thioacetic acid S-[1-(5-tert-butyl-isoxazol-3-ylcarbamoyl)-1-methyl-ethyl] ester (synthesized according to Method F, step 1) in ethanol (2
 30







5 mL) were added 126 mg (0.704 mmol) of 2-(bromomethyl)tetrahydropyran and 76 mg (1.41 mmol) of sodium methoxide at room temperature. The reaction was heated to 130 °C for 0.5 h within a microwave (CEM Discover). The reaction mixture was concentrated under reduced pressure. The residue was dissolved in DCM (5ml) and washed with saturated aqueous NaHCO₃ solution (5ml). The organic phase was dried (Na₂SO₄), filtered and concentrated under reduced pressure to yield 119 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethylsulfanyl)-propionamide.








According to this procedure the following thioethers were synthesized:








Table 12






MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethylsulfanyl)-propionamide	100	354
	N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexylmethylsulfanyl-2-methyl-propionamide	100	339
	N-(5-tert-Butyl-isoxazol-3-yl)-2-ethylsulfanyl-2-methyl-propionamide	90	271

	N-(5-tert-Butyl-isoxazol-3-yl)-2-cyanomethylsulfanyl-2-methyl-propionamide	96	282
	N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydropyran-2-ylmethylsulfanyl)-propionamide	100	341
	N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-(2,2-dimethyl-propylsulfanyl)-2-methyl-propionamide	63	326
	N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cycloheptylsulfanyl-2-methyl-propionamide	68	352
	N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclohexylmethylsulfanyl-2-methyl-propionamide	95	352
	2-Methyl-2-(tetrahydrofuran-2-ylmethylsulfanyl)-N-(5-(trifluoromethyl)-pyridin-2-yl)-propionamide	53	349

	2-Methyl-2-(tetrahydro- pyran-2-ylmethylsulfanyl)- N-(5-trifluoromethyl- pyridin-2-yl)-propionamide	54	363
	2-(3,3-Dimethyl-2-oxo- butylsulfanyl)-2-methyl-N- (5-trifluoromethyl-pyridin- 2-yl)-propionamide	45	363
	N-(5-tert-Butyl-isoxazol-3- yl)-2-(3,3-dimethyl-2-oxo- butylsulfanyl)-2-methyl- propionamide	75	341
	N-(5-tert-Butyl-2-methyl- 2H-pyrazol-3-yl)-2-methyl- 2-((S)-5-oxo-pyrrolidin-2- ylmethylsulfanyl)- propionamide	85	353
	N-(5-tert-Butyl-2-methyl- 2H-pyrazol-3-yl)-2-methyl- 2-((R)-5-oxo-pyrrolidin-2- ylmethylsulfanyl)- propionamide	97	353
	N-(5-tert-Butyl-isoxazol-3- yl)-2-methyl-2-(3-methyl- butylsulfanyl)-propionamide	70	313

	N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(3-methyl-butylsulfanyl)-propionamide	90	326
	N-(5-tert-Butyl-isoxazol-3-yl)-2-cycloheptylsulfanyl-2-methyl-propionamide	89	339
	2-Cycloheptylsulfanyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	83	361
	2-Cyclopropylmethylsulfanyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	100	319
	2-Cyclobutylmethylsulfanyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	97	333
	2-Methyl-2-(3-phenyl-propylsulfanyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	82	383
	N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropylmethylsulfanyl-2-methyl-propionamide	74	297

	N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclobutylmethylsulfanyl-2-methyl-propionamide	77	311
	N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-phenyl-propylsulfanyl)-propionamide	86	361
	2-Cycloheptylsulfanyl-2-methyl-N-(5-(2,2,2-trifluoromethyl)-pyridin-2-yl)-propionamide	83	361
	N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclobutylmethylsulfanyl-2-methyl-propionamide	55	311
	N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopropylmethylsulfanyl-2-methyl-propionamide	70	297
	N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(3-methyl-butylsulfanyl)-propionamide	50	313
	N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexylmethylsulfanyl-2-methyl-propionamide	50	339

	<i>N</i> -(3- <i>tert</i> -Butyl-isoxazol-5-yl)-2- <i>sec</i> -butylsulfanyl-2-methyl-propionamide	73	299
	<i>N</i> -(3- <i>tert</i> -Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethylsulfanyl)-propionamide	75	341
	<i>N</i> -(3- <i>tert</i> -Butyl-isoxazol-5-yl)-2-isobutylsulfanyl-2-methyl-propionamide	73	299
	<i>N</i> -(3- <i>tert</i> -Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethylsulfanyl)-propionamide	100	327
	<i>N</i> -(3- <i>tert</i> -Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-ylmethylsulfanyl)-propionamide	100	327

5

Step 3: Synthesis of N-(5-*tert*-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide


To a solution of 119 mg (0.365 mmol) of N-(5-*tert*-butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethylsulfanyl)-propionamide in acetic acid (1 mL) were added 103 μ L (1.83 mmol) of hydrogen peroxide (50% solution, stabilized in water) and heated to 80 °C for 50 minutes. The colorless solution was quenched with ethanol (2 mL) and concentrated under reduced pressure. The crude material was purified by mass-triggered preparative LCMS and residual TFA was removed by filtration through a small plug of Ambersep 900-OH resin to

- 5 yield 27 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide.

Examples listed in Table 17, Method G were made according to this procedure.

10 **Method H**

Synthesis of 2-Methyl-2-(piperidin-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide (Example 98, Table 17)







15 **Step 1: Synthesis of 4-[1-Methyl-1-(5-trifluoromethyl-pyridin-2-ylcarbamoyl)-ethylsulfanylmethyl]-piperidine-1-carboxylic acid tert-butyl ester**

- To a solution of 200 mg (0.654 mmol) of thioacetic acid S-[1-methyl-1-(5-trifluoromethyl-pyridin-2-ylcarbamoyl)-ethyl] ester (synthesized according to Method F, step 1) in ethanol (2 mL) were added 363 mg (1.31 mmol) of 4-bromomethyl-piperidine-1-carboxylic acid tert-butyl ester and 141 mg (2.61 mmol) of sodium methoxide at room temperature. The reaction was heated to 130 °C for 0.5 h within a microwave (CEM Discover). The reaction mixture was concentrated under reduced pressure. The residue was dissolved in DCM (4 mL) and washed with saturated aqueous NaHCO₃ solution (5mL). The organic phase was dried over Na₂SO₄, filtered and the filtrate was concentrated under reduced pressure to yield 375 mg of 4-[1-methyl-1-(5-trifluoromethyl-pyridin-2-ylcarbamoyl)-ethylsulfanylmethyl]-piperidine-1-carboxylic acid tert-butyl ester.

According to this procedure the following thioethers were synthesized:

30 **Table 13**

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
--------------	------	-----------	-------------------------

	4-[1-(5-tert-Butyl-isoxazol-3-ylcarbamoyl)-1-methyl-ethylsulfanylmethyl]-piperidine-1-carboxylic acid tert-butyl ester	89	462 [M+Na ⁺], 340 [M+H ⁺ -C ₅ H ₈ O ₂]
	4-[1-Methyl-1-(5-trifluoromethyl-pyridin-2-ylcarbamoyl)-ethylsulfanylmethyl]-piperidine-1-carboxylic acid tert-butyl ester	100	462 [M+H ⁺], 485 [M+Na ⁺]
	3-[1-(5-tert-Butyl-isoxazol-3-ylcarbamoyl)-1-methyl-ethylsulfanylmethyl]-piperidine-1-carboxylic acid tert-butyl ester	88	462 [M+Na ⁺], 340 [M+H ⁺ -C ₅ H ₈ O ₂]
	3-[1-(5-tert-Butyl-isoxazol-3-ylcarbamoyl)-1-methyl-ethylsulfanylmethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester	30	426 [M+H ⁺], 448 [M+Na ⁺], 326 [M+H ⁺ -C ₅ H ₈ O ₂]

5

Step 2: Synthesis of 2-Methyl-2-(piperidin-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide

To a solution of 375 mg (0.81 mmol) of 4-[1-methyl-1-(5-trifluoromethyl-pyridin-2-ylcarbamoyl)-ethylsulfanylmethyl]-piperidine-1-carboxylic acid tert-butyl ester in acetic acid (1ml) were added 241 μ L (4.26 mmol) of hydrogen peroxide (50% solution, stabilized in water) and heated to 80 °C for 50 minutes. The colorless solution was quenched with ethanol (2 mL) and concentrated under reduced pressure. The crude material was treated with a solution of 10% TFA in DCM (5 mL) for 16 h. The solvents were removed under reduced


5 pressure. The residue was purified by mass triggered preparative LC and residual TFA was removed by filtration through a small plug of Ambersep 900-OH resin to yield 24.8 mg of 2-methyl-2-(piperidin-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide.

Examples listed in Table 17, Method H were made according to this procedure.

10

Method I

Synthesis of 2-(1-Acetyl-piperidin-3-ylmethanesulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide (Example 39, Table 17)



15



Step 1: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(piperidin-3-ylmethanesulfonyl)-propionamide

20 To a solution of 288 mg (0.65 mmol) of 3-[1-(5-tert-butyl-isoxazol-3-ylcarbamoyl)-1-methyl-ethylsulfanylmethyl]-piperidine-1-carboxylic acid tert-butyl ester (synthesized according to Method H, step 1) in DCM (5 mL) were added 1.1 mL of HCl (2M solution in dioxane). The reaction was stirred at room temperature for 16 h. The mixture was diluted with saturated aqueous NaHCO₃ solution (5 mL), the organic layer was separated and dried over Na₂SO₄.

25 Filtration and concentration of the filtrate under reduced pressure afforded 175 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(piperidin-3-ylmethanesulfonyl)-propionamide.

5 According to this procedure the following thioethers were synthesized

Table 14

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(piperidin-3-ylmethylsulfanyl)-propionamide	79	340
	N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(pyrrolidin-3-ylmethylsulfanyl)-propionamide	89	326

Step 2: Synthesis of 2-(1-Acetyl-piperidin-3-ylmethylsulfanyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide


10


A solution of 175 mg (0.52 mmol) of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(piperidin-3-ylmethylsulfanyl)-propionamide 0.18 mL (1.04 mmol) of N,N-diisopropylethylamine and acetic anhydride (1.5 mL) were heated for 0.5 h to 100 °C within a microwave (CEM discover). The solvent was removed under reduced pressure. The residue was dissolved in DCM (5 mL) and washed with saturated aqueous NaHCO₃ solution (5 mL). The organic phase was dried over Na₂SO₄, filtered and the filtrate concentrated under reduced pressure to afford 183 mg of 2-(1-acetyl-piperidin-3-ylmethylsulfanyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide, which was used in the next step without further purification.

15

20 According to this method the following amides were synthesized.

Table 15

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	2-(1-Acetyl-piperidin-3-ylmethylsulfanyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide	92	382

	2-(1-Acetyl-pyrrolidin-3-ylmethanesulfanyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide	80	368
---	--	----	-----

5


Step 3: Synthesis of 2-(1-Acetyl-piperidin-3-ylmethanesulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide

To a solution of 183 mg (0.48 mmol) of 2-(1-acetyl-piperidin-3-ylmethanesulfanyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide in DCM (5 mL) were added 123 mg (0.72 mmol) of *m*-chloroperoxybenzoic acid. The reaction was stirred at room temperature until completion. Then 774 mg (2.4 mmol) of aminomethylene polystyrene were added to the reaction and the mixture was shaken for 18h. The resins were separated by filtration and rinsed with DCM. The filtrate was concentrated under reduced pressure and purified by mass triggered preparative LCMS (at neutral pH) to afford 34 mg of 2-(1-acetyl-piperidin-3-ylmethanesulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide.

Examples listed in Table 17, Method I were made according to this procedure.

Method J



Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-methyl-propionamide (Example 127, Table 17)



Step 1: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-methyl-propionamide

- 5 To a solution of 172 mg (0.52 mmol) of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(pyrrolidin-3-ylmethylsulfanyl)-propionamide (prepared according to Method I, step 1) and 0.18 mL (1.04 mmol) of N,N-diisopropylethylamine in THF were added 0.16 mL (2.11 mmol) of methanesulfonyl chloride. The reaction was heated within a microwave to 90 °C for 0.5 h. The mixture was concentrated under reduced pressure. The residue was dissolved in DCM (5 mL) and
- 10 washed with saturated aqueous NaHCO₃ solution (5 mL). The organic layer was dried over Na₂SO₄, filtered and the filtrate was concentrated under reduced pressure to afford 168 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethylsulfanyl)-2-methyl-propionamide, which was used in the next step without further purification.
- 15 According to this method, the following sulfonamides were synthesized.

Table 16

MOLSTRUCTURE	NAME	YIELD [%]	m/z [M+H ⁺]
	N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-piperidin-3-ylmethylsulfanyl)-2-methyl-propionamide	73	418
	N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethylsulfanyl)-2-methyl-propionamide	78	404

Step 2: Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-methyl-propionamide

20

To a solution of 168 mg (0.41 mmol) of N-(5-tert-butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethylsulfanyl)-2-methyl-propionamide in DCM (5 mL) were added 107 mg (0.63 mmol) of *m*-chloroperoxybenzoic acid. The reaction was stirred at room temperature until completion. Then 672 mg (2.08 mmol) of aminomethylene polystyrene were added to the reaction


5 and the mixture was shaken for 18 h. The resins were separated by filtration and rinsed with DCM. The filtrate was concentrated under reduced pressure and the residue was purified by mass triggered preparative LCMS (at neutral pH) to afford 37 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-methyl-propionamide.

10 Examples listed in Table 17, Method J were made according to this procedure.

Method K

Synthesis of N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(4-methyl-tetrahydro-pyran-4-sulfonyl)-propionamide (Example 100, Table 17)

15


















To a solution of 202 mg (0.56 mmol) of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide (prepared according to Method E) in anhydrous THF (10 mL) at -78 °C under nitrogen atmosphere were added 0.8 mL (1.4 mmol) of lithium diisopropylamide (1.8 M solution in THF/heptane/ethylbenzene) and the reaction was stirred at -78 °C for 0.5 h. Then 0.1 mL (1.6 mmol) of methyl iodide were added in one portion and stirring was continued for further 0.5 h at -78 °C, before the reaction was warmed to room temperature. After 18 h at room temperature the reaction was quenched by the addition of saturated aqueous NH₄Cl solution (25 mL). The mixture was extracted with ethyl acetate (3 x 25 mL). The organic extracts were combined, dried over Na₂SO₄ and filtered. The filtrate was concentrated under reduced pressure and the residue purified by mass-triggered preparative LCMS to afford 31 mg of N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-2-(4-methyl-tetrahydro-pyran-4-sulfonyl)-propionamide.

Examples listed in Table 17, Method K were made according to this procedure.

30









5 **Table 17: Examples**

#	MOLSTRUCTURE	NAME	m/z [M+H ⁺]	Method
1		N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide	357	A
2		N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide	333	A
3		N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide	347	A
4		1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (5-tert-butyl-isoxazol-3-yl)-amide	369	A
5		1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (4-tert-butyl-thiazol-2-yl)-amide	385	A
6		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide	317	A
7		N-(4-tert-Butyl-thiazol-2-yl)-2-(4-chlorophenylmethanesulfonyl)-2-methyl-propionamide	415/417	A









8		N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide	429/431	A
9		N-(5-tert-Butyl-isoxazol-3-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide	399/401	A
10		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	317	C
11		2-Methyl-2-(propane-1-sulfonyl)-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide	339	A
12		2-Methyl-N-naphthalen-2-yl-2-(propane-2-sulfonyl)-propionamide	320.25	A
13		2-Methyl-2-(propane-2-sulfonyl)-N-(4,5,6,7-tetrahydro-benzothiazol-2-yl)-propionamide	331.35	A
14		2-Methyl-N-(2-methyl-5-thiophen-2-yl-2H-pyrazol-3-yl)-2-(propane-2-sulfonyl)-propionamide	356.21	A
15		N-Isoquinolin-3-yl-2-methyl-2-(propane-2-sulfonyl)-propionamide	321.27	A









16		2-Methyl-2-(propane-2-sulfonyl)-N-(4-trifluoromethyl-thiazol-2-yl)-propionamide	345.13	A
17		N-Benzothiazol-2-yl-2-methyl-2-(propane-2-sulfonyl)-propionamide	327.23	A
18		N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	333	A
19		2-Methyl-2-(propane-2-sulfonyl)-N-quinolin-3-yl-propionamide	321.36	A
20		N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide	326.27	A
21		2-Methyl-N-(4-phenyl-thiazol-2-yl)-2-(propane-2-sulfonyl)-propionamide	353.26	A
22		2-Methyl-2-(propane-2-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	339.23	A
23		2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(propane-2-sulfonyl)-propionamide	350.33	A









24		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide	330.29	A
25		N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide	326.27	A
26		2-Methyl-N-(5-phenyl-pyridin-2-yl)-2-(propane-2-sulfonyl)-propionamide	347.31	A
27		2-Methyl-2-(propane-2-sulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-propionamide	339.18	A
28		N-(4-tert-Butyl-thiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide	359	A
29		2-Cyclohexanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	379	A
30		N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide	343	A
31		N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropanesulfonyl-2-methyl-propionamide	315	C

32		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide	331	A
33		N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide	347	A
34		N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide	361	A
35		2-Methyl-2-(2-methyl-propane-2-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	353	A
36		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide	344	A
37		2-Methyl-2-(2-methyl-propane-2-sulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-propionamide	353	A
38		2-Methyl-2-(2-methyl-propane-2-sulfonyl)-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide	353	A
39		2-(1-Acetyl-piperidin-3-ylmethanesulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide	414	I

40		N-(5-tert-Butyl-isoxazol-3-yl)-2-(propane-2-sulfonyl)-propionamide	303	A
41		N-(5-tert-Butyl-isoxazol-3-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide	331	A
42		2-(Butane-2-sulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide	331	A
43		2-(Butane-2-sulfonyl)-N-(5-tert-butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-propionamide	344	A
44		2-(Butane-2-sulfonyl)-N-(4-tert-butyl-thiazol-2-yl)-2-methyl-propionamide	347	A
45		N-(4-tert-Butyl-thiazol-2-yl)-2-(propane-2-sulfonyl)-propionamide	319	A
46		N-(4-tert-Butyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide	347	A
47		N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide	361	A









48		2-(Butane-2-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	353	A
49		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-ethanesulfonyl-2-methyl-propionamide	316	A
50		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methanesulfonyl-2-methyl-propionamide	302	B
51		N-(4-tert-Butyl-thiazol-2-yl)-2-methanesulfonyl-2-methyl-propionamide	305	B
52		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide	331	D
53		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2,2,2-trifluoro-ethanesulfonyl)-propionamide	357	B
54		N-(5-tert-Butyl-isoxazol-3-yl)-2-methanesulfonyl-2-methyl-propionamide	289	B
55		2-Methanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	311	B









56		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	373	E
57		N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	389	E
58		N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	403	E
59		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	386	E
60		2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	395	E
61		2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-propionamide	395	E
62		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide	356	A
63		2-Cyclopentanesulfonyl-2-methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-propionamide	376	A

64		2-Cyclopentanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	365	A
65		2-Cyclopentanesulfonyl-2-methyl-N-(4-trifluoromethyl-thiazol-2-yl)-propionamide	371	A
66		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	359	E
67		N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	375	E
68		2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	381	E
69		2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	392	E
70		N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	389	E
71		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	372	E









72		N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-cyclopentanesulfonyl-2-methyl-propionamide	413/415	A
73		2-Cyclopentanesulfonyl-2-methyl-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide	365	A
74		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide	359	F
75		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide	356	F
76		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-(hexane-3-sulfonyl)-2-methyl-propionamide	372	F
77		N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide	357	A
78		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-propionamide	386	G
79		N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide	371	G









80		N-(5-tert-Butyl-isoxazol-3-yl)-2-ethanesulfonyl-2-methyl-propionamide	303	G
81		N-(5-tert-Butyl-isoxazol-3-yl)-2-cyanomethanesulfonyl-2-methyl-propionamide	314	G
82		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-propionamide	373	G
83		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-(2,2-dimethyl-propane-1-sulfonyl)-2-methyl-propionamide	358	G
84		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide	384	G
85		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide	384	G
86		2-Methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	381	G
87		2-Methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	395	G









88		2-(3,3-Dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	395	G
89		N-(5-tert-Butyl-isoxazol-3-yl)-2-(3,3-dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-propionamide	373	G
90		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-((S)-5-oxo-pyrrolidin-2-ylmethanesulfonyl)-propionamide	385	G
91		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-((R)-5-oxo-pyrrolidin-2-ylmethanesulfonyl)-propionamide	385	G
92		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide	345	G
93		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide	358	G
94		N-(5-tert-Butyl-isoxazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide	371	G
95		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(piperidin-4-ylmethanesulfonyl)-propionamide	372	H









96		2-Cyclohexylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	393	G
97		2-Methyl-2-(3-methyl-butane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	367	G
98		2-Methyl-2-(piperidin-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	394	H
99		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide	372	G
100		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(4-methyl-tetrahydro-pyran-4-sulfonyl)-propionamide	373	K
101		2-Cycloheptanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	393	G
102		2-Cyclopropylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	351	G
103		2-Cyclobutylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	365	G









104		N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide	329	G
105		N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide	343	G
106		N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-phenyl-propane-1-sulfonyl)-propionamide	393	G
107		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide	317	A
108		N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-2-sulfonyl)-propionamide	387/389	A
109		2-Methyl-2-(propane-2-sulfonyl)-N-(4-pyridin-3-yl-thiazol-2-yl)-propionamide	354	A
110		2-Methyl-2-(propane-2-sulfonyl)-N-(4-pyridin-2-yl-thiazol-2-yl)-propionamide	354	A
111		2-Methyl-2-(propane-2-sulfonyl)-N-(4-pyridin-4-yl-thiazol-2-yl)-propionamide	354	A









112		2-Methyl-2-(propane-2-sulfonyl)-N-quinolin-2-yl-propionamide	321	A
113		2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-3-yl-propionamide	321	A
114		N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	326	A
115		2-Methyl-2-(propane-1-sulfonyl)-N-(4,5,6,7-tetrahydro-benzothiazol-2-yl)-propionamide	331	A
116		2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-2-yl-propionamide	321	A
111 7		N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide	387/389	A
118		2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(propane-1-sulfonyl)-propionamide	350	A
119		N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	330	A

120		2-Methyl-N-(2-methyl-5-thiophen-2-yl-2H-pyrazol-3-yl)-2-(propane-1-sulfonyl)-propionamide	356	A
121		N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	326	A
122		2-Methyl-2-(propane-1-sulfonyl)-N-(4-pyridin-4-yl-thiazol-2-yl)-propionamide	354	A
123		2-Methyl-2-(propane-1-sulfonyl)-N-(4-trifluoromethyl-thiazol-2-yl)-propionamide	345	A
124		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	317	A
125		N-Benzothiazol-2-yl-2-methyl-2-(propane-1-sulfonyl)-propionamide	327	A
126		2-Methyl-2-(propane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	339	A
127		N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-methyl-propionamide	436	J









128		2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(propane-1-sulfonyl)-propionamide	367	A
129		N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	381	A
130		2-Methyl-N-(5-phenyl-pyridin-2-yl)-2-(propane-1-sulfonyl)-propionamide	347	A
131		N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	359	A
132		N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	347	A
133		N-(2,4-Dimethyl-5-phenyl-2H-pyrazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	364	A
134		N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide	331	A
135		N-(3-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	382	E









136		2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-thiazol-2-yl)-propionamide	401	E
137		N-(4-Cyclopropyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	373	E
138		N-Benzothiazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	383	E
139		2-Methyl-N-(4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	409	E
140		N-(5-tert-Butyl-[1,3,4]thiadiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	390	E
141		2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide	395	E
142		N-(4-Ethyl-pyridin-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	355	E
143		2-Methyl-N-(5-phenyl-[1,2,4]thiadiazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	410	E



144		N-Benzooxazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	367	E
145		2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	423	E
146		N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	437	E
147		N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	415	E
148		2-Methyl-N-(3-propyl-isoxazol-5-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	359	E
149		N-(5-Isopropyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	372	E
150		N-(2,4-Dimethyl-5-phenyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	420	E
151		N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	387	E

152		2-Methyl-N-naphthalen-1-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	376	E
153		2-Methyl-N-(5-methyl-pyridin-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	341	E
154		2-Methyl-N-quinolin-3-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	377	E
155		2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(3-trifluoromethyl-phenyl)-propionamide	394	E
156		N-(4-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	382	E
157		2-Methyl-N-quinolin-2-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	377	E
158		N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	443	E
159		2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-phenyl)-propionamide	394	E

160		2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	406	E
161		N-Isoquinolin-3-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	377	E
162		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	373	E
163		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	359	E
164		N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide	360	A
165		2-Cyclopentanesulfonyl-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide	363	A
166		N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide	343	G
167		N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide	329	G

168		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide	345	G
169		2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	393	E
170		N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide	343	A
171		N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide	371	G
172		2-(Butane-2-sulfonyl)-N-(3-tert-butyl-isoxazol-5-yl)-2-methyl-propionamide	331	G
173		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-propionamide	373	G
174		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide	331	G
175		1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-5-yl)-amide	385	E





176		N-(3-tert-Butyl-isoxazol-5-yl)-2-(4-fluorophenylmethanesulfonyl)-2-methylpropionamide	383	D
177		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide	359	G
178		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-ylmethanesulfonyl)-propionamide	359	G
179		1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-5-yl)-amide	373	E
180		2-Methyl-N-(4-pyridin-3-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	410	E
181		2-Methyl-N-(4-pyridin-2-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	410	E
182		2-Methyl-N-(4-pyridin-4-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	410	E
183		N-[4-(4-Fluoro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	427	E




184		N-(3-Isopropyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	359	E
185		2-Methyl-N-(3-phenyl-1,2,4-thiadiazol-5-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	410	E

5

The following compounds can also be made according the procedures described above:

Table 18: Examples

#	MOLSTRUCTURE	NAME
186		1-(Tetrahydro-pyran-4-sulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-5-yl)-amide
187		N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-sulfonyl)-propionamide
188		1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (5-phenyl-4H-1,2,4-triazol-3-yl)-amide
189		2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-sulfonyl)-propionamide

190		1-(Tetrahydro-pyran-4-sulfonyl)-cyclobutanecarboxylic acid (5-phenyl-4H-1,2,4-triazol-3-yl)-amide
191		N-(3-tert-Butyl-isoxazol-5-yl)-2-(1-methanesulfonyl-pyrrolidin-3-yl)methanesulfonyl)-2-methyl-propionamide
192		2-(1-Methanesulfonyl-pyrrolidine-3-sulfonyl)-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide

5

Assessment of Biological Properties

The biological properties of the compounds of the formula I were assessed using the assays described below.

10

A. Human CB1 and CB2 Receptor Binding:

Experimental Method:

CB2 membranes were purchased and made from HEK293 EBNA cells stably transfected with human CB2 receptor cDNA (Perkin Elmer Life and Analytical Sciences). CB1 membranes were isolated from HEK cells stably co-transfected with human CB1 receptor and Gα16 cDNA's. The membrane preparation was bound to scintillation beads (Ysi-Poly-L-lysine SPA beads, GE Healthcare) for 4 hours at room temperature in assay buffer containing 50mM Tris, pH 7.5, 2.5mM EDTA, 5mM MgCl₂, 0.8% fatty acid free Bovine Serum Albumin. Unbound membrane was removed by washing in assay buffer. Membrane-bead mixture was added to 96-well assay plates in the amounts of 15ug membrane per well (CB2) or 2.5ug per well (CB1) and 1mg SPA bead per well. Compounds were added to the membrane-bead mixture in

20

5 dose-response concentrations ranging from 1×10^{-5} M to 1×10^{-10} M with 0.25% DMSO, final. The competition reaction was initiated with the addition of ^3H -CP55940 (Perkin Elmer Life and Analytical Sciences) at a final concentration of 1.5nM (CB2) or 2.5nM (CB1). The reaction was incubated at room temperature for 18hours and read on TopCount NXT plate reader. Total and non-specific binding was determined in the absence and presence of 1.25uM
10 Win 55212 (Sigma). IC50 values for each compound were calculated as the concentration of compound that inhibits the specific binding of the radioactively labeled ligand to the receptor by 50% using the XLFit 4.1 four parameter logistic model. IC50 values were converted to inhibition constant (Ki) values using Cheng-Prusoff equation.

15 B. CB2R mediated modulation of cAMP synthesis:

Compounds of the invention were evaluated for their CB2 agonist or inverse agonistic activity in accordance with the following experimental method. Compounds which were shown to bind to CB2 by the binding assay described above but which were not shown to exhibit CB2R-mediated modulation of cAMP synthesis by this assay were presumed to be CB2
20 antagonists.

Experimental Method:

CHO cells expressing human CB2R (Euroscreen) were plated at a density of 5000 cells per well in 384 well plates and incubated overnight at 37°C. After removing the media, the cells
25 were treated with test compounds diluted in stimulation buffer containing 1mM IBMX, 0.25% BSA and 10uM Forskolin. The assay was incubated for 30 minutes at 37°C. Cells were lysed and the cAMP concentration was measured using DiscoverX –XS cAMP kit, following the manufacturer's protocol. In this setting, agonists will decrease forskolin induced production of cAMP while inverse agonists will further increase forskolin induced production of cAMP.
30 EC50 of agonists were calculated as follows. The maximal amount of cAMP produced by forskolin compared to the level of cAMP inhibited by 1uM CP55940 is defined as 100%. The EC50 value of each test compound was determined as the concentration at which 50% of the

- 5 forskolin-stimulated cAMP synthesis was inhibited. Data was analyzed using a four-parameter logistic model. (Model 205 of XLfit 4.0).

C. CB1R mediated modulation of cAMP synthesis:

- Compounds of the invention were evaluated for their CB1 agonist or inverse agonistic activity
10 in accordance with the following experimental method. Compounds which were shown to bind to CB1 by the binding assay described above but which were not shown to exhibit CB1R-mediated modulation of cAMP synthesis by this assay were presumed to be CB1 antagonists.

15 Experimental Method:

- CHO cells expressing human CB1R (Euroscreen) were plated at a density of 5000 cells per well in 384 well plates and incubated overnight at 37°C. After removing the media, the cells were treated with test compounds diluted in stimulation buffer containing 1mM IBMX, 0.25% BSA and 10uM Forskolin. The assay was incubated for 30 minutes at 37°C. Cells were lysed
20 and the cAMP concentration was measured using DiscoverX –XS cAMP kit, following the manufacturer's protocol. In this setting, agonists will decrease forskolin induced production of cAMP while inverse agonists will further increase forskolin induced production of cAMP. EC50 of agonists were calculated as follows. The maximal amount of cAMP produced by forskolin compared to the level of cAMP inhibited by 1uM CP55940 is defined as 100%. The
25 EC50 value of each test compound was determined as the concentration at which 50% of the forskolin-stimulated cAMP synthesis was inhibited. Data was analyzed using a four-parameter logistic model. (Model 205 of XLfit 4.0).

Compounds Having Agonist Activity

NAME	CB2 EC ₅₀ [nM]
------	---------------------------------

N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide	0.2
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide	25.8
1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (5-tert-butyl-isoxazol-3-yl)-amide	0.5
1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (4-tert-butyl-thiazol-2-yl)-amide	11.7
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide	12.1
N-(5-tert-Butyl-isoxazol-3-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide	16.6
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	61.9
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	19.2
N-(4-tert-Butyl-thiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide	19.0
2-Cyclohexanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	11.8
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide	2.4
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide	61.1
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide	10.7
2-(Butane-2-sulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide	18.7
2-(Butane-2-sulfonyl)-N-(4-tert-butyl-thiazol-2-yl)-2-methyl-propionamide	14.4
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide	44.6
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	4.5
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	23.0
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	0.2
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	72.3
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	0.7
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	12.4

2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	62.4
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	0.3
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-cyclopentanesulfonyl-2-methyl-propionamide	1.5
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide	70.5
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide	33.0
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide	0.0
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide	0.6
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-propionamide	0.2
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide	6.5
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide	23.8
2-(3,3-Dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	41.5
N-(5-tert-Butyl-isoxazol-3-yl)-2-(3,3-dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-propionamide	39.8
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide	0.2
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide	15.4
N-(5-tert-Butyl-isoxazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide	0.1
2-Cyclohexylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	15.4
2-Methyl-2-(3-methyl-butane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	19.9

N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(4-methyl-tetrahydro-pyran-4-sulfonyl)-propionamide	9.4
2-Cycloheptanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide	7.1
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide	34.3
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide	3.4
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-2-sulfonyl)-propionamide	44.9
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	12.3
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide	13.0
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	89.7
N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-methyl-propionamide	44.1
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(propane-1-sulfonyl)-propionamide	52.2
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	10.8
N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	31.1
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide	7.2
N-(3-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	14.3
N-Benzothiazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	27.9
2-Methyl-N-(4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	7.1
2-Methyl-N-(5-phenyl-[1,2,4]thiadiazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	17.6
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	2.5
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	2.0
N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	8.3
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(tetrahydro-pyran-4-	74.3

ylmethanesulfonyl)-propionamide	
2-Methyl-N-quinolin-3-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	86.0
N-(4-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	3.3
2-Methyl-N-quinolin-2-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	5.8
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	2.2
N-Isoquinolin-3-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	99.1
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	4.0
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide	0.1
N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide	18.6
2-Cyclopentanesulfonyl-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide	3.8
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide	3.7
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide	12.5
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide	0.04
2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	1.7
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide	0.9
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide	0.1
2-(Butane-2-sulfonyl)-N-(3-tert-butyl-isoxazol-5-yl)-2-methyl-propionamide	3.3
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-propionamide	0.2
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide	1.6
1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-5-yl)-amide	6.2
N-(3-tert-Butyl-isoxazol-5-yl)-2-(4-fluoro-phenylmethanesulfonyl)-2-methyl-propionamide	17.0
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-	31.5

propionamide	
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-ylmethanesulfonyl)-propionamide	18.5
N-(3-sec-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	20.0
N-[4-(4-Fluoro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide	5.5

5

Through the use of the above described assays the following compounds were found to exhibit agonistic activity and thus to be particularly well suited for the treatment of pain as well as for the treatment of inflammation.

10

N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide

N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide

N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide

1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (5-tert-butyl-isoxazol-3-yl)-amide

15 1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (4-tert-butyl-thiazol-2-yl)-amide

N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide

N-(4-tert-Butyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide

N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide

20 N-(5-tert-Butyl-isoxazol-3-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide

N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide

2-Methyl-2-(propane-1-sulfonyl)-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide

2-Methyl-N-naphthalen-2-yl-2-(propane-2-sulfonyl)-propionamide

2-Methyl-2-(propane-2-sulfonyl)-N-(4,5,6,7-tetrahydro-benzothiazol-2-yl)-propionamide

25 2-Methyl-N-(2-methyl-5-thiophen-2-yl-2H-pyrazol-3-yl)-2-(propane-2-sulfonyl)-propionamide

- 5 N-Isoquinolin-3-yl-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-Benzothiazol-2-yl-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-quinolin-3-yl-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
- 10 2-Methyl-N-(4-phenyl-thiazol-2-yl)-2-(propane-2-sulfonyl)-propionamide
2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(propane-2-sulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-propionamide
- 15 N-(4-tert-Butyl-thiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclohexanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
- 20 N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-
propionamide
- 25 2-Methyl-2-(2-methyl-propane-2-sulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(2-methyl-propane-2-sulfonyl)-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide
2-(1-Acetyl-piperidin-3-ylmethanesulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-
propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
- 30 2-(Butane-2-sulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide
2-(Butane-2-sulfonyl)-N-(5-tert-butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-propionamide
2-(Butane-2-sulfonyl)-N-(4-tert-butyl-thiazol-2-yl)-2-methyl-propionamide

- 5 N-(4-tert-Butyl-thiazol-2-yl)-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
2-(Butane-2-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-ethanesulfonyl-2-methyl-propionamide
10 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methanesulfonyl-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2,2,2-trifluoro-ethanesulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methanesulfonyl-2-methyl-propionamide
15 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
20 propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-
ylmethanesulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
25 2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
30 2-Cyclopentanesulfonyl-2-methyl-N-(4-trifluoromethyl-thiazol-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide

- 5 2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide
- 10 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-
15 propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-(hexane-3-sulfonyl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
- 20 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-
ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-ethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyanomethanesulfonyl-2-methyl-propionamide
- 25 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-(2,2-dimethyl-propane-1-sulfonyl)-2-methyl-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
- 30 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-
propionamide

- 5 2-Methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-(3,3-Dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
- 10 N-(5-tert-Butyl-isoxazol-3-yl)-2-(3,3-dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-((S)-5-oxo-pyrrolidin-2-ylmethanesulfonyl)-propionamide
- 15 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-((R)-5-oxo-pyrrolidin-2-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
- 20 N-(5-tert-Butyl-isoxazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(piperidin-4-ylmethanesulfonyl)-propionamide
2-Cyclohexylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(3-methyl-butane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(piperidin-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
- 25 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(4-methyl-tetrahydro-pyran-4-sulfonyl)-propionamide
2-Cycloheptanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
- 30 2-Cyclopropylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cyclobutylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide

- 5 N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-phenyl-propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-2-sulfonyl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-(4-pyridin-3-yl-thiazol-2-yl)-propionamide
10 2-Methyl-2-(propane-2-sulfonyl)-N-(4-pyridin-2-yl-thiazol-2-yl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-(4-pyridin-4-yl-thiazol-2-yl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-quinolin-2-yl-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-3-yl-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
15 2-Methyl-2-(propane-1-sulfonyl)-N-(4,5,6,7-tetrahydro-benzothiazol-2-yl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-2-yl-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
20 2-Methyl-N-(2-methyl-5-thiophen-2-yl-2H-pyrazol-3-yl)-2-(propane-1-sulfonyl)-
propionamide
N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-(4-pyridin-4-yl-thiazol-2-yl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-(4-trifluoromethyl-thiazol-2-yl)-propionamide
25 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-Benzothiazol-2-yl-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-
methyl-propionamide
30 2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(propane-1-sulfonyl)-propionamide
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-N-(5-phenyl-pyridin-2-yl)-2-(propane-1-sulfonyl)-propionamide

- 5 N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(2,4-Dimethyl-5-phenyl-2H-pyrazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
10 2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-thiazol-2-yl)-
propionamide
N-(4-Cyclopropyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-Benzothiazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
15 2-Methyl-N-(4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-[1,3,4]thiadiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-pyridin-2-yl)-
propionamide
20 N-(4-Ethyl-pyridin-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-phenyl-[1,2,4]thiadiazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-Benzooxazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
25 propionamide
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
30 2-Methyl-N-(3-propyl-isoxazol-5-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(5-Isopropyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-
ylmethanesulfonyl)-propionamide

- 5 N-(2,4-Dimethyl-5-phenyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-naphthalen-1-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 10 2-Methyl-N-(5-methyl-pyridin-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-3-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(3-trifluoromethyl-phenyl)-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 15 2-Methyl-N-quinolin-2-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-phenyl)-propionamide
- 20 2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-Isoquinolin-3-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 25 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
- 30 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide

- 5 N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
 N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
 2-(Butane-2-sulfonyl)-N-(3-tert-butyl-isoxazol-5-yl)-2-methyl-propionamide
 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-
 propionamide
- 10 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide
 1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-
 5-yl)-amide
 N-(3-tert-Butyl-isoxazol-5-yl)-2-(4-fluoro-phenylmethanesulfonyl)-2-methyl-propionamide
 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-
 15 propionamide
 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-ylmethanesulfonyl)-
 propionamide
 N-(3-sec-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
 propionamide
- 20 N-[4-(4-Fluoro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
 propionamide
 2-Methyl-N-(4-pyridin-3-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
 propionamide
 2-Methyl-N-(4-pyridin-2-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
 25 propionamide
 2-Methyl-N-(4-pyridin-4-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
 propionamide
 N-(3-Isopropyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
 propionamide
- 30 2-Methyl-N-(3-phenyl-1,2,4-thiadiazol-5-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
 propionamide

Of the above compounds, the following are preferred:

- 5
- N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (5-tert-butyl-isoxazol-3-yl)-amide
- 10 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
- 15 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
- 20 2-Cyclohexanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
- 25 N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
2-(Butane-2-sulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide
2-(Butane-2-sulfonyl)-N-(4-tert-butyl-thiazol-2-yl)-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
- 30 N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide

- 5 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
10 propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-
ylmethanesulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
15 2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-
propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
20 2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide
25 N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-
30 propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-(hexane-3-sulfonyl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide

- 5 N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-
10 propionamide
2-Methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
2-(3,3-Dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
15 N-(5-tert-Butyl-isoxazol-3-yl)-2-(3,3-dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-
propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-
propionamide
20 N-(5-tert-Butyl-isoxazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
2-Cyclohexylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(3-methyl-butane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cycloheptanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cyclopropylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
25 2-Cyclobutylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-2-sulfonyl)-propionamide
30 2-Methyl-2-(propane-2-sulfonyl)-N-quinolin-2-yl-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-3-yl-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide

- 5 2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-2-yl-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-Benzothiazol-2-yl-2-methyl-2-(propane-1-sulfonyl)-propionamide
- 10 N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-methyl-propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(propane-1-sulfonyl)-propionamide
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
- 15 N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-Benzothiazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-[1,3,4]thiadiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
- 20 propionamide
2-Methyl-N-(5-phenyl-[1,2,4]thiadiazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 25 N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(3-propyl-isoxazol-5-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 30 N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-3-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide

- 5 2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(3-trifluoromethyl-phenyl)-
propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-2-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
10 propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-phenyl)-
propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
15 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
20 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
25 2-(Butane-2-sulfonyl)-N-(3-tert-butyl-isoxazol-5-yl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-
propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide
1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-
30 5-yl)-amide
N-(3-tert-Butyl-isoxazol-5-yl)-2-(4-fluoro-phenylmethanesulfonyl)-2-methyl-propionamide

- 5 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-ylmethanesulfonyl)-propionamide
N-(3-sec-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
10 propionamide
N-[4-(4-Fluoro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(4-pyridin-3-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
15 2-Methyl-N-(4-pyridin-2-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(4-pyridin-4-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 20 Of the above compounds, the following are most preferred:
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (5-tert-butyl-isoxazol-3-yl)-amide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-
25 propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Cyclohexanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
30 N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide

- 5 N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-
propionamide
2-(Butane-2-sulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide
2-(Butane-2-sulfonyl)-N-(4-tert-butyl-thiazol-2-yl)-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide
- 10 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
- 15 propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
- 20 2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-
- 25 propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-
propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
- 30 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide

- 5 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
2-(3,3-Dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(3,3-dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-
- 10 propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
- 15 2-Cyclohexylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(3-methyl-butane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cycloheptanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
- 20 N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-
- 25 methyl-propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(propane-1-sulfonyl)-propionamide
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
- 30 N-Benzothiazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-phenyl-[1,2,4]thiadiazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide

- 5 2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(tetrahydro-pyran-4-
10 ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-3-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-2-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
15 propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
20 2-Cyclopentanesulfonyl-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
25 propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
2-(Butane-2-sulfonyl)-N-(3-tert-butyl-isoxazol-5-yl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-
30 propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide

- 5 1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-5-yl)-amide
N-(3-tert-Butyl-isoxazol-5-yl)-2-(4-fluoro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide
10 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-ylmethanesulfonyl)-propionamide
N-(3-sec-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[4-(4-Fluoro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
15 propionamide

Therapeutic Use

As can be demonstrated by the assays described above, the compounds of the invention are useful in modulating the CB2 receptor function. By virtue of this fact, these compounds have
20 therapeutic use in treating disease-states and conditions mediated by the CB2 receptor function or that would benefit from modulation of the CB2 receptor function.

As the compounds of the invention modulate the CB2 receptor function, they have very useful anti-inflammatory and immune-suppressive activity and they can be used in patients as drugs,
25 particularly in the form of pharmaceutical compositions as set forth below, for the treatment of disease-states and conditions.

As noted before, those compounds which are CB2 agonists can also be employed for the treatment of pain.

30

5 The agonist, antagonist and inverse agonist compounds according to the invention can be used
in patients as drugs for the treatment of the following disease-states or indications that are
accompanied by inflammatory processes:

- 10 (i) Lung diseases: e.g. asthma, bronchitis, allergic rhinitis, emphysema, adult
respiratory distress syndrome (ARDS), pigeon fancier's disease, farmer's lung, chronic
obstructive pulmonary disease (COPD), asthma including allergic asthma (atopic or
non-atopic) as well as exercise-induced bronchoconstriction, occupational asthma,
viral- or bacterial exacerbation of asthma, other non-allergic asthmas and "wheezy-
15 infant syndrome", pneumoconiosis, including aluminosis, anthracosis, asbestosis,
chalicosis, ptilosis, siderosis, silicosis, tabacosis and byssinosis;
- (ii) Rheumatic diseases or autoimmune diseases or musculoskeletal diseases: all
forms of rheumatic diseases, especially rheumatoid arthritis, acute rheumatic fever, and
polymyalgia rheumatica; reactive arthritis; rheumatic soft tissue diseases;
inflammatory soft tissue diseases of other genesis; arthritic symptoms in degenerative
20 joint diseases (arthroses); tendinitis, bursitis, osteoarthritis, traumatic arthritis;
collagenoses of any genesis, e.g., systemic lupus erythematosus, scleroderma,
polymyositis, dermatomyositis, Sjögren syndrome, Still disease, Felty syndrome; and
osteoporosis and other bone resorption diseases;
- (iii) Allergic diseases: all forms of allergic reactions, e.g., angioneurotic edema, hay
25 fever, insect bites, allergic reactions to drugs, blood derivatives, contrast agents, etc.,
anaphylactic shock (anaphylaxis), urticaria, angioneurotic edema, and contact
dermatitis;
- (iv) Vascular diseases: panarteritis nodosa, polyarteritis nodosa, periarteritis
nodosa, arteritis temporalis, Wegner granulomatosis, giant cell arthritis,
30 atherosclerosis, reperfusion injury and erythema nodosum;
- (v) Dermatological diseases: e.g. dermatitis, psoriasis; sunburn, burns, eczema;

- 5 (vi) Renal diseases: e.g. nephrotic syndrome; and all types of nephritis, e.g., glomerulonephritis; pancreatitis;
- (vii) Hepatic diseases: e.g. acute liver cell disintegration; acute hepatitis of various genesis, e.g., viral, toxic, drug-induced; and chronically aggressive and/or chronically intermittent hepatitis;
- 10 (viii) Gastrointestinal diseases: e.g. inflammatory bowel diseases, irritable bowel syndrome, regional enteritis (Crohn's disease), colitis ulcerosa; gastritis; aphthous ulcer, celiac disease, regional ileitis, gastroesophageal reflux disease;
- (ix) Neuroprotection: e.g. in the treatment of neurodegeneration following stroke; cardiac arrest; pulmonary bypass; traumatic brain injury; spinal cord injury or the like;
- 15 (x) Eye diseases: allergic keratitis, uveitis, or iritis; conjunctivitis; blepharitis; neuritis nervi optici; choroiditis; glaucoma and sympathetic ophthalmia;
- (xi) Diseases of the ear, nose, and throat (ENT) area: e.g. tinnitus; allergic rhinitis or hay fever; otitis externa; caused by contact eczema, infection, etc.; and otitis media;
- (xii) Neurological diseases: e.g. brain edema, particularly tumor-related brain edema; multiple sclerosis; acute encephalomyelitis; meningitis; acute spinal cord injury; trauma; dementia, particularly degenerative dementia (including senile dementia, Alzheimer's disease; Parkinson's disease and Creutzfeldt-Jacob disease; Huntington's chorea, Pick's disease; motor neuron disease), vascular dementia (including multi-infarct dementia) as well as dementia associated with intracranial space occupying lesions; infections and related conditions (including HIV infection);
- 20 Guillain-Barre syndrome; myasthenia gravis, stroke; and various forms of seizures, e.g., nodding spasms;
- (xiii) Blood diseases: acquired hemolytic anemia; aplastic anemia, and idiopathic thrombocytopenia;
- 25 (xiv) Tumor diseases: acute lymphatic leukemia; Hodgkin's disease, malignant lymphoma; lymphogranulomatosis; lymphosarcoma; solid malignant tumors; extensive metastases,;
- 30

- 5 (xv) Endocrine diseases: endocrine ophthalmopathy; endocrine orbitopathy;
thyrotoxic crisis; Thyroiditis de Quervain; Hashimoto thyroiditis; Morbus Basedow;
granulomatous thyroiditis; struma lymphomatosa; and Graves disease; type I diabetes
(insulin-dependent diabetes);
- (xvi) Organ and tissue transplantations and graft-versus-host diseases;
- 10 (xvii) Severe states of shock, e.g., septic shock, anaphylactic shock, and systemic
inflammatory response syndrome (SIRS);
- (xviii) Acute pain such as dental pain, perioperative, post-operative pain, traumatic
pain, muscle pain, pain in burned skin, sun burn, trigeminal neuralgia, sun burn; spasm
of the gastrointestinal tract or uterus, colics;
- 15 (xix) Visceral pain such as pain associated with chronic pelvic pain, pancreatitis,
peptic ulcer, interstitial cystitis, renal colic, angina, dysmenorrhoea, menstruation,
gynaecological pain, irritable bowel syndrome (IBS), non-ulcer dyspepsia, non-cardiac
chest pain, myocardial ischemia;
- (xx) Neuropathic pain such as low back pain, non-herpetic neuralgia, post herpetic
20 neuralgia, diabetic neuropathy, nerve injury, acquired immune deficiency syndrome
(AIDS) related neuropathic pain, head trauma, painful traumatic mononeuropathy,
toxin and chemotherapy induced pain, phantom limb pain, painful polyneuropathy,
thalamic pain syndrome, post-stroke pain, central nervous system injury, post surgical
pain, stump pain, repetitive motion pain, pain induced by post mastectomy syndrome,
25 multiple sclerosis, root avulsions, postthoracotomy syndrome, neuropathic pain
associated hyperalgesia and allodynia.
- (xxi) Inflammatory/nociceptive pain induced by or associated with disorders such as
osteoarthritis, rheumatoid arthritis, rheumatic disease, teno-synovitis, gout, vulvodynia,
myofascial pain (muscular injury, fibromyalgia), tendonitis, osteoarthritis, juvenile
30 arthritis, spondylitis, gouty arthritis, psoriatic arthritis, musculoskeletal pain,
fibromyalgia, sprains and strains, sympathetically maintained pain, myositis, pain

- 5 associated with migraine, toothache, influenza and other viral infections such as the common cold, rheumatic fever, systemic lupus erythematosus;
- (xxii) Cancer pain induced by or associated with tumors such as lymphatic leukemia; Hodgkin's disease, malignant lymphoma; lymphogranulomatoses; lymphosarcoma; solid malignant tumors; extensive metastases;
- 10 (xxiii) Headache such as cluster headache, migraine with and without aura, tension type headache, headache with different origins, headache disorders including prophylactic and acute use;
- (xxiv) various other disease-states or conditions including, restenosis following percutaneous transluminal coronary angioplasty, acute and chronic pain,
- 15 atherosclerosis, reperfusion injury, congestive heart failure, myocardial infarction, thermal injury, multiple organ injury secondary to trauma, necrotizing enterocolitis and syndromes associated with hemodialysis, leukopheresis, and granulocyte transfusion, sarcoidosis, gingivitis, pyrexia. edema resulting from trauma associated with burns, sprains or fracture, cerebral oedema and angioedema, Diabetes such as
- 20 diabetic vasculopathy, diabetic neuropathy, diabetic retinopathy, post capillary resistance or diabetic symptoms associated with insulinitis (e.g. hyperglycemia, diuresis, proteinuria and increased nitrite and kallikrein urinary excretion).

Other indications include: epilepsy, septic shock e.g. as antihypovolemic and/or

25 antihypotensive agents, cancer, sepsis, osteoporosis, benign prostatic hyperplasia and hyperactive bladder, pruritis, vitiligo, general gastrointestinal disorders, disturbances of visceral motility at respiratory, genitourinary, gastrointestinal or vascular regions, wounds, burns, tissue damage and postoperative fever, syndromes associated with Itching.

30 Besides being useful for human treatment, these compounds are also useful for veterinary treatment of companion animals, exotic animals and farm animals, including mammals, rodents, and the like.

5

For treatment of the above-described diseases and conditions, a therapeutically effective dose will generally be in the range from about 0.01 mg to about 100 mg/kg of body weight per dosage of a compound of the invention; preferably, from about 0.1 mg to about 20 mg/kg of body weight per dosage. For example, for administration to a 70 kg person, the dosage range
10 would be from about 0.7 mg to about 7000 mg per dosage of a compound of the invention, preferably from about 7.0 mg to about 1400 mg per dosage. Some degree of routine dose optimization may be required to determine an optimal dosing level and pattern. The active ingredient may be administered from 1 to 6 times a day.

15 Combination Therapy

These compounds may also be employed in combination therapies with the following compounds:

non-steroidal antiinflammatory drugs (NSAIDs) including COX-2 inhibitors such as propionic acid derivatives (alminoprofen, benoxaprofen, bucloxic acid, carprofen, fenhufen, fenoprofen,
20 flurbiprofen, ibuprofen, indoprofen, ketoprofen, mioprofen, naproxen, oxaprozin, pirprofen, pranoprofen, suprofen, tiaprofenic acid, and tioxaprofen), acetic acid derivatives (indomethacin, acemetacin, alclofenac, clidanac, diclofenac, fenclofenac, fenclozic acid, fentiazac, furofenac, ibufenac, isoxepac, oxpinac, sulindac, tiopinac, tolmetin, zidometacin, and zomepirac), fenamic acid derivatives (meclofenamic acid, mefe- namic acid, and
25 tolfenamic acid), biphenyl- carboxylic acid derivatives, oxicams (isoxicam, meloxicam, piroxicam, sudoxicam and tenoxicam), salicylates (acetyl salicylic acid, sulfasalazine) and the pyrazolones (apazone, bezpiperylon, feprazone, mofebutazone, oxyphenbutazone, phenylbutazone), and the coxibs (celecoxib, valecoXID, rofecoxib and etoricoxib);
opiate receptor agonists such as morphine, propoxyphene (Darvon), tramadol, buprenorphin;
30 sodium channel blockers such as carbamazepine, mexiletine, lamotrigine, pregabalin, tectin, NW-1029, CGX-1002;

- 5 N-type calcium channel blockers such as Ziconotide, NMED-160, SPI-860;
serotonergic and noradrenergic modulators such as SR-57746, paroxetine, duloxetine,
clonidine, amitriptyline, citalopram;
corticosteroids such as betamethasone, budesonide, cortisone, dexamethasone, hydrocortisone,
methylprednisolone, prednisolone, prednisone and triamcinolone;
- 10 histamine H1 receptor antagonists such as brompheniramine, chlorpheniramine,
dexchlorpheniramine, triprolidine, clemastine, diphenhydramine, diphenylpyraline,
tripelennamine, hydroxyzine, methdiazine, promethazine, trimeprazine, azatadine,
cypheptadine, antazoline, pheniramine pyrilamine, astemizole, terfenadine, loratadine,
cetirizine, desloratadine, fexofenadine and levocetirizine;
- 15 histamine H2 receptor antagonists such as cimetidine, famotidine and ranitidine;
proton pump inhibitors such as omeprazole, pantoprazole and esomeprazole;
leukotriene antagonists and 5-lipoxygenase inhibitors such as zafirlukast, montelukast,
pranlukast and zileuton;
local anesthetics such as ambroxol, lidocaine;
- 20 VR1 agonists and antagonists such as NGX-4010, WL-1002, ALGRX-4975, WL-10001,
AMG-517;
nicotinic acetylcholine receptor agonists such as ABT-202, A-366833, ABT-594; BTG-102,
A-85380, CGX1204;
P2X3 receptor antagonists such as A-317491, ISIS-13920, AZD-9056;
- 25 NGF agonists and antagonists such as RI-724, RI-1024, AMG-819, AMG-403, PPH 207;
NK1 and NK2 antagonists such as DA-5018, R-116301; CP-728663, ZD-2249;
NMDA antagonist such as NER-MD-11, CNS-5161, EAA-090, AZ-756, CNP-3381;
potassium channel modulators such as CL-888, ICA-69673, retigabine;

5 GABA modulators such as lacosamide;
serotonergic and noradrenergic modulators such as SR-57746, paroxetine, duloxetine,
clonidine, amitriptyline, citalopram, flibanserin; and
combination with anti-migraine drugs like sumatriptan, zolmitriptan, naratriptan, eletriptan.

10

General Administration and Pharmaceutical Compositions

When used as pharmaceuticals, the compounds of the invention are typically administered in the form of a pharmaceutical composition. Such compositions can be prepared using procedures well known in the pharmaceutical art and comprise at least one compound of the
15 invention. The compounds of the invention may also be administered alone or in combination with adjuvants that enhance stability of the compounds of the invention, facilitate administration of pharmaceutical compositions containing them in certain embodiments, provide increased dissolution or dispersion, increased inhibitory activity, provide adjunct therapy, and the like. The compounds according to the invention may be used on their own or
20 in conjunction with other active substances according to the invention, optionally also in conjunction with other pharmacologically active substances. In general, the compounds of this invention are administered in a therapeutically or pharmaceutically effective amount, but may be administered in lower amounts for diagnostic or other purposes.

25 Administration of the compounds of the invention, in pure form or in an appropriate pharmaceutical composition, can be carried out using any of the accepted modes of administration of pharmaceutical compositions. Thus, administration can be, for example, orally, buccally (e.g., sublingually), nasally, parenterally, topically, transdermally, vaginally, or rectally, in the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such
30 as, for example, tablets, suppositories, pills, soft elastic and hard gelatin capsules, powders, solutions, suspensions, or aerosols, or the like, preferably in unit dosage forms suitable for simple administration of precise dosages. The pharmaceutical compositions will generally

5 include a conventional pharmaceutical carrier or excipient and a compound of the invention as the/an active agent, and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, vehicles, or combinations thereof. Such pharmaceutically acceptable excipients, carriers, or additives as well as methods of making pharmaceutical compositions for various modes or administration are well-known to those of skill in the art.

10 The state of the art is evidenced, e.g., by *Remington: The Science and Practice of Pharmacy*, 20th Edition, A. Gennaro (ed.), Lippincott Williams & Wilkins, 2000; *Handbook of Pharmaceutical Additives*, Michael & Irene Ash (eds.), Gower, 1995; *Handbook of Pharmaceutical Excipients*, A.H. Kibbe (ed.), American Pharmaceutical Ass'n, 2000; H.C. Ansel and N.G. Popovich, *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 5th ed.,

15 Lea and Febiger, 1990; each of which is incorporated herein by reference in their entireties to better describe the state of the art.

As one of skill in the art would expect, the forms of the compounds of the invention utilized in a particular pharmaceutical formulation will be selected (e.g., salts) that possess suitable

20 physical characteristics (e.g., water solubility) that is required for the formulation to be efficacious.

Pharmaceutical compositions suitable for buccal (sub-lingual) administration include lozenges comprising a compound of the present invention in a flavored base, usually sucrose, and

25 acacia or tragacanth, and pastilles comprising the compound in an inert base such as gelatin and glycerin or sucrose and acacia.

Pharmaceutical compositions suitable for parenteral administration comprise sterile aqueous preparations of a compound of the present invention. These preparations are preferably

30 administered intravenously, although administration can also be effected by means of subcutaneous, intramuscular, or intradermal injection. Injectable pharmaceutical formulations are commonly based upon injectable sterile saline, phosphate-buffered saline, oleaginous

5 suspensions, or other injectable carriers known in the art and are generally rendered sterile and isotonic with the blood. The injectable pharmaceutical formulations may therefore be provided as a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, including 1,3-butanediol, water, Ringer's solution, isotonic sodium chloride solution, fixed oils such as synthetic mono- or diglycerides, fatty acids such as oleic acid, and
10 the like. Such injectable pharmaceutical formulations are formulated according to the known art using suitable dispersing or setting agents and suspending agents. Injectable compositions will generally contain from 0.1 to 5% w/w of a compound of the invention.

Solid dosage forms for oral administration of the compounds include capsules, tablets, pills,
15 powders, and granules. For such oral administration, a pharmaceutically acceptable composition containing a compound(s) of the invention is formed by the incorporation of any of the normally employed excipients, such as, for example, pharmaceutical grades of mannitol, lactose, starch, pregelatinized starch, magnesium stearate, sodium saccharine, talcum, cellulose ether derivatives, glucose, gelatin, sucrose, citrate, propyl gallate, and the
20 like. Such solid pharmaceutical formulations may include formulations, as are well-known in the art, to provide prolonged or sustained delivery of the drug to the gastrointestinal tract by any number of mechanisms, which include, but are not limited to, pH sensitive release from the dosage form based on the changing pH of the small intestine, slow erosion of a tablet or capsule, retention in the stomach based on the physical properties of the formulation,
25 bioadhesion of the dosage form to the mucosal lining of the intestinal tract, or enzymatic release of the active drug from the dosage form.

Liquid dosage forms for oral administration of the compounds include emulsions, microemulsions, solutions, suspensions, syrups, and elixirs, optionally containing
30 pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol and the like. These compositions can also contain additional adjuvants such as wetting, emulsifying, suspending, sweetening, flavoring, and perfuming agents.

5

Topical dosage forms of the compounds include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants, eye ointments, eye or ear drops, impregnated dressings and aerosols, and may contain appropriate conventional additives such as preservatives, solvents to assist drug penetration and emollients in ointments and creams. Topical application may be once or more than once per day depending upon the usual medical considerations. Furthermore, preferred compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles. The formulations may also contain compatible conventional carriers, such as cream or ointment bases and ethanol or oleyl alcohol for lotions. Such carriers may be present as from about 1% up to about 98% of the formulation, more usually they will form up to about 80% of the formulation.

Transdermal administration is also possible. Pharmaceutical compositions suitable for transdermal administration can be presented as discrete patches adapted to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen. Such patches suitably contain a compound of the invention in an optionally buffered, aqueous solution, dissolved and/or dispersed in an adhesive, or dispersed in a polymer. A suitable concentration of the active compound is about 1% to 35%, preferably about 3% to 15%.

For administration by inhalation, the compounds of the invention are conveniently delivered in the form of an aerosol spray from a pump spray device not requiring a propellant gas or from a pressurized pack or a nebulizer with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, tetrafluoroethane, heptafluoropropane, carbon dioxide, or other suitable gas. In any case, the aerosol spray dosage unit may be determined by providing a valve to deliver a metered amount so that the

5 resulting metered dose inhaler (MDI) is used to administer the compounds of the invention in a reproducible and controlled way. Such inhaler, nebulizer, or atomizer devices are known in the prior art, for example, in PCT International Publication Nos. WO 97/12687 (particularly Figure 6 thereof, which is the basis for the commercial RESPIMAT® nebulizer); WO 94/07607; WO 97/12683; and WO 97/20590, to which reference is hereby made and each of
10 which is incorporated herein by reference in their entireties.

Rectal administration can be effected utilizing unit dose suppositories in which the compound is admixed with low-melting water-soluble or insoluble solids such as fats, cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various
15 molecular weights, or fatty acid esters of polyethylene glycols, or the like. The active compound is usually a minor component, often from about 0.05 to 10% by weight, with the remainder being the base component.

In all of the above pharmaceutical compositions, the compounds of the invention are
20 formulated with an acceptable carrier or excipient. The carriers or excipients used must, of course, be acceptable in the sense of being compatible with the other ingredients of the composition and must not be deleterious to the patient. The carrier or excipient can be a solid or a liquid, or both, and is preferably formulated with the compound of the invention as a unit-dose composition, for example, a tablet, which can contain from 0.05% to 95% by weight of
25 the active compound. Such carriers or excipients include inert fillers or diluents, binders, lubricants, disintegrating agents, solution retardants, resorption accelerators, absorption agents, and coloring agents. Suitable binders include starch, gelatin, natural sugars such as glucose or β -lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like.
30 Lubricants include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators include starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

5

Pharmaceutically acceptable carriers and excipients encompass all the foregoing additives and the like.

Examples of Pharmaceutical Formulations

10

A. TABLETS	
Component	Amount per tablet (mg)
active substance	100
lactose	140
corn starch	240
polyvinylpyrrolidone	15
magnesium stearate	5
TOTAL	500

The finely ground active substance, lactose, and some of the corn starch are mixed together. The mixture is screened, then moistened with a solution of polyvinylpyrrolidone in water, kneaded, wet-granulated and dried. The granules, the remaining corn starch and the
 15 magnesium stearate are screened and mixed together. The mixture is compressed to produce tablets of suitable shape and size.

B. TABLETS	
Component	Amount per tablet (mg)
active substance	80
lactose	55
corn starch	190
polyvinylpyrrolidone	15
magnesium stearate	2

microcrystalline cellulose	35
sodium-carboxymethyl starch	23
TOTAL	400

5

The finely ground active substance, some of the corn starch, lactose, microcrystalline cellulose, and polyvinylpyrrolidone are mixed together, the mixture is screened and worked with the remaining corn starch and water to form a granulate which is dried and screened. The sodium-carboxymethyl starch and the magnesium stearate are added and mixed in and the mixture is compressed to form tablets of a suitable size.

10

C. COATED TABLETS	
Component	Amount per tablet (mg)
active substance	5
lactose	30
corn starch	41.5
polyvinylpyrrolidone	3
magnesium stearate	0.5
TOTAL	90

The active substance, corn starch, lactose, and polyvinylpyrrolidone are thoroughly mixed and moistened with water. The moist mass is pushed through a screen with a 1 mm mesh size, dried at about 45°C and the granules are then passed through the same screen. After the magnesium stearate has been mixed in, convex tablet cores with a diameter of 6 mm are compressed in a tablet-making machine. The tablet cores thus produced are coated in known manner with a covering consisting essentially of sugar and talc. The finished coated tablets are polished with wax.

20

D. CAPSULES

Component	Amount per capsule (mg)
active substance	50
corn starch	268.5
magnesium stearate	1.5
TOTAL	320

5

The substance and corn starch are mixed and moistened with water. The moist mass is screened and dried. The dry granules are screened and mixed with magnesium stearate. The finished mixture is packed into size 1 hard gelatine capsules.

E. AMPOULE SOLUTION	
Component	Amount per ampoule
active substance	50 mg
sodium chloride	50 mg
water for inj.	5 mL

10

The active substance is dissolved in water at its own pH or optionally at pH 5.5 to 6.5 and sodium chloride is added to make it isotonic. The solution obtained is filtered free from pyrogens and the filtrate is transferred under aseptic conditions into ampoules which are then sterilized and sealed by fusion. The ampoules contain 5 mg, 25 mg, and 50 mg of active substance.

15

F. SUPPOSITORIES	
Component	Amount per suppository (mg)
active substance	50
solid fat	1650
TOTAL	1700

- 5 The hard fat is melted. At 40°C, the ground active substance is homogeneously dispersed therein. The mixture is cooled to 38°C and poured into slightly chilled suppository molds.

G. METERING AEROSOL	
Component	Amount
active substance	0.005
sorbitan trioleate	0.1
Monofluorotrichloromethane and difluorodichloromethane (2:3)	To 100

The suspension is transferred into a conventional aerosol container with a metering valve.

- 10 Preferably, 50 µL of suspension are delivered per spray. The active substance may also be metered in higher doses if desired (e.g., 0.02% by weight).

H. POWDER FOR INHALATION	
Component	Amount
active substance	1.0 mg
lactose monohydrate	to 25 mg

I. POWDER FOR INHALATION	
Component	Amount
active substance	2.0 mg
lactose monohydrate	to 25 mg

5


J. POWDER FOR INHALATION	
Component	Amount
active substance	1.0 mg
lactose monohydrate	to 5 mg

K. POWDER FOR INHALATION	
Component	Amount
active substance	2.0 mg
lactose monohydrate	to 5 mg

In Examples H, I, J, and K, the powder for inhalation is produced in the usual way by mixing
10 the individual ingredients together.

5 **WHAT IS CLAIMED IS:**

1. A compound of the formula (I):



10

wherein:

R¹ is C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, C₃₋₁₀ cycloalkyl, 3-10 membered saturated heterocyclic ring, benzyl or phenethyl each optionally independently substituted with 1-3 substituents
 15 chosen from C₁₋₁₀ alkyl optionally substituted with 1 to 3 halogen atoms, C₃₋₁₀ cycloalkyl, 3-10 membered saturated heterocyclic ring optionally substituted with acyl, oxo or methyl sulfone, acyl, cyano, phenyl, oxo, hydroxyl and halogen;

R² and **R³** are independently hydrogen or C₁₋₆ alkyl; or **R²** and **R³** together with the
 20 carbon atom to which they are attached form a 3- to 6-membered cycloalkyl or heterocyclic ring;

R⁴ is hydrogen or methyl;

R⁵ is aryl or heteroaryl each optionally independently substituted with 1 to 3 substituents
 25 chosen from C₁₋₆ alkyl optionally substituted with 1 to 3 halogen atoms or with a heterocyclyl group, C₁₋₆ alkoxy optionally substituted with 1 to 3 halogen atoms, C₃₋₆ cycloalkyl, phenoxy, halogen, cyano, dimethylaminoC₁₋₄alkyl, phenyl optionally substituted with 1 to 2 halogen atoms or C₁₋₄ alkyl optionally substituted with halogen, thienyl
 30 optionally substituted with 1 to 2 halogen atoms or C₁₋₄ alkyl optionally substituted with

5 halogen and pyridinyl optionally substituted with 1 to 2 C₁-C₄ alkyl optionally substituted with halogen;

or a tautomer or pharmaceutically acceptable salt thereof.

10

2. The compound according to claim 1 wherein

R¹ is methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, tetrahydropyranyl, bicyclo(3.3.0)octanyl, bicyclo[4.3.0]nonyl, benzyl or phenethyl each optionally independently substituted with 1 to 3 substituents chosen from methyl, ethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, tetrahydrofuranlyl, tetrahydropyranyl, pyrrolidinyl (optionally substituted with acyl, oxo or methylsulfonyl), piperidinyl (optionally substituted with acyl, oxo or methylsulfone, acyl, cyano, phenyl, oxo, fluoro, chloro, bromo and hydroxyl, ;

20

R² and **R³** are independently hydrogen, methyl, ethyl, n-propyl, isopropyl, isobutyl, tert-butyl, or **R²** and **R³** together with the carbon to which they are attached form a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or tetrahydropyranyl ring;

25 **R⁴** is hydrogen;

R⁵ is phenyl, naphthyl, pyridinyl, quinolinyl, isoquinolinyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, indolyl, benzoxazolyl, benzopyrazolyl, benzoisoxazolyl, benzothiazolyl, benzoisothiazolyl, benzimidazolyl, tetrahydrobenzoxazolyl, tetrahydrobenzothiazolyl or tetrahydrobenzimidazolyl each optionally independently substituted with 1 to 3 substituents chosen from C₁-C₆ alkyl optionally substituted with 1 to 3 halogen atoms, C₁-C₆ alkoxy

30

5 optionally substituted with 1 to 3 halogen atoms, C₃-C₆ cycloalkyl, phenoxy, halogen, cyano, dimethylaminoC₁-C₄alkyl, phenyl optionally substituted with 1 to 2 halogen atoms or C₁-C₄ alkyl optionally substituted with halogen, thienyl optionally substituted with 1 to 2 halogen atoms or C₁-C₄ alkyl optionally substituted with halogen and pyridinyl optionally substituted with 1 to 2 C₁-C₄ alkyl optionally substituted with halogen.

10

3. The compound according to claim 2 wherein

R¹ is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, tetrahydropyranyl, , bicyclo(3.3.0)octanyl, or bicyclo[4.3.0]nonyl each optionally independently substituted with 1 to 3 substituents chosen from methyl, ethyl, pivaloyl, cyclopropyl, cyclobutyl, cyclohexyl, tetrahydropyranyl, tetrahydrofuranyl, pyrrolidinyl optionally substituted with methylsulfone, fluoro, chloro, bromo, oxo and hydroxy;

20

R² and **R³** are independently hydrogen, methyl, ethyl, n-propyl, isopropyl, isobutyl, tert-butyl, or **R²** and **R³** together with the carbon to which they are attached form a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or tetrahydropyranyl ring;

25

R⁴ is hydrogen;

R⁵ is phenyl, naphthyl, pyridinyl, quinolinyl, isoquinolinyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, indolyl, benzoxazolyl, benzopyrazolyl, benzoisoxazolyl, benzothiazolyl, benzoisothiazolyl, benzimidazolyl, tetrahydrobenzoxazolyl, tetrahydrobenzothiazolyl, or tetrahydrobenzimidazolyl each optionally independently substituted with 1 to 3 substituents

30

5 chosen from C₁-C₆ alkyl optionally substituted with 1 to 3 halogen atoms, C₁-C₆ alkoxy
optionally substituted with 1 to 3 halogen atoms, C₃-C₆ cycloalkyl, phenoxy, halogen, cyano,
dimethylaminoC₁-C₄alkyl, phenyl optionally substituted with 1 to 2 halogen atoms or C₁-C₄
alkyl optionally substituted with halogen, thienyl optionally substituted with 1 to 2 halogen
atoms or C₁-C₄ alkyl optionally substituted with halogen and pyridinyl optionally substituted
10 with 1 to 2 C₁-C₄ alkyl optionally substituted with halogen.

4. The compound according to claim 3 wherein

15 **R¹** is benzyl or phenethyl each optionally independently substituted with 1 to 3
substituents chosen from methyl, fluoro, chloro and bromo.

5. The compound according to claim 3 wherein

20

R¹ is methyl, ethyl, n-propyl, isopropyl, tert-butyl, n-butyl, cyclopropyl, cyclopentyl,
cyclohexyl, cycloheptyl, tetrahydropyranyl or benzyl each optionally substituted with methyl,
pivaloyl, cyclopropyl, cyclobutyl, cyclohexyl tetrahydropyranyl, tetrahydrofuranyl,
pyrrolidinyl (optionally substituted with methylsulfonyl) or chloro;

25

R² and **R³** are independently hydrogen, methyl, isopropyl, tert-butyl, or **R²** and **R³** together
with the carbon to which they are attached form a cyclopropyl or cyclobutyl, ring;

R⁴ is hydrogen;

30

R⁵ is phenyl, pyridinyl, quinolinyl, isoquinolinyl, pyrazolyl, isoxazolyl, benzothiazolyl,
thiadiazolyl, or thiazolyl each optionally independently substituted with 1 to 2 substituents

- 5 chosen from, methyl, ethyl, tert-butyl, neopentyl, cyclohexyl, trifluoromethyl or phenyl optionally substituted with a chlorine atom.

6. A compound chosen from

- 10 N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (5-tert-butyl-isoxazol-3-yl)-amide
1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (4-tert-butyl-thiazol-2-yl)-amide
- 15 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
- 20 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-N-naphthalen-2-yl-2-(propane-2-sulfonyl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-(4,5,6,7-tetrahydro-benzothiazol-2-yl)-propionamide
2-Methyl-N-(2-methyl-5-thiophen-2-yl-2H-pyrazol-3-yl)-2-(propane-2-sulfonyl)-
- 25 propionamide
N-Isoquinolin-3-yl-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-Benzothiazol-2-yl-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-quinolin-3-yl-propionamide
- 30 N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
2-Methyl-N-(4-phenyl-thiazol-2-yl)-2-(propane-2-sulfonyl)-propionamide
2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(propane-2-sulfonyl)-propionamide

- 5 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclohexanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
- 10 N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-
- 15 propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-
propionamide
2-Methyl-2-(2-methyl-propane-2-sulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(2-methyl-propane-2-sulfonyl)-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide
- 20 2-(1-Acetyl-piperidin-3-ylmethanesulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-
propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
2-(Butane-2-sulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide
2-(Butane-2-sulfonyl)-N-(5-tert-butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-propionamide
- 25 2-(Butane-2-sulfonyl)-N-(4-tert-butyl-thiazol-2-yl)-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
2-(Butane-2-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
- 30 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-ethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methanesulfonyl-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methanesulfonyl-2-methyl-propionamide

- 5 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2,2,2-trifluoro-ethanesulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
- 10 N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-
15 ylmethanesulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-
propionamide
- 20 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(4-trifluoromethyl-thiazol-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
- 25 N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide
- 30 N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide

- 5 N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-
10 propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-(hexane-3-sulfonyl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-
ylmethanesulfonyl)-propionamide
15 N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-ethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyanomethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-
propionamide
20 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-(2,2-dimethyl-propane-1-sulfonyl)-2-methyl-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-
propionamide
25 2-Methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
2-Methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
2-(3,3-Dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-
30 propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(3,3-dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-
propionamide

- 5 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-((S)-5-oxo-pyrrolidin-2-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-((R)-5-oxo-pyrrolidin-2-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
- 10 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(piperidin-4-ylmethanesulfonyl)-propionamide
2-Cyclohexylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
- 15 2-Methyl-2-(3-methyl-butane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(piperidin-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(4-methyl-tetrahydro-pyran-4-sulfonyl)-
- 20 propionamide
2-Cycloheptanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cyclopropylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cyclobutylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
- 25 N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-phenyl-propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-2-sulfonyl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-(4-pyridin-3-yl-thiazol-2-yl)-propionamide
- 30 2-Methyl-2-(propane-2-sulfonyl)-N-(4-pyridin-2-yl-thiazol-2-yl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-(4-pyridin-4-yl-thiazol-2-yl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-quinolin-2-yl-propionamide

- 5 2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-3-yl-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-(4,5,6,7-tetrahydro-benzothiazol-2-yl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-2-yl-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
- 10 2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-N-(2-methyl-5-thiophen-2-yl-2H-pyrazol-3-yl)-2-(propane-1-sulfonyl)-
propionamide
N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
- 15 2-Methyl-2-(propane-1-sulfonyl)-N-(4-pyridin-4-yl-thiazol-2-yl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-(4-trifluoromethyl-thiazol-2-yl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-Benzothiazol-2-yl-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
- 20 N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-
methyl-propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(propane-1-sulfonyl)-propionamide
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-N-(5-phenyl-pyridin-2-yl)-2-(propane-1-sulfonyl)-propionamide
- 25 N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(2,4-Dimethyl-5-phenyl-2H-pyrazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 30 2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-thiazol-2-yl)-
propionamide

- 5 N-(4-Cyclopropyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-Benzothiazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-[1,3,4]thiadiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
10 propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-pyridin-2-yl)-
propionamide
N-(4-Ethyl-pyridin-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-phenyl-[1,2,4]thiadiazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
15 propionamide
N-Benzooxazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
20 propionamide
N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
2-Methyl-N-(3-propyl-isoxazol-5-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(5-Isopropyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-
25 ylmethanesulfonyl)-propionamide
N-(2,4-Dimethyl-5-phenyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-
ylmethanesulfonyl)-propionamide
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(tetrahydro-pyran-4-
ylmethanesulfonyl)-propionamide
30 2-Methyl-N-naphthalen-1-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-methyl-pyridin-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-3-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide

- 5 2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(3-trifluoromethyl-phenyl)-
propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-2-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
10 propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-phenyl)-
propionamide
2-Methyl-N-(2-methyl-5-phenyl-2H-pyrazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
15 N-Isoquinolin-3-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
20 2-Cyclopentanesulfonyl-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
25 propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
2-(Butane-2-sulfonyl)-N-(3-tert-butyl-isoxazol-5-yl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-
30 propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide

- 5 1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-5-yl)-amide
N-(3-tert-Butyl-isoxazol-5-yl)-2-(4-fluoro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide
- 10 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-ylmethanesulfonyl)-propionamide
N-(3-sec-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[4-(4-Fluoro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 15 2-Methyl-N-(4-pyridin-3-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(4-pyridin-2-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 20 2-Methyl-N-(4-pyridin-4-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(3-Isopropyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide and
2-Methyl-N-(3-phenyl-1,2,4-thiadiazol-5-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 25

or the tautomers or the pharmaceutically acceptable salts thereof.

- 30 7. A compound chosen from
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide

- 5 N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (5-tert-butyl-isoxazol-3-yl)-amide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-
10 propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
15 N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclohexanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropanesulfonyl-2-methyl-propionamide
20 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-
propionamide
2-(Butane-2-sulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide
25 2-(Butane-2-sulfonyl)-N-(4-tert-butyl-thiazol-2-yl)-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-3-methyl-2-(propane-2-sulfonyl)-butyramide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
30 propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide

- 5 N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-
10 propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(6-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
15 N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-
20 propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(4-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide
25 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-(hexane-3-sulfonyl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
30 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide

- 5 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
2-Methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-(3,3-Dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-
10 propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(3,3-dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-
15 propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
2-Cyclohexylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(3-methyl-butane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cycloheptanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
20 2-Cyclopropylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Cyclobutylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
25 N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-2-sulfonyl)-propionamide
2-Methyl-2-(propane-2-sulfonyl)-N-quinolin-2-yl-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-3-yl-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Methyl-2-(propane-1-sulfonyl)-N-quinolin-2-yl-propionamide
30 N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide

- 5 N-Benzothiazol-2-yl-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-methyl-propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(propane-1-sulfonyl)-propionamide
N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
- 10 N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-Benzothiazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 15 N-(5-tert-Butyl-[1,3,4]thiadiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-phenyl-[1,2,4]thiadiazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 20 N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 25 2-Methyl-N-(3-propyl-isoxazol-5-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-3-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(3-trifluoromethyl-phenyl)-propionamide
- 30 N-(4-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-2-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide

- 5 N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(4-trifluoromethyl-phenyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
10 propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
15 N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
20 N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
2-(Butane-2-sulfonyl)-N-(3-tert-butyl-isoxazol-5-yl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide
25 1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-5-yl)-amide
N-(3-tert-Butyl-isoxazol-5-yl)-2-(4-fluoro-phenylmethanesulfonyl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide
30 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-ylmethanesulfonyl)-propionamide

- 5 N-(3-sec-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-[4-(4-Fluoro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
2-Methyl-N-(4-pyridin-3-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
10 propionamide
2-Methyl-N-(4-pyridin-2-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide and
2-Methyl-N-(4-pyridin-4-yl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
15 or the tautomers or the pharmaceutically acceptable salts thereof.

8. A compound chosen from

- N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
20 N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-2-sulfonyl)-propionamide
1-Cyclohexanesulfonyl-cyclobutanecarboxylic acid (5-tert-butyl-isoxazol-3-yl)-amide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-
propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(4-chloro-phenylmethanesulfonyl)-2-methyl-propionamide
25 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
2-Cyclohexanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-propionamide
30 N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(2-methyl-propane-2-sulfonyl)-
propionamide
2-(Butane-2-sulfonyl)-N-(5-tert-butyl-isoxazol-3-yl)-2-methyl-propionamide

- 5 2-(Butane-2-sulfonyl)-N-(4-tert-butyl-thiazol-2-yl)-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
10 propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide
2-Methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
15 N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(4-tert-Butyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
2-Methyl-2-(tetrahydro-pyran-4-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-
propionamide
20 N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-cyclopentanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-
propionamide
25 N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-
propionamide
N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
30 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-cyclohexylmethanesulfonyl-2-methyl-
propionamide

- 5 2-(3,3-Dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-
propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(3,3-dimethyl-2-oxo-butane-1-sulfonyl)-2-methyl-
propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
- 10 N-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-
propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cycloheptanesulfonyl-2-methyl-propionamide
2-Cyclohexylmethanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
2-Methyl-2-(3-methyl-butane-1-sulfonyl)-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
- 15 2-Cycloheptanesulfonyl-2-methyl-N-(5-trifluoromethyl-pyridin-2-yl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-2-sulfonyl)-propionamide
N-(4-tert-Butyl-phenyl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
- 20 N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-isoxazol-3-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-
methyl-propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(propane-1-sulfonyl)-propionamide
- 25 N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(4-Cyclohexyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-(5-tert-Butyl-4-methyl-thiazol-2-yl)-2-methyl-2-(propane-1-sulfonyl)-propionamide
N-Benzothiazol-2-yl-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-(5-phenyl-[1,2,4]thiadiazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
- 30 propionamide
2-Methyl-N-(5-methyl-4-phenyl-thiazol-2-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
propionamide

- 5 N-(5-Ethyl-4-phenyl-thiazol-2-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[3-(2,2-Dimethyl-propyl)-isoxazol-5-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-3-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
- 10 N-(4-tert-Butyl-phenyl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
2-Methyl-N-quinolin-2-yl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-[4-(4-Chloro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
- 15 propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
2-Cyclopentanesulfonyl-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclobutylmethanesulfonyl-2-methyl-propionamide
- 20 N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopropylmethanesulfonyl-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(3-methyl-butane-1-sulfonyl)-propionamide
2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclopentanesulfonyl-2-methyl-propionamide
- 25 N-(3-tert-Butyl-isoxazol-5-yl)-2-cyclohexylmethanesulfonyl-2-methyl-propionamide
2-(Butane-2-sulfonyl)-N-(3-tert-butyl-isoxazol-5-yl)-2-methyl-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-2-ylmethanesulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(2-methyl-propane-1-sulfonyl)-propionamide
- 30 1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-5-yl)-amide
N-(3-tert-Butyl-isoxazol-5-yl)-2-(4-fluoro-phenylmethanesulfonyl)-2-methyl-propionamide

- 5 N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-2-ylmethanesulfonyl)-propionamide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-ylmethanesulfonyl)-propionamide
N-(3-sec-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-
10 propionamide and
N-[4-(4-Fluoro-phenyl)-thiazol-2-yl]-2-methyl-2-(tetrahydro-pyran-4-ylmethanesulfonyl)-propionamide
or the tautomers or the pharmaceutically acceptable salts thereof.
- 15 9. A compound chosen from
1-(Tetrahydro-pyran-4-sulfonyl)-cyclobutanecarboxylic acid (3-tert-butyl-isoxazol-5-yl)-amide
N-(3-tert-Butyl-isoxazol-5-yl)-2-methyl-2-(tetrahydro-furan-3-sulfonyl)-propionamide
1-(Tetrahydro-pyran-4-ylmethanesulfonyl)-cyclobutanecarboxylic acid (5-phenyl-4H-1,2,4-
20 triazol-3-yl)-amide
2-Methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-2-(tetrahydro-pyran-4-sulfonyl)-propionamide
1-(Tetrahydro-pyran-4-sulfonyl)-cyclobutanecarboxylic acid (5-phenyl-4H-1,2,4-triazol-3-yl)-amide
N-(3-tert-Butyl-isoxazol-5-yl)-2-(1-methanesulfonyl-pyrrolidin-3-ylmethanesulfonyl)-2-
25 methyl-propionamide and
2-(1-Methanesulfonyl-pyrrolidine-3-sulfonyl)-2-methyl-N-(5-phenyl-4H-1,2,4-triazol-3-yl)-propionamide
or the tautomers or the pharmaceutically acceptable salts thereof.
- 30 10. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to claim 1.

5 11. A method of treating a CB2 receptor-mediated disease or condition in an animal subject comprising administering to said animal subject a therapeutically effective dose of the compound according to claim 1.

12. The method according to claim 11 wherein said CB2 receptor-mediated disease or
10 condition is selected from the group consisting of an inflammatory disease and an autoimmune disease.

13. The method according to claim 11 wherein said CB2 receptor-mediated disease or condition is pain.

15

14. The method according to claim 10 wherein said CB2 receptor-mediated disease or condition is a lung disease, a rheumatic disease, an autoimmune disease, a musculoskeletal disease, an allergic disease, an allergic reaction, a vascular disease, a dermatological disease, a renal disease, a hepatic disease, a gastrointestinal disease, neurodegeneration eye disease,
20 diseases of the ear, nose, and throat, neurological disease blood disease, tumors, endocrine diseases, organ and tissue transplantations and graft-versus-host diseases, severe states of shock, acute pain, visceral pain, spasm of the gastrointestinal tract or uterus, colics, neuropathic pain, inflammatory and nociceptive pain, cancer pain, headache, restenosis, atherosclerosis, reperfusion injury, congestive heart failure, myocardial infarction, thermal
25 injury, multiple organ injury secondary to trauma, necrotizing enterocolitis and syndromes associated with hemodialysis, leukopheresis, and granulocyte transfusion, sarcoidosis, gingivitis, and pyrexia.