




162162



G00002

ABSTRACT

The present invention relates to a compound or a pharmacologically acceptable salt thereof having superior glucokinase activating activity, and is a compound represented by general formula (I), or pharmacologically acceptable salt thereof:



[wherein,

A represents, for example, an oxygen atom or sulfur atom, R^1 represents, for example, a C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group or a C_1 - C_6 halogenated alkyl group, A and R^1 together with the carbon atom bonded thereto form a heterocyclic group that may be substituted with 1 to 3 group(s) independently selected from Substituent Group α , R^2 represents a phenyl group that may be substituted with 1 to 5 group(s) independently selected from Substituent Group α or a heterocyclic group that may be substituted with 1 to 3 group(s) independently selected from Substituent Group α , R^3 represents a hydroxy group or a C_1 - C_6 alkoxy group, and Substituent Group α consists of, for example, a halogen atom, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group substituted with 1 or 2 hydroxy group(s), a C_1 - C_6 alkylsulfonyl group, and a group represented by the formula $-V-NR^5R^6$ (wherein, V represents a carbonyl group or a sulfonyl group, and R^5 and R^6 may be the same or different and respectively represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^5 and R^6 together with the nitrogen atom bonded thereto form a 4- to 6-membered saturated heterocycle that may be substituted with 1 or 2 group(s) independently selected from a C_1 - C_6 alkyl group and a hydroxy group, and the 4- to 6-membered saturated heterocycle may further contain one oxygen atom or nitrogen atom)].