

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
9 November 2006 (09.11.2006)

PCT

(10) International Publication Number
WO 2006/117368 A1

(51) International Patent Classification:

A61K 31/47 (2006.01) C07D 409/12 (2006.01)
C07D 215/20 (2006.01) C07D 417/12 (2006.01)
C07D 401/12 (2006.01) A61K 31/4709 (2006.01)
C07D 405/12 (2006.01) A61P 5/24 (2006.01)

(74) Agent: KRAAK, Hajo; N.V. Organon, Global Patent Department, P.O. Box 20, NL-5340 BH Oss (NL).

(21) International Application Number:

PCT/EP2006/061972

(22) International Filing Date: 2 May 2006 (02.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

05103735.6 4 May 2005 (04.05.2005) EP

(71) Applicant (for all designated States except US): N.V. ORGANON [NL/NL]; Kloosterstraat 6, NL-5349 AB Oss (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GRIMA POVEDA, Pedro, Manuel [ES/NL]; N.V. Organon, P.O. Box 20, NL-5340 BH Oss (NL). KARSTENS, Willem, Frederik, Johan [NL/NL]; N.V. Organon, P.O. Box 20, NL-5340 BH Oss (NL). TIMMERS, Cornelis, Marius [NL/NL]; N.V. Organon, P.O. Box 20, NL-5340 BH Oss (NL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 4-PHENYL-5-OXO-1,4,5,6,7,8-HEXAHYDROQUINOLINE DERIVATIVES AS MEDICAMENTS FOR THE TREATMENT OF INFERTILITY



(57) Abstract: The present invention relates to 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivatives according to Formula I, Formula I or a pharmaceutically acceptable salt thereof, wherein R^1 is (1-6C)alkyl, (2-6C)alkenyl or (2-6C)alkynyl; R^2 , R^3 are independently halogen, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)-alkynyl, (1-4C)alkoxy, (3-4C)alkenyloxy or (3-4C)alkynyoxy; R^4 is phenyl or (2-5C)-heteroaryl, both substituted with R and optionally substituted on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylthio and (di)(1-4C)-alkylamino. The invention also relates to pharmaceutical compositions comprising said derivatives, as well as to the use of these 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivatives in therapy, more specifically for the treatment of infertility.

4-PHENYL-5-OXO-1,4,5,6,7,8-HEXAHYDROQUINOLINE DERIVATIVES AS MEDICAMENTS FOR THE TREATMENT OF INFERTILITY

The present invention relates to 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivatives, to pharmaceutical compositions comprising the same and to the use of said
5 derivatives for the manufacture of medicaments for the treatment of infertility.

Gonadotropins serve important functions in a variety of bodily functions including metabolism, temperature regulation and the reproductive process. Gonadotropins act on specific gonadal cell types to initiate ovarian and testicular differentiation and steroidogenesis. The pituitary gonadotropin FSH (follicle stimulating hormone) for
10 example plays a pivotal role in the stimulation of follicle development and maturation whereas LH (luteinizing hormone) induces ovulation (Sharp, R.M. Clin Endocrinol. 33:787-807, 1990; Dorrington and Armstrong, Recent Prog. Horm. Res. 35:301-342,1979). Currently, FSH is applied clinically for ovarian stimulation i.e. ovarian hyperstimulation for in vitro fertilisation (IVF) and induction of ovulation in infertile
15 anovulatory women (Insler, V., Int. J. Fertility 33:85-97, 1988, Navot and Rosenwaks, J. Vitro Fert. Embryo Transfer 5:3-13, 1988), as well as for male hypogonadism and male infertility.

The gonadotropin FSH is released from the anterior pituitary under the influence of gonadotropin-releasing hormone and estrogens, and from the placenta during
20 pregnancy. In the female, FSH acts on the ovaries promoting development of follicles and is the major hormone regulating secretion of estrogens. In the male, FSH is responsible for the integrity of the seminiferous tubules and acts on Sertoli cells to support gametogenesis. Purified FSH is used clinically to treat infertility in females and for some types of failure of spermatogenesis in males. Gonadotropins destined for
25 therapeutic purposes can be isolated from human urine sources and are of low purity (Morse et al, Amer. J. Reproduct. Immunol. and Microbiology 17:143, 1988). Alternatively, they can be prepared as recombinant gonadotropins. Recombinant human FSH is available commercially and is being used in assisted reproduction (Olijve et al. Mol. Hum. Reprod. 2:371, 1996; Devroey et al. Lancet 339:1170, 1992).

The actions of the FSH hormone are mediated by a specific plasma membrane receptor that is a member of the large family of G-protein coupled receptors. These receptors consist of a single polypeptide with seven transmembrane domains and are able to interact with the Gs protein, leading to the activation of adenylate cyclase.

- 5 The FSH receptor is a highly specific target in the ovarian follicle growth process and is exclusively expressed in the ovary. Blocking of the receptor or inhibiting the signalling which is normally induced after FSH-mediated receptor activation will disturb follicle development and thus ovulation and fertility. Low molecular weight FSH antagonists could form the basis for new contraceptives, while low molecular
- 10 weight FSH agonists can be used for the same clinical purposes as native FSH, i.e. for the treatment of infertility and for ovarian hyperstimulation on behalf of in vitro fertilisation.

Low molecular weight FSH mimetics with agonistic properties were disclosed in the International Application WO 2000/08015 (Applied Research Systems ARS Holding


15 N.V.) and in WO 2002/09706 (Affymax Research Institute).

Certain tetrahydroquinoline derivatives have recently been disclosed in the International Application WO 2003/004028 (AKZO NOBEL N.V.) as FSH modulating substances, either having agonistic or antagonistic properties.

There remains a need for low molecular weight hormone mimetics that selectively

20 activate the FSH receptor.

In one aspect, there is provided 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivatives of general formula I



Formula I

or a pharmaceutically acceptable salt thereof, wherein

R^1 is (1-6C)alkyl, (2-6C)alkenyl or (2-6C)alkynyl;

R^2 , R^3 are independently halogen, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)-alkoxy, (3-4C)alkenyloxy or (3-4C)alkynyloxy;

5 R^4 is phenyl or (2-5C)heteroaryl, both substituted with R^7 and optionally substituted on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylthio and (di)(1-4C)alkylamino;

R^7 is H, (1-4C)alkylthio, (1-4C)alkylsulfonyl, (di)(1-4C)alkylamino, R^8R^9 -amino,

10 $R^{10}R^{11}$ -aminocarbonyl, $R^{12}R^{13}$ -amino(1-4C)alkylcarbonylamino, $R^{14}R^{15}$ -amino-(1-4C)-alkyl, R^{16} -oxy, $R^{17}R^{18}$ -aminocarbonyl(1-4C)alkoxy, R^{19} -oxy(1-4C)alkyl, R^{19} -oxycarbonyl(1-4C)alkyl, $R^{20}R^{21}$ -aminosulfonyl, R^{20} -oxysulfonyl, aminoiminomethyl, (di)(1-4C)alkylaminoiminomethyl or (2-6C)heterocycloalkyliminomethyl, trifluoromethylsulfonyl; R^{23} -oxycarbonyl, R^{23} -carbonyl or $R^{23}R^{24}$ -aminocarbonyl;

15 R^8 is H or (1-4C)alkyl;

R^9 is (1-4C)alkylsulfonyl, (1-6C)alkylcarbonyl, (2-6C)alkenylcarbonyl, (2-6C)-alkynylcarbonyl, (3-6C)cycloalkylcarbonyl, (3-6C)cycloalkyl(1-4C)alkylcarbonyl, (1-4C)alkoxycarbonyl, (3-4C)alkenyloxycarbonyl, (3-4C)alkynyloxycarbonyl, (di)(1-4C)alkylaminocarbonyl, (2-6C)heterocycloalkylcarbonyl, (5-8C)alkyl, (3-6C)-

20 cycloalkyl, (3-6C)cycloalkyl(1-4C)alkyl, (di)(1-4C)alkylamino(2-4C)alkyl, (2-6C)-heterocycloalkyl(2-4C)alkyl or phenylcarbonyl, phenylsulfonyl, phenyl(1-4C)-alkoxy(1-4C)alkylcarbonyl, phenyl(1-4C)alkyl, (2-5C)heteroarylcarbonyl, (2-5C)-heteroarylsulfonyl, (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino,

25 halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (1-4C)alkoxy or (di)(1-4C)-alkylamino;

R^{10} is H or (1-4C)alkyl;

R^{11} is hydroxy(2-4C)alkyl, amino(2-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl or (di)-(1-4C)-alkylamino(2-4C)alkyl; or

30 $R^{10}R^{11}$ in $R^{10}R^{11}$ -aminocarbonyl may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring substituted with one or more substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy(1-4C)alkyl;

- R^{12} , R^{13} are independently H, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (3-6C)-cycloalkyl, hydroxy(2-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl, (3-6C)cycloalkyl-(1-4C)-alkyl, (2-6C)heterocycloalkyl(1-4C)alkyl, amino(2-4C)alkyl, (di)(1-4C)alkylamino-(2-4C)alkyl or phenyl(1-4C)alkyl, (2-5C)heteroaryl(1-4C)alkyl, optionally substituted
- 5 on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (1-4C)alkoxy and (di)(1-4C)alkylamino; or
- $R^{12}R^{13}$ in $R^{12}R^{13}$ -amino(1-4C)alkylcarbonylamino may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or
- 10 more substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy(1-4C)alkyl;
- R^{14} , R^{15} are independently H, (1-6C)alkyl, (2-6C)heterocycloalkyl(1-4C)alkyl, (3-6C)-cycloalkyl(1-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl, hydroxy(2-4C)alkyl, (di)(1-4C)-alkylamino(2-4C)alkyl, amino(2-4C)alkyl, (1-4C)alkoxycarbonyl-(1-4C)alkyl, (1-6C)-
- 15 alkylcarbonyl, (3-6C)cycloalkylcarbonyl, (1-4C)alkoxycarbonyl, (3-4C)alkenyloxy-carbonyl, (di)(1-4C)alkylaminocarbonyl, (2-6C)heterocycloalkylcarbonyl or (2-5C)-heteroaryl(1-4C)alkyl, phenyl(1-4C)alkyl, (2-5C)heteroarylcarbonyl, phenylcarbonyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl or
- 20 (1-4C)alkoxy, (di)(1-4C)alkylamino; or
- $R^{14}R^{15}$ in $R^{14}R^{15}$ -amino(1-4C)alkyl may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy(1-4C)alkyl;
- R^{16} is (2-6C)heterocycloalkyl(1-4C)alkyl, (di)(1-4C)alkylamino(2-4C)alkyl, (2-4C)-
- 25 alkoxy(1-4C)alkyl, hydroxy(2-4C)alkyl, amino(2-4C)alkyl, hydroxycarbonyl(1-4C)-alkyl, (1-4C)alkoxycarbonyl(1-4C)alkyl, (1-4C)alkoxycarbonyl, (3-4C)alkenyloxy-carbonyl, (3-4C)alkynyloxy-carbonyl, (di)(1-4C)alkylaminocarbonyl, (2-6C)hetero-cycloalkylcarbonyl, or phenyl(1-4C)alkyl, (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from
- 30 hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (1-4C)alkoxy and (di)(1-4C)alkylamino;
- R^{17} , R^{18} are independently H, (1-6C)alkyl, (3-6C)cycloalkyl(1-4C)alkyl, (1-4C)-

- alkoxy(2-4C)alkyl, hydroxy(2-4C)alkyl, amino(2-4C)alkyl, (di)(1-4C)alkylamino-(2-4C)alkyl, (2-6C)heterocycloalkyl(2-4C)alkyl, or phenyl(1-4C)alkyl, (2-5C)-heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano,
- 5 (1-4C)alkyl, (1-4C)alkoxy and (di)(1-4C)alkylamino; or
- $R^{17}R^{18}$ in $R^{17}R^{18}$ -aminocarbonyl(1-4C)alkoxy may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy-(1-4C)alkyl;
- 10 R^{19} is H or (1-6C)alkyl;
- R^{20}, R^{21} are independently H, (1-6C)alkyl, (1-6C)alkenyl, (1-6C)alkynyl or (1-4C)-alkoxy(1-4C)alkyl; or
- $R^{20}R^{21}$ in $R^{20}R^{21}$ -aminosulfonyl may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents
- 15 selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy- (1-4C)alkyl;
- X is O or N- R^{22} ;
- Y is CH_2 , C(O) or SO_2 ;
- Z is CN or NO_2 ;
- R^{22} is H, (1-4C)alkyl;
- 20 R^{23}, R^{24} are independently H, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (3-6C)cycloalkyl, (3-6C)cycloalkyl(1-4C)alkyl, (2-6C)heterocycloalkyl, (2-6C)heterocycloalkyl-(1-4C)alkyl, (1-4C)alkoxycarbonyl(1-4C)alkyl, (di)(1-4C)alkylaminocarbonyl(1-4C)-alkyl or phenylaminocarbonyl(1-4C)alkyl, (2-5C)heteroarylaminocarbonyl(1-4C)alkyl, phenyl, (2-5C)heteroaryl, phenyl(1-4C)alkyl, (2-5C)heteroaryl(1-4C)alkyl, optionally
- 25 substituted at the (hetero)atom with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)-alkynyl, (1-4C)alkoxy and (di)(1-4C)alkylamino; or
- $R^{23}R^{24}$ in $R^{23}R^{24}$ -aminocarbonyl may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents
- 30 selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy- (1-4C)alkyl; with the proviso that the compounds of formula I wherein X is O, R^4 is phenyl and R^7 is selected from H, (1-4C)alkylthio, (1-4C)alkylsulfonyl, di(1-4C)alkylamino, R^{23} -

oxycarbonyl, R^{23} -carbonyl and $R^{23}R^{24}$ -aminocarbonyl, and the compounds of formula I wherein X is O, R^4 is (2-5C)heteroaryl and R^7 is H or (di)(1-4C)alkylamino, are excluded.

5 The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivatives disclosed herein are potent FSH receptor activators and can be used for the same clinical purposes as native FSH since they behave like agonists, with the advantage that they may be prepared synthetically, may display altered stability properties and may be administered differently.

10 Thus, the FSH-receptor agonists disclosed herein may be used for the treatment of fertility disorders *e.g.* controlled ovarian hyperstimulation and IVF procedures.

In a second aspect, there is provided a 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative or a pharmaceutically acceptable salt thereof selected from the group of

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydroquinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3,4,5-trimethoxy-benzamide;

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydroquinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[(pyridin-4-ylmethyl)-amino]-acetamide;

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydroquinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-acetamide;

4-{3-Bromo-4-[3-(3,6-dihydro-2*H*-pyridine-1-carbonyl)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;

3-[Bis-(2-methoxy-ethyl)-amino]-*N*-{3-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-propionamide;

2-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-

quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-*N,N*-dimethyl-acetamide;
4-{3-Bromo-5-ethoxy-4-[3-(2-morpholin-4-yl-2-oxo-ethoxy)-benzyloxy]-
phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-
carbonitrile;

5 Furan-2-carboxylic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-
1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide;
N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-
quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-acrylamide;

Cyclopropanecarboxylic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-
10 1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide;
2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-
quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid methyl ester;
1-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-
quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3-methyl-urea

15 {3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-
quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-pyridin-2-yl}-carbamic acid methyl
ester

4-(3-Bromo-5-ethoxy-4-{3-[(1*H*-imidazol-4-ylmethyl)-amino]-benzyloxy}-
phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-
20 carbonitrile;

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-
quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-methanesulfonamide;

{4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-
quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid allyl ester;

25 4-[3-Bromo-5-ethoxy-4-(1-methanesulfonyl-1*H*-pyrrol-2-ylmethoxy)-phenyl]-2-
methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;

4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-
quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-*N*-methyl-benzamidine;

4-{3-Bromo-5-ethoxy-4-[(pyridin-3-ylmethyl)-amino]-phenyl}-2-methyl-5-oxo-
30 7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;

N-(2-{[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-
quinolin-4-yl)-6-ethoxy-phenylamino]-methyl}-phenyl)-methanesulfonamide;

4-{3-Bromo-4-[2-(cyclopropylmethyl-amino)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile; or
N-{2-[2-Bromo-4-((4*R*,7*S*)-3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-4,5-difluoro-phenyl}-
5 methanesulfonamide

In a third aspect, there is provided a 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative as defined above for use in therapy.

In a fourth aspect, there is provided a pharmaceutical composition comprising a 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative as defined above, and
10 pharmaceutically suitable auxiliaries.

In a fifth aspect, there is provided the use of the 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative as defined above, or a pharmaceutically acceptable salt or solvate thereof for the manufacture of a medicament for the treatment of fertility disorders.

15 The term (1-4C)alkyl as used in the definition means a branched or unbranched alkyl group having 1-4 carbon atoms, being methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl.

The term (1-6C)alkyl means a branched or unbranched alkyl group having 1-6 carbon atoms, for example methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, n-pentyl and n-
20 hexyl. (1-5C)Alkyl groups are preferred, (1-4C)alkyl being the most preferred.

The term (5-8C)alkyl as used in the definition means a branched or unbranched alkyl group having 5-8 carbon atoms.

The term (2-6C)alkenyl means a branched or unbranched alkenyl group having 2-6 carbon atoms, such as ethenyl, 2-butenyl, and n-pentenyl.

25 The term (2-4C)alkenyl means a branched or unbranched alkenyl group having 2-4 carbon atoms, such as ethenyl, n-propenyl and 2-butenyl.

The term (2-6C)alkynyl means a branched or unbranched alkynyl group having 2-6 carbon atoms, such as ethynyl, propynyl and n-pentynyl.

The term (2-4C)alkynyl means an alkynyl group having 2-4 carbon atoms, such as ethynyl and propynyl.

The term (3-6C)cycloalkyl means a cycloalkyl group having 3-6 carbon atoms, being cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term (3-6C)cycloalkyl(1-4C)alkyl means a cycloalkylalkyl group, the cycloalkyl group of which has 3-6 carbon atoms with the same meaning as previously defined and the alkyl group having 1-4 carbon atoms with the same meaning as previously defined.

The term (3-6C)cycloalkyl(1-4C)alkylcarbonyl means a cycloalkyl moiety having 3-6 carbon atoms as previously defined, attached to the alkyl moiety, having 1-4 carbon atoms, of an alkylcarbonyl group.

The term (3-6C)cycloalkylcarbonyl means a cycloalkyl group having 3-6 carbon atoms as previously defined, attached to a carbonyl group.

The term (2-6C)heterocycloalkyl means a heterocycloalkyl group having 2-6 carbon atoms, preferably 3-5 carbon atoms, and at least including one heteroatom selected from N, O and/or S, which may be attached via a heteroatom if feasible, or a carbon atom. Preferred heteroatoms are N or O. Most preferred are piperidin-1-yl, morpholin-4-yl, pyrrolidin-1-yl and piperazin-1-yl.

The term (4-6C)heterocycloalkenyl means a heterocycloalkenyl group having 4-6 carbon atoms, preferably 3-5 carbon atoms, and at least including one heteroatom selected from N, O and/or S, which may be attached via a heteroatom if feasible, or a carbon atom. Preferred heteroatoms are N or O.

The term (2-6C)heterocycloalkyl(1-4C)alkyl means a heterocycloalkylalkyl group, the heterocycloalkyl group of which has 2-6 carbon atoms, preferably 3-5 carbon atoms, with the same meaning as previously defined and the alkyl group having 1-4 carbon atoms with the same meaning as previously defined.

The term (2-6C)heterocycloalkyl(2-4C)alkyl means a heterocycloalkylalkyl group, the heterocycloalkyl group of which has 2-6 carbon atoms, preferably 3-5 carbon atoms, with the same meaning as previously defined and the alkyl group having 2-4 carbon atoms.

The term (2-6C)heterocycloalkylcarbonyl means a heterocycloalkylcarbonyl group, the heterocycloalkyl group of which has 2-6 carbon atoms, preferably 3-5 carbon atoms, with the same meaning as defined previously.

The term (2-5C)heteroaryl means a substituted or unsubstituted aromatic group having 2-5 carbon atoms and at least including one heteroatom selected from N, O and S, like

imidazolyl, thiadiazolyl, pyridinyl, thienyl or furyl. Preferred heteroaryl groups are thienyl, furyl and pyridinyl. The (2-5C)heteroaryl group may be attached via a carbon atom or a heteroatom, if feasible.

5 The term (2-5C)heteroaryl(1-4C)alkyl means a heteroarylalkyl group, the heteroaryl group of which contains 2-5 carbon atoms with the same meaning and preferences as previously defined and the alkyl group contains 1-4 carbon atoms with the same meaning as previously defined.

10 The term (2-5C)heteroarylamincarbonyl(1-4C)alkyl means a heteroarylamincarbonyl group, the heteroaryl group of which contains 2-5 carbon atoms with the same meaning and preferences as previously defined attached to an alkyl group containing 1-4 carbon atoms with the same meaning as previously defined.

The term (2-5C)heteroarylcarbonyl means a heteroarylcarbonyl group, the heteroaryl group of which contains 2-5 carbon atoms with the same meaning and preferences as previously defined.

15 The term (2-5C)heteroarylsulfonyl means a heteroarylsulfonyl group, the heteroaryl group of which contains 2-5 carbon atoms with the same meaning and preferences as previously defined.

The term (1-6C)alkylcarbonyl means an alkylcarbonyl group, the alkyl group of which contains 1-6 carbon atoms with the same meaning as previously defined.

20 The term (2-6C)alkenylcarbonyl means an alkenylcarbonyl group, the alkenyl group of which contains 2-6 carbon atoms with the same meaning as previously defined.

The term (2-6C)alkynylcarbonyl means an alkynylcarbonyl group, the alkyl group of which contains 2-6 carbon atoms with the same meaning as previously defined.

25 The term (1-4C)alkylsulfonyl means an alkylsulfonyl group, the alkyl group of which contains 1-4 carbon atoms with the same meaning as previously defined.

The term (1-4C)alkylthio means an alkylthio group having 1-4 carbon atoms, the alkyl moiety having the same meaning as previously defined.

The term (1-4C)alkoxy means an alkoxy group having 1-4 carbon atoms, the alkyl moiety having the same meaning as previously defined. (1-2C)Alkoxy groups are preferred.

5 The term (3-4C)alkenyloxy means an alkenyloxy group having 3-4 carbon atoms, the alkenyl moiety having the same meaning as previously defined.

The term (3-4C)alkynyloxy means an alkynyloxy group having 3-4 carbon atoms, the alkynyl moiety having the same meaning as previously defined.

The term (1-4C)alkoxycarbonyl means an alkoxycarbonyl group, the alkoxy group of which contains 1-4 carbon atoms with the same meaning as previously defined.
10 (1-2C)Alkoxycarbonyl groups are preferred.

The term (3-4C)alkenyloxycarbonyl means an alkenyloxycarbonyl group, the alkenyl group of which contains 3-4 carbon atoms.

The term (3-4C)alkynyloxycarbonyl means an alkynyloxycarbonyl group, the alkynyl group of which contains 3-4 carbon atoms.

15 The term (1-4C)alkoxy(1-4C)alkyl means an alkoxy group, the alkyl group of which contains 1-4 carbon atoms which is attached to an alkyl group having 1-4 carbon atoms.

The term (1-4C)alkoxy(2-4C)alkyl means an alkoxy group, the alkyl group of which contains 1-4 carbon atoms which is attached to an alkyl group having 2-4 carbon
20 atoms.

The term (1-4C)alkoxycarbonyl(1-4C)alkyl means an alkoxycarbonylalkyl group, the alkyl groups of which contain 1-4 carbon atoms with the same meaning as previously defined.

The term phenyl(1-4C)alkyl means a phenyl group attached to an alkyl group having
25 1-4 carbon atoms as defined previously.

The term phenyl(1-4C)alkoxy(1-4C)alkylcarbonyl means a phenylalkoxy moiety, the alkyl group of which contains 1-4 carbon atoms attached to the alkyl group of a alkylcarbonyl moiety, the alkyl group also containing 1-4 carbon atoms.

The term phenylaminocarbonyl(1-4C)alkyl means a phenylaminocarbonyl group attached to an alkyl group having 1-4 carbon atoms as defined previously.

The term hydroxy(2-4C)alkyl as used herein means an hydroxyalkyl group having 2-4 carbon atoms, the alkyl moiety having the same meaning as previously defined.

- 5 The term hydroxycarbonyl(1-4C)alkyl means a hydroxycarbonylalkyl group, the alkyl group of which contains 1-4 carbon atoms with the same meaning as previously defined.

The term (di)(1-4C)alkylamino as used herein means an amino group, monosubstituted or disubstituted with alkyl groups, each of which contains 1-4 carbon atoms and has the
10 same meaning as previously defined.

The term (di)(1-4C)alkylaminocarbonyl means a (di)alkylaminocarbonyl group, the (di)alkylamino group of which is as defined previously.

The term (di)(1-4C)alkylaminocarbonyl(1-4C)alkyl means a (di)alkylaminocarbonyl-alkyl group, the alkyl groups of which contain 1-4 carbon atoms with the same
15 meaning as previously defined.

The term aminocarbonyl(1-4C)alkoxy means an aminocarbonylalkoxy group, the alkyl group of which contains 1-4 carbon atoms with the same meaning as previously defined.

The term (di)(1-4C)alkylamino(2-4C)alkyl as used herein means an (di)alkylamino
20 group having 1-4 carbon atoms, connected via the amino group to an alkyl group having 2-4 carbon atoms, the alkyl moieties having the same meaning as previously defined.

The term amino(1-4C)alkylcarbonylamino means an aminoalkylcarbonylamino group, the alkyl group of which contains 1-4 carbon atoms with the same meaning as
25 previously defined.

The term amino(2-4C)alkyl as used herein means an aminoalkyl group having 2-4 carbon atoms, the alkyl moiety having the same meaning as previously defined.

The term (di)(1-4C)alkylaminoiminomethyl as used herein means an alkylaminoiminomethyl group, the amino group of which is monosubstituted or

disubstituted with alkyl groups having 1-4 carbon atoms and having the same meaning as previously defined.

The term (2-6C)heterocycloalkyliminomethyl as used herein means a heterocycloalkyliminomethyl group, the heterocycloalkyl moiety of which is as
5 previously defined.

The term halogen means fluorine, chlorine, bromine or iodine; chlorine, bromine or iodine being preferred.

The term pharmaceutically acceptable salt represents those salts which are, within the scope of medical judgement, suitable for use in contact for the tissues of humans and
10 lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. They may be obtained during the final isolation and purification of the compounds of the invention, or separately by reacting the free base function with a suitable mineral acid such as hydrochloric acid, phosphoric acid, or
15 sulfuric acid, or with an organic acid such as for example ascorbic acid, citric acid, tartaric acid, lactic acid, maleic acid, malonic acid, fumaric acid, glycolic acid, succinic acid, propionic acid, acetic acid, methanesulfonic acid, and the like. The acid function can be reacted with an organic or a mineral base, like sodium hydroxide, potassium hydroxide or lithium hydroxide.

20 The invention also relates to compounds of formula I, wherein R^1 is (1-6C)alkyl. More in particular, the invention relates to compounds wherein R^1 is (1-4C)alkyl.

Another aspect of the invention are compounds according to formula I wherein R^2 , R^3 is halogen and/or (1-4C)alkoxy.

In yet another aspect, the invention concerns compounds of formula I, wherein Y is
25 CH_2 .

Another aspect of the invention is a compound wherein Z is CN.

In another aspect the invention concerns compounds wherein $X = O$.

The invention also relates to compounds according to general Formula I wherein R^7 is R^8R^9 -amino, $R^{10}R^{11}$ -aminocarbonyl, $R^{12}R^{13}$ -amino(1-4C)alkylcarbonylamino, $R^{14}R^{15}$ --
30 amino(1-4C)alkyl or $R^{17}R^{18}$ -aminocarbonyl(1-4C)alkoxy.

In another aspect the invention concerns compounds wherein

R⁸ is H,

R⁹ is (1-4C)alkylsulfonyl, (1-6C)alkylcarbonyl, (3-6C)cycloalkylcarbonyl, (1-4C)-alkoxycarbonyl, (3-4C)alkenyloxycarbonyl, (di)(1-4C)alkylaminocarbonyl, (2-6C)-
5 heterocycloalkylcarbonyl, or phenylcarbonyl, phenyl(1-4C)alkoxy(1-4C)alkylcarbonyl, (2-5C)heteroarylcarbonyl, (2-5C)heteroarylsulfonyl or (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from, halogen or (1-4C)alkoxy;

R¹⁰ is H or (1-4C)alkyl;

10 R¹¹ is hydroxy(2-4C)alkyl, amino(2-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl or (di)(1-4C)-alkylamino(2-4C)alkyl; or

R¹⁰ and R¹¹ form together with the N to which they are bonded a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl;

15 R¹², R¹³ are independently H, (1-6C)alkyl, (3-6C)cycloalkyl, hydroxy(2-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl, (2-6C)heterocycloalkyl(1-4C)alkyl or (di)(1-4C)-alkylamino(2-4C)alkyl or phenyl(1-4C)alkyl, (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from halogen; or

20 R¹² and R¹³ form together with the N to which they are bonded a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, both optionally substituted with one or more substituents selected from (1-4C)alkyl or hydroxy(1-4C)alkyl;

R¹⁴, R¹⁵ are independently H, (1-6C)alkyl, hydroxy(2-4C)alkyl, (1-4C)alkoxycarbonyl-
(1-4C)alkyl, (1-6C)alkylcarbonyl, (3-6C)cycloalkylcarbonyl, (1-4C)alkoxycarbonyl or
25 (2-5C)heteroaryl(1-4C)alkyl or (2-5C)heteroarylcarbonyl, phenylcarbonyl;

R¹⁶ is (2-6C)heterocycloalkyl(1-4C)alkyl, (di)(1-4C)alkylamino(2-4C)alkyl, (2-4C)-alkoxy(1-4C)alkyl, hydroxycarbonyl(1-4C)alkyl, (1-4C)alkoxycarbonyl(1-4C)alkyl; or
phenyl(1-4C)alkyl or (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the
(hetero)aromatic ring with one or more substituents selected from halogen or (1-4C)-
30 alkoxy;

R¹⁷, R¹⁸ are independently H, (1-6C)alkyl, (3-6C)cycloalkyl(1-4C)alkyl, (1-4C)alkoxy-
(2-4C)alkyl, (2-6C)heterocycloalkyl(2-4C)alkyl; or phenyl(1-4C)alkyl or (2-5C)hetero-

aryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from halogen; or

- R^{17} and R^{18} form together with the N to which they are bonded a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl or hydroxy(1-4C)alkyl.

The invention also relates to compounds according to Formula I wherein R^7 is R^8R^9 -amino.

Another aspect of the invention concerns compounds wherein R^8 is H and R^9 is (1-4C)alkylsulfonyl.

- Still another aspect of the invention concerns compounds wherein one or more of the specific definitions of the groups R^1 through R^{24} and X, Y and Z as defined here above are combined in the definition of the 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivatives of formula I.

- The invention also concerns compounds selected from the group of *N*-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3,4,5-trimethoxy-benzamide;

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[(pyridin-4-ylmethyl)-amino]-acetamide;

- N*-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-acetamide;

4-{3-Bromo-4-[3-(3,6-dihydro-2*H*-pyridine-1-carbonyl)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;

- 3-[Bis-(2-methoxy-ethyl)-amino]-*N*-{3-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-propionamide;


2-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-*N,N*-dimethyl-acetamide;

- 4-{3-Bromo-5-ethoxy-4-[3-(2-morpholin-4-yl-2-oxo-ethoxy)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;


Furan-2-carboxylic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide;

- N*-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-acrylamide;
- Cyclopropanecarboxylic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide;
- 5 2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid methyl ester;
- 1-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3-methyl-urea
- {3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-pyridin-2-yl}-carbamic acid methyl ester
- 10 4-(3-Bromo-5-ethoxy-4-{3-[(1*H*-imidazol-4-ylmethyl)-amino]-benzyloxy}-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;
- N*-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-methanesulfonamide;
- 15 {4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid allyl ester;
- 4-[3-Bromo-5-ethoxy-4-(1-methanesulfonyl-1*H*-pyrrol-2-ylmethoxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;
- 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-*N*-methyl-benzamidine;
- 20 4-{3-Bromo-5-ethoxy-4-[(pyridin-3-ylmethyl)-amino]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;
- N*-(2-{[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenylamino]-methyl}-phenyl)-methanesulfonamide;
- 25 4-{3-Bromo-4-[2-(cyclopropylmethyl-amino)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile; or
- N*-{2-[2-Bromo-4-((4*R*,7*S*)-3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-4,5-difluoro-phenyl}-methanesulfonamide

Suitable methods for the preparation of the 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivatives of the invention are outlined below.



5 The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivatives I of the present




invention can be prepared starting from appropriately functionalised 1,4-dihydropyridine derivatives of general structure II, wherein R¹, R², R³, X, Y and Z are as previously defined and A is a phenyl or a heteroaryl ring.

- 10 For example, acylation or sulfonylation of compounds of general formula II-b yields compounds of general formula I-a, wherein R¹, R², R³, X, Y and Z are as previously defined, R⁸ is H, R⁹ is an acyl or sulfonyl group and A is a (substituted) phenyl or a (substituted) heteroaryl ring.

- 16 -


In a typical experiment, compounds II-b are reacted in a solvent, such as dichloromethane, *N,N*-dimethylformamide, dimethyl sulfoxide, ethanol, tetrahydrofuran, 1,4-dioxane, toluene, 1-methyl-pyrrolidin-2-one or pyridine with an appropriately substituted acyl halide, acid anhydride or sulfonyl halide in the presence
5 of a base such as triethylamine, *N,N*-diisopropylethylamine (DiPEA) or pyridine, to



give *N*-acylated or *N*-sulfonylated derivatives of formula I-a, respectively. Alternatively, *N*-acylated compounds of general formula I-a can be obtained by reaction of a (hetero)aromatic carboxylic acid in the presence of a coupling reagent such as diisopropyl carbodiimide (DIC), (3-dimethylaminopropyl)-ethyl-carbodiimide
10 (EDCI), *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (TBTU) or *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU) and a tertiary amine base (*e.g.* DiPEA) in a solvent such as *N,N*-dimethylformamide or dichloromethane at ambient or elevated temperature.

Compounds of general formula I-b, wherein R¹, R², R³, X, Y and Z are as previously
15 defined, R⁹ is a (substituted) alkyl group and A is a (substituted) phenyl or a (substituted) heteroaryl ring, can be prepared by reductive alkylation of the amino group with appropriately substituted aldehydes of general formula E-C(O)H (E = alkyl,

cycloalkyl, cycloalkylalkyl, (di)alkylaminoalkyl, heterocycloalkylalkyl, (hetero)aryl, (hetero)arylalkyl) and a suitable reducing agent such as sodium cyanoborohydride or sodium triacetoxyborohydride. Likewise, appropriately substituted ketones can be used in this procedure. Alternatively, compounds of general formula II-b can be converted to the corresponding (hetero)arylimine II-c upon reaction with appropriately substituted aldehydes of general formula E-C(O)H (see above) by methods known to those skilled in the art, followed by reduction with a reducing agent such as sodium borohydride to give compounds of general formula I-b. In this case, appropriately substituted ketones can be used as well.




10

Alternatively, compounds of general formula II-b can be converted to 2,4-dinitrobenzenesulfonamide derivatives II-d by sulfonylation with 2,4-dinitrobenzenesulfonyl chloride. The sulfonamide can be alkylated to give compounds of general formula II-e by using art known Mitsunobu reactions with appropriately substituted primary or secondary alcohols of formula R⁹-OH (R⁹ = alkyl, cycloalkyl, cycloalkylalkyl, (di)alkylaminoalkyl, heterocycloalkylalkyl or (hetero)arylalkyl), triphenylphosphine (optionally resin bound) and a dialkyl azodicarboxylate in appropriate solvents such as 1,4-dioxane, tetrahydrofuran or dichloromethane at elevated or ambient temperature. Alternatively, the sulfonamide can be alkylated using alkyl halides of formula R⁹-Hal (Hal = Cl, Br, I) and a suitable base such as K₂CO₃ in a

20


solvent such as *N,N*-dimethylformamide, tetrahydrofuran or 1,4-dioxane. Cleavage of the N-S sulfonamide bond with a primary amine such as propylamine in a suitable solvent such as dichloromethane then gives compounds of formula I-b. Alternatively, the N-S sulfonamide bond can be cleaved using mercaptoacetic acid and a tertiary amine base in a solvent such as dichloromethane. Precedence for these types of reactions can be found in literature. For example, see: Tetrahedron Lett. 38 (1997) 5831-5834, Bioorg. Med. Chem. Lett. 10 (2000) 835-838.

Compounds of general formula I-c, wherein R^1 , R^2 , R^3 , R^9 , X, Y and Z are as previously defined, R^8 is (1-4C)alkyl and A is a (substituted) (hetero)aryl group, can be prepared from compounds I-a, I-b or II-f. Alkylation of compounds I-a with R^8 -Hal (Hal = Cl, Br, I) or R^8 -OH by the same methods described for the alkylations of compounds II-d yields the desired *N,N*-disubstituted aniline derivatives I-c. Additionally, alkylation of compounds of formula I-b by the same methods described for the reductive alkylation of compounds of formula II-b also yields *N,N*-disubstituted aniline derivatives of formula I-c.




Compounds of general formula II-f, prepared by the same methods described for the synthesis of compounds of formula I-b, can be acylated, sulfonylated or reductively alkylated as described for the preparation of I-a and I-b, respectively, to give compounds of general formula I-c.

The nitro group in compounds of general formula II-a can be reduced to the corresponding amino group to give compounds of general formula II-b. Typically, compounds II-a are treated with zinc and acetic acid in a suitable solvent such as THF or dioxane at temperatures between 0 °C and reflux temperature. Alternative methods include treatment with iron, SnCl₂ or hydrogen in the presence of a transition metal catalyst such palladium or platinum on charcoal, using methods and reagents well known to those skilled in the art. Alternatively, compounds of general formula II-b can be obtained by cleavage of known *N*-protecting groups such as an Allyloxycarbonyl (Alloc), Fluoren-9-yl-methoxycarbonyl (Fmoc) or *tert*-butoxycarbonyl (Boc) group in compounds of general formula II-g to give the corresponding derivatives II-b. Related protective group manipulations can be found in *Protective groups in Organic Synthesis*, T.W. Greene and P.G.M. Wuts, John Wiley & sons, Inc., New York, 1999 .




Carboxylic acid derivatives of general formula II-i, accessible by saponification of corresponding alkyl esters II-h, can be condensed with amines of general structure R¹⁰R¹¹NH using a coupling reagent as described before for the preparation of derivatives I-a to give compounds of formula I-d, wherein R¹, R², R³, R¹⁰, R¹¹, X, Y and Z are as previously defined and A is a (substituted) phenyl or a (substituted) heteroaryl ring. A similar coupling method has been described before for the preparation of derivatives I-a. Alternatively, compounds of general formula II-i can be converted to the corresponding acid chlorides II-j by art known methods. Treatment of carboxylic acids of general formula II-i with thionyl chloride or oxalyl chloride and DMF in a suitable solvent such as dichloromethane or toluene gives the corresponding



acid chlorides II-j. Subsequent reaction with amines of general structure $R^{10}R^{11}NH$, optionally in the presence of a suitable tertiary amine base yields compounds of general formula I-d.


- Compounds of general formula I-e, wherein R^1 , R^2 , R^3 , R^{12} , R^{13} , X,Y and Z are as previously defined and A is a (substituted) phenyl or a (substituted) heteroaryl ring, can be obtained by acylation of compounds of general formula II-b with appropriate acyl halides to give compounds II-k, followed by standard substitution of the chloride or bromide with amines of general formula $R^{12}R^{13}NH$. Additionally, compounds of general formula I-e with a two-carbon spacer between the carbonyl and the $NR^{12}R^{13}$ group can be obtained by acylation of compounds of general formula II-b with α,β -unsaturated acyl chlorides to give compounds of general formula II-l, followed by art known Michael addition reactions with appropriately substituted amines of general formula $R^{12}R^{13}NH$.



- Compounds of general formula I-f, wherein R^1 , R^2 , R^3 , R^{14} , R^{15} , X,Y and Z are as previously defined and A is a (substituted) phenyl or a (substituted) heteroaryl ring, can be prepared by alkylation of amines of general formula $R^{14}R^{15}NH$ with halides of general formula II-m, under the agency of a tertiary base such as *N,N*-diisopropyl ethylamine or triethylamine in an appropriate solvent such as *N,N*-dimethylformamide, tetrahydrofuran or dichloromethane.


Derivatives of general formula I-g, wherein R^1 , R^2 , R^3 , R^{16} , X,Y and Z are as previously defined and A is a (substituted) phenyl or a (substituted) heteroaryl ring, can

be prepared by alkylation of the hydroxyl group in compounds of general formula II-o with an alkyl halide of general formula R^{16} -Hal, in which Hal may be Br, Cl or I. Typically, such a reaction is carried out in an aprotic solvent such as *N,N*-dimethylformamide, 1,4-dioxane or tetrahydrofuran in the presence of a base, such as sodium hydride, potassium carbonate, cesium carbonate or triethyl amine at ambient or elevated temperature. Alternatively, conversion of compounds of general formula II-o into aryl ethers of general formula I-g may be accomplished under Mitsunobu-type alkylation conditions. In such a transformation, alkylation of the hydroxyl group in compounds of formula II-o is effected with alcohols of general formula R^{16} -OH under the agency of (resin bound) triphenyl phosphine and diethyl azodicarboxylate or its derivatives in a suitable aprotic solvent such as tetrahydrofuran or dichloromethane.



Derivatives of general formula II-o can be obtained by cleavage of the hydroxyl-protecting group in compounds of general formula II-n. Suitable protective groups, well-known to those skilled in the art, are the tetrahydropyranyl (THP) or *tert*-butyl dimethylsilyl (TBS) protective groups. Cleavage of the THP and TBS groups is generally accomplished by treatment with acids, such as hydrochloric acid, trifluoromethanesulfonic acid or trifluoroacetic acid in a suitable solvent, such as tetrahydrofuran or methanol. Alternatively, the TBS group can be removed by treatment with tetra-*n*-butylammonium fluoride in tetrahydrofuran. Related protective group manipulations can be found in *Protective groups in Organic Synthesis*, T.W. Greene and P.G.M. Wuts, John Wiley & sons, Inc., New York, 1999.


Compounds of the present invention with general formula I-h can be prepared by condensation of derivatives II-q with amines of general formula $R^{17}R^{18}NH$ under the agency of a coupling agent and a tertiary amine base, as described e.g. for the



preparation of derivatives of general formula I-d. The requisite derivatives of formula

- 5 I-q are accessible in a two-step synthetic protocol from compounds of general formula II-o, i.e. alkylation of the phenolic moiety with an appropriate halogenated alkyl ester to obtain derivatives of formula II-p, followed by acid or base-mediated cleavage of the ester function, well known to those skilled in the art.

- 10 Derivatives of general formula I-i are accessible from nitriles of general formula II-r by the art known Pinner amidine synthesis. This synthetic procedure comprises the selective conversion of the nitrile function on the (hetero)aryl ring of derivatives II-r



into imidates II-s by treatment with an appropriate acid such as hydrochloric acid, using a suitable alcohol as a solvent, optionally in the presence of cosolvents such as


1,4-dioxane or diethyl ether, at ambient or elevated temperature. Reaction with the appropriate acyclic or cyclic amine yields the amidine derivatives I-i. Related amidine formations are well known to those skilled in the art. For example, see: J. Med. Chem. 42 (1999) 5415-5425, Bioorg. Med. Chem. Lett. 13 (2003) 697-700, J. Org. Chem. 62
5 (1997), 8449-8454.

Compounds of general formula III-a wherein X is O can be used to prepare compounds I-j, wherein R¹, R², R³, R⁴, Y and Z are as previously defined and X is O, by *O*-alkylation, *O*-acylation or *O*-sulfonylation using standard conditions, well known to those skilled in the art. The substitution pattern of the (hetero)aryl ring in R⁴ is as
10 previously defined. In a typical experiment, compounds III-a are reacted in a solvent, such as dichloromethane, *N,N*-dimethylformamide, dimethyl sulfoxide, ethanol, tetrahydrofuran, 1,4-dioxane, toluene, 1-methyl-pyrrolidin-2-one or pyridine with an appropriately substituted (hetero)aromatic alkyl halide of formula IV, acyl chloride of formula V, or sulfonyl chloride of formula VII in the presence of a base such as
15 triethylamine, *N,N*-diisopropylethylamine (DiPEA), potassium carbonate, cesium carbonate or sodium hydride, optionally in the presence of a catalytic amount of potassium iodide or tetrabutylammonium iodide, to give *O*-alkylated, *O*-acylated or *O*-sulfonylated derivatives of formula I-j, respectively.

Alternatively, *O*-alkylated compounds of general formula I-j in which Y is CH₂ can be
20 obtained by using art known Mitsunobu reactions with alcohols of formula VIII, triphenylphosphine (optionally resin bound) and a dialkyl azodicarboxylate (e.g. diethyl azodicarboxylate) in appropriate solvents such as 1,4-dioxane, tetrahydrofuran or dichloromethane at elevated or ambient temperature.


Additionally, *O*-acylated compounds of general formula I-j, wherein Y is C(O) can be
25 obtained by reaction of a (hetero)aromatic carboxylic acid of formula VI in the presence of a coupling reagent such as diisopropyl carbodiimide (DIC), (3-dimethylaminopropyl)-ethyl-carbodiimide (EDCI), *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (TBTU) or *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU) and a tertiary amine base
30 (e.g. DiPEA) in a solvent such as *N,N*-dimethylformamide or dichloromethane at ambient or elevated temperature.

Likewise, compounds of general formula I-k can be prepared from compounds III-b by *N*-alkylation, *N*-acylation or *N*-sulfonylation using the same methods described for the



synthesis of compounds I-j using the reagents of formula IV-X. Additionally, compounds of general formula I-k in which Y is CH₂ can be prepared by reductive amination of (hetero)aromatic aldehydes of formula IX with compounds III-b and a suitable reducing agent such as sodium cyanoborohydride or sodium triacetoxyborohydride.

Alternatively, compounds of general formula III-b can be converted to the corresponding benzimine upon reaction with (hetero)aromatic aldehydes IX by methods known to those skilled in the art, followed by reduction with a reducing agent




such as sodium borohydride to give compounds I-k in which Y = CH₂.


Compounds of general formula I-k wherein R¹, R², R³, R⁴ and Z are as previously defined and X is NH can be *N*-alkylated by the same methods as described for the preparation of derivatives I-b to afford compounds of general formula I-l, wherein R²²

5 is a (1-4C)alkyl group.

Compounds of general formula II are also accessible from derivatives of general formula III-a and III-b using the same methods as described for the preparation of compounds of general formula I-j and I-k using reagents of formula X-XV.



10 Similar to the *N*-alkylations of compounds of general formula I-k to give I-l, compounds of general formula II-t, wherein R¹, R², R³ and Z are as previously defined and A is a (substituted) phenyl or a (substituted) heteroaryl can be *N*-alkylated to give compounds of general formula II-u, wherein R²² is a (1-4C)alkyl group.




Compounds of general formula I, wherein R¹, R², R³, R⁴, X, Y and Z are as previously defined, can also be prepared starting from cyclohexane-1,3-diones of general formula XVI, enamines of general formula XVII and benzaldehydes of general formula XVIII-

15

a-b, by the well-documented three component Hantzsch-type cyclo-condensation reaction.


Related Hantzsch-type cyclo-condensation reactions can be found in: Bioorg. Med. Chem. Lett. 12 (2002) 1481-1484, J. Chem. Soc., Perkin Trans. 1 (2002) 1141-1156, Synlett (2002) 89-92, Drug Dev. Res. 51 (2000) 225-232, Drug Dev. Res. 51 (2000) 233-243, J. Med. Chem. 42 (1999) 1422-1427, *ibid.* 5266-5271, *ibid.* 41 (1998) 2643-2650, WO 9408966, Arzneim.-Forsch./Drug Res. 45 (1995) 1054-1056, J. Med. Chem. 34 (1991) 2248-2260, *ibid.* 17 (1974) 956-65, Indian J. Chem., Sect B (1994) 526-531, Chem. Rev. 72 (1972), 1-42. The above mentioned reaction is typically conducted at elevated temperature in suitable solvents such as acetic acid, (iso)propanol, ethanol, methanol or mixtures thereof.



Compounds of general formula II and III-a-b are also prepared by the previously mentioned Hantzsch-type cyclo-condensation, by using substituted benzaldehydes of general formula XIX or XX, respectively.

Compounds of general formula III-c-d in which R² is Br can also be obtained by *ortho*-bromination of phenols or anilines, which are well known to those skilled in the art. Thus, compounds of formula III-e-f afford compounds of formula III-c-d upon treatment with bromine in a suitable solvent such as acetic acid, ethanol or dichloromethane or mixtures thereof, optionally in the presence of sodium acetate. Alternatively, *N*-bromosuccinimide in *N,N*-dimethylformamide or acetonitrile can be


used to achieve this conversion. For example, see: J. Chem. Soc. Perkin Trans.2 6 (2000) 1113-1118, J. Org. Chem. 44 (1979), 4733-4735.



Compounds of general formula III-e-f are prepared by the aforementioned Hantzsch reaction using benzaldehydes of general formula XXI-a-b.


- 5 Compounds of general formula IV to XV are either commercially available, documented in literature or readily synthesized by those skilled in the art.

The substituted cyclohexane-1,3-diones of general formula XVI are commercially available or can be prepared by literature procedures. Relevant examples are found in: J. Med. Chem. 43 (2000) 4678-4693, Tetrahedron 56 (2000) 4753-4758, J. Med. Chem.
10 35 (1992) 3429-3447, *ibid.* 24 (1981) 1026-1034, Org. Synt. Coll. Vol. V (1973) 400, Chem. Ber. 88 (1955) 316-327, Justus Liebig Ann. Chem. 570 (1950) 15-31.




The compound of formula XVII-a is commercially available and compound XVII-b can be prepared by art-known methods, see for example: Drug Dev. Res. 51 (2000) 225-232.

- 15 Benzaldehydes of general formula XVIII-a, wherein R², R³, R⁴ and Y are as previously defined and X = O, are readily prepared from benzaldehydes of general formula XX-a using the same methods as described for the conversion of compounds of formula III-a to I-j. Likewise compounds of general formula XVIII-b are prepared from XX-b using the same methods as described for the conversion of compounds of formula III-b to I-k.



Benzaldehydes of general formula XX-a and XX-b are commercially available or can be prepared according to literature procedures: J. Chem. Soc., Perkin Trans. 2 (2000) 1119-1124, J. Chem. Soc., Chem. Commun. 4 (1993) 419-420, Synth. Commun. 20
 5 (1990) 2659-2666, Chem. Pharm. Bull. 34 (1986) 121-129, Indian J. Chem. Sect. B 20 (1981) 1010-1013, Monatsh. Chem. 106 (1975) 1191-1201, DE 1070162, J. Org. Chem. 23 (1958) 120, Tetrahedron Lett. 25 (1984), 2901-2904, J. Org. Chem. 25 (1960), 2053-2055, J. Chem. Soc., Perkin Trans. 2 (1992), 2235-2242. Additionally, benzaldehydes of general formula XX-c wherein R² is bromide and X is N-H can be
 10 obtained by bromination of compounds of general formula XXI using the same procedures described for the conversion of compounds of general formula III-f to III-d.




Compounds of general formula XXI are easily prepared from compounds of general formula XXII using the same reduction methods that were described for the preparation of compounds of general formula II-b from II-a. Compounds of general formula XXII

are commercially available, reported in literature or can be readily be prepared by those skilled in the art.

Similarly, derivatives of general formula XIX-a-b are accessible from compounds XX-a-b upon reaction with compounds X to XV, as previously described for the preparation of derivatives XVIII-a-b.


The compounds of the present invention possess at least two chiral carbon atoms and may therefore be obtained as pure enantiomers, or as a mixture of enantiomers, or as a mixture of diastereomers. Methods for obtaining the pure enantiomers are well known in the art, e.g. crystallization of salts which are obtained from optically active acids and the racemic mixture, or chromatography using chiral columns. For separation of diastereomers, straight phase or reversed phase columns may be used.



Additionally, enantiomerically pure hexahydroquinoline derivatives of general formula XXIII may be obtained in enantiomerically pure form by dehydration of enantiomerically pure amides XXIV with trifluoroacetic anhydride and a suitable base such as triethylamine or pyridine in a suitable solvent such as dichloromethane, 1,4-dioxane or tetrahydrofuran at 0 °C or ambient temperature. Related dehydrations of amides to give nitriles can be found in literature, for example, see: Org. Prep. Proced. Int. 26 (1994) 429-438, Acta Chem. Scand. 53 (1999) 714-720, J. Org. Chem. 57 (1992) 2700-2705. Compounds of general formula XXIII may then be converted – if necessary – to compounds of general formula I by the syntheses outlined above. Amides of general formula XXIV can be prepared by cleavage of the chiral benzyl group of amides of general formula XXV (indicated for XXV-a). This reaction can be effected by stirring with an acid such as trifluoroacetic acid in dichloromethane.

Compounds of general formula XXV are obtained by a Hantzsch-type cyclocondensation as described previously, starting from XVI, enamine XXVI and an appropriately substituted benzaldehyde. This reaction gives a mixture of 4 stereoisomers of general formulas XXVa-d which may be separated by chromatographic techniques, such as flash column chromatography using silica gel and/or HPLC.

Enamine XXVI can be prepared by methods known to those of skill in the art in two steps from the commercially available (S)-1-(4-Methoxy-phenyl)-ethylamine and diketene or 2,2,6-trimethyl-1,3-dioxin-4-one.



10

The compounds of the invention may form hydrates or solvates. It is known to those of skill in the art that charged compounds form hydrated species when lyophilized with water, or form solvated species when concentrated in a solution with an appropriate organic solvent. The compounds of this invention include the hydrates or solvates of the compounds listed.

The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivatives of the invention were found to agonists of the FSH receptor. Methods to determine receptor binding, as well as *in vitro* and *in vivo* assays to determine biological activity, of gonadotropins are well known. In general, expressed receptor is contacted with the compound to be tested and binding or stimulation or inhibition of a functional response is measured.

20

To measure a functional response, isolated DNA encoding the FSH receptor gene, preferably the human receptor, is expressed in suitable host cells. Such a cell might be the Chinese Hamster Ovary cell, but other cells are also suitable. Preferably the cells are of mammalian origin (Jia et al, Mol.Endocrin., 5:759-776, 1991).

- 5 Methods to construct recombinant FSH expressing cell lines are well known in the art (Sambrook et al., Molecular Cloning: a Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, latest edition). Expression of receptor is attained by expression of the DNA encoding the desired protein. Techniques for site directed mutagenesis, ligation of additional sequences, PCR, and construction of suitable
- 10 expression systems are all, by now, well known in the art. Portions, or all, of the DNA encoding the desired protein can be constructed synthetically using standard solid phase techniques, preferably to include restriction sites for ease of ligation. Suitable control elements for transcription and translation of the included coding sequence can be provided to the DNA coding sequences. As is well known, expression systems are
- 15 now available which are compatible with a wide variety of hosts, including prokaryotic hosts such as bacteria and eukaryotic hosts such as yeast, plant cells, insect cells, mammalian cells, avian cells and the like.

Cells expressing the receptor are then contacted with the test compound to observe binding, or stimulation or inhibition of a functional response.

- 20 Alternatively, isolated cell membranes containing the expressed receptor may be used to measure binding of the test compound.

For measurement of binding, radioactive or fluorescent compounds may be used. As reference compound human recombinant FSH can be used.

In the alternative also competition binding assays may be performed.

- 25 Another assay involves screening for FSH receptor agonist compounds by determining stimulation of receptor mediated cAMP accumulation. Thus, such a method involves expression of the receptor on the cell surface of a host cell and exposing the cell to the test compound. The amount of cAMP is then measured. The level of cAMP will be increased, by the stimulating effect of the test compound upon binding to the receptor.

In addition to direct measurement of e.g. cAMP levels in the exposed cell, cells lines can be used which in addition to transfection with receptor encoding DNA are also transfected with a second DNA encoding a reporter gene the expression of which responds to the level of cAMP. Such reporter genes might be cAMP inducible or might
5 be constructed in such a way that they are connected to novel cAMP responsive elements. In general, reporter gene expression might be controlled by any response element reacting to changing levels of cAMP. Suitable reporter genes are e.g. LacZ, alkaline phosphatase, firefly luciferase and green fluorescence protein. The principles of such transactivation assays are well known in the art and are described e.g. in
10 Stratowa, Ch., Himmler, A. and Czernilofsky, A.P., (1995) Curr. Opin. Biotechnol. 6:574.

The present invention also relates to a pharmaceutical composition comprising a 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative or pharmaceutically acceptable salts thereof having the general formula I in admixture with
15 pharmaceutically acceptable auxiliaries and optionally other therapeutic agents. The auxiliaries must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipients thereof.

Compositions include e.g. those suitable for oral, sublingual, subcutaneous,
20 intravenous, intramuscular, nasal, local, or rectal administration, and the like, all in unit dosage forms for administration.

For oral administration, the active ingredient may be presented as discrete units, such as tablets, capsules, powders, granulates, solutions, suspensions, and the like.

For parenteral administration, the pharmaceutical composition of the invention may be
25 presented in unit-dose or multi-dose containers, e.g. injection liquids in predetermined amounts, for example in sealed vials and ampoules, and may also be stored in a freeze dried (lyophilized) condition requiring only the addition of sterile liquid carrier, e.g. water, prior to use.

Mixed with such pharmaceutically acceptable auxiliaries, e.g. as described in the
30 standard reference, Gennaro, A.R. et al., Remington: *The Science and Practice of Pharmacy* (20th Edition., Lippincott Williams & Wilkins, 2000, see especially Part 5:

Pharmaceutical Manufacturing), the active agent may be compressed into solid dosage units, such as pills, tablets, or be processed into capsules or suppositories. By means of pharmaceutically acceptable liquids the active agent can be applied as a fluid composition, e.g. as an injection preparation, in the form of a solution, suspension, 5 emulsion, or as a spray, e.g. a nasal spray.

For making solid dosage units, the use of conventional additives such as fillers, colorants, polymeric binders and the like is contemplated. In general any pharmaceutically acceptable additive which does not interfere with the function of the active compounds can be used. Suitable carriers with which the active agent of the invention 10 can be administered as solid compositions include lactose, starch, cellulose derivatives and the like, or mixtures thereof, used in suitable amounts. For parenteral administration, aqueous suspensions, isotonic saline solutions and sterile injectable solutions may be used, containing pharmaceutically acceptable dispersing agents and/or wetting agents, such as propylene glycol or butylene glycol.

15 The invention further includes a pharmaceutical composition, as hereinbefore described, in combination with packaging material suitable for said composition, said packaging material including instructions for the use of the composition for the use as hereinbefore described.

The exact dose and regimen of administration of the active ingredient, or a 20 pharmaceutical composition thereof, may vary with the particular compound, the route of administration, and the age and condition of the individual subject to whom the medicament is to be administered.

In general parenteral administration requires lower dosages than other methods of administration which are more dependent upon absorption. However, a suitable dosage 25 for humans may be 0.05-25 mg per kg body weight. The desired dose may be presented as one dose or as multiple subdoses administered at appropriate intervals throughout the day, or, in case of female recipients, as doses to be administered at appropriate daily intervals throughout the menstrual cycle. The dosage as well as the regimen of administration may differ between a female and a male recipient.

30 Thus, the compounds according to the invention can be used in therapy.

A further aspect of the invention resides in the use of a 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative having the general formula I for the manufacture of a medicament to be used for the treatment of disorders responsive to FSH receptor mediated pathways, preferably for the treatment of fertility disorders. Thus, patients in need thereof can be administered with suitable amounts of the compounds according to the invention.

In yet another aspect the invention resides in the use of a 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative having the general formula I for the manufacture of a medicament to be used for the treatment of infertility.

The invention is illustrated by the following examples.

General Comments:

The following abbreviations are used in the examples: DMA = *N,N*-dimethylaniline, DIPEA = *N,N*-diisopropylethylamine, TFA = trifluoroacetic acid, HATU = *O*-(7-azabenzotriazole-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate, Fmoc = 9-fluorenylmethoxycarbonyl, Fmoc-Cl = 9-fluorenylmethoxycarbonylchloride, Alloc = allyloxycarbonyl, DMF = *N,N*-dimethylformamide, THF = tetrahydrofuran.

Unless stated otherwise, all final products of the examples below were lyophilized from water/1,4-dioxane mixtures, water/*tert*-butanol or water/acetonitrile mixtures. If the compound was prepared as a TFA salt, the acid was added in an appropriate amount to the solvent mixture before lyophilization.

The names of the final products described in the examples were generated using the Beilstein Autonom program (version: 2.02.304).

The following analytical HPLC methods were used for determination of retention times:

- Method 1: Column: 5 μ m Luna C-18(2) 150 $\times$ 4.6 mm; flow: 1 ml/min; detection: 210 nm; column temperature: 40 $^{\circ}$ C; solvent A: CH₃CN/H₂O = 1/9 (v/v); solvent B: CH₃CN; solvent C: 0.1 M aqueous trifluoroacetic acid; gradient: solvent A/B/C = 65/30/5 to 10/85/5 (v/v/v) in 30.00 min, then constant for an additional 10.00 min at A/B/C = 10/85/5 (v/v/v).
- Method 2: Identical to method 1, except for the gradient used: Gradient: solvent A/B/C = 75/20/5 to 15/80/5 (v/v/v) in 30.00 min, then constant for an additional 10.00 min at A/B/C = 15/80/5 (v/v/v).

Method 3: Identical to method 1, except for the gradient used: Gradient: solvent A/B/C = 35/60/5 to 10/85/5 (v/v/v) in 30.00 min, then constant for an additional 10.00 min at A/B/C = 10/85/5 (v/v/v).

Method 4: Identical to method 1, except for the gradient used: Gradient: solvent A/B/C = 95/0/5 to 15/80/5 (v/v/v) in 30.00 min, then constant for an additional 10.00 min at A/B/C = 15/80/5 (v/v/v).

Method 5: Identical to method 1, except for the gradient used: Gradient: solvent A/B/C = 75/25/0 to 0/100/0 (v/v/v) in 20.00 min, then constant for an additional 10.00 min at A/B/C = 0/100/0 (v/v/v).

Method 6: Identical to method 1, except for the gradient used: Gradient: solvent A/B/C = 60/40/0 to 0/100/0 (v/v/v) in 20.00 min, then constant for an additional 10.00 min at A/B/C = 0/100/0 (v/v/v).

Method 7: Identical to method 1, except for the gradient used: Gradient: solvent A/B/C = 70/30/3 to 40/60/3 (v/v/v) in 30.00 min, then constant for an additional 10.00 min at A/B/C = 0/100/3 (v/v/v).

Method 8: Column: 3 μ m Luna C-18(2) 100 $\times$ 2 mm (Phenomenex); flow: 0.25 ml/min; detection: 210 nm; column temperature: 40 $^{\circ}$ C; solvent A: CH₃CN/H₂O = 1/9 (v/v); solvent B: CH₃CN; gradient: solvent A/B = 65/35 to 10/90 (v/v) in 30.00 min, in 2.00 minutes to A/B = 0/100 (v/v), then constant for an additional 8.00 min at A/B = 0/100 (v/v), then in 1.00 minute to A/B = 65/35 (v/v) and finally constant for an additional 15.00 min at A/B = 65/35 (v/v).

Method 9: Column: Chiralpak AD-H 25 x 0.46 cm; detection: Chirality (+/-) and 210 nm; Conditions : isocratic heptane/isopropylalcohol 80/20 (v/v).

The diastereomeric ratio (Diast. ratio:) was determined if baseline separation of the individual diastereomers was observed using the appropriate analytical HPLC method. Alternatively, the diastereomeric ratio was determined by ¹H NMR analysis when distinct signals corresponding to the diastereomers were identified.

Preparative HPLC-purifications were performed on a Luna C-18(2) column (5 μ m) (150 $\times$ 21.2 mm) with water/acetonitrile mixtures, optionally in the presence of 0.1% aqueous TFA, using the indicated gradient: Flow: 20 ml/min: Detection: 210 nm:

Example 1

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3,4,5-trimethoxy-benzamide

(a). 3-Bromo-5-ethoxy-4-(3-nitro-benzyloxy)-benzaldehyde

- 5 A mixture of 3-nitrobenzylbromide (2 g), 3-bromo-5-ethoxy-4-hydroxy benzaldehyde (2.06 g), K₂CO₃ (2.55 g) and Bu₄NI (186 mg) in DMF (50 ml) was stirred at 60°C for 2 h. The reaction mixture was diluted with ethyl acetate and washed with water and brine. The organic layer was dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was recrystallized from heptane.

- 10 Yield: 2.98 g. MS-ESI: [M+H]⁺ = 380/382

(b). 4-[4-(3-Nitro-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

A mixture of the product of example 1a (2g), 3-aminocrotonitrile (431 mg) and 5-propylcyclohexane-1,3-dione (810 mg) in ethanol (20 ml) was stirred at 80°C for 4 h.

- 15 The reaction mixture was concentrated *in vacuo* and the residue was chromatographed on silica gel in heptane/ethyl acetate = 1/1 (v/v) as eluent.

Yield: 2.7 g. MS-ESI: [M+H]⁺ = 580/582

(c). 4-[4-(3-Amino-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- 20 SnCl₂ (1.62 g) was added to a solution of the product of example 1b (500 mg) in ethanol (10 ml). After stirring for 18 h, the reaction mixture was poured in water and extracted with ethyl acetate. The organic layer was separated and washed with 0.5N NaOH and water, dried (MgSO₄), filtered and concentrated *in vacuo*.

Yield: 440 mg. MS-ESI: [M+H]⁺ = 550/552

- 25 (d). N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3,4,5-trimethoxy-benzamide

A mixture of the product of example 1c (44 mg), 3,4,5-trimethoxybenzoylchloride (39 mg) and DIPEA (70 µl) in dichloromethane (10 ml) was stirred for 18 h. The reaction mixture was diluted with dichloromethane and washed with water and brine. The organic layer was separated, dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (0 → 90% acetonitrile). Yield: 39 mg. MS-ESI: [M+H]⁺ = 744.4/746.4; anal. HPLC: Rt = 20.73 min 1 (method 1)

Example 2

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3,3-dimethyl-butyramide

Acylation of the product described in example 1c (44 mg) with 3,3-dimethyl-butyryl chloride (16 mg) was performed according to the method described in example 1d.
Yield: 36 mg. MS-ESI: $[M+H]^+ = 648/650$; anal. HPLC: $R_t = 22.58$ min (method 1)

Example 3

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[(pyridin-4-ylmethyl)-amino]-acetamide

(a). 2-Bromo-N-{3-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-acetamide

Bromoacetyl chloride (600 μ l) in dichloromethane (5 ml) was added dropwise to a solution of the product of example 1c (1.32 g) and DIPEA (2.1 ml) in dichloromethane (25 ml). After stirring for 4 h, the reaction mixture was washed with water. The organic layer was separated, dried ($MgSO_4$), filtered and concentrated *in vacuo*. Yield: 900 mg.
MS-ESI: $[M+H]^+ = 670/672/674$

(b). N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[(pyridin-4-ylmethyl)-amino]-acetamide

A mixture of the product of example 3a (50 mg) and 4-picolylamine (81 mg) in dichloromethane/THF = 4:1 (v/v) (2.5 ml) was stirred for 18 h. The reaction mixture was diluted with dichloromethane and washed with water and brine. The organic layer was separated, dried ($MgSO_4$), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (0 $\rightarrow$ 90% acetonitrile, 0.1% TFA). Yield: 17.9 mg (as TFA salt). MS-ESI: $[M+H]^+ = 698.4/700.4$; anal. HPLC: $R_t = 8.85$ min (diast1), $R_t = 9.07$ min (diast2) (method 2)

Example 4

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-(cyclopentyl-methyl-amino)-acetamide

Reaction of the product described in example 3a (50mg) with cyclopentyl-methylamine (74 mg) was performed according to the method described in example 3b. Yield: 53 mg (as TFA salt). MS-ESI: $[M+H]^+ = 689/691$; anal. HPLC: $R_t = 13.45$ min (diast1), $R_t = 13.72$ min (diast2) (method 2).

Example 5

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-acetamide

- 5 Reaction of the product described in example 3a (50mg) with 2-piperazin-1-yl-ethanol (98mg) was performed according to the method described in example 3b. Yield: 41 mg (as TFA salt). MS-ESI: $[M+H]^+ = 720/721$; anal. HPLC: $R_t = 10.40$ min (diast1) $R_t = 10.67$ min (diast2) (method 2).

Example 6

- 10 3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-ethyl-N-(2-hydroxy-ethyl)-benzamide

(a). 3-(2-Bromo-6-ethoxy-4-formyl-phenoxy-methyl)-benzoic acid methyl ester

- Alkylation of 3-bromo-5-ethoxy-4-hydroxy benzaldehyde (2 g) with 3-bromomethyl-benzoic acid methyl ester (2.06 g) was performed according to the method described in
 15 example 1a. The residue was chromatographed on silica gel in heptane/ethyl acetate = 7/3 (v/v) as eluent. Yield: 2.42 g. MS-ESI: $[M+H]^+ = 393/395$

(b). 3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzoic acid methyl ester

- The title compound was prepared according to the method described in example 1b
 20 starting from the product of example 6a (2.4 g), 3-aminocrotonitrile (504 mg) and 5-propylcyclohexane-1,3-dione (947 mg). Yield: 2.96 g. MS-ESI: $[M+H]^+ = 593/595$

(c). 3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzoic acid

- A mixture of the product of example 6b (2.96 g) and 2N NaOH (5 ml) in dioxane (100
 25 ml) was stirred for 3 days. The reaction mixture was poured in water followed by addition of 4N HCl until pH 2 and extraction with ethyl acetate. The organic layer was washed with brine, dried ($MgSO_4$), filtered and concentrated *in vacuo*. Yield: 3.0 g. MS-ESI: $[M+H]^+ = 579/581$

- (d). 3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-ethyl-N-(2-hydroxy-ethyl)-benzamide
 30

A mixture of the product of example 6c (90 mg), 2-ethylamino-ethanol (49 μ l) and TBTU (82 mg) in dichloromethane (5 ml) was stirred for 4h. The reaction mixture was washed with sat. $NaHCO_3$, dried ($MgSO_4$) and concentrated *in vacuo*. The residue was

purified by preparative HPLC (0 → 90% acetonitrile, 0.1% TFA). Yield: 43 mg. MS-ESI: $[M+H]^+ = 650/652$; anal. HPLC: $R_t = 19.96$ min (diast 1), $R_t = 20.22$ min (diast 2) (method 2). Diast. ratio: 3:1

Example 7

- 5 4-{3-Bromo-4-[3-(3,6-dihydro-2H-pyridine-1-carbonyl)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

Coupling of the compound described in example 6c with 1,2,3,6-tetrahydro-pyridine (47 μ l) and TBTU (82 mg) was performed according to the method described in example 6d. Yield: 61 mg. MS-ESI: $[M+H]^+ = 644/646$; anal. HPLC: $R_t = 23.48$ min (diast 1), $R_t = 23.75$ min (diast 2) (method 2) Diast. ratio: 3:1

Example 8

{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-acetic acid tert-butyl ester

(a). 3-Bromomethyl-phenol

- 15 At 0 °C and under a nitrogen atmosphere carbontetrabromide (28.05 g) was added over a period of 20 min to a suspension of 3-hydroxybenzylalcohol (7 g) and triphenylphosphine (22.18 g) in dichloromethane (250 ml). After stirring for 1 h at 0°C, the reaction mixture was concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 2/1 (v/v) as eluent. Yield: 10 g. MS-ESI: $[M-H]^- =$
20 184.8/186.8

(b). 2-(3-Bromomethyl-phenoxy)-tetrahydro-pyran

- At 0°C 3,4-dihydro-2H-pyran (9.8 ml) and a catalytic amount of pyridinium-*p*-toluenesulfonate were added to a solution of the product of example 8a (10 g) in dichloromethane (300 ml). After stirring for 1.5 h at room temperature, the reaction
25 mixture was diluted with dichloromethane and washed with sat. NaHCO_3 and brine. The organic layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 4/1 (v/v). Yield: 13.6 g.

(c). 3-Bromo-5-ethoxy-4-[3-(tetrahydro-pyran-2-yloxy)-benzyloxy]-benzaldehyde

- 30 Alkylation of 3-ethoxy-4-hydroxy-5-bromo-benzaldehyde (11.19 g) with the product of example 8b (13.6 g) was performed according to the method described in example 1a. The residue was chromatographed on silica gel in heptane/ethyl acetate = 3/1 (v/v) as eluent. Yield: 17.8 g. MS-ESI: $[M+H]^+ = 435/437$

(d). 4-{3-Bromo-5-ethoxy-4-[3-(tetrahydro-pyran-2-yloxy)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was prepared according to the method described in example 1b, using the product of example 8c (5.5 g), 3-aminocrotonitrile (1 g) and 5-propylcyclohexane-1,3-dione (1.95 g). Yield: 6.4 g. MS-ESI: $[M+H]^+ = 635.4/637.4$

(e). 4-[3-Bromo-5-ethoxy-4-(3-hydroxy-benzyloxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

Oxalic acid was added to a solution of the product of example 8d (6.4 g) in methanol (135 ml) and water (15 ml) to pH 2. After stirring at 55°C for 2 h, the reaction mixture was concentrated *in vacuo*. The residue was dissolved in ethyl acetate and washed with water, sat. NaHCO_3 and brine. The organic layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. Yield: 5.5 g. MS-ESI: $[M+H]^+ = 551.2/553.2$

(f). {3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-acetic acid *tert*-butyl ester

A mixture of the product of example 8e (100 mg), *tert*-butyl bromoacetate (105 mg) and CsCO_3 (355 mg) in dioxane (7 ml) was stirred at 80°C under a nitrogen atmosphere for 18 h. The reaction mixture was diluted with ethyl acetate and washed with water, sat. NaHCO_3 and brine. The organic layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (50 → 100% acetonitrile). Yield: 34.5 mg. MS-ESI: $[M+H]^+ = 665.4/667.4$; anal. HPLC: $R_t = 10.16$ min (diast1), 10.55 min (diast2) (method 3) Diast. ratio: 5:1

Example 9

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-(3-pyrrolidin-1-yl-propylamino)-acetamide

Reaction of the product described in example 3a (100mg) with 3-pyrrolidin-1-yl-propylamine (191 mg) was performed according to the method described in example 3b. Yield: 32 mg (as TFA salt). MS-ESI: $[M+H]^+ = 718.6/720.6$; anal. HPLC: $R_t = 7.24$ min (method 2).

Example 10

{3-(Benzyl-methyl-amino)-*N*-{3-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-propionamide

(a). *N*-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-acrylamide

To a solution of the product of example 1c (1.5 g) in dichloromethane was added acryloylchloride (222 μ l) and DIPEA (2.38 ml). After stirring for 18 h, the reaction mixture was washed with water. The organic layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. Yield: 1.65 g. ESI-MS: $[\text{M}+\text{H}]^+ = 604.2/606.2$

- 5 (b). {3-(Benzyl-methyl-amino)-N-[3-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl]-propionamide

A mixture of the product of example 10a (150 mg) and N-methyl-benzylamine (300 mg) in THF (2.5 ml) was stirred for 18 h. The reaction mixture was concentrated *in vacuo*. The residue was dissolved in ethyl acetate and washed with water. The organic
10 layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (10 $\rightarrow$ 90% acetonitrile, 0.1% TFA). Yield: 34 mg (as TFA salt). MS-ESI: $[\text{M}+\text{H}]^+ = 725.4/727.4$; anal. HPLC: $R_t = 14.32$ min (diast1), $R_t = 14.58$ min (diast2) (method 2)

Example 11

- 15 3-[Bis-(2-methoxy-ethyl)-amino]-N-[3-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl]-propionamide

The title compound was obtained according to the method described for example 10b, starting from Bis-(2-methoxy-ethyl)-amine (319 mg) and the product of example 10a
20 (150 mg). Yield: 30 mg (as TFA salt). MS-ESI: $[\text{M}+\text{H}]^+ = 737.4/739.4$; anal. HPLC: $R_t = 13.24$ min (method 2).

Example 12

4-{3-Bromo-5-ethoxy-4-[3-(pyridin-2-ylmethoxy)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- 25 A mixture of the product of example 8e (100 mg), 2-picolychloride hydrochloride (31.2 mg) and CsCO_3 (124 mg) in DMF (7 ml) was stirred at 60°C under a nitrogen atmosphere for 18 h. The reaction mixture was diluted with ethyl acetate and washed with water, sat. NaHCO_3 and brine. The organic layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (0
30 $\rightarrow$ 100% acetonitrile, 0.1% TFA). Yield: 19.6 mg (as TFA salt). MS-ESI: $[\text{M}+\text{H}]^+ = 642.1/644.1$; anal. HPLC: $R_t = 17.24$ min (method 2).

Example 13

4-{3-Bromo-5-ethoxy-4-[3-(4-fluoro-benzyloxy)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was obtained according to the method described for example 12, starting from 4-fluoro-benzylbromide (23.7 μ l) and the product of example 8e (100 mg). The residue was purified by preparative HPLC (50 $\rightarrow$ 100% acetonitrile). Yield: 58.9 mg. MS-ESI: $[M+H]^+ = 659.2/661.2$; anal. HPLC: $R_t = 14.38$ min (diast1), $R_t =$
5 14.90 min (diast2) (method 3). Diast. ratio: 5:1

Example 14

4-{3-Bromo-4-[3-(2-diethylamino-ethoxy)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was obtained according to the method described for example 12,
10 starting from 2-diethylaminoethylchloride hydrochloride (32.7 mg) and the product of example 8e (100 mg). The residue was purified by preparative HPLC (0 $\rightarrow$ 100% acetonitrile, 0.1% TFA). Yield: 62.9 mg (as TFA salt). MS-ESI: $[M+H]^+ = 650.4/652.4$; anal. HPLC: $R_t = 13.35$ min (diast 1), $R_t = 13.70$ (diast 2) (method 2).

Example 15

15 2-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-N,N-dimethyl-acetamide

The title compound was obtained according to the method described for example 12, starting from 2-chloro-N,N-dimethylacetamide (19.6 μ l) and the product of example 8e (100 mg). The residue was purified by preparative HPLC (0 $\rightarrow$ 100% acetonitrile).
20 Yield: 53.8 mg. MS-ESI: $[M+H]^+ = 636.4/638.4$; anal. HPLC: $R_t = 20.44$ min (diast1), $R_t = 20.77$ min (diast2) (method 2). Diast. ratio: 5:1

Example 16

4-{3-Bromo-5-ethoxy-4-[3-(2-methoxy-ethoxy)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

25 A mixture of the product of example 8e (100 mg), 2-methoxyethanol (29 μ l), diethylazodicarboxylate (57 μ l) and triphenylphosphine polymer bound (122 mg) in dichloromethane (6 ml) was stirred under a nitrogen atmosphere for 54 h. The reaction mixture was filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (50 $\rightarrow$ 100% acetonitrile). Yield: 15 mg. MS-ESI: $[M+H]^+ =$
30 609.2/611.2; anal. HPLC: $R_t = 7.66$ min (method 3).

Example 17

4-{3-Bromo-5-ethoxy-4-[3-(2-morpholin-4-yl-2-oxo-ethoxy)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

(a). {3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-acetic acid

A mixture of the compound of example 8f (620 mg) in dichloromethane (45 ml) and trifluoroacetic acid (5 ml) was stirred for 4 h. The reaction mixture was diluted with
5 dichloromethane, washed with water and brine. The organic layer was separated, dried (MgSO₄), filtered and concentrated *in vacuo*. Yield: 553 mg. MS-ESI: [M+H]⁺ = 609.2/611.2

(b). 4-{3-Bromo-5-ethoxy-4-[3-(2-morpholin-4-yl-2-oxo-ethoxy)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

10 Coupling of the compound described in example 17a (79 mg) with morpholine (34 µl) and TBTU (62 mg) was performed according to the method described in example 6d. Yield: 50.5 mg. MS-ESI: [M+H]⁺ = 678.4/680.4; anal. HPLC: Rt = 16.29 min (diast1), Rt = 16.65 (diast2) (method 1). Diast. ratio: 5:1

Example 18

15 2-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-N-thiophen-2-ylmethyl-acetamide

Coupling of the compound described in example 17a (75 mg) with 2-thiophenemethylamine (38 µl) and TBTU (59 mg) was performed according to the method described in example 6d. The residue was purified by preparative HPLC (40 →
20 100% acetonitrile). Yield: 45.7 mg. MS-ESI: [M+H]⁺ = 704.4/706.4; anal. HPLC: Rt = 6.96 min (method 3).

Example 19

N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-pyrrolidin-1-yl-acetamide

25 (a). 3-Bromo-5-ethoxy-4-(2-nitro-benzyloxy)-benzaldehyde

Alkylation of 3-bromo-5-ethoxy-4-hydroxybenzaldehyde (5 g) with 2-nitrobenzylbromide (4.85 g) was performed according to the method described in example 1a. Yield: 7.88 g. MS-ESI: [M+H]⁺ = 380/382

(b). 4-[3-Bromo-5-ethoxy-4-(2-nitro-benzyloxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile
30

The title compound was prepared according to the method described in example 1b starting from the product of example 19a (7.88 g), 3-aminocrotonitrile (1.7 g) and 5-propylcyclohexane-1,3-dione (3.2 g). Yield: 8.54 g. MS-ESI: [M+H]⁺ = 580.2/582.2

(c). 4-[4-(2-Amino-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

At 0°C Zinc dust (9.01 g) was added portion wise to a solution of the product of example 19c (6.9 g) and acetic acid (5.9 ml) in THF (150 ml). After stirring at 0°C for 1 h and at room temperature for 2 h, the reaction mixture was filtered, diluted with dichloromethane and washed with sat. NaHCO₃ and brine. The organic layer was separated, dried (MgSO₄), filtered and concentrated *in vacuo*. Yield: 3.66 g. MS-ESI: [M+H]⁺ = 550.2/552.2

(d). 2-Bromo-N-{2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-acetamide

Reaction of bromoacetyl chloride (0.546 ml) with the product of example 19c (1.2 g) was performed according to the method described for example 3a. Yield: 1.36 g. MS-ESI: [M+H]⁺ = 626.2/628.2

(e). N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-pyrrolidin-1-yl-acetamide

A mixture of the product of example 19d (136 mg) and pyrrolidine (181 µl) in THF (5 ml) was stirred for 54 h. The reaction mixture was diluted with ethyl acetate and washed with water, sat. NaHCO₃ and brine. The organic layer was dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (10 → 100% acetonitrile, 0.1% TFA). Yield: 51.8 mg (as TFA salt). MS-ESI: [M+H]⁺ = 661.4/663.4; anal. HPLC: Rt = 13.58 (method 2)

Example 20

N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[(2-dimethylamino-ethyl)-methyl-amino]-acetamide

The title compound was prepared according to the method described in example 19e starting from the product of example 19d (136 mg) and *N,N,N'*-trimethyl-ethane-1,2-diamine (277 µl). Yield: 41 mg (as TFA salt). MS-ESI: [M+H]⁺ = 690.4/692.4 ; anal. HPLC: Rt = 12.14 min (diast 1), Rt = 12.46 (diast 2) (method 2). Diast. ratio: 7:1

Example 21

4-[3-Bromo-5-ethoxy-4-(2-methylsulfonyl-pyrimidin-4-ylmethoxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

(a). 5-Bromo-2-methylsulfonyl-pyrimidine-4-carboxylic acid

At 5°C triethylamine (16.9 ml) was added dropwise to a solution of mucobromic acid (10.31 g) and 2-methyl-isothioureia (11.13 g) in water (200 ml). After stirring at 5°C for 3 h, the reaction mixture was allowed to warm up (room temperature) and stirring was continued for another 54 h. The reaction mixture was acidified with conc. HCl (33%).

- 5 The title product was obtained by filtration. Yield: 4.1 g. MS-ESI: $[M+H]^+ = 249/251$

(b). 2-Methylsulfanyl-pyrimidine-4-carboxylic acid

The product of example 21a (517 mg) in methanol (25 ml) was hydrogenated in a PARR apparatus in the presence of KOH (260 mg) and 10% Pd on BaSO₄ (260 mg) for 4 h. The reaction mixture was filtered over decalite and washed with methanol (warm).

- 10 The filtrate was concentrated to 50% of its volume followed by addition of conc. HCl (33%) to pH 1. The precipitate (KBr) was filtered off and the mother liquor was concentrated *in vacuo*. The residue was recrystallized from dioxane. Yield: 200 mg. MS-ESI: $[M+H]^+ = 170$

(c). 2-Methylsulfanyl-pyrimidin-4-yl)-methanol

- 15 At 0°C oxalyl chloride (295 µl) was added to a solution of DMF (91.2 µl) in dichloromethane (2 ml). After stirring at 0°C for 1 h, the reaction mixture was concentrated *in vacuo*. The residue was dissolved in acetonitrile (2 ml) and THF (3 ml), followed by portion wise addition of the product of example 21b (200 mg) at 0°C in a nitrogen atmosphere. After stirring at 0°C for 15 min, the reaction mixture was cooled
20 (-78°C) and 2M NaBH₄ in DMF (590 µl) was added. After stirring at -20°C for 3 h, the reaction mixture was quenched with 2N HCl and concentrated *in vacuo*. The residue was dissolved in water and the remaining solution was adjusted to pH 11 with 2N NaOH, followed by extraction with dichloromethane. The organic layer was separated, dried (MgSO₄), filtered and concentrated *in vacuo*. Yield: 180 mg. MS-ESI: $[M+H]^+ =$
25 156

(d). 3-Bromo-5-ethoxy-4-(2-methylsulfanyl-pyrimidin-4-ylmethoxy)-benzaldehyde

A mixture of the product of example 21c (180 mg), 3-bromo-5-ethoxy-4-hydroxy benzaldehyde (289 mg), triphenylphosphine polymer bound (393 mg) and diethylazodicarboxylate (186 µl) in dichloromethane (25 ml) was stirred for 18 h. The
30 reaction mixture was filtered off and the filtrate concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 3/2 (v/v) as eluent.

Yield: 98 mg. MS-ESI: $[M+H]^+ = 383/385$

(e). 4-[3-Bromo-5-ethoxy-4-(2-methylsulfanyl-pyrimidin-4-ylmethoxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was prepared according to the method described in example 1b starting from the product of example 22d (49 mg), 3-aminocrotonitrile (11 mg) and 5-propylcyclohexane-1,3-dione (20 mg). The residue was purified by preparative HPLC (0 → 100% acetonitrile, 0.1% TFA). Yield: 24 mg (as TFA salt). MS-ESI: $[M+H]^+ = 583.2/585.2$; anal. HPLC: $R_t = 25.06$ min (diast1), $R_t = 25.51$ min (diast2) (method 2). Diast. ratio: 4:1

Example 22

Furan-2-carboxylic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide

10 A mixture of the product of example 19c (100 mg), 2-furoylchloride (36 μ l) and DIPEA (159 μ l) in dichloromethane (5 ml) was stirred for 16 h. The reaction mixture was diluted with dichloromethane and washed with sat. NaHCO_3 . The organic layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (20 → 100% acetonitrile). Yield: 67 mg. MS-ESI: $[M+H]^+ = 644.4/646.4$; anal. HPLC: $R_t = 20.19$ min (method 1).

Example 23

N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-acrylamide

20 Reaction of the product of example 19c (100 mg) with acryloylchloride (29 μ l) in the presence of DIPEA (159 μ l) was performed according to the method described in example 22. Yield: 51.1 mg. MS-ESI: $[M+H]^+ = 604.4/606.4$; anal. HPLC: $R_t = 19.01$ min (diast1), $R_t = 19.42$ min (diast2) (method 1). Diast. ratio: 9:1

Example 24

25 Cyclopropanecarboxylic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide

Reaction of the product of example 19c (100 mg) with cyclopropanecarbonylchloride (33 μ l) in the presence of DIPEA (159 μ l) was performed according to the method described in example 22. Yield: 52.1 mg. MS-ESI: $[M+H]^+ = 618.2/620.2$; anal. HPLC: $R_t = 20.30$ min (diast1), $R_t = 20.71$ min (diast2) (method 1). Diast. ratio: 7:1

30 Example 25

2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid methyl ester

Reaction of the product of example 19c (100 mg) with methylchloroformate (28 μ l) in the presence of DIPEA (159 μ l) was performed according to the method described in example 22. Yield: 65.9 mg. MS-ESI: $[M+H]^+ = 608.2/610.2$; anal. HPLC: $R_t = 21.37$ min (diast1), $R_t = 21.87$ min (diast2) (method 1). Diast. ratio: 11:1

5 **Example 26**

2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid isobutyl ester

Reaction of the product of example 19c (100 mg) with isobutyl chloroformate (47 μ l) in the presence of DIPEA (159 μ l) was performed according to the method described in
10 example 22. Yield: 70 mg. MS-ESI: $[M+H]^+ = 650.4/652.4$; anal. HPLC: $R_t = 26.84$ min (diast), $R_t = 27.26$ min (diast2) (method 1). Diast. ratio: 7:1

Example 27

2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-furan-3-carboxylic acid methyl ester

15 (a). 2-(2-Bromo-6-ethoxy-4-formyl-phenoxy-methyl)-furan-3-carboxylic acid methyl ester

Alkylation of 3-bromo-5-ethoxy-4-hydroxy benzaldehyde (123 mg) with 2-bromomethyl-furan-3-carboxylic acid methyl ester (109 mg) was performed according to the method described in example 1a. Yield: 191 mg. MS-ESI: $[M+H]^+ = 383/385$

20 (b). 2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-furan-3-carboxylic acid methyl ester

The title compound was prepared according to the method described in example 1b starting from the product of example 27a (191 mg), 3-aminocrotonitrile (41 mg) and 5-propylcyclohexane-1,3-dione (77 mg). The residue was purified by preparative HPLC
25 (10 $\rightarrow$ 90% acetonitrile). Yield: 168 mg. MS-ESI: $[M+H]^+ = 583/585$; anal. HPLC: $R_t = 23.06$ min (diast1), $R_t = 23.43$ min (diast2) (method 2). Diast. ratio: 4:1

Example 28

Propane-1-sulfonic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide

30 Propane-1-sulfonyl chloride (31 μ l) and pyridine (44 μ l) were added to a solution of the product of example 19c (100 mg) in dichloromethane (3 ml). After stirring for 18 h, the reaction mixture was diluted with dichloromethane and washed with sat. NaHCO_3 . The organic layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. The

residue was purified by preparative HPLC (20 → 100% acetonitrile). Yield: 52.7 mg. MS-ESI: $[M+H]^+$ = 656.4/658.4; anal. HPLC: R_t = 22.36 min (diast1), R_t = 22.72 min (diast2) (method 1). Diast. ratio: 7:1

Example 29

- 5 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-(2-methoxy-ethyl)-benzamide

(a). 4-(2-Bromo-6-ethoxy-4-formyl-phenoxy-methyl)-benzoic acid methyl ester

Alkylation of 3-bromo-5-ethoxy-4-hydroxy benzaldehyde (4 g) with 4-bromomethyl-benzoic acid methyl ester (3.73 g) was performed according to the method described in
10 example 1a. Yield: 6.4 g. MS-ESI: $[M+H]^+$ = 393/395

(b). 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzoic acid methyl ester

The title compound was prepared according to the method described in example 1b starting from the product of example 29a (6.4 g), 3-aminocrotonitrile (1.34 g) and 5-
15 propylcyclohexane-1,3-dione (2.51 g). Yield: 6.74 g. MS-ESI: $[M+H]^+$ = 593/595

(c). 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzoic acid

A mixture of the product of example 29b (6.5 g) and 2N NaOH (11.3 ml) in dioxane (400 ml) was stirred at 50°C for 18 h. The reaction mixture was poured in water and in
20 addition 4N HCl was added until pH 2, followed by extraction with ethyl acetate. The organic layer was washed with brine, dried ($MgSO_4$), filtered and concentrated *in vacuo*. Yield: 6.4 g. MS-ESI: $[M+H]^+$ = 579/581

(d). 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-(2-methoxy-ethyl)-benzamide

25 Coupling of the compound described in example 29c (100 mg) with 2-methoxy-ethylamine (44 μ l) and TBTU (82 mg) was performed according to the method described in example 6d. Yield: 70 mg. MS-ESI: $[M+H]^+$ = 636.4/638.4; anal. HPLC: R_t = 19.25 min (diast1), 19.55 min (diast2) (method 2). Diast. ratio: 4:1 (HPLC)

Example 30

- 30 4-{3-Bromo-5-ethoxy-4-[3-(isobutylamino-methyl)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

(a). 4-(3-Bromo-5-ethoxy-4-hydroxy-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was prepared according to the method described in example 1b starting from 3-bromo-5-ethoxy-4-hydroxy-benzaldehyde (6 g), 3-aminocrotonitrile (2.01 g) and 5-propylcyclohexane-1,3-dione (3.8 g). The reaction mixture was cooled (0°C) and the title compound was filtered off and washed with ethanol (cold). Yield:

5 6.3 g. MS-ESI: $[M+H]^+ = 445/447$

(b). 4-[3-Bromo-4-(3-bromomethyl-benzyloxy)-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

A mixture of the product of example 30a (1.96 g), α,α -dibromo-m-xylene (9.31 g) and K_2CO_3 (1.22 g) in DMF (100 ml) was stirred at 60°C under a nitrogen atmosphere for 2
10 h. The reaction mixture was diluted with ethyl acetate and washed with water, sat. $NaHCO_3$ and brine. The organic layer was dried ($MgSO_4$), filtered and concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 1/2 (v/v). Yield: 2.52 g MS-ESI: $[M+H]^+ = 627.4/629.4/631.4$; anal. HPLC: $R_t = 17.71$ min (diast1), 18.07 min (diast2) (method 6). Diast. ratio: 4:1

15 (c). 4-{3-Bromo-5-ethoxy-4-[3-(isobutylamino-methyl)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

A mixture of the product of example 30b (100 mg) and iso-butylamine (79 μ l) in acetonitrile (5 ml) was stirred for 18 h. The reaction mixture was concentrated *in vacuo*; the residue was dissolved in dichloromethane and washed with sat. $NaHCO_3$.
20 The organic layer was separated, dried ($MgSO_4$), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (0 $\rightarrow$ 100% acetonitrile, 0.1% TFA). Yield: 76 mg (as TFA salt). MS-ESI: $[M+H]^+ = 620.4/622.4$; anal. HPLC: $R_t = 13.56$ min (diast1), $R_t = 13.85$ min (diast2) (method 2). Diast. ratio: 3:1

Example 31

25 4-[3-Bromo-5-ethoxy-4-(3-{[(pyridin-4-ylmethyl)-amino]-methyl}-benzyloxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was prepared according to the method described in example 30c starting from the product of example 30b (100 mg) and 4-(aminomethyl)pyridine (81 μ l) in acetonitrile (5 ml). Yield: 41 mg (as TFA salt). MS-ESI: $[M+H]^+ =$
30 655.2/657.2; anal. HPLC: $R_t = 8.08$ min (method 2). Diast. ratio: 3:1

Example 32

4-[4-(3-Aminomethyl-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

(a). 4-[4-(3-Azidomethyl-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

A mixture of the product of example 30b (250 mg) and NaN_3 (78 mg) in DMF (10 ml) was stirred for 2 h. The reaction mixture was poured in water and the precipitate was
5 filtered off, washed with water and dried *in vacuo* (50°C). Yield: 197 mg. MS-ESI: $[\text{M}+\text{H}]^+ = 590.4/592.4$; anal. HPLC: $R_t = 23.72$ min (diast1), $R_y = 24.12$ min (diast2) (method 1). Diast. ratio: 3.5:1

(b). 4-[4-(3-Aminomethyl-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

10 A mixture of the product of example 32a (133 mg), triphenylphosphine polymer bound (150 mg) and 10 drops of water in THF/dichloromethane = 2:1 (v/v) was stirred at 40°C for 18 h. The reaction mixture was filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (0 → 100% acetonitrile, 0.1% TFA). Yield: 197 mg (as TFA salt). MS-ESI: $[\text{M}+\text{H}]^+ = 564.2/566.2$; anal. HPLC: $R_t = 10.31$ min
15 (diast1), $R_t = 10.58$ min (diast2) (method 2). Diast. ratio: 3:1

Example 33

4-[3-Bromo-5-ethoxy-4-(3-hydroxymethyl-benzyloxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

A mixture of the product of example 30b (100 mg) and CaCO_3 (80 mg) in
20 dioxane/water = 1/1 (v/v) was stirred at 100°C for 18 h. The reaction mixture was diluted with dichloromethane followed by addition of 2N HCl. The organic layer was separated, washed with sat. NaHCO_3 , dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (20 → 100% acetonitrile). Yield: 44 mg. MS-ESI: $[\text{M}+\text{H}]^+ = 565.2/567.2$; anal. HPLC: $R_t = 15.96$ min (diast1), $R_t =$
25 16.33 min (diast2) (method 1). Diast. ratio: 4:1

Example 34

1-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3-methyl-urea

A mixture of the compound of example 19c and 4-nitrophenyl chloroformate (55 mg)
30 in dichloromethane (4 ml) was stirred for 2 h. Methylamine (2.7 ml, 2M in THF) was added and stirring was continued for another 3 h. The reaction mixture was diluted with dichloromethane and washed with sat. NaHCO_3 . The organic layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (20 → 100% acetonitrile). Yield: 54 mg. MS-ESI: $[\text{M}+\text{H}]^+ =$

607.2/609.2; anal. HPLC: Rt = 15.88 min (diast1), Rt = 16.18 min (diast2) (method 1).
Diast. ratio: 8:1

Example 35

Piperazine-1-carboxylic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide

The title compound was prepared according to the method described in example 34 starting from the product of example 19c (150 mg), 4-nitrophenyl chloroformate (55 mg) and piperazine (469 mg). The residue was purified by preparative HPLC (0 → 100% acetonitrile, 0.1% TFA). Yield: 69 mg (as TFA salt). MS-ESI: $[M+H]^+ = 662.2/664.2$; anal. HPLC: Rt = 11.70 min (diast1), Rt = 12.04 min (diast2) (method 1).
Diast. ratio: 5:1

Example 36

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzyl}-3,3-dimethyl-butyramide

15 Acylation of the product of example 32b (100 mg) with *tert*-butylacetyl chloride (27 μ l) was performed according to the method described in example 1d. Yield: 33.5 mg. MS-ESI: $[M+H]^+ = 662.4/664.4$; anal. HPLC: Rt = 21.26 min (diast1), Rt = 21.54 min (diast2) (method 1).

Example 37

20 {3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzyl}-carbamic acid ethyl ester

Reaction of the product of example 32b (100 mg) with ethylchloroformate (19 μ l) in the presence of DIPEA (93 μ l) was performed according to the method described in example 22. Yield: 37 mg. MS-ESI: $[M+H]^+ = 636.2/638.2$; anal. HPLC: Rt = 19.81 min (diast1), Rt = 20.14 min (diast2) (method 1). Diast. ratio: 4.5:1

Example 38

{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-pyridin-2-yl}-carbamic acid methyl ester

(a). 4-(2-Amino-pyridin-3-ylmethoxy)-3-bromo-5-ethoxy-benzaldehyde

30 Mitsunobu reaction of 3-bromo-5-ethoxy-4-hydroxy benzaldehyde (490 mg) with (2-amino-pyridin-3-yl)-methanol (250 mg) was performed according to the method described for example 21d. The residue was chromatographed on silica gel in

dichloromethane/methanol = 99/1 (v/v) as eluent. Yield: 500 mg. MS-ESI: $[M+H]^+ = 351/353$.

(b). 4-[4-(2-Amino-pyridin-3-ylmethoxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- 5 The title compound was prepared according to the method described in example 1b starting from the product of example 38a (500 mg), 3-aminocrotonitrile (116mg) and 5-propylcyclohexane-1,3-dione (218 mg). Yield: 290 mg. MS-ESI: $[M+H]^+ = 551/553$

(c). {3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-pyridin-2-yl}-carbamic acid methyl ester

- 10 Reaction of the product of example 38b (50 mg) with methylchloroformate (8 μ l) in the presence of DIPEA (47 μ l) was performed according to the method described in example 22. The residue was purified by preparative HPLC (10 $\rightarrow$ 90% acetonitrile). Yield: 17 mg. MS-ESI: $[M+H]^+ = 609/611$; anal. HPLC: $R_t = 13.82$ min (diast1), $R_t = 14.08$ min (diast2) (method 5). Diast. ratio: 7:2

15 **Example 39**

4-(3-Bromo-5-ethoxy-4-{3-[(1H-imidazol-4-ylmethyl)-amino]-benzyloxy}-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- A mixture of the product of example 1c (100 mg), 1H-imidazol-4-carbaldehyde (21 mg), NaCNBH₄ (25 mg) and acetic acid (114 μ l) in methanol (3 ml) was stirred for 18 h. The reaction mixture was diluted with ethyl acetate, washed with sat. NaHCO₃ and brine. The organic layer was separated, dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (40 $\rightarrow$ 100% acetonitrile). Yield: 13 mg. MS-ESI: $[M+H]^+ = 550.2/552.2$; anal. HPLC: $R_t = 7.27$ min (method 1). Diast. ratio: 4:1.

25 **Example 40**

4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-(2-methoxy-ethyl)-benzenesulfonamide

(a). 4-bromomethyl-benzenesulfonyl chloride

- At 20 °C benzoyl peroxide (600 mg) was added to a suspension of 4-methyl-benzenesulfonyl chloride (9.5 g) and N-bromosuccinimide (8.9 g) in 1,2-dichloropropane (60 ml). The reaction mixture was stirred at 80 °C for 2 h. The reaction mixture was concentrated and the title compound was obtained as a white solid after recrystallization from heptane. Yield: 2.26 g.

(b). 4-Bromomethyl-N-(2-methoxy-ethyl)-benzenesulfonamide

At 20 °C triethylamine (61 mg) was added to a solution of the product of example 40a (135 mg) in diethyl ether (2 ml). After 5 minutes, 2-methoxy-ethylamine (37 mg) was added. The reaction mixture was stirred at 20 °C for 3h and concentrated *in vacuo*. The residue was dissolved in dichloromethane (20 ml) and washed several times with sat. NaHCO₃. The organic phase was separated, dried (MgSO₄) and concentrated *in vacuo*. The title product was obtained as brown oil. Yield: 120 mg. MS-ESI: [M+H]⁺ = 308/310.

(c). 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-(2-methoxy-ethyl)-benzenesulfonamide

At 20 °C a solution of the product of example 40b (120 mg) in DMF (3 ml) was added to a suspension of the product of 30a (222 mg), K₂CO₃ (200 mg) and KI (10 mg) in DMF (7 ml). The reaction mixture was stirred at 20 °C for 16 h and poured into 10 ml sat. NaHCO₃ and extracted with ethyl acetate. The organic layer was concentrated, dissolved in dichloromethane and washed with water, dried (MgSO₄) and concentrated *in vacuo*. The title product was obtained via preparative HPLC (0→90% CH₃CN, 0.1% TFA). Yield: 22 mg. MS-ESI: [M+H]⁺ = 672/674; anal. HPLC: Rt = 21.31 min (method 2). Diast. ratio: 4:1

Example 41

20 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N,N-diethyl-benzenesulfonamide

(a). 4-Bromomethyl-N,N-diethyl-benzenesulfonamide

The title compound was obtained analogously to example 40 b, starting from diethylamine (37 mg) and the product of example 40a (135 mg). Yield: 122 mg. R_f (heptane/ethyl acetate (1/1, v/v)) = 0.45. MS-ESI: [M+H]⁺ = 306/308.

(b). 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N,N-diethyl-benzenesulfonamide

The title compound was obtained analogously to example 40 c, starting from product of example 41a (122 mg) and the product of example 30a (222mg). Yield: 79 mg. MS-ESI: [M+H]⁺ = 670/672; anal. HPLC: Rt= 26.63 min (diast1), 26.93 min (diast2) (method 2). Diast. ratio: 4:1

Example 42

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-methanesulfonamide

- Sulfonylation of the product of example 1c (100 mg) with methanesulfonyl chloride (15 μ l) was preformed according to the method described in example 28. Yield: 46 mg. MS-ESI: $[M+H]^+ = 628/630$; anal. HPLC: $R_t = 20.76$ min (diast1), $R_t = 21.01$ min (diast2) (method 2). Diast. ratio: 4:1

Example 43

- Thiophene-2-sulfonic acid {3-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide

Sulfonylation of the product of example 1c (100 mg) with thiophene-2-sulfonyl chloride (36 μ l) was preformed according to the method described in example 28. Yield: 70 mg. MS-ESI: $[M+H]^+ = 696.2/698.2$; anal. HPLC: $R_t = 23.75$ min (method 2). Diast. ratio: 4:1

Example 44

4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzenesulfonic acid isopropyl ester

(a). Toluene-4-sulfonic acid isopropyl ester

- At 0 °C, 4-methyl-benzenesulfonyl chloride (3.8 g) was added to a solution of 2-propanol (6.12 ml) in pyridine (6.6 ml). The reaction mixture was stirred at 0 °C for 2 h and then at 20 °C for 16 h. The reaction mixture was poured into water (150 ml) and extracted with dichloromethane. The combined organic phases were washed with 3M aqueous HCl and sat. NaHCO_3 . After drying (MgSO_4) and concentrating *in vacuo*, the title product was obtained as a pale yellow oil. Yield: 2.89 g. MS-ESI: $[M+H]^+ = 215$.

- (b). 4-Bromomethyl-benzenesulfonic acid isopropyl ester

- At 20 °C benzoyl peroxide (48 mg) was added to a solution of the product of example 44a (956 mg) and *N*-bromosuccinimide (712 mg) in deuteriochloroform (5 ml). The reaction mixture was stirred at 80 °C for 3 h. The reaction mixture was poured into 20 ml of sat. NaHCO_3 and extracted several times with dichloromethane. The combined organic layers were dried (MgSO_4) and concentrated *in vacuo*. The title product was obtained as pale yellow oil. Yield: 1.06 g.

(c). 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzenesulfonic acid isopropyl ester

The title compound was obtained analogously to example 40c, starting from the product of example 44b (293 mg) and product of example 30a (400 mg). Yield: 38 mg. MS-ESI: $[M+H]^+$ = 657/659; anal. HPLC: R_t = 22.64 min (diast1), R_t = 22.64 min (diast 2) (method 2). Diast. ratio: 10:1

5 **Example 45**

3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-methyl-benzenesulfonamide

(a). 3 - bromomethyl-benzenesulfonyl chloride

The title compound was obtained analogously to example 40a, starting from 3-methyl-
10 benzenesulfonyl chloride (8.5 g), *N*-bromosuccinimide (8.9 g) and benzoylperoxide (600 mg) in deuteriochloroform (20 ml) as solvent. Yield: 3.1 g.

(b). 3-Bromomethyl-N-methyl-benzenesulfonamide

The title compound was obtained analogously to example 40b, starting from methylamine (105 mg) and the product of example 45a (269 mg) and the product of
15 example 30a (200 mg). Yield: 220 mg. MS-ESI: $[M-Me]^-$ = 250/252

(c). 3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-methyl-benzenesulfonamide

The title compound was obtained analogously to example 40 c, starting from the product of example 45b (220 mg) and the product of example 30a (200 mg). Yield: 17
20 mg. MS-ESI: $[M+H]^+$ = 628/630; anal. HPLC: R_t = 20.7 min (diast1), R_t = 20.7 min (diast2) (method 2). Diast. ratio: 10:1

Example 46

4-{3-Bromo-5-ethoxy-4-[3-(morpholine-4-sulfonyl)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

25 (a). 4-(3-Bromomethyl-benzenesulfonyl)-morpholine

The title compound was obtained analogously to example 40 b, starting from morpholine (137 μ l), and the product of example 45a (269 mg). Yield: 280 mg.

(b). 4-{3-Bromo-5-ethoxy-4-[3-(morpholine-4-sulfonyl)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

30 The title compound was obtained analogously to example 40c, starting from the product of example 46a (280 mg) and the product of example 30a (200 mg). Yield: 63

mg. MS-ESI: $[M+H]^+ = 684/686$; anal. HPLC: $R_t = 22.84$ min (method 2). Diast. ratio: 10:1

Example 47

5 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzenesulfonic acid

At 20 °C a solution of KI (91 mg) in water (0.5 ml) was added to a solution of the product of example 44c (37 mg) in acetone (0.5 ml). The reaction mixture was stirred at 60 °C for 16 h and poured into water (15 ml). After evaporation of acetone, the reaction mixture was extracted several times with dichloromethane. The organic phases
10 were combined dried ($MgSO_4$) and concentrated. The title product was obtained via preparative HPLC (0→90% CH_3CN , 1% TFA). Yield: 7 mg. MS-ESI: $[M+H]^+ = 617/615$; anal. HPLC: $R_t = 14.31$ min (diast1), $R_t = 14.65$ min (diast2) (method 2). Diast. ratio: 6:4 (HPLC)

Example 48

15 N-{4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-methanesulfonamide

(a). (4-Hydroxymethyl-phenyl)-carbamic acid 9H-fluoren-9-ylmethyl ester

A mixture of (4-amino-phenyl)-methanol (500 mg), Fmoc-Cl (1.2 g) and pyridine (1 ml) in THF (15 ml) was stirred for 1 h. The reaction mixture was diluted with ethyl
20 acetate and washed with water. The organic layer was separated, dried ($MgSO_4$), filtered and concentrated *in vacuo*. The residue was recrystallized from dichloromethane. Yield: 470 mg. MS-ESI: $[M+H]^+ = 346$.

(b). (4-Chloromethyl-phenyl)-carbamic acid 9H-fluoren-9-ylmethyl ester

Thionyl chloride (1.03 ml) in dichloromethane (5 ml) was added dropwise to a solution
25 of the product of example 48a (460 mg) in dichloromethane (10 ml). After stirring for 2 h, the reaction mixture was concentrated *in vacuo*. Yield: 476 mg. $R_f = 0.75$ (heptane/ethyl acetate = 1/2 (v/v))

(c). {4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid 9H-fluoren-9-ylmethyl ester
30

A mixture of the product of example 48b (100 mg), the product of example 30a (122 mg) and K_2CO_3 (114 mg) in DMF (5 ml) was stirred for 4 h. The reaction mixture was diluted with ethyl acetate and washed with water and brine. The organic layer was

separated, dried (MgSO₄), filtered and concentrated *in vacuo*. Yield: 233 mg MS-ESI: [M+H]⁺ = 772.4 / 774.4

(d). 4-[4-(4-Amino-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- 5 Piperidine (335 µl) in dichloromethane (5 ml) was added dropwise to a solution of the product of example 48c (223 mg) in dichloromethane (5 ml). After stirring for 2 h, the reaction mixture was concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 1/4 (v/v) as eluent. Yield: 55 mg. MS-ESI: [M+H]⁺ = 550.2/552.2

- 10 (e). N-{4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-methanesulfonamide

Methanesulfonyl chloride (8 µl) and pyridine (21 µl) were added to a solution of the product of example 48d (50 mg) in dichloromethane (2 ml). After stirring for 2 h, the reaction mixture was diluted with ethyl acetate and washed with water. The organic
15 layer was separated, dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (10 → 100% acetonitrile). Yield: 11 mg. MS-ESI: [M+H]⁺ = 628/630; anal. HPLC: Rt = 19.78 min. Diast. ratio: 8:1

Example 49

- 20 N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-methoxy-phenoxy-methyl]-phenyl}-methanesulfonamide

(a). 4-(3-Bromo-4-hydroxy-5-methoxy-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was obtained analogously to example 1b, starting from 3-Bromo-4-hydroxy-5-methoxy-benzaldehyde (3.0 g). 3-aminocrotonitrile (1.08 g) and 5-propylcyclohexane-1,3-dione (1.98 g). Yield: 4.8 g. MS-ESI: [M+H]⁺ = 431/433.
25

(b). 4-[3-Bromo-5-methoxy-4-(2-nitro-benzyloxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was obtained analogously to example 30b, starting from 1-bromomethyl-2-nitro-benzene (200 mg) and the product of example 49a (400 mg).
30 Yield: 384 mg. MS-ESI: [M+H]⁺ = 566/568; anal. HPLC: Rt = 24.50 min (diast1), Rt = 24.95 min (diast2) (method 2). Diast. ratio: 10:1

(c). 4-[4-(2-Amino-benzyloxy)-3-bromo-5-methoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was obtained analogously to example 19c, starting from the product of example 49b (300 mg). Yield: 289 mg. MS-ESI: $[M+H]^+ = 536/538$; anal. HPLC: $R_t = 17.45$ min (diast1), $R_t = 17.91$ min (diast2) (method 2). Diast. ratio: 10:1

- 5 (d). N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-methoxy-phenoxy-methyl]-phenyl}-methanesulfonamide

The title compound was obtained analogously to example 28, starting from the product of example 49c (100 mg) and methanesulfonyl chloride (20 μ l). Yield: 61 mg. MS-ESI: $[M+H]^+ = 614/616$; anal. HPLC: $R_t = 22.20$ min (method 2). Diast. ratio: 15:1

Example 50

- 10 {4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid allyl ester

- (a). (4-Hydroxymethyl-phenyl)-carbamic acid allyl ester

- Alloc-Cl (537 g) in dichloromethane (5 ml) was added dropwise to a solution of (4-amino-phenyl)-methanol (500 mg) and pyridine (5 ml) in dichloromethane (5 ml).
15 After stirring for 2 h, the reaction mixture was diluted with dichloromethane and washed with water. The organic layer was separated, dried ($MgSO_4$), filtered and concentrated *in vacuo*. The title compound (243 mg) was obtained by recrystallization from dichloromethane. The mother liquor was chromatographed on silica gel in heptane/ethyl acetate = 1/1 (v/v) as eluent which yielded another 245 mg of the title
20 compound. Yield: 488 mg.

- (b). (4-Chloromethyl-phenyl)-carbamic acid allyl ester

The title compound was prepared analogously to example 48b starting from the product of example 50a (480 mg) and thionyl chloride (1.7 ml). The residue was chromatographed on silica gel in heptane/ethyl acetate = 1/1 (v/v) as eluent. Yield: 310 mg.

- 25 (c). {4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid allyl ester

The title compound was prepared analogously to example 48c starting from the product of example 50b (300 mg) and the product of example 30a (593 mg). Yield: 333 mg. MS-ESI: $[M+H]^+ = 634.4/636.4$; anal. HPLC: $R_t = 24.99$ min (method 2).

30 Example 51

4-[3-Bromo-5-ethoxy-4-(1-methanesulfonyl-1H-pyrrol-2-ylmethoxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- (a). 1-Methanesulfonyl-1H-pyrrole-2-carbaldehyde

A solution of pyrrole-2-carboxyaldehyde (500 mg) in THF (5 ml) was added to a suspension of NaH (252 mg, 60% dispersion on oil) in THF (15 ml). After stirring for 15 min, a solution of methanesulfonyl chloride (570 μ l) in THF (5 ml) was added dropwise. After stirring for 1 h, water (25 ml) was added and the THF was evaporated.

- 5 The residue was diluted with dichloromethane and washed with water, sat. NaHCO_3 and brine. The organic layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 1/1 (v/v) as eluent. Yield: 265 mg.

(b). (1-Methanesulfonyl-1H-pyrrol-2-yl)-methanol

- 10 At 0°C LiBH_4 (1.15 ml, 2.0M in THF) was added dropwise to a solution of the product of example 51a (265 mg) in diethylether (10 ml). After stirring at 0°C for 30 min, water (2 ml) and acetic acid (2 ml, 10% in water) were added. The water layer was separated and extracted with dichloromethane. The organic layer was washed with water and brine, separated, dried (MgSO_4), filtered and concentrated *in vacuo*. Yield: 205 mg.

- 15 (c). 2-Chloromethyl-1-methanesulfonyl-1H-pyrrole

- At 0°C methanesulfonyl chloride (136 μ l) was added dropwise to a solution of the product of example 51b (205 mg) and DIPEA (307 μ l) in dichloromethane (6 ml). After stirring for 30 min, the reaction mixture was diluted with dichloromethane (30 ml) and washed with ice water (30 ml), 10% HCl (30 ml) and sat. NaHCO_3 . The organic
20 layer was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. Yield: 252 mg.

(d). 4-[3-Bromo-5-ethoxy-4-(1-methanesulfonyl-1H-pyrrol-2-ylmethoxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- Alkylation of the product of example 30a (539 mg) with the product of example 51c (252 mg) was performed according to the method described in example 1a. The residue
25 was purified by preparative HPLC (40 $\rightarrow$ 100% acetonitrile). Yield: 363 mg. MS-ESI: $[\text{M}+\text{H}]^+ = 602.4/604.4$; anal. HPLC: $R_t = 15.53$ min (diast1), $R_t = 15.99$ min (diast2) (method 1). Diast. ratio: 11:1

Example 52

- N-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenyl]-2-nitro-benzamide
30

(a). 3-Ethoxy-4-nitro-benzaldehyde

At 20 °C iodoethane (3.78 g) was added to a suspension of K_2CO_3 (1.09 g) and 3-hydroxy-4-nitro-benzaldehyde (1.0 g) in DMF (5 ml). The reaction mixture was stirred at 70 °C for 18 h, poured into water and then extracted with ethyl acetate. The organic

phases were combined, dried (MgSO₄) and concentrated *in vacuo*. The title product was obtained as a yellow solid. Yield: 1.17 g. MS-ESI: [M+H]⁺ = 196.

(b). 4-(3-Ethoxy-4-nitro-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- 5 The title compound was obtained analogously to example 1b, starting from the product of example 52a (1.16 g), 3-aminocrotonitrile (502 mg) and 5-propylcyclohexane-1,3-dione (920mg). Yield: 1.24 g. MS-ESI: [M+H]⁺ = 396.

(c). 4-(4-Amino-3-ethoxy-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- 10 The title compound was obtained analogously to example 19c starting from the product of example 52b (1.2 g). Yield: 1.1 g. MS-ESI: [M+H]⁺ = 366.

(d). 4-(4-Amino-3-bromo-5-ethoxy-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- At 0 °C a solution of bromine (56 µl) in dichloromethane (5 ml) was added slowly to a mixture of the product example 52c (365 mg) and sodium acetate (89 mg) in acetic acid (10 ml) and dichloromethane (5 ml). The reaction mixture was stirred at 0 °C for 1 h, poured into water and then extracted with dichloromethane. The organic phases were combined, dried (MgSO₄) and concentrated *in vacuo*. The title compound was obtained after flash column chromatography (silica gel, heptane/ethyl acetate (1/4, v/v), R_f = 0.55). Yield: 1.17 g. MS-ESI: [M+H]⁺ = 444/446.

(e). N-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenyl]-2-nitro-benzamide

- At 20 °C, 2-nitro-benzoyl chloride (125 mg) was slowly added to a solution of the product of example 52d (200 mg) and *N,N*-dimethylaniline (172 µl) in THF (5 ml). The reaction mixture was stirred at 20 °C until total conversion and then poured in sat. NaHCO₃ and extracted with ethyl acetate. The organic phases were combined, dried (MgSO₄) and concentrated *in vacuo*. The title compound was obtained by recrystallization from acetonitrile. Yield: 37 mg. MS-ESI: [M+H]⁺ = 593/595; anal. HPLC: R_t = 17.98 min (method 2).

30 **Example 53**

4-[3-Bromo-5-ethoxy-4-(1-trifluoromethanesulfonyl-1H-pyrrol-2-ylmethoxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

(a). 1-Trifluoromethanesulfonyl-1H-pyrrole-2-carbaldehyde

At -78°C and under a nitrogen atmosphere, trifluoromethanesulfonic anhydride (3.2 ml) was added dropwise to a solution of pyrrole-2-carboxyaldehyde (1 g) and DIPEA (3.67 ml) in dichloromethane (100 ml). After stirring for 5 min at -78°C, the reaction mixture was poured into sat. NaHCO₃ (200 ml) and extracted with dichloromethane. The organic layer was separated, washed with water, dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 2/1 (v/v) as eluent. Yield: 675 mg.

(b). (1-Trifluoromethanesulfonyl-1H-pyrrol-2-yl)-methanol

The title compound was obtained analogously to example 51b starting from the product of example 53a (300 mg). Yield: 361 mg

(c). 2-Chloromethyl-1-trifluoromethanesulfonyl-1H-pyrrole

The title compound was obtained analogously to example 51c starting from the product of example 53b (361 mg). Yield: 391 mg

(d). 3-Bromo-5-ethoxy-4-(1-trifluoromethanesulfonyl-1H-pyrrol-2-ylmethoxy)-benzaldehyde

Alkylation of 3-bromo-5-ethoxy-4-hydroxy benzaldehyde (387 mg) with the product of example 53c (391 mg) was performed according to the method described in example 1a. The residue was chromatographed on silica gel in heptane/ethyl acetate = 3/1 (v/v) as eluent. Yield: 208 mg. MS-ESI: [M+H]⁺ = 456.2/458.2

(e). 4-[3-Bromo-5-ethoxy-4-(1-trifluoromethanesulfonyl-1H-pyrrol-2-ylmethoxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was obtained analogously to example 1b starting from the product of example 53d (208 mg). The residue was purified by preparative HPLC (20 → 100% acetonitrile). Yield: 166 mg. MS-ESI: [M+H]⁺ = 656.2/658.2; anal. HPLC: Rt = 23.99 min (diast1), Rt = 24.47 (diast.2) (method 1). Diast. ratio: 5:1.

Example 54

4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-methyl-benzamidine

(a). 4-[3-Bromo-4-(4-cyano-benzyloxy)-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

Alkylation of the product of example 30a (2.5 g) with α-bromo-*p*-tolunitrile (1.21 g) was performed according to the method described in example 30b. The reaction mixture was poured in water and filtered over decalite. The residue was washed with

water en dichloromethane. The organic layer was separated, washed with water and brine, dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 3/1 (v/v) as eluent. Yield: 2.6 g. MS-ESI: [M+H]⁺ = 560 /562

- 5 (b). 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzimidic acid ethyl ester

HCl (g) was bubbled through a solution of the product of example 54a (1.5 g) in ethanol (10 ml) for 2 h. After additional stirring for 2 h, the reaction mixture was concentrated *in vacuo*. Yield: 1.62 g.

- 10 (c). 4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-N-methyl-benzamidine

A mixture of the product of example 54b (180 mg), methylamine (126 µl, 8 M in EtOH) and triethylamine (126 µl) in ethanol (1 ml) was stirred for 54 h. The reaction mixture was concentrated *in vacuo*. The residue was purified by preparative HPLC (10
15 → 90% acetonitrile. 0.1% TFA). Yield: 56.5 mg (as TFA salt). MS-ESI: [M+H]⁺ = 591/593; anal. HPLC: Rt = 17.8 min (method 4).

Example 55

2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-pyrrole-1-carboxylic acid tert-butyl ester

- 20 (a). 2-Formyl-pyrrole-1-carboxylic acid tert-butyl ester

Bocylation of pyrrole-2-carboxyaldehyde (750 mg) in the presence of NaH (410 mg, 60% dispersion on oil) and Boc-On (2.72 g) was performed according to the method described in example 51a. The residue was chromatographed on silica gel in heptane/ethyl acetate = 2/1 (v/v) as eluent. Yield: 196.2 mg.

- 25 (b). 2-Hydroxymethyl-pyrrole-1-carboxylic acid tert-butyl ester

Reduction of the product of example 55a (500 mg) with LiBH₄ was performed according to the method described in example 51b. The residue was chromatographed on silica gel in heptane/ethyl acetate = 2/1 (v/v) as eluent. Yield: 350 mg.

(c). 2-Methanesulfonyloxymethyl-pyrrole-1-carboxylic acid tert-butyl ester

- 30 Sulfonylation of the product of example 55b (350 mg) with methanesulfonyl chloride (210 µl) was performed according to the method described in example 51c. The reaction mixture was washed with water, sat. NaHCO₃ and brine. The organic layer was separated, dried (MgSO₄), filtered and concentrated *in vacuo*. Yield: 487 mg.

(d). 2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-pyrrole-1-carboxylic acid *tert*-butyl ester

Alkylation of the product of example 30a (787 mg) with the product of example 55c (487 mg) was performed according to the method described in example 1a. The residue
5 was purified by preparative HPLC (20 → 100% acetonitrile). Yield: 61 mg. MS-ESI: $[M+H]^+ = 624.2/626.2$; anal. HPLC: $R_t = 29.35$ min (diast1), $R_t = 29.70$ min (diast2) (method 2). Diast. ratio: 4:1.

Example 56

4-{3-Bromo-5-ethoxy-4-[(pyridin-3-ylmethyl)-amino]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile
10

At 20 °C pyridine-3-carbaldehyde (210 µl) was added to a solution of the product of example 52d (100 mg) and acetic acid (127 µl) in methanol (4 ml). The reaction mixture was stirred at 20 °C for 16 h and then NaCNBH_4 (142 mg) was added. The reaction was stirred at 20 °C for 24 h, poured in water and extracted with ethyl acetate.
15 The title product was obtained via preparative HPLC (0→90% CH_3CN , 0.1% TFA). Yield: 27 mg (as TFA salt). MS-ESI: $[M+H]^+ = 535/537$; anal. HPLC: $R_t = 8.56$ min (method 2).

Example 57

4-[3-Bromo-5-ethoxy-4-(2-nitro-benzylamino)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile
20

The title compound was obtained analogously to example 56, starting from 2-nitro-benzaldehyde (1.93 g) and the product of example 52d (570 mg). The residue was purified by preparative HPLC (0→90% CH_3CN). Yield: 610 mg. MS-ESI: $[M+H]^+ = 579/581$; HPLC: $R_t = 24.35$ min (diast1), $R_t = 24.85$ min (diast2) (method 2).
25 Diast. ratio: 9:1 (HPLC)

Example 58

N-(2-{[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenylamino]-methyl}-phenyl)-methanesulfonamide
(a). 4-[4-(2-Amino-benzylamino)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile
30

The title compound was obtained analogously to example 19c, starting from the product of example 57 (550 mg). Yield: 510 mg. $[M+H]^+ = 549/551$.

(b). N-(2-{[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenylamino]-methyl}-phenyl)-methanesulfonamide

The title compound was obtained analogously to example 28, starting from the product of example 58a (130 mg) and methanesulfonyl chloride (20 μ l). Yield: 31 mg. MS-ESI: $[M+H]^+ = 627/629$; anal. HPLC: $R_t = 20.82$ min (diast1), $R_t = 21.35$ min (diast2) (method 2). Diast. ratio: 4:1 (HPLC)

Example 59

(3-{[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenylamino]-methyl}-phenyl)-carbamic acid methyl ester

(a). 4-[3-Bromo-5-ethoxy-4-(3-nitro-benzylamino)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was obtained analogously to example 56, starting from 3-nitro-benzaldehyde (1.93 g) and the product of example 52d (570 mg). Yield: 630 mg. MS-ESI: $[M+H]^+ = 579/581$; anal. HPLC: $R_t = 23.91$ min (diast1), $R_t = 24.32$ min (diast2) (method 2). Diast. ratio: 6:1 (HPLC)

(b). 4-[4-(3-Amino-benzylamino)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was obtained analogously to example 19c, starting from the product of example 59a (570 mg). Yield: 510 mg. $[M+H]^+ = 549/551$.

(c). (3-{[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenylamino]-methyl}-phenyl)-carbamic acid methyl ester

The title compound was obtained analogously to example 25, starting from the product of example 59b (130 mg) and methylchloroformate (24 μ l). Yield: 29 mg. MS-ESI: $[M+H]^+ = 607/609$; anal. HPLC: $R_t = 17.89$ min (method 2).

Example 60

2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzamidine; compound with hydrochloric acid

(a). 4-[3-Bromo-4-(2-cyano-benzyloxy)-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

Alkylation of the product of example 30a (2.5 g) with α -bromo-*o*-tolunitrile (1.21 g) was performed according to the method described in example 54a. Yield: 2.44 g. MS-ESI: $[M+H]^+ = 560/562$

(b.) 2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzamidine; compound with hydrochloric acid

HCl (g) was bubbled through a solution of the product of example 60a (200 mg) in ethanol (1 ml) for 1 h. After additional stirring for 2 weeks, the reaction mixture was concentrated *in vacuo*. The residue was dissolved in ethanol and NH₄OAc (83 mg) was added. After stirring for 18 h, the reaction mixture was concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 1/1 (v/v) as eluent. Yield: 17.4 mg (HCl-salt). MS-ESI: [M+H]⁺ = 577/579; anal. HPLC: Rt = 16.8 min (diast1) (method 4).

10 Example 61

4-{3-Bromo-5-ethoxy-4-[3-(imino-morpholin-4-yl-methyl)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

(a.) 4-[3-Bromo-4-(3-cyano-benzyloxy)-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

15 Alkylation of the product of example 30a (2.5 g) with α -bromo-*m*-tolunitrile (1.21 g) was performed according to the method described in example 54a. Yield: 2.6 g. MS-ESI: [M+H]⁺ = 560 /562

(b.) 3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-benzimidic acid ethyl ester

20 The title compound was prepared analogously to example 54b, starting from the product of example 61a (1.5 g). Yield: 1.62 g.

(c.) 4-{3-Bromo-5-ethoxy-4-[3-(imino-morpholin-4-yl-methyl)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was prepared analogously to example 54c, starting from the product of example 61b (180 mg) and morpholine (77.8 mg). Yield: 86.7 mg (as TFA salt). MS-ESI: [M+H]⁺ = 647/649; anal. HPLC: Rt = 17.7 min (method 4).

Example 62

4-{3-Bromo-4-[2-(cyclopropylmethyl-amino)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

30 (a.) N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2,4-dinitro-benzenesulfonamide

2,4-Dinitro-benzenesulfonyl chloride (3.2 g) and pyridine (3.8 ml) were added to a solution of the product of example 19c (5 g) in dichloromethane (30 ml). After stirring

for 3 h, the reaction mixture was poured in water and acidified with 2N HCl. The organic layer was separated and washed with sat. NaHCO₃ and brine. The organic layer was separated, dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 1/2 (v/v) as eluent. Yield: 5 g.

5 MS-ESI: $[M+H]^+ = 780.2/782.2$

(b). *N*-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-*N*-cyclopropylmethyl-2,4-dinitro-benzenesulfonamide

Diisopropylazodicarboxylate (102 μ l) was added dropwise to a solution of the product of example 62a (200 mg), cyclopropanemethanol (38.2 mg) and triphenylphosphine (134 mg) in THF (6 ml). After stirring for 10 min, the reaction mixture was concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 1/2 (v/v) as eluent. Yield: 158 mg. ESI-MS: $[M+H]^+ = 834.4/836.4$

(c). 4-{3-Bromo-4-[2-(cyclopropylmethyl-amino)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

A mixture of the product of example 62b (158 mg) and propylamine (200 μ l) in dichloromethane (10 ml) was stirred for 1 h. The reaction mixture was concentrated *in vacuo* and the residue was purified by preparative HPLC (10 $\rightarrow$ 90% CH₃CN). Yield: 72 mg. MS-ESI: $[M+H]^+ = 604.4/606.4$; anal. HPLC: Rt = 21.8 min (diast1), Rt = 22.39 min (diast2) (method 1). Diast. ratio: 6:1

Example 63

4-{3-Bromo-5-ethoxy-4-[2-(3-methyl-butylamino)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

(a). *N*-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-*N*-(3-methyl-butyl)-2,4-dinitro-benzenesulfonamide

The title compound was obtained analogously to example 62b, starting from the product of example 62a (200 mg) and isoamylalcohol (56 μ l). Yield: 205 mg. MS-ESI: $[M+H]^+ = 850.4/852.4$

(b). 4-{3-Bromo-5-ethoxy-4-[2-(3-methyl-butylamino)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

The title compound was prepared analogously to example 62c, starting from the product of compound 63a (205 mg). Yield: 66.7 mg. MS-ESI: $[M+H]^+ = 620.4/622.4$;

anal. HPLC: R_t = 23.61 min (diast1), R_t = 24.05 min (diast2) (method 5). Diast. ratio: 5:1

Example 64

5 {2-[2-Bromo-6-ethoxy-4-(2-methyl-3-nitro-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-phenoxy-methyl]-phenyl}-carbamic acid methyl ester

(a). 1-Nitro-propan-2-one

At 0 °C sodium hydride (60 % dispersion on oil) (1.28 g) was added to a solution of nitromethane (1.95 g) in THF (25 ml). The reaction mixture was stirred at 20°C for 30 minutes and then added to a solution of 1-pyrazol-1-yl-ethanone (2.72 g) in THF (25 ml). The reaction mixture was stirred at 60 °C for 18 h. The solid formed was filtered,
10 dissolved in water (50 ml), acidified to pH=3 with 1 M HCl and then extracted several times with ethyl acetate. The organic phases were combined dried (MgSO₄) and concentrated. The title product was obtained as a yellow oil. Yield: 2.47 g.

(b). 1-Methyl-2-nitro-vinylamine

15 A mixture of the product of example 64a (1.6 g) and NH₄OAc (1.3 g) in toluene (25 ml) was heated under reflux for 18 h with azeotropic removal of water using a Dean-Stark apparatus. The reaction mixture was concentrated *in vacuo* and chromatographed on silica gel in heptane/ethyl acetate = 2/1 as eluent. Yield: 10.7 g

(c). (2-Hydroxymethyl-phenyl)-carbamic acid methyl ester

20 Reaction of aminobenzylalcohol (3 g) with methylchloroformate (1.9 ml) in the presence of DIPEA (12.8 ml) was performed according to the method described in example 22. The residue was chromatographed on silica gel in dichloromethane/ethyl acetate = 25/1 as eluent. Yield: 4.19 g.

(d). (2-Chloromethyl-phenyl)-carbamic acid methyl ester

25 The title compound was prepared analogously to example 48b starting from the product of example 64c (4.19 g) and thionyl chloride (10 ml). Yield: 1.98 g.

(e). [2-(2-Bromo-6-ethoxy-4-formyl-phenoxy-methyl)-phenyl]-carbamic acid methyl ester

30 Alkylation of 3-bromo-5-ethoxy-4-hydroxy benzaldehyde (614 mg) with the product of example 64d (500 mg) was performed according to the method described in example 1b. Yield: 845 mg.

(f). {2-[2-Bromo-6-ethoxy-4-(2-methyl-3-nitro-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-phenoxy-methyl]-phenyl}-carbamic acid methyl ester

Reaction of the product of example 64e (100 mg) with the product of example 64b (28 mg) and 5-propylcyclohexane-1,3-dione (41 g) was performed according to the method described in example 1b. The residue was purified by preparative HPLC (10 → 90% acetonitrile). Yield: 75 mg. MS-ESI: $[M+H]^+ = 628/630$; anal. HPLC: $R_t = 26.22$ min (diast1), $R_t = 26.67$ min (diast2) (method 2). Diast. ratio: 4:1

Example 65

4-[3-Bromo-5-ethoxy-4-(3-methoxy-benzyloxy)-phenyl]-2-methyl-3-nitro-7-propyl-4,6,7,8-tetrahydro-1H-quinolin-5-one

(a). 4-(3-Bromo-5-ethoxy-4-hydroxy-phenyl)-2-methyl-3-nitro-7-propyl-4,6,7,8-tetrahydro-1H-quinolin-5-one

Reaction of the product of example 64b (1 g) with 3-bromo-5-ethoxy-4-hydroxy benzaldehyde (2.57 g) and 5-propylcyclohexane-1,3-dione (1.61 g) was performed according to the method described in example 1b. The residue was recrystallized from dichloromethane. Yield: 2.8 g. MS-ESI: $[M+H]^+ = 465/467$

(b). 4-[3-Bromo-5-ethoxy-4-(3-methoxy-benzyloxy)-phenyl]-2-methyl-3-nitro-7-propyl-4,6,7,8-tetrahydro-1H-quinolin-5-one

The title compound was obtained analogously to example 30b, starting from the product of example 65a (100 mg) and 1-bromomethyl-3-methoxy-benzene (50 mg). Yield: 36 mg. MS-ESI: $[M+H]^+ = 554/556$; anal. HPLC: $R_t = 27.18$ min (diast1), $R_t = 27.61$ min (diast2) (method 2). Diast. ratio: 1:1

Example 66

4-[3-Bromo-5-ethoxy-4-(2-methyl-thiazol-4-ylmethoxy)-phenyl]-2-methyl-3-nitro-7-propyl-4,6,7,8-tetrahydro-1H-quinolin-5-one

The title compound was obtained analogously to example 30b, starting from 4-chloromethyl-2-methyl-thiazole (37 mg) and the product of example 65a (100 mg). Yield: 27 mg. MS-ESI: $[M+H]^+ = 574/576$; anal. HPLC: $R_t = 22.01$ min (diast), $R_t = 22.50$ min (diast2) (method 2). Diast. ratio: 1:1

Example 67

{4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-acetic acid methyl ester

Alkylation of the product of example 30a (196 mg) with (4-bromomethyl-phenyl)-acetic acid methyl ester (107 mg) was performed according to the method described in

example 30b. The residue was purified by preparative HPLC (10→90% acetonitrile). Yield: 116 mg. MS-ESI: $[M+H]^+ = 607/609$; anal. HPLC: $R_t = 28.41$ min (method 4).

Example 68

5 {4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-acetic acid

(a). {4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-acetic acid methyl ester

Alkylation of the product of example 30a (445 mg) with (4-bromomethyl-phenoxy)-acetic acid methyl ester (259 mg) was performed according to the method described in
10 example 67. Yield: 308 mg. MS-ESI: $[M+H]^+ = 623/625$; anal. HPLC: $R_t = 23.67$ min (method 2).

(b). {4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-acetic acid

Saponification of the product of example 68a (230 mg) was performed according to the
15 method described in example 6c. Yield: 166 mg. MS-ESI: $[M+H]^+ = 609/611$; anal. HPLC: $R_t = 19.32$ min.

Example 69

N-(3-{[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenylamino]-methyl}-pyridin-2-yl)-methanesulfonamide

20 (a). N-(3-Cyano-pyridin-2-yl)-methanesulfonamide

A mixture of 2-chloro-nicotinonitrile (2 g), H_2NSO_2Me (1.73 g) and K_2CO_3 (4.44 g) in DMF (100 ml) was stirred at 100°C for 18 h. The reaction mixture was concentrated *in vacuo*. The residue was dissolved in ethyl acetate and washed with water and brine. The organic layer was separated, dried ($MgSO_4$), filtered and concentrated *in vacuo*.
25 The residue was chromatographed on silica gel in dichloromethane/methanol = 98/2 (v/v) as eluent. Yield: 1.32 g. MS-ESI: $[M+H]^+ = 198.2$

(b). N-(3-Formyl-pyridin-2-yl)-methanesulfonamide

Raney nickel (2.64 ml, 50% suspension in water) was added to a solution of the product of example 69a (1.32 g) in formic acid (55 ml). After stirring at 100°C for 3 h,
30 the reaction mixture was filtered over decalite and washed with formic acid. After concentration of the filtrate, the residue was dissolved in ethyl acetate and washed with sat. $NaHCO_3$ and brine. The organic layer was separated, dried ($MgSO_4$), filtered and

concentrated *in vacuo*. The residue was chromatographed on silica gel in heptane/ethyl acetate = 2/1 (v/v) as eluent. Yield: 461 mg. MS-ESI: $[M+H]^+ = 201.2$

(c). N-(3-([2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenylamino]-methyl)-pyridin-2-yl)-methanesulfonamide

- 5 A mixture of the product of example 52d (200 mg), the product of example 69b (99 mg) and acetic acid (129 μ l) in methanol (8 ml) was stirred for 18 h followed by addition of NaCNBH₃ (141 mg). Stirring was continued for another 4 h. The reaction mixture was diluted with ethyl acetate and washed with water, sat. NaHCO₃ and brine. The organic layer was separated, dried (MgSO₄), filtered and concentrated *in vacuo*.
- 10 The residue was purified by preparative HPLC (10 $\rightarrow$ 90% CH₃CN). Yield: 145 mg. MS-ESI: $[M+H]^+ = 628/630$; anal. HPLC: Rt = 10.77 min (diast1), Rt = 11.17 min (diast2) (method 2). Diast. ratio: 9:1.

Example 70

- 15 4-[3-Bromo-4-(2,5-dimethyl-2H-pyrazol-3-ylmethoxy)-5-ethoxy-phenyl]-2-methyl-3-nitro-7-propyl-4,6,7,8-tetrahydro-1H-quinolin-5-one

(a). 5-Chloromethyl-1,3-dimethyl-1H-pyrazole

The title compound was prepared analogously to example 48b starting from (2,5-dimethyl-2H-pyrazol-3-yl)-methanol (778 mg) and thionyl chloride (2.35 ml). Yield: 900 mg.

- 20 (b). 4-[3-Bromo-4-(2,5-dimethyl-2H-pyrazol-3-ylmethoxy)-5-ethoxy-phenyl]-2-methyl-3-nitro-7-propyl-4,6,7,8-tetrahydro-1H-quinolin-5-one

- Alkylation of the product of example 65a (100 mg) with the product of example 70a (48 mg) was performed according to the method described in example 30b. The residue was purified by preparative HPLC (10 $\rightarrow$ 90% acetonitrile). Yield: 77.3 mg. MS-ESI:
- 25 $[M+H]^+ = 573/575$; anal. HPLC: Rt = 20.14 min (diast1), Rt = 20.64 min (diast2) (method 2). Diast. ratio: 4:1

Example 71

- N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-5-trifluoromethyl-phenyl}-methanesulfonamide
- 30 (a). 4-[3-Bromo-5-ethoxy-4-(2-nitro-4-trifluoromethyl-benzyloxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

Alkylation of the product of example 30a (778 mg) with 1-chloromethyl-2-nitro-4-trifluoromethyl-benzene (419 mg) was performed according to the method described in

example 30b. The residue was chromatographed on silica gel in heptane/ethyl acetate = 2/1 (v/v) as eluent. Yield: 493.9 mg. MS-ESI: $[M+H]^+ = 648/650$

(b). 4-[4-(2-Amino-4-trifluoromethyl-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- 5 Reduction of the product of example 71a (431 mg) with zinc dust (1.08 g) was performed according to the method described in example 19c. Yield: 400 mg. MS-ESI: $[M+H]^+ = 618/620$

(c). N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-5-trifluoromethyl-phenyl}-

- 10 di(methanesulfon)amide

A mixture of the product of example 71c (120 mg), methanesulfonyl chloride (23 μ l) and triethylamine (81 μ l) in dichloromethane (2 ml) was stirred at 40°C for 54 h. The reaction mixture was concentrated *in vacuo*. Yield: 150 mg (crude). ESI-MS: $[M+H]^+ = 774/776$

- 15 (d). N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-5-trifluoromethyl-phenyl}-methanesulfonamide

- A mixture of the crude product of example 71c (150 mg) and 2M NaOH (1 ml) in dioxane (2 ml) was stirred for 18 h. The reaction mixture was acidified with 0.5N HCl, diluted with ethyl acetate and washed with sat. NaHCO_3 and brine. The organic layer
20 was separated, dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (10 $\rightarrow$ 90% CH_3CN). Yield: 32.4 mg. MS-ESI: $[M+H]^+ = 696/698$; anal. HPLC: $R_t = 26.7$ min (diast1), $R_t = 27.0$ min (diast2) (method 2). Diast. ratio: 1:1

Example 72

- 25 N-{2-[2-Bromo-4-((4R,7S)-3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-4,5-difluoro-phenyl}-methanesulfonamide

(a). (1S)-N-[1-(4-Methoxy-phenyl)-ethyl]-3-oxo-butylamide

- To a solution of (S)-1-(4-methoxy-phenyl)-ethylamine (25 g) in dichloromethane (60 mL) was added TEA (20 g) and DMPA (200 mg). Then a solution of 4-methylene-oxetan-2-one (16.7 g) in dichloromethane (60 ml) was added over a period of 30
30 minutes and then stirred at 20 °C for 18 h. The mixture was washed with 1M HCl, aq. NaHCO_3 and aq. NaCl. The organic layer was dried (MgSO_4), filtered and concentrated *in vacuo*.

Yield: 39.8 g. MS-ESI: $[M+H]^+ = 236$

(b). (1S)- 3-Amino-but-2-enoic acid [1-(4-methoxy-phenyl)-ethyl]-amide

To a suspension of (S)-N-[1-(4-methoxy-phenyl)-ethyl]-3-oxo-butyramide (10 g) in toluene (800 ml) was added at 20 °C ammonium acetate (10 g) and the reaction was
5 heated for 20 h at 100 °C in a Dean Stark system. The mixture was concentrated in vacuo and the product obtained was used without further purification in the following reaction step.

(c). 4-(3-Bromo-5-ethoxy-4-hydroxy-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carboxylic acid (1S)-[1-(4-methoxy-phenyl)-ethyl]-amide

10 A mixture of (S)- 3-amino-but-2-enoic acid [1-(4-methoxy-phenyl)-ethyl]-amide (10 g), 3-bromo-5-ethoxy-4-hydroxy-benzaldehyde (10.4 g) and 5-propylcyclohexane-1,3-dione (6.5 g) in ethanol (150 ml) was stirred at 80°C for 18 hr. The mixture was concentrated in vacuo. The residue was dissolved in toluene (100 ml) and dichloromethane (50 ml) and evaporated until a precipitated formed. The solid obtained
15 was then crystallized in EtOAc.

Yield: 12.9 g. MS-ESI: $[M+H]^+ = 597.4/599.4$; anal. HPLC $R_t = 16.70$ (diast.1) $R_t = 17.55$ (diast.2) $R_t = 18.81$ (diast.3) $R_t = 19.74$ (diast.4) (method 7). Diast. ratio diast.1:diast2:diast3:diast4 = 10:1:10:1.

(d). (4S, 7S)-4-(3-Bromo-5-ethoxy-4-hydroxy-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carboxylic acid (1S)-[1-(4-methoxy-phenyl)-ethyl]-amide
20

The mixture of stereoisomers obtained in Example 72c was then separated by chromatography on silicagel in heptane/ethyl acetate 2/8 -> 0/1 (v/v) as eluent.

Yield: 2.6 g MS-ESI: $[M+H]^+ = 597.4/599.4$; anal. HPLC $R_t = 18.81$ (diast.3) (method
25 7). Diast. ratio: 0:0:1:0

(e). (4R, 7S)-4-(3-Bromo-5-ethoxy-4-hydroxy-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

A solution of (4S,7S)-4-(3-Bromo-5-ethoxy-4-hydroxy-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carboxylic acid [1-(4-methoxy-phenyl)-ethyl]-amide (Example 72d, 1.43 g) was dissolved in TFA (30 mL) and stirred at 75°C
30 for 1 hr. The TFA was evaporated in vacuo and the residue was then dissolved in dichloromethane (30 ml). Triethylamine (1.04 g) and trifluoroacetic acid anhydride (0.95 g) were added dropwise at 0°C for 15 minutes. The reaction was stirred for 1 hr and the reaction mixture was washed with water (20 ml). The organic layer was dried

(MgSO₄), filtered, concentrated in vacuo and concentrated. The residue was purified by chromatography on silicagel in heptane/ethyl acetate 1/0 → 0/1 (v/v) as eluent.

Yield: 777 mg. MS-ESI: [M+H]⁺ = 445.4/447.4

(f). (4,5-Difluoro-2-nitro-phenyl)-methanol

- 5 To a solution of 4,5-difluoro-2-nitrobenzoic acid (25.9 g) in THF (90 ml), cooled to 0 °C, was carefully added borane-tetrahydrofuran complex (319 ml, 1M in THF). After the addition was complete, the mixture was stirred at 60°C for 3 h. Methanol (150 ml) was added to the mixture at 0°C and after 10 minutes the mixture was concentrated in vacuo. The residue was poured into water and extracted with ethyl acetate. The organic
10 layer was washed with sat. aq. NaCl, dried (MgSO₄), filtered and concentrated in vacuo.

Yield: 24.1 g

(g). (4R,7S)-4-[3-Bromo-4-(4,5-difluoro-2-nitro-benzyloxy)-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- 15 To a mixture of (4,5-difluoro-2-nitro-phenyl)-methanol (Example 72f, 5.1 g), (4R, 7S)-4-(3-Bromo-5-ethoxy-4-hydroxy-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile (Example 72e, 10.0 g) and polymer-supported triphenylphosphine (13.5 g, 3 mmol/g loading) in THF (500 ml), cooled to 0°C, was added diisopropyl azodicarboxylate (DIAD) (5.3 ml). After 1.5 h, the mixture was
20 filtered and the filtrate was concentrated in vacuo. The residue was purified by chromatography on silicagel in heptane/ethyl acetate 4/1 → 2/3 (v/v) as eluent.

Yield: 13.5 g. MS-ESI: [M+H]⁺ = 616.10/618.10

(h). (4R,7S)-4-[4-(2-Amino-4,5-difluoro-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile

- 25 A solution of (4R,7S)-4-[3-Bromo-4-(4,5-difluoro-2-nitro-benzyloxy)-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile (Example 72 g, 10.6 g) and acetic acid (19.6 ml) in THF (400 ml) was cooled to 0°C. Zinc dust (27.6 g) was added in portions under vigorous stirring. The mixture was stirred for 2 h at 0°C and then filtered. The solids were washed with dichloromethane
30 and the combined filtrates were washed with sat. aq. NaHCO₃. The organic layer was dried (MgSO₄), filtered and concentrated in vacuo.

Yield: 10.9 g. MS-ESI: [M+H]⁺ = 586.05/588.05

(i). *N,N*-{2-[2-Bromo-4-((4*R*,7*S*)-3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-4,5-difluoro-phenyl}-bismethanesulfonamide

A solution of (4*R*,7*S*)-4-[4-(2-Amino-4,5-difluoro-benzyloxy)-3-bromo-5-ethoxy-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile (Example 72 h, 10.9 g) and triethylamine (11.9 ml) in dichloromethane (140 ml), cooled to 0°C, was treated with methanesulfonyl chloride (3.3 ml). After stirring for 2 h at 0°C, the mixture was diluted with 0.2N hydrochloric acid and extracted with dichloromethane. The organic layer was washed with sat. aq. NH₄Cl, dried (MgSO₄), filtered and concentrated in vacuo. The residue was purified by chromatography on silicagel in heptane/ethyl acetate 7/3 → 2/5 (v/v) as eluent.

Yield: 11.0g. MS-ESI: [M+H]⁺ = 742.08/744.08

(j). *N*-{2-[2-Bromo-4-((4*R*,7*S*)-3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-4,5-difluoro-phenyl}-methanesulfonamide

N,N-{2-[2-Bromo-4-((4*R*,7*S*)-3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-4,5-difluoro-phenyl}-bismethanesulfonamide (Example 72 i, 11.0 g) was dissolved in dioxane (190 ml) and treated with 2N NaOH (80 ml). After 2 h the mixture was quenched with 2N hydrochloric acid and extracted with ethyl acetate. The organic layer was washed with sat. aq. NH₄Cl, dried (MgSO₄), filtered and concentrated in vacuo. The residue was purified by chromatography on silicagel in toluene/acetone 1/0 → 3/1 (v/v) as eluent.

Yield: 7.98 g. MS-ESI: [M+H]⁺ = 664.06/666.06; anal. HPLC R_t = 17.18 min (method 8)

Chiral purity : 99.2% (levorotatory or -) (method 9)

Example 73

Agonistic activity of compounds at the human FSH receptor expressed in CHO cells


Agonist activity of the compounds at the human FSH receptor was tested in Chinese Hamster Ovary (CHO) cells stably transfected with the human FSH receptor and cotransfected with a cAMP responsive element (CRE)/promotor directing the expression of a firefly luciferase reporter gene. Binding of the compound to the Gs-coupled FSH receptor will result in an increase of cAMP, which in turn will induce an increased transactivation of the luciferase reporter construct. The luciferase activity

was quantified using a luminescence counter. The compounds were tested in the concentration range of 0.1 nM to 10 μ M. This assay was used to determine the EC₅₀ (concentration of test compound causing half-maximal (50 %) luciferase stimulation) and efficacy of the compounds compared to recombinant human FSH. For this, the software program XLfit (Excel version 2.0, built 30, ID Business Solutions Limited) was used.

Compounds of all examples had an activity (EC₅₀) of less than 10⁻⁶ M. Some of the compounds, such as those of examples 1, 3, 5, 7, 11, 15, 17, 22, 23, 24, 25, 34, 38, 39, 42, 50, 51, 54, 56, 58, 62 and 72 showed an EC₅₀ of less than 10⁻⁸ M.

Claims

1. A 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to Formula I,



Formula I

- or a pharmaceutically acceptable salt thereof, wherein
- R^1 is (1-6C)alkyl, (2-6C)alkenyl or (2-6C)alkynyl;
- R^2 , R^3 are independently halogen, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (3-4C)alkenyloxy or (3-4C)alkynyloxy;
- R^4 is phenyl or (2-5C)heteroaryl, both substituted with R^7 and optionally substituted on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylthio and (di)(1-4C)alkylamino;
- R^7 is H, (1-4C)alkylthio, (1-4C)alkylsulfonyl, (di)(1-4C)alkylamino, R^8R^9 -amino, $R^{10}R^{11}$ -aminocarbonyl, $R^{12}R^{13}$ -amino(1-4C)alkylcarbonylamino, $R^{14}R^{15}$ -amino(1-4C)alkyl, R^{16} -oxy, $R^{17}R^{18}$ -aminocarbonyl(1-4C)alkoxy, R^{19} -oxy(1-4C)alkyl, R^{19} -oxycarbonyl(1-4C)alkyl, $R^{20}R^{21}$ -aminosulfonyl, R^{20} -oxysulfonyl, aminoiminomethyl, (di)(1-4C)alkylaminoiminomethyl or (2-6C)-heterocycloalkyliminomethyl, trifluoromethylsulfonyl; R^{23} -oxycarbonyl, R^{23} -carbonyl or $R^{23}R^{24}$ -aminocarbonyl;
- R^8 is H or (1-4C)alkyl;
- R^9 is (1-4C)alkylsulfonyl, (1-6C)alkylcarbonyl, (2-6C)alkenylcarbonyl, (2-6C)-alkynylcarbonyl, (3-6C)cycloalkylcarbonyl, (3-6C)cycloalkyl(1-4C)alkylcarbonyl, (1-4C)alkoxycarbonyl, (3-4C)alkenyloxycarbonyl, (3-4C)alkynyloxy-carbonyl, (di)(1-4C)alkylaminocarbonyl, (2-6C)heterocycloalkylcarbonyl, (5-8C)alkyl, (3-6C)cycloalkyl, (3-6C)cycloalkyl(1-4C)alkyl, (di)(1-4C)alkyl-

amino(2-4C)alkyl, (2-6C)heterocycloalkyl(2-4C)alkyl or phenylcarbonyl, phenylsulfonyl, phenyl(1-4C)alkoxy(1-4C)alkylcarbonyl, phenyl(1-4C)alkyl, (2-5C)heteroarylcarbonyl, (2-5C)heteroarylsulfonyl, (2-5C)heteroaryl(1-4C)-alkyl, optionally substituted on the (hetero)aromatic ring with one or more
5 substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (1-4C)alkoxy or (di)(1-4C)alkylamino;
 R^{10} is H or (1-4C)alkyl;
 R^{11} is hydroxy(2-4C)alkyl, amino(2-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl or (di)-(1-4C)alkylamino(2-4C)alkyl; or
10 $R^{10}R^{11}$ in $R^{10}R^{11}$ -aminocarbonyl may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring substituted with one or more substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy(1-4C)alkyl;
 R^{12} , R^{13} are independently H, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (3-6C)-cycloalkyl, hydroxy(2-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl, (3-6C)cycloalkyl-
15 (1-4C)alkyl, (2-6C)heterocycloalkyl(1-4C)alkyl, amino(2-4C)alkyl, (di)(1-4C)-alkylamino(2-4C)alkyl or phenyl(1-4C)alkyl, (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)-alkyl, (1-4C)alkoxy and (di)(1-4C)alkylamino; or
20 $R^{12}R^{13}$ in $R^{12}R^{13}$ -amino(1-4C)alkylcarbonylamino may be joined in a (4-6C)-heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)-alkyl and hydroxy(1-4C)alkyl;
 R^{14} , R^{15} are independently H, (1-6C)alkyl, (2-6C)heterocycloalkyl(1-4C)alkyl, (3-6C)cycloalkyl(1-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl, hydroxy(2-4C)alkyl, (di)(1-4C)alkylamino(2-4C)alkyl, amino(2-4C)alkyl, (1-4C)alkoxycarbonyl-
25 (1-4C)alkyl, (1-6C)alkylcarbonyl, (3-6C)cycloalkylcarbonyl, (1-4C)alkoxy-carbonyl, (3-4C)alkenyloxycarbonyl, (di)(1-4C)alkylaminocarbonyl, (2-6C)-heterocycloalkylcarbonyl or (2-5C)heteroaryl(1-4C)alkyl, phenyl(1-4C)alkyl, (2-5C)heteroarylcarbonyl, phenylcarbonyl, optionally substituted on the
30 (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl or (1-4C)alkoxy,

(di)(1-4C)alkylamino; or

$R^{14}R^{15}$ in $R^{14}R^{15}$ -amino(1-4C)alkyl may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy(1-4C)alkyl;

R^{16} is (2-6C)heterocycloalkyl(1-4C)alkyl, (di)(1-4C)alkylamino(2-4C)alkyl, (2-4C)alkoxy(1-4C)alkyl, hydroxy(2-4C)alkyl, amino(2-4C)alkyl, hydroxycarbonyl(1-4C)alkyl, (1-4C)alkoxycarbonyl(1-4C)alkyl, (1-4C)alkoxycarbonyl, (3-4C)alkenyloxycarbonyl, (3-4C)alkynyloxycarbonyl, (di)(1-4C)-alkylaminocarbonyl, (2-6C)heterocycloalkylcarbonyl, or phenyl(1-4C)alkyl, (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (1-4C)alkoxy and (di)(1-4C)alkylamino;

R^{17}, R^{18} are independently H, (1-6C)alkyl, (3-6C)cycloalkyl(1-4C)alkyl, (1-4C)-alkoxy(2-4C)alkyl, hydroxy(2-4C)alkyl, amino(2-4C)alkyl, (di)(1-4C)alkyl-amino(2-4C)alkyl, (2-6C)heterocycloalkyl(2-4C)alkyl, or phenyl(1-4C)alkyl, (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl, cyano, (1-4C)alkyl, (1-4C)alkoxy and (di)(1-4C)alkylamino; or $R^{17}R^{18}$ in $R^{17}R^{18}$ -aminocarbonyl(1-4C)alkoxy may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy(1-4C)alkyl;

R^{19} is H or (1-6C)alkyl;

R^{20}, R^{21} are independently H, (1-6C)alkyl, (1-6C)alkenyl, (1-6C)alkynyl or (1-4C)alkoxy(1-4C)alkyl; or

$R^{20}R^{21}$ in $R^{20}R^{21}$ -aminosulfonyl may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy(1-4C)alkyl;

X is O or N- R^{22} ;

Y is CH_2 , C(O) or SO_2 ;

Z is CN or NO₂;

R²² is H, (1-4C)alkyl;

R²³, R²⁴ are independently H, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (3-6C)-cycloalkyl, (3-6C)cycloalkyl(1-4C)alkyl, (2-6C)heterocycloalkyl, (2-6C)hetero-
 5 cycloalkyl(1-4C)alkyl, (1-4C)alkoxycarbonyl(1-4C)alkyl, (di)(1-4C)alkylamino-
 carbonyl(1-4C)alkyl or phenylaminocarbonyl(1-4C)alkyl, (2-5C)heteroaryl-
 aminocarbonyl(1-4C)alkyl, phenyl, (2-5C)heteroaryl, phenyl(1-4C)alkyl, (2-
 5C)heteroaryl(1-4C)alkyl, optionally substituted at the (hetero)atom with one or
 more substituents selected from hydroxy, amino, halogen, nitro, trifluoromethyl,
 10 cyano, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy and (di)(1-4C)-
 alkylamino; or

R²³R²⁴ in R²³R²⁴-aminocarbonyl may be joined in a (4-6C)heterocycloalkenyl
 ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more
 substituents selected from (1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl and hydroxy-
 15 (1-4C)alkyl; with the proviso that the compounds of formula I wherein X is O, R⁴
 is phenyl and R⁷ is selected from H, (1-4C)alkylthio, (1-4C)alkylsulfonyl,
 di(1-4C)alkylamino, R²³-oxycarbonyl, R²³-carbonyl and R²³R²⁴-aminocarbonyl,
 and the compounds of formula I wherein X is O, R⁴ is (2-5C)heteroaryl and R⁷ is
 H or (di)(1-4C)alkylamino, are excluded.

- 20 2. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to
 claim 1 wherein R¹ is (1-6C)alkyl.
3. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to
 claims 1-2 wherein R² is halogen.
4. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to
 25 claims 1-3 wherein R³ is (1-4C)alkoxy.
5. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to
 claims 1-4 wherein Z is CN.
6. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to
 claims 1-5 wherein X is O.
- 30 7. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to
 claims 1-6 wherein Y is CH₂.

8. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to claims 1-7 wherein R^7 is R^8R^9 -amino, $R^{10}R^{11}$ -aminocarbonyl, $R^{12}R^{13}$ -amino- (1-4C)alkylcarbonylamino, $R^{14}R^{15}$ -amino(1-4C)alkyl or $R^{17}R^{18}$ -aminocarbonyl(1-4C)alkoxy.
- 5 9. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to claims 1-8 wherein
 R^8 is H;
 R^9 is (1-4C)alkylsulfonyl, (1-6C)alkylcarbonyl, (3-6C)cycloalkylcarbonyl, (1-4C)alkoxycarbonyl, (3-4C)alkenyloxycarbonyl, (di)(1-4C)alkylamino-
10 carbonyl, (2-6C)heterocycloalkylcarbonyl, or phenylcarbonyl, phenyl(1-4C)-alkoxy(1-4C)alkylcarbonyl, (2-5C)heteroarylcarbonyl, (2-5C)heteroarylsulfonyl or (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from, halogen or (1-4C)alkoxy;
 R^{10} is H or (1-4C)alkyl;
15 R^{11} is hydroxy(2-4C)alkyl, amino(2-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl or (di)-(1-4C)alkylamino(2-4C)alkyl; or
 $R^{10}R^{11}$ in $R^{10}R^{11}$ -aminocarbonyl may be joined in a (4-6C)heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl;
20 R^{12} , R^{13} are independently H, (1-6C)alkyl, (3-6C)cycloalkyl, hydroxy(2-4C)-alkyl, (1-4C)alkoxy(2-4C)alkyl, (2-6C)heterocycloalkyl(1-4C)alkyl or (di)-(1-4C)alkylamino(2-4C)alkyl or phenyl(1-4C)alkyl or (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from halogen; or
25 $R^{12}R^{13}$ in $R^{12}R^{13}$ -amino(1-4C)alkylcarbonylamino may be joined in a (4-6C)-heterocycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl or hydroxy(1-4C)alkyl;
 R^{14} , R^{15} are independently H, (1-6C)alkyl, hydroxy(2-4C)alkyl, (1-4C)alkoxy-carbonyl(1-4C)alkyl, (1-6C)alkylcarbonyl, (3-6C)cycloalkylcarbonyl, (1-4C)-alkoxycarbonyl, (2-5C)heteroaryl(1-4C)alkyl, (2-5C)heteroarylcarbonyl or
30 phenylcarbonyl;
 R^{16} is (2-6C)heterocycloalkyl(1-4C)alkyl, (di)(1-4C)alkylamino(2-4C)alkyl,

(2-4C)alkoxy(1-4C)alkyl, hydroxycarbonyl(1-4C)alkyl, (1-4C)alkoxycarbonyl(1-4C)alkyl; or phenyl(1-4C)alkyl or (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from halogen or (1-4C)alkoxy;

5 R^{17}, R^{18} are independently H, (1-6C)alkyl, (3-6C)cycloalkyl(1-4C)alkyl, (1-4C)alkoxy(2-4C)alkyl or (2-6C)heterocycloalkyl(2-4C)alkyl; or phenyl(1-4C)alkyl or (2-5C)heteroaryl(1-4C)alkyl, optionally substituted on the (hetero)aromatic ring with one or more substituents selected from halogen; or $R^{17}R^{18}$ in $R^{17}R^{18}$ -aminocarbonyl(1-4C)alkoxy may be joined in a (4-6C)hetero-
10 cycloalkenyl ring or a (2-6C)heterocycloalkyl ring, optionally substituted with one or more substituents selected from (1-4C)alkyl or hydroxy(1-4C)alkyl.

10. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to claim 8 wherein R^7 is R^8R^9 -amino.

11. The 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to
15 claim 10 wherein R^8 is H and R^9 is (1-4C)alkylsulfonyl.

12. A 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative or a pharmaceutically acceptable salt thereof selected from the group of
N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3,4,5-trimethoxy-benzamide;
20 *N*-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[(pyridin-4-ylmethyl)-amino]-acetamide;

N-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-2-[4-(2-hydroxy-ethyl)-
25 piperazin-1-yl]-acetamide;
4-{3-Bromo-4-[3-(3,6-dihydro-2*H*-pyridine-1-carbonyl)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;

3-[Bis-(2-methoxy-ethyl)-amino]-*N*-{3-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-
30 propionamide;

2-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-

quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenoxy}-*N,N*-dimethyl-acetamide;
4-{3-Bromo-5-ethoxy-4-[3-(2-morpholin-4-yl-2-oxo-ethoxy)-benzyloxy]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;

5 Furan-2-carboxylic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide;
N-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-acrylamide;

Cyclopropanecarboxylic acid {2-[2-bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-amide;
10 2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid methyl ester;
1-{2-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-3-methyl-urea

15 {3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-pyridin-2-yl}-carbamic acid methyl ester

4-(3-Bromo-5-ethoxy-4-{3-[(1*H*-imidazol-4-ylmethyl)-amino]-benzyloxy}-phenyl)-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;

20 *N*-{3-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-methanesulfonamide;

{4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-phenyl}-carbamic acid allyl ester;

25 4-[3-Bromo-5-ethoxy-4-(1-methanesulfonyl-1*H*-pyrrol-2-ylmethoxy)-phenyl]-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;

4-[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-*N*-methyl-benzamidine;

4-{3-Bromo-5-ethoxy-4-[(pyridin-3-ylmethyl)-amino]-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile;

30 *N*-(2-{[2-Bromo-4-(3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenylamino]-methyl}-phenyl)-methanesulfonamide;

4-{3-Bromo-4-[2-(cyclopropylmethyl-amino)-benzyloxy]-5-ethoxy-phenyl}-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinoline-3-carbonitrile; or
N-{2-[2-Bromo-4-((4*R*,7*S*)-3-cyano-2-methyl-5-oxo-7-propyl-1,4,5,6,7,8-hexahydro-quinolin-4-yl)-6-ethoxy-phenoxy-methyl]-4,5-difluoro-phenyl}-methanesulfonamide

- 5
13. A 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative according to claims 1-12 for use in therapy.
14. A pharmaceutical composition comprising a 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative of any one of claims 1-12, and pharmaceutically
- 10 suitable auxiliaries.
15. Use of the 4-phenyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline derivative of any one of claims 1-12, or a pharmaceutically acceptable salt or solvate thereof for the manufacture of a medicament for the treatment of fertility disorders.