SEQUENCE LISTING

<110> Novo Nordisk A/S

<120> SOLID COMPOSITIONS COMPRISING A GLP-1 AGONIST AND A SALT OF N-(8-(2-HYDROXYBENZOYL)AMINO)CAPRYLIC ACID

<130> 180033WO01

<160> 6

<170> PatentIn version 3.5

<210> 1
<211> 31
<212> PRT
<213> Homo sapiens

<400> 1

His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly
1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly
20 25 30

<210> 2
<211> 40
<212> PRT
<213> Heloderma suspectum

<400> 2

His Gly Glu Gly Thr Phe Ile Thr Ser Asp Leu Ser Lys Gln Met Glu
1 5 10 15

Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro
20 25 30

Ser Ser Gly Ala Pro Pro Pro Ser
35 40

<210> 3
<211> 29
<212> PRT

<213> Homo sapiens

<400> 3

His Ser Gln Gly Thr Phe Thr Ser Asp Tyr Ser Lys Tyr Leu Asp Ser
1   5   10   15

Arg Arg Ala Gln Asp Phe Val Gln Trp Leu Met Asn Thr
            20   25

<210> 4
<211> 31
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic

<220>
<221> MOD_RES
<222> (2)..(2)
<223> Aib

<220>
<221> MOD_RES
<222> (20)..(20)
<223> This amino acid residue is covalently attached to a substituent

<400> 4

His Xaa Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly
1   5   10   15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Arg Gly Arg Gly
            20   25   30

<210> 5
<211> 31
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic

<220>
<221> MOD_RES
<222> (2)..(2)
<223> Aib

<220>
<221> MOD_RES
<222> (20)..(20)
<223> The amino acid residue is modified with a subsitutent

<220>
<221> MOD_RES
<222> (31)..(31)
<223> The amino acid residue is modified with a subsitutent

<400> 5

His Xaa Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly
1   5   10   15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Arg Gly Arg Lys
            20   25   30

<210> 6
<211> 33
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic

<220>
<221> MOD_RES
<222> (2)..(2)
<223> Aib

<220>
<221> MOD_RES
<222> (21)..(21)
<223> The amino acid residue is modified with a substituent

<220>
<221> MOD_RES
<222> (30)..(30)
<223> The amino acid residue is modified with a substituent

<400> 6

```
His Xaa Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu
1               5                   10                  15


Gln Ala Ala Arg Lys Phe Ile Gly Trp Leu Val Arg Gly Lys Gly Glu
            20                  25                  30


Gly
```