the solvent (B), and 1% to 30% by mass of the emulsifier (C), with respect to a total mass of the agrochemical emulsifiable composition.

[3] The agrochemical emulsifiable composition according to [1] or [2], wherein the agrochemical active ingredient (A) is at least one selected from the group consisting of flometoquin, carbosulfan, benfuracarb, cadusafos, chlorpyrifos, prothiofos, profenofos, dienochlor, fipronil, fluxametamide, broflanilide, acrinathrin, allethrin, etofenprox, cyhalothrin, cyfluthrin, cypermethrin, silafluofen, tefluthrin, tralomethrin, bifenthrin, pyrethrin, fenvalerate, fenpropathrin, flucythrinate, fluvalinate, permethrin, indoxacarb, metaflumizone, abamectin, emamectin benzoate, milbemectin, lepimectin, acynonapyr, amitraz, flubendiamide, tebufenpyrad, tolfenpyrad, pyridaben, fenazaquin, pyrimidifen, fenpyroximate, cyenopyrafen, cyflumetofen, pyflubumide, acequinocyl, fluacrypyrim, diafenthiuron, propargite, tetradifon, chlorfenapyr, pyriproxyfen, tebufenozide, chlorfluazuron, teflubenzuron, novaluron, flufenoxuron, lufenuron, clofentezine, etoxazole, buprofezin, spirodiclofen, spiromesifen, and pyridalyl.

[4] The agrochemical emulsifiable composition according to [3], wherein the agrochemical active