

HAEMOPOIETIC GROWTH FACTOR ANTAGONISTS

The present invention relates to modified and variant forms of haemopoietic growth factors capable of acting as antagonists to the corresponding native haemopoietic growth factors and their use in ameliorating aberrant effects caused by the native molecules.

Bibliographic details of the publications numerically referred to in this specification are collected at the end of the description. Sequence Identity Numbers (SEQ ID NOs.) for the nucleotide and amino acid sequences referred to in the specification are defined following the bibliography.

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

Granulocyte-macrophage colony-stimulating factor (GM-CSF) is one member of a family of haemopoietic growth factors (HGFs) which have a similar predicted tertiary configuration (Parry *et al*, 1988) and whose receptors also belong to a common family (Gearing *et al*, 1989, Bazan, 1990). This family of haemopoietic growth factors includes, for example, in addition to GM-CSF, the cytokines IL-2, IL-3, IL-5, IL-6 and IL-10. A distinct subfamily comprising GM-CSF, IL-3 and IL-5 can be discerned based on structural similarities (Goodall *et al*, 1993) and on their ability to interact with a common receptor component (Lopez *et al*, 1992).

Human GM-CSF (hGM-CSF) comprises 127 amino acids and is available in recombinant form (rhGM-CSF). The hGM-CSF receptor has also been cloned and shown to comprise a binding (α) chain exhibiting low affinity binding to GM-CSF (Gearing *et al*, 1989) and a second (β) chain which does not measurably bind GM-CSF by itself but it allows the formation of a high affinity receptor when co-expressed with the α chain (Hayashide *et al*, 1990).

GM-CSF exhibits a range of activities extending over neutrophil, eosinophil and monocyte lineages. Specifically, GM-CSF stimulates the progenitors of these cells to proliferate and differentiate to become mature cells. In addition, it stimulates mature cells to greater function. The stimulation of mature cells results in greater capacity to phagocytose and kill micro-organisms, kill antibody-coated tumour cells and parasites and generate superoxide anion (O_2^-) in response to various stimuli. The purpose of this activation is presumed to enable the mature cells to become better effector cells in inflammatory reactions.

10 Therapeutically, the HGFs form an important group of molecules for their actual or potential properties. For example, the main indications for GM-CSF are for its effects on progenitor cells or mature cells. Using its effects on progenitor cells, GM-CSF is used in the treatment of bone marrow failure as seen in aplastic anaemia or chemotherapy or AIDS-induced marrow suppression. In the treatment of infections, the capacity to
15 stimulate mature cells is especially relevant. The capacity of GM-CSF-activated neutrophils and eosinophils to kill tumour cells that have bound antibody is especially remarkable and could be used in tumour therapy.

However, despite the actual and potential benefits of HGFs, they can exhibit some
20 detrimental side effects. For example, GM-CSF can exhibit toxicity due to stimulation of mature cells causing blood vessel damage or thrombosis. The eosinophilia caused by GM-CSF appears especially damaging in this regard. The molecule can also have detrimental effects by stimulating growth of leukaemia cells and tumour cells of non-haemopoietic origin and stimulating production of inflammatory mediators.

25

International Patent Application No. PCT/AU89/00177 and an article by Lopez *et al.* (1992) disclose amino acid variants of GM-CSF which have exhibited reduced potency. These variants were investigated further for their potential as GM-CSF antagonists. However, the variants cause classical stimulation at concentrations 100 fold greater
30 compared to the native GM-CSF molecule. Furthermore, attempts to find antagonistic properties failed since mixing large concentrations of one of the variants with suboptimal concentrations of native GM-CSF resulted in stronger GM-CSF stimulation with no

evidence of inhibition being observed.

There is a need, therefore, to develop antagonists to HGFs and in particular GM-CSF which are capable of ameliorating the aberrant effects of the corresponding native
5 molecules. There is also a need for such antagonists not to exhibit agonist properties in respect of the corresponding HGFs.

Accordingly, one aspect of the present invention provides a haemopoietic growth factor characterised by being in unglycosylated form and comprising a sequence of amino acids
10 within a first α -helix wherein one or more exposed amino acids in said first α -helix having acidic or acidic-like properties are substituted with a basic amino acid residue.

In accordance with the present invention, it is proposed that the modified HGFs defined above act as antagonists of the native form of the corresponding HGF but not other
15 HGFs. The term "modified" is considered herein synonymous with terms such as "variant", "derivative" and "mutant".

The HGFs are preferably GM-CSF, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-10, granulocyte colony-stimulating factor (G-CSF) and erythropoietin (EPO) modified in
20 accordance with the present invention. Most preferably, the HGF is GM-CSF. The HGFs are preferably in recombinant or synthetic form and, with the exception of the amino acid substitution(s) in the first α -helix, the amino acid sequence of the HGF may be the same as the naturally occurring molecule (i.e. native molecule) or may carry single or multiple amino acid substitutions, deletions and/or additions to the native amino acid
25 sequence. The HGF sequences are preferably of mammalian origin such as from humans, livestock animals, companion animals or laboratory test animals. Most preferably, the HGFs are of human origin or of a mammalian origin capable of functioning in humans.

The first α -helix of GM-CSF has been determined at 2.4 angstrom resolution by X-ray
30 crystallography and encompasses amino acid residues 13 to 28. Similar procedures may be adopted to determine the first α -helix in other haemopoietic growth factors. The position may also be determined by analogy to GM-CSF structure.

Reference to "unglycosylated form" herein means that the molecule is completely unglycosylated such as when expressed in recombinant form in a prokaryotic organism (e.g. *E. coli*). Alternatively, a glycosylation-deficient mammalian cell may be used or complete deglycosylation may occur *in vitro* using appropriate enzymes. Accordingly, the present invention extends to chemically synthesised GM-CSF which is in unglycosylated form.

An "exposed" amino acid is taken herein to refer to an amino acid on an exposed or outer portion of an α -helix compared to those amino acids orientated towards the inside of the molecule.

An acidic amino acid includes, for example, Glu and Asp. Preferred basic amino acids are Arg and Lys.

According to another aspect of the present invention, there is provided a haemopoietic growth factor characterised by:

- (i) being in unglycosylated form;
- (ii) comprising a sequence of amino acids within the first α -helix;
- (iii) one or more exposed amino acids in said α -helix which have acidic or acidic-like properties being substituted by a basic amino acid residue;
- (iv) being in recombinant or synthetic form;
- (v) being capable of acting as an antagonist for at least one property of the corresponding native HGF.

This aspect of the present invention is predicated in part on the surprising discovery that a mutation in amino acid 21 (Glu) of hGM-CSF to Arg or Lys together with the variant in GM-CSF being in unglycosylated form results in the hGM-CSF variant being unable to detectably exhibit classical GM-CSF function. The variants, referred to herein "GM-CSF Arg²¹" and "GM-CSF Lys²¹" are unable to bind to high affinity receptors but are still able to fully bind the low affinity α chain of the GM-CSF receptor. Importantly, the non-glycosylated GM-CSF Arg²¹ and GM-CSF Lys²¹ act as antagonists, preventing the stimulatory effect of native GM-CSF. For convenience, the numbering of amino acid

residues in hGM-CSF is taken from Wong *et al.* (1985).

By way of a shorthand notation the following three letter abbreviations for amino acid residues are used in the specification as defined in Table 1.

5

Where a specific amino residue is referred to by its position in the polypeptide of an HGF, the amino acid abbreviation is used with the residue number given in superscript (i.e. Xaaⁿ, wherein Xaa is the amino acid residue).

10

Table 1

15

20

25

30

35

Amino acid	Three-letter abbreviation	Corresponding single-letter abbreviation
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartic acid	Asp	D
Cysteine	Cys	C
Glutamine	Gln	Q
Glutamic acid	Glu	E
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L
Lysine	Lys	K
Methionine	Met	M
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S
Threonine	Thr	T
Tryptophan	Trp	W
Tyrosine	Tyr	Y
Valine	Val	V

The present invention is exemplified using GM-CSF and in particular hGM-CSF Arg²¹ and hGM-CSF Lys²¹. This is done, however, with the understanding that the present invention extends to all HGFs as hereinbefore described.

5

For example, given that there is a Glu at position 22 of IL-3 and position 13 in IL-5 which in the three dimensional structure (in the case of IL-5) occupies an equivalent position to Glu²¹ of GM-CSF, then the present invention provides the basis for the creation of antagonists in IL-3 and IL-5. The substitution of Glu at these positions in IL-
10 3 and IL-5 may be the sole mutation or it may be in combination with other amino acid mutations (including substitutions, deletions and/or additions) for the development of effective antagonists. This similarly applies to other HGFs based on the acidic or acidic-like amino acid residues in the first α -helix of the molecule. The location of the N-terminal helix can in each case be readily determined on comparable motifs from
15 predicted helices. Such HGFs are listed in Table 2 showing the acidic or acidic-like amino acid residue in bold in the equivalent position to Glu²¹ of hGM-CSF. The acidic or acidic-like amino acid residues are readily substituted by, for example, recombinant DNA technology.

20 According to another aspect of the present invention there is provided a modified variant including HGF comprising an amino acid sequence in the first α -helix of said HGF selected from the group consisting of:

- 25 i) His Val Asn Ala Ile Gln Xaa Ala Arg Arg Leu Leu Asn Leu (SEQ ID No. 1);
- ii) Ala Leu Val Lys Xaa Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu (SEQ ID No. 2);
- iii) Asn Met Ile Xaa Xaa Ile Ile Thr His Leu (SEQ ID No. 3);
- iv) Leu Leu Leu Xaa Leu Gln Met Ile Leu (SEQ ID No. 4);
- 30 v) Ile Thr Leu Gln Xaa Ile Ile Lys Thr Leu (SEQ ID No. 5);
- vi) Arg Tyr Ile Leu Xaa Gly Ile Ser Ala Leu Arg Lys (SEQ ID No. 6);
- vii) Gly Asp Gln Tyr Xaa Ser Val Leu Met Val Ser Ile (SEQ ID No. 7);

- viii) Ala Gly Ile Leu Xaa Ile Asn Phe Leu Ile Asn Lys Met Gln
Glu Asp (SEQ ID No. 8);
- ix) Asn Met Leu Arg Xaa Leu Arg Asp Ala Phe Ser
(SEQ ID No. 9);
- 5 x) Phe Leu Leu Lys Cys Leu Xaa Gln Val Arg Lys Ile
(SEQ ID No. 10); and
- xi) Tyr Leu Leu Glu Ala Lys Xaa Ala Glu Asn Ile Thr Thr Gly
(SEQ ID No. 11);

10 wherein Xaa is a basic amino acid, preferably selected from the group consisting of Arg and Lys, and wherein said variant HGF is in unglycosylated form and acts as an antagonist for at least one property of the corresponding native HGF. Preferably, the haemopoietic growth factor is hGM-CSF and Xaa is Arg or Lys at position 21 of the first α -helix.

15

The HGF antagonists of the present invention and in particular GM-CSF Arg²¹ and GM-CSF Lys²¹ are useful *inter alia* in the treatment of myeloid and lymphocyte leukaemias, some tumours of non-haemopoietic origins and acute and chronic inflammation such as asthma and rheumatoid arthritis. These and other conditions are considered herein to

20 result from or be facilitated by the aberrant effects of an endogenous HGF such as GM-CSF. hGM-CSF Arg²¹ and hGM-CSF Lys²¹ will also be useful in mobilising stem cells and progenitor cells into the circulation without the risk of activating neutrophils and monocytes. Other related molecules may have different useful properties.

25 The present invention, therefore, contemplates a method of treatment comprising the administration to a mammal of an effective amount of a modified HGF as hereinbefore defined and in particular hGM-CSF Arg²¹ or hGM-CSF Lys²¹ or both for a time and under conditions sufficient for effecting said treatment. Generally, the mammal is a human, livestock animal, companion animal or laboratory test animal. Most preferably,

30 the mammal is a human. A single modified HGF may be administered or a combination of variants of the same HGF. For example, a range of hGM-CSF variants could be used such as a combination of hGM-CSF Arg²¹ and hGM-CSF Lys²¹.

TABLE 2

Cytokines related to GM-CSF exhibit a conserved acidic residue analogous to E21 in GM-CSF

CYTOKINE ²	HELIX ¹ (Amino Acid Residue No.)	AMINO ACID SEQUENCE	SEQ ID No.
hGM-CSF	15-28	His Val Asn Ala Ile Gln Glu Ala Arg Arg Leu Leu Asn Leu	12
hIL-5	9-24	Ala Leu Val Lys Glu Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu	13
hIL-3	18-27	Asn Met Ile Asp Glu Ile Ile Thr His Leu	14
hIL-2	17-25	Leu Leu Leu Asp Leu Gln Met Ile Leu	15
hIL-4	8-17	Ile Thr Leu Gln Asp Ile Ile Lys Thr Leu	16
hIL-6	18-43	Arg Tyr Ile Leu Asp Gly Ile Ser Ala Leu Arg Lys	17
hIL-7	9-20	Gly Asp Gln Tyr Glu Ser Val Leu Met Val Ser Ile	18
hIL-9	7-22	Ala Gly Ile Leu Asp Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp	19
hIL-10	21-31	Asn Met Leu Arg Asp Leu Arg Asp Ala Phe Ser	20
hG-CSF	13-24	Phe Leu Leu Lys Cys Leu Glu Gln Val Arg Lys Ile	21
hEPO	4-28	Tyr Leu Leu Glu Ala Lys Glu Ala Glu Asn Ile Thr Thr Gly	22

¹ Only the pertinent portion of each helix is shown.

² Predicted helices are from: for IL-2, (Brandhuber et al, 1987; Zurawski and Zurawski, 1989); for hIL-3 (Parry et al, 1988); for hIL-5, (Parry et al, 1988); for hIL-6 (Bazan, 1990a); for hG-CSF, (Parry et al, 1988); for hEPO, (Bazan, 1990); for hGM-CSF the first helix was determined from the crystal structure (Karpplus, 1991). The location of the N-terminal helix in the other cytokines was based on comparable motifs from these secondary structure predictions.

The present invention also provides a pharmaceutical composition comprising the variant HGFs as hereinbefore defined or combinations thereof. Most particularly, the pharmaceutical composition comprises hGM-CSF Arg²¹ or hGM-CSF Lys²¹ or both.

5

Methods for preparing pharmaceutical compositions are well known in the art and reference can conveniently be made to Remington's Pharmaceutical Sciences, Mack Publishing Company, Eaton, Pennsylvania, USA and may also include one or more pharmaceutical acceptable carriers and/or diluents.

10

The present invention is further described by reference to the following non-limiting Examples and/or Figures.

In the Figures:

15

Figure 1 is a graphical representation showing titration of *E. coli* derived GM-CSF Arg²¹ for its ability to affect O₂⁻ production in human neutrophils (□) and to antagonise the enhancement of O₂⁻ by wild type GM-CSF tested at 1.0 ng/ml (Δ) and at 0.1 ng/ml (closed triangle).

20

Figure 2 is a graphical representation showing failure of *E. coli*-derived GM-CSF Arg²¹ to antagonise the enhancement of O₂⁻ production in human neutrophils stimulated with tumour necrosis factor-α (TNF-α).

25

Figure 3 is a graphical representation showing competitive inhibition of ¹²⁵I-GM-CSF binding to COS cells transfected with the GM-CSF receptor α chain alone (top) or α and β chains (bottom) by GM-CSF Arg²¹.

30

Figure 4 is a graphical representation showing titration of GM-CSF Arg²¹ for its ability to antagonise GM-CSF (A) in contrast to no effect on IL-3 (B)-mediated proliferation of TF-1 cells.

Figure 5 is a graphical representation showing the titration of GM-CSF Arg²¹ for its ability to antagonise TF-1 proliferation stimulated by either *E. coli*-derived GM-CSF (A), yeast-derived GM-CSF (B) or CHO-derived GM-CSF (C).

- 5 Figure 6 is a graphical representation showing the titration of GM-CSF Arg²¹ for its ability to antagonise three primary human myeloid leukaemia (A, B & C) *ex vivo*.

Figure 7 is a graphical representation showing that E21R antagonises both GM-CSF but not TNF- α -mediated stimulation of human neutrophils (A), and both E21R and E21K
10 also antagonise neutrophil stimulation by CHO cell-derived GM-CSF (B). In panel A, titrations of *E. coli*-derived wild type GM-CSF ($\bullet$), TNF- α ($\blacklozenge$) and E21R ($\blacksquare$) are shown. In antagonistic experiments, E21R was titrated against 1ng/ml of *E. coli*-derived GM-CSF ($\square$) or 3ng/ml TNF- α ($\circ$). In panel B, titrations of CHO cell-derived wild type GM-CSF ($\circ$), E21R ($\blacksquare$) and E21K ($\blacktriangle$) are shown. In antagonistic experiments, E21R
15 ($\square$) or E21K ($\blacktriangle$) were titrated against 3ng/ml CHO cell-derived GM-CSF. Each value represents the mean of triplicate determinations and error bars represent the SEM.

EXAMPLE 1

Expression of wild type and GM-CSF Arg²¹ and GM-CSF Lys²¹ in an
20 *E. coli* expressed system

Wild type GM-CSF was expressed in *E. coli* using a plasmid (designated pshGM-CSF) containing a synthetic human GM-CSF cDNA cloned into the *E. coli* expression vector pIN-III-OmpH3, a derivative of the vector pIN-III-OmpA2 (Ghrayeb *et al*, 1984). GME21R was expressed from the plasmid pSGM21.1 containing Glu²¹ $\rightarrow$ Arg²¹
25 substitution and was derived from the pSGM-CSF parental plasmid. GME21K was expressed from the plasmid pSGM21.4 containing Glu²¹ $\rightarrow$ Lys²¹ substitution and was derived from the pSGM-CSF parental plasmid.

pSGM21 was generated by initially eliminating a SacII site from the wild type GM-CSF
30 using oligonucleotide cassette mutagenesis to generate plasmid pSGMV1. A 64 bp Nco I/SacII fragment was then excised from the pSGMV1 plasmid and replaced by double-stranded 64bp oligonucleotides containing the appropriate mutation in the DNA sequence.

pSGM21.4 was generated by excising an 88 bp Bgl11/SacII fragment from pSGF1 and replacing it with 88bp oligonucleotides containing the appropriate mutation site directed mutagenesis (Zoller & Smith, 1984).

- 5 Protein was expressed in either MC1061, for wild type GM-CSF or BL21 for GME21R or GME21K, after induction by isopropyl β -D-thiogalactoside and recovered from the periplasmic space by osmotic shock (Koshland and Botstein, 1980).

- 10 GM-CSF protein was purified using a monoclonal antibody 4A12 generated in the laboratory coupled to Sepharose beads. Further purification was achieved by reversed phase HPLC using a BioRad controller and a Brownlee Aquaport C8 100 x 10mm column. GM-CSF was eluted using a 30-50% gradient of acetonitrile in 0.1% trifluoroacetic acid.

- 15 Resultant purified GM-CSF was lyophilised and resuspended in 1 x PBS before being quantitated by HPLC gel filtration. Samples were fractionated on a Beckman Ultraspherogel SEC3000 7.5 x 300mm using a 0.1M Na Phosphate pH 7.0/0.1M Na₂SO₄ mobile phase. Purity was estimated at >95% and area under peaks corresponding to GM-CSF integrated as the extinction coefficient of 0.95 absorbance units.ml.mg⁻¹.

20

EXAMPLE 2

Visualisation of mutant GM-CSF protein

- GM-CSF unpurified or purified from *E. coli* was size-fractionated by NaDodSo₄/12.5% w/v polycarylamide gel electrophoresis (Laemmli, 1970). For Western blot analysis, 25 protein was transferred to nitrocellulose as described (Towbin *et al*, 1979). Filters were probed with a sheep anti-GM-CSF followed by a second layer of biotinylated-rabbit anti-sheep IgG. After a further incubation with an avidin-biotinylated-horseradish peroxidase conjugate, the complex was visualised using a diaminobenzidine substrate solution. For silver staining, the method of Morrissey (1981) was used.

30

EXAMPLE 3

Stimulation of haemopoietic cell proliferation

The human erythroleukaemia cell line TF-1 (and myeloid leukaemia) cells were used to
5 measure the proliferative function of GM-CSF and GM-CSF Arg²¹. Proliferation of TF-1
cells were measured by the ability to incorporate [³H]-thymidine in response to increasing
doses of GM-CSF. This assay was performed as described by Lopez *et al* (1988).

EXAMPLE 4

10 Functional activation of human granulocytes and monocytes

The superoxide anion production assay was carried out as previously described (Lopez
et al, 1986).

EXAMPLE 5

15 Radioreceptor assay

(a) Radioiodination of GM-CSF.

Yeast derived human GM-CSF or *E. coli*-derived human GM-CSF was radioiodinated by
the ICl method (Contreras *et al*, 1983). Iodinated protein was separated from free ¹²⁵I
by chromatography on a Sephadex G-25 PD10 column (Pharmacia, Uppsala, Sweden),
20 equilibrated in phosphate buffered saline (PBS) containing 0.02% w/v Tween 20, and
stored at 4°C for up to 4 weeks. Before use, the iodinated protein was purified from
Tween and non-protein-associated radioactivity by cation exchange chromatography on
a 0.3ml CM-Sepharose CL-6B column (Pharmacia) and stored at 4°C for up to 5 days.
The radiolabelled GM-CSF retained >90% biological activity as judged from titration
25 curves using non-iodinated GM-CSF as controls.

(b) Competition binding assays.

Competition for binding to high affinity and low affinity receptors used stably transfected
CHO cell lines expressing either the α and β chains, or the α chain alone. The cells were
30 suspended in binding medium consisting of RPMI-1640 supplemented with 20mmol/l
HEPES and 0.5% w/v bovine serum albumin (BSA) and 0.1% w/v sodium azide.
Typically, equal volumes (50 μ l) of 4 x 10⁴ CHO cells, iodinated GM-CSF and different

- concentrations of GM-CSF and GM-CSF Arg²¹ were mixed in siliconised glass tubes for 3 hr at 4°C. At the end of the incubation period, cell suspensions were overlaid on 0.2ml foetal calf serum (FCS) at 4°C, centrifuged in a Beckman Microfuge 12, and the tip of each tube containing the visible cell pellet cut off and counted in a gamma counter.
- 5 Specific counts were determined by first subtracting the counts, obtained in the presence of excess wild type GM-CSF.

EXAMPLE 6

Generation of hGM-CSF Variants

- 10 By way of example only, the generation of GM-CSF Arg²¹ is hereinafter described in detail. A human GM-CSF cDNA was subjected to mutagenesis to introduce the amino acid Arg for Glu at position 21. Two mutants were obtained, one containing the Glu²¹→Arg mutation and a second one containing a double mutation X¹⁰→Ile and Glu²¹→Arg. These mutants were cloned into the expression system p1N OMPIII and
- 15 expressed in *E. coli*. Wild type (WT) GM-CSF was expressed in MC1061. GM-CSF Arg²¹ could not be expressed in MC1061. Of twenty strains tested for GM-CSF Arg²¹ expression, BL21 was the highest producer and used for subsequent studies.

- To obtain purified GM-CSF Arg²¹ in high yields a two-step purification procedure was
- 20 devised. In the first step, GM-CSF Arg²¹ was purified by an affinity column constructed with a monoclonal antibody (4A12) that binds to GM-CSF in solution and with high affinity. Affinity-purified GM-CSF Arg²¹ was then purified by reverse-phase HPLC and quantitated by HPLC before being analysed for biological and binding activities.

- 25 It was found that *E. coli*-derived GM-CSF Arg²¹ was unable to enhance neutrophil O₂ production up to a concentration of 3,000 ng/ml (Figure 1). This is different to the inventors' previous results with CHO-derived (i.e. glycosylated) GM-CSF Arg²¹ which was able to enhance neutrophil function at approximately 30 ng/ml which represents a 300 fold reduced potency compared to wild type GM-CSF (Lopez *et al*, 1992).

Despite its inability to activate neutrophils, *E. coli*-derived GM-CSF Arg²¹ was able to bind as well as wild type GM-CSF to the α -chain of the GM-CSF receptor (Figure 3). In contrast, GM-CSF Arg²¹ exhibited an approximate 100-fold reduction in binding to the $\alpha\beta$ GM-CSF receptor complex (Figure 3) indicating that the influence of the β chain has been selectively lost.

This is the first time that a GM-CSF mutant is shown to have unaltered binding to the GM-CSF receptor α -chain as well as being devoid of agonistic activity. This indicates that whilst binding to the α chain is necessary for GM-CSF activity it is not sufficient, and that GM-CSF binding to the β chain is required for GM-CSF-mediated activation.

Since *E. coli*-derived GM-CSF Arg²¹ was not able to stimulate neutrophils yet fully bound the GM-CSF receptor α chain, it was tested for antagonistic activity. The inventors found that this mutant fully inhibited the effect of wild type GM-CSF with an approximately 300-fold excess required to induce 50% inhibition of *E. coli*-derived wild type GM-CSF (Figure 1). This antagonistic effect was specific for GM-CSF as judged by the lack of antagonistic effect of GM-CSF Arg²¹ on TNF enhancement of neutrophil O_2^- production (Figure 2).

20

The antagonistic effect of *E. coli*-derived GM-CSF Arg²¹ was present whether wild type GM-CSF was expressed in *E. coli* (Figure 5A), yeast (Figure 5B) or CHO cells (Figure 5C). In keeping with the differences in binding affinity between heavily glycosylated CHO GM-CSF (lower affinity), partially glycosylated yeast GM-CSF (intermediate affinity) and unglycosylated *E. coli* GM-CSF (higher affinity), *E. coli*-derived GM-CSF Arg²¹ antagonised better the CHO wild type GM-CSF (Figure 5).

The antagonism of *E. coli*-derived GM-CSF Arg²¹ was not restricted to proliferation of the established TF-1 cell line but was also seen in primary myeloid leukaemias. In three different leukaemias, *E. coli*-derived GM-CSF Arg²¹ antagonised the proliferative effect of wild type *E. coli* GM-CSF with an EC50 that varied with each leukaemia (Figure 6)

30

These results show that an unglycosylated GM-CSF molecule with a mutated Glu for an Arg in position 21 of the first α -helix is able to antagonise native GM-CSF.

Since the mutated Glu in GM-CSF is in a position where the same acidic residue or similar acidic residues (e.g. Asp) are present in related growth factors (see Table 2), this invention extends to antagonistic molecules for these growth factors constructed by incorporating the analogous charge reversal mutation. In particular, given that the GM-CSF, IL-3 and IL-5 receptor share the β chain, the Glu²² in IL-3 and the Glu¹³ in IL-5 are predicted to play a similar role. Other variant HGFs are shown in Examples 3 to 22 in which the equivalent or similar amino acid residue to Glu²¹ of hGM-CSF is replaced by either Arg or Lys. In these Examples, the amino acid sequences are provided for the relevant portion of the first α -helix carrying the substitution (see Table 2).

EXAMPLE 7

hGM-CSF Lys²¹

His Val Asn Ala Ile Gln Lys Ala Arg Arg Leu Leu Asn Leu (SEQ ID NO. 23)

EXAMPLE 8

hIL-5 Lys¹³

Ala Leu Val Lys Lys Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu (SEQ ID NO. 24)

EXAMPLE 9

hIL-5 Arg¹³

Ala Leu Val Lys Arg Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu (SEQ ID NO. 25)

EXAMPLE 10

hIL-3 Variants

Asn Met Ile Asp Lys Ile Ile Thr His Leu	hIL-3 Lys ²² (SEQ ID NO. 26)
Asn Met Ile Lys Glu Ile Ile Thr His Leu	hIL-3 Lys ²¹ (SEQ ID NO. 27)
Asn Met Ile Asp Arg Ile Ile Thr His Leu	hIL-3 Arg ²² (SEQ ID NO. 28)
Asn Met Ile Arg Glu Ile Ile Thr His Leu	hIL-3 Arg ²¹ (SEQ ID NO. 29)
Asn Met Ile Lys Lys Ile Ile Thr His Leu	hIL-3 Lys ²¹ Lys ²² (SEQ ID NO. 30)

Asn Met Ile Arg Arg Ile Ile Thr His Leu hIL-3 Arg²¹ Arg²² (SEQ ID NO. 31)

EXAMPLE 11

hIL-2 Lys²⁰

5 Leu Leu Leu Lys Leu Gln Met Ile Leu (SEQ ID NO. 32)

EXAMPLE 12

hIL-2 Arg²⁰

Leu Leu Leu Arg Leu Gln Met Ile Leu (SEQ ID NO. 33)

10

EXAMPLE 13

hIL-4 Lys¹²

Ile Thr Leu Gln Lys Ile Ile Lys Thr Leu (SEQ ID NO. 34)

15

EXAMPLE 14

hIL-4 Arg¹²

Ile Thr Leu Gln Arg Ile Ile Lys Thr Leu (SEQ ID NO. 35)

EXAMPLE 15

hIL-6 Lys²²

20

Arg Tyr Ile Leu Lys Gly Ile Ser Ala Leu Arg Lys (SEQ ID NO. 36)

EXAMPLE 16

hIL-6 Arg²²

25 Arg Tyr Ile Leu Arg Gly Ile Ser Ala Leu Arg Lys (SEQ ID NO. 37)

EXAMPLE 17

hIL-7 Lys¹³

Gly Asp Gln Tyr Lys Ser Val Leu Met Val Ser Ile (SEQ ID NO. 38)

30

EXAMPLE 18

hIL-7 Arg¹³

Gly Asp Gln Tyr Arg Ser Val Leu Met Val Ser Ile (SEQ ID NO. 39)

5

EXAMPLE 19

hIL-9 Lys¹¹

Ala Gly Ile Leu Lys Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp (SEQ ID NO. 40)

10

EXAMPLE 20

hIL-9 Arg¹¹

Ala Gly Ile Leu Arg Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp (SEQ ID NO. 41)

EXAMPLE 21

hIL-10 Lys²⁵

15

Asn Met Leu Arg Lys Leu Arg Asp Ala Phe Ser (SEQ ID NO. 42)

EXAMPLE 22

hIL-10 Arg²⁵

20 Asn Met Leu Arg Arg Leu Arg Asp Ala Phe Ser (SEQ ID NO. 43)

EXAMPLE 23

hG-CSF Lys¹⁹

Phe Leu Leu Lys Cys Leu Lys Gln Val Arg Lys Ile (SEQ ID NO. 44)

25

EXAMPLE 24

hG-CSF Arg¹⁹

Phe Leu Leu Lys Cys Leu Arg Gln Val Arg Lys Ile (SEQ ID NO. 45)

30

EXAMPLE 25

hEPO Lys¹⁹

Tyr Leu Leu Glu Ala Lys Lys Ala Glu Asn Ile Thr Thr Gly (SEQ ID NO. 46)

5

EXAMPLE 26

hEPO Arg¹⁰

Tyr Leu Leu Glu Ala Lys Arg Ala Glu Asn Ile Thr Thr Gly (SEQ ID NO. 47)

- 10 Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all
- 15 combinations of any two or more of said steps or features.

REFERENCES

- Bazan JF. (1990) *Immunol Today* 11, 350-354.
- Brandhuber BJ *et al.* (1987) *Science* 238, 1707-1709.
- Contreras MA *et al.* (1983) *Methods Enzymol* 92, 277-292.
- Diederich *et al.* (1991) *Science* 254, 1779-1782.
- Elliott MJ *et al.* (1990) *J Immunol* 145, 167-176.
- Gasson JC *et al.* (1986) *Proc Natl Acad Sci USA* 83, 669-673.
- Gearing DP *et al.* (1989) *EMBO J* 8, 3667-3676.
- Ghrayeb J *et al.* (1984) *EMBO J* 3, 2437-2442.
- Goodall GJ *et al.* (1993) *Growth Factors* 8, 87-97.
- Hayashide K *et al.* (1990) *Proc Natl Acad Sci USA* 87, 9655-9659.
- Koshland D and Botstein D. (1980) *Cell* 20, 749-760.
- Laemmli UK. (1970) *Nature* 227, 680-685.
- Lopez AF *et al.* (1986) *J Clin Invest* 78, 1202-1228.
- Lopez AF *et al.* (1988) *Blood* 72, 1797-1804.
- Lopez AF *et al.* (1992) *EMBO J* 11, 909-916.
- Morrissey JH. (1981) *Anal Biochem* 117, 307-310.
- Parry, DAD *et al.* (1988) *J Mol Recogn* 1, 107-110.
- Towbin H *et al.* (1979) *Proc Natl Acad Sci USA* 76, 4350-4354.
- Wong G *et al.* (1985) *Science* 228, 810-815.
- Zoller MJ and Smith M. (1984) *DNA* 3, 479-488.
- Zurawski SM and Zurawski G. (1989) *EMBO J* 8, 2583-2590.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) **APPLICANT**
(other than the US): MEDVET SCIENCE PTY LTD
APPLICANT/INVENTORS
(US only): LOPEZ, A.F., SHANNON, M.F. and VADAS, M.A.
- (ii) **TITLE OF INVENTION:** "Haemopoietic Growth Factor Antagonists"
- (iii) **NUMBER OF SEQUENCES:** 47
- (iv) **CORRESPONDENCE ADDRESS:**
 - (A) **ADDRESSEE:** Davies Collison Cave
 - (B) **STREET:** 1 Little Collins Street
 - (C) **CITY:** Melbourne
 - (D) **STATE:** Victoria
 - (E) **COUNTRY:** Australia
 - (F) **ZIP:** 3000
- (v) **COMPUTER READABLE FORM:**
 - (A) **MEDIUM TYPE:** floppy disc
 - (B) **COMPUTER:** IBM PC compatible
 - (C) **OPERATING SYSTEM:** PC-DOS/MS-DOS
 - (D) **SOFTWARE:** PatentIn Release #1.0, Version #1.25
- (vi) **CURRENT APPLICATION DATA:**
 - (A) **APPLICATION NUMBER:** International PCT App.
 - (B) **FILING DATE:** 28-JUL-1994
- (vii) **PRIOR APPLICATION DATA:**
 - (A) **APPLICATION NUMBER:** Provisional AU PM0186
 - (B) **FILING DATE:** 28-JUL-1993
 - (C) **APPLICATION NUMBER:** Provisional AU PM4772
 - (D) **FILING DATE:** 30-MAR-1993
- (viii) **ATTORNEY/AGENT INFORMATION:**
 - (A) **NAME:** HUGHES, DR E JOHN L
 - (C) **REFERENCE/DOCKET NUMBER:** EJH/EK
- (ix) **TELECOMMUNICATION INFORMATION:**
 - (A) **TELEPHONE:** +61 3 254 2777
 - (B) **TELEFAX:** +61 3 254 2770

(2) INFORMATION FOR SEQ ID NO. 1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14 amino acid residues
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 1:

His Val Asn Ala Ile Gln Xaa Ala Arg Arg Leu Leu Asn Leu

(2) INFORMATION FOR SEQ ID NO. 2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 2:

Ala Leu Val Lys Xaa Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu

(2) INFORMATION FOR SEQ ID NO. 3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 amino acid residues
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 3:

Asn Met Ile Xaa Xaa Ile Ile Thr His Leu

(2) INFORMATION FOR SEQ ID NO. 4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 4:

Leu Leu Leu Xaa Leu Gln Met Ile Leu

(2) INFORMATION FOR SEQ ID NO. 5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 5:

Ile Thr Leu Gln Xaa Ile Ile Lys Thr Leu

(2) INFORMATION FOR SEQ ID NO. 6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 6:

Arg Tyr Ile Leu Xaa Gly Ile Ser Ala Leu Arg Lys

(2) INFORMATION FOR SEQ ID NO. 7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 7:

Gly Asp Gln Tyr Xaa Ser Val Leu Met Val Ser Ile

(2) INFORMATION FOR SEQ ID NO. 8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 8:

Ala Gly Ile Leu Xaa Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp

(2) INFORMATION FOR SEQ ID NO. 9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 9:

Asn Met Leu Arg Xaa Leu Arg Asp Ala Phe Ser

(2) INFORMATION FOR SEQ ID NO. 10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 10:

Phe Leu Leu Lys Cys Leu Xaa Gln Val Arg Lys Ile

(2) INFORMATION FOR SEQ ID NO. 11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 11:

Tyr Leu Leu Glu Ala Lys Xaa Ala Glu Asn Ile Thr Thr Gly

(2) INFORMATION FOR SEQ ID NO. 12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 12:

His Val Asn Ala Ile Gln Glu Ala Arg Arg Leu Leu Asn Leu

(2) INFORMATION FOR SEQ ID NO. 13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 13:

Ala Leu Val Lys Glu Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu

(2) INFORMATION FOR SEQ ID NO. 14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 14:

Asn Met Ile Asp Glu Ile Ile Thr His Leu

(2) INFORMATION FOR SEQ ID NO. 15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 15:

Leu Leu Leu Asp Leu Gln Met Ile Leu

(2) INFORMATION FOR SEQ ID NO. 16:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 16:

Ile Thr Leu Gln Asp Ile Ile Lys Thr Leu

(2) INFORMATION FOR SEQ ID NO. 17:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 17:

Arg Tyr Ile Leu Asp Gly Ile Ser Ala Leu Arg Lys

(2) INFORMATION FOR SEQ ID NO. 18:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 18:

Gly Asp Gln Tyr Glu Ser Val Leu Met Val Ser Ile

(2) INFORMATION FOR SEQ ID NO. 19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 19:

Ala Gly Ile Leu Asp Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp

(2) INFORMATION FOR SEQ ID NO. 20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 20:

Asn Met Leu Arg Asp Leu Arg Asp Ala Phe Ser

(2) INFORMATION FOR SEQ ID NO. 21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 21:

Phe Leu Leu Lys Cys Leu Glu Gln Val Arg Lys Ile

(2) INFORMATION FOR SEQ ID NO. 22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 22:

Tyr Leu Leu Glu Ala Lys Glu Ala Glu Asn Ile Thr Thr Gly

(2) INFORMATION FOR SEQ ID NO. 23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 23:

His Val Asn Ala Ile Gln Lys Ala Arg Arg Leu Leu Asn Leu

(2) INFORMATION FOR SEQ ID NO. 24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 24:

Ala Leu Val Lys Lys Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu

(2) INFORMATION FOR SEQ ID NO. 25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 25:

Ala Leu Val Lys Arg Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu

(2) INFORMATION FOR SEQ ID NO. 26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 26:

Asn Met Ile Asp Lys Ile Ile Thr His Leu

(2) INFORMATION FOR SEQ ID NO. 27:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 27:

Asn Met Ile Lys Glu Ile Ile Thr His Leu

(2) INFORMATION FOR SEQ ID NO. 28:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 28:

Asn Met Ile Asp Arg Ile Ile Thr His Leu

(2) INFORMATION FOR SEQ ID NO. 29:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 29:

Asn Met Ile Arg Glu Ile Ile Thr His Leu

(2) INFORMATION FOR SEQ ID NO. 30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 30:

Asn Met Ile Lys Lys Ile Ile Thr His Leu

(2) INFORMATION FOR SEQ ID NO. 31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 31:

Asn Met Ile Arg Arg Ile Ile Thr His Leu

(2) INFORMATION FOR SEQ ID NO. 32:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 32:

Leu Leu Leu Lys Leu Gln Met Ile Leu

(2) INFORMATION FOR SEQ ID NO. 33:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 33:

Leu Leu Leu Arg Leu Gln Met Ile Leu

(2) INFORMATION FOR SEQ ID NO. 34:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 34:

Ile Thr Leu Gln Lys Ile Ile Lys Thr Leu

(2) INFORMATION FOR SEQ ID NO. 35:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 35:

Ile Thr Leu Gln Arg Ile Ile Lys Thr Leu

(2) INFORMATION FOR SEQ ID NO. 36:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 36:

Arg Tyr Ile Leu Lys Gly Ile Ser Ala Leu Arg Lys

(2) INFORMATION FOR SEQ ID NO. 37:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 37:

Arg Tyr Ile Leu Arg Gly Ile Ser Ala Leu Arg Lys

(2) INFORMATION FOR SEQ ID NO. 38:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 38:

Gly Asp Gln Tyr Lys Ser Val Leu Met Val Ser Ile

(2) INFORMATION FOR SEQ ID NO. 39:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 39:

Gly Asp Gln Tyr Arg Ser Val Leu Met Val Ser Ile

(2) INFORMATION FOR SEQ ID NO. 40:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 40:

Ala Gly Ile Leu Lys Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp

(2) INFORMATION FOR SEQ ID NO. 41:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 41:

Ala Gly Ile Leu Arg Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp

(2) INFORMATION FOR SEQ ID NO. 42:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 42:

Asn Met Leu Arg Lys Leu Arg Asp Ala Phe Ser

(2) INFORMATION FOR SEQ ID NO. 43:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 43:

Asn Met Leu Arg Arg Leu Arg Asp Ala Phe Ser

(2) INFORMATION FOR SEQ ID NO. 44:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 44:

Phe Leu Leu Lys Cys Leu Lys Gln Val Arg Lys Ile

(2) INFORMATION FOR SEQ ID NO. 45:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 45:

Phe Leu Leu Lys Cys Leu Arg Gln Val Arg Lys Ile

(2) INFORMATION FOR SEQ ID NO. 46:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 46:

Tyr Leu Leu Glu Ala Lys Lys Ala Glu Asn Ile Thr Thr Gly

(2) INFORMATION FOR SEQ ID. NO. 47:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: polypeptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 47:

Tyr Leu Leu Glu Ala Lys Arg Ala Glu Asn Ile Thr Thr Gly

CLAIMS:

1. A modified haemopoietic growth factor (HGF) characterised by being in unglycosylated form and comprising a sequence of amino acids within a first α -helix wherein one or more exposed amino acids in said first α -helix having acidic or acidic-like properties are substituted with a basic amino acid residue.
2. A modified HGF according to claim 1, wherein said HGF is a modified form of an HGF selected from granulocyte-macrophage colony-stimulating factor (GM-CSF), interleukin (IL) -2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-10, G-CSF, erythropoietin (EPO).
3. A modified HGF according to claim 2 wherein said HGF is a modified form of GM-CSF.
4. A modified HGF according to claim 1 or 2 or 3 wherein said HGF is of human, livestock animal, companion animal or laboratory test animal origin.
5. A modified HGF according to claim 4 wherein said HGF is of human origin.
6. A modified HGF according to any one of the preceding claims wherein the acidic amino acid residue on the first α -helix is Glu and/or Asp and the basic amino acid residue substituted therefor is Arg and/or Lys.
7. A modified HGF comprising an amino acid sequence in the first α -helix of said HGF selected from the group consisting of:
 - i) His Val Asn Ala Ile Gln Xaa Ala Arg Arg Leu Leu Asn Leu;
 - ii) Ala Leu Val Lys Xaa Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu;
 - iii) Asn Met Ile Xaa Xaa Ile Ile Thr His Leu;
 - iv) Leu Leu Leu Xaa Leu Gln Met Ile Leu;
 - v) Ile Thr Leu Gln Xaa Ile Ile Lys Thr Leu;



- vi) Arg Tyr Ile Leu Xaa Gly Ile Ser Ala Leu Arg Lys;
- vii) Gly Asp Gln Tyr Xaa Ser Val Leu Met Val Ser Ile;
- viii) Ala Gly Ile Leu Xaa Ile Asn Phe Leu Ile Asn Lys Met Gln
Glu Asp;
- ix) Asn Met Leu Arg Xaa Leu Arg Asp Ala Phe Ser;
- x) Phe Leu Leu Lys Cys Leu Xaa Gln Val Arg Lys Ile;
- xi) Tyr Leu Leu Glu Ala Lys Xaa Ala Glu Asn Ile Thr Thr Gly;

wherein Xaa is a basic amino acid, selected from the group consisting of Arg and Lys and wherein said modified HGF is in unglycosylated form and acts as an antagonist for at least one property of the corresponding native HGF.

- 8. A modified HGF according to claim 7 wherein the HGF is a modified form of an HGF selected from the list consisting of GM-CSF, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-10, G-CSF, EPO.
- 9. A modified HGF according to claim 8, wherein the variant HGF is a modified form of GM-CSF.
- 10. A modified HGF according to claim 7 or 8 wherein Xaa is Arg.
- 11. A modified human GM-CSF in unglycosylated form having an amino acid sequence in a first α -helix comprising His Val Asn Ala Ile Gln Arg Ala Arg Arg Leu Leu Asn Leu.
- 12. A modified human GM-CSF in unglycosylated form having an amino acid sequence in a first α -helix comprising His Val Asn Ala Ile Gln Lys Ala Arg Arg Leu Leu Asn Leu.
- 13. A modified human IL-5 in unglycosylated form having an amino acid sequence in a first α -helix comprising
Ala Leu Val Lys Lys Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu.



14. A modified human IL-5 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Ala Leu Val Lys Arg Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu.

15. A modified human IL-3 in unglycosylated form having an amino acid sequence in a first α -helix selected from the list consisting of:

Asn Met Ile Asp Lys Ile Ile Thr His Leu;

Asn Met Ile Lys Glu Ile Ile Thr His Leu;

Asn Met Ile Asp Arg Ile Ile Thr His Leu;

Asn Met Ile Arg Glu Ile Ile Thr His Leu;

Asn Met Ile Lys Lys Ile Ile Thr His Leu; and

Asn Met Ile Arg Arg Ile Ile Thr His Leu.

16. A modified human IL-2 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Leu Leu Leu Lys Leu Gln Met Ile Leu.

17. A modified human IL-2 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Leu Leu Leu Arg Leu Gln Met Ile Leu.

18. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Ile Thr Leu Gln Lys Ile Ile Lys Thr Leu.

19. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Ile Thr Leu Gln Arg Ile Ile Lys Thr Leu.





20. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Arg Tyr Ile Leu Lys Gly Ile Ser Ala Leu Arg Lys.

21. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Arg Tyr Ile Leu Arg Gly Ile Ser Ala Leu Arg Lys.

22. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Gly Asp Gln Tyr Lys Ser Val Leu Met Val Ser Ile.

23. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Gly Asp Gln Tyr Arg Ser Val Leu Met Val Ser Ile.

24. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Ala Gly Ile Leu Lys Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp.

25. A modified human IL-4 having an amino acid sequence in a first α -helix comprising

Ala Gly Ile Leu Arg Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp.

26. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Asn Met Leu Arg Lys Leu Arg Asp Ala Phe Ser.

27. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Asn Met Leu Arg Arg Leu Arg Asp Ala Phe Ser.



28. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Phe Leu Leu Lys Cys Leu Lys Gln Val Arg Lys Ile.

29. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Phe Leu Leu Lys Cys Leu Arg Gln Val Arg Lys Ile.

30. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Tyr Leu Leu Glu Ala Lys Lys Ala Glu Asn Ile Thr Thr Gly.

31. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Tyr Leu Leu Glu Ala Lys Arg Ala Glu Asn Ile Thr Thr Gly.

32. A method of ameliorating the aberrant effects of an endogenous HGF in a mammal, said method comprising administering to said mammal an effective amount of a modified HGF characterised by being in unglycosylated form and comprising a sequence of amino acids within a first α -helix wherein one or more exposed amino acids in said first α -helix having acidic or acidic-like properties are substituted with a basic amino acid residue.

33. A method according to claim 32 wherein the modified HGF is selected from the group consisting of:

- i) His Val Asn Ala Ile Gln Xaa Ala Arg Arg Leu Leu Asn Leu;
- ii) Ala Leu Val Lys Xaa Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu;
- iii) Asn Met Ile Xaa Xaa Ile Ile Thr His Leu;
- iv) Leu Leu Leu Xaa Leu Gln Met Ile Leu;
- v) Ile Thr Leu Gln Xaa Ile Ile Lys Thr Leu;
- vi) Arg Tyr Ile Leu Xaa Gly Ile Ser Ala Leu Arg Lys;
- vii) Gly Asp Gln Tyr Xaa Ser Val Leu Met Val Ser Ile;



- viii) Ala Gly Ile Leu Xaa Ile Asn Phe Leu Ile Asn Lys Met Gln
Glu Asp;
- ix) Asn Met Leu Arg Xaa Leu Arg Asp Ala Phe Ser;
- x) Phe Leu Leu Lys Cys Leu Xaa Gln Val Arg Lys Ile; and
- xi) Tyr Leu Leu Glu Ala Lys Xaa Ala Glu Asn Ile Thr Thr Gly;

wherein Xaa is a basic amino acid selected from the group consisting of Arg and Lys and wherein said variant HGF is in unglycosylated form and acts as an antagonist for at least one property of the corresponding native HGF.

34. A method according to claim 32 or 33 wherein the HGF is a modified form of an HGF selected from GM-CSF, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-10, G-CSF, EPO.

35. A method according to claim 32 or 33 wherein the modified HGF is a modified human GM-CSF having an amino acid sequence in a first α -helix comprising His Val Asn Ala Ile Gln Arg Ala Arg Arg Leu Leu Asn Leu.

36. A method according to claim 32 or 33 wherein the modified HGF is a modified human GM-CSF having an amino acid sequence in a first α -helix comprising His Val Asn Ala Ile Gln Lys Ala Arg Arg Leu Leu Asn Leu.

37. Use of one or more modified HGFs each as defined in claim 1 or 7 in the manufacture of a medicament for the treatment of the affects of an aberrant endogenous HGF.

38. An agent comprising one or more modified HGFs each as defined in claim 1 or 7 for treating the affects of an aberrant endogenous HGF.

39. A pharmaceutical composition comprising one or more modified HGFs each as defined in claim 1 or 7 together with one or more pharmaceutically acceptable carriers and/or diluents.



This Sheet Has Been Replaced



- 43 -

THIS SHEET HAS BEEN
REPLACED

ABSTRACT

HAEMOPOIETIC GROWTH FACTOR ANTAGONISTS

The present invention relates to modified and variant forms of haemopoietic growth factors capable of acting as antagonists to the corresponding native haemopoietic growth factors and their use in ameliorating aberrant effects caused by the native molecules.

(No figure to accompany the abstract)




FIGURE 1




FIGURE 2



**COMPETITION OF Wt-GMCSF AND E21R FOR LOW
AFFINITY RECEPTORS (1nM)****COMPETITION OF Wt-GMCSF AND E21R FOR HIGH
AFFINITY RECEPTORS (100pM)**

FIGURE 3

FIG 4A




FIG 4B




FIG 5A




FIG 5B




FIG 5C




FIG 6A




FIG 6B




FIG 6C




FIG 7A





FIG 7B


Box III

TEXT OF THE ABSTRACT (Continuation of Item 5 of the first sheet)

The present invention relates to modified and variant forms of haemopoietic growth factors (HGF) capable of acting as antagonists to the corresponding native haemopoietic growth factors and their use in ameliorating aberrant effects caused by the native molecules.

A modified haemopoietic growth factor (HGF) is characterised by being in unglycosylated form and comprising a sequence of amino acids within a first α -helix wherein one or more exposed amino acids in said first α -helix having acidic or acidic-like properties are substituted with a basic amino acid residue.

The preferred HGF are granulocyte-macrophage colony-stimulating factor (GM-CSF), interleukin (IL)-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-10, G-CSF and erythropoietin (EPO).





20. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Arg Tyr Ile Leu Lys Gly Ile Ser Ala Leu Arg Lys.

21. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Arg Tyr Ile Leu Arg Gly Ile Ser Ala Leu Arg Lys.

22. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Gly Asp Gln Tyr Lys Ser Val Leu Met Val Ser Ile.

23. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Gly Asp Gln Tyr Arg Ser Val Leu Met Val Ser Ile.

24. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Ala Gly Ile Leu Lys Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp.

25. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Ala Gly Ile Leu Arg Ile Asn Phe Leu Ile Asn Lys Met Gln Glu Asp.

26. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Asn Met Leu Arg Lys Leu Arg Asp Ala Phe Ser.

27. A modified human IL-4 in unglycosylated form having an amino acid sequence in a first α -helix comprising

Asn Met Leu Arg Arg Leu Arg Asp Ala Phe Ser.

REPLACEMENT SHEET

9 JUN 1998