NASAL ADMINISTRATION OF AGENTS FOR TREATMENT OF DELAYED ONSET EMESIS

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention is directed to a method for treating an emetogenic reaction. More particularly, the present invention is directed to a method for treating a delayed onset emetogenic reaction, most typically associated with chemotherapy.

Description of the Related Art

Emetogenic reaction refers to vomiting (*i.e.* actual vomiting or dry heaving or dry retching) and/or to nausea. The emetogenic reaction is most commonly encountered in response to chemotherapeutic agents such as bleomycin, vincristine, vinblastine, adriamycin, etoposide, methotrexate, doxorubicin, cyclophosphamide, 5-fluorouracil, cisplatin and some combinations thereof. Among these chemotherapeutic agents, the most severe emetogenic reaction is usually associated with cisplatin therapy.

Generally, there are acute onset emetogenic reactions (hereinafter synonymous with "acute emesis" or "acute onset emesis") and delayed onset emetogenic reactions (hereinafter synonymous with "delayed emesis" or "delayed onset emesis"). Acute emesis and delayed emesis typically include vomiting episodes (actual and/or dry heaving or dry retching) and/or feelings of nausea. However, in acute emesis, the vomiting and/or nausea occur within the first 24 hours after

chemotherapy. By contrast, in delayed emesis, the vomiting and/or nausea occur after the first 24 hours of chemotherapy. Delayed emesis is a distinct syndrome from acute emesis occurring more than 24 hours after the administration of anticancer agents, especially, cisplatin.

In a study by Kris et al. entitled Controlling Delayed Vomiting: Double-Blind, Randomized Trial Comparing Placebo, Dexamethasone Alone, and Metoclopramide Plus Dexamethosone in Patients Receiving Cisplatin published in the *Journal of Clinical Oncology*, Vol. 7, No. 1, at pp. 108-114 (January 1989), the authors report that delayed vomiting was observed in a 4 day period post cisplatin therapy (at a dosage level of 120mg/m^2 of body surface area) in 74% of patients and that delayed nausea was observed in 87% of the same patients over the same 4-day post cisplatin period. Kris et al. further reported that the severity and incidence of delayed emesis was most prevalent (*i.e.* worst) during the period of 48 - 72 hours (*i.e.* days 2 - 3 post cisplatin administration) after cisplatin therapy was given. Further, according to Kris et al.

> As the control of acute emesis following high doses of cisplatin has improved, the need to address the remaining problem of delayed nausea and vomiting [i.e. delayed emesis] beginning or persisting for more than 24 hours after cisplatin has become apparent. (Emphasis added.) Id at p. 113.

The patients in the Kris et al. study received a two drug regimen of oral dexamethasone (*i.e.* DXM po) and oral metoclopramide (i.e. MCP po). Side effects observed in the Kris et al. study patients included hiccoughs, loose bowel

movements (LBMs), heartburn, restlessness, sleepiness, insomnia and acute dystonic reactions. Among these symptoms, restlessness and sleepiness were the most common.

In another study, the side effects of oral MCP for the prophylaxis of delayed emesis (*i.e.* delayed nausea and/or vomiting) induced by cisplatin were reported by Grunberg et al., in Oral Metoclopramide With or Without Diphenhydramine: Potential for Prevention of Late Nausea and Vomiting Induced by Cisplatin, published in the *Journal of the National Cancer Institute*, Vol. 80, No. 11, pp. 864-868 (August 3, 1988). The most severe toxic effects observed were extrapyramidal symptoms, agitation and depression.

Side effects may interfere with patient **compliance** (compliance = following dosage regimen prescribed) with the drug regimen prescribed as well as interfere with the patient's ability to effectively communicate the nature and severity of this and other side effects. Even with a short-term medication regimen, poor compliance or non-compliance is observed in about 25% of patients.

Additionally, oral therapy may be unsuitable for some patients experiencing a delayed emetogenic reaction. Due to nausea and/or vomiting, the patient may be even more reluctant to comply with the oral drug regimen. Therefore, there is a need to provide treatment or prophylaxis (*i.e.* complete or partial prophylaxis) for delayed emesis in an appropriate dosage form wherein the problems associated with non-compliance (*e.g.* with oral or intravenous dosage forms) are avoided.

U.S. Patent No. 4,624,965 (hereinafter Wenig) discusses nasal administration of MCP. In Wenig, at column 1, lines 40 - 52, it is stated that

> A number of antinausea and antiemetic agents are already known. Such agents are widely used therapeutically, chiefly in the treatment of emesis and nausea . . . . Unfortunately, many of these agents when used: (1) cause undesirable side effects, (2) are inefficiently and variably absorbed from current dosage forms, (3) are difficult or inconvenient to administer in the current dosage forms [*e.g.* oral, intravenous, intramuscular or subcutaneous dosage forms] . . . . (Emphasis added.)

Further, in Examples 8 and 9 of Wenig (See columns 11 and 12 therein), human subjects are studied. However, a gel intranasal formulation is used as opposed to a nasal spray. No experience with human subjects using a nasal spray formulation of MCP (MCP ns) is disclosed within Wenig.

Since chronic or severe nausea and/or vomiting can significantly influence a patient's ability to maintain an acceptable level of health, for example, after chemotherapy, it is desirable to provide a method for treating and sufficiently controlling (prophylactically controlling) such emetogenic reactions. In the case of chemotherapy-induced delayed emesis, the control and treatment of the nausea and/or vomiting can not only help to maintain an acceptable level of health but can also aid in maintaining an aggressive cancer treatment program.

To that end, while advances have been made in the area of controlling and treating acute emesis, treatment for

adequate control of delayed emesis was heretofore not available.

Summary of the Invention

It is therefore an object of the present invention to provide a method for treating delayed emesis by using a dosage form of MCP that avoids or reduces the incidence of patient non-compliance.

It is therefore another object of the present invention to provide a method for treating delayed emesis by using MCP nasal spray that avoids or reduces the incidence of side-effects experienced by patients.

It is therefore yet another object of the present invention to provide a method for treating delayed emesis by using MCP nasal spray administered intranasally.

It the therefore even yet another object of the present invention to provide a method for treating delayed emesis induced by chemotherapy by using a dosage form of MCP that avoids or reduces the problem of patient non-compliance.

It is still another object of the present invention to provide a method for treating delayed emesis induced by cisplatin chemotherapy by using a dosage form of MCP that avoids or reduces the problem of patient non-compliance.

It is yet still another object of the present invention to provide a method for sufficiently treating delayed emesis induced by chemotherapy using a dosage form of MCP that avoids or reduces the incidence of side-effects experienced by patients.

It is even yet still another object of the present invention to provide a method for sufficiently treating delayed emesis induced by cisplatin chemotherapy using MCP nasal spray that avoids or reduces the severity of side-effects experienced by patients.

It is a further object of the present invention to provide a method for controlling delayed emesis induced by chemotherapy using MCP nasal spray administered intranasally after the first 24 hours of completing chemotherapy.

It is an even further object of the present invention to provide a method for sufficiently controlling delayed emesis induced by cisplatin chemotherapy using MCP nasal spray that avoids or reduces the problems associated with patient non-compliance.

It is an even yet further object of the present invention to provide a method for prophylaxis of delayed emesis by intranasally administering a pharmaceutically acceptable MCP nasal spray dosage form at a therapeutic dosage level of between about 1.95 mg/kg to about 3.9 mg/kg on days 1-7 wherein chemotherapy was administered on day 0.

It is still another object of the present invention to provide a method for prophylaxis of delayed emesis by intranasally administering a pharmaceutically acceptable MCP nasal spray dosage form at a therapeutic dosage level of between about 1.95 mg/kg to about 3.9 mg/kg on days 1-6 wherein cisplatin was administered.

It is still another object of the present invention to provide a method for prophylaxis of delayed emesis by

intranasally administering a pharmaceutically acceptable MCP nasal spray dosage form at a therapeutic dosage level of between about 0.8 mg/kg to about 1.0 mg/kg on days 1-5 wherein cisplatin was administered.

It is still another object of the present invention to provide a method for prophylaxis of delayed emesis by intranasally administering a pharmaceutically acceptable MCP nasal spray dosage form at a therapeutic dosage level of between about 0.9 mg/kg to about 1.1 mg/kg on days 1-4 wherein cisplatin was administered.

It is still another object of the present invention to provide a method for prophylaxis of delayed emesis by intranasally administering a pharmaceutically acceptable MCP nasal spray at a therapeutic dosage level of between about 40 mg/day - 120 mg/day on days 1-7 wherein chemotherapy was administered by 0.

It is still another object of the present invention to provide a method for prophylaxis of delayed emesis by intranasally administering a pharmaceutically acceptable MCP nasal spray at a therapeutic dosage level of between about 40 mg/day - 120 mg/day on days 1-6 wherein cisplatin was administered on day 0.

It is still another object of the present invention to provide a method for prophylaxis of delayed emesis by intranasally administering a pharmaceutically acceptable MCP nasal spray at a therapeutic dosage level of about 60 mg/day on days 1-6 wherein chemotherapy was administered on day 0.

It is still another object of the present invention to provide a method for prophylaxis of delayed emesis by intranasally administering a pharmaceutically acceptable MCP nasal spray at a therapeutic dosage level of about 60 mg/day on days 1-6 wherein cisplatin was administered on day 0.

It is still another object of the present invention to provide a method for prophylaxis of delayed emesis by intranasally administering a pharmaceutically acceptable MCP nasal spray at a therapeutic dosage level of about 60 mg/day on days 1-5 wherein cisplatin was administered on day 0.

It is still another object of the present invention to provide a method for prophylaxis of delayed emesis by intranasally administering a pharmaceutically acceptable MCP nasal spray at a therapeutic dosage level of about 60 mg/day on days 1-4 wherein cisplatin was administered on day 0.

These and other objects of the present invention are accomplished by administering intranasally to patients experiencing delayed emesis a therapeutically effective dosage of MCP nasal spray in a pharmaceutically acceptable dosage form which is therapeutically and medically acceptable.

Detailed Description of the Preferred Embodiments

The presently disclosed invention is directed to a method for treating and controlling delayed emesis by using a MCP nasal spray dosage formulation. The MCP nasal spray dosage form is formulated to contain a therapeutically effective amount of MCP such that upon administration by the intranasal route, a therapeutically effective amount of MCP

is delivered to the patient. In addition, the therapeutically effective amount of MCP is chosen to minimize the severity and incidence of untoward side-effects and drug-interactions encountered with MCP.

The MCP nasal spray formulation administered to deliver a dose of 20mg three times a day is indicated below (formulation per 0.1 ml of MCP nasal spray (MCP ns = metoclopramide nasal spray dosage form)):

10mg/0.1ml	metoclopramide hydrochloride
1.5 mg	benzyl alcohol
0.8 mg	NaCl
0.320 mg	glacial acetic acid
0.077 mg	sodium acetate
6.425 mg	sorbitol
0.1 ml	purified water (qs ad to 0.1 ml).

The MCP nasal spray formulation was given (to the patients indicated in Figures 1(a) and 1(b)) as 1 puff per nostril (*i.e.* 2 puffs at 10mg/puff (10mg/0.1ml and 0.1ml/puff)) three times a day (2 puffs TID for 6 days). The above formulation was sterile with a bacteria count of 10 below the level allowed by the U.S.P. on a per ml basis. In addition, pathogens were absent. The pH of the above formulation was about 4.0.

Typical MCP nasal spray dosage forms are solutions or suspensions that can be administered as a nasal spray. However, nasal drops may also be used. The MCP nasal spray dosage formulation contains the active agent in any suitable form *e.g.* salt, as the hydrochloride etc. The MCP nasal

spray dosage formulation typically also contains pH adjusters, emulsifiers or dispersing agents, buffering agents, preservatives and wetting agents as are known to those skilled in the art. See REMINGTON'S PHARMACEUTICAL SCIENCES, 14th Edition, 1970. Typically, the MCP nasal spray dosage formulation is isotonic.

A typical MCP nasal spray formulation is in solution form having a light amber color and being non-cloudy to the naked eye with an pH of between about 3.0 - 5.0. The typical formulation may contain benzyl alcohol of at least about 13.5 mg/ml containing practically no impurities as determined by high pressure liquid chromatography (HPLC) and having a bacterial count of less than 250 ufc/ml and free of pathogens sufficient to form an acceptable pharmaceutical nasal spray dosage form. The solvent is typically purified water suitable for use in nasal spray dosage forms or any equivalent water (e.g. injectable water) that is allowed for use in such nasal dosage forms. See REMINGTON'S PHARMACEUTICAL SCIENCES, any edition from 1980 - 1996. For the adequate and/or sufficient treatment and control of delayed emesis (e.g. chemotherapy induced, cisplatin induced or induced by other causes), a typical dose is that dose which is therapeutically effective and which minimizes side-effects and drug interactions, for example, in patients receiving chemotherapy.

A typical dosage of MCP nasal spray for the treatment and control of delayed onset emesis depends upon the degree and severity of delayed emesis experienced by a typical

patient (e.g. chemotherapy induced delayed emesis). The dosage of MCP nasal spray may be varied between about 40 mg/day to about 120 mg/day. Above about 120 mg/day, the dosage may be undesirable due to untoward side effects experienced by patients receiving more than about 120 mg/day from the MCP nasal spray dosage form. A preferred dosage of MCP nasal spray is 60 mg/day typically given as 20mg three times a day (i.e. 2 puffs of 10 mg/0.1 ml of MCP nasal spray, one puff per nostril).

The weight of the patient may also affect the dosage to be administered. Typically, a dose of between about 1.95 mg/kg to about 3.90 mg/kg is given to an patient experiencing delayed emesis (e.g. chemotherapy induced delayed emesis). A preferred dosage is about 0.8 mg/kg.

The aforementioned dosages for the treatment and control of delayed emesis are given after 24 hours from the time that the chemotherapeutic regimen cycle has been administered. Thus, for example, if the chemotherapeutic regimen cycle is given between 2:00 to 4:00 pm on day zero, then the MCP nasal spray is administered starting at about 4:00 pm on day 1 - day 7, typically, in three or four equal dosages at three or four equal time intervals, respectively. However, the dosage may be given during waking hours so as not to wake the patient. Note that the MCP nasal spray dosage form is administered intranasally and is given in addition to medications administered for the treatment and control of acute emesis. All references cited within this patent

application are incorporated herein by reference in their entirety and for all purposes.

Having described the invention, the following examples are included to illustrate the benefits of the present invention. The examples are only illustrative and are not meant to unduly limit the scope of the present invention.

EXAMPLES

Example 1

Table I indicates the characteristics of patients that received metoclopramide nasal spray alone or metoclopramide nasal spray (MCP ns) and dexamethasone (DXM) for controlling delayed emeses induced by chemotherapy. Primary disease site indicates primary location of cancer.

TABLE I - PATIENTS' CHARACTERISTICS

Number of patients	12
Sex	
Male	2
Female	10
Age (years)	
Mean	50
Range	23-63
Primary disease site	
Ovary	8 patients
Testis	1 patient
Bladder	3 patients

Table II indicates the chemotherapy regimen that each of the patients received on day 0.

TABLE II - CHEMOTHERAPY REGIMEN (administered on day 0)

Cisplatin	1 patient
Cisplatin + Cyclophosphamide	5 patients
Cisplatin + Cyclophosphamide + Adriamycin	3 patients
Cisplatin + Cyclophosphamide + Vinblastine + Methotrexate	2 patients
Cisplatin + Etoposide + Bleomycin	1 patient

Example 2

Table III indicates the vomiting, retching and nausea reported in each patient during days 1 through 6 following chemotherapy on day 0. Note that MCP nasal spray (60mg/day) indicates administration of a MCP nasal spray formulation per 0.1 ml of:

10mg/0.1ml	metoclopramide hydrochloride
1.5 mg	benzyl alcohol
0.8 mg	NaCl
0.320 mg	glacial acetic acid
0.077 mg	sodium acetate
6.425 mg	sorbitol
0.1 ml	purified water (qs to 0.1 ml).

The MCP nasal spray formulation was given as 1 puff per nostril (i.e. 2 puffs at 10mg/puff (10mg/0.1ml and 0.1ml/puff)) three times a day (2 puffs TID for 6 days). The above formulation was sterile for bacteria. In addition, pathogens were absent. The pH of the above formulation was about 4.0.

TABLE III - VOMITING, RETCHING AND NAUSEA REPORTED IN EACH PATIENT DURING DAYS 1 THROUGH 6 FOLLOWING CHEMOTHERAPY (Chemotherapy administered on day 0)

TREATMENT	PATIENT NO.	VOMITING No. of emetic episodes/day						RETCHING No. of episodes/day						NAUSEA Intensity*/day					
		DAY						DAY						DAY					
		1	2	3	4	5	6	1	2	3	4	5	6	1	2	3	4	5	6
MCP ns (60mg/day)	1	0	3	0	0	0	0	0	1	1	0	0	0	1	2	2	1	1	0
	3	0	0	0	0	0	0	0	0	0	0	0	0	3	2	3	3	2	2
	4	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0
	5	0	6	3	4	2	0	0	0	0	0	0	0	0	2	2	2	2	2
	6	0	1	4	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1
MCP ns (60mg/day) + DXM (8mg/day)	7	10	10	9	10	10	10	0	0	0	0	0	0	3	3	3	3	2	2
	8	0	0	0	1	0	0	0	0	0	0	0	0	1	1	1	3	1	1
	9	2	3	1	0	0	0	0	0	0	0	0	0	2	2	2	2	1	1
	10	2	3	3	4	3	2	0	1	2	0	0	0	1	1	2	2	3	3
	11	0	1	0	0	0	0	0	0	0	0	0	0	1	1	1	0	0	0
	12	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0

* 1=mild, 2=moderate, 3=severe

Patient No. 2 was not included because he received fractionation of chemotherapy, he was treated with MCP ns and he had no vomiting retching or nausea (complete response)

Example 3

The results of MCP nasal spray therapy (given on day 1-6) for the treatment and control (partial or complete prophylaxis) of delayed emesis is indicated in Table IV.

TABLE IV(a) - COMPLETE RESPONSE (no emetic episodes) BY STUDY DAY

COMPLETE RESPONSE† DAY	MCP ns number of patients	MCP ns + DXM number of patients
1	5/5	3/6
2	1/5	2/6
3	3/5	3/6
4	4/5	3/6
5*	8/11	---
6*	9/11	---

Thus, on day 1 all 5 of 5 patients that received MCP ns had no emetic episodes (complete response) while only 3 of 6 patients that received MCP ns + DXM showed no emetic episodes.

TABLE IV(b) - MAJOR RESPONSE (1 - 2 emetic episodes)

MAJOR RESPONSE‡ DAY	MCP ns number of patients	MCP ns + DXM number of patients
1	---	2/6
2	2/5	1/6
3	---	1/6
4	---	1/6
5*	1/11	---
6*	1/11	---

† = complete response = 0 emetic episodes

‡ = major response = 1-2 emetic episodes

* = on days 5 and 6 all patients received only MCP ns (60 mg/day); and --- = 0.

Similarly on day 1, no patients that received MCP ns had any emetic episodes (major response) while 2 of 6 patients who received MCP ns + DXM had 1-2 emetic episodes.

This Sheet Has Been Replaced

EET HAS BEEN PLACED

CLAIMS

What is claimed is:

1. A method for prophylaxis of delayed emesis comprising the step of:

administering a pharmaceutically acceptable metoclopramide nasal spray formulation via an intranasal route to deliver a therapeutically effective and medically acceptable daily dosage for said prophylaxis.

2. The process of claim 1 wherein said daily dosage is between about 40 mg/day to about 120 mg/day.

3. The process of claim 2 wherein said daily dosage is between 40 mg/day to 120 mg/day.

4. The process of claim 2 wherein said daily dosage is about 60 mg/day.

5. The process of claim 4 wherein said daily dosage is 60 mg/day.

6. The process of claim 1 wherein said daily dosage is divided into 3 or 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

7. The process of claim 6 wherein said smaller doses are 20 mg and administered every 8 hours.

8. The process of claim 1 wherein said daily dosage is between about 1.95 mg/kg to about 3.90/kg.

9. The process of claim 8 wherein said daily dosage is between 1.95 mg/kg to 3.90 mg/kg.

10. The process of claim 9, wherein said daily dosage is about 0.8 mg/kg.

* C L M S *

11. The process of claim 10 wherein said daily dosage is 0.8 mg/kg.

12. The process of claim 11 wherein said daily dosage is divided into 3 or 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

13. The process of claim 12 wherein said daily dosage is divided into 3 equal smaller doses and administered at equally spaced intervals within 24 hours.

14. The process of claim 12 wherein daily dosage is divided into 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

15. The process of claim 1 wherein said dosage is administered for prophylaxis of delayed emesis induced by chemotherapy.

16. The process of claim 15 wherein said dosage is induced by chemotherapeutic agents selected from the group consisting of cisplatin, cyclophosphamide, adriamycin, vinblastine, methotrexate, etoposide, bleomycin, vincristine and combinations thereof.

17. The process of claim 16 wherein said daily dosage is administered during 24-168 hours after completion of said chemotherapy.

18. The process of claim 16 wherein said daily dosage is administered during 24-144 hours after completion of said chemotherapy.

19. The process of claim 16 wherein said daily dosage is administered during 24-120 hours after completion of said chemotherapy.

* C L M S *

THIS SHEET HAS BEEN REPLACED

20. The process of claim 16 wherein said daily dosage is administered during 24-96 hours after completion of said chemotherapy.

This Sheet Has Been Replaced





AMENDED CLAIMS

[received by the International Bureau on 14 August 1997 (14.08.97); original claim 8 amended; remaining claims unchanged (1 page)]

What is claimed is:

1. A method for prophylaxis of delayed emesis comprising the step of:

administering a pharmaceutically acceptable metoclopramide nasal spray formulation via an intranasal route to deliver a therapeutically effective and medically acceptable daily dosage for said prophylaxis.

2. The process of claim 1 wherein said daily dosage is between about 40 mg/day to about 120 mg/day.

3. The process of claim 2 wherein said daily dosage is between 40 mg/day to 120 mg/day.

4. The process of claim 2 wherein said daily dosage is about 60 mg/day.

5. The process of claim 4 wherein said daily dosage is 60 mg/day.

6. The process of claim 1 wherein said daily dosage is divided into 3 or 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

7. The process of claim 6 wherein said smaller doses are 20 mg and administered every 8 hours.

8. The process of claim 1 wherein said daily dosage is between about 1.95 mg/kg to about 3.90 mg/kg.

9. The process of claim 8 wherein said daily dosage is between 1.95 mg/kg to 3.90 mg/kg.

10. The process of claim 9, wherein said daily dosage is about 0.8 mg/kg.

* C L M S *

This Sheet Has Been Replaced

* C L M S *

CLAIMS

What is claimed is:

1. A composition for prophylaxis of delayed emesis comprising a pharmaceutically acceptable metoclopramide compound in a nasal spray formulation for administration of the compound via an intranasal route to deliver a therapeutically effective and medically acceptable daily dosage for said prophylaxis.

2. The composition of claim 1 wherein said daily dosage is between about 40 mg/day to about 120 mg/day.

3. The composition of claim 2 wherein said daily dosage is between 40 mg/day to 120 mg/day.

4. The composition of claim 2 wherein said daily dosage is about 60 mg/day.

5. The composition of claim 4 wherein said daily dosage is 60 mg/day.

6. The composition of claim 1 wherein said daily dosage is divided into 3 or 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

7. The composition of claim 6 wherein said smaller doses are 20 mg and administered every 8 hours.

8. The composition of claim 1 wherein said daily dosage is between about 1.95 mg/kg to about 3.90/kg.

9. The composition of claim 8 wherein said daily dosage is between 1.95 mg/kg to 3.90 mg/kg.

10. The composition of claim 9, wherein said daily dosage is about 0.8 mg/kg.

REPLACED SHEETS SG
☐ Description : paged ..... to paged .....
☑ Claim(s) : paged ..... to paged .....
☐ Drawing(s) : paged ..... to paged .....
☐ Abstract : paged ..... to paged .....

REPLACEMENT SHEETS SG
☐ Description : paged ..... to paged .....
☑ Claim(s) : paged ..... to paged .....
☐ Drawing(s) : paged ..... to paged .....
☐ Abstract : paged ..... to paged .....
DATE OF RECEIPT 16/3/99

This Sheet Has Been Replaced

11. The composition of claim 10 wherein said daily dosage is 0.8 mg/kg.

12. The composition of claim 11 wherein said daily dosage is divided into 3 or 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

13. The composition of claim 12 wherein said daily dosage is divided into 3 equal smaller doses and administered at equally spaced intervals within 24 hours.

14. The composition of claim 12 wherein daily dosage is divided into 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

15. The composition of claim 1 wherein said dosage is administered for prophylaxis of delayed emesis induced by chemotherapy.

16. The composition of claim 15 wherein said dosage is induced by chemotherapeutic agents selected from the group consisting of cisplatin, cyclophosphamide, adriamycin, vinblastine, methotrexate, etoposide, bleomycin, vincristine and combinations thereof.

17. The composition of claim 16 wherein said daily dosage is administered during 24-168 hours after completion of said chemotherapy.

18. The composition of claim 16 wherein said daily dosage is administered during 24-144 hours after completion of said chemotherapy.

19. The composition of claim 16 wherein said daily dosage is administered during 24-120 hours after completion of said chemotherapy.

This is the replacement sheet

* C L M S *

Date: 16 MAR 1999

20. The composition of claim 16 wherein said daily dosage is administered during 24-96 hours after completion of said chemotherapy.

This Sheet Has Been Replaced

* C L M S *

This is the replacement sheet
Date: 16 MAR 1999

CLAIMS

What is claimed is:

1. A composition for prophylaxis of delayed emesis comprising a pharmaceutically acceptable metoclopramide compound in a nasal spray formulation for administration of the compound via an intranasal route to deliver a therapeutically effective and medically acceptable daily dosage for said prophylaxis.

2. The composition of claim 1 wherein said daily dosage is between about 40 mg/day to about 120 mg/day.

3. The composition of claim 2 wherein said daily dosage is between 40 mg/day to 120 mg/day.

4. The composition of claim 2 wherein said daily dosage is about 60 mg/day.

5. The composition of claim 4 wherein said daily dosage is 60 mg/day.

6. The composition of claim 1 wherein said daily dosage is divided into 3 or 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

7. The composition of claim 6 wherein said smaller doses are 20 mg and administered every 8 hours.

8. The composition of claim 1 wherein said daily dosage is between about 1.95 mg/kg to about 3.90/kg.

9. The composition of claim 8 wherein said daily dosage is between 1.95 mg/kg to 3.90 mg/kg.

10. The composition of claim 9, wherein said daily dosage is about 0.8 mg/kg.

REPLACEMENT SHEETS SG
☐ Description : paged ..... to paged .....
☑ Claim(s) : paged 17 to paged 19
☐ Drawing(s) : paged ..... to paged .....
☐ Abstract : paged ..... to paged .....
DATE OF RECEIPT 17/11/99

* C L M S *

11. The composition of claim 10 wherein said daily dosage is 0.8 mg/kg.

12. The composition of claim 11 wherein said daily dosage is divided into 3 or 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

13. The composition of claim 12 wherein said daily dosage is divided into 3 equal smaller doses and administered at equally spaced intervals within 24 hours.

14. The composition of claim 12 wherein daily dosage is divided into 4 equal smaller doses and administered at equally spaced intervals within 24 hours.

15. The composition of claim 1 wherein said dosage is administered for prophylaxis of delayed emesis induced by chemotherapy.

16. The composition of claim 15 wherein said dosage is induced by chemotherapeutic agents selected from the group consisting of cisplatin, cyclophosphamide, adriamycin, vinblastine, methotrexate, etoposide, bleomycin, vincristine and combinations thereof.

17. The composition of claim 16 wherein said daily dosage is administered during 24-168 hours after completion of said chemotherapy.

18. The composition of claim 16 wherein said daily dosage is administered during 24-144 hours after completion of said chemotherapy.

19. The composition of claim 16 wherein said daily dosage is administered during 24-120 hours after completion of said chemotherapy.

* C L M S *

This is the replacement sheet

Date: 17 NOV 1999

20. The composition of claim 16 wherein said daily dosage is administered during 24-96 hours after completion of said chemotherapy.

* C L M S *

This is the replacement sheet

Date: 17 NOV 1999

IPOS Copy

9804933-1

PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 31/165 — A1

(11) International Publication Number: **WO 97/33573**

(43) International Publication Date: 18 September 1997 (18.09.97)

(21) International Application Number: PCT/US97/03974

(22) International Filing Date: 13 March 1997 (13.03.97)

(30) Priority Data:
08/616,121 14 March 1996 (14.03.96) US

(71) Applicant: RIBOGENE, INC. [US/US]; 21375 Cabot Boulevard, Hayward, CA 94545 (US).

(72) Inventor: PSILOGENIS, Mary; Via Nazario Sauro, 28, I-22038 Tavernerio (IT).

(74) Agents: POISSANT, Brian, M. et al.; Pennie & Edmonds L.L.P., 1155 Avenue of the Americas, New York, NY 10036 (US).

(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, HU, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published
With international search report.
With amended claims.

(54) Title: NASAL ADMINISTRATION OF AGENTS FOR TREATMENT OF DELAYED ONSET EMESIS

(57) Abstract

The present invention is directed to a method for the prophylactic management of delayed emesis by the use of metoclopramide nasal spray.