

PROCESS FOR PROVIDING PEGYLATED PROTEIN COMPOSITION

Field of the Invention

The present invention relates to processes for providing PEGylated protein compositions, particularly processes for providing PEGylated protein compositions comprising mono-PEGylated protein and oligo-PEGylated protein and processes for providing mono-PEGylated protein compositions. In particular the present invention relates to processes for providing PEGylated erythropoietin (EPO) compositions comprising mono-PEGylated EPO and oligo-PEGylated EPO and processes for providing mono-PEGylated erythropoietin (EPO) compositions.

Background

- PEGylation, or pegylation, of proteins refers to the addition of one or more PEG (polyethylene glycol) groups to a protein. PEGylation is particularly useful for therapeutic proteins, for example because it increases *in vivo* circulation half-life. However, PEGylation may also reduce the biological activity of therapeutic proteins, thereby reducing their effectiveness. There is therefore a balance to be struck between increased circulation time and reduced therapeutic efficacy.
- For certain therapeutic proteins, mono-PEGylation is particularly desirable because it provides improved stability without significantly compromising therapeutic efficacy. Mono-PEGylated therapeutic proteins include Mircera®, PegIntron® and Pegasys®. Micera® is a mono-PEGylated form of erythropoietin (EPO) and is used to treat anaemia.

- PEGylation reactions tend to produce mixtures comprising non-PEGylated protein (unreacted protein), mono-PEGylated protein, and oligo-PEGylated protein. Reaction conditions that favour a high degree of PEGylation (such as long reaction times, high PEG/protein molar ratio) tend to produce mixture with high a proportion of oligo-PEGylated protein, which results in low yields when mono-PEGylated protein is the desired product. Reaction conditions that favour a low degree of PEGylation (such as short reaction times, low PEG/protein molar ratio) tend to produce mixtures with a high proportion of unreacted (non-PEGylated) protein, which also results in low yields when mono-PEGylated protein is the desired product.

Therapeutic proteins are often expensive to manufacture and therefore processes that provide good yields of mono-PEGylated therapeutic protein (minimal waste of therapeutic protein as unreacted and/or oligo-PEGylated) are economically favourable and therefore are particularly desirable.

- Methods of PEGylating a protein of interest while it is bound to an ion exchange chromatography column have been developed in attempts to manipulate the specificity of PEGylation and thereby improve yields of proteins with a desired degree of PEGylation (so-called "on column" methods). Such methods are technically complex and time consuming and consequently have low productivity. Such methods may also consume relatively large amounts of PEGylation reagent, making them economically unfavourable.

Various "on column" PEGylation methods are discussed in Pfister (Reac React. Chem. Eng., 2016, 1,204), Ingold (2016) and Fee & Van Alstine (2006).

Methods of providing mono-PEGylated proteins may comprise performing a PEGylation reaction to provide a mixture comprising non-PEGylated protein (unreacted protein), mono-PEGylated protein, and oligo-PEGylated protein, and then performing a purification step to purify the mono-PEGylated protein.

WO 2009/010270 and WO 2012/035037 relate to methods for purifying mono-PEGylated EPO from a mixture comprising non-PEGylated protein, mono-PEGylated protein, and oligo-PEGylated protein. The purification methods involve subjecting the mixture that results from a PEGylation reaction to at least one cation exchange chromatography (CEC) step. The CEC step is performed in bind and elute mode and different elution fractions comprise mostly non-PEGylated, mono-PEGylated or oligo-PEGylated EPO. Such CEC methods must be carried out below the isoelectric point of the protein, which in the case of EPO (isoelectric point in the range 4.0 to 5.5) involves CEC at a pH of around 3.0.

Methods for producing mono-PEGylated proteins by performing a PEGylation reaction and then purifying the mono-PEGylated protein from the resultant mixture may involve recycling unreacted (non-PEGylated) protein. These methods involve recovering unreacted protein and adding it to a subsequent PEGylation reaction. Such recycling improves the overall yield of mono-PEGylated protein.

Pfister (Biotech and Bioeng, 2016) describes a process in which a PEGylation reaction is performed followed by purification of mono-PEGylated protein from a mixture comprising unreacted (non-PEGylated), mono-PEGylated, and oligo-PEGylated protein. In the process unreacted protein is recovered and used in a subsequent PEGylation reaction. The purification process comprises cation exchange chromatography (CEC) in bind and elute mode, wherein unreacted protein, mono-PEGylated protein and oligo-PEGylated protein are separated in sequential elution. Unreacted protein is eluted last, using a high salt elution buffer. Recycling of unreacted protein therefore requires removal of salt by diafiltration before it can be subject to a further PEGylation reaction. The requirement for removal of salt reduces productivity and increases complexity of the process.

The present invention has been devised in light of the above considerations.

Summary

The present disclosure relates to processes for providing PEGylated protein compositions comprising mono-PEGylated protein and oligo-PEGylated protein and processes for providing mono-PEGylated protein compositions. The processes of the disclosure are particularly suitable for providing PEGylated EPO compositions comprising mono-PEGylated EPO and oligo-PEGylated EPO, as well as mono-PEGylated EPO compositions. Advantages of the process described herein include high yield and productivity.

In a first aspect, the present disclosure provides a process for producing a PEGylated protein mixture the process comprising: (a) reacting a non-PEGylated protein with a PEGylation reagent to produce a mixture of reaction products comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein; (b) subjecting the mixture of reaction products to an ion exchange chromatography (IEC) step to provide an IEC flow-through solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products; the IEC step comprising applying the mixture of reaction products to an IEC material under conditions suitable for binding non-PEGylated protein; and (c) collecting the IEC flow-through solution from step b) to provide a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein, wherein non-PEGylated protein is recovered in step b) by eluting an IEC eluate from the IEC material and the eluted non-PEGylated protein is used in a subsequent PEGylation reaction.

The non-PEGylated protein may be recovered in step b) by elution and used in a subsequent PEGylation reaction. That is, step b) may further comprise eluting non-PEGylated protein in an IEC eluate. The IEC eluate comprises non-PEGylated protein.

The processes may involve two or more cycles of steps (a), (b) and (c), in which non-PEGylated protein is recovered in step b) by eluting an IEC eluate from the IEC material and the eluted non-PEGylated protein is used in a subsequent PEGylation reaction.

The IEC step may be an anion exchange chromatography (AEC) step, or a cation exchange chromatography (CEC) step.

The present disclosure provides a process for producing a PEGylated protein mixture, the process comprising: (a) reacting a non-PEGylated protein with a PEGylation reagent to produce a mixture of reaction products comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein; (b) subjecting the mixture of reaction products to an anion exchange chromatography (AEC) step to provide an AEC flow-through solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products; the AEC step comprising applying the mixture of reaction products to an AEC material under conditions suitable for binding non-PEGylated protein; and (c) collecting the AEC flow-through solution from step b) to provide a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein, wherein non-PEGylated protein is recovered in step b) by eluting an AEC eluate from the AEC material and the eluted non-PEGylated protein is used in a subsequent PEGylation reaction.

The processes may involve two or more cycles of steps (a), (b) and (c), in which non-PEGylated protein is recovered in step b) by eluting an AEC eluate from the AEC material and the eluted non-PEGylated protein is used in a subsequent PEGylation reaction.

The mixture of reaction products may contain a relatively low proportion of oligo-PEGylated protein. For example, the mixture of reaction products may comprise less than 25%, 20%, 15%, 10% or 5% oligo-PEGylated protein.

The mixture of reaction products may comprise at least about 20%, 30%, 40%, 45%, 50%, 55%, 60% or 70% of non-PEGylated protein.

The IEC flow-through solution (AEC or CEC flow-through solution) may contain a relatively high proportion of PEGylated protein. For example, the IEC flow-through solution may comprise at least about 90%, 95%, 98%, 99% or at least about 99.9% PEGylated protein. The AEC flow-through solution may contain a relatively high proportion of PEGylated protein. For example, the AEC flow-through solution may comprise at least about 90%, 95%, 98%, 99% or at least about 99.9% PEGylated protein.

The protein may be erythropoietin (EPO). The protein may be a hormone. The protein may be a hormone, a cytokine, an enzyme or an antibody.

The processes of the disclosure involve performing a PEGylation reaction and recycling non PEGylated protein. Such processes are advantageous because they enable relatively high yield.

The processes of the disclosure may involve performing a PEGylation reaction at about pH 7.0 to 9.0, which may be performed, for example, using an N-Hydroxysuccinimide (NHS) activated PEG reagent. The PEGylation reaction may be performed at about pH 7.5 to 8.5, or at about pH 8.0, and may be performed using an NHS activated PEG reagent. This provides a relatively rapid PEGylation step, which contributes to the overall rapidity of and high productivity of the processes described herein. The protein to be PEGylated may be erythropoietin.

The processes involve performing a PEGylation reaction to provide the mixture of reaction products, and recycling the non-PEGylated protein (unreacted protein) from that PEGylation reaction. The non PEGylated protein is recovered from the mixture of reaction products as part of an IEC step. The IEC step involves contacting the mixture of reaction products with an ion exchange material. The IEC flow-through comprises a relatively high proportion of PEGylated protein (mono-PEGylated and oligo-PEGylated protein) because most of the non-PEGylated protein binds to the anion exchange material. Thus non-PEGylated protein may be recovered from the mixture of reaction products by eluting it from the IEC material. The recovered non-PEGylated protein is added to a subsequent PEGylation reaction, thereby recycling the non-PEGylated protein.

The processes involve performing a PEGylation reaction to provide the mixture of reaction products, and recycling the non-PEGylated protein (unreacted protein) from that PEGylation reaction. The non PEGylated protein is recovered from the mixture of reaction products as part of an AEC step. The AEC step involves contacting the mixture of reaction products with an anion exchange material. The AEC flow-through comprises a relatively high proportion of PEGylated protein (mono-PEGylated and oligo-PEGylated protein) because most of the non-PEGylated protein binds to the anion exchange material. Thus non-PEGylated protein may be recovered from the mixture of reaction products by eluting it from the AEC material. The recovered non-PEGylated protein is added to a subsequent PEGylation reaction, thereby recycling the non-PEGylated protein.

The recycling of non-PEGylated protein in the processes disclosed herein is achieved by eluting non-PEGylated protein in the IEC step. The recycling of non-PEGylated protein in the processes disclosed herein is achieved by eluting non-PEGylated protein in the AEC step. Elution of non-PEGylated protein from an anion exchange material is relatively straightforward and fast, thereby enabling simple and rapid recovery of unreacted protein for recycling. The rapidity of the elution for non-PEGylated protein recycling contributes to the overall rapidity and productivity of the processes disclosed herein.

The processes are performed using a mixture of reaction products that contains non-PEGylated, mono-PEGylated and oligo-PEGylated protein. The proportion of oligo-PEGylated protein in the mixture of reaction products may be relatively low. The processes involve an ion exchange chromatography (IEC) step that provides an IEC flow through solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products. The processes involve an ion exchange chromatography (IEC) step that provides an IEC flow through solution in which the fraction of non-PEGylated protein is decreased relative to the mixture of reaction products. The IEC flow-through solution may contain a high proportion of PEGylated protein, wherein the PEGylated protein consists of mono-PEGylated protein and oligo-PEGylated protein. That is, the processes of the disclosure involve an IEC step that is performed in flow through mode. The IEC step in flow through mode is relatively rapid. The relatively fast IEC step of the disclosure contributes to the relatively overall high productivity of the processes of the disclosure. The IEC step may be an AEC step or a CEC step.

The IEC flow-through solution has an increased fraction of PEGylated protein relative to the mixture of reaction products. The AEC flow-through solution has a decreased fraction of non-PEGylated protein relative to the mixture of reaction products

The processes may involve an anion exchange chromatography (AEC) step that provides an AEC flow through solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products. The processes involve an anion exchange chromatography (AEC) step that provides an AEC flow through solution in which the fraction of non-PEGylated protein is decreased relative to the mixture of reaction products. The AEC flow-through solution may contain a high proportion of PEGylated protein, wherein the PEGylated protein consists of mono-PEGylated protein and oligo-PEGylated protein. That is, the processes of the disclosure involve an AEC step that is performed in flow through mode. The AEC step in flow through mode is relatively rapid. The relatively fast AEC step of the disclosure contributes to the relatively overall high productivity of the processes of the disclosure.

An AEC flow-through solution has an increased fraction of PEGylated protein relative to the mixture of reaction products. The AEC flow-through solution has a decreased fraction of non-PEGylated protein relative to the mixture of reaction products.

An AEC step comprises applying the mixture of reaction products to an anion exchange material. The anion exchange material may be a strong anion exchange material. Suitable materials are well known in

the art and include Toyopearl SuperQ 650M, which is a strong anion exchange material. The anion exchange material may have a low binding capacity for the PEGylated form of the protein.

- The IEC material may have a binding capacity for PEGylated protein of less than about 1.5 g/L. The IEC material may have a binding capacity for PEGylated protein of less than about 1.0 g/L, 0.75 g/L, 0.5 g/L, 0.25 g/L, 0.10 g/L, or 0.05 g/L. The binding capacity may be the dynamic binding capacity. The anion exchange material may have a binding capacity for PEGylated protein of less than about 1.5 g/L, less than about 1.0 g/L, less than about 0.75 g/L, less than about 0.5 g/L, less than about 0.1 g/L, less than about 0.05 g/L, less than about 0.01 g/L or less than about 0.001 g/L. The anion exchange material may have a binding capacity for PEGylated protein of close to 0 g/L (i.e. close to 0 g of PEGylated protein / L AEC resin). In particular, the anion exchange material may have a binding capacity for PEGylated EPO of less than about 0.5 g/L, less than about 0.05 g/L, less than about 0.01 g/L or less than about 0.001 g/L. The anion exchange material may have a binding capacity for PEGylated EPO of close to 0 g/L (i.e. close to 0 g of PEGylated EPO / L AEC resin). . The IEC material may be an AEC material and the PEGylated protein may be PEGylated erythropoietin.
- The IEC material may have a relatively high binding capacity for non-PEGylated protein (e.g. non-PEGylated EPO) of 20 – 50 g/L, 30 – 40 g/L, or about 35 g/L. The IEC material may have a binding capacity for non-PEGylated protein (e.g. non-PEGylated EPO) of at least about 20 g/L, 25 g/L, 30 g/L or about 35 g/L. The binding capacity may be the dynamic binding capacity. The IEC material may be an AEC material and the non-PEGylated protein may be non-PEGylated erythropoietin. The protein may be erythropoietin (EPO). The protein may be a hormone. The protein may be a hormone, a cytokine, an enzyme or an antibody.

The process may comprise an anion exchange chromatography (AEC) step and the protein may be erythropoietin.

- The use of an ion exchange material with a low binding capacity for the PEGylated form of the protein is advantageous because it facilitates rapid separation of non-PEGylated protein from PEGylated protein. The IEC step can be performed in flow through mode to rapidly provide an IEC flow-through solution from which most, or almost all, of the unreacted (non-PEGylated) protein has been removed. Hence the flow through solution from the IEC step comprises a relatively high proportion of PEGylated protein. It may comprise at least about 80%, 85%, 90% or 95% PEGylated protein, or 80-95% PEGylated protein. It may comprise at least about 90% PEGylated protein. It may comprise at least about 95% PEGylated protein. It may comprise a low proportion, close to zero, or zero non-PEGylated protein. It may comprise less than about 20%, 15%, 10% or 5% non-PEGylated protein, or about 20-5% non-PEGylated protein. It may comprise less than 10% non-PEGylated protein. It may comprise less than 5% non-PEGylated protein.

- A further advantage of using an ion exchange material with a low binding capacity for the PEGylated form of the protein is that it facilitates rapid recovery of unreacted (non-PEGylated) protein from the ion

exchange material. The non PEGylated protein may be recovered rapidly by step elution, which may provide an eluate comprising the non-PEGylated protein in a relatively concentrated form which may be suitable for direct addition to a PEGylation reaction. The non PEGylated protein may be eluted from the anion exchange material in an elution step performed at a relatively low conductivity, or low salt

5 concentration, which facilitates recycling of non-PEGylated protein for example because removal of high salt concentrations (for example by diafiltration, reconcentration or buffer exchange) is not required before the eluate is added to a subsequent PEGylation reaction. The use of an ion exchange material with a low binding capacity for the PEGylated form of the protein is thus advantageous because it provides for rapid recovery of unreacted (non-PEGylated) protein in a form that is particularly suitable for recycling in a
10 subsequent PEGylation reaction.

The use of an anion exchange material with a low binding capacity for the PEGylated form of the protein is advantageous because it facilitates rapid separation of non-PEGylated protein from PEGylated protein. The AEC step can be performed in flow through mode to rapidly provide an AEC flow-through solution from which most, or almost all, of the unreacted (non-PEGylated) protein has been removed. Hence the
15 flow through solution from the AEC step comprises a relatively high proportion of PEGylated protein. It may comprise at least about 80%, 85%, 90% or 95% PEGylated protein, or 80-95% PEGylated protein. It may comprise at least about 90% PEGylated protein. It may comprise at least about 95% PEGylated protein. It may comprise a low proportion, close to zero, or zero non-PEGylated protein. It may comprise less than about 20%, 15%, 10% or 5% non-PEGylated protein, or about 20-5% non-PEGylated protein. It
20 may comprise less than 10% non-PEGylated protein. It may comprise less than 5% non-PEGylated protein.

A further advantage of using an anion exchange material with a low binding capacity for the PEGylated form of the protein is that it facilitates rapid recovery of unreacted (non-PEGylated) protein from the anion exchange material. The non PEGylated protein may be recovered rapidly by step elution, which may
25 provide an eluate comprising the non-PEGylated protein in a relatively concentrated form which may be suitable for direct addition to a PEGylation reaction. The non PEGylated protein may be eluted from the anion exchange material in an elution step performed at a relatively low conductivity, or low salt concentration, which facilitates recycling of non-PEGylated protein for example because removal of high salt concentrations (for example by diafiltration, reconcentration or buffer exchange) is not required before
30 the eluate is added to a subsequent PEGylation reaction. The use of an anion exchange material with a low binding capacity for the PEGylated form of the protein is thus advantageous because it provides for rapid recovery of unreacted (non-PEGylated) protein in a form that is particularly suitable for recycling in a subsequent PEGylation reaction.

The processes of the disclosure involve providing a PEGylated protein mixture comprising mono-
35 PEGylated protein and oligo-PEGylated protein, which comprises collecting the flow-through solution from the IEC step. Collecting the IEC flow-through solution may comprise collecting batches of IEC flow-

through solution from separate IEC cycles, in this way a PEGylated protein mixture is provided which is a pooled PEGylated protein mixture. The IEC step may be an AEC step or a CEC step.

The processes of the disclosure involve providing a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein, which comprises collecting the flow-through solution from the AEC step. Collecting the AEC flow-through solution may comprise collecting batches of AEC flow-through solution from separate AEC cycles, in this way a PEGylated protein mixture is provided which is a pooled PEGylated protein mixture. The present disclosure provides processes for producing a mono-PEGylated protein composition comprising at least about 90% mono-PEGylated protein, the process comprising: subjecting the PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein produced by a process according to the first aspect of the disclosure to a purification process separating mono-PEGylated protein and oligo-PEGylated protein; and recovering a mono-PEGylated protein composition in which the fraction of mono-PEGylated protein is increased relative to the PEGylated protein mixture, wherein the mono-PEGylated protein composition comprises at least about 90% mono-PEGylated protein.

The mono-PEGylated protein composition may contain a relatively high proportion of mono-PEGylated protein. For example, the protein composition may comprise at least about 95%, 98%, 99%, or 99.9% mono-PEGylated protein.

PEGylation reactions tend to produce mixtures comprising non-PEGylated protein (unreacted protein), mono-PEGylated protein, and oligo-PEGylated protein. Mono-PEGylated proteins may be desirable because they provide a balance between increased half-life and comparable biological activity relative to non-PEGylated and oligo-PEGylated versions of the protein. Proteins, especially therapeutic proteins, are often expensive to manufacture and therefore processes that provide good yields of mono-PEGylated therapeutic protein are particularly desirable. Processes that provide good yields of mono-PEGylated EPO (Mircera®) are particularly desirable.

The processes disclosed herein are advantageous because they are relatively rapid, and therefore enable relatively high productivity. The processes disclosed herein also enable high yield, in that a high proportion of the starting protein becomes PEGylated, and in particular mono-PEGylated. The processes disclosed herein also provide mono-PEGylated protein compositions of relatively high purity.

The flow-through solution from the IEC step is used to provide a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein. For example the flow-through solution from an AEC step is used to provide a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein. This mixture is subjected to a purification process to separate the mono-PEGylated protein from the oligo-PEGylated protein and thereby provide a composition comprising a relatively high proportion of mono-PEGylated protein. The purification process may comprise subjecting the PEGylated protein mixture to a hydrophobic interaction chromatography (HIC) step. Both the anion exchange step

and HIC step are relatively rapid, thereby providing a relatively rapid process for producing a mono-PEGylated protein composition. The HIC step may be performed in flow through mode, which is particularly rapid, to provide a rapid process for producing a mono-PEGylated protein composition.

The processes of the disclosure involve collecting the flow-through solution from the IEC step (which may be an AEC or a CEC step). For example the processes of the disclosure may involve collecting an IEC flow-through solution. Collecting the IEC (AEC or CEC) flow-through solution may comprise collecting batches or fractions of IEC (AEC or CEC) flow-through solution from separate IEC (AEC or CEC) cycles, in this way a PEGylated protein mixture is provided which is a pooled PEGylated protein mixture.

Collecting the IEC (AEC or CEC) flow-through solution may comprise delivering the IEC (AEC or CEC) flow-through solution to a HIC material via an in-line connection, in an "in-line conditioning" operation. Collecting the IEC (AEC or CEC) flow-through solution may comprise delivering the IEC (AEC or CEC) flow-through solution to a vessel for conditioning the IEC (AEC or CEC) flow-through solution. Collecting the IEC (AEC or CEC) flow-through solution may comprise delivering the IEC (AEC or CEC) flow-through solution directly to the HIC material. When the IEC (AEC or CEC) flow-through solution is delivered directly to the HIC material the IEC (AEC or CEC) flow-through solution may be conditioned for example by in-line conditioning. There may be a continuous flow of IEC (AEC or CEC) flow-through solution (PEGylated protein mixture) from the IEC (AEC or CEC) material to the HIC material which is in-line conditioned to provide a continuous low of flow-through solution to the HIC material. The IEC (AEC or CEC) flow-through solution may comprise a low proportion of non-PEGylated protein and a high proportion of PEGylated protein (mono-PEGylated and oligo-PEGylated protein). The processes of the disclosure may further involve separating the mono-PEGylated protein from the oligo-PEGylated protein in the PEGylated protein mixture by performing a hydrophobic interaction chromatography (HIC) step.

The processes of the disclosure are particularly suitable for producing mono-PEGylated EPO compositions. In processes for producing mono-PEGylated EPO, the AEC step is performed at a pH of about 7.0 to 10.0, about 7.0 to 9.0, about 7.5 to 8.5, or about 8.0. This is advantageous over processes involving cation exchange chromatography (CEC) which is generally performed at a pH of 3.0 or lower. Acid conditions of about pH 3.0 or lower may have adverse effects on EPO quality, such adverse effects are reduced in the processes of the present disclosure, which do not require CEC and thus do not require the low pH conditions necessary for CEC-based purification of EPO.

Furthermore, acidic forms (acidic variants) of EPO may have higher therapeutic activity, making them particularly useful. However, recovery of mono-PEGylated acidic forms of EPO by CEC may be poor because their elution conditions in CEC are similar to those of di-PEGylated non-acidic forms of EPO. Acidic conditions result in the oxidation of the side chains of amino acids in a protein molecule. As CEC separates protein molecules according to their charge, basic variants of a species will elute later than acidic variants. For this reason, the CEC elution profile of di-pegylated non-acidic forms of EPO tends to overlap with mono-PEGylated acidic forms of EPO. The poor recovery of these useful mono-PEGylated

acidic EPO forms by CEC is reduced in the processes disclosed herein, which do not rely on CEC to separate mono-PEGylated from oligo-PEGylated EPO.

The advantages associated with the disclosure to provide mono-PEGylated EPO may apply to other proteins, particularly other therapeutic proteins. That is, avoidance of low pH conditions and recovery of acidic variants may render the disclosure useful for providing mono-PEGylated proteins other than mono-PEGylated EPO. The processes may comprise carrying out the IEC and HIC steps at substantially the same pH. The processes may comprise carrying out the PEGylation reaction, IEC and HIC steps at substantially the same pH. The processes may comprise carrying out the AEC step and purification process at substantially the same pH. The processes may comprise carrying out the AEC and HIC steps at substantially the same pH. The processes may comprise carrying out the PEGylation reaction, AEC and HIC steps at substantially the same pH. The pH may be about 7.0 to 9.0, or about 7.5 to 8.5, or about 8.0. Substantially the same pH may refer to ± 0.5 pH units. Performing steps of the process at substantially the same pH is advantageous because it avoids the need for pH adjustments between steps, which pH adjustments may be time-consuming and therefore productivity-reducing. . The process may comprise carrying out a PEGylation reaction to provide a mixture of reaction products and applying the mixture of reaction products directly to the IEC material. That is, the process may comprise carrying out a PEGylation reaction and applying the resultant mixture to the IEC material without adjusting the pH of the mixture.

The processes of the disclosure involve providing a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein, which comprises collecting the flow-through solution from the IEC step. Collecting the IEC flow-through solution may comprise collecting batches of IEC flow-through solution from separate IEC cycles, in this way a PEGylated protein mixture is provided which is a pooled PEGylated protein mixture. The processes of the disclosure involve providing a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein, which comprises collecting the flow-through solution from the AEC step. Collecting the AEC flow-through solution may comprise collecting batches of AEC flow-through solution from separate AEC cycles, in this way a PEGylated protein mixture is provided which is a pooled PEGylated protein mixture.

The processes of the disclosure involve subjecting a PEGylated protein mixture to a purification process. This may involve subjecting a pooled PEGylated protein mixture to a purification process. Collecting the IEC flow-through solution may comprise delivering the IEC flow-through solution to a HIC material via an in-line connection, in an "in-line conditioning operation". Collecting the IEC flow-through solution may comprise delivering the IEC flow-through solution to a vessel for conditioning the IEC flow-through solution to provide a PEGylated protein mixture. The IEC flow-through solution may comprise a low proportion of non-PEGylated protein and a high proportion of PEGylated protein (mono-PEGylated and oligo-PEGylated protein). The purification process may comprise subjecting the PEGylated protein mixture to a hydrophobic interaction chromatography (HIC) step. The HIC step may be performed in flow

through mode. The HIC step in flow through mode may comprise contacting the PEGylated protein mixture with a hydrophobic interaction material, and collecting the flow-through. The flow-through comprises a relatively high proportion of mono-PEGylated protein. Performing the HIC step in flow through mode is advantageous because it can be performed relatively rapidly, thereby contributing to the overall rapidity and productivity of the process. Performing the HIC step in flow through mode, rather than bind and elute mode, is advantageous because it can be performed using a column of relatively small size.

The step of providing the PEGylated protein mixture may comprise pooling the flow-through solution from two IEC steps, or three IEC steps, or more than three IEC steps, or four IEC steps, or five IEC steps.

Such pooling of IEC flow-through solution enables the HIC step to produce a mono-PEGylated protein composition from the products of multiple PEGylation reactions, which is relatively efficient. The IEC steps may be a series of AEC steps, or a series of CEC steps. The processes of the disclosure involve subjecting a PEGylated protein mixture to a purification process. This may involve subjecting a pooled PEGylated protein mixture to a purification process. Collecting the AEC flow-through solution may comprise delivering the AEC flow-through solution to a HIC material via an in-line connection, in an "in-line conditioning operation". Collecting the AEC flow-through solution may comprise delivering the AEC flow-through solution to a vessel for conditioning the AEC flow-through solution to provide a PEGylated protein mixture. Collecting the AEC flow-through solution may comprise delivering the AEC flow-through solution directly to the HIC material. When the AEC flow-through solution is delivered directly to the HIC material the AEC flow-through solution may be conditioned for example by in-line conditioning. There may be a continuous flow of AEC flow-through solution from the AEC material to the HIC material which is in-line conditioned to provide a continuous flow of conditioned flow-through solution to the HIC material. The AEC flow-through solution may comprise a low proportion of non-PEGylated protein and a high proportion of PEGylated protein (mono-PEGylated and oligo-PEGylated protein). The purification process may comprise subjecting the PEGylated protein mixture to a hydrophobic interaction chromatography (HIC) step.

Conditioning the IEC flow-through solution may be referred to as providing a conditioned second mixture. Conditioned second mixtures are described in more detail below. Conditioning an IEC flow-through solution (or providing a conditioned second mixture) may comprise the addition of salt. Conditioning an IEC flow-through solution the addition of bicine. Conditioning an IEC flow-through solution (or providing a conditioned second mixture) may comprise the addition of salt and bicine. Conditioning an IEC flow-through solution (or providing a conditioned second mixture) may comprise the addition of salt and/or bicine to provide a conditioned second mixture as described below.

The HIC step may be performed in flow through mode. The HIC step in flow through mode may comprise contacting the PEGylated protein mixture with a hydrophobic interaction material, and collecting the

flow-through. The flow-through comprises a relatively high proportion of mono-PEGylated protein. Performing the HIC step in flow through mode is advantageous because it can be performed relatively rapidly, thereby contributing to the overall rapidity and productivity of the process. Performing the HIC step in flow through mode, rather than bind and elute mode, is advantageous because it can be
5 performed using a column of relatively small size.

The step of providing the PEGylated protein mixture may comprise pooling the flow-through solution from two IEC steps, or three IEC steps, or more than three IEC steps, or four IEC steps, or five IEC steps. Such pooling of IEC flow-through solution enables the HIC step to produce a mono-PEGylated protein composition from the products of multiple PEGylation reactions, which is relatively efficient. For example
10 the step of providing the PEGylated protein mixture may comprise pooling the flow-through solution from two AEC steps, or three AEC steps, or more than three AEC steps, or four AEC steps, or five AEC steps. Such pooling of AEC flow-through solution enables the HIC step to produce a mono-PEGylated protein composition from the products of multiple PEGylation reactions, which is relatively efficient.

15 ***Summary of the Figures***

In the figures "EPO" refers to un-reacted EPO (non-PEGylated EPO), "mono" refers to mono-PEGylated EPO, "oligo" refers to oligo-PEGylated EPO. "F/T" refers to flow-through solution.

Figure 1: Schematic sequence and key data of an embodiment of a process in accordance with the present invention. Non-PEGylated starting material is used in two further cycles without supplementing
20 with fresh EPO. Productivity is calculated to be increased relative to a known production process for mono-PEGylated EPO (Mircera®) disclosed in WO 2009/010270, in which two CEC chromatographic columns are used. The dynamic binding capacity of the AEC material is 35 g/L (35 g of non-PEGylated protein / L resin) (a factor of 30 higher than the CEC material of the process of WO 2009/010270. The HIC column can be loaded in flow through mode with dynamic binding capacity 5 g/L (5 g of oligo-
25 PEGylated protein / L resin) (a factor of ~4 higher than the CEC material used in the process of WO 2009/010270). Compared with the process disclosed in WO 2009/010270, the first column can be made smaller, and the combined fractions (flow-through) from the 3 cycles can be processed in one run on a second column of equal size.

Figure 2: Schematic sequence and key data of an embodiment of a process in accordance with the
30 present invention. The batch size is increased by replacing the reacted EPO with an identical quantity of fresh EPO for the subsequent cycle. The number of cycles can be varied here as desired. Here the batch size was selected such that it can be completed in one week. The calculated productivity is increased by a factor of 10 relative to a known production process for mono-PEGylated EPO disclosed in WO 2009/010270.

Figure 3: Schematic sequence and key data of an embodiment of a process in accordance with the present invention. The batch size is increased because the EPO consumed in the PEGylation reaction is replaced with an identical quantity of fresh EPO in the next cycle. In addition, the concentration of the EPO for the PEGylation reaction was increased to 10 g/L. In this way, twice the quantity can be processed at the same reaction volume. The number of cycles can be varied as desired in this case. Here the batch size was selected such that it can be completed in one week. Productivity is calculated to be increased by a factor of 20 relative to a known production process for mono-PEGylated EPO disclosed in WO 2009/010270.

Figure 4: Chromatogram of samples from AEC in flow through mode, showing recovery of EPO from the PEGylation reaction mixture in the flow-through and eluate fractions. Mono-PEGylated EPO and oligo-PEGylated EPO are present in the flow-through solution. EPO can be recovered by step elution almost quantitatively and in high purity for a further PEGylation reaction. The composition of fractions was determined by RP-HPLC.

Figure 5: Example AEC chromatograms for the recovery of EPO from the PEGylation reaction mixture in three sequential cycles. The total amount of EPO in cycles 2 and 3 is reduced, because only unreacted EPO recovered from the previous cycle was used for PEGylation in cycles 2 and 3. Mono-PEGylated and oligo-PEGylated EPO are present in the flow-through solution. Unreacted EPO can be recovered by step elution almost quantitatively and in high purity for further PEGylation reactions. The composition of fractions was determined by RP-HPLC.

Figure 6: Example HIC chromatogram for the performance of the purification on Toyopearl Phenyl 650 M in bind and elute mode. Non-PEGylated EPO was removed in the flow-through solution at 500 mM Na_2SO_4 . In a falling Na_2SO_4 gradient, mono-PEGylated EPO elutes at approximately 300 mM Na_2SO_4 with good resolution from the oligo species. The composition of fractions was determined by RP-HPLC.

Figure 7: Example HIC chromatogram for the performance capacity of the purification on Toyopearl Phenyl 650 M in flow through mode. Mono-PEGylated EPO is present in the flow-through solution at about 300 mM Na_2SO_4 . The oligo-PEGylated EPO species remain on the column until regeneration. The composition of fractions was determined by RP-HPLC.

Figure 8: Figures 8A, 8B and 8C show AEC chromatograms for recovery of EPO from the PEGylation reaction mixture in three sequential cycles.

Figure 9: HIC chromatogram for the performance capacity of the purification on Toyopearl Phenyl 650 M in bind and elute mode.

Figure 10: HIC chromatogram for the performance capacity of the purification on Toyopearl Phenyl 650 M in flow through mode.

Detailed Description of the Invention

Embodiments and experiments illustrating the principles of the invention will now be discussed with reference to the accompanying figures.

Mono-PEGylated protein compositions

- 5 The present invention relates to processes for providing mono-PEGylated protein compositions. The present invention relates to processes for providing a protein composition comprising at least 90% mono-PEGylated protein. In particular, the present invention provides processes for providing an EPO composition comprising at least 90% mono-PEGylated EPO.

10 In the present context, a mono-PEGylated protein composition is a composition comprising a protein, wherein a relatively high proportion of the protein present in the composition is present as mono-PEGylated protein. A mono-PEGylated protein composition may comprises at least about 90%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.9% or 99.99% mono-PEGylated protein

A protein composition comprising at least x% mono-PEGylated protein refers to a protein composition in which at least x% of the protein present in that composition is mono-PEGylated. For example, a protein
15 composition comprising at least 99% mono-PEGylated protein is a protein composition in which at least 99% of that protein present in the composition is mono-PEGylated. Protein compositions produced by the processes of the invention may comprise at least about 90%, at least about 95%, 96%, 97%, 98%, 99%, 99.5%, 99.9% or 99.99% mono-PEGylated protein.

In particular, in the present context, a mono-PEGylated EPO composition is a composition comprising
20 EPO, wherein a relatively high proportion of the EPO present in the composition is present as mono-PEGylated EPO. An EPO composition comprising at least "x%" mono-PEGylated EPO refers to an EPO composition in which at least "x%" of the EPO present in that composition is mono-PEGylated. For example, an EPO composition comprising at least 99% mono-PEGylated EPO is an EPO composition in which at least 99% of the EPO present in the composition is mono-PEGylated. EPO compositions
25 produced by the processes of the invention may comprise at least about 90%, at least about 95%, 96%, 97%, 98%, 99%, 99.5%, 99.9%, or 99.99% mono-PEGylated EPO. EPO compositions produced by the processes of the invention may comprise at least about 98% mono-PEGylated EPO. EPO compositions produced by the processes of the invention may comprise at least about 99% mono-PEGylated EPO.

Methods for the determination of purity are known to those of skill in the art. Purity of a non-PEGylated,
30 mono-PEGylated or oligo-PEGylated protein may be determined by any suitable method of analysis (e.g. band intensity on gel, ELISA, HPLC and the like). Determination of purity may involve using a standard curve generated using a reference material of known purity. Purity may also be determined on a weight-by-weight basis. The purity of a non-PEGylated or PEGylated protein expressed herein in percentage terms (%) may be determined for example using relative "area under the curve" values, which can

typically be obtained for peaks in a chromatogram, such as an HPLC chromatogram. Methods of determining purity include RP-HPLC and size exclusion chromatography (SEC).

Chromatography

5 The "isoelectric point" or "pI" of a protein is the pH at which the protein has zero net overall charge. That is, the pI of a protein is the pH at which the protein has an equal number of positive and negative charges. Determination of the pI for any given protein can be done according to well-established techniques, such as isoelectric focusing. The pI of EPO is in the range 4.0 to 5.5. Alternative methods of measuring pI may give slightly different values, for example microchip isoelectric focusing gives an apparent pI of about 3.5 – 4.0 (Vickova 2008). The precise pI of EPO may depend for example on the
10 degree of glycosylation and sialic acid residues, or the existence of charge variants, which may in turn depend on the means by which it has been produced (e.g. host cells used for recombinant expression).

Ion exchange chromatography separates molecules on the basis of differences in their net surface charge. It can be used to separate protein molecules. In ion exchange chromatography a mixture including a protein of interest may be passed over an ion exchange material, which carries a charge.
15 When the ion exchange material has a negative charge the process is termed cation exchange chromatography (CEC), and when it has a positive charge the process is termed anion exchange chromatography (AEC). The protein of interest may be separated from the remainder of the mixture by manipulating charge-based interactions between the protein and the ion exchange material. This is typically done by manipulating the ionic strength, or conductivity, of the mixture and or of buffers passed
20 over the ion exchange material.

The term ion exchange chromatography (IEC) used herein may refer to cation exchange chromatography (CEC), or anion exchange chromatography (AEC). Where a process involves multiple IEC steps, they are either all AEC steps or all CEC steps. Likewise reference to an IEC flow-through solution or an IEC material may refer to either CEC or an AEC flow-through solution, and to a CEC or an AEC material.
25 Where a process involves an IEC step which is an AEC step, all IEC steps are AEC steps, the IEC materials is an AEC materials, the IEC flow-through solution is an AEC flow-through solution, any IEC eluate is an AEC eluate. Where a process involves an IEC step which is a CEC step, the IEC step is a CEC step, the IEC materials is a CEC material, the IEC flow-through solution is a CEC flow-through solution, and any IEC eluate is a CEC eluate.

30 For example, a mixture containing a protein of interest may be loaded onto an ion exchange material. The loading conditions (and wash conditions, if used) are selected to promote binding of only certain components of the mixture to the ion exchange material.

For example the loading conditions may promote binding of the protein of interest to the ion exchange material, and ion exchange material may then be washed with a wash buffer, to remove unwanted
35 components of the mixture (e.g. contaminants), and finally the protein of interest may be eluted (removed

from the ion exchange material) by increasing the ionic strength (conductivity) of the buffer. This type of procedure is known as “bind and elute” mode, because the protein of interest is bound to ion exchange material and then eluted. The solution that comes off the matrix in the elution step and contains the protein of interest may be known as the eluant or eluate.

- 5 Alternatively, the loading conditions may promote binding of unwanted components of the mixture (e.g. contaminants). In this mode, the protein of interest may flow through a matrix of the ion exchange material and be collected. This type of procedure is known as “flow through” mode. The composition that is collected after flowing through a matrix of ion exchange material is the “flow-through” or “flow-through solution”. The flow-through solution that comes off the matrix and contains the protein of interest may also
- 10 be termed “effluent”. Unlike “bind and elute” mode, which involves a change in conductivity and/or pH to elute the protein of interest from the column, “flow through” mode is carried out under isocratic conditions. The flow-through solution may be collected as fractions, which may be pooled to provide a flow-through pool. Proteins, such as EPO, are comprised of amino acids which include acidic and basic residues. At low pH (high H^+ concentration) the carboxylic acid groups of proteins tend to be uncharged ($-COOH$) and
- 15 their nitrogen-containing basic groups fully charged ($-NH_3^+$) giving most proteins a net positive charge. At high pH the carboxylic acid groups are negatively charged ($-COO^-$) and the basic groups tend to be uncharged ($-NH_2$), giving most proteins a net negative charge.

The isoelectric point (pI) of a protein is the pH at which that protein has no net charge because the positive and negative charges balance.

- 20 Ion exchange chromatography takes advantage of the fact that the relationship between net surface charge and pH is unique for a particular protein. At a pH below its isoelectric point a protein will have an overall positive charge and will therefore bind to a negatively charged material (i.e. in CEC). At a pH above its isoelectric point a protein will have an overall negative charge and therefore will bind to a positively charged material (i.e. in AEC).
- 25 This is why in AEC protein mixtures (load compositions) and wash buffers of a relatively high pH are generally used, in order that the protein of interest (or contaminant, e.g. unwanted protein) has a net negative charge and therefore binds to the positively charged anion exchange material. When a protein of interest is bound to the anion exchange material, it may optionally be washed to remove contaminants, and can then be eluted. When a contaminant (such as unwanted protein) is bound to the anion exchange
- 30 material it is removed from the mixture, thereby purifying one or more proteins of interest in the flow-through.

- Hydrophobic interaction chromatography (HIC) separates molecules on the basis of differences in their surface hydrophobicity. It can be used to separate protein molecules. In HIC a mixture including a protein of interest may be passed over an HIC material. The interaction between proteins and a HIC
- 35 material is altered by the presence of certain salts. Increasing the salt concentration increases the

interaction and reducing the salt concentration reduces the interaction. For selective elution, the salt concentration may be lowered and the components of the mixture elute in order of hydrophobicity, the most hydrophobic components eluting last. HIC may be performed in flow through mode or in bind and elute mode as described for ion exchange chromatography above.

- 5 A chromatography "material" in the present context, such as an anion exchange material or a HIC material, refers to the stationary phase or solid phase. This may also be referred to as a resin or matrix. The "material" in this context provides a matrix to which a component of a mixture, such as protein of interest, may bind. The material may be or may comprise a column, for example an expanded bed or packed bed column. The material may be in the form of discrete particles or beads. The material may in
10 the form of a membrane. In the present context the mobile phase flows through the stationary phase and carries the substances to be separated by chromatography with it. The mobile phase is a mixture, such as a first or second mixture, or a solution such as a flow through solution. A buffer, such as an elution buffer or wash buffer, is also a mobile phase.

- 15 In the present context the term "loading" refers to a step of contacting a mixture or composition onto chromatography material. The chromatography material may be equilibrated before the loading step.

Equilibration involves applying an equilibration buffer to the chromatography material. The pH, ionic strength, conductivity, and/or salt concentration of the equilibration buffer are selected to ensure that when a protein mixture is loaded onto the chromatography matrix, the desired binding and/or flow through or specific proteins or contaminants is achieved.

- 20 Elution from a chromatography material, such an anion exchange material or a cation exchange material or a hydrophobic interaction material, may be achieved by changing the ionic strength, pH, conductivity, or salt concentration of a buffer using gradient elution or step elution. Gradient elution, or linear gradient elution, may be used when many components of individual interest are bound to the material and may be eluted differently, and for high resolution separation. Step elution is useful for removing a single
25 component (or removing a specific group of components together) from a chromatography material. Step elution is relatively fast, and consumes less buffer. Step elution may be used to elute a protein of interest from a chromatography material in a relatively concentrated form. A rising gradient of ionic strength is commonly used for elution in AEC. A falling gradient of salt concentration is commonly used for elution in HIC.

- 30 Producing a mono-PEGylated protein composition

The present invention provides a process for producing a mono-PEGylated protein composition.

Compared with prior art processes using CEC to provide a mono-PEGylated protein composition, the processes disclosed herein are more productive. A prior art process for production of mono-PEGylated EPO is described in WO 2009/010270.

The present invention provides a process for producing a PEGylated protein mixture, the process comprising: a) reacting a non-PEGylated protein with a PEGylation reagent to produce a mixture of reaction products comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein; (b) subjecting the mixture of reaction products to an anion exchange chromatography (AEC) step to provide an AEC flow-through solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products; the AEC step comprising applying the mixture of reaction products to an AEC material under conditions suitable for binding non-PEGylated protein; and (c) collecting the AEC flow-through solution from step b) to provide a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein, wherein non-PEGylated protein is recovered in step b) by eluting an AEC eluate from the AEC material and the eluted non-PEGylated protein is used in a subsequent PEGylation reaction.

The mixture of reaction products comprising non-PEGylated protein and PEGylated protein produced by reacting a non-PEGylated protein with a PEGylation reagent may also be referred to herein as a "first mixture". The PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein produced by this method may also be referred to herein as a "second mixture".

The present invention also provides a process for producing a mono-PEGylated protein composition comprising at least about 90% mono-PEGylated protein, the process comprising: subjecting the PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein produced by the process described above to a purification process separating mono-PEGylated protein and oligo-PEGylated protein; and recovering a mono-PEGylated protein composition in which the fraction of mono-PEGylated protein is increased relative to the PEGylated protein mixture, wherein the mono-PEGylated protein composition comprises at least about 90% mono-PEGylated protein.

The present invention provides a process for producing a protein composition comprising at least about 90% mono-PEGylated protein, the process comprising: (a) providing a first mixture comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein (b) subjecting the first mixture to an anion exchange chromatography (AEC) step to provide an AEC flow-through solution in which the fraction of PEGylated protein is increased relative to the first mixture; the AEC step comprising applying the first mixture to an AEC material under conditions suitable for binding non-PEGylated protein; (c) collecting the AEC flow-through solution from step b) to provide a second mixture comprising mono-PEGylated protein and oligo-PEGylated protein; and (d) subjecting the second mixture to a hydrophobic interaction chromatography (HIC) step to provide a protein composition in which the fraction of mono-PEGylated protein is increased relative to the second mixture, wherein the protein composition comprises at least about 90% mono-PEGylated protein.

Figures 1 to 3 show embodiments of processes in accordance with the present disclosure.

Producing a mixture of reaction products

The processes of the invention comprise providing a mixture of reaction products (step a). The processes of the invention are performed using a mixture of reaction products (or "first mixture") that comprises non-PEGylated, mono-PEGylated and oligo-PEGylated protein. The mixture of reaction products comprises non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein. The protein may be EPO.

The processes of the invention are performed using a mixture of reaction products (or "first mixture") that comprises non-PEGylated, mono-PEGylated and oligo-PEGylated protein, wherein the proportion of oligo-PEGylated protein is relatively low. The mixture of reaction products may comprise less than about 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, or 1% oligo-PEGylated protein, the oligo-PEGylated protein may be oligo-PEGylated EPO. The mixture of reaction products may comprise less than about 10% oligo-PEGylated protein. The mixture of reaction products may comprise less than about 10% oligo-PEGylated EPO.

The mixture of reaction products (or "first mixture") may comprise at least about 20%, 30%, 40%, 45%, 50%, 55%, 60% or 70% non-PEGylated protein. The mixture of reaction products may comprise about 20-70%, 40-60%, 45-55% non-PEGylated protein.

The mixture of reaction products (or "first mixture") may comprise at least about 30%, 40%, 45%, 50%, 55%, 60% or 70% mono-PEGylated protein. The mixture of reaction products may comprise about 30-70%, 40-60%, or 45-55% mono-PEGylated protein.

The mixture of reaction products (or "first mixture") may comprise (A) non-PEGylated protein in the range 40-60%, and (B) mono-PEGylated protein in the range 40-60%, and (C) oligo-PEGylated protein in the range 1-10%, wherein the total of (A), (B) and (C) is 100%. The mixture of reaction products may comprise (A) non-PEGylated protein in the range 45-55%, and (B) mono-PEGylated protein in the range 45-55%, and (C) oligo-PEGylated protein in the range 1-5%, wherein the total of (A), (B) and (C) is 100%.

The mixture of reaction products (or "first mixture") may comprise (A) non-PEGylated protein, and (B) mono-PEGylated protein, and (C) oligo-PEGylated protein. The non-PEGylated protein, mono-PEGylated protein, and oligo-PEGylated protein may be present in the ratio A:B:C, wherein A is about 40-60, B is about 30-50, and C is about 1-10; or wherein A is about 45-55, B is about 35-45, and C is about 1-10; or wherein A is about 35-55, B is about 35-45, and C is about 1-25. The non-PEGylated protein, mono-PEGylated protein, and oligo-PEGylated protein may be present in the ratio (A+B):C wherein (A+B):C is about 19:1; about 9:1 or about 9-19:1.

The ratio may be a weight or mass ratio. The ratio may be a molar ratio.

The mixture of reaction products (or "first mixture") is provided by performing a PEGylation reaction. Performing a PEGylation reaction involves reacting the protein (which may be the non-PEGylated protein)

with a PEGylation reagent. The protein may be EPO. The PEGylation reaction may be performed as described above in relation to PEGylation of EPO. The PEGylation reaction may be performed at a pH of about 7.0 to 9.0, wherein the PEG/protein molar ratio is about 0.6 – 1.0.

The PEGylation reaction may be performed at a pH of about 7.0 to 9.0, or about 7.5 to 8.5, or about 8.0.

5 The PEGylation reaction may be performed at a pH of about 7.0 to 9.0, or about 7.5 to 8.5, or about 8.0 using (NHS) activated PEG reagent. The pH at which the PEGylation reaction is performed may depend on the PEG reagent used. The PEG reagent may be mPEG-NHS, mPEG-SPA, mPEG-SVA or mPEG-Cl. The relationship between PEGylation reaction rates and pH is reviewed in Pfister 2016. Other reagents for PEGylation reactions are known in the art.

10 The molar ratio of PEG/protein in the PEGylation reaction may be ≤ 1.0 . The molar ratio of PEG/protein in the PEGylation reaction may be about 0.8. The PEG/protein molar ratio may be about 0.25 to 1.0, 0.3 to 1.0, 0.4 to 1.0, 0.5 to 1.0, 0.6 to 1.0 or 0.7 to 1.0. The PEG/protein molar ratio may be about 0.25 to 1.2, 0.3 to 1.2, 0.4 to 1.2, 0.5 to 1.2, 0.6 to 1.2 or 0.7 to 1.2. The PEG/protein molar ratio may be about 0.25 to 1.5, 0.3 to 1.5, 0.4 to 1.5, 0.5 to 1.5, 0.6 to 1.5 or 0.7 to 1.5. The PEG/protein molar ratio may have a
15 lower limit of about 0.4, 0.5, 0.6 or 0.7 and may have an upper limit of 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, or 1.5.

Ion exchange chromatography

The first mixture is subjected to an ion exchange chromatography (AEC) step (step b). This provides an IEC flow-through solution in which the fraction of PEGylated protein is increased relative to the first
20 mixture. This provides an IEC flow-through solution in which the fraction of non-PEGylated protein is decreased relative to the first mixture. The IEC step may be an AEC step or a CEC step.

Anion exchange chromatography

The mixture of reaction products (or "first mixture") may be subjected to an anion exchange chromatography (AEC) step (step b). This provides an AEC flow-through solution in which the fraction of
25 PEGylated protein is increased relative to the mixture of reaction products. This provides an AEC flow-through solution in which the fraction of non-PEGylated protein is decreased relative to the mixture of reaction products.

In the context of the present disclosure a fraction which is increased may be increased by an amount of at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100%. A fraction which is
30 increased may be increased by at least about 1.5-fold, 2-fold, 3-fold, 5-fold or 10-fold.

In the context of the present disclosure a fraction which is decreased may be decreased by an amount of at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100%. A fraction which is decreased may be increased by at least about 1.5-fold, 2-fold, 3-fold, 5-fold or 10-fold.

The AEC step comprises contacting the mixture of reaction products with an anion exchange material, or applying the mixture of reaction products to an anion exchange material. The AEC step may comprise loading the mixture of reaction products onto a column comprising an anion exchange material. The anion exchange material has a low binding capacity for PEGylated protein, such as PEGylated EPO.

5 That is, the AEC material has a relatively low binding capacity for the PEGylated version of the protein species for which a mono-PEGylated composition is desired. AEC conditions are used which are suitable for binding of the non-PEGylated protein to the anion exchange material. The solution that flows through the anion exchange material is the AEC flow-through solution. The low binding capacity of the anion exchange material for PEGylated protein has the effect that most of the PEGylated protein flows through

10 the anion exchange material and is therefore present in the AEC flow-through solution. The binding conditions suitable for binding of the non-PEGylated protein to the anion exchange material have the effect that most of the non-PEGylated protein is removed (by binding to the anion exchange material) and is not present in the flow-through solution.

The AEC material may be a material that does not significantly bind PEGylated protein. This may mean

15 that the AEC material binds less than about 10%, 5%, 2%, 1%, 0.5%, 0.1% or 0.01% of the PEGylated protein that is applied to it. Binding of PEGylated protein to the anion exchange material can also be reduced by applying high amounts of non-PEGylated protein to the anion exchange material. In this way, bound PEGylated protein is displaced by non-PEGylated protein. The amount of non-PEGylated protein applied to the anion exchange material may be about 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%,

20 100%, 110% or 120% of the dynamic break through capacity of the anion exchange material. The amount of non-PEGylated protein applied to the anion exchange material may be in the range of about 70-110%, about 80-100%, or about 85-95% of the dynamic break through capacity of the anion exchange material. The amount of non-PEGylated protein applied to the anion exchange material may be in the range of about 80-95% of the dynamic break through capacity of the anion exchange material.

25 The AEC flow-through solution comprises mono-PEGylated and oligo-PEGylated protein. The AEC flow-through solution may also comprise non-PEGylated protein, wherein the proportion of non-PEGylated protein is relatively low or close to zero. The AEC flow-through solution may comprise zero non-PEGylated protein, or amounts of non-PEGylated protein below the limit of detection. The protein may be EPO. The flow-through solution may comprise less than about 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1.5%,

30 1.0%, 0.5% or 0.1% non-PEGylated protein. The flow-through solution may comprise less than about 2% non-PEGylated protein. The flow-through solution may comprise less than about 2% non-PEGylated EPO.

The AEC flow-through solution may comprise at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or at least about 99.9% PEGylated protein.

35 The AEC flow-through may comprise about 80% mono-PEGylated protein and about 20 % oligo-PEGylated protein. The AEC flow-through solution may comprise at least about 60%, 65%, 70%, 75%, or

80% mono-PEGylated protein. The flow-through solution may comprise about 60-90%, 70-90%, or 75-85% mono-PEGylated protein. The AEC flow-through solution may comprise at least about 20%, 25%, 30%, 35%, or 40% oligo-PEGylated protein. The flow-through solution may comprise about 10-40%, 10-30%, or 15-25% oligo-PEGylated protein.

- 5 The AEC flow-through solution may comprise (A) non-PEGylated protein in the range 0.1-5%, and (B) mono-PEGylated protein in the range 60-90%, and (C) oligo-PEGylated protein in the range 10-40%, wherein the total of (A), (B) and (C) is 100%. The flow-through solution may comprise (A) non-PEGylated protein in the range 0.1-1.5%, and (B) mono-PEGylated protein in the range 75-85%, and (C) oligo-PEGylated protein in the range 15-25%, wherein the total of (A), (B) and (C) is 100%.
- 10 The AEC flow-through solution may comprise (A) non-PEGylated protein, and (B) mono-PEGylated protein, and (C) oligo-PEGylated protein. The non-PEGylated protein, mono-PEGylated protein, and oligo-PEGylated protein may be present in the ratio A : B : C, wherein A is about 1-5, B is about 10-20, and C is about 5-15; or wherein A is about 1-3, B is about 25-35 and C is about 5-10. The non-PEGylated protein, mono-PEGylated protein, and oligo-PEGylated protein may be present in the ratio
- 15 A:(B+C) wherein A:(B+C) is about 1:25; 1:50; 1:100; about 1:150, about 1:200, or about 1:100-200. The ratio may be a weight or mass ratio. The ratio may be a molar ratio.

The AEC step is performed under conditions suitable for non-PEGylated protein to bind to the AEC material. Conditions suitable for non-PEGylated protein binding may be determined empirically by screening one or more of the AEC material, or the pH, conductivity, salt content or buffer content of the

20 equilibration buffer.

The functional groups on an ion exchange material determine its charge. The terms "strong" and "weak" in this context refer to the extent that the ionisation state of the functional groups varies with varying pH. Strong ion exchangers show little or no variation in ion exchange capacity with change in pH. Strong ion exchangers remain fully charged over a broad pH range. Weak ion exchangers display pH-dependent

25 function and deliver optimal performance over a narrower pH range.

Strong anionic exchange materials may include a quaternary ammonium group (Q), or a quaternary amino ethyl group (QAE). A weak anionic exchange material may include diethylaminoethyl residues (DEAE). An advantage of using strong anionic exchange materials is that the binding capacity is maintained at high or low pH since there is no loss of charge.

- 30 AEC materials may comprise a resin that has been functionalised with a positively charged ligand or functional group, that is, with a basic ligand or functional group.

The anion exchange material used in the processes of the invention may be a strong anion exchange material. The anion exchange material is preferably a strong anion exchange material. It may have a quaternary ammonium group (Q). It may be a hydroxylated polymethacrylic polymer bead functionalised

with a Q groups. It may have a 50-150 nm, 70-120 nm, 70-100 nm or a 90-110 nm pore size. Pore size herein may refer to mean pore size. It may have a particle size of 40-150 μm , 40-120 μm , 40-100 μm , 40-80 μm , or about 60, 65, 70, 80, or 90 100 μm . Particle size (or bead size) may refer to mean particle size. It may have a 70-100 nm pore size and a 35-100 μm bead size. It may have a 100 nm pore size and a 65 μm bead size. It may be Toyopearl Super Q 650M (Tosoh Bioscience LLC; product number 43205). The anion exchange material used in the processes of the invention may be used in accordance with the manufacturer's instructions.

The anion exchange material used in the processes of the invention may be a weak anion exchange material. It may have tertiary and/or secondary amine functional groups. It may not have primary amine functional groups. It may have only one type of amine group or may have two or more types of amine group.

The anion exchange material may be or may comprise a column, for example an expanded bed or packed bed column. The anion exchange material may be or may comprise a continuous countercurrent tangential chromatography system. The anion exchange material may be in the form of discrete particles or beads. The anion exchange material may be in the form of a membrane. The anion exchange material may be in the form of a functionalised filter, or a functionalised fibre, fleece, or mesh. The anion exchange material may be in the form of any other solid support able to carry functional groups or exhibiting anion exchange properties.

Examples of anion exchangers include Toyopearl Super Q 650C, Toyopearl Super Q 650S, Toyopearl GigaCap Q 650M, GigaCap Q 650S, Toyopearl Q-600C AR, Toyopearl DEAE 650S, Toyopearl DEAE 650M and Toyopearl DEAE 650C, Toyopearl Super Q 650, Toyopearl Super Q 650 QAE, Toyopearl Super Q 550C, Macrorep High Q, Fractoprep TMAE, Q Hyper DF, Capto Q, Q-Sepharose FF, Q-Sepharose BB, Q-Sepharose XL, Q-Sepharose HP, MiniQ, MonoQ, MonoP, DEAE Sepharose FF, Source SQ, Source 30Q, ANX Sepharose 4 FF (high sub), Streamline DEAE, Streamline QXL, Poros HQ, Poros PI, Poros D, DEAEHyperD, Q Ceramic Hyper D, Fractogel DMAE.

The AEC material in the processes of the invention may be an anion exchange material having a low binding capacity for the PEGylated form of the protein. In particular, it may be an anion exchange material having low binding capacity for PEGylated EPO. The anion exchange material may be Toyopearl Super Q 650 M having low binding capacity for PEGylated EPO, for example a binding capacity of less than about 0.5 g/L, 0.05 g/L, 0.01 g/L or 0.001 g/L, or close to 0 g/L. The anion exchange material may be an anion exchange material having substantially the same binding capacity for PEGylated EPO as Toyopearl Super Q 650 M equilibrated with 25 mM bicine, 7.5 mM Na_2SO_4 , at pH 8.0.

The AEC material may have a relatively high binding capacity for non-PEGylated protein, e.g. non-PEGylated EPO, of 20 – 50 g/L, 30 – 40 g/L, or about 35 g/L. The AEC material may have a binding capacity for non-PEGylated protein, e.g. non-PEGylated EPO, of at least about 35 g/L.

An anion exchange material may be equilibrated before the mixture of reaction products (or "first mixture") is applied to it. An AEC equilibration buffer may comprise a salt such as Na_2SO_4 and/or bicine. An AEC equilibration buffer may comprise a phosphate salt. An AEC equilibration buffer may not comprise compounds having primary amine groups, for example tris(hydroxymethyl)aminomethane (Tris). An AEC equilibration buffer may have a pH of about 6.5 to 9.5, 7.0 to 9.0, 7.5 to 8.5, or 8.0. An AEC equilibration buffer may comprise about 10-40 mM, 15-35 mM, 20-30 mM bicine or about 25 mM bicine. An AEC equilibration buffer may comprise may comprise about 1-20 mM, 1-10 mM, 2.5-10 mM, 5-10 mM salt or about 7.5 mM salt. An equilibration buffer for use in AEC may comprise about 25 mM bicine, about 7.5 mM Na_2SO_4 , pH 8.0. An equilibration buffer may have a conductivity of about 1 to about 8 mS/cm, about 1 to about 5 mS/cm, about 1.5 to about 3.0 mS/cm, about 2.0 to about 3.0 mS/cm, or about 1.0 to 3.0 mS/cm. If the mixture of reaction products has a conductivity outside the conductivity range of the equilibration buffer, then it may be conditioned (for example by addition of salt and/or buffer) to provide a mixture of reaction products having a conductivity that is within the conductivity range of the equilibration buffer, or is within 0.5 mS/cm or 0.05 mS/cm of the equilibration buffer.

The mixture of reaction products (or "first mixture"), comprising non-PEGylated, mono-PEGylated and oligo-PEGylated protein may be conditioned to provide a mixture of reaction products which has the pH, buffer and salt values set out above for the equilibration buffer. A conditioned mixture of reaction products may also be referred to as an AEC load composition or load solution. An AEC load solution may be an aqueous solution of non-PEGylated, mono-PEGylated and oligo-PEGylated protein in about 10 - 30 mM bicine, about 5.0 – 10.0 mM Na_2SO_4 , pH 7.0 – 9.0. An AEC load solution may be an aqueous solution of non-PEGylated, mono-PEGylated and oligo-PEGylated protein in about 25 mM bicine, about 7.5 mM Na_2SO_4 , pH 8.0.

The AEC step may be performed at a pH at least 1.0 to 2.0 pH units above the pI of the protein. Many proteins have a pI in the range 5.5 to 7.5 and therefore the processes of the invention may be used for producing a mono-PEGylated protein wherein the AEC step is performed at pH of about 6.5 to 9.5, or 6.5 to 8.5, or 7.5 to 8.5.

The pI of EPO is in the range 4.0 to 5.5. Processes in which the protein is EPO are may be performed at least 2.0, 2.5, or 3.0 pH units above the pI of the protein. This enables efficient EPO binding to the AEC material. Processes for producing a mono-PEGylated EPO composition may involve an AEC step performed at a pH of at least about 7.0, or about pH 7.0 to 10.0, or pH 7.0 to 9.0. The pH may be about 7.5 to 8.5, or about 7.8 to 8.2, or about 8.0.

The AEC step may be performed at a conductivity of about 1.0 – 3.0 mS/cm, 1.5 – 2.5 mS/cm or about 2.2 mS/cm. The conductivity of the AEC step may be adjusted by adjusting the concentration of a salt, such as NaCl or Na_2SO_4 in the AEC equilibration buffer or AEC wash buffer. The AEC step may be performed at Na_2SO_4 concentration of about 5.0 – 15 mM, 5.0 – 10 mM, or about 7.5 mM. The AEC step

may be performed using other salts, at concentrations that provide ionic strengths equivalent to those mentioned here for Na_2SO_4 .

After the mixture of reaction products (or "first mixture"), or AEC load solution, has been applied to the AEC material, the AEC material may be washed. The AEC material may be washed with equilibration
5 buffer. The AEC material may be washed with a wash buffer. A wash buffer may comprise about 10-40 mM, 15-35 mM, 20-30 mM bicine or about 25 mM bicine. A wash buffer may comprise may comprise about 1-20 mM, 1-10 mM, 2.5-10 mM, 5-10 mM salt or about 7.5 mM salt. It may comprise about 25 mM bicine, about 7.5 mM Na_2SO_4 , pH 8.0. A wash buffer buffer may have a conductivity of about 1 to about 8 mS/cm, about 1 to about 5 mS/cm, about 1.5 to about 3.0 mS/cm, about 2.0 to about 3.0 mS/cm. The
10 AEC material may be washed with 1 to 10, 1 to 5, or about 1, 2, 3, 4, or 5 column volumes of equilibration buffer or wash buffer. The AEC material may be washed with a volume of equilibration buffer or wash buffer that is determined by monitoring protein content of the flow-through solution (e.g. by UV spectrometry) and stopping the wash process once the protein content is below a threshold value (e.g. when UV signal returns to baseline). The flow-through solution, or effluent, may be collected as fractions,
15 and some or all of these flow-through fractions may be pooled.

Processes involving anion exchange chromatography are particularly preferred for production of mono-PEGylated EPO because they allow use of relatively high pH conditions under which EPO is relatively stable.

The process of the invention may comprise an AEC step wherein the AEC material is Toyopearl Super Q
20 650 M; the AEC step is performed at pH of about 7.0 to 9.0; the AEC step is performed at a conductivity of about 1.0 to 3.0 mS/cm; and the first mixture is applied to the AEC material as a AEC load solution comprising about 10 – 30 mM bicine and about 1 – 10 mM Na_2SO_4 . The protein may be erythropoietin.

Cation exchange chromatography

25 The IEC step may be a CEC step. The first mixture may be subjected to a cation exchange chromatography (CEC) step (step b). This provides an CEC flow-through solution in which the fraction of PEGylated protein is increased relative to the first mixture. This provides an CEC flow-through solution in which the fraction of non-PEGylated protein is decreased relative to the first mixture.

In the context of the present disclosure a fraction which is increased may be increased by an amount of at
30 least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100%. A fraction which is increased may be increased by at least about 1.5-fold, 2-fold, 3-fold, 5-fold or 10-fold.

In the context of the present disclosure a fraction which is decreased may be decreased by an amount of at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100%. A fraction which is decreased may be increased by at least about 1.5-fold, 2-fold, 3-fold, 5-fold or 10-fold.

A CEC step comprises contacting the first mixture with a cation exchange material, or applying the first mixture to an cation exchange material. The CEC step may comprise loading the first mixture onto a column comprising a cation exchange material. The cation exchange material has a low binding capacity for PEGylated protein, such as PEGylated EPO. That is, the CEC material has a relatively low binding capacity for the PEGylated version of the protein species for which a mono-PEGylated composition is desired. CEC conditions are used which are suitable for binding of the non-PEGylated protein to the cation exchange material. The solution that flows through the cation exchange material is the CEC flow-through solution. The low binding capacity of the cation exchange material for PEGylated protein has the effect that most of the PEGylated protein flows through the cation exchange material and is therefore present in the CEC flow-through solution. The binding conditions suitable for binding of the non-PEGylated protein to the cation exchange material have the effect that most of the non-PEGylated protein is removed (by binding to the cation exchange material) and is not present in the flow-through solution.

The CEC material may be a material that does not significantly bind PEGylated protein. This may mean that the CEC material binds less than about 10%, 5%, 2%, 1%, 0.5%, 0.1% or 0.01% of the PEGylated protein that is applied to it. Binding of PEGylated protein to the cation exchange material can also be reduced by applying high amounts of non-PEGylated protein to the cation exchange material. In this way, bound PEGylated protein is displaced by non-PEGylated protein. The amount of non-PEGylated protein applied to the cation exchange material may be about 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 110% or 120% of the dynamic break through capacity of the cation exchange material. The amount of non-PEGylated protein applied to the cation exchange material may be in the range of about 70-110%, about 80-100%, or about 85-95% of the dynamic break through capacity of the cation exchange material. The amount of non-PEGylated protein applied to the cation exchange material may be in the range of about 80-95% of the dynamic break through capacity of the cation exchange material.

The CEC flow-through solution comprises mono-PEGylated and oligo-PEGylated protein. The CEC flow-through solution may also comprise non-PEGylated protein, wherein the proportion of non-PEGylated protein is relatively low or close to zero. The CEC flow-through solution may comprise zero non-PEGylated protein, or amounts of non-PEGylated protein below the limit of detection. The protein may be EPO. The flow-through solution may comprise less than about 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1.5%, 1.0%, 0.5% or 0.1% non-PEGylated protein. The flow-through solution may comprise less than about 2% non-PEGylated protein. The flow-through solution may comprise less than about 2% non-PEGylated EPO.

The CEC flow-through solution may comprise at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or at least about 99.9% PEGylated protein.

The CEC flow-through may comprise about 80% mono-PEGylated protein and about 20 % oligo-PEGylated protein. The CEC flow-through solution may comprise at least about 60%, 65%, 70%, 75%, or 80% mono-PEGylated protein. The flow-through solution may comprise about 60-90%, 70-90%, or 75-

85% mono-PEGylated protein. The CEC flow-through solution may comprise at least about 20%, 25%, 30%, 35%, or 40% oligo-PEGylated protein. The flow-through solution may comprise about 10-40%, 10-30%, or 15-25% oligo-PEGylated protein.

5 The CEC flow-through solution may comprise (A) non-PEGylated protein in the range 0.1-5%, and (B) mono-PEGylated protein in the range 60-90%, and (C) oligo-PEGylated protein in the range 10-40%, wherein the total of (A), (B) and (C) is 100%. The flow-through solution may comprise (A) non-PEGylated protein in the range 0.1-1.5%, and (B) mono-PEGylated protein in the range 75-85%, and (C) oligo-PEGylated protein in the range 15-25%, wherein the total of (A), (B) and (C) is 100%.

10 The CEC flow-through solution may comprise (A) non-PEGylated protein, and (B) mono-PEGylated protein, and (C) oligo-PEGylated protein. The non-PEGylated protein, mono-PEGylated protein, and oligo-PEGylated protein may be present in the ratio A : B : C, wherein A is about 1-5, B is about 10-20, and C is about 5-15; or wherein A is about 1-3, B is about 25-35 and C is about 5-10. The non-PEGylated protein, mono-PEGylated protein, and oligo-PEGylated protein may be present in the ratio A:(B+C) wherein A:(B+C) is about 1:25; 1:50; 1:100; about 1:150, about 1:200, or about 1:100-200. The
15 ratio may be a weight or mass ratio. The ratio may be a molar ratio.

The CEC step is performed under conditions suitable for non-PEGylated protein to bind to the CEC material. Conditions suitable for non-PEGylated protein binding may be determined empirically by screening one or more of the CEC material, or the pH, conductivity, salt content or buffer content of the equilibration buffer.

20 The functional groups on an ion exchange material determine its charge. The terms "strong" and "weak" in this context refer to the extent that the ionisation state of the functional groups varies with varying pH. Strong ion exchangers show little or no variation in ion exchange capacity with change in pH. Strong ion exchangers remain fully charged over a broad pH range. Weak ion exchangers display pH-dependent function and deliver optimal performance over a narrower pH range.

25 Strong cation exchange materials may include a sulfonic acid group. A weak cationic exchange material may include carboxylic or phosphonic acid functional groups. An advantage of using strong cationic exchange materials is that the binding capacity is maintained at high or low pH since there is no loss of charge.

30 CEC materials may comprise a resin that has been functionalised with a negatively charged ligand or functional group, that is, with an acidic ligand or functional group.

CEC materials include POROS HS materials, Fractogel S materials, Fractogel EMD SO₃-(S), Fractogel EMD SO₃-(M), Fractogel SMD SE HiCap (M), Fractogel EMD COO- (M), and Capto S materials.

The cation exchange material used in the processes of the invention may be a strong cation exchange material. It may have a 50-150 nm, 70-120 nm, 70-100 nm or a 90-110 nm pore size. Pore size herein

may refer to mean pore size. It may have a particle size of 40-150 µm, 40-120 µm, 40-100 µm, 40-80 µm, or about 60, 65, 70, 80, or 90 100 µm. Particle size (or bead size) may refer to mean particle size. It may have a 70-100 nm pore size and a 35-100 µm bead size. It may have a 100 nm pore size and a 65 µm bead size. It may be SP Toyopearl 650 M.

- 5 The cation exchange material may be or may comprise a column, for example an expanded bed or packed bed column. The cation exchange material may be or may comprise a continuous countercurrent tangential chromatography system. The cation exchange material may be in the form of discrete particles or beads. The cation exchange material may be in the form of a membrane. The cation exchange material may be in the form of a functionalised filter, or a functionalised fibre, fleece, or mesh. The anion
10 exchange material may be in the form of any other solid support able to carry functional groups or exhibiting anion exchange properties.

- The CEC material in the processes of the invention may be a cation exchange material having a low binding capacity for the PEGylated form of the protein. In particular, it may be a cation exchange material having low binding capacity for PEGylated EPO. The cation exchange material may be SP Toyopearl
15 650 M having low binding capacity for PEGylated EPO, for example a binding capacity of less than about 0.5 g/L, 0.05 g/L, 0.01 g/L or 0.001 g/L, or close to 0 g/L.

The CEC material may have a relatively high binding capacity for non-PEGylated protein, e.g. non-PEGylated EPO, of 20 – 50 g/L, 30 – 40 g/L, or about 35 g/L. The CEC material may have a binding capacity for non-PEGylated protein, e.g. non-PEGylated EPO, of at least about 35 g/L.

20 Providing a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein

- As described above, the processes of the invention involve subjecting a mixture of reaction products ("first mixture") to an ion exchange chromatography (IEC) step (step b). The flow-through from the ion exchange chromatography step is used to provide a PEGylated protein mixture (or "second mixture") (step c). For example, the PEGylated protein mixture may comprise the collected flow-through from one
25 or more IEC steps. The step of providing a PEGylated protein mixture may comprise providing a batch of PEGylated protein mixture to an HIC apparatus (e.g. HIC column). Alternatively, the step of providing a PEGylated protein mixture may comprise providing the flow-through to an HIC apparatus (e.g. HIC column) via an in-line connection.

- The step of providing a PEGylated protein mixture (or "second mixture") (step c) comprises collecting the
30 flow-through solution from the IEC step. The step of providing a PEGylated protein mixture may comprise pooling the flow-through solution from two, three, four or five IEC steps, or more than three IEC steps. The step of providing a PEGylated protein mixture may comprise conditioning the second mixture. Conditioning the PEGylated protein mixture may provide a conditioned PEGylated protein mixture having improved suitability for separating the mono-PEGylated protein from the oligo-PEGylated protein
35 contained therein. Conditioning the PEGylated protein mixture may provide a conditioned PEGylated

protein mixture having improved suitability for separating the mono-PEGylated protein from the oligo-PEGylated protein contained therein by HIC. A conditioned PEGylated protein mixture may be referred to as a HIC load composition or load solution. Conditioning the PEGylated protein mixture may comprise addition of salt, and/or addition of bicine.

- 5 For example, as described above, the processes of the invention involve subjecting a mixture of reaction products ("first mixture") to an anion exchange chromatography (AEC) step (step b). The flow-through from the anion exchange chromatography step is used to provide a PEGylated protein mixture (or "second mixture") (step c). For example, the PEGylated protein mixture may comprise the collected flow-through from one or more AEC steps. The step of providing a PEGylated protein mixture may comprise
- 10 providing a batch of PEGylated protein mixture to an HIC apparatus (e.g. HIC column). Alternatively, the step of providing a PEGylated protein mixture may comprise providing the flow-through to an HIC apparatus (e.g. HIC column) via an in-line connection.

- The step of providing a PEGylated protein mixture (or "second mixture") (step c) comprises collecting the flow-through solution from the AEC step. The step of providing a PEGylated protein mixture may
- 15 comprise pooling the flow-through solution from two, three, four or five AEC steps, or more than three AEC steps. The step of providing a PEGylated protein mixture may comprise conditioning the second mixture. Conditioning the PEGylated protein mixture may provide a conditioned PEGylated protein mixture having improved suitability for separating the mono-PEGylated protein from the oligo-PEGylated protein contained therein. Conditioning the PEGylated protein mixture may provide a conditioned
- 20 PEGylated protein mixture having improved suitability for separating the mono-PEGylated protein from the oligo-PEGylated protein contained therein by HIC. A conditioned PEGylated protein mixture may be referred to as a HIC load composition or load solution. Conditioning the PEGylated protein mixture may comprise addition of salt, and/or addition of bicine.

Recycling unreacted protein

- 25 The processes involve performing a PEGylation reaction to provide a mixture of reaction products. The non-PEGylated (unreacted) protein is recycled by including it in a subsequent PEGylation reaction. Non-PEGylated protein is recovered in step b) by eluting an IEC eluate from the IEC material. The eluted non-PEGylated protein is then used in a subsequent PEGylation reaction. The eluted non-PEGylated protein may be used in a subsequent step a) PEGylation reaction.
- 30 The processes may involve performing two or more cycles of steps a), b) and c), in which non-PEGylated protein is recovered in step b) by eluting an IEC eluate from the IEC material. The eluted non-PEGylated protein is then used in a subsequent PEGylation reaction. The eluted non-PEGylated protein may be used in a subsequent step a) PEGylation reaction. The eluted non-PEGylated protein may be used in the PEGylation reaction of step a) of the next cycle. That is, further cycles of steps a), b) and c) may be
- 35 performed in which non-PEGylated protein from a previous cycle is used in the PEGylation reaction of that further cycle. Alternatively, a cycle of steps a), b) and c) may be followed by a subsequent

PEGylation reaction (which may be a step a) PEGylation reaction) and a processing step which provides a solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products and which allows recovery of non-PEGylated protein. That is, cycles of steps a), b) and c) may be followed by, or alternated with, a PEGylation reaction and a processing step different to but having similar effects as the step b) IEC step. Recovered non-PEGylated protein may be used in a subsequent PEGylation reaction. Recovered non-PEGylated protein may be used in a subsequent step a) PEGylation reaction. Recovered non-PEGylated protein may be used in a subsequent cycle of steps a), b) and c).

For example, the processes involve performing a PEGylation reaction to provide a mixture of reaction products. The non-PEGylated (unreacted) protein is recycled by including it in a subsequent PEGylation reaction. Non-PEGylated protein is recovered in step b) by eluting an AEC eluate from the AEC material. The eluted non-PEGylated protein is then used in a subsequent PEGylation reaction. The eluted non-PEGylated protein may be used in a subsequent step a) PEGylation reaction.

The processes may involve performing two or more cycles of steps a), b) and c), in which non-PEGylated protein is recovered in step b) by eluting an IEC eluate from the IEC material. The eluted non-PEGylated protein is then used in a subsequent PEGylation reaction. The eluted non-PEGylated protein may be used in a subsequent step a) PEGylation reaction. The eluted non-PEGylated protein may be used in the PEGylation reaction of step a) of the next cycle. That is, further cycles of steps a), b) and c) may be performed in which non-PEGylated protein from a previous cycle is used in the PEGylation reaction of that further cycle. Alternatively, a cycle of steps a), b) and c) may be followed by a subsequent PEGylation reaction (which may be a step a) PEGylation reaction) and a processing step which provides a solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products and which allows recovery of non-PEGylated protein. That is, cycles of steps a), b) and c) may be followed by, or alternated with, a PEGylation reaction and a processing step different to but having similar effects as the step b) IEC step. Recovered non-PEGylated protein may be used in a subsequent PEGylation reaction. Recovered non-PEGylated protein may be used in a subsequent step a) PEGylation reaction. Recovered non-PEGylated protein may be used in a subsequent cycle of steps a), b) and c).

Various processing steps suitable for providing a solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products and which allow recovery of non-PEGylated protein exist and will be known to the skilled person. These include, for example, chromatographic methods such as cation exchange chromatography, ion exchange chromatography, and hydrophobic interaction chromatography.

A "cycle" as used herein refers to a full cycle of steps a), b), and c).

The processes may comprise three, four or five cycles, wherein step b) of each cycle comprises eluting non-PEGylated protein from the IEC material to provide an IEC eluate, and wherein the non-PEGylated

protein eluted in step b) is used in a subsequent PEGylation reaction. The non-PEGylated protein eluted in step b) may be added to the PEGylation reaction of step a) in a subsequent cycle.

The processes may comprise pooling the non-PEGylated protein recovered in step b) from multiple IEC steps before it is used in a subsequent PEGylation reaction. The processes may comprise pooling the non-PEGylated protein recovered in step b) from multiple IEC steps before it is used in a subsequent step a) PEGylation reaction. The processes may comprise pooling the non-PEGylated protein recovered from two, three, four or five step b) IEC steps, or more than five step b) IEC steps.

The processes may comprise adding the step b) IEC eluate directly to a subsequent PEGylation reaction. The processes may comprise adding the IEC eluate from step b) of a cycle directly to the PEGylation reaction of step a) of the next cycle. For example the process may comprise adding the IEC eluate from step b) of a first cycle directly to the PEGylation reaction of step a) of the second cycle. Adding directly in this context means that no intervening step of purifying or cleaning the non-PEGylated protein in the eluate (e.g. by diafiltration or ultrafiltration). Rather, the eluate from the IEC material that contains the non-PEGylated protein is added to a subsequent PEGylation reaction.

The non-PEGylated protein used in a subsequent PEGylation reaction may be supplemented with fresh protein. The process may comprise adding fresh protein to the PEGylation reaction in addition to the non-PEGylated protein recovered from the IEC material. The step of recycling non-PEGylated protein comprises adding non-PEGylated protein recovered from the IEC material to a subsequent PEGylation reaction, and this may further comprise adding fresh protein to the PEGylation reaction such that the starting concentration of protein in the PEGylation reactions is substantially constant. Substantially constant in this context may mean that the starting concentration of protein in the second and subsequent PEGylation reactions is substantially the same as that in the first PEGylation reaction. This may mean that the starting concentration of protein in the second and subsequent PEGylation reactions is substantially the same as that in the PEGylation reaction of the first cycle. Substantially the same may mean $\pm 10\%$. Fresh protein refers to protein that has not previously been subject to a PEGylation reaction.

The processes may involve cycles of steps a), b) and c), such that a first and a second cycle are performed and the second cycle is the final cycle. Alternatively the processes of the invention may involve two further cycles of steps a), b), and c), such that a first, a second and a third cycle is performed, and the third cycle is the final cycle. The processes may involve more than two further cycles of steps a), b) and c). The processes may involve a total of one, two, three, four, or five cycles of steps a), b) and c).

The final cycle may comprise performing a PEGylation reaction wherein the PEG/protein molar ratio is from about 1.4 to 1 to about 2 to 1. The molar ratio of PEG/protein may be about 1.4 to 1.0, 1.5 to 1.0, 1.6 to 1.0, 1.7 to 1.0, 1.8 to 1.0, 1.9 to 1.0, 2.0 to 1.0, 2.1 to 1.0, 2.2 to 1.0, 2.3 to 1.0, 2.4 to 1.0 or 2.5 to 1.0. A relatively high PEG/protein ratio may be advantageous in the final cycle. After the final cycle there is no recycling of unreacted (non-PEGylated) protein and therefore it may be advantageous to minimise

waste of protein by minimising unreacted protein. That is, in the final cycle it may be advantageous to maximise protein PEGylation, and in particular mono-PEGylation.

The final cycle may comprise performing a PEGylation reaction at about pH 7.0 to 9.0. The PEGylation reaction may be performed at about pH 7.5 to 8.5, or at about pH 8.0. The PEGylation reaction may be performed at about 15-25°C, or about 20°C. The PEGylation reaction may have a reaction time of 30 - 120 minutes or about 60 minutes. The PEGylation may use an NHS-activated PEG.

The non-PEGylated protein may be recovered in the IEC step by eluting it from the IEC material. Recovering the non-PEGylated protein may therefore involve providing an eluate comprising the non-PEGylated protein in a relatively concentrated form and/or a relatively pure form. The eluate may comprise at least about 95% non-PEGylated protein.

Eluting non-PEGylated protein from the IEC material may comprise contacting the IEC material with an elution buffer. The IEC elution buffer may comprise about 40-100 mM, 50-90 mM, 60-80 mM NaCl, or about 70 mM NaCl, or at least about 40 mM, 50 mM, 60 mM or 70 mM NaCl, or about 20-60 mM, 40-50 mM, or about 45 mM NaCl. The elution buffer may comprise less than or equal to about 45 mM NaCl. The elution buffer may comprise about 20-50 mM, 25-45 mM, 30-40 mM Na₂SO₄, or about 35 mM Na₂SO₄, or at least about 20 mM, 25 mM, 30 mM or 35 mM Na₂SO₄. Other salts may be suitable for use in the IEC elution buffer, which may provide the same amount of negatively charged ions as the concentration ranges indicated for NaCl and Na₂SO₄. For example the elution buffer may contain about 70 mM NaCl or about 35 mM Na₂SO₄. The elution buffer may comprise about 20-30 mM, or about 25 mM bicine. The elution buffer may have a pH of about 7.0 to 9.0, 7.5 to 8.5, or about 8.0. The elution buffer may comprise about 35 mM Na₂SO₄, about 25 mM bicine, pH about 8.0.

The elution buffer for the IEC material may contain salt in an amount of less than or equal to about 60 mM, 55 mM, 50 mM, 45 mM, 40 mM or 35 mM. The elution buffer may contain salt in an amount of less than about 60 mM, 55 mM, 50 mM, 45 mM, or 40 mM. The salt may be Na₂SO₄. The salt may be a mixture of salts comprising Na₂SO₄. The elution buffer for the IEC material may have a conductivity of about 5-20 mS/cm, 5-15 mS/cm, or 8-10 mS/cm. The elution buffer may have a conductivity of less than or equal to about 20 mS/cm, 15mS/cm, or 10 mS/cm.

Alternatives to bicine, for use the IEC elution buffer (or in other buffer solutions discussed herein), include for example other "Good's buffers". Buffers having a pK_a from about 6 to 10 may be used. Phosphate or HEPES may be used. Buffers that do not contain a primary amine are particularly suitable (because a primary amine may act as a partner for the PEG reagent resulting in some PEGylated buffer molecules).

Alternatives to Na₂SO₄, for use in the IEC elution buffer (or in other buffer solutions discussed herein) may include for example salts comprising an anion selected from: PO₄³⁻, SO₄²⁻, CH₃COO⁻, Cl⁻, Br⁻, NO₃⁻, ClO₄⁻, I⁻, SCN⁻; and a cation selected from: NH₄⁺, Rb⁺, K⁺, Na⁺, Cs⁺, Li⁺, Mg²⁺, Ca²⁺, Ba²⁺. Alternative salts to Na₂SO₄ may include NaCl, KCl, NaHPO₄, LiCl, and KSCN.

For example, the processes may involve performing two or more cycles of steps a), b) and c), in which non-PEGylated protein is recovered in step b) by eluting an AEC eluate from the AEC material. The eluted non-PEGylated protein is then used in a subsequent PEGylation reaction. The eluted non-PEGylated protein may be used in a subsequent step a) PEGylation reaction. The eluted non-PEGylated protein may be used in the PEGylation reaction of step a) of the next cycle. That is, further cycles of steps a), b) and c) may be performed in which non-PEGylated protein from a previous cycle is used in the PEGylation reaction of that further cycle. Alternatively, a cycle of steps a), b) and c) may be followed by a subsequent PEGylation reaction (which may be a step a) PEGylation reaction) and a processing step which provides a solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products and which allows recovery of non-PEGylated protein. That is, cycles of steps a), b) and c) may be followed by, or alternated with, a PEGylation reaction and a processing step different to but having similar effects as the step b) AEC step. Recovered non-PEGylated protein may be used in a subsequent PEGylation reaction. Recovered non-PEGylated protein may be used in a subsequent step a) PEGylation reaction. Recovered non-PEGylated protein may be used in a subsequent cycle of steps a), b) and c).

Various processing steps suitable for providing a solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products and which allow recovery of non-PEGylated protein exist and will be known to the skilled person. These include, for example, chromatographic methods such as cation exchange chromatography, anion exchange chromatography, and hydrophobic interaction chromatography.

A "cycle" as used herein refers to a full cycle of steps a), b), and c).

The processes may comprise three, four or five cycles, wherein step b) of each cycle comprises eluting non-PEGylated protein from the AEC material to provide an AEC eluate, and wherein the non-PEGylated protein eluted in step b) is used in a subsequent PEGylation reaction. The non-PEGylated protein eluted in step b) may be added to the PEGylation reaction of step a) in a subsequent cycle.

The processes may comprise pooling the non-PEGylated protein recovered in step b) from multiple AEC steps before it is used in a subsequent PEGylation reaction. The processes may comprise pooling the non-PEGylated protein recovered in step b) from multiple AEC steps before it is used in a subsequent step a) PEGylation reaction. The processes may comprise pooling the non-PEGylated protein recovered from two, three, four or five step b) AEC steps, or more than five step b) AEC steps.

The processes may comprise adding the step b) AEC eluate directly to a subsequent PEGylation reaction. The processes may comprise adding the AEC eluate from step b) of a cycle directly to the PEGylation reaction of step a) of the next cycle. For example the process may comprise adding the AEC eluate from step b) of a first cycle directly to the PEGylation reaction of step a) of the second cycle. Adding directly in this context means that no intervening step of purifying or cleaning the non-PEGylated

protein in the eluate (e.g. by diafiltration or ultrafiltration). Rather, the eluate from the AEC material that contains the non-PEGylated protein is added to a subsequent PEGylation reaction.

- The non-PEGylated protein used in a subsequent PEGylation reaction may be supplemented with fresh protein. The process may comprise adding fresh protein to the PEGylation reaction in addition to the non-PEGylated protein recovered from the AEC material. The step of recycling non-PEGylated protein comprises adding non-PEGylated protein recovered from the AEC material to a subsequent PEGylation reaction, and this may further comprise adding fresh protein to the PEGylation reaction such that the starting concentration of protein in the PEGylation reactions is substantially constant. Substantially constant in this context may mean that the starting concentration of protein in the second and subsequent PEGylation reactions is substantially the same as that in the first PEGylation reaction. This may mean that the starting concentration of protein in the second and subsequent PEGylation reactions is substantially the same as that in the PEGylation reaction of the first cycle. Substantially the same may mean $\pm 10\%$. Fresh protein refers to protein that has not previously been subject to a PEGylation reaction.
- The processes may involve cycles of steps a), b) and c), such that a first and a second cycle are performed and the second cycle is the final cycle. Alternatively the processes of the invention may involve two further cycles of steps a), b), and c), such that a first, a second and a third cycle is performed, and the third cycle is the final cycle. The processes may involve more than two further cycles of steps a), b) and c). The processes may involve a total of one, two, three, four, or five cycles of steps a), b) and c).
- The final cycle may comprise performing a PEGylation reaction wherein the PEG/protein molar ratio is from about 1.4 to 1 to about 2 to 1. The molar ratio of PEG/protein may be about 1.4 to 1.0, 1.5 to 1.0, 1.6 to 1.0, 1.7 to 1.0, 1.8 to 1.0, 1.9 to 1.0, 2.0 to 1.0, 2.1 to 1.0, 2.2 to 1.0, 2.3 to 1.0, 2.4 to 1.0 or 2.5 to 1.0. A relatively high PEG/protein ratio may be advantageous in the final cycle. After the final cycle there is no recycling of unreacted (non-PEGylated) protein and therefore it may be advantageous to minimise waste of protein by minimising unreacted protein. That is, in the final cycle it may be advantageous to maximise protein PEGylation, and in particular mono-PEGylation.

- The final cycle may comprise performing a PEGylation reaction at about pH 7.0 to 9.0. The PEGylation reaction may be performed at about pH 7.5 to 8.5, or at about pH 8.0. The PEGylation reaction may be performed at about 15-25°C, or about 20°C. The PEGylation reaction may have a reaction time of 30 - 120 minutes or about 60 minutes. The PEGylation may use an NHS-activated PEG.

The non-PEGylated protein may be recovered in the AEC step by eluting it from the AEC material. Recovering the non-PEGylated protein may therefore involve providing an eluate comprising the non-PEGylated protein in a relatively concentrated form and/or a relatively pure form. The eluate may comprise at least about 95% non-PEGylated protein.

Eluting non-PEGylated protein from the AEC material may comprise contacting the AEC material with an elution buffer. The AEC elution buffer may comprise about 40-100 mM, 50-90 mM, 60-80 mM NaCl, or about 70 mM NaCl, or at least about 40 mM, 50 mM, 60 mM or 70 mM NaCl. The elution buffer may comprise about 20-50 mM, 25-45 mM, 30-40 mM Na₂SO₄, or about 35 mM Na₂SO₄, or at least about 20 mM, 25 mM, 30 mM or 35 mM Na₂SO₄. Other salts may be suitable for use in the AEC elution buffer, which may provide the same amount of negatively charged ions as the concentration ranges indicated for NaCl and Na₂SO₄. For example the elution buffer may contain about 70 mM NaCl or about 35 mM Na₂SO₄. The elution buffer may comprise about 20-30 mM, or about 25 mM bicine. The elution buffer may have a pH of about 7.0 to 9.0, 7.5 to 8.5, or about 8.0. The elution buffer may comprise about 35 mM Na₂SO₄, about 25 mM bicine, pH about 8.0.

The elution buffer for the AEC material may contain salt in an amount of less than or equal to about 60 mM, 55 mM, 50 mM, 45 mM, 40 mM or 35 mM. The elution buffer may contain salt in an amount of less than about 60 mM, 55 mM, 50 mM, 45 mM, or 40 mM. The salt may be Na₂SO₄. The salt may be a mixture of salts comprising Na₂SO₄. The elution buffer for the AEC material may have a conductivity of about 5-20 mS/cm, 5-15 mS/cm, or 8-10 mS/cm. The elution buffer may have a conductivity of less than or equal to about 20 mS/cm, 15mS/cm, or 10 mS/cm.

Alternatives to bicine, for use the AEC elution buffer (or in other buffer solutions discussed herein), include for example other "Good's buffers". Buffers having a pK_a from about 6 to 10 may be used. Phosphate or HEPES may be used. Buffers that do not contain a primary amine are particularly suitable (because a primary amine may act as a partner for the PEG reagent resulting in some PEGylated buffer molecules).

Alternatives to Na₂SO₄, for use in the AEC elution buffer (or in other buffer solutions discussed herein) may include for example salts comprising an anion selected from: PO₄³⁻, SO₄²⁻, CH₃COO⁻, Cl⁻, Br⁻, NO₃⁻, ClO₄⁻, I⁻, SCN⁻; and a cation selected from: NH₄⁺, Rb⁺, K⁺, Na⁺, Cs⁺, Li⁺, Mg²⁺, Ca²⁺, Ba²⁺. Alternative salts to Na₂SO₄ may include NaCl, KCl, NaHPO₄, LiCl, and KSCN.

Performance of AEC and HIC steps

The IEC and HIC steps in the processes of the invention may be performed sequentially. For example the AEC and HIC steps in the processes of the invention may performed sequentially. That is the process may comprise sequential steps of a) providing a first mixture comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein b) subjecting the first mixture to an anion exchange chromatography (AEC) step to provide an AEC flow-through solution in which the fraction of PEGylated protein is increased relative to the first mixture; the AEC step comprising applying the first mixture to an AEC material under conditions suitable for binding non-PEGylated protein; c) collecting the AEC flow-through solution from step b) to provide a second mixture comprising mono-PEGylated protein and oligo-PEGylated protein; and d) subjecting the second mixture to a hydrophobic interaction chromatography (HIC) step to provide a mono-

PEGylated protein composition in which the fraction of mono-PEGylated protein is increased relative to the second mixture, wherein the mono-PEGylated protein composition comprises at least about 90% mono-PEGylated protein.

The term "sequential" means that no intervening chromatography step occurs between any of steps (a) to (d) (no chromatography step between steps (a) and (b), (b) and (c), (c) and (d)). The term "sequential" means that no intervening chromatography step occurs between any of the steps recited in any one of the claims. The steps of the processes of the invention may be performed directly, meaning that each of steps (b) to (d) is performed directly following the previous step. The process may consist of steps (a) to (d). The process may consist of the steps recited in any one of the claims.

Steps (b) to (d) may be performed discontinuously or continuously. Steps (b) to (d) may be performed sequentially and discontinuously or sequentially and continuously. Continuously means that the AEC material and the HIC material are connected directly or there is some other mechanism which allows for continuous flow between the AEC material and the HIC material. Discontinuously means that there is no continuous flow between the AEC material and the HIC material, for example the AEC flow-through solution or effluent is collected and pooled to provide the second mixture. Performance of the steps continuously may mean that the same flow rate is used for the entire process.

The IEC flow-through solution may be applied directly to the HIC material. Preferably a conditioning step is performed to directly provide the HIC material with a conditioned second mixture. The IEC material and HIC material may be directly connected in series. For example the IEC material may be comprised in an IEC column and the HIC material may be comprised in a HIC column, and the IEC column is directly connected to the HIC column. The IEC flow-through solution from the IEC column may be directly delivered to the HIC column. There may be an in-line connection between the IEC column and the HIC column. There may be a mechanism that permits continuous flow between the IEC column and the HIC column. The steps of the process may be performed continuously. The steps of the process may be performed in parallel, that is, step (d) may be begun before step (b) is completed, such that steps (b) and (d) run in parallel. That is HIC step and IEC step may be run in parallel. Performing the process continuously (or in parallel) is advantageous because it is relatively fast and efficient.

When the IEC step is a CEC step, the CEC and HIC steps can be performed sequentially, and/or can be performed continuously or discontinuously, as described for the AEC and HIC steps above.

Purification processes separating mono-PEGylated protein and oligo-PEGylated protein

The processes of the invention use an IEC material with a relatively low binding capacity for the PEGylated form of the protein and a relatively high binding capacity for the non-PEGylated form of the protein. This is advantageous because it facilitates rapid separation of non-PEGylated protein from PEGylated protein. Much of the binding capacity of the column is used for binding to the non-PEGylated form of the protein which facilitates the processing of relatively large volumes of first mixture (the mixture comprising non-PEGylated mono-PEGylated and oligo-PEGylated protein) using a relatively small

column. This is because the first mixture is generally relatively low in non-PEGylated protein (which the IEC binding material has a relatively high binding capacity for) and high in PEGylated protein (which the IEC binding material has a relatively low binding capacity for) – this means that in operation the processes disclosed herein can process relatively large volumes of first mixture in the IEC step. The relatively rapid separation of non-PEGylated protein from PEGylated protein facilitates rapid recycling of non-PEGylated protein and overall process efficiency.

As described above, the present invention provides processes for producing a mono-PEGylated protein composition comprising at least about 90% mono-PEGylated protein, the process comprising: subjecting the PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein produced as an IEC flow-through solution described above to a purification process separating mono-PEGylated protein and oligo-PEGylated protein. The IEC flow-through solution may be an AEC flow-through solution.

Various suitable purification processes exist and will be known to the skilled person, for example, precipitation, size dependent separation, and filtration methods, as well as chromatographic methods such as size exclusion chromatography, cation exchange chromatography, anion exchange chromatography, and hydrophobic interaction chromatography. In the processes described herein hydrophobic interaction chromatography may be a preferred purification process.

Hydrophobic interaction chromatography

HIC materials may comprise a resin that has been functionalised with a hydrophobic ligand. Hydrophobic ligands on HIC materials may interact with hydrophobic surfaces of proteins. The ligand and the degree of substitution (high or low substitution “sub”) on a HIC material may contribute to its final hydrophobicity and thereby to its selectivity. The ligand may contain alkyl or aryl groups. The hydrophobicity of HIC materials increases through the ligand series: ether, polypropyleneglycol (PPG), phenyl, butyl and hexyl.

The HIC material used in the processes of the invention may comprise a phenyl ligand. It may comprise a methacrylic resin functionalised with a phenyl ligand. It may be a hydroxylated polymethacrylic polymer bead functionalised with a phenyl ligand. It may have a 50-150 nm, 70-120 nm, 70-100 nm, 90-110 nm pore size. It may have a particle size of 40-150 µm, 40-120 µm, 40-100 µm, 40-80 µm, or about 60, 65, 70, 80, or 90 100 µm. It may have a 70-100 nm pore size and a 30-65 µm bead size. It may have a 100 nm pore size and a 65 µm bead size. It may be Toyopearl Phenyl 650M (Tosoh Bioscience LLC; product number 14478). The HIC material used in the processes of the invention may be used in accordance with the manufacturer’s instructions.

The HIC material may be or may comprise a column, for example an expanded bed or packed bed column. The HIC material may be in the form of a porous monolithic material. The HIC material may be in the form of a functionalised fibre, fleece, or mesh. The HIC material may be in the form of any other solid support able to carry functional groups or exhibiting HIC properties. HIC materials include

Toyopearl Phenyl 650M, Toyopearl Phenyl 650S, Toyopearl PPG 600M, TSKgel Phenyl 5PW, Butyl Sepharose 4 FF, Butyl-S Sepharose FF, Octyl Sepharose 4 FF, Phenyl Sepharose BB, Phenyl Sepharose HP, Phenyl Sepharose 6 FF High Sub, Phenyl Sepharose 6 FF Low Sub, Source I SETH, Source 151 SO, Source 1 SPHE, Phenyl Sepharose BB, Phenyl Sepharose HP, Phenyl Sepharose 6 FF High Sub, Phenyl Sepharose 6 FF Low Sub, Source I SETH, Source 151 SO, Source 1 SPHE, Cellufine Butyl, Cellufine Octyl, Cellufine Phenyl, WP HI-Propyl (C3), Macroprep t-Butyl, Macroprep methyl.

The processes of the invention may comprise a HIC step in flow through mode. In flow through mode the PEGylated protein mixture (or "second mixture") is applied to a HIC material under conditions suitable for mono-PEGylated protein to flow through the HIC material and is therefore present in the HIC flow-through solution. The conditions are suitable for oligo-PEGylated protein to bind to the HIC material.

The process of the invention may comprise a HIC step in bind and elute mode. In bind and elute mode the PEGylated protein mixture (or "second mixture") is applied to a HIC material under conditions suitable for mono-PEGylated protein and oligo-PEGylated protein to bind to the HIC material. The mono-PEGylated protein is eluted from the HIC material. A gradient of decreasing salt is used to elute the mono-PEGylated protein. The oligo-PEGylated protein is eluted at a lower salt concentration and therefore in a separate fraction from the mono-PEGylated protein.

A HIC material may be equilibrated before the PEGylated protein mixture (or "second mixture" or HIC load solution), is loaded on to it. An HIC equilibration buffer may comprise a salt such as Na_2SO_4 and/or bicine. An HIC equilibration buffer may have a pH of about 6.5 to 9.0, 7.5 to 8.5, or 8.0. An HIC equilibration buffer may comprise about 10-40 mM, 15-35 mM, 20-30 mM bicine or about 25 mM bicine. An HIC equilibration buffer may comprise about 10-40 mM, 15-35 mM, 20-30 mM bicine or about 25 mM NaPO_4 .

Conditions suitable for mono-PEGylated protein to flow through the HIC material while oligo-PEGylated protein is bound to the HIC material can be readily determined. The pH, ionic strength, and composition of the equilibration buffer may be selected to ensure that when a protein mixture is loaded onto the chromatography matrix, the desired binding and/or flow through or specific proteins or contaminants is achieved. For example, the pH, ionic strength, and composition of the equilibration buffer may be selected to ensure that mono-PEGylated protein flows through the HIC material while oligo-PEGylated protein is bound to the HIC material.

For equilibration before a HIC step in flow through mode is carried out, an equilibration buffer may comprise about 200-500 mM, 250-450 mM, 300-450 mM, 300-400 mM, 350-400 mM salt, or about 390 mM salt or about 300 mM salt. An equilibration buffer for use in flow through mode HIC may comprise about 25 mM bicine, about 390 mM Na_2SO_4 , pH 8.0. An equilibration buffer for use in flow through mode HIC may comprise about 25 mM bicine, about 300 mM Na_2SO_4 , pH 8.0. An equilibration buffer for use in flow through mode HIC may comprise about 25 mM NaPO_4 , about 300 mM Na_2SO_4 , pH 8.0. An

equilibration buffer for use in flow through mode HIC may have a conductivity of about 30-70 mS/cm, 40-60 mS/cm, or about 50 mS/cm.

When the process comprises a HIC step in flow through mode, the process may involve providing a conditioned PEGylated protein mixture (or "second mixture") comprising salt (such as Na₂SO₄) and
5 bicine, which may be present as about 300 mM Na₂SO₄ and about 25 mM bicine, or which may be present as about 390 mM Na₂SO₄ and about 25 mM bicine. A conditioned PEGylated protein mixture may be referred to as a HIC load solution. A conditioned PEGylated protein mixture for use in HIC flow
through mode may comprise about 200-500 mM, 250-450 mM, 250-350 mM salt, or about 300 mM salt, or about 350 – 450 mM salt, or about 390 mM salt. The salt may be Na₂SO₄. A conditioned PEGylated
10 protein mixture may comprise about 10-40 mM, 15-35 mM, 20-30 mM bicine or about 25 mM bicine. A conditioned PEGylated protein mixture may comprise about 300 mM Na₂SO₄ and about 25 mM bicine.

In HIC flow through mode, after the PEGylated protein mixture has been applied to the HIC material the HIC material may be washed. The HIC material may be washed with equilibration buffer that is suitable for use in flow through mode. The HIC material may be washed with a HIC wash buffer, which may
15 comprise about 200-500 mM, 250-450 mM, 300-450 mM, 300-400mM, 350-400 mM salt, or about 390 mM salt, or about 300 mM salt. A wash buffer for use in flow through mode HIC may comprise about 25 mM bicine, about 390 mM Na₂SO₄, pH 8.0. A wash buffer for use in flow through mode HIC may comprise about 25 mM bicine, about 300 mM Na₂SO₄, pH 8.0. The HIC material may be washed with 1
to 10, 1 to 5, or about 1, 2, 3, 4, or 5 column volumes of buffer. The HIC material may be washed with a
20 volume of equilibration buffer or wash buffer that is determined by monitoring protein content of the flow-through solution (e.g. by UV spectrometry) and stopping the wash process once the protein content is below a threshold value (e.g. when UV signal returns to baseline). The flow-through solution, or effluent, may be collected as fractions, and some or all of these flow-through fractions may be pooled.

The PEGylated protein mixture (or "second mixture") may comprise zero, or close to zero, non-PEGylated
25 protein. The PEGylated protein mixture may comprise less than about 10%, 5%, 4%, 3%, 2%, 1.5%, 1.0%, 0.5% or 0.1% non-PEGylated protein. The PEGylated protein mixture may comprise less than about 2% non-PEGylated protein. The PEGylated protein mixture may comprise less than about 2% non-PEGylated EPO.

When the process of the invention comprises a HIC step in flow through mode the non-PEGylated protein
30 content of the PEGylated protein mixture should be below the threshold specification for the final therapeutic product.

When the HIC step is performed in flow-through mode, the HIC flow-through provides the mono-PEGylated protein composition. That is the HIC flow-through is a mono-PEGylated protein composition as described herein. The HIC flow-through may comprise at least 90% mono-PEGylated protein.

When the HIC step is performed in flow through mode, oligo-PEGylated protein may be eluted from the HIC material. For example by a gradient or step elution at a conductivity of 0 mS/cm.

For equilibration before an HIC step in bind and elute mode is carried out, an equilibration buffer may comprise about 400-600 mM, 450-550 mM, salt, or about 500 mM salt. An equilibration buffer for use in
5 bind and elute mode HIC may comprise about 25 mM bicine, about 500 mM Na₂SO₄, pH 8.0. An equilibration buffer for use in bind and elute mode HIC may have a conductivity of about 40-80 mS/cm, 50-70 mS/cm, or about 60 mS/cm or about 58 mS/cm.

When the process comprises a HIC step in bind and elute mode, the process may involve providing a conditioned PEGylated protein mixture (or "second medium") comprising salt (such as Na₂SO₄) and
10 bicine, which may be in any combination of the previously-mentioned concentrations, and which may be present as about 500 mM Na₂SO₄ and about 25 mM bicine, and the elution step may comprise elution with a decreasing gradient of 500 mM to 0 mM Na₂SO₄. . A conditioned PEGylated protein mixture may be referred to as a HIC load solution. A conditioned PEGylated protein mixture for use in HIC bind and elute mode may comprise about 250-750 mM, 400-600 mM, 450-550 mM salt, or about 500 mM salt.
15 The salt may be Na₂SO₄. Other salts may be used, such as NaCl. Other salts may be used at concentrations that provide ionic strength equivalent to the Na₂SO₄ concentrations. A conditioned PEGylated protein mixture may comprise about 10-40 mM, 15-35 mM, 20-30 mM bicine or about 25 mM bicine. A conditioned PEGylated protein mixture may comprise about 500 mM Na₂SO₄ and about 25 mM bicine.

When the HIC step is performed in bind and elute mode, the mono-PEGylated protein is eluted from the HIC material. Elution is by decreasing salt gradient. Elution may comprise applying a linear elution gradient from 500 mM to 0 mM salt, such as Na₂SO₄. Elution may comprise applying a linear elution gradient from about 60 mS/cm to 0 mS/cm. In a decreasing salt gradient mono-PEGylated EPO elutes at approximately 300 mM Na₂SO₄ (see Figure 6). Eluting the mono-PEGylated protein from the HIC
20 material to provides a HIC eluate, wherein the HIC eluate provides the mono-PEGylated protein.
25

When the HIC step is performed in flow-through mode, the HIC flow-through provides the mono-PEGylated protein composition. That is the HIC flow-through is a mono-PEGylated protein composition as described herein. The HIC flow-through may comprise at least 90% mono-PEGylated protein.

The PEGylated protein mixture (or "second mixture") may be conditioned for the HIC step in one or more
30 batches. The PEGylated protein mixture may be conditioned in-line. The step of providing a second mixture (step c) may comprise conditioning the mixture in-line, to provide a conditioned medium directly from the IEC apparatus (e.g. a AEC or CEC column) to the HIC apparatus (e.g. HIC column). For example, the step of providing a PEGylated protein mixture (step c) may comprise conditioning the mixture in-line, to provide a conditioned medium directly from the AEC apparatus (e.g. AEC column) to
35 the HIC apparatus (e.g. HIC column).

Room temperature may be about 18-25°C, 20-22°C, about 20°C, about 21°C or about 22°C. The processes disclosed herein may be performed at room temperature. HIC may be performed at room temperature. HIC may be performed at a temperature of 15-25°C. For reproducibility processes may perform HIC at a specific and stable temperature. A stable temperature may be a specific temperature $\pm$ 1.0°C or $\pm$ 1.0°C.

The process of the invention may comprise subjecting the second mixture to a HIC step in flow through mode to provide a HIC flow-through solution in which the fraction of mono-PEGylated protein is increased relative to the second mixture, the HIC step comprising applying the second mixture to a HIC material under conditions suitable for binding oligo-PEGylated protein, wherein the HIC flow-through provides the mono-PEGylated protein composition. In such processes, the HIC material may be Toyopearl Phenyl 650M, the HIC step is performed at a pH of about 7.0 to 9.0 and a conductivity of about 30-40 mS/cm; and the second mixture is applied to the HIC material as a HIC load solution comprising about 25 mM bicine and about 390 mM Na₂SO₄. The protein may be erythropoietin.

Processes disclosed herein are advantageous

form of the protein and a relatively high binding capacity for the non-PEGylated form of the protein. This is advantageous because it facilitates rapid separation of non-PEGylated protein from PEGylated protein. Much of the binding capacity of the column is used for binding to the non-PEGylated form of the protein which facilitates the processing of relatively large volumes of first mixture (the mixture comprising non-PEGylated mono-PEGylated and oligo-PEGylated protein) using a relatively small column. This is because the first mixture is generally relatively low in non-PEGylated protein (which the IEC binding material has a relatively high binding capacity for) and high in PEGylated protein (which the IEC binding material has a relatively low binding capacity for) – this means that in operation the processes disclosed herein can process relatively large volumes of first mixture in the IEC step. The relatively rapid separation of non-PEGylated protein from PEGylated protein facilitates rapid recycling of non-PEGylated protein and overall process efficiency.

The processes of the invention comprise recycling non-PEGylated protein without performing a diafiltration or ultrafiltration step. This provides a faster process enabling higher productivity compared with processes that involve recycling of non-PEGylated protein but which require removal or reduction of salt from non-PEGylated protein fraction.

The processes of the invention also have a relatively high yield compared with prior art process such as that disclosed in WO 2009/010270. Recycling of non-PEGylated (unreacted) protein increases overall yield. As discussed in the Examples below, the processes disclosed herein provide a yield of mono-PEGylated protein that is more than 40% higher than that of the process disclosed in WO 2009/010270. The productivity of the processes disclosed herein is expected to be at least double that of prior art

processes of the type disclosed in WO 2009/010270, and may be many fold higher. The productivity of the processes disclosed herein is expected to be at least double that of prior art processes of the type disclosed in WO 2009/010270, when using equipment of equal size, and may be many fold higher.

5 Compared with the process disclosed in Pfister (Biotech and BioEng 2016; 113) the processes disclosed herein are calculated to be around twice as productive (around 1.7 fold more productive). The increased productivity is mostly the result of avoiding the need for time-consuming diafiltration of non-PEGylated (unreacted) EPO before it can be recycled by addition to a subsequent PEGylation reaction. Because the unreacted EPO is recovered faster in the presently-disclosed processes the overall productivity is greater. The processes disclosed herein also involve a faster PEGylation reaction. The PEGylation processes
10 disclosed herein are calculated to produce 20-33% more mono-PEGylated EPO per reaction than the process disclosed in Pfister.

Compared with the process disclosed in WO 2009/010270, the processes disclosed herein are more productive and have higher yields of mono-PEGylated protein.

Binding capacity

15 The binding capacity of a chromatography material refers to the amount of protein which can bind to the medium under defined experimental conditions. Binding capacity herein may refer to dynamic binding capacity. The dynamic binding capacity refers to the amount of protein which can bind to the medium under defined experimental conditions that include the flow rate of the mobile phase (or buffer solution). The binding capacity may refer to the static binding capacity, which refers defined experimental
20 conditions that do not include the flow rate.

Dynamic binding capacity for a specific protein may be determined using conventional techniques. Dynamic binding capacity for a specific protein may be determined by loading a sample containing a known concentration of that specific protein and measuring protein concentration in the flow-through to solution to establish the amount of protein can protein be bound before a significant amount of protein
25 starts to "break through". This may be achieved by generating a "break through curve" for the chromatography material under defined experimental conditions. The experimental conditions under which the dynamic binding capacity is measured may be the operating conditions. The experimental conditions may be the conditions used in the process used to provide the mono-PEGylated protein composition. In processes employing chromatography materials, loading conditions may thus be
30 adjusted such that a protein of interest is applied in a range less than the "break through" capacity of the chromatography material, to avoid overloading of the chromatography material.

Reference to dynamic binding capacity herein refers to the dynamic binding capacity under the conditions chosen for the chromatography process or step. The dynamic binding capacity of a chromatography medium under particular chromatography conditions for a particular protein can be thought of as the
35 maximum amount of that protein that can be loaded onto the chromatography medium without causing

unnecessary protein loss. That is, the maximum amount of protein of interest that can be loaded without causing "breakthrough". Protein breakthrough may be monitored by spectrophotometry for example at 280 nm (A_{280}). The breakthrough threshold may be 10%. The chromatography conditions may be the pH, conductivity and flow rate, and the concentrations of any salts or other additives to the mobile phase.

5 . The dynamic binding capacity chromatography material are determined under the operating conditions (the conditions chosen for) for the relevant chromatography step. For example the dynamic binding capacity of the AEC material is determined under the operating conditions for the AEC step of the processes of the invention

10 In the present context the binding capacity may refer to the dynamic binding capacity under the chromatography conditions used to prepare the mono-PEGylated protein composition. For example, the binding capacity of AEC material may refer to the dynamic binding capacity of an AEC material in 25 mM bicine, about 7.5 mM Na_2SO_4 , pH 8, at a flow rate of 200 cm/h, residence time 3.3 minutes.

The binding capacity of HIC material may be about 5 g/L PEGylated protein in flow through mode and 2 g/L PEGylated protein in bind and elute mode. The dynamic binding capacity of the HIC material used in

15 flow-through mode for oligo-PEGylated protein may be at least about 2 g/L, 3 g/L, 4 g/L, or 5 g/L, or about 2-8 g/L, 3-7 g/L, 4-6 g/L, or about 5 g/L. The dynamic binding capacity of the HIC material used in bind and elute mode for PEGylated protein may be at least about 0.5 g/L, 1 g/L, 2 g/L, 3 g/L, 4 g/L, or 5 g/L, or about 0.5-4 g/L, 1-3 g/L, or about 5 g/L. The binding capacity may refer to the dynamic binding capacity for PEGylated protein of an HIC material in 25 mM bicine, about 300 mM Na_2SO_4 , pH 8.0, at a

20 flow rate of 150 cm/h, residence time 3.3 minutes, or in 25 mM bicine, about 390 mM Na_2SO_4 , pH 8.0, at a flow rate of 150 cm/h, residence time 3.3 minutes.

In embodiments where the ion exchange chromatography step is anion exchange chromatography, the AEC step is performed under conditions suitable for non-PEGylated protein to bind to the AEC material. The suitability of the conditions can be considered in terms of dynamic binding capacities. The AEC step

25 is performed in flow through mode. In flow through mode for AEC in the processes of the invention PEGylated protein is bound relatively weakly by the AEC material and non-PEGylated protein is bound relatively strongly (tightly). In such conditions the dynamic binding capacity of the AEC material for non-PEGylated protein is relatively high and the dynamic binding capacity of the AEC material for PEGylated protein is relatively low. The chromatography conditions can be selected to maximise dynamic binding capacity for non-PEGylated protein. The chromatography conditions can be selected to minimise

30 dynamic binding capacity for PEGylated protein. The chromatography conditions can be selected to adjust, or optimise, the dynamic binding capacity of the AEC material for PEGylated protein and non-PEGylated protein in order that non-PEGylated protein is retained on the AEC material and PEGylated protein is recovered in the flow-through solution. The AEC material may have a dynamic binding capacity

35 for the PEGylated protein of less than about 2.5 g/L, 1.5 g/L or 1.0 g/L, or about 1.0 – 2.5 g/L. For example, the AEC material may have a dynamic binding capacity for the PEGylated protein of less than about 1.5 g/L.

Thus in the processes of the invention subjecting the first mixture to an AEC step may comprise loading the first mixture onto the AEC material at a load that is about equal to, or at a load that is lower than, the dynamic binding capacity of the AEC material for non-PEGylated protein. It may comprise loading the first mixture onto the AEC material at a load at least 2, 5, 10, 20, 25 or 30 times lower than the dynamic binding capacity of the AEC material for non-PEGylated protein (such that non-PEGylated material is retained on the AEC material). The amount of non-PEGylated protein applied to the anion exchange material may be in the range of about 80-95% of the dynamic binding capacity of the anion exchange material. The first mixture may be loaded onto the AEC material at a load that is higher than, or at least 2, 5, 20, 25 or 30 times higher than the dynamic binding capacity of the AEC material for PEGylated protein (such that PEGylated protein is recovered in the flow-through solution or effluent from the AEC material).

Subsequent elution of non-PEGylated protein from the AEC material is carried out under conditions in which the dynamic binding capacity of the AEC material for the non-PEGylated protein is relatively low. This may be achieved by applying an elution buffer of relatively high conductivity compared with the load and/or wash buffers.

The HIC step may be performed in flow through mode. In flow through mode of an HIC step in a process of the invention the protein of interest (in this case mono-PEGylated protein) is bound relatively weakly by the HIC material and the unwanted protein (or contaminant, in this case oligo-PEGylated protein) is bound relatively strongly (tightly). In such conditions the dynamic binding capacity of the HIC material for oligo-PEGylated protein is higher than the dynamic binding capacity of the HIC material for mono-PEGylated protein. The chromatography conditions can be selected to maximise strength of binding for oligo-PEGylated protein and/or minimise strength of binding for mono-PEGylated protein.

The HIC step may be performed in bind and elute mode. In bind and elute mode in an HIC step in a process of the invention the protein of interest (in this case mono-PEGylated protein) is bound relatively weakly by the HIC material and the unwanted protein (or contaminant, in this case oligo-PEGylated protein) is bound relatively strongly (tightly) by the HIC material. The mono-PEGylated protein can be eluted from the HIC material before the oligo-PEGylated protein, using a decreasing salt gradient.

In embodiments where the ion exchange chromatography step is cation exchange chromatography, the CEC step is performed under conditions suitable for non-PEGylated protein to bind to the CEC material. The suitability of the conditions can be considered in terms of dynamic binding capacities. The CEC step is performed in flow through mode. In flow through mode for CEC in the processes of the invention PEGylated protein is bound relatively weakly by the CEC material and non-PEGylated protein is bound relatively strongly (tightly). In such conditions the dynamic binding capacity of the CEC material for non-PEGylated protein is relatively high and the dynamic binding capacity of the CEC material for PEGylated protein is relatively low. The chromatography conditions can be selected to maximise dynamic binding capacity for non-PEGylated protein and/or minimise dynamic binding capacity for PEGylated (or mono-PEGylated) protein. The chromatography conditions can be selected to adjust, or optimise, the dynamic

binding capacity of the CEC material for PEGylated protein and non-PEGylated protein in order that non-PEGylated protein is retained on the CEC material and PEGylated protein is recovered in the flow-through solution. The CEC material may have a dynamic binding capacity for the PEGylated protein of less than about 2.5g/L, 1.5 g/L or 1.0 g/L, or about 1.0 – 2.5 g/L. For example, the CEC material may
5 have a dynamic binding capacity for the PEGylated protein of less than about 1.5 g/L.

Thus in the processes of the invention subjecting the first mixture to an CEC step may comprise loading the first mixture onto the CEC material at a load that is about equal to, or at a load that is lower than, the dynamic binding capacity of the CEC material for non-PEGylated protein. It may comprise loading the first mixture onto the CEC material at a load at least 2, 5, 10, 20, 25 or 30 times lower than the dynamic
10 binding capacity of the CEC material for non-PEGylated protein (such that non-PEGylated material is retained on the CEC material). The amount of non-PEGylated protein applied to the anion exchange material may be in the range of about 80-95% of the dynamic binding capacity of the anion exchange material. The first mixture may be loaded onto the CEC material at a load that is higher than, or at least 2, 5, 20, 25 or 30 times higher than the dynamic binding capacity of the CEC material for PEGylated
15 protein (such that PEGylated protein is recovered in the flow-through solution or effluent from the CEC material).

Subsequent elution of non-PEGylated protein from the CEC material is carried out under conditions in which the dynamic binding capacity of the CEC material for the non-PEGylated protein is relatively low. This may be achieved by applying an elution buffer of relatively high conductivity compared with the load
20 and/or wash buffers.

Chromatography conditions

The amount of protein loaded on to a chromatography material may depend on the binding capacity of the material. For example if an AEC material has a binding capacity for non-PEGylated EPO of about 35
25 g/L, and about 50% of the EPO in the mixture of reaction products is non-PEGylated then the maximum load of the mixture of reaction products (in terms of total protein) applied to the AEC material would be about 70 g/L.

Flow rates for chromatography steps may be selected and adjusted according to conventional
30 techniques. Faster flow rates will decrease binding capacity, which means that a balance may be reached between achieving maximum dynamic binding capacity and a fast separation, particularly when applying large volumes of protein mixtures to be separated. Suitable flow rates may be 50 – 400 cm/h. Residence times may be 3-5 minutes, or possibly less, depending on the chromatography material used. Faster flow rates and shorter residence times may be desirable for improving overall process productivity.

Chromatography steps may be performed under conditions "suitable for binding" a particular protein (or PEGylation form of a protein). The skilled person is familiar with chromatography techniques and is able to find conditions suitable for binding a particular protein empirically, using his common general knowledge and guided by the present disclosure. Parameters such as chromatography material, type
5 and/or concentration of salt, pH, buffers, temperature and flow rate can all be altered to provide conditions suitable for binding a particular protein (or PEGylation form of a protein) in chromatography.

As noted above, in AEC protein mixtures (load compositions) and wash buffers of a relatively high pH are generally used, in order that the protein of interest (or contaminant, e.g. unwanted protein) has a net negative charge and therefore binds to the positively charged anion exchange material. Conversely in
10 CEC protein mixtures and wash buffers of a relatively low pH are generally used, in order that the protein of interest (or contaminant, e.g. unwanted protein) has a net positive charge and therefore binds to the negatively charged cation exchange material.

The processes of the present invention may be used for producing a mono-PEGylated protein composition wherein the protein has a pI of about 8.0 or lower, or 7.0 or lower, or 6.0 or lower. The
15 processes of the invention are particularly suitable for proteins having a pI of 6.0 or lower. In this case, the ion exchange chromatography step may be an AEC step and the AEC conditions are at a pH greater than the pI of the protein, preferably at least 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, or 4.0 pH units greater than the pI of the protein. In this context the pI may refer to the pI of the most basic isoform of the protein, or the pI of the most abundant isoform of the protein, in the protein composition.

20 The processes of the present invention may be used for producing a mono-PEGylated protein composition wherein the protein has a pI of about 8.0 or higher. In this case, the ion exchange chromatography step may be a CEC step and the CEC conditions are at a pH lower than the pI of the protein, preferably at least 0.5, 1.0, 2.0 or 3.0 pH units lower than the pI of the protein. In this context the pI may refer to the pI of the most acidic isoform of the protein, or the pI of the most abundant isoform of
25 the protein, in the protein composition. The protein may have a pI of about 2.0 - 6.0, 2.5 - 5.5, 3.0 - 5.5, 3.5 - 5.0, or about 3.5-4.5.

The processes of the present invention may be used for producing a mono-PEGylated protein composition wherein the protein has a pI of about 6.0 - 8.0. In this case, the ion exchange chromatography step may be an AEC or a CEC step the pH conditions selected to be pH lower than the
30 pI of the protein for AEC, and higher than the pI of the protein for CEC. Preferably the pH of the ion exchange step is at least 0.5, 1.0, 2.0 or 3.0 pH units away from (different than) the pI of the protein. In this context the pI may refer to the pI of the most acidic or basic isoform of the protein, or the pI of the most abundant isoform of the protein, in the protein composition. Ion exchange chromatography is an established technique and therefore the skilled person would have no difficulty in selecting pH conditions
35 suitable for practicing the processes of the present invention (i.e. selecting pH conditions that favour

binding of the non-PEGylated protein to the ion exchange material and allow PEGylated protein to pass through/over the material to be recovered in the flow-through solution).

The pH of the PEGylation reaction may be the substantially same as the pH of the ion exchange step (the AEC step or CEC step). The pH of the PEGylation reaction may be selected to be substantially the same as the pH of the ion exchange step, such that the protein mixture resulting from the PEGylation reaction is loaded directly onto the ion exchange material. In this context substantially the same means within 1.0, 0.9, 0.8, 0.6, 0.7, 0.5, 0.4, 0.3, 0.2, or 0.1 pH units. In such processes the first mixture is the mixture of reaction products that results from the PEGylation reaction. Such processes are relatively efficient and fast. In this context "loaded directly" means that no pH adjustment of the mixture of reaction products is carried out before it is subjected to the ion exchange chromatography step. The buffer for the PEGylation reaction may be the same as the buffer for the ion exchange step.

A PEGylation reaction may be performed at a pH of about 6.5 to 9.5. The pH of the PEGylation reaction may be selected to be at least 0.5, 1.0, 2.0 or 3.0 pH units away from the pI of the protein, such that the mixture of reaction products can be loaded directly on to the ion exchange material at a pH that favours binding of the protein. In this context, a protein having a relatively neutral pH (e.g. pI about 7.5) could be PEGylated in a reaction at about pH 9.5 and the mixture of reaction products subjected to an AEC step. Alternatively, a protein having a pI of about 7.5 could be PEGylated in a reaction at about pH 6.5 and the mixture of reaction products subjected to a CEC step. The pH of the PEGylation reaction can be selected such that it is at least 0.5, 1.0, 2.0 or 3.0 pH units away from the pI of the protein, and the mode of ion exchange chromatography (AEC or CEC) selected accordingly. In this way the mixture of reaction products can be subjected to the ion exchange chromatography step.

In the present context, chromatography steps (such as AEC, CEC or HIC) may remove a "contaminant" from a mixture (such as a mixture of reaction products, also referred to herein as a "first mixture"; or PEGylated protein mixture, also referred to herein as a "second mixture"), wherein the contaminant is an undesired form of protein. A contaminant may also be referred to as an impurity. For example, since the desired form of protein in the present context is mono-PEGylated protein, the chromatography step may remove non-PEGylated protein or oligo-PEGylated protein. The chromatography step may also remove other contaminants, such as protein aggregates, or PEGylation reactants.

The present disclosure provides the use of an AEC medium, or a CEC medium, for removing non-PEGylated protein from a mixture comprising non-PEGylated protein and PEGylated protein. The protein may be EPO. The use of the AEC medium may be carried out in the processes for preparing a mono-PEGylated protein, as disclosed herein.

Chromatography conditions and chromatography materials for carrying out the present invention may be selected using a screening process. The screening process may allow screening for selectivity of binding. Selectivity of binding is advantageous. For example ion exchange materials and conditions can

be screened for selectivity which favours binding of non-PEGylated protein to the AEC material and non-binding of PEGylated protein to the ion exchange material. For example HIC materials and flow-through conditions can be screened for selectivity which favours binding of oligo-PEGylated protein and non-binding of mono-PEGylated protein. A screening process for an ion exchange material, such as an AEC material or a CEC material, may involve applying a mixture of non-PEGylated, mono-PEGylated and oligo-PEGylated proteins to the ion exchange material at relatively low conductivity, then applying an increasing conductivity (e.g. salt) gradient to the material and monitoring the appearance of the non-PEGylated, mono-PEGylated and oligo-PEGylated in the eluate. This process could be conducted at several different pH values. A screening process for a HIC material may involve applying a mixture of non-PEGylated, mono-PEGylated and oligo-PEGylated proteins to the HIC material under relatively high salt conditions, then applying a decreasing salt gradient to the material and monitoring the appearance of the non-PEGylated, mono-PEGylated and oligo-PEGylated in the eluate. Chromatography materials and conditions may be assessed for selectivity of binding by analysing the eluate by UV chromatography and screening for materials and conditions that provide good separation of peaks in the chromatograms.

15

PEG

Poly(ethylene glycol) or PEG is a neutral hydrophilic polyether. The term "molecular weight" (in kDa) in the present context is to be understood as the mean molecular weight of the PEG because PEG as polymeric compound is not obtained with a defined molecular weight but in fact has a molecular weight distribution; the term "about" indicates that some PEG molecules, or residues, will weigh more and some less than the indicated molecular weight, i.e. the term "about" in this context may refer to a molecular weight distribution in which 95 % of the PEG molecules have a molecular weight within +/- 10 % of the indicated molecular weight. For example, a molecular weight of 30 kDa may denote a range of from 27 kDa to 33 kDa.

25 A PEG residue can contain further chemical groups which are necessary for binding reactions, which results from the chemical synthesis of the PEGylated molecule, or which is a spacer for optimal distance of parts of the molecule. These further chemical groups are not used for the calculation of the molecular weight of the PEG residue. In addition, such a PEG residue can consist of one or more PEG chains which are covalently linked together. PEG residues with more than one PEG chain are called multiarmed or
30 branched PEG residues. Branched PEG residues can be prepared, for example, by the addition of polyethylene oxide to various polyols, including glycerol, pentaerythriol, and sorbitol. Branched PEG residues are reported in, for example, EP 0 473 084, US 5,932,462.

A PEG molecule used in a PEGylation reaction, and a PEG residue on a PEGylated protein, may each have a molecular weight of at least about 12 kDa, or at least about 20kDa, about 20 kDa to 40 kDa, or
35 about 30 kDa. A PEG residue may have a molecular weight of 20 kDa to 35 kDa and be a linear PEG

residue. A PEG residue may have a molecular weight of 35 kDa to 40 kDa and be a branched PEG residue.

A mono-PEGylated EPO may comprise a single PEG residue having a molecular weight of at least about 12 kDa, or at least about 20 kDa, about 20 kDa to 40 kDa, about 20 kDa, or about 30 kDa. A PEG
5 residue may have a molecular weight of at least about 20 kDa.

A mono-PEGylated protein is a protein comprising a single PEG residue. That is, the mono-PEGylated protein has one PEG residue only. An oligo-PEGylated protein comprises at least two PEG residues. For example an oligo-PEGylated protein may be a di-, tri or tetra-PEGylated protein. The term oligo-PEGylated protein (or poly-PEGylated protein) may refer to a group of oligo-PEGylated protein molecules
10 having varying degrees of PEGylation (two, three, or more PEG residues). The term PEGylated protein refers to mono-PEGylated and oligo-PEGylated proteins. The term PEGylated protein may refer to protein consisting of mono-PEGylated and oligo-PEGylated protein. A non-PEGylated protein is a protein that does not comprise any PEG residues. A non-PEGylated protein may also be referred to in the context of a PEGylation reaction as an unreacted protein. A non-PEGylated protein may be referred to as
15 a native protein, or an un-PEGylated protein or a free protein.

The term "PEGylation" means a covalent linkage of a PEG residue with a protein. In particular it may refer to a covalent linkage at the N-terminus of the polypeptide and/or an internal lysine residue. PEGylation of proteins is widely known in the state of the art and reviewed by, for example, Veronese, F.M., *Biomaterials* 22 (2001) 405-417. PEG can be linked using different functional groups and
20 polyethylene glycols with different molecular weight, linear and branched PEGs as well as different linking groups (see also Francis, G.E., et al., *Int. J. Hematol.* 68 (1998) 1-18; Delgado, C., et al., *Crit. Rev. Ther. Drug Carrier* 30 Systems 9 (1992) 249-304). PEGylation of erythropoietin can be performed in aqueous solution with PEGylation reagents as described, for example, in WO 00/44785, in one embodiment by using NHS-activated linear or branched PEG molecules of a molecular weight between 5kDa and 40 kDa.
25 PEGylation can also be performed at the solid phase according to Lu, Y., et al., *Reactive Polymers* 22 (1994) 221-229. Not randomly, N-terminally PEGylated polypeptide can also be produced according to WO 94/01451. PEGylation reactions are also reviewed in WO 2009/010270 and WO 2012/035037.

Suitable PEG derivatives are activated PEG molecules with an average molecular weight of from about 5 to about 40 kDa, or from about 20 to about 40 kDa. The PEG derivative is in one embodiment a linear or
30 a branched PEG. A wide variety of PEG derivatives suitable for use in the preparation of PEG-protein and PEG-peptide conjugates can be obtained from Shearwater Polymers (Huntsville, AL, U.S.A.; www.nektar.com). Activated PEG derivatives are known in the art and are described in, for example, Morpurgo, M., et al., *J. Bioconjug. Chem.* 7 (1996) 363-368.

A PEGylation reaction may be performed at a pH of about 6.5 to 9.5, about 7.0 to 9.0, or about 7.5 to 8.5,
35 or about 8.0. The pH at which the PEGylation reaction is performed may depend on the PEG reagent

used. The PEG reagent may be mPEG-NHS, mPEG-SPA, mPEG-SVA or mPEG-Cl. The PEGylation reaction may be performed at a pH of about 7.0 to 9.0, or about 7.5 to 8.5, or about 8.0 using (NHS) activated PEG reagent. The PEGylation reaction may be performed at about 15-25°C, or about 18-22°C, or about 20°C. The PEGylation reaction may be carried out for at least about 20, 30, 40, 50 or 60 minutes, or about 30-90, or 30-60 minutes, or about 40, 50, or 60 minutes.

A PEGylation reaction may be performed in a solution comprising a salt and a buffer. The salt may be Na₂SO₄ and the buffer may be bicine. The salt may be present in an amount of 5-10 mM or about 7.5 mM. The buffer may be present in an amount of about 10-50 mM, 20-30 mM or about 25 mM. The PEGylation reaction may be performed in 7.5 mM Na₂SO₄ and 25 mM bicine.

10 EPO

The term "erythropoietin" and its abbreviation "EPO" refer to a protein having the amino acid sequence of SEQ ID NO: 1 or of SEQ ID NO: 2, or a protein or polypeptide substantially homologous thereto, whose biological properties relate to the stimulation of red blood cell production and the stimulation of the division and differentiation of committed erythroid progenitors in the bone marrow. Recombinant erythropoietin may be prepared via expression in eukaryotic cells, for example in CHO cells, or BHK cells, or HeLa cells by recombinant DNA technology or by endogenous gene activation, i.e. the erythropoietin glycoprotein is expressed by endogenous gene activation, see for example US 5,733,761, US 5,641,670, US 5,733,746, WO 93/09222, WO 94/12650, WO 95/31560, WO 90/11354, WO 91/06667, and WO 91/09955. The EPO may be human EPO. The EPO may be glycosylated EPO.

20 The human EPO may have the amino acid sequence set out in SEQ ID NO: 1 or SEQ ID NO: 2. The human EPO may have the amino acid sequence set out in SEQ ID NO: 1. The term "EPO" also denotes variants of the protein of SEQ ID NO: 1 or of SEQ ID NO: 2, in which one or more amino acid residues have been changed, deleted, or inserted, and which has comparable biological activity as the not modified protein, such as e.g. reported in EP 1 064 951 or US 6,583,272. The number of amino acids changed, deleted or inserted may be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 1-50, 1-40, 1-30, 1-20, or 1-10. The term EPO denotes proteins that comprise or consist of the amino acid sequence set out in SEQ ID NO:1 or SEQ ID NO:2 or variants thereof. A variant may have the amino acid sequence of human erythropoietin having from 1 to 6 additional sites for glycosylation. The specific activity of PEGylated erythropoietin can be determined by various assays known in the art. The biological activity of the purified PEGylated erythropoietin are such that administration of the protein by injection to human patients results in bone marrow cells increasing production of reticulocytes and red blood cells compared to noninjected or control groups of subjects. The biological activity of PEGylated erythropoietin obtained and purified in accordance with the method as reported herein can be tested by methods according to Bristow, A, *Pharmeuropa Spec. Issue Biologicals BRP Erythropoietin Bio 97-2* (1997) 31-48.

35 Amino acid sequence variants of EPO can be prepared by introducing appropriate modifications into the nucleotide sequence encoding the EPO, or by peptide synthesis. Such modifications include, for

example, deletions from, and/or insertions into, and/or substitutions of residues within the amino acid sequences of the erythropoietin. Any combination of deletion, insertion, and substitution can be made to arrive at the final construct, provided that the final construct possesses comparable biological activity to the human EPO.

- 5 Conservative amino acid substitutions are shown in the table below under the heading of "preferred substitutions". More substantial changes are provided under the heading of "exemplary substitutions", and as described below in reference to amino acid side chain classes. Amino acid substitutions may be introduced into human erythropoietin and the products screened for retention of the biological activity of human erythropoietin.
- 10 Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

The EPO may be a variant EPO. The EPO may be comprised in a fusion protein with another protein, or may be conjugated to another moiety in addition to PEG.

The chemical PEGylation of erythropoietin generally results in a protein preparation comprising erythropoietin which is PEGylated at one or more ϵ -amino groups of lysine residues and/or at the

15 N-terminal amino group. Selective PEGylation at the N-terminal amino acid can be performed according to Felix (1997). Selective N-terminal PEGylation can be achieved during solid-phase synthesis by coupling of a N $^{\alpha}$ -PEGylated amino acid derivative to the N-1 terminal amino acid of the peptide chain. Side chain PEGylation can be performed during solid-phase synthesis by coupling of N $^{\epsilon}$ -PEGylated lysine derivatives to the growing chain. Combined N-terminal and side chain PEGylation is feasible either

20 as described above within solid-phase synthesis or by solution phase synthesis by applying activated PEG reagents to an amino deprotected peptide.

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp; Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu

Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu;Phe;Ile	Leu
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Val; Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

Amino acids may be grouped according to common side-chain properties:

- (1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;
- (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;
- 5 (3) acidic: Asp, Glu;
- (4) basic: His, Lys, Arg;
- (5) residues that influence chain orientation: Gly, Pro;
- (6) aromatic: Trp, Tyr, Phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

10 Proteins

The processes of the invention are especially suitable for producing mono-PEGylated EPO compositions. Reference to a protein or protein of interest herein may refer to EPO.

- The processes of the invention may be suitable for producing mono-PEGylated compositions of other proteins, particularly therapeutic proteins. For example interleukin-2 (IL-2), peginterferon alfa-2a, human
- 15 growth hormone (hGH), bone morphogenetic protein 2 (BMP-2), bone morphogenetic protein 7 (BMP-7), bone morphogenetic protein 15 (BMP-15), neurotrophin-3 (NT-3), von Willebrand factor (vWF) protease, granulocyte colony stimulating factor (G-CSF), granulocyte-macrophage colonystimulating factor (GM-CSF), interferon alpha, interferon beta, interferon gamma, tissue-type plasminogen activator (IPA), leptin, hirudin, urokinase, human DNase, insulin, hepatitis B surface protein (HbsAg), chimeric diphtheria toxin-
 - 20 IL-2, human chorionic gonadotropin (hCG), thyroid peroxidase (TPO), alpha-galactosidase, alpha-L-iduronidase, beta-glucosidase, alpha-galactosidase A, acid a-glucosidase (acid maltase), anti-thrombin III

(AT III), follicle stimulating hormone (FSH), glucagon-like peptide-I (GLP-1), glucagonlike peptide-2 (GLP-2), fibroblast growth factor 7 (FGF-7), fibroblast growthfactor21 (FGF-21), fibroblast growth factor 23 (FGF-23), Factor X, Factor XIII, prokinetisin, extendin-4, CD4, tumor necrosis factor receptor (TNF-R), a-CD₂₀, P-selectin glycoprotein ligand-I (PSGL-I), complement, transferrin, glycosylation-dependent cell
5 adhesion molecule (GlyCAM), neural-cell adhesion molecule (N-CAM), INF receptor-IgG Fe region fusion protein. Such proteins also include antibodies such as monoclonal antibodies against any one of: respiratory syncytial virus, protein F of respiratory syncytial virus, INF- α , glycoprotein IIb/IIIa, CD20, VEGF-A, PSGL-1, CD4, a-CD3, EGF, carcinoembryonic antigen (CEA), TNF α and IL-2 receptor.

The processes of the invention may be suitable for producing mono-PEGylated compositions of proteins
10 such as hormones, cytokines or enzymes. Such a protein may be erythropoietin. The protein may be interferon- α -2a or interferon- α -2b. The protein may be granulocyte colony-stimulating factor, human growth hormone, or urate oxidase.

The invention is particularly useful for therapeutic proteins having a relatively short half-life, because PEGylation increases *in vivo* circulation half-life. Generally hormones and cytokines have a relatively
15 short half-life. Smaller biological molecules tend to have a relatively short half-life. The pharmacokinetic profile of relatively small biological molecules may be improved by PEGylation and so the present invention may also be particularly useful for relatively small therapeutic proteins. Generally proteins and peptides smaller than approximately 70 kDa are more likely to be eliminated by kidney filtration than are larger proteins. Smaller biological molecules, or proteins, may be defined as those having a molecular
20 weight of less than about 70 kDa. Erythropoietin has a molecular weight of about 37 kDa. The invention is useful for therapeutic proteins having a molecular weight less than about 70 kDa (in their non-PEGylated form). The invention is useful for therapeutic proteins having a molecular weight less than about 70 kDa, 60 kDa, 50 kDa, or 40 kDa, or having a molecular weight of about 10-70 kDa, 20-60 kDa, 20-50 kDa, or 30-40 kDa. The invention is useful for hormones or cytokines having a molecular weight
25 less than about 70 kDa, 60 kDa, 50 kDa, or 40 kDa, or having a molecular weight of about 10-70 kDa, 20-60 kDa, 20-50 kDa, or 30-40 kDa.

The protein may be an antibody. An antibody may be a polyclonal antibody or a monoclonal antibody. An antibody may be a biologically functional antibody fragment. Antibody fragments include Fab, Fab', F(ab')₂, scFv, (scFv)₂, single-domain antibodies (sdAb, or dAB), complementarity determining region
30 fragments, linear antibodies, single-chain antibody molecules, minibodies, diabodies and multispecific antibodies formed from antibody fragments. Many antibody fragments have a relatively short half-life *in vivo*, which results from their relatively small size and their no longer having an Fc region. The invention is particularly useful for antibody fragments having a relatively small size, such as single-domain antibodies, Fabs, Fab's, and scFvs. The invention is useful for antibody fragments having a molecular
35 weight less than about 70 kDa, 60 kDa, 50 kDa, or 40 kDa, or having a molecular weight of about 10-70 kDa, 20-60 kDa, 20-50 kDa, or 30-40 kDa.

- The mono-PEGylated protein compositions may be formulated as pharmaceutical compositions. The mono-PEGylated protein compositions produced by the processes disclosed herein may be formulated with one or more pharmaceutically acceptable excipients, and/or may be formulated in a physiologically acceptable buffer such as physiological saline. The salts and/or buffers used in the equilibration, wash, or elution buffers of the HIC step may be removed before formulating the mono-PEGylated protein composition as a pharmaceutical composition. For example the mono-PEGylated protein compositions may be dialysed. The mono-PEGylated protein of the mono-PEGylated protein composition may be lyophilised. The mono-PEGylated protein of the mono-PEGylated protein composition may be isolated and reformulated in a pharmaceutical composition.
- The processes disclosed herein may be industrial scale processes. An industrial scale process may be a process that produces at least about 5g, 10g, 25g, 50g, 100g, 250g or 500g per batch or per cycle. A batch or cycle in this context may be a process comprising all of the steps a) to c) as disclosed herein. An industrial scale process may be a process in which the volume of the first mixture (comprising non-PEGylated, mono-PEGylated and oligo-PEGylated protein) that is subjected to the AEC step is at least 100L, 500L, 1000L, 5000L, 10000L, 50000L, or 100000L.

The invention includes the combination of the aspects and preferred features described except where such a combination is clearly impermissible or expressly avoided.

- The features disclosed in the foregoing description, or in the following claims, or in the accompanying drawings, expressed in their specific forms or in terms of a means for performing the disclosed function, or a method or process for obtaining the disclosed results, as appropriate, may, separately, or in any combination of such features, be utilised for realising the invention in diverse forms thereof.

- While the invention has been described in conjunction with the exemplary embodiments described above, many equivalent modifications and variations will be apparent to those skilled in the art when given this disclosure. Accordingly, the exemplary embodiments of the invention set forth above are considered to be illustrative and not limiting. Various changes to the described embodiments may be made without departing from the spirit and scope of the invention.

For the avoidance of any doubt, any theoretical explanations provided herein are provided for the purposes of improving the understanding of a reader. The inventors do not wish to be bound by any of these theoretical explanations.

- The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

Aspects and embodiments of the present invention will now be illustrated, by way of example, with reference to the accompanying figures. Further aspects and embodiments will be apparent to those skilled in the art. All documents mentioned in this text are incorporated herein by reference.

Throughout this specification, including the claims which follow, unless the context requires otherwise, the word “comprise” and “include”, and variations such as “comprises”, “comprising”, and “including” will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

- 5 It must be noted that, as used in the specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. Ranges may be expressed herein as from “about” one particular value, and/or to “about” another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by the use of the antecedent “about,” it
- 10 will be understood that the particular value forms another embodiment. The term “about” in relation to a numerical value is optional and means for example +/- 10%.

All references mentioned above are hereby incorporated by reference.

Examples

- The following examples, including the experiments conducted and the results achieved, are provided for illustrative purposes only and should not be construed to include all possible embodiments along with the full scope of equivalents to which such claims are entitled.
- 15

EXAMPLE 1

- EPO was PEGylated using NHS-activated PEG reagent (molecular weight of about 30 kDa) at pH 8.0 in 25 mM bicine and 7.5 mM Na₂SO₄ to provide a mixture of reaction products (“first mixture”) comprising non-PEGylated EPO, mono-PEGylated EPO and oligo-PEGylated EPO.
- 20

- The first mixture was subjected to AEC at pH 8.0, 25 mM bicine, 7.5 mM Na₂SO₄. The AEC resin was Toyopearl SuperQ 650M. This has a dynamic binding capacity for EPO of about 34 g/L. Elution was carried out by a step elution to 25 mM bicine, 35 mM sodium sulphate at pH 8. Non-PEGylated EPO recovered by elution was recycled into a subsequent PEGylation reaction, the product of which (a subsequent “first mixture”) was subjected to a second AEC step. This process was repeated again, such that the process included three cycles of PEGylation and AEC steps.
- 25

- A chromatograph showing protein elution in an AEC step is shown in Figure 4. Three chromatographs showing the three AEC steps are shown in Figure 5, which shows that the amount of EPO in the second cycle is less than in the first, and the amount of EPO in the third cycle is less than in the second.
- 30

Table 1 shows the contents of the various intermediate compositions. The PEGylation product is the “first mixture”, the product pool is the AEC flow-through solution, and the epo pool is the AEC eluate.

Table 1

	Conc [g/l]	Vol [ml]	Mass [mg]	Composition [%]			Mass [mg]		
				Epo	Mono	Oligo	Epo	Mono	Oligo
EPO start	8.71	5.74	50.0	100.0	0	0	50.0	0	0
PEGylation 1									
PEGylation product	5	9.5	47.5	51.12	40.1	8.79	24.28	19.05	4.18
Chromatography 1									
Product pool	1.97	14	27.58	0.94	80.01	19.06	0.26	22.07	5.26
EPO pool	8.9	2	17.8	100	0	0	17.8	0	0
PEGylation 2									
PEGylation product	2.48	7.0	17.37	50.46	40.19	9.35	8.77	6.98	1.62
Chromatography 2									
Product pool	0.68	14	9.52	0.57	78.89	19.55	0.05	7.61	1.86
EPO pool	3.21	2	6.42	100	0	0	6.42	0	0
PEGylation 3									
PEGylation product	0.80	6.7	5.3	57.62	37.03	5.35	3.05	1.96	0.28
Chromatography 3									
Product pool	0.19	14	2.62	0	86.18	13.82	0	2.26	0.36
EPO pool	1.34	2	2.68	96.98	2.6	0	2.6	0.07	0
Pooled Product (Chromatography 1-3)	0.95	42.0	39.8	0.9	79.7	19.5	0.3	31.7	7.7
Yield (EPO conversion to mono-PEG product) % after 3 cycles				63.4					

The ability of HIC to separate mono-PEGylated EPO from oligo-PEGylated EPO was then investigated in both bind and elute mode and flow-through mode.

In bind and elute mode the HIC resin was Toyopearl Phenyl-650M. The resin was equilibrated with 25 mM bicine and 500 mM Na₂SO₄ at pH 8.0. The mixture applied to the resin comprised non-PEGylated EPO, mono-PEGylated EPO and oligo-PEGylated EPO for testing purposes. The mixture was conditioned with 25 mM bicine and 500 mM Na₂SO₄ at pH 8.0 and applied to the HIC resin. An elution gradient of decreasing salt concentration was applied, from 500 mM to 0 mM Na₂SO₄. Figure 6 is a chromatogram showing that EPO is in the flow-through, and that mono-PEGylated EPO is eluted at relatively high salt whereas oligo-PEGylated EPO is eluted at relatively low salt.

In flow-through mode the HIC resin was Toyopearl Phenyl-650M. The resin was equilibrated with 25 mM bicine and 390 mM Na₂SO₄ at pH 8.0. The mixture applied to the resin comprised mono-PEGylated EPO and oligo-PEGylated EPO, it did not contain significant amounts of non-PEGylated EPO and is therefore representative of a "second mixture" in accordance with the invention. The mixture was conditioned with bicine and Na₂SO₄ at pH 8.0 and applied to the HIC resin. The salt concentration was adjusted to 390 mM. Figure 7 is a chromatogram showing that mono-PEGylated EPO is in the flow-through. Oligo-PEGylated EPO was eluted using a decreasing salt gradient.

EXAMPLE 2

PEGylation of EPO

EPO stock solution was thawed, concentrated to 6.5 g/l using 3 kDa Centricon centrifugal filters, and buffered in 25 mM bicine, pH 8. 7.7 ml of this solution was then transferred into a 50 ml Falcon tube and mixed with 0.3 ml of 25 mM bicine, pH 8 buffer. 96 mg of PEG reagent was weighed and dissolved in 3 ml of 1 mM HCl to prepare the PEGylation solution.

The PEGylation reaction was started by addition of 2 ml of PEGylation solution (molecular weight of about 30 kDa) to the EPO solution. The reaction was carried out in a water bath at 20° C for 50 minutes.

Cyclic PEGylation and purification using ion chromatography

As a first step, PEGylated EPO was produced as outlined above. 1 ml of the PEGylated EPO solution was transferred to an Eppendorf cup, and the remaining 9 ml was injected into a 150 ml Superloop using a 50 ml syringe and anion exchange chromatography started.

A 16 ml Toyopearl SuperQ-650M with a diameter of 1 cm was used as the anion exchange chromatography column. The column was equilibrated with a 25 mM bicine, 7.5 mM Na₂SO₄, pH 8 buffer. Elution was carried out by a step elution to 25 mM bicine, 35 mM sodium sulphate at pH 8. The corresponding chromatogram is shown in Figure 8A.

1 ml of fraction 1B3 was removed and its composition measured by HPLC. Subsequently, fractions 1B3 to 1C1 were combined, and the composition of a 1 ml sample of this mixed fraction also measured by RP HPLC. The protein concentration was determined photometrically, and at an extinction coefficient of 1.25 was 0.097 g/l. This mixed fraction (AEC eluate) is from the elution step.

5

The remaining 21 ml of the mixed fraction was transferred into a 50 ml Falcon tube and placed in a 20 °C water bath for renewed PEGylation. 1 ml of a 26.89 g/l PEGylation reagent solution was added and the second PEGylation reaction started.

10 The AEC eluate (EPO-fraction) is at the same conditions (35 mM sodium sulphate) as the elution buffer of the first chromatography, outlined above. The reaction solution was diluted by a factor of 4.66 with 25 mM bicine buffer after 50 min and thus adjusted to the equilibration conditions. The resulting 97.9 ml sample was again injected into the 150 ml Superloop and the second chromatography was started. The chromatography method was changed only in terms of the larger sample volume. All other parameters and ingredients remained unchanged. The corresponding chromatogram is shown in Figure 8B.

15 Fractions 2A3 to 2A5 were pooled again and the procedure above repeated as a third cycle.

Fractions 1A1 to 1C5 of the third chromatography were combined and concentrated using 3 kDa Centricon centrifugal filters. The concentrate and the pools of the fractions 2A1 to 2A3 were measured by RP HPLC. The corresponding chromatogram is shown in Figure 8C.

20 All pipetted volumes of the PEGylation reactions and the dilution steps are listed in Table 2.

Table 2 - Volumes of starting ingredients and samples

Cycle	Vsample [ml]	Cprotein [g/l]	Vdiluted [ml]	VPEG reagent [ml]	CPEG reagent [g/l]	stoichiometry [MPEG/MEPO]
1	10	5.00	-	2	33.00	0.8
2	21	0.97	98.7	1	26.89	0.8
3	30	0.47	141.0	1	33.50	1.44

Results and discussion

25 The measured results of the RP HPLC analysis are shown in Table 3. The percentage by mass of EPO in the cyclically PEGylated EPO pools is on average about 95%.

Table 3 - Composition before and after PEGylation reaction and chromatographic separation

Sample description	Fraction	Epo [%]	Mono-PEG Epo [%]	Oligo Forms [%]
Starting material	-	100	0	0
PEGylation product 1	-	49.06	41.38	9.55
PEGylation product 2	-	62.36	32.88	4.77
PEGylation product 3	-	51.95	39.82	8.24
Cycle 1 Pool	1A1 – 1C1	96.59	2.05	1.3
Cycle 1 B3	1B3	35.22	64.23	0.55
Cycle 2 Pool	2A3 – 2A5	94.68	3.05	2.27
Cycle 3 Pool	2A1 – 2A3	94.24	3.35	2.18

The mass balance for each cycle is listed in Table 4, as well as for the entire process. It should be noted that the information on the product pools is not based on measured values, but has been calculated.

Table 4 - Mass balances

Description and process step	Composition [%]				Composition [mg]		
	Mass [mg]	Epo	Mono-PEG Epo	Oligo	Epo	Mono-PEG Epo	Oligo
PEGylation 1	50	49.06	41.38	9.55	24.53	20.69	4.78
Chromatography 1							
Product Pool 1	24.91	0.7	80.07	19.23	0.17	19.94	4.79
1B3	0.78	35.22	64.23	0.55	0.28	0.50	0.00
Epo Pool 1	21.37	96.59	2.05	1.35	20.64	0.44	0.29
PEGylation 2	21.37	62.36	32.88	4.77	13.33	7.03	1.02
Chromatography 2							
Product pool 2	7.72	5.21	85.60	9.19	0.40	6.61	0.71
Epo Pool 2	13.65	94.68	3.05	2.27	12.92	0.42	0.31

PEGylation 3	13.65	51.95	39.82	8.24	7.09	5.44	1.12
Chromatography 3							
Product pool 3	6.10	0.00	84.54	15.46	0.00	5.16	0.94
Epo Pool 3	8.35	94.24	3.35	2.18	7.87	0.28	0.18
Balance (1B3 + Sum Product Pools + EPO Pool 3)	47.83	23.99	63.62	12.38	8.72	32.49	6.62
Sum of Product pools	38.7	1.4	82.0	16.6	0.57	31.71	6.44

There was a 95.5% recovery over all samples at the end of the process. A loss of 4.5% of total protein was observed due to sampling and analytics. EPO Pool 1 and EPO Pool 2 were consumed in PEGylation 2 and 3. 63.6% of initial EPO was converted to mono-PEG-EPO. 82.0% mono-PEG-EPO content in intermediate pool.

- On the basis of the mass balance it can be seen that 32.5 mg of mono-PEGylated EPO could be prepared from the 50 mg unPEGylated EPO used, which exists as a mixture of 1.4% Epo, 82.0% mono-PEG Epo and 16.6% oligo-forms. Thus the AEC step produced a solution (AEC flow through solution) comprising 82 % mono-PEGylated EPO (82% purity).
- 10 The reaction yield of mono-PEG-EPO is 63.6%. That is, of the amount of EPO protein at the start of the process (50 mg) 63.6% was converted to mono-PEGylated EPO (32.5 mg). Compared to the prior art process of WO 2009/010270, in which the PEGylation reaction is carried out only once with a yield of about 44%, this corresponds to a yield increase of about 44%.

- For technical reasons in this experiment it was not possible to remove all EPO from the flow through on the AEC after the 2nd PEGylation reaction. The resulting composition of the 2nd reaction is therefore different from the expected values. PEGylation 1 and 3 delivered approximately 50 % EPO, 40 % mono-PEG-EPO and <10% Oligo-PEG-EPO. The 2nd PEGylation reaction was less effective, producing only 33% of mono-PEG EPO and 5% of oligo-PEG-EPO.

- These results demonstrate that AEC can isolate and recover unreacted EPO from the PEGylation reaction mixture without much of the PEGylated forms. The sum of the PEGylated forms in the recovered EPO fraction is about 5% - this might be due to the fact that the AEC column employed in this particular experiment was oversized. When sized to fit, the capacity for PEGylated species is expected to diminish due to displacement effects or competition for ligands.

- These results also show that in cycle 1 and cycle 3 that it is possible to remove EPO almost quantitatively by applying anion exchange chromatography. The intermediate product passes through the column in the flow through, and is usually of a composition about 80% mono-PEG-EPO and 20 % oligo-PEG-EPO.

EXAMPLE 3**Cyclic PEGylation, removal of erythropoietin by AEC chromatography, and purification of mono-PEG EPO by HIC**

The aim of this experiment is to PEGylate EPO in 3 cycles and to separate the reaction products (non-
5 PEGylated EPO, single PEGylated EPO (mono-PEG EPO) and oligo-PEGylated forms of EPO) using anion exchange chromatography (AEC) and HIC.

To remove the EPO in the AEC stage, Toyopearl SuperQ-650M from Tosoh Bioscience was used as the adsorbent. This is a strong anion exchanger. To separate mono-PEG EPO from the oligo forms in the HIC stage, Toyopearl Phenyl-650M from Tosoh Bioscience was used.

10 Design of the AEC column

A column with an inside diameter of 7 mm and a bed height of 100 mm was packed with Toyopearl SuperQ-650M adsorber material and operated at a flow rate of 200 cm/h (1.28 ml/min), resulting in a process step time of about 30 min at a column volume of 3.847 ml.

Cyclic PEGylation and AEC

15 An EPO stock solution with a concentration of 12.5 g/l in 25 mM bicine, 7.5 mM Na₂SO₄ was used. A starting volume of 18 ml of this stock solution, corresponding to 225 mg EPO, was used.

For the first PEGylation reaction, 18 ml of EPO stock solution was mixed with 22.5 ml of 25 mM bicine, 7.5 mM Na₂SO₄ and placed in a 50 ml Falcon tube.

328 mg of PEG reagent (molecular weight of about 30 kDa) was weighed into a 15 ml Falcon tube and
20 dissolved in 4.97 ml of 1 mM HCl.

To start the reaction, 4.5 ml of the PEG reagent solution was pipetted into the 40.5 ml EPO solution.

The reaction mixture was mixed at 300 rpm, and the temperature adjusted to 20 °C by means of a cryostat.

The reaction was run for 60 min. The concentration was then measured photometrically at 280 nm and a
25 250 µl sample was taken. The concentration was 5.56 g/l. The conductivity was measured from the remaining solution. It was 2.19 mS/cm, so no conditioning to equilibration conditions of the AEX Chromatography (2.21 mS/cm) was performed. The remaining 44.8 ml solution was purified by means of AEC. It was equilibrated with 25 mM bicine, 7.5 mM Na₂SO₄, pH 8.

Elution was carried out stepwise with 25 mM bicine, 35 mM Na₂SO₄. Fractions 1B1-1B4 were combined
30 with each other to make 16.5 ml and hereinafter will be referred to as EPO pool 1. The concentration of EPO pool 1 was photometrically determined at 5.263 g/l. A 200 µl sample was taken.

Fractions 1A1-1A5 were combined to make 62.17 ml and hereinafter are referred to as product pool 1. The concentration of this was 2.27 g/l. A 500 µl sample was taken.

The remaining volume of the EPO pool 1 was transferred to a 50 ml Falcon tube for second PEGylation. 164.73 mg of PEG reagent was weighed and dissolved with 2.4 ml of 1 mM HCl. 1.62 ml of this solution was added to the EPO Pool 1 and the second PEGylation reaction started. The reaction was again carried out at 20 °C for 60 min. Then the concentration was measured, a 250 µl sample was taken. The concentration was 4.73 g/l.

The conductivity of the remaining solution was adjusted to a conductivity of 2.1 mS/cm using 22 ml of 25 mM bicine, pH 8 buffer and purified by chromatography.

Fractions 1B2-1B4 of the second chromatography were combined to make EPO pool 2. The volume was 15 ml. A 500 µl sample was taken. The concentration measurement gave a concentration of 2.5 g/l.

10 Fractions 1A1-1A5 were combined with 62.7 ml of product pool 2. A 2 ml sample was taken. The concentration was 0.75 g/l.

The remaining volume of EPO pool 2 was transferred to a 50 ml Falcon tube. 119.4 mg of PEG reagent was weighed and dissolved with 1.9 ml of 1 mM HCl. 1.72 ml of this solution was added to EPO pool 2 and the third PEGylation reaction was started. It was again carried out at 20 °C. After a reaction time of 15 60 min, the concentration was determined and a 500 µl sample was taken. The concentration was 2.8 g/l.

The reaction product was again adjusted to equilibration conditions by means of 25 mM bicine buffer and purified by chromatography.

Fractions 1C2-1C5 were combined to make 15 ml EPO pool 3. A 4 ml sample was taken from this. The concentration was 0.429 g/l.

Fractions 1A1-1C1 were combined to 160.5 ml of Product pool 3. The concentration was 0.15 g/l. A 10 ml sample was taken.

Purification of mono-PEG EPO using HIC

A 220 mm high column having an inner diameter of 10 mm was used. The adsorbent used was Toyopearl 25 Phenyl-650M from Tosoh. The volume of the column was 17.27 ml. It was operated with a flow of 150 cm / h. This column was used to extract the target product mono-PEG EPO from a mixture of mono-PEG EPO and oligo.

For this purpose, two experiments were carried out. In the first, mono-PEG EPO and oligo were bound to the column material and then selectively eluted via a gradient. In the second, the oligo forms were bound 30 to the column material while the mono-PEG EPO was not.

Separation in HIC Bind & Elute mode

From Product pool 1 described above (a "second mixture" in accordance with the processes disclosed herein; protein concentration of 2.27 g/l) 15 ml was taken and adjusted to a conductivity of 56 mS/cm with 20 ml, 25 mM bicine, 1 M Na₂SO₄ pH 8 solution. This sample was completely dispensed through a 150 35 ml Superloop.

To equilibrate the Phenyl-650M column, 25 mM bicine, 500 mM Na₂SO₄, pH 8 buffer was used. Elution was carried out using a gradient of 500 mM Na₂SO₄ to 0 mM Na₂SO₄ over 15 CV. The corresponding chromatogram is shown in Figure 9.

Fractions 1A1– 1B1, 1B5– 2A3 and 2A4 to 2B4 were each pooled and analysed for their composition by RP-HPLC.

Separation in HIC Flow-through mode

The complete 62.7 ml of Product pool 2 (a “second mixture” in accordance with the processes disclosed herein) was used as a sample for separation, corresponding to a protein level of 47.03 mg, composed of 0.58% EPO, 79.9% mono-PEG EPO and 19.52% oligo forms. The conductivity of the sample was adjusted by means of 25 mM bicine, 1 M Na₂SO₄, pH 8 buffer, to a target to a conductivity of 50 mS/cm, and injected via a 150 ml Superloop.

The HIC resin was equilibrated with 25 mM bicine, 390 mM Na₂SO₄. Elution was carried out as a gradient of 390 mM Na₂SO₄ to 0 mM Na₂SO₄. The corresponding chromatogram is shown in Figure 10.

In practice the sample was unintentionally adjusted to 56 mS/cm (rather than the target 50 mS/cm).

Therefore a stepwise breakthrough curve can be seen in Figure 10. In the beginning only EPO is in the flow through. After about 80 ml the UV signal rises to approximately 120 mAU due to a breakthrough of the mono-PEG-EPO. When the conductivity decreases at the transition from load to post load equilibration a peak of earlier bound mono-PEG-EPO is eluted. The oligo-PEG-EPO forms are eluted in the following gradient. Nevertheless separation of mono-PEG-EPO from oligo-PEG-EPO has been shown. Flow through mode is not suitable if unreacted EPO levels are higher than the target value for the product specifications.

Results and discussion

Cyclic PEGylation and AEC

Table 5 shows the composition of the individual samples, as well as a balancing of the protein quantities.

Table 5 - Composition, masses and volumes of the individual samples during the process

	Conc [g/l]	Vol [ml]	Mass [mg]	Fracti on	Composition [%]			Mass [mg]		
					Epo	Mono	Oligo	Epo	Mono	Oligo
PEGylation 1										
Sample 1	12.5	1	12.5	-	100	0	0	12.5	0	0
Epo stock soln	12.5	18	225	-	100	0	0	225	0	0
PEG Solution	66	4.5	297	-	-	-	-	-	-	-

	Conc [g/l]	Vol [ml]	Mass [mg]	Fracti on	Composition [%]			Mass [mg]		
					Epo	Mono	Oligo	Epo	Mono	Oligo
25 mM bicine, 7.5 mM Na2SO4	-	22.5	-	-	-	-	-	-	-	-
Sample 2	5.56	0.25	1.39	-	49.8	40.59	9.61	0.692	0.564	0.134
Chromatography 1										
Sample 1	5.56	40.5	225	-	49.8	40.59	9.61	112	91.3	21.6
Epo Pool	5.263	16.5	86.84	1B1- 1B4	100	0	0	86.840	0.000	0.000
Sample 3	5.263	0.2	1.0526	-	100	0	0	1.053	0.000	0.000
Product pool	2.27	62.17	141.13	1A1- 1A5	10.16	72.45	16.23	14.338	102.246	22.905
Sample 4	2.27	0.5	1.135	-	10.16	72.45	16.23	0.115.	0.822.	0.184.
PEGylation 2										
Epo Solution	5.263	16.3	85.79	-	100	0	0	85.787	0	0
PEG Solution	68.6	1.62	111.13	-	-	-	-	-	-	-
Sample 5	4.73	0.25	1.18	-	50.88	39.91	8.61	0.602	0.472	0.102
Chromatography 2										
Sample 2	2.22	38.05	84.60	-	50.88	39.91	8.61	43.047	33.766	7.284
Epo pool	2.5	15	37.5	1B2- 1B4	98.82	1.18	0	37.058	0.443	0.000
Sample 6	2.5	0.5	1.25	-	98.82	1.18	0	1.235	0.015	0.000
Product pool	0.75	62.7	47.03	1A1- 1A5	0.58	79.9	19.52	0.273	37.573	9.179
Sample 7	0.75	2	1.5	-	0.58	79.9	19.52	0.009	1.199	0.293
PEGylation 3										

	Conc [g/l]	Vol [ml]	Mass [mg]	Fracti on	Composition [%]			Mass [mg]		
					Epo	Mono	Oligo	Epo	Mono	Oligo
Epo Solution	2.5	14.5	36.25	-	98.82	1.18	0	35.822	0.428	0.000
PEG Solution	62.84	1.72	108.08 48	-	-	-	-	-	-	-
Sample 8	2.	0.5	1.4	-	19.22	46.1	34.69	0.269	0.645	0.486
Chromatography 3										
Sample 2	-	150	-	-	19.22	46.1	34.69	-	-	-
Epo pool	0.42	15	6.3	1C2- 1C5	98.99	1.01	0	6.236	0.064	0
Sample 9	0.42	4	1.68	-	98.99	1.01	0	1.663	0.017	0
Product pool	0.15	160.5	24.08	1A1- 1C1	0.29	58.45	41.25	0.070	14.072	9.931
Sampling 10	0.15	5	0.75	-	0.29	58.45	41.25	0.002	0.438	0.309
Total balance Product pools vs Load Sample 1					2.77	68.40	18.67	6.24	153.89	42.01

The first cycle on the AEC was inadvertently overloaded. Due to this, the recovery of unreacted EPO in this experiment is only about 78%. Because the column has been overloaded with EPO there is no PEGylated form in the elution pool. That corroborates that proper sizing of the AEC column reduces the content of PEGylated forms in the AEC Elution Pool and thereby maximizes the yield of PEGylated forms in the intermediate pool. The AEC runs after PEGylation 2 and 3 are not fully loaded with EPO and therefore there is a small amount of PEGylated-EPO in the elution pool, in both cases <2%.

The results of PEGylation 1 and 2 show the consistency of the reaction outcome. Under the chosen conditions the results are about 50% EPO, 40% mono-PEG-EPO and <10% oligo-PEG-EPO.

PEGylation 3 as the final PEGylation uses different conditions to favour the formation of the maximum mono-PEG-EPO content that is possible. This worked well with 46% mono-PEG-EPO content.

The composition each of the intermediate product pools are different. The Pool of the first flow through operation consists of about 10 % EPO (due to the breakthrough mentioned above), 72% of mono-PEG-EPO and 16% of oligo-PEG EPO. Without the breakthrough of EPO the composition would have been

identical to Run 2 (80% mono-PEG-EPO and 20% oligo-PEG-EPO). The 3rd run employed different PEGylation conditions, which resulted in a higher content of oligo-PEG-EPO in the reaction mixture. This translates into a pool composition of 59% mono-PEG-EPO and 41% oligo Peg EPO.

- 5 The total yield of mono-PEG-EPO in the intermediate pool after 3 cycles of "PEGylation reaction plus EPO recovery by AEC" is 68.4%. This is already > 50% better than the original process but could be even better if there hasn't been a significant amount of EPO lost during the flow through loading of the first AEC run.

Purification of Mono-PEG EPO by HIC – Separation in Bind & Elute mode

Table 5 shows the results of the RP HPLC analysis.

10 Table 5- RP-HPLC analysis data for pools of HIC in Bind & Elute mode

Description	Fraction	Epo [%]	Mono-PEG Epo [%]	Oligo forms [%]
Starting sample	-	10.16	72.45	16.23
Epo pool	1A1-1B1	100	0	0
Mono-PEG Epo Pool	1B5-2A3	0	100	0
Oligo pool	2A4-2B4	0	2.33	97.67

- 15 The use of the intermediate pool from AEC Run 1 which contained EPO was used to show the performance of the final purification step by hydrophobic interaction chromatography. The EPO and the mono-PEG-EPO Pool are 100% pure. A small amount of mono-PEG-EPO can still be found in the Oligo-PEG-EPO Pool.

Purification of Mono-PEG EPO by HIC – Separation in Flow-Through mode

Due to the low protein concentrations of the individual fractions, no meaningful results could be generated by means of RP HPLC. However the chromatogram shows that separation of mono-PEGylated EPO from non-PEGylated and oligo-PEGylated forms was achieved.

20 **References**

A number of publications are cited above in order to more fully describe and disclose the invention and the state of the art to which the invention pertains. Full citations for these references are provided below. The entirety of each of these references is incorporated herein by reference for all purposes.

Bristow, A, *Pharmeuropa Spec. Issue Biologicals BRP Erythropoietin Bio 97-2* (1997) 31-48

- 25 Delgado, C., et al., *Crit. Rev. Ther. Drug Carrier 30 Systems 9* (1992) 249-304)

Fee & Van Alstine, *Chemical Engineering Science*, 2016, 61,924-939

- Francis, G.E., et al., *Int. J. Hematol.* 68 (1998) 1-18
 Ingold et al, *React. Chem. Eng.*, 2016, 1,218.
 Lu, Y., et al., *Reactive Polymers* 22 (1994) 221-229
 Morpurgo, M., et al., *J. Bioconjug. Chem.* 7 (1996) 363-368.
 5 Pfister et al, *Reac React. Chem. Eng.*, 2016, 1,204
 Pfister et al. *Biotechnology and Bioengineering*, 2016, 113, 1711-1718.
 Veronese, F.M., *Biomaterials* 22 (2001) 405-417.
 WO 2009/010270
 WO 2012/035037
 10 For standard molecular biology techniques, see Sambrook, J., Russel, D.W. *Molecular Cloning, A Laboratory Manual*. 3 ed. 2001, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press.

The following numbered statements relate to aspects of the present disclosure and form part of the description.

- 15 1. A process for producing a mono-PEGylated protein composition comprising at least about 90% mono-PEGylated protein, the process comprising:
- a) providing a first mixture comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein
 - b) subjecting the first mixture to an ion exchange chromatography (IEC) step to provide an IEC flow-
 20 through solution in which the fraction of PEGylated protein is increased relative to the first mixture; the IEC step comprising
 applying the first mixture to an IEC material under conditions suitable for binding non-PEGylated protein;
 - c) collecting the IEC flow-through solution from step b) to provide a second mixture comprising
 25 mono-PEGylated protein and oligo-PEGylated protein; and
 - d) subjecting the second mixture to a hydrophobic interaction chromatography (HIC) step to provide a mono-PEGylated protein composition in which the fraction of mono-PEGylated protein is increased relative to the second mixture, wherein the mono-PEGylated protein composition comprises at least about 90% mono-PEGylated protein.
- 30 2. The process according to statement 1, wherein the protein is a hormone, a cytokine, an enzyme or an antibody.
3. The process according to statement 1 or statement 2, wherein the protein is erythropoietin.
4. The process according to any one of the preceding statements, wherein the IEC step is an AEC step.
5. A process for producing a mono-PEGylated protein composition comprising at least about 90%
 35 mono-PEGylated protein, wherein the protein is erythropoietin, the process comprising:

- e) providing a first mixture comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein
- f) subjecting the first mixture to an anion exchange chromatography (AEC) step to provide an AEC flow-through solution in which the fraction of PEGylated protein is increased relative to the first mixture; the AEC step comprising
- 5 applying the first mixture to an AEC material under conditions suitable for binding non-PEGylated protein;
- g) collecting the AEC flow-through solution from step b) to provide a second mixture comprising mono-PEGylated protein and oligo-PEGylated protein; and
- 10 h) subjecting the second mixture to a hydrophobic interaction chromatography (HIC) step to provide a mono-PEGylated protein composition in which the fraction of mono-PEGylated protein is increased relative to the second mixture, wherein the mono-PEGylated protein composition comprises at least about 90% mono-PEGylated protein.
6. The process according to any one of the preceding statements, wherein the IEC material has a
- 15 binding capacity for the PEGylated protein of less than about 1.5 g/L.
7. The process according to any one of statements 4 to 6, wherein the AEC material has a binding capacity for the PEGylated protein of less than about 1.5 g/L.
8. The process according to any one of the preceding statements wherein:
- i. the first mixture comprises less than 25% oligo-PEGylated protein; and/or
- 20 ii. the IEC flow-through solution comprises at least 90% PEGylated protein; and/or
- iii. the mono-PEGylated protein composition comprises at least about 95%, 98%, 99%, or 99.9% mono-PEGylated protein.
9. The process according to any one of the preceding statements wherein:
- i. the first mixture comprises less than 25% oligo-PEGylated protein; and/or
- 25 ii. the AEC flow-through solution comprises at least 90% PEGylated protein; and/or
- iii. the mono-PEGylated protein composition comprises at least about 95%, 98%, 99%, or 99.9% mono-PEGylated protein.
10. The process according to any one of the preceding statements, wherein step a) further comprises performing a PEGylation reaction comprising reacting the non-PEGylated protein with a
- 30 PEGylation reagent.

11. The process according to statement 10, wherein the PEGylation reaction is performed at a pH of about 7.0 to 9.0, and wherein the PEG/protein molar ratio is about 0.6 – 1.0.
12. The process according to statement 10 or 11, comprising
 - performing a first cycle comprising steps a), b) and c), wherein step b) further comprises eluting
5 non-PEGylated protein from the IEC material to provide an IEC eluate, and
 - performing a second cycle of steps a), b) and c), in which the non-PEGylated protein eluted in step b) of the first cycle is added to the PEGylation reaction of step a).
13. The process according to statement 10 or 11, wherein the IEC step is an AEC step, comprising
 - performing a first cycle comprising steps a), b) and c), wherein step b) further comprises eluting
10 non-PEGylated protein from the AEC material to provide an AEC eluate, and
 - performing a second cycle of steps a), b) and c), in which the non-PEGylated protein eluted in step b) of the first cycle is added to the PEGylation reaction of step a).
14. The process according to statement 12, wherein eluting non-PEGylated protein from the IEC material uses an elution buffer comprising less than or equal to about 45 mM salt
15. 15. The process according to statement 13, wherein eluting non-PEGylated protein from the AEC material uses an elution buffer comprising less than or equal to about 45 mM salt.
16. The process according to any one of statements 12 to 15, wherein the process comprises three, four or five cycles, and wherein step b) of each cycle comprises eluting non-PEGylated protein from the IEC material to provide an IEC eluate, and wherein the non-PEGylated protein eluted in step b) is
20 added to the PEGylation reaction of step a) in the next cycle.
17. The process according to any one of statements 12 to 15, wherein the IEC step is an AEC step, and wherein the process comprises three, four or five cycles, and wherein step b) of each cycle comprises eluting non-PEGylated protein from the AEC material to provide an AEC eluate, and wherein the non-PEGylated protein eluted in step b) is added to the PEGylation reaction of step a) in the next cycle
- 25 18. The process according to any one of statements 12 to 17, wherein the IEC eluate from step b) of a cycle is added directly to the PEGylation reaction of step a) of the next cycle.
19. The process according to any one of statements 12 to 17, wherein the IEC step is an AEC step, and wherein the AEC eluate from step b) of a cycle is added directly to the PEGylation reaction of step a) of the next cycle.
- 30 20. The process according to any one of statements 12 to 17, in which the non-PEGylated protein eluted in step b) of a cycle is added to the PEGylation reaction of step a) of the next cycle, and wherein fresh non-PEGylated protein is also added to step a) in order to maintain substantially constant PEGylation reaction conditions in step a) of each cycle.

21. The process according to any one of the preceding statements, comprising performing two or more cycles of steps a), b) and c), wherein
- step c) further comprises pooling the flow-through solution collected from of each IEC step to provide a second mixture which is a pooled second mixture, and wherein
- step d) comprises subjecting the second mixture to an HIC step.
22. The process according to any one of the preceding statements, comprising performing two or more cycles of steps a), b) and c), wherein
- step c) further comprises pooling the flow-through solution collected from of each AEC step to provide a second mixture which is a pooled second mixture, and wherein
- step d) comprises subjecting the second mixture to an HIC step.
23. The process according to any one of the preceding statements, wherein the IEC step is an AEC step, and wherein
- i. the AEC material is Toyopearl Super Q 650 M; and/or
 - ii. the AEC step is performed at pH of about 7.0 to 9.0; and/or
 - iii. the AEC step is performed at a conductivity of about 1.0 to 3.0 mS/cm; and/or
 - iv. the first mixture is applied to the AEC material as a AEC load solution comprising about 10 – 30 mM bicine and about 1 – 10 mM Na₂SO₄.
24. The process according to any one of the preceding statements, wherein step d) comprises subjecting the second mixture to a HIC step in flow-through mode to provide a HIC flow-through solution in which the fraction of mono-PEGylated protein is increased relative to the second mixture,
- the HIC step comprising applying the second mixture to a HIC material under conditions suitable for binding oligo-PEGylated protein, wherein the HIC flow-through provides the mono-PEGylated protein composition.
25. The process according to any one of statements 1 to 23, wherein step d) comprises subjecting the second mixture to a HIC step in bind and elute mode to provide a HIC eluate in which the fraction of mono-PEGylated protein is increased relative to the second mixture, the HIC step comprising
- applying the second mixture to a HIC material under conditions suitable for binding mono-PEGylated protein and oligo-PEGylated protein,
- eluting the mono-PEGylated protein from the HIC material to provide a HIC eluate, wherein the HIC eluate provides the mono-PEGylated protein.

26. The process according to statement 24, wherein
- i. the HIC material is Toyopearl Phenyl 650M; and/or
 - ii. the HIC step is performed at a pH of about 7.0 to 9.0; and/or
 - iii. the HIC step is performed at a conductivity of about 30-40 mS/cm; and/or
 - 5 iv. the second mixture is applied to the HIC material as a HIC load solution comprising about 25 mM bicine and about 390 mM Na₂SO₄.
27. The process according to statement 25, wherein
- i. the HIC material is Toyopearl Phenyl 650M; and/or
 - ii. the HIC step is performed at a pH of about 7.0 to 9.0; and/or
 - 10 iii. the step of eluting the mono-PEGylated protein from the HIC material comprises applying a gradient of from about 50 mS/cm to 0 mS/cm; and/or
 - iv. the second mixture is applied to the HIC material as a HIC load solution comprising about 25 mM bicine and about 500 mM Na₂SO₄.
28. The process according to any one of the preceding statements, wherein
- 15 i. the IEC step and HIC step are performed at substantially the same pH; or
 - ii. the PEGylation reaction, IEC step and HIC step are performed at substantially the same pH.
29. The process according to any one of the preceding statements, wherein
- i. the AEC step and HIC step are performed at substantially the same pH; or
 - ii. the PEGylation reaction, AEC step and HIC step are performed at substantially the same pH.
- 20 30. The process according to any one of the preceding statements, wherein mono-PEGylated protein comprises a PEG residue having a molecular weight of at least about 20kDa.
31. The process according to any one of the preceding statements, further comprising formulating the protein composition with a pharmaceutically acceptable carrier to provide a pharmaceutical composition.
- 25 32. A process for producing a PEGylated protein mixture, the process comprising:
- a) reacting a non-PEGylated protein with a PEGylation reagent to produce a mixture of reaction products comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein;
 - b) 30 subjecting the mixture of reaction products to an ion exchange chromatography (IEC) step to provide an IEC flow-through solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products;

the IEC step comprising applying the mixture of reaction products to an IEC material under conditions suitable for binding non-PEGylated protein; and

c) collecting the IEC flow-through solution from step b) to provide a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein,

5 wherein two or more cycles of steps a), b) and c) are performed, in which non-PEGylated protein is recovered in step b) by eluting an IEC eluate from the IEC material and the eluted non-PEGylated protein is used in the PEGylation reaction of step a) of the next cycle.

33. The process according to statement 32, wherein the protein is a hormone, a cytokine, an enzyme or an antibody.

10 34. The process according to statement 32 or 33, wherein the protein is erythropoietin.

35. The process according to any one of statements 32 to 34, wherein the IEC step is an anion exchange chromatography (AEC) step.

36. A process for producing a PEGylated protein mixture, wherein the protein is erythropoietin, the process comprising:

15 a) reacting a non-PEGylated protein with a PEGylation reagent to produce a mixture of reaction products comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein;

b) subjecting the mixture of reaction products to an anion exchange chromatography (AEC) step to provide an AEC flow-through solution in which the fraction of PEGylated protein is
20 increased relative to the mixture of reaction products;

the AEC step comprising applying the mixture of reaction products to an AEC material under conditions suitable for binding non-PEGylated protein; and

c) collecting the AEC flow-through solution from step b) to provide a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein,

25 wherein two or more cycles of steps a), b) and c) are performed, in which non-PEGylated protein is recovered in step b) by eluting an AEC eluate from the AEC material and the eluted non-PEGylated protein is used in the PEGylation reaction of step a) of the next cycle.

37. The process according to any one of statements 32 to 36, wherein the mixture of reaction products comprises at least about 20% non-PEGylated protein.

30 38. The process according to any one of statements 32 to 37 wherein the process comprises three, four or five cycles, in which in step b) of each cycle non-PEGylated protein is recovered and used in the PEGylation reaction of step a) in the next cycle.

39. The process according to any one of statements 32 to 38, wherein step c) further comprises pooling the flow through solution collected from each IEC step to provide a PEGylated protein mixture which is a pooled PEGylated protein mixture.
40. The process according to any one of statements 35 to 39, wherein step c) further comprises pooling the flow through solution collected from each AEC step to provide a PEGylated protein mixture which is a pooled PEGylated protein mixture.
41. The process according to any one of statements 32 to 40, wherein eluting non-PEGylated protein from the IEC material uses an elution buffer comprising less than or equal to about 45 mM salt.
42. The process according to any one of statements 35 to 40, wherein eluting non-PEGylated protein from the AEC material uses an elution buffer comprising less than or equal to about 45 mM salt.
43. The process according to any one of statements 32 to 42, wherein the IEC eluate of a cycle is added directly to the PEGylation reaction of step a) of the next cycle.
44. The process according to any one of statements 35 to 42, wherein the AEC eluate of a cycle is added directly to the PEGylation reaction of step a) of the next cycle.
45. The process according to any one of statements 32 to 44, in which the non-PEGylated protein is recovered and used in the PEGylation reaction of step a) in the next cycle, and wherein fresh non-PEGylated protein is also added to step a) in order to maintain substantially constant PEGylation reaction conditions in step a) of each cycle.
46. The process according to any one of statements 32 to 45 wherein the IEC material has a binding capacity for the PEGylated protein of less than about 1.5 g/L.
47. The process according to any one of statements 35 to 45 wherein the AEC material has a binding capacity for the PEGylated protein of less than about 1.5 g/L.
48. The process according to any one of statements 32 to 47, wherein:
- i. the mixture of reaction products comprises less than 25% oligo-PEGylated protein; and/or
 - ii. the IEC flow-through solution comprises at least 90% PEGylated protein.
49. The process according to any one of statements 35 to 47, wherein:
- i. the mixture of reaction products comprises less than 25% oligo-PEGylated protein; and/or
 - ii. the AEC flow-through solution comprises at least 90% PEGylated protein.
50. The process according to any one of statements 35 to 49, wherein
- i. the AEC material is Toyopearl Super Q 650 M; and/or

- ii. the AEC step is performed at pH of about 7.0 to 9.0; and/or
 - iii. the AEC step is performed at a conductivity of about 1.0 to 3.0 mS/cm; and/or
 - iv. the mixture of reaction products is applied to the AEC material as a AEC load solution comprising about 10 – 30 mM bicine and about 1 – 10 mM Na₂SO₄.
- 5 51. The process according to statement 50 wherein the protein is erythropoietin.
52. The process according to any one of statements 32 to 51, wherein the PEGylation reaction is performed at a pH of about 7.0 to 9.0, and wherein the PEG/protein molar ratio is about 0.6 – 1.0.
53. A process for producing a mono-PEGylated protein composition comprising at least about 90% mono-PEGylated protein, the process comprising:
- 10 subjecting the PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein produced by any one of statements 32 to 51 to a purification process separating mono-PEGylated protein and oligo-PEGylated protein; and
- recovering a mono-PEGylated protein composition in which the fraction of mono-PEGylated protein is increased relative to the PEGylated protein mixture, wherein the mono-
- 15 PEGylated protein composition comprises at least about 90% mono-PEGylated protein.
54. The process of statement 53, wherein the purification process comprises subjecting the PEGylated protein mixture to a hydrophobic interaction chromatography (HIC) step in flow-through mode to provide a HIC flow-through solution in which the fraction of mono-PEGylated protein is increased relative to the PEGylated protein mixture,
- 20 the HIC step comprising applying the PEGylated protein mixture to a HIC material under conditions suitable for binding oligo-PEGylated protein, wherein the HIC flow-through provides the mono-PEGylated protein composition.
55. The process of statement 53, wherein the purification process comprises subjecting the PEGylated protein mixture to a hydrophobic interaction chromatography (HIC) step in bind and elute mode to
- 25 provide a HIC eluate in which the fraction of mono-PEGylated protein is increased relative to the PEGylated protein mixture, the HIC step comprising
- applying the second mixture to a HIC material under conditions suitable for binding mono-PEGylated protein and oligo-PEGylated protein,
- eluting the mono-PEGylated protein from the HIC material to provide a HIC eluate, wherein the
- 30 HIC eluate provides the mono-PEGylated protein.
56. The process according to any one of statements 32 to 55 wherein the mono-PEGylated protein composition comprises at least about 95%, 98%, 99%, or 99.9% mono-PEGylated protein.

Claims:

1. A process for producing a PEGylated protein mixture the process comprising:

a) reacting a non-PEGylated protein with a PEGylation reagent to produce a mixture of reaction products comprising non-PEGylated protein and PEGylated protein, wherein the PEGylated protein comprises mono-PEGylated protein and oligo-PEGylated protein;

b) subjecting the mixture of reaction products to an ion exchange chromatography (IEC) step to provide an IEC flow-through solution in which the fraction of PEGylated protein is increased relative to the mixture of reaction products;

the IEC step comprising applying the mixture of reaction products to an IEC material under conditions suitable for binding non-PEGylated protein; and

c) collecting the IEC flow-through solution from step b) to provide a PEGylated protein mixture comprising mono-PEGylated protein and oligo-PEGylated protein,

wherein non-PEGylated protein is recovered in step b) by eluting an IEC eluate from the IEC material and the eluted non-PEGylated protein is used in a subsequent PEGylation reaction.

2. The process according to claim 1, wherein the protein is a hormone, a cytokine, an enzyme or an antibody.

3. The process according to claim 1 or claim 2, wherein the protein is erythropoietin.

4. The process according to any one of the preceding claims, wherein the IEC step is an anion exchange (AEC) step.

5. The process according to any one of the preceding claims, wherein two or more cycles of steps a), b) and c) are performed, in which non-PEGylated protein is recovered in step b) by eluting an IEC eluate from the IEC material and the eluted non-PEGylated protein is used in a subsequent PEGylation reaction.

6. The process according to claim 5 wherein the process comprises three, four or five cycles, in which in step b) of each cycle non-PEGylated protein is recovered and used in a subsequent PEGylation reaction.

7. The process according to claims 5 or claim 6, wherein step c) further comprises pooling the flow-through solution collected from each IEC step to provide a PEGylated protein mixture which is a pooled PEGylated protein mixture.

5 8. The process according to any one of the preceding claims, wherein eluting non-PEGylated protein from the IEC material uses an elution buffer comprising less than or equal to 45 mM salt.

9. The process according to any one of the preceding claims, wherein the step b) IEC eluate is added directly to the subsequent PEGylation reaction.

10

10. The process according to any one of the preceding claims, in which the non-PEGylated protein is recovered and used in a subsequent PEGylation reaction, and wherein fresh non-PEGylated protein is also added to the subsequent PEGylation reaction in order to maintain substantially constant PEGylation reaction conditions in each PEGylation reaction.

15

11. The process according to any one of the preceding claims wherein the IEC material has a binding capacity for the PEGylated protein of less than 1.5 g/L.

12. The process according to any one of the preceding claims, wherein:

- 20 i. the mixture of reaction products comprises less than 25% oligo-PEGylated protein; and/or
- ii. the IEC flow-through solution comprises at least 90% PEGylated protein.

13. The process according to any one of the preceding claims, wherein the IEC step is an AEC step and wherein,

25

- i. the AEC material is Toyopearl Super Q 650 M; and/or
- ii. the AEC step is performed at pH of 7.0 to 9.0; and/or
- iii. the AEC step is performed at a conductivity of 1.0 to 3.0 mS/cm; and/or
- iv. the mixture of reaction products is applied to the AEC material as a AEC load solution
- 30 comprising 10 – 30 mM bicine and 1 – 10 mM Na₂SO₄.

14. The process according to any one of the preceding claims, wherein the PEGylation reaction is performed at a pH of 7.0 to 9.0, and wherein the PEG/protein molar ratio is 0.6 – 1.0.

15. The process according to any one of the preceding claims, further comprising:

- 5 (d) subjecting the PEGylated protein mixture produced in step (c) of claim 1 and comprising mono-PEGylated protein and oligo-PEGylated protein to a purification process separating mono-PEGylated protein and oligo-PEGylated protein; and
- (e) recovering a mono-PEGylated protein composition in which the fraction of mono-PEGylated protein is increased relative to the PEGylated protein mixture and wherein the mono-PEGylated protein composition comprises at least 90% mono-PEGylated protein.
- 10

16. The process of claim 15, wherein the purification process comprises subjecting the PEGylated protein mixture to a hydrophobic interaction chromatography (HIC) step in flow-through mode to provide a HIC flow-through solution in which the fraction of mono-PEGylated protein is increased relative to the PEGylated protein mixture,

15

the HIC step comprising applying the PEGylated protein mixture to a HIC material under conditions suitable for binding oligo-PEGylated protein, wherein the HIC flow-through provides the mono-PEGylated protein composition.

20 17. The process of claim 15, wherein the purification process comprises subjecting the PEGylated protein mixture to a hydrophobic interaction chromatography (HIC) step in bind and elute mode to provide a HIC eluate in which the fraction of mono-PEGylated protein is increased relative to the PEGylated protein mixture, the HIC step comprising

applying the second mixture to a HIC material under conditions suitable for binding mono-PEGylated protein and oligo-PEGylated protein,

25

eluting the mono-PEGylated protein from the HIC material to provide a HIC eluate, wherein the HIC eluate provides the mono-PEGylated protein.

18. The process according to any one of claims 15 to 17 wherein the mono-PEGylated protein composition comprises at least 95%, 98%, 99%, or 99.9% mono-PEGylated protein.

30

19. The process according to any one of the preceding claims, wherein

- i. the IEC step and HIC step are performed at substantially the same pH; or

ii. the PEGylation reaction, IEC step and HIC step are performed at substantially the same pH.

20. The process according to any one of the preceding claims, wherein mono-PEGylated protein comprises a PEG residue having a molecular weight of at least 20kDa.

5

21. The process according to any one of the preceding claims, further comprising formulating the protein composition with a pharmaceutically acceptable carrier to provide a pharmaceutical composition.