

According to the above route, a 22-step reaction is required from the starting material to the diphenylmethane derivative, and 14 steps and 15 steps are required respectively from the starting material to the key intermediates compound 1 and compound 2, with complicated overall operations and a high production cost, therefore the route is difficult to meet the needs of the pharmaceutical industry.

SUMMARY

Regarding characteristics of a long synthetic route, complicated operations and a high production cost for the existing diphenylmethane derivatives, the present invention provides an improved method for preparing compound 1 and compound 2.



1



2

The present invention provides a method for preparing compound 1 and compound 2, comprising the following steps.

Compound 1:

step 1-1. taking 2,3-dihydrobenzofuran-7-amine~~aniline~~ as a raw material, and carrying out selective dibromination by using a bromination reagent to obtain 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~;

step 1-2. subjecting the 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~ to a Sandmeyer reaction for chlorination to obtain 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran;

step 1-3. selectively removing bromine from the 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran by selectively using a strong base, followed by adding a formylation reagent to obtain the compound 1.

The synthetic route is as follows.




1

DETAILED DESCRIPTION OF THE EMBODIMENTS

Preferred embodiments of the present invention are described in detail below so as to make those skilled in the art to understand the advantages and features of the present invention more easily and thus to define the protection scope of the present invention more clearly.

Example 1: Synthesis of 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~




8.7 g of 2,3-dihydrobenzofuran-7-amine~~aniline~~ and 200 mL of ethyl acetate were added to a 1000 mL of reaction flask, and then 41.4 g of bromine was added dropwise at a controlled temperature between 10°C to 20°C, in which the addition amount of bromine was optimized via experiments. If too little bromine is added, the reaction will be incomplete; and if too much bromine is added, the system will become complicated and generate increased by-products. The reaction temperature was controlled mainly for the prevention of the increase of by-products. After the completion of the dropwise addition, the system was filtered. The filter cake was added to a mixture solvent of 100 mL water and 100 mL ethyl acetate, followed by dropwise adding 10% sodium hydroxide solution to adjust the pH to a value between 8 to 10. In which, the system is acidic after the reaction and some products forms salts. The pH was adjusted to the value of 8 to 10 for freeing of salts. If the pH is too low, the freeing of salts is incomplete; and if the pH is too high, more alkali will be consumed. After that, the organic layer was separated out, dried with anhydrous sodium sulfate and concentrated under reduced pressure to give 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~ (15.73 g, 83.4% yield) as a white solid.

Example 2: Synthesis of 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~

5 g of 2,3-dihydrobenzofuran-7-amine~~aniline~~ and 100 mL of N, N-dimethylformamide were added to a 100 mL of reaction flask, and cooled down to a temperature between -5°C and 0°C, followed by dropwise adding 13.17 g of N-bromosuccinimide in batches and reaction under stirring for 30 minutes, in which the N-bromosuccinimide as a solid has the advantages of being calculated and controlled easily while bromine is a corrosive liquid, irritating and inconvenient to be weighed and used. After that, the reaction solution was added to 200 mL of water, and extracted with ethyl acetate twice. The organic phases were combined, washed with 10% sodium

hydroxide solution, dried with anhydrous sodium sulfate and concentrated under reduced pressure to give 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~ (8.47 g, 78.2% yield).

Example 3: Synthesis of 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran



10 g of 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~ and 50 mL of concentrated hydrochloric acid were added to a 250 mL of reaction flask, cooled down to 0°C to 5°C, and a solution of 2.47 g sodium nitrite in 8 mL water was dropwise added under a kept temperature. After the completion of the dropwise addition, the system was reacted for 30 minutes under the kept temperature, and then 6.76 g of cuprous chloride was added in batches, warmed to a room temperature and reacted for 2 hours. After that, ethyl acetate was added for extraction, the organic layer was washed with 5% sodium hydroxide solution, and concentrated to obtain a crude product, which was subjected to silica gel column chromatography to give 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran (9.01 g, 84.5% yield) as a white solid.

Example 4: Synthesis of 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran

10 g of 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~, 100 mL of acetonitrile and 9.18 g of cupric chloride were added to a 250 mL of reaction flask, heated to a temperature to 50°C, and 8.0 g of isoamyl nitrite was dropwise added. After the completion of dropwise addition, the system was reacted for 30 minutes under the kept temperature. After that, the system was cooled down to a room temperature and concentrated to remove the solvent, followed by adding ethyl acetate and filtration. The filtrate solution was washed with 5% sodium hydroxide solution and concentrated to obtain a crude product, which was subjected to silica gel column chromatography to give 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran (8.78 g, 82.3% yield) as a white solid.


Example 5: Synthesis of compound 1




1

CLAIMS

[CLAIM 1] A method for preparing compound 1 and compound 2 having a structure as shown below, wherein the method comprises the following steps:



1



2

1) subjecting 2,3-dihydrobenzofuran-7-amine~~aniline~~ as a raw material to a selective -dibromination with a brominating reagent to obtain 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~;

2) subjecting 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~ obtained in the step 1) to Sandmeyer reaction for chlorination to obtain 4,6-dibromo -7-chloro-2,3-dihydrobenzofuran;

3-1) selectively debrominating from 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran obtained in the step 2) using a strong base, and then adding a formylation reagent to obtain compound 1;

3-2) selectively debrominating from 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran obtained in the step 2) using a strong base, and then reacting with 4-cyclopropyl benzaldehyde to obtain compound 2.

[CLAIM 2] The method according to claim 1, characterized in that, in the step 1), the brominating reagent ~~comprises~~ is at least one of bromine, N -bromosuccinimide (NBS), and dibromohydantoin~~, preferably N -bromosuccinimide;~~.

[CLAIM 3] The method according to claim 1~~preferably~~, wherein the selective-dibromination is carried out at a reaction temperature in a range from 10°C to 20°C.

[CLAIM ~~3~~4] The method according to claim 1, characterized in that, in the step 2), a reagent used in the Sandmeyer reaction comprises at least one of hydrochloric acid/sodium nitrite/ cuprous chloride system, isoamyl nitrite ester/cupric chloride system and t-butyl nitrite/cupric chloride system.

[CLAIM ~~4~~5] The method according to claim 1, characterized in that, in the step 3-1), the strong base ~~comprises~~ is at least one of n-butyllithium, tert-butyllithium, and isopropylmagnesium chloride~~, preferably isopropylmagnesium chloride~~.

[CLAIM ~~5~~6] The method according to claim 1, characterized in that, in the step 3-1), the formylation reagent ~~comprises~~ is one of N,N -dimethylformamide, N- formylmorpholine , and ethyl formate~~, preferably N,N -dimethylformamide~~.

[CLAIM ~~6~~7] The method according to claim 1, characterized in that, in the step 3-2), the strong base is at least one selected from n-butyllithium, tert-butyllithium, or isopropylmagnesium chloride.

[CLAIM ~~7~~8] The method according to claim 2, characterized in that, the step 1) comprises: filtering after the dibromination, adding the filter cake to a mixed solvent of water and ethyl acetate, followed by sodium hydroxide solution dropwise to adjust pH in a range from 8 to 10, separating and drying the organic layer over anhydrous sodium sulfate, and concentrating the residue under reduced pressure to obtain 4,6-dibromo-2,3-dihydrobenzofuran-7- amine~~aniline; further preferably, in the mixed solvent of water and ethyl acetate, water and ethyl acetate has a volume ratio of 1:1, and sodium hydroxide has a mass concentration of 10%~~.

[CLAIM 8] The method according to claim 3, characterized in that, the step 2) comprises: adding 4,6-dibromo-2,3-dihydrobenzofuran-7-amine~~aniline~~ and concentrated hydrochloric acid to a reaction flask and cooling down to a temperature of 0-5℃; adding sodium nitrite solution dropwise while maintaining at a constant temperature; after the dropwise addition, reacting for 20-60 minutes while maintaining at a constant temperature; then adding cuprous chloride in

batches and reacting for 1-3 hours at room temperature; extracting with ethyl acetate, washing the organic layer with sodium hydroxide solution and concentrating to obtain a crude product; and separating and purifying the crude product to obtain 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran.

[CLAIM ~~9~~10] The method according to claim ~~4~~3, characterized in that, the step 3-1) comprises: to a reaction flask, adding 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran and tetrahydrofuran, followed by isopropyl magnesium chloride in tetrahydrofuran dropwise under nitrogen protection; after the dropwise addition, stirring for 0.5-1.5 hours while maintaining at a constant temperature; then adding N,N -dimethylformamide dropwise while maintaining at a constant temperature, and continuing to react for 20-60 minutes while maintaining at a constant temperature; pouring the reaction solution into hydrochloric acid, extracting with ethyl acetate, drying the organic layer over anhydrous sodium sulfate; and concentrating and separating to obtain compound 1.

[CLAIM ~~10~~11] The method according to claim ~~6~~7, characterized in that, the step 3-2) comprises: adding 4,6-dibromo-7-chloro-2,3-dihydrobenzofuran and tetrahydrofuran to a reaction flask, cooling down to -30℃ under nitrogen protection; adding isopropyl magnesium chloride in tetrahydrofuran dropwise, followed by n-butyllithium in n-hexane dropwise; after the dropwise addition, stirring for 20-60 minutes while maintaining at a constant temperature; then adding 4-cyclopropylbenzaldehyde in tetrahydrofuran dropwise while maintaining at a constant temperature; after the dropwise addition, continuing to react for 20-60 minutes while maintaining at a constant temperature; pouring the reaction solution into hydrochloric acid, extracting with ethyl acetate, and drying the organic layer over anhydrous sodium sulfate; and concentrating, separating and purifying to obtain compound 2.