

COMPOSITIONS AND RELATED METHODS FOR CONTROLLING VECTOR-BORNE DISEASES**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims priority to U.S. Provisional Application No. 62/450,032, filed on January 24, 2017, and U.S. Provisional Application No. 62/583,925, filed on November 9, 2017, the contents of which are hereby incorporated herein by reference in their entireties.

BACKGROUND

Insects function as vectors for pathogens causing severe disease in humans and animals such as dengue, trypanosomiasis, and malaria. Vector-borne diseases that infect animals, such as livestock, represent a major global public health burden. Thus, there is need in the art for methods and compositions to control insects that carry vector-borne diseases.

SUMMARY OF THE INVENTION

Disclosed herein are compositions and methods for modulating the fitness of insects for controlling the spread of vector-borne diseases in animals. The composition includes an agent that alters a level, activity, or metabolism of one or more microorganisms resident in a host, the alteration resulting in a modulation in the host's fitness.

In one aspect, provided herein is a method of decreasing fitness of a vector (e.g., insect vector) for an animal pathogen, the method including delivering an antimicrobial peptide having at least 90% sequence identity (e.g., at least 90%, 92%, 94%, 96%, 98%, or 100% sequence identity) with one or more of the following: cecropin (SEQ ID NO: 82), melittin, copsin, drosomycin (SEQ ID NO: 93), dermcidin (SEQ ID NO: 81), andropin (SEQ ID NO: 83), moricin (SEQ ID NO: 84), ceratotoxin (SEQ ID NO: 85), abaecin (SEQ ID NO: 86), apidaecin (SEQ ID NO: 87), prophenin (SEQ ID NO: 88), indolicidin (SEQ ID NO: 89), protegrin (SEQ ID NO: 90), tachyplesin (SEQ ID NO: 91), or defensin (SEQ ID NO: 92) to the vector.

In some embodiments, the delivery includes delivering the antimicrobial peptide to at least one habitat where the vector grows, lives, reproduces, feeds, or infests.

In some embodiments, the antimicrobial peptide may be delivered in an insect comestible composition for ingestion by the vector.

In some embodiments, the antimicrobial peptide may be formulated as a liquid, a solid, an aerosol, a paste, a gel, or a gas composition.

In some embodiments, the insect may be at least one of a mosquito, midge, louse, sandfly, tick, triatomine bug, tsetse fly, or flea.

In another aspect, provided herein is a composition including an antimicrobial peptide having at least 90% sequence identity (e.g., at least 90%, 92%, 94%, 96%, 98%, or 100% sequence identity) with one or more of the following: cecropin (SEQ ID NO: 82), melittin, copsin, drosomycin (SEQ ID NO: 93), dermcidin (SEQ ID NO: 81), andropin (SEQ ID NO: 83), moricin (SEQ ID NO: 84), ceratotoxin (SEQ ID NO: 85), abaecin (SEQ ID NO: 86), apidaecin (SEQ ID NO: 87), prophenin (SEQ ID NO: 88), indolicidin

(SEQ ID NO: 89), protegrin (SEQ ID NO: 90), tachyplesin (SEQ ID NO: 91), or defensin (SEQ ID NO: 92) formulated for targeting a microorganism in a vector (e.g., an insect vector) for an animal pathogen.

In some embodiments of the second aspect, the antimicrobial peptide may be at a concentration of about 0.1 ng/g to about 100 mg/g (about 0.1 ng/g to about 1 ng/g, about 1 ng/g to about 10 ng/g, about 10 ng/g to about 100 ng/g, about 100 ng/g to about 1000 ng/g, about 1 mg/g to about 10 mg/g, about 10 mg/g to about 100 mg/g) or about 0.1 ng/mL to about 100 mg/mL (about 0.1 ng/mL to about 1 ng/mL, about 1 ng/mL to about 10 ng/mL, about 10 ng/mL to about 100 ng/mL, about 100 ng/mL to about 1000 ng/mL, about 1 mg/mL to about 10 mg/mL, about 10 mg/mL to about 100 mg/mL) in the composition.

In some embodiments of the second aspect, the antimicrobial peptide may further include a targeting domain.

In some embodiments of the second aspect, the antimicrobial peptide may further include a cell penetrating peptide.

In yet another aspect, the composition includes an agent that alters a level, activity, or metabolism of one or more microorganisms resident in an insect host, the alteration resulting in a decrease in the insect host's fitness.

In some embodiments of any of the above compositions, the one or more microorganisms may be a bacterium or fungus resident in the host. In some embodiments, the bacterium resident in the host is at least one selected from the group consisting of *Candidatus spp*, *Buchenera spp*, *Blattabacterium spp*, *Baumania spp*, *Wigglesworthia spp*, *Wolbachia spp*, *Rickettsia spp*, *Orientia spp*, *Sodalis spp*, *Burkholderia spp*, *Cupriavidus spp*, *Frankia spp*, *Snirrhizobium spp*, *Streptococcus spp*, *Wolinella spp*, *Xylella spp*, *Erwinia spp*, *Agrobacterium spp*, *Bacillus spp*, *Paenibacillus spp*, *Streptomyces spp*, *Micrococcus spp*, *Corynebacterium spp*, *Acetobacter spp*, *Cyanobacteria spp*, *Salmonella spp*, *Rhodococcus spp*, *Pseudomonas spp*, *Lactobacillus spp*, *Enterococcus spp*, *Alcaligenes spp*, *Klebsiella spp*, *Paenibacillus spp*, *Arthrobacter spp*, *Corynebacterium spp*, *Brevibacterium spp*, *Thermus spp*, *Pseudomonas spp*, *Clostridium spp*, and *Escherichia spp*. In some embodiments, the fungus resident in the host is at least one selected from the group consisting of *Candida*, *Metschnikowia*, *Debaromyces*, *Starmerella*, *Pichia*, *Cryptococcus*, *Pseudozyma*, *Symbiotaphrina buchneri*, *Symbiotaphrina kochii*, *Scheffersomyces shehatae*, *Scheffersomyces stipites*, *Cryptococcus*, *Trichosporon*, *Amylostereum areolatum*, *Epichloe spp*, *Pichia pinus*, *Hansenula capsulate*, *Daldinia decipien*, *Ceratocytis spp*, *Ophiostoma spp*, and *Attamyces bromatificus*. In certain embodiments, the bacteria is a *Wolbachia spp*. (e.g., in a mosquito host). In certain emobdiments, the bacteria is a *Rickettsia spp*. (e.g., in a tick host).

In any of the above compositions, the agent, which hereinafter may also be referred to as a modulating agent, may alter the growth, division, viability, metabolism, and/or longevity of the microorganism resident in the host. In any of the above embodiments, the modulating agent may decrease the viability of the one or more microorganisms resident in the host. In some embodiments, the modulating agent increases growth or viability of the one or more microorganisms resident in the host.

In any of the above embodiments, the modulating agent is a phage, a polypeptide, a small molecule, an antibiotic, a bacterium, or any combination thereof.

In some embodiments, the phage binds a cell surface protein on a bacterium resident in the host. In some embodiments, the phage is virulent to a bacterium resident in the host. In some embodiments,

the phage is at least one selected from the group consisting of *Myoviridae*, *Siphoviridae*, *Podoviridae*, *Lipothrixviridae*, *Rudiviridae*, *Ampullaviridae*, *Bicaudaviridae*, *Clavaviridae*, *Corticoviridae*, *Cystoviridae*, *Fuselloviridae*, *Gluboloviridae*, *Guttaviridae*, *Inoviridae*, *Leviviridae*, *Microviridae*, *Plasmaviridae*, and *Tectiviridae*.

5 In some embodiments, the polypeptide is at least one of a bacteriocin, R-type bacteriocin, nodule C-rich peptide, antimicrobial peptide, lysin, or bacteriocyte regulatory peptide.

In some embodiments, the small molecule is a metabolite.

In some embodiments, the antibiotic is a broad-spectrum antibiotic.

10 In some embodiments, the modulating agent is a naturally occurring bacteria. In some embodiments, the bacteria is at least one selected from the group consisting of *Bartonella apis*, *Parasaccharibacter apium*, *Frischella perrara*, *Snodgrassella alvi*, *Gilliamella apicola*, *Bifidobacterium spp*, and *Lactobacillus spp*. In some embodiments, the bacterium is at least one selected from the group consisting of *Candidatus spp*, *Buchenera spp*, *Blattabacterium spp*, *Baumania spp*, *Wigglesworthia spp*, *Wolbachia spp*, *Rickettsia spp*, *Orientia spp*, *Sodalis spp*, *Burkholderia spp*, *Cupriavidus spp*, *Frankia*
15 *spp*, *Snirrhizobium spp*, *Streptococcus spp*, *Wolinella spp*, *Xylella spp*, *Erwinia spp*, *Agrobacterium spp*, *Bacillus spp*, *Paenibacillus spp*, *Streptomyces spp*, *Micrococcus spp*, *Corynebacterium spp*, *Acetobacter spp*, *Cyanobacteria spp*, *Salmonella spp*, *Rhodococcus spp*, *Pseudomonas spp*, *Lactobacillus spp*, *Enterococcus spp*, *Alcaligenes spp*, *Klebsiella spp*, *Paenibacillus spp*, *Arthrobacter spp*, *Corynebacterium spp*, *Brevibacterium spp*, *Thermus spp*, *Pseudomonas spp*, *Clostridium spp*, and *Escherichia spp*.

20 In any of the above compositions, host fitness may be measured by survival, reproduction, or metabolism of the host. In any of the above embodiments, the modulating agent may modulate the host's fitness by increasing pesticidal susceptibility of the host (e.g., susceptibility to a pesticide listed in Table 12). In some embodiments, the modulating agent modulates the host's fitness by increasing pesticidal susceptibility of the host. In some embodiments, the pesticidal susceptibility is bactericidal or fungicidal
25 susceptibility. In some embodiments, the pesticidal susceptibility is insecticidal susceptibility.

In any of the above compositions, the composition may include a plurality of different modulating agents. In some embodiments, the composition includes a modulating agent and a pesticidal agent (e.g., a pesticide listed in Table 12). In some embodiments, the pesticidal agent is a bactericidal or fungicidal agent. In some embodiments, the pesticidal agent is an insecticidal agent.

30 In any of the above compositions, modulating agent may be linked to a second moiety. In some embodiments, the second moiety is a modulating agent.

In any of the above compositions, the modulating agent may be linked to a targeting domain. In some embodiments, the targeting domain targets the modulating agent to a target site in the host. In some embodiments, the targeting domain targets the modulating agent to the one or more
35 microorganisms resident in the host.

In any of the above compositions, the modulating agent may include an inactivating pre- or pro-sequence, thereby forming a precursor modulating agent. In some embodiments, the precursor modulating agent is converted to an active form in the host.

40 In any of the above compositions, the modulating agent may include a linker. In some embodiments, the linker is a cleavable linker.

In any of the above compositions, the composition may further include a carrier. In some instances, the carrier may be an agriculturally acceptable carrier.

In any of the above compositions, the composition may further include a host bait, a sticky agent, or a combination thereof. In some embodiments, the host bait is a comestible agent and/or a
5 chemoattractant.

In any of the above compositions, the composition may be at a dose effective to modulate host fitness. I

In any of the above compositions, the composition may be formulated for delivery to a microorganism inhabiting the gut of the host. In any of the above compositions, the composition may be
10 formulated for delivery to a microorganism inhabiting a bacteriocyte of the host and/or the gut of the host. In some embodiments, the composition may be formulated for delivery to a plant. In some embodiments, the composition may be formulated for use in a host feeding station.

In any of the above compositions, the composition may be formulated as a liquid, a powder, granules, or nanoparticles. In some embodiments, the composition is formulated as one selected from
15 the group consisting of a liposome, polymer, bacteria secreting peptide, and synthetic nanocapsule. In some embodiments, the synthetic nanocapsule delivers the composition to a target site in the host. In some embodiments, the target site is the gut of the host. In some embodiments, the target site is a bacteriocyte in the host.

In a further aspect, also provided herein are hosts that include any of the above compositions. In
20 some embodiments, the host is an insect. In some embodiments, the insect is a mosquito, midge, louse, sandfly, tick, triatomine bug, tsetse fly, or flea. In certain embodiments, the insect is a mosquito. In certain embodiments, the insect is a tick. In certain embodiments, the insect is a mite. In certain embodiments, the insect is a louse.

Also provided herein is a system for modulating a host's fitness comprising a modulating agent
25 that targets a microorganism that is required for a host's fitness, wherein the system is effective to modulate the host's fitness, and wherein the host is an insect. The modulating agent may include any of the compositions described herein. In some embodiments, the modulating agent is formulated as a powder. In some embodiments, the modulating agent is formulated as a solvent. In some embodiments, the modulating agent is formulated as a concentrate. In some embodiments, the modulating agent is
30 formulated as a diluent. In some embodiments, the modulating agent is prepared for delivery by combining any of the previous compositions with a carrier.

In yet a further aspect, also provided herein are methods for modulating the fitness of an insect using any of the compositions described herein. In one instance, the method of modulating the fitness of an insect host includes delivering the composition of any one of the previous claims to the host, wherein
35 the modulating agent targets the one or more microorganisms resident in the host, and thereby modulates the host's fitness. In another instance, the method of modulating microbial diversity in an insect host includes delivering the composition of any one of the previous claims to the host, wherein the modulating agent targets the one or more microorganisms resident in the host, and thereby modulates microbial diversity in the host.

In some embodiments of any of the above methods, the modulating agent may alter the levels of the one or more microorganisms resident in the host. In some embodiments of any of the above methods, the modulating agent may alter the function of the one or more microorganisms resident in the host. In some embodiments, the one or more microorganisms may be a bacterium and/or fungus. In some embodiments, the one or more microorganisms are required for host fitness. In some
5 embodiments, the one or more microorganisms are required for host survival.

In some embodiments of any of the above methods, the delivering step may include providing the modulating agent at a dose and time sufficient to effect the one or more microorganisms, thereby modulating microbial diversity in the host. In some embodiments, the delivering step includes topical
10 application of any of the previous compositions to a plant. In some embodiments, the delivering step includes providing the modulating agent through a genetically engineered plant. In some embodiments, the delivering step includes providing the modulating agent to the host as a comestible. In some embodiments, the delivering step includes providing a host carrying the modulating agent. In some embodiments the host carrying the modulating agent can transmit the modulating agent to one or more
15 additional hosts.

In some embodiments of any of the above methods, the composition may be effective to increase the host's sensitivity to a pesticidal agent (e.g., a pesticide listed in Table 12). In some embodiments, the host is resistant to the pesticidal agent prior to delivery of the modulating agent. In some embodiments, the pesticidal agent is an allelochemical agent. In some embodiments, the allelochemical agent is
20 caffeine, soyacystatin N, monoterpenes, diterpene acids, or phenolic compounds. In some embodiments, the composition is effective to selectively kill the host. In some embodiments, the composition is effective to decrease host fitness. In some embodiments, the composition is effective to decrease the production of essential amino acids and/or vitamins in the host.

In some embodiments of any of the above methods, the host is an insect. In some embodiments, the host is a vector for an animal pathogen. In some embodiments, the vector is a mosquito, midge, louse, sandfly, tick, triatomine bug, tsetse fly, or flea. In certain embodiments, the vector is a mosquito. In certain embodiments, the vector is a tick. In certain embodiments, the vector is a mite. In certain
25 embodiments, the vector is a louse. In some embodiments, the animal pathogen is a virus, a protozoan, a bacterium, a protist, or a nematode. In some embodiments, the virus is one belonging to the group
30 *Togaviridae*, *Flaviviridae*, *Bunyaviridae*, *Rhabdoviridae*, or *Orbiviridae*. In some embodiments, the bacterium is one belonging to the genus *Yersinia*, *Francisella*, *Rickettsia*, *Orientia*, or *Borrelia*. In some embodiments, the protozoan is one belonging to the genus *Plasmodium*, *Trypanosoma*, *Leishmania*, or *Babesia*. In some embodiments, the nematode is one belonging to the genus *Brugia*. In some
35 embodiments, the composition is effective to prevent or decrease transmission of the pathogen to animals. In some embodiments, the composition is effective to prevent or decrease horizontal or vertical transmission of the pathogen between hosts. In some embodiments, the composition is effective to decrease host fitness, host development, or vectorial competence.

In another aspect, also provided herein are screening assays to identify modulating agent that modulate the fitness of a host. In one instance, the screening assay to identify a modulating agent that
40 modulates the fitness of a host, includes the steps of (a) exposing a microorganism that can be resident

in the host to one or more candidate modulating agents and (b) identifying a modulating agent that decreases the fitness of the host.

In some embodiments of the screening assay, the modulating agent is a microorganism resident in the host. In some embodiments, the microorganism is a bacterium. In some embodiments, the bacterium, when resident in the host, decreases host fitness. In some embodiments of the screening assay, the modulating agent affects an allelochemical-degrading microorganism. In some embodiments, the modulating agent is a phage, an antibiotic, or a test compound. In some embodiments, the antibiotic is timentin or azithromycin.

In some embodiments of the screening assay, the host may be an invertebrate. In some embodiments, the invertebrate is an insect. In some embodiments, the insect is a mosquito. In some embodiments, the insect is a tick. In certain embodiments, the insect is a mite. In certain embodiments, the insect is a louse.

In any of the above embodiments of the screening assay, host fitness may be modulated by modulating the host microbiota.

15

Definitions

As used herein, the term “animals” refers to livestock or farm animals and other mammalian veterinary animals.

As used herein, the term “bacteriocin” refers to a peptide or polypeptide that possesses antimicrobial properties. Naturally occurring bacteriocins are produced by certain prokaryotes and act against organisms related to the producer strain, but not against the producer strain itself. Bacteriocins contemplated herein include, but are not limited to, naturally occurring bacteriocins, such as bacteriocins produced by bacteria, and derivatives thereof, such as engineered bacteriocins, recombinantly expressed bacteriocins, and chemically synthesized bacteriocins. In some instances, the bacteriocin is a functionally active variant of the bacteriocins described herein. In some instances, the variant of the bacteriocin has at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity, e.g., over a specified region or over the entire sequence, to a sequence of a bacteriocin described herein or a naturally occurring bacteriocin.

As used herein, the term “bacteriocyte” refers to a specialized cell found in certain insects where intracellular bacteria reside with symbiotic bacterial properties.

As used herein, the term “effective amount” refers to an amount of a modulating agent (e.g., a phage, lysin, bacteriocin, small molecule, or antibiotic) or composition including said agent sufficient to effect the recited result, e.g., to decrease or reduce the fitness of a host organism (e.g., insect, e.g., mosquito, tick, mite, louse); to reach a target level (e.g., a predetermined or threshold level) of a modulating agent concentration inside a target host; to reach a target level (e.g., a predetermined or threshold level) of a modulating agent concentration inside a target host gut; to reach a target level (e.g., a predetermined or threshold level) of a modulating agent concentration inside a target host bacteriocyte; to modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host.

As used herein, the term “fitness” refers to the ability of a host organism to survive, and/or to produce surviving offspring. Fitness of an organism may be measured by one or more parameters, including, but not limited to, life span, reproductive rate, mobility, body weight, and metabolic rate. Fitness may additionally be measured based on measures of activity (e.g., biting animals) or disease transmission (e.g., vector-vector transmission or vector-animal transmission).

As used herein, the term “gut” refers to any portion of a host’s gut, including, the foregut, midgut, or hindgut of the host.

As used herein, the term “host” refers to an organism (e.g., insect, e.g., mosquito, louse, mite, or tick) carrying resident microorganisms (e.g., endogenous microorganisms, endosymbiotic microorganisms (e.g., primary or secondary endosymbionts), commensal organisms, and/or pathogenic microorganisms).

As used herein “decreasing host fitness” or “decreasing host fitness” refers to any disruption to host physiology, or any activity carried out by said host, as a consequence of administration of a modulating agent, including, but not limited to, any one or more of the following desired effects: (1) decreasing a population of a host by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more; (2) decreasing the reproductive rate of a host (e.g., insect, e.g., mosquito, tick, mite, louse) by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more; (3) decreasing the mobility of a host (e.g., insect, e.g., mosquito, tick, mite, louse) by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more; (4) decreasing the body weight of a host (e.g., insect, e.g., mosquito, tick, mite, louse) by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more; (5) increasing the metabolic rate or activity of a host (e.g., insect, e.g., mosquito, tick, mite, louse) by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more; (6) decreasing vector-vector pathogen transmission (e.g., vertical or horizontal transmission of a pathogen from one insect to another) by a host (e.g., insect, e.g., mosquito, tick, mite, louse) by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more; (7) decreasing vector-animal pathogen transmission (e.g., insect, e.g., mosquito, tick, mite, louse) by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more; (8) decreasing host (e.g., insect, e.g., mosquito, tick, mite, louse) lifespan by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more; (9) increasing host (e.g., insect, e.g., mosquito, tick, mite, louse) susceptibility to pesticides (e.g., insecticides) by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more; or (10) decreasing vector competence by a host (e.g., insect, e.g., mosquito, tick, mite, louse) by about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 100% or more. A decrease in host fitness can be determined in comparison to a host organism to which the modulating agent has not been administered.

The term “insect” includes any organism belonging to the phylum *Arthropoda* and to the class *Insecta* or the class *Arachnida*, in any stage of development, i.e., immature and adult insects.

As used herein, “lysin” also known as endolysin, autolysin, murein hydrolase, peptidoglycan hydrolase, or cell wall hydrolase refers to a hydrolytic enzyme that can lyse a bacterium by cleaving peptidoglycan in the cell wall of the bacterium. Lysins contemplated herein include, but are not limited to, naturally occurring lysins, such as lysins produced by phages, lysins produced by bacteria, and

derivatives thereof, such as engineered lysins, recombinantly expressed lysins, and chemically synthesized lysins. A functionally active variant of the bacteriocin may have at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity, e.g., over a specified region or over the
 5 entire sequence, to a sequence of a synthetic, recombinant, or naturally derived bacteriocin, including any described herein.

As used herein, the term “microorganism” refers to bacteria or fungi. Microorganisms may refer to microorganisms resident in a host organism (e.g., endogenous microorganisms, endosymbiotic microorganisms (e.g., primary or secondary endosymbionts)) or microorganisms exogenous to the host,
 10 including those that may act as modulating agents. As used herein, the term “target microorganism” refers to a microorganism that is resident in the host and impacted by a modulating agent, either directly or indirectly.

As used herein, the term “agent” or “modulating agent” refers to an agent that is capable of altering the levels and/or functioning of microorganisms resident in a host organism (e.g., insect, e.g.,
 15 mosquito, tick, mite, louse), and thereby modulate (e.g., decrease) the fitness of the host organism (e.g., insect, e.g., mosquito, tick, mite, louse).

As used herein, the term “pesticide” or “pesticidal agent” refers to a substance that can be used in the control of agricultural, environmental, or domestic/household pests, such as insects, fungi, bacteria, or viruses. The term “pesticide” is understood to encompass naturally occurring or synthetic insecticides
 20 (larvicides or adulticides), insect growth regulators, acaricides (miticides), nematocides, ectoparasitocides, bactericides, fungicides, or herbicides (substance which can be used in agriculture to control or modify plant growth). Further examples of pesticides or pesticidal agents are listed in Table 12. In some instances, the pesticide is an allelochemical. As used herein, “allelochemical” or “allelochemical agent” is a substance produced by an organism that can effect a physiological function (e.g., the germination,
 25 growth, survival, or reproduction) of another organism (e.g., a host insect).

As used herein, the term “peptide,” “protein,” or “polypeptide” encompasses any chain of naturally or non-naturally occurring amino acids (either D- or L-amino acids), regardless of length (e.g., at least 2,
 3, 4, 5, 6, 7, 10, 12, 14, 16, 18, 20, 25, 30, 40, 50, 100, or more amino acids), the presence or absence of post-translational modifications (e.g., glycosylation or phosphorylation), or the presence of, e.g., one or
 30 more non-amino acyl groups (for example, sugar, lipid, etc.) covalently linked to the peptide, and includes, for example, natural proteins, synthetic, or recombinant polypeptides and peptides, hybrid molecules, peptoids, and peptidomimetics.

As used herein, “percent identity” between two sequences is determined by the BLAST 2.0 algorithm, which is described in Altschul et al., (1990) J. Mol. Biol. 215:403-410. Software for performing
 35 BLAST analyses is publicly available through the National Center for Biotechnology Information.

As used herein, the term “bacteriophage” or “phage” refers to a virus that infects and replicates in bacteria. Bacteriophages replicate within bacteria following the injection of their genome into the cytoplasm and do so using either a lytic cycle, which results in bacterial cell lysis, or a lysogenic (non-
 40 lytic) cycle, which leaves the bacterial cell intact. The phage may be a naturally occurring phage isolate, or an engineered phage, including vectors, or nucleic acids that encode either a partial phage genome

(e.g., including at least all essential genes necessary to carry out the life cycle of the phage inside a host bacterium) or the full phage genome.

As used herein, the term "plant" refers to whole plants, plant organs, plant tissues, seeds, plant cells, seeds, and progeny of the same. Plant cells include, without limitation, cells from seeds,
 5 suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. Plant parts include differentiated and undifferentiated tissues including, but not limited to the following: roots, stems, shoots, leaves, pollen, seeds, tumor tissue, and various forms of cells and culture (e.g., single cells, protoplasts, embryos, and callus tissue). The plant tissue may be in a plant or in a plant organ, tissue, or cell culture. In addition, a plant may be genetically
 10 engineered to produce a heterologous protein or RNA, for example, of any of the modulating agents in the methods or compositions described herein.

As used herein, the term "vector" refers to an insect that can carry or transmit an animal pathogen from a reservoir to an animal. Exemplary vectors include insects, such as those with piercing-sucking mouthparts, as found in Hemiptera and some Hymenoptera and Diptera such as mosquitoes,
 15 bees, wasps, midges, lice, tsetse fly, fleas and ants, as well as members of the Arachnidae such as ticks and mites.

Other features and advantages of the invention will be apparent from the following Detailed Description and the Claims.

20 BRIEF DESCRIPTION OF THE DRAWINGS

The figures are meant to be illustrative of one or more features, aspects, or embodiments of the invention and are not intended to be limiting.

Fig. 1A-1G show images of different antibiotic delivery systems. First instar LSR-1 aphids were treated with different therapeutic solutions by delivery through plants (Fig. 1A), leaf coating (Fig. 1B),
 25 microinjection (Fig. 1C), and topical delivery (Fig. 1D).

Fig. 2A-2C show the delay in aphid development during rifampicin treatment in first instar LSR-1 aphids treated by delivery through plants with three different conditions: artificial diet without essential amino acids (AD only), artificial diet without essential amino acids with 100 µg/ml rifampicin (AD + Rif), and artificial diet with 100 µg/ml rifampicin and essential amino acids (AD + Rif + EAA). Fig. 2A is a
 30 series of graphs showing the percentage of living aphids at each developmental stage (sample size=33 aphids/group). Fig 2B shows representative images from each treatment taken at 12 days. Scale bars 2.5 mm. Fig 2C shows area measurements from aphid bodies showing the drastic effect of rifampicin treatment. Adding back essential amino acids partially rescues development defects.

Fig. 3 shows that rifampicin treatment resulted in aphid death. Survival was monitored daily for
 35 LSR-1 aphids treated by delivery through plants with artificial diet without essential amino acids (AD only), artificial diet without essential amino acids with 100 ug/ml rifampicin (AD + Rif), and artificial diet with 100 ug/ml rifampicin and (AD + Rif + EAA). Number in parentheses represents number of aphids in each group. Statistical significance was determined by Log-Rank Test and the following statistically significant differences were determined: AD only vs. AD + Rif, $p < 0.0001$ and AD + Rif vs. AD + Rif + EAA, $p = 0.017$.

Fig. 4 is a graph showing that rifampicin treatment resulted in loss of reproduction in aphids. First instar LSR-1 aphids were treated by delivery through plants with artificial diet without essential amino acids (AD only), artificial diet without essential amino acids with 100 ug/ml rifampicin (AD + Rif), and artificial diet with 100 ug/ml rifampicin and (AD + Rif + EAA) and the number of offspring produced each day after aphid reached adulthood was measured. Shown is the mean number of offspring produced per day after aphid reached adulthood $\pm$ S.D.

Fig. 5 is a graph showing that rifampicin treatment eliminated endosymbiotic *Buchnera*. Symbiont titer was determined for the different conditions at 7 days post-treatment. DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA $\pm$ SD of 3 aphids per group. Statistically significant differences were determined using a one-way-ANOVA followed by Tukey's Post-Test; *, $p < 0.05$.

Fig. 6A and 6B show that rifampicin treatment delivered through leaf coating delayed aphid development. First instar eNASCO aphids were treated by coating leaves with 100 μ l of two different solutions: solvent control (0.025% Silwet L-77), and 50 μ g/ml rifampicin. Fig. 6A is a series of graphs showing the developmental stage over time for each condition. Shown is the percentage of living aphids at each developmental stage (sample size=20 aphids/group). Fig. 6B is a graph showing area measurements from aphid bodies showing the drastic effect of rifampicin coated leaves on aphid size. Statistically significant differences were determined using a one-way-ANOVA followed by Tukey's Post-Test; *, $p < 0.05$.

Fig. 7 shows that rifampicin treatment delivered through leaf coating resulted in aphid death. Survival was monitored daily for eNASCO aphids treated by coating leaves with 100 μ l of two different solutions: solvent control (Silwet L-77), and 50 μ g/ml rifampicin. Treatment affects survival rate of aphids.

Fig. 8 shows that rifampicin treatment delivered through leaf coating eliminated endosymbiotic *Buchnera*. Symbiont titer was determined for the two conditions at 6 days post-treatment. DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA $\pm$ SD. Statistically significant differences were determined using a one-way-ANOVA followed by Tukey's Post-Test; *, $p < 0.05$.

Fig. 9 is a graph showing rifampicin treatment by microinjection eliminated endosymbiotic *Buchnera*. Symbiont titer was determined 4 days post-injection with the indicated conditions. Control sample is the solvent, 0.025% Silwet L-77 described before. DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA $\pm$ SD. Statistically significant differences were determined using a one-way-ANOVA followed by Tukey's Post-Test; *, $p < 0.05$.

Fig. 10 is a graph showing that rifampicin treatment delivered through topical treatment eliminated endosymbiotic *Buchnera*. Symbiont titer was determined 3 days post-spraying with: solvent (silwet L-77) or the rifampicin solution diluted in solvent. DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA $\pm$ SD. Statistically significant differences were determined using a one-way-ANOVA followed by Tukey's Post-Test; *, $p < 0.05$.

Fig. 11 shows a panel of graphs demonstrating that 1st and 2nd instar LSR-1 aphids were placed on leaves perfused with water plus food coloring or 50 µg/ml rifampicin in water plus food coloring. Developmental stage was measured over time for each condition. Shown is the percentage of living aphids at each developmental stage (sample size=74-81 aphids/group).

5 Fig. 12 shows a graph demonstrating survival of 1st and 2nd instar LSR-1 aphids placed on leaves perfused with water plus food coloring or 50 µg/ml rifampicin in water plus food coloring. Number in parentheses represents the number of aphids in each group. Statistical significance was determined by Log-Rank Test.

10 Fig. 13 shows a graph demonstrating symbiont titer determined 8 days post-treatment with leaves perfused with water and food coloring or rifampicin plus water and food coloring. DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA ± SD. Number in box indicates the median of the experimental group.

15 Fig. 14 shows a panel of graphs demonstrating 1st and 2nd instar LSR-1 aphids treated via leaf injection and through the plant with water plus food coloring or 100 µg/ml rifampicin in water plus food coloring. Developmental stage was measured over time for each condition. Shown is the percentage of living aphids at each developmental stage (sample size=49-50 aphids/group).

20 Fig. 15 is a graph demonstrating survival of 1st and 2nd instar LSR-1 aphids placed on leaves perfused and treated with water plus food coloring or 100 µg/ml rifampicin in water plus food coloring. Number in parentheses represents the number of aphids in each group. A Log-Rank Test was performed and determined that there were no statistically significant differences between groups.

25 Fig. 16A and 16B are graphs showing symbiont titer determined 6 (16A) and 8 (16B) days post-treatment in aphids feeding on leaves perfused and treated with water and food coloring or rifampicin plus water and food coloring. DNA was extracted from aphids and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA ± SD. Number in box indicates the median of the experimental group.

30 Fig. 17 is a panel of graphs showing that 1st and 2nd instar LSR-1 aphids were treated with control solutions (water and Silwet L-77) or a combination of treatments with 100 µg/ml rifampicin. Developmental stage was measured over time for each condition. Shown is the percentage of living aphids at each developmental stage (sample size=76-80 aphids/group).

Fig. 18 is a graph showing 1st and 2nd instar LSR-1 aphids were treated with control solutions of a combination of treatments containing rifampicin. Number in parentheses represents the number of aphids in each group. A Log-Rank Test was performed and determined that there were no statistically significant differences between groups.

35 Fig. 19 is a graph showing symbiont titer determined at 7 days post-treatment with control or rifampicin solutions. DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA ± SD. Number in box indicates the median of the experimental group. Statistically significant differences were determined by t-test.

Fig. 20 is an image showing the chitosan delivery system. *A. pisum* aphids were treated with a therapeutic solution by delivery through leaf perfusion and through the plants as shown.

Fig. 21 is a panel of graphs showing that chitosan treatment resulted in delayed aphid development. First and second instar *A. pisum* aphids were treated by delivery through plants and leaf perfusion with the control solution (Water), and 300 ug/ml chitosan in water. Developmental stage was monitored throughout the experiment. Shown are the percent of aphids at each developmental stage (1st instar, 2nd instar, 3rd instar, 4th instar, 5th instar, or 5R which represents a reproducing 5th instar) per treatment group.

Fig. 22 is a graph showing there was a decrease in insect survival upon treatment with chitosan. First and second instar *A. pisum* aphids were treated by delivery through plants and leaf perfusion with just water or chitosan solution and survival was monitored daily over the course of the experiment. Number in parentheses represents the total number of aphids in the treatment group.

Fig. 23 is a graph showing treatment with chitosan reduced endosymbiotic *Buchnera*. First and second instar *A. pisum* aphids were treated by delivery through plants and leaf perfusion with water or 300 ug/ml chitosan in water. At 8 days post-treatment, DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA $\pm$ SD of 6 aphids/group. The median value for each group is shown in box.

Fig. 24 is a panel of graphs showing treatment with nisin resulted in delayed aphid development. First and second instar LSR-2 *A. pisum* aphids were treated with water (control) or 1.6 or 7 mg/ml nisin via delivery by leaf injection and through the plant and development was measured over time. Shown are the percent of aphids at each life stage (1st, 2nd, 3rd, 4th, 5th, and 5R (reproducing 5th) instar) at the indicated time point. N=56-59 aphids/group.

Fig. 25 is a graph showing there was a dose dependent decrease in insect survival upon treatment with nisin. First and second instar LSR-1 *A. pisum* aphids were treated with water (control) or 1.6 or 7 mg/ml nisin via delivery by leaf injection and through the plant and survival was monitored over time. Number in parentheses indicates the number of aphids/group. Statistically significant differences were determined by Log Rank (Mantel-Cox) test.

Fig. 26 is a graph showing treatment with nisin reduced endosymbiotic *Buchnera*. First and second instar LSR-1 *A. pisum* aphids were treated with water (control) or 1.6 mg/ml nisin via delivery by leaf injection and through the plant and DNA was extracted from select aphids at eight days post-treatment and used for qPCR to determine *Buchnera* copy numbers. Shown are the mean *Buchnera*/aphid ratios for each treatment $\pm$ SEM. Number in the box above each experimental group indicates the median value for that group. Each data point represents a single aphid.

Fig. 27 is a panel of graphs showing treatment with levulinic acid resulted in delayed aphid development. First and second instar eNASCO *A. pisum* aphids were treated with water (control) or 0.03 or 0.3% levulinic acid via delivery by leaf injection and through the plant and development was measured over time. Shown are the percent of aphids at each life stage (1st, 2nd, 3rd, 4th, and 5th instar) at the indicated time point. N=57-59 aphids/group.

Fig. 28 is a graph showing there was a decrease in insect survival upon treatment with levulinic acid. First and second instar eNASCO *A. pisum* aphids were treated with water (control) or 0.03 or 0.3%

levulinic acid via delivery by leaf injection and through the plant and survival was monitored over time. N=57-59 aphids/group. Statistically significant differences were determined by Log Rank (Mantel-Cox) test; **, $p < 0.01$.

Fig. 29 is a panel of graphs showing treatment with levulinic acid reduced endosymbiotic *Buchnera*. First and second instar eNASCO *A. pisum* aphids were treated with water (control) or 0.03 or 0.3% levulinic acid via delivery by leaf injection and through the plant and DNA was extracted from select aphids at seven and eleven days post-treatment and used for qPCR to determine *Buchnera* copy numbers. Shown are the mean *Buchnera*/aphid ratios for each treatment +/- SEM. Statistically significant differences were determined by One-way ANOVA and Dunnett's Multiple Comparison Test; *, $p < 0.05$. Each data point represents a single aphid.

Fig. 30A and 30B show graphs demonstrating that gossypol treatment resulted in delayed aphid development. First and second instar *A. pisum* aphids were treated by delivery through plants with artificial diet without essential amino acids (AD only), and artificial diet without essential amino acids with different concentrations of gossypol (0.05%, 0.25% and 0.5%). Developmental stage was monitored throughout the experiment. Fig. 30A is a series of graphs showing the mean number of aphids at each developmental stage (1st instar, 2nd instar, 3rd instar, 4th instar, 5th instar, or 5R which represents a reproducing 5th instar) per treatment group. At the indicated time, aphids were imaged and their size was determined using Image J. Fig. 30B is a graph showing the mean aphid area $\pm$ SD of artificial diet treated (Control) or gossypol treated aphids. Statistical significance was determined using a One-Way ANOVA followed by Tukey's post-test. *, $p < 0.05$. **, $p < 0.01$.

Fig. 31 is a graph showing a dose-dependent decrease in survival of aphids upon treatment with the allelochemical gossypol. First and second instar *A. pisum* aphids were treated by delivery through plants with artificial diet without essential amino acids (AD no EAA), artificial diet without essential amino acids with 0.5% gossypol acetic acid (0.5% gossypol), artificial diet without essential amino acids with 0.25% gossypol acetic acid (0.25% gossypol), and artificial diet without essential amino acids and 0.05% gossypol acetic acid (0.05% gossypol) and survival was monitored daily over the course of the experiment. Number in parentheses represents the essential amino acids number of aphids in each group. Statistically significant differences were determined by Log-Rank test and AD no EAA and 0.5% gossypol are significantly different, $p = 0.0002$.

Fig. 32A and 32B are two graphs showing that treatment with 0.25% gossypol resulted in decreased fecundity. First and second instar *A. pisum* aphids were treated by delivery through plants with artificial diet without essential amino acids (AD5-2 no EAA), or artificial diet without essential amino acids with 0.25% gossypol acetic acid (AD5-2 no EAA + 0.25% gossypol), and fecundity was determined throughout the time course of the experiment. Fig. 32A shows the mean day $\pm$ SD at which aphids began producing offspring was measured and gossypol treatment delayed production of offspring. Fig. 32B shows the mean number of offspring produced after the aphid began a reproducing adult $\pm$ SD was measured and gossypol treatment results in decreased number of offspring produced. Each data point represents one aphid.

Fig. 33 is a graph showing that treatment with different concentrations of gossypol reduced endosymbiotic *Buchnera*. First and second instar *A. pisum* aphids were treated by delivery through

plants with artificial diet without essential amino acids (Control)) or artificial diet without essential amino acids with 0.5%, 0.25%, or 0.05% gossypol. At 5 or 13 days post-treatment, DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA $\pm$ SD of 2-6 aphids/group. Statistically significant differences were determined by Unpaired T-test; *, $p < 0.05$.

Fig. 34 is a graph showing that microinjection of gossypol resulted in decreased *Buchnera* levels in aphids. *A. pisum* LSR-1 aphids <3rd instar stage (nymphs) were injected with 20 nl of artificial diet without essential amino acids (AD) or artificial diet without essential amino acids with 0.05% gossypol (gossypol (0.05%)). Three days after injection, DNA was extracted from aphids and *Buchnera* levels were assessed by qPCR. Shown are the mean ratios of *Buchnera*/aphid DNA $\pm$ SD. Each data point represents one aphid.

Fig. 35 is a panel of graphs showing Trans-cinnamaldehyde treatment resulted in delayed aphid development. First and second instar *A. pisum* aphids were treated by delivery through plants with water and water with different concentrations of trans-cinnamaldehyde (TC, 0.05%, 0.5%, and 5%). Developmental stage was monitored throughout the experiment. Shown are the mean number of aphids at each developmental stage (1st instar, 2nd instar, 3rd instar, 4th instar, 5th instar, or 5R which represents a reproducing 5th instar) per treatment group. N=40-49 aphids/experimental group.

Fig. 36 is a graph showing there was a dose-dependent decrease in survival upon treatment the natural antimicrobial trans-cinnamaldehyde. First and second instar *A. pisum* aphids were treated by delivery through plants with water and water with different concentrations of trans-cinnamaldehyde (TC, 0.05%, 0.5%, and 5%). Survival was monitored throughout the course of the treatment. Statistically significant differences were determined by Log-Rank test. N=40-49 aphids/group.

Fig. 37 is a graph showing treatment with different concentrations of trans-cinnamaldehyde reduced endosymbiotic *Buchnera*. First and second instar *A. pisum* aphids were treated by delivery through plants with water and water with different concentrations of trans-cinnamaldehyde (0.05%, 0.5%, and 5%). At 3 days post-treatment, DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA $\pm$ SD of 2-11 aphids/group. The median of each treatment group is shown in the box above the data points. Statistically significant differences were determined by Unpaired T-test; *, $p < 0.05$. There was a statistically significant difference between the water control and the 0.5% trans-cinnamaldehyde group.

Fig. 38 is a panel of graphs showing treatment with scorpion peptide Uy192 resulted in delayed aphid development. First and second instar *A. pisum* aphids were treated by delivery through plants and leaf perfusion with the control solution (water), and 100 ug/ml Uy192 in water. a) developmental stage was monitored throughout the experiment. Shown are the percent of aphids at each developmental stage (1st instar, 2nd instar, 3rd instar, 4th instar, 5th instar, or 5R which represents a reproducing 5th instar) per treatment group.

Fig. 39 is a graph showing there was a decrease in insect survival upon treatment with the scorpion AMP Uy192. First and second instar *A. pisum* aphids were treated by delivery through plants and leaf perfusion with just water or Uy192 solution and survival was monitored daily over the course of the experiment. Number in parentheses represents the total number of aphids in the treatment group.

Fig. 40 is a graph showing treatment with Uy192 reduced endosymbiotic *Buchnera*. First and second instar *A. pisum* aphids were treated by delivery through plants and leaf perfusion with water or 100 µg/ml Uy192 in water, at 8 days post-treatment, DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA $\pm$ SD of 2-6 aphids/group. The median value for each group is shown in box.

Fig. 41 is a graph showing a decrease in survival in aphids microinjected with scorpion peptides D10 and D3. LSR-1 *A. pisum* aphids were microinjected with water (control) or with 100 ng of either scorpion peptide D3 or D10. After injection, aphids were released to fava bean leaves and survival was monitored throughout the course of the experiment. The number in parentheses indicates the number of aphids in each experimental treatment group.

Fig. 42 is a graph showing a decrease in endosymbiont titers upon injection with scorpion peptides D3 and D10. LSR-1 *A. pisum* aphids were microinjected with water (control) or with 100 ng of either scorpion peptide D3 or D10. After injection, aphids were released to fava bean leaves and at 5 days post-treatment, DNA was extracted from the remaining living aphids and qPCR was performed to determine the ratio of *Buchnera*/aphid DNA. Shown are the mean $\pm$ SD of each treatment group. N=2-9 aphids/group. The number above each treatment group in the box represents the median of the dataset.

Fig. 43 is a graph showing a decrease in insect survival upon treatment with a cocktail of scorpion AMPs. First and second instar eNASCO aphids were treated by delivery through leaf perfusion and through plants with a cocktail of scorpion peptides (40 µg/ml of each of Uy17, D3, UyCt3, and D10) and survival was monitored over the course of the experiment. The number in parentheses represents the number of aphids in each treatment group.

Fig. 44 is a panel of graphs showing treatment with scorpion peptide fused to a cell penetrating peptide resulted in delayed aphid development. First instar LSR-2 *A. pisum* aphids were treated with water (control) or 100 µg/ml Uy192+CPP+FAM via delivery by leaf injection and through the plant and development was measured over time. Shown are the percent of aphids at each life stage (1st, 2nd, 3rd, 4th, 5th, and 5R (reproducing 5th) instar) at the indicated time point. N=90 aphids/group.

Fig. 45 is a graph showing treatment of aphids with a scorpion peptide fused to a cell penetrating peptide increased mortality. First instar LSR-1 *A. pisum* aphids were treated with water or 100 µg/ml Uy192+CPP+FAM (peptide) in water delivered by leaf injection and through the plant. Survival was monitored over time. The number in parentheses indicates the number of aphids/group. Statistically significant differences were determined by Log Rank (Mantel-Cox) test and there is a significant difference between the two experimental groups ($p=0.0036$).

Fig. 46 is a graph showing treatment with Uy192+CPP+FAM reduced endosymbiotic *Buchnera*. First instar LSR-1 *A. pisum* aphids were treated with water or 100 µg/ml Uy192+CPP+FAM (peptide) in water delivered by leaf injection and through the plant. DNA was extracted from select aphids at five days post-treatment and used for qPCR to determine *Buchnera* copy numbers. Shown are the mean *Buchnera*/aphid ratios for each treatment $\pm$ SEM. Number in the box above each experimental group indicates the median value for that group. Each data point represents a single aphid. Statistically significant differences were determined by Student's T-test; ****, $p<0.0001$.

Fig. 47 is a panel of images showing Uy192+CPP+FAM penetrated bacteriocyte membranes. Bacteriocytes were dissected from the aphids and incubated with 250ug/ml of the Uy192+CPP+FAM peptide for 30 min. Upon washing and imaging, the Uy192+CPP+FAM can be seen at high quantities inside the bacteriocytes.

5 Fig. 48A and Fig. 48B are a panel of graphs showing Pantothenol treatment delayed aphid development. First instar and second eNASCO aphids were treated by delivery through plants with three different conditions: artificial diet without essential amino acids (AD no EAA), artificial diet without essential amino acids with 10 uM pantothenol (10 uM pantothenol), and artificial diet without essential amino acids with 100 uM pantothenol (100 uM pantothenol), artificial diet without essential amino acids with 100 uM pantothenol, and artificial diet without essential amino acids with 10 uM pantothenol. Fig. 48A shows developmental stage monitored over time for each condition. Fig. 48B shows relative area measurements from aphid bodies showing the drastic effect of pantothenol treatment.

Fig. 49 is a graph showing that treatment with pantothenol increased aphid mortality. Survival was monitored daily for eNASCO aphids treated by delivery through plants with artificial diet without essential amino acids, or artificial diet without essential amino acids containing 10 or 100 uM pantothenol. Number in parentheses represents number of aphids in each group.

Fig. 50A, 50B, and 50C are a panel of graphs showing Pantothenol treatment resulted in loss of reproduction. First and second instar eNASCO aphids were treated by delivery through plants with artificial diet without essential amino acids or with artificial diet without essential amino acids with 10 or 100 uM pantothenol. Fig. 50A shows the fraction of aphids surviving to maturity and reproducing. Fig. 50B shows the mean day aphids in each group began reproducing. Shown is the mean day an aphid began reproducing $\pm$ SD. Fig. 50C shows the mean number of offspring produced per day after an aphid began reproducing. Shown are the mean number of offspring/day $\pm$ SD.

Fig. 51 is a graph showing Pantothenol treatment did not affect endosymbiotic *Buchnera*. Symbiont titer was determined for the different conditions at 8 days post-treatment. DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown is the mean ratio of *Buchnera* DNA to aphid DNA $\pm$ SD of 6 aphids per group.

Fig. 52 is a panel of graphs showing Pantothenol treatment delivered through plants did not affect aphid development. First instar eNASCO aphids were treated by coating leaves with 100 μ l of two different solutions: solvent control (0.025% Silwet L-77), and 10 uM pantothenol and the developmental stage was measured over time for each condition. Shown is the percentage of living aphids at each developmental stage (sample size=20 aphids/group).

Fig. 53 is a graph showing Pantothenol treatment delivered through leaf coating resulted in aphid death. Survival was monitored daily for eNASCO aphids treated by coating leaves with 100 μ l of two different solutions: solvent control (Silwet L-77), and 10 uM pantothenol. Treatment affects survival rate of aphids. Sample size = 20 aphids/group. Log-Rank Mantel Cox test was used to determine whether there were statistically significant differences between groups and identified that the two group are significantly different ($p=0.0019$).

Fig. 54A and 54B are a panel of graphs showing treatment with a cocktail of amino acid analogs delayed aphid development. First instar LSR-1 aphids were treated by delivery through leaf perfusion

and through plants with water or a cocktail of amino acid analogs in water (AA cocktail). Fig. 54A shows the developmental stage measured over time for each condition. Shown are the percentage of living aphids at each developmental stage. Fig. 54B shows the area measurements from aphid bodies showing the drastic effect of treatment with an amino acid analog cocktail (AA cocktail). Statistically significant differences were determined using a Student's T-test; ****, $p < 0.0001$.

Fig. 55 is a graph showing treatment with a cocktail of amino acid analogs eliminated endosymbiotic *Buchnera*. Symbiont titer was determined for the different conditions at 6 days post-treatment. DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown are the mean ratios of *Buchnera* DNA to aphid DNA $\pm$ SD of 19-20 aphids per group. Each data point represents an individual aphid. Statistically significant differences were determined using a Student's T-test; *, $p < 0.05$.

Fig. 56A and 56B is a panel of graphs showing treatment with a combination of three agents delayed aphid development. First instar LSR-1 aphids were treated by delivery through leaf perfusion and through plants with water or a combination of three agents in water (Pep-Rif-Chitosan). Fig. 56A shows the developmental stage measured over time for each condition. Shown are the percentage of living aphids at each developmental stage. Fig. 56B shows the area measurements from aphid bodies showing the drastic effect of treatment with a combination of three treatments (Pep-Rif-Chitosan). Statistically significant differences were determined using a Student's T-test; ****, $p < 0.0001$.

Fig. 57 is a graph showing treatment with a combination of a peptide, antibiotic, and natural antimicrobial agent increased aphid mortality. LSR-1 aphids were treated with water or a combination of three treatments (Pep-Rif-Chitosan) and survival was monitored daily after treatment.

Fig. 58 is a graph showing treatment with a combination of a peptide, antibiotic, and natural antimicrobial agent eliminated endosymbiotic *Buchnera*. Symbiont titer was determined for the different conditions at 6 days post-treatment. DNA from aphids was extracted and qPCR was performed to determine the ratio of *Buchnera* DNA to aphid DNA. Shown are the mean ratios of *Buchnera* DNA to aphid DNA $\pm$ SD of 20-21 aphids per group. Each data point represents an individual aphid.

Fig. 59A and 59B are a panel of images showing ciprofloxacin coated and penetrated corn kernels. Corn kernels were soaked in water (no antibiotic) or the indicated concentration of ciprofloxacin in water and whole kernels or kernel were tested to see whether they can inhibit the growth of *E. coli* DH5 α . Fig. 59A shows bacterial growth in the presence of a corn kernel soaked in water without antibiotics and Fig. 59B shows the inhibition of bacterial growth when whole or half corn kernels that have been soaked in antibiotics are placed on a plate spread with *E. coli*.

Fig. 60 is a graph showing that adult *S. zeamais* weevils were treated with ciprofloxacin (250 μ g/ml or 2.5 mg/ml) or mock treated with water. After 18 days of treatment, genomic DNA was isolated from weevils and the amount of *Sitophilus* primary endosymbiont was determined by qPCR. Shown is the mean $\pm$ SEM of each group. Each data point represents one weevil. The median of each group is listed above the dataset.

Fig. 61A and 61B are graphs showing weevil development after treatment with ciprofloxacin. Fig. 61A shows individual corn kernels cut open 25 days after adults were removed from one replicate each of the initial corn kernels soaked/coated with water (control) or ciprofloxacin (250 μ g/ml or 2.5 mg/ml) and

examined for the presence of larvae, pupae, or almost fully developed (adult) weevils. Shown is the percent of each life stage found in kernels from each treatment group. The total number of offspring found in the kernels from each treatment group is indicated above each dataset. Fig. 61B shows genomic DNA isolated from offspring dissected from corn kernels from the control (water) and 2.5 mg/ml ciprofloxacin treatment groups and qPCR was done to measure the amount of *Sitophilus* primary endosymbiont present. Shown are the mean $\pm$ SD for each group. Statistically significant differences were determined by unpaired t-test; ***, $p \leq 0.001$.

Fig. 62A and 62B are graphs showing the two remaining replicates of corn kernels mock treated (water) or treated with 250 μ g/ml or 2.5 mg/ml ciprofloxacin monitored for the emergence of offspring after mating pairs were removed (at 7 days post-treatment). Fig. 62A shows the mean number of newly emerged weevils over time $\pm$ SD for each treatment group. Fig. 62B shows the mean number $\pm$ SEM of emerged weevils for each treatment group at 43 days after mating pairs were removed.

Fig. 63 is a panel of graphs showing rifampicin and doxycycline treatment resulted in mite mortality. Survival was monitored daily for untreated two-spotted spider mites and mites treated with 250 μ g/ml rifampicin and 500 μ g/ml doxycycline in 0.025% Silwet L-77.

Fig. 64 is a panel of graphs showing the results of a Seahorse flux assay for bacterial respiration. Bacteria were grown to logarithmic phase and loaded into Seahorse XFe96 plates for temporal measurements of oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) as described in methods. Treatments were injected into the wells after approximately 20 minutes and bacteria were monitored to detect changes in growth. Rifampicin = 100 μ g/mL; Chloramphenicol = 25 μ g/mL; Phages (T7 for *E. coli* and Φ SmVL-C1 for *S. marcescens*) were lysates diluted either 1:2 or 1:100 in SM Buffer. The markers on each line are solely provided as indicators of the condition to which each line corresponds, and are not indicative of data points.

Fig. 65 is a graph showing phage against *S. marcescens* reduced fly mortality. Flies that were pricked with *S. marcescens* were all dead within a day, whereas a sizeable portion of the flies that were pricked with both *S. marcescens* and the phage survived for five days after the treatment. Almost all of the control flies which were not treated in anyway survived till the end of the experiment. Log-rank test was used to compare the curves for statistical significance, asterisk denotes $p < 0.0001$.

DETAILED DESCRIPTION

Provided herein are methods and compositions useful for animal health, e.g., for altering a level, activity, or metabolism of one or more microorganisms resident in a host insect (e.g., arthropod, e.g., insect, e.g., an animal pathogen vector, e.g., mosquito, mite, louse, or tick), the alteration resulting in a decrease in the fitness of the host. The invention features a composition that includes a modulating agent (e.g., phage, peptide, small molecule, antibiotic, or combinations thereof) that can alter the host's microbiota in a manner that is detrimental to the host. By disrupting microbial levels, microbial activity, microbial metabolism, or microbial diversity, the modulating agent described herein may be used to

decrease the fitness of a variety of insects that carry vector-borne pathogens that cause disease in animals.

The methods and compositions described herein are based in part on the examples provided herein, which illustrate how modulating agents, for example antibiotics (e.g., oxytetracycline, doxycycline, or a combination thereof) can be used to target symbiotic microorganisms in a host (e.g., endosymbionts in insect vectors of animal pathogens, e.g., endosymbiotic *Wolbachia* in mosquitos or *Rickettsia* in ticks) to decrease the fitness of the host by altering the level, activity, or metabolism of the microorganisms within the hosts. Oxytetracycline and doxycycline are representative examples of antibiotics useful for this purpose. On this basis the present disclosure describes a variety of different approaches for the use of agents that alter a level, activity, or metabolism of one or more microorganisms resident in a host (e.g., a vector of an animal pathogen, e.g., a mosquito, mite, louse or a tick) the alteration resulting in a decrease in the host's fitness.

I. Hosts

i. Hosts

The methods and compositions provided herein may be used with any insect host that is considered a vector for a pathogen that is capable of causing disease in animals.

For example, the insect host may include, but is not limited to those with piercing-sucking mouthparts, as found in Hemiptera and some Hymenoptera and Diptera such as mosquitoes, bees, wasps, midges, lice, tsetse fly, fleas and ants, as well as members of the Arachnidae such as ticks and mites; order, class or family of Acarina (ticks and mites) e.g. representatives of the families Argasidae, Dermanyssidae, Ixodidae, Psoroptidae or Sarcoptidae and representatives of the species *Amblyomma* spp., *Anocenton* spp., *Argas* spp., *Boophilus* spp., *Cheyletiella* spp., *Chorioptes* spp., *Demodex* spp., *Dermacentor* spp., *Dermanyssus* spp., *Haemophysalis* spp., *Hyalomma* spp., *Ixodes* spp., *Lynxacarus* spp., *Mesostigmata* spp., *Notoednes* spp., *Ornithodoros* spp., *Ornithonyssus* spp., *Otobius* spp., *otodectes* spp., *Pneumonyssus* spp., *Psoroptes* spp., *Rhipicephalus* spp., *Sarcoptes* spp., or *Trombicula* spp.; Anoplura (sucking and biting lice) e.g. representatives of the species *Bovicola* spp., *Haematopinus* spp., *Linognathus* spp., *Menopon* spp., *Pediculus* spp., *Pemphigus* spp., *Phylloxera* spp., or *Solenopotes* spp.; Diptera (flies) e.g. representatives of the species *Aedes* spp., *Anopheles* spp., *Calliphora* spp., *Chrysomyia* spp., *Chrysops* spp., *Cochliomyia* spp., *Cw/ex* spp., *Culicoides* spp., *Cuterebra* spp., *Dermatobia* spp., *Gastrophilus* spp., *Glossina* spp., *Haematobia* spp., *Haematopota* spp., *Hippobosca* spp., *Hypoderma* spp., *Lucilia* spp., *Lyperosia* spp., *Melophagus* spp., *Oestrus* spp., *Phaenicia* spp., *Phlebotomus* spp., *Phormia* spp., Acari (sarcoptic mange) e.g., *Sarcoptidae* spp., *Sarcophaga* spp., *Simulium* spp., *Stomoxys* spp., *Tabanus* spp., *Tannia* spp. or *Zzpu/alpha* spp.; Mallophaga (biting lice) e.g. representatives of the species *Damalina* spp., *Felicola* spp., *Heterodoxus* spp. or *Trichodectes* spp.; or Siphonaptera (wingless insects) e.g. representatives of the species *Ceratophyllus* spp., *Xenopsylla* spp.; Cimicidae (true bugs) e.g. representatives of the species *Cimex* spp., *Tritominae* spp., *Rhodnius* spp., or *Triatoma* spp.

In some instances, the insect is a blood-sucking insect from the order Diptera (e.g., suborder Nematocera, e.g., family Colicidae). In some instances, the insect is from the subfamilies Culicinae,

Corethrinae, Ceratopogonidae, or Simuliidae. In some instances, the insect is of a *Culex* spp., *Theobaldia* spp., *Aedes* spp., *Anopheles* spp., *Aedes* spp., *Forcipomyia* spp., *Culicoides* spp., or *Helea* spp.

In certain instances, the insect is a mosquito. In certain instances, the insect is a tick. In certain instances, the insect is a mite. In certain instances, the insect is a biting louse.

ii. *Host Fitness*

The methods and compositions provided herein may be used to decrease the fitness of any of the hosts described herein. The decrease in fitness may arise from any alterations in microorganisms resident in the host, wherein the alterations are a consequence of administration of a modulating agent and have detrimental effects on the host.

In some instances, the decrease in host fitness may manifest as a deterioration or decline in the physiology of the host (e.g., reduced health or survival) as a consequence of administration of a modulating agent. In some instances, the fitness of an organism may be measured by one or more parameters, including, but not limited to, reproductive rate, lifespan, mobility, fecundity, body weight, metabolic rate or activity, or survival in comparison to a host organism to which the modulating agent has not been administered. For example, the methods or compositions provided herein may be effective to decrease the overall health of the host or to decrease the overall survival of the host. In some instances, the decreased survival of the host is about 2%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or greater than 100% greater relative to a reference level (e.g., a level found in a host that does not receive a modulating agent). In some instances, the methods and compositions are effective to decrease host reproduction (e.g., reproductive rate) in comparison to a host organism to which the modulating agent has not been administered. In some instances, the methods and compositions are effective to decrease other physiological parameters, such as mobility, body weight, life span, fecundity, or metabolic rate, by about 2%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or greater than 100% relative to a reference level (e.g., a level found in a host that does not receive a modulating agent).

In some instances, the decrease in host fitness may manifest as a decrease in the production of one or more nutrients in the host (e.g., vitamins, carbohydrates, amino acids, or polypeptides). In some instances, the methods or compositions provided herein may be effective to decrease the production of nutrients in the host (e.g., vitamins, carbohydrates, amino acids, or polypeptides) by about 2%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or greater than 100% relative to a reference level (e.g., a level found in a host that does not receive a modulating agent). In some instances, the methods or compositions provided herein may decrease nutrients in the host by decreasing the production of nutrients by one or more microorganisms (e.g., endosymbiont) in the host in comparison to a host organism to which the modulating agent has not been administered.

In some instances, the decrease in host fitness may manifest as an increase in the host's sensitivity to a pesticidal agent (e.g., a pesticide listed in Table 12) and/or a decrease in the host's resistance to a pesticidal agent (e.g., a pesticide listed in Table 12) in comparison to a host organism to which the modulating agent has not been administered. In some instances, the methods or compositions

provided herein may be effective to increase the host's sensitivity to a pesticidal agent (e.g., a pesticide listed in Table 12) by about 2%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or greater than 100% relative to a reference level (e.g., a level found in a host that does not receive a modulating agent). The pesticidal agent may be any pesticidal agent known in the art, including insecticidal agents. In some instances, the methods or compositions provided herein may increase the host's sensitivity to a pesticidal agent (e.g., a pesticide listed in Table 12) by decreasing the host's ability to metabolize or degrade the pesticidal agent into usable substrates in comparison to a host organism to which the modulating agent has not been administered.

In some instances, the decrease in host fitness may manifest as an increase in the host's sensitivity to an allelochemical agent and/or a decrease in the host's resistance to an allelochemical agent in comparison to a host organism to which the modulating agent has not been administered. In some instances, the methods or compositions provided herein may be effective to decrease the host's resistance to an allelochemical agent by about 2%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or greater than 100% relative to a reference level (e.g., a level found in a host that does not receive a modulating agent). In some instances, the allelochemical agent is caffeine, soyacystatin N, monoterpenes, diterpene acids, or phenolic compounds. In some instances, the methods or compositions provided herein may increase the host's sensitivity to an allelochemical agent by decreasing the host's ability to metabolize or degrade the allelochemical agent into usable substrates in comparison to a host organism to which the modulating agent has not been administered.

In some instances, the methods or compositions provided herein may be effective to decrease the host's resistance to parasites or pathogens (e.g., fungal, bacterial, or viral pathogens or parasites) in comparison to a host organism to which the modulating agent has not been administered. In some instances, the methods or compositions provided herein may be effective to decrease the host's resistance to a pathogen or parasite (e.g., fungal, bacterial, or viral pathogens; or parasitic mites) by about 2%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or greater than 100% relative to a reference level (e.g., a level found in a host that does not receive a modulating agent).

In some instances, the decrease in host fitness may manifest as other fitness disadvantages, such as decreased tolerance to certain environmental factors (e.g., a high or low temperature tolerance), decreased ability to survive in certain habitats, or a decreased ability to sustain a certain diet in comparison to a host organism to which the modulating agent has not been administered. In some instances, the methods or compositions provided herein may be effective to decrease host fitness in any plurality of ways described herein. Further, the modulating agent may decrease host fitness in any number of host classes, orders, families, genera, or species (e.g., 1 host species, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 500, or more host species). In some instances, the modulating agent acts on a single host class, order, family, genus, or species.

Host fitness may be evaluated using any standard methods in the art. In some instances, host fitness may be evaluated by assessing an individual host. Alternatively, host fitness may be evaluated by assessing a host population. For example, a decrease in host fitness may manifest as a decrease in successful competition against other insects, thereby leading to a decrease in the size of the host population.

iii. Host insects in disease transmission

By decreasing the fitness of host insects that carry animal pathogens, the modulating agents provided herein are effective to reduce the spread of vector-borne diseases. The modulating agent may be delivered to the insects using any of the formulations and delivery methods described herein, in an amount and for a duration effective to reduce transmission of the disease, e.g., reduce vertical or horizontal transmission between vectors and/or reduce transmission to animals. For example, the modulating agent described herein may reduce vertical or horizontal transmission of a vector-borne pathogen by about 2%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or more in comparison to a host organism to which the modulating agent has not been administered. As an another example, the modulating agent described herein may reduce vectorial competence of an insect vector by about 2%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or more in comparison to a host organism to which the modulating agent has not been administered.

Non-limiting examples of diseases that may be controlled by the compositions and methods provided herein include diseases caused by Togaviridae viruses (e.g., Chikungunya, Ross River fever, Mayaro, Onyon-nyong fever, Sindbis fever, Eastern equine encephalomyelitis, Western equine encephalomyelitis, Venezuelan equine encephalomyelitis, or Barmah forest); diseases caused by Flaviviridae viruses (e.g., Dengue fever, Yellow fever, Kyasanur Forest disease, Omsk haemorrhagic fever, Japanese encephalitis, Murray Valley encephalitis, Rocio, St. Louis encephalitis, West Nile encephalitis, or Tick-borne encephalitis); diseases caused by Bunyaviridae viruses (e.g., Sandly fever, Rift Valley fever, La Crosse encephalitis, California encephalitis, Crimean-Congo haemorrhagic fever, or Oropouche fever); disease caused by Rhabdoviridae viruses (e.g., Vesicular stomatitis); disease caused by Orbiviridae (e.g., Bluetongue); diseases caused by bacteria (e.g., Plague, Tularemia, Q fever, Rocky Mountain spotted fever, Murine typhus, Boutonneuse fever, Queensland tick typhus, Siberian tick typhus, Scrub typhus, Relapsing fever, or Lyme disease); or diseases caused by protozoa (e.g., Malaria, African trypanosomiasis, Nagana, Chagas disease, Leishmaniasis, Piroplasmosis, Bancroftian filariasis, or Brugian filariasis).

II. Target Microorganisms

The microorganisms targeted by the modulating agent described herein may include any microorganism resident in or on the host, including, but not limited to, any bacteria and/or fungi described herein. Microorganisms resident in the host may include, for example, symbiotic (e.g., endosymbiotic microorganisms that provide beneficial nutrients or enzymes to the host), commensal, pathogenic, or parasitic microorganisms. An endosymbiotic microorganism may be a primary endosymbiont or a secondary endosymbiont. A symbiotic microorganism (e.g., bacteria or fungi) may be an obligate symbiont of the host or a facultative symbiont of the host. Microorganisms resident in the host may be acquired by any mode of transmission, including vertical, horizontal, or multiple origins of transmission.

i. Bacteria

Exemplary bacteria that may be targeted in accordance with the methods and compositions provided herein, include, but are not limited to, *Xenorhabdus spp*, *Photorhabdus spp*, *Candidatus spp*, *Buchnera spp*, *Blattabacterium spp*, *Baumania spp*, *Wigglesworthia spp*, *Wolbachia spp*, *Rickettsia spp*, *Orientia spp*, *Sodalis spp*, *Burkholderia spp*, *Cupriavidus spp*, *Frankia spp*, *Snirrhizobium spp*, *Streptococcus spp*, *Wolinella spp*, *Xylella spp*, *Erwinia spp*, *Agrobacterium spp*, *Bacillus spp*, *Paenibacillus spp*, *Streptomyces spp*, *Micrococcus spp*, *Corynebacterium spp*, *Acetobacter spp*, *Cyanobacteria spp*, *Salmonella spp*, *Rhodococcus spp*, *Pseudomonas spp*, *Lactobacillus spp*, *Enterococcus spp*, *Alcaligenes spp*, *Klebsiella spp*, *Paenibacillus spp*, *Arthrobacter spp*, *Corynebacterium spp*, *Brevibacterium spp*, *Thermus spp*, *Pseudomonas spp*, *Clostridium spp*, and *Escherichia spp*. Non-limiting examples of bacteria that may be targeted by the methods and compositions provided herein are shown in Table 1. In some instances, the 16S rRNA sequence of the bacteria targeted by the modulating agent has at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99%, 99.9%, or 100% identity with a sequence listed in Table 1.

Table 1: Examples of Target Bacteria and Host Insects

Primary endosymbiont	Host	Location	16S rRNA
Gamma proteobacteria			
<i>Carsonella ruddii</i>	Psyllids (Psylloidea)	bacteriocytes	TATCCAGCCACAGGTTCCCCTA CAGCTACCTTGTTACGACTTCA CCCCAGTTACAAATCATACCGT TGTAATAGTAAAATTACTTATGA TACAATTTACTTCCATGGTGTGA CGGGCGGTGTGTACAAGGCTC GAGAACGTATTCACCGTAACAT TCTGATTTACGATTACTAGCGAT TCCAACCTTCATGAAATCGAGTT ACAGATTTCAATCCGAACATAAG AATATTTTTTAAGATTAGCATTA TGTTGCCATATAGCATATAACTT TTTGTAACTCATTGTAGCACG TGTGTAGCCCTACTTATAAGGG CCATGATGACTTGACGTCGTCC TCACCTTCCTCCAATTTATCATT GGCAGTTTCTTATTAGTTCTAAT ATATTTTTAGTAAAATAAGATAA GGGTTGCGCTCGTTATAGGACT TAACCCAACATTTACAACACG AGCTGACGACAGCCATGCAGC ACCTGTCTCAAAGCTAAAAAAG CTTTATTATTCTAATAAATTCTT

			<p>TGGATGTCAAAAGTAGGTAAGA TTTTTCGTGTTGTATCGAATTAA ACCACATGCTCCACCGCTTGTG CGAGCCCCCGTCAATTCATTG AGTTTTAACCTTGCGGTCGTAA TCCCCAGGCGGTCAACTTAACG CGTTAGCTTTTTCTACTAAAAATA TATACTTTTTTTCATAAAACAA AATTACAATTATAATTTTAATA AATAGTTGACATCGTTTACTGC ATGGACTACCAGGGTATCTAAT CCTGTTTGCTCCCCATGCTTTC GTGTATTAGTGTGAGTATTAATA TAGAAATACGCCTTCGCCACTA GTATTCTTTCAGATATCTAAGCA TTTCACTGCTACTCCTGAAATTC TAATTTCTTCTTTTATACTCAAG TTTATAAGTATTAATTTCAATATT AAATTACTTTAATAAAATTTAAAA ATTAATTTTAAAAACAACCTGC ACACCCTTACGCCCAATAATT CCGATTAACGCTTGACCCCTC GTATTACCGCGGCTGCTGGCA CGAAGTTAGCCGGTGCTTCTTT TACAAATAACGTCAAAGATAATA TTTTTTTATTATAAAATCTCTTCT TACTTTGTTGAAAGTGTTTTACA ACCCTAAGGCCTTCTTCACACA CGCGATATAGCTGGATCAAGCT TTCGCTCATTGTCCAATATCCC CCTGCTGCCTTCCGTAAAAG TTTGGGCGGTGTCTCAGTCCCA ATGTGGTTGTTTCATCCTCTAAG ATCAACTACGAATCATAGTCTT GTTAAGCTTTTACTTTAACAACT AACTAATTCGATATAAGCTCTTC TATTAGCGAACGACATTCTCGT TCTTTATCCATTAGGATACATAT TGAATTACTATACATTTCTATAT ACTTTTCTAATACTAATAGGTAG</p>
--	--	--	---

			ATTCTTATATATTACTCACCCGT TCGCTGCTAATTATTTTTTAAT AATTCGCACAACCTTGCATGTGT TAAGCTTATCGCTAGCGTTCAA TCTGAGCTATGATCAAACCTCA (SEQ ID NO: 1)
<i>Portiera aleyrodidarum</i> BT-B	whiteflies (Aleyrodoidea)	bacteriocytes	AAGAGTTTGATCATGGCTCAGA TTGAACGCTAGCGGCAGACATA ACACATGCAAGTCGAGCGGCA TCATACAGGTTGGCAAGCGGC GCACGGGTGAGTAATACATGTA AATATACCTAAAAGTGGGGAAT AACGTACGGAAACGTACGCTAA TACCGCATAATTATTACGAGAT AAAGCAGGGGCTTGATAAAAAA AATCAACCTTGCGCTTTTAGAA AATTACATGCCGATTAGCTAG TTGGTAGAGTAAAAGCCTACCA AGGTAACGATCCGTAGCTGGTC TGAGAGGATGATCAGCCACACT GGGACTGAGAAAAGGCCCAGA CTCCTACGGGAGGCAGCAGTG GGGAATATTGGACAATGGGGG GAACCCTGATCCAGTCATGCCG CGTGTGTGAAGAAGGCCTTTGG GTTGTAAAGCACTTTCAGCGAA GAAGAAAAGTTAGAAAATAAAA AGTTATAACTATGACGGTACTC GCAGAAGAAGCACCGGCTAAC TCCGTGCCAGCAGCCGCGGTA AGACGGAGGGTGCAAGCGTTA ATCAGAATTACTGGGCGTAAAG GGCATGTAGGTGGTTTGTAAAG CTTTATGTGAAAGCCCTATGCT TAACATAGGAACGGAATAAAGA ACTGACAACTAGAGTGCAGAA GAGGAAGGTAGAATTCCCGGT GTAGCGGTGAAATGCGTAGATA TCTGGAGGAATACCAGTTGCGA

			<p>AGGCGACCTTCTGGGCTGACA CTGACACTGAGATGCGAAAGC GTGGGGAGCAAACAGGATTAG ATACCCTGGTAGTCCACGCTGT AAACGATATCAACTAGCCGTTG GATTCTTAAAGAATTTTGTGGC GTAGCTAACGCGATAAGTTGAT CGCCTGGGGAGTACGGTCCGA AGGCTAAAACTCAAATGAATTG ACGGGGGGCCCGCACAAAGCGGT GGAGCATGTGGTTTAATTGAT GCAACGCGCAAAACCTTACCTA CTCTTGACATCCAAAGTACTTTC CAGAGATGGAAGGGTGCCTTA GGGAACTTTGAGACAGGTGCT GCATGGCTGTCGTCAGCTCGT GTTGTGAAATGTTGGGTTAAGT CCCGTAACGAGCGCAACCCTT GTCCTTAGTTGCCAACGCATAA GGCGGGAACTTTAAGGAGACT GCTGGTGATAAACCGGAGGAA GGTGGGGACGACGTCAAGTCA TCATGGCCCTTAAGAGTAGGGC AACACACGTGCTACAATGGCAA AAACAAAGGGTCGCAAAATGGT AACATGAAGCTAATCCCAAAAA AATTGTCTTAGTTCGGATTGGA GTCTGAAACTCGACTCCATAAA GTCGGAATCGCTAGTAATCGTG AATCAGAATGTCACGGTGAATA CGTTCTCGGGCCTTGACACAC CGCCCGTCACACCATGGAAGT GAAATGCACCAGAAGTGGCAA GTTTAACCAAAAAACAGGAGAA CAGTCACTACGGTGTGGTTCAT GACTGGGGTGAAGTCGTAACA AGGTAGCTGTAGGGGAACCTG TGGCTGGATCACCTCCTTAA (SEQ ID NO: 2)</p>
--	--	--	--

<i>Buchnera aphidicola str. APS (Acyrthosiphon pisum)</i>	Aphids (Aphidoidea)	bacteriocytes	AGAGTTTGATCATGGCTCAGAT TGAACGCTGGCGGCAAGCCTA ACACATGCAAGTCGAGCGGCA GCGAGAAGAGAGCTTGCTCTCT TTGTGCGCAAGCGGCAAACGG GTGAGTAATATCTGGGGATCTA CCCCAAAGAGGGGGATAACTA CTAGAAATGGTAGCTAATAACCG CATAATGTTGAAAAACCAAAGT GGGGGACCTTTTGGCCTCATG CTTTTGGATGAACCCAGACGAG ATTAGCTTGTTGGTAGAGTAAT AGCCTACCAAGGCAACGATCTC TAGCTGGTCTGAGAGGATAACC AGCCACACTGGAAGTGAAGACA CGGTCCAGACTCCTACGGGAG GCAGCAGTGGGGAATATTGCA CAATGGGCGAAAGCCTGATGC AGCTATGCCGCGTGTATGAAGA AGGCCTTAGGGTTGTAAAGTAC TTTCAGCGGGGAGGAAAAAAT AAAATAATAATTTTATTTCTGTG ACGTTACCCGCAGAAGAAGCA CCGGCTAACTCCGTGCCAGCA GCCGCGGTAATACGGAGGGTG CAAGCGTTAATCAGAATTACTG GGCGTAAAGAGCGGTAGGTG GTTTTTTAAGTCAGGTGTGAAAT CCCTAGGCTCAACCTAGGAACT GCATTTGAACTGGAAACTAG AGTTTCGTAGAGGGAGGTAGAA TTCTAGGTGTAGCGGTGAAATG CGTAGATATCTGGAGGAATACC CGTGGCGAAAGCGGCCTCCTA AACGAAACTGACACTGAGGC GCGAAAGCGTGGGGAGCAAAC AGGATTAGATACCCTGGTAGTC CATGCCGTAAACGATGTCGACT TGGAGGTTGTTTCCAAGAGAAG TGACTTCCGAAGCTAACGCATT
---	------------------------	---------------	---

			AAGTCGACCGCCTGGGGAGTA CGGCCGCAAGGCTAAACTCA AATGAATTGACGGGGGCCCGC ACAAGCGGTGGAGCATGTGGT TTAATTCGATGCAACGCGAAAA ACCTTACCTGGTCTTGACATCC ACAGAATTCTTTAGAAATAAAGA AGTGCCTTCGGGAGCTGTGAG ACAGGTGCTGCATGGCTGTCGT CAGCTCGTGTTGTGAAATGTTG GGTTAAGTCCCGCAACGAGCG CAACCCTTATCCCTGTTGCCA GCGGTTGCGCCGGGAAGTCA AGGAGACTGCCGTTATAAACC GGAGGAAGGTGGGGACGACGT CAAGTCATCATGGCCCTTACGA CCAGGGCTACACACGTGCTAC AATGGTTTATACAAAGAGAAGC AAATCTGCAAAGACAAGCAAAC CTCATAAAGTAAATCGTAGTCC GGACTGGAGTCTGCAACTCGA CTCCACGAAGTCGGAATCGCTA GTAATCGTGGATCAGAATGCCA CGGTGAATACGTTCCCGGGCC TTGTACACACCGCCCGTCACAC CATGGGAGTGGGTTGCAAAG AAGCAGGTATCCTAACCCTTTA AAAGGAAGGCGCTTACCACTTT GTGATTCATGACTGGGGTGAAG TCGTAACAAGGTAACCGTAGGG GAACCTGCGGTTGGATCACCTC CTT (SEQ ID NO: 3)
<i>Buchnera aphidicola str. Sg (Schizaphis graminum)</i>	Aphids (Aphidoidea)	bacteriocytes	AAACTGAAGAGTTTGATCATGG CTCAGATTGAACGCTGGCGGC AAGCCTAACACATGCAAGTCGA GCGGCAGCGAAAAGAAAGCTT GCTTTCTTGTCGGCGAGCGGC AAACGGGTGAGTAATATCTGGG GATCTGCCCAAAAGAGGGGGA

			TAAGTACTAGAAATGGTAGCTA ATACCGCATAAAGTTGAAAAAC CAAAGTGGGGGACCTTTTTTAA AGGCCTCATGCTTTTGGATGAA CCCAGACGAGATTAGCTTGTG GTAAGGTAAAAGCTTACCAAGG CAACGATCTCTAGCTGGTCTGA GAGGATAACCAGCCACACTGG AACTGAGACACGGTCCAGACTC CTACGGGAGGCAGCAGTGGGG AATATTGCACAATGGGCGAAAAG CCTGATGCAGCTATGCCGCGT GTATGAAGAAGGCCTTAGGGTT GTAAAGTACTTTCAGCGGGGAG GAAAAAATTAAACTAATAATTT TATTTTGTGACGTTACCCGCAG AAGAAGCACCGGCTAACTCCGT GCCAGCAGCCGCGGTAATACG GAGGGTGCGAGCGTTAATCAG AATTACTGGGCGTAAAGAGCAC GTAGGTGGTTTTTAAGTCAGA TGTGAAATCCCTAGGCTTAACC TAGGAACTGCATTTGAAACTGA AATGCTAGAGTATCGTAGAGGG AGGTAGAATTCTAGGTGTAGCG GTGAAATGCGTAGATATCTGGA GGAATACCCGTGGCGAAAGCG GCCTCCTAAACGAATACTGACA CTGAGGTGCGAAAGCGTGGGG AGCAAACAGGATTAGATACCCT GGTAGTCCATGCCGTAAACGAT GTCGACTTGGAGGTTGTTTCCA AGAGAAGTGACTTCCGAAGCTA ACGCGTTAAGTCGACCGCCTG GGGAGTACGGCCGCAAGGCTA AAACTCAAATGAATTGACGGGG GCCCCGACAAGCGGTGGAGCA TGTGGTTTAATTGATGCAACG CGAAAAACCTTACCTGGTCTTG ACATCCACAGAATTTTTTAGAAA
--	--	--	---

			TAAAAAAGTGCCTTCGGGAACT GTGAGACAGGTGCTGCATGGC TGTCGTCAGCTCGTGTGTGAA ATGTTGGGTAAAGTCCCGCAAC GAGCGCAACCCTTATCCCCTGT TGCCAGCGGTTCCGGCCGGGAA CTCAGAGGAGACTGCCGGTTAT AAACCGGAGGAAGGTGGGGAC GACGTCAAGTCATCATGGCCCT TACGACCAGGGCTACACACGT GCTACAATGGTTTATACAAAGA GAAGCAAATCTGTAAAGACAAG CAAACCTCATAAAGTAAATCGT AGTCCGGACTGGAGTCTGCAA CTCGACTCCACGAAGTCGGAAT CGCTAGTAATCGTGGATCAGAA TGCCACGGTGAATACGTTCCCG GGCCTTGTACACACCGCCCGT CACACCATGGGAGTGGGTTGC AAAAGAAGCAGATTTCTAACC ACGAAAGTGGAAGGCGTCTAC CACTTTGTGATTCATGACTGGG GTGAAGTCGTAACAAGGTAACC GTAGGGGAACCTGCGGTTGGA TCACCTCCTTA (SEQ ID NO: 4)
<i>Buchnera aphidicola str. Bp (Baizongia pistaciae)</i>	Aphids (Aphidoidea)	bacteriocytes	ACTTAAAATTGAAGAGTTTGATC ATGGCTCAGATTGAACGCTGGC GGCAAGCTTAACACATGCAAGT CGAGCGGCATCGAAGAAAAGT TTACTTTTCTGGCGGCGAGCGG CAAACGGGTGAGTAACATCTGG GGATCTACCTAAAAGAGGGGG ACAACCATTGGAAACGATGGCT AATACCGCATAATGTTTTAAAT AAACCAAAGTAGGGGACTAAAA TTTTTAGCCTTATGCTTTTAGAT GAACCCAGACGAGATTAGCTTG ATGGTAAGGTAATGGCTTACCA AGGCGACGATCTCTAGCTGGTC TGAGAGGATAACCAGCCACACT

		<p>GGAAGTACGATACGGTCCAGA CTCCTACGGGAGGCAGCAGTG GGGAATATTGCACAATGGGCTA AAGCCTGATGCAGCTATGCCG CGTGTATGAAGAAGGCCTTAGG GTTGTAAAGTACTTTTCAGCGGG GAGGAAAGAATTATGTCTAATA TACATATTTTGTGACGTTACCC GAAGAAGAAGCACCGGCTAAC TCCGTGCCAGCAGCCGCGGTA ATACGGAGGGTGCGAGCGTTA ATCAGAATTACTGGGCGTAAAG AGCACGTAGGCGGTTTATTAAG TCAGATGTGAAATCCCTAGGCT TAACTTAGGAACTGCATTTGAA ACTAATAGACTAGAGTCTCATA GAGGGAGGTAGAATTCTAGGT GTAGCGGTGAAATGCGTAGATA TCTAGAGGAATACCCGTGGCG AAAGCGACCTCCTAAATGAAAA CTGACGCTGAGGTGCGAAAGC GTGGGGAGCAAACAGGATTAG ATACCCTGGTAGTCCATGCTGT AAACGATGTCGACTTGGAGGTT GTTTCCTAGAGAAGTGCTTCC GAAGCTAACGCATTAAGTCGAC CGCCTGGGGAGTACGGTCGCA AGGCTAAAACTCAAATGAATTG ACGGGGGGCCCGCACAAGCGGT GGAGCATGTGGTTTAATTGAT GCAACGCGAAGAACCTTACCTG GTCTTGACATCCATAGAATTTTT TAGAGATAAAAGAGTGCCTTAG GGAAGTATGAGACAGGTGCTG CATGGCTGTGTCAGCTCGTGT TGTGAAATGTTGGGTTAAGTCC CGCAACGAGCGCAACCCCTAT CCTTTGTTGCCATCAGGTTATG CTGGGAACTCAGAGGAGACTG CCGGTTATAAACCGGAGGAAG</p>
--	--	--

			GTGGGGATGACGTCAAGTCAT CATGGCCCTTACGACCAGGGC TACACACGTGCTACAATGGCAT ATACAAAGAGATGCAACTCTGC GAAGATAAGCAAACCTCATAAA GTATGTCGTAGTCCGGACTGGA GTCTGCAACTCGACTCCACGAA GTAGGAATCGCTAGTAATCGTG GATCAGAATGCCACGGTGAATA CGTTCCCGGGCCTTGTACACAC CGCCCGTCACACCATGGGAGT GGGTTGCAAAAAGAAGCAGGTA GCTTAACCAGATTATTTTATTGG AGGGCGCTTACCACTTTGTGAT TCATGACTGGGGTGAAGTCGTA ACAAGGTAACCGTAGGGGAAC CTGCGGTTGGATCACCTCCTTA (SEQ ID NO: 5)
<i>Buchnera aphidicola</i> BCc	Aphids (Aphidoidea)	bacteriocytes	ATGAGATCATTAATATATAAAAA TCATGTTCCAATTAATAAATTAG GACAAAATTTTTTACAGAATAAA GAAATTATTAATCAGATAATTAA TTTAATAAATATTAATAAAAAATG ATAATATTATTGAAATAGGATCA GGATTAGGAGCGTTAACTTTTC CTATTTGTAGAATCATTAAAAAA ATGATAGTATTAGAAATTGATGA AGATCTTGTTGTTTTTTAACTC AAAGTTTATTTATTAATAAAATTA CAAATTATAATTGCTGATATTAT AAAATTTGATTTTTGTTGTTTTTT TTCTTTACAGAAATATAAAAAAT ATAGGTTTATTGGTAATTTACCA TATAATATTGCTACTATATTTTTT TTAAAAACAATTAATTTCTTTA TAATATAATTGATATGCATTTTA TGTTCAAAAAGAAGTAGCAAA GAGATTATTAGCTACTCCTGGT ACTAAAGAATATGGTAGATTAA GTATTATTGCACAATATTTTTAT

			AAGATAGAACTGTTATTAATGT TAATAAATTTAATTTTTTCCTAC TCCTAAAGTAGATTCTACTTTTT TACGATTTACTCCTAAATATTTT AATAGTAAATATAAAATAGATAA ACATTTTTCTGTTTTAGAATTAA TACTAGATTTTCTTTCAACAT AGAAGAAAATTTTAAATAATAA TTAATATCTTTATTTTCTACAAA AGAATTAATTTCTTTAGATATTG ATCCATATTCAAGAGCAGAAAA TGTTTCTTTAATTCAATATTGTA AATTAATGAAATATTATTTGAAA AGAAAAATTTTATGTTTAGATTA A (SEQ ID NO: 6)
<i>Buchnera aphidicola</i> (<i>Cinara tujaefilina</i>)	Aphids (Aphidoidea)	bacteriocytes	TTATCTTATTTACATATACGTA ATATTGCGCTGCGTGACGAG GATTTTTTTGAATTCAGATATA TTTGGTTTAATACGTTTAATAAA ACGTATTTTTTTTTTATTTTTCT TATTTGCAATTCAGTAATAGGAA GTTTTTTAGGTATATTTGGATAA TACTGTAATTCCTAATAAAGTT TTTTACAATCCTATCTTCAATAG AATGAAAATAATAATAGCAATT TTTGATCCGGAATGTAATATGTT AATAATAATTTTAATATTTTATG TAATTCATTTATTTCTTGGTTAA TATATATTCGAAAAGCTTGAAAT GTTCTCGTAGCTGGATGTTTAA ATTTGTCATATTTGGGATTGAT TTTTTTATGATTTGAACTAACTC TAACGTGCTTGTTATGGTTTTTT TTTTTATTTGTAATATGATGGCT CGGGATATTTTTTTGCGTATTT TTCTTCGCCAAAATTTTATTA CCTGTTCTATTGTTTTTTGGTTT GTTTTTTTAACCATTTGACTAAC TGATATTCCAGATTTAGGGTTC ATACGCATATCTAAAGGTCCAT

			CATTCAATAATGAAAATCCTCG GATACTAGAATTTAACTGTATTG AAGAAATACCTAAATCTAATAAT ATTCCATCTATTTTATCTCTATTT TTTTCTTTTTTTAATATTTTTTCA ATATTAGAAAATTTACCTAAAAA TATTTTAAATCGCGAATCTTTTA TTTTTTTTCCGATTTTTATAGATT GTGGGTCTTGATCAATACTATA TAACTTTCCATTAACCCCTAATT CTTGAAGAATTGCTTTTGAATGA CCACCACCTCCAAATGTACAAT CAACATATGTACCGTCTTTTTTT ATTTTAAAGTATTGTATGATTTC TTTTGTAAAAACAGGTTTATGAA TCAT (SEQ ID NO: 7)
<i>Buchnera aphidicola str. G002 (Myzus persicae)</i>	Aphids (Aphidoidea)	bacteriocytes	ATGAAAAGTATAAAAACTTTTAA AAAACACTTTCTGTGAAAAAAT ATGGACAAAATTTTCTTATTAAT AAAGAGATCATAAAAAATATTGT TAAAAAATTAATCCAAATATAG AACAAACATTAGTAGAAATCGG ACCAGGATTAGCTGCATTAACT GAGCCCATATCTCAGTTATTAA AAGAGTTAATAGTTATTGAAATA GACTGTAATCTATTATTTTTTT AAAAAACAACCATTTTATTCAA AATTAATAGTTTTTTGTCAAGAT GCTTTAACTTTAATTATACAAA TTTATTTTATAAAAAAATAAATT AATTCGTATTTTTGGTAATTTAC CATATAATATCTCTACATCTTTA ATTATTTTTTTATTTCAACACATT AGAGTAATTCAAGATATGAATTT TATGCTTCAAAAAGAAGTTGCT GCAAGATTAATTGCATTACCTG GAAATAAATATTACGGTCGTTT GAGCATTATATCTCAATATTATT GTGATATCAAAATTTTATTAAAT GTTGCTCCTGAAGATTTTTGGC

			CTATTCCGAGAGTTCATTCTATA TTTGTAATTTAACACCTCATCA TAATTCTCCTTATTTTGTATG ATATTAATATTTTAAGCCTTATT ACAAATAAGGCTTTCCAAAATA GAAGAAAAATATTACGTCATAG TTTAAAAATTTATTTTCTGAAA CAACTTTATTAATTTAGATATT AATCCCAGATTAAGAGCTGAAA ATATTTCTGTTTTTCAGTATTGT CAATTAGCTAATTATTTGTATAA AAAAAATTATACTAAAAAAATT AA (SEQ ID NO: 8)
<i>Buchnera aphidicola str.</i> <i>Ak (Acyrthosiphon</i> <i>kondoi)</i>	Aphids (Aphidoidea)	bacteriocytes	ATTATAAAAAATTTTAAAAACA TTTTCCTTTAAAAAGGTATGGAC AAAATTTTCTTGTCATACAAAA ACTATTCAAAAGATAATTAATAT AATTAATCCAAACACCAAACAA ACATTAGTGGAAATTGGACCTG GATTAGCTGCATTAACAAAACC AATTTGTCAATTATTAGAAGAAT TAATTGTTATTGAAATAGATCCT AATTTATTGTTTTATTAAAAAAA CGTTCATTTTATTCAAAATTAAC AGTTTTTTATCAAGACGCTTTAA ATTTCAATTATACAGATTTGTTT TATAAGAAAAATCAATTAATTGCG TGTTTTTGGAACTTGCCATATA ATATTTCTACATCTTTAATTATTT CTTTATTCAATCATATTAAAGTT ATTCAAGATATGAATTTTATGTT ACAGAAAGAGGTTGCTGAAAGA TTAATTTCTATTCCTGAAAATAA ATCTTATGGCCGTTTAAGCATT TTTCTCAGTATTATTGTAAATT AAAATATTATTAAATGTTGTACC TGAAGATTTTCGACCTATACCG AAAGTGCACTCTGTTTTATCAA TTAACTCCTCATACCAATTCTC CATATTTGTTTATGATACAAAT

			ATCCTCAGTTCTATCACAAGAA ATGCTTTTCAAATAGAAGGAA AATTTTGCATCATAGTTTAAAAA ATTTATTTCTGAAAAAGAACTA ATTCAATTAGAAATTAATCCAAA TTTACGAGCTGAAAATATTTCTA TCTTTCAGTATTGTCAATTAGCT GATTATTTATATAAAAAATTA TAATCTTGTAATAATCAATTAA (SEQ ID NO: 9)
<i>Buchnera aphidicola str.</i> <i>Ua (Uroleucon</i> <i>ambrosiae)</i>	Aphids (Aphidoidea)	bacteriocytes	ATGATACTAAATAAATATAAAAA ATTTATTCCTTTAAAAAGATACG GACAAAATTTCTTGTAATAGA GAAATAATCAAAAATATTATCAA AATAATTAATCCTAAAAAACGC AAACATTATTAGAAATTGGACC GGGTTTAGGTGCGTTAACAAAA CCTATTTGTGAATTTTAAATGA ACTTATCGTCATTGAAATAGATC CTAATATATTATCTTTTTAAAG AAATGTATATTTTGTAAATTT AAAAATATATTGTCATAATGCTT TAGATTTTAATTATAAAAAATATA TTCTATAAAAAAAGTCAATTAAT TCGTATTTTGGAAATTTACCAT ATAATATTTCTACATCTTAATA ATATATTTATTTCCGAATATTGA TATTATTCAAGATATGAATTTTA TGTTACAACAAGAAGTGGCTAA AAGATTAGTTGCTATTCCTGGT GAAAACTTTATGGTCGTTTAA GTATTATATCTCAATATTATTGT AATATTAATAATTATTACATATT CGACCTGAAAAATTTCAACCTA TTCCTAAAGTTAATTCAATGTTT GTAAATTTAACTCCGCATATTCA TTCTCCTTATTTGTTTATGATA TTAATTTATTAAGTAGTATTACA AAACATGCTTTTCAACATAGAA GAAAAATATTGCGTCATAGTTTA

			AGAAATTTTTTTCTGAGCAAGA TTTAATTCATTTAGAAATTAATC CAAATTTAAGAGCTGAAAATGT TTCTATTATTCAATATTGTCAAT TGGCTAATAATTTATATAAAAAA CATAAACAGTTTATTAATAATTA A (SEQ ID NO: 10)
<i>Buchnera aphidicola</i> (<i>Aphis glycines</i>)	Aphids (Aphidoidea)	bacteriocytes	ATGAAAAAGCATATTCCTATAAA AAAATTTAGTCAAAATTTCTTG TAGATTTGAGTGTGATTAaaaaa ATAATTAAATTTATTAATCCGCA GTTAAATGAAATATTGGTTGAAA TTGGACCGGGATTAGCTGCTAT CACTCGACCTATTTGTGATTTG ATAGATCATTTAATTGTGATTGA AATTGATAAAATTTATTAGATA GATTAaaACAGTTCTCATTTTAT TCAAAATTAACAGTATATCATCA AGATGCTTTAGCATTTGATTACA TAAAGTTATTTAATAAAAAAAT AAATTAGTTCGAATTTTGGTAA TTTACCATATCATGTTTCTACGT CTTTAATATTGCATTTATTTAAA AGAATTAATATTATTAAAGATAT GAATTTTATGCTACAAAAAGAA GTTGCTGAACGTTTAATTGCAA CTCCAGGTAGTAAATTATATGG TCGTTTAAGTATTATTTCTCAAT ATTATTGTAATATAAAAGTTTTA TTGCATGTGTCTTCAAAATGTTT TAAACCAGTTCCTAAAGTAGAA TCAATTTTCTTAATTTGACACC TTATACTGATTATTTCCCTTATT TACTTATAATGTAAACGTTCTT AGTTATATTACAAATTTAGCTTT TCAAAAAAGAAGAAAAATATTAC GTCATAGTTTAGGTAAATATTT TCTGAAAAAGTTTTATAAAATT AAATATTAATCCCAAATTAAGAC CTGAGAATATTTCTATATTACAA

			TATTGTCAGTTATCTAATTATAT GATAGAAAATAATATTCATCAG GAACATGTTTGTATTTAA (SEQ ID NO: 11)
<i>Annandia pinicola</i>	(Phylloxeroidea)	bacteriocytes	AGATTGAACGCTGGCGGCATG CCTTACACATGCAAGTCGAACG GTAACAGGTCTTCGGACGCTGA CGAGTGGCGAACGGGTGAGTA ATACATCGGAACGTGCCCAGTC GTGGGGGATAACTACTCGAAA GAGTAGCTAATACCGCATACGA TCTGAGGATGAAAGCGGGGGA CCTTCGGGCCTCGCGCGATTG GAGCGGCCGATGGCAGATTAG GTAGTTGGTGGGATAAAAGCTT ACCAAGCCGACGATCTGTAGCT GGTCTGAGAGGACGACCAGCC ACACTGGAAGTGAATACGGTC CAGACTCTTACGGGAGGCAGC AGTGGGGAATATTGCACAATGG GCGCAAGCCTGATGCAGCTAT GTCGCGTGTATGAAGAAGACCT TAGGGTTGTAAAGTACTTTTGA TAGCATAAGAAGATAATGAGAC TAATAATTTTATTGTCTGACGTT AGCTATAGAAGAAGCACCGGCT AACTCCGTGCCAGCAGCCGCG GTAATACGGGGGGTGCTAGCG TTAATCGGAATTACTGGGCGTA AAGAGCATGTAGGTGGTTTATT AAGTCAGATGTGAAATCCCTGG ACTTAATCTAGGAAGTGCATTT GAAACTAATAGGCTAGAGTTTC GTAGAGGGAGGTAGAATTCTAG GTGTAGCGGTGAAATGCATAGA TATCTAGAGGAATATCAGTGGC GAAGGCGACCTTCTGGACGAT AACTGACGCTAAAATGCGAAAG CATGGGTAGCAAACAGGATTAG ATACCCTGGTAGTCCATGCTGT

			AAACGATGTCGACTAAGAGGTT GGAGGTATAACTTTTAATCTCT GTAGCTAACGCGTTAAGTCGAC CGCCTGGGGAGTACGGTCGCA AGGCTAAAACTCAAATGAATTG ACGGGGGCCTGCACAAGCGGT GGAGCATGTGGTTTAATTCGAT GCAACGCGTAAACCTTACCTG GTCTTGACATCCACAGAATTTTA CAGAAATGTAGAAGTGCAATTT GAACTGTGAGACAGGTGCTGC ATGGCTGTCGTCAGCTCGTGTT GTGAAATGTTGGGTTAAGTCCC GCAACGAGCGCAACCCTTGTC CTTTGTTACCATAAGATTTAAGG AACTCAAAGGAGACTGCCGGT GATAAACTGGAGGAAGGCGGG GACGACGTCAAGTCATCATGGC CCTTATGACCAGGGCTACACAC GTGCTACAATGGCATATACAAA GAGATGCAATATTGCGAAATAA AGCCAATCTTATAAAATATGTCC TAGTTCGGACTGGAGTCTGCAA CTCGACTCCACGAAGTCGGAAT CGCTAGTAATCGTGGATCAGCA TGCCACGGTGAATATGTTTCCA GGCCTTGTTACACACCGCCCGT CACACCATGGAAGTGGATTGCA AAAGAAGTAAGAAAATTAACCT TCTTAACAAGGAAATAACTTAC CACTTTGTGACTCATAACTGGG GTGA (SEQ ID NO: 12)
<i>Moranella endobia</i>	(Coccoidea)	bacteriocytes	TCTTTTGGTAAGGAGGTGATC CAACCGCAGGTTCCCCTACGGT TACCTTGTTACGACTTCACCCC AGTCATGAATCACAAAGTGGTA AGCGCCCTCCTAAAAGGTTAGG CTACCTACTTCTTTTGCAACCCA CTTCCATGGTGTGACGGGCGG

		<p>TGTGTACAAGGCCCGGGAACG TATTCACCGTGGCATTCTGATC CACGATTACTAGCGATTCCTAC TTCATGGAGTCGAGTTGCAGAC TCCAATCCGGACTACGACGCAC TTTATGAGGTCCGCTAACTCTC GCGAGCTTGCTTCTCTTTGTAT GCGCCATTGTAGCACGTGTGTA GCCCTACTCGTAAGGGCCATG ATGACTTGACGTCATCCCCACC TTCCTCCGGTTTATCACCGGCA GTCTCCTTTGAGTTCCCGACCG AATCGCTGGCAAAAAAGGATAA GGGTTGCGCTCGTTGCGGGAC TTAACCCAACATTTACAACAC GAGCTGACGACAGCCATGCAG CACCTGTCTCAGAGTTCCCGAA GGTACCAAAACATCTCTGCTAA GTTCTCTGGATGTCAAGAGTAG GTAAGGTTCTTCGCGTTGCATC GAATTAAACCACATGCTCCACC GCTTGTGCGGGCCCCCGTCAA TTCATTTGAGTTTTAACCTTGCG GCCGTACTCCCCAGGCGGTGCG ATTTAACGCGTTAACTACGAAA GCCACAGTTCAAGACCACAGCT TTCAAATCGACATAGTTTACGG CGTGGACTACCAGGGTATCTAA TCCTGTTTGCTCCCCACGCTTT CGTACCTGAGCGTCAGTATTCG TCCAGGGGGCCGCTTCGCCA CTGGTATTCCTCCAGATATCTA CACATTTACCGCTACACCTGG AATTCTACCCCCCTCTACGAGA CTCTAGCCTATCAGTTTCAAAT GCAGTTCCTAGGTAAAGCCCAG GGATTTACATCTGACTTAATAA ACCGCCTACGTACTCTTTACGC CCAGTAATTCCGATTAACGCTT GCACCTCCGTATTACCGCGG</p>
--	--	--

			CTGCTGGCACGGAGTTAGCCG GTGCTTCTTCTGTAGGTAACGT CAATCAATAACCGTATTAAGGA TATTGCCTTCCTCCCTACTGAA AGTGCTTTACAACCCGAAGGCC TTCTTCACACACGCGGCATGGC TGCATCAGGGTTTCCCCATTG TGCAATATTCCCCACTGCTGCC TCCCGTAGGAGTCTGGACCGT GTCTCAGTTCCAGTGTGGCTGG TCATCCTCTCAGACCAGCTAGG GATCGTCGCCTAGGTAAGCTAT TACCTCACCTACTAGCTAATCC CATCTGGGTTCATCTGAAGGTG TGAGGCCAAAAGGTCCCCAC TTTGGTCTTACGACATTATGCG GTATTAGCTACCGTTTCCAGCA GTTATCCCCCTCCATCAGGCAG ATCCCCAGACTTTACTCACCCG TTCGCTGCTCGCCGGCAAAAAA GTAAACTTTTTTCCGTTGCCGC TCAACTTGCATGTGTTAGGCCT GCCGCCAGCGTTCAATCTGAG CCATGATCAAACCTCTTCAATTAA A (SEQ ID NO: 13)
<i>Ishikawaella capsulata</i> <i>Mpkobe</i>	(Heteroptera)	bacteriocytes	AAATTGAAGAGTTTGATCATGG CTCAGATTGAACGCTAGCGGCA AGCTTAACACATGCAAGTCGAA CGGTAACAGAAAAAAGCTTGCT TTTTTGCTGACGAGTGGCGGAC GGGTGAGTAATGTCTGGGGAT CTACCTAATGGCGGGGGATAA CTACTGGAACGGTAGCTAATA CCGCATAATGTTGTAAAACCAA AGTGGGGGACCTTATGGCCTC ACACCATTAGATGAACCTAGAT GGGATTAGCTTGTAGGTGGGG TAAAGGCTCACCTAGGCAACGA TCCCTAGCTGGTCTGAGAGGAT

		<p>GACCAGCCACACTGGAAGCTGA GATACGGTCCAGACTCCTACG GGAGGCAGCAGTGGGGAATCT TGCACAATGGGCGCAAGCCTG ATGCAGCTATGTCGCGTGTATG AAGAAGGCCTTAGGGTTGTAAA GTACTTTCATCGGGGAAGAAGG ATATGAGCCTAATATTCTCATAT ATTGACGTTACCTGCAGAAGAA GCACCGGCTAACTCCGTGCCA GCAGCCGCGGTAACACGGAGG GTGCGAGCGTTAATCGGAATTA CTGGGCGTAAAGAGCACGTAG GTGGTTTATTAAGTCATATGTGA AATCCCTGGGCTTAACCTAGGA ACTGCATGTGAAACTGATAAAC TAGAGTTTCGTAGAGGGAGGT GGAATTCCAGGTGTAGCGGTG AAATGCGTAGATATCTGGAGGA ATATCAGAGGCGAAGGCGACC TTCTGGACGAAAACTGACACTC AGGTGCGAAAGCGTGGGGAGC AAACAGGATTAGATACCCTGGT AGTCCACGCTGTAAACAATGTC GACTAAAAAACTGTGAGCTTGA CTTGTGGTTTTTGTAGCTAACG CATTAAAGTCGACCGCCTGGGG AGTACGGCCGCAAGGTTAAAC TCAAATGAATTGACGGGGGTCC GCACAAGCGGTGGAGCATGTG GTTTAATTCGATGCAACGCGAA AAACCTTACCTGGTCTTGACAT CCAGCGAATTATATAGAAATAT ATAAGTGCCTTTCGGGGAAGTC TGAGACGCTGCATGGCTGTCGT CAGCTCGTGTTGTGAAATGTTG GGTTAAGTCCCGCAACGAGCG CCCTTATCCTCTGTTGCCAGCG GCATGGCCGGGAAGTCAGAGG AGACTGCCAGTATTAACTGGA</p>
--	--	---

			GGAAGGTGGGGATGACGTCAA GTCATCATGGCCCTTATGACCA GGGCTACACACGTGCTACAATG GTGTATACAAAGAGAAGCAATC TCGCAAGAGTAAGCAAACTCA AAAAGTACATCGTAGTTCGGAT TAGAGTCTGCAACTCGACTCTA TGAAGTAGGAATCGCTAGTAAT CGTGGATCAGAATGCCACGGT GAATACGTTCTCTGGCCTTGTA CACACCGCCCGTCACACCATG GGAGTAAGTTGCAAAAGAAGTA GGTAGCTTAACCTTTATAGGAG GGCGCTTACCACTTTGTGATTT ATGACTGGGGTGAAGTCGTAAC AAGGTAAGTGTAGGGGAACCT GTGGTTGGATTACCTCCTTA (SEQ ID NO: 14)
<i>Baumannia cicadellincola</i>	sharpshooter leafhoppers (Cicadellinae)	bacteriocytes	TTCAATTGAAGAGTTTGATCATG GCTCAGATTGAACGCTGGCGG TAAGCTTAACACATGCAAGTCG AGCGGCATCGGAAAGTAAATTA ATTACTTTGCCGGCAAGCGGCG AACGGGTGAGTAATATCTGGG GATCTACCTTATGGAGAGGGAT AACTATTGGAAACGATAGCTAA CACCGCATAATGTCGTCAGACC AAAATGGGGGACCTAATTTAGG CCTCATGCCATAAGATGAACCC AGATGAGATTAGCTAGTAGGTG AGATAATAGCTCACCTAGGCAA CGATCTCTAGTTGGTCTGAGAG GATGACCAGCCACACTGGAAC GAGACACGGTCCAGACTCCTA CGGGAGGCAGCAGTGGGGAAT CTTGCACAATGGGGGAAACCCCT GATGCAGCTATACCGCGTGTGT GAAGAAGGCCTTCGGGTTGTAA AGCACTTTCAGCGGGGAAGAA AATGAAGTTACTAATAATAATTG

		<p>TCAATTGACGTTACCCGCAAAA GAAGCACCGGCTAACTCCGTG CCAGCAGCCGCGGTAAGACGG AGGGTGCAAGCGTTAATCGGA ATTACTGGGCGTAAAGCGTATG TAGGCGGTTTATTTAGTCAGGT GTGAAAGCCCTAGGCTTAACCT AGGAATTGCATTTGAACTGGT AAGCTAGAGTCTCGTAGAGGG GGGGAGAATTCCAGGTGTAGC GGTGAAATGCGTAGAGATCTG GAAGAATACCAGTGGCGAAGG CGCCCCCTGGACGAAAACCTG ACGCTCAAGTACGAAAGCGTG GGGAGCAAACAGGATTAGATAC CCTGGTAGTCCACGCTGTAAAC GATGTGATTTGAAGGTTGTAG CCTTGAGCTATAGCTTTCGAAG CTAACGCATTAAATCGACCGCC TGGGGAGTACGACCGCAAGGT TAAAACTCAAATGAATTGACGG GGGCCCCGACAAGCGGTGGAG CATGTGGTTTAATTCGATACAA CGCGAAAAACCTTACCTACTCT TGACATCCAGAGTATAAAGCAG AAAAGCTTTAGTGCCTTCGGGA ACTCTGAGACAGGTGCTGCATG GCTGTGTCAGCTCGTGTTGTG AAATGTTGGGTTAAGTCCCGCA ACGAGCGCAACCCTTATCCTTT GTTGCCAACGATTAAGTCGGGA ACTCAAAGGAGACTGCCGGTG ATAAACCGGAGGAAGGTGAGG ATAACGTCAAGTCATCATGGCC CTTACGAGTAGGGCTACACACG TGCTACAATGGTGATACAAAG AGAAGCAATCTCGTAAGAGTTA GCAAACCTCATAAAGTGCATCG TAGTCCGGATTAGAGTCTGCAA CTCGACTCTATGAAGTCGGAAT</p>
--	--	---

			CGCTAGTAATCGTGGATCAGAA TGCCACGGTGAATACGTTCCCG GGCCTTGTAACACACCGCCCGT CACACCATGGGAGTGTATTGCA AAAGAAGTTAGTAGCTTAAGTC ATAATACGAGAGGGCGCTTACC ACTTTGTGATTCACTGAGGG TGAAGTCGTAACAAGGTAACCG TAGGGGAACCTGCGGTTGGAT CACCTCCTTACACTAAA (SEQ ID NO: 15)
<i>Sodalis like</i>	Rhopalus sapporensis	wider tissue tropism	ATTGAACGCTGGCGGCAGGCC TAACACATGCAAGTCGAGCGG CAGCGGGAAGAAGCTTGCTTCT TTGCCGGCGAGCGGCGGACGG GTGAGTAATGTCTGGGGATCTG CCCGATGGAGGGGGATAACTA CTGGAAACGGTAGCTAATACCG CATAACGTCGCAAGACCAAAGT GGGGGACCTTCGGGCCTCACA CCATCGGATGAACCCAGGTGG GATTAGCTAGTAGGTGGGGTAA TGGCTCACCTAGGCGACGATC CCTAGCTGGTCTGAGAGGATG ACCAGTCACACTGGAAGTGA CACGGTCCAGACTCCTACGGG AGGCAGCAGTGGGGAATATTG CACAATGGGGGAAACCCTGAT GCAGCCATGCCGCGTGTGTGA AGAAGGCCCTTCGGGTTGTAAAG CACTTTCAGCGGGGAGGAAGG CGATGGCGTTAATAGCGCTATC GATTGACGTTACCCGCAGAAGA AGCACCGGCTAACTCCGTGCC AGCAGCCGCGGTAATACGGAG GGTGCGAGCGTTAATCGGAATT ACTGGGCGTAAAGCGTACGCA GGCGGTCTGTAAAGTCAGATGT GAAATCCCCGGGCTCAACCTG GGAAGTGCATTTGAAACTGGCA

			<p>GGCTAGAGTCTCGTAGAGGGG GGTAGAATTCCAGGTGTAGCG GTGAAATGCGTAGAGATCTGGA GGAATACCGGTGGCGAAGGCG GCCCCCTGGACGAAGACTGAC GCTCAGGTACGAAAGCGTGGG GAGCAAACAGGATTAGATACCC TGGTAGTCCACGCTGTAAACGA TGTCGATTTGAAGGTTGTGGCC TTGAGCCGTGGCTTTCGGAGCT AACGTGTTAAATCGACCGCCTG GGGAGTACGGCCGCAAGGTTA AAACTCAAATGAATTGACGGGG GCCCCGACAAGCGGTGGAGCA TGTGGTTTAATTCGATGCAACG CGAAGAACCTTACCTACTCTTG ACATCCAGAGAACTTGGCAGAG ATGCTTTGGTGCCTTCGGGAAC TCTGAGACAGGTGCTGCATGG CTGTCGTCAGCTCGTGTTGTGA AATGTTGGGTAAAGTCCCGCAA CGAGCGCAACCCTTATCCTTTA TTGCCAGCGATTCCGGTCGGGA ACTCAAAGGAGACTGCCGGTG ATAAACCGGAGGAAGGTGGGG ATGACGTCAAGTCATCATGGCC CTTACGAGTAGGGCTACACACG TGCTACAATGGCGCATACAAAG AGAAGCGATCTCGCGAGAGTC AGCGGACCTCATAAAGTGCGTC GTAGTCCGGATTGGAGTCTGCA ACTCGACTCCATGAAGTCGGAA TCGCTAGTAATCGTGGATCAGA ATGCCACGGTGAATACGTTCCC GGGCCTTGTACACACCGCCCG TCACACCATGGGAGTGGGTTG CAAAAGAAGTAGGTAGCTTAAC CTTCGGGAGGGCGCTTACCAC TTTGTGATTCATGACTGGGGTG (SEQ ID NO: 16)</p>
--	--	--	---

<i>Hartigia pinicola</i>	The pine bark adelgid	bacteriocytes	AGATTTAACGCTGGCGGCAGG CCTAACACATGCAAGTCGAGCG GTACCAGAAGAAGCTTGCTTCT TGCTGACGAGCGGCGGACGGG TGAGTAATGTATGGGGATCTGC CCGACAGAGGGGGATAACTATT GGAAACGGTAGCTAATACCGCA TAATCTCTGAGGAGCAAAGCAG GGGAACTTCGGTCCTTGCGCTA TCGGATGAACCCATATGGGATT AGCTAGTAGGTGAGGTAATGG CTCCCCTAGGCAACGATCCCTA GCTGGTCTGAGAGGATGATCA GCCACACTGGGACTGAGACAC GGCCCAGACTCCTACGGGAGG CAGCAGTGGGGAATATTGCACA ATGGGCGAAAGCCTGATGCAG CCATGCCGCGTGTATGAAGAA GGCTTTAGGGTTGTAAAGTACT TTCAGTCGAGAGGAAAACATTG ATGCTAATATCATCAATTATTGA CGTTTCCGACAGAAGAAGCACC GGCTAACTCCGTGCCAGCAGC CGCGGTAATACGGAGGGTGCA AGCGTTAATCGGAATTACTGGG CGTAAAGCGCACGCAGGCGGT TAATTAAGTTAGATGTGAAAGC CCCGGGCTTAACCCAGGAATA GCATATAAACTGGTCAACTAG AGTATTGTAGAGGGGGGTAGA ATTCCATGTGTAGCGGTGAAAT GCGTAGAGATGTGGAGGAATA CCAGTGGCGAAGGCGGCCCCC TGGACAAAACTGACGCTCAAA TGCGAAAGCGTGGGGAGCAAA CAGGATTAGATACCCTGGTAGT CCATGCTGTAAACGATGTGCGAT TTGGAGGTTGTTCCCTTGAGGA GTAGCTTCCGTAGCTAACGCGT TAAATCGACCGCCTGGGGGAG
--------------------------	-----------------------	---------------	---

			TACGACTGCAAGGTTAAACTC AAATGAATTGACGGGGGCCCG CACAAGCGGTGGAGCATGTGG TTTAATTCGATGCAACGCGAAA AACCTTACCTACTCTTGACATC CAGATAATTTAGCAGAAATGCT TTAGTACCTTCGGGAAATCTGA GACAGGTGCTGCATGGCTGTC GTCAGCTCGTGTTGTGAAATGT TGGGTAAAGTCCCGCAACGAG CGCAACCCTTATCCTTTGTTGC CAGCGATTAGGTCGGGAAGTC AAAGGAGACTGCCGGTGATAAA CCGGAGGAAGGTGGGGATGAC GTCAAGTCATCATGGCCCTTAC GAGTAGGGCTACACACGTGCT ACAATGGCATATACAAAGGGAA GCAACCTCGCGAGAGCAAGCG AAACTCATAAATTATGTCGTAGT TCAGATTGGAGTCTGCAACTCG ACTCCATGAAGTCGGAATCGCT AGTAATCGTAGATCAGAATGCT ACGGTGAATACGTTCCCGGGC CTTGTACACACCGCCCGTCACA CCATGGGAGTGGGTTGCAAAA GAAGTAGGTAACCTTAACCTTAT GGAAAGCGCTTACCACTTTGTG ATTCATAACTGGGGTG (SEQ ID NO: 17)
<i>Wigglesworthia glossinidia</i>	tsetse fly (Diptera: Glossinidae)	bacteriocytes	
Beta proteobacteria			
<i>Tremblaya phenacola</i>	Phenacoccus avenae (TPPAVE).	bacteriomes	AGGTAATCCAGCCACACCTTCC AGTACGGCTACCTTGTTACGAC TTCACCCCAGTCACAACCCTTA CCTTCGGAAGTGCCCTCCTCAC AACTCAAACCACCAAACACTTT TAAATCAGGTTGAGAGAGGTTA

			GGCCTGTTACTTCTGGCAAGAA TTATTTCCATGGTGTGACGGGC GGTGTGTACAAGACCCGAGAA CATATTCACCGTGGCATGCTGA TCCACGATTACTAGCAATTCCA ACTTCATGCACTCGAGTTTCAG AGTACAATCCGAAGTGAAGGCC GGCTTTGTGAGATTAGCTCCCT TTTGCAAGTTGGCAACTCTTTG GTCCGGCCATTGTATGATGTGT GAAGCCCCACCCATAAAGGCC ATGAGGACTTGACGTCATCCCC ACCTTCCTCCAATTATCGCTG GCAGTCTCTTTAAGGTAAGTGA CTAATCCAGTAGCAATTAAAGA CAGGGGTTGCGCTCGTTACAG GACTTAACCCAACATCTCACGA CACGAGCTGACGACAGCCATG CAGCACCTGTGCACTAATTCTC TTTCAAGCACTCCCGCTTCTCA ACAGGATCTTAGCCATATCAAA GGTAGGTAAGGTTTTTCGCGTT GCATCGAATTAATCCACATCAT CCACTGCTTGTGCGGGTCCCC GTCAATTCCTTTGAGTTTTAACC TTGCGGCCGTAAGTCCCCAGGC GGTCGACTTGTGCGTTAGCTGC ACCACTGAAAAGGAAAAGTCC CAATGGTTAGTCAACATCGTTT AGGGCATGGACTACCAGGGTA TCTAATCCTGTTTGCTCCCAT GCTTTAGTGTCTGAGCGTCAGT AACGAACCAGGAGGCTGCCTA CGCTTTCGGTATTCCTCCACAT CTCTACACATTTCACTGCTACAT GCGGAATTCTACCTCCCCCTCT CGTACTCCAGCCTGCCAGTAAC TGCCGCATTCTGAGGTTAAGCC TCAGCCTTTCACAGCAATCTTA ACAGGCAGCCTGCACACCCTTT
--	--	--	--

			ACGCCCAATAAATCTGATTAAC GCTCGCACCCCTACGTATTACCG CGGCTGCTGGCACGTAGTTTG CCGGTGCTTATTCTTTCGGTAC AGTCACACCACCAAATTGTTAG TTGGGTGGCTTTCTTTCGGAAC AAAAGTGCTTTACAACCCAAAG GCCTTCTTCACACACGCGGCAT TGCTGGATCAGGCTTCGCCCCA TTGTCCAAGATTCTCACTGCT GCCTTCCTCAGAAGTCTGGGCC GTGTCTCAGTCCCAGTGTGGCT GGCCGTCCTCTCAGACCAGCTA CCGATCATTGCCTTGGAAGCC ATTACCTTTCCAACAAGCTAATC AGACATCAGCCAATCTCAGAGC GCAAGGCAATTGGTCCCCTGCT TTCATTCTGCTTGGTAGAGAAC TTTATGCGGTATTAATTAGGCTT TCACCTAGCTGTCCCCACTCT GAGGCATGTTCTGATGCATTAC TCACCCGTTTGCCACTTGCCAC CAAGCCTAAGCCCGTGTTGCC GTTGACTTGATGTGTAAGGC ATGCCGCTAGCGTTCAATCTGA GCCAGGATCAAACCTCT (SEQ ID NO: 18)
<i>Tremblaya princeps</i>	citrus mealybug Planococcus citri	bacteriomes	AGAGTTTGATCCTGGCTCAGAT TGAACGCTAGCGGCATGCATTA CACATGCAAGTCGTACGGCAG CACGGGCTTAGGCCTGGTGGC GAGTGGCGAACGGGTGAGTAA CGCCTCGGAACGTGCCTTGTA GTGGGGGATAGCCTGGCGAAA GCCAGATTAATACCGCATGAAG CCGCACAGCATGCGCGGTGAA AGTGGGGGATTCTAGCCTCAC GCTACTGGATCGGCCGGGGTC TGATTAGCTAGTTGGCGGGGTA ATGGCCCACCAAGGCTTAGATC

		<p>AGTAGCTGGTCTGAGAGGACG ATCAGCCACACTGGGACTGAG ACACGGCCCAGACTCCTACGG GAGGCAGCAGTGGGGAATCTT GGACAATGGGCGCAAGCCTGA TCCAGCAATGCCGCGTGTGTGA AGAAGGCCTTCGGGTCGTAAA GCACTTTTGTTCGGGATGAAGG GGGGCGTGCAAACACCATGCC CTCTTGACGATACCGAAAGAAT AAGCACCGGCTAACTACGTGC CAGCAGCCGCGGTAATACGTA GGGTGCGAGCGTTAATCGGAA TCACTGGGCGTAAAGGGTGCG CGGGTGTTTGCCAAGACCCC TGTAATCCTACGGCCCAACC GTAGTGCTGCGGAGGTTACTG GTAAGCTTGAGTATGGCAGAG GGGGGTAGAATTCCAGGTGTA GCGGTGAAATGCGTAGATATCT GGAGGAATACCGAAGGCGAAG GCAACCCCTGGGCCATCACT GACACTGAGGCACGAAAGCGT GGGAGCAAACAGGATTAGAT ACCCTGGTAGTCCACGCCCTAA ACCATGTCGACTAGTTGTCGGG GGGAGCCCTTTTTCCTCGGTGA CGAAGCTAACGCATGAAGTCGA CCGCCTGGGGAGTACGACCGC AAGGTTAAACTCAAAGGAATT GACGGGGACCCGACACAAGCGG TGGATGATGTGGATTAATTGGA TGCAACGCGAAAAACCTTACCT ACCCTTGACATGGCGGAGATTG TGCCGAGAGGCGGAAGTGCTC GAAAGAGAATCCGTGCACAGG TGCTGCATGGCTGTCGTCAGCT CGTGTCGTGAGATGTTGGGTTA AGTCCCATAACGAGCGCAACC CCCGTCTTTAGTTGCTACCACT</p>
--	--	---

			GGGGCACTCTATAGAGACTGC CGGTGATAAACCGGAGGAAGG TGGGGACGACGTCAAGTCATC ATGGCCTTTATGGGTAGGGCTT CACACGTCATACAATGGCTGGA GCAAAGGGTCGCCAACTCGAG AGAGGGAGCTAATCCCACAAAC CCAGCCCCAGTTCGGATTGCAC TCTGCAACTCGAGTGCATGAAG TCGGAATCGCTAGTAATCGTGG ATCAGCATGCCACGGTGAATAC GTTCTCGGGTCTTGTACACACC GCCCCGCACACCATGGGAGTA AGCCGCATCAGAAGCAGCCTC CCTAACCTATGCTGGGAAGGA GGCTGCGAAGGTGGGGTCTAT GACTGGGGTGAAGTCGTAACA AGGTAGCCGTACCGGAAGGTG CGGCTGGATTACCT (SEQ ID NO: 19)
Vidania		bacteriomes	
<i>Nasuia deltocephalinicola</i>	pestiferous insect host, Macrosteles quadripunctulatus (Hemiptera: Cicadellidae)	bacteriomes	AGTTTAATCCTGGCTCAGATTTA ACGCTTGCGACATGCCTAACAC ATGCAAGTTGAACGTTGAAAAT ATTTCAAAGTAGCGTATAGGTG AGTATAACATTTAAACATACCTT AAAGTTCGGAATACCCGATGA AAATCGGTATAATACCGTATAA AAGTATTTAAGAATTAAAGCGG GGAAAACCTCGTGCTATAAGAT TGTTAAATGCCTGATTAGTTTGT TGGTTTTTAAGGTAAAAGCTTAC CAAGACTTTGATCAGTAGCTAT TCTGTGAGGATGTATAGCCACA TTGGGATTGAAATAATGCCCAA ACCTCTACGGAGGGCAGCAGT GGGGAATATTGGACAATGAGC GAAAGCTTGATCCAGCAATGTC GCGTGTGCGATTAAGGGAAACT

		<p>GTAAAGCACTTTTTTTTAAGAAT AAGAAATTTTAATTAATAATTAA AATTTTTGAATGTATTAAGAA TAAGTACCGACTAATCACGTGC CAGCAGTCGCGTAATACGTG GGGTGCGAGCGTTAATCGGATT TATTGGGCGTAAAGTGATTCA GGCTGCTTAAAAAGATTTATATT AAATATTTAAATTAATTTAAAA AATGTATAAATTACTATTAAGCT AGAGTTTAGTATAAGAAAAAG AATTTTATGTGTAGCAGTGAAAT GCGTTGATATATAAGGAACGC CGAAAGCGAAAGCATTCTG TAATAGAACTGACGCTTATATA CGAAAGCGTGGTAGCAAACA GGATTAGATACCCTGGTAGTCC ACGCCCTAACTATGTCAATTA ACTATTAGAATTTTTTTTAGTGG TG TAGCTAACGCGTTAAATTGA CCGCCTGGGTATTACGATCGCA AGATTA AAACTCAAAGGAATTG ACGGGGACCAGCACAAGCGGT GGATGATGTGGATTAATTGAT GATACGCGAAAAACCTTACCTG CCCTTGACATGGTTAGAATTTTA TTGAAAAATAAAAGTGCTTGA AAAGAGCTAACACACAGGTGCT GCATGGCTGTCGTCAGCTCGT GTCGTGAGATGTTGGGTTAAGT CCCGCAACGAGCGCAACCCCT ACTCTTAGTTGCTAATTAAAGAA CTTTAAGAGAACAGCTAACAAT AAGTTTAGAGGAAGGAGGGGA TGACTTCAAGTCCTCATGGCCC TTATGGGCAGGGCTTCACACGT CATACAATGGTTAATACAAAAA GTTGCAATATCGTAAGATTGAG CTAATCTTTAAATTAATCTTAG TTCGGATTGTA CTCTGCAACTC</p>
--	--	---

			GAGTACATGAAGTTGGAATCGC TAGTAATCGCGGATCAGCATGC CGCGGTGAATAGTTTAACTGGT CTTGTACACACCGCCCGTCACA CCATGGAAATAAATCTTGTTTTA AATGAAGTAATATATTTTATCAA AACAGGTTTTGTAACCGGGGTG AAGTCGTAACA (SEQ ID NO: 20)
<i>Zinderia insecticola</i> CARI	spittlebug <i>Clastoptera</i> <i>arizonana</i>	bacteriocytes	ATATAAATAAGAGTTTGATCCTG GCTCAGATTGAACGCTAGCGGT ATGCTTTACACATGCAAGTCGA ACGACAATATTAAAGCTTGCTTT AATATAAAGTGGCGAACGGGTG AGTAATATATCAAACGTACCTT AAAGTGGGGGATAACTAATTGA AAAATTAGATAATACCGCATATT AATCTTAGGATGAAAATAGGAA TAATATCTTATGCTTTTAGATCG GTTGATATCTGATTAGCTAGTT GGTAGGGTAAATGCTTACCAAG GCAATGATCAGTAGCTGGTTTT AGCGAATGATCAGCCACACTG GAACTGAGACACGGTCCAGAC TTCTACGGAAGGCAGCAGTGG GGAATATTGGACAATGGGAGAA ATCCTGATCCAGCAATACCGCG TGAGTGATGAAGGCCTTAGGGT CGTAAACTCTTTTGTTAGGAA AGAAATAATTTTAAATAATATTT AAAATTGATGACGGTACCTAAA GAATAAGCACCGGCTAACTACG TGCCAGCAGCCGCGGTAATAC GTAGGGTGCAAGCGTTAATCG GAATTATTGGGCGTAAAGAGTG CGTAGGCTGTTATATAAGATAG ATGTGAAATACTTAAGCTTAAC TAAGAACTGCATTTATTACTGTT TAACTAGAGTTTATTAGAGAGA AGTGGAATTTTATGTGTAGCAG

			<p>TGAAATGCGTAGATATATAAAG GAATATCGATGGCGAAGGCAG CTTCTTGAATAATACTGACGC TGAGGCACGAAAGCGTGGGGA GCAAACAGGATTAGATACCTG GTAGTCCACGCCCTAACTATG TCTACTAGTTATTAAATTAAAA TAAATTTAGTAACGTAGCTAAC GCATTAAGTAGACCGCCTGGG GAGTACGATCGCAAGATTAAAA CTCAAAGGAATTGACGGGGAC CCGCACAAGCGGTGGATGATG TGGATTAATTCGATGCAACACG AAAAACCTTACCTACTCTTGAC ATGTTTGGAATTTTAAAGAAATT TAAAAGTGCTTGAAAAAGAACC AAAACACAGGTGCTGCATGGCT GTCGTCAGCTCGTGTCTGAGA TGTTGGGTAAAGTCCCGCAACG AGCGCAACCCTTGTTATTATTT GCTAATAAAAAGAAGCTTTAATAA GACTGCCAATGACAAATTGGAG GAAGGTGGGGATGACGTCAAG TCCTCATGGCCCTTATGAGTAG GGCTTCACACGTCATACAATGA TATATACAATGGGTAGCAAATTT GTGAAAATGAGCCAATCCTTAA AGTATATCTTAGTTCGGATTGTA GTCTGCAACTCGACTACATGAA GTTGGAATCGCTAGTAATCGCG GATCAGCATGCCGCGGTGAAT ACGTTCTCGGGTCTTGACACA CCGCCCCGTCACACCATGGAAG TGATTTTACCAGAAATTATTTG TTAACCTTTATTGAAAAAAAT AATTAAGGTAGAATTCATGACT GGGGTGAAGTCGTAACAAGGT AGCAGTATCGGAAGGTGCGGC TGGATTACATTTTAAAT (SEQ ID NO: 21)</p>
--	--	--	--

<i>Proffella armatura</i>	<i>Diaphorina citri</i> , the Asian citrus psyllid	bacteriomes	
Alpha proteobacteria			
Hodgkinia	Cicada <i>Diceroprocta</i> <i>semicineta</i>	bacteriome	AATGCTGGCGGCAGGCCTAAC ACATGCAAGTCGAGCGGACAA CGTTCAAACGTTGTTAGCGGCG AACGGGTGAGTAATACGTGAGA ATCTACCCATCCCAACGTGATA ACATAGTCAACACCATGTCAAT AACGTATGATTCTGCAACAGG TAAAGATTTTATCGGGGATGGA TGAGCTCAGCTAGATTAGCTA GTTGGTGAGATAAAAGCCCACC AAGGCCAAGATCTATAGCTGGT CTGGAAGGATGGACAGCCACA TTGGGACTGAGACAAGGCCCA ACCCTCTAAGGAGGGCAGCAG TGAGGAATATTGGACAATGGGC GTAAGCCTGATCCAGCCATGCC GCATGAGTGATTGAAGGTCCAA CGGACTGTAAACTCTTTTCTC CAGAGATCATAAATGATAGTAT CTGGTGATATAAGCTCCGGCCA ACTTCGTGCCAGCAGCCGCGG TAATACGAGGGGAGCGAGTATT GTTTCGGTTTTATTGGGCGTAAA GGGTGTCCAGGTTGCTAAGTAA GTTAACAACAAAATCTTGAGATT CAACCTCATAACGTTTCGGTTAA TACTACTAAGCTCGAGCTTGGA TAGAGACAAACGGAATTCCGAG TGTAAGGTTGAAATTCGTTGAT ACTTGGAGGAACACCAGAGGC GAAGGCGGTTTGTACATACCAAG CTGACACTGAAGACACGAAAGC ATGGGGAGCAAACAGGATTAG ATACCCTGGTAGTCCATGCCCT AAACGTTGAGTGCTAACAGTTC

			GATCAAGCCACATGCTATGATC CAGGATTGTACAGCTAACGCGT TAAGCACTCCGCCTGGGTATTA CGACCGCAAGGTTAAACTCAA AGGAATTGACGGAGACCCGCA CAAGCGGTGGAGCATGTGGTTT AATTCGAAGCTACACGAAGAAC CTTACCAGCCCTTGACATACCA TGGCCAACCATCCTGGAAACAG GATGTTGTTCAAGTTAAACCCTT GAAATGCCAGGAACAGGTGCT GCATGGCTGTTGTCAGTTCGTG TCGTGAGATGTATGGTTAAGTC CCAAAACGAACACAACCCTCAC CCATAGTTGCCATAAACACAAT TGGGTTCTCTATGGGTACTGCT AACGTAAGTTAGAGGAAGGTGA GGACCACAACAAGTCATCATGG CCCTTATGGGCTGGGCCACAC ACATGCTACAATGGTGGTTACA AAGAGCCGCAACGTTGTGAGA CCGAGCAAATCTCCAAAGACCA TCTCAGTCCGGATTGTA CTG CAACCCGAGTACATGAAGTAGG AATCGCTAGTAATCGTGGATCA GCATGCCACGGTGAATACGTTT TCGGGTCTTGACACGCCGCC CGTCACACCATGGGAGCTTCG CTCCGATCGAAGTCAAGTTACC CTTGACCACATCTTGGCAAGTG ACCGA (SEQ ID NO: 22)
<i>Wolbachia sp. wPip</i>	Mosquito <i>Culex</i> <i>quinquefasciatus</i>	bacteriome	AAATTTGAGAGTTTGATCCTGG CTCAGAATGAACGCTGGCGGC AGGCCTAACACATGCAAGTCGA ACGGAGTTATATTGTAGCTTGC TATGGTATAACTTAGTGGCAGA CGGGTGAGTAATGTATAGGAAT CTACCTAGTAGTACGGAATAAT TGTTGGAAACGACAACATAAC

			CGTATACGCCCTACGGGGGAA AAATTTATTGCTATTAGATGAGC CTATATTAGATTAGCTAGTTGGT GGGGTAATAGCCTACCAAGGTA ATGATCTATAGCTGATCTGAGA GGATGATCAGCCACACTGGAA CTGAGATACGGTCCAGACTCCT ACGGGAGGCAGCAGTGGGGAA TATTGGACAATGGGCGAAAGCC TGATCCAGCCATGCCGCATGA GTGAAGAAGGCCTTTGGGTTGT AAAGCTCTTTTAGTGAGGAAGA TAATGACGGTACTCACAGAAGA AGTCCTGGCTAACTCCGTGCCA GCAGCCGCGGTAATACGGAGA GGGCTAGCGTTATTCGGAATTA TTGGGCGTAAAGGGCGCGTAG GCTGGTTAATAAGTTAAAAGTG AAATCCCGAGGCTTAACCTTGG AATTGCTTTTAAACTATTAATC TAGAGATTGAAAGAGGATAGAG GAATTCCTGATGTAGAGGTAAA ATTCGTAAATATTAGGAGGAAC ACCAGTGGCGAAGGCGTCTAT CTGGTTCAAATCTGACGCTGAA GCGCGAAGGCGTGGGAGCAA ACAGGATTAGATACCCTGGTAG TCCACGCTGTAAACGATGAATG TTAAATATGGGGAGTTTACTTTC TGTATTACAGCTAACGCGTTAA ACATTCCGCCTGGGGACTACG GTCGCAAGATTAAACTCAAAG GAATTGACGGGGACCCGCACA AGCGGTGGAGCATGTGGTTTAA TTCGATGCAACGCGAAAAACCT TACCACTTCTTGACATGAAAAT CATACCTATTGAAGGGATAGG GTCGGTTCGGCCGATTTTACA CAAGTGTTGCATGGCTGTCGTC AGCTCGTGTCGTGAGATGTTGG
--	--	--	---

			GTTAAGTCCCGCAACGAGCGC AACCCATCATCCTTAGTTGCCAT CAGGTAATGCTGAGTACTTTAA GGAAACTGCCAGTGATAAGCTG GAGGAAGGTGGGGATGATGTC AAGTCATCATGGCCTTTATGGA GTGGGCTACACACGTGCTACAA TGGTGTCTACAATGGGCTGCAA GGTGCGCAAGCCTAAGCTAATC CCTAAAAGACATCTCAGTTCGG ATTGTACTCTGCAACTCGAGTA CATGAAGTTGGAATCGCTAGTA ATCGTGGATCAGCATGCCACG GTGAATACGTTCTCGGGTCTTG TACACACTGCCCGTCACGCCAT GGAATTGGTTTCACTCGAAGC TAATGGCCTAACCGCAAGGAAG GAGTTATTTAAAGTGGGATCAG TGAAGTGGGGTGAAGTCGTAACA AGGTAGCAGTAGGGGAATCTG CAGCTGGATTACCTCCTTA (SEQ ID NO: 23)
Bacteroidetes			
<i>Uzinura diaspidicola</i>	armoured scale insects	bacteriocytes	AAAGGAGATATTCCAACCACAC CTTCCGGTACGGTTACCTTGTT ACGACTTAGCCCTAGTCATCAA GTTTACCTTAGGCAGACCACTG AAGGATTACTGACTTCAGGTAC CCCCAGCTCCCATGGCTTGAC GGGCGGTGTGTACAAGGTTCTG AGAACATATTCACCGCGCCATT GCTGATGCGCGATTACTAGCGA TTCCTGCTTCATAGAGTCGAAT TGCAGACTCCAATCCGAAGTGA GACTGGTTTTAGAGATTAGCTC CTGATCACCAGTGGCTGCCCT TTGTAACCAGCCATTGTAGCAC GTGTGTAGCCCAAGGCATAGA GGCCATGATGATTTGACATCAT

			CCCCACCTTCCTCACAGTTTAC ACCGGCAGTTTTGTTAGAGTCC CCGGCTTTACCCGATGGCAACT AACAATAGGGGTTGCGCTCGTT ATAGGACTTAACCAAACACTTC ACAGCACGAACTGAAGACAACC ATGCAGCACCTTGTAATACGTC GTATAGACTAAGCTGTTTCCAG CTTATTCGTAATACATTTAAGCC TTGGTAAGGTTCTCGCGTATC ATCGAATTAAACCACATGCTCC ACCGCTTGTCGAACCCCGT CAATTCCTTTGAGTTTCAATCTT GCGACTGTACTTCCCAGGTGGA TCACTTATCGCTTTCGCTAAGC CACTGAATATCGTTTTTCCAATA GCTAGTGATCATCGTTTAGGGC GTGGACTACCAGGGTATCTAAT CCTGTTTGCTCCCCACGCTTTC GTGCACTGAGCGTCAGTAAAGA TTTAGCAACCTGCCTTCGCTAT CGGTGTTCTGTATGATATCTAT GCATTTACCGCTACACCATAC ATTCCAGATGCTCCAATCTTACT CAAGTTTACCAGTATCAATAGC AATTTTACAGTTAAGCTGTAAG CTTTCACTACTGACTTAATAAAC AGCCTACACACCCTTTAAACCC AATAAATCCGA ATAACGCTTGTGTCATCCGTAT TGCCGCGGCTGCTGGCACGGA ATTAGCCGACACTTATTCGTATA GTACCTTCAATCTCCTATCACG TAAGATATTTATTTCTATACAA AAGCAGTTTACAACCTAAAAGA CCTTCATCCTGCACGCGACGTA GCTGGTTCAGAGTTTCCTCCAT TGACCAATATTCCTCACTGCTG CCTCCCGTAGGAGTCTGGTCC GTGTCTCAGTACCAGTGTGGAG
--	--	--	--

			GTACACCCTCTTAGGCCCCCTA CTGATCATAGTCTTGGTAGAGC CATTACCTCACCACTAACTAAT CAAACGCAGGCTCATCTTTTGC CACCTAAGTTTTAATAAAGGCT CCATGCAGAACTTTATATTATG GGGGATTAATCAGAATTTCTTC TGGCTATACCCCAGCAAAAGGT AGATTGCATACGTGTTACTCAC CCATTGCGCGGTCGCCGACAA ATTAAAAATTTTCGATGCCCCCT CGACTTGCATGTGTTAAGCTCG CCGCTAGCGTTAATTCTGAGCC AGGATCAAACCTCTTCGTTGTAG (SEQ ID NO: 24)
<i>Sulcia muelleri</i>	Blue-Green Sharpshooter and several other leafhopper species	bacteriocytes	CTCAGGATAAACGCTAGCGGA GGGCTTAACACATGCAAGTCGA GGGGCAGCAAAAATAATTATTT TTGGCGACCGGCAAACGGGTG AGTAATACATACGTAACCTTCCT TATGCTGAGGAATAGCCTGAGG AAACCTGGATTAATACCTCATAA TACAATTTTTTAGAAAGAAAAAT TGTTAAAGTTTTATTATGGCATA AGATAGGCGTATGTCCAATTAG TTAGTTGGTAAGGTAATGGCTT ACCAAGACGATGATTGGTAGG GGGCCTGAGAGGGGCGTTCCC CCACATTGGTACTGAGACACGG ACCAACTTCTACGGAAGGCTG CAGTGAGGAATATTGGTCAATG GAGGAACTCTGAACCAGCCA CTCCGCGTGCAGGATGAAAGA AAGCCTTATTGGTTGTAACTG CTTTTGTATATGAATAAAAAATT CTAATTATAGAAATAATTGAAGG TAATATACGAATAAGTATCGACT AACTCTGTGCCAGCAGTCGCG GTAAGACAGAGGATACAAGCGT TATCCGGATTTATTGGGTTTAA

			GGGTGCGTAGGCGGTTTTTAAA GTCAGTAGTGAAATCTTAAAGC TTAACTTTAAAAGTGCTATTGAT ACTGAAAACTAGAGTAAGGTT GGAGTAACTGGAATGTGTGGT GTAGCGGTGAAATGCATAGATA TCACACAGAACACCGATAGCGA AAGCAAGTTACTAACCCTATAC TGACGCTGAGTCACGAAAGCAT GGGGAGCAAACAGGATTAGAT ACCCTGGTAGTCCATGCCGTAA ACGATGATCACTAACTATTGGG TTTTATACGTTGTAATTCAGTGG TGAAGCGAAAGTGTTAAGTGAT CCACCTGAGGAGTACGACCGC AAGGTTGAAACTCAAAGGAATT GACGGGGGCCCCGCACAATCGG TGGAGCATGTGGTTTAATTCGA TGATACACGAGGAACCTTACCA AGACTTAAATGTACTACGAATA AATTGGAAACAATTTAGTCAAG CGACGGAGTACAAGGTGCTGC ATGGTTGTCGTCAGCTCGTGCC GTGAGGTGTAAGGTTAAGTCCT TTAAACGAGCGCAACCCTTATT ATTAGTTGCCATCGAGTAATGT CAGGGGACTCTAATAAGACTGC CGGCGCAAGCCGAGAGGAAGG TGGGGATGACGTCAAATCATCA CGGCCCTTACGTCTTGGGCCA CACACGTGCTACAATGATCGGT ACAAAAGGGAGCGACTGGGTG ACCAGGAGCAAATCCAGAAAG CCGATCTAAGTTCGGATTGGAG TCTGAAACTCGACTCCATGAAG CTGGAATCGCTAGTAATCGTGC ATCAGCCATGGCACGGTGAATA TGTTCCCGGGCCTTGACACAC CGCCCGTCAAGCCATGGAAGT TGGAAGTACCTAAAGTTGGTTC
--	--	--	---

			GCTACCTAAGGTAAGTCTAATA ACTGGGGCTAAGTCGTAACAAG GTA (SEQ ID NO: 25)
Yeast like			
<i>Symbiotaphrina buchneri</i> voucher JCM9740	Anobiid beetles <i>Stegobium</i> <i>paniceum</i>	mycetome between the foregut and midgut	AGATTAAGCCATGCAAGTCTAA GTATAAGNAATCTATACNGTGA AACTGCGAATGGCTCATTAAAT CAGTTATCGTTTATTTGATAGTA CCTTACTACATGGATAACCGTG GTAATTCTAGAGCTAATACATG CTAAAAACCCCGACTTCGGAAG GGGTGTATTTATTAGATAAAAAA CCAATGCCCTTCGGGGCTCCTT GGTGATTCATGATAACTTAACG AATCGCATGGCCTTGCGCCGG CGATGGTTCATTCAAATTTCTG CCCTATCAACTTTTCGATGGTAG GATAGTGGCCTACCATGGTTTT AACGGGTAACGGGGAATTAGG GTTTCGATTCCGGAGAGGGAGC CTGAGAAACGGCTACCACATCC AAGGAAGGCAGCAGGCGCGCA AATTACCCAATCCCGACACGGG GAGGTAGTGACAATAAATACTG ATACAGGGCTCTTTTGGGTCTT GTAATTGGAATGAGTACAATTT AAATCCCT TAACGAGGAACAATTGGAGGG CAAGTCTGGTGCCAGCAGCCG CGGTAATTCCAGCTCCAATAGC GTATATTAAAGTTGTTGCAGTTA AAAAGCTCGTAGTTGAACCTTG GGCCTGGCTGGCCGGTCCGCC TAACCGCGTGTACTGGTCCGG CCGGGCCCTTTCCTTCTGGGGA GCCGCATGCCCTTCACTGGGT GTGTCGGGGAACCAGGACTTTT ACTTTGAAAAAATTAGAGTGTTT

			<p>AAAGCAGGCCTATGCTCGAATA CATTAGCATGGAATAATAGAAT AGGACGTGCGGTTCTATTTTGT TGGTTTCTAGGACCGCCGTAAT GATTAATAGGGATAGTCGGGG GCATCAGTATTCAATTGTCAGA GGTGAAATTCTTGGATTTATTGA AGACTAACTACTGCGAAAGCAT TTGCCA AGGATGTTTTTCATTAATCAGTGA ACGAAAGTTAGGGGATCGAAG ACGATCAGATACCGTCGTAGTC TTAACCATAAACTATGCCGACT AGGGATCGGGCGATGTTATTAT TTGACTCGCTCGGCACCTTAC GAGAAATCAAAGTCTTTGGGT CTGGGGGGAGTATGGTCGCAA GGCTGAAACTTAAAGAAATTGA CGGAAGGGCACCACCAGGAGT GGAGCCTGCGGCTTAATTTGAC TCAACACGGGGAACTCACCA GGTCCAGACACATTAAGGATTG ACAGATTGAGAGCTCTTTCTTG ATTATGTGGGTGGTGGTGCATG GCCGTTCTTAGTTGGTGGAGTG ATTTGTCTGCTTAATTGCGATAA CGAACGAGACCTTAACCTGCTA AATAGCCCGGTCCGCTTTGGC GGGCCGCTGGCTTCTTAGAGG GACTATCGGCTCAAGCCGATG GAAGTTTGAGGCAATAACAGGT CTGTGATGCCCTTAGATGTTCT GGGCCGCACGCGCGCTACACT GACAGAGCCAACGAGTAAATCA CCTTGGCCGGAAGGTCTGGGT AATCTTGTTAACTCTGTCGTG CTGGGGATAGAGCATTGCAATT ATTGCTCTTCAACGAGGAATTC CTAGTAAGCGCAAGTCATCAGC TTGCGCTGATTACGTCCCTGCC</p>
--	--	--	--

			CTTTGTACACACCGCCCGTCGC TACTACCGATTGAATGGCTCAG TGAGGCCTTCGGACTGGCACA GGGACGTTGGCAACGACGACC CAGTGCCGG AAAGTTGGTCAAACCTTGGTCAT TTAGAGGAAGTAAAAGTCGTAA CAAGGTTTCCGTAGGTGAACCT GCGGAAGGATCATT (SEQ ID NO: 26)
<i>Symbiotaphrina kochii</i> voucher CBS 589.63	Anobiid beetles <i>Lasioderma</i> <i>serricorne</i>	mycetome	TACCTGGTTGATTCTGCCAGTA GTCATATGCTTGTCTCAAAGATT AAGCCATGCAAGTCTAAGTATA AGCAATCTATACGGTGAACTG CGAATGGCTCATTAAATCAGTT ATCGTTTATTTGATAGTACCTTA CTACATGGATAACCGTGGAAT TCTAGAGCTAATACATGCTAAA AACCTCGACTTCGGAAGGGGT GTATTTATTAGATAAAAAACCAA TGCCCTTCGGGGCTCCTTGGT GATTCATGATAACTTAACGAAT CGCATGGCCTTGCGCCGGCGA TGGTTCATTCAAATTTCTGCCCT ATCAACTTTTCGATGGTAGGATA GTGGCCTACCATGGTTTCAACG GGTAACGGGGAATTAGGGTTC GATTCCGGAGAGGGAGCCTGA GAAACGGCTACCACATCCAAG GAAGGCAGCAGGCGCGCAAAT TACCCAATCCCGACACGGGGA GGTAGTGACAATAAATACTGAT ACAGGGCTCTTTGGGTCTTGT AATTGGAATGAGTACAATTTAAA TCCCTTAACGAGGAACAATTGG AGGGCAAGTCTGGTGCCAGCA GCCGCGTAATTCCAGCTCCAA TAGCGTATATTAAAGTTGTTGCA GTTAAAAAGCTCGTAGTTGAAC CTTGGGCCTGGCTGGCCGGTC

			CGCCTAACCGCGTGTACTGGTC CGGCCGGGCCTTTCCTTCTGG GGAGCCGCATGCCCTTCACTG GGTGTGTCGGGGAACCAGGAC TTTTACTTTGAAAAAATTAGAGT GTTCAAAGCAGGCCTATGCTCG AATACATTAGCATGGAATAATA GAATAGGACGTGTGGTTCTATT TTGTTGGTTTCTAGGACCGCCG TAATGATTAATAGGGATAGTCG GGGGCATCAGTATTCAATTGTC AGAGGTGAAATTCTTGGATTTA TTGAAGACTAACTACTGCGAAA GCATTTGCCAAGGATGTTTTCA TTAATCAGTGAACGAAAGTTAG GGGATCGAAGACGATCAGATA CCGTCGTAGTCTTAACCATAAA CTATGCCGACTAGGGATCGGG CGATGTTATTATTTGACTCGCT CGGCACCTTACGAGAAATCAAA GTCTTTGGGTTCTGGGGGGAG TATGGTCGCAAGGCTGAACTT AAAGAAATTGACGGAAGGGCA CCACCAGGAGTGGAGCCTGCG GCTTAATTTGACTCAACACGGG GAAACTCACCAGGTCCAGACAC ATTAAGGATTGACAGATTGAGA GCTCTTTCTTGATTATGTGGGT GGTGGTGCATGGCCGTTCTTAG TTGGTGGAGTGATTTGTCTGCT TAATTGCGATAACGAACGAGAC CTTAACCTGCTAAATAGCCCGG TCCGCTTTGGCGGGCCGCTGG CTTCTTAGAGGGACTATCGGCT CAAGCCGATGGAAGTTTGAGG CAATAACAGGTCTGTGATGCCC TTAGATGTTCTGGGCCGCACGC GCGCTACACTGACAGAGCCAA CGAGTACATCACCTTGGCCGG AAGGTCTGGGTAATCTTGTTAA
--	--	--	--

			ACTCTGTCGTGCTGGGGATAGA GCATTGCAATTATTGCTCTTCAA CGAGGAATTCCTAGTAAGCGCA AGTCATCAGCTTGCGCTGATTA CGTCCCTGCCCTTTGTACACAC CGCCCGTCGCTACTACCGATTG AATGGCTCAGTGAGGCCTTCG GACTGGCACAGGGACGTTGGC AACGACGACCCAGTGCCGGA AGTTCGTCAAACCTTGGTCATTTA GAGGAAGNNNAAGTCGTAACA AGGTTTCCGTAGGTGAACCTGC GGAAGGATCATT (SEQ ID NO: 27)
Primary extracellular symbiont	Host	Location	16 rRNA
fenitrothion-degrading bacteria			
<i>Burkholderia</i> sp. SFA1	<i>Riptortus pedestris</i>	Gut	AGTTTGATCCTGGCTCAGATTG AACGCTGGCGGCATGCCTTAC ACATGCAAGTCGAACGGCAGC ACGGGGGCAACCCTGGTGGCG AGTGGCGAACGGGTGAGTAAT ACATCGGAACGTGTCCTGTAGT GGGGGATAGCCCGCGAAAGC CGGATTAATACCGCATACGACC TAAGGGAGAAAGCGGGGGATC TTCGGACCTCGCGCTATAGGG GCGGCCGATGGCAGATTAGCT AGTTGGTGGGGTAAAGGCCTA CCAAGGCGACGATCTGTAGCT GGTCTGAGAGGACGACCAGCC ACACTGGGACTGAGACACGGC CCAGACTCCTACGGGAGGCAG CAGTGGGGAATTTTGACAATG GGGGCAACCCTGATCCAGCAA TGCCGCGTGTGTGAAGAAGGC TTCGGGTTGTAAAGCACTTTG TCCGGAAGAAAACCTTCGTCCC

		<p>TAATATGGATGGAGGATGACGG TACCGGAAGAATAAGCACCGG CTAACTACGTGCCAGCAGCCG CGGTAATACGTAGGGTGCGAG CGTTAATCGGAATTACTGGGCG TAAAGCGTGCGCAGGCGGTCT GTTAAGACCGATGTGAAATCCC CGGGCTTAACCTGGGAACTGC ATTGGTGACTGGCAGGCTTTGA GTGTGGCAGAGGGGGGTAGAA TTCCACGTGTAGCAGTGAAATG CGTAGAGATGTGGAGGAATAC CGATGGCGAAGGCAGCCCCCT GGGCCAACTACTGACGCTCAT GCACGAAAGCGTGGGGAGCAA ACAGGATTAGATACCCTGGTAG TCCACGCCCTAAACGATGTCAA CTAGTTGTTGGGGATTCAATTC CTTAGTAACGTAGCTAACGCGT GAAGTTGACCGCCTGGGGAGT ACGGTCGCAAGATTAATACTCA AAGGAATTGACGGGGACCCGC ACAAGCGGTGGATGATGTGGA TTAATTCGATGCAACGCGAAAA ACCTTACCTACCCTTGACATGG TCGGAACCCTGCTGAAAGGTG GGGGTGCTCGAAAGAGAACCG GCGCACAGGTGCTGCATGGCT GTCGTCAGCTCGTGTCTGAGA TGTGGGTTAAGTCCCGCAACG AGCGCAACCCTTGTCCTTAGTT GCTACGCAAGAGCACTCTAAG GAGACTGCCGGTGACAAACCG GAGGAAGGTGGGGATGACGTC AAGTCCTCATGGCCCTTATGGG TAGGGCTTCACACGTCATACAA TGGTCGGAACAGAGGGTTGCC AAGCCGCGAGGTGGAGCCAAT CCCAGAAAACCGATCGTAGTCC GGATCGCAGTCTGCAACTCGA</p>
--	--	--

			CTGCGTGAAGCTGGAATCGCTA GTAATCGCGGATCAGCATGCC GCGGTGAATACGTTCCCGGGT CTTGTACACACCGCCCGTCACA CCATGGGAGTGGGTTTCACCA GAAGTAGGTAGCCTAACCGCAA GGAGGGCGCTTACCACGGTGG GATTCATGACTGGGGTGAAGTC GTAACAAGGTAGC (SEQ ID NO: 28)
<i>Burkholderia</i> sp. KM-A	<i>Riptortus pedestris</i>	Gut	GCAACCCTGGTGGCGAGTGGC GAACGGGTGAGTAATACATCG GAACGTGTCCTGTAGTGGGGG ATAGCCCGGCGAAAGCCGGAT TAATACCGCATACGATCTACGG AAGAAAGCGGGGGATCCTTCG GGACCTCGCGCTATAGGGGCG GCCGATGGCAGATTAGCTAGTT GGTGGGGTAAAGGCCTACCAA GGCGACGATCTGTAGCTGGTCT GAGAGGACGACCAGCCACACT GGGACTGAGACACGGCCAGA CTCCTACGGGAGGCAGCAGTG GGGA ATTTTGGACAATGGGGGCAACC CTGATCCAGCAATGCCGCGTGT GTGAAGAAGGCCTTCGGGTTGT AAAGCACTTTTGTCCGAAAGA AAACGTCTTGTTAATACCTGA GGCGGATGACGGTACCGGAAG AATAAGCACCGGCTAACTACGT GCCAGCAGCCGCGGTAATACG TAGGGTGCGAGCGTTAATCGG AATTACTGGGCGTAAAGCGTGC GCAGGCGGTCTGTTAAGACCG ATGTGAAATCCCCGGGCTTAAC CTGGGAACTGCATTGGTGACTG GCAGGCTTTGAGTGTGGCAGA GGGGGGTAGAATTCCACGTGT AGCAGTGAAATGCGTAGAGATG

			<p>TGGA</p> <p>GGAATACCGATGGCGAAGGCA GCCCCCTGGGCCAACACTGAC GCTCATGCACGAAAGCGTGGG GAGCAAACAGGATTAGATACCC TGGTAGTCCACGCCCTAAACGA TGTCAACTAGTTGTTGGGGATT CATTTCCCTTAGTAACGTAGCTAA CGCGTGAAGTTGACCGCCTGG GGAGTACGGTCGCAAGATTAAA ACTCAAAGGAATTGACGGGGA CCCGCACAAAGCGGTGGATGAT GTGGATTAATTCGATGCAACGC GAAAAACCTTACCTACCCTTGA CATGGTCGGAAGTCTGCTGAG AGGTGGACGTGCTCGAAAGAG AACCGGCGCACAGGTGCTGCA TGGCTGTCGTCAGCTCGTGTCCG TGAGATGTTGGGTAAAGTCCCG CAACGAGCGCAACCCTTGTCTT TAGTTGCTACGCAAGAGCACTC TAAGGAGACTGCCGGTGACAA ACCGGAGGAAGGTGGGGATGA CGTCAAGTCCTCATGGCCCTTA TGGGTAGGGCTTCACACGTCAT ACAATGGTCGGAACAGAGGGT TGCCAAGCCGCGAGGTGGAGC CAATCCCAGAAAACCGATCGTA GTCCGGATCGCAGTCTGCAACT CGACTGCGTGAAGCTGGAATC GCTAG</p> <p>TAATCGCGGATCAGCATGCCG CGGTGAATACGTTCCCGGGTCT TGTACACACCGCCCGTCACACC ATGGGAGTGGGTTTACCAGAA GTAGGTAGCCTAACCGCAAGG AGGGCGCTTACCACGGTGGGA TTCATGACTGGGGTGAAGT</p> <p>(SEQ ID NO: 29)</p>
--	--	--	---

Burkholderia sp. KM-G	Riptortus pedestris	Gut	GCAACCCTGGTGGCGAGTGGC GAACGGGTGAGTAATACATCG GAACGTGTCCTGTAGTGGGGG ATAGCCCGGCGAAAGCCGGAT TAATACCGCATACGACCTAAGG GAGAAAGCGGGGGATCTTCGG ACCTCGCGCTATAGGGGCGGC CGATGGCAGATTAGCTAGTTGG TGGGGTAAAGGCCTACCAAGG CGACGATCTGTAGCTGGTCTGA GAGGACGACCAGCCACACTGG GACTGAGACACGGCCCAGACT CCTACGGGAGGCAGCAGTGGG GAATTTTGGACAATGGGGGCAA CCCTGATCCAGCAATGCCGCGT GTGTGAAGAAGGCCTTCGGGTT GTAAAGCACTTTTGTCCGAAA GAAAACCTTCGAGGTTAATACCC TTGGAGGATGACGGTACCGGA AGAATAAGCACCGGCTAACTAC GTGCCAGCAGCCGCGGTAATA CGTAGGGTGCGAGCGTTAATC GGAATTACTGGGCGTAAAGCGT GCGCAGGCGGTCTGTTAAGAC CGATGTGAAATCCCCGGGCTTA ACCTGGGAACTGCATTGGTGAC TGGCAGGCTTTGAGTGTGGCA GAGGGGGGTAGAATTCCACGT GTAGCAGTGAAATGCGTAGAGA TGTGGAGGAATACCGATGGCG AAGGCAGCCCCCTGGGCCAAC ACTGACGCTCATGCACGAAAGC GTGGGGAGCAAACAGGATTAG ATACCCTGGTAGTCCACGCCCT AAACGATGTCAACTAGTTGTTG GGGATTCATTTCTTAGTAACG TAGCTAACGCGTGAAGTTGACC GCCTGGGGAGTACGGTCGCAA GATTA AAACTCAAAGGAATTGA CGGGGACCCGCACAAGCGGTG
-----------------------	---------------------	-----	---

			GATGATGTGGATTAATTCGATG CAACGCGAAAAACCTTACCTAC CCTTGACATGGTCGGAAGTCTG CTGAGAGGTGGACGTGCTCGA AAGAGAACCGGCGCACAGGTG CTGCATGGCTGTC GTCAGCTCGTGTCTGAGATGT TGGGTAAAGTCCCGCAACGAG CGCAACCCTTGTCTTAGTTGC TACGCAAGAGCACTCTAAGGAG ACTGCCGGTGACAAACCGGAG GAAGGTGGGGATGACGTCAAG TCCTCATGGCCCTTATGGGTAG GGCTTCACACGTCATACAATGG TCGGAACAGAGGGTTGCCAAG CCGCGAGGTGGAGCCAATCCC AGAAAACCGATCGTAGTCCGGA TCGCAGTCTGCAACTCGACTGC GTGAAGCTGGAATCGCTAGTAA TCGCGGATCAGCATGCCGCGG TGAATACGTTCCCGGGTCTTGT ACACACCGCCCGTCACACCAT GGGAGTGGGTTTCACCAGAAG TAGGTAGCCTAACCTGCAAAGG AGGGCGCTTACCACG (SEQ ID NO: 30)
<i>Snodgrassella alvi</i>	Honeybee (<i>Apis mellifera</i>) and <i>Bombus</i> spp.	Ileum	GAGAGTTTGATCCTGGCTCAGA TTGAACGCTGGCGGCATGCCTT ACACATGCAAGTCGAACGGCA GCACGGAGAGCTTGCTCTCTG GTGGCGAGTGGCGAACGGGTG AGTAATGCATCGGAACGTACCG AGTAATGGGGGATAACTGTCCG AAAGGATGGCTAATACCGCATA CGCCCTGAGGGGGAAAGCGGG GGATCGAAAGACCTCGCGTTAT TTGAGCGGCCGATGTTGGATTA GCTAGTTGGTGGGGTAAAGGC CTACCAAGGCGACGATCCATAG

		CGGGTCTGAGAGGATGATCCG CCACATTGGGACTGAGACACG GCCCAAACCTCCTACGGGAGGC AGCAGTGGGGAATTTGGACAA TGGGGGGAACCCTGATCCAGC CATGCCGCGTGTCTGAAGAAG GCCTTCGGGTTGTAAAGGACTT TTGTTAGGGAAGAAAAGCCGG GTGTTAATACCATCTGGTGCTG ACGGTACCTAAAGAATAAGCAC CGGCTAACTACGTGCCAGCAG CCGCGGTAATACGTAGGGTGC GAGCGTTAATCGGAATTACTGG GCGTAAAGCGAGCGCAGACGG TTAATTAAGTCAGATGTGAAATC CCCGAGCTCAACTTGGGACGT GCATTTGAAACTGGTAACTAG AGTGTGTCAGAGGGAGGTAGA ATTCCACGTGTAGCAGTGAAAT GCGTAGAGATGTGGAGGAATA CCGATGGCGAAGGCAGCCTCC TGGGATAACACTGACGTTTCATG CTCGAAAGCGTGGGTAGCAA CAGGATTAGATACCCTGGTAGT CCACGCCCTAAACGATGACAAT TAGCTGTTGGGACACTAGATGT CTTAGTAGCGAAGCTAACGCGT GAAATTGTCCGCTGGGGAGT ACGGTCGCAAGATTAAACTCA AAGGAATTGACGGGGACCCGC ACAAGCGGTGGATGATGTGGA TTAATTCGATGCAACGCGAAGA ACCTTACCTGGTCTTGACATGT ACGGAATCTCTTAGAGATAGGA GAGTGCCTTCGGAACCGTAA CACAGGTGCTGCATGGCTGTC GTCAGCTCGTGTCTGAGATGT TGGGTAAAGTCCCGCAACGAG CGCAACCCTTGTCATTAGTTGC CATCATTAAGTTGGGCACTCTA
--	--	--

			ATGAGACTGCCGGTGACAAAC CGGAGGAAGGTGGGGATGACG TCAAGTCCTCATGGCCCTTATG ACCAGGGCTTCACACGTCATAC AATGGTCGGTACAGAGGGTAG CGAAGCCGCGAGGTGAAGCCA ATCTCAGAAAGCCGATCGTAGT CCGGATTGCACTCTGCAACTCG AGTGCATGAAGTCGGAATCGCT AGTAATCGCAGGTCAGCATACT GCGGTGAATACGTTCCCGGGT CTTGTACACACCGCCCGTCACA CCATGGGAGTGGGGGATACCA GAATTGGGTAGACTAACCGCAA GGAGGTCGCTTAACACGGTAT GCTTCATGACTGGGGTGAAGTC GTAACAAGGTAGCCGTAG (SEQ ID NO: 33)
<i>Gilliamella apicola</i>	honeybee (<i>Apis mellifera</i>) and <i>Bombus</i> spp.	Ileum	TTAAATTGAAGAGTTTGATCATG GCTCAGATTGAACGCTGGCGG CAGGCTTAACACATGCAAGTCG AACGGTAACATGAGTGCTTGCA CTTGATGACGAGTGGCGGACG GGTGAGTAAAGTATGGGGATCT GCCGAATGGAGGGGGACAACA GTTGGAACGACTGCTAATACC GCATAAAGTTGAGAGACCAAAG CATGGGACCTTCGGGCCATGC GCCATTTGATGAACCCATATGG GATTAGCTAGTTGGTAGGGTAA TGGCTTACCAAGGCGACGATCT CTAGCTGGTCTGAGAGGATGA CCAGCCACACTGGAAGTGA CACGGTCCAGACTCCTACGGG AGGCAGCAGTGGGGAATATTG CACAATGGGGGAAACCCTGAT GCAGCCATGCCGCGTGTATGA AGAAGGCCTTCGGGTTGTAAAG TACTTTCGGTGATGAGGAAGGT GGTGTATCTAATAGGTGCATCA

			ATTGACGTTAATTACAGAAGAA GCACCGGCTAACTCCGTGCCA GCAGCCGCGGTAATACGGAGG GTGCGAGCGTTAATCGGAATGA CTGGGCGTAAAGGGCATGTAG GCGGATAATTAAGTTAGGTGTG AAAGCCCTGGGCTCAACCTAG GAATTGCACTTAAACTGGTTA ACTAGAGTATTGTAGAGGAAGG TAGAATTCCACGTGTAGCGGTG AAATGCGTAGAGATGTGGAGG AATACCGGTGGCGAAGGCGGC CTTCTGGACAGATACTGACGCT GAGATGCGAAAGCGTGGGGAG CAAACAGGATTAGATACCCTGG TAGTCCACGCTGTAAACGATGT CGATTTGGAGTTTGTTCCTAG AGTGATGGGCTCCGAAGCTAA CGCGATAAATCGACCGCCTGG GGAGTACGGCCGCAAGGTAA AACTCAAATGAATTGACGGGGG CCCGCACAAGCGGTGGAGCAT GTGGTTTAATTGATGCAACGC GAAGAACCTTACCTGGTCTTGA CATCCACAGAATCTTGACAGAG TGCGGGAGTGCCTTCGGGAAC TGTGAGACAGGTGCTGCATGG CTGTCGTCAGCTCGTGTGTGA AATGTTGGGTTAAGTCCCGCAA CGAGCGCAACCCTTATCCTTTG TTGCCATCGGTTAGGCCGGGA ACTCAAAGGAGACTGCCGTTGA TAAAGCGGAGGAAGGTGGGGA CGACGTCAAGTCATCATGGCCC TTACGACCAGGGCTACACACGT GCTACAATGGCGTATACAAAGG GAGGCGACCTCGCGAGAGCAA GCGGACCTCATAAAGTACGTCT AAGTCCGGATTGGAGTCTGCAA CTCGACTCCATGAAGTCGGAAT
--	--	--	---

			CGCTAGTAATCGTGAATCAGAA TGTCACGGTGAATACGTTCCCG GGCCTTGTAACACACCGCCCGT CACACCATGGGAGTGGGTTGC ACCAGAAGTAGATAGCTTAACC TTCGGGAGGGCGTTTACCACG GTGTGGTCCATGACTGGGGTG AAGTCGTAACAAGGTAACCGTA GGGGAACCTGCGGTTGGATCA CCTCCTTAC (SEQ ID NO: 34)
<i>Bartonella apis</i>	honeybee (<i>Apis mellifera</i>)	Gut	AAGCCAAAATCAAATTTTCAACT TGAGAGTTTGATCCTGGCTCAG AACGAACGCTGGCGGCAGGCT TAACACATGCAAGTCGAACGCA CTTTTCGGAGTGAGTGGCAGAC GGGTGAGTAACGCGTGGGAAT CTACCTATTTCTACGGAATAAC GCAGAGAAATTTGTGCTAATAC CGTATACGTCCTTCGGGAGAAA GATTTATCGGAGATAGATGAGC CCGCGTTGGATTAGCTAGTTGG TGAGGTAATGGCCCACCAAGG CGACGATCCATAGCTGGTCTGA GAGGATGACCAGCCACATTGG GACTGAGACACGGCCAGACT CCTACGGGAGGCAGCAGTGGG GAATATTGGACAATGGGCGCAA GCCTGATCCAGCCATGCCGCG TGAGTGATGAAGGCCCTAGGG TTGTAAAGCTCTTTCACCGGTG AAGATAATGACGGTAACCGGAG AAGAAGCCCCGGCTAACTTCGT GCCAGCAGCCGCGGTAATACG AAGGGGGCTAGCGTTGTTCCG ATTTACTGGGCGTAAAGCGCAC GTAGGCGGATATTTAAGTCAGG GGTGAAATCCCGGGGCTCAAC CCCGGAACTGCCTTTGATACTG GATATCTTGAGTATGGAAGAGG

		TAAGTGGAATTCCGAGTGTAGA GGTGAAATTCGTAGATATTCGG AGGAACACCAGTGGCGAAGGC GGCTTACTGGTCCATTACTGAC GCTGAGGTGCGAAAGCGTGGG GAGCAAACAGGATTAGATACCC TGGTAGTCCACGCTGTAAACGA TGAATGTTAGCCGTTGGACAGT TACTGTTCCGTGGCGCAGCTA ACGCATTAAACATTCCGCCTGG GGAGTACGGTCGCAAGATTA ACTCAAAGGAATTGACGGGGG CCCGCACAAAGCGGTGGAGCAT GTGGTTTAATTGAAGCAACGC GCAGAACCTTACCAGCCCTTGA CATCCCGATCGCGGATGGTGG AGACACCGTCTTTCAGTTCGGC TGGATCGGTGACAGGTGCTGC ATGGCTGTCGTCAGCTCGTGTC GTGAGATGTTGGGTTAAGTCCC GCAACGAGCGCAACCCTCGCC CTTAGTTGCCATCATTTAGTTG GGCACTCTAAGGGGACTGCCG GTGATAAGCCGAGAGGAAGGT GGGGATGACGTCAAGTCCTCAT GGCCCTTACGGGCTGGGCTAC ACACGTGCTACAATGGTGGTGA CAGTGGGCAGCGAGACCGCGA GGTCGAGCTAATCTCCAAAAGC CATCTCAGTTCGGATTGCACTC TGCAACTCGAGTGCATGAAGTT GGAATCGCTAGTAATCGTGGAT CAGCATGCCACGGTGAATACGT TCCCGGGCCTTGTACACACCG CCCGTCACACCATGGGAGTTG GTTTTACCCGAAGGTGCTGTGC TAACCGCAAGGAGGCAGGCAA CCACGGTAGGGTCAGCGACTG GGGTGAAGTCGTAACAAGGTA GCCGTAGGGGAACCTGCGGCT
--	--	--

			GGATCACCTCCTTTCTAAGGAA GATGAAGAATTGGAA (SEQ ID NO: 35)
<i>Parasaccharibacter apium</i>	honeybee (<i>Apis mellifera</i>)	Gut	CTACCATGCAAGTCGCACGAAA CCTTTCGGGGTTAGTGCGGA CGGGTGAGTAACGCGTTAGGA ACCTATCTGGAGGTGGGGGAT AACATCGGGAAACTGGTGCTAA TACCGCATGATGCCTGAGGGC CAAAGGAGAGATCCGCCATTG GAGGGGCCTGCGTTCGATTAG CTAGTTGGTTGGGTAAAGGCTG ACCAAGGCGATGATCGATAGCT GGTTTGAGAGGATGATCAGCCA CACTGGGACTGAGACACGGCC CAGACTCCTACGGGAGGCAGC AGTGGGGAATATTGGACAATGG GGGCAACCCTGATCCAGCAAT GCCGCGTGTGTGAAGAAGGTC TTCGATTGTAAAGCACTTTCA CTAGGGAAGATGATGACGGTA CCTAGAGAAGAAGCCCCGGCT AACTTCGTGCCAGCAGCCGCG GTAATACGAAGGGGGCTAGCG TTGCTCGGAATGACTGGGCGTA AAGGGCGCGTAGGCTGTTTGTA CAGTCAGATGTGAAATCCCCGG GCTTAACCTGGGAAC TGCATTT GATACGTGCAGACTAGAGTCC GAGAGAGGGTTGTGGAATTCC CAGTGTAGAGGTGAAATTCGTA GATATTGGGAAGAACACCGGTT GCGAAGGCGGCAACCTGGCTN NNNNNNNNNNNNNNNNNNNNNN NNNNNNNNNNNNNNNNNNNNNN NNNNNNNNNNNNNNNNNNNNNN NNNNNNNNNNNNNNNNNNNNNN NNNNNNNNNNNNNNNNNNNNNN NNNNNNNNNGAGCTAACGCGTT AAGCACACCGCCTGGGGAGTA

			CGGCCGCAAGGTTGAACTCA AAGGAATTGACGGGGGCCCGC ACAAGCGGTGGAGCATGTGGT TTAATTCGAAGCAACGCGCAGA ACCTTACCAGGGCTTGCATGGG GAGGCTGTATTTCAGAGATGGAT ATTTCTTCGGACCTCCCGCACA GGTGCTGCATGGCTGTCTGCA GCTCGTGTCTGAGATGTTGG GTAAAGTCCCGCAACGAGCGC AACCCTTGTCTTTAGTTGCCAT CACGTCTGGGTGGGCACTCTA GAGAGACTGCCGGTGACAAGC CGGAGGAAGGTGGGGATGACG TCAAGTCCTCATGGCCCTTATG TCCTGGGCTACACACGTGCTAC AATGGCGGTGACAGAGGGATG CTACATGGTGACATGGTGCTGA TCTCAAAAACCGTCTCAGTTC GGATTGTACTCTGCAACTCGAG TGCATGAAGGTGGAATCGCTAG TAATCGCGGATCAGCATGCCG CGGTGAATACGTTCCCGGGCC TTGTACACACCGCCCGTCACAC CATGGGAGTTGGTTTGACCTTA AGCCGGTGAGCGAACCGCAAG GAACGCAGCCGACCACCGGTT CGGGTTCAGCGACTGGGGGA (SEQ ID NO: 36)
<i>Lactobacillus kunkeei</i>	honeybee (<i>Apis mellifera</i>)	Gut	TTCCTTAGAAAGGAGGTGATCC AGCCGCAGGTTCTCCTACGGCT ACCTTGTTACGACTTCACCCTA ATCATCTGTCCCACCTTAGACG ACTAGCTCCTAAAAGGTTACCC CATCGTCTTTGGGTGTTACAAA CTCTCATGGTGTGACGGGCGG TGTGTACAAGGCCCGGGAACG TATTCACCGTGGCATGCTGATC CACGATTACTAGTGATTCCAAC TTCATGCAGGCGAGTTGCAGCC

			TGCAATCCGAACTGAGAATGGC TTTAAGAGATTAGCTTGACCTC GCGGTTTCGCGACTCGTTGTAC CATCCATTGTAGCACGTGTGTA GCCCAGCTCATAAGGGGCATG ATGATTTGACGTGTCCTCCACC TTCCTCCGGTTTATCACCGGCA GTCTCACTAGAGTGCCCAACTA AATGCTGGCAACTAATAATAAG GGTTGCGCTCGTTGCGGGACT TAACCCAACATCTCACGACACG AGCTGACGACAACCATGCACCA CCTGTCACTCTGTCCCGAAGG GAACGCCCAATCTCTTGGGTTG GCAGAAGATGTCAAGAGCTGG TAAGGTTCTTCGCGTAGCATCG AATTAAACCACATGCTCCACCA CTTGTGCGGGCCCCCGTCAATT CCTTTGAGTTTCAACCTTGCGG TCGTACTCCCCAGGCGGAATAC TTAATGCGTTAGCTGCGGCACT GAAGGGCGGAAACCCTCCAAC ACCTAGTATTCATCGTTTACGG CATGGACTACCAGGGTATCTAA TCCTGTTGCTACCCATGCTTT CGAGCCTCAGCGTCAGTAACA GACCAGAAAGCCGCTTCGCC ACTGGTGTTCTTCATATATCTA CGCATTTACCGCTACACATGG AGTTCCACTTTCCTCTTGTAC TCAAGTTTTGTAGTTTCCACTGC ACTTCCTCAGTTGAGCTGAGGG CTTTCACAGCAGACTTACAAAA CCGCCTGCGCTCGCTTTACGC CCAATAAATCCGGACAACGCTT GCCACCTACGTATTACCGCGGC TGCTGGCACGTAGTTAGCCGTG GCTTCTGGTTAAATACCGTCA AAGTGTTAACAGTTACTCTAAC ACTTGTTCTTCTTAACAACAGA
--	--	--	--

			GTTTTACGATCCGAAAACCTTC ATCACTCACGCGCGTTGCTCC ATCAGACTTTTGTCCATTGTGG AAGATTCCCTACTGCTGCCTCC CGTAGGAGTCTGGGCCGTGTC TCAGTCCCAATGTGGCCGATTA CCCTCTCAGGTCGGCTACGTAT CATCGTCTTGGTGGGCTTTTAT CTCACCAACTAACTAATACGGC GCGGGTCCATCCCAAAGTGATA GCAAAGCCATCTTTCAAGTTGG AACCATGCGGTTCCAATAATT ATGCGGTATTAGCACTTGTTTC CAAATGTTATCCCCGCTTCGG GGCAGGTACCCACGTGTTACT CACCAGTTCGCCACTCGCTCCG AATCCAAAAATCATTTATGCAAG CATAAAATCAATTTGGGAGAAC TCGTTGCACTTGATGTATTAG GCACGCCGCCAGCGTTCGTCC TGAGCCAGGATCAAACCTCAT CTTAA (SEQ ID NO: 37)
<i>Lactobacillus Firm-4</i>	honeybee (<i>Apis mellifera</i>)	Gut	ACGAACGCTGGCGGCGTGCCT AATACATGCAAGTCGAGCGCG GGAAGTCAGGGAAGCCTTCGG GTGGAAGTGGTGAACGAGCG GCGGATGGGTGAGTAACACGT AGGTAACCTGCCCTAAAGCGG GGGATACCATCTGGAAACAGGT GCTAATACCGCATAAACCCAGC AGTCACATGAGTGCTGGTTGAA AGACGGCTTCGGCTGTCACTTT AGGATGGACCTGCGGCGTATT AGCTAGTTGGTGGAGTAACGGT TCACCAAGGCAATGATACGTAG CCGACCTGAGAGGGTAATCGG CCACATTGGGACTGAGACACG GCCCAAACCTCCTACGGGAGGC AGCAGTAGGGAATCTTCCACAA

		<p>TGGACGCAAGTCTGATGGAGC AACGCCGCGTGGATGAAGAAG GTCTTCGGATCGTAAAATCCTG TTGTTGAAGAAGAACGGTTGTG AGAGTAACTGCTCATAACGTGA CGGTAATCAACCAGAAAAGTCAC GGCTAACTACGTGCCAGCAGC CGCGGTAATACGTAGGTGGCA AGCGTTGTCCGATTATTGGG CGTAAAGGGAGCGCAGGCGGT CTTTTAAGTCTGAATGTGAAAG CCCTCAGCTTAACTGAGGAAGA GCATCGGAAACTGAGAGACTTG AGTGCAGAAGAGGAGAGTGGA ACTCCATGTGTAGCGGTGAAAT GCGTAGATATATGGAAGAACAC CAGTGGCGAAGGCGGCTCTCT GGTCTGTTACTGACGCTGAGGC TCGAAAGCATGGGTAGCGAAC AGGATTAGATACCCTGGTAGTC CATGCCGTAAACGATGAGTGCT AAGTGTTGGGAGGTTTCCGCCT CTCAGTGCTGCAGCTAACGCAT TAAGCACTCCGCCTGGGGAGT ACGACCGCAAGGTTGAACTCA AAGGAATTGACGGGGGCCCGC ACAAGCGGTGGAGCATGTGGT TTAATTCGAAGCAACGCGAAGA ACCTTACCAGGTCTTGACATCT CCTGCAAGCCTAAGAGATTAGG GGTTCCTTCGGGGACAGGAA GACAGGTGGTGCATGGTTGTC GTCAGCTCGTGTCTGAGATGT TGGGTTAAGTCCCGCAACGAG CGCAACCTTGTTACTAGTTGC CAGCATTAAAGTTGGGCACTCTA GTGAGACTGCCGGTGACAAAC CGGAGGAAGGTGGGGACGACG TCAAATCATCATGCCCTTATG ACCTGGGCTACACACGTGCTAC</p>
--	--	---

			AATGGATGGTACAATGAGAAGC GAACTCGCGAGGGGAAGCTGA TCTCTGAAAACCATTCTCAGTTC GGATTGCAGGCTGCAACTCGC CTGCATGAAGCTGGAATCGCTA GTAATCGCGGATCAGCATGCC GCGGTGAATACGTTCCCGGGC CTTGTACACACCGCCC (SEQ ID NO: 38)
Silk worm			
Enterococcus	<i>Bombyx mori</i>	Gut	AGGTGATCCAGCCGCACCTTCC GATACGGCTACCTTGTTACGAC TTCACCCCAATCATCTATCCCA CCTTAGGCGGCTGGCTCCAAA AAGGTTACCTCACCGACTTCGG GTGTTACAACTCTCGTGGTGT GACGGGCGGTGTGTACAAGGC CCGGGAACGTATTCACCGCGG CGTGCTGATCCGCGATTACTAG CGATTCCGGCTTCATGCAGGC GAGTTGCAGCCTGCAATCCGAA CTGAGAGAAGCTTTAAGAGATT TGCATGACCTCGCGGTCTAGC GACTCGTTGTACTTCCCATTGT AGCACGTGTGTAGCCCAGGTC ATAAGGGGCATGATGATTGAC GTCATCCCCACCTTCCTCCGGT TTGTCACCGGCAGTCTCGCTAG AGTGCCCAACTAAATGATGGCA ACTAACAATAAGGGTTGCGCTC GTTGCGGGACTTAACCCAACAT CTCACGACACGAGCTGACGAC AACCATGCACCACCTGTCACTT TGTCCCCGAAGGGAAAGCTCTA TCTCTAGAGTGGTCAAAGGATG TCAAGACCTGGTAAGGTTCTTC GCGTTGCTTCGAATTAACCCAC ATGCTCCACCGCTTGTGCGGG CCCCCGTCAATTCCTTTGAGTT

		<p>TCAACCTTGCGGTCGTA CTCCC CAGGCGGAGTGCTTAATGCGTT TGCTGCAGCACTGAAGGGCGG AAACCCTCCAACACTTAGCACT CATCGTTTACGGCGTGGACTAC CAGGGTATCTAATCCTGTTTGC TCCCCACGCTTTCGAGCCTCAG CGTCAGTTACAGACCAGAGAG CCGCCTTCGCCACTGGTGTTCC TCCATATATCTACGCATTTACC GCTACACATGGAATTCCACTCT CCTCTTCTGCACTCAAGTCTCC CAGTTTCCAATGACCCTCCCCG GTTGAGCCGGGGGCTTTCACAT CAGACTTAAGAAACCGCCTGCG CTCGCTTTACGCCAATAAATC CGGACAACGCTTGCCACCTAC GTATTACCGCGGCTGCTGGCA CGTAGTTAGCCGTGGCTTTCTG GTTAGATACCGTCAGGGGACG TTCAGTTACTAACGTCCTTGTT TTCTCTAACAACAGAGTTTTACG ATCCGAAAACCTTCTTCACTCA CGCGGCGTTGCTCGGTCAGAC TTTCGTCCATTGCCGAAGATTC CCTACTGCTGCCTCCCGTAGGA GTCTGGGCGGTGTCTCAGTCC CAGTGTGGCCGATCACCCTCTC AGGTCGGCTATGCATCGTGGC CTTGGTGAGCCGTTACCTCACC AACTAGCTAATGCACCGCGGGT CCATCCATCAGCGACACCCGAA AGCGCCTTTCACTCTTATGCCA TGCGGCATAAACTGTTATGCGG TATTAGCACCTGTTTCCAAGTG TTATCCCCCTCTGATGGGTAGG TTACCCACGTGTTACTCACCCG TCCGCCACTCCTCTTTCCAATT GAGTGCAAGCACTCGGGAGGA AAGAAGCGTTCGACTTGCATGT</p>
--	--	--

			ATTAGGCACGCCGCCAGCGTT CGTCCTGAGCCAGGATCAAACCT CT (SEQ ID NO: 39)
<i>Delftia</i>	<i>Bombyx mori</i>	Gut	CAGAAAGGAGGTGATCCAGCC GCACCTTCCGATACGGCTACCT TGTTACGACTTCACCCAGTCA CGAACCCCGCCGTGGTAAGCG CCCTCCTTGCGGTTAGGCTACC TACTTCTGGCGAGACCCGCTCC CATGGTGTGACGGGCGGTGTG TACAAGACCCGGAACGTATTC ACCGCGGCATGCTGATCCGCG ATTACTAGCGATTCCGACTTCA CGCAGTCGAGTTGCAGACTGC GATCCGGACTACGACTGGTTTT ATGGGATTAGCTCCCCCTCGCG GGTTGGCAACCCTCTGTACCAG CCATTGTATGACGTGTGTAGCC CCACCTATAAGGGCCATGAGG ACTTGACGTCATCCCCACCTTC CTCCGGTTTGTACCGGCAGTC TCATTAGAGTGCTCAACTGAAT GTAGCAACTAATGACAAGGTT GCGCTCGTTGCGGGACTTAAC CCAACATCTCACGACACGAGCT GACGACAGCCATGCAGCACCT GTGTGCAGGTTCTTTTCGAGC ACGAATCCATCTCTGGAACTT CCTGCCATGTCAAAGGTGGGTA AGGTTTTTCGCGTTGCATCGAA TTAAACCACATCATCCACCGCT TGTGCGGGTCCCCGTCAATTCC TTTGAGTTTCAACCTTGCGGCC GTACTCCCCAGGCGGTCAACTT CACGCGTTAGCTTCGTTACTGA GAAACTAATTCCCAACAACCA GTTGACATCGTTTAGGGCGTGG ACTACCAGGGTATCTAATCCTG TTTGCTCCCCACGCTTTTCGTGC

			ATGAGCGTCAGTACAGGTCCA GGGGATTGCCTTCGCCATCGG TGTTCTCCGCATATCTACGCA TTTCAGTCTACACGCGGAATT CCATCCCCCTCTACCGTACTCT AGCCATGCAGTCACAAATGCAG TTCCCAGGTTGAGCCCGGGGA TTTCACATCTGTCTTACATAACC GCCTGCGCACGCTTTACGCCC AGTAATTCCGATTAACGCTCGC ACCCTACGTATTACCGCGGCTG CTGGCACGTAGTTAGCCGGTG CTTATTCTTACGGTACCGTCAT GGGCCCCCTGTATTAGAAGGA GCTTTTTCGTTCCGTACAAAAG CAGTTTACAACCCGAAGGCCTT CATCCTGCACGCGGCATTGCTG GATCAGGCTTTGCCCCATTGTC CAAAATTCCCCACTGCTGCCTC CCGTAGGAGTCTGGGCCGTGT CTCAGTCCCAGTGTGGCTGGTC GTCCTCTCAGACCAGCTACAGA TCGTCGGCTTGGTAAGCTTTTA TCCCACCAACTACCTAATCTGC CATCGGCCGCTCCAATCGCGC GAGGCCCCGAAGGGCCCCCGCT TTCATCCTCAGATCGTATGCGG TATTAGCTACTCTTTCGAGTAGT TATCCCCACGACTGGGCACGT TCCGATGTATTACTACCCGTT CGCCACTCGTCAGCGTCCGAA GACCTGTTACCGTTGACTTGC ATGTGTAAGGCATGCCGCCAG CGTTCAATCTGAGCCAGGATCA AACTCTACAGTTTCGATCT (SEQ ID NO: 40)
<i>Pelomonas</i>	<i>Bombyx mori</i>	Gut	ATCCTGGCTCAGATTGAACGCT GGCGGCATGCCTTACACATGC AAGTCGAACGGTAACAGGTAA GCTGACGAGTGGCGAACGGGT

		<p>GAGTAATATATCGGAACGTGCC CAGTCGTGGGGGATAACTGCT CGAAAGAGCAGCTAATACCGCA TACGACCTGAGGGTGAAAGCG GGGGATCGCAAGACCTCGCNN GATTGGAGCGGCCGATATCAG ATTAGGTAGTTGGTGGGGTAAA GGCCCACCAAGCCAACGATCT GTAGCTGGTCTGAGAGGACGA CCAGCCACACTGGGACTGAGA CACGGCCCAGACTCCTACGGG AGGCAGCAGTGGGGAATTTTG GACAATGGGCGCAAGCCTGAT CCAGCCATGCCGCGTGCGGGA AGAAGGCCTTCGGGTTGTAAAC CGCTTTTGTGAGGGAAGAAAAG GTTCTGGTTAATACCTGGGACT CATGACGGTACCTGAAGAATAA GCACCGGCTAACTACGTGCCA GCAGCCGCGGTAATACGTAGG GTGCAAGCGTTAATCGGAATTA CTGGGCGTAAAGCGTGCGCAG GCGGTTATGCAAGACAGAGGT GAAATCCCCGGGCTCAACCTG GGAAGTGCCTTTGTGACTGCAT AGCTAGAGTACGGTAGAGGGG GATGGAATTCCGCGTGTAGCA GTGAAATGCGTAGATATGCGGA GGAACACCGATGGCGAAGGCA ATCCCCTGGACCTGTACTGACG CTCATGCACGAAAGCGTGGGG AGCAAACAGGATTAGATACCCT GGTAGTCCACGCCCTAAACGAT GTCAACTGGTTGTTGGGAGGGT TTCTTCTCAGTAACGTANNTAAC GCGTGAAGTTGACCGCCTGGG GAGTACGGCCGCAAGGTTGAA ACTCAAAGGAATTGACGGGGA CCCGCACAAGCGGTGGATGAT GTGGTTTAATTGATGCAACGC</p>
--	--	--

			GAAAAACCTTACCTACCCTTGA CATGCCAGGAATCCTGAAGAGA TTTGGGAGTGCTCGAAAGAGAA CCTGGACACAGGTGCTGCATG GCCGTCGTCAGCTCGTGTCGT GAGATGTTGGGTAAAGTCCCGC AACGAGCGCAACCCTTGTCATT AGTTGCTACGAAAGGGCACTCT AATGAGACTGCCGGTGACAAAC CGGAGGAAGGTGGGGATGACG TCAGGTCATCATGGCCCTTATG GGTAGGGCTACACACGTCATAC AATGGCCGGGACAGAGGGCTG CCAACCCGCGAGGGGGAGCTA ATCCCAGAAACCCGGTCGTAGT CCGGATCGTAGTCTGCAACTCG ACTGCGTGAAGTCGGAATCGCT AGTAATCGCGGATCAGCTTGCC GCGGTGAATACGTTCCCGGGT CTTGTACACACCGCCCGTCACA CCATGGGAGCGGGTTCTGCCA GAAGTAGTTAGCCTAACCGCAA GGAGGGCGATTACCACGGCAG GGTTCGTGACTGGGGTGAAGT CGTAACAAGGTAGCCGTATCG GAAGGTGCGGCTGGATCAC (SEQ ID NO: 41)
--	--	--	--

For example, a mosquito (e.g., *Aedes spp.* or *Anopheles spp.*) harbors symbiotic bacteria that modulate the mosquito's immune response and influence vectorial competence to pathogens. The modulating agent described herein may be useful in targeting bacteria resident in the mosquito, including, but not limited to, *EspZ*, *Serratia spp.* (e.g., *Serratia marcescens*), *Enterobacteriaceae spp.*, *Enterobacter spp.* (e.g., *Enterobacter cloacae*, *Enterobacter amnigenus*, *Enterobacter ludwigii*), *Proteus spp.*, *Acinetobacter spp.*, *Wigglesworthia spp.* (*Wigglesworthia gloosinidia*), *Xanthomonas spp.* (e.g., *Xanthomonas maltophilia*), *Pseudomonas spp.* (e.g., *Pseudomonas aeruginosa*, *Pseudomonas stutzeri*, *Pseudomonas rhodesiae*), *Escherichia spp.* (e.g., *Escherichia coli*), *Cedecea spp.* (e.g., *Cedecea lapagei*), *Ewingella spp.* (e.g., *Ewingella americana*), *Bacillus spp.* (e.g., *Bacillus pumilus*), *Comamonas spp.*, or *Vagococcus spp.* (e.g., *Vagococcus salmoninarum*), or *Wolbachia spp.* (e.g., *Wolbachia – wMel*, *Wolbachia – wAlbB*, *Wolbachia – wMelPop*, *Wolbachia – wMelPop-CLA*).

Any number of bacterial species may be targeted by the compositions or methods described herein. For example, in some instances, the modulating agent may target a single bacterial species. In

some instances, the modulating agent may target at least about any of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 500, or more distinct bacterial species. In some instances, the modulating agent may target any one of about 1 to about 5, about 5 to about 10, about 10 to about 20, about 20 to about 50, about 50 to about 100, about 100 to about 200, about 200 to about 500, about 500 to about 1000, or more distinct bacterial species. In some instances, the modulating agent may target at least about any of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, or more phyla, classes, orders, families, or genera of bacteria.

In some instances, the modulating agent may increase a population of one or more bacteria (e.g., pathogenic bacteria, toxin-producing bacteria) by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in the host in comparison to a host organism to which the modulating agent has not been administered. In some instances, the modulating agent may reduce the population of one or more bacteria (e.g., symbiotic bacteria, pesticide-degrading bacteria, e.g., a bacterium that degrades any one of the pesticides listed in Table 12) by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more in the host in comparison to a host organism to which the modulating agent has not been administered. In some instances, the modulating agent may eradicate the population of a bacterium (e.g., symbiotic bacteria, pesticide-degrading bacteria, e.g., a bacterium that degrades any one of the pesticides listed in Table 12) in the host. In some instances, the modulating agent may increase the level of one or more pathogenic bacteria by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in the host and/or decreases the level of one or more symbiotic bacteria by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more in the host in comparison to a host organism to which the modulating agent has not been administered.

In some instances, the modulating agent may alter the bacterial diversity and/or bacterial composition of the host. In some instances, the modulating agent may increase the bacterial diversity in the host relative to a starting diversity by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more in comparison to a host organism to which the modulating agent has not been administered. In some instances, the modulating agent may decrease the bacterial diversity in the host relative to a starting diversity by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more in comparison to a host organism to which the modulating agent has not been administered.

In some instances, the modulating agent may alter the function, activity, growth, and/or division of one or more bacterial cells. For example, the modulating agent may alter the expression of one or genes in the bacteria. In some instances, the modulating agent may alter the function of one or more proteins in the bacteria. In some instances, the modulating agent may alter the function of one or more cellular structures (e.g., the cell wall, the outer or inner membrane) in the bacteria. In some instances, the modulating agent may kill (e.g., lyse) the bacteria.

The target bacterium may reside in one or more parts of the insect. Further, the target bacteria may be intracellular or extracellular. In some instances, the bacteria reside in or on one or more parts of the host gut, including, e.g., the foregut, midgut, and/or hindgut. In some instances, the bacteria reside as an intracellular bacteria within a cell of the host insect. In some instances, the bacteria reside in a bacteriocyte of the host insect.

Changes to the populations of bacteria in the host may be determined by any methods known in the art, such as microarray, polymerase chain reaction (PCR), real-time PCR, flow cytometry, fluorescence microscopy, transmission electron microscopy, fluorescence *in situ* hybridization (e.g., FISH), spectrophotometry, matrix-assisted laser desorption ionization-mass spectrometry (MALDI-MS), and DNA sequencing. In some instances, a sample of the host treated with a modulating agent is sequenced (e.g., by metagenomics sequencing of 16S rRNA or rDNA) to determine the microbiome of the host after delivery or administration of the modulating agent. In some instances, a sample of a host that did not receive the modulating agent is also sequenced to provide a reference.

ii. Fungi

Exemplary fungi that may be targeted in accordance with the methods and compositions provided herein, include, but are not limited to *Amylostereum areolatum*, *Epichloe spp*, *Pichia pinus*, *Hansenula capsulate*, *Daldinia decipien*, *Ceratocytis spp*, *Ophiostoma spp*, and *Attamyces bromatificus*. Non-limiting examples of yeast and yeast-like symbionts found in insects include *Candida*, *Metschnikowia*, *Debaromyces*, *Scheffersomyces shehatae* and *Scheffersomyces stipites*, *Starmerella*, *Pichia*, *Trichosporon*, *Cryptococcus*, *Pseudozyma*, and yeast-like symbionts from the subphylum *Pezizomycotina* (e.g., *Symbiotaphrina buchneri* and *Symbiotaphrina kochii*). Non-limiting examples of yeast that may be targeted by the methods and compositions herein are listed in Table 2.

Table 2

Insect Species	Order: Family	Yeast Location (Species)
<i>Stegobium paniceum</i>	Coleoptera: Anobiidae	Mycetomes
(= <i>Sitodrepa panicea</i>)		(<i>Saccharomyces</i>)
		Cecae (<i>Torulopsis buchnerii</i>)
		Mycetome between foregut and midgut
		Mycetomes (<i>Symbiotaphrina buchnerii</i>)
		Mycetomes and digestive tube (<i>Torulopsis buchnerii</i>)
		Gut cecae (<i>Symbiotaphrina buchnerii</i>)
<i>Lasioderma serricorne</i>	Coleoptera: Anobiidae	Mycetome between foregut and midgut
		(<i>Symbiotaphrina kochii</i>)
<i>Ernobius abietis</i>	Coleoptera: Anobiidae	Mycetomes (<i>Torulopsis karawaiewii</i>)
		(<i>Candida karawaiewii</i>)
<i>Ernobius mollis</i>	Coleoptera: Anobiidae	Mycetomes (<i>Torulopsis ernobii</i>)
		(<i>Candida ernobii</i>)
<i>Hemicoelus gibbicollis</i>	Coleoptera: Anobiidae	Larval mycetomes

<i>Xestobium plumbeum</i>	Coleoptera: Anobiidae	Mycetomes (<i>Torulopsis xestobii</i>)
		(<i>Candida xestobii</i>)
<i>Criocephalus rusticus</i>	Coleoptera: Cerambycidae	Mycetomes
<i>Phoracantha semipunctata</i>	Coleoptera: Cerambycidae	Alimentary canal (<i>Candida guilliermondii</i> , <i>C. tenuis</i>)
		Cecae around midgut (<i>Candida guilliermondii</i>)
<i>Harpium inquisitor</i>	Coleoptera: Cerambycidae	Mycetomes (<i>Candida rhagii</i>)
<i>Harpium mordax</i> <i>H. sycophanta</i>	Coleoptera: Cerambycidae	Cecae around midgut (<i>Candida tenuis</i>)
<i>Gaurotes virginea</i>	Coleoptera: Cerambycidae	Cecae around midgut (<i>Candida rhagii</i>)
<i>Leptura rubra</i>	Coleoptera: Cerambycidae	Cecae around midgut (<i>Candida tenuis</i>)
		Cecae around midgut (<i>Candida parapsilosis</i>)
<i>Leptura maculicornis</i>	Coleoptera: Cerambycidae	Cecae around midgut (<i>Candida parapsilosis</i>)
<i>L. cerambyciformis</i>		
<i>Leptura sanguinolenta</i>	Coleoptera: Cerambycidae	Cecae around midgut (<i>Candida sp.</i>)
<i>Rhagium bifasciatum</i>	Coleoptera: Cerambycidae	Cecae around midgut (<i>Candida tenuis</i>)
<i>Rhagium inquisitor</i>	Coleoptera: Cerambycidae	Cecae around midgut (<i>Candida guilliermondii</i>)
<i>Rhagium mordax</i>	Coleoptera: Cerambycidae	Cecae around midgut (<i>Candida</i>)
<i>Carpophilus hemipterus</i>	Coleoptera: Nitidulidae	Intestinal tract (10 yeast species)
<i>Odontotaenius disjunctus</i>	Coleoptera: Passalidae	Hindgut (<i>Enteroramus dimorphus</i>)
<i>Odontotaenius disjunctus</i>	Coleoptera: Passalidae	Gut (<i>Pichia stipitis</i> , <i>P. segobiensis</i> , <i>Candida shehatae</i>)
<i>Verres sternbergianus</i>		(<i>C. ergatensis</i>)
<i>Scarabaeus semipunctatus</i>	Coleoptera: Scarabaeidae	Digestive tract (10 yeast species)
<i>Chironitis furcifer</i>		
<i>Unknown species</i>	Coleoptera: Scarabaeidae	Guts (<i>Trichosporon cutaneum</i>)

<i>Dendroctonus and Ips spp.</i>	Coleoptera: Scolytidae	Alimentary canal (13 yeast species)
<i>Dendroctonus frontalis</i>	Coleoptera: Scolytidae	Midgut (<i>Candida sp.</i>)
<i>Ips sexdentatus</i>	Coleoptera: Scolytidae	Digestive tract (<i>Pichia bovis</i> , <i>P. rhodanensis</i>)
		Hansenula holstii (<i>Candida rhagii</i>)
		Digestive tract
		(<i>Candida pulcherina</i>)
<i>Ips typographus</i>	Coleoptera: Scolytidae	Alimentary canal
		Alimentary tracts (<i>Hansenula capsulata</i> , <i>Candida parapsilosis</i>)
		Guts and beetle homogenates (<i>Hansenula holstii</i> , <i>H. capsulata</i> , <i>Candida diddensii</i> , <i>C. mohshtana</i> , <i>C. nitratophila</i> , <i>Cryptococcus albidus</i> , <i>C. laurentii</i>)
<i>Trypodendron lineatum</i>	Coleoptera: Scolytidae	Not specified
<i>Xyloterinus politus</i>	Coleoptera: Scolytidae	Head, thorax, abdomen (<i>Candida</i> , <i>Pichia</i> , <i>Saccharomycopsis</i>)
<i>Periplaneta americana</i>	Dictyoptera: Blattidae	Hemocoel (<i>Candida sp. nov.</i>)
<i>Blatta orientalis</i>	Dictyoptera: Blattidae	Intestinal tract (<i>Kluyveromyces blattae</i>)
<i>Blatella germanica</i>	Dictyoptera: Blattellidae	Hemocoel
<i>Cryptocercus punctulatus</i>	Dictyoptera: Cryptocercidae	Hindgut (1 yeast species)
<i>Philophylla heraclei</i>	Diptera: Tephritidae	Hemocoel
<i>Aedes (4 species)</i>	Diptera: Culicidae	Internal microflora (9 yeast genera)
<i>Drosophila pseudoobscura</i>	Diptera: Drosophilidae	Alimentary canal (24 yeast species)
<i>Drosophila (5 spp.)</i>	Diptera: Drosophilidae	Crop (42 yeast species)
<i>Drosophila melanogaster</i>	Diptera: Drosophilidae	Crop (8 yeast species)
<i>Drosophila (4 spp.)</i>	Diptera: Drosophilidae	Crop (7 yeast species)
<i>Drosophila (6 spp.)</i>	Diptera: Drosophilidae	Larval gut (17 yeast species)
<i>Drosophila (20 spp.)</i>	Diptera: Drosophilidae	Crop (20 yeast species)

<i>Drosophila</i> (8 species groups)	Diptera: Drosophilidae	Crop (<i>Kloeckera</i> , <i>Candida</i> , <i>Kluyveromyces</i>)
<i>Drosophila serido</i>	Diptera: Drosophilidae	Crop (18 yeast species)
<i>Drosophila</i> (6 spp.)	Diptera: Drosophilidae	Intestinal epithelium (<i>Coccidiascus legeri</i>)
<i>Protaxymia melanoptera</i>	Diptera	Unknown (<i>Candida</i> , <i>Cryptococcus</i> , <i>Sporobolomyces</i>)
<i>Astegopteryx styraci</i>	Homoptera: Aphididae	Hemocoel and fat body
<i>Tuberaphis sp.</i>	Homoptera: Aphididae	Tissue sections
<i>Hamiltonaphis styraci</i>		
<i>Glyphinaphis bambusae</i>		
<i>Cerataphis sp.</i>		
<i>Hamiltonaphis styraci</i>	Homoptera: Aphididae	Abdominal hemocoel
<i>Cofana unimaculata</i>	Homoptera: Cicadellidae	Fat body
<i>Leofa unicolor</i>	Homoptera: Cicadellidae	Fat body
<i>Lecaniines, etc.</i>	Homoptera: Coccoidea d	Hemolymph, fatty tissue, etc.
<i>Lecanium sp.</i>	Homoptera: Coccidae	Hemolymph, adipose tissue
<i>Ceroplastes</i> (4 sp.)	Homoptera: Coccidae	Blood smears
<i>Laodelphax striatellus</i>	Homoptera: Delphacidae	Fat body
		Eggs
		Eggs (<i>Candida</i>)
<i>Nilaparvata lugens</i>	Homoptera: Delphacidae	Fat body
		Eggs (2 unidentified yeast species)
		Eggs, nymphs (<i>Candida</i>)
		Eggs (7 unidentified yeast species)
		Eggs (<i>Candida</i>)
<i>Nisia nervosa</i>	Homoptera: Delphacidae	Fat body
<i>Nisia grandiceps</i>		
<i>Perkinsiella spp.</i>		
<i>Sardia rostrata</i>		

<i>Sogatella furcifera</i>		
<i>Sogatodes orizicola</i>	Homoptera: Delphacidae	Fat body
<i>Amrasca devastans</i>	Homoptera: Jassidae	Eggs, mycetomes, hemolymph
<i>Tachardina lobata</i>	Homoptera: Kerriidae	Blood smears (<i>Torulopsis</i>)
<i>Laccifer (=Lakshadia)</i> <i>sp.</i>	Homoptera: Kerriidae	Blood smears (<i>Torula variabilis</i>)
<i>Comperia merceti</i>	Hymenoptera: Encyrtidae	Hemolymph, gut, poison gland
<i>Solenopsis invicta</i>	Hymenoptera: Formicidae	Hemolymph (<i>Myrmecomycetes annellisae</i>)
<i>S. quinquecupis</i>		
<i>Solenopsis invicta</i>	Hymenoptera: Formicidae	Fourth instar larvae (<i>Candida parapsilosis</i> , <i>Yarrowia lipolytica</i>)
		Gut and hemolymph (<i>Candida parapsilosis</i> , <i>C. lipolytica</i> , <i>C. guillermontii</i> , <i>C. rugosa</i> , <i>Debaryomyces hansenii</i>)
<i>Apis mellifera</i>	Hymenoptera: Apidae	Digestive tracts (<i>Torulopsis sp.</i>)
		Intestinal tract (<i>Torulopsis apicola</i>)
		Digestive tracts (8 yeast species)
		Intestinal contents (12 yeast species)
		Intestinal contents (7 yeast species)
		Intestines (14 yeast species)
		Intestinal tract (<i>Pichia melissophila</i>)
		Intestinal tracts (7 yeast species)
		Alimentary canal (<i>Hansenula silvicola</i>)
		Crop and gut (13 yeast species)
<i>Apis mellifera</i>	Hymenoptera: Apidae	Midguts (9 yeast genera)
<i>Anthophora occidentalis</i>	Hymenoptera:Anthophoridae	
<i>Nomia melanderi</i>	Hymenoptera:Halictidae	
<i>Halictus rubicundus</i>	Hymenoptera:Halictidae	
<i>Megachile rotundata</i>	Hymenoptera:Megachilidae	

Any number of fungal species may be targeted by the compositions or methods described herein. For example, in some instances, the modulating agent may target a single fungal species. In some instances, the modulating agent may target at least about any of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 500, or more distinct fungal species. In some instances, the modulating agent may target any one of about 1 to about 5, about 5 to about 10, about 10 to about 20, about 20 to about 50, about 50 to about 100, about 100 to about 200, about 200 to about 500, about 10 to about 50, about 5 to about 20, or about 10 to about 100 distinct fungal species. In some instances, the modulating agent may target at least about any of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, or more phyla, classes, orders, families, or genera of fungi.

In some instances, the modulating agent may increase a population of one or more fungi (e.g., pathogenic or parasitic fungi) by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in the host in comparison to a host organism to which the modulating agent has not been administered. In some instances, the modulating agent may reduce the population of one or more fungi (e.g., symbiotic fungi) by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in the host in comparison to a host organism to which the modulating agent has not been administered. In some instances, the modulating agent may eradicate the population of a fungi (e.g., symbiotic fungi) in the host. In some instances, the modulating agent may increase the level of one or more symbiotic fungi by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in the host and/or may decrease the level of one or more symbiotic fungi by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in the host in comparison to a host organism to which the modulating agent has not been administered.

In some instances, the modulating agent may alter the fungal diversity and/or fungal composition of the host. In some instances, the modulating agent may increase the fungal diversity in the host relative to a starting diversity by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more in comparison to a host organism to which the modulating agent has not been administered. In some instances, the modulating agent may decrease the fungal diversity in the host relative to a starting diversity by at least about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more in comparison to a host organism to which the modulating agent has not been administered.

In some instances, the modulating agent may alter the function, activity, growth, and/or division of one or more fungi. For example, the modulating agent may alter the expression of one or more genes in the fungus. In some instances, the modulating agent may alter the function of one or more proteins in the fungus. In some instances, the modulating agent may alter the function of one or more cellular components in the fungus. In some instances, the modulating agent may kill the fungus.

Further, the target fungus may reside in one or more parts of the insect. In some instances, the fungus resides in or on one or more parts of the insect gut, including, e.g., the foregut, midgut, and/or hindgut. In some instances, the fungus lives extracellularly in the hemolymph, fat bodies or in specialized structures in the host.

Changes to the population of fungi in the host may be determined by any methods known in the art, such as microarray, polymerase chain reaction (PCR), real-time PCR, flow cytometry, fluorescence microscopy, transmission electron microscopy, fluorescence *in situ* hybridization (e.g., FISH),

spectrophotometry, matrix-assisted laser desorption ionization-mass spectrometry (MALDI-MS), and DNA sequencing. In some instances, a sample of the host treated with a modulating agent is sequenced (e.g., by metagenomics sequencing) to determine the microbiome of the host after delivery or administration of the modulating agent. In some instances, a sample of a host that did not receive the modulating agent is also sequenced to provide a reference.

III. Modulating Agents

The modulating agent of the methods and compositions provided herein may include a phage, a polypeptide, a small molecule, an antibiotic, a secondary metabolite, a bacterium, a fungus, or any combination thereof.

i. Phage

The modulating agent described herein may include a phage (e.g., a lytic phage or a non-lytic phage). In some instances, an effective concentration of any phage described herein may alter a level, activity, or metabolism of one or more microorganisms (as described herein) resident in a host described herein (e.g., a vector of an animal pathogen, e.g., a mosquito, a mite, a biting louse, or a tick), the modulation resulting in a decrease in the host's fitness (e.g., as outlined herein). In some instances, the modulating agent includes at least one phage selected from the order *Tectiviridae*, *Myoviridae*, *Siphoviridae*, *Podoviridae*, *Caudovirales*, *Lipothrixviridae*, *Rudoviridae*, or *Ligamenvirales*. In some instances, the composition includes at least one phage selected from the family *Myoviridae*, *Siphoviridae*, *Podoviridae*, *Lipothrixviridae*, *Rudoviridae*, *Ampullaviridae*, *Bicaudaviridae*, *Clavaviridae*, *Corticoviridae*, *Cystoviridae*, *Fuselloviridae*, *Gluboloviridae*, *Guttaviridae*, *Inoviridae*, *Leviviridae*, *Microviridae*, *Plasmaviridae*, and *Tectiviridae*. Further non-limiting examples of phages useful in the methods and compositions are listed in Table 3.

Table 3: Examples of Phage and Targeted Bacteria

Phage and Accession #	Target bacteria	Target host
SA1(NC_027991), phiP68 (NC_004679)	<i>Staphylococcus</i> sp.	<i>Apidae</i> family
WO (AB036666.1)	<i>Wolbachia</i> sp.	<i>Aedes aegypti</i> ; <i>Drosophila melanogaster</i> ; <i>Plasmodium</i> sp; <i>Teleogryllus taiwanemima</i> ; <i>Bombyx mori</i>
KL1 (NC_018278), BcepNazgul (NC_005091) PhiE125 (NC_003309)	<i>Burkholderia</i> sp.	<i>Riptortus</i> sp.; <i>Pyrrhocoris apterus</i> .
Fern (NC_028851), Xenia (NC_028837), HB10c2 (NC_028758)	<i>Paenibacillus</i> larvae	bumble bees: <i>Bombus</i> sp.; honey bees: <i>A. mellifera</i>

CP2 (NC_020205), XP10 (NC_004902), XP15 (NC_007024), phiL7 (NC_012742)	<i>Xanthomonas</i> sp.	<i>Plebeina denoiti</i> ; <i>Apidae</i> family; <i>Apis mellifera</i> ; <i>Drosophilidae</i> family; and <i>Chloropidae</i> family
PP1 (NC_019542), PM1 (NC_023865)	<i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i>	<i>Apidae</i> family
ΦRSA1 (NC_009382), ΦRSB1 (NC_011201), ΦRSL1 (NC_010811), RSM1 (NC_008574)	<i>Ralstonia solanacearum</i>	<i>Bombyx mori</i>
SF1(NC_028807)	<i>Streptomyces scabies</i>	<i>Philantus</i> sp.; <i>Trachypus</i> sp
ECML-4 (NC_025446), ECML-117 (NC_025441), ECML-134 (NC_025449)	<i>Escherichia coli</i>	<i>Apidae</i> family; <i>Varroa destructor</i>
SSP5(JX274646.1), SSP6 (NC_004831), SFP10 (NC_016073), F18SE (NC_028698)	<i>Salmonella</i> sp.	<i>Drosophilidae</i> family
γ (NC_001416), Bcp1 (NC_024137)	<i>Bacillus</i> sp.	Gypsy moth; <i>Lymantria dispar</i> ; <i>Varroa destructor</i>
Phi1 (NC_009821)	<i>Enterococcus</i> sp.	<i>Schistocerca gregaria</i>
ΦKMV (NC_005045), ΦEL(AJ697969.1), ΦKZ (NC_004629)	<i>Pseudomonas</i> sp.	<i>Lymantria dispar</i> ; <i>Apidae</i> family
A2 (NC_004112), phig1e (NC_004305)	<i>Lactobacilli</i> sp.	<i>Apidae</i> family; <i>Drosophila</i> family; <i>Varroa destructor</i>
KLPN1 (NC_028760)	<i>Klebsiella</i> sp	<i>C. capitata</i>
vB_AbaM_Acibel004 (NC_025462), vB_AbaP_Acibel007 (NC_025457)	<i>Acinetobacter</i> sp.	<i>Schistocerca gregaria</i>

In some instances, a modulating agent includes a lytic phage. Thus, after delivery of the lytic phage to a bacterial cell resident in the host, the phage causes lysis in the target bacterial cell. In some instances, the lytic phage targets and kills a bacterium resident in a host insect to decrease the fitness of the host. Alternatively or additionally, the phage of the modulating agent may be a non-lytic phage (also referred to as lysogenic or temperate phage). Thus, after delivery of the non-lytic phage to a bacterial cell resident in the host, the bacterial cell may remain viable and able to stably maintain expression of genes encoded in the phage genome. In some instances, a non-lytic phage is used to alter gene expression in a bacterium resident in a host insect to decrease the fitness of the host. In some instances, the modulating agent includes a mixture of lytic and non-lytic phage.

In certain instances, the phage is a naturally occurring phage. For example, a naturally occurring phage may be isolated from an environmental sample containing a mixture of different phages. The naturally occurring phage may be isolated using methods known in the art to isolate, purify, and identify phage that target a particular microorganism (e.g., a bacterial endosymbiont in an insect host).

5 Alternatively, in certain instances, the phage may be engineered based on a naturally occurring phage.

The modulating agent described herein may include phage with either a narrow or broad bacterial target range. In some instances, the phage has a narrow bacterial target range. In some instances, the phage is a promiscuous phage with a large bacterial target range. For example, the promiscuous phage may target at least about any of 5, 10, 20, 30, 40, 50, or more bacterium resident in the host. A phage
10 with a narrow bacterial target range may target a specific bacterial strain in the host without affecting another, e.g., non-targeted, bacterium in the host. For example, the phage may target no more than about any of 50, 40, 30, 20, 10, 8, 6, 4, 2, or 1 bacterium resident in the host. For example, the phage described herein may be useful in targeting one or more bacteria resident in the mosquito, including, but not limited to, *EspZ*, *Serratia* spp. (e.g., *Serratia marcescens*), *Enterobacteriaceae* spp., *Enterobacter* spp.
15 (e.g., *Enterobacter cloacae*, *Enterobacter amnigenus*, *Enterobacter ludwigii*), *Proteus* spp., *Acinetobacter* spp., *Wigglesworthia* spp. (*Wigglesworthia gloosinidia*), *Xanthomonas* spp. (e.g., *Xanthomonas maltophilia*), *Pseudomonas* spp. (e.g., *Pseudomonas aeruginosa*, *Pseudomonas stutzeri*, *Pseudomonas rhodesiae*), *Escherichia* spp. (e.g., *Escherichia coli*), *Cedecea* spp. (e.g., *Cedecea lapagei*), *Ewingella* spp. (e.g., *Ewingella americana*), *Bacillus* spp. (e.g., *Bacillus pumilus*), *Comamonas* spp., or *Vagococcus* spp. (e.g., *Vagococcus salmoninarum*), or *Wolbachia* spp. (e.g., *Wolbachia* – *wMel*, *Wolbachia* – *wAlbB*, *Wolbachia* – *wMelPop*, *Wolbachia* – *wMelPop-CLA*).

The compositions described herein may include any number of phage, such as at least about any one of 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, 100, or more phage. In some instances, the composition includes phage from one or more (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phage) families, one or more
25 orders (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phage), or one or more species (e.g., 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, 100, or more phage). Compositions including one or more phage are also referred herein as "phage cocktails." Phage cocktails are useful because they allow for targeting of a wider host range of bacteria. Furthermore, they allow for each bacterial strain (i.e. subspecies) to be targeted by multiple orthogonal phages, thereby preventing or significantly delaying the onset of resistance. In some
30 instances, a cocktail includes multiple phages targeting one bacterial species. In some instances, a cocktail includes multiple phages targeting multiple bacterial species. In some instances, a one-phage "cocktail" includes a single promiscuous phage (i.e. a phage with a large host range) targeting many strains within a species.

Suitable concentrations of the phage in the modulating agent described herein depends on
35 factors such as efficacy, survival rate, transmissibility of the phage, number of distinct phage, and/or lysin types in the compositions, formulation, and methods of application of the composition. In some instances, the phage is in a liquid or a solid formulation. In some instances, the concentration of each phage in any of the compositions described herein is at least about any of 10^2 , 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or more pfu/ml. In some instances, the concentration of each phage in any of the
40 compositions described herein is no more than about any of 10^2 , 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 pfu/ml.

In some instances, the concentration of each phage in the composition is any of about 10^2 to about 10^3 , about 10^3 to about 10^4 , about 10^4 to about 10^5 , about 10^5 to about 10^6 , about 10^7 to about 10^8 , about 10^8 to about 10^9 , about 10^2 to about 10^4 , about 10^4 to about 10^6 , about 10^6 to about 10^9 , or about 10^3 to about 10^8 pfu/ml. In some instances, wherein the composition includes at least two types of phages, the

5 concentration of each type of the phages may be the same or different. For example, in some instances, the concentration of one phage in the cocktail is about 10^8 pfu/ml and the concentration of a second phage in the cocktail is about 10^6 pfu/ml.

A modulating agent including a phage as described herein can be contacted with the target host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold level)

10 of phage concentration inside a target host; (b) reach a target level (e.g., a predetermined or threshold level) of phage concentration inside a target host gut; (c) reach a target level (e.g., a predetermined or threshold level) of phage concentration inside a target host bacteriocyte; (d) modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host; or/and (e) modulate fitness of the target host.

15 As illustrated by Examples 5-7 and 28, phages (e.g., one or more naturally occurring phage) can be used as modulating agents that target an endosymbiotic bacterium in an insect host to decrease the fitness of the host (e.g., as outlined herein).

ii. Polypeptides

20 Numerous polypeptides (e.g., a bacteriocin, R-type bacteriocin, nodule C-rich peptide, antimicrobial peptide, lysin, or bacteriocyte regulatory peptide) may be used in the compositions and methods described herein. In some instances, an effective concentration of any peptide or polypeptide described herein may alter a level, activity, or metabolism of one or more microorganisms (as described herein, e.g., a *Wolbachia* spp. or a *Rickettsia* spp.) resident in a host (e.g., a vector of an animal

25 pathogen, e.g., a mosquito, mite, biting louse, or tick), the modulation resulting in a decrease in the host's fitness (e.g., as outlined herein). Polypeptides included herein may include naturally occurring polypeptides or recombinantly produced variants. For example, the polypeptide may be a functionally active variant of any of the polypeptides described herein with at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%,

30 94%, 95%, 96%, 97%, 98%, or 99% identity, e.g., over a specified region or over the entire sequence, to a sequence of a polypeptide described herein or a naturally occurring polypeptide.

A modulating agent comprising a polypeptide as described herein can be contacted with the target host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold level) of concentration inside a target host; (b) reach a target level (e.g., a predetermined or

35 threshold level) of concentration inside a target host gut; (c) reach a target level (e.g., a predetermined or threshold level) of concentration inside a target host bacteriocyte; (d) modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host; or/and (e) modulate fitness of the target host.

The polypeptide modulating agents discussed hereinafter, namely bacteriocins, lysins,

40 antimicrobial peptides, nodule C-rich peptides, and bacteriocyte regulatory peptides, can be used to alter

the level, activity, or metabolism of target microorganisms (e.g., *Rickettsia* or *Wolbochia*) as indicated in the section for decreasing the fitness of host insects (e.g., a vector of an animal pathogen, e.g., a mosquito, a mite, a biting louse, or a tick).

5 (a) *Bacteriocins*

The modulating agent described herein may include a bacteriocin. In some instances, the bacteriocin is naturally produced by Gram-positive bacteria, such as *Pseudomonas*, *Streptomyces*, *Bacillus*, *Staphylococcus*, or lactic acid bacteria (LAB, such as *Lactococcus lactis*). In some instances, the bacteriocin is naturally produced by Gram-negative bacteria, such as *Hafnia alvei*, *Citrobacter*
 10 *freundii*, *Klebsiella oxytoca*, *Klebsiella pneumonia*, *Enterobacter cloacae*, *Serratia plymthicum*, *Xanthomonas campestris*, *Erwinia carotovora*, *Ralstonia solanacearum*, or *Escherichia coli*. Exemplary bacteriocins include, but are not limited to, Class I-IV LAB antibiotics (such as lantibiotics), colicins, microcins, and pyocins. Non-limiting examples of bacteriocins are listed in Table 4.

15 Table 4: Examples of Bacteriocins

Class	Name	Producer	Targets	Sequence
Class I	Nisin	<i>Lactococcus lactis</i>	Active on Gram-positive bacteria: <i>Enterococcus</i> , <i>Lactobacillus</i> , <i>Lactococcus</i> , <i>Leuconostoc</i> , <i>Listeria</i> , <i>Clostridium</i>	ITSISLCTPGCKT GALMGCNMKTA TCHCSIHVSK (SEQ ID NO: 42)
	Epidermin	<i>Staphylococcus epidermis</i>	Gram-positive bacteria	IASKFICTPGCA KTGSFNSYCC (SEQ ID NO: 43)

Class II				
Class II a	Pediocin PA-1	<i>Pediococcus acidilactici</i>	<i>Pediococci</i> , <i>Lactobacilli</i> , <i>Leuconostoc</i> , <i>Brochothrix thermosphacta</i> , <i>Propionibacteria</i> , <i>Bacilli</i> , <i>Enterococci</i> , <i>Staphylococci</i> , <i>Listeria clostridia</i> , <i>Listeria monocytogenes</i> , <i>Listeria innocua</i>	KYYGNGVTCTG KHSCSVDWVK ATTCIINNGAMA WATGGHQGNH KC (SEQ ID NO: 44)
Class II b	Enterocin P	<i>Enterococcus faecium</i>	<i>Lactobacillus sakei</i> , <i>Enterococcus faecium</i>	ATRSYGNGVYC NNSKCWVNWG EAKENIAGIVISG WASGLAGMGH (SEQ ID NO: 45)
Class II c	Lactococcin G	<i>Streptococcus lactis</i>	Gram-positive bacteria	GTWDDIGQGIG RVAYWVGKAM GNMSDVNQAS RINRKKKH (SEQ ID NO: 46)
Class II d	Lactacin-F	<i>Lactobacillus johnsonii</i>	<i>Lactobacilli</i> , <i>Enterococcus faecalis</i>	NRWGDTVLSAA SGAGTGKACK SFGPWGMAICG VGGAAIGGYFG YTHN (SEQ ID NO: 47)

Class III				
Class III a	Enterocin AS-48	<i>Enterococcus faecalis</i>	Broad spectrum: Gram positive and Gram negative bacteria.	MAKEFGIPAAVA GTVLNVEAGG WVTTIVSILTAV GSGGLSLLAAA GRESIKAYLKKE IKKKGKRAVIAW (SEQ ID NO: 48)
Class III b	aureocin A70	<i>Staphylococcus aureus</i>	Broad spectrum: Gram positive and Gram negative bacteria.	MSWLNFLKYIAK YGKKAWSAAWK YKGKVLEWLVN GPTLEWVWQKL KKIAGL (SEQ ID NO: 49)
Class IV	Garvicin A	<i>Lactococcus garvieae</i>	Broad spectrum: Gram positive and Gram negative bacteria.	IGGALGNALNGL GTWANMMNGG GFVNQWQVYA NKGKINQYRPY (SEQ ID NO: 50)
Unclassified	Colicin V	<i>Escherichia coli</i>	Active against <i>Escherichia coli</i> (also closely related bacteria), Enterobacteriaceae	MRTLTLNELDS VSGGASGRDIA MAIGTLGQFV AGGIGAAAGGV AGGAIYDYAST HKPNPAMSPSG LGGTIKQKPEGI PSEAWNYAAGR LCNWSPNNLSD VCL (SEQ ID NO: 51)

In some instances, the bacteriocin is a colicin, a pyocin, or a microcin produced by Gram-negative bacteria. In some instances, the bacteriocin is a colicin. The colicin may be a group A colicin (e.g., uses the Tol system to penetrate the outer membrane of a target bacterium) or a group B colicin (e.g., uses the Ton system to penetrate the outer membrane of a target bacterium). In some instances, the bacteriocin is a microcin. The microcin may be a class I microcin (e.g., < 5 kDa, has post-translational modifications) or a class II microcin (e.g., 5-10 kDa, with or without post-translational modifications). In some instances, the class II microcin is a class IIa microcin (e.g., requires more than one genes to synthesize and assemble functional peptides) or a class IIb microcin (e.g., linear peptides with or without post-translational modifications at C-terminus). In some instances, the bacteriocin is a pyocin. In some instances, the pyocin is an R-pyocin, F-pyocin, or S-pyocin.

In some instances, the bacteriocin is a class I, class II, class III, or class IV bacteriocin produced by Gram-positive bacteria. In some instances, the modulating agent includes a Class I bacteriocin (e.g., lanthionine-containing antibiotics (lantibiotics) produced by a Gram-positive bacteria). The class I bacteriocins or lantibiotic may be a low molecular weight peptide (e.g., less than about 5 kDa) and may possess post-translationally modified amino acid residues (e.g., lanthionine, β -methylanthionine, or dehydrated amino acids).

In some instances, the bacteriocin is a Class II bacteriocin (e.g., non-lantibiotics produced by Gram-positive bacteria). Many are positively charged, non-lanthionine-containing peptides, which unlike lantibiotics, do not undergo extensive post-translational modification. The Class II bacteriocin may belong to one of the following subclasses: "pediocin-like" bacteriocins (e.g., pediocin PA-1 and carnobacteriocin X (Class IIa)); two-peptide bacteriocins (e.g., lactacin F and ABP-118 (Class IIb)); circular bacteriocins (e.g., carnocyclin A and enterocin AS-48 (Class IIc)); or unmodified, linear, non-pediocin-like bacteriocins (e.g., epidermicin NI01 and lactococcin A (Class IId)).

In some instances, the bacteriocin is a Class III bacteriocin (e.g., produced by Gram-positive bacteria). Class III bacteriocins may have a molecular weight greater than 10 kDa and may be heat unstable proteins. The Class III bacteriocins can be further subdivided into Group IIIA and Group IIIB bacteriocins. The Group IIIA bacteriocins include bacteriolytic enzymes which kill sensitive strains by lysis of the cell wall, such as Enterolisin A. Group IIIB bacteriocins include non-lytic proteins, such as Caseicin 80, Helveticin J, and lactacin B.

In some instances, the bacteriocin is a Class IV bacteriocin (e.g., produced by Gram-positive bacteria). Class IV bacteriocins are a group of complex proteins, associated with other lipid or carbohydrate moieties, which appear to be required for activity. They are relatively hydrophobic and heat stable. Examples of Class IV bacteriocins leuconocin S, lactocin 27, and lactocin S.

In some instances, the bacteriocin is an R-type bacteriocin. R-type bacteriocins are contractile bacteriocidal protein complexes. Some R-type bacteriocins have a contractile phage-tail-like structure. The C-terminal region of the phage tail fiber protein determines target-binding specificity. They may attach to target cells through a receptor-binding protein, e.g., a tail fiber. Attachment is followed by sheath contraction and insertion of the core through the envelope of the target bacterium. The core penetration results in a rapid depolarization of the cell membrane potential and prompt cell death. Contact with a single R-type bacteriocin particle can result in cell death. An R-type bacteriocin, for example, may be thermolabile, mild acid resistant, trypsin resistant, sedimentable by centrifugation, resolvable by electron microscopy, or a combination thereof. Other R-type bacteriocins may be complex molecules including multiple proteins, polypeptides, or subunits, and may resemble a tail structure of bacteriophages of the myoviridae family. In naturally occurring R-type bacteriocins, the subunit structures may be encoded by a bacterial genome, such as that of *C. difficile* or *P. aeruginosa* and form R-type bacteriocins to serve as natural defenses against other bacteria. In some instances, the R-type bacteriocin is a pyocin. In some instances, the pyocin is an R-pyocin, F-pyocin, or S-pyocin.

In some instances, the bacteriocin is a functionally active variant of the bacteriocins described herein. In some instances, the variant of the bacteriocin has at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%,

94%, 95%, 96%, 97%, 98%, or 99% identity, e.g., over a specified region or over the entire sequence, to a sequence of a bacteriocin described herein or a naturally occurring bacteriocin.

In some instances, the bacteriocin may be bioengineered, according to standard methods, to modulate their bioactivity, e.g., increase or decrease or regulate, or to specify their target
5 microorganisms. In other instances, the bacteriocin is produced by the translational machinery (e.g. a ribosome, etc.) of a microbial cell. In some instances, the bacteriocin is chemically synthesized. Some bacteriocins can be derived from a polypeptide precursor. The polypeptide precursor can undergo cleavage (e.g., processing by a protease) to yield the polypeptide of the bacteriocin itself. As such, in some instances, the bacteriocin is produced from a precursor polypeptide. In some other instances, the
10 bacteriocin includes a polypeptide that has undergone post-translational modifications, for example, cleavage, or the addition of one or more functional groups.

The bacteriocins described herein may be formulated in a composition for any of the uses described herein. The compositions disclosed herein may include any number or type (e.g., classes) of bacteriocins, such as at least about any one of 1 bacteriocin, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, 100, or
15 more bacteriocins. Suitable concentrations of each bacteriocin in the compositions described herein depends on factors such as efficacy, stability of the bacteriocin, number of distinct bacteriocin types in the compositions, formulation, and methods of application of the composition. In some instances, each bacteriocin in a liquid composition is from about 0.01 ng/ml to about 100 mg/mL. In some instances, each bacteriocin in a solid composition is from about 0.01 ng/g to about 100 mg/g. In some instances, wherein
20 the composition includes at least two types of bacteriocins, the concentration of each type of the bacteriocins may be the same or different. In some instances, the bacteriocin is provided in a composition including a bacterial cell that secretes the bacteriocin. In some instances, the bacteriocin is provided in a composition including a polypeptide (e.g., a polypeptide isolated from a bacterial cell).

Bacteriocins may neutralize (e.g., kill) at least one microorganism other than the individual
25 bacterial cell in which the polypeptide is made, including cells clonally related to the bacterial cell and other microbial cells. As such, a bacterial cell may exert cytotoxic or growth-inhibiting effects on a plurality of microbial organisms by secreting bacteriocins. In some instances, the bacteriocin targets and kills one or more species of bacteria resident in the host via cytoplasmic membrane pore formation, cell wall interference (e.g., peptidoglycanase activity), or nuclease activity (e.g., DNase activity, 16S rRNase
30 activity, or tRNase activity).

In some instances, the bacteriocin has a neutralizing activity. Neutralizing activity of bacteriocins may include, but is not limited to, arrest of microbial reproduction, or cytotoxicity. Some bacteriocins have cytotoxic activity, and thus can kill microbial organisms, for example bacteria, yeast, algae, and the like. Some bacteriocins can inhibit the reproduction of microbial organisms, for example bacteria, yeast, algae,
35 and the like, for example by arresting the cell cycle.

In some instances, the bacteriocin has killing activity. The killing mechanism of bacteriocins is specific to each group of bacteriocins. In some instances, the bacteriocin has narrow-spectrum bioactivity. Bacteriocins are known for their very high potency against their target strains. Some bacteriocin activity is limited to strains that are closely related to the bacteriocin producer strain (narrow-

spectrum bioactivity). In some instances, the bacteriocin has broad-spectrum bioactivity against a wide range of genera.

In some instances, bacteriocins interact with a receptor molecule or a docking molecule on the target bacterial cell membrane. For example, nisin is extremely potent against its target bacterial strains, showing antimicrobial activity even at a single-digit nanomolar concentration. The nisin molecule has been shown to bind to lipid II, which is the main transporter of peptidoglycan subunits from the cytoplasm to the cell wall

In some instances, the bacteriocin has anti-fungal activity. A number of bacteriocins with anti-yeast or anti-fungal activity have been identified. For example, bacteriocins from *Bacillus* have been shown to have neutralizing activity against some yeast strains (see, for example, Adetunji and Olaoye, *Malaysian Journal of Microbiology* 9:130-13, 2013). In another example, an *Enterococcus faecalis* peptide has been shown to have neutralizing activity against *Candida* species (see, for example, Shekh and Roy, *BMC Microbiology* 12:132, 2012). In another example, bacteriocins from *Pseudomonas* have been shown to have neutralizing activity against fungi, such as *Curvularia lunata*, *Fusarium species*, *Helminthosporium* species, and *Biopolaris* species (see, for example, Shalani and Srivastava, *The Internet Journal of Microbiology* Volume 5 Number 2, 2008). In another example, botrycidin AJ1316 and alirin B1 from *B. subtilis* have been shown to have antifungal activities.

A modulating agent including a bacteriocin as described herein can be contacted with the target host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold level) of bacteriocin concentration inside a target host; (b) reach a target level (e.g., a predetermined or threshold level) of bacteriocin concentration inside a target host gut; (c) reach a target level (e.g., a predetermined or threshold level) of bacteriocin concentration inside a target host bacteriocyte; (d) modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host; or/and (e) modulate fitness of the target host.

As illustrated by Examples 8, 9, or 16, bacteriocins (e.g., colA or nisin) can be used as modulating agents that target an endosymbiotic bacterium in an insect host to decrease the fitness of the host (e.g., as outlined herein).

(b) Lysins

The modulating agent described herein may include a lysin (e.g., also known as a murein hydrolase or peptidoglycan autolysin). Any lysin suitable for inhibiting a bacterium resident in the host may be used. In some instances, the lysin is one that can be naturally produced by a bacterial cell. In some instances, the lysin is one that can be naturally produced by a bacteriophage. In some instances, the lysin is obtained from a phage that inhibits a bacterium resident in the host. In some instances, the lysin is engineered based on a naturally occurring lysin. In some instances, the lysin is engineered to be secreted by a host bacterium, for example, by introducing a signal peptide to the lysin. In some instances, the lysin is used in combination with a phage holin. In some instances, a lysin is expressed by a recombinant bacterium host that is not sensitive to the lysin. In some instances, the lysin is used to inhibit a Gram-positive or Gram-negative bacterium resident in the host.

The lysin may be any class of lysin and may have one or more substrate specificities. For example, the lysin may be a glycosidase, an endopeptidase, a carboxypeptidase, or a combination thereof. In some instances, the lysin cleaves the β -1-4 glycosidic bond in the sugar moiety of the cell wall, the amide bond connecting the sugar and peptide moieties of the bacterial cell wall, and/or the peptide bonds between the peptide moieties of the cell wall. The lysin may belong to one or more specific lysin groups, depending on the cleavage site within the peptidoglycan. In some instances, the lysin is a *N*-acetyl- β -D-muramidase (e.g., lysozyme), lytic transglycosylase, *N*-acetyl- β -D-glucosaminidase, *N*-acetylmuramyl-L-alanine amidase, L,D-endopeptidase, D,D-endopeptidase, D,D-carboxypeptidase, L,D-carboxypeptidase, or L,D-transpeptidase. Non-limiting examples of lysins and their activities are listed in Table 5.

Table 5: Examples of Lysins

Target Bacteria	Producer	Lysins	Activity	Sequence
<i>S. pneumoniae</i>	Cp1	Cpl-1	Muramidase	MVKKNDLFVDVSSH NGYDITGILEQMGTT NTIIKISESTTYLNPCL SAQVEQSNPIGFYHF ARFGGDVAEAAREA QFFLDNVPMQVKYLV LDYEDDPGDAQAN TNACLRFMQMIADAG YKPIYYSYKPFTHDN VDYQQILAQFPNSLW IAGYGLNDGTANFEY FPSMDGIRWWQYSS NPFDKNIVLLDDEED DKPKTAGTWKQDSK GWWFRRNNGSFPY NKWEKIGGVWYYFD SKGYCLTSEWLKDN EKWYYLKDNAMAT GWVLVGSEWYYMD DSGAMVTGWVKYKN NWYYMTNERGNMV SNEFIKSGKGWYFM NTNGELADNPSFTKE PDGLITVA (SEQ ID NO: 52)
<i>S. pneumoniae</i>	Dp-1	Pal	Amidase	MGVDIEKGVAWMQA RKGRVSYSMDFRDG PDSYDCSSSMYYAL

				RSAGASSAGWAVNT EYMHAWLIENGYELI SENAPWDAKRGDIFI WGRKGASAGAGGH TGMFIDSDNIIHCNYA YDGISVNDHDERWY YAGQPYYYVYRLTNA NAQPAEKKLGWQKD ATGFWYARANGTYP KDEFEYIEENKSWFY FDDQGYMLAEKWLK HTDGNWYWFDRDG YMATSWKRIGESWY YFNRDGSMVTGWIK YYDNWYYCDATNGD MKSNAFIRYNDGWY LLLPDGRLADKPQFT VEPDGLITAKV (SEQ ID NO: 53)
<i>S. pyogenes</i>	C1	C1	Amidase	N/A
<i>B. anthracis</i>	γ	PlyG	Amidase	MEIQKKLVDPISKYGT KCPYTMKPKYITVHN TYNDAPAENEVSIMI SNNNEVSFHIAVDDK KAIQGIPLERNAWAC GDGNGSGNRQSSIV EICYSKSGGDRYYKA EDNAVDVVRQLMSM YNIPINVRTHQSWS GKYCPHRMLAEGRW GAFIQKVKNNGNVATT SPTKQNIQSGAFSPY ETPDVMGALTSLKMT ADFILQSDGLTYFISK PTSDAQLKAMKEYLD RKGWWYEVK (SEQ ID NO: 54)
<i>B. anthracis</i>	Ames prophage	PlyPH	Amidase	N/A
<i>E. faecalis</i> and <i>E. faecium</i>	Phi1	PlyV12	Amidase	N/A

<i>S. aureus</i>	ΦMR11	MV-L	Endopeptidase and amidase	MQAKLTKEFIEWLK TSEGKQFNVDLWYG FQCFDYANAGWKVL FGLLLKGLGAKDIPFA NNFDGLATVYQNTF DFAQPGDMVVFGS NYGAGYGHVAVVIE ATLDYIIVYEQNWL GGWTDRIEQPGWG WEKVTRRQHAYDFP MWFIRPNFKSETAPR SIQSPTQASKKETAK PQPKAVELKIIDVVK GYDLPKRGGNPKGIV IHNDAGSKGATAEAY RNGLVNAPLSRLEAG IAHSYVSGNTVWQAL DESQVGWHTANQLG NKYYYGIEVCQSMG ADNATFLKNEQATFQ ECARLLKKWGLPAN RNTIRLHNEFTSTSC PHRSSVLHTGFDPVT RGLLPEDKQLQLKDY FIKQIRVYMDGKIPVA TVSNESSASSNTVKP VASAWKRNKYGTYY MEENARFTNGNQPIT VRKIGPFLSCPVA YQ FQPGGYCDYTEVML QDGHVWVGWYTWEG QRYYLPIRTWNGSAP PNQILGDLWGEIS (SEQ ID NO: 55)
<i>S. pyogenes</i>	C1	PlyC	Amidase	N/A
<i>S. agalactiae</i>	B30	GBS lysin	Muramidase and endopeptidase	MVINIEQAIWMA SR KGKVTYSMDYR NGP SSYDCSSSVYF ALRS AGASDNGWAVN TEY EHDWLIKNGYV LIAE

				NTNWNAQRGDIFIW GKRASAGAFGHTG MFVDPDNIHCNYGY NSITVNNHDEIWGYN GQPYVYAYRYSKQ SNAKVDNKSVMKFE KELDVNTPLSNSNMP YYEATISEDYVESK PDVNSTDKELLVAGT RVRVYEKVKGWARI GAPQSNQWVEDAYL IDATDM (SEQ ID NO: 56)
<i>S. aureus</i>	P68	Lys16	Endopeptidase	N/A
<i>S. aureus</i>	K	LysK	Amidase and endopeptidase	MAKQAEINKRLDAY AKGTVDSPYRVKAT SYDPSFGVMEAGID ADGYYHAQCQDLITD YVLWLTDNKVRTWG NAKDQIKQSYGTGFK IHENKPSTVPKKGWI AVFTSGSYEQWGH GIVYDGGNTSTFTILE QNWNGYANKKPTKR VDNYYGLTHFIEIPVK AGTTVKKETAKKSAS KTPAPKKKATLKVSK NHINYTMDKRGKKPE GMVIHNDAGRSSGQ QYENSLANAGYARY ANGIAHYGSEGYV WEIDAQNKIAWHTG DGTGANSNFRFAGI EVCQSMSASDAQFL KNEQAVFQFTAEEKFK EWGLTPNRKTVRLH MEFVPTACPHRSMV LHTGFNPVTQGRPS QAIMNKLKDYFIKQIK NYMDKGTSSSTVVK

				DGKTSSASTPATRPV TGSWKKNQYGTWYK PENATFVNGNQPIVT RIGSPFLNAPVGGNL PAGATIVYDEVCIQA GHIWIGYNAYNGNRV YCPVRTCQGVPPNQI PGVAWGVFK (SEQ ID NO: 57)
<i>L. monocytogenes</i>	A118	Ply118	Amidase	MTSYYYSRSLANVNK LADNTKAAARKLLDW SENGIEVLIYETIRTK EQQAANVNSGASQT MRSYHLVGQALDFV MAKGKTVDWGAYRS DKGKKFVAKAKSLGF EWGGDWSGFVDNP HLQFNKYGYGTDTF GKGASTSNSSKPSA DTNTNSLGLVDYMN NKLDSSFANRKKLAT SYGIKNYSGTATQNT TLLAKLKAGKPHTPA SKNTYYTENPRKVKT LVQCDLYKSVDFTTK NQTGGTFPPGTVFTI SGMGKTKGGTPRLK TKSGYYLTANTKFVK KI (SEQ ID NO: 58)
<i>L. monocytogenes</i>	A511	Ply511	Amidase	MVKYTVENKIIAGLPK GKLKGANFVIAHETA NSKSTIDNEVSymTR NWKNAFVTHFVGGG GRVVQVANVNYVSW GAGQYANSYSYAQV ELCRTSNATTFKKDY EVYCQLLVDLAKKAG IPITLDGSKTSDKGI KSHKWVADKLGGTT HQDPYAYLSSWGISK

				AQFASDLAKVSGGG NTGTAPAKPSTPAPK PSTPSTNLDKLGGLVD YMNAKKMDSSYSNR DKLAKQYGIANYSGT ASQNTTLLSKIKGGA PKPSTPAPKPSTSTA KKIYFPPNKGWWSVY PTNKAPVKANAIGAIN PTKFGGLTYTIQKDR GNGVYEIQTDQFGR VQVYGAPSTGAVIKK (SEQ ID NO: 59)
<i>L. monocytogenes</i>	A500	Ply500	Endopeptidase	MALTEAWLIEKANRK LNAGGMYKITSKTR NVIKKMAKEGIYLCVA QGYRSTAEQNALYA QGRTPGAIVTNAKG GQSNHNYGVAVDLC LYTNDGKDVWESTT SRWKKVVAAMKAEG FKWGGDWKSFKDYP HFELCDAVSGEKIPA ATQNTNTNSNRYEG KVIDSAPLLPKMDFK SSPFRMYKVGTEFLV YDHNQYWYKTYIDD KLYYMYKSFCDVVAK KDAKGRIKVRISAK DLRIPVWNNIKLNSG KIKWYAPNVKLAWYN YRRGYLELWYPNDG WYYTAEYFLK (SEQ ID NO: 60)
<i>S. pneumoniae</i>	Φ Dp-1	Pal, S	Endopeptidase and amidase	N/A
<i>S. agalactiae</i>	LambdaSa1 prophage	LambdaSa1	Glycosidase	MVINIEQAIWMA SR KGKVTYSMDYRNGP SSYDCSSSVYFALRS AGASDNGWAVNTEY EHDWLIKNGYVLIAE

				NTNWNAQRGDIFIW GKRASAGAFGHTG MFVDPDNIHCNYGY NSITVNNHDEIWGYN GQPYVYAYRYARKQ SNAKVDNQSVVSKF EKELDVNTPLSNSNM PYYEATISEDYYVES KPDVNSTDKELLVAG TRVRVYEKVKGWARI GAPQSNQWVEDAYL IDATDM (SEQ ID NO: 61)
<i>S. agalactiae</i>	LambdaSa2 prophage	LambdaSa2	Glycosidase and endopeptidase	MEINTEIAIAWMSAR QGKVSYSMDYRDGP NSYDCSSSVYYALRS AGASSAGWAVNTEY MHDWLIKNGYELIAE NVDWNAVRGDIW GMRGHSSGAGGHV VMFIDPENIIHCNWA NNGITVNNYNQTA SGWMYCYVYRLKSG ASTQGKSLDTLVKET LAGNYGNGEARKAV LGNQYEAVMSVINGK TTTNQKTVDQLVQEV IAGKHGNGEARKKSL GSQYDAVQKRVELL KKQPSEPFKAQEVN KPTETKTSQTELTGQ ATATKEEGDLSFNGT ILKKAVLDKILGNCKK HDILPSYALTILHYEG LWGTSVAVGKADNNW GGMTWTGQGNRPS GVTVTQGSARPSNE GGHYMHYASVDDFL TDWFFYLLRAGGSYK VSGAKTFSEAIKGMF

				KVGGAVDYAASGF DSYIVGASSRLKAIEA ENGSLDKFDKATDIG DGSKDKIDITIEGIEVT INGITYELTKKPV (SEQ ID NO: 62)
<i>S. uberis</i>	(ATCC700407) prophage	Ply700	Amidase	MTDSIQEMRKLQSIP VRYDMGDRYGNAD RDGRIEMDCSSAVSK ALGISMTNNTETLQQ ALPAIGYGKIHDADV GTFDMQAYDVIIWAP RDGSSSLGAFGHVLI ATSPTTAIHCNYGSD GITENDYNYWEING RPREIVFRKGVQTQ ATVTSQFERELDVNA RLTVSDKPYEATLS EDYYVEAGPRIDSQD KELIKAGTRVRVYEK LNGWSRINHPESAQ WVEDSYLVDATEM (SEQ ID NO: 63)
<i>S. suis</i>	SMP	LySMP	Glycosidase and endopeptidase	N/A
<i>B. anthracis</i>	Bcp1	PlyB	Muramidase	N/A
<i>S. aureus</i>	Phi11 and Phi12	Phi11 lysin	Amidase and endopeptidase	MQAKLTKNEFIEWLK TSEGKQFNVDLWYG FQCFDYANAGWKVL FGLLLKGLGAKDIPFA NNFDGLATVYQNTP DFLAQPGDMVVFGS NYGAGYGHVAWVIE ATLDYIIVYEQNWL GGWTDGIEQPGWG WEKVTRRRQHAYDFP MWFIRPNFKSETAPR SVQSPTQAPKKETAK PQPKAVELKIIDVVK GYDLPKRGSNPKGIV IHNDAGSKGATAEAY

				RNGLVNAPLSRLEAG IAHSYVSGNTVWQAL DESQVGWHTANQIG NKYYYGIEVCQSMG ADNATFLKNEQATFQ ECARLLKKWGLPAN RNTIRLHNEFTSTSC PHRSSVLHTGFDPT RGLLPEDKRLQLKDY FIKQIRAYMDGKIPVA TVSNESSASSNTVKP VASAWKRNKYGTYY MEESARFTNGNQPIT VRKVGPFLLSCPVG QFQPGGYCDYTEVM LQDGHVWVGYTWE GQRYYLPIRTWNGS APPNQILGDLWGEIS (SEQ ID NO: 64)
<i>S. aureus</i>	ΦH5	LysH5	Amidase and endopeptidase	MQAKLTKKEFIEWLK TSEGKQYNADGWYG FQCFDYANAGWKAL FGLLLKGVGAKDIPF ANNFDGLATVYQNT DFLAQPGDMVVFGS NYGAGYGHVAVVIE ATLDYIIVYEQNWLG GGWTDGVQQPGSG WEKVTRRQHAYDFP MWFIRPNFKSETAPR SVQSPTQASKKETAK PQPKAVELKIIDVVK GYDLPKRGSNPNFIVI HNDAGSKGATAEAY RNGLVNAPLSRLEAG IAHSYVSGNTVWQAL DESQVGWHTANQIG NKYGYGIEVCQSMG ADNATFLKNEQATFQ ECARLLKKWGLPAN

				RNTIRLHNEFTSTSC PHRSSVLHTGFDPVT RGLLPEDKRLQLKDY FIKQIRAYMDGKIPVA TVSNDSSASSNTVKP VASAWKRNKYGTYY MEESARFTNGNQPIT VRKVGPFLLSCPVG QFQPGGYCDYTEVM LQDGHVWVGYTWE GQRYLPIRTWNGS APPNQILGDLWGEIS (SEQ ID NO: 65)
<i>S. warneri</i>	ΦWMY	LysWMY	Amidase and endopeptidase	MKTAKAQAKSWINSKI GKGIDWDGMYGYQC MDEAVDYIHHVTDGK VTMWGNAIDAPKNN FQGLCTVYTNTPEFR PAYGDVIVWSYGTFA TYGHIAIVVNPDPYG DLQYITVLEQNWNGN GIYKTEFATIRTHDYT GVSHFIRPKFADEVK ETAKTVNKL SVQKKI VTPKNSVERIKNYVK TSGYINGEHYELYNR GHKPKGVIHNTAGT ASATQEGQRLTNMT FQQLANGVAHVYIDK NTIYETLPEDRIAWHV AQQYGNTEFYGIEVC GSRNTDKEQFLANE QVAFQEAARRLKSW GMRANRNTVRLHHT FSSTECPDMSMLLHT GYSMKNGKPTQDITN KCADYFMKQINAYID GKQPTSTVVGSSSS NKLKAKNKDKSTGW NTNEYGTLWKKKEHA

				TFTCGVRQGIVTRTT GPFTSCPQAGVLYY GQSVNYDTVCKQDG YVWISWTTSDGYDV WMPIRTWDRSTDKV SEIWGTIS (SEQ ID NO: 66)
<i>Streptococci (GBS)</i>	ΦNCTC 11261	PlyGBS	Muramidase and endopeptidase	MATYQEYKSRSENGN AYDIDGSFGAQCWD GYADYCKYLGLPYA NCTNTGYARDIWEQ RHENGILNYFDEVEV MQAGDVAIFMVVDG VTPYSHVAIFDSDAG GGYGWFLGQNQGG ANGAYNIVKIPYSATY PTAFRPKVFKNAVTV TGNIGLNKGDYFIDV SAYQQADLTTCQQ AGTTKTIKVSSEIAW LSDRHQQQANTSDPI GYYHFGRFGGDSAL AQREADLFLSNLPSK KVSYLVIDYEDSASA DKQANTNAVIAFMDK IASAGYKPIYYSYKPF TLNNIDYQKIIAKYPN SIWIAGYPDYEVRTE PLWEFFPSMDGVRW WQFTSVGVAGGLDK NIVLLADDSSKMDIPK VDKPQELTFYQKLAT NTKLDNSNPYYEAT LSTDYYVESKPNASS ADKEFIKAGTRVRVY EKVNGWSRINHPES AQWVEDSYLVNATD M (SEQ ID NO: 67)
<i>C. perfringens</i>	Φ3626	Ply3626	Amidase	N/A
<i>C. difficile</i>	ΦCD27	CD27 lysin	Amidase	N/A

<i>E. faecalis</i>	Φ1	PlyV12	Amidase	N/A
<i>A. naeslundii</i>	ΦAv-1-	Av-1 lysin	Putative amidase/murami dase	N/A
<i>L. gasseri</i>	ΦgaY	LysgaY	Muramidase	N/A
<i>S. aureus</i>	ΦSA4	LysSA4	Amidase and endopeptidase	N/A
<i>S. haemolyticus</i>	ΦSH2	SH2	Amidase and endopeptidase	N/A
<i>B. thuringiensis</i>	ΦBtCS33	PlyBt33	Amidase	N/A
<i>L. monocytogenes</i>	ΦP40	PlyP40	Amidase	N/A
<i>L. monocytogenes</i>	ΦFWLLm3	LysZ5	Amidase	MVKYTVENKIIAGLPK GKLKGANFVIAHETA NSKSTIDNEVS YMTR NWQNAFVTHFVGGG GRVVQVANVNYVSW GAGQYANSYSYAQV ELCRTSNATTFKKDY EVCQQLLDLAKKAG IPITLDGSKTSDKGI KSHKWVADKLGTT HQDPYAYLSSWGISK AQFASDLAKVSGGG NTGTAPAKPSTPSTN LDKLGLVDYMNACK MDSSYSNRAKLAKQ YGIANYS GTASQNTT LLSKIKGGAPKPSTP APKPSTSTAKKIYFPP NKG NWSVYPTNKAP VKANAIGAINPTKFG GLTYTIQKDRGNGVY EIQTDQFGRVQVYGA PSTGAVIKK (SEQ ID NO: 68)
<i>B. cereus</i>	ΦBPS13	LysBPS13	Amidase	MAKREKYIFDVEAEV GKAAKSIKSLEAELS KLQKLNKEIDATGGD RTEKEMLATLKAKE VNAEYQKMQRILKDL

				SKYSGKVS RKEFND SKVINNAKTSVQGGK VTDSFGQMLKNMER QINSV NKQFDNHRKA MVDRGQQYTPHLKT NRKDSQGN SNPSMM GRNKSTTQDMEKAV DKFLNGQNEATTGLN QALYQLKEISKLNRR SESLRRASASGYM SFQQYSNFTGDRRT VQQTYGGLKTANRE RVLELSGQATGISKE LDRLNSKKGLTAREG EERKKLMRQLEGIDA ELTARKKLNSSLDET TSNMEKFNQSLVDA QVSVKPERGTMRGM MYERAPAIALAIGGAI TATIGKLYSEGGNHS KAMRPDEMYVGQQT GAVGANWRPNRTAT MRSGLG NHLGFTGQ EMMEFQSNYLSANG YHGAEDMKAATTGQ ATFARATGLGSDEVK DFFNTAYRSGGIDGN QTKQFQNAFLGAMK QSGAVGREKDQLKA LNGILSSMSQNRTVS NQDMMRTVGLQSAI SSSGVASLQGTKGG ALMEQLDNGIREGFN DPQMRVLFQGQTKY QGMGGRAALRKQM EKGISDPDNLNTLIDA SKASAGQDPAEQAE VLATLASKMGVNMS SDQARGLIDLQPSGK LTKENIDKVMKEGLK
--	--	--	--	---

				EGSIESAKRDKAYSE SKASIDNSSEAATAK QATELNDMGSKLRQ ANAALGGLPAPLYTA IAAVVAFTAAVAGSA LMFKGASWLKGGMA SKYGGKGGKGGKG GGTGGGGGAGGAA ATGAGAAAGAGGVG AAAAGEVGAGVAAG GAAAGAAAGGSKLA GVGKGFMMKGAGKLM LPLGILMGASEIMQA PEEAKGSAIGSAVGG IGGGIAGGAATGAIA GSFLGPIGTAVGGIA GGIAGGFAGSSLGET IGGWFDSPKEDAS AADKAKADASAAALA AAAGTSGAVGSSAL QSQMAQGITGAPNM SQVGSMASALGISSG AMASALGISSGQENQ IQTMTDKENTNTKKA NEAKKGDNLSYERE NISMVERVLTRAEQIL AQARAQNGIMGVGG GGTAGAGGGINGFT GGGKLQFLAAGQKW SSSNLQQHDLGFTD QNLTAEDLDKWIDSK APQGSMMRGMGAT FLKAGQEYGLDPRYL IAHAAEESGWGTSKI ARDKGNFFGIGAFDD SPYSSAYEFKDGTS AAERGIMGGAKWISE KYYGKGNTTLDKMK AAGYATNASWAPNIA SIMAGAPTGSGSGN
--	--	--	--	--

				VTATINVNVKGDEKV SDKLKNSSDMKKAG KDIGSLLGFYSREMTI A (SEQ ID NO: 69)
<i>S. aureus</i>	ΦGH15	LysGH15	Amidase and endopeptidase	MAKTQAEINKRLDAY AKGTVDSPYRIKKAT SYDPSFGVMEAGID ADGYYHAQCQDLITD YVLWLTDNKVRTWG NAKDQIKQSYGTGFK IHENKPSTVPPKGWI AVFTSGSYQQWGH GIVYDGGNTSTFTILE QNWNGYANKKPTKR VDNYYGLTHFIEIPVK AGTTVKKETAKKSAS KTPAPKKKATLKVSK NHINYTMDKRGKKPE GMVIHNDAGRSSGQ QYENSLANAGYARY ANGIAHYGSEGYV WEAIDAKNQIAWHTG DGTGANSNFRFAGI EVCQSMSASDAQFL KNEQAVFQFTAEEKFK EWGLTPNRKTVRLH MEFVPTACPHRSMV LHTGFNPVTQGRPS QAIMNKLKDYFIKQIK NYMDKGTSSSTVVK DGKTSSASTPATRPV TGSWKKNQYGTWYK PENATFVNGNQPIVT RIGSPFLNAPVGGNL PAGATIVYDEVCIQA GHIWIGYNAYNGDRV YCPVRTCQGVPPNHI PGVAWGVFK (SEQ ID NO: 70)

<i>S. aureus</i>	ΦvB SauS-PLA88	HydH5	Endopeptidase and glycosidase	N/A
<i>E. faecalis</i>	ΦF168/08	Lys168	Endopeptidase	N/A
<i>E. faecalis</i>	ΦF170/08	Lys170	Amidase	N/A
<i>S. aureus</i>	ΦP-27/HP	P-27/HP	Nonspecified	N/A
<i>C. perfringens</i>	ΦSM101	Psm	Muramidase	N/A
<i>C. sporogenes</i>	Φ8074-B1	CS74L	Amidase	MKIGIDMGHTLSGAD YGVVGLRPESVLTRE VGTKVIYKQLKLGHV VVNCTVDKASSVSES LYTRYRANQANVDL FISIHFNATPGGTGTE VYTYAGRQLGEATRI RQEFKSLGLRDRGT KDGSGLAVIRNTKAK AMLVECCFCNDPND MKLYNSESFSNAIVK GITGKLPNGESGNNN QGGNKVKAVVIYNEG ADRRGAEYLADYLN CPTISNSRTFDYSCV EHVYAVGGKKEQYT KYLKTLLSGANRYDT MQQILNFINGGK (SEQ ID NO: 71)
<i>S. typhimurium</i>	ΦSPN1S	SPN1S	Glycosidase	MDINQFRRASGINEQ LAARWFPHITTAMNE FGITKPDDQAMFIAQ VGHESGGFTRLQEN FNYSVNLGSGFIRAG RITPDQANALGRKTY EKSLPLERQRAIANL VYSKRMGNNPGDGG WNYRGRGLIQITGLN NYRDCGNGLKVDLV AQPELLAQDEYAARS AAWFFSSKGCMKYT GDLVRVTQIINGGQN

				GIDDRRTRYAAARKV LAL (SEQ ID NO: 72)
<i>C. michiganensis</i>	ΦCMP1	CMP1	Peptidase	N/A
<i>C. michiganensis</i>	ΦCN77	CN77	Peptidase	MGYWGYPNGQIPND KMALYRGCLLRADAA AQAYALQDAYTRAT GKPLVILEGYRDLTR QKYLRLNLYLSGRGNI AAVPGLSNHGWGLA CDFAAPLNSSGSEEH RWMRQNAPLFGFD WARGKADNEPWHW EYGNVPVSRWASLD VTPIDRNDMADITEG QMQRIAVILLDEIQT PLGPRLVKHALGDAL LLGQANANSIAEVPD KTWDVLVDHPLAKN EDGTPLKVRLGDVAK YEPLEHQNTRDAIAK LGTQFTDKQLATIG AGVKPIDEASLVKKIV DGVRALFGRAAA (SEQ ID NO: 73)
<i>A. baumannii</i>	ΦAB2	LysAB2	Glycosidase	MILTKDGFISIIRNELF GGKLDQTQVDAINFI VAKATESGLTYPEAA YLLATIHETGLPSGY RTMQPIKEAGSDSYL RSKKYYPYIGYGYVQ LTWKENYERIGKLIG VDLIKNPEKALEPLIAI QIAIKGMLNGWFTGV GFRRKRVPVSKYNKQ QYVAARNIINGKDKA ELIAKYAIIIFERALRSL (SEQ ID NO: 74)
<i>B. cereus</i>	ΦB4	LysB4	Endopeptidase	MAMALQTLIDKANRK LNVSGMRKDVADRT

				RAVITQMHAQGIYICV AQGFRSFAEQNALY AQGRTPGSIVTNAR GGQSNHNYGVAVDL CLYTQDGSDDVIWTV GNFRKVIAAMKAQGF KWGGDWVSFKDYP HFELYDVVGGQKPP ADNGGAVDNGGGS GSTGGSGGGSTGG GSTGGGYDSSWFTK ETGTFVTNTSIKLR PFTSADVIATLPAGSP VNYNGFGIEYDGYV WIRQPRSNYGYLA TGESKGGKRQNYW GTFK (SEQ ID NO: 75)
<i>P. aeruginosa</i>	ΦKMV	KMV45	Nonspecified	N/A
<i>C. tyrobutyricum</i>	ΦCTP1	Ctp1l	Glycosidase	MKKIADISNLNGNVD VKLLFNLGYIGIIAKAS EGGTFVDKYYKQNY TNTKAQKITGAYHF ANFSTIAKAQQEANF FLNCIAGTTPDFVVD LEQQCTGDITDACLA FLNIVAKKFKCVVYC NSSFIKEHLNSKICAY PLWIANYGVATPAFT LWTKYAMWQFTEKG QVSGISGYIDFSYITD EFIKYIKGEDEVENLV VYNDGADQRAAEYL ADRLACPTINNARKF DYSNVKNVYAVGGN KEQYTSYLTTLIAGST RYTTMQAVLDYIKNL K (SEQ ID NO: 76)
<i>P. aeruginosa</i>	ΦEL	EL188	Transglycosylase	N/A
<i>P. aeruginosa</i>	ΦKZ	KZ144	Transglycosylase	N/A

<i>S. aureus</i>		Ply187	Cell Wall Hydrolase	MALPKTGKPTAKQVV DWAINLIGSGVDVDG YYGRQCWDLPNYIF NRYWNFKTPGNARD MAWYRYPEGFKVFR NTSDFVPKPGDIAVW TGGNYNWNTWGHT GIVVGPSTKSYFYVS DQNWNNNSYVGSP AAKIKHSYFGVTHFV RPAYKAEPKPTPPAQ NNPAPKDPEPSKKP ESNKPIYKVVTILFT TAHIEHVKANRFVHYI TKSDNHNNKPNKIVIK NTNTALSTIDVYRYR DELDKDEIPHFFVDR LNVWACRPIEDSING YHDSVLSITETRTAL SDNFKMNEIECLSLA ESILKANNKKMSASNI IVDNKAWRTFKLHTG KDSLKSSSFTSKDYQ KAVNELIKLFNDKDKL LNNKPKDVVERIRIRT IVKENTKFVPSELKPR NNIRDKQDSKIDRVIN NYTLKQALNIQYKLN PKPQTSNGVSWYNA SVNQIKSAMDTTKIFN NNVQVYQFLKLNQY QGIPVDKLNKLLVGK GTLANQGHAFADGC KKYNINEIYLIAHRFLE SANGTSFFASGKTGV YNYFGIGAFDNNPNN AMAFARSHGWTSP KAIIGGAEFVGKGYF NVGQNTLYRMRWNP QKPGTHQYATDISW
------------------	--	--------	------------------------	---

				AKVQAQMISAMYKEI GLTGDYFIYDQYKK (SEQ ID NO: 77)
<i>P. uorescens</i>	ΦOBP	OBPgp279	Glycosidase	N/A
<i>L. monocytogenes</i>	ΦP35	PlyP35	Amidase	MARKFTKAELVAKAE KKVGGLKPDVKKAVL SAVKEAYDRYGIGIIV SQGYRSIAEQNGLYA QGRTPGNIVTNAKG GQSNHNFVAVDFAI DLIDDGKIDSWQPSA TIVNMMKRRGFKWG GDWKSFTDLPHFEA CDWYRGERKYKVDV SEWKKKENINIVIKDV GYFQDKPQFLNSKS VRQWKHGTKVKLTK HNSHWYTGTVKDG KSVRGYIYHSMKVT SKNSDGSVNATINAH AFCWDNKKLNGGDFI NLKRGFKGITHPASD GFYPLYFASRKCTFYI PRYMFDIKK (SEQ ID NO: 78)
<i>L. fermentum</i>	ΦPYB5	Lyb5	Muramidase	N/A
<i>S. pneumoniae</i>	ΦCP-7	Cpl-7	Muramidase	MVKKNDLFDVASH QGYDISGILEEAGTT NTIHKVSESTSYLNP LSAQVSQSNPIGFYH FAWFGGNEEEAEAE ARYFLDNVPTQVKYL VLDYEDHASASVQR NTTACLRFMQIIAEAG YTPYYSYKPFTLDNV DYQQILAQFPNSLWI AGYGLNDGTANFEY FPSMDGIRWWQYSS NPFDKNIVLLDDEKE DNINNENTLKSLTTVA

				NEVIQGLWGNGQER YDSLANAGYDPQAV QDKVNEILNAREIADL TTVANEVIQGLWGN GQERYDSLANAGYD PQAVQDKVNEILNAR EIADLTTVANEVIQGL WGNGQERYDSLANA GYDPQAVQDKVNEL LS (SEQ ID NO: 79)
<i>P. chlororaphis</i> 201	Φ2-1	201φ2- 1gp229	Glycosidase	N/A
<i>S. enterica</i>	ΦPVP-SE1)	PVP- SE1gp146	Glycosidase	N/A
<i>Corynebacterium</i>	ΦBKF20	BKF20	Amidase	N/A
<i>E. faecalis</i>	ΦEFAP-1	EFAL-1	Amidase	MKLKGILLSVVTTFGL LFGATNVQAYEVNNE FNLQPWEGSQQLAY PNKIILHETANPRATG RNEATYMKNNWFNA HTTAIVGDGGIVYKV APEGNSVWGAGNAN PYAPVQIELQHTNDP ELFKANYKAYVDYTR DMGKKFGIPMTLDQ GGSLWEKGVVSHQ WVTDFVWGDHTDPY GYLAKMGISKAQLAH DLANGVSGNTATPTP KPDKPKPTQPSKPSN KKRFNYRVDGLEIV NGMWQIYNEHLGKID FNWTENGIPVEVVDK VNPATGQPTKDQVL KVG DYFN FQENSTG VVQE QTPYMGYTLS HVQLPNEFIWLFTDS KQALMYQ (SEQ ID NO: 80)

<i>Lactobacilli</i>	lamdaSA2	LysA, LysA2, and Lysga Y	Nonspecified	N/A
<i>S. aureus</i>		SAL-1	Nonspecified	N/A

In some instances, the lysin is a functionally active variant of the lysins described herein. In some instances, the variant of the lysin has at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity, e.g., over a specified region or over the entire sequence, to a sequence of a lysin described herein or a naturally occurring lysin.

In some instances, the lysin may be bioengineered to modulate its bioactivity, e.g., increase or decrease or regulate, or to specify a target microorganism. In some instances, the lysin is produced by the translational machinery (e.g. a ribosome, etc.) of a microbial cell. In some instances, the lysin is chemically synthesized. In some instances, the lysin is derived from a polypeptide precursor. The polypeptide precursor can undergo cleavage (for example, processing by a protease) to yield the polypeptide of the lysin itself. As such, in some instances, the lysin is produced from a precursor polypeptide. In some instances, the lysin includes a polypeptide that has undergone post-translational modifications, for example, cleavage, or the addition of one or more functional groups.

The lysins described herein may be formulated in a composition for any of the uses described herein. The compositions disclosed herein may include any number or type (e.g., classes) of lysins, such as at least about any one of 1 lysin, 2, 3, 4, 5, 10, 15, 20, or more lysins. A suitable concentration of each lysin in the composition depends on factors such as efficacy, stability of the lysin, number of distinct lysin, the formulation, and methods of application of the composition. In some instances, each lysin in a liquid composition is from about 0.1 ng/mL to about 100 mg/mL. In some instances, each lysin in a solid composition is from about 0.1 ng/g to about 100 mg/g. In some instances, wherein the composition includes at least two types of lysins, the concentration of each type of lysin may be the same or different.

A modulating agent including a lysin as described herein can be contacted with the target host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold level) of lysin concentration inside a target host; (b) reach a target level (e.g., a predetermined or threshold level) of lysin concentration inside a target host gut; (c) reach a target level (e.g., a predetermined or threshold level) of lysin concentration inside a target host bacteriocyte; (d) modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host; or/and (e) modulate fitness of the target host.

(c) Antimicrobial Peptides

The modulating agent described herein may include an antimicrobial peptide (AMP). Any AMP suitable for inhibiting a microorganism resident in the host may be used. AMPs are a diverse group of molecules, which are divided into subgroups on the basis of their amino acid composition and structure. The AMP may be derived or produced from any organism that naturally produces AMPs, including AMPs derived from plants (e.g., cospin), insects (e.g., drosocin, scorpion peptide (e.g., Uy192, UyCT3, D3, D10,

Uy17, Uy192), mastoparan, poneratoxin, cecropin, moricin, melittin), frogs (e.g., magainin, dermaseptin, aurein), and mammals (e.g., cathelicidins, defensins and protegrins). For example, the AMP may be a scorpion peptide, such as Uy192 (5'- FLSTIWNGIKGLL-3'; SEQ ID NO: 227), UyCT3 (5'- LSAIWSGIKSLF-3; SEQ ID NO: 228), D3 (5'- LWGKLWEGVKSLI-3'; SEQ ID NO: 229), and D10 (5'-

5 FPFLKLSLKIPKSAIKSAIKRL-3'; SEQ ID NO: 230), Uy17 (5'- ILSAIWSGIKGLL-3'; SEQ ID NO: 231), or a combination thereof. In some instances, the antimicrobial peptide may be one having at least 90% sequence identity (e.g., at least 90%, 92%, 94%, 96%, 98%, or 100% sequence identity) with one or more of the following: cecropin (SEQ ID NO: 82), melittin, copsin, drosomycin (SEQ ID NO: 93), dermcidin (SEQ ID NO: 81), andropin (SEQ ID NO: 83), moricin (SEQ ID NO: 84), ceratotoxin (SEQ ID NO: 85),

10 abaecin (SEQ ID NO: 86), apidaecin (SEQ ID NO: 87), prophenin (SEQ ID NO: 88), indolicidin (SEQ ID NO: 89), protegrin (SEQ ID NO: 90), tachyplesin (SEQ ID NO: 91), or defensin (SEQ ID NO: 92) to a vector of an animal pathogen. Non-limiting examples of AMPs are listed in Table 6.

Table 6: Examples of Antimicrobial Peptides

Type	Characteristic	Example AMP	Sequence
Anionic peptides	rich in glutamic and aspartic acid	dermcidin	SSLLEKGLDGAKKAVGGLGKL GKDAVEDLESVGKGAVHDVKD VLDSVL (SEQ ID NO: 81)
Linear cationic α -helical peptides	lack cysteine	cecropin A	KWKLFKKIEKVGQNIRDGIKAG PAVAVVGQATQIAK (SEQ ID NO: 82)
		andropin	MKYFSVLVVLTLILAIVDQSDAFI NLLDKVEDALHTGAQAGFKLIR PVERGATPKKSEKPEK (SEQ ID NO: 83)
		moricin	MNILKFFFVIVAMSLVSCSTAA PAKIPIKAIKTVGKAVGKGLRAI NIASTANDVFNFLKPKKRKH (SEQ ID NO: 84)
		ceratotoxin	MANLKAVFLICIVAFIALQCVVA EPAAEDSVVVKRSIGSALKKAL PVAKKIGKIALPIAKAALPVAAG LVG (SEQ ID NO: 85)
Cationic peptide enriched for specific amino acid	rich in proline, arginine, phenylalanine, glycine, tryptophan	abaecin	MKVVIFIFALLATICAAYVPLP NVPQPGRRPFPTFGQGPFNP KIKWPQGY (SEQ ID NO: 86)

		apidaecins	KNFALAILVVTFFVAVFGNTNLD PPTRPTRLRREAKPEAEPGNN RPVYIPQPRPPHPRLRREAEP AEPGNNRPVYIPQPRPPHPRL RREAELAEAGNNRPVYISQP RPPHPRLRREAEP AEPGNNR PVYIPQPRPPHPRLRREAEL EPGNNRPVYISQPRPPHPRLR REAEP AEPGNNRPVYIPQPR PPHPRLRREAEP AEPGNNR VYIPQPRPPHPRLRREAEP PGNNRPVYIPQPRPPHPRLR EAKPEAKPGNNRPVYIPQPR PHPRI (SEQ ID NO: 87)
		prophenin	METQRASLCLGRWSLWLLLLA LVVPSASAQALS YREAVLRAVD RLNEQSSEANLYRLELDQPPK ADEDPGTPKPVSTVKETVCP RPTRRPPELCDFKENG RVKQC VGTVTLDQIKDPLDITCNEGVR RFPWWPFLRRPRLRRQAFP PPNVPGPRFPPPNVPGPRFPP PNFPGPRFPPPNFPGPRFPP NFGPPFPPIFGPWFPFPP FRPPFPGPRFPGRR (SEQ ID NO: 88)
		indolicidin	MQTQRASLSLGRWSLWLLLLG LVVPSASAQALS YREAVLRAVD QLNELSSEANLYRLELDPPK DNEDLGTRKPVSTVKETVCP RTIQQPAEQCDFKEKGRVKQC VGTVTLDPSNDQFDLNCNELQ SVILPWKWPWWPWRRG (SEQ ID NO: 89)
Anionic and cationic peptides that contain cysteine and	contain 1-3 disulfide bond	protegrin	METQRASLCLGRWSLWLLLLA LVVPSASAQALS YREAVLRAVD RLNEQSSEANLYRLELDQPPK ADEDPGTPKPVSTVKETVCP RPTRRPPELCDFKENG RVKQC

form disulfide bonds			VGTVTLDQIKDPLDITCNEVQG VRGGRLCYCRRRFCVCVGRG (SEQ ID NO: 90)
		tachyplesins	KWCFRVCYRGICYRRCR (SEQ ID NO: 91)
		defensin	MKCATIVCTIAVVLAATLLNGSV QAAPQEEAALSGGANLNTLLD ELPEETHHAALENYRAKRATC DLASGFGVGSSLCAAHCIARR YRGGYCNSKAVCVCRN (SEQ ID NO: 92)
		drosomycin	MMQIKYLFALFAVLMLVVLGAN EADADCLSGRYKGPCAVWDN ETCRRVCKEEGRSSGHCSPL KCWCEGC (SEQ ID NO: 93)

The AMP may be active against any number of target microorganisms. In some instances, the AMP may have antibacterial and/or antifungal activities. In some instances, the AMP may have a narrow-spectrum bioactivity or a broad-spectrum bioactivity. For example, some AMPs target and kill only a few species of bacteria or fungi, while others are active against both gram-negative and gram-positive bacteria as well as fungi.

Further, the AMP may function through a number of known mechanisms of action. For example, the cytoplasmic membrane is a frequent target of AMPs, but AMPs may also interfere with DNA and protein synthesis, protein folding, and cell wall synthesis. In some instances, AMPs with net cationic charge and amphipathic nature disrupt bacterial membranes leading to cell lysis. In some instances, AMPs may enter cells and interact with intracellular target to interfere with DNA, RNA, protein, or cell wall synthesis. In addition to killing microorganisms, AMPs have demonstrated a number of immunomodulatory functions that are involved in the clearance of infection, including the ability to alter host gene expression, act as chemokines and/or induce chemokine production, inhibit lipopolysaccharide induced pro-inflammatory cytokine production, promote wound healing, and modulating the responses of dendritic cells and cells of the adaptive immune response.

In some instances, the AMP is a functionally active variant of the AMPs described herein. In some instances, the variant of the AMP has at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity, e.g., over a specified region or over the entire sequence, to a sequence of an AMP described herein or a naturally derived AMP.

In some instances, the AMP may be bioengineered to modulate its bioactivity, e.g., increase or decrease or regulate, or to specify a target microorganism. In some instances, the AMP is produced by the translational machinery (e.g. a ribosome, etc.) of a cell. In some instances, the AMP is chemically synthesized. In some instances, the AMP is derived from a polypeptide precursor. The polypeptide precursor can undergo cleavage (for example, processing by a protease) to yield the polypeptide of the

AMP itself. As such, in some instances, the AMP is produced from a precursor polypeptide. In some instances, the AMP includes a polypeptide that has undergone post-translational modifications, for example, cleavage, or the addition of one or more functional groups.

The AMPs described herein may be formulated in a composition for any of the uses described herein. The compositions disclosed herein may include any number or type (e.g., classes) of AMPs, such as at least about any one of 1 AMP, 2, 3, 4, 5, 10, 15, 20, or more AMPs. For example, the compositions may include a cocktail of AMPs (e.g., a cocktail of scorpion peptides, e.g., UyCT3, D3, D10, and Uy17). A suitable concentration of each AMP in the composition depends on factors such as efficacy, stability of the AMP, number of distinct AMP in the composition, the formulation, and methods of application of the composition. In some instances, each AMP in a liquid composition is from about 0.1 ng/mL to about 100 mg/mL (about 0.1 ng/mL to about 1 ng/mL, about 1 ng/mL to about 10 ng/mL, about 10 ng/mL to about 100 ng/mL, about 100 ng/mL to about 1000 ng/mL, about 1 mg/mL to about 10 mg/mL, about 10 mg/mL to about 100 mg/mL). In some instances, each AMP in a solid composition is from about 0.1 ng/g to about 100 mg/g (about 0.1 ng/g to about 1 ng/g, about 1 ng/g to about 10 ng/g, about 10 ng/g to about 100 ng/g, about 100 ng/g to about 1000 ng/g, about 1 mg/g to about 10 mg/g, about 10 mg/g to about 100 mg/g). In some instances, wherein the composition includes at least two types of AMPs, the concentration of each type of AMP may be the same or different.

A modulating agent including an AMP as described herein can be contacted with the target host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold level) of AMP concentration inside a target host; (b) reach a target level (e.g., a predetermined or threshold level) of AMP concentration inside a target host gut; (c) reach a target level (e.g., a predetermined or threshold level) of AMP concentration inside a target host bacteriocyte; (d) modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host; or/and (e) modulate fitness of the target host.

As illustrated by Examples 16 and 20-22, AMPs, such as scorpion peptides, can be used as modulating agents that target an endosymbiotic bacterium in an insect host to decrease the fitness of the host (e.g., as outlined herein).

(d) Nodule C-rich Peptides

The modulating agent described herein may include a nodule C-rich peptide (NCR peptide). NCR peptides are produced in certain leguminous plants and play an important role in the mutualistic, nitrogen-fixing symbiosis of the plants with bacteria from the *Rhizobiaceae* family (rhizobia), resulting in the formation of root nodules where plant cells contain thousands of intracellular endosymbionts. NCR peptides possess anti-microbial properties that direct an irreversible, terminal differentiation process of bacteria, e.g., to permeabilize the bacterial membrane, disrupt cell division, or inhibit protein synthesis. For example, in *Medicago truncatula* nodule cells infected with *Sinorhizobium meliloti*, hundreds of NCR peptides are produced which direct irreversible differentiation of the bacteria into large polyploid nitrogen-fixing bacteroids. Non-limiting examples of NCR peptides are listed in Table 7.

Table 7: Examples of NCR Peptides

NAME	Peptide sequence	Producer
>gi 152218086 gb ABS31477. 1 NCR 340	MTKIVVFYVILLTIFHVS AKKKRYI ECETHEDCSQVFMPPFVMRCVIHE CKIFNGEHLRY (SEQ ID NO: 94)	<i>Medicago truncatula</i>
>gi 152218084 gb ABS31476. 1 NCR 339	MAKIMKFVYNMIPFLSIFIITLQVNVV VCEIDADCPQICMPPEVRCVNHRC GWVNTDDSLFLTQEFTRSKQYIIS (SEQ ID NO: 95)	<i>Medicago truncatula</i>
>gi 152218082 gb ABS31475. 1 NCR 338	MYKVVESIFIRYMRKPNMTKFFKF VYTMFILISFLVVTNANAHNCTDISD CSSNHCSYEGVSLCMNGQCICIYE (SEQ ID NO: 96)	<i>Medicago truncatula</i>
>gi 152218080 gb ABS31474. 1 NCR 337	MVETLRLFYIMLFVSLCLVVDGES KLEQTCSEDFECYIKNPHVPFGHLR CFEGFCQQLNGPA (SEQ ID NO: 97)	<i>Medicago truncatula</i>
>gi 152218078 gb ABS31473. 1 NCR 336	MAKIVNFVYSMIVFLFLVATKAAR GYLCVTDSHCPPHMCPGMEPRCV RRMCKCLPIGWRYFVP (SEQ ID NO: 98)	<i>Medicago truncatula</i>
>gi 152218076 gb ABS31472. 1 NCR 335	MQIGKNMVETPKLDYVIFFLYFFF RQMILRLNTTFRPLNFKMLRFWGQ NRNIMKHRGQKVHFSLILSDCKTNK DCPKLRRANVRCKSYCVPI (SEQ ID NO: 99)	<i>Medicago truncatula</i>
>gi 152218074 gb ABS31471. 1 NCR 334	MLRLYLVSFYLLKRTLVSYSYFST YIECKTDNDPISQLKIYAWKCVKN GCHLFDVIPMMYE (SEQ ID NO: 100)	<i>Medicago truncatula</i>
>gi 152218072 gb ABS31470. 1 NCR 333	MAEILKFVYIVILFVSLLLIVVASEREC VTDDDCEKLYPTNEYRMMCDSGYC MNLLNGKIIYLLCLKKKKFLIISVLL (SEQ ID NO: 101)	<i>Medicago truncatula</i>
>gi 152218070 gb ABS31469. 1 NCR 332	MAEIIKFVYIMILCVSLLIEVAGEECV TDADCDKLYPDIRKPLMCSIGECYSL YKGKFSLSIISKTSFSLMVYNVVTLVI CLRLAYISLLKFL (SEQ ID NO: 102)	<i>Medicago truncatula</i>
>gi 152218068 gb ABS31468. 1 NCR 331	MAEILKDFYAMNLFIFLIILPAKIRGET LSLTHPKCHHIMLPSLFITEVFQRT	<i>Medicago truncatula</i>

	DDGCPKPVNHLRVVKCIEHICEYGY NYRPDFASQIPESTKMMPKRE (SEQ ID NO: 103)	
>gi 152218066 gb ABS31467. 1 NCR 330	MVEILKNFYAMNLFIFLILAVKIRGAH FPCVTDDDCPKPVNKLRLVIKCIDHIC QYARNLPDFASEISESTKMPCKGE (SEQ ID NO: 104)	<i>Medicago truncatula</i>
>gi 152218064 gb ABS31466. 1 NCR 329	MFHAQAENMAKVSFVCIMILFLALF FITMNDAAARFECREDSHCVTIRKCV LPRKPECRNYACGCYDSNKYR (SEQ ID NO: 105)	<i>Medicago truncatula</i>
>gi 152218062 gb ABS31465. 1 NCR 328	MQMRQNMATILNFVFIILFISLLLTV TKGYREPFSSTFTEGPTCKEDIDCPSI SCVNPQVPKCMFECHCKYIPTTLK (SEQ ID NO: 106)	<i>Medicago truncatula</i>
>gi 152218060 gb ABS31464. 1 NCR 327	MATILMYVYITILFISILTVLTEGLYEPL YNFRDPDCRRNIDCPSYLCVAPKV PRCIMFECHCKDIPSDH (SEQ ID NO: 107)	<i>Medicago truncatula</i>
>gi 152218058 gb ABS31463. 1 NCR 326	MTTSLKFVYVAILFLSLLLVMGGIR RFECRQSDCPSYFCEKLTVPKCF WSKCYCK (SEQ ID NO: 108)	<i>Medicago truncatula</i>
>gi 152218056 gb ABS31462. 1 NCR 325	MTTSLKFVYVAILFLSLLLVMGGIR KKECRQSDCPSYFCEKLTIAKCIHS TCLCK (SEQ ID NO: 109)	<i>Medicago truncatula</i>
>gi 152218054 gb ABS31461. 1 NCR 324	MQIGKNMVETPKLVYFIILFISFLCIT VSNSSFSQIFNSACKTDKDCPKFGR VNVRCRKGNCVPI (SEQ ID NO: 110)	<i>Medicago truncatula</i>
>gi 152218046 gb ABS31457. 1 NCR 320	MTAILKKFINAVFLFIVLFLATTNVED FVGGSNDECYVPDVFQCINNICKCV SHHRT (SEQ ID NO: 111)	<i>Medicago truncatula</i>
>gi 152218044 gb ABS31456. 1 NCR 319	MQKRKNMAQIIFYVYALILFSPFLAA RLVFVNPEKPCVTDADCDRYRHES AIYSDMFCKDGYCFIDYHHPYP (SEQ ID NO: 112)	<i>Medicago truncatula</i>
>gi 152218042 gb ABS31455. 1 NCR 318	MQMRKNMAQILFYVYALLILFTPFLV ARIMVVNPNNPCVTDADCQRYRHK	<i>Medicago truncatula</i>

	LATRMICNQGFLMDFTHPYAPSL P (SEQ ID NO: 113)	
>gi 152218040 gb ABS31454. 1 NCR 317	MNHISKFVYALIIFLSIYLVLDGLPIS CKDHFECRRKINILRCIYRQEKPMCI NSICTCVKLL (SEQ ID NO: 114)	<i>Medicago truncatula</i>
>gi 152218038 gb ABS31453. 1 NCR 316	MQREKNMAKIFEFVYAMIIIFILLFLVE KNVVAYLKFECKTDDDCQKSLLKTY VWKCVKNECYFFAKK (SEQ ID NO: 115)	<i>Medicago truncatula</i>
>gi 152218036 gb ABS31452. 1 NCR 315	MAGIIKFVHVLIIFLSLFHVKNDDGS FCFKDSDCPDEMCPSPLEKEMCYFL QCKCGVDTIA (SEQ ID NO: 116)	<i>Medicago truncatula</i>
>gi 152218034 gb ABS31451. 1 NCR 314	MANTHKLVSMLIFLFLASNNVEGY VNCETDADCPPSTRVKRFKCVKGE CRWTRMSYA (SEQ ID NO: 117)	<i>Medicago truncatula</i>
>gi 152218032 gb ABS31450. 1 NCR 313	MQRRKKKAQVVMFVHDLIICIYLFIVI TTRKTDIRCFYYDCPRLEYHFCECI EDFCAYIRLN (SEQ ID NO: 118)	<i>Medicago truncatula</i>
>gi 152218030 gb ABS31449. 1 NCR 312	MAKVYMFVYALIIFVSPFLLATFRTRL PCEKDDDCPEAFLPPVMKCVNRFC QYEILE (SEQ ID NO: 119)	<i>Medicago truncatula</i>
>gi 152218028 gb ABS31448. 1 NCR 310	MIKQFSVCYIQMRRNMTTILKFPYIM VICLLLLHVAAYEDFEKEIFDCKKD DCDHMCVTPGIPKCTGYVCFCFENL (SEQ ID NO: 120)	<i>Medicago truncatula</i>
>gi 152218026 gb ABS31447. 1 NCR 309	MQRSRNMTTIFKFAYIMIICVFLNIA AQEIENGHPCKKNEDCNHMCVMP GLPWCHENNLFCYENAYGNTR (SEQ ID NO: 121)	<i>Medicago truncatula</i>
>gi 152218024 gb ABS31446. 1 NCR 304	MTIIKFVNVLIIFLSLFHVAKNDDNKL LLSFIEEGFLCFKSDCPYNMCPSP LKEMCYFIKCVCGVYGPIRERRLYQ SHNPMIQ (SEQ ID NO: 122)	<i>Medicago truncatula</i>

>gi 152218022 gb ABS31445. 1 NCR 303	MRKNMTKILMIGYALMIFILSIASIT GNLARASRKPPVDVPCYDHDGPR KLYFLERCVRVCKYL (SEQ ID NO: 123)	<i>Medicago truncatula</i>
>gi 152218020 gb ABS31444. 1 NCR 301	MAHKLVYAITLFIFLFIANNIEDDIFCI TDNDCPPNTLVQRYRCINGKCNLSF VSYG (SEQ ID NO: 124)	<i>Medicago truncatula</i>
>gi 152218018 gb ABS31443. 1 NCR 300	MDETLKFVYILILFVSLCLVVADGVK NINRECTQTSDCYKKYPFIPWGKVR CVKGRCLDM (SEQ ID NO: 125)	<i>Medicago truncatula</i>
>gi 152218016 gb ABS31442. 1 NCR 290	MAKIIKFVYVLAIFFSLFLVAKNVNG WTCVEDSDCPANICQPPMQRMCFY GECACVRSKFCT (SEQ ID NO: 126)	<i>Medicago truncatula</i>
>gi 152218014 gb ABS31441. 1 NCR 289	MVKIIFVYFMTLFLSMLLVTTKEDG SVECIANIDCPQIFMLPFVMRCINFR CQIVNSED (SEQ ID NO: 127)	<i>Medicago truncatula</i>
>gi 152218012 gb ABS31440. 1 NCR 286	MDEILKFVYTLIIFFSLFFAANNVDANI MNCQSTFDCPRDMCSHIRDVICIFK KCKCAGGRYPQVP (SEQ ID NO: 128)	<i>Medicago truncatula</i>
>gi 152218008 gb ABS31438. 1 NCR 278	MQRRKNMANNHMLIYAMIICLFPYL VVTFKTAITCDCNEDCLNFFTPLDNL KCIDNVCEVFM (SEQ ID NO: 129)	<i>Medicago truncatula</i>
>gi 152218006 gb ABS31437. 1 NCR 266	MVNILKFIYVIFFILMFFVLIDVDGHV LVECIENRDCEKGMCKFPFIVRCLM DQCKCVRIHNLI (SEQ ID NO: 130)	<i>Medicago truncatula</i>
>gi 152218004 gb ABS31436. 1 NCR 265	MIIQFSIYYMQRRKLNLMVEILKFSHA LIIFLFLSALVTNANIFFCSTDEDCTW NLCRQPWWQKCRMHMCSEKN (SEQ ID NO: 131)	<i>Medicago truncatula</i>
>gi 152218002 gb ABS31435. 1 NCR 263	MDEVFKFVYVMIIFFLILDVATNAEK IRRCFNDAHCPPDMCTLGVIPKCSR FTICIC (SEQ ID NO: 132)	<i>Medicago truncatula</i>

>gi 152218000 gb ABS31434.1 NCR 244	MHRKPNMTKFFKFVYTMFILISLFLV VTNANANNCTDTSDCSSNHCSYEG VSLCMNGQCICIYE (SEQ ID NO: 133)	<i>Medicago truncatula</i>
>gi 152217998 gb ABS31433.1 NCR 239	MQMKKMATILKFVYLILLIYPLLVVTE ESHYMKFSICKDDTDCPTLFCVLPN VPKCIGSKCHCKLMVN (SEQ ID NO: 134)	<i>Medicago truncatula</i>
>gi 152217996 gb ABS31432.1 NCR 237	MVETLRLFYIMILFVSLYLVVVDGVS KLAQSCSEDFECYIKNPHAPFGQLR CFEGYCQRLDKPT (SEQ ID NO: 135)	<i>Medicago truncatula</i>
>gi 152217994 gb ABS31431.1 NCR 228	MTTFLKVAYIMIICVFLHLAAQVDS QKRLHGCCKEDRDCDNICSVHAVTK CIGNMCRCLANVK (SEQ ID NO: 136)	<i>Medicago truncatula</i>
>gi 152217992 gb ABS31430.1 NCR 224	MRINRTPAIFKFVYTIILYFLLRVVAK DLFPNICEKDEDCLEFCAHDKVAKC MLNICFCF (SEQ ID NO: 137)	<i>Medicago truncatula</i>
>gi 152217990 gb ABS31429.1 NCR 221	MAEILKILYVFIIFLSLILAVISQHPFTP CETNADCKCRNHKRPDCLWHKCYC Y (SEQ ID NO: 138)	<i>Medicago truncatula</i>
>gi 152217988 gb ABS31428.1 NCR 217	MRKSMATILKFVYVIMLFYISLFVIES FGHRFLIYNCKNDTECPNDGCPHE QAKCILYACYCVE (SEQ ID NO: 139)	<i>Medicago truncatula</i>
>gi 152217986 gb ABS31427.1 NCR 209	MNTILKFIFVVFLFLSIFLSAGNSKSY GPCTTLQDCETHNWFEVCSCIDFEC KCWSLL (SEQ ID NO: 140)	<i>Medicago truncatula</i>
>gi 152217984 gb ABS31426.1 NCR 206	MAEIKFVYIMILCVSLLIAEASGKEC VTDADCENLYPGNKKPMFCNNTGY CMSLYKEPSRYM (SEQ ID NO: 141)	<i>Medicago truncatula</i>
>gi 152217982 gb ABS31425.1 NCR 201	MAKIIKFVYIMILCVSLLLIVEAGGKEC VTDVDCEKIYPGNKKPLICSTGYCYS LYEEPTRYHK (SEQ ID NO: 142)	<i>Medicago truncatula</i>

>gi 152217980 gb ABS31424. 1 NCR 200	MAKVTKFGYIIHFLSLFFLAMNVAG GRECHANSCHCVGKITCVLPQKPEC WNYACVCYDSNKYR (SEQ ID NO: 143)	<i>Medicago truncatula</i>
>gi 152217978 gb ABS31423. 1 NCR 192	MAKIFNYVYALIMFLSLFLMGTSGMK NGCKHTGHCPRKMCAGAKTTKCRN NKCQCV (SEQ ID NO: 144)	<i>Medicago truncatula</i>
>gi 152217976 gb ABS31422. 1 NCR 189	MTEILKFVCMIIFFISFIVSKSLNGG GKDKCFRSDCPKHMCPSSLVAKCI NRLCRCRRPELQVQLNP (SEQ ID NO: 145)	<i>Medicago truncatula</i>
>gi 152217974 gb ABS31421. 1 NCR 187	MAHIIMFYVYALIIFSSLFVRDGIP CLSDDECPESHYSFKCINNICEYD LGEMSDDDYLEMSRE (SEQ ID NO: 146)	<i>Medicago truncatula</i>
>gi 152217972 gb ABS31420. 1 NCR 181	MYREKNMAKTLKFVYVIVLFLSLFLA AKNIDGRVSYNSFIALPVCQTAADC PEGTRGRTYKCINNCRYPKLLKPI Q (SEQ ID NO: 147)	<i>Medicago truncatula</i>
>gi 152217970 gb ABS31419. 1 NCR 176	MAHIFNYVYALLVFLSLFLMVTNGIHI GCDKDRDCPKQMCHLNQTPKCLKN ICKCV (SEQ ID NO: 148)	<i>Medicago truncatula</i>
>gi 152217968 gb ABS31418. 1 NCR 175	MAEILKCFYTMNLFIFLIIPAKIREHI QCVIDDDCPKSLNKLIIKCINHVQY VGNLPDFASQIPKSTKMPYKGE (SEQ ID NO: 149)	<i>Medicago truncatula</i>
>gi 152217966 gb ABS31417. 1 NCR 173	MAYISRIFYVLIIFLSLFFVINGVKSL LLIKVRSFIPQCRSDDCPRNLCVDQII PTCVWAKCKCKNYND (SEQ ID NO: 150)	<i>Medicago truncatula</i>
>gi 152217964 gb ABS31416. 1 NCR 172	MANVTKFVYIAIYFLSLFFIAKNDA TFCHDDSHCVTKIKCVLPRTQCRN EACGCYHSNKFR (SEQ ID NO: 151)	<i>Medicago truncatula</i>
>gi 152217962 gb ABS31415. 1 NCR 171	MGEIMKFVYVMIIYLFMFNVATGSEF IFTKKLTSCDSSKDCRSFLCYSPKFP VCKRGICECI (SEQ ID NO: 152)	<i>Medicago truncatula</i>

>gi 152217960 gb ABS31414.1 NCR 169	MGEMFKFIYTFILFVHLFLVVIFEDIG HIKYCGIVDDCYKSKKPLFKIWKCVE NVCVLWYK (SEQ ID NO: 153)	<i>Medicago truncatula</i>
>gi 152217958 gb ABS31413.1 NCR 165	MARTLKFVYSMILFSLFLVANGLKIF CIDVADCPKDLYPILLYKCIYNKCIVFT RIPFPFDWI (SEQ ID NO: 154)	<i>Medicago truncatula</i>
>gi 152217956 gb ABS31412.1 NCR 159	MANITKFVYIAILFSLFFIGMNDAAIL ECREDSHCVTKIKCVLPRKPECRNN ACTCYKGGFSFHH (SEQ ID NO: 155)	<i>Medicago truncatula</i>
>gi 152217954 gb ABS31411.1 NCR 147	MQRVKKMSETLKFVYVILFISIFHV IVCDSIYFPVSRPCITDKDCPNMKHY KAKCRKGFCISSRVR (SEQ ID NO: 156)	<i>Medicago truncatula</i>
>gi 152217952 gb ABS31410.1 NCR 146	MQIRKIMSGVLKFVYAILFLFLVA REVGLETIECETDGDGCPRSMIKM WNKNYRHKCIDGKCEWIKKLP (SEQ ID NO: 157)	<i>Medicago truncatula</i>
>gi 152217950 gb ABS31409.1 NCR 145	MFVYDLILFISLILVVTGINAEADTSC HSFDDCPWWAHYRECIEGLCAYRI LY (SEQ ID NO: 158)	<i>Medicago truncatula</i>
>gi 152217948 gb ABS31408.1 NCR 144	MQRRKKSMAKMLKFFFAIILLSLFL VATEVGGAYIECEVDDDCPKPMKN SHPDTYYKCVKHRCQWAWK (SEQ ID NO: 159)	<i>Medicago truncatula</i>
>gi 152217946 gb ABS31407.1 NCR 140	MFVYTLIIFLPSHVITNKIAIYCVSDD DCLKTFTPLDLKCDNVCEFNLRCK GKCGERDEKFVFLKALKKMDQKLVL EEQGNAREVKIPKLLFDRIQVPTPA TKDQVEEDDYDDDDDEEEEEEDDV DMWFHLPDVVCH (SEQ ID NO: 160)	<i>Medicago truncatula</i>
>gi 152217944 gb ABS31406.1 NCR 138	MAKFSMFVYALINFLSLFLVETAITNI RCVSDDDCPKVIKPLVMKCIGNYCY FFMIYEGP (SEQ ID NO: 161)	<i>Medicago truncatula</i>
>gi 152217942 gb ABS31405.1 NCR 136	MAHKFVYAILFIFLFLVAKNVKGYVV CRTVDDCPPDTRDLRYRCLNGKCK SYRLSYG (SEQ ID NO: 162)	<i>Medicago truncatula</i>

>gi 152217940 gb ABS31404.1 NCR 129	MQRKKNMGQILIFVFALINFLSPILVE MTTTTIPCTFIDDCPKMPLVVKCIDN FCNYFEIK (SEQ ID NO: 163)	<i>Medicago truncatula</i>
>gi 152217938 gb ABS31403.1 NCR 128	MAQTLMLVYALIIFTSLFLVVISRQTD IPCKSDDACPRVSSHIECVKGFCT YWKLD (SEQ ID NO: 164)	<i>Medicago truncatula</i>
>gi 152217936 gb ABS31402.1 NCR 127	MLRRKNTVQILMFVSALLIYIFLFLVIT SSANIPCNSDSDCPWKIYYTYRCND GFCVYKSIDPSTIPQYMTDLIFPR (SEQ ID NO: 165)	<i>Medicago truncatula</i>
>gi 152217934 gb ABS31401.1 NCR 122	MAVILKFVYIMIIIFLLYVVNGTRCN RDEDCPFICTGPQIPKCVSHICFCLS SGKEAY (SEQ ID NO: 166)	<i>Medicago truncatula</i>
>gi 152217932 gb ABS31400.1 NCR 121	MDAILKFIYAMFLFLFLVTTTRNVEAL FECNRDFVCGNDDECVPYAVQCI HRYCKCLKSRN (SEQ ID NO: 167)	<i>Medicago truncatula</i>
>gi 152217930 gb ABS31399.1 NCR 119	MQIGRKKGGETPKLVYVILFLSIFLC TNSSFSQMINFRGCKRDKDCPQFR GVNIRCRSGFCTPIDS (SEQ ID NO: 168)	<i>Medicago truncatula</i>
>gi 152217928 gb ABS31398.1 NCR 118	MQMRKNMAQILFYVYALLILFSPFLV ARIMVVNPNNPCVTDADCQRYRHK LATRMVCNIGFCLMDFTHDPYAPSL P (SEQ ID NO: 169)	<i>Medicago truncatula</i>
>gi 152217926 gb ABS31397.1 NCR 111	MYVYIYQMGKNMAQRFMFIYALIIFL SQFFVINTSDIPNNSNRNSPKEDVF CNSNDDCPTILYYVSKCVYNFCEYW (SEQ ID NO: 170)	<i>Medicago truncatula</i>
>gi 152217924 gb ABS31396.1 NCR 103	MAKIVNFVYSMIIFVSLFLVATKGGG KPFLTRPYPCNTGSDCPQNMCPG YKPGCEDGYCNHCYKRW (SEQ ID NO: 171)	<i>Medicago truncatula</i>
>gi 152217922 gb ABS31395.1 NCR 101	MVRTLKFYVYVILILSLFLVAKGGGKK IYCENAASCPRMLYPLVYKCLDNKC VKFMMKSRFV (SEQ ID NO: 172)	<i>Medicago truncatula</i>

>gi 152217920 gb ABS31394.1 NCR 96	MARTLKFFVYAVILFLSLFLVAKGDDV KIKCVVAANCPDLMYPLVYKCLNGIC VQFTLTFFPV (SEQ ID NO: 173)	<i>Medicago truncatula</i>
>gi 152217918 gb ABS31393.1 NCR 94	MSNTLMFVITFIVLVTFLGPKNVYA FQPCVTTADCMKTLKTDENIWYECI NDFCIPFPIPKGRK (SEQ ID NO: 174)	<i>Medicago truncatula</i>
>gi 152217916 gb ABS31392.1 NCR 93	MKRNVNMAKIVKYVYVIIIIFLSLFLVA TKIEGYYYKCFKSDCVKLLCRIPLR PKCMYRHICKCKVVLQTNNYVLT (SEQ ID NO: 175)	<i>Medicago truncatula</i>
>gi 152217914 gb ABS31391.1 NCR 90	MKRGKNMSKILFIYATLVLYLFLVV TKASDDECKIDGDCPISWQKFHTYK CINQKCKWVLRFFEY (SEQ ID NO: 176)	<i>Medicago truncatula</i>
>gi 152217912 gb ABS31390.1 NCR 88	MAKTLNFMFALILFISLFLVSKNVAID FVCQTDADCPKSELSTWTKCIDN ECNLFKVMQGMV (SEQ ID NO: 177)	<i>Medicago truncatula</i>
>gi 152217910 gb ABS31389.1 NCR 86	MANTHKLVSMLFIFLFLVANNVEGY VNCETDADCPPSTRVKRFKCVKGE CRWTRMSYA (SEQ ID NO: 178)	<i>Medicago truncatula</i>
>gi 152217908 gb ABS31388.1 NCR 77	MAHFLMFVYALITCLSLFLVEMGHLS IHCVSDDCPKVEKPITMKCINNYCK YFVDHKL (SEQ ID NO: 179)	<i>Medicago truncatula</i>
>gi 152217906 gb ABS31387.1 NCR 76	MNQIPMFGYTLIIFFSLFPVITNGDRI PCVTNGDCPVMRLPLYMRCITYSCE LFFDGPNLCAVERI (SEQ ID NO: 180)	<i>Medicago truncatula</i>
>gi 152217904 gb ABS31386.1 NCR 74	MRKDMARISLFFVYALIIFFSLFFVLTN GELEIRCVSDADCPFLPLPLHNRCID DVCHLFTS (SEQ ID NO: 181)	<i>Medicago truncatula</i>
>gi 152217902 gb ABS31385.1 NCR 68	MAQILMFVYFLIIIFLSLFLVESIKIFTE HRCRTDADCPARELPEYLKQGGM CRLLIKKD (SEQ ID NO: 182)	<i>Medicago truncatula</i>

>gi 152217900 gb ABS31384.1 NCR 65	MARVISLFYALIIFFLFLVATNGDLS PCLRSGDCSKDECPHLPKICIGLT CYCI (SEQ ID NO: 183)	<i>Medicago truncatula</i>
>gi 152217898 gb ABS31383.1 NCR 62	MQRRKNMAQILLFAYVFIISLFLVV TNGVKIPCVKDTDCPTLPCPLYSKC VDGFCKMLSI (SEQ ID NO: 184)	<i>Medicago truncatula</i>
>gi 152217896 gb ABS31382.1 NCR 57	MNHISKFVYALIIFLSVYLVLDGRPV SCKDHYDCRRKVKIVGCIFPQEKPM CINSMCTCIREIVP (SEQ ID NO: 185)	<i>Medicago truncatula</i>
>gi 152217894 gb ABS31381.1 NCR 56	MKSQNHAKFISFYKNDLFKIFQNND SHFKVFFALIIFLYTLHVTNGVFVSC NSHIHCRVNNHKIGCNIPEQYLLCVN LFCLWLDY (SEQ ID NO: 186)	<i>Medicago truncatula</i>
>gi 152217892 gb ABS31380.1 NCR 54	MTYISKVVYALIIFLSIYVGVNDCMLV TCEDHFDQRQNVQVGCSEIPQ CINSICKCMKG (SEQ ID NO: 187)	<i>Medicago truncatula</i>
>gi 152217890 gb ABS31379.1 NCR 53	MTHISKFVFALIIFLSIYVGVNDCRIP CKDNNDCCNNWQLLACRFEREVPR CINSICKCMPM (SEQ ID NO: 188)	<i>Medicago truncatula</i>
>gi 152217888 gb ABS31378.1 NCR 43	MVQTPKLVYVIVLLLSIFLGMTICNSS FSHFFEGACKSDKDCPKLHRSNVR CRKGQCVQI (SEQ ID NO: 189)	<i>Medicago truncatula</i>
>gi 152217886 gb ABS31377.1 NCR 28	MTKILMLFYAMIVFHSIFLVASYTDEC STDADCEYILCLFPIIKRCIHNHCKCV PMGSIEPMSTIPNGVHKFHIINN (SEQ ID NO: 190)	<i>Medicago truncatula</i>
>gi 152217884 gb ABS31376.1 NCR 26	MAKTLNFVCAMILFISLFLVSKNVAL YIIECKTDADCPISKLNMYNWRCIKS SCHLYKVIQFMV (SEQ ID NO: 191)	<i>Medicago truncatula</i>
>gi 152217882 gb ABS31375.1 NCR 24	MQKEKNMAKTFEFVYAMIIIFILLFLVE NNFAAYIIECQTDDDCPKSQLEMFA WKCVKNGCHLFGMYEDDDDP (SEQ ID NO: 192)	<i>Medicago truncatula</i>

>gi 152217880 gb ABS31374.1 NCR 21	MAATRKFYVLSHFLFLVTKITDAR VCKSDKDCKDIIYRYILKCRNGECV KIKI (SEQ ID NO: 193)	<i>Medicago truncatula</i>
>gi 152217878 gb ABS31373.1 NCR 20	MQRLDNMAKNVKFIYVILLFIFLVII VCDSAFVPNSGPCTTDKDKQVKG YIARCRKGYCMQSVKRTWSSYSR (SEQ ID NO: 194)	<i>Medicago truncatula</i>
>gi 152217876 gb ABS31372.1 NCR 19	MKFIYIMILFSLFLVQFLTCKGLTVP CENPTTCTPEDFCTPPMITRCINFICL CDGPEYAEPEYDGPEPEYDHKGDF LSVKPKIINENMMMRERHMMKEIEV (SEQ ID NO: 195)	<i>Medicago truncatula</i>
>gi 152217874 gb ABS31371.1 NCR 12	MAQFLMFIYVLIIFLYFYVEAMFEL TKSTIRCVTDADCPNVVKPLKPKCV DGFCEYT (SEQ ID NO: 196)	<i>Medicago truncatula</i>
>gi 152217872 gb ABS31370.1 NCR 10	MKMRIHMAQIIMFFYALIIFLSPFLVD RRSFPSSFVSPKSYTSEIPCKATRD CPYELYYETKCVDLSCTY (SEQ ID NO: 197)	<i>Medicago truncatula</i>

Any NCR peptide known in the art is suitable for use in the methods or compositions described herein. NCR peptide-producing plants include but are not limited to *Pisum sativum* (pea), *Astragalus sinicus* (IRLC legumes), *Phaseolus vulgaris* (bean), *Vigna unguiculata* (cowpea), *Medicago truncatula* (barrelclover), and *Lotus japonicus*. For example, over 600 potential NCR peptides are predicted from the *M. truncatula* genome sequence and almost 150 different NCR peptides have been detected in cells isolated from root nodules by mass spectrometry.

The NCR peptides described herein may be mature or immature NCR peptides. Immature NCR peptides have a C-terminal signal peptide that is required for translocation into the endoplasmic reticulum and cleaved after translocation. The N-terminus of a NCR peptide includes a signal peptide, which may be cleavable, for targeting to a secretory pathway. NCR peptides are generally small peptides with disulfide bridges that stabilize their structure. Mature NCR peptides have a length in the range of about 20 to about 60 amino acids, about 25 to about 55 amino acids, about 30 to about 50 amino acids, about 35 to about 45 amino acids, or any range therebetween. NCR peptides may include a conserved sequence of cysteine residues with the rest of the peptide sequence highly variable. NCR peptides generally have about four or eight cysteines.

NCR peptides may be anionic, neutral, or cationic. In some instances, synthetic cationic NCR peptides having a pI greater than about eight possess antimicrobial activities. For example, NCR247 (pI = 10.15) (RNGCIVDPRCPYQQCRRPLYCRRR; SEQ ID NO: 198) and NCR335 (pI = 11.22) (MAQFLLFVYSLIIFLSLFFGEAAFERTETRMLTIPCTSDDNCPKVISPCHTKCFDGFCEGWYIEGSYEGP; SEQ ID NO: 199) are both effective against gram-negative and gram-positive bacteria as well as fungi. In

some instances, neutral and/or anionic NCR peptides, such as NCR001, do not possess antimicrobial activities at a pI greater than about 8.

In some instances, the NCR peptide is effective to kill bacteria. In some instances, the NCR peptide is effective to kill *S. meliloti*, *Xenorhabdus spp*, *Photorhabdus spp*, *Candidatus spp*, *Buchnera*
 5 *spp*, *Blattabacterium spp*, *Baumania spp*, *Wigglesworthia spp*, *Wolbachia spp*, *Rickettsia spp*, *Orientia*
spp, *Sodalis spp*, *Burkholderia spp*, *Cupriavidus spp*, *Frankia spp*, *Snirrhizobium spp*, *Streptococcus spp*,
Wolinella spp, *Xylella spp*, *Erwinia spp*, *Agrobacterium spp*, *Bacillus spp*, *Paenibacillus spp*,
Streptomyces spp, *Micrococcus spp*, *Corynebacterium spp*, *Acetobacter spp*, *Cyanobacteria spp*,
Salmonella spp, *Rhodococcus spp*, *Pseudomonas spp*, *Lactobacillus spp*, *Enterococcus spp*, *Alcaligenes*
 10 *spp*, *Klebsiella spp*, *Paenibacillus spp*, *Arthrobacter spp*, *Corynebacterium spp*, *Brevibacterium spp*,
Thermus spp, *Pseudomonas spp*, *Clostridium spp*, or *Escherichia spp*.

In some instances, the NCR peptide is a functionally active variant of a NCR peptide described herein. In some instances, the variant of the NCR peptide has at least 70%, 71%, 72%, 73%, 74%, 75%,
 15 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%,
 94%, 95%, 96%, 97%, 98%, or 99% identity, e.g., over a specified region or over the entire sequence, to
 a sequence of a NCR peptide described herein or naturally derived NCR peptide.

In some instances, the NCR peptide may be bioengineered to modulate its bioactivity, e.g.,
 increase or decrease or regulate, or to specify a target microorganism. In some instances, the NCR
 peptide is produced by the translational machinery (e.g. a ribosome, etc.) of a cell. In some instances,
 20 the NCR peptide is chemically synthesized. In some instances, the NCR peptide is derived from a
 polypeptide precursor. The polypeptide precursor can undergo cleavage (for example, processing by a
 protease) to yield the NCR peptide itself. As such, in some instances, the NCR peptide is produced from
 a precursor polypeptide. In some instances, the NCR peptide includes a polypeptide that has undergone
 post-translational modifications, for example, cleavage, or the addition of one or more functional groups.

25 The NCR peptide described herein may be formulated in a composition for any of the uses
 described herein. The compositions disclosed herein may include any number or type of NCR peptides,
 such as at least about any one of 1 NCR peptide, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, 100, or more NCR
 peptides. A suitable concentration of each NCR peptide in the composition depends on factors such as
 efficacy, stability of the NCR peptide, number of distinct NCR peptide, the formulation, and methods of
 30 application of the composition. In some instances, each NCR peptide in a liquid composition is from
 about 0.1 ng/mL to about 100 mg/mL. In some instances, each NCR peptide in a solid composition is
 from about 0.1 ng/g to about 100 mg/g. In some instances, wherein the composition includes at least two
 types of NCR peptides, the concentration of each type of NCR peptide may be the same or different.

A modulating agent including a NCR peptide as described herein can be contacted with the target
 35 host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold
 level) of NCR peptide concentration inside a target host; (b) reach a target level (e.g., a predetermined or
 threshold level) of NCR peptide concentration inside a target host gut; (c) reach a target level (e.g., a
 predetermined or threshold level) of NCR peptide concentration inside a target host bacteriocyte; (d)
 modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host;
 40 or/and (e) modulate fitness of the target host.

(e) Bacteriocyte Regulatory Peptides

The modulating agent described herein may include a bacteriocyte regulatory peptide (BRP). BRPs are peptides expressed in the bacteriocytes of insects. These genes are expressed first at a developmental time point coincident with the incorporation of symbionts and their bacteriocyte-specific expression is maintained throughout the insect's life. In some instances, the BRP has a hydrophobic amino terminal domain, which is predicted to be a signal peptide. In addition, some BRPs have a cysteine-rich domain. In some instances, the bacteriocyte regulatory peptide is a bacteriocyte-specific cysteine rich (BCR) protein. Bacteriocyte regulatory peptides have a length between about 40 and 150 amino acids. In some instances, the bacteriocyte regulatory peptide has a length in the range of about 45 to about 145, about 50 to about 140, about 55 to about 135, about 60 to about 130, about 65 to about 125, about 70 to about 120, about 75 to about 115, about 80 to about 110, about 85 to about 105, or any range therebetween. Non-limiting examples of BRPs and their activities are listed in Table 8.

Table 8: Examples of Bacteriocyte Regulatory Peptides

Name	Peptide Sequence
Bacteriocyte-specific cysteine rich proteins BCR family, peptide BCR1	MKLLHGFLIIMLTMHLSIQYAYGGPFLTKYLCDRVC HKLCGDEFVCSCIQYKSLKGLWFPHCPTGKASVV LHNFLTSP (SEQ ID NO: 200)
Bacteriocyte-specific cysteine rich proteins BCR family, peptide BCR2	MKLLYGFLIIMLTIHLSVQYFESPFETKYNCIDTHCN KLCGKIDHCSCIQYHSM EGLWFPHCRTGSAAQML HDFLSNP (SEQ ID NO: 201)
Bacteriocyte-specific cysteine rich proteins BCR family, peptide BCR3	MSVRKNVLPTMFVLLIMSPVTPTSVFISAVCYSG CGSLALVCFVSNGITNGLDYFKSSAPLSTSETSCG EAFDCTDHCLANFKF (SEQ ID NO: 202)
Bacteriocyte-specific cysteine rich proteins BCR family, peptide BCR4	MRLLYGFLIIMLTIYLSVQDFDPTEFKGPFTIEICS KYCAVVCNYTSRPCYCVEAAKERDQWFPYCYD (SEQ ID NO: 203)
Bacteriocyte-specific cysteine rich proteins BCR family, peptide BCR5	MRLLYGFLIIMLTIHLSVQDIDPNTLRGPYPTKEICS KYCEYNNVCGASLPCICVQDARQLDHWFACCYD GGPEMLM (SEQ ID NO: 204)
Secreted proteins SP family, peptide SP1	MKLFVVVVLVAVGIMFVFASDTAAAPTDYEDTND MISLSSLVGDNSPYVRVSSADSGGSSKTSSKNPIL GLLKSVIKLLTKIFGTYSDAAPAMPPIPPALRKNRG MLA (SEQ ID NO: 205)
Secreted proteins SP family, peptide SP2	MVACKVILAVAVVFVAAVQGRPGGEPEWAAPIFA ELKSVSDNITNLVGLDNAGEYATAAKNNLNAFAES LKTEAAVFSKSFEGKASASDVFKESTKNFQAVVD

	TYIKNLPKDLTLKDFTEKSEQALKYMVEHGTEITKK AQGNTETEKEIKEFFKKQIENLIGQGKALQAKIAEA KKA (SEQ ID NO: 206)
Secreted proteins SP family, peptide SP3	MKTSSSKVFASCVAIVCLASVANALPVQKSVAATT ENPIVEKHGCRANKLVRQNVVDLKYDSMLITNE VVQKQSNEVQSSEQSNEGQNSEQSNEGQNSEQ SNEVQSSEHSNEGQNSKQSNEGQNSEQSNEVQ SSEHSNEGQNSEQSNEVQSSEHSNEGQNSKQS NEGQNSKQSNEVQSSEHWNNEGQNSKQSNEDQN SEQSNEGQNSKQSNEGQNSKQSNEDQNSEQSN EGQNSKQSNEVQSSEQSNEGQNSKQSNEGQSS EQSNEGQNSKQSNEVQSPPEHYDLPDESSYES EETKGSHEGDDSEHR (SEQ ID NO: 207)
Secreted proteins SP family, peptide SP4	MKTILGLCLFGALFWSTQSMPVGEVAPVPAVPS EAVPQKQVEAKPETNAASPVSDAKPESDSKPVDA EVKPTVSEVKAESQKPSGEPKPESDAKPVVASE SKPESDPKPAAVVESKPENDAVAPETNNDAPEN AAPVSENKPATDAKAETELIAQAKPESKPASDLK AEPEAAKPNSEVPVALPLNPTETKATQQSVETNQ VEQAAPAAAQADPAAAPAADPAPAPAAAPVAAEE AKLSESAPSTENKAAEPPSKPAEQQSAKPVEDAV PAASEISETKVSPAVPAVPEVPASPSAPAVADPVS APEAEKNAEPAKAANSAEPAVQSEAKPAEDIQKS GAVVSAENPKPVVEEQKPAEVAKPAEQSKSEAPAE APKPTEQSAAEPPKPESANDEKKEQHSVNKRDA TKEKKPTDSIMKKQKQKKAN (SEQ ID NO: 208)
Secreted proteins SP family, peptide SP5a	MNGKIVLCFAVVFIGQAMSAATGTTPEVEDIKKVA EQMSQTFMSVANHLVGITPNSADAQKSIEKIRTIM NKGFTDMETEANKMKDIVRKNADPKLVEKYDELE KELKKHLSTAKDMFEDKVVKPIGEKVELKKITENVI KTTKDMEATMNKAIDGFKKQ (SEQ ID NO: 209)
Secreted proteins SP family, peptide SP6	MHLFLALGLFIVCGMVDATFYNPRSQTFNQLMER RQRSIPYSGYHYNPIEPSINVLDLSEGLDSRI NTFKPIYQNVKMSTQDVNSVPRTQYQPKNSLYDS EYISAKDIPSLFPEEDSYDYKLGSPLNKYLTRPST QESGIAINLVAIKETSVFDYGFPTYKSPYSSDSVW NFGSKIPNTVFEDPQSVESDPNTFKVSSPTIKIVKL LPETPEQESIITTTKNYELNYKTTQETPTEAELYPIT SEEFQTEDEWHPMPKENTTKDESSFITTEEPLTE DKSNSITIEKTQTEDESNSIEFNSIRTEEKSNSITTE

	ENQKEDDESMSTTSQETTTAFNLNDFDTNRYSS SHESLMLRIELMKNIADQQNKSQFRTVDNIPAKS QSNLSSDESTNQQFEPQLVNGADTYK (SEQ ID NO: 210)
Colepotericin A, ColA peptide	MTRTMLFLACVAALYVCISATAGKPEEFAKLSDEA PSNDQAMYESIQRYRRFVDGNRYNGGQQQQQQ PKQWEVRPDLSDRQRGNTKAQVEINKKGDNDHI NAGWGKNINGPDSHKDTWHVGGSVRW (SEQ ID NO: 211)
RlpA type I	MKETTVVWAKLFLILILAKPLGLKAVNECKRLGNN SCRSHGECCSGFCFIEPGWALGVCKRLGTPKKS DDSNNGKNIKNNGVHERIDDVFERGVCSYYKGP SITANGDVFDEEMTAHRTLFPNTMVKVEGMGT SVVVKINDRKTAADGKVMLLSRAAAESLNIDENTG PVQCQLKFVLDGSGCTPDYGDTCVLHHECCSQN CFREMFSDKGFCLPK (SEQ ID NO: 212)

In some instances, the BRP alters the growth and/or activity of one or more bacteria resident in the bacteriocyte of the host. In some instances, the BRP may be bioengineered to modulate its bioactivity (e.g., increase, decrease, or regulate) or to specify a target microorganism. In some instances, the BRP is produced by the translational machinery (e.g. a ribosome, etc.) of a cell. In some instances, the BRP is chemically synthesized. In some instances, the BRP is derived from a polypeptide precursor. The polypeptide precursor can undergo cleavage (for example, processing by a protease) to yield the polypeptide of the BRP itself. As such, in some instances, the BRP is produced from a precursor polypeptide. In some instances, the BRP includes a polypeptide that has undergone post-translational modifications, for example, cleavage, or the addition of one or more functional groups.

Functionally active variants of the BRPs as described herein are also useful in the compositions and methods described herein. In some instances, the variant of the BRP has at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity, e.g., over a specified region or over the entire sequence, to a sequence of a BRP described herein or naturally derived BRP.

The BRP described herein may be formulated in a composition for any of the uses described herein. The compositions disclosed herein may include any number or type (e.g., classes) of BRPs, such as at least about any one of 1 BRP, 2, 3, 4, 5, 10, 15, 20, or more BRPs. A suitable concentration of each BRP in the composition depends on factors such as efficacy, stability of the BRP, number of distinct BRP, the formulation, and methods of application of the composition. In some instances, each BRP in a liquid composition is from about 0.1 ng/mL to about 100 mg/mL. In some instances, each BRP in a solid composition is from about 0.1 ng/g to about 100 mg/g. In some instances, wherein the composition includes at least two types of BRPs, the concentration of each type of BRP may be the same or different.

A modulating agent including a BRP as described herein can be contacted with the target host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold level) of

BRP concentration inside a target host; (b) reach a target level (e.g., a predetermined or threshold level) of BRP concentration inside a target host gut; (c) reach a target level (e.g., a predetermined or threshold level) of BRP concentration inside a target host bacteriocyte; (d) modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host; or/and (e) modulate fitness of the target host.

iii. Small Molecules

Numerous small molecules (e.g., an antibiotic or a metabolite) may be used in the compositions and methods described herein. In some instances, an effective concentration of any small molecule described herein may alter the level, activity, or metabolism of one or more microorganisms (as described herein) resident in a host, the alteration resulting in a decrease in the host's fitness.

A modulating agent comprising a small molecule as described herein can be contacted with the target host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold level) of a small molecule concentration inside a target host; (b) reach a target level (e.g., a predetermined or threshold level) of small molecule concentration inside a target host gut; (c) reach a target level (e.g., a predetermined or threshold level) of a small molecule concentration inside a target host bacteriocyte; (d) modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host; or/and (e) modulate fitness of the target host.

The small molecules discussed hereinafter, namely antibiotics and secondary metabolites, can be used to alter the level, activity, or metabolism of target microorganisms as indicated in the sections for decreasing the fitness of a host insect (e.g., vector of an animal pathogen), such as a mosquito, a mite, a louse, or a tick.

(a) Antibiotics

The modulating agent described herein may include an antibiotic. Any antibiotic known in the art may be used. Antibiotics are commonly classified based on their mechanism of action, chemical structure, or spectrum of activity.

The antibiotic described herein may target any bacterial function or growth processes and may be either bacteriostatic (e.g., slow or prevent bacterial growth) or bactericidal (e.g., kill bacteria). In some instances, the antibiotic is a bactericidal antibiotic. In some instances, the bactericidal antibiotic is one that targets the bacterial cell wall (e.g., penicillins and cephalosporins); one that targets the cell membrane (e.g., polymyxins); or one that inhibits essential bacterial enzymes (e.g., rifamycins, lipiarmycins, quinolones, and sulfonamides). In some instances, the bactericidal antibiotic is an aminoglycoside. In some instances, the antibiotic is a bacteriostatic antibiotic. In some instances the bacteriostatic antibiotic targets protein synthesis (e.g., macrolides, lincosamides and tetracyclines). Additional classes of antibiotics that may be used herein include cyclic lipopeptides (such as daptomycin), glycylcyclines (such as tigecycline), oxazolidinones (such as linezolid), or lipiarmycins (such as fidaxomicin). Examples of antibiotics include oxytetracycline, doxycycline, rifampicin, ciprofloxacin, ampicillin, and polymyxin B. Other non-limiting examples of antibiotics are found in Table 9.

Table 9: Examples of Antibiotics

Antibiotics	Action
Penicillins, cephalosporins, vancomycin	Cell wall synthesis
Polymixin, gramicidin	Membrane active agent, disrupt cell membrane
Tetracyclines, macrolides, chloramphenicol, clindamycin, spectinomycin	Inhibit protein synthesis
Sulfonamides	Inhibit folate-dependent pathways
Ciprofloxacin	Inhibit DNA-gyrase
Isoniazid, rifampicin, pyrazinamide, ethambutol, (myambutol), streptomycin	Antimycobacterial agents

The antibiotic described herein may have any level of target specificity (e.g., narrow- or broad-spectrum). In some instances, the antibiotic is a narrow-spectrum antibiotic, and thus targets specific types of bacteria, such as gram-negative or gram-positive bacteria. Alternatively, the antibiotic may be a broad-spectrum antibiotic that targets a wide range of bacteria.

The antibiotics described herein may be formulated in a composition for any of the uses described herein. The compositions disclosed herein may include any number or type (e.g., classes) of antibiotics, such as at least about any one of 1 antibiotic, 2, 3, 4, 5, 10, 15, 20, or more antibiotics (e.g., a combination of rifampicin and doxycycline, or a combination of ampicillin and rifampicin). A suitable concentration of each antibiotic in the composition depends on factors such as efficacy, stability of the antibiotic, number of distinct antibiotics, the formulation, and methods of application of the composition. In some instances, wherein the composition includes at least two types of antibiotics, the concentration of each type of antibiotic may be the same or different.

A modulating agent including an antibiotic as described herein can be contacted with the target host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold level) of antibiotic concentration inside a target host; (b) reach a target level (e.g., a predetermined or threshold level) of antibiotic concentration inside a target host gut; (c) reach a target level (e.g., a predetermined or threshold level) of antibiotic concentration inside a target host bacteriocyte; (d) modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host; or/and (e) modulate fitness of the target host.

As illustrated by Examples 1-4, 14, 26, and 27, antibiotics (e.g., doxycycline, oxytetracycline, azithromycin, ciprofloxacin, or rifampicin) can be used as modulating agents that target an endosymbiotic bacterium, such as a *Wolbachia* spp., in an insect host (e.g., an insect vector of an animal pathogen), such as a mosquito or mite or tick or biting louse, to decrease the fitness of the host (e.g., as outlined herein). As illustrated by Example 3, antibiotics such as oxytetracycline can be used as modulating agents that target an endosymbiotic bacterium, such as a *Rickettsia* spp., in an insect host, such as ticks, to decrease the fitness of the host (e.g., as outlined herein).

(b) Secondary Metabolites

In some instances, the modulating agent of the compositions and methods described herein includes a secondary metabolite. Secondary metabolites are derived from organic molecules produced by an organism. Secondary metabolites may act (i) as competitive agents used against bacteria, fungi, amoebae, plants, insects, and large animals; (ii) as metal transporting agents; (iii) as agents of symbiosis between microbes and plants, insects, and higher animals; (iv) as sexual hormones; and (v) as differentiation effectors. Non-limiting examples of secondary metabolites are found in Table 10.

Table 10: Examples of Secondary Metabolites

Phenyl-propanoids	Alkaloids	Terpenoids	Quinones	Steroids	Polyketides
Anthocyanins	Acridines	Carotenes	Anthro-quinones	Cardiac	Erythromycin
Coumarins	Betalaines	Monoterpenes	Bezo-quinones	Glycosides	Lovastatin and other statins
Flavonoids	Quinolozidines	Sesquiterpenes	Naphtho-quinones	Pregnen-olone	Discoder-molide
Hydroxy-cinnamoyl	Furono-quinones	Diterpenes		Derivatives	Aflatoxin B1
Derivatives	Harring-tonines	Triterpenes			Avermectins
Isoflavonoids	Isoquino-lines				Nystatin
Lignans	Indoles				Rifamycin
Phenolenones	Purines				
Proantho-cyanidins	Pyridines				
Stilbenes	Tropane				
Tanins	Alkaloids				

10

The secondary metabolite used herein may include a metabolite from any known group of secondary metabolites. For example, secondary metabolites can be categorized into the following groups: alkaloids, terpenoids, flavonoids, glycosides, natural phenols (e.g., gossypol acetic acid), enals (e.g., trans-cinnamaldehyde), phenazines, biphenols and dibenzofurans, polyketides, fatty acid synthase peptides, nonribosomal peptides, ribosomally synthesized and post-translationally modified peptides, polyphenols, polysaccharides (e.g., chitosan), and biopolymers. For an in-depth review of secondary metabolites see, for example, Vining, *Annu. Rev. Microbiol.* 44:395-427, 1990.

15

Secondary metabolites useful for compositions and methods described herein include those that alter a natural function of an endosymbiont (e.g., primary or secondary endosymbiont), bacteriocyte, or extracellular symbiont. In some instances, one or more secondary metabolites described herein is isolated from a high throughput screening (HTS) for antimicrobial compounds. For example, a HTS screen identified 49 antibacterial extracts that have specificity against gram positive and gram negative bacteria from over 39,000 crude extracts from organisms growing in diverse ecosystems of one specific region. In some instances, the secondary metabolite is transported inside a bacteriocyte.

20

In some instances, the small molecule is an inhibitor of vitamin synthesis. In some instances, the vitamin synthesis inhibitor is a vitamin precursor analog. In certain instances, the vitamin precursor analog is pantothenol.

In some instances, the small molecule is an amino acid analog. In certain instances, the amino acid analog is L-canvanine, D-arginine, D-valine, D-methionine, D-phenylalanine, D-histidine, D-tryptophan, D-threonine, D-leucine, L-NG-nitroarginine, or a combination thereof.

In some instances the small molecule is a natural antimicrobial compound, such as propionic acid, levulinic acid, trans-cinnamaldehyde, nisin, or low molecular weight chitosan. The secondary metabolite described herein may be formulated in a composition for any of the uses described herein. The compositions disclosed herein may include any number or type (e.g., classes) of secondary metabolites, such as at least about any one of 1 secondary metabolite, 2, 3, 4, 5, 10, 15, 20, or more secondary metabolites. A suitable concentration of each secondary metabolite in the composition depends on factors such as efficacy, stability of the secondary metabolite, number of distinct secondary metabolites, the formulation, and methods of application of the composition. In some instances, wherein the composition includes at least two types of secondary metabolites, the concentration of each type of secondary metabolite may be the same or different.

A modulating agent including a secondary metabolite as described herein can be contacted with the target host in an amount and for a time sufficient to: (a) reach a target level (e.g., a predetermined or threshold level) of secondary metabolite concentration inside a target host; (b) reach a target level (e.g., a predetermined or threshold level) of secondary metabolite concentration inside a target host gut; (c) reach a target level (e.g., a predetermined or threshold level) of secondary metabolite concentration inside a target host bacteriocyte; (d) modulate the level, or an activity, of one or more microorganism (e.g., endosymbiont) in the target host; or/and (e) modulate fitness of the target host.

As illustrated by Example 15, secondary metabolites (e.g., gossypol) can be used as modulating agents that target an endosymbiotic bacterium in an insect host to decrease the fitness of the host (e.g., as outlined herein). As further illustrated by Examples 11-13, 15-19, 23, and 24, small molecules, such as trans-cinnamaldehyde, levulinic acid, chitosan, vitamin analogs, or amino acid transport inhibitors, can be used as modulating agents that target an endosymbiotic bacterium in an insect host to decrease the fitness of the host (e.g., as outlined herein).

iv. Bacteria as modulating agents

In some instances, the modulating agent described herein includes one or more bacteria. Numerous bacteria are useful in the compositions and methods described herein. In some instances, the agent is a bacterial species endogenously found in the host. In some instances, the bacterial modulating agent is an endosymbiotic bacterial species. Non-limiting examples of bacteria that may be used as modulating agents include all bacterial species described herein in Section II of the detailed description and those listed in Table 1. For example, the modulating agent may be a bacterial species from any bacterial phyla present in insect guts, including *Gammaproteobacteria*, *Alphaproteobacteria*, *Betaproteobacteria*, *Bacteroidetes*, *Firmicutes* (e.g., *Lactobacillus* and *Bacillus* spp.), *Clostridia*, *Actinomycetes*, *Spirochetes*, *Verrucomicrobia*, and *Actinobacteria*.

In some instances, the modulating agent is a bacterium that disrupts microbial diversity or otherwise alters the microbiota of the host in a manner detrimental to the host. In one instance, bacteria may be provided to disrupt the microbiota of mosquitos. For example, the bacterial modulating agent may compete with, displace, and/or reduce a population of symbiotic bacteria in a mosquito.

5 In another instance, bacteria may be provided to disrupt the microbiota of mites. For example, the bacterial modulating agent may compete with, displace, and/or reduce a population of symbiotic bacteria in a mite.

In another instance, bacteria may be provided to disrupt the microbiota of biting lice. For example, the bacterial modulating agent may compete with, displace, and/or reduce a population of symbiotic bacteria in a biting louse.

10 In another instance, bacteria may be provided to disrupt the microbiota of ticks. For example, the bacterial modulating agent may compete with, displace, and/or reduce a population of symbiotic bacteria in a tick.

The bacterial modulating agents discussed herein can be used to alter the level, activity, or metabolism of target microorganisms as indicated in the sections for decreasing the fitness of a host insect (e.g., a vector of an animal pathogen), such as a mosquito a mite, a biting louse, or a tick.

v. Modifications to modulating agents

(a) Fusions

20 Any of the modulating agents described herein may be fused or linked to an additional moiety. In some instances, the modulating agent includes a fusion of one or more additional moieties (e.g., 1 additional moiety, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more additional moieties). In some instances, the additional moiety is any one of the modulating agents described herein (e.g., a peptide, polypeptide, small molecule, or antibiotic). Alternatively, the additional moiety may not act as modulating agent itself but may instead

25 serve a secondary function. For example, the additional moiety may to help the modulating agent access, bind, or become activated at a target site in the host (e.g., at a host gut or a host bacteriocyte) or at a target microorganism resident in the host (e.g., a vector of an animal pathogen, e.g., a mosquito, a mite, a biting louse, or a tick).

In some instances, the additional moiety may help the modulating agent penetrate a target host cell or target microorganism resident in the host. For example, the additional moiety may include a cell penetrating peptide. Cell penetrating peptides (CPPs) may be natural sequences derived from proteins; chimeric peptides that are formed by the fusion of two natural sequences; or synthetic CPPs, which are synthetically designed sequences based on structure–activity studies. In some instances, CPPs have the capacity to ubiquitously cross cellular membranes (e.g., prokaryotic and eukaryotic cellular membranes)

30 with limited toxicity. Further, CPPs may have the capacity to cross cellular membranes via energy-dependent and/or independent mechanisms, without the necessity of a chiral recognition by specific receptors. CPPs can be bound to any of the modulating agents described herein. For example, a CPP can be bound to an antimicrobial peptide (AMP), e.g., a scorpion peptide, e.g., UY192 fused to a cell penetrating peptide (e.g., YGRKKRRQRRRFLSTIWNIGIKLLFAM; SEQ ID NO: 232). Non-limiting

35 examples of CPPs are listed in Table 11.

40

Table 11: Examples of Cell Penetrating Peptides (CPPs)

Peptide	Origin	Sequence
<i>Protein-derived</i>		
Penetratin	Antennapedia	RQIKIWFQNRRMKWKK (SEQ ID NO: 213)
Tat peptide	Tat	GRKKRRQRRRPQ (SEQ ID NO: 214)
pVEC	Cadherin	LLIILRRRIRKQAHASK (SEQ ID NO: 215)
<i>Chimeric</i>		
Transportan	Galanine/Mastoparan	GWTLNSAGYLLGKINLKALAALAKKIL (SEQ ID NO: 216)
MPG	HIV-gp41/SV40 T-antigen	GALFLGFLGAAGSTMGAWSQPKKKRKV (SEQ ID NO: 217)
Pep-1	HIV-reverse transcriptase/SV40 T-antigen	KETWWETWWTEWSQPKKKRKV (SEQ ID NO: 218)
<i>Synthetic</i>		
Polyarginines	Based on Tat peptide	$(R)_n; 6 < n < 12$
MAP	de novo	KLALKLALKALKALKLA (SEQ ID NO: 219)
R ₆ W ₃	Based on penetratin	RRWWRRWRR (SEQ ID NO: 220)

In other instances, the additional moiety helps the modulating agent bind a target microorganism (e.g., a fungi or bacterium) resident in the host. The additional moiety may include one or more targeting domains. In some instances, the targeting domain may target the modulating agent to one or more microorganisms (e.g., bacterium or fungus) resident in the gut of the host. In some instances, the targeting domain may target the modulating agent to a specific region of the host (e.g., host gut or bacteriocyte) to access microorganisms that are generally present in said region of the host. For example, the targeting domain may target the modulating agent to the foregut, midgut, or hindgut of the host. In other instances, the targeting domain may target the modulating agent to a bacteriocyte in the host and/or one or more specific bacteria resident in a host bacteriocyte. For example, the targeting domain may be Galanthus nivalis lectin or agglutinin (GNA) bound to a modulating agent described herein, e.g., an AMP, e.g., a scorpion peptide, e.g., Uy192.

15

(b) Pre- or Pro-domains

In some instances, the modulating agent may include a pre- or pro- amino acid sequence. For example, the modulating agent may be an inactive protein or peptide that can be activated by cleavage or post-translational modification of a pre- or pro-sequence. In some instances, the modulating agent is

engineered with an inactivating pre- or pro-sequence. For example, the pre- or pro-sequence may obscure an activation site on the modulating agent, e.g., a receptor binding site, or may induce a conformational change in the modulating agent. Thus, upon cleavage of the pre- or pro-sequence, the modulating agent is activated.

5 Alternatively, the modulating agent may include a pre- or pro-small molecule, e.g., an antibiotic. The modulating agent may be an inactive small molecule described herein that can be activated in a target environment inside the host. For example, the small molecule may be activated upon reaching a certain pH in the host gut.

10 (c) Linkers

In instances where the modulating agent is connected to an additional moiety, the modulating agent may further include a linker. For example, the linker may be a chemical bond, e.g., one or more covalent bonds or non-covalent bonds. In some instances, the linker may be a peptide linker (e.g., 2, 3, 4, 5, 6, 8, 10, 12, 14, 16, 20, 25, 30, 35, 40, or more amino acids longer). The linker may include any 15 flexible, rigid, or cleavable linkers described herein.

A flexible peptide linker may include any of those commonly used in the art, including linkers having sequences having primarily Gly and Ser residues ("GS" linker). Flexible linkers may be useful for joining domains that require a certain degree of movement or interaction and may include small, non-polar (e.g. Gly) or polar (e.g. Ser or Thr) amino acids.

20 Alternatively, a peptide linker may be a rigid linker. Rigid linkers are useful to keep a fixed distance between moieties and to maintain their independent functions. Rigid linkers may also be useful when a spatial separation of the domains is critical to preserve the stability or bioactivity of one or more components in the fusion. Rigid linkers may, for example, have an alpha helix-structure or Pro-rich sequence, (XP)_n, with X designating any amino acid, preferably Ala, Lys, or Glu.

25 In yet other instances, a peptide linker may be a cleavable linker. In some instances, linkers may be cleaved under specific conditions, such as the presence of reducing reagents or proteases. *In vivo* cleavable linkers may utilize the reversible nature of a disulfide bond. One example includes a thrombin-sensitive sequence (e.g., PRS) between two Cys residues. *In vitro* thrombin treatment of CPRSC results in the cleavage of the thrombin-sensitive sequence, while the reversible disulfide linkage remains intact. 30 Such linkers are known and described, e.g., in Chen et al., *Adv. Drug Deliv. Rev.* 65(10):1357-1369, 2013. Cleavage of linkers in fusions may also be carried out by proteases that are expressed *in vivo* under conditions in specific cells or tissues of the host or microorganisms resident in the host. In some instances, cleavage of the linker may release a free functional, modulating agent upon reaching a target site or cell.

35 Fusions described herein may alternatively be linked by a linking molecule, including a hydrophobic linker, such as a negatively charged sulfonate group; lipids, such as a poly (--CH₂--) hydrocarbon chains, such as polyethylene glycol (PEG) group, unsaturated variants thereof, hydroxylated variants thereof, amidated or otherwise N-containing variants thereof, non-carbon linkers; carbohydrate linkers; phosphodiester linkers, or other molecule capable of covalently linking two or more molecules, 40 e.g., two modulating agents. Non-covalent linkers may be used, such as hydrophobic lipid globules to

which the modulating agent is linked, for example, through a hydrophobic region of the modulating agent or a hydrophobic extension of the modulating agent, such as a series of residues rich in leucine, isoleucine, valine, or perhaps also alanine, phenylalanine, or even tyrosine, methionine, glycine, or other hydrophobic residue. The modulating agent may be linked using charge-based chemistry, such that a
5 positively charged moiety of the modulating agent is linked to a negative charge of another modulating agent or an additional moiety.

IV. Formulations and Compositions

The compositions described herein may be formulated either in pure form (e.g., the composition
10 contains only the modulating agent) or together with one or more additional agents (such as excipient, delivery vehicle, carrier, diluent, stabilizer, etc.) to facilitate application or delivery of the compositions. Examples of suitable excipients and diluents include, but are not limited to, lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, saline solution, syrup,
15 methylcellulose, methyl- and propylhydroxybenzoates, talc, magnesium stearate, and mineral oil.

In some instances, the composition includes a delivery vehicle or carrier. In some instances, the delivery vehicle includes an excipient. Exemplary excipients include, but are not limited to, solid or liquid carrier materials, solvents, stabilizers, slow-release excipients, colorings, and surface-active substances (surfactants). In some instances, the delivery vehicle is a stabilizing vehicle. In some instances, the
20 stabilizing vehicle includes a stabilizing excipient. Exemplary stabilizing excipients include, but are not limited to, epoxidized vegetable oils, antifoaming agents, e.g. silicone oil, preservatives, viscosity regulators, binding agents and tackifiers. In some instances, the stabilizing vehicle is a buffer suitable for the modulating agent. In some instances, the composition is microencapsulated in a polymer bead delivery vehicle. In some instances, the stabilizing vehicle protects the modulating agent against UV
25 and/or acidic conditions. In some instances, the delivery vehicle contains a pH buffer. In some instances, the composition is formulated to have a pH in the range of about 4.5 to about 9.0, including for example pH ranges of about any one of 5.0 to about 8.0, about 6.5 to about 7.5, or about 6.5 to about 7.0.

Depending on the intended objectives and prevailing circumstances, the composition may be formulated into emulsifiable concentrates, suspension concentrates, directly sprayable or dilutable
30 solutions, coatable pastes, diluted emulsions, spray powders, soluble powders, dispersible powders, wettable powders, dusts, granules, encapsulations in polymeric substances, microcapsules, foams, aerosols, carbon dioxide gas preparations, tablets, resin preparations, paper preparations, nonwoven fabric preparations, or knitted or woven fabric preparations. In some instances, the composition is a liquid. In some instances, the composition is a solid. In some instances, the composition is an aerosol, such as
35 in a pressurized aerosol can. In some instances, the composition is present in the waste (such as feces) of the pest. In some instances, the composition is present in or on a live pest.

In some instances, the delivery vehicle is the food or water of the host. In other instances, the delivery vehicle is a food source for the host. In some instances, the delivery vehicle is a food bait for the host. In some instances, the composition is a comestible agent consumed by the host. In some
40 instances, the composition is delivered by the host to a second host, and consumed by the second host.

In some instances, the composition is consumed by the host or a second host, and the composition is released to the surrounding of the host or the second host via the waste (such as feces) of the host or the second host. In some instances, the modulating agent is included in food bait intended to be consumed by a host or carried back to its colony.

5 In some instances, the delivery vehicle is a bacterial vector. The modulating agent can be incorporated in a bacterial vector using any suitable cloning methods and reagents known in the art, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989. "Bacterial vector" as used herein refers to any genetic element, such as plasmids, bacteriophage vectors, transposons, cosmids, and chromosomes, which is
10 capable of replication inside bacterial cells and which is capable of transferring genes between cells. Exemplary bacterial vectors include, but are not limited to, lambda vector system gtl 1, gt WES.tB, Charon 4, and plasmid vectors such as pBR322, pBR325, pACYC177, pACYC184, pUC8, pUC9, pUC18, pUC19, pLG339, pR290, pKC37, pKCIOL, SV 40, pBluescript II SK +/- or KS +/- (see "Stratagene Cloning Systems" Catalog, Stratagene, La Jolla, California, 1993), pQE, pIH821, pGEX, pET series (see Studier
15 et al., "Use of T7 RNA Polymerase to Direct Expression of Cloned Genes," *Gene Expression Technology* Vol. 185, 1990), and any derivatives thereof.

Each bacterial vector may encode one or more modulating agents. In some instances, the bacterial vector includes a phage genome to be expressed and packaged in the target symbiotic bacterium. In some instances, the bacterial vector includes a nucleic acid molecule encoding a lysin to
20 be expressed in the target symbiotic bacterium or a host bacterium. In some instances, the lysin is co-expressed with a holin, or the lysin is engineered to have a signal peptide for secretion from the host bacterium. In some instances, the bacterial vector includes a nucleic acid molecule encoding a bacteriocin to be expressed in the target symbiotic bacterium. In some instances, the bacterial vector further includes one or more regulatory elements, such as promoters, termination signals, and
25 transcription and translation elements. In some instances, the regulatory sequence is operably linked to a nucleic acid encoding a gene (such as a bacteriocin, lysin, or other polypeptides) to be expressed in the target symbiotic bacterium.

In some instances, the bacterial vector is introduced into a bacterium to be consumed by the host or a member in the colony of the host. In some instances, the bacterium is the target symbiotic
30 bacterium. In some instances, the bacterium is a naturally occurring bacterium of the gut of the host, or a genetically modified derivative thereof, which can be easily introduced to the host through ingestion. Exemplary bacteria for use in carrying the bacterial vector include, but are not limited to, *Proteobacter*, including the genus *Pseudomonas*; *Actinobacter*, including *Priopionibacterium* and *Corynebacterium*; *Firmicutes*, including the any species of the genera *Mycoplasma*, *Bacillus*, *Streptococcus*,
35 *Staphylococcus*; *Fibrobacteres*; *Spirochaetes*, including *Treponema* and *Borrelia*; *Bacteroides*, including the genera *Bacteroides* and *Flavobacterium*. Also suitable are any bacteria of the *Enterobacteriaceae*, including the genus *Serratia*, including, but not limited to *S. marcescens*, *S. entomophila*, *S. proteamaculans*, *S. marcescens*; any species of *Enterobacter*, including, but not limited to, *E. cloacae*, *E. amnigenus*, *E. aerogenes*, *E. dissolvens*, *E. agglomerans*, *E. hafiiiae*; and any species belonging to the

following genera: *Citrobacter*, *Escherichia*, *Klebsiella*, *Kluyvera*, *Parabacterium*, *Proteus*, *Salmonella*, *Xenorhabdus*, and *Yokenella*.

In some instances, the modulating agent may make up about 0.1% to about 100% of the composition, such as any one of about 0.01% to about 100%, about 1% to about 99.9%, about 0.1% to about 10%, about 1% to about 25%, about 10% to about 50%, about 50% to about 99%, or about 0.1% to about 90% of active ingredients (such as phage, lysin or bacteriocin). In some instances, the composition includes at least any of 0.1%, 0.5%, 1%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or more active ingredients (such as phage, lysin or bacteriocin). In some instances, the concentrated agents are preferred as commercial products, the final user normally uses diluted agents, which have a substantially lower concentration of active ingredient.

Any of the formulations described herein may be used in the form of a bait, a coil, an electric mat, a smoking preparation, a fumigant, or a sheet.

i. Liquid Formulations

The compositions provided herein may be in a liquid formulation. Liquid formulations are generally mixed with water, but in some instances may be used with crop oil, diesel fuel, kerosene or other light oil as a carrier. The amount of active ingredient often ranges from about 0.5 to about 80 percent by weight.

An emulsifiable concentrate formulation may contain a liquid active ingredient, one or more petroleum-based solvents, and an agent that allows the formulation to be mixed with water to form an emulsion. Such concentrates may be used in agricultural, ornamental and turf, forestry, structural, food processing, livestock, and public health pest formulations. These may be adaptable to application equipment from small portable sprayers to hydraulic sprayers, low-volume ground sprayers, mist blowers, and low-volume aircraft sprayers. Some active ingredients are readily dissolve in a liquid carrier. When mixed with a carrier, they form a solution that does not settle out or separate, e.g., a homogenous solution. Formulations of these types may include an active ingredient, a carrier, and one or more other ingredients. Solutions may be used in any type of sprayer, indoors and outdoors.

In some instances, the composition may be formulated as an invert emulsion. An invert emulsion is a water-soluble active ingredient dispersed in an oil carrier. Invert emulsions require an emulsifier that allows the active ingredient to be mixed with a large volume of petroleum-based carrier, usually fuel oil. Invert emulsions aid in reducing drift. With other formulations, some spray drift results when water droplets begin to evaporate before reaching target surfaces; as a result the droplets become very small and lightweight. Because oil evaporates more slowly than water, invert emulsion droplets shrink less and more active ingredient reaches the target. Oil further helps to reduce runoff and improve rain resistance. It further serves as a sticker-spreader by improving surface coverage and absorption. Because droplets are relatively large and heavy, it is difficult to get thorough coverage on the undersides of foliage. Invert emulsions are most commonly used along rights-of-way where drift to susceptible non-target areas can be a problem.

A flowable or liquid formulation combines many of the characteristics of emulsifiable concentrates and wettable powders. Manufacturers use these formulations when the active ingredient is a solid that

does not dissolve in either water or oil. The active ingredient, impregnated on a substance such as clay, is ground to a very fine powder. The powder is then suspended in a small amount of liquid. The resulting liquid product is quite thick. Flowables and liquids share many of the features of emulsifiable concentrates, and they have similar disadvantages. They require moderate agitation to keep them in suspension and leave visible residues, similar to those of wettable powders.

Flowables/liquids are easy to handle and apply. Because they are liquids, they are subject to spilling and splashing. They contain solid particles, so they contribute to abrasive wear of nozzles and pumps. Flowable and liquid suspensions settle out in their containers. Because flowable and liquid formulations tend to settle, packaging in containers of five gallons or less makes remixing easier.

Aerosol formulations contain one or more active ingredients and a solvent. Most aerosols contain a low percentage of active ingredients. There are two types of aerosol formulations—the ready-to-use type commonly available in pressurized sealed containers and those products used in electrical or gasoline-powered aerosol generators that release the formulation as a smoke or fog.

Ready to use aerosol formulations are usually small, self-contained units that release the formulation when the nozzle valve is triggered. The formulation is driven through a fine opening by an inert gas under pressure, creating fine droplets. These products are used in greenhouses, in small areas inside buildings, or in localized outdoor areas. Commercial models, which hold five to 5 pounds of active ingredient, are usually refillable.

Smoke or fog aerosol formulations are not under pressure. They are used in machines that break the liquid formulation into a fine mist or fog (aerosol) using a rapidly whirling disk or heated surface.

ii. Dry or Solid Formulations

Dry formulations can be divided into two types: ready-to-use and concentrates that must be mixed with water to be applied as a spray. Most dust formulations are ready to use and contain a low percentage of active ingredients (less than about 10 percent by weight), plus a very fine, dry inert carrier made from talc, chalk, clay, nut hulls, or volcanic ash. The size of individual dust particles varies. A few dust formulations are concentrates and contain a high percentage of active ingredients. Mix these with dry inert carriers before applying. Dusts are always used dry and can easily drift to non-target sites.

iii. Granule or Pellet Formulations

In some instances, the composition is formulated as granules. Granular formulations are similar to dust formulations, except granular particles are larger and heavier. The coarse particles may be made from materials such as clay, corncobs, or walnut shells. The active ingredient either coats the outside of the granules or is absorbed into them. The amount of active ingredient may be relatively low, usually ranging from about 0.5 to about 15 percent by weight. Granular formulations are most often used to apply to the soil, insects living in the soil, or absorption into plants through the roots. Granular formulations are sometimes applied by airplane or helicopter to minimize drift or to penetrate dense vegetation. Once applied, granules may release the active ingredient slowly. Some granules require soil moisture to release the active ingredient. Granular formulations also are used to control larval

mosquitoes and other aquatic pests. Granules are used in agricultural, structural, ornamental, turf, aquatic, right-of-way, and public health (biting insect) pest-control operations.

In some instances, the composition is formulated as pellets. Most pellet formulations are very similar to granular formulations; the terms are used interchangeably. In a pellet formulation, however, all the particles are the same weight and shape. The uniformity of the particles allows use with precision application equipment.

iv. Powders

In some instances, the composition is formulated as a powder. In some instances, the composition is formulated as a wettable powder. Wettable powders are dry, finely ground formulations that look like dusts. They usually must be mixed with water for application as a spray. A few products, however, may be applied either as a dust or as a wettable powder—the choice is left to the applicator. Wettable powders have about 1 to about 95 percent active ingredient by weight; in some cases more than about 50 percent. The particles do not dissolve in water. They settle out quickly unless constantly agitated to keep them suspended. They can be used for most pest problems and in most types of spray equipment where agitation is possible. Wettable powders have excellent residual activity. Because of their physical properties, most of the formulation remains on the surface of treated porous materials such as concrete, plaster, and untreated wood. In such cases, only the water penetrates the material.

In some instances, the composition is formulated as a soluble powder. Soluble powder formulations look like wettable powders. However, when mixed with water, soluble powders dissolve readily and form a true solution. After they are mixed thoroughly, no additional agitation is necessary. The amount of active ingredient in soluble powders ranges from about 15 to about 95 percent by weight; in some cases more than about 50 percent. Soluble powders have all the advantages of wettable powders and none of the disadvantages, except the inhalation hazard during mixing.

In some instances, the composition is formulated as a water-dispersible granule. Water-dispersible granules, also known as dry flowables, are like wettable powders, except instead of being dust-like, they are formulated as small, easily measured granules. Water-dispersible granules must be mixed with water to be applied. Once in water, the granules break apart into fine particles similar to wettable powders. The formulation requires constant agitation to keep it suspended in water. The percentage of active ingredient is high, often as much as 90 percent by weight. Water-dispersible granules share many of the same advantages and disadvantages of wettable powders, except they are more easily measured and mixed. Because of low dust, they cause less inhalation hazard to the applicator during handling

v. Bait

In some instances, the composition includes a bait. The bait can be in any suitable form, such as a solid, paste, pellet or powdered form. The bait can also be carried away by the host back to a population of said host (e.g., a colony or hive). The bait can then act as a food source for other members of the colony, thus providing an effective modulating agent for a large number of hosts and potentially an entire host colony.

The baits can be provided in a suitable "housing" or "trap." Such housings and traps are commercially available and existing traps can be adapted to include the compositions described herein. The housing or trap can be box-shaped for example, and can be provided in pre-formed condition or can be formed of foldable cardboard for example. Suitable materials for a housing or trap include plastics and cardboard, particularly corrugated cardboard. The inside surfaces of the traps can be lined with a sticky substance in order to restrict movement of the host once inside the trap. The housing or trap can contain a suitable trough inside which can hold the bait in place. A trap is distinguished from a housing because the host cannot readily leave a trap following entry, whereas a housing acts as a "feeding station" which provides the host with a preferred environment in which they can feed and feel safe from predators.

vi. Attractants

In some instances, the composition includes an attractant (e.g., a chemoattractant). The attractant may attract an adult host or immature host (e.g., larva) to the vicinity of the composition. Attractants include pheromones, a chemical that is secreted by an animal, especially an insect, which influences the behavior or development of others of the same species. Other attractants include sugar and protein hydrolysate syrups, yeasts, and rotting meat. Attractants also can be combined with an active ingredient and sprayed onto foliage or other items in the treatment area.

Various attractants are known which influence host behavior as a host's search for food, oviposition or mating sites, or mates. Attractants useful in the methods and compositions described herein include, for example, eugenol, phenethyl propionate, ethyl dimethylisobutyl-cyclopropane carboxylate, propyl benzodioxanecarboxylate, cis-7,8-epoxy-2-methyloctadecane, trans-8,trans-0-dodecadienol, cis-9-tetradecenal (with cis-11-hexadecenal), trans-11-tetradecenal, cis-11-hexadecenal, (Z)-11,12-hexadecadienal, cis-7-dodecenyl acetate, cis-8-dodecenyl acetate, cis-9-dodecenyl acetate, cis-9-tetradecenyl acetate, cis-11-tetradecenyl acetate, trans-11-tetradecenyl acetate (with cis-11), cis-9,trans-11-tetradecadienyl acetate (with cis-9,trans-12), cis-9,trans-11-tetradecadienyl acetate, cis-7,cis-11-hexadecadienyl acetate (with cis-7,trans-11), cis-3,cis-13-octadecadienyl acetate, trans-3,cis-13-octadecadienyl acetate, anethole and isoamyl salicylate.

Means other than chemoattractants may also be used to attract insects, including lights in various wavelengths or colors.

vii. Nanocapsules/Microencapsulation/Liposomes

In some instances, the composition is provided in a microencapsulated formulation. Microencapsulated formulations are mixed with water and sprayed in the same manner as other sprayable formulations. After spraying, the plastic coating breaks down and slowly releases the active ingredient.

viii. Carriers

Any of the compositions described herein may be formulated to include the modulating agent described herein and an inert carrier. Such carrier can be a solid carrier, a liquid carrier, a gel carrier, and/or a gaseous carrier. In certain instances, the carrier can be a seed coating. The seed coating is

any non-naturally occurring formulation that adheres, in whole or part, to the surface of the seed. The formulation may further include an adjuvant or surfactant. The formulation can also include one or more modulating agents to enlarge the action spectrum.

A solid carrier used for formulation includes finely-divided powder or granules of clay (e.g. kaolin clay, diatomaceous earth, bentonite, Fubasami clay, acid clay, etc.), synthetic hydrated silicon oxide, talc, ceramics, other inorganic minerals (e.g., sericite, quartz, sulfur, activated carbon, calcium carbonate, hydrated silica, etc.), a substance which can be sublimated and is in the solid form at room temperature (e.g., 2,4,6-triisopropyl-1,3,5-trioxane, naphthalene, p-dichlorobenzene, camphor, adamantane, etc.); wool; silk; cotton; hemp; pulp; synthetic resins (e.g., polyethylene resins such as low-density polyethylene, straight low-density polyethylene and high-density polyethylene; ethylene-vinyl ester copolymers such as ethylene-vinyl acetate copolymers; ethylene-methacrylic acid ester copolymers such as ethylene-methyl methacrylate copolymers and ethylene-ethyl methacrylate copolymers; ethylene-acrylic acid ester copolymers such as ethylene-methyl acrylate copolymers and ethylene-ethyl acrylate copolymers; ethylene-vinylcarboxylic acid copolymers such as ethylene-acrylic acid copolymers; ethylene-tetracyclododecene copolymers; polypropylene resins such as propylene homopolymers and propylene-ethylene copolymers; poly-4-methylpentene-1, polybutene-1, polybutadiene, polystyrene; acrylonitrile-styrene resins; styrene elastomers such as acrylonitrile-butadiene-styrene resins, styrene-conjugated diene block copolymers, and styrene-conjugated diene block copolymer hydrides; fluororesins; acrylic resins such as poly(methyl methacrylate); polyamide resins such as nylon 6 and nylon 66; polyester resins such as polyethylene terephthalate, polyethylene naphthalate, polybutylene terephthalate, and polycyclohexylenedimethylene terephthalate; polycarbonates, polyacetals, polyacrylsulfones, polyarylates, hydroxybenzoic acid polyesters, polyetherimides, polyester carbonates, polyphenylene ether resins, polyvinyl chloride, polyvinylidene chloride, polyurethane, and porous resins such as foamed polyurethane, foamed polypropylene, or foamed ethylene, etc.), glasses, metals, ceramics, fibers, cloths, knitted fabrics, sheets, papers, yarn, foam, porous substances, and multifilaments.

A liquid carrier may include, for example, aromatic or aliphatic hydrocarbons (e.g., xylene, toluene, alkyl naphthalene, phenylxylethane, kerosene, gas oil, hexane, cyclohexane, etc.), halogenated hydrocarbons (e.g., chlorobenzene, dichloromethane, dichloroethane, trichloroethane, etc.), alcohols (e.g., methanol, ethanol, isopropyl alcohol, butanol, hexanol, benzyl alcohol, ethylene glycol, etc.), ethers (e.g., diethyl ether, ethylene glycol dimethyl ether, diethylene glycol monomethyl ether, diethylene glycol monoethyl ether, propylene glycol monomethyl ether, tetrahydrofuran, dioxane, etc.), esters (e.g., ethyl acetate, butyl acetate, etc.), ketones (e.g., acetone, methyl ethyl ketone, methyl isobutyl ketone, cyclohexanone, etc.), nitriles (e.g., acetonitrile, isobutyronitrile, etc.), sulfoxides (e.g., dimethyl sulfoxide, etc.), amides (e.g., N,N-dimethylformamide, N,N-dimethylacetamide, cyclic imides (e.g. N-methylpyrrolidone) alkylidene carbonates (e.g., propylene carbonate, etc.), vegetable oil (e.g., soybean oil, cottonseed oil, etc.), vegetable essential oils (e.g., orange oil, hyssop oil, lemon oil, etc.), or water.

A gaseous carrier may include, for example, butane gas, flon gas, liquefied petroleum gas (LPG), dimethyl ether, and carbon dioxide gas.

ix. Adjuvants

In some instances, the composition provided herein may include an adjuvant. Adjuvants are chemicals that do not possess activity. Adjuvants are either pre-mixed in the formulation or added to the spray tank to improve mixing or application or to enhance performance. They are used extensively in products designed for foliar applications. Adjuvants can be used to customize the formulation to specific needs and compensate for local conditions. Adjuvants may be designed to perform specific functions, including wetting, spreading, sticking, reducing evaporation, reducing volatilization, buffering, emulsifying, dispersing, reducing spray drift, and reducing foaming. No single adjuvant can perform all these functions, but compatible adjuvants often can be combined to perform multiple functions simultaneously.

Among nonlimiting examples of adjuvants included in the formulation are binders, dispersants and stabilizers, specifically, for example, casein, gelatin, polysaccharides (e.g., starch, gum arabic, cellulose derivatives, alginic acid, etc.), lignin derivatives, bentonite, sugars, synthetic water-soluble polymers (e.g., polyvinyl alcohol, polyvinylpyrrolidone, polyacrylic acid, etc.), PAP (acidic isopropyl phosphate), BHT (2,6-di-t-butyl-4-methylphenol), BHA (a mixture of 2-t-butyl-4-methoxyphenol and 3-t-butyl-4-methoxyphenol), vegetable oils, mineral oils, fatty acids and fatty acid esters.

x. Surfactants

In some instances, the composition provided herein includes a surfactant. Surfactants, also called wetting agents and spreaders, physically alter the surface tension of a spray droplet. For a formulation to perform its function properly, a spray droplet must be able to wet the foliage and spread out evenly over a leaf. Surfactants enlarge the area of formulation coverage, thereby increasing the pest's exposure to the chemical. Surfactants are particularly important when applying a formulation to waxy or hairy leaves. Without proper wetting and spreading, spray droplets often run off or fail to cover leaf surfaces adequately. Too much surfactant, however, can cause excessive runoff and reduce efficacy.

Surfactants are classified by the way they ionize or split apart into electrically charged atoms or molecules called ions. A surfactant with a negative charge is anionic. One with a positive charge is cationic, and one with no electrical charge is nonionic. Formulation activity in the presence of a nonionic surfactant can be quite different from activity in the presence of a cationic or anionic surfactant. Selecting the wrong surfactant can reduce the efficacy of a pesticide product and injure the target plant. Anionic surfactants are most effective when used with contact pesticides (pesticides that control the pest by direct contact rather than being absorbed systemically). Cationic surfactants should never be used as stand-alone surfactants because they usually are phytotoxic.

Nonionic surfactants, often used with systemic pesticides, help pesticide sprays penetrate plant cuticles. Nonionic surfactants are compatible with most pesticides, and most EPA-registered pesticides that require a surfactant recommend a nonionic type. Adjuvants include, but are not limited to, stickers, extenders, plant penetrants, compatibility agents, buffers or pH modifiers, drift control additives, defoaming agents, and thickeners.

Among nonlimiting examples of surfactants included in the compositions described herein are alkyl sulfate ester salts, alkyl sulfonates, alkyl aryl sulfonates, alkyl aryl ethers and polyoxyethylenated products thereof, polyethylene glycol ethers, polyvalent alcohol esters and sugar alcohol derivatives.

xi. Combinations

In formulations and in the use forms prepared from these formulations, the modulating agent may be in a mixture with other active compounds, such as pesticidal agents (e.g., insecticides, sterilants, acaricides, nematocides, molluscicides, or fungicides; see, e.g., pesticides listed in table 12), attractants, growth-regulating substances, or herbicides. As used herein, the term "pesticidal agent" refers to any substance or mixture of substances intended for preventing, destroying, repelling, or mitigating any pest. A pesticide can be a chemical substance or biological agent used against pests including insects, pathogens, weeds, and microbes that compete with humans for food, destroy property, spread disease, or are a nuisance. The term "pesticidal agent" may further encompass other bioactive molecules such as antibiotics, antivirals pesticides, antifungals, antihelminthics, nutrients, pollen, sucrose, and/or agents that stun or slow insect movement.

In instances where the modulating agent is applied to plants, a mixture with other known compounds, such as herbicides, fertilizers, growth regulators, safeners, semiochemicals, or else with agents for improving plant properties is also possible.

V. Delivery

A host described herein can be exposed to any of the compositions described herein in any suitable manner that permits delivering or administering the composition to the insect. The modulating agent may be delivered either alone or in combination with other active or inactive substances and may be applied by, for example, spraying, microinjection, through plants, pouring, dipping, in the form of concentrated liquids, gels, solutions, suspensions, sprays, powders, pellets, briquettes, bricks and the like, formulated to deliver an effective concentration of the modulating agent. Amounts and locations for application of the compositions described herein are generally determined by the habits of the host, the lifecycle stage at which the microorganisms of the host can be targeted by the modulating agent, the site where the application is to be made, and the physical and functional characteristics of the modulating agent. The modulating agents described herein may be administered to the insect by oral ingestion, but may also be administered by means which permit penetration through the cuticle or penetration of the insect respiratory system.

In some instances, the insect can be simply "soaked" or "sprayed" with a solution including the modulating agent. Alternatively, the modulating agent can be linked to a food component (e.g., comestible) of the insect for ease of delivery and/or in order to increase uptake of the modulating agent by the insect. Methods for oral introduction include, for example, directly mixing a modulating agent with the insect's food, spraying the modulating agent in the insect's habitat or field, as well as engineered approaches in which a species that is used as food is engineered to express a modulating agent, then fed to the insect to be affected. In some instances, for example, the modulating agent composition can be incorporated into, or overlaid on the top of, the insect's diet. For example, the modulating agent composition can be sprayed onto a field of crops which an insect inhabits.

In some instances, the composition is sprayed directly onto a plant e.g., crops, by e.g., backpack spraying, aerial spraying, crop spraying/dusting etc. In instances where the modulating agent is delivered

to a plant, the plant receiving the modulating agent may be at any stage of plant growth. For example, formulated modulating agents can be applied as a seed-coating or root treatment in early stages of plant growth or as a total plant treatment at later stages of the crop cycle. In some instances, the modulating agent may be applied as a topical agent to a plant, such that the host insect ingests or otherwise comes
5 in contact with the plant upon interacting with the plant.

Further, the modulating agent may be applied (e.g., in the soil in which a plant grows, or in the water that is used to water the plant) as a systemic agent that is absorbed and distributed through the tissues (e.g., stems or leaflets) of a plant or animal host, such that an insect feeding thereon will obtain an effective dose of the modulating agent. In some instances, plants or food organisms may be genetically
10 transformed to express the modulating agent such that a host feeding upon the plant or food organism will ingest the modulating agent.

Delayed or continuous release can also be accomplished by coating the modulating agent or a composition containing the modulating agent(s) with a dissolvable or bioerodable coating layer, such as gelatin, which coating dissolves or erodes in the environment of use, to then make the modulating agent
15 available, or by dispersing the agent in a dissolvable or erodable matrix. Such continuous release and/or dispensing means devices may be advantageously employed to consistently maintain an effective concentration of one or more of the modulating agents described herein in a specific host habitat.

The modulating agent can also be incorporated into the medium in which the insect grows, lives, reproduces, feeds, or infests. For example, a modulating agent can be incorporated into a food
20 container, feeding station, protective wrapping, or a hive. For some applications the modulating agent may be bound to a solid support for application in powder form or in a "trap" or "feeding station." As an example, for applications where the composition is to be used in a trap or as bait for a particular host insect, the compositions may also be bound to a solid support or encapsulated in a time-release material. For example, the compositions described herein can be administered by delivering the composition to at
25 least one habitat where a vector (e.g., a vector of an animal pathogen, e.g., a mosquito, mite, biting louse, or tick) grows, lives, reproduces, feeds, or infests.

VI. Screening

Included herein are methods for screening for modulating agents that are effective to alter the
30 microbiota of a host (e.g., insect) and thereby decrease host fitness. The screening assays provided herein may be effective to identify one or more modulating agents (e.g., phage) that target symbiotic microorganisms resident in the host and thereby decrease the fitness of the host. For example, the identified modulating agent (e.g., phage) may be effective to decrease the viability of pesticide- or allelochemical-degrading microorganisms (e.g., bacteria, e.g., a bacterium that degrades a pesticide listed
35 in Table 12), thereby increasing the host's sensitivity to a pesticide (e.g., sensitivity to a pesticide listed in Table 12) or allelochemical agent.

For example, a phage library may be screened to identify a phage that targets a specific endosymbiotic microorganism resident in a host. In some instances, the phage library may be provided in the form of one or more environmental samples (e.g., soil, pond sediments, or sewage water).
40 Alternatively, the phage library may be generated from laboratory isolates. The phage library may be co-

- cultured with a target bacterial strain. After incubation with the bacterial strain, phage that successfully infect and lyse the target bacteria are enriched in the culture media. The phage-enriched culture may be sub-cultured with additional bacteria any number of times to further enrich for phage of interest. The phage may be isolated for use as a modulating agent in any of the methods or compositions described
- 5 herein, wherein the phage alters the microbiota of the host in a manner that decreases host fitness.

Table 12. Pesticides

Aclonifen	Fenchlorazole-ethyl	Pendimethalin
Acetamiprid	Fenothiocarb	Penflufen
Alanycarb	Fenitrothion	Penflufen
Amidosulfuron	Fenpropidin	Pentachlorobenzene
Aminocyclopyrachlor	Fluazolate	Penthiopyrad
Amisulbrom	Flufenoxuron	Penthiopyrad
Anthraquinone	Flumetralin	Pirimiphos-methyl
Asulam, sodium salt	Fluxapyroxad	Prallethrin
Benfuracarb	Fuberidazole	Profenofos
Bensulide	Glufosinate-ammonium	Proquinazid
beta-HCH; beta-BCH	Glyphosate	Prothiofos
Bioresmethrin	Group: Borax, borate salts (see	Pyraclofos
Blasticidin-S	Group: Paraffin oils, Mineral	Pyrazachlor
Borax; disodium tetraborate	Halfenprox	Pyrazophos
Boric acid	Imiprothrin	Pyridaben
Bromoxynil heptanoate	Imidacloprid	Pyridalyl
Bromoxynil octanoate	Ipconazole	Pyridiphenthion
Carbosulfan	Isopyrazam	Pyrifenox
Chlorantraniliprole	Isopyrazam	Quinmerac
Chlordimeform	Lenacil	Rotenone
Chlorfluazuron	Magnesium phosphide	Sedaxane
Chlorphropham	Metaflumizone	Sedaxane
Climbazole	Metazachlor	Silafluofen
Clopyralid	Metazachlor	Sintofen
Copper (II) hydroxide	Metobromuron	Spinetoram
Cyflufenamid	Metoxuron	Sulfoxaflor
Cyhalothrin	Metsulfuron-methyl	Temephos
Cyhalothrin, gamma	Milbemectin	thiocloprid
Decahydrate	Naled	Thiamethoxam
Diafenthiuron	Napropamide	Tolfenpyrad
Dimefuron	Nicosulfuron	Tralomethrin
Dimoxystrobin	Nitenpyram	Tributyltin compounds
Dinotefuran	Nitrobenzene	Tridiphane
Diquat dichloride	o-phenylphenol	Triflumizole
Dithianon	oils	Validamycin

E-Phosphamidon	Oxadiargyl	Zinc phosphide
EPTC	Oxycarboxin	
Ethaboxam	Paraffin oil	
Ethirimol	Penconazole	

EXAMPLES

The following is an example of the methods of the invention. It is understood that various other embodiments may be practiced, given the general description provided above.

5

Example 1: Treatment of the *Aedes vexans* mosquito with an antibiotic solution

This Example demonstrates the ability to kill or decrease the fitness of the *Aedes vexans* mosquitoes by treatment with doxycycline, a broad spectrum antibiotic that inhibits protein production. The effect of doxycycline on mosquitoes is mediated through the modulation of bacterial populations endogenous to the mosquito that are sensitive to doxycycline. One targeted bacterial strain is *Wolbachia*.

Successful control and eradication of porcine reproductive and respiratory syndrome virus (PRRSV) is of great importance to the global swine industry today. To reduce the risk of PRRSV entry, swine producers utilize stringent measures to enhance the biosecurity of their farms; however, infection of PRRSV in swine herds still frequently occurs. One vector of transmission of PRRSV is the *Aedes vexans* mosquito. *Aedes vexans* is a cosmopolitan and common pest mosquito. On top of PRRSV, it is also a known vector of *Dirofilaria immitis* (dog heartworm); *Myxomatosis* (deadly rabbit virus disease) and Eastern equine encephalitis (deadly horse virus disease in the USA). *Aedes vexans* is the most common mosquito in Europe, often composing more than 80% the European mosquito community. Its abundance depends upon availability of floodwater pools. In summer, mosquito traps can collect up to 8,000 mosquitoes per trap per night.

Therapeutic design: Blood meals mixed with doxycycline solutions are formulated with final antibiotic concentrations of 0 (negative control), 1, 10, or 50 µg/ml in 1 mL of blood

Experimental design:

To prepare for the treatment, mosquitoes are grown in a lab environment and medium. Experiments are performed with female mosquitoes from an *Aedes vexans*, originally established from field mosquitoes collected on a field of the University of Minnesota St. Paul campus, maintained on human blood and fed as adults with 5% fructose. Doxycycline solutions are made by dissolving doxycycline (SIGMA-ALDRICH, D9891) in sterile water. Different volumes of a doxycycline solution are added to fresh blood to total 1 mL in preparation for blood meals. The final doxycycline concentrations in the blood are approximately 0 (control solution), 1, 10 or 50 µg/ml.

For each replicate, age-matched, 2- to 3-day-old mosquitoes are offered a control or experimental blood meal from a membrane-feeding device (2 ml Eppendorf tube) covered with parafilm and kept at 37 °C. Nonengorged mosquitoes are discarded. Meals are given every four days for a total of three blood meals. Between the blood meals, mosquitoes are provided with a cotton pad moistened with distilled water for oviposition. Unfed mosquitoes are not removed after the second and later blood meals. Deaths are counted daily and carcasses are removed and stored for *Wolbachia* analysis as

described herein. At least 50 mosquitoes per concentration of doxycycline are used for each replicate. At the end of the last blood meal, mosquitoes are kept for 12 hours before dissection.

Microbiota Analysis by Quantitative Polymerase Chain Reaction:

Before dissection, mosquitoes are immersed in 70% ethanol for 5 minutes then rinsed 3 times in sterile phosphate-buffered saline (PBS) to kill and remove surface bacteria, thus minimizing sample contamination with cuticle bacteria during dissection. The midgut of each mosquito (control and doxycycline treatment) is removed and frozen immediately on dry ice and stored at 20°C until processing. Midguts are only excluded from analysis if they burst and a substantial amount of the gut content is lost. Samples are homogenized in phenol-chloroform in a Precellys 24 homogenizer (Bertin) using 0.5 mm wide glass beads (Bertin) for 30 seconds at 6800 rpm and deoxy-ribonucleic acid (DNA) is extracted with phenol-chloroform. The 16S ribosomal DNA (rDNA) is used for *Wolbachia* quantification and is shown as a ratio of the *Aedes* housekeeping gene 40S ribosomal protein S7 (Vector- Base gene ID AAEL009496). Primer sequences for *Wolbachia* are: forward primer 5'- TCAGCCACACTGGAAGTGAAG -3' (SEQ ID NO: 221) and reverse primer 5'- TAACGCTAGCCCTCTCCGTA -3' (SEQ ID NO: 222), and for S7: forward 5'- AAGGTCGACACCTTCACGTC-3' (SEQ ID NO: 223) and reverse 5'-CCGTTTGGTGAGGGTCTTTA-3' (SEQ ID NO: 224). Quantitative polymerase chain reaction (qPCR) is performed on a 7500 Fast Real-Time thermocycler (Applied Biosystems) using the SYBR Premix Ex Taq kit (Takara), following the manufacturer's instructions. Doxycycline treated mosquitoes show a reduction of *Wolbachia* specific genes.

The survival rates of mosquitoes treated with doxycycline solution are compared to the mosquitoes treated with the negative control. The survival rate of mosquitoes treated with doxycycline solution is decreased compared to the control.

Example 2: Treatment of the *Anopheles* mosquito with azithromycin solutions

This Example demonstrates the ability to kill or decrease the fitness of the *Anopheles coluzzii* mosquitoes and decrease the transmission rate of parasites by treatment with azithromycin, relatively broad but shallow antibacterial activity. It inhibits some Gram-positive bacteria, some Gram-negative bacteria, and many atypical bacteria. The effect of azithromycin on mosquitoes is mediated through the modulation of bacterial populations endogenous to the mosquito that are sensitive to azithromycin. One targeted bacterial strain is *Asaia*.

The mosquito has been described as the most dangerous animal in the world and malaria is one mosquito-borne disease that detrimentally impacts humans. There are about 3,500 mosquito species and those that transmit malaria all belong to a sub-set called the *Anopheles*. Approximately 40 *Anopheles* species are able to transmit malaria that significantly impacts human health.

Therapeutic design: Blood meals mixed with azithromycin solutions are formulated with final antibiotic concentrations of 0 (negative control), 0.1, 1, or 5 µg/ml in 1 mL of blood.

Experimental design:

To prepare for the treatment, mosquitoes are grown in a lab environment and medium. Experiments are performed with female mosquitoes from an *Anopheles coluzzii* Ngousso colony,

originally established from field mosquitoes collected in Cameroon, maintained on human blood and fed as adults with 5% fructose. Larvae are fed tetramin fish food. Temperature is maintained at 28 °C (± 1 °C), 70–80% humidity on a 12 hr light/dark cycle.

5 *Human Blood Feeding and Plasmodium Infections:*

Plasmodium falciparum NF54 gametocytes are cultured in RPMI medium (GIBCO) including 300 mg. L-1 L-glutamine supplemented with 50 mg/L hypoxanthine, 25 mM HEPES plus 10% heat-inactivated human serum without antibiotics. Two 25-mL cultures are started 17 and 14 days before the infection at 0.5% parasitemia in 6% v/v washed O+ red blood cells (RBCs). Media is changed daily.

10 Before mosquito infection, centrifuged RBCs are pooled and supplemented with 20% fresh washed RBCs and human serum (2:3 v/v ratio between RBCs and serum). Mosquitoes are offered a blood meal from a membrane-feeding device (2 ml Eppendorf tube) covered with Parafilm and kept at 37 °C.

Azithromycin solutions are made by dissolving azithromycin (SIGMA-ALDRICH, PZ0007) in DMSO. Different volumes of azithromycin solution are added to fresh blood to total 1 mL in preparation
15 for blood meals. The final azithromycin concentrations in the blood are 0 (just solvent as control solution), 0.1, 1, or 5 µg/mL.

For each *Plasmodium* infection, at least 100 age-matched, 2- to 3-day-old, mosquitoes per condition are offered a control or experimental blood meal from a membrane-feeding device (2 ml Eppendorf tube) covered with parafilm and kept at 37 °C and nonengorged mosquitoes are removed.

20 Meals are given every four days for a total of three blood meals. Between the blood meals, mosquitoes are provided with a cotton pad moistened with distilled water for oviposition. Unfed mosquitoes are not removed after the second and later blood meals. Deaths are counted daily and carcasses are removed and stored for *Asaia* analysis as described herein. At least 50 mosquitoes per concentration of azithromycin are used for each replicate. At the end of the last blood meal, mosquitoes are kept for 12
25 hours before dissection.

Microbiota Analysis by Quantitative Polymerase Chain Reaction:

Before dissection, mosquitoes are immersed in 70% ethanol for 5 minutes then rinsed 3 times in sterile phosphate-buffered saline (PBS) to kill and remove surface bacteria, thus minimizing sample
30 contamination with cuticle bacteria during dissection. The midgut of each mosquito (control and azithromycin treatment) is removed and frozen immediately on dry ice and stored at 20 °C until processing. Midguts are only excluded from analysis if they burst and a substantial amount of the gut content is lost. Samples are homogenized in phenol-chloroform in a Precellys 24 homogenizer (Bertin) using 0.5 mm- wide glass beads (Bertin) for 30 seconds at 6800 rpm and deoxy-ribonucleic acid (DNA) is
35 extracted with phenol-chloroform. The 16S ribosomal DNA (rDNA) is used for *Asaia* quantification and is shown as a ratio of the *Anopheles* housekeeping gene 40S ribosomal protein S7 (Vector- Base gene ID AGAP010592). Primer sequences for *Asaia* are: forward 5'-GTGCCGATCTCTAAAAGCCGTCTCA-3' (SEQ ID NO:248) and reverse 5'-TTCGCTCACCGGCTTCGGGT-3' (SEQ ID NO: 249), and for S7: forward 5'- GTGCGCGAGTTGGAGAAGA -3' (SEQ ID NO: 250) and reverse 5'-
40 ATCGGTTTGGGCAGAATGC -3' (SEQ ID NO: 251). Quantitative polymerase chain reaction (qPCR) is

performed on a 7500 Fast Real- Time thermocycler (Applied Biosystems) using the SYBR Premix Ex Taq kit (Takara), following the manufacturer's instructions. Azithromycin treated mosquitoes show a reduction of *Asaia* specific genes.

The survival rates of mosquitoes treated with azithromycin are compared to the mosquitoes
 5 treated with the negative control. The survival rate of mosquitoes treated with azithromycin solution is decreased compared to the control.

Example 3: Treatment of the *Dermacentor andersoni*, with an antibiotic solution

This Example demonstrates the ability to kill or decrease the fitness of the tick, *Dermacentor*
 10 *andersoni*, by treatment with Liqueamycin LA-200 oxytetracycline, a broad spectrum antibiotic commonly used to treat a broad range of bacterial infections in cattle. The effect of Liqueamycin LA-200 oxytetracycline on ticks is mediated through the modulation of bacterial populations endogenous to the tick that are sensitive to Liqueamycin LA-200 oxytetracycline. One targeted bacterial strain is *Rickettsia*.

Ticks are obligate hematophagous arthropods that feed on vertebrates and cause great
 15 economic losses to livestock due to their ability to transmit diseases to humans and animals. In particular, ticks transmit pathogens throughout all continents and are labeled as principle vectors of zoonotic pathogens. In fact, 415 new tick-borne bacterial pathogens have been discovered since Lyme disease was characterized in 1982. *Dermacentor andersoni*, the Rocky Mountain wood tick, has been labeled a 'veritable Pandora's box of disease-producing agents' and transmits several pathogens,
 20 including *Rickettsia rickettsii* and *Francisella tularensis*. It is also a vector of *Anaplasma marginale*, the agent of anaplasmosis, and the most widespread tick-borne pathogen of livestock worldwide (Gall et al., *The ISME Journal* 10:1846-1855, 2016). Economic losses due to anaplasmosis in cattle are estimated to be \$300 million per year in the United States (Rochon et al., *J. Med. Entomol.* 49:253-261, 2012).

Therapeutic design: A therapeutic dose (11 mg/kg of body weight) of Liqueamycin LA-200
 25 oxytetracycline injection on -4, -1, +3 and +5 days post application of ticks.

Experimental design:

Questing adult *D. andersoni* are collected by flag and drag techniques at sites in Burns, Oregon and Lake Como, Montana as described in (Scoles et al., *J. Med. Entomol.* 42:153-162, 2005). Field
 30 collected ticks are used to establish laboratory colonies. For tick bacteria analysis, a cohort of adult F1 or F2 male ticks from each colony is fed on a Holstein calf and dissected to collect midguts (MG) and salivary glands (SG) for genomic DNA isolation and bacteria quantification as follows:

A cohort of F1 ticks are fed on either antibiotic-treated calves or untreated calves (control). The antibiotic-treated calves received a therapeutic dose (11 mg/kg of body weight) of Liqueamycin LA-200
 35 oxytetracycline injections on - 4, - 1, +3 and +5 days post application of ticks, whereas untreated calves did not receive any injections (untreated control). Females ticks are allowed to oviposit to continue a second generation of the untreated and treated ticks (F2 generation). The F2 treated generation arose from F1 adults that are exposed to antibiotics. The F2 ticks are not subjected to antibiotics.

F1 and F2 generation adult male ticks are fed for 7 days and then dissected within 24 h. Deaths
 40 are counted daily and ticks are removed and stored for *Rickettsia* analysis as described herein. Before

dissection, the ticks are surface sterilized and all dissection tools are sterilized between each dissection. Tick MG and SG are dissected and pooled in groups of 30 with three biological replicates. Tissues are stored in Cell Lysis Solution (Qiagen, Valencia, CA, USA) and Proteinase K (1.25 mg/ml). Genomic DNA is isolated using the PureGene Extraction kit (Qiagen) according to the manufacturer's specifications.

5

Quantitative analysis of Rickettsia bellii:

To quantify Rickettsia, rickA gene copy numbers are measured using SYBR Green quantitative PCR of non-treated and antibiotic treated in F1 and F2 ticks. The quantity of Rickettsia is determined using Forward (5'-TACGCCACTCCCTGTGT CA-3'; SEQ ID NO: 225) and Reverse (5'-

10 GATGTAACGGTATTAC ACCAACAG-3'; SEQ ID NO: 226) primers. The bacterial quantity is measured in F1 and F2 MG and SG of the pooled samples. Quantitative polymerase chain reaction (qPCR) is performed on a 7500 Fast Real- Time thermocycler (Applied Biosystems) using the SYBR Premix Ex Taq kit (Takara), following the manufacturer's instructions. Liqueamycin LA-200 oxytetracycline treated ticks show a reduction of Rickettsia specific genes.

15 The survival rates of ticks treated with antibiotic solution are compared to the ticks untreated. The survival rate of ticks treated with Liqueamycin LA-200 oxytetracycline solution is decreased compared to the untreated.

Example 4: Treatment of mites that infect livestock with rifampicin solutions

20 This Example demonstrates the ability to kill or decrease the fitness of mites by treating them with an antibiotic solution. This Example demonstrates that the effect of oxytetracycline on mites is mediated through the modulation of bacterial populations endogenous, such as *Bacillus*, to the mites that are sensitive to oxytetracycline.

Sarcoptic mange is caused by mites that infest animals by burrowing deeply into the skin and

25 laying eggs inside the burrows. The eggs hatch into the larval stage. The larval mites then leave the burrows, move up to the skin surface, and begin forming new burrows in healthy skin tissue. Development from egg to adult is completed in about 2 weeks. The lesions resulting from infestations by these mites are a consequence of the reaction of the animals' immune system to the mites' presence. Because of the intensity of the animals' immunological response, it takes only a small number of mites to

30 produce widespread lesions and generalized dermatitis. While mites can be killed with chemically synthesized miticides, these types of chemicals must be sprayed on every animal in the herd with high-pressure hydraulic spray equipment to ensure penetration by the spray into the skin. Furthermore, these types of chemical pesticides may have detrimental ecological and/or agricultural effects.

Therapeutic design: Oxytetracycline solution is formulated with 0 (negative control), 1, 10, or 50

35 µg/ml in 10 mL of sterile water with 0.5% sucrose and essential amino acids.

Experimental design:

To determine whether adult mites at the reproductive stage have different susceptibility compared to phoretic mites or their offspring because their cuticle is not hardened, mites living on livestock and

40 mites associated with larvae and pupae are collected. This assay tests antibiotic solutions on different

types of mites and determines how their fitness is altered by targeting endogenous microbes, such as *Bacillus*.

The brood mites are collected from mite-infested pigs. Skin samples are collected by gently scraping and lifting off encrusted areas from the inner ear area of the pig with a sharpened teaspoon and subsequently examined for mites.

Mites are grouped per age and assayed separately. The age is determined based on the morphology and pigmentation of the larva or the pupa as follows: mites collected from spinning larvae that are small enough to move around are grouped into Group 1; mites collected from stretched larvae, which are too big to lay in the cell and start to stretch upright with their mouth in the direction of the cell opening, are grouped into Group 2; and mites collected from pupae are grouped into Group 3. Mites are stored on their host larva or pupa in glass Petri dishes until 50 units are collected. This ensures their feeding routine and physiological status remains unchanged. To prevent mites from straying from their host larva or pupa or climbing onto one another, only hosts at the same development stage are kept in the same dish.

The equipment – a stainless steel ring (56 mm inner diameter, 2-3 mm height) and 2 glass circles (62 mm diameter) – is cleaned with acetone and hexane or pentane to form the testing arena. The oxytetracycline solutions and control solution are applied on the equipment by spraying the glass disks and ring of the arena homogeneously. For this, a reservoir is loaded with 1 ml of the solutions; the distance of the sprayed surface from the bottom end of the tube is set at 11 mm and a 0.0275 inch nozzle is used. The pressure is adjusted (usually in the range 350–500 hPa) until the amount of solution deposited is 1 ± 0.05 mg/cm². The antibiotic solutions are poured in their respective dishes, covering the whole bottom of the dishes, and residual liquid is evaporated under a fume hood. The ring is placed between the glass circles to build a cage. The cages are used within 60 hr of preparation, for not more than three assays, in order to control the exposure of mites to antibiotic solutions. 10 to 15 mites are introduced in this cage and the equipment pieces are bound together with droplets of melted wax. Mites collected from spinning larvae, stretched larvae, white eyed pupae and dark eyed with white and pale body are used.

After 4 hours, mites are transferred into a clean glass Petri dish (60 mm diameter) with two or three white eye pupae (4-5 days after capping) to feed on. The mites are observed under a dissecting microscope at 4hr, 24hr, and 48hr after being treated with the antibiotic or the control solutions, and classified according to the following categories:

- Mobile: they walk around when on their legs, whether after being poked by a needle.
- Paralyzed: they move one or more appendages, unstimulated or after stimulation, but they cannot move around.
- Dead: immobile and do not react to 3 subsequent stimulations.

A sterile toothpick or needle is used to stimulate the mites by touching their legs. New tooth picks or sterile needles are used for stimulating each group to avoid contamination between mite groups.

The assays are carried out at 32.5°C and 60-70% relative humidity. If the mortality in the controls exceeds 30%, the replicate is excluded. Each experiment is replicated with four series of cages.

The status of *Bacillus* in mite groups is assessed by PCR. Total DNA is isolated from control (non-oxytetracycline treated) and oxytetracycline treated individuals (whole body) using a DNA Kit (OMEGA, Bio-tek) according to the manufacturer's protocol. The primers for *Bacillus*, forward primer 5'-GAGGTAGACGAAGCGACCTG -3' (SEQ ID NO: 233) and reverse primer 5'-TTCCCTCACGGTACTGGTTC -3' (SEQ ID NO: 234), are designed based on *23S-5S rRNA* sequences obtained from the *Bacillus* genome (Accession Number: AP007209.1) (Takeno et al., *J. Bacteriol.* 194(17):4767-4768, 2012) using Primer 5.0 software (Primer-E Ltd., Plymouth, UK). The PCR amplification cycles included an initial denaturation step at 95°C for 5min, 35 cycles of 95°C for 1min, 59°C for 1min, and 72°C for 2min, and a final extension step of 5min at 72°C. Amplification products from oxytetracycline treated and control samples are analyzed on 1% agarose gels, stained with SYBR safe, and visualized using an imaging System.

The survival rates of mites treated with an oxytetracycline solution are compared to the Varroa mites treated with the negative control.

The survival rate and the mobility of mites treated with oxytetracycline solution are expected to be decreased compared to the control.

Example 5: Production of a phage library

This Example demonstrates the acquisition of a phage collection from environmental samples.

Therapeutic design: Phage library collection having the following phage families: *Myoviridae*, *Siphoviridae*, *Podoviridae*, *Lipothruxviridae*, *Rudoviridae*, *Ampullaviridae*, *Bicaudaviridae*, *Clavaviridae*, *Corticoviridae*, *Cystoviridae*, *Fuselloviridae*, *Gluboloviridae*, *Guttaviridae*, *Inoviridae*, *Leviviridae*, *Microviridae*, *Plasmaviridae*, *Tectiviridae*

Experimental design:

Multiple environmental samples (soil, pond sediments, sewage water) are collected in sterile 1L flasks over a period of 2 weeks and are immediately processed as described below after collection and stored thereafter at 4°C. Solid samples are homogenized in sterile double-strength difco luria broth (LB) or tryptic soy broth (TSB) supplemented with 2mM CaCl₂ to a final volume of 100mL. The pH and phosphate levels are measured using phosphate test strips. For purification, all samples are centrifuged at 3000–6000 g in a Megafuge 1.0R, Heraeus, or in Eppendorf centrifuge 5702 R, for 10–15 min at +4°C, and filtered through 0.2-µm low protein filters to remove all remaining bacterial cells. The supernatant is stored at 4°C in the presence of chloroform in a glass bottle.

Example 6: Identification of target specific phage

This Example demonstrates the isolation, purification, and identification of single target specific phages from a heterogeneous phage library.

Experimental design:

20–30 ml of the phage library described in Example 5 is diluted to a volume of 30–40 ml with LB-broth. The target bacterial strain, e.g., *Buchnera*, is added (50–200 µl overnight culture grown in LB-

broth) to enrich phages that target this specific bacterial strain in the culture. This culture is incubated overnight at +37°C, shaken at 230 rpm. Bacteria from this enrichment culture are removed by centrifugation (3000–6000 g in Megafuge 1.0R, Heraeus, or in Eppendorf centrifuge 5702 R, 15–20 min, +4°C) and filtered (0.2 or 0.45 µm filter). 2.5 ml of the bacteria free culture is added to 2.5 ml of LB-broth and 50–100 µl of the target bacteria to enrich the phages. The enrichment culture is grown overnight as above. A sample from this enrichment culture is centrifuged at 13,000 g for 15 min at room temperature and 10 µl of the supernatant is plated on a LB-agar containing petri dish along with 100–300 µl of the target bacteria and 3 ml of melted 0.7% soft-agar. The plates are incubated overnight at +37°C. Each of the plaques observed on the bacterial lawn are picked and transferred into 500 µl of LB-broth. A sample from this plaque-stock is further plated on the target bacteria. Plaque-purification is performed three times for all discovered phages in order to isolate a single homogenous phage from the heterogeneous phage mix.

Lysates from plates with high-titer phages ($>1 \times 10^{10}$ PFU/ml) are prepared by harvesting overlay plates of a host bacterium strain exhibiting confluent lysis. After being flooded with 5 ml of buffer, the soft agar overlay is macerated, clarified by centrifugation, and filter sterilized. The resulting lysates are stored at 4°C. High-titer phage lysates are further purified by isopycnic CsCl centrifugation, as described in (Summer et al., *J. Bacteriol.* 192:179-190, 2010).

DNA is isolated from CsCl-purified phage suspensions as described in (Summer, *Methods Mol. Biol.* 502:27-46, 2009). An individual isolated phage is sequenced as part of two pools of phage genomes by using a 454 pyrosequencing method. Phage genomic DNA is mixed in equimolar amounts to a final concentration of about 100 ng/L. The pooled DNA is sheared, ligated with a multiplex identifier (MID) tag specific for each of the pools, and sequenced by pyrosequencing using a full-plate reaction on a Roche FLX Titanium sequencer according to the manufacturer's protocols. The pooled phage DNA is present in two sequencing reactions. The trimmed FLX Titanium flow-gram output corresponding to each of the pools is assembled individually by using Newbler Assembler version 2.5.3 (454 Life Sciences), by adjusting the settings to include only reads containing a single MID per assembly. The identity of individual contigs is determined by PCR using primers generated against contig sequences and individual phage genomic DNA preparations as the template. Sequencher 4.8 (Gene Codes Corporation) is used for sequence assembly and editing. Phage chromosomal end structures are determined experimentally. Cohesive (cos) ends for phages are determined by sequencing off the ends of the phage genome and sequencing the PCR products derived by amplification through the ligated junction of circularized genomic DNA, as described in (Summer, *Methods Mol. Biol.* 502:27-46, 2009). Protein-coding regions are initially predicted using GeneMark.hmm (Lukashin et al. *Nucleic Acids Res.* 26:1107-1115, 1998), refined through manual analysis in Artemis (Rutherford et al., *Bioinformatics* 16:944-945, 2000.), and analyzed through the use of BLAST (E value cutoff of 0.005) (Camacho et al., *BMC Bioinformatics* 10:421, 2009). Proteins of particular interest are additionally analyzed by InterProScan (Hunter et al., *Nucleic Acids Res.* 40:D306-D312, 2012).

Electron microscopy of CsCl-purified phage ($>1 \times 10^{11}$ PFU/ml) that lysed the endosymbiotic bacteria, *Buchnera*, is performed by diluting stock with the tryptic soy broth buffer. Phages are applied onto thin 400-mesh carbon-coated Formvar grids, stained with 2% (wt/vol) uranyl acetate, and air dried.

Specimens are observed on a JEOL 1200EX transmission electron microscope operating at an acceleration voltage of 100 kV. Five virions of each phage are measured to calculate mean values and standard deviations for dimensions of capsid and tail, where appropriate.

5 Example 7: Treatment of aphids with a solution of purified phages

This Example demonstrates the ability to kill or decrease the fitness of aphids by treating them with a phage solution. This Example demonstrates that the effect of phage on aphids is mediated through the modulation of bacterial populations endogenous to the aphid that are sensitive to phages. One targeted bacterial strain is *Buchnera* with the phage identified in Example 6.

10 Aphids are representative species for testing microbiota modulating agents and effects on fitness of the aphids.

Therapeutic design:

15 Phage solutions are formulated with 0 (negative control), 10^2 , 10^5 , or 10^8 plaque-forming units (pfu)/ml phage from Example 6 in 10 mL of sterile water with 0.5% sucrose and essential amino acids.

Experimental design:

To prepare for the treatment, aphids are grown in a lab environment and medium. In a climate-controlled room (16 h light photoperiod; $60 \pm 5\%$ RH; 20 ± 2 °C), fava bean plants are grown in a mixture of vermiculite and perlite at 24 °C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants are distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, second and third instar aphids are collected from healthy plants and divided into treatments so that each treatment receives approximately the same number of individuals from each of the collection plants.

25 Phage solutions are prepared as described herein. Wells of a 96-well plate are filled with 200 μ l of artificial aphid diet (Febvay et al., *Canadian Journal of Zoology* 66(11):2449-2453, 1988) and the plate is covered with parafilm to make a feeding sachet. Artificial diet is either mixed with sterile water and with 0.5% sucrose and essential amino acids as a negative control or phage solutions with varying concentrations of phages. Phage solutions are mixed with artificial diet to get final concentrations of phages between 10^2 to 10^8 (pfu)/ml.

30 For each replicate treatment, 30–50 second and third instar aphids are placed individually in wells of a 96-well plate and a feeding sachet plate is inverted above them, allowing the insects to feed through the parafilm and keeping them restricted to individual wells. Experimental aphids are kept under the same environmental conditions as aphid colonies. After the aphids are fed for 24 hr, the feeding sachet is replaced with a new one containing sterile artificial diet and a new sterile sachet is provided every 24 h for 4 days. At the time that the sachet is replaced, aphids are also checked for mortality. An aphid is counted as dead if it had turned brown or is at the bottom of the well and does not move during the observation. If an aphid is on the parafilm of the feeding sachet but not moving, it is assumed to be feeding and alive.

The status of *Buchnera* in aphid samples is assessed by PCR. Aphids adults from the negative control (non-phage treated) and phage treated groups are first surface-sterilized with 70% ethanol for 1 min, 10% bleach for 1 min and three washes of ultrapure water for 1 min. Total DNA is extracted from each individual (whole body) using an Insect DNA Kit (OMEGA, Bio-tek) according to the manufacturer's protocol. The primers for *Buchnera*, forward primer 5'-GTCGGCTCATCACATCC-3' (SEQ ID NO: 235) and reverse primer 5'-TTCCGTCTGTATTATCTCCT-3' (SEQ ID NO: 236), are designed based on 23S-5S rRNA sequences obtained from the *Buchnera* genome (Accession Number: GCA_000009605.1) (Shigenobu et al., *Nature* 407:81-86, 2000) using Primer 5.0 software (Primer-E Ltd., Plymouth, UK). The PCR amplification cycles included an initial denaturation step at 95 °C for 5min, 35 cycles of 95 °C for 30s, 55 °C for 30s, and 72 °C for 60s, and a final extension step of 10min at 72 °C. Amplification products from rifampicin treated and control samples are analyzed on 1% agarose gels, stained with SYBR safe, and visualized using an imaging System. Phage treated aphids show a reduction of *Buchnera* specific genes.

The survival rates of aphids treated with *Buchnera* specific phages are compared to the aphids treated with the negative control. The survival rate of aphids treated with *Buchnera* specific phages is decreased as compared to the control treated aphids.

Example 8: Production of a *colA* bacteriocin solution

This Example demonstrates the production and purification of *colA* bacteriocin.

Construct sequence:

catatgatgacccgcaccatgctgtttctggcgtgctggcgccgctgtatgtgtgcattagcgcgaccgcgggcaaacgggaagaatttgcgaaac
tgagcgtatgaagcgcgcagcaacgatcaggcgtatgaaagcattcagcgctatcgccgctttgtgatggcaaccgctataacggcgccag
cagcagcagcagcagccgaacagtgggaagtgcgcccggatctgagccgcgatcagcgcggcaacaccaaagcgcaggtggaaattaac
aaaaaaggcgataaccatgatattaacggggctggggcaaaaacattaacggcccgatagccataaagatactggcatgtggcgccgagc
gtgcgctgctcgag (SEQ ID NO: 237)

Experimental design:

DNA is generated by PCR with specific primers with upstream (NdeI) and downstream (XhoI) restriction sites. Forward primer GTATCTATTCCCGTCTACGAACATATGGAATTCC (SEQ ID NO: 238) and reverse primer CCGCTCGAGCCATCTGACACTTCCTCCAA (SEQ ID NO: 239). Purified PCR fragments (Nucleospin Extract II-Macherey Nagel) are digested with NdeI or XhoI and then the fragments are ligated. For *colA* cloning, the ligated DNA fragment is cloned into pcr2.1 (GenBank database accession number EY122872) vector (Anselme et al., *BMC Biol.* 6:43, 2008). The nucleotide sequence is systematically checked (Cogenics).

The plasmid with *colA* sequence is expressed in BL21 (DE3)/pLys. Bacteria are grown in LB broth at 30 °C. At an OD600 of 0.9, isopropyl β-D-1-thiogalactopyranoside (IPTG) is added to a final concentration of 1mM and cells were grown for 6h. Bacteria are lysed by sonication in 100mM NaCl, 1% Triton X-100, 100mM Tris-base pH 9.5, and proteins are loaded onto a HisTrap HP column (GE Healthcare). The column is washed successively with 100mM NaCl, 100mM Tris-HCl pH 6.8, and PBS. Elution is performed with 0.3M imidazol in PBS. Desalting PD-10 columns (GE Healthcare) are used to eliminate imidazol and PBS solubilized peptides are concentrated on centrifugal filter units (Millipore).

ColA Protein sequence:

MTRTMLFLAC VAALYVCISA TAGKPEEFAK LSDEAPSNDQ AMYESIQRYR RFVDGNRYNG
 GQQQQQPKQ WEVRPDLSDR QRGNTKAQVE INKKGDNHDI NAGWGKNING PDSHKDTWHV
 GGSVRW (SEQ ID NO: 211)

5 Example 9: Treatment of aphids with a solution of *colA* bacteriocin

This Example demonstrates the ability to kill or decrease the fitness of aphids by treating them with a bacteriocin solution. This Example demonstrates that the effect of bacteriocins on aphids is mediated through the modulation of bacterial populations endogenous to the aphid that are sensitive to ColA bacteriocin. One targeted bacterial strain is *Buchnera* with the bacteriocin produced in Example 8.

10

Therapeutic design:

ColA solutions are formulated with 0 (negative control), 0.6, 1, 50 or 100 mg/ml of ColA from Example 8 in 10 mL of sterile water with 0.5% sucrose and essential amino acids.

15

Experimental design:

To prepare for the treatment, aphids are grown in a lab environment and medium. In a climate-controlled room (16 h light photoperiod; 60±5% RH; 20±2 °C), plants are grown in a mixture of vermiculite and perlite and are infested with aphids. In the same climatic conditions, *E. balteatus* larvae are obtained from a mass production; the hoverflies are reared with sugar, pollen and water; and the oviposition is induced by the introduction of infested host plants in the rearing cage during 3 h. The complete life cycle takes place on the host plants that are daily re-infested with aphids.

20

Wells of a 96-well plate are filled with 200 µl of artificial aphid diet (Febvay et al., *Canadian Journal of Zoology* 66(11):2449-2453, 1988) and the plate is covered with parafilm to make a feeding sachet. Artificial diet is either mixed with the solution of sterile water with 0.5% sucrose and essential amino acids as a negative control or ColA solutions with varying concentrations of ColA. ColA solutions are mixed with artificial diet to obtain final concentrations between 0.6 to 100 mg/ml.

25

For each replicate treatment, 30–50 second and third instar aphids are placed individually in wells of a 96-well plate and a feeding sachet plate is inverted above them, allowing the insects to feed through the parafilm and keeping them restricted to individual wells. Experimental aphids are kept under the same environmental conditions as aphid colonies. After the aphids are fed for 24 hr, the feeding sachet is replaced with a new one containing sterile artificial diet and a new sterile sachet is provided every 24 h for 4 days. At the time that the sachet is replaced, aphids are also checked for mortality. An aphid is counted as dead if it had turned brown or is at the bottom of the well and does not move during the observation. If an aphid is on the parafilm of the feeding sachet but not moving, it is assumed to be feeding and alive.

35

The status of *Buchnera* in aphid samples is assessed by PCR. Aphids adults from the negative control and phage treated are first surface-sterilized with 70% ethanol for 1 min, 10% bleach for 1 min and three washes of ultrapure water for 1 min. Total DNA is extracted from each individual (whole body) using an Insect DNA Kit (OMEGA, Bio-tek) according to the manufacturer's protocol. The primers for *Buchnera*, forward primer 5'-GTCGGCTCATCACATCC-3' (SEQ ID NO: 235) and reverse primer 5'-

40

TTCCGTCTGTATTATCTCCT-3' (SEQ ID NO: 236), are designed based on *23S-5S rRNA* sequences obtained from the *Buchnera* genome (Accession Number: GCA_000009605.1) (Shigenobu, et al., *Nature* 200.407, 81–86) using Primer 5.0 software (Primer-E Ltd., Plymouth, UK). The PCR amplification cycles included an initial denaturation step at 95°C for 5min, 35 cycles of 95°C for 30s, 55°C for 30s, and 72°C
 5 for 60s, and a final extension step of 10min at 72°C. Amplification products from rifampicin treated and control samples are analyzed on 1% agarose gels, stained with SYBR safe, and visualized using an imaging System. ColA treated aphids show a reduction of *Buchnera* specific genes.

The survival rates of aphids treated with *Buchnera* specific ColA bacteriocin are compared to the aphids treated with the negative control. The survival rate of aphids treated with *Buchnera* specific ColA
 10 bacteriocin is decreased as compared to the control treated aphids.

Example 10: Treatment of aphids with rifampicin solutions

This Example demonstrates the ability to kill or decrease the fitness of aphids by treating them with rifampicin, a narrow spectrum antibiotic that inhibits DNA-dependent RNA synthesis by inhibiting a
 15 bacterial RNA polymerase. This Example demonstrates that the effect of rifampicin on aphids is mediated through the modulation of bacterial populations endogenous to the aphid that are sensitive to rifampicin. One targeted bacterial strain is *Buchnera*.

Therapeutic design:

20 The antibiotic solutions are formulated with 0 (negative control), 1, 10, or 50 µg/ml of rifampicin in 10 mL of sterile water with 0.5% sucrose and essential amino acids.

Experimental design:

To prepare for the treatment, aphids are grown in a lab environment and medium. In a climate-
 25 controlled room (16 h light photoperiod; 60±5% RH; 20±2 °C), fava bean plants are grown in a mixture of vermiculite and perlite at 24 °C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants are distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, second and third instar aphids are collected from healthy plants and divided into treatments so that each treatment receives
 30 approximately the same number of individuals from each of the collection plants.

Rifampicin solutions are made by dissolving rifampicin (SIGMA-ALDRICH, 557303) in sterile water with 0.5% sucrose and essential aminoacids. Wells of a 96-well plate are filled with 200 µl of artificial aphid diet (Febvay et al., *Canadian Journal of Zoology* 66(11):2449-2453, 1988) and the plate is covered with parafilm to make a feeding sachet. Artificial diet is either mixed with sterile water and with
 35 0.5% sucrose and essential aminoacids as a negative control or a rifampicin solution with one of the concentrations of rifampicin. Rifampicin solutions are mixed with artificial diet to get final concentrations of the antibiotic between 1 and 50 µg/mL.

For each replicate treatment, 30–50 second and third instar aphids are placed individually in wells of a 96-well plate and a feeding sachet plate is inverted above them, allowing the insects to feed through
 40 the parafilm and keeping them restricted to individual wells. Experimental aphids are kept under the

same environmental conditions as aphid colonies. After the aphids are fed for 24 hr, the feeding sachet is replaced with a new one containing sterile artificial diet and a new sterile sachet is provided every 24 h for four days. At the time that the sachet is replaced, aphids are also checked for mortality. An aphid is counted as dead if it had turned brown or is at the bottom of the well and does not move during the observation. If an aphid is on the parafilm of the feeding sachet but not moving, it is assumed to be feeding and alive.

The status of *Buchnera* in aphid samples is assessed by PCR. Total DNA is isolated from control (non-rifampicin treated) and rifampicin treated individuals using an Insect DNA Kit (OMEGA, Bio-tek) according to the manufacturer's protocol. The primers for *Buchnera*, forward primer 5'-GTCCGGCTCATCACATCC-3' (SEQ ID NO: 235) and reverse primer 5'-TTCCGTCTGTATTATCTCCT-3' (SEQ ID NO: 236), are designed based on 23S-5S rRNA sequences obtained from the *Buchnera* genome (Accession Number: GCA_000009605.1) (Shigenobu et al., *Nature* 407:81-86, 2000) using Primer 5.0 software (Primer-E Ltd., Plymouth, UK). The PCR amplification cycles included an initial denaturation step at 95°C for 5min, 35 cycles of 95°C for 30s, 55°C for 30s, and 72°C for 60s, and a final extension step of 10min at 72°C. Amplification products from rifampicin treated and control samples are analyzed on 1% agarose gels, stained with SYBR safe, and visualized using an imaging System. Rifampicin treated aphids show a reduction of *Buchnera* specific genes.

The survival rates of aphids treated with rifampicin solution are compared to the aphids treated with the negative control. The survival rate of aphids treated with rifampicin solution is decreased compared to the control.

Example 11: High throughput screening (HTS) for *Buchnera* targeting molecules

This Example demonstrates the identification of molecules that target *Buchnera*.

Experimental design: A HTS to identify inhibitors of targeted bacterial strains, *Buchnera*, uses sucrose fermentation in pH-MMSuc medium (Ymele-Leki et al., *PLoS ONE* 7(2):e31307, 2012) to decrease the pH of the medium. pH indicators in the medium monitor medium acidification spectrophotometrically through a change in absorbance at 615 nm (A615). A target bacterial strain, *Buchnera*, derived from a glycerol stock, is plated on an LB-agar plate and incubated overnight at 37 °C. A loopful of cells is harvested, washed three times with PBS, and then resuspended in PBS at an optical density of 0.015.

For the HTS, 10 µL of this bacterial cell suspension is aliquoted into the wells of a 384-well plate containing 30 µL of pH-MMSuc medium and 100 nL of a test compound fraction from a natural product library of pre-fractionated extracts (39,314 extracts arrayed in 384-well plates) from microbial sources, such as fungal endophytes, bacterial endophytes, soil bacteria, and marine bacteria, described in (Ymele-Leki et al., *PLoS ONE* 7(2):e31307, 2012). For each assay, the A615 is measured after incubation at room temperature at 6 hr and 20 hr. This step is automated and validated in the 384-well plate format using an EnVision™ multi-well spectrophotometer to test all fractions from the library. Fractions that demonstrate delayed medium acidification by sucrose fermentation and inhibited cell growth are selected for further purification and identification.

Example 12: Isolation and identification of *Buchnera* specific molecules

This Example demonstrates the isolation and identification of an isolate from the fraction described in Example 11 that blocks sucrose fermentation and inhibits cell growth of *Buchnera*.

5 *Experimental design:*

The fraction described in Example 11 is resuspended in 90% water/methanol and passed over a C18 SPE column to get fraction I. The column is then washed with methanol to get fraction II. Fraction II is separated on an Agilent 1100 series HPLC with a preparative Phenyl-hexyl column (Phenomenex, Luna, 25 cm610 mm, 5 mm particle size) using an elution buffer with 20% acetonitrile/water with 0.1% formic acid at a flow rate of 2 mL/min for 50 minutes. This yields multiple compounds at different elution times. Spectra for each compound is obtained on an Alpha FT-IR mass spectrometer (Bruker), an UltrospecTM 5300 pro UV/ Visible Spectrophotometer (Amersham Biosciences), and an INOVA 600 MHz nuclear magnetic resonance spectrometer (Varian).

15 **Example 13: Treatment of aphids with a solution of a *Buchnera* specific molecule**

This Example demonstrates the ability to kill or decrease the fitness of aphids by treating them with one of the compounds identified in Example 12 through the modulation of bacterial populations endogenous to the aphid that are sensitive to this compound. One targeted bacterial strain is *Buchnera*.

20 *Therapeutic design:*

Each compound from Example 12 is formulated at 0 (negative control), 0.6, 1, 20 or 80 µg/ml in 10 mL of sterile water with 0.5% sucrose and essential amino acids.

Experimental design:

25 To prepare for the treatment, aphids are grown in a lab environment and medium. In a climate-controlled room (16 h light photoperiod; 60±5% RH; 20±2 °C), fava bean plants are grown in a mixture of vermiculite and perlite at 24 °C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants are distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, second and third instar aphids are collected from healthy plants and divided into treatments so that each treatment receives approximately the same number of individuals from each of the collection plants.

30 Wells of a 96-well plate are filled with 200 µl of artificial aphid diet (Febvay et al., *Canadian Journal of Zoology* 66(11):2449-2453, 1988) and the plate is covered with parafilm to make a feeding sachet. Artificial diet is either mixed with sterile water with 0.5% sucrose and essential amino acids as a negative control or solutions with varying concentrations of the compound.

35 For each replicate treatment, 30–50 second and third instar aphids are placed individually in wells of a 96-well plate and a feeding sachet plate is inverted above them, allowing the insects to feed through the parafilm and keeping them restricted to individual wells. Experimental aphids are kept under the same environmental conditions as aphid colonies. After the aphids are fed for 24 hr, the feeding sachet is replaced with a new one containing sterile artificial diet and a new sterile sachet is provided every 24 h for

4 days. At the time that the sachet is replaced, aphids are also checked for mortality. An aphid is counted as dead if it had turned brown or is at the bottom of the well and does not move during the observation. If an aphid is on the parafilm of the feeding sachet but not moving, it is assumed to be feeding and alive.

5 The status of *Buchnera* in aphid samples is assessed by PCR. Aphids from the negative control and compound 1 treated are first surface-sterilized with 70% ethanol for 1 min, 10% bleach for 1 min and three washes of ultrapure water for 1 min. Total DNA is extracted from each individual (whole body) using an Insect DNA Kit (OMEGA, Bio-tek) according to the manufacturer's protocol. The primers for *Buchnera*, forward primer 5'-GTCGGCTCATCACATCC-3' (SEQ ID NO: 235) and reverse primer 5'-
10 TTCCGTCTGTATTATCTCCT-3' (SEQ ID NO: 236), are designed based on *23S-5S rRNA* sequences obtained from the *Buchnera* genome (Accession Number: GCA_000009605.1) (Shigenobu et al., *Nature* 407:81-86, 2000) using Primer 5.0 software (Primer-E Ltd., Plymouth, UK). The PCR amplification cycles included an initial denaturation step at 95°C for 5min, 35 cycles of 95°C for 30s, 55°C for 30s, and 72°C for 60s, and a final extension step of 10min at 72°C. Amplification products from compound 1 treated and
15 control samples are analyzed on 1% agarose gels, stained with SYBR safe, and visualized using an imaging System. Reduction of *Buchnera* specific genes indicates antimicrobial activity of compound 1.

The survival rate of aphids treated with the compound is compared to the aphids treated with the negative control. A decrease in the survival rate of aphids treated with the compound is expected to indicate antimicrobial activity of the compound.

20

Example 14: Insects treated with an antibiotic solution

This Example demonstrates the treatment of aphids with rifampicin, a narrow spectrum antibiotic that inhibits DNA-dependent RNA synthesis by inhibiting a bacterial RNA polymerase. This Example demonstrates that the effect of rifampicin on a model insect species, aphids, was mediated through the
25 modulation of bacterial populations endogenous to the insect that were sensitive to rifampicin. One targeted bacterial strain is *Buchnera*.

Therapeutic Design

30 The antibiotic solution was formulated according to the means of delivery, as follows (Fig. 1A-1G):

1) Through the plants: with 0 (negative control) or 100 µg/ml of rifampicin formulated in an artificial diet (based on Akey and Beck, 1971; see *Experimental Design*) with and without essential amino acids (2 mg/ml histidine, 2 mg/ml isoleucine, 2 mg/ml leucine, 2 mg/ml lysine, 1 mg/ml methionine, 1.62 mg/ml phenylalanine, 2 mg/ml threonine, 1 mg/ml tryptophan, and 2 mg/ml valine).

35 2) Leaf coating: 100 µl of 0.025% nonionic organosilicone surfactant solvent Silwet L-77 in water (negative control), or 100 µl of 50 µg/ml of rifampicin formulated in solvent solution was applied directly to the leaf surface and allowed to dry.

3) Microinjection: injection solutions were either 0.025% nonionic organosilicone surfactant solvent Silwet L-77 in water (negative control), or 50 µg/ml of rifampicin formulated in solvent solution.

4) Topical delivery: 100 μ l of 0.025% nonionic organosilicone surfactant solvent Silwet L-77 (negative control), or 50 μ g/ml of rifampicin formulated in solvent solution were sprayed using a 30 mL spray bottle.

5) Leaf injection method A – Leaf perfusion and cutting: leaves were injected with approximately 1 ml of 50 μ g/ml of rifampicin in water with food coloring or approximately 1 ml of negative control with water and food coloring. Leaves were cut into 2x2 cm squared pieces and aphids were placed on the leaf pieces.

6) Leaf perfusion and delivery through plant: Leaves were injected with approximately 1 ml of 100 μ g/ml of rifampicin in water plus food coloring or approximately 1 ml of negative control with water and food coloring. The stem of injected leaf was then placed into an Eppendorf tube with 1 ml of 100 μ g/ml of rifampicin plus water and food coloring or 1 ml of negative control with only water and food coloring.

7) Combination delivery method: a) Topical delivery to aphid and plant: via spraying both aphids and plants with 0.025% nonionic organosilicone surfactant solvent Silwet L-77 in water (negative control) or 100 μ g/ml of rifampicin formulated in solvent solution using a 30 mL, b) Delivery through plant: water only (negative control) or 100 μ g/ml of rifampicin formulated in water.

Plant Delivery Experimental Design:

Aphids (LSR-1 strain, *Acyrtosiphon pisum*) were grown on fava bean plants (*Vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60 $\pm$ 5% RH; 25 $\pm$ 2 $^{\circ}$ C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24 $^{\circ}$ C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first instar aphids were collected from healthy plants and divided into 3 different treatment groups: 1) artificial diet alone without essential amino acids, 2) artificial diet alone without essential amino acids and 100 μ g/ml rifampicin, and 3) artificial diet with essential amino acids and 100 μ g/ml rifampicin. Each treatment group received approximately the same number of individuals from each of the collection plants.

The artificial diet used was made as previously published (Akey and Beck, 1971 Continuous Rearing of the Pea Aphid, *Acyrtosiphon pisum*, on a Holidic Diet), with and without the essential amino acids (2 mg/ml histidine, 2 mg/ml isoleucine, 2 mg/ml leucine, 2 mg/ml lysine, 1 mg/ml methionine, 1.62 mg/ml phenylalanine, 2 mg/ml threonine, 1 mg/ml tryptophan, and 2 mg/ml valine), except neither diet included homoserine or beta-alanyltyrosine. The pH of the diets was adjusted to 7.5 with KOH and diets were filter sterilized through a 0.22 μ m filter and stored at 4 $^{\circ}$ C for short term (<7 days) or at -80 $^{\circ}$ C for long term.

Rifampicin (Tokyo Chemical Industry, LTD) stock solutions were made at 25 mg/ml in methanol, sterilized by passing through a 0.22 μ m syringe filter, and stored at -20 $^{\circ}$ C. For treatments (see *Therapeutic design*), the appropriate amount of stock solution was added to the artificial diet with or without essential amino acids to obtain a final concentration of 100 μ g/ml rifampicin. The diet was then placed into a 1.5 ml Eppendorf tube with a fava bean stem with a leaf and the opening of the tube was

closed using parafilm. This artificial diet feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

For each treatment, 33 aphids were placed onto each leaf. Artificial diet feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead
5 aphids were removed from the deep petri dish housing the artificial feeding system when they were discovered.

In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th instar) was determined daily throughout the experiment. Once an aphid reached the 4th instar stage, they were given their own artificial feeding system in a deep petri dish so that fecundity could be monitored once they reached adulthood.

10 For adult aphids, new nymphs were counted daily and then discarded. At the end of the experiments, fecundity was determined as the mean number of offspring produced daily once the aphid reached adulthood. Pictures of aphids were taken throughout the experiment to evaluate size differences between treatment groups.

After 7 days of treatment, DNA was extracted from multiple aphids from each treatment group.
15 Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F
20 (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 240) and Buch_groES_98R (CTGTTCTCGAGTCGATTTCC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for
25 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

Antibiotic treatment delays and stops progression of aphid development

30 LSR-1 1st instar aphids were divided into three separate treatment groups as defined in *Experimental Design* (above). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with artificial diet alone without essential amino acids began reaching maturity (5th instar stage) at approximately 6 days (Fig. 2A). Development was delayed in aphids treated with rifampicin, and by 6 days of treatment, most aphids did not mature further
35 than the 3rd instar stage, even after 12 days and their size is drastically affected (Figs. 2A-2C).

In contrast, aphids treated with artificial diet with rifampicin supplemented with essential amino acids developed faster and to higher instar stages as compared to aphids treated with rifampicin alone, but not as quickly as aphids treated with artificial diet without essential amino acids (Figs. 2A-2C). These data indicate that treatment with rifampicin impaired aphid development. Adding back essential amino
40 acids partially rescued this defect in development.

Antibiotic treatment increased aphid mortality

Survival rate of aphids was also measured during the treatments. The majority of the aphids treated with artificial diet alone without essential amino acids were alive at 5 days post-treatment (Fig. 3).

5 After 5 days, aphids began gradually dying, and some survived beyond 13 days post-treatment.

In contrast, aphids treated with rifampicin without essential amino acids had lower survival rates than aphids treated with artificial diet alone ($p < 0.00001$). Rifampicin-treated aphids began dying after 1 day of treatment and all aphids succumbed to treatment by 9 days. Aphids treated with both rifampicin and essential amino acids survived longer than those treated with rifampicin alone ($p = 0.017$). These data
10 indicate that rifampicin treatment affected aphid survival.

Antibiotic treatment decreased aphid reproduction

Fecundity was also monitored in aphids during the treatments. By days 7 and 8 post-treatment, the majority of the adult aphids treated with artificial diet without essential amino acids began
15 reproducing. The mean number of offspring produced daily after maturity by aphids treated with artificial diet without essential amino acids was approximately 4 (Fig. 4). In contrast, aphids treated with rifampicin with or without essential amino acids were unable to reach adulthood and produce offspring. These data indicate that rifampicin treatment resulted in a loss of aphid reproduction.

Antibiotic treatment decreased Buchnera in aphids

To test whether rifampicin, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 7 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids treated with artificial diet alone without essential amino acids had high ratios of *Buchnera*/aphid DNA copies. In contrast,
25 aphids treated with rifampicin had ~4-fold less *Buchnera*/aphid DNA copies (Fig. 5), indicating that rifampicin treatment decreased *Buchnera* levels.

Leaf Coating Delivery Experimental Design

Rifampicin stock solution was added to 0.025% of a nonionic organosilicone surfactant solvent, Silwet L-77, to obtain a final concentration of 50 µg/ml rifampicin. Aphids (eNASCO strain, *Acyrtosiphon pisum*) were grown on fava bean plants as described in a previous Example. For experiments, first instar aphids were collected from healthy plants and divided into 2 different treatment groups: leaves were sprayed with 1) negative control (solvent solution only), 2) 50 µg/ml rifampicin in solvent. Solutions were absorbed onto a 2x2 cm piece of fava bean leaf.

35 Each treatment group received approximately the same number of individuals from each of the collection plant. For each treatment, 20 aphids were placed onto each leaf. Aphids were monitored daily for survival and dead aphids were removed when they were discovered. In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th instar, and 5R, representing a reproducing 5th instar) was determined daily throughout the experiment. Pictures of aphids were taken throughout the experiment to evaluate size
40 differences between treatment groups.

After 6 days of treatment, DNA was extracted from multiple aphids from each treatment group and qPCR for quantifying *Buchnera* levels was done as described in the previous Example.

Antibiotic treatment delivered through leaf coating delays and stops progression of aphid development

5 LSR-1 1st instar aphids were divided into two separate treatment groups as defined in the *Experimental Design* described herein. Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids placed on coated leaves treated with control began reaching maturity (5th instar reproducing stage; 5R) at approximately 6 days (Fig. 6A). Development was delayed in aphids placed on coated leaves with rifampicin, and by 6 days of treatment, most aphids did not mature further than the 3rd instar stage, even after 12 days, and their size is drastically affected (Fig. 6A and 6B).

These data indicate that leaf coating with rifampicin impaired aphid development.

Antibiotic treatment delivered through leaf coating increased aphid mortality

15 Survival rate of aphids was also measured during the leaf coating treatments. Aphids placed on coated leaves with rifampicin had lower survival rates than aphids placed on coated leaves with the control (Fig. 7). These data indicate that rifampicin treatment delivered through leaf coating affected aphid survival.

20 *Antibiotic treatment delivered through leaf coating decreased Buchnera in aphids*

To test whether rifampicin delivered through leaf coating, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 6 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers.

25 Aphids placed on leaves treated with control had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids placed on leaves treated with rifampicin had a drastic reduction of *Buchnera*/aphid DNA copies (Fig. 8), indicating that rifampicin leaf coating treatment eliminated endosymbiotic *Buchnera*.

Microinjection Delivery Experimental Design:

30 Microinjection was performed using NanoJet III Auto-Nanoliter Injector with the in-house pulled borosilicate needle (Drummond Scientific; Cat# 3-000-203-G/XL). Aphids (eNASCO strain, *Acyrtosiphon pisum*) were grown on fava bean plants as described in a previous Example. Aphids are transferred using a paint brush to a tubing system connected to vacuum (Fig. 1C). The injection site was at the ventral thorax of the aphid. The injection solutions were either the organosilicone surfactant solvent 0.025% Silwet L-77 (Lehle Seeds, Cat No VIS-01) in water (negative control), or 50 µg/ml of rifampicin formulated in solvent solution. The injection volume was 10 nl for nymph and 20 nl for adult (both at a rate of 2 nl/sec). Each treatment group had approximately the same number of individuals injected from each of the collection plants. After injection, aphids were released into a petri dish with fava bean leaves, whose stems are in 2% agar.

40

Microinjection with antibiotic treatment decreased Buchnera in aphids

To test whether rifampicin delivered by microinjection results in loss of *Buchnera* in aphids, and that this loss impacts aphid fitness as demonstrated in previous Examples, DNA was extracted from aphids in each treatment group after 4 days of treatment and qPCR was performed as described in a previous Example to determine the *Buchnera*/aphid copy numbers.

Aphids microinjected with negative control had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphid nymphs and adults microinjected with rifampicin had a drastic reduction of *Buchnera*/aphid DNA copies (Fig. 9), indicating that rifampicin microinjection treatment decreased the presence of endosymbiotic *Buchnera*.

10

Topical Delivery Experimental Design:

Aphids (LSR-1 strain, *Acyrtosiphon pisum*) were grown on fava bean plants as described in a previous Example. Spray bottles were filled with 2 ml of control (0.025% Silwet L-77) or rifampicin solutions (50 µg/ml of in solvent solution). Aphids (number = 10) were transferred to the bottom of a clean petri dish and gathered to the corner of the petri dish using a paint brush. Subsequently, aphids were separated into two cohorts and sprayed with ~100 µl of either control or rifampicin solutions. Immediately after spraying, the petri dish was covered with a lid. Five minutes after spraying, aphids were released into a petri dish with fava bean leaves with stems in 2% agar.

Topical delivery of antibiotic treatment decreased Buchnera in aphids

To test whether rifampicin delivered by topical delivery results in loss of *Buchnera* in aphids, and that this loss impacts aphid fitness as demonstrated in previous Examples, DNA was extracted from aphids in each treatment group after 3 days of treatment and qPCR as described in a previous Example was performed to determine the *Buchnera*/aphid copy numbers.

Aphids sprayed with the control solution had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids sprayed with rifampicin had a drastic reduction of *Buchnera*/aphid DNA copies (Fig. 10), indicating that rifampicin treatment delivered through topical treatment decreases the presence of endosymbiotic *Buchnera*.

Leaf injection method A – Leaf perfusion and cutting

Experimental Design:

Aphids LSR-1 (which harbor only *Buchnera*), *Acyrtosiphon pisum* were grown on fava bean plants (*Vroma vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24°C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first and second instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) negative control (leaf injected with water plus blue food coloring) and 2) leaf injected with 50 µg/ml rifampicin in water plus blue food coloring. Each treatment group received approximately the same number of

40

individuals from each of the collection plants. For treatment, rifampicin stock solution (25 mg/ml in 100% methanol) was diluted to 50 µg/ml in water plus blue food coloring. The solution was then placed into a 1.5 ml Eppendorf tube with a fava bean stem perfused with the solutions and the opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat#
5 FB0875711) and aphids were applied to the leaves of the plant. For each treatment, 74-81 aphids were placed onto each leaf. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered. In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th, and 5R (5th that has reproduced) instar) was determined daily throughout the experiment.

10 *Antibiotic treatment delivered through leaf injection method A delays and stops progression of aphid development*

LSR-1 1st and 2nd instar aphids were divided into two separate treatment groups as defined in Leaf injection method A – Leaf perfusion and cutting Experimental Design (described herein). Aphids
15 were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with water plus food coloring began reaching maturity (5th instar stage) at approximately 6 days (Fig. 11). Development was delayed in aphids feeding on rifampicin injected leaves, and by 6 days of treatment, most aphids did not mature further than the 4th instar stage. Even after 8 days, the development of aphids feeding on rifampicin injected leaves was drastically delayed (Fig. 11). These
20 data indicate that rifampicin treatment via leaf perfusion impaired aphid development.

Antibiotic treatment delivered through leaf injection method A increased aphid mortality

Survival rate of aphids was also measured during the leaf perfusion experiments. Aphids placed on leaves injected with rifampicin had lower survival rates than aphids placed on leaves injected with the
25 control solution (Fig. 12). These data indicate that rifampicin treatment delivered through leaf injection affected aphid survival.

Antibiotic treatment delivered thorough leaf injection method A results in decreased levels of Buchnera

To test whether rifampicin delivered via leaf perfusion results in loss of *Buchnera* in aphids, and
30 that this loss impacts aphid fitness as demonstrated in previous Examples, DNA was extracted from aphids in each treatment group after 8 days of treatment and qPCR as described in a previous Example was performed to determine the *Buchnera*/aphid copy numbers.

Aphids feeding on leaves injected with the control solution had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids feeding on leaves injected with rifampicin had a reduction of
35 *Buchnera*/aphid DNA copies (Fig. 13), indicating that rifampicin treatment delivered via leaf injection decreases the presence of endosymbiotic *Buchnera*, as shown in previous Examples, and resulted in a fitness decrease.

*Leaf perfusion and delivery through plant**Experimental Design:*

Aphids LSR-1 (which harbor only *Buchnera*), *Acyrtosiphon pisum* were grown on fava bean plants (*Vroma vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were

grown in potting soil at 24°C with 16 h of light and 8 h of darkness.

To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first and second instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) aphids placed on leaves injected with the negative control solution (water and food coloring) and placed into an Eppendorf tube with the negative control solution, or 2) aphids placed on leaves that were injected with 100 ug/ml rifampicin in water plus food coloring and put into an Eppendorf tube with 100 ug/ml rifampicin in water. Each treatment group received approximately the same number of individuals from each of the collection plants.

For treatment, rifampicin stock solution (25 mg/ml in 100% methanol) was diluted to 100 µg/ml in water plus blue food coloring. The solution was then placed into a 1.5 ml Eppendorf tube with a fava bean stem with a leaf also perfused with the solutions and the opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

For each treatment, 49-50 aphids were placed onto each leaf. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered.

In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th, and 5R (5th that has reproduced) instar) was determined daily throughout the experiment.

25

Antibiotic treatment delivered through leaf injection and delivery through plant delays and stops progression of aphid development

LSR-1 1st and 2nd instar aphids were divided into two separate treatment groups as defined in *Leaf perfusion and delivery through plant Experimental Design* (described herein). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with the control solution (water plus food coloring only) began reaching maturity (5th instar stage) at approximately 6 days (Fig. 14).

Development was delayed in aphids treated with rifampicin, and by 6 days of treatment, most aphids did not mature further than the 3rd instar stage. Even after 8 days, their development was drastically delayed (Fig. 14). These data indicate that rifampicin treatment via leaf perfusion impaired aphid development.

Antibiotic treatment delivered through leaf injection and delivery through plant increased aphid mortality

Survival rate of aphids was also measured during the experiments where aphids were treated with either control solution or rifampicin delivered via leaf perfusion and through the plant. Aphids feeding

40

on leaves perfused and treated with rifampicin had lower survival rates than aphids feeding on leaves perfused and treated with the control solution (Fig. 15). These data indicate that rifampicin treatment delivered through leaf perfusion and through the plant negatively impacted aphid survival.

5 *Antibiotic treatment delivered via leaf injection and through the plant results in decreased levels of Buchnera*

To test whether rifampicin delivered via leaf perfusion and through the plant results in loss of *Buchnera* in aphids, and that this loss impacts aphid fitness as demonstrated in previous Examples, DNA was extracted from aphids in each treatment group after 6 and 8 days of treatment and qPCR was
10 performed to determine the *Buchnera*/aphid copy numbers, as described in previous Examples.

Aphids feeding on leaves injected and treated with the control solution had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids feeding on leaves perfused and treated with rifampicin had a statistically significant reduction of *Buchnera*/aphid DNA copies at both time points ($p=0.0037$ and $p=0.0025$ for days 6 and 8, respectively) (Figs. 16A and 16B), indicating that rifampicin treatment
15 delivered via leaf perfusion and through the plant decreased the presence of endosymbiotic *Buchnera*, and as shown in previous Examples, and resulted in a fitness decrease.

Combination delivery method

Experimental Design:

20 Aphids LSR-1 (which harbor only *Buchnera*), *Acyrtosiphon pisum* were grown on fava bean plants (*Vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; $60\pm5\%$ RH; 20 ± 2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24 °C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old
25 plants, and allowed to multiply to high density for 5–7 days.

For experiments, first and second instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) treatment with Silwet-L77 or water control solutions or 2) treatment with rifampicin diluted in silwet L-77 or water. Each treatment group received approximately the same number of individuals from each of the collection plants. The combination of delivery methods was as
30 follows: a) Topical delivery to aphid and plant by spraying 0.025% nonionic organosilicone surfactant solvent Silwet L-77 (negative control) or 100 µg/ml of rifampicin formulated in solvent solution using a 30 mL spray bottle and b) Delivery through plant with either water (negative control) or 100 µg/ml of rifampicin formulated in water. For treatment, rifampicin stock solution (25 mg/ml in 100% methanol) was diluted to 100 µg/ml in Silwet L-77 (for topical treatment to aphid and coating the leaf) or water (for
35 delivery through plant). The solution was then placed into a 1.5 ml Eppendorf tube with a fava bean stem with a leaf also perfused with the solutions and the opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

For each treatment, 76-80 aphids were placed onto each leaf. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered.

In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th, and 5R (5th that has reproduced) instar) was determined daily throughout the experiment.

Combination antibiotic treatment delays aphid development

LSR-1 1st and 2nd instar aphids were divided into two separate treatment groups as defined in *Combination delivery method Experimental Design* (described herein). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Control treated aphids began reaching maturity (5th instar stage) at approximately 6 days (Fig. 17). Development was delayed in aphids treated with rifampicin, and by 6 days of treatment, most aphids did not mature further than the 3rd instar stage, even after 7 days their development was drastically delayed (Fig. 17). These data indicate that a combination of rifampicin treatments impaired aphid development.

Combination antibiotic treatment results in increased aphid mortality

Survival rate of aphids was also measured during the experiments where aphids were treated with a combination of rifampicin treatments. Rifampicin treated aphids had slightly lower survival rates than aphids treated with control solutions (Fig. 18). These data indicate that rifampicin treatment delivered through a combination of treatments affected aphid survival.

Combination antibiotic treatment in decreased levels of Buchnera

To test whether rifampicin delivered via a combination of methods results in loss of *Buchnera* in aphids, and that this loss impacts aphid fitness as demonstrated in previous Examples, DNA was extracted from aphids in each treatment group after 7 days of treatment and qPCR as described in a previous Example was performed to determine the *Buchnera*/aphid copy numbers.

Aphids treated with the control solutions had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids treated with rifampicin had a statistically significant and drastic reduction of *Buchnera*/aphid DNA copies ($p=0.0227$; Fig. 19), indicating that rifampicin treatment delivered via a combination of methods decreases the presence of endosymbiotic *Buchnera*, and as shown in previous Examples, this resulted in a fitness decrease.

Together this data described in the previous Examples demonstrated the ability to kill and decrease the development, reproductive ability, longevity, and endogenous bacterial populations, e.g., fitness, of aphids by treating them with an antibiotic through multiple delivery methods.

Example 15: Insects treated with a natural antimicrobial polysaccharide

This Example demonstrates the treatment of aphids with Chitosan, a natural cationic linear polysaccharide of deacetylated beta-1,4-D-glucosamine derived from chitin. Chitin is the structural element in the exoskeleton of insects, crustaceans (mainly shrimp and crabs) and cell walls of fungi, and the second most abundant natural polysaccharide after cellulose. This Example demonstrates that the

effect of chitosan on insects was mediated through the modulation of bacterial populations endogenous to the insect that were sensitive to chitosan. One targeted bacterial strain is *Buchnera aphidicola*.

Therapeutic Design

5 The chitosan solution was formulated using a combination of leaf perfusion and delivery through plants (Fig. 20). The control solution was leaf injected with water + blue food coloring and water in tube. The treatment solution with 300 ug/ml chitosan in water (low molecular weight) via leaf injection (with blue food coloring) and through plant (in Eppendorf tube).

10 *Leaf perfusion-Plant Delivery Experimental Design:*

Aphids LSR-1 (which harbor only *Buchnera*), *Acyrtosiphon pisum* were grown on fava bean plants (*Vroma vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24 °C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first and second instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) negative control (water treated), 2) The treatment solution included 300 ug/ml chitosan in water (low molecular weight). Each treatment group received approximately the same number of individuals from each of the collection plants.

Chitosan (Sigma, catalog number 448869-50G) stock solution was made at 1% in acetic acid, sterilized autoclaving, and stored at 4 °C. For treatment (see *Therapeutic design*), the appropriate amount of stock solution was diluted with water to obtain the final treatment concentration of chitosan. The solution was then placed into a 1.5 ml Eppendorf tube with a fava bean stem with a leaf also perfused with the solutions and the opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

For each treatment, 50-51 aphids were placed onto each leaf. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered.

In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th, and 5R (5th that has reproduced) instar) was determined daily throughout the experiment.

After 8 days of treatment, DNA was extracted from multiple aphids from each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 240) and Buch_groES_98R

(CTGTTCTCGAGTCGATTTCC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

10 *There was a negative response on insect fitness upon treatment with the natural antimicrobial polysaccharide*

LSR-1 *A. pisum* 1st and 2nd instar aphids were divided into two separate treatment groups as defined in *Experimental Design* (above). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with the negative control alone began reaching maturity (5th instar stage) at approximately 6 days (Fig. 21). Development was delayed in aphids treated with chitosan solution, and by 6 days of treatment with chitosan, most aphids did not mature further than the 4th instar stage. These data indicate that treatment with chitosan delayed and stopped progression of aphid development.

Chitosan treatment increased aphid mortality

Survival rate of aphids was also measured during the treatments. The majority of the aphids treated with the control alone were alive at 3 days post-treatment (Fig. 22). After 4 days, aphids began gradually dying, and some survived beyond 7 days post-treatment.

- 5 In contrast, aphids treated with chitosan solution had lower survival rates than aphids treated with control. These data indicate that there was a decrease in survival upon treatment with the natural antimicrobial polysaccharide.

Chitosan treatment decreased Buchnera in aphids

- 10 To test whether the chitosan solution treatment, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 8 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids treated with control alone had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids treated with 300 ug/ml chitosan in water had ~2-5-fold less *Buchnera*/aphid DNA copies (Fig. 23),
15 indicating that chitosan treatment decreased *Buchnera* levels.

Together this data described previously demonstrated the ability to kill and decrease the development, longevity, and endogenous bacterial populations, e.g., fitness, of aphids by treating them with a natural antimicrobial polysaccharide.

20 **Example 16: Insects treated with nisin, a natural antimicrobial peptide**

- This Example demonstrates the treatment of aphids with the natural, "broad spectrum", polycyclic antibacterial peptide produced by the bacterium *Lactococcus lactis* that is commonly used as a food preservative. The antibacterial activity of nisin is mediated through its ability to generate pores in the bacterial cell membrane and interrupt bacterial cell-wall biosynthesis through a specific lipid II interaction.
25 This Example demonstrates that the negative effect of nisin on insect fitness is mediated through the modulation of bacterial populations endogenous to the insect that were sensitive to nisin. One targeted bacterial strain is *Buchnera aphidicola*.

Therapeutic Design:

- 30 Nisin was formulated using a combination of leaf perfusion and delivery through plants. The control solution (water) or treatment solution (nisin) was injected into the leaf and placed in the Eppendorf tube. The treatment solutions consisted of 1.6 or 7 mg/ml nisin in water.

Leaf perfusion-Plant Delivery Experimental Design:

- 35 LSR-1 aphids, *Acyrtosiphon pisum* were grown on fava bean plants (Vroma vicia faba from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24°C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to
40 high density for 5–7 days. For experiments, first and second instar aphids were collected from healthy

plants and divided into 2 different treatment groups: 1) negative control (water treated), 2) nisin treated with either 1.6 or 7 mg/ml nisin in water. Each treatment group received approximately the same number of individuals from each of the collection plants.

For treatment (see Therapeutic design), nisin (Sigma, product number: N5764) solution was made at 1.6 or 7 mg/ml (w/v) in water and filter sterilized using a 0.22 μ m syringe filter. The solution was then injected into the leaf of the plant and the stem of the plant was placed into a 1.5 ml Eppendorf tube. The opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

For each treatment, 56-59 aphids were placed onto each leaf. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered.

In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th, and 5R (5th instar aphids that are reproducing) instar) was determined daily throughout the experiment.

After 8 days of treatment, DNA was extracted from the remaining aphids in each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 240) and Buch_groES_98R (CTGTTCTCGAGTCGATTTCC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

There was a dose-dependent negative response on insect fitness upon treatment with nisin

LSR-1 *A. pisum* 1st and 2nd instar aphids were divided into three separate treatment groups as defined in Experimental Design (above). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with the negative control solution (water) began reaching maturity (5th instar stage) at approximately 6 days, and reproducing (5R stage) by 7 days (Fig. 24). Development was severely delayed in aphids treated with 7 mg/ml nisin. Aphids treated with 7 mg/ml nisin only developed to the 2nd instar stage by day 3, and by day 6, all aphids in the group were dead (Fig. 24). Development was slightly delayed in aphids treated with the lower concentration of nisin (1.6 mg/ml) and at each time point assessed, there were more less-developed aphids compared to water-treated controls (Fig. 24). These data indicate that treatment with nisin delayed and stopped progression of aphid development and this delay in development was dependent on the dose of nisin administered.

However, it is important to note that treatment with 7 mg/ml of nisin also had a negative impact on the health of the leaves used in the assay. Shortly after leaf perfusion of 7mg/ml of nisin it started turning brown. After two days, the leaves perfused with 7mg/ml turned black. There was no noticeable difference in the condition of the leaves treated with 1.6 mg/ml nisin.

5

Treatment with nisin resulted in increased aphid mortality

Survival rate of aphids was also measured during the treatments. Approximately 50% of aphids treated with the control alone survived the 8-day experiment (Fig. 25). In contrast, survival was significantly less for aphids treated with 7 mg/ml nisin ($p < 0.0001$, by Log-Rank Mantel Cox test), and all aphids in this group succumbed to the treatment by 6 days (Fig. 25). Aphids treated with the lower dose of nisin (1.6 mg/ml) had higher mortality compared to control treated aphids, although the difference did not reach statistical significance ($p = 0.0501$ by Log-Rank Mantel Cox test). These data indicate that there was a dose-dependent decrease in survival upon treatment with nisin.

15 *Treatment with nisin resulted in decreased Buchnera in aphids*

To test whether treatment with nisin, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 8 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids treated with control alone had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids treated with 1.6 mg/ml nisin had ~1.4-fold less *Buchnera*/aphid DNA copies after 8 days of treatment (Fig. 26). No aphids were alive in the group treated with 7 mg/ml nisin, and therefore, *Buchnera*/aphid DNA copies was not assessed in this group. These data indicate that nisin treatment decreased *Buchnera* levels.

Together this data described previously demonstrate the ability to kill and decrease the development, longevity, and endogenous bacterial populations, e.g., fitness, of aphids by treating them with the antimicrobial peptide nisin.

25

Example 17: Insects treated with levulinic acid decreases insect fitness

This Example demonstrates the treatment of aphids with levulinic acid, a keto acid produced by heating a carbohydrate with hexose (e.g., wood, starch, wheat, straw, or cane sugar) with the addition of a dilute mineral acid reduces insect fitness. Levulinic acid has previously been tested as an antimicrobial agent against *Escherichia coli* and *Salmonella* in meat production (Carpenter et al., 2010, Meat Science; Savannah G. Hawkins, 2014, University of Tennessee Honors Thesis). This Example demonstrates that the effect of levulinic acid on insects was mediated through the modulation of bacterial populations endogenous to the insects that were sensitive to levulinic acid. One targeted bacterial strain is *Buchnera aphidicola*.

35

Therapeutic Design:

The levulinic acid was formulated using a combination of leaf perfusion and delivery through plants. The control solution was leaf injected with water and water was placed in the Eppendorf tube.

The treatment solutions included 0.03 or 0.3% levulinic acid in water via leaf injection and through plant (in Eppendorf tube).

Leaf perfusion-Plant Delivery

5 *Experimental Design:*

eNASCO aphids, *Acyrtosiphon pisum* were grown on fava bean plants (*Vroma vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24°C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first and second instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) negative control (water treated), 2) The treatment solution included 0.03 or 0.3% levulinic acid in water. Each treatment group received approximately the same number of individuals from each of the collection plants.

15 For treatment (see *Therapeutic design*), levulinic acid (Sigma, product number: W262706) was diluted to either 0.03 or 0.3% in water. The solution was then placed into a 1.5 ml Eppendorf tube with a fava bean stem with a leaf also perfused with the solutions and the opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

20 For each treatment, 57-59 aphids were placed onto each leaf. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered.

In addition, the developmental stage (1st, 2nd, 3rd, 4th, and 5th instar) was determined daily throughout the experiment.

25 After 7 of treatment, DNA was extracted from the remaining aphids in each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch groES_18F (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 240) and Buch groES_98R (CTGTTCCCTCGAGTCGATTTC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

There was a dose-dependent negative response on insect fitness upon treatment with levulinic acid

eNASCO *A. pisum* 1st and 2nd instar aphids were divided into three separate treatment groups as defined in *Experimental Design* (above). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with the negative control alone began reaching maturity (5th instar stage) at approximately 7 days (Fig. 27). Development was delayed in aphids treated with levulinic acid and by 11 days post-treatment, nearly all control treated aphids reached maturity while ~23 and 63% aphids treated with 0.03 and 0.3% levulinic acid, respectively, did not mature further than the 4th instar stage. These data indicate that treatment with levulinic acid delayed and stopped progression of aphid development and this delay in development is dependent on the dose of levulinic acid administered.

Treatment with levulinic acid results in increased aphid mortality

Survival rate of aphids was also measured during the treatments. Approximately 50% of aphids treated with the control alone survived the 11-day experiment (Fig. 28). In contrast, survival was significantly less for aphids treated with 0.3% levulinic acid ($p < 0.01$). Aphids treated with the low dose of levulinic acid (0.03%) had higher mortality compared to control treated aphids, although the difference did not reach statistical significance. These data indicate that there was a dose-dependent decrease in survival upon treatment with levulinic acid.

Treatment with levulinic acid results in decreased Buchnera in aphids

To test whether treatment with levulinic acid, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 7 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids treated with control alone had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids treated with 0.03 or 0.3% levulinic acid in water had ~6-fold less *Buchnera*/aphid DNA copies after 7 days of treatment (Fig. 29, left panel). These data indicate that levulinic acid treatment decreased *Buchnera* levels.

Together this data described previously demonstrated the ability to kill and decrease the development, longevity, and endogenous bacterial populations, e.g., fitness, of aphids by treating them with levulinic acid.

30

Example 18: Insects treated with a plant derived secondary metabolite solution

This Example demonstrates the treatment of aphids with gossypol acetic acid, a natural phenol derived from the cotton plant (genus *Gossypium*) that permeates cells and acts as an inhibitor for several dehydrogenase enzymes. This Example demonstrates that the effect of gossypol on insects was mediated through the modulation of bacterial populations endogenous to the insect that were sensitive to gossypol. One targeted bacterial strain is *Buchnera aphidicola*.

Therapeutic Design: The gossypol solution was formulated depending on the delivery method:

1) Through the plants: with 0 (negative control) or 0.5%, 0.25%, and 0.05% of gossypol formulated in an artificial diet (based on Akey and Beck, 1971; see *Experimental Design*) without

40

essential amino acids (histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, and valine).

2) Microinjection: injection solutions were either 0.5% of gossypol or artificial diet only (negative control).

Plant Delivery Experimental Design:

5 Aphids (either eNASCO (which harbor both *Buchnera* and *Serratia* primary and secondary symbionts, respectively) or LSR-1 (which harbor only *Buchnera*) strains, *Acyrtosiphon pisum*) were grown on fava bean plants (*Vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24 °C with 16 h of light and 8 h of darkness. To limit
10 maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first and second instar aphids were collected from healthy plants and divided into 4 different treatment groups: 1) artificial diet alone without essential amino acids, 2) artificial diet alone without essential amino acids and 0.05% of gossypol, 3) artificial diet alone without essential amino acids and 0.25% of gossypol,
15 and 4) artificial diet alone without essential amino acids and 0.5% of gossypol. Each treatment group received approximately the same number of individuals from each of the collection plants.

The artificial diet used was made as previously published (Akey and Beck, 1971 Continuous Rearing of the Pea Aphid, *Acyrtosiphon pisum*, on a Holidic Diet), with and without the essential amino acids (2 mg/ml histidine, 2 mg/ml isoleucine, 2 mg/ml leucine, 2 mg/ml lysine, 1 mg/ml methionine, 1.62
20 mg/ml phenylalanine, 2 mg/ml threonine, 1 mg/ml tryptophan, and 2 mg/ml valine), except neither diet included homoserine or beta-alanyltyrosine. The pH of the diets was adjusted to 7.5 with KOH and diets were filter sterilized through a 0.22 µm filter and stored at 4 °C for short term (<7 days) or at -80 °C for long term.

Gossypol acetic acid (Sigma, Cat#G4382-250MG) stock solution was made at 5% in methanol
25 and sterilized by passing through a 0.22 µm syringe filter, and stored at 4 °C. For treatments (see *Therapeutic design*), the appropriate amount of stock solution was added to the artificial diet to obtain the different final concentrations of gossypol. The diet was then placed into a 1.5 ml Eppendorf tube with a fava bean stem with a leaf and the opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to
30 the leaves of the plant.

For each treatment, 15-87 aphids were placed onto each leaf. Artificial diet feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish housing the artificial feeding system when they were discovered.

35 In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th, and 5R (5th that has reproduced) instar) was determined daily throughout the experiment. Once an aphid reached the 4th instar stage, they were given their own artificial feeding system in a deep petri dish so that fecundity could be monitored once they reached adulthood.

For adult aphids, new nymphs were counted daily and then discarded. At the end of the
40 experiments, fecundity was measured in two ways: 1) the mean day at which the first offspring for the

treatment group was determined and 2) the mean number of offspring produced daily once the aphid reached adulthood. Pictures of aphids were taken throughout the experiment to evaluate size differences between treatment groups.

After 5 or 13 days of treatment, DNA was extracted from multiple aphids from each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGCTTGCTTAAG; SEQ ID NO: 240) and Buch_groES_98R (CTGTTCTCGAGTCGATTTC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

There was a dose-dependent negative response on insect fitness upon treatment with the allelochemical gossypol

eNASCO and LSR-1 *A. pisum* 1st and 2nd instar aphids were divided into four separate treatment groups as defined in *Experimental Design* (described herein). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with artificial diet alone began reaching maturity (5th instar stage) at approximately 3 days (Fig. 30A). Development was delayed in aphids treated with gossypol, and by 5 days of treatment with 0.5% of gossypol, most aphids did not mature further than the 3rd instar stage, and their size is also affected (Fig. 30A and 30B). These data indicate that treatment with gossypol delayed and stopped progression of aphid development, and that this response was dose dependent.

Gossypol treatment increased aphid mortality

Survival rate of aphids was also measured during the treatments. The majority of the aphids treated with artificial diet alone without essential amino acids were alive at 2 days post-treatment (Fig. 31). After 4 days, aphids began gradually dying, and some survived beyond 7 days post-treatment.

In contrast, aphids treated with 0.25 (not significantly different than control, $p=0.2794$) and 0.5% of gossypol had lower survival rates than aphids treated with artificial diet alone, with 0.5% gossypol treatment being significantly different than AD no EAA control ($p=0.002$). 0.5% gossypol-treated aphids began dying after 2 days of treatment and all aphids succumbed to treatment by 4 days. Aphids treated with 0.25% survived a bit longer than those treated with 0.5% but were also drastically affected. These data indicate that there was a dose-dependent decrease in survival upon treatment with the allelochemical gossypol.

Gossypol treatment decreased aphid reproduction

Fecundity was also monitored in aphids during the treatments. By days 7 and 8 post-treatment, the majority of the adult aphids treated with artificial diet without essential amino acids began
 5 reproducing. The mean number of offspring produced daily after maturity by aphids treated with artificial diet without essential amino acids was approximately 5 (Fig. 32A and 32B).

In contrast, aphids treated with 0.25% of gossypol show a reduction to reach adulthood and produce offspring. These data indicate that gossypol treatment resulted in a decrease of aphid reproduction.

Gossypol treatment decreased Buchnera in aphids

To test whether different concentrations of gossypol, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 5 or 13 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy
 15 numbers. Aphids treated with artificial diet alone without essential amino acids (control) had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids treated with 0.25 and 0.5% of gossypol had ~4-fold less *Buchnera*/aphid DNA copies (Fig. 33), indicating that gossypol treatment decreased *Buchnera* levels, and that this decrease was concentration dependent.

Microinjection delivery experimental design:

Microinjection was performed using NanoJet III Auto-Nanoliter Injector with the in-house pulled borosilicate needle (Drummond Scientific; Cat# 3-000-203-G/XL). Aphids (LSR-1 strain, *A. pisum*) were grown on fava bean plants as described in a previous Example. Each treatment group had approximately the same number of individuals injected from each of the collection plants. Nymph aphids (<3rd instar
 25 stage) were transferred using a paint brush to a tubing system connected to vacuum and microinjected into the ventral thorax with 20 nl of either artificial diet without essential amino acids (negative control) or 0.05% of gossypol solution in artificial diet without essential amino acids. After injection, aphids were placed in a deep petri dish with a fava bean leaf with stem in 2% agar.

Microinjection with antibiotic treatment decreased Buchnera in aphids

To test whether gossypol delivered by microinjection results in loss of *Buchnera* in aphids, and that this loss impacts aphid fitness as demonstrated in previous Examples, DNA was extracted from aphids in each treatment group after 4 days of treatment and qPCR was performed as described in a previous Example to determine the *Buchnera*/aphid copy numbers.

35 Aphids microinjected with negative control had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphid nymphs and adults microinjected with gossypol had a drastic reduction of *Buchnera*/aphid DNA copies (Fig. 34), indicating that gossypol microinjection treatment decreases the presence of endosymbiotic *Buchnera*, and as shown in previous Examples this resulted in a fitness decrease.

Together this data described in the previous Examples demonstrated the ability to kill and
 40 decrease the development, reproductive ability, longevity, and endogenous bacterial populations, e.g.,

fitness, of aphids by treating them with plant secondary metabolite solution through multiple delivery methods.

Example 19: Insects treated with natural plant derived antimicrobial compound, trans-cinnamaldehyde

This Example demonstrates the treatment of aphids with trans-cinnamaldehyde, a natural aromatic aldehyde that is the major component of bark extract of cinnamon (*Cinnamomum zeylandicum*) results in decreased fitness. Trans-cinnamaldehyde has been shown to have antimicrobial activity against both gram-negative and gram-positive organisms, although the exact mechanism of action of its antimicrobial activity remains unclear. Trans-cinnamaldehyde may damage bacterial cell membranes and inhibit of specific cellular processes or enzymes (Gill and Holley, 2004 Applied Environmental Microbiology). This Example demonstrates that the effect of trans-cinnamaldehyde on insects was mediated through the modulation of bacterial populations endogenous to the insect that were sensitive to trans-cinnamaldehyde. One targeted bacterial strain is *Buchnera aphidicola*.

Therapeutic Design:

Trans-cinnamaldehyde was diluted to 0.05%, 0.5%, or 5% in water and was delivered through leaf perfusion (~1 ml was injected into leaves) and through plants.

Experimental Design:

Aphids (LSR-1 (which harbor only *Buchnera*) strains, *Acyrtosiphon pisum*) were grown on fava bean plants (Vroma vicia faba from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24 °C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first and second instar aphids were collected from healthy plants and divided into four different treatment groups: 1) water treated controls, 2) 0.05% trans-cinnamaldehyde in water, 3) 0.5% trans-cinnamaldehyde in water, and 4) 5% trans-cinnamaldehyde in water. Each treatment group received approximately the same number of individuals from each of the collection plants.

Trans-cinnamaldehyde (Sigma, Cat#C80687) was diluted to the appropriate concentration in water (see Therapeutic design), sterilized by passing through a 0.22 µm syringe filter, and stored at 4 °C. Fava bean leaves were injected with approximately 1 ml of the treatment and then the leaf was placed in a 1.5 ml Eppendorf tube containing the same treatment solution. The opening of the tube where the fava bean stem was placed was closed using parafilm. This treatment feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

For each treatment, 40-49 aphids were placed onto each leaf. Treatment feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish housing the treatment feeding system when they were discovered.

In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th, and 5R (5th that has reproduced) instar) was determined daily throughout the experiment.

After 3 days of treatment, DNA was extracted from the remaining living aphids from each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 240) and Buch_groES_98R (CTGTTCTCGAGTCGATTTCC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

There was a dose-dependent negative response on insect fitness upon treatment with the natural antimicrobial trans-cinnamaldehyde

LSR-1 *A. pisum* 1st and 2nd instar aphids were divided into four separate treatment groups as defined in Experimental Design (described herein). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with water alone began reaching the 3rd instar stage at 3 days post-treatment (Fig. 35). Development was delayed in aphids treated with trans-cinnamaldehyde, and by 3 days of treatment with each the three of the trans-cinnamaldehyde concentrations, none of the aphids matured past the second instar stage (Fig. 35). Moreover, all the aphids treated with the highest concentration of trans-cinnamaldehyde (5%) remained at the 1st instar stage throughout the course of the experiment. These data indicate that treatment with trans-cinnamaldehyde delays and stops progression of aphid development, and that this response is dose dependent.

Trans-cinnamaldehyde treatment increased aphid mortality

Survival rate of aphids was also measured during the treatments. Approximately 75 percent of the aphids treated with water alone were alive at 3 days post-treatment (Fig. 36). In contrast, aphids treated with 0.05%, 0.5%, and 5% trans-cinnamaldehyde had significantly lower ($p < 0.0001$ for each treatment group compared to water treated control) survival rates than aphids treated with water alone. These data indicate that there was a dose-dependent decrease in survival upon treatment with the natural antimicrobial trans-cinnamaldehyde.

Trans-cinnemaldehyde treatment decreased Buchnera in aphids

To test whether different concentrations of trans-cinnemaldehyde, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 3 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids treated with water alone (control) had high ratios of *Buchnera*/aphid DNA copies. Similarly, aphids treated with the lowest concentration of trans-cinnemaldehyde (0.5%) had high ratios of *Buchnera*/aphid DNA copies.

In contrast, aphids treated with 0.5 and 5% of trans-cinnemaldehyde had ~870-fold less *Buchnera*/aphid DNA copies (Fig. 37), indicating that trans-cinnemaldehyde treatment decreased *Buchnera* levels, and that this decrease was concentration dependent.

Together this data described in the previous Examples demonstrate the ability to kill and decrease the development, reproductive ability, longevity and endogenous bacterial populations, e.g., fitness, of aphids by treating them with plant secondary metabolite solution through multiple delivery methods.

15

Example 20: Insects treated with scorpion antimicrobial peptides

This Example demonstrates the treatment of aphids with multiple scorpion antimicrobial peptides (AMP), of which several are identified AMPs in the venom gland transcriptome of the scorpion *Urodacus yaschenkoi* (Luna-Ramirez et al., 2017, Toxins). AMPs typically have a net positive charge and hence, a high affinity for bacterial membranes. This Example demonstrates that the effect of the AMP on insects was mediated through the modulation of bacterial populations endogenous to the insect that were sensitive to AMP peptides. One targeted bacterial strain is *Buchnera aphidicola*, an obligate symbiont of aphids.

Therapeutic Design:

The Uy192 solution was formulated using a combination of leaf perfusion and delivery through plants. The control solution was leaf injected with water + blue food coloring and water in tube. The treatment solution consisted of 100 ug/ml Uy192 in water via leaf injection (with blue food coloring) and through plant (in Eppendorf tube).

30

Leaf perfusion-Plant Delivery Experimental Design:

Aphids LSR-1 (which harbor only *Buchnera*), *Acyrtosiphon pisum* were grown on fava bean plants (*Vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 20±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24°C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first and second instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) negative control (water treated), 2) The treatment solution of 100 ug/ml AMP in water. Each treatment group received approximately the same number of individuals from each of the collection plants.

40

Uy192 was synthesized by Bio-Synthesis at >75% purity. 1 mg of lyophilized peptide was reconstituted in 500 μ l of 80% acetonitrile, 20% water, and 0.1% TFA, 100 μ l (100 μ g) was aliquoted into 10 individual Eppendorf tubes, and allowed to dry. For treatment (see *Therapeutic design*), 1 ml of water was added to a 100 μ g aliquot of peptide to obtain the final concentration of Uy192 (100 μ g/ml). The solution was then placed into a 1.5 ml Eppendorf tube with a fava bean stem with a leaf also perfused with the solutions and the opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

For each treatment, 50 aphids were placed onto each leaf. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered.

In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th, and 5R (5th that has reproduced) instar) was determined daily throughout the experiment.

After 8 days of treatment, DNA was extracted from the remaining aphids in each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGTGCTTGTAAAG; SEQ ID NO: 240) and Buch_groES_98R (CTGTTCTCGAGTCGATTTC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

There was a negative response on insect fitness upon treatment with the scorpion AMPs

LSR-1 *A. pisum* 1st and 2nd instar aphids were divided into two separate treatment groups as defined in *Experimental Design* (above). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with the negative control alone began reaching maturity (5th instar stage) at approximately 6 days (Fig. 38). Development was delayed in aphids treated with Uy192, and after 8 days of treatment, aphids did not mature further than the 4th instar stage. These data indicate that treatment with Uy192 delayed and stopped progression of aphid development.

Treatment with scorpion AMPs results in increased aphid mortality

Survival rate of aphids was also measured during the treatments. The majority of the aphids treated with the control alone were alive at 3 days post-treatment (Fig. 39). After 4 days, aphids began gradually dying, and some survived beyond 7 days post-treatment.

In contrast, aphids treated with Uy192 had lower survival rates than aphids treated with control. These data indicate that there was a decrease in survival upon treatment with the scorpion AMP Uy192.

Treatment with scorpion AMP Uy192 results in decreased Buchnera in aphids

5 To test whether treatment with Uy192, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 8 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids treated with control alone had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids treated with 100 ug/ml Uy192 in water had ~7-fold less *Buchnera*/aphid DNA copies (Fig. 40), indicating that Uy192 treatment
10 decreased *Buchnera* levels.

Together this data described previously demonstrated the ability to kill and decrease the development, longevity and endogenous bacterial populations, e.g., fitness, of aphids by treating them with a natural scorpion antimicrobial peptide.

15

Example 21: Insects treated with scorpion antimicrobial peptides

This Example demonstrates the treatment of aphids with several scorpion antimicrobial peptides (AMPs) D10, D3, Uyct3, and Uy17, which have been recently identified AMPs in the venom gland transcriptome of the scorpion *Urodoacus yaschenkoi* (Luna-Ramirez et al., 2017, Toxins). AMPs typically
20 have a net positive charge and hence, a high affinity for bacterial membranes. This Example demonstrates that the effect of the AMPs on insects was mediated through the modulation of bacterial populations endogenous to the insect that were sensitive to AMP peptides. One targeted bacterial strain is *Buchnera aphidicola*, an obligate symbiont of aphids.

Aphids are agricultural insect pests causing direct feeding damage to the plant and serving as
25 vectors of plant viruses. In addition, aphid honeydew promotes the growth of sooty mold and attracts nuisance ants. The use of chemical treatments, unfortunately still widespread, leads to the selection of resistant individuals whose eradication becomes increasingly difficult.

Therapeutic Design:

30 The indicated peptide or peptide cocktail (see Aphid Microinjection Experimental Design and Leaf perfusion-Plant Experimental Design sections for details below) was directly microinjected into aphids or delivered using a combination of leaf perfusion and delivery through plants. As a negative control, aphids were microinjected with water (for microinjection experiments) or placed on leaves injected with water and water in tube (for leaf perfusion and plant delivery experiments). The treatment solutions consisted of 20
35 nl of 5 µg/µl of D3 or D10 dissolved in water (for aphid microinjections) or 40 µg/ml of a cocktail of D10, Uy17, D3, and UyCt3 in water via leaf injection and through plant (in Eppendorf tube).

Aphid Microinjection Experimental Design

Microinjection was performed using NanoJet III Auto-Nanoliter Injector with the in-house pulled
40 borosilicate needle (Drummond Scientific; Cat# 3-000-203-G/XL). Aphids (LSR-1 strain, *Acyrtosiphon*

pisum) were grown on fava bean plants (*Vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24 °C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. Each treatment group had approximately the same number of individuals injected from each of the collection plants. Adult aphids were microinjected into the ventral thorax with 20 nl of either water or 100 ng (20 µl of a 5 µg/ml solution of peptide D3 or D10. The microinjection rate as 5 nl/sec. After injection, aphids were placed in a deep petri dish containing a fava bean leaf with stem in 2% agar.

Peptides were synthesized by Bio-Synthesis at >75% purity. 1 mg of lyophilized peptide was reconstituted in 500 µl of 80% acetonitrile, 20% water, and 0.1% TFA, 100 µl (100 µg) was aliquoted into 10 individual Eppendorf tubes, and allowed to dry.

After 5 days of treatment, DNA was extracted from the remaining aphids in each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 240) and Buch_groES_98R (CTGTTCTCGAGTCGATTTCC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95 °C for 10 minutes, 2) 95 °C for 15 seconds, 3) 55 °C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95 °C for 15 seconds, 6) 55 °C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95 °C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

Microinjection of aphids with scorpion peptides D3 and D10 results in decreased insect survival

LSR-1 *A. pisum* 1st and 2nd instar aphids were divided into three separate treatment groups as defined in *Experimental Design* (described herein). Aphids were monitored daily and survival rate was determined. After 5 days of treatment, the aphids injected with the scorpion peptides had lower survival rates compared to water injected controls (9, 35, and 45% survival for injection with D3, D10, and water, respectively) (Fig. 41). These data indicate that there was a decrease in survival upon treatment with the scorpion AMPs D3 and D10.

Microinjection of aphids with scorpion peptides D3 and D10 results in a reduction of Buchnera endosymbionts

To test whether injection with the scorpion AMPs D3 and D10, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group 5 days after injection and qPCR was performed to determine the *Buchnera*/aphid copy

numbers. Aphids injected with water alone had high ratios of *Buchnera*/aphid DNA (47.4) while aphids injected with D3 and D10 had lower ratios of *Buchnera*/aphid DNA (25.3 and 30.9, respectively) (Fig. 42). These data indicate that treatment with scorpion peptides D3 and D10 resulted in decreased levels of the bacterial symbiont *Buchnera*.

5

Leaf perfusion-Plant Delivery Experimental Design:

eNASCO Aphids (which harbor *Buchnera* and *Serratia*), *Acyrtosiphon pisum* were grown on fava bean plants (*Vicia faba* from Johnny's Selected Seeds) as described above. For experiments, first and second instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) negative control (water treated), 2) The treatment solution consisted of 40 µg/ml of each D10, Uy17, D3, and UyCt3 in water. Each treatment group received approximately the same number of individuals from each of the collection plants.

Peptides were synthesized, dissolved, and aliquoted, as described above. For treatment (see Therapeutic design), water was added to a 100 µg aliquot of peptide to obtain the desired final concentration (40 µg/ml). The four peptides were combined to make the peptide cocktail solution. This solution was used to perfuse into leaves via injection. Following injection, the stems of the injected leaves were placed into a 1.5 ml Eppendorf tube which was then sealed with parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

For each treatment, 41-49 aphids were placed onto each leaf. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered.

Treatment with cocktail of scorpion peptides results in increased aphid mortality

Survival rate of aphids was also measured during the treatments. After 6 days of treatment, aphids treated with the peptide cocktail had lower survival rates compared to those treated with water, and after 9 days the effect is more evident (16 and 29% survival for peptide cocktail and water treated, respectively) (Fig. 43). These data indicate that there was a decrease in survival upon treatment with the cocktail of scorpion AMPs, and as shown in previous Examples these AMP decreased the endosymbiont levels in the aphids.

Together this data described previously demonstrated the ability to kill and decrease the longevity and endogenous bacterial populations, e.g., fitness, of aphids by treating them with single natural scorpion antimicrobial peptides or a peptide cocktail.

Example 22: Insects treated with an antimicrobial peptide fused to a cell penetrating peptide

This Example demonstrates the treatment of aphids with a fused scorpion antimicrobial peptide (AMP) (Uy192) to a cell penetrating peptide derived from a virus. The AMP Uy192 is one of several recently identified AMPs in the venom gland transcriptome of the scorpion *Urodacus yaschenkoi* (Luna-Ramirez et al., 2017, Toxins). AMPs typically have a net positive charge and hence, a high affinity for bacterial membranes. To enhance the delivery of the scorpion peptide Uy192 into aphid cells, the

peptide was synthesized fused to a portion of the transactivator of transcription (TAT) protein of human immunodeficiency virus I (HIV-1). Previous studies have shown that conjugating this cell penetrating peptide (CPP) to other molecules increased their uptake into cells via transduction (Zhou et al., 2015 Journal of Insect Science and Cermenati et al., 2011 Journal of Insect Physiology). This Example demonstrates that the effect of the fused peptide on insects was mediated through the modulation of bacterial populations endogenous to the insect that were sensitive to the antimicrobial peptide. One targeted bacterial strain is *Buchnera*.

Therapeutic Design

The scorpion peptide conjugated to the cell penetrating peptide and fluorescently tagged with 6FAM (Uy192+CPP+FAM) was formulated using a combination of leaf perfusion and delivery through plants. The control solution (water) or treatment solution (Uy192+CPP+FAM) was injected into the leaf and placed in the Eppendorf tube. The treatment solution included 100 µg/ml Uy192+CPP+FAM in water.

Leaf perfusion-Plant Delivery Experimental Design

LSR-1 aphids, *Acyrtosiphon pisum* were grown on fava bean plants (*Vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24°C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) negative control (water treated), 2) Uy192+CPP+FAM treated with 100 µg/ml Uy192+CPP+FAM in water. Each treatment group received approximately the same number of individuals from each of the collection plants.

For treatment (see Therapeutic design), Uy192+CPP+FAM (amino acid sequence: YGRKKRRQRRRFLSTIWNGIKGLL-FAM) was synthesized by Bio-Synthesis at >75% purity. 5 mg of lyophilized peptide was reconstituted in 1 ml of 80% acetonitrile, 20% water, and 0.1% TFA, 50 µl (100 µg) was aliquoted into individual Eppendorf tubes, and allowed to dry. For treatment (see Therapeutic design), 1 ml of sterile water was added to a 100 µg aliquot of peptide to obtain the final concentration of Uy192+CPP+FAM (100 µg/ml). The solution was then injected into the leaf of the plant and the stem of the plant was placed into a 1.5 ml Eppendorf tube. The opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant. Epi fluorescence imaging of the leaf confirmed that the leaves contained the green fluorescently tagged peptide in their vascular system.

For each treatment, 30 aphids were placed onto each leaf in triplicate. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered. In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th, and 5R (5th instar aphids that are reproducing) instar) was determined daily throughout the experiment.

At 5 days post-treatment, DNA was extracted from several aphids in each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions.

5 DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 240) and Buch_groES_98R (CTGTTCTCGAGTCGATTTCC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

15

Treatment with scorpion peptide Uy192 fused to a cell penetrating peptide delayed and stopped progression of aphid development

LSR-1 *A. pisum* 1st instar aphids were divided into three separate treatment groups as defined in Experimental Design (above). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Development for both aphids treated with water and those treated with the scorpion peptide fused to the cell penetrating peptide was similar for days 0 and 1 (Fig. 44). By day 2, however, control treated aphids developed to either in the second or third instar stage, while some aphids treated with the scorpion peptide fused to the cell penetrating peptide remained in the first instar stage (Fig. 44). Even at 3 days post-treatment, some aphids treated with the scorpion peptide fused to the cell penetrating peptide remained in the first instar stage (Fig. 44). By 7 days post-treatment, the majority of the water treated aphids developed to the 5th or 5th reproducing instar stage. In contrast, only 50 percent of aphids treated with the scorpion peptide fused to the cell penetrating peptide developed to the 5th instar stage, while ~42 and ~8 percent of aphids remained at the 3rd or 4th instar stage, respectively (Fig. 44). These data indicate that treatment with the scorpion peptide Uy192 fused to the cell penetrating peptide delayed and stopped progression of aphid development.

30

Treatment with the scorpion peptide Uy192 fused to a cell penetrating peptide resulted in increased aphid mortality

Survival rate of aphids was also measured during the treatments. Approximately 40% of aphids treated with the control alone survived the 7-day experiment (Fig. 45). In contrast, survival was significantly less for aphids treated with 100 µg/ml Uy192+CPP+FAM ($p=0.0036$, by Log-Rank Mantel Cox test), with only 16% of aphids surviving to day 7 (Fig. 45). These data indicate that there was a decrease in survival upon treatment with the scorpion peptide Uy192 fused to a cell penetrating peptide.

35

Treatment with a scorpion peptide fused to a cell penetrating peptide resulted in decreased Buchnera/aphid DNA ratios

To test whether treatment with treatment with Uy192+CPP+FAM, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each group after 5 days of treatment, and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids treated with water had high ratios (~134) of *Buchnera*/aphid DNA. In contrast, aphids treated with the scorpion peptide fused to the cell penetrating peptide had ~1.8-fold less *Buchnera*/aphid DNA copies after 5 days of treatment (Fig. 46). These data indicate that treatment with the scorpion peptide fused to a cell penetrating peptide decreased endosymbiont levels.

The scorpion peptide fused to a cell penetrating peptide freely entered the bacteriocytes to act against Buchnera

To test whether the cell penetrating peptide aids in the delivery of the scorpion peptide into the bacteriocytes directly, isolated bacteriocytes were directly exposed to the fusion protein and imaged. The bacteriocytes were dissected from the aphids in Schneider's medium supplemented with 1% w/v BSA (Schneider-BSA medium), and placed in an imaging well containing 20ul of schneider's medium. A 100ug lyophilized aliquot of the scorpion peptide was resuspended in 100ul of the schneider's medium to produce 1mg/ml solution, and 5ul of this solution was mixed in to the well containing bacteriocytes. After 30 min of incubation at room temperature, the bacteriocytes were thoroughly washed to eliminate any excess free peptide in the solution. Images of the bacteriocytes were captured before and after the incubation (Fig. 47). The fusion peptide penetrated the bacteriocyte membranes and was directly available to *Buchnera*.

Together, this data demonstrates the ability to kill and decrease the development, longevity, and endogenous bacterial populations, e.g., fitness, of aphids by treating them with the scorpion antimicrobial peptide Uy192 fused to a cell penetrating peptide.

Example 23: Insects treated with vitamin analogs

This Example demonstrates the treatment of aphids with the provitamin pantothenol which is the alcohol analog of pantothenic acid (Vitamin B5). Aphids have obligate endosymbiont bacteria, *Buchnera*, that help them make essential amino acids and vitamins, including Vitamin B5. A previous study has shown that pantothenol inhibits the growth of *Plasmodium falciparum* by inhibition of the specific parasite kinases (Saliba et al., 2005). It was hypothesized that treating insects with pantothenol would be detrimental for the bacterial-insect symbiosis therefore affecting insect fitness. This Example demonstrates that the treatment with pantothenol decreased insect fitness.

Therapeutic Design:

Pantothenol solutions were formulated depending on the delivery method:

1) In artificial diet through the plants: with 0 (negative control) or 10 or 100 uM pantothenol formulated in an artificial diet (based on Akey and Beck, 1971; see Experimental Design) without

essential amino acids (2 mg/ml histidine, 2 mg/ml isoleucine, 2 mg/ml leucine, 2 mg/ml lysine, 1 mg/ml methionine, 1.62 mg/ml phenylalanine, 2 mg/ml threonine, 1 mg/ml tryptophan, and 2 mg/ml valine).

2) Leaf coating: 100 μ l of 0.025% nonionic organosilicone surfactant solvent Silwet L-77 in water (negative control) or 100 μ l of 50 μ g/ml of rifampicin formulated in solvent solution was applied directly to the leaf surface and allowed to dry.

Plant Delivery Experimental Design

Aphids (eNASCO, *Acyrtosiphon pisum*) were grown on fava bean plants (*Vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60 $\pm$ 5% RH; 25 $\pm$ 2 $^{\circ}$ C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24 $^{\circ}$ C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first and second instar aphids were collected from healthy plants and divided into 3 different treatment groups: 1) artificial diet alone without essential amino acids, 2) artificial diet alone without essential amino acids and 10 μ M pantothenol, and 3) artificial diet alone without essential amino acids and 100 μ M pantothenol. Each treatment group received approximately the same number of individuals from each of the collection plants.

The artificial diet used was made as previously published (Akey and Beck, 1971 Continuous Rearing of the Pea Aphid, *Acyrtosiphon pisum*, on a Holidic Diet), with and without the essential amino acids (2 mg/ml histidine, 2 mg/ml isoleucine, 2 mg/ml leucine, 2 mg/ml lysine, 1 mg/ml methionine, 1.62 mg/ml phenylalanine, 2 mg/ml threonine, 1 mg/ml tryptophan, and 2 mg/ml valine), except neither diet included homoserine or beta-alanyltyrosine. The pH of the diets was adjusted to 7.5 with KOH and diets were filter sterilized through a 0.22 μ m filter and stored at 4 $^{\circ}$ C for short term (<7 days) or at -80 $^{\circ}$ C for long term.

Pantothenol (Sigma Cat# 295787) solutions were made at 10 μ M and 100 μ M in artificial diet without essential amino acids, sterilized by passing through a 0.22 μ m syringe filter, and stored at -20 $^{\circ}$ C. For treatments (see Therapeutic design), the appropriate amount of stock solution was added to the artificial diet without essential amino acids to obtain a final concentration of 10 or 100 μ M pantothenol. The diet was then placed into a 1.5 ml Eppendorf tube with a fava bean stem with a leaf and the opening of the tube was closed using parafilm. This artificial diet feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant.

For each treatment, 16-20 aphids were placed onto each leaf. Artificial diet feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish housing the artificial feeding system when they were discovered.

In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th instar) was determined daily throughout the experiment. Once an aphid reached the 4th instar stage, they were given their own artificial feeding system in a deep petri dish so that fecundity could be monitored once they reached adulthood.

For adult aphids, new nymphs were counted daily and then discarded. At the end of the experiments, fecundity was determined as the mean number of offspring produced daily once the aphid reached adulthood. Pictures of aphids were taken throughout the experiment to evaluate size differences between treatment groups.

5 After 8 days of treatment, DNA was extracted from multiple aphids from each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid
10 DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 240) and Buch_groES_98R (CTGTTCTCGAGTCGATTTCC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was
15 performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

20 *Vitamin analog treatment delays aphid development*

eNASCO 1st and 2nd instar aphids were divided into three separate treatment groups as defined in Plant Delivery Experimental Design (described herein). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with artificial diet alone without essential amino acids began reaching maturity (5th instar stage) at approximately 5 days (Fig. 48A).
25 Development was delayed in aphids treated with pantothenol, especially at days two and three post-treatment (Fig. 48A), indicating that treatment with pantothenol impairs aphid development. Eventually, most aphids from each treatment group reached maturity and began reproducing. In addition to monitoring developmental stage of aphids over time, aphids were also imaged and aphid area was determined. All aphids were the same size after 1 day of treatment, however, by 3 days post-treatment,
30 aphids treated with pantothenol were smaller in area than untreated controls. Moreover, aphids treated with pantothenol had much less of an increase in body size (as determined by area) over the course of the experiment, compared to aphids treated with artificial diet alone without essential amino acids (Fig. 48B).

35 *Vitamin analog treatment increased aphid mortality*

Survival rate of aphids was also measured during the treatments. Aphids reared on artificial diet alone without essential amino acids had higher survival rates compared to aphids treated with 10 or 100 μ M pantothenol (Fig. 49), indicating that pantothenol treatment negatively affected aphid fitness.

Treatment with pantothenol decreases aphid fecundity

Fecundity was also monitored in aphids during the treatments. The fraction of aphids surviving to maturity and reproducing was determined. Approximately one quarter of aphids treated with artificial diet without essential amino acids survived to reach maturity by 8 days post-treatment (Fig. 50A). In contrast, only a little over 1/10th of aphids treated with 10 or 100 uM pantothenol survived to reach maturity and reproduce by 8 days post-treatment. The mean day aphids in each treatment group began reproducing was also measured and for all treatment groups, the mean day aphids began reproducing was 7 days (Fig. 50B). Additionally, the mean number of offspring per day produced by mature, reproducing aphids was also monitored. Aphids treated with artificial diet alone without essential amino acids produced approximately 7 offspring/day. In contrast, aphids treated with 10 and 100 uM pantothenol only produced approximately 4 and 5 offspring/day, respectively, shown in Fig. 50C. Taken together, these data indicate that pantothenol treatment resulted in a loss of aphid reproduction.

Pantothenol treatment does not affect Buchnera in aphids

To test whether treatment with pantothenol, specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 8 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids treated with artificial diet alone without essential amino acids had high ratios of *Buchnera*/aphid DNA copies as did aphids treated with each of the two concentrations of pantothenol (Fig. 51). These data indicate that pantothenol treatment does not affect *Buchnera*/aphid DNA copy number directly.

Leaf Coating Delivery Experimental Design:

Pantothenol powder was added to 0.025% of a nonionic organosilicone surfactant solvent, Silwet L-77, to obtain a final concentration of 10 uM pantothenol. The treatment was filter sterilized using a 0.22 um filter and stored at 4 degrees C. Aphids (eNASCO strain, *Acyrtosiphon pisum*) were grown on fava bean plants as described in a previous Example. For experiments, first instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) negative control (solvent solution only) and 2) 10 uM pantothenol in solvent. 100 ul of the solution was absorbed onto a 2x2 cm piece of fava bean leaf.

Each treatment group received approximately the same number of individuals from each of the collection plant. For each treatment, 20 aphids were placed onto each leaf. Aphids were monitored daily for survival and dead aphids were removed when they were discovered. In addition, the developmental stage (1st, 2nd, 3rd, 4th, 5th instar, and 5R, representing a reproducing 5th instar) was determined daily throughout the experiment.

Pantothenol treatment delivered through leaf coating does not affect aphid development

eNASCO 1st instar aphids were divided into two separate treatment groups as defined in the Experimental Design described herein. Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids placed on coated leaves treated with either the control or

pantothenol solution matured at similar rates up to two days post-treatment (Fig. 52). These data indicate that leaf coating with pantothenol did not affect aphid development.

Pantothenol treatment delivered through leaf coating increased aphid mortality

5 Survival rate of aphids was also measured during the leaf coating treatments. Aphids placed on coated leaves with pantothenol had lower survival rates than aphids placed on coated leaves with the control solution (Fig. 53). These data indicate that pantothenol treatment delivered through leaf coating significantly ($p=0.0019$) affected aphid survival. All aphids died in this experiment and there were no remaining aphids left to extract DNA from and determine *Buchnera*/aphid DNA ratios.

10 Together this data described in the previous Examples demonstrate the ability to kill and decrease the development, reproductive ability, longevity, and endogenous bacterial populations, e.g., fitness, of aphids by treating them with pantothenol through multiple delivery methods.

Example 24: Insects treated with a cocktail of amino acid transporters inhibitors

15 This Example demonstrates the treatment of aphids with a cocktail of amino acid analogs. The objective of this treatment was to inhibit uptakes of glutamine into the bacteriocytes through the ApGLNT1 glutamine transporter. It has previously been shown that arginine inhibits glutamine uptake by the glutamine transporter (Price et al., 2014 PNAS), and we hypothesized that treatment with analogs of arginine, or other amino acid analogs, may also inhibit uptake of essential amino acids into the aphid bacteriocytes. This Example demonstrates that the decrease in fitness upon treatment was mediated through the modulation of bacterial populations endogenous to the insect that were sensitive to amino acid analogs. One targeted bacterial strain is *Buchnera*.

Therapeutic Design:

25 The amino acid cocktail was formulated for delivery through leaf perfusion and through the plant. This delivery method consisted of injecting leaves with approximately 1 ml of the amino acid cocktail in water (see below for list of components in the cocktail) or 1 ml of the negative control solution containing water only.

30 *Leaf perfusion and delivery through plants experimental design:*

Aphids LSR-1 (which harbor only *Buchnera*), *Acyrtosiphon pisum* were grown on fava bean plants (*Vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; $60\pm5\%$ RH; 25 ± 2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24°C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) negative control (water treatment) and 2) amino acid cocktail treatment. The amino acid cocktail contained each of the following agents at the indicated final concentrations: 330 μ M L-NNA (N-nitro-L-Arginine; Sigma), 0.1 mg/ml L-canavanine (Sigma), 0.5 mg/ml D-arginine (Sigma), 0.5 mg/ml D-phenylalanine (Sigma), 0.5 mg/ml D-

histidine (Sigma), 0.5 mg/ml D-tryptophan (Sigma), 0.5 mg/ml D-threonine (Sigma), 0.5 mg/ml D-valine (Sigma), 0.5 mg/ml D-methionine (Sigma), 0.5 mg/ml D-leucine, and 6 μ M L-NMMA (citrate) (Cayman Chemical). ~ 1 ml of the treatment solution was perfused into the fava bean leaf via injection and the stem of the plant was put into a 1.5 ml Eppendorf tube containing the treatment solution. The opening of the tube was closed using parafilm. This feeding system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant. For each treatment, a total of 56-58 aphids were placed onto each leaf (each treatment consisted of two replicates of 28-31 aphids). Each treatment group received approximately the same number of individuals from each of the collection plants. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered. The aphid developmental stage (1st, 2nd, 3rd, 4th, and 5th instar) was determined daily throughout the experiment and microscopic images were taken of the aphids on day 5 to determine aphid area measurements.

Stock solutions of L-NNA were made at 5 mM in water, sterilized by passing through a 0.22 μ m syringe filter, and stored at -20°C. Stock solutions of L-canavanine were made at 100 mg/ml in water, sterilized by passing through a 0.22 μ m syringe filter, and stored at 4°C. Stock solutions of D-arginine and D-threonine were made at 50 mg/ml in water, sterilized by passing through a 0.22 μ m syringe filter, and stored at 4°C. Stock solutions of D-valine and D-methionine were made at 25 mg/ml in water, sterilized by passing through a 0.22 μ m syringe filter, and stored at 4°C. Stock solutions of D-leucine were made at 12 mg/ml in water, sterilized by passing through a 0.22 μ m syringe filter, and stored at 4°C. Stock solutions of D-phenylalanine and D-histidine were made at 50 mg/ml in 1M HCl, sterilized by passing through a 0.22 μ m syringe filter, and stored at 4°C. Stock solutions of D-tryptophan were made at 50 mg/ml in 0.5M HCl, sterilized by passing through a 0.22 μ m syringe filter, and stored at 4°C. Stock solutions of L-NMMA were made at 6 mg/ml in sterile PBS, sterilized by passing through a 0.22 μ m syringe filter, and stored at -20°C. For treatments (see Therapeutic design), the appropriate amount of stock solution was added to water to obtain the final concentration of the agent in the cocktail as indicated above.

After 6 days of treatment, DNA was extracted from multiple aphids from each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 240) and Buch_groES_98R (CTGTTCTCGAGTCGATTTC; SEQ ID NO: 241) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 242) and ApEF1a 246R (TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 243) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15

seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

Treatment with a cocktail of amino acid analogs delayed and stopped progression of aphid development

5 LSR-1 1st instar aphids were divided into two separate treatment groups as defined in Leaf perfusion and delivery through plants experimental design (described herein). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with water began reaching maturity (5th instar stage) at day 5 post-treatment (Fig. 54A). By 6 days post-treatment, ~20 percent of aphids treated with water reached the 5th instar stage. In contrast, less than 3 percent of 10 the aphids treated with the amino acid cocktail reached the 5th instar stage, even after 6 days (Fig. 54A). This delay in development upon treatment with the amino acid cocktail was further exemplified by aphid size measurements taken at 5 days post-treatment. Aphids treated with water alone were approximately 0.45 mm², whereas aphids treated with the amino acid cocktail were approximately 0.33 mm² (Fig. 54B). These data indicate that treatment with the amino acid cocktail delayed aphid development, negatively 15 impacting aphid fitness.

Treatment with an amino acid analog cocktail resulted in decreased Buchnera in aphids

To test whether treatment with the amino acid analog cocktail specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each 20 treatment group after 6 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids placed on control solution had high ratios of *Buchnera*/aphid DNA copies. In contrast, aphids placed on AA cocktail treatment had a drastic reduction of *Buchnera*/aphid DNA copies (Fig. 55), indicating that the AA analog cocktail treatment eliminated endosymbiotic *Buchnera*.

Together, this data demonstrates the ability to decrease the development and endogenous 25 bacterial populations, e.g., fitness, of aphids by treating them with a cocktail of amino acid analogs.

Example 25: Insects treated with a combination of agents (antibiotic, peptide, and natural antimicrobial)

This Example demonstrates the treatment of insects with a combination of three antimicrobial 30 agents – an antibiotic (rifampicin), a peptide (the scorpion peptide Uy192), and a natural antimicrobial (low molecular weight chitosan). In other Examples, each of these agents administered individually resulted in decreased aphid fitness and reduced endosymbiont levels. This Example demonstrates that through the delivery of a combination of treatments, insect fitness and endosymbiont levels were reduced as well as, or better than, treatment with each individual agent alone.

35

Therapeutic Design

The combination treatment was formulated for delivery through leaf perfusion and through the plant. This delivery method consisted of injecting leaves with approximately 1 ml of the combination treatment in water (with final concentrations of 100 µg/ml rifampicin, 100 µg/ml Uy192, and 300 µg/ml 40 chitosan) or 1 ml of the negative control solution containing water only.

Leaf perfusion and delivery through plants experimental design

Aphids LSR-1 (which harbor only *Buchnera*), *Acyrtosiphon pisum* were grown on fava bean plants (*Vroma vicia faba* from Johnny's Selected Seeds) in a climate-controlled incubator (16 h light/8 h dark photoperiod; 60±5% RH; 25±2 °C). Prior to being used for aphid rearing, fava bean plants were grown in potting soil at 24°C with 16 h of light and 8 h of darkness. To limit maternal effects or health differences between plants, 5–10 adults from different plants were distributed among 10 two-week-old plants, and allowed to multiply to high density for 5–7 days. For experiments, first instar aphids were collected from healthy plants and divided into 2 different treatment groups: 1) negative control (water treatment) and 2) a combination of 100 µg/ml rifampicin, 100 µg/ml Uy192, and 300 µg/ml chitosan treatment. ~1 ml of the treatment solution was perfused into the fava bean leaf via injection and the stem of the plant was put into a 1.5 ml Eppendorf tube containing the treatment solution. The opening of the tube was closed using parafilm. This treatment system was then placed into a deep petri dish (Fisher Scientific, Cat# FB0875711) and aphids were applied to the leaves of the plant. For each treatment, a total of 56 aphids were placed onto each leaf (each treatment consisted of two replicates of 28 aphids). Each treatment group received approximately the same number of individuals from each of the collection plants. The feeding systems were changed every 2-3 days throughout the experiment. Aphids were monitored daily for survival and dead aphids were removed from the deep petri dish when they were discovered. The aphid developmental stage (1st, 2nd, 3rd, 4th, and 5th instar) was determined daily throughout the experiment and microscopic images were taken of the aphids on day 5 to determine aphid area measurements.

Rifampicin (Tokyo Chemical Industry, LTD) stock solution was made at 25 mg/ml in methanol, sterilized by passing through a 0.22 µm syringe filter, and stored at -20°C. For treatment, the appropriate amount of stock solution was added to water to obtain a final concentration of 100 µg/ml rifampicin. Uy192 was synthesized by Bio-Synthesis at >75% purity. 1 mg of lyophilized peptide was reconstituted in 500 µl of 80% acetonitrile, 20% water, and 0.1% TFA. 100 µl (100 µg) was aliquoted into 10 individual Eppendorf tubes and allowed to dry. For treatment, 1 ml of water was added to a 100 µg aliquot of peptide to obtain the final concentration of 100 µg/ml Uy192. Chitosan (Sigma, catalog number 448869-50G) stock solution was made at 1% in acetic acid, sterilized autoclaving, and stored at 4°C. For treatments the appropriate amount of stock solution was added to water to obtain the final concentration of 300 µg/ml chitosan.

After 6 days of treatment, DNA was extracted from multiple aphids from each treatment group. Briefly, the aphid body surface was sterilized by dipping the aphid into a 6% bleach solution for approximately 5 seconds. Aphids were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *Buchnera* and aphid DNA copy numbers were measured by qPCR. The primers used for *Buchnera* were Buch_groES_18F (CATGATCGTGTGCTTGTTAAG; SEQ ID NO: 228) and Buch_groES_98R (CTGTTCTCGAGTCGATTTC; SEQ ID NO: 229) (Chong and Moran, 2016 PNAS). The primers used for aphid were ApEF1a 107F (CTGATTGTGCCGTGCTTATTG; SEQ ID NO: 230) and ApEF1a 246R

(TATGGTGGTTCAGTAGAGTCC; SEQ ID NO: 231) (Chong and Moran, 2016 PNAS). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 55°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

Treatment with a combination of three antimicrobial agents delayed and stopped progression of aphid development

LSR-1 1st instar aphids were divided into two separate treatment groups as defined in *Leaf perfusion and delivery through plants experimental design* (described herein). Aphids were monitored daily and the number of aphids at each developmental stage was determined. Aphids treated with water began reaching maturity (5th instar stage) at day 5 post-treatment (Fig. 56A). By 6 days post-treatment, ~20 percent of aphids treated with water reached the 5th instar stage. In contrast, no aphids treated with the combination of three agents reached the 5th instar stage, even after 6 days (Fig. 56A). This delay in development upon combination treatment was further exemplified by aphid size measurements taken at 5 days post-treatment. Aphids treated with water alone were approximately 0.45 mm², whereas aphids treated with the 3-agent combination were approximately 0.26 mm² (Fig. 56B). These data indicate that treatment with a combination of agents delayed aphid development, negatively impacting aphid fitness.

Treatment with a combination of three antimicrobial agents increased aphid mortality

Survival was also monitored daily after treatment. At 2 days post-treatment, approximately 75 percent of aphids treated with water were alive, whereas only 62 percent of aphids treated with the combination of agents were alive. This trend of more aphids surviving treatment in the control (water-treated) group continued for the duration of the experiment. At 6 days post-treatment, 64 percent of control (water-treated) aphids survived, whereas 58 percent of aphids treated with a combination of rifampicin, Uy192, and chitosan survived (Fig. 57). These data indicate that the combination of treatments negatively affected aphid survival.

Treatment with a combination of three agents resulted in decreased Buchnera in aphids

To test whether treatment with a combination of a peptide, antibiotic, and natural antimicrobial specifically resulted in loss of *Buchnera* in aphids, and that this loss impacted aphid fitness, DNA was extracted from aphids in each treatment group after 6 days of treatment and qPCR was performed to determine the *Buchnera*/aphid copy numbers. Aphids treated with water alone ratios of approximately 2.3 *Buchnera*/aphid DNA (Fig. 58). In contrast, aphids treated with the combination of a peptide, antibiotic, and natural antimicrobial had approximately 2-fold lower ratios of *Buchnera*/aphid DNA (Fig. 58). These data indicate that combination treatment reduced endosymbiont levels, which resulted in decreased aphid fitness.

Together, this data demonstrates the ability to decrease the development and endogenous bacterial populations, e.g., fitness, of aphids by treating them with a combination of a peptide, antibiotic, and natural antimicrobial.

Example 26: Insects treated with an antibiotic solution

This Example demonstrates the effects of treatment of weevils with ciprofloxacin, a bactericidal antibiotic that inhibits the activity of DNA gyrase and topoisomerase, two enzymes essential for DNA replication. This Example demonstrates that the phenotypic effect of ciprofloxacin on another model insect, weevils, was mediated through the modulation of bacterial populations endogenous to the insects that were sensitive to ciprofloxacin. One targeted bacterial strain is *Sitophilus* primary endosymbiont (SPE, *Candidatus Sodalis pierantonius*).

Experimental Design:

Sitophilus maize weevils (*Sitophilus zeamais*) were reared on organic corn at 27.5°C and 70% relative humidity. Prior to being used for weevil rearing, corn was frozen for 7 days and then tempered to 10% humidity with sterile water. For experiments, adult male/female mating pairs were divided into 3 different treatment groups that were done in triplicate: 1) water control, 2) 250 µg/ml ciprofloxacin, and 3) 2.5 mg/ml ciprofloxacin. Ciprofloxacin (Sigma) stock solutions were made at 25 mg/ml in 0.1N HCl, sterilized by passing through a 0.22 µm syringe filter, and stored at -20°C. For treatments, the appropriate amount of stock solution was diluted in sterile water.

The weevils were subjected to three successive treatments:

1. The first treatment included soaking 25 g of corn with each of the three treatment groups listed above: 1) water control, 2) 250 µg/ml ciprofloxacin, and 3) 2.5 mg/ml ciprofloxacin. Briefly, 25 g of corn was placed into a 50ml conical tube and each of the treatment was added to fill the tube completely. The tube was put on a shaker for 1.5 hours after which, the corn was removed and placed into a deep petri dish and air dried. Male/Female mating pairs were then added to each treatment group and allowed to feed for 4 days.
2. After 4 days, mating pairs were removed and subjected to a second treatment by putting them onto 25 g of new corn treated with 1) water control, 2) 250 µg/ml ciprofloxacin, and 3) 2.5 mg/ml ciprofloxacin. Mating pairs fed and laid eggs on this corn for 7 days. The corn from the second treatment was assessed for the emergence of offspring (see *assessment of offspring*, below)
3. Mating pairs were subjected to a final treatment which included a combination of submerging them into the treatment (1) water control, 2) 250 µg/ml ciprofloxacin, and 3) 2.5 mg/ml ciprofloxacin for 5 seconds and then placing them in a vial with 10 corn kernels that had been coated with 1 ml of 1) water control, 2) 250 µg/ml ciprofloxacin, and 3) 2.5 mg/ml ciprofloxacin.

Weevil survival was monitored daily for 18 days, after which DNA was extracted from the remaining weevils in each group. Briefly, the weevil body was surface sterilized by dipping the weevil into a 6% bleach solution for approximately 5 seconds. Weevils were then rinsed in sterile water and DNA was extracted from each individual aphid using a DNA extraction kit (Qiagen, DNeasy kit) according to manufacturer's instructions. DNA concentration was measured using a nanodrop nucleic acid quantification, and *SPE* and weevil DNA copy numbers were measured by qPCR. The primers used for

SPE were qPCR_Sod_F (ATAGCTGTCCAGACGCTTCG; SEQ ID NO: 244) and qPCR_Sod_R (ATGTCGTCGAGGCGATTACC; SEQ ID NO: 245). The primers used for weevil (β -actin) were SACT144_FOR (GGTGTGGCGTACAAGTCCT; SEQ ID NO: 246) and SACT314_REV (GAATTGCCTGATGGACAGGT; SEQ ID NO: 247) (Login et al., 2011). qPCR was performed using a qPCR amplification ramp of 1.6 degrees C/s and the following conditions: 1) 95°C for 10 minutes, 2) 95°C for 15 seconds, 3) 57°C for 30 seconds, 4) repeat steps 2-3 40x, 5) 95°C for 15 seconds, 6) 55°C for 1 minute, 7) ramp change to 0.15 degrees C/s, 8) 95°C for 1 second. qPCR data was analyzed using analytic (Thermo Fisher Scientific, QuantStudio Design and Analysis) software.

10 *Assessment of offspring:*

After 25 days, one replicate of the corn kernels from the second treatment of the adult mating pairs was dissected (see *Experimental Design*, above) to check for the presence of any developing larvae, pupae, or adult weevils. Most of the development of *Sitophilus weevils* takes place within the grain/rice/corn and adults emerge from the kernels once their development is complete. Corn kernels were gently dissected open with a scalpel and any developing weevils were collected and the percent of adults, pupae, and larvae were determined. The weevils from the dissection were then surface sterilized and the levels of SPE were determined by qPCR. Corn kernels from the remaining two replicates of each of the groups from the second treatment were not dissected but checked daily for the emergence of adult weevils.

20

Assessment of antibiotic penetration into corn

25 mg of corn kernels was placed into a 50 ml conical tube and water or 2.5 mg/ml or 0.25 mg/ml ciprofloxacin in water was added to fill the tube. The kernels were soaked for 1.5 hours as described herein. After soaking, kernels were air dried and assayed to determine whether the antibiotic was able to coat and penetrate the kernel. To test this, a concentrated sample of *Escherichia coli* DH5 α in water was spread onto 5 Luria Broth (LB) plates. To each plate the following was done, 1) a corn kernel soaked in water was added, 2) an entire corn kernel that had been soaked with 2.5 or 0.25 mg/ml ciprofloxacin was added, and 3) a half of corn kernel that had been soaked with 2.5 or 0.25 mg/ml ciprofloxacin was added and placed inside down on the plate. The plates were incubated overnight at 37 degrees C and bacterial growth and/or zone(s) of inhibition were assessed the next day.

30

Soaking corn kernels in antibiotics allowed antibiotics to coat the surface and penetrate corn kernels.

To test whether ciprofloxacin could coat the surface of a corn kernel after a kernel, corn kernels were soaked in water without antibiotics or water with 2.5 or 0.25 mg/ml ciprofloxacin (as described above). A concentrated culture of *E. coli* was then spread onto LB plates and one of the coated kernels was then placed onto the center of the plate. The plates were incubated overnight, and bacterial growth was assessed the next day.

35

A lawn of bacteria grew on the entire plate with the corn kernel that had been coated in water without any antibiotics (Fig. 56A). In contrast, no bacteria grew on plates with entire corn kernels that had been soaked in either of the two concentrations of ciprofloxacin (Fig. 56B, left panels). These data show

40

that the coating method employed in these experiments allowed for ciprofloxacin to successfully coat the surface of corn kernels and inhibit bacterial growth.

To test whether ciprofloxacin could penetrate the corn kernel, corn kernels soaked in 2.5 or 0.25 mg/ml ciprofloxacin were cut in half and placed cut side down on an LB plate with a concentrated culture of *E. coli*. The plates were incubated overnight and the next day bacterial growth was assessed. No bacterial growth was present on the plates with the kernels soaked in either concentration of antibiotic, indicating that ciprofloxacin penetrated the corn kernel (Fig. 56B, right panels). Together, these data indicate that the method of corn kernel soaking used for these experiments successfully coated and penetrated the kernels with the antibiotic.

Antibiotic treatment decreases SPE levels in the F0 generation.

S. zeamais mating pairs were divided into three separate treatment groups as defined in *Experimental Design* (above). Weevils were monitored daily and all weevils remained alive for the course of the experiment. After 18 days of treatment, weevils were surface sterilized, genomic DNA was extracted, and SPE levels were measured by qPCR. Weevils treated with water only had approximately 4 and 8-fold higher amounts of SPE compared to weevils treated with 250 ug/ml and 2.5 mg/ml ciprofloxacin, respectively (Fig. 57). These data indicate that treatment of weevils with ciprofloxacin resulted in decreased levels of SPE.

Antibiotic treatment delays the development and decreases the SPE levels of the F1 generation of weevils.

The development of the F1 generation of weevils was assessed by dissecting corn kernels that F0 mating pairs had oviposited on for 7 days and were subsequently removed. After 25 days, 12 offspring were found in water/control-treated corn with the majority (~67%) of offspring being in the pupae form (Fig. 58A). 13 and 20 offspring were found in weevils treated with 250 ug/ml and 2.5 mg/ml ciprofloxacin, respectively. Interestingly, weevils treated with antibiotic showed a delay in development compared to control treated weevils with the majority (38 and 65% for 250 ug/ml and 2.5 mg/ml ciprofloxacin, respectively) of the offspring being in the larval form (Fig. 58A).

Genomic DNA was extracted from weevils dissected from the corn kernels and qPCR was performed to measure the levels of SPE. Water treated F1 weevils had approximately 4-fold higher levels of SPE compared to weevils treated with 2.5 mg/ml ciprofloxacin (Fig. 58B). These data indicate that treatment with ciprofloxacin reduced the levels of the SPE in weevils which led to a delay in development.

Antibiotic treatment decreased weevil reproduction

The number of weevils that emerged over the course of 43 days after the initial mating pairs were removed from the second treatment was used as a measure for the fecundity (Fig. 59A and 59B). The first weevil emerged on day 29, and the total number of weevils that emerged till day 43 were counted. While weevils treated with water and 250 ug/ml had similar amount of F1 offspring, there were much less offspring that emerged from the 2.5 mg/ml treatment group, indicating that antibiotic treatment decreased SPE levels affected weevil fecundity.

Together with the previous Examples, this data demonstrate the ability to kill and decrease the development, reproductive ability, longevity and endogenous bacterial populations, e.g., fitness, of weevils by treating them with an antibiotic through multiple delivery methods.

5 **Example 27: Mites treated with an antibiotic solution**

This Example demonstrates the ability to kill, decrease the fitness of two-spotted spider mites by treating them with rifampicin, a narrow spectrum antibiotic that inhibits DNA-dependent RNA synthesis by inhibiting a bacterial RNA polymerase, and doxycycline, a broad-spectrum antibiotic that prevents bacterial reproduction by inhibiting protein synthesis. The effect of rifampicin and doxycycline on mites was mediated through the modulation of bacterial populations endogenous to the mites that were sensitive to the antibiotics.

Insects, such as mosquitoes, and arachnids, such as ticks, can function as vectors for pathogens causing severe diseases in humans and animals such as Lyme disease, dengue, trypanosomiasis, and malaria. Vector-borne diseases cause millions of human deaths every year. Also, vector-borne diseases that infect animals, such as livestock, represent a major global public health burden. Thus, there is a need for methods and compositions to control insects and arachnids that carry vector-borne diseases. Two-spotted spider mites are arachnids in the same subclass as ticks. Therefore, this Example demonstrates methods and compositions used to decrease the fitness of two-spotted spider mites and provide insight into decreasing tick fitness.

20

Therapeutic design

Two treatments were used for these experiments 1) 0.025% Silwet L-77 (negative control) or 2) a cocktail of antibiotics containing 250 µg/ml rifampicin and 500 µg/ml doxycycline. Rifampicin (Tokyo Chemical Industry, LTD) stock solutions were made at 25 mg/ml in methanol, sterilized by passing through a 0.22 µm syringe filter, and stored at -20 °C. Doxycycline (manufacturer) stock solutions were made at 50 mg/mL in water, sterilized by passing through a 0.22 µm syringe filter, and stored at -20 °C.

25

Experimental Design:

This assay tested an antibiotic solution on two-spotted spider mites and determined how their fitness was altered by targeting endogenous microbes.

30

Kidney plants were grown in potting soil at 24 °C with 16 h of light and 8 h of darkness. Mites were reared on kidney bean plants at 26 °C and 15-20% relative humidity. For treatments, one-inch diameter leaf disks were cut from kidney bean leaves and sprayed with either 0.025% Silwet L-77 (negative control) or the antibiotic cocktail (250 µg/ml rifampicin and 500 µg/ml doxycycline in 0.025% Silwet L-77) using a Master Airbrush Brand Compressor Model C-16-B Black Mini Airbrush Air Compressor. The compressor was cleaned with ethanol before, after, and between treatments. The liquid was feed through the compressor using a quarter inch tube. A new tube was used for each treatment.

35

After leaf discs dried, four of each treatment were placed in a cup on top of a wet cotton ball covered with a piece of kimwipe. Each treatment setup was done in duplicate. 25 adult female mites

40

were then placed in the cup. On day 4, the females were removed from the cup and the eggs and larvae were left on the leaf discs.

On day 11, mites at the protonymph stage and the deutonymph stage were taken from the cups and placed in their own tube so survival could be measured. Each tube contained a moist cotton ball
5 covered with a piece of kimwipe with a half inch leaf disc treated with the negative control or the cocktail.

The mites were observed under a dissecting microscope daily after feeding on a leaf treated with the antibiotic or the control solutions, and classified according to the following categories:

- Alive: they walked around when on their legs or moved after being poked by a paint brush.
- 10 • Dead: immobile and did not react to stimulation from a paint brush

A sterile paint brush was used to stimulate the mites by touching their legs. Mites classified as dead were kept throughout the assay and rechecked for movement daily. The assays were carried out at 26 °C and 15-20% relative humidity.

15 *Antibiotic treatment increased mite mortality*

The survival rates of the two-spotted spider mites treated with the antibiotic cocktail were compared to the mites treated with the negative control. The survival rates of the mites treated with the cocktail were decreased compared to the control (Fig. 60).

This data demonstrates the ability to decrease fitness of mites by treating them with a solution of
20 antibiotics.

Example 28: Insects treated with a solution of purified phage

This Example demonstrates the isolation and purification of phages from environmental samples that targeted specific insect bacteria. This Example also demonstrates the efficacy of isolated phages against the target bacteria in vitro by plaque assays, by measuring their oxygen consumption rate, and
25 the extracellular acidification rate. Finally, this Example demonstrates the efficacy of the phages in vivo, by measuring the ability of the phage to the target bacteria from flies by treating them with a phage isolated against the bacteria. This Example demonstrates that a pathogenic bacterium that decreased the fitness of an insect can be cleared using a phage to target the bacteria. Specifically, *Serratia marcescens* which is a pathogenic bacterium in flies can be cleared with the use of a phage that was isolated from
30 garden compost.

Experimental design

Isolation of specific bacteriophages from natural samples:

Bacteriophages against target bacteria were isolated from environmental source material. Briefly,
35 a saturated culture of *Serratia marcescens* was diluted into fresh double-strength tryptic soy broth (TSB) and grown for ~120 minutes to early log-phase at 24-26°C, or into double-strength Luria-Bertani (LB) broth and grown for ~90 min at 37°C. Garden compost was prepared by homogenization in PBS and sterilized by 0.2 µm filtration. Raw sewage was sterilized by 0.2 µm filtration. One volume of filtered source material was added to log-phase bacterial cultures and incubation was continued for 24 h.

Enriched source material was prepared by pelleting cultures and filtering supernatant fluid through 0.45 µm membranes.

Phages were isolated by plating samples onto double-agar bacterial lawns. Stationary bacterial cultures were combined with molten 0.6% agar LB or TSB and poured onto 1.5% agar LB or TSB plates.

- 5 After solidification, 2.5 µL of phage sample dilutions were spotted onto the double-agar plates and allowed to absorb. Plates were then wrapped and incubated overnight at 25°C (TSA) or 37°C (LB), then assessed for the formation of visible plaques. Newly isolated plaques were purified by serial passaging of individual plaques on the target strain by picking plaques into SM Buffer (50 mM Tris-HCl [pH 7.4], 10 mM MgSO₄, 100 mM NaCl) and incubating for 15 min at 55°C, then repeating the double-agar spotting
- 10 method from above using the plaque suspension.

Bacteriophages were successfully isolated from both sewage and compost, as detailed above. Plaque formation was clearly evident after spotting samples onto lawns of the *S. marcescens* bacteria used for the enrichments.

15 *Passaging, quantification, and propagation of bacteriophages:*

- Propagation and generation of phage lysates for use in subsequent experiments was performed using bacteriophages isolated and purified as above. Briefly, saturated bacterial cultures were diluted 100-fold into fresh medium and grown for 60-120 minutes to achieve an early-logarithmic growth state for effective phage infection. Phage suspensions or lysates were added to early log phase cultures and
- 20 incubation was continued until broth clearing, indicative of phage propagation and bacterial lysis, was observed, or until up to 24 h post-infection. Lysates were harvested by pelleting cells at 7,197 x g for 20 min, then filtering the supernatant fluid through 0.45 or 0.2 µm membranes. Filtered lysates were stored at 4°C.

- Enumeration of infective phage particles was performed using the double-agar spotting method. Briefly, a 1:10 dilution series of samples was performed in PBS and dilutions were spotted onto solidified double-agar plates prepared with the host bacteria as above. Plaque-forming units (PFU) were counted after overnight incubation to determine the approximate titer of samples.
- 25

In vitro analysis of isolated phages measuring bacterial respiration:

- 30 A Seahorse XFe96 Analyzer (Agilent) was used to measure the effects of phages on bacteria by monitoring oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) during infection. XFe96 plates were coated the day prior to experiments by 15 µL of a 1 mg/mL poly-L-lysine stock per well and dried overnight at 28°C and XFe96 probes were equilibrated by placing into wells containing 200 µL of XF Calibrant and incubating in the dark at room temperature. The following day, poly-L-lysine coated
- 35 plates were washed twice with ddH₂O. Saturated overnight cultures of *E. coli* BL21 (LB, 37°C) or *S. marcescens* (TSB, 25°C) were subcultured at 1:100 into the same media and grown with aeration for ~2.5 h at 30°C. Cultures were then diluted to O.D._{600nm} ~ 0.02 using the same media. Treatments were prepared by diluting stocks into SM Buffer at 10x final concentration and loading 20 µL of the 10x solutions into the appropriate injection ports of the probe plate. While the probes were equilibrating in the
- 40 XFe96 Flux Analyzer, bacterial plates were prepared by adding 90 µL of bacterial suspensions or media

controls and spun at 3,000 rpm for 10 min. Following centrifugation, an additional 90 μ L of the appropriate media were added gently to the wells so as not to disturb bacterial adherence, bringing the total volume to 180 μ L per well.

The XFe96 Flux Analyzer was run at $\sim 30^{\circ}\text{C}$, following a Mix, Wait, Read cycling of 1:00, 0:30, 5 3:00. Four cycles were completed to permit equilibration/normalization of bacteria, then the 20 μ L treatments were injected and cycling continued as above, for a total time of approximately 6 h. Data were analyzed using the Seahorse XFe96 Wave software package.

The effects of isolated bacteriophages were assayed by measuring oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) of bacteria with a Seahorse XFe96 Analyzer. When *E.* 10 *coli* was infected with phage T7 and *S. marcescens* infected with the newly isolated Φ SmVL-C1, dramatic decreases in OCR were observed following brief bursts in this rate (Fig. 64). For both phages with both host organisms, the Seahorse assay permitted the detection of successful phage infection without the need for plaque assays. Thus, this method is applicable for detecting phage infection of a host organism not amenable to traditional phage detection methods.

15

SYBR Gold transduction assay for infection identification:

Bacteriophage preparations were prepared for staining by pretreating with nucleases to remove 20 extraviral nucleic acids that could interfere with fluorescent signal interpretation. Briefly, MgCl_2 was added to 10 mL of phage lysate at 10 mM final concentration, and RNase A (Qiagen) and DNase I (Sigma) were both added to final concentrations of 10 $\mu\text{g/mL}$. Samples were incubated for 1 h at room temperature. After nuclease treatment, 5 mL of lysates were combined with 1 μ L of SYBR Gold (Thermo, 10,000x) and incubated at room temperature for ~ 1.5 h. Excess dye was subsequently removed from 25 samples using Amicon ultrafiltration columns. Briefly, Amicon columns (15 mL, 10k MWCO) were washed by adding 10 mL of SM Buffer and spinning at 5,000 $\times$ g, 4°C for 5 min. Labeled phage samples were then spun through the columns at 5,000 $\times$ g, 4°C until the volume had decreased by approximately 10-fold (15-30 min). To wash samples, 5 mL SM Buffer was added to each reservoir and the spin 30 repeated, followed by two additional washes. After the third wash, the retained samples were pipetted out from the Amicon reservoirs and brought up to approximately 1 mL using SM Buffer. To remove larger contaminants, washed and labeled phage samples were spun at 10,000 $\times$ g for 2 min, and the supernatants were subsequently filtered through 0.2 μm membranes into black microtubes and stored at 4°C .

Saturated bacterial cultures (*E. coli* MG1655 grown in LB at 37°C , *S. marcescens* and *S. symbiotica* grown in TSB at 26°C) were prepared by spinning down 1 mL aliquots and washing once with 1 mL PBS before a final resuspension using 1 mL PBS. Positive control labeled bacteria were stained by 35 combining 500 μ L of washed bacteria with 1 μ L of SYBR Gold and incubating for 1 h in the dark at room temperature. Bacteria were pelleted by spinning at 8,000 $\times$ g for 5 min and washed twice with an equal volume of PBS, followed by resuspension in a final volume of 500 μ L PBS. A volume of 25 μ L of stained bacteria was combined with 25 μ L of SM Buffer in a black microtube, to which 50 μ L of 10% formalin (5% final volume, $\sim 2\%$ formaldehyde) was added and mixed by flicking. Samples were fixed at room 40 temperature for ~ 3 h and then washed using Amicon ultrafiltration columns. Briefly, 500 μ L of picopure

water was added to Amicon columns (0.5 mL, 100k MWCO) and spun at 14,000 x g for 5 min to wash membranes. Fixed samples were diluted by adding 400 µL of PBS and then transferred to pre-washed spin columns and spun at 14,000 x g for 10 min. Columns were transferred to fresh collection tubes, and 500 µL of PBS was added to dilute out fixative remaining in the retentate. Subsequently, two additional
5 PBS dilutions were performed, for a total of three washes. The final retentates were diluted to roughly 100 µL, then columns were inverted into fresh collection tubes and spun at 1,000 x g for 2 min to collect samples. Washed samples were transferred to black microtubes and stored at 4°C.

For transduction experiments and controls, 25 µL of bacteria (or PBS) and 25 µL of SYBR Gold labeled phage (or SM Buffer) were combined in black microtubes and incubated static for 15-20 min at
10 room temperature to permit phage adsorption and injection into recipient bacteria. Immediately after incubation, 50 µL of 10% formalin was added to samples and fixation was performed at room temperature for ~4 h. Samples were washed with PBS using Amicon columns, as above.

Injection of bacteriophage nucleic acid was required for a phage to successfully infect a host bacterial cell. Coliphage P1kc labeled with SYBR Gold and co-incubated with *S. marcescens* revealed
15 the presence of fluorescent bacteria by microscopy, validating the use of this assay in a phage isolation pipeline. As with the Seahorse assay, this approach provided an alternative to traditional phage methods to permit expansion to organisms not amenable to plaque assay. Additionally, the SYBR Gold transduction assay did not require bacterial growth, so is applicable to analysis of phages targeting difficult or even non-culturable organisms, including endosymbionts such as *Buchnera*.

20

Testing in vivo efficacy of the phages against S. marcescens in Drosophila melanogaster flies

S. marcescens cultures were grown in Tryptic Soy Broth (TSB) at 30°C with constant shaking at 200rpm.

The media used to rear fly stocks was cornmeal, molasses and yeast medium (11 g/l yeast, 54 g/l
25 yellow cornmeal, 5 g/l agar, 66 ml/l molasses, and 4.8 ml/l propionic acid). All the components of the diet except propionic acid were heated together to 80°C in deionized water with constant mixing for 30 minutes and let to cool to 60°C. Propionic acid was then mixed in and 50ml of the diet was aliquoted into individual bottles and allowed to cool down and solidify. The flies were raised at 26°C, 16:8 hour light:dark cycle, at around 60% humidity.

30 To infect the flies with *S. marcescens*, a fine needle (About 10µm wide tip) was dipped in a dense overnight stationary phase culture and the thorax of the flies was punctured. For this experiment, four replicates of 10 males and 10 females each were infected with *S. marcescens* using the needle puncturing method as the positive control for fly mortality. For the treatment group, four replicates of 10 males and 10 females each were pricked with *S. marcescens* and a phage solution containing about 10⁸
35 phage particles/ml. Finally, two replicates of 10 males and 10 females each that were not pricked or treated in anyway were used as a negative control for mortality.

Flies in all conditions were placed in food bottles and incubated at 26°C, 16:8 light:dark cycle, at 60% humidity. The number of alive and dead flies were counted every day for four days after the pricking. All The flies pricked with *S. marcescens* alone were all dead within 24 hours of the treatment.
40 In comparison, more than 60% of the flies in the phage treatment group, and all the flies in the untreated

control group were alive at that time point (Fig. 65). Further, most of the flies in the phage treatment group and the negative control group went on to survive for four more days when the experiment was terminated.

To ascertain the reason of death of the flies, dead flies from both the *S. marcescens* and *S. marcescens* + phage pricked flies were homogenized and plated out. Four dead flies from each of the four replicates of both the *S. marcescens* and the *S. marcescens* + phage treatment were homogenized in 100ul of TSB. A 1:100 dilution was also produced by diluting the homogenate in TSB. 10ul of the concentrated homogenate as well as the 1:100 dilution was plated out onto TSA plates, and incubated overnight at 30 °C. Upon inspection of the plates for bacteria growth, all the plates from the dead *S. marcescens* pricked flies had a lawn of bacteria growing on them, whereas the plates from the dead *S. marcescens* + phage pricked flies had no bacteria on them. This shows that in the absence of the phage, *S. marcescens* likely induced septic shock in the flies leading to their fatality. However, in the presence of the phage, the mortality may have been due to injury caused by the pricking with the needle.

OTHER EMBODIMENTS

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, the descriptions and examples should not be construed as limiting the scope of the invention. The disclosures of all patent and scientific literature cited herein are expressly incorporated in their entirety by reference.

CLAIMS

1. A method of decreasing fitness of a vector for an animal pathogen, the method comprising:
delivering an antimicrobial peptide having at least 90% sequence identity with one or more of the following: cecropin (SEQ ID NO: 82), melittin, copsin, drosomycin (SEQ ID NO: 93), dermcidin (SEQ ID NO: 81), andropin (SEQ ID NO: 83), moricin (SEQ ID NO: 84), ceratotoxin (SEQ ID NO: 85), abaecin (SEQ ID NO: 86), apidaecin (SEQ ID NO: 87), prophenin (SEQ ID NO: 88), indolicidin (SEQ ID NO: 89), protegrin (SEQ ID NO: 90), tachyplesin (SEQ ID NO: 91), or defensin (SEQ ID NO: 92) to the vector.
2. The method of claim 1, wherein the delivery comprises delivering the antimicrobial peptide to at least one habitat where the vector grows, lives, reproduces, feeds, or infests.
3. The method of any one of claims 1-2, wherein the antimicrobial peptide is delivered in an insect comestible composition for ingestion by the vector.
4. The method of any one of claims 1-3, wherein the antimicrobial peptide is formulated as a liquid, a solid, an aerosol, a paste, a gel, or a gas composition.
5. The method of any one of claims 1-4, wherein the insect is at least one of a mosquito, midge, louse, sandfly, tick, triatomine bug, tsetse fly, or flea.
6. A composition comprising an antimicrobial peptide having at least 90% sequence identity with one or more of the following: cecropin, melittin, copsin, drosomycin, dermcidin, andropin, moricin, ceratotoxin, abaecin, apidaecin, prophenin, indolicidin, protegrin, tachyplesin, or defensin formulated for targeting a microorganism in a vector for an animal pathogen.
7. The composition of claim 6, wherein the antimicrobial peptide is at a concentration of about 0.1 ng/g to about 100 mg/g in the composition.
8. The composition of any one of claims 6-7, wherein the antimicrobial peptide further comprises a targeting domain.
9. The composition of any one of claims 6-8, wherein the antimicrobial peptide further comprises a cell penetrating peptide.