

ANTI-CLAUDIN 18.2 AND ANTI-4-1BB BISPECIFIC ANTIBODIES AND USES THEREOF

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of PCT Application Serial No. PCT/CN2019/100162, filed August 12, 2019, No. PCT/CN2019/104508, filed September 5, 2019, No. PCT/CN2020/071954, filed January 14, 2020, and No. PCT/CN2020/087968, filed April 30, 2020, the content of each which is hereby incorporated by reference in its entirety.

BACKGROUND

[0002] Claudins are a family of proteins that form the important components of the tight cell junctions. Claudin-18 splice variant 2 (CLDN18.2) is a gastric-specific membrane protein. In the healthy tissue, CLDN18.2 is restrictly expressed in the short-lived differentiated cells of gastric mucosa as a component of tight junction with limited accessibility of antibody treatment. However, it was ectopically expressed at significant levels in a variety of primary lesion and metastases of epithelial tumor entities, including gastric, pancreatic, esophageal, and lung adenocarcinoma cells.

[0003] 4-1BB (CD137, tumor necrosis factor receptor superfamily 9) is a member of TNF-receptor superfamily (TNFRSF) and is a costimulatory molecule which is expressed following the activation of immune cells, both innate and adaptive immune cells. 4-1BB plays an important role in modulating the activity of various immune cells. 4-1BB agonists enhance immune cell proliferation, survival, secretion of cytokines and cytolytic activity CD8 T cells. Many other studies showed that activation of 4-1BB enhances immune response to eliminate tumors in mice. Therefore, it was suggested that 4-1BB is a promising target molecule in cancer immunology.

SUMMARY

[0004] Provided are bispecific and multi-specific antibodies that target both claudin 18.2 (CLDN18.2) and 4-1BB. In some embodiments, the antibodies of the present technology include a “conditional agonist” anti-4-1BB portion which, without the anti-CLDN18.2 portion binding to CLDN18.2 proteins expressed on a cell, cannot activate 4-1BB signaling.

[0005] 4-1BB signaling activation is the expected mechanism for agonist antibodies, such as utomilumab (PF-05082566) and urelumab (BMS-663513). The anti-4-1BB portions of the presently disclosed antibodies, however, do not require such an activity. Actually, it is preferred that the anti-4-1BB portions of the present antibodies are not capable of independently activating 4-1BB in the absence of CLDN18.2 binding. As the experimental examples demonstrated, interestingly, when the anti-CLDN18.2 portion binds to CLDN18.2 proteins on a cell, such CLDN18.2 binding can trigger 4-1BB signaling activation.

[0006] Compared to the known anti-4-1BB agonist antibodies which are commonly associated with dose-limiting on-target toxicities, the antibodies of the present disclosure are contemplated to be much safer. In a tissue, such as liver, wherein CLDN18.2 is not expressed, the present antibodies are not expected to trigger cytotoxic immune response as they cannot activate 4-1BB signaling. In a tumor tissue wherein CLDN18.2 is expressed and/or accessible, by contrast, the present antibodies can initiate potent immune response to the tumor cells. Accordingly, unlike those anti-4-1BB antibodies currently being developed clinically which suffer on-target/inherent toxicities, the presently disclosed antibodies can be potent and safe at the same time in treating cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

[0007] FIG. 1A-C illustrate three different bispecific formats tested in the present disclosure.

[0008] FIG. 2A-D present ELISA results for 4-1BB and CLDN18.2 binding.

[0009] FIG. 3 shows the cell-based 4-1BB binding.

[0010] FIG. 4A-C show the binding results for CLDN18.2.

[0011] FIG. 5A-D show CLDN18.2-dependent 4-1BB signaling activation.

[0012] FIG. 6A-E show PBMC response in the presence of CLDN18.2 expressing cells.

[0013] FIG. 7A-D show CD8⁺ T response in the presence of CLDN18.2 expressing cells.

[0014] FIG. 8A-B show *in vivo* efficacy of anti-CLDN18.2-4-1BB antibody in syngeneic mouse model and *ex vivo* analysis of its impact on tumor infiltrating lymphocytes.

[0015] FIG.9A-C show *in vivo* efficacy of C-1A10 bispecific antibody in syngeneic mouse model.

[0016] FIG.10 shows dose-dependent anti-tumor efficacy of C-1A10 and D-1A10 in syngeneic mouse model.

[0017] FIG.11A-D show pharmacokinetics and pharmacodynamics relation of D-1A10 in syngeneic mouse model.

DETAILED DESCRIPTION

Definitions

[0018] It is to be noted that the term “a” or “an” entity refers to one or more of that entity; for example, “an antibody,” is understood to represent one or more antibodies. As such, the terms “a” (or “an”), “one or more,” and “at least one” can be used interchangeably herein.

[0019] As used herein, the term “polypeptide” is intended to encompass a singular “polypeptide” as well as plural “polypeptides,” and refers to a molecule composed of monomers (amino acids) linearly linked by amide bonds (also known as peptide bonds). The term “polypeptide” refers to any chain or chains of two or more amino acids, and does not refer to a specific length of the product. Thus, peptides, dipeptides, tripeptides, oligopeptides, “protein,” “amino acid chain,” or any other term used to refer to a chain or chains of two or more amino acids, are included within the definition of “polypeptide,” and the term “polypeptide” may be used instead of, or interchangeably with any of these terms. The term “polypeptide” is also intended to refer to the products of post-expression modifications of the polypeptide, including without limitation glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, or modification by non- naturally occurring amino acids. A polypeptide may be derived from a natural biological source or produced by recombinant technology, but is not necessarily translated from a designated nucleic acid sequence. It may be generated in any manner, including by chemical synthesis.

[0020] The term “isolated” as used herein with respect to cells, nucleic acids, such as DNA or RNA, refers to molecules separated from other DNAs or RNAs, respectively, that are present in the natural source of the macromolecule. The term “isolated” as used herein also

refers to a nucleic acid or peptide that is substantially free of cellular material, viral material, or culture medium when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. Moreover, an “isolated nucleic acid” is meant to include nucleic acid fragments which are not naturally occurring as fragments and would not be found in the natural state. The term “isolated” is also used herein to refer to cells or polypeptides which are isolated from other cellular proteins or tissues. Isolated polypeptides is meant to encompass both purified and recombinant polypeptides.

[0021] As used herein, the term “recombinant” as it pertains to polypeptides or polynucleotides intends a form of the polypeptide or polynucleotide that does not exist naturally, a non-limiting example of which can be created by combining polynucleotides or polypeptides that would not normally occur together.

[0022] “Homology” or “identity” or “similarity” refers to sequence similarity between two peptides or between two nucleic acid molecules. Homology can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequence is occupied by the same base or amino acid, then the molecules are homologous at that position. A degree of homology between sequences is a function of the number of matching or homologous positions shared by the sequences. An “unrelated” or “non-homologous” sequence shares less than 40% identity, though preferably less than 25% identity, with one of the sequences of the present disclosure.

[0023] A polynucleotide or polynucleotide region (or a polypeptide or polypeptide region) has a certain percentage (for example, 60 %, 65 %, 70 %, 75 %, 80 %, 85 %, 90 %, 95 %, 98 % or 99 %) of “sequence identity” to another sequence means that, when aligned, that percentage of bases (or amino acids) are the same in comparing the two sequences. This alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example those described in Ausubel *et al.* eds. (2007) *Current Protocols in Molecular Biology*. Preferably, default parameters are used for alignment. One alignment program is BLAST, using default parameters. In particular, programs are BLASTN and BLASTP, using the following default parameters: Genetic code = standard; filter = none; strand = both; cutoff = 60; expect = 10; Matrix = BLOSUM62; Descriptions = 50 sequences; sort by = HIGH SCORE; Databases = non-redundant, GenBank + EMBL + DDBJ + PDB + GenBank CDS translations + SwissProtein + SPupdate + PIR. Biologically

equivalent polynucleotides are those having the above-noted specified percent homology and encoding a polypeptide having the same or similar biological activity.

[0024] As used herein, an “antibody” or “antigen-binding polypeptide” refers to a polypeptide or a polypeptide complex that specifically recognizes and binds to an antigen. An antibody can be a whole antibody and any antigen binding fragment or a single chain thereof. Thus the term “antibody” includes any protein or peptide containing molecule that comprises at least a portion of an immunoglobulin molecule having biological activity of binding to the antigen. Examples of such include, but are not limited to a complementarity determining region (CDR) of a heavy or light chain or a ligand binding portion thereof, a heavy chain or light chain variable region, a heavy chain or light chain constant region, a framework (FR) region, or any portion thereof, or at least one portion of a binding protein.

[0025] The terms “antibody fragment” or “antigen-binding fragment”, as used herein, is a portion of an antibody such as F(ab')₂, F(ab)₂, Fab', Fab, Fv, scFv and the like. Regardless of structure, an antibody fragment binds with the same antigen that is recognized by the intact antibody. The term “antibody fragment” includes aptamers, spiegelmers, and diabodies. The term “antibody fragment” also includes any synthetic or genetically engineered protein that acts like an antibody by binding to a specific antigen to form a complex.

[0026] A “single-chain variable fragment” or “scFv” refers to a fusion protein of the variable regions of the heavy (V_H) and light chains (V_L) of immunoglobulins. In some aspects, the regions are connected with a short linker peptide of ten to about 25 amino acids. The linker can be rich in glycine for flexibility, as well as serine or threonine for solubility, and can either connect the N-terminus of the V_H with the C-terminus of the V_L, or vice versa. This protein retains the specificity of the original immunoglobulin, despite removal of the constant regions and the introduction of the linker. ScFv molecules are known in the art and are described, *e.g.*, in US patent 5,892,019.

[0027] The term antibody encompasses various broad classes of polypeptides that can be distinguished biochemically. Those skilled in the art will appreciate that heavy chains are classified as gamma, mu, alpha, delta, or epsilon (γ , μ , α , δ , ϵ) with some subclasses among them (*e.g.*, γ 1- γ 4). It is the nature of this chain that determines the “class” of the antibody as IgG, IgM, IgA IgG, or IgE, respectively. The immunoglobulin subclasses (isotypes) *e.g.*, IgG₁, IgG₂, IgG₃, IgG₄, IgG₅, etc. are well characterized and are known to confer functional

specialization. Modified versions of each of these classes and isotypes are readily discernable to the skilled artisan in view of the instant disclosure and, accordingly, are within the scope of the instant disclosure. All immunoglobulin classes are clearly within the scope of the present disclosure, the following discussion will generally be directed to the IgG class of immunoglobulin molecules. With regard to IgG, a standard immunoglobulin molecule comprises two identical light chain polypeptides of molecular weight approximately 23,000 Daltons, and two identical heavy chain polypeptides of molecular weight 53,000-70,000. The four chains are typically joined by disulfide bonds in a “Y” configuration wherein the light chains bracket the heavy chains starting at the mouth of the “Y” and continuing through the variable region.

[0028] Antibodies, antigen-binding polypeptides, variants, or derivatives thereof of the disclosure include, but are not limited to, polyclonal, monoclonal, multispecific, human, humanized, primatized, or chimeric antibodies, single chain antibodies, epitope-binding fragments, *e.g.*, Fab, Fab' and F(ab')₂, Fd, Fvs, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv), fragments comprising either a VK or VH domain, fragments produced by a Fab expression library, and anti- idiotypic (anti-Id) antibodies (including, *e.g.*, anti-Id antibodies to LIGHT antibodies disclosed herein). Immunoglobulin or antibody molecules of the disclosure can be of any type (*e.g.*, IgG, IgE, IgM, IgD, IgA, and IgY), class (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule.

[0029] Light chains are classified as either kappa or lambda (K, λ). Each heavy chain class may be bound with either a kappa or lambda light chain. In general, the light and heavy chains are covalently bonded to each other, and the “tail” portions of the two heavy chains are bonded to each other by covalent disulfide linkages or non-covalent linkages when the immunoglobulins are generated either by hybridomas, B cells or genetically engineered host cells. In the heavy chain, the amino acid sequences run from an N-terminus at the forked ends of the Y configuration to the C-terminus at the bottom of each chain.

[0030] Both the light and heavy chains are divided into regions of structural and functional homology. The terms “constant” and “variable” are used functionally. In this regard, it will be appreciated that the variable domains of both the light (VK) and heavy (VH) chain portions determine antigen recognition and specificity. Conversely, the constant domains of the light chain (CK) and the heavy chain (CH1, CH2 or CH3) confer important biological

properties such as secretion, transplacental mobility, Fc receptor binding, complement binding, and the like. By convention the numbering of the constant region domains increases as they become more distal from the antigen-binding site or amino- terminus of the antibody. The N-terminal portion is a variable region and at the C-terminal portion is a constant region; the CH3 and CK domains actually comprise the carboxy-terminus of the heavy and light chain, respectively.

[0031] As indicated above, the variable region allows the antibody to selectively recognize and specifically bind epitopes on antigens. That is, the VK domain and VH domain, or subset of the complementarity determining regions (CDRs), of an antibody combine to form the variable region that defines a three dimensional antigen-binding site. This quaternary antibody structure forms the antigen-binding site present at the end of each arm of the Y. More specifically, the antigen-binding site is defined by three CDRs on each of the VH and VK chains (*i.e.* CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2 and CDR-L3). In some instances, *e.g.*, certain immunoglobulin molecules derived from camelid species or engineered based on camelid immunoglobulins, a complete immunoglobulin molecule may consist of heavy chains only, with no light chains. *See, e.g.*, Hamers-Casterman *et al.*, *Nature* 363:446-448 (1993).

[0032] In naturally occurring antibodies, the six “complementarity determining regions” or “CDRs” present in each antigen-binding domain are short, non-contiguous sequences of amino acids that are specifically positioned to form the antigen-binding domain as the antibody assumes its three dimensional configuration in an aqueous environment. The remainder of the amino acids in the antigen-binding domains, referred to as “framework” regions, show less inter-molecular variability. The framework regions largely adopt a β -sheet conformation and the CDRs form loops which connect, and in some cases form part of, the β -sheet structure. Thus, framework regions act to form a scaffold that provides for positioning the CDRs in correct orientation by inter-chain, non-covalent interactions. The antigen-binding domain formed by the positioned CDRs defines a surface complementary to the epitope on the immunoreactive antigen. This complementary surface promotes the non-covalent binding of the antibody to its cognate epitope. The amino acids comprising the CDRs and the framework regions, respectively, can be readily identified for any given heavy or light chain variable region by one of ordinary skill in the art, since they have been precisely defined (*see* “Sequences of Proteins of Immunological Interest,” Kabat, E., *et al.*,

U.S. Department of Health and Human Services, (1983); and Chothia and Lesk, *J. Mol. Biol.*, 196:901-917 (1987)).

[0033] In the case where there are two or more definitions of a term which is used and/or accepted within the art, the definition of the term as used herein is intended to include all such meanings unless explicitly stated to the contrary. A specific example is the use of the term “complementarity determining region” (“CDR”) to describe the non-contiguous antigen combining sites found within the variable region of both heavy and light chain polypeptides. This particular region has been described by Kabat *et al.*, U.S. Dept. of Health and Human Services, “Sequences of Proteins of Immunological Interest” (1983) and by Chothia *et al.*, *J. Mol. Biol.* 196:901-917 (1987), which are incorporated herein by reference in their entireties. The CDR definitions according to Kabat and Chothia include overlapping or subsets of amino acid residues when compared against each other. Nevertheless, application of either definition to refer to a CDR of an antibody or variants thereof is intended to be within the scope of the term as defined and used herein. The appropriate amino acid residues which encompass the CDRs as defined by each of the above cited references are set forth in the table below as a comparison. The exact residue numbers which encompass a particular CDR will vary depending on the sequence and size of the CDR. Those skilled in the art can routinely determine which residues comprise a particular CDR given the variable region amino acid sequence of the antibody.

	Kabat	Chothia
CDR-H1	31-35	26-32
CDR-H2	50-65	52-58
CDR-H3	95-102	95-102
CDR-L1	24-34	26-32
CDR-L2	50-56	50-52
CDR-L3	89-97	91-96

[0034] Kabat *et al.* also defined a numbering system for variable domain sequences that is applicable to any antibody. One of ordinary skill in the art can unambiguously assign this

system of “Kabat numbering” to any variable domain sequence, without reliance on any experimental data beyond the sequence itself. As used herein, “Kabat numbering” refers to the numbering system set forth by Kabat *et al.*, U.S. Dept. of Health and Human Services, “Sequence of Proteins of Immunological Interest” (1983).

[0035] In addition to table above, the Kabat number system describes the CDR regions as follows: CDR-H1 begins at approximately amino acid 31 (*i.e.*, approximately 9 residues after the first cysteine residue), includes approximately 5-7 amino acids, and ends at the next tryptophan residue. CDR-H2 begins at the fifteenth residue after the end of CDR-H1, includes approximately 16-19 amino acids, and ends at the next arginine or lysine residue. CDR-H3 begins at approximately the thirty third amino acid residue after the end of CDR-H2; includes 3-25 amino acids; and ends at the sequence W-G-X-G, where X is any amino acid. CDR-L1 begins at approximately residue 24 (*i.e.*, following a cysteine residue); includes approximately 10-17 residues; and ends at the next tryptophan residue. CDR-L2 begins at approximately the sixteenth residue after the end of CDR-L1 and includes approximately 7 residues. CDR-L3 begins at approximately the thirty third residue after the end of CDR-L2 (*i.e.*, following a cysteine residue); includes approximately 7-11 residues and ends at the sequence F or W-G-X-G, where X is any amino acid.

[0036] Antibodies disclosed herein may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine, donkey, rabbit, goat, guinea pig, camel, llama, horse, or chicken antibodies. In another embodiment, the variable region may be chondrichthoid in origin (*e.g.*, from sharks).

[0037] As used herein, the term “heavy chain constant region” includes amino acid sequences derived from an immunoglobulin heavy chain. A polypeptide comprising a heavy chain constant region comprises at least one of: a CH1 domain, a hinge (*e.g.*, upper, middle, and/or lower hinge region) domain, a CH2 domain, a CH3 domain, or a variant or fragment thereof. For example, an antigen-binding polypeptide for use in the disclosure may comprise a polypeptide chain comprising a CH1 domain; a polypeptide chain comprising a CH1 domain, at least a portion of a hinge domain, and a CH2 domain; a polypeptide chain comprising a CH1 domain and a CH3 domain; a polypeptide chain comprising a CH1 domain, at least a portion of a hinge domain, and a CH3 domain, or a polypeptide chain comprising a CH1 domain, at least a portion of a hinge domain, a CH2 domain, and a CH3 domain. In another embodiment, a polypeptide of the disclosure comprises a polypeptide

chain comprising a CH3 domain. Further, an antibody for use in the disclosure may lack at least a portion of a CH2 domain (*e.g.*, all or part of a CH2 domain). As set forth above, it will be understood by one of ordinary skill in the art that the heavy chain constant region may be modified such that they vary in amino acid sequence from the naturally occurring immunoglobulin molecule.

[0038] The heavy chain constant region of an antibody disclosed herein may be derived from different immunoglobulin molecules. For example, a heavy chain constant region of a polypeptide may comprise a CH1 domain derived from an IgG₁ molecule and a hinge region derived from an IgG₃ molecule. In another example, a heavy chain constant region can comprise a hinge region derived, in part, from an IgG₁ molecule and, in part, from an IgG₃ molecule. In another example, a heavy chain portion can comprise a chimeric hinge derived, in part, from an IgG₁ molecule and, in part, from an IgG₄ molecule.

[0039] As used herein, the term “light chain constant region” includes amino acid sequences derived from antibody light chain. Preferably, the light chain constant region comprises at least one of a constant kappa domain or constant lambda domain.

[0040] A “light chain-heavy chain pair” refers to the collection of a light chain and heavy chain that can form a dimer through a disulfide bond between the CL domain of the light chain and the CH1 domain of the heavy chain.

[0041] As previously indicated, the subunit structures and three dimensional configuration of the constant regions of the various immunoglobulin classes are well known. As used herein, the term “VH domain” includes the amino terminal variable domain of an immunoglobulin heavy chain and the term “CH1 domain” includes the first (most amino terminal) constant region domain of an immunoglobulin heavy chain. The CH1 domain is adjacent to the VH domain and is amino terminal to the hinge region of an immunoglobulin heavy chain molecule.

[0042] As used herein the term “CH2 domain” includes the portion of a heavy chain molecule that extends, *e.g.*, from about residue 244 to residue 360 of an antibody using conventional numbering schemes (residues 244 to 360, Kabat numbering system; and residues 231-340, EU numbering system; *see* Kabat *et al.*, U.S. Dept. of Health and Human Services, “Sequences of Proteins of Immunological Interest” (1983). The CH2 domain is unique in that it is not closely paired with another domain. Rather, two N-linked branched

carbohydrate chains are interposed between the two CH2 domains of an intact native IgG molecule. It is also well documented that the CH3 domain extends from the CH2 domain to the C-terminal of the IgG molecule and comprises approximately 108 residues.

[0043] As used herein, the term “hinge region” includes the portion of a heavy chain molecule that joins the CH1 domain to the CH2 domain. This hinge region comprises approximately 25 residues and is flexible, thus allowing the two N-terminal antigen-binding regions to move independently. Hinge regions can be subdivided into three distinct domains: upper, middle, and lower hinge domains (Roux *et al.*, *J. Immunol* 161:4083 (1998)).

[0044] By “specifically binds” or “has specificity to,” it is generally meant that an antibody binds to an epitope via its antigen-binding domain, and that the binding entails some complementarity between the antigen-binding domain and the epitope. According to this definition, an antibody is said to “specifically bind” to an epitope when it binds to that epitope, via its antigen-binding domain more readily than it would bind to a random, unrelated epitope. The term “specificity” is used herein to qualify the relative affinity by which a certain antibody binds to a certain epitope. For example, antibody “A” may be deemed to have a higher specificity for a given epitope than antibody “B,” or antibody “A” may be said to bind to epitope “C” with a higher specificity than it has for related epitope “D.”

[0045] As used herein, the terms “treat” or “treatment” refer to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) an undesired physiological change or disorder, such as the progression of cancer. Beneficial or desired clinical results include, but are not limited to, alleviation of symptoms, diminishment of extent of disease, stabilized (*i.e.*, not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. “Treatment” can also mean prolonging survival as compared to expected survival if not receiving treatment. Those in need of treatment include those already with the condition or disorder as well as those prone to have the condition or disorder or those in which the condition or disorder is to be prevented.

[0046] By “subject” or “individual” or “animal” or “patient” or “mammal,” is meant any subject, particularly a mammalian subject, for whom diagnosis, prognosis, or therapy is

desired. Mammalian subjects include humans, domestic animals, farm animals, and zoo, sport, or pet animals such as dogs, cats, guinea pigs, rabbits, rats, mice, horses, cattle, cows, and so on.

[0047] As used herein, phrases such as “to a patient in need of treatment” or “a subject in need of treatment” includes subjects, such as mammalian subjects, that would benefit from administration of an antibody or composition of the present disclosure used, *e.g.*, for detection, for a diagnostic procedure and/or for treatment.

Anti-Claudin 18.2 Anti-4-1BB Antibodies

[0048] 4-1BB is an inducible costimulatory receptor expressed on activated T and natural killer (NK) cells. 4-1BB trimer clustering by 4-1BB ligand (41BBL) trimer on T cells triggers a signaling cascade that results in upregulation of antiapoptotic molecules, cytokine secretion, and enhanced effector function. On NK cells, 4-1BB signaling can increase antibody-dependent cell-mediated cytotoxicity. Agonistic monoclonal antibodies targeting 4-1BB have been developed to harness 4-1BB signaling for cancer immunotherapy. Preclinical results in a variety of induced and spontaneous tumor models suggest that targeting 4-1BB with agonist antibodies can lead to tumor clearance and durable antitumor immunity.

[0049] Two agonist antibodies, urelumab and utomilumab, are currently undergoing clinical trials. Urelumab has strong efficacy but showed inflammatory liver toxicities. The liver toxicity appears to be on-target, making it difficult to be separated from efficacy. Utomilumab is relatively safer than urelumab, but is also less effective.

[0050] The present experimental example tested a couple of anti-4-1BB antibodies which were specifically selected for their incapability to activate 4-1BB signaling independently. In their monospecific forms, they can bind to 4-1BB alone or 4-1BB on the cell surface. Their binding of 4-1BB on the cell surface, however, does not lead to 4-1BB signaling activation (see, *e.g.*, **Example 3** and **FIG. 5**).

[0051] Interestingly, when the binding fragments of one of these conditional agonist 4-1BB antibodies, 1A10, was incorporated into a bispecific antibody that further has an anti-CLDN18.2 portion, the resulting bispecific antibody was able to efficiently activate 4-1BB signaling in a CLDN18.2-binding-dependent manner (see **Example 3** and **FIG. 5**). It is worth noting that these tested bispecific antibodies employed an N297A IgG1 Fc to disable FcγR-

medicated 4-1BB agonism. Therefore, the 4-1BB signaling activation can only be attributed to the CLDN18.2 binding.

[0052] The on-target toxicity of the anti-4-1BB agonist antibody urelumab is attributed to the antibody' lack of selectivity in 4-1BB agonism. The antibodies of the present technology, however, can be readily recognized for their ability to overcome this limitation. In a tissue, such as liver, where CLDN18.2 is not expressed or accessible, the antibodies would be safe as they cannot activate 4-1BB-mediated cytotoxicity. In a tumor tissue wherein CLDN18.2 is overexpressed or accessible, by contrast, the antibodies undergo CLDN18.2 binding-dependent 4-1BB signaling activation, leading to 4-1BB-mediated immune cell activation, thereby treating the tumor.

[0053] In accordance with one embodiment of the present disclosure, therefore, provided is an antibody that includes an anti-claudin 18.2 (CLDN18.2) unit having binding specificity to a CLDN18.2 protein, and an anti-4-1BB unit having binding specificity to a 4-1BB protein. In a preferred embodiment, the anti-4-1BB unit is incapable of activating 4-1BB signaling upon binding to a 4-1BB protein, in the absence of the anti-CLDN18.2 unit binding to a CLDN18.2 protein.

[0054] The lack of 4-1BB agonism of the anti-4-1BB portion can be achieved by many different means. It is suggested that 4-1BB clustering on the cell surface is required for its activation. In some embodiments, therefore, the binding of the anti-4-1BB unit to 4-1BB proteins on a cell does not result in clustering of the 4-1BB proteins, in the absence of the anti-CLDN18.2 unit binding to a CLDN18.2 protein.

[0055] The 4-1BB protein has four extracellular cysteine-rich pseudo repeats (CRD) domains, CRD1, CRD2, CRD3 and CRD4 (see the amino acid sequence and the CRD regions in the table below). Urelumab binds to the N-terminal CDR1, and activates 4-1BB clustering in a 4-1BB ligand (4-1BBL)-dependently manner. In some embodiments, the anti-4-1BB unit of the presently disclosed antibodies does not bind to CDR1. In some embodiments, the anti-4-1BB unit of the presently disclosed antibodies binds to CDR2. In some embodiments, the anti-4-1BB unit of the presently disclosed antibodies binds to CDR3. In some embodiments, the anti-4-1BB unit of the presently disclosed antibodies binds to CDR4.

	Sequence and CRD Annotation (SEQ ID NO:39)
Human 4-1BB (NP_001552.2)	1 <u>MGN</u> SCYNIVA TLLLVLNFER TR <u>SLQDPCSN CPAGTFCDNN RNQICSPCPP</u> Signal peptide CRD1
	51 <u>NSFSSAGGQR TCDICRQCKG VFRTRKECSS TSNAECDCTP GFHCLGAGCS</u> CRD2 CRD3
	101 <u>MCEQDCKQGQ ELTKKGCKDC CFGTFNDQKR GICRPWTNCS LDGKSVLVNG</u> CRD4
	151 <u>TKERDVVCGP</u> SPADLSPGAS SVTPPAPARE PGHSPQIISF FLALTSTALL
	201 FLLFFLTLLRF SVVKRGRKKL LYIFKQPFMR PVQTTQEEDG CSCRFPEEEE
	251 GGCEL

[0056] In some instances, 4-1BB clustering may be mediated through the effector function of an anti-4-1BB antibody. In some embodiments, therefore, an antibody of the present disclosure has an Fc fragment that has reduced or no effector function. Such effector functions, in some embodiments, antibody-dependent cell-mediated cytotoxicity (ADCC), complement-dependent cytotoxicity (CDC), or antibody-dependent cellular phagocytosis (ADCP).

[0057] The effector function of an antibody can be altered using techniques known in the art. In some embodiments, the Fc fragment of the antibody is mutated, or manufactured, in a manner to reduce or eliminate its binding to FcγR. In some embodiments, the Fc fragment has reduced or no binding to FcγRI (CD64); in some embodiments, the Fc fragment has reduced or no binding to FcγRIIA (CD32); in some embodiments, the Fc fragment has reduced or no binding to FcγRIIB (CD32); in some embodiments, the Fc fragment has reduced or no binding to FcγRIIIA (CD16a); and in some embodiments, the Fc fragment has reduced or no binding to FcγRIIIB (CD16b). In some embodiments, the Fc fragment has reduced or no binding to C1q (first subcomponent of the C1 complex).

[0058] In some embodiments, the Fc fragment includes one or more mutations that reduce or eliminate the FcγR or C1q binding of the Fc fragment. Non-limiting examples of such mutations include the L235E mutation in an IgG1 Fc fragment, the L234A and/or L235A mutation in an IgG1 Fc fragment, the P329G or P329A mutation in an IgG1 Fc fragment, the F234A and/or L235A or L235E mutation in an IgG4 Fc fragment, the H268Q, V309L, A330S, and/or P331S mutation in an IgG2 Fc fragment, and the V234A, G237A, P238S, H268A, V309L, A330S, and/or P331S mutation in an IgG2 Fc fragment (EU numbering).

[0059] The effector function of an antibody can also be reduced by decreasing or inhibiting glycosylation of the Fc fragment (e.g., an aglycosylated Fc fragment). Such decrease or inhibition can be achieved, in some embodiments, by using a cell line that is incapable to glycosylate the antibody. In some embodiments, the Fc fragment is mutated.

[0060] Non-limiting examples of such mutations include a mutation at N297 (such as N297A, N297G, and N297Q) (EU numbering). In some embodiments, the mutation is N297A.

Antibody Formats and Example Sequences

[0061] In the experimental examples, three different bispecific antibody formats were tested. Among them, the format of **FIG. 1A** exhibited higher activity than those of **FIG. 1B** and **FIG. 1C**. These data suggest that a suitable format for the antibodies of the present disclosure may employ a Fab fragment for the anti-CLDN18.2 portion. In some embodiments, the format includes a single chain fragment (scFv) for the anti-4-1BB portion. In some embodiments, the anti-4-1BB portion is not located between the Fab and a Fc fragment.

[0062] In some embodiments, the anti-CLDN18.2 unit include a Fab fragment, and preferably a pair of the Fab fragment. In some embodiments, the anti-4-1BB unit includes a Fab fragment, preferably a pair of the Fab fragments. In some embodiments, the anti-4-1BB unit includes a scFv, preferably a pair of the scFv fragments.

[0063] In some embodiments, the anti-CLDN18.2 unit is located at the N-terminal side of the anti-4-1BB unit. In some embodiments, a Fc fragment is placed between the anti-CLDN18.2 unit and the anti-4-1BB unit.

[0064] Example CDR sequence and VH/VL sequences are also provided for the anti-CLDN18.2 unit and the anti-4-1BB unit. In some embodiments, the anti-4-1BB unit comprises a heavy chain variable region (VH) comprising a CDRH1, a CDRH2, and a CDRH3, and a light chain variable region (VL) comprising a CDRL1, a CDRL2 and a CDRL3, wherein: the CDRH1 comprises the amino acid sequence of SEQ ID NO:1, or an amino acid sequence having one or two amino acid substitution from SEQ ID NO:1; the CDRH2 comprises the amino acid sequence of SEQ ID NO:2, or an amino acid sequence

having one or two amino acid substitution from SEQ ID NO:2; the CDRH3 comprises the amino acid sequence of SEQ ID NO: 3, 56, 57, 58, or 59, or an amino acid sequence having one or two amino acid substitution from SEQ ID NO: 3, 56, 57, 58, or 59; the CDRL1 comprises the amino acid sequence of SEQ ID NO:4 or 60, or an amino acid sequence having one or two amino acid substitution from SEQ ID NO:4 or 60; the CDRL2 comprises the amino acid sequence of SEQ ID NO:5 or 61, or an amino acid sequence having one or two amino acid substitution from SEQ ID NO:5 or 61; and the CDRL3 comprises the amino acid sequence of SEQ ID NO:6 or 62, or an amino acid sequence having one or two amino acid substitution from SEQ ID NO:6 or 62.

Table A. CDR Sequences of an Example Anti-4-1BB Unit

	Sequence	SEQ ID NO:
CDRH1	SYDMS	1
CDRH2	WISYSGGSIYYADSVKG	2
CDRH3	DAQNSMREFDY, DGQRNSMREFDY, DAQRQSMREFDY, DGQRQSMREFDY, or DAQRNAMREFDY	3 56 57 58 59
CDRL1	SGSSSNIGNNYVT	4
CDRL2	ADSHRPS	5
CDRL3	ATWDYSLSGYV	6

	Sequence	SEQ ID NO:
CDRH1	SYDMS	1
CDRH2	WISYSGGSIYYADSVKG	2
CDRH3	DAQNSMREFDY, DGQRNSMREFDY, DAQRQSMREFDY, DGQRQSMREFDY, or DAQRNAMREFDY	3 56 57 58 59
CDRL1	GYDMS	60
CDRL2	VIYPDDGNIYYADSVKG	61
CDRL3	HGGQKPTTKSSSAYGMDG	62

[0065] In some embodiments, the CDRH1 comprises the amino acid sequence of SEQ ID NO:1; the CDRH2 comprises the amino acid sequence of SEQ ID NO:2; the CDRH3 comprises the amino acid sequence of SEQ ID NO: 3, 56, 57, 58, or 59; the CDRL1 comprises the amino acid sequence of SEQ ID NO:4 or 60; the CDRL2 comprises the amino acid sequence of SEQ ID NO:5 or 61; and the CDRL3 comprises the amino acid sequence of SEQ ID NO:6 or 62.

[0066] In some embodiments, the anti-4-1BB unit comprises a heavy chain variable region (VH) comprising an amino acid sequence selected from the group consisting of SEQ ID NO:24, 46-51 and 63-69 and a light chain variable region (VL) comprising an amino acid sequence selected from the group consisting of SEQ ID NO:25, 52-53 and 70-74.

Table B. VH/VL Sequences of an Example Anti-4-1BB Unit

1A10 VH	EVQLLES ~~GGGLVQPGGSLRL~~ SCAASGFTFS ~~SYDMS~~ WVRQAPGKCLEWVS ~~WISYS~~ ~~GGSIYYADSVKGR~~ FTISRDN SKNTLYLQMNSLRAEDTAVYYCARD ~~DAQRNSMREF~~ ~~DYWGQGT~~ LVTVSS (SEQ ID NO:24)
1A10 VL	QSVLTQPPSASGTPGQRTISC ~~SGSSSNIGNNYVT~~ WYQQLPGTAPKLLIY ~~ADSH~~ ~~RPS~~ GVPDRFSGSKSGTSASLAISGLRSEDEADYYC ~~ATWDYSLSGYV~~ FGCGTKLT VL (SEQ ID NO:25)

[0067] In some embodiments, the anti-CLDN18.2 unit comprises a heavy chain variable region (VH) comprising a CDRH1, a CDRH2, and a CDRH3, and a light chain variable region (VL) comprising a CDRL1, a CDRL2 and a CDRL3, wherein (a) the CDRH1 comprises the amino acid sequence of SEQ ID NO:7; the CDRH2 comprises the amino acid sequence of SEQ ID NO:8; the CDRH3 comprises the amino acid sequence of SEQ ID NO:9; the CDRL1 comprises the amino acid sequence of SEQ ID NO:10; the CDRL2 comprises the amino acid sequence of SEQ ID NO:11; and the CDRL3 comprises the amino acid sequence of SEQ ID NO:12.

[0068] In some embodiments, the anti-CLDN18.2 unit comprises a heavy chain variable region (VH) comprising a CDRH1, a CDRH2, and a CDRH3, and a light chain variable region (VL) comprising a CDRL1, a CDRL2 and a CDRL3, wherein (b) the CDRH1 comprises the amino acid sequence of SEQ ID NO:13; the CDRH2 comprises the amino acid sequence of SEQ ID NO:14; the CDRH3 comprises the amino acid sequence of SEQ ID NO:15; the CDRL1 comprises the amino acid sequence of SEQ ID NO:16; the CDRL2

comprises the amino acid sequence of SEQ ID NO:17; and the CDRL3 comprises the amino acid sequence of SEQ ID NO:18.

[0069] In some embodiments, the anti-CLDN18.2 unit comprises a heavy chain variable region (VH) comprising a CDRH1, a CDRH2, and a CDRH3, and a light chain variable region (VL) comprising a CDRL1, a CDRL2 and a CDRL3, wherein (c) the CDRH1 comprises the amino acid sequence of SEQ ID NO:19; the CDRH2 comprises the amino acid sequence of SEQ ID NO:20; the CDRH3 comprises the amino acid sequence of SEQ ID NO:21; the CDRL1 comprises the amino acid sequence of SEQ ID NO:22; the CDRL2 comprises the amino acid sequence of SEQ ID NO:11; and the CDRL3 comprises the amino acid sequence of SEQ ID NO:23.

Table C. CDR Sequences of Example Anti-CLDN18.2 Units

		Sequence	SEQ ID NO:
4F11E2	CDRH1	TFGMH	7
	CDRH2	YITSGESPIYFTDTVKG	8
	CDRH3	SSYYGNSMDY	9
	CDRL1	RSSQSLNAGNRKNYLT	10
	CDRL2	WASTRES	11
	CDRL3	QNAYSYFPT	12
72C1B6A3	CDRH1	TYPIE	13
	CDRH2	NFHPYNDDTKYNEKFKG	14
	CDRH3	RAYGYPYAMDY	15
	CDRL1	KSSQSLNAGNQKNYLT	16
	CDRL2	RASSRES	17
	CDRL3	QNDYIYPYT	18
120B7B2	CDRH1	GYIIQ	19
	CDRH2	FINPYNDDTKYNEQFKG	20
	CDRH3	AYFGNAFAY	21
	CDRL1	KSSQSLNAGNQKNYLT	22
	CDRL2	WASTRES	11
	CDRL3	QNAYYFPFT	23

[0070] In some embodiments, the VH of the anti-CLDN18.2 unit comprises an amino acid sequence selected from the group consisting of SEQ ID NO:26, 28 and 30, and the VH of the

anti-CLDN18.2 unit comprises an amino acid sequence selected from the group consisting of SEQ ID NO:27, 29 and 31.

[0071] In some embodiments, provided is a bispecific antibody adopting a format like illustrated in FIG. 1A. In some embodiments, provided is an antibody comprising two first polypeptides and two second polypeptides, wherein each first polypeptide comprises, from N- to C-terminus, a heavy chain variable region (VH), a CH1, a CH2, a CH3, and a single chain fragment (scFv) having specificity to a 4-1BB protein; wherein each second polypeptide comprises a light chain variable region (VL) and a CL; wherein each VH is paired with one of the VL and have specificity to a claudin 18.2 (CLDN18.2) protein; and wherein the scFv is incapable of activating 4-1BB signaling upon binding to a 4-1BB protein, in the absence of the VH/VL pairs binding to a CLDN18.2 protein.

[0072] The VH/VL pair here constitutes an anti-CLDN18.2 unit and the scFv constitutes an anti-4-1BB unit. The CH2-CH3/CH2-CH3 pair constitutes an Fc fragment. The various embodiments above relating to the anti-CLDN18.2 unit, the anti-4-1BB unit, and the Fc fragments may be applicable here as well.

[0073] The following table provides non-limiting examples of the first polypeptide (heavy component) and the second polypeptide (light component). In some embodiments, each of the first polypeptides comprises the amino acid sequence of SEQ ID NO:40 and each of the second polypeptides comprises the amino acid sequence of SEQ ID NO:41.

[0074] In some embodiments, each of the first polypeptides comprises the amino acid sequence of SEQ ID NO:42 and each of the second polypeptides comprises the amino acid sequence of SEQ ID NO:43.

[0075] In some embodiments, each of the first polypeptides comprises the amino acid sequence of SEQ ID NO:44 and each of the second polypeptides comprises the amino acid sequence of SEQ ID NO:45.

Table D. Peptide Chains of Example Bispecific Antibodies

C-1A10		
Heavy component	Heavy chain of 4F11E2	EVQLVESGGGLVQPGGSLRLSCAASGFTFS TFGMH WVRQAPGKGLEWVS YITSGESPIYFTDITVKG RFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR SSYYGNSMDY WGQGTLLTVSS (SEQ ID NO:26)

(SEQ ID NO:40)			ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWNSGALTSG VHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKV EPKSCDKTHICPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVTV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDW LNKKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQ VSLTCLVKGFPYSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:32)
	Linker		GGGGSGGGSGGGGS (SEQ ID NO:34)
	scFv of 1A10	VL	QSVLTQPPSASGTPGQRTVISC <u>SGSSSNIGNNYVT</u> WYQQLPGTAPKLLI <u>YADSHRPS</u> GVDPDRFSGSKSGTSASLAISGLRSEDEADYYC <u>ATWDYSLSG</u> <u>YV</u> FGCGTKLTVL (SEQ ID NO:25)
		Linker	GGGGSGGGSGGGSGGGGS (SEQ ID NO:35)
Light component (SEQ ID NO:41)	Light chain of 4F11E2	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFS <u>SYDMS</u> WVRQAPGKCLEWVS <u>WISYSGGSIIYADSVKG</u> RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR <u>DAQRNSMREFDY</u> WGQGT LTVSS (SEQ ID NO:24)
		DIVMTQSPDLSAVSLGERATINC <u>RSSQSLNAGNRKNYLT</u> WYQQKPGQP PKLLIY <u>WASTRES</u> GVDPDRFSGSGSGTDFTLTISLQAEADVAVYYC <u>QNAV</u> <u>SYPT</u> FGGGTKLEIK (SEQ ID NO:27)	
		RTVAAPSVFIFPPSDEQLKSGTASVVCCLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKDYSLSTLTLSKADYEKKHYACEVTHQGLSSPV TKSFNREGC (SEQ ID NO:33)	
		D-1A10	
Heavy component (SEQ ID NO:42)	Heavy chain of 72C1B6A3		QVQLVQSGAEVKKPGASVKVSCKASGYTF <u>TYP</u> IEWVRQAPGQRLWGMG <u>NFHPYND</u> DTKYNEKFKGRVTITRDTASATYME LSSLRSED TAVYYCAR <u>RAYGYPYAMDY</u> WGQGT LTVSS (SEQ ID NO:28)
	Linker		GGGGSGGGSGGGGS (SEQ ID NO:34)
	scFv of 1A10	VL	QSVLTQPPSASGTPGQRTVISC <u>SGSSSNIGNNYVT</u> WYQQLPGTAPKLLI <u>YADSHRPS</u> GVDPDRFSGSKSGTSASLAISGLRSEDEADYYC <u>ATWDYSLSG</u> <u>YV</u> FGCGTKLTVL (SEQ ID NO:25)
		Linker	GGGGSGGGSGGGSGGGGS (SEQ ID NO:35)
Heavy component (SEQ ID NO:42)	Heavy chain of 72C1B6A3	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFS <u>SYDMS</u> WVRQAPGKCLEWVS <u>WISYSGGSIIYADSVKG</u> RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR <u>DAQRNSMREFDY</u> WGQGT LTVSS (SEQ ID NO:24)
		ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWNSGALTSG VHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKV EPKSCDKTHICPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVTV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDW LNKKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQ VSLTCLVKGFPYSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:32)	
		GGGGSGGGSGGGGS (SEQ ID NO:34)	
		GGGGSGGGSGGGSGGGGS (SEQ ID NO:35)	

Light component (SEQ ID NO:43)	Light chain of 72C1B6A3	DIVMTQSPDSLAVSLGERATINC <u>KSSQSLNAGNQNLYLT</u> WYQQKPGQP PKLLIY <u>RASSRES</u> GVDPDRFSGSGSGTDFTLTISLQAEDVAVYYC <u>QNDY</u> <u>IYPYTF</u> GGGTKLEIK (SEQ ID NO:29) RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKSTYSLSTLTLSKADYEKHKVYACEVTHQGLSSPV TKSFNRGEC (SEQ ID NO:33)
E-1A10		
Heavy component (SEQ ID NO:44)	Heavy chain of 120B7B2	QVQLVQSGAEVKKPGASVKVSKASGYTFT <u>GYIIQ</u> WVRQAPGQRLEWMG <u>FINPYND</u> <u>DTKYNEQFKGR</u> VTITRDTSASTAYMELSSLRSEDVAVYYCAR <u>AYFGNAFAY</u> WGQGLTVTVSS (SEQ ID NO:30) ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSG VHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKV EPKSCDKHTHTCPPCPAPELGGPSVFLFPPKPKDTLMISRTPEVTCVTV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY <u>ASTYRVVSVLTVLHQDW</u> LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQ VSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:32)
	Linker	GGGGSGGGSGGGGS (SEQ ID NO:34)
	scFv of 1A10	VL
	Linker	GGGGSGGGSGGGSGGGGS (SEQ ID NO:35)
	VH	EVQLLESGGGLVQPGGSLRLSCAASGFTFS <u>SYDMS</u> WVRQAPGKCLEWVS <u>WISYSGGSIIYADSVKGR</u> FTISRDNKNTLYLQMNSLRAEDVAVYYCAR <u>DAQRNSMREFDY</u> WGQGLTVTVSS (SEQ ID NO:24)
Light component (SEQ ID NO:45)	Light chain of 120B7B2	DIVMTQSPDSLAVSLGERATINC <u>KSSQSLNAGNQNLYLT</u> WYQQKPGQP PKLLIY <u>WASTRES</u> GVDPDRFSGSGSGTDFTLTISLQAEDVAVYYC <u>QNAV</u> <u>YFPFT</u> GGGTKLEIK (SEQ ID NO:31) RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKSTYSLSTLTLSKADYEKHKVYACEVTHQGLSSPV TKSFNRGEC (SEQ ID NO:33)

Table D2. Integrated Sequences of Example Bispecific Antibodies

C-1A10	
Heavy component (SEQ ID NO:40)	EVQLVESGGGLVQPGGSLRLSCAASGFTFS <u>TFGMH</u> WVRQAPGKGLEWVS <u>YITSGESPIYFTD</u> <u>TVKGR</u> FTISRDNKNSLYLQMNSLRAEDVAVYYCAR <u>SSYYGNSMDY</u> WGQGLTVTVSSASTKG PSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSS VTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPELGGPSVFLFPPK PKDTLMISRTPEVTCVTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY <u>ASTYRVVSVLTV</u> LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVK GFYPSDIAVEWESNGQPENNYKTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL

	HNHYTQKSLSLSPGKGGGSGGGGSGGGGSGSVLTQPPSASGTPGQRTVITSC <u>SGSSSNIGNN</u> <u>YVTWYQQLPGTAPKLLIYADSHRPS</u> GVPDFRSGSKSGTSASLAISGLRSEDEADYYC <u>ATWDY</u> <u>SLSGYV</u> FGCGTKLTVLGGGSGGGGSGGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASG FTFS <u>SYDMS</u> WVRQAPGKCLEWVS <u>WISYSGGSIYYADSVKGR</u> FTISRDN SKNTLYLQMNSLRA EDTAVYYCARD <u>DAQRNSMREFDY</u> WGQGLTVTVSS
Light component (SEQ ID NO:41)	DIVMTQSPDSLAVSLGERATINC <u>RSSQSLNAGNRKNYLTWYQ</u> KPGQPPKLLIY <u>WASTRES</u> GVPDFRSGSGSGTDFTLTISSSLQAEDVAVYYC <u>QNAYSYPFT</u> FGGGTKLEIKRTVAAPSVFIF PPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLT LSKADYEKHKVYACEVTHQGLSSPVTKSFNREGC
D-1A10	
Heavy component (SEQ ID NO:42)	QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>TYPIE</u> WVRQAPGQRLEWMG <u>NFHPYNDDTKYNE</u> <u>KFKGR</u> VTITRDTASASTAYMELSSLRSEDTAVYYCARD <u>RAYGYPIAMDY</u> WGQGLTVTVSSASTK GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPELLGGPSVFLFPP KPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY <u>ASTYRV</u> SVSLT VLHQDWLNGKEYCKVSNKALPAPIEKTIISKAKGQPREPQVYITLPPSREEMTKNQVSLTCLV KGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEA LHNHYTQKSLSLSPGKGGGSGGGGSGGGGSGSVLTQPPSASGTPGQRTVITSC <u>SGSSSNIGN</u> <u>NYVTWYQQLPGTAPKLLIYADSHRPS</u> GVPDFRSGSKSGTSASLAISGLRSEDEADYYC <u>ATWD</u> <u>YSLSGYV</u> FGCGTKLTVLGGGSGGGGSGGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAAS GFTFS <u>SYDMS</u> WVRQAPGKCLEWVS <u>WISYSGGSIYYADSVKGR</u> FTISRDN SKNTLYLQMNSLR AEDTAVYYCARD <u>DAQRNSMREFDY</u> WGQGLTVTVSS
Light component (SEQ ID NO:43)	DIVMTQSPDSLAVSLGERATINC <u>KSSQSLNAGNQKNYLTWYQ</u> KPGQPPKLLIY <u>RASSRES</u> GVPDFRSGSGSGTDFTLTISSSLQAEDVAVYYC <u>QNDYIYPYT</u> FGGGTKLEIKRTVAAPSVFIF PPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLT LSKADYEKHKVYACEVTHQGLSSPVTKSFNREGC
E-1A10	
Heavy component (SEQ ID NO:44)	QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>GYIIQ</u> WVRQAPGQRLEWMG <u>FINPYNDDTKYNE</u> <u>QFKGR</u> VTITRDTASASTAYMELSSLRSEDTAVYYCARD <u>AYFGNAFAY</u> WGQGLTVTVSSASTKGP SVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSV VTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPKP KDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY <u>ASTYRV</u> SVSLTVL HQDWLNGKEYCKVSNKALPAPIEKTIISKAKGQPREPQVYITLPPSREEMTKNQVSLTCLVK FYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALH NHYTQKSLSLSPGKGGGSGGGGSGGGGSGSVLTQPPSASGTPGQRTVITSC <u>SGSSSNIGNNY</u> <u>VTWYQQLPGTAPKLLIYADSHRPS</u> GVPDFRSGSKSGTSASLAISGLRSEDEADYYC <u>ATWDYS</u> <u>LSGYV</u> FGCGTKLTVLGGGSGGGGSGGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASG FTFS <u>SYDMS</u> WVRQAPGKCLEWVS <u>WISYSGGSIYYADSVKGR</u> FTISRDN SKNTLYLQMNSLRAE DTAVYYCARD <u>DAQRNSMREFDY</u> WGQGLTVTVSS
Light component (SEQ ID NO:45)	DIVMTQSPDSLAVSLGERATINC <u>KSSQSLNAGNQKNYLTWYQ</u> KPGQPPKLLIY <u>WASTRES</u> GVPDFRSGSGSGTDFTLTISSSLQAEDVAVYYC <u>QNAYYFPFT</u> FGGGTKLEIKRTVAAPSVFIF PPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLT LSKADYEKHKVYACEVTHQGLSSPVTKSFNREGC

[0076] It will also be understood by one of ordinary skill in the art that antibodies as disclosed herein may be modified such that they vary in amino acid sequence from the naturally occurring binding polypeptide from which they were derived. For example, a polypeptide or amino acid sequence derived from a designated protein may be similar, *e.g.*, have a certain percent identity to the starting sequence, *e.g.*, it may be 60%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99% identical to the starting sequence.

[0077] In certain embodiments, the antibody comprises an amino acid sequence or one or more moieties not normally associated with an antibody. Exemplary modifications are described in more detail below. For example, an antibody of the disclosure may comprise a flexible linker sequence, or may be modified to add a functional moiety (*e.g.*, PEG, a drug, a toxin, or a label).

[0078] Antibodies, variants, or derivatives thereof of the disclosure include derivatives that are modified, *i.e.*, by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from binding to the epitope. For example, but not by way of limitation, the antibodies can be modified, *e.g.*, by glycosylation, acetylation, pegylation, phosphorylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the antibodies may contain one or more non-classical amino acids.

[0079] In some embodiments, the antibodies may be conjugated to therapeutic agents, prodrugs, peptides, proteins, enzymes, viruses, lipids, biological response modifiers, pharmaceutical agents, or PEG.

[0080] The antibodies may be conjugated or fused to a therapeutic agent, which may include detectable labels such as radioactive labels, an immunomodulator, a hormone, an enzyme, an oligonucleotide, a photoactive therapeutic or diagnostic agent, a cytotoxic agent, which may be a drug or a toxin, an ultrasound enhancing agent, a non-radioactive label, a combination thereof and other such agents known in the art.

[0081] The antibodies can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antigen-binding polypeptide is

then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminol, isoluminol, therrromatic acridinium ester, imidazole, acridinium salt and oxalate ester.

[0082] The antibodies can also be detectably labeled using fluorescence emitting metals such as ^{152}Eu , or others of the lanthanide series. These metals can be attached to the antibody using such metal chelating groups as diethylenetriaminepentacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA). Techniques for conjugating various moieties to an antibody are well known, *see, e.g.*, Arnon *et al.*, “Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy”, in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), pp. 243-56 (Alan R. Liss, Inc. (1985); Hellstrom *et al.*, “Antibodies For Drug Delivery”, in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.*, (eds.), Marcel Dekker, Inc., pp. 623- 53 (1987); Thorpe, “Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review”, in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera *et al.* (eds.), pp. 475-506 (1985); “Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy”, in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), Academic Press pp. 303-16 (1985), and Thorpe *et al.*, “The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates”, *Immunol. Rev.* (52:119-58 (1982)).

Polynucleotides Encoding the Antibodies and Methods of Preparing the Antibodies

[0083] The present disclosure also provides isolated polynucleotides or nucleic acid molecules encoding the antibodies, variants or derivatives thereof of the disclosure. The polynucleotides of the present disclosure may encode the entire heavy and light chain variable regions of the antigen-binding polypeptides, variants or derivatives thereof on the same polynucleotide molecule or on separate polynucleotide molecules. Additionally, the polynucleotides of the present disclosure may encode portions of the heavy and light chain variable regions of the antigen-binding polypeptides, variants or derivatives thereof on the same polynucleotide molecule or on separate polynucleotide molecules.

[0084] Methods of making antibodies are well known in the art and described herein. In certain embodiments, both the variable and constant regions of the antigen-binding polypeptides of the present disclosure are fully human. Fully human antibodies can be made

using techniques described in the art and as described herein. For example, fully human antibodies against a specific antigen can be prepared by administering the antigen to a transgenic animal which has been modified to produce such antibodies in response to antigenic challenge, but whose endogenous loci have been disabled. Exemplary techniques that can be used to make such antibodies are described in U.S. patents: 6,150,584; 6,458,592; 6,420,140 which are incorporated by reference in their entireties.

Treatment Methods

[0085] As described herein, the antibodies, variants or derivatives of the present disclosure may be used in certain treatment and diagnostic methods.

[0086] The present disclosure is further directed to antibody-based therapies which involve administering the antibodies of the disclosure to a patient such as an animal, a mammal, and a human for treating one or more of the disorders or conditions described herein. Therapeutic compounds of the disclosure include, but are not limited to, antibodies of the disclosure (including variants and derivatives thereof as described herein) and nucleic acids or polynucleotides encoding antibodies of the disclosure (including variants and derivatives thereof as described herein).

[0087] In some embodiments, provided are methods for treating a cancer in a patient in need thereof. The method, in one embodiment, entails administering to the patient an effective amount of an antibody of the present disclosure. In some embodiments, at least one of the cancer cells (e.g., stromal cells) in the patient over-express claudin 18.2.

[0088] Cellular therapies, such as chimeric antigen receptor (CAR) T-cell therapies, are also provided in the present disclosure. A suitable cell can be used, that is put in contact with an antibody of the present disclosure (or alternatively engineered to express an antibody of the present disclosure). Upon such contact or engineering, the cell can then be introduced to a cancer patient in need of a treatment. The cancer patient may have a cancer of any of the types as disclosed herein. The cell (e.g., T cell) can be, for instance, a tumor-infiltrating T lymphocyte, a CD4⁺ T cell, a CD8⁺ T cell, or the combination thereof, without limitation.

[0089] In some embodiments, the cell was isolated from the cancer patient him- or her-self. In some embodiments, the cell was provided by a donor or from a cell bank. When the cell is isolated from the cancer patient, undesired immune reactions can be minimized.

[0090] Non-limiting examples of cancers include bladder cancer, breast cancer, colorectal cancer, endometrial cancer, esophageal cancer, head and neck cancer, kidney cancer, leukemia, liver cancer, lung cancer, lymphoma, melanoma, pancreatic cancer, prostate cancer, and thyroid cancer. In some embodiments, the cancer is one or more of gastric, pancreatic, esophageal, ovarian, and lung cancers.

[0091] Additional diseases or conditions associated with increased cell survival, that may be treated, prevented, diagnosed and/or prognosed with the antibodies or variants, or derivatives thereof of the disclosure include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (*e.g.*, acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (*e.g.*, chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (*e.g.*, Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyo sarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma and retinoblastoma.

[0092] A specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the particular antibodies, variant or derivative thereof used, the patient's age, body weight, general health, sex, and diet, and the time of administration, rate of excretion, drug combination, and the severity of the particular disease being treated. Judgment of such factors by medical caregivers is within the ordinary skill in the art. The

amount will also depend on the individual patient to be treated, the route of administration, the type of formulation, the characteristics of the compound used, the severity of the disease, and the desired effect. The amount used can be determined by pharmacological and pharmacokinetic principles well known in the art.

[0093] Methods of administration of the antibodies, variants or include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The antigen-binding polypeptides or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Thus, pharmaceutical compositions containing the antigen-binding polypeptides of the disclosure may be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), buccally, or as an oral or nasal spray.

[0094] The term “parenteral” as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intra-articular injection and infusion.

[0095] Administration can be systemic or local. In addition, it may be desirable to introduce the antibodies of the disclosure into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, *e.g.*, by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

[0096] It may be desirable to administer the antigen-binding polypeptides or compositions of the disclosure locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, *e.g.*, in conjunction, with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a protein, including an antibody, of the disclosure, care must be taken to use materials to which the protein does not absorb.

[0097] The amount of the antibodies of the disclosure which will be effective in the treatment, inhibition and prevention of an inflammatory, immune or malignant disease, disorder or condition can be determined by standard clinical techniques. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease, disorder or condition, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[0098] As a general proposition, the dosage administered to a patient of the antigen-binding polypeptides of the present disclosure is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight, between 0.1 mg/kg and 20 mg/kg of the patient's body weight, or 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the disclosure may be reduced by enhancing uptake and tissue penetration (*e.g.*, into the brain) of the antibodies by modifications such as, for example, lipidation.

[0099] In an additional embodiment, the compositions of the disclosure are administered in combination with cytokines. Cytokines that may be administered with the compositions of the disclosure include, but are not limited to, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-10, IL-12, IL-13, IL-15, anti-CD40, CD40L, and TNF- α .

[0100] In additional embodiments, the compositions of the disclosure are administered in combination with other therapeutic or prophylactic regimens, such as, for example, radiation therapy.

Compositions

[0101] The present disclosure also provides pharmaceutical compositions. Such compositions comprise an effective amount of an antibody, and an acceptable carrier. In some embodiments, the composition further includes a second anticancer agent (*e.g.*, an immune checkpoint inhibitor).

[0102] In a specific embodiment, the term “pharmaceutically acceptable” means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. Further, a “pharmaceutically acceptable carrier” will generally be a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

[0103] The term “carrier” refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents such as acetates, citrates or phosphates. Antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; and agents for the adjustment of tonicity such as sodium chloride or dextrose are also envisioned. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences by E. W. Martin, incorporated herein by reference. Such compositions will contain a therapeutically effective amount of the antigen-binding polypeptide, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration. The parental preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[0104] In an embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[0105] The compounds of the disclosure can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

EXAMPLES

Example 1: Generation of Anti-CLDN18.2/4-1BB Bispecific Antibodies

[0106] Three previously identified anti-CLDN18.2 antibodies, 4F11E2, 72C1B6A3 and 120B7B2, and an anti-4-1BB antibody, 1A10 (sequences are shown in the tables below), were selected to generate anti-CLDN18.2-4-1BB bispecific antibodies in a full-length IgG X scFv form (structure illustrated in **FIG. 1A**). The anti-Claudin 18.2 portion was placed in full IgG part, while the anti-4-1BB was a scFv placed at the C-terminal side of the Fc fragment. The bispecific antibodies included an IgG1 backbone with a N297A mutation to disable Fc γ function.

Table 1. Antibody Sequences

C-1A10		
Heavy component	Heavy chain of 4F11E2	EVQLVESGGGLVQPGGSLRLSCAASGFTFS TFGMH WVRQAPGKGLEWVS YITSGESPIYFTDTVKG RFTISRDNKNSLYLQMNSLRAEDTAVYYCAR

		<u>SSYYGNSMDY</u> WGQGLTVTVSS (SEQ ID NO:26) ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY <u>A</u> STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:32)
	Linker	GGGGSGGGSGGGGS (SEQ ID NO:34)
	scFv of 1A10	VL QSVLTQPPSASGTPGQRTVISC <u>SGSSSNIGNNYVT</u> WYQQLPGTAPKLLIY <u>ADSHRPS</u> GVDPDRFSGSKSGTSASLAISGLRSEDEADYYC <u>ATWDYSLSG</u> <u>YV</u> FGCGTKLTVL (SEQ ID NO:25)
		Linker GGGGSGGGSGGGGS (SEQ ID NO:35)
		VH EVQLLESGGGLVQPGGSLRLSCAASGFTFS <u>SYDMS</u> WVRQAPGKCLEWVS <u>WISYSGGSIYYADSVKG</u> RFTISRDNKNTLYLQMNSLRAEDTAVYYCAR <u>DAQRNSMREFDY</u> WGQGLTVTVSS (SEQ ID NO:24)
Light component	Light chain of 4F11E2	DIVMTQSPDSLAVSLGERATINC <u>RSSQSLNAGNRKNYLT</u> WYQQKPGQP PKLLIY <u>WASTRES</u> GVDPDRFSGSGSGTDFTLTISSLQAEDVAVYYC <u>QONAY</u> <u>SYPFT</u> FGGGTKLEIK (SEQ ID NO:27) RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKKHYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:33)
D-1A10		
Heavy component	Heavy chain of 72C1B6A3	QVQLVQSGAEVKKPGASVKVSCKASGYTFTT <u>TYPIE</u> WVRQAPGQRLEWMG <u>NFHPYNDDTKYNEKFKG</u> RVTITRDTSASTAYMELSSLRSEDTAVYYCAR <u>RAYGYPYAMDY</u> WGQGLTVTVSS (SEQ ID NO:28) ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY <u>A</u> STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:32)
		Linker GGGGSGGGSGGGGS (SEQ ID NO:34)
		scFv of 1A10
		Linker GGGGSGGGSGGGGS (SEQ ID NO:35)

		VH	EVQLLES GGGGLVQP GGS LRLSCAASGFTFS <u>SYDMS</u> WVRQAPGKCLEWVS <u>WISYSGGSIYYADSVKGR</u> FTISRDN SKNTLYLQMNSLRAEDTAVYYCAR <u>DAQRNSMREFDY</u> WGQGLTVTVSS (SEQ ID NO:24)
Light component	Light chain of 72C1B6A3		DIVMTQSPDSLAVSLGERATINC <u>KSSQSLNAGNQKNYLT</u> WYQQKPGQP PKLLIY <u>RASSRES</u> GVPDRFSGSGSGTDFTLTISSLQAEDVAVYYC <u>QNDY</u> <u>IYPYTF</u> GGGKLEIK (SEQ ID NO:29) RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPV TKSFNRGEC (SEQ ID NO:33)
E-1A10			
Heavy component	Heavy chain of 120B7B2		QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>GYIIQ</u> WVRQAPGQRLEWMG <u>FINPYND DTKYNEQFKGR</u> VTITRDTSA STAYMELSSLRSEDTAVYYCAR <u>AYFGNAFAY</u> WGQGLTVTVSS (SEQ ID NO:30) ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSG VHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKV EPKSCDKHTICPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY <u>A</u> STYRVSVLTVLHQDW LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQ VSLTCLVKGFPYSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK (SEQ ID NO:32)
	Linker		GGGGSGGGSGGGGS (SEQ ID NO:34)
	scFv of 1A10	VL	QSVLTQPPSASGTPGQRTISCS <u>SGSSSNIGNNYVT</u> WYQQLPGTAPKLLI Y <u>ADSHRPS</u> GVPDRFSGSKSGTSASLAISGLRSEDEADYYC <u>ATWDYSLSG</u> <u>YVFGCGTKLTVL</u> (SEQ ID NO:25)
		Linker	GGGGSGGGSGGGSGGGGS (SEQ ID NO:35)
Light component	Light chain of 120B7B2	VH	EVQLLES GGGGLVQP GGS LRLSCAASGFTFS <u>SYDMS</u> WVRQAPGKCLEWVS <u>WISYSGGSIYYADSVKGR</u> FTISRDN SKNTLYLQMNSLRAEDTAVYYCAR <u>DAQRNSMREFDY</u> WGQGLTVTVSS (SEQ ID NO:24)
			DIVMTQSPDSLAVSLGERATINC <u>KSSQSLNAGNQKNYLT</u> WYQQKPGQP PKLLIY <u>WASTRES</u> GVPDRFSGSGSGTDFTLTISSLQAEDVAVYYC <u>QNAV</u> <u>YFPFT</u> GGGKLEIK (SEQ ID NO:31) RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPV TKSFNRGEC (SEQ ID NO:33)

[0107] The 4F11E2 and 1A10 sequences were also used to generate bispecific antibodies of two different formats, as illustrated in FIG. 1B and FIG. 1C, respectively. In the format of FIG. 1B, the anti-Claudin 18.2 portion also took a Fab format, while the anti-4-1BB portion was present as a scFab fragment inserted between the anti-Claudin 18.2 Fab and the Fc fragments. The IgG1 (N297A) was also used here.

Table 2. Antibody of Format of FIG. 1B

Fab-scFab-Fc			
Heavy component (SEQ ID NO:54)	Heavy chain of 4F11E2		EVQLVESGGGLVQPGGSLRLSCAASGFTFS TFGMH WVRQAPGKGLEWV SY ITSGESPIYFTDTVKGR FTISRDNAKNSLYLQMNSLRAEDTAVYYC AR SSYYGNSMDY WGQGTLLTVSS (SEQ ID NO:26) ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDK KVEPKSC (SEQ ID NO:36)
	Linker		GGGSGGGSGGGSGGGGS (SEQ ID NO:35)
	scFab of 1A10	Light chain of 1A10	QSVLTQPPSASGTPGQRTVITSC SGSSSNIGNNYVT WYQQLPGTAPKLL IY ADSHRPS GVPDRFSGSKSGTSASLAISGLRSEADYYCATWDYSL SGYV FGGGTKLTVL (SEQ ID NO:53) GQPKANPTVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADGSP VKAGVETTKPSKQSNKYAASSYLSLTPEQWKSHRYSYSCQVTHEGSTV EKTVAPECS (SEQ ID NO:37)
		Linker	GSGSGSGSGSGSGSGSGSGGGSGGGSGGGSGGGSGGGSGGGSGGGGS GSGSGSGSGSGSGSGS (SEQ ID NO:38)
		Heavy chain of 1A10	EVQLLESGGGLVQPGGSLRLSCAASGFTFS SYDMS WVRQAPGKGLEWV SW ISYSGGSIYYADSVKGR FTISRDNKNTLYLQMNSLRAEDTAVYYC AR DAQRNSMREFDY WGQGTLLTVSS (SEQ ID NO:47) ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDK KVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDITLMISRTPEVTC VVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY A STYRVVSVLTVL HQDWLNGKEYCKVSNKALPAPIEKTIKAKGQPREPQVYTLPPSREE MTKNQVSLTICLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFF LYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:32)
Light component (SEQ ID NO:41)	Light chain of 4F11E2		DIVMTQSPDLSAVSLGERATINC RSSQSLNAGNRKNYLT WYQQKPGQ PPKLLIY WASTRES GVPDRFSGSGSGTDFTLTISSLQAEDVAVYYC QCN AYSYPFT FGGGTKLEIK (SEQ ID NO:27) RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQ SGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSS PVTKSFNRGEC (SEQ ID NO:33)

[0108] In the format of FIG. 1C, the anti-Claudin 18.2 portion was present in full IgG1, while the anti-4-1BB portion was a scFab fragment placed at the C-terminal side of the Fc.

Table 3. Antibody of Format of FIG. 1C

IgG-scFab		
Heavy component (SEQ ID NO:55)	Heavy chain of 4F11E2	EVQLVESGGGLVQPGGSLRLSCAASGFTFS TFGMH WVRQAPGKGLE WVSY ITSGESPIYFTDTVKGR FTISRDNAKNSLYLQMNSLRAEDTA VYYCAR SSYYGNSMDY WGQGTLLTVSS (SEQ ID NO:26) ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGAL TSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNT

			KVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYASTYR VVSVELTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ VYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKT TPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQK SLSLSPGK (SEQ ID NO:32)
	Linker		GGGGSGGGSGGGGS (SEQ ID NO:34)
	scFab of 1A10	Light chain of 1A10	QSVLTQPPSASGTPGQRTVISC <u>SGSSSNIGNNYVT</u> WYQQLPGTAPK LLIY <u>ADSHRPS</u> GVDPDRFSGSKSGTSASLAISGLRSEDEADYYC <u>CATW</u> <u>DYSLSGYV</u> FGGGTKLTVL (SEQ ID NO:53) GQPKANPTVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADG SPVKAGVETTKPSKQSNKYYASSYLSLTPEQWKSHRSYSCQVTHE GSTVEKTVAPTECS (SEQ ID NO:37)
		Linker	GSGSGSGSGSGSGSGSGSGGGSGGGSGGGSGGGSGGGSGGGSGGG GSGSGSGSGSGSGSGSGS (SEQ ID NO:38)
		Heavy chain of 1A10	EVQLLESQGGGLVQPGGSLRLSCAASGFTFS <u>SYDMS</u> WVRQAPGKGLE WVS <u>WISYSGGSIIYADSVKGR</u> FTISRDNKNTLYLQMNSLRAEDTA VYYCAR <u>DAQRNSMREFDY</u> WGQGLTVTVSS (SEQ ID NO:47) ASTKGPSTVPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGAL TSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNT KVDKKVEPKSC (SEQ ID NO:36)
Light component (SEQ ID NO:41)	Light chain of 4F11E2		DIVMTQSPDSLAVSLGERATINC <u>RSSQSLLNAGNRKNYLT</u> WYQQKP GQPPKLLIY <u>WASTRES</u> GVDPDRFSGSGSGTDFTLTISLQAEDVAVY YC <u>QNAVSYPTFT</u> FGGGTKLEIK (SEQ ID NO:27) RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNA LQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQ GLSSPVTKSFNRGEC (SEQ ID NO:33)

[0109] The bispecific antibodies of the formats in FIG. 1B and 1C were outperformed by those of format in FIG. 1A in terms of 4-1BB binding, in preliminary testing. Thus, only those of format FIG. 1A underwent additional testing as shown below. Further, E-1A10 was slightly less potent than C-1A10 and D-1A10 and was thus also not included in further studies.

Example 2. Antigen Binding of the Anti-CLDN18.2/4-1BB Bispecific Antibodies

[0110] This example evaluated the binding activities of the bispecific antibodies of Example 1 to CLDN18.2 and 4-1BB, using protein-based and cell-based assays.

2.1 ELISA binding to 4-1BB

[0111] Briefly, microtiter plates were coated with 100µl human 4-1BB-His protein at 0.5µg/ml diluted in PBS at 4°C overnight, then blocked with 100 µl/well of 1% BSA. Three-fold dilution of antibodies starting from 100nM were added to each well and incubated for 2 hours at RT. The plates were washed with PBS/Tween and then incubated with goat-anti-human IgG antibody conjugated with Horse Radish Peroxidase (HRP) for 30 minutes at RT. After washing, the plates were developed with TMB substrate and analyzed by plate reader at OD450nm. As shown in **FIG. 2A**, the anti-CLDN18.2-4-1BB antibodies (C-1A10 and D-1A10) showed comparable binding as anti-4-1BB monoclonal antibodies (1A10, and urelumab (BMUR)).

2.2 ELISA binding to CLDN18.2

[0112] To test CLDN18.2 binding, the CLDN18.2 protein was expressed in a Virus Like Particle (VLP) to mimic the nature conformation. Similarly, microtiter plates were coated with 100µl CLDN18.2 VLP at 3µg/ml diluted in PBS at 4°C overnight, then blocked with 300 µl/well of 3% BSA. Three-fold dilution of antibodies starting from 100nM were added to each well and incubated for 2 hours at 37°C. The plates were washed with PBS/Tween and then incubated with goat-anti-human IgG antibody conjugated with Horse Radish Peroxidase (HRP) for 30 minutes at RT. After washing, the plates were developed with TMB substrate and analyzed by plate reader at OD450nm. As shown in **FIG. 2B**, the anti-CLDN18.2-4-1BB antibodies (C-1A10 and D-1A10) showed stronger binding than the anti-CLDN18.2 monoclonal antibody (I-MAB362). Meanwhile, in the same experiment setting, the binding to CLDN18.1 protein was tested as well. As shown in **FIG. 2C**, no antibody showed cross-reactivity to CLDN18.1.

2.3 ELISA binding to dual antigen

[0113] To further demonstrate that the anti-CLDN18.2-4-1BB antibodies can bind to CLDN18.2 and 4-1BB simultaneously, a DACE (Dual antigen captured ELISA) test was performed. Briefly, microtiter plate was coated with 100µl CLDN18.2 VLP at 3µg/ml diluted in PBS at 4°C overnight. Then, three-fold dilution of antibodies starting from 100nM were added to each well and incubated for 2 hours at 37°C. After washing, the plate was then incubated with 100µl human 4-1BB-biotin protein at a concentration of 1µg/ml for 1 hour at

37°C and followed with Streptavidin HRP for another 30 minutes at room temperature. After washing, the plates were developed with TMB substrate and analyzed by plate reader at OD 450nm. As shown in **FIG.2D**, D-1A10 can bind to hCLDN18.2 and 4-1BB at the same time. On the contrast, CLDN18.2 mAb 72C1B6A3 showed no signal in this assay.

2.4 Cell-based binding to 4-1BB

[0114] To evaluate the antigen binding property, the anti-CLDN18.2-4-1BB antibodies were analyzed for their binding to HEK293 expressed 4-1BB by FACS. A total number of 1×10^5 HEK293-4-1BB cells in each well were incubated with serial diluted antibodies for 30 minutes at 4°C in FACS buffer (2%FBS in PBS). After wash by FACS buffer, PE conjugated-anti-human IgG antibody was added to each well and incubated at 4°C for 30 minutes. After wash, MFI of PE was evaluated by FACS Caliber. As shown in **FIG. 3**, the anti-CLDN18.2-4-1BB antibodies (C-1A10 and D-1A10) tested showed concentration-dependent binding abilities to 4-1BB, comparable with the monoclonal antibodies (1A10 and BMUR).

2.5 Cell-based binding to CLDN18.2

[0115] To evaluate the binding capability towards CLDN18.2, a CHO-K1 cell line which stably expressed human CLDN18.2 was made. CHO-C18.2 cell line was then sorted for high expressers (CHO-K1 C18.2 High) and low expressers (CHO-K1 C18.2 Low) using flow cytometry. CHO-K1-C18.2 cells were incubated with serial diluted antibodies for 30 minutes at 4°C in FACS buffer. After wash by FACS buffer, PE conjugated-anti-human IgG antibody was added and incubated at 4°C for 30 minutes. MFI of PE was evaluated by FACS. As shown in **FIG.4A and 4B**, anti-CLDN18.2-4-1BB antibodies bound to CLDN18.2-expressed cells in a concentration-dependent manner; meanwhile, both of the bispecific antibodies showed stronger binding than IMAB362, a reference monoclonal anti-CLDN18.2 antibody currently in clinical trials.

[0116] SNU620 is a gastric carcinoma cell line with endogenous CLDN18.2 expression. As shown in **FIG. 4C**, the anti-CLDN18.2-4-1BB bispecific antibodies could bind to SNU620 as well.

Example 3. Functional Activity of Anti-CLDN18.2/4-1BB Bispecific Antibodies

3.1 Cell-line based functional characterization of CLDN18.2-4-1BB bispecific antibody

[0117] To test the ability of the bispecific antibodies to promote 4-1BB signal, a commercial 4-1BB assay was used. In this assay, GloResponse™ NFκB-luc2/4-1BB Jurkat cell line (Promega, cat# CS196004) was used as effector cells and CHO-K1-expressing or not expressing CLDN18.2 or SNU620 as target cells. GloResponse™ NFκB-luc2/4-1BB Jurkat cell line is genetically modified to stably express 4-1BB and luciferase downstream of a response element. Luciferase expression is induced upon antibody binding to the 4-1BB receptor. In brief, effector cells at a density of 2.5×10^4 cells per well were mixed with 2.5×10^4 target cells in a white 96-well plate. Antibodies 6-fold serially diluted were added to a white 96-well assay plate, at a final concentration ranging from 16.7 nM to 0.28 fm. After 6 hours' incubation at 37°C, luminescence was obtained by adding the substrate of luciferase and measured by a microplate reader (PHERAstar). Four-parameter logistic curve analysis was performed with GraphPad software.

[0118] As shown in **FIG. 5**, the BMUR monoclonal antibody can dose-dependent boost the 4-1BB signaling, while the 1A10 monoclonal antibody had no agonist activity in the same experimental settings. The activity of anti-CLDN18.2-4-1BB bispecific antibodies D-1A10 and C-1A10 was dependent on the expression of CLDN18.2.

3.2 Activity of the bispecific antibodies to promote human peripheral blood mononuclear cell (PBMC) immune response

[0119] To test the ability of bispecific antibodies to stimulate human PBMCs response, cytokine production assay was used. Human PBMCs stimulated with 0.5 μg/ml human anti-

CD3 antibody were used as the effector cells. CHO-K1 express CLDN18.2 was used as the target cells. Human PBMCs (1×10^5) were co-cultured with CHO-K1-CLDN18.2 or control CHO-K1 cells (2.5×10^4) in the presence of human anti-CD3 antibody. Serially diluted bispecific antibodies were added to the mixed culture at a final concentration starting from 20nM. 48 hours later, the level of IL2 and IFN- γ in the culture medium was measured using IL-2 (human) LANCE Ultra TR-FRET Detection Kit and IFN- γ (human) LANCE Ultra TR-FRET Detection Kit (PerkinElmer). As shown in **FIG. 6**, only bispecific antibodies can activate PBMC response in the presence of CLDN18.2 expressing cells.

[0120] To further demonstrate that the activity of bispecific antibodies was correlated with CLDN18.2 level, human PBMC from one donor were cocultured with different Gastric Adenocarcinoma (GA) PDX derived cells, which was proved to express different level of C18.2. Serially diluted bispecific antibodies were added to the mixed culture at a final concentration starting from 20nM. After 48 hours, the level of IL2 in the culture medium was measured using IL-2 (human) LANCE Ultra TR-FRET Detection Kit (PerkinElmer). As shown in **FIG. 6D and 6E**, the activity of bispecific antibody D-1A10 correlated with CLDN18.2 level. D-1A10 effectively activated T cells even in CLDN18.2 low- to middle-expressed tumors.

3.3 Activity of the bispecific antibodies to promote human CD8+ T cell response

[0121] To further test the ability of bispecific antibodies in activating human CD8+ T cell, human CD8+ T cells isolated from PBMCs were used as the effector cells. CHO-K1 expressed CLDN18.2 or SNU620 was used as the target cells. Isolated CD8+ T cells (7.5×10^4) were co-cultured with target cells (2.5×10^4) in the presence of human anti-CD3 antibody. Serially diluted bispecific antibodies were added to the mixed culture at a final concentration starting from 20nM. The level of IL2 and IFN- γ in the culture medium was measured 48 hours after coculture, using IL-2 (human) LANCE Ultra TR-FRET Detection Kit and IFN- γ (human) LANCE Ultra TR-FRET Detection Kit (PerkinElmer). As shown in **FIG. 7**, bispecific antibodies C-1A10 and D-1A10 can both increase the production of IL2

and IFN- γ of activated CD8⁺ T cells in the presence of C18.2 overexpressing CHO-K1 (FIG. 7A-7B) or SNU620 which endogenously express C18.2 (FIG. 7C-7D).

Example 4. Tumor Growth Inhibition by Anti-CLDN18.2-4-1BB Bispecific Antibodies

[0122] Humanized mice that expressed the extracellular domain of human 4-1BB were used. Mouse colon adenocarcinoma cells (MC38) were engineered to express human CLDN18.2. Humanized mice (h4-1BB) were subcutaneously implanted with MC38-hCLDN18.2 cells. Mice were intraperitoneally administered every 3 days for 5 times with following antibodies: isotype control (10 mg/kg), anti-CLDN18.2 antibody (10 mg/kg), anti-4-1BB antibody (10 mg/kg), combination of anti-CLDN18.2 (10 mg/kg) and anti-4-1BB (10 mg/kg) and anti-CLDN18.2-4-1BB bispecific antibody (13.3 mg/kg). Tumor volumes were monitored by caliper measurement twice per week for the duration of the experiment. Tumor growth inhibition induced by bispecific antibodies was significantly greater than that observed with the combination of each targeting monoclonal antibodies, shown in FIG. 8. To further understand the mechanism of the antibody, tumor infiltrated lymphocytes and peripheral lymphocytes were collected and analyzed for the percentage of CD3⁺ T cells by flow cytometry. Results showed that bispecific antibodies can specifically increase the number of CD3⁺ T cells in the tumor microenvironment, while had no impact on peripheral blood.

[0123] In another repeated experiment using the same animal model, the *in vivo* efficacy was further proved. The FIG. 9A showed the tumor growth curve of each animal in each group. The tumor growth inhibition of anti-CLDN18.2-4-1BB antibody (C-1A10) reached 105% at the end of study. Six of 7 mice in the group of bispecific were tumor free by day 25 following the first treatment.

[0124] In addition, this example re-challenged all anti-CLDN18.2-4-1BB antibody (C-1A10)-treated animals with a second dose of MC38-hCLDN18.2 tumor cells in the contralateral flank 35 days after the first tumor inoculation and monitored tumor growth without giving additional treatment. Results showed that as the tumor cells continued to grow in naïve mice, all BsAb-treated mice were resistant to tumor re-challenge and were

deemed tumor free till the end of the study (FIG. 9B), suggesting that the present bispecific antibodies can induce long-term protective immunological memory again MC38 tumors. Similarly, in a satellite group (N=3/group) where mice received the same treatment, tumors were extracted 3 days after the second dose of antibodies to quantify tumor-infiltrating lymphocytes (TILs). The percentage of CD45⁺ and CD8⁺ TILs was significantly higher in BsAb treatment group. In contrast, there was no effect on the peripheral lymphocytes (FIG.9C).

[0125] To further evaluate the anti-tumor efficacy of anti-CLDN18.2-4-1BB bispecific antibodies, different concentrations of bispecific antibodies were given to 4-1BB humanized mice implanted with MC38-hCLDN18.2 antibodies. Mice were intraperitoneally administered every 3 days for 4 times with following antibodies: isotype control (10 mg/kg), C-1A10 (13.3 mg/kg), C-1A10 (2.6mg/kg), C-1A10 (0.5mg/kg), D-1A10 (13.3mg/kg), D-1A10 (2.6mg/kg), D-1A10 (0.5mg/kg). Tumor volumes were monitored by caliper measurement twice per week for the duration of the experiment. As shown in FIG.10, both C-1A10 and D-1A10 can suppress the tumor growth in a dose-dependent manner.

[0126] To understand the pharmacokinetics (PK) and pharmacodynamics relationship of anti-CLDN18.2-4-1BB in humanized mouse mode, single administration of D-1A10 at 3 different concentrations was given to tumor bearing - humanized mice. Serum concentration was measured at various time points. As showed in FIG.11, overall, it displayed a dose-dependent PK profile. And the *in vivo* efficacy was correlated with the dose level. *Ex vivo* TIL analysis suggested a dose-dependent increase of CD8⁺ cells and CD45⁺ cells.

Example 5. Testing of Additional Conditional Agonist Full Human Anti-4-1BB Antibodies

[0127] The following anti-4-1BB antibodies were identified for their ability to bind to 4-1BB at high affinity and not to activate 4-1BB signaling upon binding.

Table 4. Additional Anti-4-1BB Antibodies

Antibody No.	VH	SEQ ID NO:
41B01	EVQLLES GGGGLVQP GGSRLRLSCAASGFTFSSYDMSWVRQAPGKGLEWVSWIS YSGGSIYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDGQRNS MREFDYWGQGTIVTVSS	46

41B01.01	EVQLLES GGGGLVQPGGSLRLS CAASGFTFS SYDMSWVRQAPGKGLEWVSWIS YSGGSIYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR <u>DAQRNS</u> MREFDYWGQGLTVTVSS	47
41B01.02	EVQLLES GGGGLVQPGGSLRLS CAASGFTFS SYDMSWVRQAPGKGLEWVSWIS YSGGSIYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR <u>DAQRQS</u> MREFDYWGQGLTVTVSS	48
41B01.03	EVQLLES GGGGLVQPGGSLRLS CAASGFTFS SYDMSWVRQAPGKGLEWVSWIS YSGGSIYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR <u>DGQRQS</u> MREFDYWGQGLTVTVSS	49
41B01.04	EVQLLES GGGGLVQPGGSLRLS CAASGFTFS SYDMSWVRQAPGKGLEWVSWIS YSGGSIYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR <u>DAQRNA</u> MREFDYWGQGLTVTVSS	50
41B02	EVQLLES GGGGLVQPGGSLRLS CAASGFTFS GYDMSWVRQAPGKGLEWVSVIY PDDGNTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDAAVYYCAK <u>HGGQKP</u> TTKSSSAYGMDGWGQGLTVTVSS	51

Antibody No.	VL	SEQ ID NO:
41B01	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGGG TKLTVL	52
41B01.01	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGGG TKLTVL	52
41B01.02	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGGG TKLTVL	52
41B01.03	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGGG TKLTVL	53
41B01.04	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGGG TKLTVL	53
41B02	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGGG TKLTVL	52

CDRs for VHs in Table 4

SEQ ID NO	VH_CDR1	SEQ ID NO	VH_CDR2	SEQ ID NO	VH_CDR3
1	SYDMS	2	WISYSGGSIYYADSVKG	3	DAQRNSMREFDY
				56	DGQRNSMREFDY
				57	DAQRQSMREFDY
				58	DGQRQSMREFDY
				59	DAQRNAMREFDY
60	GYDMS	61	VIYPDDGNTYYADSVKG	62	HGGQKPTTKSSSAYGMDG

Table 5. VHs and VLs of Anti-4-1BB scFv

Antibody No.	VH	SEQ ID NO:
--------------	----	------------

41B01	EVQLLES GGGGLVQP GGSRLRLSCAASGFTFSSYDMSWVRQAPGK <u>CLEWVSWIS</u> <u>YSGGSIYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDGQRNS</u> <u>MREFDYWGQGLTVTVSS</u>	63
41B01.01	EVQLLES GGGGLVQP GGSRLRLSCAASGFTFSSYDMSWVRQAPGK <u>CLEWVSWIS</u> <u>YSGGSIYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDAQRNS</u> <u>MREFDYWGQGLTVTVSS</u>	64
41B01.03	EVQLLES GGGGLVQP GGSRLRLSCAASGFTFSSYDMSWVRQAPGK <u>CLEWVSWIS</u> <u>YSGGSIYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDQRNS</u> <u>MREFDYWGQGLTVTVSS</u>	65
41B01.04	EVQLLES GGGGLVQP GGSRLRLSCAASGFTFSSYDMSWVRQAPGK <u>CLEWVSWIS</u> <u>YSGGSIYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDAQRNS</u> <u>MREFDYWGQGLTVTVSS</u>	66
41B02	EVQLLES GGGGLVQP GGSRLRLSCAASGFTFSGYDMSWVRQAPGK <u>CLEWVSVIY</u> <u>PDDGNTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDAAVYYCAKHGGQKP</u> <u>TTKSSSAYGMDGWGQGLTVTVSS</u>	67
41B02.01	EVQLLES GGGGLVQP GGSRLRLSCAASGFTFSGYDMSWVRQAPGK <u>CLEWVSVIY</u> <u>PDDGNTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKHGGQKP</u> <u>TTKSSSAYGMDGWGQGLTVTVSS</u>	68

Antibody No.	VL	SEQ ID NO:
41B01	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGCG TKLTVL	69
41B01.01	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGCG TKLTVL	70
41B01.03	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGCG TKLTVL	71
41B01.04	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGCG TKLTVL	72
41B02	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGCG TKLTVL	73
41B02.01	QSVLTQPPSASGTPGRRVTISCSGSSSNIGNNYVTWYQQLPGTAPKLLIYAD SHRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCATWDYSLSGYVFGCG TKLTVL	74

5.1 Antigen binding measured by ELISA

[0128] To evaluate the antigen binding activity, the antibodies were subjected to ELISA test. Briefly, microtiter plates were coated with human 4-1BB-Fc protein at 0.1 µg/ml in PBS, 100 µl/well at 4°C overnight, then blocked with 100 µl/well of 5% BSA. Five-fold dilutions of the antibodies starting from 10 µg/ml were added to each well and incubated for 1-2 hours at RT. The plates were washed with PBS/Tween and then incubate with goat-anti-human IgG

antibody conjugated with Horse Radish Peroxidase (HRP) for 1 hour at RT. After washing, the plates were developed with TMB substrate and analyzed by spectrophotometer at OD 450-630nm. The anti-4-1BB antibodies tested showed 4-1BB binding abilities.

5.2 Cell binding measured by FACS

[0129] To evaluate the antigen binding property, the antibody candidates were analyzed for its binding to mammalian expressed 4-1BB by FACS. Briefly, 4-1BB-Jurkat cells were incubated with the antibodies. After wash by FACS buffer (1%BSA in PBS), the FITC-anti-human IgG antibody was added to each well and incubated at 4°C for 1 hour. The MFI of FITC was evaluated by FACS Caliber. The anti-4-1BB antibodies tested showed binding abilities to 4-1BB which expressed on cell surface and can efficiently bind to 4-1BB expressed on mammalian cells.

5.3 Protein kinetic for 4-1BB

[0130] To explore the binding kinetics of the antibodies, this example performed the affinity ranking by using Octet Red 96. As shown in Table 5, the anti-4-1BB antibodies tested had high 4-1BB binding affinities.

Table 5. Binding Kinetics

Antibody	KD (M)	kon(1/Ms)	kdis(1/s)	Chi	R ²
41B01	1.80E-10	6.58E+05	1.19E-04	0.0392	0.9987
41B02	1.01E-09	5.95E+05	6.03E-04	0.0525	0.9973

* * *

[0131] The present disclosure is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the disclosure, and any compositions or methods which are functionally equivalent are within the scope of this disclosure. It will be apparent to those skilled in the art that various modifications and variations can be made in the methods and compositions of the present disclosure without departing from the spirit or scope of the disclosure. Thus, it is intended that the present

disclosure cover the modifications and variations of this disclosure provided they come within the scope of the appended claims and their equivalents.

[0132] All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.