

**LONG-ACTING COAGULATION FACTORS AND METHODS OF PRODUCING
SAME**

FIELD OF INVENTION

5 [001] Polypeptides comprising at least one carboxy-terminal peptide (CTP) of chorionic gonadotrophin attached to the carboxy terminus of a coagulation factor and polynucleotides encoding the same are disclosed. Pharmaceutical compositions comprising the polypeptides and polynucleotides of the invention and methods of using and producing same are also disclosed.

10

BACKGROUND OF THE INVENTION

[002] The development of coagulation factor replacement therapy has transformed the lives of many individuals with hemophilia. Hemophilia is a group of hereditary genetic disorders that impair the body's ability to control blood clotting or coagulation. Patients with hemophilia do not produce adequate amounts of Factor VIII or Factor IX proteins, which are necessary for effective blood clotting. In severe hemophiliacs even a minor injury can result in blood loss that continues for days or weeks, and complete healing may not occur, leading to the potential for debilitating permanent damage to joints and other organs, and premature death.

20 [003] One type of hemophilia, Hemophilia B, is an X-linked bleeding disorder caused by a mutation in the Factor IX (FIX) gene, resulting in a deficiency of the procoagulant activity of FIX. Hemophilia B patients have spontaneous soft tissue hemorrhages and recurrent hemarthroses that often lead to a crippling arthropathy. Current treatment for these patients includes an intravenous administration of recombinant FIX. However issues of cost and relatively rapid clearance of FIX from the circulation make developing a long-acting FIX a challenging task.

[004] Commercial availability of FVIII and FIX has lead to improved control of life-threatening bleedings episodes. Many patients receive prophylactic therapy, which reduces the risk of bleeding and its associated complications. However, a significant proportion of patients (10-30%) develop inhibitory antibodies to exogenously administered FVIII and FIX. Administration of FVIIa, which is a bypassing product, can induce homeostasis and provide an effective treatment for patients with inhibitory Abs.

30 [005] Recombinant FVIIa (NovoSeven®) is commercially available and was approved in 1996 for treatment of bleeding episodes in hemophilia patients with inhibitors. However,

rFVIIa is rapidly cleared with a terminal half-life of 2.5 hours. As a result, patients generally require multiple, frequent infusions (2-3 doses given in 2-3 hour intervals) to achieve adequate homeostasis following a mild to moderate bleed. Consequently, there is much interest in developing a long-acting form of FVIIa that would prolong the duration of haemostatic activity following a single dose and allow much less frequent dosing. A long-acting FVIIa would also increase the feasibility of long-term prophylactic therapy.

[006] Various technologies are being developed for prolonging the half-life of FVIIa. However, the challenge is to achieve a prolonged half-life of this protein while preserving its biological activity and ensuring that the modifications do not induce significant immunogenicity.

SUMMARY OF THE INVENTION

[007] In one embodiment, the present invention provides a CTP-modified Factor IX (FIX) polypeptide consisting of a FIX polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said CTP-modified FIX polypeptide.

[008] In another embodiment, the present invention provides a pharmaceutical composition comprising a CTP-modified Factor IX (FIX) polypeptide consisting of a FIX polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said CTP-modified FIX polypeptide.

[009] In another embodiment, the present invention provides a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor IX (FIX) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide.

[010] In another embodiment, the present invention provides an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor IX (FIX) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide.

[011] In another embodiment, the present invention provides a cell comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor IX (FIX) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide.

[012] In another embodiment, the present invention provides a composition comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor IX (FIX) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide.

[013] In another embodiment, the present invention provides a method of extending the biological half-life of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby extending the biological half-life of said FIX polypeptide.

[014] In another embodiment, the present invention provides a method of improving the area under the curve (AUC) of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby improving the AUC of said FIX polypeptide.

[015] In another embodiment, the present invention provides a method of reducing the dosing frequency of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby reducing the dosing frequency of said FIX polypeptide.

[016] In another embodiment, the present invention provides a method of reducing the clearance rate of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby reducing the clearance rate of said FIX polypeptide.

[017] In another embodiment, the present invention provides a method of producing a CTP-modified Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby producing a CTP-modified FIX polypeptide.

[018] In another embodiment, the present invention provides a method of treating hemophilia in a subject comprising administering a CTP-modified Factor IX (FIX) polypeptide comprising a FIX polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide to said subject, thereby treating hemophilia in said subject.

[019] In one embodiment, the present invention provides A method of preventing a blood clotting or coagulation disorder in a subject, the method comprising the step of administering to the subject a CTP-modified coagulation factor, comprising three to five chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVII polypeptide, thereby preventing hemophilia in said subject.

[020] In another embodiment, the present invention relates to a method of treating a blood clotting or coagulation disorder in a subject, the method comprising the step of administering to the subject a CTP-modified coagulation factor, comprising three to five chorionic

gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said coagulation factor, thereby preventing hemophilia in said subject.

[021] Other features and advantages of the present invention will become apparent from the following detailed description examples and figures. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[022] **Figure 1A.** Shows a bar graph showing harvests limited, diluted, transfected, and selected cells with FIX-CTP and FIX-CTP-CTP variants in the presence of 5µg/ml of Vitamin K3. The level of FIX was quantified using Human FIX ELISA kit (Affinity Biologicals; Cat. No. FIX-AG RUO), and the calculated protein concentration (µg/ml) is the average of two independent runs. Figure 1B shows SDS-PAGE gel micrographs of FIX Ab recognition. Micrograph A depicts recognition of anti-FIX antibody in Western-blot; Micrograph B depicts recognition of anti-γ carboxylation antibody in Western-blot. Lane 1 in A-B was loaded with a sample containing recombinant FIX. Lane 2 in A-B was loaded with a sample containing FIX-CTP harvests. Lane 3 in A-B was loaded with a sample containing FIX-(CTP)₂ harvest.

[023] **Figure 2.** Shows a graph showing FIX-CTP and FIX-(CTP)₂ harvests comparative chromogenic activity (measured by the EC₅₀ concentration) compared to rhFIX (American Diagnostics).

[024] **Figure 3.** Shows a graph showing PK profile of rhFIX, harvest of FIX-CTP-CTP, and harvest of FIX-CTP.

[025] **Figure 4.** Shows a bar graph showing harvests of FIX-CTP and FIX-CTP-CTP harvests and FIX-CTP-CTP purified protein FIX antigen level as determined using Human FIX ELISA kit (Affinity Biologicals; cat. No. FIX-AG RUO). The calculated protein concentration (µg/ml) is the average of two independent runs.

[026] **Figure 5.** Shows SDS-PAGE gel micrographs of FIX Ab recognition. Micrograph A depicts a coomassie blue staining; Micrograph B depicts recognition of anti-FIX antibody in Western-blot; Micrograph C depicts recognition of anti-γ carboxylation antibody in Western-blot. Lane 1 in A-C was loaded with a sample containing FIX-(CTP)₂. Lane 2 in A-C was

loaded with a sample containing unbound FIX-(CTP)₂. Lane 3 in A-C was loaded with a sample containing a concentrated elution of FIX-(CTP)₂.

[027] **Figure 6.** Shows a graph showing FIX-(CTP)₂ chromogenic activity (sample concentration/O.D.) compared to human normal pool plasma and rhFIX (American
5 Diagnostics).

[028] **Figure 7.** Shows a graph showing the PK profile of purified FIX-CTP-CTP, rhFIX, harvest of FIX-CTP-CTP, and harvest of FIX-CTP.

[029] **Figure 8.** Shows an anti-CTP and anti-gamma carboxylation antibodies Western blots of FIX fused to three, four or five CTPs. FIX-CTP₃, FIX-CTP₄, and FIX-CTP₅ harvests
10 were loaded on 12% Tris-Glycine gel using Precision plus dual color protein marker (Bio-Rad). The SDS-PAGE analysis was performed by Western immuno-blot using anti-CTP polyclonal Ab (Adar Biotech Production) or anti-Gla Ab (American Diagnostica).

[030] **Figure 9.** Shows a coomassie blue detection of FIX-CTP₃, FIX-CTP₄, and FIX-CTP₅. After a purification process utilizing Jacalin column (immunoaffinity purification of
15 glycosylated proteins), FIX-CTP₃, FIX-CTP₄, and FIX-CTP₅ were loaded on 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE was stained by Coomassie blue dye for sample detection.

[031] **Figure 10.** Shows FIX Chromogenic activity. A comparative assessment of the *in vitro* potency of fully purified (HA column) FIX-CTP₃ FIX-CTP₄ and FIX-CTP₅ versus
20 human pool normal plasma was performed using a commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221802). All samples were serially diluted and the potency was assessed by comparing a dose response curve to a reference preparation consisting of normal human plasma.

[032] **Figure 11.** Shows the comparative pharmacokinetic (PK) profile of FIX-CTP₃ FIX-CTP₄ and FIX-CTP₅. FIX concentration in plasma samples were quantified using human FIX
25 Elisa kits (Affinity Biologicals). Pharmacokinetic profile was calculated and is the mean of 3 animals at each time point. Terminal half lives were calculated using PK Solutions 2.0 software.

[033] **Figure 12.** Shows the FIX-CTP₃ SDS-PAGE analysis - Coomassie SDS-PAGE. FIX-CTP₃ γ -carboxylated enriched protein, rhFIX and rFIXa (activated FIX) were loaded on
30 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE Coomassie analysis was performed by staining the gel with Coomassie blue reagent (800 ng of protein) (Figure 12A). A Western immunoblot was performed using 100 ng of protein with anti-human FIX polyclonal Ab (Figure 12B), anti-human gamma carboxylation

monoclonal antibody (American Diagnostics Cat #499, 3570) (Figure 12C), anti-FIX pro-peptide polyclonal Ab (Figure 12D), and anti-CTP polyclonal Ab (Figure 12E).

[034] **Figure 13:** Shows the FIX-CTP₃ chromogenic activity. A comparative assessment of the *in vitro* potency of FIX-CTP₃ harvest and FIX-CTP₃ γ -carboxylated enriched protein, versus human pool normal plasma was performed using a commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221802). FIX-CTP₃ harvest and protein were serially diluted, and the potency was assessed by comparing a dose-response curve to a reference preparation consisting of normal human plasma.

[035] **Figure 14:** Shows the comparative clotting time. An *in vitro* aPTT (activated Partial Thrombin Time Assay) was performed comparing the clotting activity of FIX-CTP₃ to BeneFIX. The proteins were serially diluted and spiked into human FIX-depleted plasma, and the clotting time was evaluated.

[036] **Figure 15.** Shows FIX-CTP₃ comparative PK profile. FIX concentration was quantitated using human FIX ELISA kits (Affinity Biologicals; Cat. # FIX-AG RUO). The pharmacokinetic profile was calculated for each protein and is the mean of 3 animals at each time point.

[037] **Figure 16.** Shows the activity profile parameters. In parallel to PK sampling, FIX-deficient animals administered with either BeneFIX® or FIX-CTP₃, citrated plasma samples, were evaluated for their clotting activity by aPTT assay, which was translated to % activity. The % activity at each collection point was calculated as the current clotting time/clotting time of normal pool mice plasma*100.

[038] **Figure 17.** Shows a first challenge bleeding parameters. FIX-deficient mice were administered a single intravenous injection of 100 IU/Kg of BeneFIX® or rFIX-CTP₃. The tail vein was slightly clipped 48 hours post-dosing and tail vein bleeding time (TVBT) and bleeding intensity (hemoglobin OD) were evaluated. A second bleeding challenge was performed 15 minutes after reaching homeostasis, and the same parameters were measured.

[039] **Figure 18.** Shows a second challenge bleeding parameters. Once the first bleeding described in the legend to Figure 19 was spontaneously or manually stopped, a second bleeding challenge was performed 15 minutes following the first one, and the time and bleeding intensity were re-measured.

[040] **Figure 19.** Shows a diagram illustrating the rFVII-CTP construct (A), rFVII-CTP-CTP construct (B), rFIX-CTP construct (C), and rFIX-CTP-CTP construct (D).

[041] **Figure 20A.** Shows a bar graph showing harvests limited diluted clone transfected and selected cells with FVII-CTP variants in the presence of 5µg/ml of Vitamin K3. The level of FVII was quantified using FVII ELISA (AssayPro).

[042] **Figure 20B.** Shows a bar graph showing harvests of limited diluted transfected and selected cells with FVII-CTP variants in the presence of 5µg of Vitamin K3. activity. FVII activity was quantified using FVII chromogenic activity assay (AssayPro).

[043] **Figure 20C.** Shows a bar graph showing harvests of limited diluted transfected and selected cells with FVII-CTP variants in the presence of 5µg of Vitamin K3. The specific activity of FVII was calculated for each version by dividing the activity value by the harvest FVII concentration.

[044] **Figure 20D.** Shows a graph showing PK profile of FVII, FVII-CTP-CTP, and FVII-CTP harvests.

[045] **Figure 21.** Shows western blots of FVII fused to three, four and five CTPs, detected using anti-FVII, anti-CTP, and anti-gamma carboxylation antibodies. FVII-CTP₃, FVII-CTP₄, and FVII-CTP₅ harvests were loaded on 12% Tris -Glycine gel (*expedion*) using Precision plus dual color protein marker (Bio-Rad). The SDS-PAGE analysis was performed by Western immunoblot using anti-FVII Ab, anti-CTP polyclonal Ab (Adar Biotech Production) or anti-Gla Ab (American Diagnostica).

[046] **Figure 22.** Shows the FVII Activity – Chromogenic activity. A comparative assessment of the *in vitro* potency of HA purified (highly gamma carboxylated fraction) FVII-CTP₃, FVII-CTP₄, and FVII-CTP₅ versus normal human pool plasma was performed using a commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221304). All samples were serially diluted and the potency was assessed by comparing a dose response curve to a reference preparation consisting of normal human plasma.

[047] **Figure 23.** Shows a first comparative pharmacokinetic (PK) profile-FVII 3, 4 and 5 CTPs. FVII-CTP₃, FVII-CTP₄, and FVII-CTP₅ (Group A, B and C, respectively) were administered in a single intravenous injection to Sprague Dawley rats (six rats per treatment) in a dose of 250 µg/kg body weight. Blood samples were drawn retro-orbitally from 3 rats alternately at 0.083, 0.5 2, 5, 8, 24, 48, 72 and 96 hours post dosing. Citrated plasma (0.38%) was prepared immediately after sampling and stored at -20°C until analysis. FVII-CTP₅ demonstrated a superior profile as compared to the two other versions.

[048] **Figure 24.** Shows a second comparative PK profile-FVII 3, 4 and 5 CTPs. FVII-CTP₃, FVII-CTP₄, and FVII-CTP₅ following FVII selection and the HA purification process (Group A, B and C, respectively) were administered in a single intravenous injection to

Sprague Dawley rats (three rats per substance) in a dose of 29.45 µg/kg body weight. Blood samples were drawn retro-orbital at 0.083, 0.5 2, 8, 24, 48, and 72 hours post-dosing. Citrated plasma (0.38%) was prepared immediately after sampling and stored at -20°C until analysis.

[049] **Figure 25.** Shows a schematic diagram of FVII-CTP₃ purification process. Batch 31 was produced for the PK/PD study. Batch 38 was produced for the survival study.

[050] **Figure 26.** Shows an SDS -PAGE and Western blot of Final FVII and FVIIa. 10 µg (Batch 31) or 5 µg (Batch 38) were loaded in each lane of Coomassie stained SDS-PAGE. 1 µg protein was loaded in each lane of Western blot. 1. FVII-CTP₃ polypeptide; 2. Heavy chain, including 3x CTP; 3. Light Chain. All three antibodies detect FVII. FVIIa heavy chain was detected by α-CTP, and light chain is detected with both α-FVII and α-Gla.

[051] **Figure 27.** Shows that FVII-CTP₃ chromogenic activity is enhanced as a result of purification on ceramic hydroxyapatite (HA) column. A comparative assessment of the *in vitro* potency of FVII-CTP₃ harvest, in-process fractions, and purified FVII-CTP₃ versus human pool normal plasma was performed using a commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221304). FVII-CTP₃ harvest and protein were serially diluted and the potency was assessed by comparing a dose-response curve to a reference preparation of normal human plasma.

[052] **Figure 28.** Shows the PK profile of FVIIa-CTP₃ vs. NovoSeven® in FVIII-deficient mice. FVIIa-CTP₃ was produced following FVII selection, HA purification process and activation. FVIIa-CTP₃ or NovoSeven® was administered in a single intravenous injection to FVIII-/- hemophilic mice. Blood samples were drawn retro-orbitally at 0.083, 0.5 2, 8, 24, 48, and 72 hours post-dosing. Citrated plasma (0.38%) was prepared immediately after sampling and stored at -20°C until analysis, and a PK profile was established based on FVIIa clotting activity using a STACLOT commercial kit.

[053] **Figure 29.** Shows that FVIIa-CTP₃ was produced following FVII selection, HA purification process and activation. FVIIa-CTP₃ or NovoSeven® was administered in a single intravenous injection to FVIII-/- hemophilic mice. Blood samples were drawn retro-orbitally at 0.083, 0.5 2, 8, 24, 48, and 72 hours post-dosing. Citrated plasma (0.38%) was prepared immediately after sampling and stored at -20°C until analysis. Thrombin generation parameters were evaluated during the PK experiment, and parameters including maximal amount to peak, amount of thrombin to time point and rate of thrombin generation were evaluated.

[054] **Figure 30.** Shows hemophilic mice survival curves post tail vein transection (TVT). TVT was performed (A) 15 min, (B) 24 hours or (C) 48 hours post administration. Mice

Survival was observed for 24 hours after TVT and recorded every single hour for the first 12 hours, and after 24 hours. Figure 33D summarizes mouse survival as recorded 24 hours post TVT. Control group data (vehicle) is the sum of the 3 experiments with 5 mice/experiment.

[055] **Figure 31.** Shows FVII – 3- CTP and FVII- 5 CTP immune-blots. A) blotted for
5 GLA. B) blotted for FVIIa. C) blotted for CTP.

[056] **Figure 32.** Shows a comparative PK profile-FVII 3 & 5 CTP- from select and HA column purification (FVIIIS vs. FVII HA).

[057] **Figure 33.** Shows a comparative PK profile-FVII 3 & 5 CTP-The second study (IV vs. SC).

10 [058] **Figure 34.** Shows hemophilic mice survival curves post tail vein transection (TVT). TVT was performed 12 hours post SC administration. Mice Survival was observed for 24 hours after TVT and recorded every single hour for the first 12 hours, and after 24 hours.

[059] **Figure 35.** Shows the PK profile of MOD-5014 vs. NovoSeven® following IV or SC administration. A) shows IV administration; B) shows SC administration.

15 [060] **Figure 36.** Shows the PK profile of MOD-5014 (Clone 61 #75, #81) vs. NovoSeven® following single SC administration.

DETAILED DESCRIPTION OF THE INVENTION

[061] In one embodiment, the present invention provides long-acting coagulation factors
20 and methods of producing and using same. In another embodiment, long-acting coagulation factors comprise a carboxy terminal peptide (CTP, also referred to as CTP unit). In another embodiment, long-acting polypeptides which comprise a coagulation factor further comprise a carboxy terminal peptide (CTP) of human Chorionic Gonadotropin (hCG). In another embodiment, CTP acts as a protectant against the degradation of a coagulation factor. In
25 another embodiment, CTP extends the $C_{\max}$ of a coagulation factor. In another embodiment, CTP extends the $T_{\max}$ of a coagulation factor. In another embodiment, CTP extends the circulatory half-life of a coagulation factor. In some embodiments, CTP enhances the potency of a coagulation factor.

[062] In another embodiment, provided herein is a method of extending the biological
30 half-life of a coagulation factor, comprising the step of attaching one to ten CTPs to the carboxy terminus of the coagulation factor, thereby extending the biological half-life of the coagulation factor. In another embodiment, provided herein is a method of extending the biological half-life of a coagulation factor, comprising the step of attaching one to five CTPs to the carboxy terminus of the coagulation factor, thereby extending the biological half-life of

the coagulation factor. In another embodiment, the present invention provides a method for extending the circulatory half-life of a coagulation factor. In another embodiment, the present invention provides a method for increasing the half-life of a coagulation factor. In another embodiment, the present invention provides a method for extending the half-life of a coagulation factor.

[063] Coagulation Factor VII (FVII) is a 444 amino acid glycoprotein (50KDa) secreted by hepatocytes into the bloodstream as an inactive pro-enzyme. Upon tissue injury and exposure to circulating blood, FVII forms a complex with Tissue Factor (TF) which is a true receptor protein to FVII and is expressed by various cells localized in the deeper layers of the vessel wall. The formation of this FVII-TF complex leads to activation of FVII. Activated FVII (FVIIa) initiates the extrinsic coagulation pathway by activating Factor IX and Factor X.

[064] FVII belong to a group of Vitamin K-dependent glycoproteins associated with the coagulation system. Besides FVII, this group consists of Factor IX, Factor X, Protein C and prothrombin. These proteins have similar domain organizations and are synthesized as precursors with an N-terminal propeptide followed by a mature amino acid sequence. The propeptide contains a docking site for gammacarboxylase which converts glutamic acids (Glu) into gamma carboxy glutamic acids (Gla). This domain is followed by two epidermal growth factor-like (EGF) domains, a connecting region (CR) and a C-terminal serine protease domain. Prior to secretion, FVII propeptide is cleaved forming a 406 amino acid single chain zymogen FVII glycoprotein. After secretion, the protein can be activated into a disulfide-linked two chain heterodimer, FVIIa, by cleavage in the CR. The plasma concentration of FVII is 10 nM and approximately 1% circulates in the active form in healthy individuals.

[065] Factor IX (FIX) is a 415 Amino acid (55KDa) glycoprotein; it belongs to a group of vitamin K dependent glycoproteins associated with the coagulation system. FIX has a similar domain organization as factor FVII, Factor X, Protein C and prothrombin that are synthesized as precursors with an N-terminal propeptide followed by a mature amino acid sequence.

[066] FIX is secreted as a single chain molecule that undergoes complex post-transcriptional modifications, many of which are critical to its biochemical and pharmacokinetic properties. Among all the post-transcriptional modifications, 12 glutamic acid residues near the amino terminus of FIX that are gamma carboxylated by the vitamin K-dependent gamma carboxylase are the most crucial ones. Carboxylation is required for the interaction of FIX with the phospholipid surfaces and for optimal FIX activity. The amino terminus propeptide serves as a recognition site for the gamma carboxylase and thus, following gamma carboxylation, it is cleaved off by the Golgi apparatus serine protease

known as Paired basic Amino acid Cleave Enzyme (PACE/Furin). Four additional post-transcriptional modifications might occur at the Golgi apparatus: sulfation of tyrosine 155, phosphorylation of serine 158, O- glycosylation on Ser 63 and on 61 and finally, N-glycosylation on Asn 157 and 16, but were shown not to be necessary for proper activity of
5 FIX.

[067] FIX circulates in the plasma (average concentration of 5 µg/ml) as a single chain inactive zymogen. Upon proteolytic cleavage at two peptide bonds: Arg 145 and Arg 180 by either one or two physiological activators, FVIIa-TF complex or FIXa, the activation peptide is removed, converting FIX to a fully active enzyme consisting of a light and heavy chain
10 held together by a single disulfide bond. The N-terminal light chain contains the non-catalytic gamma carboxyglutamic acid (Gla) and two epidermal growth factor-like domains, while the C-terminal heavy chain contains the trypsin-like catalytic domain of the molecule. FIXa alone is characterized by poor catalytic activity. However when complexed with FVIII, its proteolytic activity increase by 4-5 orders of magnitude towards its natural substrate FX.

[068] In another embodiment, provided herein is a method of extending the biological half-life or a method of improving the area under the curve (AUC) of a coagulation factor, comprising the step of attaching one to ten CTPs to the carboxy terminus of the coagulation factor, thereby extending the biological half-life or improving the AUC of the coagulation factor. In another embodiment, provided herein is a method of extending the biological half-life or a method of improving the area under the curve (AUC) of a coagulation factor, comprising the step of attaching one to five CTPs to the carboxy terminus of the coagulation factor, thereby extending the biological half-life or improving the AUC of the coagulation factor. In another embodiment, provided herein is a method of extending the biological half-life or a method of improving the area under the curve (AUC) of FIX, comprising the step of attaching one to five CTPs to the carboxy terminus of the FIX, thereby extending the biological half-life or improving the AUC of the FIX. In another embodiment, provided herein is a method of extending the biological half-life or a method of improving the area under the curve (AUC) of FVII or FVIIa, comprising the step of attaching one to five CTPs to the carboxy terminus of FVII or FVIIa, thereby extending the biological half-life or improving the AUC of FVII or FVIIa.

[069] In another embodiment, the present invention provides a method of extending the biological half-life of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby extending the biological half-life of said FIX polypeptide. In

another embodiment, the present invention further provides a method of extending the biological half-life of a Factor VIIa (FVIIa) polypeptide, comprising the step of attaching up to five chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa polypeptide, thereby extending the biological half-life of said FVIIa polypeptide. In one embodiment, three chorionic gonadotrophin carboxy terminal peptides (CTPs) are attached to the carboxy terminus of said FVIIa polypeptide. In another embodiment, four chorionic gonadotrophin carboxy terminal peptides (CTPs) are attached to the carboxy terminus of said FVIIa polypeptide. In another embodiment, five chorionic gonadotrophin carboxy terminal peptides (CTPs) are attached to the carboxy terminus of said FVIIa polypeptide.

[070] In another embodiment, the present invention provides a method of improving the area under the curve (AUC) of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby improving the AUC of said FIX polypeptide. In another embodiment, the present invention provides a method of improving the area under the curve (AUC) of a Factor VIIa (FVIIa) polypeptide, comprising the step of attaching up to five chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa polypeptide, thereby improving the AUC of said FVIIa polypeptide. In one embodiment, three chorionic gonadotrophin carboxy terminal peptides (CTPs) are attached to the carboxy terminus of said FVIIa polypeptide. In another embodiment, four chorionic gonadotrophin carboxy terminal peptides (CTPs) are attached to the carboxy terminus of said FVIIa polypeptide. In another embodiment, five chorionic gonadotrophin carboxy terminal peptides (CTPs) are attached to the carboxy terminus of said FVIIa polypeptide.

[071] In another embodiment, a coagulation factor of the invention is a protein. In another embodiment, a coagulation factor of the invention is a peptide. In another embodiment, a coagulation factor of the invention is a polypeptide. In another embodiment, the coagulation factor is an enzyme. In another embodiment, the coagulation factor is a serine protease. In another embodiment, the coagulation factor is a glycoprotein. In another embodiment, the coagulation factor is a transglutaminase. In another embodiment, the coagulation factor is an inactive zymogen. In another embodiment, the coagulation factor is any coagulation factor known to one of skill in the art.

[072] In another embodiment, the coagulation factor is Factor VIII (FVIII). In another embodiment, the coagulation factor is Factor V (FV). In another embodiment, the coagulation

factor is Factor XIII (FXIII). In another embodiment, the coagulation factor is Factor X (FX). In another embodiment, the coagulation factor is fibrin.

[073] In another embodiment, the coagulation factor is Factor VIIa (FVIIa). In another embodiment, the coagulation factor is Factor VII (FVII). In another embodiment, the coagulation factor is Factor IX (FIX). In another embodiment, the coagulation factor is Factor X (FX). In another embodiment, the coagulation factor is Factor XIa (FXIa). In another embodiment, the coagulation factor is Factor XII (FXII). In another embodiment, the coagulation factor is Factor Xa (FXa). In another embodiment, the coagulation factor is Factor Va (FVa). In another embodiment, the coagulation factor is prothrombin. In another embodiment, the coagulation factor is thrombin. In another embodiment, the coagulation factor is Factor XI (FXI). In another embodiment, the coagulation factor is Von Willebrand factor (vWF). In another embodiment, the coagulation factor is Factor VIIIa (FVIIIa). In another embodiment, the coagulation factor is B-deleted Domain FVIII (FVIII_{BDD}). In another embodiment, the coagulation factor is B domain-deleted FVIII (FVIII_{BDD}). In another embodiment, the coagulation factor is Beta domain-deleted FVIII (FVIII_{BDD}). In another embodiment, the coagulation factor is Factor IXa (FIXa). In another embodiment, the coagulation factor is prekallikrein. In another embodiment, the coagulation factor is kallikrein. In another embodiment, the coagulation factor is Factor XIIa (FXIIa). In another embodiment, the coagulation factor is Fibrinogen. In another embodiment, the coagulation factor is thrombomodulin. In another embodiment, the coagulation factor is Factor II (FII).

[074] In another embodiment, the coagulation factor is a glycoprotein. In another embodiment, the coagulation factor is a vitamin K-dependent glycoprotein. In another embodiment, the coagulation factor is a vitamin K-independent glycoprotein.

[075] In another embodiment, the coagulation factor is a recombinant protein. In another embodiment, the coagulation factor is a recombinant glycoprotein. In another embodiment, the coagulation factor is a recombinant glycoprotein FV. In another embodiment, the coagulation factor is a recombinant FVI. In another embodiment, the coagulation factor is a recombinant FVII. In another embodiment, the coagulation factor is a recombinant FVIII. In another embodiment, the coagulation factor is a recombinant FIX. In another embodiment, the coagulation factor is a recombinant FX. In another embodiment, the coagulation factor is a recombinant FXI. In another embodiment, the coagulation factor is a recombinant FXII. In another embodiment, the coagulation factor is a recombinant FvW. In another embodiment, the coagulation factor is a recombinant FII. In another embodiment, the coagulation factor is a recombinant FIXa. In another embodiment, the coagulation factor is a recombinant FXIa. In

another embodiment, the coagulation factor is a recombinant fibrin. In another embodiment, the coagulation factor is a recombinant FVIIa. In another embodiment, the coagulation factor is a recombinant FXa. In another embodiment, the coagulation factor is a recombinant FVa. In another embodiment, the coagulation factor is a recombinant prothrombin. In another embodiment, the coagulation factor is a recombinant thrombin. In another embodiment, the coagulation factor is a recombinant FVIIIa. In another embodiment, the coagulation factor is a recombinant prekallikrein. In another embodiment, the coagulation factor is a recombinant kallikrein. In another embodiment, the coagulation factor is a recombinant FXIIa. In another embodiment, the coagulation factor is any known recombinant coagulation factor. In another embodiment, the coagulation factor comprising a signal peptide is any known recombinant coagulation factor.

[076] In another embodiment, a coagulation factor comprises 1-10 CTP repeats attached to the C-terminus and no CTPs attached to the N-terminus. In another embodiment, a coagulation factor comprises at least one CTP attached to the C-terminus and no CTPs attached to the N-terminus. In another embodiment, a coagulation factor comprising 1-10 CTP repeats attached to the C-terminus and no CTPs attached to the N-terminus is an engineered coagulation factor. In another embodiment, a coagulation factor comprising at least one CTP attached to the C-terminus and no CTPs attached to the N-terminus is an engineered coagulation factor. In another embodiment, a coagulation factor comprising 1-10 CTP repeats attached to the C-terminus and no CTPs attached to the N-terminus is a conjugated coagulation factor. In another embodiment, a coagulation factor comprising at least one CTP attached to the C-terminus and no CTPs attached to the N-terminus is a conjugated coagulation factor.

[077] In one embodiment, the present invention provides a CTP-modified Factor IX (FIX) polypeptide consisting of a FIX polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said CTP-modified FIX polypeptide.

[078] In another embodiment, the present invention further provides a CTP-modified Factor VIIa (FVIIa) polypeptide consisting of a FVIIa polypeptide and five gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa.

[079] In another embodiment, the coagulation factor is a coagulation factor comprising a domain organization similar or identical to the domain organization of FIX, FVII, Factor X, Protein C, or prothrombin. In another embodiment, the coagulation factor is synthesized as a precursor with an N-terminal propeptide. In another embodiment, the coagulation factor as used herein is in an inactive pro-enzyme form. In another embodiment, the coagulation factor

is produced in hepatocytes. In another embodiment, the coagulation factor comprises a docking site for gammacarboxylase which converts glutamic acids (Glu) into gamma carboxy glutamic acids (Gla). In another embodiment, the coagulation factor as used herein is a commercially available coagulation factor.

5 [080] In one embodiment, the nucleic acid sequence encoding Factor VII comprises the following nucleic acid sequence:
 ctcgaggacatggtctcccaggccctcaggctcctctgccttctgcttgggcttcagggtgcctggctgcagtcttcgtaaccaggag
 gaagcccacggcgctctgcaccggcgccggcgcgccaacgcgttcctggaggagctgcggccgggctccctggagagggagtgc
 aaggaggagcagtgctccttcgaggagggccgggagatctcaaggacgggagaggacgaagctgttctggatttctacagtgat
 10 ggggaccagtgtgcctcaagtccatgccagaatgggggctcctgcaaggaccagctccagctctatatctgcttctgcctccctgccttc
 gagggccggaactgtgagacgcacaaggatgaccagctgactgtgtgaacgagaacggcggtgtgagcagtgactgcagtacca
 cacgggcaccaagcgctcctgtcgggtgccacgaggggtactctctgctggcagacggggtgcctgcacacccacagttgaatatcc
 atgtggaaaaatacctattctagaaaaaagaaatgccagcaaaccccaaggccgaattgtggggggcaaggtgtccccaaggggg
 agtgtccatggcaggtcctgttgtgttgatggagctcagttgtgtggggggaccctgatcaacaccatctgggtggtctccgcggcc
 15 cactgtttcgacaaaatcaagaactggaggaacctgatcgcggtgctgggcgagcacgacctcagcgagcacgacgggatgagca
 gagccggcggggtggcgaggtcatcatccccagcacgtacgtccggggcaccaccaaccacgacatcgcgctgctccgctgcac
 cagcccggtggtcctcactgacctgtggtgccccctgcctgcccgaacggacgttctctgagaggacgtggccttcgtgcgttctc
 attggtcagcgggtggggccagctgctggaccgtggcgccacggccctggagctcatggtcctcaacgtgccccggctgatgacc
 aggactgcctgcagcagtcacggaaggtgggagactccccaaatcacggagtacatgttctgtgccggctactcggatggcagca
 20 aggactcctgcaagggggacagtggaggcccacatgccaccactaccggggcacgtggtacctgacgggcatcgtcagctgggg
 ccagggtgcgcaaccgtgggcccactttggggtgtacaccagggtctccagtagatcgagtggtgcaaaagctcatgcgctcaga
 gccacgcccaggagtctcctcgcgagccccatttcctgaggatgcggccgc (SEQ ID NO: 11).

[081] In another embodiment, the amino acid sequence of Factor VII comprises the following amino acid sequence:
 25 MVSQALRLCLLLGLQGCLA AVFVTQEEAHGVLHRRRRANAFLEELRPGSLERECKE
 EQCSFEEAREIFKDAERTKLFWISYSDGDQCASSPCQNGGSKDQLQSYICFLPAFE
 GRNCETHKDDQLICVNENGGCEQYCS DHTGTRSCRCHEGYSLADGV SCTPTVEY
 PCGKPILEKRNASKPQGRIVGGKVC PKGEC PWQVLLL VNGAQLCGGTLINTI W VVS
 AAHCFDKIKNWRNLIAVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTNHDIALRL
 30 HQPVVLTDHVPLCLPERTFSERTLAFVRFSLVSGWGQLLDRGATALELMVLNVPRL
 MTQDCLQQSRKVG DSPNITEYMFCAGYS D GSKD SCKGDSGGPHATHYRGTWYLTGI
 VSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSEPRPGVLLRAPFP (SEQ ID NO: 9).

[082] In another embodiment, the amino acid sequence of Factor VII comprises the following amino acid sequence:

MVSQALRLLCLLLGLQGCLA AVFVTQEEAHGVLHRRRRANAFLEELRPGSLERECKE
EQCSFEEAREIFKDAERTKLFWISYSDGDQCASSPCQNGGSKDQLQSYICFLPAFE
GRNCETHKDDQLICVNENGGCEQYCS DHTGTRSCRCHEGYSLADGV SCTPTVEY
PCGKIPILEKRNASKPQGRIVGGKVC PKGEC PWQVLLL VNGAQLCGGT LINTI WVVS
5 AAHCFDKIKNWRNLIAVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTNHDIALRL
HQPVVLT DHVPLCLPERTFSERTLAFVRFSLVSGWGQLLDRGATALELMVLNVPRL
MTQDCLQQSRKVGDS PNITEYMFCAGYS DSGSKDSC KGDSGGPHATHYRGTWYLTGI
VSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSEPRPGVLLRAPFP*GCGR (SEQ ID
NO: 10).

10 [083] In another embodiment, the nucleic acid sequence encoding Factor VII-CTP
(attached to the carboxy terminus) comprises the following nucleic acid sequence:
ctcgaggacatggtctcccaggccctcaggctcctctgccttctgcttgggcttcagggtgcctggctgcagtcttcgtaaccaggag
gaagcccacggcgtcctgcaccggcgccggcgcccaacgcgttctggaggagctgcggccgggctccctggagagggagtg
aaggaggagcagtgctcctcgaggaggccgggagatctcaaggacgcggagaggacgaagctgttctggatttcttacagtgat
15 ggggaccagtgtgcctcaagtccatgccagaatgggggctcctgcaaggaccagctccagtctatatctgcttctgcctccctgccttc
gagggccggaactgtgagacgcacaaggatgaccagctgatctgtgtaacgagaacggcggctgtgagcagtactgcagtga
cacgggccaacgcgtcctgtcggtgccacgaggggtactctctgctggcagacggggtgctcgcacaccacagttgaatattc
atgtgaaaaatacctattctagaaaaaagaaatgccagcaaaccccaaggccgaattgtggggggcaagggtgtgccccaaagggg
agtgtccatggcaggtcctgtgtgttgatggagctcagttgtgtggggggaccctgatcaacaccatctgggtgttccgcgggc
20 cactgtttcgacaaaatcaagaactggaggaaactgatcgcgggtgctgggcgagcagacacctcagcgagcagcagcgggatgagca
gagccggcggggtggcgaggtcatcatccccagcagctacgtcccgggcaccaccaaccacgacatcgcggtgctccgctgcac
cagcccgtggtcctcactgaccatgtggtgccccctgcctgcccgaacggacgttctctgagaggacgtggccttcgtgcgttctc
attggtcagcggctggggccagctgctggaccgtggcgccacggccctggagctcatggtcctcaacgtgccccggctgatgacc
aggactgcctgcagcagtcacggaaggtgggagactcccaaatatcacggagtacatgttctgtccgggtactcggatggcagca
25 aggactcctgcaagggggacagtgaggagccacatgccaccactaccggggcacgtgtgtacc
tgaccggcatcgtgagctggggccagggtgcgccaccgtggggcacttcggcgtgtacaccagggtgtcccagtagatcagtggtg
ctgcagaaactgatgagaagcgagcccagaccggcggtgctgctgagagccccctccccagcagcagctccaagggccctcccc
tagcctgccagccctagcagactgcctggggccagcgacacccccatcctccccagtgaggatccgcgggcgc (SEQ ID
NO: 12).

30 [084] In another embodiment, the amino acid sequence of Factor VII-CTP (attached to the
carboxy terminus) comprises the following amino acid sequence:
MVSQALRLLCLLLGLQGCLA AVFVTQEEAHGVLHRRRRANAFLEELRPGSLERECKE
EQCSFEEAREIFKDAERTKLFWISYSDGDQCASSPCQNGGSKDQLQSYICFLPAFE
GRNCETHKDDQLICVNENGGCEQYCS DHTGTRSCRCHEGYSLADGV SCTPTVEY

PCGKIPILEKRNASKPQGRIVGGKVC PKGEC PWQVLLL VNGAQLCGGTLINTIWVVS
 AAHCFDKIKNWRNLIAVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTNHDIALRL
 HQPVVLT DHVVPLCLPERTFSERTLAFVRFSLVSGWGQLLDRGATALELMVLNVPRL
 MTQDCLQQSRKVG DSPNITEYMFCAGYS DSGSKGDSGGPHATHYRGTWYLTGI
 5 VSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSEPRPGVLLRAPFPSSSSKAPPPSLPS
 PSRLPGPSDTPILPQ* (SEQ ID NO: 13).

[085] In another embodiment, the nucleic acid sequence encoding Factor VII-CTP-CTP (attached to the carboxy terminus) comprises the following nucleic acid sequence:

ctcaggacatggtctccaggccctcaggctcctctgccttctgcttgggcttcagggtgcctggctgcagtcttcgtaaccaggag
 10 gaagcccacggcgctctgcaccggcgccggcgcccaacgcgttctggaggagctgcggccgggctccctggagagggagtg
 aaggaggagcagtgctcctcagaggggccggagatctcaaggacgggagaggacgaagctgttctggattcttacagtgat
 ggggaccagtgtgcctcaagtccatgccagaatgggggctcctgcaaggaccagctccagctctatatctgcttctgcctccctgccttc
 gagggccggaactgtgagacgcacaaggatgaccagctgatctgtgtaacgagaacggcggtgtgagcagtactgcagtacca
 cacgggcaccaagcgctcctgtcgggtgccacgaggggtactctctgctggcagacggggtgctctgcacaccacagttgaatatcc
 15 atgtggaaaaatacctattctagaaaaaagaaatgccagcaaaccccaaggccgaattgtggggggcaaggtgtgccccaaagggg
 agtgatccatggcaggtcctgtgtgttggaatggagctcagttgtgtggggggaccctgatcaacaccatctgggtggtctccgcggcc
 cactgtttcgacaaaatcaagaactggaggaaactgatcgggtgctgggcgagcacgacctcagcgagcacgacggggatgagca
 gagccggcggggtggcgaggtcatcatccccagcacgtacgtcccgggcaccaccaaccacgacatcgcgctgctccgctgcac
 cagcccgtggctcctcactgacctgtggtgccccctcgtcctgcccgaacggacgttctctgagaggacgtggccttcgtgcgttctc
 20 attggtcagcgggtggggccagctgctggaccgtggcgccacggccctggagctcatggtcctcaacgtgccccggctgatgacct
 aggactgcctgcagcagtcacggaaggtgggagactcccaaatatcacggagtacatgttctgtgccggctactcggatggcagca
 aggactcctgcaagggggacagtggaggcccatgccaccactaccggggcacgtggtacctgaccggcatcgtgagctgggg
 ccagggtgcgccaccgtgggccacttcggcggtgacaccagggtgtcccagtcacagtggtgcagaaactgatgagaagcg
 agccagacccggcggtgctgctgagagcccccttccccagcagcagctccaaggccccctccccctagcctgccagccctagcaga
 25 ctgcctgggcccctccgacacaccaatcctgccacagagcagctcctctaaggccccctcctccatccctgccatccccctcccggtgc
 caggccccctctgacaccctatcctgcctcagtgatgaaggtctggtatccggcgccgc (SEQ ID NO: 14).

[086] In another embodiment, the amino acid sequence of Factor VII-CTP-CTP (attached to the carboxy terminus) comprises the following amino acid sequence:

MVSQALRLLCLLLGLQGCLA AVFVTQEEAHGVLHRRRRANAFLEELRPGSLERECKE
 30 EQCSFEEAREIFKDAERTKLFWISYSDGDQCASSPCQNGGSKDQLQSYICFLPAFE
 GRNCETHKDDQLICVNENGGCEQYCS DHTGTRSCRCHEGYSLLADGVSCTPTVEY
 PCGKIPILEKRNASKPQGRIVGGKVC PKGEC PWQVLLL VNGAQLCGGTLINTIWVVS
 AAHCFDKIKNWRNLIAVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTNHDIALRL
 HQPVVLT DHVVPLCLPERTFSERTLAFVRFSLVSGWGQLLDRGATALELMVLNVPRL

MTQDCLQQSRKVGDSPNITEYMFCAGYSDGSKDSCKGDSGGPHATHYRGTWYLTGI
VSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSEPRPGVLLRAPFPSSSSKAPPPSLPS
PSRLPGPSDTPILPQSSSSKAPPPSLPSRLPGPSDTPILPQ** (SEQ ID NO: 15).

[087] In another embodiment, the nucleic acid sequence encoding Factor VII-CTP-CTP-
CTP (attached to the carboxy terminus) comprises the following nucleic acid sequence:
ctcaggacatggtctcccaggccctcaggctcctctgccttctgcttgggcttcagggtgcctggctgcagtcttcgtaaccaggag
gaagccacggcgctctgcaccggcgccggcgcccaacgcgtcctggaggagctgcggccgggctccctggagagggagtg
aaggaggagcagtgctccttcgaggaggccgggagatctcaaggacgcggagaggacgaagctgttctggatttcttacagtgat
ggggaccagtgtgctcaagtccatgccagaatgggggctcctgcaaggaccagctccagctctatatctgcttctgcctccctgccttc
gagggccggaactgtgagacgcacaaggatgaccagctgatctgtgtaacgagaacggcggctgtgagcagtactgcagtacca
cacgggcaccaagcgctcctgtcgggtgccacgaggggtactctctgctggcagacggggtgctcgcacaccacagttgaatatcc
atgtggaaaaatctattctagaaaaaagaaatgccagcaaaccccaaggccgaattgtggggggcaaggtgtgccccaaagggg
agtgtccatggcaggtcctgttgttgggaatggagctcagttgtgtggggggaccctgatcaacacctctgggtgtctccgcggcc
cactgtttcgacaaaatcaagaactggaggaaactgatcgcggtgctgggcgagcacgacctcagcgagcagcagcgggatgagca
gagccggcgggtggcgaggtcatcatccccagcacgtacgtccggggcaccaccaaccacgacatcgcgctgctccgcctgcac
cagcccggtgctcactgacctgtggtgcccctctgctgcccgaacggagcttctctgagaggacgctggccttcgtgcgcttctc
attggtcagcgggtggggccagctgctggaccgtggcgccacggccctggagctcatggtcctcaacgtgccccggctgatgacc
aggactgcctgcagcagtcacggaaggtgggagactccccaatatcacggagtacatgttctgtgccgggtactcggatggcagca
aggactcctgcaagggggacagtgaggcccacatgccaccactaccggggcacgtggtacctgaccggcatcgtgagctgggg
ccagggtgcgccaccgtgggccacttcggcgtgtacaccagggtgtcccagtacatcagtggtgcagaaactgatgagaagcg
agccagacccggcgtgctgctgagagcccccttccccagcagcagctccaagccccctccctagcctgccagccctagcaga
ctgctggggccagtgacaccctatcctgctcagtcacgtccagcaaggccccccccctagcctgcttctcttctcggtgcct
ggccccagcgatactcaattctgccccagtcctccagcagtaaggctccccctccatctctgccatccccagcagactgccaggcc
cttctgatacacccatctcccacagtgatgaggatccggcgccgcttaattaa (SEQ ID NO: 24).

[088] In another embodiment, the amino acid sequence of Factor VII-CTP-CTP-CTP
(attached to the carboxy terminus) comprises the following amino acid sequence:
MVSQALRLLCLLLGLQGCLA AVFVTQEEAHGVLHRRRRANAFLEELRPGSLERECKE
EQCSFEEAREIFKDAERTKLFWISYSDGDQCASSPCQNGGSKDQLQSYICFLPAFE
GRNCETHKDDQLICVNENGGCEQYCS DHTGTRSCRCHEGYSLADGV SCTPTVEY
PCGKIPILEKRNASKPQGRIVGGKVC PKGEC PWQVLLL VNGAQLCGGTLINTI WVVS
AAHCFDKIKNWRNLI AVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTNHDIALRL
HQPVVLT DHVPLCLPERTF SERTLAFVRFSLVSGWGQLLDRGATALELMVLNVPRL
MTQDCLQQSRKVGDSPNITEYMFCAGYSDGSKDSCKGDSGGPHATHYRGTWYLTGI
VSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSEPRPGVLLRAPFPSSSSKAPPPSLPS

PSRLPGPSDTPILPQSSSSKAPPPSLPSPSRLPGPSDTPILPQSSSSKAPPPSLPSPSRLPGP
SDTPILPQ** (SEQ ID NO: 25).

[089] In another embodiment, the nucleic acid sequence encoding Factor VII-(CTP)₄ (attached to the carboxy terminus) comprises the following nucleic acid sequence:

5 ctcgaggacatggtctcccaggccctcaggctcctctgccttctgcttgggcttcagggtgcctggctgcagtcttcgtaaccaggag
gaagcccacggcgtcctgcaccggcgccggcgcccaacgcgttctggaggagctgcggccgggctccctggagagggagtgc
aaggaggagcagtgctccttcgaggaggccgggagatctcaaggacgcggagaggacgaagctgttctggattcttacagtgat
ggggaccagtgtgcctcaagtccatgccagaatgggggctcctgcaaggaccagctccagtctatatctgcttctgcctccctgccttc
gagggccgggaactgtgagacgcacaaggatgaccagctgatctgtgtgaacgagaacggcggtgtgagcagtactgcagtacca
10 cacgggcaccaagcgctcctgtcgggtgccacgaggggtactctctgctggcagacggggtgctcgcacaccacagttgaatatcc
atgtggaaaaatacctattctagaaaaaagaaatgccagcaaaccccaaggccgaattgtggggggcaaggtgtgccccaaagggg
agtgtccatggcaggtcctgttgttggtaatggagctcagttgtgtggggggaccctgatcaacaccatctgggtgtctccgcggcc
cactgtttcgacaaaatcaagaactggaggaaacctgatcgcggtgctgggcgagcacgacctcagcgagcacgacgggatgagca
gagccggcggggtggcgaggtcatcctccagcacgtacgtccggggcaccaccaaccagacatcgcgctgctccgcctgcac
15 cagcccgtggtcctcactgaccatgtggtgccccctgcctgcccgaacggacgttctctgagaggacgtggccttcgtgcgttctc
attggtcagcggtggggccagctgctggaccgtggcgccacggccctggagctcatggtcctcaacgtgccccggctgatgacc
aggactgcctgcagcagtcacggaaggtgggagactcccaaatatcacggagtacatgttctgtccgggtactcggatggcagca
aggactcctgcaagggggacagtggaggcccatgccaccactaccggggcacgtggtacctgaccggcatcgtgagctgggg
ccagggctgcgccaccgtgggccacttcggcgtgtacaccagggtgtccagtagcatcagtggtcgcagaaactgatgagaagcg
20 agcccagaccggcgtgctgctgagagcccccttcccagcagcagctccaagccccctcccctagcctgccagccctagcaga
ctgctgggcccagtgacaccctatcctgcctcagtcagctccagcaaggccccccccctagcctgccttctccttctcggtgcct
ggccccagcgatactccaattctgccccagtcctccagcagtaaggctccccctccatctctgccatccccagcagactgccaggcc
cttctgatacccatctccacagtgatgaggatccgc (SEQ ID NO: 26).

[090] In another embodiment, the amino acid sequence of Factor VII-(CTP)₄ (attached to
25 the carboxy terminus) comprises the following amino acid sequence:
LEDMVSQALRLLCLLLGLQGCLA AVFVTQEEAHGVLHRRRRANAFLEELRPGSLERE
CKEEQCSFEEAREIFKDAERTKLFWISYSDGDQCASSPCQNGGSCKDQLQSYICFCLP
AFEGRNCETHKDDQLICVNENGGCEQYCS DHTGTRKSCRCHEGYSLLADGVSTPT
VEYPCGKIPILEKRNASKPQGRIVGGKVC PKGEC PWQVLLL VNGAQLCGGTLINTIW
30 VVSAAHCFDKIKNWRNLIAVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTNHDIAL
LRLHQPVVLT DHVVPLCLPERTFSERTLAFVRFSLVSGWGQLLDRGATALELMVLNV
PRLMTQDCLQQSRKVGDS PNITEYMFCAGYSDGSKDSCKGDSGGPHATHYRGTWYL
TGIVSWGQGCATVGHFGVYTRVSQYIEWLQKL MRSEPRPGVLLRAPFPSSSSKAPPPS

LPSPSRLPGSDTPILPQSSSSKAPPPSLPSPSRLPGSDTPILPQSSSSKAPPPSLPSPSRLP
GPSTDPILPQ**G (SEQ ID NO: 27).

[091] In another embodiment, the nucleic acid sequence encoding Factor VII-(CTP)₅ (attached to the carboxy terminus) comprises the following nucleic acid sequence:

5 ctcgaggacatggtctcccaggccctcaggctcctctgccttctgcttgggcttcagggtgcctggctgcagtcttcgtaaccaggag
gaagcccacggcgctctgcaccggcgccggcgcgccaacgcgttctggaggagctgcggccgggctccctggagagggagtgc
aaggaggagcagtgctccttcgaggaggccgggagatctcaaggacgcggagaggacgaagctgttctggatttctacagtgat
ggggaccagtgtgcctcaagtccatgccagaatgggggctcctgcaaggaccagctccagtctatatctgcttctgcctccctgccttc
gagggccgggaactgtgagacgcacaaggatgaccagctgatctgtgtgaacgagaacggcggtgtgagcagtactgcagtacca
10 cacgggcaccaagcgctcctgtcgggtgccacgaggggtactctctgctggcagacggggtgctctgcacaccacagttgaatatcc
atgtggaaaaatacctattctagaaaaaagaaatgccagcaaaccccaaggccgaattgtggggggcaaggtgtgccccaaagggg
agtgtccatggcaggtcctgttgttggatggagctcagttgtgtggggggaccctgatcaacaccatctgggtgtctccgcggcc
cactgtttcgacaaaatcaagaactggaggaaacctgatcgcggtgctgggcgagcacgacctcagcgagcacgacgggatgagca
gagccggcggggtggcgaggtcatcctccagcacgtacgtccggggcaccaccaaccagacatcgcgctgctccgcctgcac
15 cagcccgtggtcctcactgaccatgtggtgccccctgcctgcccgaacggagcttctctgagaggacgtggccttcgtgcgttctc
attggtcagcggtggggccagctgctggaccgtggcgccacggccctggagctcatggtcctcaacgtgccccggctgatgaccc
aggactgcctgcagcagtcacggaaggtgggagactcccaaatatcacggagtacatgttctgtccgggtactcggatggcagca
aggactcctgcaagggggacagtgaggggccacatgccaccactaccggggcacgtggtacctgaccggcatcgtgagctgggg
ccagggctgcgccaccgtgggccacttcggcgtgtacaccagggtgtccagtagcatcagtggtcgcagaaactgatgagaagcg
20 agcccagaccggcgctgctgctgagagccccctccccagcagcagctccaagggccctccccctagcctgccagccctagcaga
ctgctggggccctctgacaccctatcctgctcagtcagctccttaaggctccaccacctccctgcttagccctcaagactgcc
ggccctagcgatacacaattctgccccagctcctccagcagcaaggctccccacctagcctgccttctccatcaaggctgcctggcc
catccgatacccaatttgcctcagagcagctctagcaaggcacctccccccagctgtccctctccaagcagactccctggccctca
gacactccaatcctccacagtcctctagctctaaagctccacctccagcctgccagccctagtagactccccggaccttctgatacc
25 cccatcttggcccagtgatgaggatccgc (SEQ ID NO: 28).

[092] In another embodiment, the amino acid sequence of Factor VII-(CTP)₅ (attached to the carboxy terminus) comprises the following amino acid sequence:

LEDMVSQALRLLCLLLGLQGCLAAVFVTQEEAHGVLHRRRRANAFLEELRPGSLERE
CKEEQCSFEEAREIFKDAERTKLFWISYSDGDQCASSPCQNGGSKDQLQSYICFLP
30 AFEGRNCETHKDDQLICVNENGGCEQYCSHTGTGRSCRCHEGYSLLADGVSTPT
VEYPCGKIPILEKRNASKPQGRIVGGKVC PKGEC PWQVLLL VNGAQLCGGT LINTIW
VVSAAHCFDKIKNWRNLIAVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTNHDIAL
LRLHQPVVLT DHVVPLCLPERTFSERTLAFVRFSLVSGWGQLLDRGATALELMVLNV
PRLMTQDCLQQSRKVGDS PNITEYMFCAGYSDGSKDSCKGDSGGPHATHYRG TWYL

TGIVSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSEPRPGVLLRAPFPSSSSSKAPPPS
 LPSPSRLPGPSDTPILPQSSSSKAPPPSLPSPSRLPGPSDTPILPQSSSSKAPPPSLPSPSRLP
 GPSDTPILPQSSSSKAPPPSLPSPSRLPGPSDTPILPQSSSSKAPPPSLPSPSRLPGPSDTPIL
 PQ**GS (SEQ ID NO: 29).

- 5 [093] In another embodiment, the nucleic acid sequence encoding Factor IX comprises the following nucleic acid sequence:
- gcgatgccatgcagcgcgtgaacatgatcatggcagaatcaccaggcctcatcaccattgccttttaggatctactcagtgcgaat
 gtacagttttctgatcatgaaaacccaacaaaattctgaatcgccaaagaggtataattcaggtaaattggaagagttgtcaaggg
 aaccttgagagagaatgatggaagaaaagtgtagtttgaagaagcacgagaagttttgaaaacactgaaagaacaactgaatttgg
 10 aagcagtatgttgatggagatcagtgtagtccaatccatgtttaaattggcggcagttgcaaggatgacattaattcctatgaattgttg
 tcccttggattgaaggaaagaactgtgaattagatgaacatgaacattaagaatggcagatgcgagcagtttgtaaaaatagtctg
 ataacaagggtgttctctgtactgagggatcgcactgcagaaaaccagaagtcctgtgaaccagcagtgccatttccatgtgaa
 gagtttctgttcacaaacttctaagctcaccgtgctgagactgttttctgatgtggactatgaaattctactgaagctgaaccattt
 gataacatcactcaaagcaccaatcatttaatgacttcactcagtggtgttggtggagaagatgccaaaccagggtcaattcccttggcag
 15 gttgttttgaatggtaaagtgtgacattctgtggaggctctatcgtaataaaaaatggattgtaactgctgccactgtgtgaaactggtg
 ttaaaattacagttgtcgcaggtgaacataatattgaggagacagaacatacagagcaaaagcgaaatgtgattcgaattattcctcacca
 caactacaatgcagctattaataagtacaacctgacattgcccttctggaactggacgaacccttagtgctaaacagctacgttacacct
 atttgactgtgacaaggaatacacgaacatcttctcaaatggatctggctatgtaagtggctggggaagagcttccacaaagggga
 gatcagcttttagttccagtagcttaggttccactgttgaccgagccacatgtcttcgatctacaaagttcaccatctataacaacatgtt
 20 ctgtgctggcttccatgaaggaggtagagattcatgtcaaggagatagtgggggaccccatgttactgaagtggaagggaccagtttct
 taactggaattattagctggggtgaagagtgatgcaatgaaaggcaaatatggaatatataccaaggtatcccggtatgtcaactggattaa
 ggaaaaaacaagctcactgaacgcggccgc (SEQ ID NO: 16).

- [094] In another embodiment, the amino acid sequence of Factor IX comprises the following amino acid sequence:
- 25 MQRVNMIMAESPLITICLLGYLLSAECTVFLDHENANKILNRPKRYNSGKLEEFVQG
 NLERECMEEKCSFEEAREVFENTERTTEFWKQYVDGDQCESNPCLNGGSKDDINSY
 ECWCPFGFEGKNCELDVTCNIKNGRCEQFCKNSADNKVVCSCTEGYRLAENQKSCE
 PAVPFPCGRVSVSQTSKLTRAETVFPDVDYVNSTEAETILDNITQSTQSFNDFTRVVG
 GEDAKPGQFPWQVVLNGKVDAFCGGSIVNEKWIVTAAHCVETGVKITVVAGEHNIE
 30 ETEHTEQKRNVIRIIPHHNYNAANKYNHDIALLELDEPLVLNSYVTPICIADKEYTNIF
 LKFGSGYVSGWGRVFHKGRSALVLQYLRVPLVDRATCLRSTKFTIYNNMFCAGFHE
 GGRDSCQGDSGGPHVTEVEGTSFLTGIISWGEECAMKGKYGIYTKVSRVYNWIKEKT
 KLT* (SEQ ID NO: 17).

[095] In another embodiment, the nucleic acid sequence encoding Factor IX-CTP (attached to the carboxy terminus) comprises the following nucleic acid sequence:

gcgatcgccatgcagcgctgaacatgatcatggcagaatcaccaggcctcatcaccatctgccttttaggatattactcagtgtgaa
 gtacagtttttctgatcatgaaaacgccaacaaaattctgaatcgccaaagaggtataattcaggtaaattggaagagttgttcaagg
 5 gaaccttgagagagaatgtatggaagaaaagtgtagtttgaagaagcacgagaagttttgaaaacactgaaagaacaactgaattttg
 gaagcagtattgtatggagatcagtgtagtccaatccatgttfaatggcggcagttgcaaggatgacattaattcctatgaatgttgt
 gtccctttggattgaaaggaaagaactgtgaattagatgaacatgaacattaagaatggcagatgcgagcagttttgaaaaatagtgt
 gataacaaggtggttctctgtactgagggatatcgactgcagaaaaccagaagtctgtgaaccagcagtgccattccatgtgga
 agagtttctgttcacaaacttctaagctcacccgtgctgagactgttttctgatgtggactatgtaatttactgaagctgaaaccattt
 10 ggataacatcactcaaagcacccaatcatttaatagacttcactcgagttgtgtggaggagaagatgccaaccagggtcaattcccttgga
 gggtgtttgaatggtaaagttgatgcattctgtggaggctctatcgtaataaaaaatggattgaactgtgcccactgtgtgaaactgg
 gtaaaattacagttgtcgcaggtgaacataatattaggagacagaacatacagagcaaaagcgaaatgtgatcgaattattcctcac
 cacaactacaatgcagctattaataagtacaacatgacattgcccttctggaactggacgaacccttagtgctaaacagctacgttacac
 ctatttgattgtgacaaggaatacacgaacatcttctcaaaatttgatctggctatgtaagtggctggggaagagttccacaaagg
 15 gagatcagcttagttcttcagctaccttagagttccactgttgaccgagccacatgtcttcgatctacaaagttcaccatctataacaacatg
 ttctgtgctggttccatgaaggaggtagagattcatgtcaaggagatagtgggggaccccatgttactgaagtgaaggaggaccagttt
 cttactggaattattagctggggtgaagagtgcaatgaaaggcaaatatggaatatataccaaggtatcccggtatgcaactggatt
 aaggaaaaacaaagctcactagctccagcagcaaggccccctccccgagcctgccctccccaagcaggctgcctgggccccccga
 cacaccaatcctgccacagtgtgaaggtctgatccgcggccgc (SEQ ID NO: 18).

[096] In another embodiment, the amino acid sequence of Factor IX-CTP (attached to the carboxy terminus) comprises the following amino acid sequence:

MQRVNMIMAESPLITICLLGYLLSAECTVFLDHENANKILNRPKRYNSGKLEEFVQG
 NLERECMEEKCSFEEAREVFENTERTTEFWKQYVDGDQCESNPCLNGGSKDDINSY
 ECWCPFGFEGKNCELDVTCNIKNGRCEQFCKNSADNKVVCSCTEGYRLAENQKSCE
 25 PAVPFPCGRVSVSQTSLTRAETVFPDVDYVNSTEAETILDNITQSTQSFNDFTRVVG
 GEDAKPGQFPWQVVLNGKVDAFCGGSIVNEKWIVTAAHCVETGVKITVVAGEHNIE
 ETEHTEQKRN VIRIIPHHNYNA AINKYNHDIALLELDEPLVLNSYVTPICIADKEYTNIF
 LKFGSGYVSGWGRVFHKGRSALVLQYLRVPLVDRATCLRSTKFTIYNNMFCAGFHE
 GGRDSCQGDSGGPHVTEVEGTSFLTGIISWGEECAMKGKYGITYTKVSRYVNWIKEKT
 30 KLTSSSSKAPPPSLPSPSRLPGPSDTPILPQ** (SEQ ID NO: 19).

[097] In another embodiment, the nucleic acid sequence encoding Factor IX-CTP-CTP (attached to the carboxy terminus) comprises the following nucleic acid sequence:

gcgatcgccatgcagcgctgaacatgatcatggcagaatcaccaggcctcatcaccatctgccttttaggatattactcagtgtgaa
 gtacagtttttctgatcatgaaaacgccaacaaaattctgaatcgccaaagaggtataattcaggtaaattggaagagttgttcaagg

gaaccttgagagagaatgtatggaagaaaagtgtagtttgaagaagcacgagaagttttgaaaacactgaaagaacaactgaattttg
 gaagcagtatgttgatggagatcagtgtagtccaatccatgtttaaattggcggcagttgcaaggatgacattaattcctatgaattgggt
 gtccctttggattgaaggaaagaactgtgaattagatgtaacatgtaacattaagaatggcagatgcgagcagttttgaaaaatagtgt
 gataacaagggtggttctcctgtactgagggatcgcactgcagaaaaccagaagtcctgtgaaccagcagtgccatttccatgtgga
 5 agagtttctgttccaaaacttctaagctcacccgtgctgagactgttttctctgatgtggactatgtaattctactgaagctgaaaccattt
 ggataacatcactcaaaagcacccaatcatttaatacttctcagtggtgtgggagaagatgccaaccagggtcaattcccttggca
 ggtgttttgaaatggtaaagtgtgacattctgtggaggctctatcgttaataaaaaatggattgtaactgctgccactgtgtgaaactggt
 gttaaaattacagttgtcgcaggtgaacataataggagagacagaacatacagagcaaaagcgaaatgtattcgaattattcctcac
 cacaactacaatgcagctattaataagtacaacatgacattgcccttctggaactggacgaacccttagtgtaaacagctacgttacac
 10 ctatttgcatgtctacaaggaatacacgaacatcttctcaaatgtgatctggctatgtaagtggctggggaagagcttccacaaaagg
 agatcagctttagttctcagtagcttagagttccactgttgaccgagccacatgtcttcgatctacaaagttcaccatctataacaacatgt
 tctgtgctggcttccatgaaggaggtagagattcatgtcaaggagatagtgggggacccatgttactgaagtgaaggaccagttc
 ttaactggaattattagctgggggtgaagagtggtcaatgaaaggcaaatatggaatatataccaaggtatcccggtatgtcaactggatta
 agggaaaaaacaagctcactagctccagcagcaagggccctccccgagcctgccctcccaagcaggtgcctgggcccctccgac
 15 acaccaatcctgccacagagcagctcctctaaggccctcctccatccctgccatccccctcccggtgcctggcccctctgacacccc
 tatcctgcctcagtgatgaaggtctggatccgcgccgc (SEQ ID NO: 20).

[098] In another embodiment, the amino acid sequence of Factor IX-CTP-CTP (attached
 to the carboxy terminus) comprises the following amino acid sequence:
 MQRVNMIMAESPLITICLLGYLLSAECTVFLDHENANKILNRPKRYNSGKLEEFVQG
 20 NLERECMEEKCSFEEAREVFENTERTTEFWKQYVDGDQCESNPCLNGGSKDDINSY
 ECWCPFGFEGKNCELDVTCNIKNGRCEQFCKNSADNKVVCSTEGYRLAENQKSCE
 PAVPFPCGRVSVSQTSLTRAETVFPDVDYVNSTEAETILDNITQSTQSFNDFTRVVG
 GEDAKPGQFPWQVVLNGKVDAFCGGSIVNEKWIVTAAHCVETGVKITVVAGEHNIE
 ETEHTEQKRN VIRIIPHHNYNAANKYNHDIALLELDEPLVLNSYVTPICIADKEYTNIF
 25 LKFGSGYVSGWGRV FHKGRSALVLQYLRVPLVDRATCLRSTKFTIYNMFCAGFHE
 GGRDSCQGD SGGPHVTEVEGTSFLTGIISWGEECAMKGKYGIYTKVSRYVNWIKEKT
 KLTSSSSKAPPPSLPSPSRLPGPSDTPILPQSSSSKAPPPSLPSPSRLPGPSDTPILPQ**
 (SEQ ID NO: 21).

[099] In another embodiment, the nucleic acid sequence encoding Factor IX-(CTP)₃
 30 (attached to the carboxy terminus) comprises the following nucleic acid sequence:
 tctagagtcgaccccgccatgcagcgcgtgaacatgatcatggcagaatcaccaggcctcatcaccatctgccttttaggatctactc
 agtgctgaatgtacagttttcttgatcatgaaaacccaacaaaattctgaatcgccaagaggtataattcaggtaaattggaagagtt
 tgttcaagggaaccttgagagagaatgtatggaagaaaagtgtagtttgaagaagcacgagaagttttgaaaacactgaaagaacaa
 ctgaattttggaagcagtatgttgatggagatcagtgtagtccaatccatgtttaaattggcggcagttgcaaggatgacattaattcctat

gaatgttggtgtcccttggatttgaaggaaagaactgtgaattagatgaacatgaacattaagaatggcagatgcgagcagtttgtaa
 aaatagtctgataacaagggtggttgcctgtactgagggatcgactgcagaaaaccagaagtcctgtgaaccagcagtgccatt
 tccatgtggaagagttctgtttcacaacttctaagctcacccgtgctgaggcagttttcctgatgtggactatgtaattctactgaagct
 gaaaccattttggataacatcactcaaagcacccaatcatttaatactcactcgagttgttggtggagaagatgccaaccaggtcaat
 5 tcccttggcaggttgttgaatggtaaagttgatgcattctgtggaggtctatcgtaataaaaaatggattgtaactgctgccactgtgt
 tgaactgggtgttaaaattacagttgtcgcaggtgaacataattgaggagacagaacatacagagcaaaagcgaaatgtgattcgaat
 tattcctcaccacaactacaatgcagctattaataagtacaacctgacattgcccttctggaactggacgaacccttagtctaaacagct
 acgttacacctatttgcattgctgacaaggaatacacgaacatctcctcaaatttggatctggctatgtaagtggctggggaagagtctc
 cacaaggagatcagctttagtctcagctaccttagagttccactgttgaccgagccacatgtcttcgatctacaaagttcacctctat
 10 aacaacatgttctgtgctggcttccatgaaggaggtagagattcatgtcaaggagatagtgggggaccccatgttactgaagtgaagg
 gaccagtttctaactggaattattagctggggtgaagagtggtcaatgaaaggcaaatatggaatatataccaaggtatcccggtatgtc
 aactggattaaggaaaaaacaagctcactagctccagcagcaaggccccctccccgagcctgccctccccagcagggtgcctgg
 gcccagtgacaccctatcctgcctcagctccagcagcaaggccccacccctagcctgccttctccttctcggtgcctggcccca
 gcgatactcaattctgccccagctcctccagcagtaaggtccccctccatctgcccacccccagcagactgccaggcccttctgata
 15 caccatectcccacagtgatgaggatccggccgc (SEQ ID NO: 30).

[0100] In another embodiment, the amino acid sequence of Factor IX-(CTP)₃ (attached to
 the carboxy terminus) comprises the following amino acid sequence:
 MQRVNMIMAESPLITICLLGYLLSAECTVFLDHENANKILNRPKRYNSGKLEEFVQG
 NLERECMEEKCSFEEAREVFENTERTTEFWKQYVDGDQCESNPCLNGGSKDDINSY
 20 ECWCPFGFEGKNCELDVTCNIKNGRCEQFCKNSADNKVVCSTEGYRLAENQKSCE
 PAVPFCGRVSVSQTSLTRAFAVFPDVEDYVNSTEAETILDNITQSTQSFNDFTRVVG
 GEDAKPGQFPWQVVLNGKVDAFCGGSIVNEKWIVTAAHCVETGVKITVVAGEHNIE
 ETEHTEQKRNIRIIPHHNYNAANKYNHDIALLEPLVLNSYVTPICIADKEYTNIF
 LKFGSGYVSGWGRVFKGRSALVLQYLRVPLVDRATCLRSTKFTIYNMFCAGFHE
 25 GGRDSCQGDGSGPHVTEVEGTSFLTGIISWGEECAMKGKYGIYTKVSRVYNWIKEKT
 KLTSSSSKAPPPSLPSPSRLPGPSDTPILPQSSSSKAPPPSLPSPSRLPGPSDTPILPQSSSS
 KAPPPSLPSPSRLPGPSDTPILPQ** (SEQ ID NO: 31).

[0101] In another embodiment, the nucleic acid sequence encoding Factor IX-(CTP)₄
 (attached to the carboxy terminus) comprises the following nucleic acid sequence:
 30 tctagatgcaccccgccatgcagcgcgtgaacatgatcatggcagaatcaccagcctcatcaccatctgccttttaggatatctactc
 agtgctgaatgtacagttttcttgatcatgaaaacccaacaaaattctgaatcgccaaagaggtataattcaggtaaattggaagagt
 tgtcaagggaaccttgagagagaatgtatggaagaaaagttagtttgaagaagcacgagaagttttgaaacactgaaagaacaa
 ctgaatttggaaagcagtatgttgatggagatcagtgtagtccaatccatgtttaaattggcggcagtgcaaggatgacattaattcctat
 gaatgttggtgtcccttggatttgaaggaaagaactgtgaattagatgaacatgaacattaagaatggcagatgcgagcagtttgtaa

aaatagtctgataacaaggtggttctctgtactgaggatatcgacttcagaaaaccagaagtcctgtgaaccagcagtgccatt
 tccatgtggaagagtttctgtttcacaaacttctaagctcacccgtgctgaggcagttttctctgatgtggactatgtaaattctactgaagct
 gaaaccattttggataacatcactcaaagcacccaatcatttaatagacttcactcgagttgttggtggagaagatgcaaaccaggtcaat
 tcccttggcaggtgtttgaatggtaaagttgatgcattctgtggaggctctatcgtaataaaaaatggattgtaactgctgccactgtgt
 5 tgaactgggtgttaaattacagttgtcgcaggtgaacataatattgaggagacagaacatacagagcaaaagcgaatgtgattcgaat
 tattctcaccacaactacaatgcagctattaataagtacaacctgacattgcccttctggaactggacgaacccttagtgctaaacagct
 acgttacacctatttgcattgctgacaaggaatacgaacatcttctcaaatttggatctggctatgtaagtggctggggaagagcttctc
 cacaaggagatcagcttagttcttcagtagcttagagttccactgttgaccgagccacatgtcttcgatctacaaagttcaccatctat
 aacaacatgttctgtgctggttccatgaaggaggtagagattcatgtcaaggagatagtgggggaccccatgttactgaagtgaagg
 10 gaccagtttcttaactggaattattagctggggtgaagagtggtcaatgaaaggcaaatatggaatatatacagggtatcccggtatgtc
 aactggattaaggaaaaaacaagctcactagctccagcagcaaggccctccccgagcctgccctccccaaagcagggtgcctgg
 gccctctgacacccctatctgcctcagtcagctcctctaaggccccaccaccttccctgcctagccctcaagactgccaggccctag
 cgatacaccaattctgccccagtcctccagcagcaaggctccccacctagcctgccttctccatcaaggctgcctggcccatccgata
 cccaattttgcctcagagcagctctagcaaggcacctccccagctgcctctccaagcagactccctggcccttcagacactccc
 15 attctgccacagtgtgaggatccgcggccgc (SEQ ID NO: 32).

[0102] In another embodiment, the amino acid sequence of Factor IX-(CTP)₄ (attached to
 the carboxy terminus) comprises the following amino acid sequence:
 SRVDPAMQRVNMIMAESPGLITICLLGYLLSAECTVFLDHENANKILNRPKRYNSGKL
 EEVQGNLERECMEEKCSFEEAREVFENTERTEFWKQYVDGDQCESNPCLNGGSC
 20 KDDINSYECWCPFGFEGKNCELDVTCNIKNRCEQFCKNSADNKVVCSCTEGYRLA
 ENQKSCEPAVFPFCGRVSVSQTSLTRAFAVFPDVDYVNSTEAETILDNITQSTQSFN
 DFTRVVGGEDAKPGQFPWQVVLNGKVDAFCGGSIVNEKWIVTAAHCVETGVKITVV
 AGEHNIEETEHTEQKRN VIRIIPHHNYNAINKYNHDIALLELDEPLVLNSYVTPICIAD
 KEYTNIFLKFGSGYVSGWGRV FHKGRSALVLQYLRVPLVDRATCLRSTKFTIYNMF
 25 CAGFHEGGRDSCQGD SGPHVTEVEGTSFLTGIISWGEECAMKGKYGIYTKVSRVYN
 WIKEKTKLTSSSSKAPPPSLPSRLPGPSDTPILQSSSSKAPPPSLPSRLPGPSDTPIL
 PQSSSSKAPPPSLPSRLPGPSDTPILQSSSSKAPPPSLPSRLPGPSDTPILQ**GSA
 A (SEQ ID NO: 33).

[0103] In another embodiment, the nucleic acid sequence encoding Factor IX-(CTP)₅
 30 (attached to the carboxy terminus) comprises the following nucleic acid sequence:
 ctagagtcgaccccgccatgcagcgcgtgaacatgatcatggcagaatcaccaggcctcatcaccatctgccttttaggatattactca
 gtgctgaatgtacagttttctgatcatgaaaacgccaacaaaattctgaatcgccaaagaggtataattcaggtaaattggaagagttt
 gttaagggaaccttgagagagaatgtatggaagaaaagtgtagttttgaagaagcacgagaagttttgaaaactgaaagaacaac
 tgaattttggaagcagtagttgatggagatcagtgtagtccaatccatgtttaaattggcggcagttgcaaggatgacattaattcctatg

aatgttggtgtcccttggattgaaggaaagaactgtgaattagatgaacatgaacattaagaatggcagatgcgagcagtttgtaa
aatagtgtgataacaagggtgttgcctgtactgaggatcgcacttgcagaaaaccagaagtcctgtgaaccagcagtgccattt
ccatgtggaagagtttctgttcacaaacttctaagctcacccgtctgaggcagttttcctgatgtggactatgtaaattctactgaagctg
aaaccatttggataacatcaactcaaagcacccaatcatttaatactcactcagttgttggtggagaagatgccaaaccaggtcaattc
5 ccttggcaggtgtttgaatggtaaagtgtgacattctgtggaggctctatcgttaataaaaaatggattgtaactgtgcccactgtgtg
aaactggtgttaaaattacagttgtcgcaggtgaacataataggaggacagaacatacagagcaaaagcgaaatgtgattcgaatta
ttcctcaccacaactacaatgcagctattaataagtacaacatgacattgcccttctggaactggacgaacccttagtgtaaacagcta
cgttacacctatttgcatgtgacaaggaatacacgaacatcttctcaaatggatctggctatgtaagtggctggggaagagtctcc
acaaaggagatcagcttagttctcagtagcttagtccactgttgaccgagccacatgtcttcgatctacaaagttcaccatctata
10 acaacatgttctgtgctggttccatgaaggaggtagagattcatgtcaaggagatagtgggggaccccatgttactgaagtggagg
accagtttcttaactggaattattagctggggtgaagagtggtcaatgaaaggcaaatatggaatatataccaaggtatccgggtatgtca
actggaltaaggaaaaaacaagctcactagctccagcagcaaggccccctccccgagcctgccctcccccaagcaggtgcctggg
ccctctgacaccctatctgcctcagtcagctcctctaaggctccaccacctccctgcctagccctcaagactgccaggccctagc
gataccaattctgccccagtcctccagcagcaaggctccccacctagcctgccttctccatcaaggctgcctggccccatccgatac
15 cccaatttgcctcagagcagctctagcaaggcacctccccagctctgccttctcaagcagactccctggcccttcagacactccaat
cctccacagtcctctagctctaaagctccacccagcctgcccagccctagtagactccccggacctctgataccccatcttgc
ccagtgatgaggatccggggccgc (SEQ ID NO: 34).

[0104] In another embodiment, the amino acid sequence of Factor IX-(CTP)₅ (attached to
the carboxy terminus) comprises the following amino acid sequence:
20 RVDPAHQVRNMIMAEPLITICLLGYLLSAECTVFLDHENANKILNRPKRYNSGKL
EEFVQGNLERECMEEKCSFEEAREVFENTERTEFWKQYVDGDQCESNPCLNGGSC
KDDINSYECWCPFGFEGKNCELDVTCNIKNRCEQFCNSADNKVVCSCTEGYRLA
ENQKSCEPAVPFPCGRVSVSQTSLTRAFAVFPDQVDYVNSTEAETILDNTQSTQSFN
DFTRVVGGEDAKPGQFPWQVVLNGKVDAFCGGSIVNEKWIVTAAHCVETGVKITVV
25 AGEHNIEETEHTEQKRNIRIIPHHNYNAANKYNHDIALLLEPLVLNSYVTPICIAD
KEYTNIFLKFSGSYVSGWGRVVFHKGRSALVLQYLRVPLVDRATCLRSTKFTIYNMF
CAGFHEGGRDSCQGDSSGPHVTEVEGTSFLTGIISWGEECAMKGKYGITKVSRYVN
WIEKTKLTSSSSKAPPPSLPSRLPGPSDTPILQSSSSKAPPPSLPSRLPGPSDTPIL
PQSSSSKAPPPSLPSRLPGPSDTPILQSSSSKAPPPSLPSRLPGPSDTPILQSSSSK
30 APPPSLPSRLPGPSDTPILQ**GSAA (SEQ ID NO: 35).

[0105] In another embodiment, furin is added to a cell expressing the coagulation factor-
CTP of the invention. In another embodiment, furin increases the production efficiency of a
coagulation factor-CTP of the invention in a cell. In another embodiment, furin is co-
transfected with the vector comprising the coding sequence of the coagulation factor-CTP of

the invention. In another embodiment, furin is encoded by a separate vector. In another embodiment, furin and a coagulation factor-CTP are encoded by one vector. In another embodiment, the coding sequence of furin is inserted into pCI-DHFR. In another embodiment, the coding sequence of furin is engineered in pCI-dhfr/smaI+NotI, Furin/AsisI
 5 F.I.+NotI.

[0106] In another embodiment, the nucleic acid sequence encoding furin comprises the following nucleic acid sequence:

tctagagtcgaccccgccatggagctgaggccctgggtgctatgggtgtagcagcaacaggaaccttggctcgtagcagctgatg
 ctgaggccagaaggtcttccaacacgtgggtgtgcgcatccctggaggccagcggaggccaacagtgtggcacggaagcat
 10 ggggttctcaacctgggccagatcttggggactattaccacttctggcatcgaggagtgcgaagcggctccctgtcgcctcaccgcc
 gcggcacagccggctgcagagggagcctcaagtacagtggctggaacagcaggtggcaaagcgacggactaaacgggacgtgta
 ccaggagccccacagacccaagtctcctcagcagtggtacctgtctggtgtcactcagcgggacctgaatgtgaaggcggcctgggc
 gcagggtacacagggcacggcattgtggtctcattctggacgatggcatcgagaagaaccacccggacttggcaggcaattatgat
 cctggggccagttttgatgtcaatgaccaggacctgacccccagcctcggtacacacagatgaatgacaacaggcacggcacagg
 15 tgtgcgggggaagtggctgcgggtggccaacaacgggtgtctggtgtagggtgtggcctacaacgcccgcattggaggggtgcgcatg
 ctggatggcgaggtgacagatgcagtgaggcacgctcgtggcctgaacccaaccacatccacatctacagtccagctgggg
 ccccgaggatgacggcaagacagtggatgggccagcccgctcgccgaggaggccttcttccgtgggggttagccaggggccgagg
 ggggctgggctccatcttctgtcggcctcggggaacgggggcccgggaacatgacagctgcaactgcgacggctacccaacagta
 tctacacgctgtccatcagcagcggcagcagtttggcaacgtgccgtggtacagcgaggcctgctcgtccactggccacgacct
 20 cagcagtggaaccagaatgagaagcagatcgtgacgactgacttgcggcagaagtgcacggagtctcacacgggcacctcagcct
 ctgcccccttagcagccggcatcattgctctaccctggaggccaataagaacctcacatggcgggacatgcaacacctgggtgtaca
 gacctcgaagccagcccacctcaatgccaacgactgggccaccaatgggtgtggccggaaagtgagccactcatatggctacgggc
 ttttggacgcaggcgccatggtggcctggccagaattggaccacagtggccccccagcgggaagtgcacatcgacatcctcaccg
 agcccaaagacatcgggaaacggctcagggtgcggaagaccgtgaccgcgtgcctgggagagcccaaccacatcactcggctgg
 25 agcacgctcaggcgcggtcaccctgtcctataatgccgtggcgacctggccatccacctggtcagcccatgggcacccgctcca
 ccctgctggcagccaggccacatgactactccgagatgggttaatgactgggccttcatgacaactattcctgggatgaggatccct
 ctggcgagtgggtcctagagattgaaaacaccagcgaagccaacaactatgggacgctgaccaagttcacctcgtactctatggcac
 cgccccctgaggggctgcccgtacctccagaagcagtggtgcaagacctcacgtccagttaggcctgtgtgtgtgcgaggaag
 gcttctcctgcaccagaagagctgtgtccagcactgccccccaggcttgcggggcagggctgctgcccacacctgcctgaggtg
 30 gacgtggagacatccgggccagcgtctgcggccctgccacgctcatgtgccacatgccaggggcccggcctgacagactgcct
 cagctgccccagccacgcctccttggacctgtggagcagacttgcctccggcaagccagagcagccgagagtccccgccacag
 cagcagccacctcggctgccccggaggtggagggcggggcaacggctgcgggcagggtgctgcccacacctgcctgaggtg
 gtggccggcctcagctgcgccttcacgtgtgtgtctcgtcactgtcttctggtcctgcagctgcgctctggttttagtttggggggt
 gaaggtgtacacatggaccgtggcctcatctcctacaaggggtgccccctgaagcctggcaggaggagtggcctgtgactcaga

agaggacgagggccggggcgagaggaccgcctttatcaaagaccagagcgccctctgaacggggccgc (SEQ ID NO: 22).

[0107] In another embodiment, the amino acid sequence of furin comprises the following amino acid sequence:

5 MELRPWLLWVVAATGTLVLLAADAQGQKVFTNTWAVRIPGGPAVANSVARKHGFL
 NLGQIFGDYYHFWHRGVTKRSLSPHRPRHSRLQREPQVQWLEQQVAKRRTKRDVY
 QEPTDPKFPQQWYLSGVTQRDLNVKAAWAQGYTGHGIVVSILDDGIEKNHPDLAGN
 YDPGASFDVNDQDPDPQPRYTQMNDNRHGTGRCAGEVA AVANNGVCGVGVAYNAR
 IGGVRMLDGEVTD AVEARSLGLNPNHIHIYSASWGPEDDGKTVDGPARLAEEAFFRG
 10 VSQGRGGLGSIFVWASGNGGREHDSNCNCDGYTNSIYTLSSSATQFGNVPWYSEACS
 STLATTYSSGNQNEKQIVTTDLRQKCTESHTGTSASAPLAAGIIALTLEANKNLTWRD
 MQHLVVQTSKPAHLNANDWATNGVGRKVSHSYGYGLLDAGAMVALAQNWTTVA
 PQRKCIIDILTEPKDIGKRLEVRKTVTACLGEPNHITRLEHAQARLTLSYNRRGDLAIH
 LVSPMGTRSTLLAARPHDYSADGFNDWAFMTTHSWDEDPGGEVWLEIENTSEANNY
 15 GTLTKFTLVLYGTAPGLPVPPESSGCKTLTSSQACVVCEEGFSLHQKSCVQHCPPGF
 APQVLDTHYSTENDVETIRASVCAPCHASCATCQGPALTDCLSCPSHASLDPVEQTCS
 RQSQSSRESPPQQPPRLPPEVEAGQRLRAGLLPSHLPEVVAGLSCAFIVLVFVTVFLV
 LQLRSGFSFRGVKVYTMDRGLISYKGLPPEAWQEECPDSEEDGREGERTAFIKDQSA
 L* (SEQ ID NO: 23).

20 [0108] In one embodiment, the term coagulation factor further includes a homologue of a known coagulation factor. In one embodiment, the homologue has a coagulating activity. In some embodiments, homology according to the present invention also encompasses deletion, insertion, or substitution variants, including an amino acid substitution, thereof and biologically active polypeptide fragments thereof. In one embodiment, the variant comprises
 25 conservative substitutions, or deletions, insertions, or substitutions that do not significantly alter the three dimensional structure of the coagulation factor. In another embodiment, the deletion, insertion, or substitution does not alter the function of interest of the coagulation factor, which in one embodiment, is binding to a particular binding partner.

[0109] In another embodiment, the invention includes a homologue of a coagulation factor.
 30 In another embodiment, the invention includes a homologue of a coagulation factor having a coagulation activity. In another embodiment, the invention includes a homologue of a coagulation factor having functional binding. In another embodiment, the invention includes homologues of a coagulation factor as described herein having a coagulation activity. In another embodiment, the invention includes homologues of a coagulation factor as described

herein having functional binding. In another embodiment, homologues e.g., polypeptides which are at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 87%, at least 89%, at least 91%, at least 93%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% homologous to a coagulation factor as determined using BlastP software of the National Center of Biotechnology Information (NCBI) using default parameters.

[0110] In another embodiment, the invention includes homologues of furin. In another embodiment, the invention includes homologues of furin maintaining a function of interest, which in one embodiment is cleaving of a precursor protein. In another embodiment, homologues e.g., polypeptides which are at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 87%, at least 89%, at least 91%, at least 93%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% homologous to a furin as determined using BlastP software of the National Center of Biotechnology Information (NCBI) using default parameters.

[0111] In another embodiment, provided herein is a polypeptide comprising a coagulation factor and one to ten gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and one to three gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and one to five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor having at least one CTP on its carboxy terminus.

[0112] In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and one to five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor.

[0113] In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and one to five CTPs attached to the carboxy terminus of the coagulation factor.

[0114] It is to be understood that the compositions and methods of the present invention comprising the elements or steps as described herein may, in another embodiment, consist of those elements or steps, or in another embodiment, consist essentially of those elements or steps. In some embodiments, the term "comprise" refers to the inclusion of the indicated active agent, such as the CTP-modified coagulation factor, as well as inclusion of other active

agents, and pharmaceutically acceptable carriers, excipients, emollients, stabilizers, etc., as are known in the pharmaceutical industry. In some embodiments, the term “consisting essentially of” refers to a composition, whose only active ingredient is the indicated active ingredient, however, other compounds may be included which are for stabilizing, preserving, etc. the formulation, but are not involved directly in the therapeutic effect of the indicated active ingredient. In some embodiments, the term “consisting essentially of” may refer to components which facilitate the release of the active ingredient. In some embodiments, the term “consisting” refers to a composition, which contains the active ingredient and a pharmaceutically acceptable carrier or excipient.

10 [0115] In one embodiment, the present invention provides a polypeptide comprising a coagulation factor and two gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and two to three CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide
15 comprising a coagulation factor and two to four CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and two to five CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and two to six CTPs attached to the carboxy terminus of the coagulation factor. In
20 another embodiment, provided herein is a polypeptide comprising a coagulation factor and two to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and two to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and two to nine
25 CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and two to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0116] In one embodiment, the present invention provides a polypeptide comprising a coagulation factor and three gonadotrophin carboxy terminal peptides (CTPs) attached to the
30 carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and three to four CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and three to five CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a

coagulation factor and three to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and three to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and three to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and three to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and three to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0117] In one embodiment, the present invention provides a polypeptide comprising a coagulation factor and four gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and four to five CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and four to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and four to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and four to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and four to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and four to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0118] In one embodiment, the present invention provides a polypeptide comprising a coagulation factor and five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and five to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and five to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and five to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide comprising a coagulation factor and five to nine CTPs attached to the carboxy terminus of the coagulation factor. In

another embodiment, provided herein is a polypeptide comprising a coagulation factor and five to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0119] In one embodiment, the present invention provides a polypeptide consisting of a coagulation factor and two gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and two to three CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and two to four CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and two to five CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and two to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and two to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and two to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and two to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and two to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0120] In one embodiment, the present invention provides a polypeptide consisting of a coagulation factor and three gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and three to four CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and three to five CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and three to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and three to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and three to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and three to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment,

provided herein is a polypeptide consisting of a coagulation factor and three to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0121] In one embodiment, the present invention provides a polypeptide consisting of a coagulation factor and four gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and four to five CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and four to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and four to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and four to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and four to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and four to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0122] In one embodiment, the present invention provides a polypeptide consisting of a coagulation factor and five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and five to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and five to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and five to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and five to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting of a coagulation factor and five to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0123] In one embodiment, the present invention provides a polypeptide consisting essentially of a coagulation factor and two gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and two to three CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and two to four CTPs

attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and two to five CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and two to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and two to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and two to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and two to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and two to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0124] In one embodiment, the present invention provides a polypeptide consisting essentially of a coagulation factor and three gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and three to four CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and three to five CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and three to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and three to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and three to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and three to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and three to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0125] In one embodiment, the present invention provides a polypeptide consisting essentially of a coagulation factor and four gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and four to five CTPs

attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and four to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and four to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and four to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and four to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and four to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0126] In one embodiment, the present invention provides a polypeptide consisting essentially of a coagulation factor and five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and five to six CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and five to seven CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and five to eight CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and five to nine CTPs attached to the carboxy terminus of the coagulation factor. In another embodiment, provided herein is a polypeptide consisting essentially of a coagulation factor and five to ten CTPs attached to the carboxy terminus of the coagulation factor.

[0127] In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of a coagulation factor having no CTPs on its amino terminus. In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of a coagulation factor lacking a CTP on its amino terminus. In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of a coagulation factor having at least one CTP on its carboxy terminus and no CTPs on its amino terminus. In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of a coagulation factor having the number of CTPs on its carboxy terminus as described herein and no CTPs on its amino terminus.

[0128] In another embodiment, the present invention provides a polynucleotide encoding a polypeptide as described hereinabove.

[0129] In another embodiment, the present invention further provides a composition comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor IX (FIX) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide.

[0130] In another embodiment, the present invention further provides a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor VIIa (FVIIa) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide.

[0131] In one embodiment, the present invention provides a recombinant coagulation factor as described hereinabove. In one embodiment, the present invention provides an engineered coagulation factor as described hereinabove. In one embodiment, the engineered coagulation factor as described hereinabove is referred to as a CTP-modified coagulation factor.

[0132] In one embodiment, the CTPs that are attached to the carboxy terminus of the coagulation factor are attached in tandem to the carboxy terminus.

[0133] In one embodiment, an engineered coagulation factor as described herein has equivalent or improved biological activity compared to the non-CTP-modified coagulation factor. In another embodiment, an engineered coagulation factor as described herein has equivalent or improved pharmacological measurements compared to the non-CTP-modified coagulation factor. In another embodiment, an engineered coagulation factor as described herein has equivalent or improved pharmacokinetics compared to the non-CTP-modified coagulation factor. In another embodiment, an engineered coagulation factor as described herein has equivalent or improved pharmacodynamics compared to the non-CTP-modified coagulation factor.

[0134] In one embodiment, the present invention provides a method of preventing or treating a clotting or coagulation disorder. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising administering a CTP-modified coagulation factor of the present invention. In another embodiment, the present invention provides a method of preventing and treating hemophilia in a subject comprising administering a CTP-modified coagulation factor of the present invention. In another embodiment, the present invention provides a method of treating hemophilia in a subject comprising administering a CTP-modified Factor VII of the present invention.

[0135] In another embodiment, the present invention provides a method of treating hemophilia in a subject comprising administering a CTP-modified Factor IX of the present invention. In one embodiment, hemophilia is hemophilia B. In one embodiment, hemophilia B is known as factor IX deficiency or Christmas disease. In one embodiment, the hemophilia is severe hemophilia, which in one embodiment, describes hemophilia in which the coagulation factor levels are 0-1%. In another embodiment, the hemophilia is moderate hemophilia, which in one embodiment, describes hemophilia in which the coagulation factor levels are 1-5%. In another embodiment, the hemophilia is mild hemophilia, which in one embodiment, describes hemophilia in which the coagulation factor levels are 5-50%.

[0136] In another embodiment, the present invention provides a method of preventing or treating a clotting or coagulation disorder in a subject comprising administering a CTP-modified Factor IX (FIX) polypeptide comprising a FIX polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide to said subject, thereby preventing or treating a clotting or coagulation disorder in said subject. In another embodiment, the present invention provides a method of preventing or treating a clotting or coagulation disorder in a subject comprising administering a CTP-modified Factor VII (FVII) polypeptide comprising a FVII polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVII polypeptide to said subject, thereby preventing or treating a clotting or coagulation disorder in said subject.

[0137] In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising administering a CTP-modified Factor IX (FIX) polypeptide comprising a FIX polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising administering a CTP-modified Factor VIIa (FVIIa) polypeptide comprising a FVIIa polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide to said subject, thereby preventing or treating hemophilia in said subject.

[0138] In another embodiment, the present invention provides a method of treating hemophilia in a subject comprising administering one or more CTP-modified coagulation factors as described herein to said subject. Thus, in one embodiment, the present invention provides a method of treating hemophilia in a subject comprising administering a CTP-

modified Factor IX (FIX) polypeptide comprising a FIX polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide and a CTP-modified Factor VIIa (FVIIa) polypeptide comprising a FVIIa polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide to said subject, thereby treating hemophilia in said subject. In one embodiment, the CTP-modified FIX and the CTP-modified FVIIa are administered in the same composition at the same time. In another embodiment, the CTP-modified FIX and the CTP-modified FVIIa are administered in separate compositions at the same time. In another embodiment, the CTP-modified FIX and the CTP-modified FVIIa are administered in separate compositions at separate times.

[0139] In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising administering a CTP-modified Factor IX (FIX) or a CTP-modified Factor VII polypeptide comprising a FIX or a FVII polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX or said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising administering a CTP-modified Factor IX (FIX) or a CTP-modified Factor VII polypeptide comprising a FIX or a FVII polypeptide and four chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX or said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising administering a CTP-modified Factor IX (FIX) or a CTP-modified Factor VII polypeptide comprising a FIX or a FVII polypeptide and five chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX or said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising administering a CTP-modified Factor IX (FIX) or a CTP-modified Factor VII polypeptide comprising a FIX or a FVII polypeptide and three to five chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX or said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising administering a CTP-modified Factor IX (FIX) and a CTP-modified Factor VII polypeptide comprising a FIX and a FVII polypeptide and three

chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX and said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising administering a CTP-modified Factor IX (FIX) and a CTP-modified Factor VII polypeptide comprising a FIX and a FVII polypeptide and three to five chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX and said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject.

[0140] In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising subcutaneously or intravenously administering a CTP-modified Factor IX (FIX) or a CTP-modified Factor VII polypeptide comprising a FIX or a FVII polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX or said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising subcutaneously or intravenously administering a CTP-modified Factor IX (FIX) or a CTP-modified Factor VII polypeptide comprising a FIX or a FVII polypeptide and four chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX or said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising subcutaneously or intravenously administering a CTP-modified Factor IX (FIX) or a CTP-modified Factor VII polypeptide comprising a FIX or a FVII polypeptide and five chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX or said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising subcutaneously or intravenously administering a CTP-modified Factor IX (FIX) or a CTP-modified Factor VII polypeptide comprising a FIX or a FVII polypeptide and three to five chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX or said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising subcutaneously or intravenously administering a CTP-modified Factor IX (FIX) and a CTP-modified Factor VII polypeptide comprising a FIX and a FVII polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX and said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject. In another embodiment, the present invention provides a method of preventing or treating hemophilia in a subject comprising subcutaneously or intravenously administering a CTP-modified Factor IX (FIX) and a CTP-modified Factor VII polypeptide comprising a FIX and a FVII polypeptide and three to five chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX and said FVII polypeptide to said subject, thereby preventing or treating hemophilia in said subject.

[0141] In some embodiments, provided herein is a method of preventing or treating a hemophilia in a subject, the method comprising the step of administering to the subject a CTP-modified coagulation factor, comprising three to five chorionic gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said coagulation factor polypeptide, wherein the sequence of said CTP-modified coagulation factor is selected from the group consisting of SEQ ID NO: 25, 27, or 29, thereby preventing hemophilia in said subject.

[0142] In some embodiments, provided herein is a method of preventing or treating a hemophilia in a subject, the method comprising the step of subcutaneously administering to the subject a CTP-modified Factor VII, comprising three to five chorionic gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVII polypeptide, wherein the sequence of said CTP-modified FVII is selected from the group consisting of SEQ ID NO: 25, 27, or 29, thereby preventing hemophilia in said subject.

[0143] In other embodiments, the engineered coagulation factor is for the treatment of hemophilia B patients. In one embodiment, coagulation Factor IX comprising 3 CTPs in tandem in its carboxy terminus is for the treatment of hemophilia B patients. In one embodiment, coagulation Factor IX comprising 4 CTPs in tandem in its carboxy terminus is for the treatment of hemophilia B patients. In one embodiment, coagulation Factor IX comprising 5 CTPs in tandem in its carboxy terminus is for the treatment of hemophilia B patients. In another embodiment, coagulation Factor IX comprising 2 CTPs in tandem in its carboxy terminus is for the treatment of hemophilia B patients. In another embodiment, coagulation Factor IX comprising 1 CTP repeat in its carboxy terminus is for the treatment of hemophilia B patients. In other embodiments, the engineered coagulation factor can reduce the number of infusions required for a patient, reduce the required doses for a patient, or a combination thereof.

[0144] In one embodiment, coagulation Factor IX comprising 3 CTPs in tandem in its carboxy terminus exhibits an improved PK profile while maintaining its coagulation activity vs. FIX-CTP-CTP harvest, FIX-CTP harvest or rhFIX. In one embodiment, the elimination half-life of rFIX-CTP3 is 2.5- to 4-fold longer than rFIX in rats and in FIX-deficient mice. In one embodiment, the administration of rFIX-CTP3 significantly prolonged the procoagulatory effect in FIX-deficient mice for at least 76 hr after dosing. In one embodiment, the administration of rFIX-CTP3 produced a higher activity peak than rFIX in FIX-deficient mice. In another embodiment, coagulation Factor IX comprising 2 CTPs in tandem in its carboxy terminus exhibits an improved PK profile while maintaining its coagulation activity

vs. FIX-CTP harvest or rhFIX. In another embodiment, coagulation Factor IX comprising 2 CTPs in tandem in its carboxy terminus exhibits 3-fold increase in half-life and 4.5-fold higher AUC compared to rhFIX.

[0145] In another embodiment, SC administration results in higher bioavailability of CTP-
5 modified FVII as compared to recombinant FVII. In another embodiment, half life is longer and bioavailability (AUC SC/AUC IV) is higher following FVIIa-CTP3 and 5 SC administration when compared to SC administration of NovoSeven®. In another embodiment, subcutaneously injected MOD-5014 and MOD-5019 shows improved mice survival in comparison to recombinant FVII (NovoSeven®) (see Example 8 below).

10 [0146] In another embodiment, the terms “CTP peptide,” “carboxy terminal peptide” and “CTP sequence” are used interchangeably herein. In another embodiment, the carboxy terminal peptide is a full-length CTP. Each possibility represents a separate embodiment of the invention.

[0147] In another embodiment, a signal peptide is attached to the amino terminus of the
15 CTP, as described in US 7,553,940, which is incorporated by reference herein in its entirety.

[0148] In other embodiments, the term engineered coagulation factor refers to the amino acid sequence of a matured coagulation factor. In other embodiments, the term engineered coagulation factor refers to the amino acid sequence of the coagulation factor including its signal sequence or signal peptide.

20 [0149] In another embodiment, “signal sequence” and “signal peptide” are used interchangeably herein. In another embodiment, “sequence” when in reference to a polynucleotide molecule can refer to a coding portion. Each possibility represents a separate embodiment of the present invention.

[0150] In another embodiment, an engineered coagulation factor comprising at least one
25 CTP as described herein has enhanced in vivo biological activity compared the same coagulation factor without at least one CTP. In one embodiment, the enhanced biological activity stems from the longer half-life of the engineered coagulation factor while maintaining at least some biological activity. In another embodiment, the enhanced biological activity stems from enhanced biological activity resulting from the CTP modification. In another
30 embodiment, the enhanced biological activity stems from both a longer half life and from enhanced functionality of the CTP-modified coagulation factor.

[0151] In some embodiments, at least one CTP sequence at the carboxy terminal end of the coagulation factor provides enhanced protection against degradation of a coagulation factor. In some embodiments, at least one CTP sequence at the carboxy terminal end of the

coagulation factor provides enhanced protection against clearance. In some embodiments, at least one CTP sequence at the carboxy terminal end of the coagulation factor provides prolonged clearance time. In some embodiments, at least one CTP sequence at the carboxy terminal end of the coagulation factor enhances its C_{max}. In some embodiments, at least one CTP sequence at the carboxy terminal end of the coagulation factor enhances its T_{max}. In some embodiments, at least one CTP sequence at the carboxy terminal end of the coagulation factor prolongs its T_{1/2}.

[0152] In another embodiment, a conjugated coagulation factor of this invention is used in the same manner as an unmodified conjugated coagulation factor. In another embodiment, a conjugated coagulation factor of this invention has an increased circulating half-life and plasma residence time, decreased clearance, and increased clinical activity in vivo. In another embodiment, due to the improved properties of the conjugated coagulation factor as described herein, this conjugate is administered less frequently than the unmodified form of the same coagulation factor.

[0153] In another embodiment, decreased frequency of administration will result in improved treatment strategy, which in one embodiment, will lead to improved patient compliance leading to improved treatment outcomes, as well as improved patient quality of life. In another embodiment, compared to conventional conjugates of coagulation factors, it has been found that conjugates having the molecular weight and linker structure of the conjugates of this invention have an improved potency, improved stability, elevated AUC levels, and enhanced circulating half-life.

[0154] In another embodiment, the present invention further provides a pharmaceutical composition comprising a CTP-modified Factor IX (FIX) polypeptide consisting of a FIX polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said CTP-modified FIX polypeptide.

[0155] In another embodiment, the present invention further provides a pharmaceutical composition comprising a CTP-modified Factor VIIa (FVIIa) polypeptide consisting of a FVIIa polypeptide and five gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa.

[0156] In another embodiment, provided herein is a composition comprising a conjugated coagulation factor as described herein. In another embodiment, provided herein is a pharmaceutical composition comprising the conjugated coagulation factor as described herein. In another embodiment, provided herein is a pharmaceutical composition comprising a therapeutically effective amount of the conjugated coagulation factor as described herein. In

one embodiment, a therapeutically effective amount of a conjugated coagulation factor is determined according to factors such as the specific condition being treated, the condition of the patient being treated, as well as the other ingredients in the composition.

[0157] In another embodiment, a conjugated coagulation factor as described herein is useful
5 in the treatment of subjects afflicted with Hemophilia. In another embodiment, a conjugated coagulation factor as described herein is useful in the prophylactic therapy of Hemophilia thus reducing the risk of bleeding and associated complications. In another embodiment, a conjugated coagulation factor as described herein is useful in the treatment of subjects afflicted with Hemophilia while reducing the risk of developing inhibitory antibodies to
10 exogenously administered coagulation factors. In another embodiment, a conjugated coagulation factor as described herein is useful in the treatment of subjects afflicted with Hemophilia thus inducing homeostasis.

[0158] In one embodiment, a CTP-modified coagulation factor of the present invention has
15 therapeutic uses. In another embodiment, a CTP-modified coagulation factor of the present invention has prophylactic uses.

[0159] In another embodiment, a conjugated coagulation factor as described herein is useful
in the treatment of subjects experiencing excessive bleeding or bruising or having a prolonged Prothrombin Time (PT) or Partial Thromboplastin Time (PTT). In another embodiment, a conjugated coagulation factor as described herein is useful in the treatment of subjects having
20 an acquired condition that is causing bleeding, such as vitamin K deficiency or liver disease. In another embodiment, a conjugated coagulation factor as described herein is useful in the treatment of subjects having deficiencies in coagulation factors that are acquired (due to other diseases) or inherited, mild or severe, permanent or temporary. In another embodiment, a conjugated coagulation factor as described herein is useful in the treatment of subjects
25 afflicted with hemophilia A. In another embodiment, a conjugated coagulation factor as described herein is useful in the treatment of subjects afflicted with hemophilia B. In another embodiment, a conjugated coagulation factor as described herein is useful in the treatment of subjects having acquired deficiencies due to chronic diseases, such as liver disease or cancer; to an acute condition such as disseminated intravascular coagulation (DIC), which uses up
30 clotting factors at a rapid rate; or to a deficiency in vitamin K or treatment with a vitamin K antagonist like warfarin (the production of factors II, VII, IX, and X require vitamin K). In another embodiment, a conjugated coagulation factor as described herein is useful in the treatment of subjects afflicted with a disease in which causes clotting imbalances such as but not limited to: a liver disease, uremia, a cancer, a bone marrow disorder, an exposure to snake

venom, a vitamin K deficiency, an anticoagulation therapy, an accidental ingestion of the anticoagulant warfarin, multiple blood transfusions (stored units of blood lose some of their clotting factors), or a combination thereof. In another embodiment, the present invention provides a method of treating deep vein thrombosis in a subject comprising administering a CTP-modified coagulation factor of the present invention. In another embodiment, the present invention provides a method of preventing uncontrolled bleeding in a subject with hemophilia comprising administering a CTP-modified coagulation factor of the present invention. In another embodiment, the present invention provides a method of preventing bleeding episodes in a subject with hemophilia comprising administering a CTP-modified coagulation factor of the present invention. In another embodiment, the present invention provides a method of controlling bleeding episodes in a subject with hemophilia B (congenital factor IX deficiency).

[0160] In another embodiment, the compositions and methods of the present invention are for the treatment of bleeding episodes in hemophilia A or B patients with inhibitors to FVIII or FIX and in patients with acquired hemophilia; prevention of bleeding in surgical interventions or invasive procedures in hemophilia A or B patients with inhibitors to FVIII or FIX and in patients with acquired hemophilia; treatment of bleeding episodes in patients with congenital FVII deficiency and prevention of bleeding in surgical interventions or invasive procedures in patients with congenital FVII deficiency. In another embodiment, the compositions and methods of the present invention are for the treatment or prevention of muscle bleeds. In another embodiment, the compositions and methods of the present invention are for the treatment or prevention of joint bleeds. In another embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of epistaxis and gum bleeding, mucous membrane bleeding, bleeding into the central nervous system. In another embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of gastrointestinal or cerebral bleeding. In another embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of low frequency mild bleeds. In another embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of low frequency moderate bleeds. In another embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of high frequency mild bleeds. In another embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of high frequency moderate bleeds.

[0161] In one embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of asymptomatic hemophilia. In another embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of mild to moderate hemophilia. In another embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of severe hemophilia.

[0162] In one embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of hemorrhage, which in one embodiment, is uncontrollable hemorrhage, and, in another embodiment, intracerebral hemorrhage. In another embodiment, the compositions and methods of the present invention provide therapeutic or prophylactic treatment of neonatal coagulopathies; severe hepatic disease; high-risk surgical procedures; traumatic blood loss; bone marrow transplantation; thrombocytopenias and platelet function disorders; urgent reversal of oral anticoagulation; congenital deficiencies of factors V, VII, X, and XI; or von Willebrand disease, in one embodiment, von Willebrand disease with inhibitors to von Willebrand factor.

[0163] In one embodiment, a CTP-modified coagulation factor of the present invention is for the treatment of hemophilia or a related disease as described herein in a subject. In one embodiment, the subject is human. In another embodiment, the subject is a domesticated animal. In another embodiment, the subject is a mammal. In another embodiment, the subject is a farm animal. In another embodiment, the subject is a monkey. In another embodiment, the subject is a horse. In another embodiment, the subject is a cow. In another embodiment, the subject is a mouse. In another embodiment, the subject is a rat. In another embodiment, the subject is canine. In another embodiment, the subject is feline. In another embodiment, the subject is bovine, ovine, porcine, equine, murine, or cervine. In one embodiment, the subject is male. In another embodiment, the subject is female. In one embodiment, the subject is a child, in another embodiment, an adolescent, in another embodiment, an adult or, in another embodiment, an elderly subject. In another embodiment, the subject is a pediatric subject, in another embodiment, a geriatric subject.

[0164] In another embodiment, a [(CTP)_n>1-coagulation factor] as described herein comprises a full length coagulation factor or an active fragment thereof connected via a peptide bond on its carboxy terminus to at least one CTP unit with no CTPs on its amino terminus. In another embodiment, a [(CTP)_n>1-coagulation factor] as described herein comprises a coagulation factor or an active fragment thereof connected via a peptide bond to at least one CTP unit which is connected to an additional CTP unit via a peptide bond with no

CTPs on its amino terminus. In another embodiment, one nucleic acid molecule encodes an engineered coagulation factor comprising at least one CTP attached to its C-terminus and no CTPs on its amino terminus.

[0165] In another embodiment, the CTP is attached to the coagulation factor via a linker. In another embodiment, the linker which connects the CTP sequence to the coagulation factor is a covalent bond. In another embodiment, the linker which connects the CTP sequence to the coagulation factor is a peptide bond. In another embodiment, the linker which connects the CTP sequence to the coagulation factor is a substituted peptide bond. In another embodiment, the CTP sequence comprises: DPRFQDSSSSKAPPPSLPSPSRLPGPSDTPIL (SEQ ID NO: 1). In another embodiment, the CTP sequence comprises: SSSSKAPPPSLPSPSRLPGPSDTPILPQ (SEQ ID NO: 2). In another embodiment, the CTP sequence comprises an amino acid sequence selected from the sequences set forth in SEQ ID NO: 1 and SEQ ID NO: 2.

[0166] In another embodiment, the carboxy terminal peptide (CTP) peptide of the present invention comprises the amino acid sequence from amino acid 112 to position 145 of human chorionic gonadotrophin, as set forth in SEQ ID NO: 1. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 118 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 2. In another embodiment, the CTP sequence also commences from any position between positions 112-118 and terminates at position 145 of human chorionic gonadotrophin. In some embodiments, the CTP sequence peptide is 28, 29, 30, 31, 32, 33 or 34 amino acids long and commences at position 112, 113, 114, 115, 116, 117 or 118 of the CTP amino acid sequence.

[0167] In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 1-5 conservative amino acid substitutions as described in U.S. Pat. No. 5,712,122, which is incorporated herein by reference. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 1 conservative amino acid substitution. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 2 conservative amino acid substitutions. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 3 conservative amino acid substitutions. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 4 conservative amino acid substitutions. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 5 conservative amino acid substitutions.

[0168] In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 70% homologous to the native CTP amino acid sequence or a peptide thereof. In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 80% homologous to the native CTP amino acid sequence or a peptide thereof. In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 90% homologous to the native CTP amino acid sequence or a peptide thereof. In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 95% homologous to the native CTP amino acid sequence or a peptide thereof. In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 98% homologous to the native CTP amino acid sequence or a peptide thereof.

[0169] In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 70% homologous to the native human CTP DNA sequence or a peptide thereof. In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 80% homologous to the native human CTP DNA sequence or a peptide thereof. In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 90% homologous to the native CTP DNA sequence or a peptide thereof. In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 95% homologous to the native CTP DNA sequence or a peptide thereof. In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 98% homologous to the native CTP DNA sequence or a peptide thereof.

[0170] In one embodiment, at least one of the chorionic gonadotrophin CTP amino acid sequences is truncated. In another embodiment, both of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, 2 of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, 3 of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, 4 of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, 5 of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, 2 or more of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, all of the chorionic gonadotrophin CTP amino acid sequences are truncated. In one embodiment, the truncated CTP comprises the first 10 amino acids of SEQ ID NO: 3. In another embodiment, SEQ ID NO: 3 comprises the following amino acid (AA) sequence: SSSSKAPPSLP.

[0171] In one embodiment, the truncated CTP comprises the first 10 amino acids of SEQ ID NO: 4. In another embodiment, SEQ ID NO: 4 comprises the following amino acid (AA) sequence: SSSSKAPPSLPSPSRLPGPSDTPILPQ.

[0172] In one embodiment, the truncated CTP comprises the first 11 amino acids of SEQ ID NO: 4. In one embodiment, the truncated CTP comprises the first 12 amino acids of SEQ ID NO: 4. In one embodiment, the truncated CTP comprises the first 8 amino acids of SEQ ID NO: 4 or SEQ ID NO: 3. In one embodiment, the truncated CTP comprises the first 13 amino acids of SEQ ID NO: 4. In one embodiment, the truncated CTP comprises the first 14 amino acids of SEQ ID NO: 4. In one embodiment, the truncated CTP comprises the first 6 amino acids of SEQ ID NO: 4 or SEQ ID NO: 3. In one embodiment, the truncated CTP comprises the first 5 amino acids of SEQ ID NO: 4 or SEQ ID NO: 3.

[0173] In one embodiment, at least one of the chorionic gonadotrophin CTP amino acid sequences is glycosylated. In another embodiment, both of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, 2 of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, 3 of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, 4 of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, 5 of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, 2 or more of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, all of the chorionic gonadotrophin CTP amino acid sequences are glycosylated.

[0174] In one embodiment, the CTP sequence of the present invention comprises at least one glycosylation site. In one embodiment, the CTP sequence of the present invention comprises 2 glycosylation sites. In one embodiment, the CTP sequence of the present invention comprises 3 glycosylation sites. In one embodiment, the CTP sequence of the present invention comprises 4 glycosylation sites. In one embodiment, one or more of the chorionic gonadotrophin CTP amino acid sequences is fully glycosylated. In another embodiment, one or more of the chorionic gonadotrophin CTP amino acid sequences is partially glycosylated. In one embodiment, partially glycosylated indicates that one of the CTP glycosylation sites is glycosylated. In another embodiment, two of the CTP glycosylation sites are glycosylated. In another embodiment, three of the CTP glycosylation sites are glycosylated.

[0175] In some embodiments, the CTP sequence modification is advantageous in permitting the usage of lower dosages. In some embodiments, the CTP sequences

modification is advantageous in permitting fewer dosages. In some embodiments, the CTP sequences modification is advantageous in permitting a safe, long-acting effect.

[0176] In some embodiments, "polypeptide", "engineered coagulation factor", or "protein" as used herein encompasses native polypeptides (either degradation products, synthetically synthesized polypeptides or recombinant polypeptides) and peptidomimetics (typically, synthetically synthesized polypeptides), as well as peptoids and semipeptoids which are polypeptide analogs, which have, in some embodiments, modifications rendering the polypeptides comprising a coagulation factor even more stable while in a body or more capable of penetrating into cells.

[0177] In some embodiments, modifications include, but are limited to C terminus modification, polypeptide bond modification, including, but not limited to, CH₂-NH, CH₂-S, CH₂-S=O, O=C-NH, CH₂-O, CH₂-CH₂, S=C-NH, CH=CH or CF=CH, backbone modifications, and residue modification. Methods for preparing peptidomimetic compounds are well known in the art and are specified, for example, in Quantitative Drug Design, C.A. Ramsden Gd., Chapter 17.2, F. Choplin Pergamon Press (1992), which is incorporated by reference as if fully set forth herein. Further details in this respect are provided hereinunder.

[0178] In some embodiments, polypeptide bonds (-CO-NH-) within the polypeptide are substituted. In some embodiments, the polypeptide bonds are substituted by N-methylated bonds (-N(CH₃)-CO-). In some embodiments, the polypeptide bonds are substituted by ester bonds (-C(R)H-C-O-O-C(R)-N-). In some embodiments, the polypeptide bonds are substituted by ketomethylen bonds (-CO-CH₂-). In some embodiments, the polypeptide bonds are substituted by α -aza bonds (-NH-N(R)-CO-), wherein R is any alkyl, e.g., methyl, carba bonds (-CH₂-NH-). In some embodiments, the polypeptide bonds are substituted by hydroxyethylene bonds (-CH(OH)-CH₂-). In some embodiments, the polypeptide bonds are substituted by thioamide bonds (-CS-NH-). In some embodiments, the polypeptide bonds are substituted by olefinic double bonds (-CH=CH-). In some embodiments, the polypeptide bonds are substituted by retro amide bonds (-NH-CO-). In some embodiments, the polypeptide bonds are substituted by polypeptide derivatives (-N(R)-CH₂-CO-), wherein R is the "normal" side chain, naturally presented on the carbon atom. In some embodiments, these modifications occur at any of the bonds along the polypeptide chain and in one embodiment at several (2-3 bonds) at the same time.

[0179] In some embodiments, natural aromatic amino acids of the polypeptide such as Trp, Tyr and Phe, are substituted for synthetic non-natural acid such as Phenylglycine, TIC,

naphthylelanine (Nol), ring-methylated derivatives of Phe, halogenated derivatives of Phe or o-methyl-Tyr. In some embodiments, the polypeptides of the present invention include one or more modified amino acid or one or more non-amino acid monomers (e.g. fatty acid, complex carbohydrates etc).

5 [0180] In one embodiment, "amino acid" or "amino acid sequence" is understood to include the 20 naturally occurring amino acid; those amino acid often modified post-translationally in vivo, including, for example, hydroxyproline, phosphoserine and phosphothreonine; and other unusual amino acid including, but not limited to, 2-aminoadipic acid, hydroxylysine, isodesmosine, nor-valine, nor-leucine and ornithine. In one embodiment, "amino acid"
10 includes both D- and L-amino acids.

[0181] In some embodiments, the polypeptides of the present invention are utilized in therapeutics which requires the polypeptides comprising a coagulation factor to be in a soluble form. In some embodiments, the polypeptides of the present invention include one or more non-natural or natural polar amino acid, including but not limited to serine and
15 threonine which are capable of increasing polypeptide solubility due to their hydroxyl-containing side chain.

[0182] In some embodiments, the engineered coagulation factor of the present invention is utilized in a linear form, although it will be appreciated by one skilled in the art that in cases where cyclicization does not severely interfere with engineered coagulation factors
20 characteristics, cyclic forms of the engineered coagulation factors can also be utilized.

[0183] In some embodiments, the engineered coagulation factors of the present invention are biochemically synthesized such as by using standard solid phase techniques. In some embodiments, these biochemical methods include exclusive solid phase synthesis, partial solid phase synthesis, fragment condensation, or classical solution synthesis.

25 [0184] In some embodiments, recombinant protein techniques are used to generate the engineered coagulation factors of the present invention. In some embodiments, recombinant protein techniques are used for the generation of relatively long polypeptides (e.g., longer than 18-25 amino acids). In some embodiments, recombinant protein techniques are used for the generation of large amounts of the engineered coagulation factors of the present invention.

30 In some embodiments, recombinant techniques are described by Bitter et al., (1987) Methods in Enzymol. 153:516-544, Studier et al. (1990) Methods in Enzymol. 185:60-89, Brisson et al. (1984) Nature 310:511-514, Takamatsu et al. (1987) EMBO J. 6:307-311, Coruzzi et al. (1984) EMBO J. 3:1671-1680 and Brogli et al., (1984) Science 224:838-843, Gurley et al. (1986) Mol. Cell. Biol. 6:559-565 and Weissbach & Weissbach, 1988, Methods for Plant

Molecular Biology, Academic Press, NY, Section VIII, pp 421-463, which are incorporated herein by reference in their entirety.

[0185] In another embodiment, the invention provides a polynucleotide molecule comprising the coding portion of a gene encoding a polypeptide comprising a coagulation factor and gonadotrophin carboxy terminal peptides attached to the carboxy terminus of the coagulation factor, as described hereinabove. In another embodiment, the invention provides a polynucleotide molecule consisting of the coding portion of a gene encoding a polypeptide comprising a coagulation factor and gonadotrophin carboxy terminal peptides attached to the carboxy terminus of the coagulation factor, as described hereinabove. In another embodiment, the invention provides a polynucleotide molecule consisting essentially of the coding portion of a gene encoding a polypeptide comprising a coagulation factor and gonadotrophin carboxy terminal peptides attached to the carboxy terminus of the coagulation factor, as described hereinabove.

[0186] In another embodiment, the invention provides a polynucleotide encoding a polypeptide comprising a coagulation factor and three gonadotrophin carboxy terminal peptides attached to the carboxy terminus of the coagulation factor, as described hereinabove. In another embodiment, the invention provides a polynucleotide encoding a polypeptide consisting of a coagulation factor and three gonadotrophin carboxy terminal peptides attached to the carboxy terminus of the coagulation factor, as described hereinabove. In another embodiment, the invention provides a polynucleotide encoding a polypeptide consisting essentially of a coagulation factor and three gonadotrophin carboxy terminal peptides attached to the carboxy terminus of the coagulation factor, as described hereinabove. In one embodiment, the polynucleotide is a polynucleotide sequence. In one embodiment, the polynucleotide is a polynucleotide molecule.

[0187] In another embodiment, the invention provides an expression vector comprising a polynucleotide molecule as described herein. In another embodiment, the present invention provides an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor IX (FIX) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide. In another embodiment, the present invention provides an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor VIIa (FVIIa) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide.

[0188] In another embodiment, the invention provides a cell comprising the expression vector as described herein. In another embodiment, the present invention provides a cell comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor IX (FIX) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide. In another embodiment, the present invention provides a cell comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor VIIa (FVIIa) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide.

[0189] In another embodiment, the invention provides a composition comprising the expression vector as described herein. In another embodiment, the present invention provides a composition comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor IX (FIX) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide.

In another embodiment, the present invention provides a composition comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor VIIa (FVIIa) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide.

[0190] In another embodiment, the invention provides a composition comprising the cell as described herein. In another embodiment, the cell is a eukaryotic cell. In another embodiment, the cell is a prokaryotic cell.

[0191] In another embodiment, the present invention provides a method of producing a CTP-modified coagulation factor, comprising the step of attaching one to ten chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said coagulation factor, thereby producing a CTP-modified coagulation factor. In another embodiment, the present invention provides a method of producing a CTP-modified coagulation factor, comprising the step of attaching one to ten polynucleotide sequences encoding a chorionic gonadotrophin carboxy terminal peptide (CTP) to the carboxy terminus of a polynucleotide sequence encoding said coagulation factor, thereby producing a CTP-modified coagulation factor. In another embodiment, the present invention provides a method of producing a CTP-modified Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby producing a CTP-modified FIX polypeptide. In another embodiment, the present invention provides a method of producing a CTP-modified Factor VIIa (FVIIa)

polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa polypeptide, thereby producing a CTP-modified FVIIa polypeptide.

[0192] In another embodiment, the engineered coagulation factors of the present invention are synthesized using a polynucleotide molecule encoding a polypeptide of the present invention. In some embodiments, the polynucleotide molecule encoding the engineered coagulation factors of the present invention is ligated into an expression vector, comprising a transcriptional control of a cis-regulatory sequence (e.g., promoter sequence). In some embodiments, the cis-regulatory sequence is suitable for directing constitutive expression of an engineered coagulation factor of the present invention. In some embodiments, the cis-regulatory sequence is suitable for directing tissue-specific expression of the engineered coagulation factors of the present invention. In some embodiments, the cis-regulatory sequence is suitable for directing inducible expression of the engineered coagulation factors of the present invention.

[0193] In some embodiment, tissue-specific promoters suitable for use with the present invention include sequences which are functional in one or more specific cell populations. Examples include, but are not limited to, promoters such as albumin that is liver-specific [Pinkert et al., (1987) *Genes Dev.* 1:268-277], lymphoid-specific promoters [Calame et al., (1988) *Adv. Immunol.* 43:235-275]; in particular promoters of T-cell receptors [Winoto et al., (1989) *EMBO J.* 8:729-733] and immunoglobulins; [Banerji et al. (1983) *Cell* 33729-740], neuron-specific promoters such as the neurofilament promoter [Byrne et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5473-5477], pancreas-specific promoters [Edlunch et al. (1985) *Science* 230:912-916] or mammary gland-specific promoters such as the milk whey promoter (U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). Inducible promoters suitable for use with the present invention include, for example, the tetracycline-inducible promoter (Srouf, M.A., et al., 2003. *Thromb. Haemost.* 90: 398-405).

[0194] In one embodiment, the phrase "a polynucleotide molecule" refers to a single or double stranded nucleic acid sequence which is isolated and provided in the form of an RNA sequence, a complementary polynucleotide sequence (cDNA), a genomic polynucleotide sequence and/or a composite polynucleotide sequences (e.g., a combination of the above).

[0195] In one embodiment, a "complementary polynucleotide sequence" refers to a sequence, which results from reverse transcription of messenger RNA using a reverse transcriptase or any other RNA-dependent DNA polymerase. In one embodiment, the sequence can be subsequently amplified in vivo or in vitro using a DNA polymerase.

[0196] In one embodiment, a "genomic polynucleotide sequence" refers to a sequence derived (isolated) from a chromosome and thus it represents a contiguous portion of a chromosome.

[0197] In one embodiment, a "composite polynucleotide sequence" refers to a sequence, which is at least partially complementary and at least partially genomic. In one embodiment, a composite sequence can include some exonal sequences required to encode the polypeptide of the present invention, as well as some intronic sequences interposing therebetween. In one embodiment, the intronic sequences can be of any source, including of other genes, and typically will include conserved splicing signal sequences. In one embodiment, intronic sequences include cis-acting expression regulatory elements.

[0198] In one embodiment, following expression and secretion, the signal peptides are cleaved from the precursor engineered coagulation factors resulting in the mature engineered coagulation factors.

[0199] In some embodiments, polynucleotides of the present invention are prepared using PCR techniques, or any other method or procedure known to one skilled in the art. In some embodiments, the procedure involves the ligation of two different DNA sequences (See, for example, "Current Protocols in Molecular Biology", eds. Ausubel et al., John Wiley & Sons, 1992).

[0200] In one embodiment, polynucleotides of the present invention which encode the engineered coagulation factors are inserted into expression vectors (i.e., a nucleic acid construct) to enable expression of the recombinant polypeptide. In one embodiment, the expression vector of the present invention includes additional sequences which render this vector suitable for replication and integration in prokaryotes. In one embodiment, the expression vector of the present invention includes additional sequences which render this vector suitable for replication and integration in eukaryotes. In one embodiment, the expression vector of the present invention includes a shuttle vector which renders this vector suitable for replication and integration in both prokaryotes and eukaryotes. In some embodiments, cloning vectors comprise transcription and translation initiation sequences (e.g., promoters, enhances) and transcription and translation terminators (e.g., polyadenylation signals).

[0201] In one embodiment, a variety of prokaryotic or eukaryotic cells can be used as host-expression systems to express the coagulation factors of the present invention. In some embodiments, these include, but are not limited to, microorganisms, such as bacteria transformed with a recombinant bacteriophage DNA, plasmid DNA or cosmid DNA

expression vector containing the polypeptide coding sequence; yeast transformed with recombinant yeast expression vectors containing the polypeptide coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression
5 vectors, such as Ti plasmid, containing the polypeptide coding sequence.

[0202] In some embodiments, non-bacterial expression systems are used (e.g. mammalian expression systems such as CHO cells) to express the coagulation factors of the present invention. In one embodiment, the expression vector used to express polynucleotides of the present invention in mammalian cells is pCI-DHFR vector comprising a CMV promoter and a
10 neomycin resistance gene. Construction of the pCI-dhfr vector is described, according to one embodiment, in Example 1.

[0203] In some embodiments, in bacterial systems of the present invention, a number of expression vectors can be advantageously selected depending upon the use intended for the polypeptide expressed. In one embodiment, large quantities of polypeptide are desired. In one
15 embodiment, vectors that direct the expression of high levels of the protein product, possibly as a fusion with a hydrophobic signal sequence, which directs the expressed product into the periplasm of the bacteria or the culture medium where the protein product is readily purified are desired. In one embodiment, certain fusion proteins are engineered with a specific cleavage site to aid in recovery of the polypeptide. In one embodiment, vectors adaptable to
20 such manipulation include, but are not limited to, the pET series of E. coli expression vectors [Studier et al., Methods in Enzymol. 185:60-89 (1990)].

[0204] In one embodiment, yeast expression systems are used. In one embodiment, a number of vectors containing constitutive or inducible promoters can be used in yeast as disclosed in U.S. Pat. Application. No: 5,932,447, which is incorporated by reference herein
25 in its entirety. In another embodiment, vectors which promote integration of foreign DNA sequences into the yeast chromosome are used.

[0205] In one embodiment, the expression vector of the present invention can further include additional polynucleotide sequences that allow, for example, the translation of several proteins from a single mRNA such as an internal ribosome entry site (IRES) and sequences
30 for genomic integration of the promoter-chimeric polypeptide.

[0206] In some embodiments, mammalian expression vectors include, but are not limited to, pcDNA3, pcDNA3.1(+/-), pGL3, pZeoSV2(+/-), pSecTag2, pDisplay, pEF/myc/cyto, pCMV/myc/cyto, pCR3.1, pSinRep5, DH26S, DHBB, pNMT1, pNMT41, pNMT81, which are available from Invitrogen, pCI which is available from Promega, pMbac, pPbac, pBK-

RSV and pBK-CMV which are available from Strategene, pTRES which is available from Clontech, and their derivatives.

[0207] In some embodiments, expression vectors containing regulatory elements from eukaryotic viruses such as retroviruses are used in the present invention. SV40 vectors include pSVT7 and pMT2. In some embodiments, vectors derived from bovine papilloma virus include pBV-1MTHA, and vectors derived from Epstein Bar virus include pHEBO, and p2O5. Other exemplary vectors include pMSG, pAV009/A+, pMTO10/A+, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of the SV-40 early promoter, SV-40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

[0208] In some embodiments, recombinant viral vectors are useful for in vivo expression of the coagulation factors of the present invention since they offer advantages such as lateral infection and targeting specificity. In one embodiment, lateral infection is inherent in the life cycle of, for example, a retrovirus and is the process by which a single infected cell produces many progeny virions that bud off and infect neighboring cells. In one embodiment, the result is that a large area becomes rapidly infected, most of which was not initially infected by the original viral particles. In one embodiment, viral vectors are produced that are unable to spread laterally. In one embodiment, this characteristic can be useful if the desired purpose is to introduce a specified gene into only a localized number of targeted cells.

[0209] In one embodiment, various methods can be used to introduce the expression vector of the present invention into cells. Such methods are generally described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Springs Harbor Laboratory, New York (1989, 1992), in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, Md. (1989), Chang et al., *Somatic Gene Therapy*, CRC Press, Ann Arbor, Mich. (1995), Vega et al., *Gene Targeting*, CRC Press, Ann Arbor Mich. (1995), *Vectors: A Survey of Molecular Cloning Vectors and Their Uses*, Butterworths, Boston Mass. (1988) and Gilboa et al. [*Biotechniques* 4 (6): 504-512, 1986] and include, for example, stable or transient transfection, lipofection, electroporation and infection with recombinant viral vectors. In addition, see U.S. Pat. Nos. 5,464,764 and 5,487,992, incorporated herein by reference, for positive-negative selection methods.

[0210] In some embodiments, introduction of nucleic acid by viral infection offers several advantages over other methods such as lipofection and electroporation, since higher transfection efficiency can be obtained due to the infectious nature of viruses.

[0211] In one embodiment, it will be appreciated that the engineered coagulation factors of the present invention can also be expressed from a nucleic acid construct administered to the individual employing any suitable mode of administration, described hereinabove (i.e., in vivo gene therapy). In one embodiment, the nucleic acid construct is introduced into a suitable cell via an appropriate gene delivery vehicle/method (transfection, transduction, homologous recombination, etc.) and an expression system as needed and then the modified cells are expanded in culture and returned to the individual (i.e., ex vivo gene therapy).

[0212] In one embodiment, plant expression vectors are used. In one embodiment, the expression of a polypeptide coding sequence is driven by a number of promoters. In some embodiments, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV [Brisson et al., *Nature* 310:511-514 (1984)], or the coat protein promoter to TMV [Takamatsu et al., *EMBO J.* 6:307-311 (1987)] are used. In another embodiment, plant promoters are used such as, for example, the small subunit of RUBISCO [Coruzzi et al., *EMBO J.* 3:1671-1680 (1984); and Brogli et al., *Science* 224:838-843 (1984)] or heat shock promoters, e.g., soybean hsp17.5-E or hsp17.3-B [Gurley et al., *Mol. Cell. Biol.* 6:559-565 (1986)]. In one embodiment, constructs are introduced into plant cells using Ti plasmid, Ri plasmid, plant viral vectors, direct DNA transformation, microinjection, electroporation and other techniques well known to the skilled artisan. See, for example, Weissbach & Weissbach [Methods for Plant Molecular Biology, Academic Press, NY, Section VIII, pp 421-463 (1988)]. Other expression systems such as insects and mammalian host cell systems, which are well known in the art, can also be used by the present invention.

[0213] It will be appreciated that other than containing the necessary elements for the transcription and translation of the inserted coding sequence (encoding the polypeptide), the expression construct of the present invention can also include sequences engineered to optimize stability, production, purification, yield or activity of the expressed polypeptide.

[0214] In some embodiments, transformed cells are cultured under effective conditions, which allow for the expression of high amounts of recombinant engineered coagulation factors. In some embodiments, effective culture conditions include, but are not limited to, effective media, bioreactor, temperature, pH and oxygen conditions that permit protein production. In one embodiment, an effective medium refers to any medium in which a cell is cultured to produce the recombinant polypeptide of the present invention. In some embodiments, a medium typically includes an aqueous solution having assimilable carbon, nitrogen and phosphate sources, and appropriate salts, minerals, metals and other nutrients, such as vitamins. In some embodiments, cells of the present invention can be cultured in

conventional fermentation bioreactors, shake flasks, test tubes, microtiter dishes and petri plates. In some embodiments, culturing is carried out at a temperature, pH and oxygen content appropriate for a recombinant cell. In some embodiments, the determination of culturing conditions are within the expertise of one of ordinary skill in the art.

5 [0215] In some embodiments, depending on the vector and host system used for production, resultant engineered coagulation factors of the present invention either remain within the recombinant cell, are secreted into the fermentation medium, are secreted into a space between two cellular membranes, such as the periplasmic space in *E. coli*; or are retained on the outer surface of a cell or viral membrane.

10 [0216] In one embodiment, following a predetermined time in culture, recovery of the recombinant engineered coagulation factor is effected.

[0217] In one embodiment, the phrase "recovering the recombinant engineered coagulation factor" used herein refers to collecting the whole fermentation medium containing the polypeptide and need not imply additional steps of separation or purification.

15 [0218] In one embodiment, engineered coagulation factors of the present invention are purified using a variety of standard protein purification techniques, such as, but not limited to, affinity chromatography, ion exchange chromatography, filtration, electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and differential
20 solubilization.

[0219] In one embodiment, to facilitate recovery, the expressed coding sequence can be engineered to encode the engineered coagulation factor of the present invention and fused cleavable moiety. In one embodiment, a fusion protein can be designed so that the polypeptide can be readily isolated by affinity chromatography; e.g., by immobilization on a
25 column specific for the cleavable moiety. In one embodiment, a cleavage site is engineered between the engineered coagulation factor and the cleavable moiety and the polypeptide can be released from the chromatographic column by treatment with an appropriate enzyme or agent that specifically cleaves the fusion protein at this site [e.g., see Booth et al., *Immunol. Lett.* 19:65-70 (1988); and Gardella et al., *J. Biol. Chem.* 265:15854-15859 (1990)].

30 [0220] In one embodiment, the engineered coagulation factor of the present invention is retrieved in "substantially pure" form.

[0221] In one embodiment, the phrase "substantially pure" refers to a purity that allows for the effective use of the protein in the applications described herein.

[0222] In one embodiment, the engineered coagulation factor of the present invention can also be synthesized using in vitro expression systems. In one embodiment, in vitro synthesis methods are well known in the art and the components of the system are commercially available.

5 [0223] In some embodiments, the recombinant engineered coagulation factors are synthesized and purified; their therapeutic efficacy can be assayed either in vivo or in vitro. In one embodiment, the binding activities of the recombinant engineered coagulation factors of the present invention can be ascertained using various assays as known to one of skill in the art.

10 [0224] In another embodiment, the engineered coagulation factor of the present invention can be provided to the individual per se. In one embodiment, the engineered coagulation factor of the present invention can be provided to the individual as part of a pharmaceutical composition where it is mixed with a pharmaceutically acceptable carrier.

[0225] In another embodiment, a "pharmaceutical composition" refers to a preparation of
15 one or more of the active ingredients described herein with other chemical components such as physiologically suitable carriers and excipients. The purpose of a pharmaceutical composition is to facilitate administration of a compound to an organism.

[0226] In another embodiment, "active ingredient" refers to the polypeptide sequence of interest, which is accountable for the biological effect.

20 [0227] In another embodiment, any of the compositions of the present invention will comprise at least one CTP sequence bound only to the carboxy terminus of an engineered coagulation factor of interest, in any form. In one embodiment, the present invention provides combined preparations. In one embodiment, "a combined preparation" defines especially a "kit of parts" in the sense that the combination partners as defined above can be dosed
25 independently or by use of different fixed combinations with distinguished amounts of the combination partners i.e., simultaneously, concurrently, separately or sequentially. In some embodiments, the parts of the kit of parts can then, e.g., be administered simultaneously or chronologically staggered, that is at different time points and with equal or different time intervals for any part of the kit of parts. The ratio of the total amounts of the combination
30 partners, in some embodiments, can be administered in the combined preparation. In one embodiment, the combined preparation can be varied, e.g., in order to cope with the needs of a patient subpopulation to be treated or the needs of the single patient which different needs can be due to a particular disease, severity of a disease, age, sex, or body weight as can be readily made by a person skilled in the art.

[0228] In another embodiment, the phrases "physiologically acceptable carrier" and "pharmaceutically acceptable carrier" which are interchangeably used refer to a carrier or a diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered compound. An adjuvant is included under these phrases. In one embodiment, one of the ingredients included in the pharmaceutically acceptable carrier can be for example polyethylene glycol (PEG), a biocompatible polymer with a wide range of solubility in both organic and aqueous media (Mutter et al. (1979)).

[0229] In another embodiment, "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate administration of an active ingredient. In one embodiment, excipients include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

[0230] Techniques for formulation and administration of drugs are found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition, which is incorporated herein by reference.

[0231] Various embodiments of dosage ranges are contemplated by this invention. The dosage of the engineered coagulation factor of the present invention, in one embodiment, is in the range of 0.005-100 mg/day. In another embodiment, the dosage is in the range of 0.005-5 mg/day. In another embodiment, the dosage is in the range of 0.01-50 mg/day. In another embodiment, the dosage is in the range of 0.1-20 mg/day. In another embodiment, the dosage is in the range of 0.1-10 mg/day. In another embodiment, the dosage is in the range of 0.01-5 mg/day. In another embodiment, the dosage is in the range of 0.001-0.01 mg/day. In another embodiment, the dosage is in the range of 0.001-0.1 mg/day. In another embodiment, the dosage is in the range of 0.1-5 mg/day. In another embodiment, the dosage is in the range of 0.5-50 mg/day. In another embodiment, the dosage is in the range of 0.2-15mg/day. In another embodiment, the dosage is in the range of 0.8-65 mg/day. In another embodiment, the dosage is in the range of 1-50 mg/day. In another embodiment, the dosage is in the range of 5-10 mg/day. In another embodiment, the dosage is in the range of 8-15 mg/day. In another embodiment, the dosage is in a range of 10-20mg/day. In another embodiment, the dosage is in the range of 20-40 mg/day. In another embodiment, the dosage is in a range of 60-120 mg/day. In another embodiment, the dosage is in the range of 12-40 mg/day. In another embodiment, the dosage is in the range of 40-60 mg/day. In another embodiment, the dosage is in a range of 50-100mg/day. In another embodiment, the dosage is in a range of 1-60 mg/day. In another embodiment, the dosage is in the range of 15-25 mg/day. In another

embodiment, the dosage is in the range of 5-10 mg/day. In another embodiment, the dosage is in the range of 55-65 mg/day.

[0232] In another embodiment, the dosage is in a range of 50-500 mg/day. In another embodiment, the dosage is in a range of 50-150 mg/day. In another embodiment, the dosage is in a range of 100-200 mg/day. In another embodiment, the dosage is in a range of 150-250 mg/day. In another embodiment, the dosage is in a range of 200-300 mg/day. In another embodiment, the dosage is in a range of 250-400 mg/day. In another embodiment, the dosage is in a range of 300-500 mg/day. In another embodiment, the dosage is in a range of 350-500 mg/day.

[0233] In one embodiment, the dosage is 20 mg/day. In one embodiment, the dosage is 30 mg/day. In one embodiment, the dosage is 40 mg/day. In one embodiment, the dosage is 50 mg/day. In one embodiment, the dosage is 0.01 mg/day. In another embodiment, the dosage is 0.1 mg/day. In another embodiment, the dosage is 1 mg/day. In another embodiment, the dosage is 0.530 mg/day. In another embodiment, the dosage is 0.05 mg/day. In another embodiment, the dosage is 50 mg/day. In another embodiment, the dosage is 10 mg/day. In another embodiment, the dosage is 20-70 mg/day. In another embodiment, the dosage is 5 mg/day.

[0234] In one embodiment, the dosage of the CTP-modified coagulation factor is 1-5 mg/day. In one embodiment, the dosage of the CTP-modified coagulation factor is 1-3 mg/day. In another embodiment, the dosage of the CTP-modified coagulation factor is 2 mg/day.

[0235] In another embodiment, the dosage is 1-90 mg/day. In another embodiment, the dosage is 1-90 mg/2 days. In another embodiment, the dosage is 1-90 mg/3 days. In another embodiment, the dosage is 1-90 mg/4 days. In another embodiment, the dosage is 1-90 mg/5 days. In another embodiment, the dosage is 1-90 mg/6 days. In another embodiment, the dosage is 1-90 mg/week. In another embodiment, the dosage is 1-90 mg/9 days. In another embodiment, the dosage is 1-90 mg/11 days. In another embodiment, the dosage is 1-90 mg/14 days.

[0236] In another embodiment, the coagulation factor dosage is 10-50 mg/day. In another embodiment, the dosage is 10-50 mg/2 days. In another embodiment, the dosage is 10-50 mg/3 days. In another embodiment, the dosage is 10-50 mg/4 days. In another embodiment, the dosage is 10-50 micrograms mg/5 days. In another embodiment, the dosage is 10-50 mg/6 days. In another embodiment, the dosage is 10-50 mg/week. In another embodiment, the

dosage is 10-50 mg/9 days. In another embodiment, the dosage is 10-50 mg/11 days. In another embodiment, the dosage is 10-50 mg/14 days.

[0237] In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is formulated in an intranasal dosage form. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is formulated in an injectable dosage form. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 0.0001 mg to 0.6 mg. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 0.001 mg to 0.005 mg. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 0.005 mg to 0.01 mg. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 0.01 mg to 0.3 mg. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 0.2 mg to 0.6 mg. In another embodiment, the coagulation factor is free of CTPs on its amino terminus.

[0238] In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 1-100 micrograms. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 10-80 micrograms. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 20-60 micrograms. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 10-50 micrograms. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 40-80 micrograms. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 10-30 micrograms. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 30-60 micrograms.

[0239] In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 0.2 mg to 2 mg. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 2 mg to 6 mg. In another embodiment, a

polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 4 mg to 10 mg. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject in a dose ranging from 5 mg and 15 mg.

5 [0240] In one embodiment, the dosage of the CTP-modified FIX comprises 50% of the amount of FIX administered in the recommended dosage of recombinant FIX (e.g., Benefix®, Wyeth or Mononine®, CSL Behring) to patients over the same period of time. In one embodiment, the dosage of the CTP-modified FVIIa comprises 50% of the amount of FVIIa administered in the recommended dosage of recombinant FVIIa (e.g., NovoSeven®) to
10 patients over the same period of time. In one embodiment, the dosage of the CTP-modified FVII comprises 50% of the amount of FVII administered in the recommended dosage of recombinant FVII to patients over the same period of time. For example, if NovoSeven® is given at a dose of 90 mcg/kg every two hours to a patient pre- or post-operatively (i.e., 7.65 mg every two hours or 45.9 mg in six doses over a 12 hour period, for an 85 kg patient), a
15 CTP-modified coagulation factor of the present invention may be given at a dose that is 50% of the patient's 12-hour dose of recombinant FVIIa (i.e., at a dose of 23 mg given once over a 12-hour period).

[0241] In another embodiment, the dosage of CTP-modified coagulation factor is such that it contains 45% of the amount of the coagulation factor than that administered using the non-
20 CTP-modified coagulation factor. In another embodiment, the dosage of CTP-modified coagulation factor is such that it contains 10% of the amount of the coagulation factor than that administered using the non-CTP-modified coagulation factor. In another embodiment, the dosage of CTP-modified coagulation factor is such that it contains 25% of the amount of the coagulation factor than that administered using the non-CTP-modified coagulation factor. In
25 another embodiment, the dosage of CTP-modified coagulation factor is such that it contains 35% of the amount of the coagulation factor than that administered using the non-CTP-modified coagulation factor. In another embodiment, the dosage of CTP-modified coagulation factor is such that it contains 75% of the amount of the coagulation factor than that administered using the non-CTP-modified coagulation factor. In another embodiment, the
30 dosage of CTP-modified coagulation factor is such that it contains 100% of the amount of the coagulation factor than that administered using the non-CTP-modified coagulation factor. However, even if the dosage contains the same amount of coagulation factor (e.g. FIX) as non-CTP-modified coagulation factor, it is still advantageous to subjects in that it will be

administered less frequently because of its increased half-life compared to recombinant coagulation factors.

[0242] In another embodiment, a therapeutically effective amount of a conjugated coagulation factor is between 50-500 IU per kg body weight administered once a day to once a week for FIX or 10µg/Kg-500µg/Kg for FVIIa. In another embodiment, a therapeutically effective amount of a conjugated coagulation factor is 150-250 IU per kg body weight, administered once a day. In another embodiment, a pharmaceutical composition comprising a conjugated coagulation factor is formulated at a strength effective for administration by various means to a human patient.

[0243] In one embodiment, FIX is administered in an amount effective to bring circulating Factor IX activity to 20-30 IU/dL in a subject. In another embodiment, FIX is administered in an amount effective to bring circulating Factor IX activity to 25-50 IU/dL in a subject. In another embodiment, FIX is administered in an amount effective to bring circulating Factor IX activity to 50-100 IU/dL in a subject. In another embodiment, FIX is administered in an amount effective to bring circulating Factor IX activity to 100-200 IU/dL in a subject. In another embodiment, FIX is administered in an amount effective to bring circulating Factor IX activity to 10-50 IU/dL in a subject. In another embodiment, FIX is administered in an amount effective to bring circulating Factor IX activity to 20-100 IU/dL in a subject.

[0244] In one embodiment, the CTP-modified coagulation factor is administered to a subject on a weekly basis. In another embodiment, the CTP-modified coagulation factor is administered to a subject twice a week. In another embodiment, the CTP-modified coagulation factor is administered to a subject on a fortnightly (once every two weeks) basis. In another embodiment, the CTP-modified coagulation factor is administered to a subject twice a month. In another embodiment, the CTP-modified coagulation factor is administered to a subject once a month. In another embodiment, the CTP-modified coagulation factor is administered to a subject on a daily basis. In another embodiment, the CTP-modified coagulation factor is administered to a subject every two days.

[0245] In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject once every three days. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject once every four days. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject once every five days. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject once every six days. In another embodiment, a polypeptide

comprising a coagulation factor and at least one CTP unit is administered to a subject once every 7-14 days. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject once every 10-20 days. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject once every 5-15 days. In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is administered to a subject once every 15-30 days.

[0246] In another embodiment, the methods of the invention include increasing the compliance in the use of coagulation factor therapy, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the carboxy terminus of the coagulation factor, thereby increasing compliance in the use of coagulation factor therapy.

[0247] In another embodiment, the methods of the invention include increasing the compliance of patients afflicted with chronic illnesses that are in need of a coagulation factor therapy. In another embodiment, the methods of the invention enable reduction in the dosing frequency of a coagulation factor by modifying the coagulation factor with CTPs as described hereinabove.

[0248] In another embodiment, the present invention provides a method of reducing the dosing frequency of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby reducing the dosing frequency of said FIX polypeptide. In another embodiment, the present invention provides a method of reducing the dosing frequency of a Factor VIIa (FVIIa) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa polypeptide, thereby reducing the dosing frequency of said FVIIa polypeptide.

[0249] In another embodiment, the term compliance comprises adherence. In another embodiment, the methods of the invention include increasing the compliance of patients in need of a coagulation factor therapy by reducing the frequency of administration of the coagulation factor. In another embodiment, reduction in the frequency of administration of the coagulation factor is achieved due to the CTP modifications which render the CTP-modified coagulation factor more stable. In another embodiment, reduction in the frequency of administration of the coagulation factor is achieved as a result of increasing $T_{1/2}$ of the coagulation factor. In another embodiment, reduction in the frequency of administration of the

coagulation factor is achieved as a result of increasing the clearance time or reducing the clearance rate of the coagulation factor.

[0250] In another embodiment, the present invention provides a method of reducing the clearance rate of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said
5 FIX polypeptide, thereby reducing the clearance rate of said FIX polypeptide. In another embodiment, the present invention provides a method of reducing the clearance rate of a Factor VIIa (FVIIa) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa
10 polypeptide, thereby reducing the clearance rate of said FVIIa polypeptide.

[0251] In another embodiment, reduction in the frequency of administration of the coagulation factor is achieved as a result of increasing the AUC measure of the coagulation factor.

[0252] In another embodiment, provided herein is a method of reducing the dosing
15 frequency of a coagulation factor, comprising the step of attaching one to ten CTPs to the carboxy terminus of the coagulation factor, thereby reducing a dosing frequency of the coagulation factor. In another embodiment, provided herein is a method of reducing the dosing frequency of a coagulation factor, comprising the step of attaching one to five CTPs to the carboxy terminus of the coagulation factor, thereby reducing a dosing frequency of the
20 coagulation factor. In another embodiment, provided herein is a method of reducing the dosing frequency of a coagulation factor, comprising the step of attaching three CTPs to the carboxy terminus of the coagulation factor, thereby reducing a dosing frequency of the coagulation factor. In another embodiment, provided herein is a method of reducing the dosing frequency of a coagulation factor, comprising the step of attaching three to five CTPs
25 to the carboxy terminus of the coagulation factor, thereby reducing a dosing frequency of the coagulation factor.

[0253] In another embodiment, provided herein is a method of increasing compliance in the use of coagulation factor therapy, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and one to ten chorionic gonadotrophin carboxy
30 terminal peptides attached to the carboxy terminus of a coagulation factor, thereby increasing compliance in the use of coagulation factor therapy. In another embodiment, provided herein is a method of increasing compliance in the use of coagulation factor therapy, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and one to five chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of

a coagulation factor, thereby increasing compliance in the use of coagulation factor therapy. In another embodiment, provided herein is a method of increasing compliance in the use of coagulation factor therapy, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and three chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby increasing compliance in the use of coagulation factor therapy. In another embodiment, provided herein is a method of increasing compliance in the use of coagulation factor therapy, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and three to five chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby increasing compliance in the use of coagulation factor therapy.

[0254] In another embodiment, provided herein is a method of preventing or treating a blood clotting or coagulation disorder in a subject, comprising providing to said subject a polypeptide comprising a coagulation factor and one to ten chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby treating a blood clotting or coagulation disorder in said subject. In another embodiment, provided herein is a method of preventing or treating a blood clotting or coagulation disorder in a subject, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and one to five chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby preventing or treating a blood clotting or coagulation disorder in said subject. In another embodiment, provided herein is a method of preventing or treating a blood clotting or coagulation disorder in a subject, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and three chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby preventing or treating a blood clotting or coagulation disorder in said subject. In another embodiment, provided herein is a method of preventing or treating a blood clotting or coagulation disorder in a subject, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and three to five chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby preventing or treating a blood clotting or coagulation disorder in said subject.

[0255] In another embodiment, provided herein is a method of preventing hemophilia in a subject, comprising providing to said subject a polypeptide comprising a coagulation factor and one to ten chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby preventing hemophilia in said subject. In another embodiment, provided herein is a method of preventing hemophilia in a subject, comprising

providing to a subject in need thereof, a polypeptide comprising a coagulation factor and one to five chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby preventing hemophilia in said subject. In another embodiment, provided herein is a method of preventing hemophilia in a subject, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and three chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby preventing hemophilia in said subject. In another embodiment, provided herein is a method of preventing hemophilia in a subject, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and three to five chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby preventing hemophilia in said subject.

[0256] In another embodiment, the present invention shows that the compositions provided herein are surprisingly more effectively absorbed into the bloodstream after SC administration (see Examples 7-9 herein). To be able to administer FVIIa subcutaneously serves as an advantage as it can be used for prophylactic applications. Subcutaneous injections are also much easier for patients to self-inject, and are advantage when the patients are very young and their veins are small and difficult to find.

[0257] In another embodiment, provided herein is a method of treating hemophilia in a subject, comprising providing to said subject a polypeptide comprising a coagulation factor and one to ten chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby treating hemophilia in said subject. In another embodiment, provided herein is a method of treating hemophilia in a subject, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and one to five chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby treating hemophilia in said subject. In another embodiment, provided herein is a method of treating hemophilia in a subject, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and three chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby treating hemophilia in said subject. In another embodiment, provided herein is a method of treating hemophilia in a subject, comprising providing to a subject in need thereof, a polypeptide comprising a coagulation factor and three to five chorionic gonadotrophin carboxy terminal peptides attached to the carboxy terminus of a coagulation factor, thereby treating hemophilia in said subject.

[0258] Oral administration, in one embodiment, comprises a unit dosage form comprising tablets, capsules, lozenges, chewable tablets, suspensions, emulsions and the like. Such unit dosage forms comprise a safe and effective amount of the desired coagulation factor of the invention, each of which is in one embodiment, from about 0.7 or 3.5 mg to about 280 mg/70 kg, or in another embodiment, about 0.5 or 10 mg to about 210 mg/70 kg. The pharmaceutically-acceptable carriers suitable for the preparation of unit dosage forms for peroral administration are well-known in the art. In some embodiments, tablets typically comprise conventional pharmaceutically-compatible adjuvants as inert diluents, such as calcium carbonate, sodium carbonate, mannitol, lactose and cellulose; binders such as starch, gelatin and sucrose; disintegrants such as starch, alginic acid and croscarmellose; lubricants such as magnesium stearate, stearic acid and talc. In one embodiment, glidants such as silicon dioxide can be used to improve flow characteristics of the powder-mixture. In one embodiment, coloring agents, such as the FD&C dyes, can be added for appearance. Sweeteners and flavoring agents, such as aspartame, saccharin, menthol, peppermint, and fruit flavors, are useful adjuvants for chewable tablets. Capsules typically comprise one or more solid diluents disclosed above. In some embodiments, the selection of carrier components depends on secondary considerations like taste, cost, and shelf stability, which are not critical for the purposes of this invention, and can be readily made by a person skilled in the art.

[0259] In one embodiment, the oral dosage form comprises predefined release profile. In one embodiment, the oral dosage form of the present invention comprises an extended release tablets, capsules, lozenges or chewable tablets. In one embodiment, the oral dosage form of the present invention comprises a slow release tablets, capsules, lozenges or chewable tablets. In one embodiment, the oral dosage form of the present invention comprises an immediate release tablets, capsules, lozenges or chewable tablets. In one embodiment, the oral dosage form is formulated according to the desired release profile of the pharmaceutical active ingredient as known to one skilled in the art.

[0260] Peroral compositions, in some embodiments, comprise liquid solutions, emulsions, suspensions, and the like. In some embodiments, pharmaceutically-acceptable carriers suitable for preparation of such compositions are well known in the art. In some embodiments, liquid oral compositions comprise from about 0.001% to about 0.933% of the desired compound or compounds, or in another embodiment, from about 0.01% to about 10%.

[0261] In some embodiments, compositions for use in the methods of this invention comprise solutions or emulsions, which in some embodiments are aqueous solutions or emulsions comprising a safe and effective amount of the compounds of the present invention

and optionally, other compounds, intended for topical intranasal administration. In some embodiments, the compositions comprise from about 0.001% to about 10.0% w/v of a subject compound, more preferably from about 0.1% to about 2.0, which is used for systemic delivery of the compounds by the intranasal route.

5 [0262] In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is injected into the muscle (intramuscular injection). In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is injected below the skin (subcutaneous injection). In another embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is injected into the muscle. In another
10 embodiment, a polypeptide comprising a coagulation factor and at least one CTP unit is injected into the skin. In another embodiment, a coagulation factor as described herein is administered via systemic administration. In another embodiment, a coagulation factor as described herein is administered by intravenous injection. In another embodiment, administration can be parenteral, pulmonary, oral, topical, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, transnasal, intraocular, ophthalmic,
15 epidural, buccal, rectal, transmucosal, intestinal or parenteral delivery, including intramedullary injections as well as intrathecal or direct intraventricular administration.

[0263] In another embodiment, the preparation is administered in a local rather than systemic manner, for example, via injection of the preparation directly into a specific region
20 of a patient's body.

[0264] In one embodiment, the route of administration may be enteral. In another embodiment, the route may be conjunctival, transdermal, intradermal, intra-arterial, vaginal, rectal, intratumoral, paracancerous, transmucosal, intramuscular, intravascular, intraventricular, intracranial, intra-nasal, sublingual, or a combination thereof.

25 [0265] In another embodiment, the pharmaceutical compositions are administered by intravenous, intra-arterial, or intramuscular injection of a liquid preparation. In some embodiments, liquid formulations include solutions, suspensions, dispersions, emulsions, oils and the like. In one embodiment, the pharmaceutical compositions are administered intravenously, and are thus formulated in a form suitable for intravenous administration. In
30 another embodiment, the pharmaceutical compositions are administered intra-arterially, and are thus formulated in a form suitable for intra-arterial administration. In another embodiment, the pharmaceutical compositions are administered intramuscularly, and are thus formulated in a form suitable for intramuscular administration.

[0266] Further, in another embodiment, the pharmaceutical compositions are administered topically to body surfaces, and are thus formulated in a form suitable for topical administration. Suitable topical formulations include gels, ointments, creams, lotions, drops and the like. For topical administration, the compounds of the present invention are combined
5 with an additional appropriate therapeutic agent or agents, prepared and applied as solutions, suspensions, or emulsions in a physiologically acceptable diluent with or without a pharmaceutical carrier.

[0267] In one embodiment, pharmaceutical compositions of the present invention are manufactured by processes well known in the art, e.g., by means of conventional mixing,
10 dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

[0268] In one embodiment, pharmaceutical compositions for use in accordance with the present invention is formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the
15 active ingredients into preparations which, can be used pharmaceutically. In one embodiment, formulation is dependent upon the route of administration chosen.

[0269] In one embodiment, injectables of the invention are formulated in aqueous solutions. In one embodiment, injectables of the invention are formulated in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological salt buffer. In some
20 embodiments, for transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

[0270] In one embodiment, the preparations described herein are formulated for parenteral administration, e.g., by bolus injection or continuous infusion. In some embodiments, formulations for injection are presented in unit dosage form, e.g., in ampoules or in multidose
25 containers with optionally, an added preservative. In some embodiments, compositions are suspensions, solutions or emulsions in oily or aqueous vehicles, and contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

[0271] The compositions also comprise, in some embodiments, preservatives, such as benzalkonium chloride and thimerosal and the like; chelating agents, such as edetate sodium
30 and others; buffers such as phosphate, citrate and acetate; tonicity agents such as sodium chloride, potassium chloride, glycerin, mannitol and others; antioxidants such as ascorbic acid, acetylcystine, sodium metabisulfite and others; aromatic agents; viscosity adjustors, such as polymers, including cellulose and derivatives thereof; and polyvinyl alcohol and acid and bases to adjust the pH of these aqueous compositions as needed. The compositions also

comprise, in some embodiments, local anesthetics or other actives. The compositions can be used as sprays, mists, drops, and the like.

[0272] In some embodiments, pharmaceutical compositions for parenteral administration include aqueous solutions of the active preparation in water-soluble form. Additionally, suspensions of the active ingredients, in some embodiments, are prepared as appropriate oil or water based injection suspensions. Suitable lipophilic solvents or vehicles include, in some embodiments, fatty oils such as sesame oil, or synthetic fatty acid esters such as ethyl oleate, triglycerides or liposomes. Aqueous injection suspensions contain, in some embodiments, substances, which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol or dextran. In another embodiment, the suspension also contains suitable stabilizers or agents which increase the solubility of the active ingredients to allow for the preparation of highly concentrated solutions.

[0273] In another embodiment, the active compound can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez- Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; J. E. Diederichs and al., *Pharm./nd.* 56 (1994) 267- 275).

[0274] In another embodiment, the pharmaceutical composition delivered in a controlled release system is formulated for intravenous infusion, implantable osmotic pump, transdermal patch, liposomes, or other modes of administration. In one embodiment, a pump is used (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989). In another embodiment, polymeric materials can be used. In yet another embodiment, a controlled release system can be placed in proximity to the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138 (1984). Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

[0275] In some embodiments, the active ingredient is in powder form for constitution with a suitable vehicle, e.g., sterile, pyrogen-free water based solution, before use. Compositions are formulated, in some embodiments, for atomization and inhalation administration. In another embodiment, compositions are contained in a container with attached atomizing means.

[0276] In one embodiment, the preparation of the present invention is formulated in rectal compositions such as suppositories or retention enemas, using, e.g., conventional suppository bases such as cocoa butter or other glycerides.

[0277] In some embodiments, pharmaceutical compositions suitable for use in context of the present invention include compositions wherein the active ingredients are contained in an amount effective to achieve the intended purpose. In some embodiments, a therapeutically effective amount means an amount of active ingredients effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated.

[0278] In one embodiment, determination of a therapeutically effective amount is well within the capability of those skilled in the art.

[0279] Some examples of substances which can serve as pharmaceutically-acceptable carriers or components thereof are sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose, and methyl cellulose; powdered tragacanth; malt; gelatin; talc; solid lubricants, such as stearic acid and magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols such as propylene glycol, glycerine, sorbitol, mannitol, and polyethylene glycol; alginic acid; emulsifiers, such as the Tween™ brand emulsifiers; wetting agents, such sodium lauryl sulfate; coloring agents; flavoring agents; tableting agents, stabilizers; antioxidants; preservatives; pyrogen-free water; isotonic saline; and phosphate buffer solutions. The choice of a pharmaceutically-acceptable carrier to be used in conjunction with the compound is basically determined by the way the compound is to be administered. If the subject compound is to be injected, in one embodiment, the pharmaceutically-acceptable carrier is sterile, physiological saline, with a blood-compatible suspending agent, the pH of which has been adjusted to about 7.4.

[0280] In addition, the compositions further comprise binders (e.g. acacia, cornstarch, gelatin, carbomer, ethyl cellulose, guar gum, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, povidone), disintegrating agents (e.g. cornstarch, potato starch, alginic acid, silicon dioxide, croscarmellose sodium, crospovidone, guar gum, sodium starch glycolate), buffers (e.g., Tris-HCl, acetate, phosphate) of various pH and ionic strength, additives such as albumin or gelatin to prevent absorption to surfaces, detergents (e.g., Tween 20, Tween 80, Pluronic F68, bile acid salts), protease inhibitors, surfactants (e.g. sodium lauryl sulfate), permeation enhancers, solubilizing agents (e.g., glycerol, polyethylene glycerol), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite, butylated hydroxyanisole), stabilizers (e.g.

hydroxypropyl cellulose, hydroxypropylmethyl cellulose), viscosity increasing agents (e.g. carbomer, colloidal silicon dioxide, ethyl cellulose, guar gum), sweeteners (e.g. aspartame, citric acid), preservatives (e.g., Thimerosal, benzyl alcohol, parabens), lubricants (e.g. stearic acid, magnesium stearate, polyethylene glycol, sodium lauryl sulfate), flow-aids (e.g. colloidal silicon dioxide), plasticizers (e.g. diethyl phthalate, triethyl citrate), emulsifiers (e.g. carbomer, hydroxypropyl cellulose, sodium lauryl sulfate), polymer coatings (e.g., poloxamers or poloxamines), coating and film forming agents (e.g. ethyl cellulose, acrylates, polymethacrylates) and/or adjuvants.

[0281] Typical components of carriers for syrups, elixirs, emulsions and suspensions include ethanol, glycerol, propylene glycol, polyethylene glycol, liquid sucrose, sorbitol and water. For a suspension, typical suspending agents include methyl cellulose, sodium carboxymethyl cellulose, cellulose (e.g. Avicel™, RC-591), tragacanth and sodium alginate; typical wetting agents include lecithin and polyethylene oxide sorbitan (e.g. polysorbate 80). Typical preservatives include methyl paraben and sodium benzoate. In another embodiment, peroral liquid compositions also contain one or more components such as sweeteners, flavoring agents and colorants disclosed above.

[0282] The compositions also include incorporation of the active material into or onto particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, hydrogels, etc, or onto liposomes, microemulsions, micelles, unilamellar or multilamellar vesicles, erythrocyte ghosts, or spheroplasts.) Such compositions will influence the physical state, solubility, stability, rate of in vivo release, and rate of in vivo clearance.

[0283] Also comprehended by the invention are particulate compositions coated with polymers (e.g. poloxamers or poloxamines) and the compound coupled to antibodies directed against tissue-specific receptors, ligands or antigens or coupled to ligands of tissue-specific receptors.

[0284] In some embodiments, compounds modified by the covalent attachment of water-soluble polymers such as polyethylene glycol, copolymers of polyethylene glycol and polypropylene glycol, carboxymethyl cellulose, dextran, polyvinyl alcohol, polyvinylpyrrolidone or polyproline. In another embodiment, the modified compounds exhibit substantially longer half-lives in blood following intravenous injection than do the corresponding unmodified compounds. In one embodiment, modifications also increase the compound's solubility in aqueous solution, eliminate aggregation, enhance the physical and chemical stability of the compound, and greatly reduce the immunogenicity and reactivity of the compound. In another embodiment, the desired in vivo biological activity is achieved by

the administration of such polymer-compound adducts less frequently or in lower doses than with the unmodified compound.

[0285] In some embodiments, preparation of effective amount or dose can be estimated initially from in vitro assays. In one embodiment, a dose can be formulated in animal models and such information can be used to more accurately determine useful doses in humans.

[0286] In one embodiment, toxicity and therapeutic efficacy of the active ingredients described herein can be determined by standard pharmaceutical procedures in vitro, in cell cultures or experimental animals. In one embodiment, the data obtained from these in vitro and cell culture assays and animal studies can be used in formulating a range of dosage for use in human. In one embodiment, the dosages vary depending upon the dosage form employed and the route of administration utilized. In one embodiment, the exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. [See e.g., Fingl, et al., (1975) "The Pharmacological Basis of Therapeutics", Ch. 1 p.1].

[0287] In one embodiment, depending on the severity and responsiveness of the condition to be treated, dosing can be of a single or a plurality of administrations, with course of treatment lasting from several days to several weeks or until cure is effected or diminution of the disease state is achieved.

[0288] In one embodiment, the amount of a composition to be administered will, of course, be dependent on the subject being treated, the severity of the affliction, the manner of administration, the judgment of the prescribing physician, etc.

[0289] In one embodiment, compositions including the preparation of the present invention formulated in a compatible pharmaceutical carrier are also prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

[0290] In another embodiment, a coagulation factor as described herein is lyophilized (i.e., freeze-dried) preparation in combination with complex organic excipients and stabilizers such as nonionic surface active agents (i.e., surfactants), various sugars, organic polyols and/or human serum albumin. In another embodiment, a pharmaceutical composition comprises a lyophilized coagulation factor as described in sterile water for injection. In another embodiment, a pharmaceutical composition comprises a lyophilized coagulation factor as described in sterile PBS for injection. In another embodiment, a pharmaceutical composition comprises a lyophilized coagulation factor as described in sterile 0.9% NaCl for injection.

[0291] In another embodiment, the pharmaceutical composition comprises a coagulation factor as described herein and complex carriers such as human serum albumin, polyols,

sugars, and anionic surface active stabilizing agents. In another embodiment, the pharmaceutical composition comprises a coagulation factor as described herein and lactobionic acid and an acetate/glycine buffer. In another embodiment, the pharmaceutical composition comprises a coagulation factor as described herein and amino acids, such as arginine or glutamate that increase the solubility of interferon compositions in water. In another embodiment, the pharmaceutical composition comprises a lyophilized coagulation factor as described herein and glycine or human serum albumin (HSA), a buffer (e.g. acetate) and an isotonic agent (e.g. NaCl). In another embodiment, the pharmaceutical composition comprises a lyophilized coagulation factor as described herein and phosphate buffer, glycine and HSA.

[0292] In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein is stabilized when placed in buffered solutions having a pH between about 4 and 7.2. In another embodiment, the pharmaceutical composition comprising a coagulation factor is in a buffered solution having a pH between about 4 and 8.5. In another embodiment, the pharmaceutical composition comprising a coagulation factor is in a buffered solution having a pH between about 6 and 7. In another embodiment, the pharmaceutical composition comprising a coagulation factor is in a buffered solution having a pH of about 6.5. In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein is stabilized with an amino acid as a stabilizing agent and in some cases a salt (if the amino acid does not contain a charged side chain).

[0293] In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein is a liquid composition comprising a stabilizing agent at between about 0.3% and 5% by weight which is an amino acid.

[0294] In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein provides dosing accuracy and product safety. In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein provides a biologically active, stable liquid formulation for use in injectable applications. In another embodiment, the pharmaceutical composition comprises a non-lyophilized coagulation factor as described herein.

[0295] In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein provides a liquid formulation permitting storage for a long period of time in a liquid state facilitating storage and shipping prior to administration.

[0296] In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein comprises solid lipids as matrix material. In another embodiment,

the injectable pharmaceutical composition comprising a coagulation factor as described herein comprises solid lipids as matrix material. In another embodiment, the production of lipid microparticles by spray congealing was described by Speiser (Speiser and al., Pharm. Res. 8 (1991) 47-54) followed by lipid nanopellets for peroral administration (Speiser EP 0167825 (1990)). In another embodiment, lipids, which are used, are well tolerated by the body (e. g. glycerides composed of fatty acids which are present in the emulsions for parenteral nutrition).

[0297] In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein comprises polymeric microparticles. In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein comprises nanoparticles. In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein comprises liposomes. In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein comprises lipid emulsion. In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein comprises microspheres. In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein comprises lipid nanoparticles. In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein comprises lipid nanoparticles comprising amphiphilic lipids. In another embodiment, the pharmaceutical composition comprising a coagulation factor as described herein comprises lipid nanoparticles comprising a drug, a lipid matrix and a surfactant. In another embodiment, the lipid matrix has a monoglyceride content which is at least 50% w/w.

[0298] In one embodiment, compositions of the present invention are presented in a pack or dispenser device, such as an FDA approved kit, which contain one or more unit dosage forms containing the active ingredient. In one embodiment, the pack, for example, comprise metal or plastic foil, such as a blister pack. In one embodiment, the pack or dispenser device is accompanied by instructions for administration. In one embodiment, the pack or dispenser is accommodated by a notice associated with the container in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions or human or veterinary administration. Such notice, in one embodiment, is labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert.

[0299] In one embodiment, it will be appreciated that the coagulation factors of the present invention can be provided to the individual with additional active agents to achieve an

improved therapeutic effect as compared to treatment with each agent by itself. In another embodiment, measures (e.g., dosing and selection of the complementary agent) are taken to avoid adverse side effects which are associated with combination therapies.

[0300] In another embodiment, the present invention provides a CTP-modified Factor VIIa (FVIIa) polypeptide consisting of a FVIIa polypeptide and five gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa.

[0301] In another embodiment, the present invention provides a pharmaceutical composition comprising a CTP-modified Factor VIIa (FVIIa) polypeptide consisting of a FVIIa polypeptide and five gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa.

[0302] In another embodiment, the present invention provides a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor VIIa (FVIIa) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide.

[0303] In another embodiment, the present invention provides an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor VIIa (FVIIa) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide.

[0304] In another embodiment, the present invention provides a cell comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor VIIa (FVIIa) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide.

[0305] In another embodiment, the present invention provides a composition comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor VIIa (FVIIa) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide.

[0306] In another embodiment, the present invention provides a method of extending the biological half-life of a Factor VIIa (FVIIa) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa polypeptide, thereby extending the biological half-life of said FVIIa polypeptide.

[0307] In another embodiment, the present invention provides a method of improving the area under the curve (AUC) of a Factor VIIa (FVIIa) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa polypeptide, thereby improving the AUC of said FVIIa polypeptide.

[0308] In another embodiment, the present invention provides a method of reducing the dosing frequency of a Factor VIIa (FVIIa) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa polypeptide, thereby reducing the dosing frequency of said FVIIa polypeptide.

5 [0309] In another embodiment, the present invention provides a method of reducing the clearance rate of a Factor VIIa (FVIIa) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa polypeptide, thereby reducing the clearance rate of said FVIIa polypeptide.

[0310] In another embodiment, the present invention provides a method of producing a
10 CTP-modified Factor VIIa (FVIIa) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FVIIa polypeptide, thereby producing a CTP-modified FVIIa polypeptide.

[0311] In another embodiment, the present invention provides a method of treating hemophilia in a subject comprising administering a CTP-modified Factor VIIa (FVIIa)
15 polypeptide comprising a FVIIa polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FVIIa polypeptide to said subject, thereby treating hemophilia in said subject.

[0312]

[0313] In one embodiment, the present invention provides a CTP-modified Factor IX (FIX)
20 polypeptide consisting of a FIX polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said CTP-modified FIX polypeptide. In another embodiment, the present invention provides a CTP-modified FIX polypeptide, wherein the sequence of said CTP-modified FIX polypeptide is the sequence set forth in SEQ ID NO: 31. In another embodiment, the present invention provides a CTP-modified FIX
25 polypeptide, wherein at least one CTP is encoded by an amino acid sequence selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2. In another embodiment, the present invention provides a CTP-modified FIX polypeptide, wherein at least one CTP is glycosylated. In another embodiment, the present invention provides a CTP-modified FIX polypeptide, wherein at least one CTP is truncated. In another embodiment, the present
30 invention provides a CTP-modified FIX polypeptide, wherein at least one CTP is attached to said FIX polypeptide via a linker. In another embodiment, the present invention provides a CTP-modified FIX polypeptide, wherein said linker is a peptide bond.

[0314] In one embodiment, the present invention provides a pharmaceutical composition comprising the CTP-modified FIX polypeptide.

[0315] In one embodiment, the present invention provides a polynucleotide encoding a CTP-modified polypeptide consisting of a Factor IX (FIX) polypeptide and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide. In another embodiment, the present invention provides a polynucleotide, wherein the sequence of said polynucleotide is as set forth in SEQ ID NO: 30. In another embodiment, the present invention provides a polynucleotide, wherein at least one CTP is encoded by an amino acid sequence selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2. In another embodiment, the present invention provides a polynucleotide, wherein at least one CTP is glycosylated. In another embodiment, the present invention provides a polynucleotide, wherein at least one CTP is truncated. In another embodiment, the present invention provides a polynucleotide, wherein at least one CTP is attached to said FIX polypeptide via a linker. In another embodiment, the present invention provides a polynucleotide, wherein said linker is a peptide bond. An expression vector comprising the polynucleotide.

[0316] In one embodiment, the present invention provides a cell comprising the expression vector.

[0317] In one embodiment, the present invention provides a composition comprising the expression vector.

[0318] In one embodiment, the present invention provides a method of extending the biological half-life of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby extending the biological half-life of said FIX polypeptide. In another embodiment, the present invention provides a method, wherein at least one CTP is encoded by an amino acid sequence selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2. In another embodiment, the present invention provides a method, wherein at least one CTP is glycosylated. In another embodiment, the present invention provides a method, wherein at least one CTP is truncated. In another embodiment, the present invention provides a method, wherein at least one CTP is attached to said FIX polypeptide via a linker. In another embodiment, the present invention provides a method, wherein said linker is a peptide bond.

[0319] In one embodiment, the present invention provides a method of improving the area under the curve (AUC) of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby improving the AUC of said FIX polypeptide. In another

embodiment, the present invention provides a method, wherein at least one CTP is encoded by an amino acid sequence selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2. In another embodiment, the present invention provides a method, wherein at least one CTP is glycosylated. In another embodiment, the present invention provides a method, wherein at least one CTP is truncated. In another embodiment, the present invention provides a method, wherein at least one CTP is attached to said FIX polypeptide via a linker. In another embodiment, the present invention provides a method, wherein said linker is a peptide bond.

[0320] In one embodiment, the present invention provides a method of reducing the dosing frequency of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby reducing the dosing frequency of said FIX polypeptide. In another embodiment, the present invention provides a method, wherein at least one CTP is encoded by an amino acid sequence selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2. In another embodiment, the present invention provides a method, wherein at least one CTP is glycosylated. In another embodiment, the present invention provides a method, wherein at least one CTP is truncated. In another embodiment, the present invention provides a method, wherein at least one CTP is attached to said FIX polypeptide via a linker. In another embodiment, the present invention provides a method, wherein said linker is a peptide bond.

[0321] In one embodiment, the present invention provides a method of reducing the clearance rate of a Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby reducing the clearance rate of said FIX polypeptide. In another embodiment, the present invention provides a method, wherein at least one CTP is encoded by an amino acid sequence selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2. In another embodiment, the present invention provides a method, wherein at least one CTP is glycosylated. In another embodiment, the present invention provides a method, wherein at least one CTP is truncated. In another embodiment, the present invention provides a method, wherein at least one CTP is attached to said FIX polypeptide via a linker. In another embodiment, the present invention provides a method, wherein said linker is a peptide bond.

[0322] In one embodiment, the present invention provides a method of producing a CTP-modified Factor IX (FIX) polypeptide, comprising the step of attaching three chorionic

gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of said FIX polypeptide, thereby producing a CTP-modified FIX polypeptide. In another embodiment, the present invention provides a method, wherein the sequence of said CTP-modified FIX polypeptide is the sequence set forth in SEQ ID NO: 31. In another embodiment, the present invention provides a method, wherein at least one CTP is encoded by an amino acid sequence selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2. In another embodiment, the present invention provides a method, wherein at least one CTP is glycosylated. In another embodiment, the present invention provides a method, wherein at least one CTP is truncated. In another embodiment, the present invention provides a method, wherein at least one CTP is attached to said FIX polypeptide via a linker. In another embodiment, the present invention provides a method, wherein said linker is a peptide bond.

[0323] In one embodiment, the present invention provides a method of treating hemophilia in a subject comprising administering a CTP-modified Factor IX (FIX) polypeptide comprising a FIX polypeptide and three chorionic gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of said FIX polypeptide to said subject, thereby treating hemophilia in said subject. In another embodiment, the present invention provides a method, wherein the sequence of said CTP-modified FIX polypeptide is the sequence set forth in SEQ ID NO: 31. In another embodiment, the present invention provides a method, wherein at least one CTP is encoded by an amino acid sequence selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2. In another embodiment, the present invention provides a method, wherein at least one CTP is glycosylated. In another embodiment, the present invention provides a method, wherein at least one CTP is truncated. In another embodiment, the present invention provides a method, wherein at least one CTP is attached to said FIX polypeptide via a linker. In another embodiment, the present invention provides a method, wherein said linker is a peptide bond.

[0324] As is generally known in the art, the modified peptides and proteins of the invention may be coupled to labels, drugs, targeting agents, carriers, solid supports, and the like, depending on the desired application. The labeled forms of the modified biologicals may be used to track their metabolic fate; suitable labels for this purpose include, especially, radioisotope labels such as iodine 131, technetium 99, indium 111, and the like. The labels may also be used to mediate detection of the modified proteins or peptides in assay systems; in this instance, radioisotopes may also be used as well as enzyme labels, fluorescent labels, chromogenic labels, and the like. The use of such labels is particularly helpful if the peptide or protein is itself a targeting agent such as an antibody or a receptor ligand.

[0325] Similar linking techniques, along with others, may be employed to couple the modified peptides and proteins of the invention to solid supports. When coupled, these modified peptides and proteins can then be used as affinity reagents for the separation of desired components with which specific reaction is exhibited.

5 [0326] Finally, the modified peptides and proteins of the invention may be used to generate antibodies specifically immunoreactive with these new compounds. These antibodies are useful in a variety of diagnostic and therapeutic applications, depending on the nature of the biological activity of the unmodified peptide or protein. It is to be understood that the invention provides antibodies that are immunoreactive with CTP-modified FIX, FVII, or
10 FVIIa as described herein. In one embodiment, such antibodies may be used to distinguish or identify CTP-modified coagulation factors that were administered from endogenous coagulation factors. In another embodiment, the antibodies may be used to localize administered CTP-modified coagulation factors.

[0327] Additional objects, advantages, and novel features of the present invention will
15 become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

20

EXAMPLES

[0328] Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example,
25 "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989); "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. (1994); Ausubel et al., "Current Protocols in Molecular Biology", John Wiley and Sons, Baltimore, Maryland (1989); Perbal, "A Practical Guide to Molecular Cloning", John Wiley & Sons, New York (1988); Watson et al., "Recombinant DNA", Scientific American Books, New York; Birren et al. (eds) "Genome
30 Analysis: A Laboratory Manual Series", Vols. 1-4, Cold Spring Harbor Laboratory Press, New York (1998); methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057; "Cell Biology: A Laboratory Handbook", Volumes I-III Cellis, J. E., ed. (1994); "Culture of Animal Cells - A Manual of Basic Technique" by Freshney, Wiley-Liss, N. Y. (1994), Third Edition; "Current Protocols in Immunology"

Volumes I-III Coligan J. E., ed. (1994); Stites et al. (eds), "Basic and Clinical Immunology" (8th Edition), Appleton & Lange, Norwalk, CT (1994); Mishell and Shiigi (eds), "Selected Methods in Cellular Immunology", W. H. Freeman and Co., New York (1980); available immunoassays are extensively described in the patent and scientific literature, see, for example, U.S. Pat. Nos. 3,791,932; 3,839,153; 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; 4,098,876; 4,879,219; 5,011,771 and 5,281,521; "Oligonucleotide Synthesis" Gait, M. J., ed. (1984); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J., eds. (1985); "Transcription and Translation" Hames, B. D., and Higgins S. J., eds. (1984); "Animal Cell Culture" Freshney, R. I., ed. (1986); "Immobilized Cells and Enzymes" IRL Press, (1986); "A Practical Guide to Molecular Cloning" Perbal, B., (1984) and "Methods in Enzymology" Vol. 1-317, Academic Press; "PCR Protocols: A Guide To Methods And Applications", Academic Press, San Diego, CA (1990); Marshak et al., "Strategies for Protein Purification and Characterization - A Laboratory Course Manual" CSHL Press (1996); all of which are incorporated by reference.

Other general references are provided throughout this document.

EXAMPLE 1

Generation and Utilization of Coagulation Factor IX

- [0329] **Cloning and expression of recombinant FIX molecule:**
- [0330] Factor IX clones were constructed in our eukaryotic expression vector pCI-neo (Promega, catalog no. E1841). ORF Clone of Homo sapiens coagulation factor IX was ordered from "OriGene" (RC219065). Primers were ordered from **Sigma-Genosys**.
- [0331] **Construction of 301-1-pCI-neo-p200-11 (Factor IX-ctp x2):**
- [0332] Primer 101: 5' GTTTAGTGAACCGTCAGAAT 3' (SEQ ID NO: 36)
- [0333] Primer 103^R: 5' TTGAGGAAGATGTTTCGTGTA 3' (contains the SspI site of factor IX) (SEQ ID NO: 37)
- [0334] A PCR reaction was conducted with primer 101 and primer 103^R and plasmid DNA, cDNA clone of Factor IX (OriGene" RC219065) as a template; as a result of the PCR amplification, a ~ 1085 bp (per 10) product was formed and purified from the gel (the fragment containing the amino terminus of Factor IX sequence).
- [0335] Primer 98: 5' ATTACAGTTGTCGCAGGTGA 3' (SEQ ID NO: 38)
- [0336] Primer 99^R: 5' GCTGGAGCTAGTGAGCTTTGTTTTTTCCTT 3' (SEQ ID NO: 39)
- [0337] Primer 100: 5' GCTCACTAGCTCCAGCAGCAAGGCC 3' (SEQ ID NO: 40)

[0338] Primer 27^R: 5' TTTTCACTGCATTCTAGTTGTGG 3' (SEQ ID NO: 41)

[0339] Three PCR reactions were performed. The first reaction was conducted with primer 98 and primer 99^R and plasmid DNA, cDNA clone of Factor IX (OriGene[®], RC219065) as a template; as a result of the PCR amplification, a ~ 540 bp product was formed.

5 [0340] The second reaction was conducted with primer 100 and primer 27^R and plasmid DNA of 402-2-p72-3 (hGH-CTP-CTP) as a template; as a result of the PCR amplification, a ~ 258 bp product was formed.

[0341] The last reaction (pcr 3) was conducted with primers 98 and 27^R and a mixture of the products of the previous two reactions as a template; as a result of the PCR amplification,
10 a ~ 790 bp product was formed and ligated into TA cloning vector (Invitrogen, catalog K2000-01). SspI –EcoRI fragment was isolated (TA 3-3).

[0342] Another PCR reaction was conducted (pcr 12) with primer 101 and primer 27^R and a mixture of the products of pcr 10 and SspI-EcoRI fragment from pcr 3 as a template; as a result of the PCR amplification, a ~ 1700 bp product was formed (Factor IX-ctp-ctp) and
15 ligated into TA cloning vector (Invitrogen, catalog K2000-01) (lig 180).

[0343] A mistake was found in the Factor IX sequence so fragments were replaced in order to form an insert of Factor IX-ctp-ctp with the correct DNA sequence.

[0344] TA- pcr 3-3 was digested with SspI and XbaI and the large fragment was isolated (vector). TA 180-4 was digested with SspI and XbaI and the small fragment (insert) was
20 isolated and ligated to the isolated large fragment of TA-pcr-3-3 digested with SspI and XbaI. The new plasmid TA-183-2 was digested with Sal I and NotI, and the Factor IX-CTP-CTP insert was isolated (~1575 bp). This fragment was inserted into eukaryotic expression vector pCI-neo (digested with Sal I and Not I) to yield the 301-2-p200-11 clone.

[0345] **pCI-dhfr –Factor 9- ctpx2 (p223-4) construction:** Vector pCI-dhfr (p6-1) was
25 digested with SmaI and NotI. Factor IX-CTP-CTP (p200-11) was digested with ASiI F.I. and NotI. The two fragments were ligated.

[0346] **pCI-dhfr Factor 9-ctp x3 (p225-7) construction:** Vector pCI-dhfr OXM-CTP×3 (p216-4) was digested with XbaI and ApaI. Factor IX-CTP-CTP (223-4) was digested with XbaI and ApaI. The two fragments were ligated.

30 [0347] **pCI-dhfr Factor 9-ctp x3 T148A (p243-2) construction:** Plasmid p225-7 contained Threonine at position 148, since the more common version of FIX contains Alanine at this position, Thr was replaced to Ala using site directed mutagenesis method.

[0348] Primer 75: ctccaggtcaattacagct (SEQ ID NO: 42)

[0349] Primer 122r: ggaaaaactgcctcagcacgggtgagc (SEQ ID NO: 43)

[0350] Primer 123: gtgctgaggcagttttcctgatgtggactat (SEQ ID NO: 44)

[0351] Primer 124r: caacacagtgggcagcag (SEQ ID NO: 45)

[0352] Three PCR reactions were performed. The first reaction was conducted with primer 75 and primer 122r and plasmid DNA p225-7 as a template; as a result of the PCR amplification, a ~ 692 bp product was formed and purified from the gel. A second PCR reaction was conducted with primer 123 and primer 124r and plasmid DNA p225-7 as a template; as a result of the PCR amplification, a ~237 bp product was formed and purified from the gel. The third - overlap PCR reaction reaction was conducted with primers 75 and 124r, and a mixture of the products of the previous two reactions as a template; as a result of the PCR amplification, a ~ 910 bp product was formed. This overlap PCR product was digested with XbaI and NsiI and re ligated into p225-7 plasmid (digested with XbaI and NsiI) to yield Factor IX-ctpx3 T148A designated p243-2.

[0353] **FIX-4CTP (p259-4) construction:** 3.5CTP fragment was isolated from oxym-4CTP (p254-3) by restriction enzymes ApaI and XbaI. FIX+0.5CTP fragment was isolated from FIX-3CTP (p243-2) with restriction enzymes ApaI and XbaI. The two fragments were ligated.

[0354] **FIX-5CTP (p260-18) construction:** 4.5CTP fragment was isolated from oxym-5CTP (255-1) by restriction enzymes ApaI and XbaI. FIX+0.5CTP fragment was isolated from FIX-3CTP (p243-2) using enzymes ApaI and XbaI. The two fragments were ligated.

[0355] Dg44 cells were plated in 100mm tissue culture dishes and grown to 50-60% confluence. A total of 2 µg (microgram) of FIX cDNA was used for the transfection of one 100mm plate using the FuGene reagent (Roche) in protein-free medium (Invitrogene CD Dg44). The media was removed 48 hours after transfection and replaced with a protein-free medium (Invitrogene CD Dg44) without nucleosides and in the presence of 800 µg/ml of G418 (Neomycin). After 14 days, the transfected cell population was transferred into T25 tissue culture flasks, and selection continued for an additional 10-14 days until the cells began to grow as stable clones. High expressing clones were selected. Approximately 2×10^7 cells were used to inoculate 300 ml of growth medium in a 1700 cm² roller bottle (Corning, Corning NY) supplemented with 5 ng/ml of Vitamin K3 (menadione sodium bisulfate; Sigma). The production medium (harvest) was collected after a rapid decrease in cell viability to about 70%. The production medium was first clarified and then concentrated approximately 20-fold and dialyzed with PBS using flow filtration cassette (10KDa MWCO; Millipore Corp.).

[0356] **Determination of FIX antigen level:** FIX-CTP harvest antigen levels were determined using AssayMax Human FIX ELISA kit (AssayPro-EF1009-1). The calculated protein concentration is the average of three different dilutions in two independent runs (Figure 1A, Table 1).

Table 1: Calculated protein concentration

	FIX-CTP	FIX-CTP-CTP
FIX Ag level (µg/ml)	41.9	19.2
SD	8.76	3.67
%CV	20.92	19.15

[0357] **FIX SDS-PAGE - immune blot:** FIX-CTP harvests or purified rhFIX (American Diagnostics), 100 ng of protein, were loaded on 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE analysis was performed by Western immunoblot using anti-human FIX polyclonal antibody and anti-human gamma carboxylation monoclonal antibody (American Diagnostics). As previously reported, rhFIX migrated at 55KDa, while FIX fused to two CTPs migrated at 75KDa. Both variants of FIX-CTP proteins were shown to be gamma carboxylated, an essential post-translational modification for FIX activity and function (Figure 1B).

[0358] **Determination of FIX chromogenic activity:** A comparative assessment of the *in vitro* potency of FIX-CTP harvests versus rhFIX protein (American Diagnostics) was performed using the commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221802). In the presence of thrombin, phospholipids, calcium, excess amounts of FXIa activates sampled FIX into FIXa. FIXa forms an enzymatic complex with thrombin, activated FVIII:C (supplied in an excess amounts), phospholipids, and calcium and activates Factor X, present in the assay system, into FXa. The activity directly correlates with the amount of FIX, which is the limiting factor. The generated FXa is then measured by its specific activity on FXa chromogenic substrate (pNA). The amount of pNA generated is directly proportional to FIXa activity. rhFIX and FIX-CTP harvests were serially diluted, and the potency was assessed by comparing a dose-response curve of the FIX harvests to a reference preparation consisting of rhFIX or human plasma. The average EC₅₀ of FIX was 21 ng/ml, while the FIX-(CTP)₂ harvest calculated EC₅₀ was 382 ng/ml, and the FIX-CTP harvest calculated EC₅₀ was 1644 ng/ml. An approximately 15-fold decrease in the enzymatic activity of the FIX-(CTP)₂ harvest was observed (Figure 2).

[0359] **FIX Clotting activity (aPTT):** The activated partial thromboplastin time (aPTT) is a measure of the integrity of the intrinsic and common pathways of the coagulation cascade. The aPTT is the time, in seconds, for plasma to clot following the addition of an intrinsic pathway activator, phospholipid and calcium. The aPTT reagent is called a partial thromboplastin because tissue factor is not included with the phospholipid as it is with the protime (PT) reagent. The activator initiates the system and then the remaining steps of the intrinsic pathway take place in the presence of phospholipid. Reference aPTT range varies from laboratory to laboratory, but is usually in the range of 27-34 seconds.

[0360] The principal of the assay was to quantitate the ability of FIX-CTP harvests to restore the clotting activity of FIX-depleted human plasma by the addition of rhFIX. 300 µl of FIX-deficient human plasma was mixed with 100µl of rhFIX or FIX-CTP harvests and serially diluted. Following a 60 second incubation at 37°C, thromboplastin, CaCl₂, and phospholipids were added to the mixture, and clotting time in seconds was determined (performed by American Medical Laboratories). The potency was assessed by comparing a dose-response curve of the FIX harvests to a reference preparation consisting of rhFIX or human plasma. One unit of FIX activity corresponds to the FIX concentration that equals the activity of one ml normal human plasma. The presented aPTT results indicate that FIX-(CTP)₂ exhibit a 5.7-fold reduction in its specific coagulation activity compared to rhFIX (Table 2). Moreover, the aPTT results together with the chromogenic activity *in vitro* assay suggest that FIX-(CTP)₂ harvest has an improved enzymatic activity vs. FIX-CTP harvest (Table 2). An improved activity of FIX-CTP proteins can be obtained following optimization of the expression system (i.e. co-transfection with Furin and optimization of Vitamin K3 medium concentration), which was strengthened following super-transfection with Furin (data not shown).

Table 2: FIX clotting activity

rhFIX(A D) (µg/ml)	PTT(Sec)	FIX-CTP (µg/ml)	PTT (Sec)	FIX-CTP-CTP (µg/ml)	PTT (Sec)
5	31.3	9	45.2	4	47.5
1.25	35.7	2.25	53.3	1	55.9
0.3125	43	0.5625	64.1	0.25	67
0.07812	52.1	0.140625	76.3	0.0625	77.4

[0361] **Pharmacokinetic study:** rhFIX (American Diagnostic) and FIX-CTP harvests were administered in a single intravenous injection to Sprague-Dawley rats (six rats per substance) at a dose of 75 µg/kg body weight (Table 3).

Table 3: PK study plan of operation

Treated Groups	Test Article	No. of animals / group	Dose Route	Gender	Dose Level (µg/kg)	Dose Level (µg per animal)	Injected Vol. (µl)	Con. (µg/ml)	*Time-Points (hours post-dose)
1	rhFIX	6	IV	M	75	15	500	30	0 (Pre-dose), 0.083, 0.5, 1.5, 4, 8, 24, 48, 72.
2	rhFIX-CTP	6	IV	M	75	15	500	30	0 (Pre-dose), 0.083, 0.5, 1.5, 4, 8, 24, 48, 72.
3	rhFIX-CTP-CTP	6	IV	M	75	15	1000	15	0 (Pre-dose), 0.083, 0.5, 1.5, 4, 8, 24, 48, 72.

5

[0362] Blood samples were drawn retro-orbitally from 3 rats alternately at 0.083, 0.5, 1.5, 4, 8, 24, 48, and 72 hours post-dosing. Plasma was prepared immediately after sampling and stored at -20°C until analysis. FIX concentration was quantitated by FIX ELISA-specific assay (AssayPro). A pharmacokinetic profile was calculated for each protein and represents the mean of 3 animals at each time point (Figure 3). The terminal half-lives were calculated using PK solutions 2.0 software. Table 4 summarizes the observed FIX concentrations at the different sampling time points.

10

Table 4: Observed FIX concentrations

Time (Hr)	FIX-AD (ng/ml)	FIX-CTP (ng/ml)	FIX-CTP-CTP (ng/ml)
-----------	----------------	-----------------	---------------------

0.083	1506.7	1477.5	1914.8
0.5	1949.8	1150.1	1830.1
1.5	2189.4	1009.0	1264.3
4	733.90	709.33	1000.00
8	319.80	167.20	1234.67
24	BLQ	54.625	230
48	BLQ	BLQ	120.9

[0363] The PK profile and summary of the terminal half-lives are summarized in Table 5. FIX-CTP harvests exhibit an improved $T_{1/2\beta}$ values compared to rhFIX (2- and 5-fold increases, respectively). Since in FIX dosing collection, animal serum concentrations of FIX at 24hr were below limit of quantitation (BLQ), additional PK parameters were not calculated.

Table 5: Summary of PK parameters

Product	Terminal half-life- (hr)	Ratio (FIX-(CTP) _x /rhFIX)
rhFIX (American Diagnostics)	2.62	—
FIX-CTP	5.55	2.11
FIX-CTP (FIX-CTP-CTP)	12.9	4.92

[0364] In this study, a novel approach was described for prolonging FIX half-life while retaining the therapeutic potency. Adding a CTP peptide to an active protein has a harmful potential in interfering with the protein's activity. Therefore, the generation of an active recombinant FIX-CTP by adding a CTP sequence at the C-terminus of the FIX is unexpected.

Characterization of an immunoaffinity purified FIX-CTP-CTP

FIX-CTP-CTP purification

[0365] In order to evaluate a protein at high grade content with increased activity whose PK profile mimics and can be extrapolated to a clinical setting, FIX-CTP-CTP is a FIX modified with 2 CTP units in tandem in its carboxy-terminal. FIX-CTP-CTP was purified using matrix-bound monoclonal antibody against γ carboxyglutamyl (Gla) residues present in the N-terminal region of FIX (American Diagnostics Cat. # 3570MX). The monoclonal antibody was bound to Sepharose CL-4B. The FIX-CTP-CTP harvest at a concentration of 88 μ g/ml

was dialyzed against 20mM Tris, 150mM NaCl and 10mM EDTA at PH =7.4. The loading rate was 0.5 ml/min, elution was performed using 20mM Tris-HCl, 350 mM NaCl and 50 mM CaCl, and the unbound fraction was recycled five times. Finally, the elution fraction was dialyzed with PBS, pulled and concentrated.

- 5 [0366] **Determination of FIX antigen level:** FIX-CTP harvests, FIX-(CTP)₂ harvests, and FIX-(CTP)₂ purified protein levels were determined using the Human FIX ELISA kit (Affinity Biologicals; Cat. #FIX-AG RUO). The calculated protein concentration (µg/ml) is the average of two independent runs (Figure 4, Table 6).

Table 6: Calculated protein concentration

	FIX-CTP	FIX-CTP-CTP	FIX-CTP-CTP (purified)
FIX Ag level (µg/ml)	125.78	88.53	172.9
SD	17.28	21.31	2.63
%CV	13.74	24.08	1.52

10

[0367] Additionally, FIX-CTP-CTP was quantitated by Bradford assay. The calculated concentration was 202 µg/ml, which is similar to the concentration obtained by human FIX ELISA.

- 15 [0368] **SDS-PAGE blots:** FIX-CTP-CTP harvest, unbound fraction and purified protein, were loaded on a 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE Coomassie analysis was performed by staining the gel with Coomassie blue reagent (800ng of protein). A Western immunoblot was performed with 100 ng of protein, anti-human FIX polyclonal antibody (Ab), and anti-human gamma carboxylation monoclonal Ab (American Diagnostics Cat #499 and #3570). The
- 20 immunoaffinity purification procedure significantly enriched the FIX-CTP-CTP portion while reduced impurity (Figure 5).

- [0369] **N-terminal sequencing:** FIX-CTP-CTP purified protein was separated by 12% Tris-Glycine SDS-PAGE and subsequently electro-blotted to PVDF membrane. The band of interest was cut out and put on a purified Biobrene treated glass fiber filter. The N-terminal
- 25 sequence analysis was carried out by Edmann degradation using a pulsed liquid protein sequencer equipped with a 140 C HPLC micro-gradient system. N-terminal sequencing revealed that FIX-CTP-CTP is a mixture of incomplete and complete pro-peptide cleaved proteins. Inadequate pro-peptide cleavage was shown to reduce FIX coagulation activity. By co-transfection with Furin, the pro-peptide cleavage process can be an improved.

[0370] **Determination of FIX chromogenic activity:** A comparative assessment of the *in vitro* potency of FIX-CTP-CTP purified protein versus rhFIX (American Diagnostics) and a pool of human normal plasma was performed using the commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221802). In the presence of thrombin, phospholipids and calcium, excess amounts of FXIa activates FIX into FIXa. FIXa forms an enzymatic complex with thrombin (supplied in excess amounts), phospholipids and calcium activates Factor X, present in the assay system, into FXa. The activity directly correlates with the amount of FIX, which is the limiting factor. The generated FXa was measured by its specific activity on FXa chromogenic substrate (pNA). The amount of pNA generated was directly proportional to FIXa activity. rhFIX, human plasma and FIX-CTP-CTP were serially diluted, and potency was assessed by comparing a dose-response curve (Figure 6). The average EC₅₀ of rhFIX was 68.74 ng/ml while FIX-CTP-CTP calculated EC₅₀ was 505 ng/ml. An approximately 7-fold decrease in the enzymatic activity of FIX-CTP-CTP was observed vs. recombinant FIX and a 16.5-fold decrease versus normal human pulled plasma. This reduced activity could be explained by inadequate cleavage of N-terminal pro-peptide, which was identified by N-terminal analysis.

[0371] **FIX Clotting activity (aPTT):** The activated partial thromboplastin time (aPTT) is a measure of the integrity of the intrinsic and common pathways of the coagulation cascade. The aPTT is the time (measured in seconds) it takes plasma to clot following the addition of an intrinsic pathway activator, phospholipid and calcium.

[0372] The assay quantitated the ability of the FIX-CTP-CTP protein to restore the clotting activity of FIX depleted human plasma by the addition of rhFIX. 300 µl of FIX-deficient human plasma was mixed with 100 µl of rhFIX, FIX-CTP-CTP (FIX-CTP-CTP (the CTP are in tandem at the C-terminal)), or normal pool human plasma which was further diluted. Following a 60 second incubation at 37°C, Tissue Factor (TF), CaCl₂, and phospholipids were added to the mixture. Clotting time in seconds was determined. Potency was assessed by comparing a dose-response curve of FIX-CTP-CTP to a reference preparation of rhFIX or human plasma. One unit of FIX was defined as the amount of FIX which equals to the activity of 1 ml human normal plasma.

[0373] The aPTT results indicate that FIX-CTP-CTP coagulation activity is only 1.4 less than normal pool human plasma and similar to the rhFIX. The aPTT results together with the chromogenic activity *in vitro* assay suggest that FIX-CTP-CTP purification did not damage its activity.

[0374] **Pharmacokinetic activity of FIX-CTP-CTP:** Purified FIX-CTP-CTP, rhFIX (American Diagnostic) and harvests containing FIX-CTP-CTP and FIX-CTP were administered in a single intravenous injection to Sprague-Dawley rats (eight rats per substance) in a dose of 100µg/kg body weight (Table 7).

5

Table 7: PK study outline

Treated Groups	Test Article	No.of animals/ group/ time point	Dose Level (µg/kg)	Dose Level (µg per animal)	Injecte d Vol. (µl)	Con. (µg/ml)	Time-Points (hours post-dose)
A	rFIX	8	100	20	500	40	0 (Pre-dose) 0.083, 0.5, 1, 2, 4, 7, 10, 24, 48, 72.
B	rFIX-CTP (harvest)	8	100	20	500	40	0 (Pre-dose) 0.083, 0.5, 1, 2, 4, 7, 10, 24, 48, 72.
C	rFIX-CTP-CTP(harvest)	6	100	20	500	40	0 (Pre-dose) 0.083, 0.5, 1, 2, 4, 7, 10, 24, 48, 72.
D	rFIX-CTP-CTP (purified)	4	100	20	500	40	0.083, 0.5 1, 2, 4, 7, 10, 24, 4, 8, 72.

[0375] Blood samples were drawn retro-orbitally from 4 rats alternately at 0.083, 0.5, 2, 4, 7 10, 24, 48, and 72 hours post-dosing. Citrated plasma (0.32%) was prepared immediately after sampling and stored at -20°C until analysis. FIX concentration was quantitated using a human FIX ELISA kit (Affinity Biologicals). The pharmacokinetic profile was calculated for each protein as the mean of 4 animals at each time point (Figure 7). The terminal half-life was calculated using PK Solutions 2.0 Software. Table 8 summarizes the observed FIX concentrations at different sampling time points.

10

Table 8: Observed FIX concentrations

Time (hr)	FIX-CTP harvest ng/ml	FIX-(CTP) ₂ harvest ng/ml	rhFIX ng/ml	Purified FIX-CTP-CTP ng/ml
0.085	1038.97	1123.62	325.05	886.48
0.5	939.12	956.80	274.58	670.92

1	791.97	843.85	222.90	674.17
2	304.98	673.31	186.00	503.91
4	315.37	525.50	109.69	357.36
7	171.45	384.36	67.62	257.02
10	50.34	250.73	40.20	158.66
24	10.07	78.50	BLQ	52.13
48	BLQ	23.40	BLQ	18.07

[0376] A summary of the PK parameters are presented in Table 9.

Table 9: Summary of PK parameters

	T _{1/2} (hr)	AUC hr/ml	ng- MRT (hr)	Vd ml/Kg	CL ml/hr/Kg
FIX-CTP harvest	4.17	3622	4.5	155.1	27.6
FIX-(CTP)₂ harvest	10.44	9105.7	12	165.4	10.9
rhFIX	3.72	1416.8	5.1	373.8	70.183
Purified FIX-CTP-CTP	11.14	6314.2	12.3	254.5	15.83

[0377] The FIX-CTP-CTP harvest demonstrated an improved PK profile compared to FIX-CTP harvest. Furthermore, purified FIX-CTP-CTP exhibited a 3-fold increase in T_{1/2β} value and a 4.5-fold increase in AUC compared to rhFIX.

[0378] The reduced amount of secreted FIX fused to tandem CTP molecules versus fusion of a single CTP appears to be due to the addition of an extra CTP and not to reduced detection by ELISA, because the Bradford-purified FIX-CTP-CTP calculated concentration was similar to the ELISA-calculated concentration.

[0379] FIX-CTP-CTP clotting activity was similar to pooled human plasma; however, its *in vitro* chromogenic activity was significantly lower when compared to rhFIX or pooled human plasma. The chromogenic activity assay was reported as a very sensitive assay compared to the coagulation assay. The reason for reduced activity of FIX-CTP-CTP may vary. Addition of CTP may decrease the affinity of FIX to FXIa or reduce post-transcriptional modifications (e.g. 12-10 GLA residues and pro-peptide cleavage). N-terminal analysis revealed that the proteolytic cleavage of the FIX-CTP-CTP pro-peptide was not fully

completed prior to secretion. Since this post-transcriptional modification is crucial for the normal enzymatic activity of the protein, co-transfection with Furin-PACE plasmid is favorable and may improve FIX-CTP-CTP activity.

[0380] Finally, FIX-CTP-CTP comparative PK study in rats demonstrated that fusion of two tandem CTPs to the C-terminal of FIX generated a FIX with an extended half-life.

[0381] **FIX depleted mouse model:** In order to assess the *in vivo* activity, FIX knockout mice are obtained, and a breeding colony is established. 10µg of either commercial recombinant hFIX (BeneFIX®) or rFIX-(CTP)₂ (FIX-CTP-CTP) are injected into the tail vein of an anaesthetized FIX knockout mouse (22-28g). The amount of injected protein equals to the required concentration of FIX in normal plasma (5µg/ml). Blood samples are taken from the clipped tail into heparinized capillary tubes at specific time points. Plasma samples are assessed for FIX levels by ELISA and efficacy is measured by aPTT coagulation assay.

[0382] **Increasing FIX Propeptide cleavage efficacy:** CTP peptide cDNA was fused to the 3' end of human FIX cDNA. The corresponding rFIX and Furin expressing constructs were co-transfected into Dg44 cells; a human rFIX cDNA was also co-transfected with the Furin plasmid as a control. Secretion of high level of FIX leads to secretion of a mixture of pro-factor and a mature factor FIX, due to limited amount of the Furin protease in the cell. Co-transfection of a Furin expressing vector with a pro-factor expressing vector increases the recovery and result in the secretion of fully processed FIX in to the medium.

[0383] Following FIX-(CTP)₂ and Furin co-transfection, stable clones are generated and harvest is collected for pro-peptide cleavage evaluation. 100 ng of protein, are loaded on 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE analysis is performed by Western immunoblot using anti-human FIX polyclonal Ab (American Diagnostics) and anti-pro-peptide polyclonal antibody. As previously reported, rhFIX migrated at 55KDa, while FIX fused to two CTPs migrated at 75 kDa. Both variants of FIX proteins are shown to undergo a proper, full pro-peptide cleavage.

[0384] To determine whether proper pro-peptide cleavage improves FIX-(CTP)₂ enzymatic activity, a comparative assessment of chromogenic and coagulation activity of FIX-(CTP)₂ harvest co transfected with Furin is performed. A significant improvement in FIX-(CTP)₂ specific activity is observed, which is similar to rhFIX.

[0385] In conclusion, the results described herein suggest that FIX-CTP-CTP can be used efficiently for treating Hemophilia B patients. FIX fused to CTP constructs benefit from improved *in vivo* pharmacologic performance that overcomes the drawback in certain *in vitro*

measures. This proposed treatment is advantageous over previous treatments as the rate of infusions and the amount of required doses are reduced.

[0386] It is important to notice that when an albumin-fused molecule strategy was used to improve the FIX half-life, the recombinant FIX became inactive. The present novel approach lead to the design and purification of a novel recombinant FIX-fused protein that presents an improved long-lasting activity. Since mere size modifications did not improve the pharmacokinetics of injected FIX, the finding that CTP fused to FIX facilitates pharmacokinetic parameters was unexpected. The presence of highly glycosylated peptide-sialic acid residues stabilized the protein and protected it from interactions with vascular receptors without abrogating key determinants of FIX function.

[0387] FIX-CTP has a similar therapeutic efficacy to rFIX in hemophilia B patients and required less frequent dosing. A single injection of FIX-CTP is sufficient to control bleeding episodes and reduce the number of injections that are needed during surgical intervention in hemophilia B patients.

[0388] The CTP technology was utilized for the development of a long-acting FIX. Specifically, extending the half-life of recombinant rFIX molecule was performed by fusion of at least one human CTP to FIX. The recombinant FIX-CTP was expressed in mammalian cells and characterized *in vitro* and *in vivo*. It was demonstrated that the *in vitro* activity of rFIX-CTP was comparable to rFIX. Pharmacokinetics and efficacy studies in rats demonstrated improved properties of the rFIX-CTP. The results of this study demonstrate that it is feasible to develop a half-life extended rFIX molecule having similar haemostatic properties to the wild type enzyme.

EXAMPLE 2

Comparative Assessment of Purified FIX-CTP₃ vs. FIX-CTP₄ and FIX-CTP₅

[0389] **2.1 Study objective**

[0390] A comparative assessment of the pharmacokinetic parameters of FIX-CTP₄ and FIX-CTP₅ versus FIX-CTP₃ following a partial purification process.

[0391] **2.2 Production of FIX-CTP₄ and FIX-CTP₅ harvests**

[0392] FIX cDNA (OriGene RC219065) fused at the C-terminal to four or five tandem CTP sequences was expressed in Dg44 cells using Excellgene expression system in the presence of 10 ng/L of vitamin K3 (Sigma, Mennadion). The harvests were collected (300ml), filtered and frozen.

[0393] **2.3 Production of FIX-CTP₃ harvest**

[0394] FIX-CTP₃ was expressed in-house in CHO cells using pCI-DHFR vector, clone 196, BR-9 in the presence of 25 ng/L of vitamin K3 (Sigma). The harvests were collected and filtered.

[0395] All FIX-CTP samples (3, 4 and 5 CTP) were purified only by Jacalin column because of a lack of material.

[0396] 2.4 Determination of FIX antigen level

[0397] FIX antigen level was determined using Human FIX ELISA kit (Affinity Biologicals; Cat. # FIX-AG RUO). The calculated protein concentration is the average of four independent runs. FIX-CTP₃ concentration was slightly higher as compared to the two additional versions (Table 10).

Table 10: FIX antigen level

	3 CTP	4 CTP	5 CTP
	Final	Final	Final
	Jacalin40	Jacalin40	Jacalin40
Av.	1016.69	4644.11	1686.82
(ng/ml)			
SD	225.41	925.63	160.07
%CV	22.17	19.93	9.49

[0398] 2.5 FIX-CTP Coomassie stain and immune-blot

[0399] FIX-CTP₃, FIX-CTP₄, and FIX-CTP₅ harvests were loaded on 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE analysis was performed by Western immuno-blot using anti-CTP polyclonal Ab (Adar Biotech Production) or anti-Gla Ab (American Diagnostica).

[0400] As previously reported, FIX fused to three CTPs migrated at 80 kDa while FIX fused to four or five CTPs migrated at 85 KDa or 90 KDa, respectively. As expected, FIX-CTP₄ and FIX-CTP₅ harvests from Excellgene showed very low levels of gamma carboxylation compared to FIX-CTP₃ harvest, which was produced at Prolor (Figure 8).

[0401] After a purification process utilizing Jacalin column (immunoaffinity purification of glycosylated proteins), FIX-CTP₃, FIX-CTP₄, and FIX-CTP₅ were loaded on 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE was stained by Coomassie blue Dye for samples detection. All variants showed much cleaner band profiles (Figure 9), suggesting an improved purity.

[0402] **2.6 Determination of FIX chromogenic activity**

[0403] A comparative assessment of the *in vitro* potency of fully purified (HA column) FIX-CTP₃, FIX-CTP₄, and FIX-CTP₅ versus human pool normal plasma was performed using a commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221802).

- 5 All samples were serially diluted, and the potency was assessed by comparing a dose-response curve to a reference preparation of normal human plasma. The reduced chromogenic activity of FIX-CTP₄ and FIX-CTP₅ (Figure 10) as compared to plasma can be a consequence of improper post-transcriptional modifications of FIX proteins, e.g. inappropriate gamma carboxylation and pro-peptide cleavage or, alternatively, due to the addition of CTP cassettes.
- 10 The fluctuation in the FIX-CTP₄ and FIX-CTP₅ activity (Table 11) might be caused by inappropriate quantitation capabilities of the FIX ELISA due to CTP masking of the antigen site.

Table 11: Sample/plasma EC50 ratio

Sample	Sample/plasma EC 50 ratio
Plasma	1
3 CTP Final HA	2
4 CTP Final HA	5.35
5 CTP Final HA	2.73

15 [0404] **2.7 Pharmacokinetic study**

- [0405] Jacalin-purified FIX-CTP₃, FIX-CTP₄, and FIX-CTP₅ (Group A, B and C, respectively) were administered in a single intravenous injection to Sprague-Dawley rats (six rats per treatment group) at a dose of 250 µg/kg body weight. Blood samples were drawn retro-orbitally from 3 rats alternately at 0.083, 0.5, 2, 5, 8, 24, 48, 72 and 96 hours post-dosing
- 20 (Table 12). Citrated plasma (0.38%) was prepared immediately after sampling and stored at -20°C until analysis.

Table 12: PK study plan of operation

Treatm ent Group	Treatmen t	No. of animals/ group	Dose Route	Dose Level (µg per animal)	Injecte d Vol. (µl)	Conc. (µg/ml)	Time-Points (hr post-dose)
------------------------	---------------	-----------------------------	---------------	-------------------------------------	---------------------------	------------------	-------------------------------

A	FIX- CTP*3 Jacalin 40	6	IV	50	200	250	0.083, 0.5, 2, 5, 8, 24, 48, 72, 96
B	FIX- CTP*4 Jacalin 40	6	IV	50	200	250	0.083, 0.5, 2, 5, 8, 24, 48, 72, 96
C	FIX- CTP*5 Jacalin 40	6	IV	50	200	250	0.083, 0.5, 2, 5, 8, 24, 48, 72, 96

[0406] FIX concentration in plasma samples were quantified using human FIX ELISA kits (Affinity Biologicals). The pharmacokinetic profile was calculated and is the mean of 3 animals at each time point. Terminal half-lives were calculated using PK Solutions 2.0 Software. Table 13 below summarizes the calculated FIX concentrations at the different sampling time points.

Table 13: Calculated FIX concentrations

Time (hr)	Av. 3 CTP ng/ml	SD 3 CTP	Av. 4 CTP ng/ml	SD 4 CTP	Av. 5 CTP ng/ml	SD 5 CTP
0.083	1087.82	72.39	904.54	21.06	1097.23	82.24
0.5	774.18	86.31	736.82	66.93	998.79	70.43
2	562.23	3.70	627.09	32.47	747.85	14.02
5	357.44	8.63	431.23	29.41	576.49	27.36
8	239.20	7.82	327.46	30.26	394.96	36.48
24	77.08	4.26	107.38	5.18	142.42	16.13
48	27.73	2.02	39.83	1.85	53.66	3.33
72	12.55	1.48	21.53	1.55	23.54	3.32
96	6.66	1.23	10.63	0.13	18.54	3.39

[0407] The PK profile and a summary of the PK parameters are presented in Table 14 below and in Figure 11. A full PK analysis profile at all time points suggested that addition of 4 or 5 CTP cassettes to FIX did not increase its half-life as compared to FIX-CTP₃. The AUC

following FIX-CTP₅ administration increased by 1.4- to 1.6-fold versus FIX-CTP₃, which was not statistically significant.

Table 14: PK profile and a summary of the PK parameters

24- 96hr	3 CTP	4 CTP	5 CTP
Half-life (hr)	20.43	22.02	23.96
AUC (ng-hr/ml)	8218.38	10504.49	13329.41
Vd (ml/kg)	700.76	586.02	494.89
CL (ml/hr/kg)	23.77	18.45	14.32

[0408] Since 96 hr post-dosing samples were shown to have very low FIX concentrations, which were at the lower limit of quantification of the assay, the terminal half-life was recalculated providing a more precise and scientifically appropriate calculation (Table 15). According to this calculation, even smaller differences were obtained between the half-life of FIX-CTP₃, FIX-CTP₄, and FIX-CTP₅.

Table 15: Recalculated terminal half-life

8-72 hr	3 CTP	4 CTP	5 CTP
Half-life (hr)	15.38	16.63	16.04

[0409] 2.8 Conclusions:

[0410] In this study, the pharmacokinetic parameters and potential clotting activity of FIX-CTP₃, FIX-CTP₄, and FIX-CTP₅ were assessed. Fusion of 4 and 5 CTPs to FIX did not provide a superior or improved half-life extension, as compared to FIX-CTP₃, and reduced chromogenic activity was observed. Table 16 below summarizes the percent improvement of half-life for the different FIX-CTP fused variants (1 to 5 CTPs). Fusion of CTP to FIX improved its pharmacokinetic behavior, but, unpredictably, this improvement was limited. Surprisingly, following fusion of 3, 4 or 5 CTPs in tandem to FIX, a similar half-life value was calculated.

Table 16: Summary of the percent improvement of half-life

4CTP vs. 5CTP	0
	% increase
rhFIX vs. 1CTP	112
1CTP vs. 2CTP	141
2CTP vs. 3CTP	37
3CTP vs. 4CTP	6

[0411] These data suggest that fusion of 3 CTPs to FIX produces a maximal improvement in protein half-life, confirming that FIX-CTP₃ is the optimal variant in terms of half-life, structure and potential clotting activity for further clinical development.

EXAMPLE 3

FIX-CTP₃ TREATMENT OF FIX-/- HEMOPHILIC MOUSE MODEL

[0412] As described above, a study testing FIX-CTP, FIX-CTP₂ and FIX-CTP₃ harvest PK profile and coagulation activity vs. rhFIX was conducted. FIX-CTP₃ exhibited an improved PK profile while maintaining its coagulation activity vs. FIX-CTP₁ and FIX-CTP₂ harvests or rhFIX. To further evaluate this result, FIX-CTP₃ γ -Carboxyglutamate protein was purified. FIX-CTP₃ exhibits a 3-fold increase in half-life and 4.5-fold higher AUC compared to rhFIX in normal rats following a single IV administration. FIX-CTP₃ demonstrated a reduced *in vitro* chromogenic and clotting activity, most likely due to insufficient cleavage of N-terminal pro-peptide and in appropriate post-transcriptional modifications (PTMs), such as appropriate gamma carboxylation.

[0413] In the current study, the pharmacokinetic and pharmacodynamic properties of human recombinant FIX fused to three tandem CTPs were tested in FIX-deficient mice.

[0414] Study purpose:

[0415] To determine the pharmacokinetic and pharmacodynamic parameters of rFIX-(CTP)₃ vs. commercial rhFIX (BeneFIX®) in FIX-deficient mice following a single IV administration of FIX-(CTP)₃ at a similar specific activity and dose (similar specific activity to PD and similar FIX constant for PK).

[0416] Production of FIX-CTP₃ harvest:

[0417] FIX cDNA (OriGene RC219065-Thr 148) fused at the C-terminal to three tandem CTP sequences was expressed in Dg44 cells using Excellgene expressing system in the presence of 25 ng/ml of Vitamin K3 (Sigma, Mennadion). Five separate batches containing 5 liters of cell suspension was cultured (total of twenty-five liters) and harvested following viability decline to 60-70%. The harvest was filtered and frozen at -70°C.

[0418] Determination of harvest FIX antigen level:

[0419] Harvest FIX antigen level was determined using a human FIX ELISA kit (Affinity Biologicals; Cat. # FIX-AG RUO). The antigen level was calculated per each batch. The FIX concentration was maintained through the different batches (Table 17).

Table 17: FIX antigen level

FIX antigen level			
Batch	#1	Bat #2	Bat #3
Av (µg/ml)	28.81	32.74	42.9
STD	2.5	2.69	4.0
%CV	8.84	8.38.2	9.4

[0420] **FIX-CTP₃ purification process:**

[0421] Following a short purification study, a purification process using the following 3 columns was performed: DEAE Sepharose, Heparin Sepharose and HA Bio Rad Ceramic Hydroxyapatite type 1 (40 µm), FIX-CTP₃. γ-carboxylated enriched protein was purified. In brief: Five liters of clarified harvest was thawed at 4°C over a 4 day period. For each purification batch, the clarified harvest (2 liters) was concentrated 4-fold and dialyzed against 20 mM Tris-HCl pH 8.2 using a disposable hollow fiber cartridge with a nominal molecular weight cutoff size of 10 KDa. This process (UFDF1) was performed twice, and one liter of UFDF1 was loaded on DEAE Sepharose column, and Factor IX was eluted with 20 mM Tris-HCl, 200 mM NaCl, 10 mM CaCl₂ pH 8.2. The product was diluted 1:1 with 20 mM Tris-HCl, 10 mM CaCl₂ pH 7.5, and the pH was adjusted to 7.5 before loading on Heparin Sepharose column. The elution was performed with 20 mM Tris-HCl, 300 mM NaCl, and 10 mM CaCl₂ pH 7.5. The eluted product was concentrated and dialyzed against 10 mM phosphate pH 6.8 using a Pellicon XL cassette 10 KDa cutoff membrane (UFDF2). The product was loaded on an HA column, and the activated fraction of Factor IX was eluted with 150 mM phosphate pH 6.8. The purification product was concentrated to a target concentration of 2 mg/ml and dialyzed against TBS pH 7.45, divided in aliquots and stored at -70°C.

[0422] The purification process was repeated five times, on a weekly basis in order to purify the total volume (25 liters). The purification processes were named HA# 6-10. Each purification product was separately evaluated (App # 1-5). At the end of the purification process, the different batches were pooled and further concentrated to a target concentration of 4 mg/ml.

[0423] **FIX-CTP₃ analytical properties:**

[0424] Determination of FIX antigen level

[0425] FIX-CTP₃ γ-carboxylated enriched protein antigen level was determined using a human FIX ELISA kit (Affinity Biologicals; Cat. # FIX-AG RUO). The calculated protein concentration is the average of two independent runs (Table 18).

Table 18: FIX-CTP₃ antigen level

FIX-CTP ₃ HA purified pool ELISA #1				FIX-CTP ₃ HA purified pool- ELISA #2				Final Av.
Dil.	1	2	Av.	Dil.	1	2	Av.	
130000	3412240	3781830	3597035	130000	3692260	3568240	3630250	3613643
260000	3915600	4158440	4037020	260000	3706820	3595540	3651180	3844100
520000	4158544	4334096	4246320	520000	3831464	3530748	3681106	3963713
1040000	4096352	4004104	4050228	1040000	3863392	3684304	3773848	3912038
Av. (ng/ml)	3895684	4069618	3982651	Av. (ng/ml)	3773484	3594708	3684096	3833373
STD	338367.5	234486.7	274313.5	STD	86576.66	65369.65	63369.86	154459.6
%CV	8.685703	5.761884	6.887712	%CV	2.294343	1.818497	1.720092	4.029338
Av. (mg/ml)	3.895684	4.069618	3.982651	Av. (mg/ml)	3.773484	3.594708	3.684096	3.833373

FIX-CTP ₃ HA purified pool ELISA #1				FIX-CTP ₃ HA purified pool- ELISA #2				Final Av.
Dil.	1	2	Av.	Dil.	1	2	Av.	
130000	3412240	3781830	3597035	130000	3692260	3568240	3630250	3613643
260000	3915600	4158440	4037020	260000	3706820	3595540	3651180	3844100
520000	4158544	4334096	4246320	520000	3831464	3530748	3681106	3963713
1040000	4096352	4004104	4050228	1040000	3863392	3684304	3773848	3912038
Av. (ng/ml)	3895684	4069618	3982651	Av. (ng/ml)	3773484	3594708	3684096	3833373
STD	338367.5	234486.7	274313.5	STD	86576.66	65369.65	63369.86	154459.6
%CV	8.685703	5.761884	6.887712	%CV	2.294343	1.818497	1.720092	4.029338
Av. (mg/ml)	3.895684	4.069618	3.982651	Av. (mg/ml)	3.773484	3.594708	3.684096	3.833373

5 [0426] **SDS-PAGE blots:**

[0427] FIX-CTP₃ γ -carboxylated enriched protein, rhFIX and rFIXa (activated FIX) were loaded on 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE Coomassie analysis was performed by staining the gel with Coomassie blue reagent (800 ng of protein) (Figure 12). A Western immunoblot was performed using 100 ng of protein with anti-human FIX polyclonal Ab (Figure 12B), anti-human gamma carboxylation monoclonal antibody (American Diagnostics Cat #499, 3570) (Figure 12C), anti-FIX pro-peptide polyclonal Ab (Figure 12D), and anti-CTP polyclonal Ab (Figure 12E). As previously reported, FIX-CTP₃ migrated at 75KDa.

[0428] The purification procedure significantly enriched FIX-CTP₃ portion while reducing impurities. The purification process yield was very low ranging around 2-3% (data not shown) due to the requirement to collect only the γ -carboxylated FIX-CTP₃ fractions, as demonstrated in the anti-Gla immunoblot (Figure 12B). Based on the Coomassie and FIX immunoblot, the FIX-CTP₃ portion is only around 60-70%, and additional lower molecular weight bands, presumably with lower glycosylation forms, were also detected.

20

[0429] **FIX-CTP₃ clotting activity:**

[0430] FIX-CTP₃ chromogenic activity:

[0431] A comparative assessment of the *in vitro* potency of FIX-CTP₃ harvest and FIX-CTP₃ γ -carboxylated enriched protein, versus human pool normal plasma was performed using a commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221802). FIX-CTP₃ harvest and protein were serially diluted, and the potency was assessed by comparing a dose-response curve to a reference preparation consisting of normal human plasma. As previously demonstrated, FIX-CTP₃ harvest was 50 times less active than human pool plasma (Table 19, Figure 13). Following FIX-CTP₃ purification, the chromogenic activity was significantly improved and was only 4.72 times less active than human pool plasma (Table 19, Figure 13). Harvest reduced chromogenic activity can be a consequence of improper post-transcriptional modifications of FIX protein variants, e.g. inappropriate gamma carboxylation and pro-peptide cleavage. Following purification and enrichment of the FIX-CTP₃ γ -carboxylated fraction, the activity was improved, demonstrating the important contribution of γ -carboxylation to FIX activity.

15 **Table 19: FIX-CTP₃ chromogenic activity**

Sample	EC ₅₀ (ng/ml)	Sample /plasma EC ₅₀ ratio
FIX-CTP₃ Harvest	741.3	54.4
Pur. FIX-CTP₃	64.6	4.72
Plasma	13.63	1

20 [0432] One stage clotting assay (aPTT):

[0433] The activated partial thromboplastin time (aPTT) is a measure of the integrity of the intrinsic and common pathways of the coagulation cascade. The aPTT is the time, in seconds, for plasma to clot following the addition of an intrinsic pathway activator, phospholipid and calcium. The principal of the assay was to quantitate the ability of FIX-CTP₃ to restore the clotting activity of FIX-depleted human plasma by the addition of rhFIX. 200 μ l of FIX-deficient human plasma was mixed with 25 μ g/ml of FIX-CTP₃ and further diluted in TBS. Following a 60 second incubation at 37°C, 50 μ l of PTT activator (Actin FS) and 50 μ l of calcium 25 mM were added to the mixture, and the clotting time in seconds was determined

using a Sysmex® CA 1500 Coagulator (performed by Sheba hospital, National Coagulation Center using validated aPTT assay). The potency was assessed by comparison of FIX-CTP₃ to the dose-response curve of a reference preparation of normal human pool plasma. The results are expressed in percent of activity interpolated from the standard curve covering FIX levels of <1-110%. FIX-CTP₃ exhibited a 15-20-fold reduction in its coagulation activity versus normal human pool plasma since the activity at 5 µg/ml, which is the normal value of FIX in the body, was shown to be 6.5% (Table 20).

Table 20: FIX-CTP₃ clotting activity

FIX-CTP ₃	FIX Concentration by provider (mg/ml)	Concentration in tested sample (µg/ml)	FIX % of activity (normalized to human normal pool plasma)
	3.83	25	34.7
		5	6.5

[0434] FIX-CTP₃ also exhibited increased clotting time compared to BeneFIX® (Table 21 and Figure 14).

Table 21: Comparative clotting time (aPTT)

Clotting time		
	FIX-CTP ₃	BeneFIX®
38ug/ml	77.6	
19ug/ml	83.4	
7.6ug/ml	93.2	50.6
3.8ug/ml	104.8	57.6
1.9ug/ml	112.2	63.7
0.95ug/ml	122.6	71.5
0.475ug/ml		83.7
0.238ug/ml		94.3

[0435] An additional clotting assay was performed independently in FIX-deficient mice by Dr. Paul Monahan at University of North Carolina prior to the initiation of the PK-PD study. The aPTT results suggested that FIX-CTP₃ coagulation activity is 40 times less than normal

pooled human plasma as demonstrated by the longer period (as measured in seconds) and higher concentration that are required for proper clotting activity (Table 22).

Table 22: Comparative clotting activity

FIX activity (Units)		
	FIX-CTP ₃	BeneFIX®
38ug/ml	13.9	
19ug/ml	8.8	
7.6ug/ml	4	116.8
3.8ug/ml	1.6	67.4
1.9ug/ml	0.9	41.7
0.95ug/ml	0.4	22.4
0.475ug/ml		8.5
0.238ug/ml		3.7

[0436] The specific activity (u/ml), which was based on FIX antigen level as calculated by ELISA for FIX-CTP₃ and BeneFIX®, was 4.46 and 198.9 respectively.

[0437] The inconsistency in the calculated FIX-CTP₃ activity as demonstrated in the chromogenic vs. aPTT assays can be explained by the superior sensitivity of the aPTT assay and *in vivo* relevance. In the chromogenic activity assay, an excess amount of reagents and enzymes are present which can activate less potent FIX versions. The difference in the FIX-CTP specific activity values can be explained by the use of different reagents and automated machines. The activity value as calculated at University of North Carolina was used for the PK-PD study design.

FIXa Protein detection:

[0438] In order to confirm that following the purification process, FIX activation (FIXa) did not occur, a FIXa detection assay was performed using FIXa Biophen Chromogenic Assay (Cat. # Ref. 221812). The assay measures the amount of FIXa present in a specific sample using the chromogenic activity cascade, as previously described. FIX-CTP₃ and rhFIX were diluted and FIXa levels were evaluated. FIX-CTP₃ wasn't activated through purification or storage (Table 23).

Table 23: FIXa detection

Sample	FIX-CTP ₃	rhFIX
Initial	1000	5.7

Con.(mg/ml)		
rFIXa	BLQ	0.00487
(mg/ml)		
%FIXa in sample	BLQ	0.085

[0439] **FIX-CTP₃ PK-PD study:** FIX-CTP₃ and rhFIX (BeneFIX®) were administered in a single intravenous injection to C57BI FIX-deficient mice in a dose of 625 µg/kg body weight containing 100 IU FIX/kg body weight. Blood samples were drawn retro-orbitally from 3 mice alternately at 0.25, 4, 24, 48, 72, and 96 hours post-dosing. Citrated plasma (0.32%) was prepared immediately after sampling and stored at -20°C until analysis. hFIX antigen level was evaluated, and a detailed PK analysis was performed. In order to evaluate the ability of FIX-CTP₃ to elongate the clotting activity of FIX-deficient animals compared to BeneFIX®, FIX activity in citrated plasma samples, collected from the FIX-/- treated mice, was calculated using an automated FIX activity assay (Table 24).

Table 24: Study outline

	Product	Administ ration	Dose	# mic e	Collection Points (hr post-dosing)	Required amount
**Coh ort 1	FIX-CTP ₃	Single dose: IV	100IU/Kg 2.5IU/mous e (553µg/mou se)	12 mic e,	0.25, 1, 4, 8, 16, 24, 48	6636µg
Cohort 2	FIX-CTP ₃	Single dose: IV	**472µg/K g 12.57µg/m ouse	18 mice	*0.25, 1*, 4*, 8 *, 16*, 24*, 48*, 72*, 96*	200µg 12.57µg/mou se
**Coh ort 3	BeneFIX ®	Single dose: IV	100IU/Kg 2.5IU/mou se	18 mice,	0.25, 1, 4, 8, 16, 24, 48, *72, *96	226.3 µg 12.57µg/mou se

* PK collection points only

** Tail vein bleeding at T=48 post-dosing; cohorts 1 & 3

[0440] FIX-CTP₃ Pharmacokinetic profile in FIX^{-/-} mice

[0441] FIX concentration was quantitated using human FIX ELISA kits (Affinity Biologicals; Cat. # FIX-AG RUO). The pharmacokinetic profile was calculated for each protein and is the mean of three animals at each time point. Table 25 below and Figure 15 summarize the calculated FIX concentrations at the different sampling time points for Cohorts 1 & 3. The PK profile and a summary of the PK parameters are presented below (Tables 26 & 27). A PK analysis was also performed for Cohort #2 in order to verify exposure (data not shown).

Table 25: FIX concentrations

Time point(hr)	FIX-CTP ₃ ng/ml	BeneFIX® ng/ml
0.25	3645.397	2823.023
1	2411.09	2416.248
4	1703.205	1506.228
8	1139.736	864.764
16	415.32	347.465
24	238.37	158.7973
36	141.0105	94.40067
48	95.461	42.28833
72	76.90953	11.87567
96	24.955	BLQ

[0442] A two-compartmental module was used (WinLin software) to determine AUC_{0-inf}, T_{terminal} and clearance (CL). The PK parameters are described below in Table 26.

Table 26: PK properties

FIX Version	T _{1/2α} (1/hr)	T _{1/2β} (1/hr)	AUC ng/ml*hr	CL ml/Kg/hr	MRT (hr)	V _{ss} (ml/Kg)
BeneFIX®	3.4	12.7	22428	29	11.5	320.8
FIX-CTP ₃	4	28.7	31770	19	22	425.2

[0443] The addition of the three CTP “cassettes” to rhFIX elongated FIX half-life *in vivo* by at least 2.5-fold. AUC following *in vivo* FIX-CTP₃ administration increased 2-fold versus rhFIX. FIX-CTP₃-injected mice demonstrated an improved PK profile compared to BeneFIX®-injected mice.

[0444] FIX-CTP₃ Pharmacodynamic profile in FIX-deficient mice:

[0445] In parallel to PK sampling, FIX-deficient animals administered with either BeneFIX® or FIX-CTP₃, citrated plasma samples, were evaluated for their clotting activity by aPTT assay, which was translated to % activity. The % activity at each collection point was

5 calculated as the current clotting time/clotting time of normal pool mice plasma*100. Table 27 summarizes the activity values following administration of either BeneFIX® or FIX-CTP₃.

[0446] Following FIX-CTP₃ administration, significant clotting activity was detected one hour after administration reaching 96% activity at four hours post-dosing, while BeneFIX® highest activity value was 40% (Table 27, Figure 16). FIX-CTP₃ clotting activity was

10 maintained for a longer period of time, demonstrating elongated activity. Clotting activity for the BeneFIX®-treated mice was undetectable at time points later than 36 hours, while FIX-CTP₃-treated mice continued to retain measurable activity at 72 hours post-dosing (Table 27, Figure 16). Analysis of the % clotting pharmacokinetic profile suggest that FIX-CTP₃ clotting activity is maintained for a significantly longer period and its half life is almost 2-fold higher

15 than Benefix® (Table 28).

Table 27: FIX % of activity

Hr post-dosing	BeneFIX® % of activity	FIX-CTP ₃ % of activity
0.25	39.9	1.0
1	33.4	15.5
4	24.9	93.6
8	18.8	65.2
16	10.3	39.9
24	1.7	11.9
36	1.4	11.0
48	<1	4.6
72	<1	1.4

Table 28: Clotting Activity

FIX Version	T _{1/2α} (1/hr)	T _{1/2 β} (1/hr)
BeneFIX®	5.7	-----
FIX-CTP ₃	7.3	16

20 [0447] 9.3 FIX-deficient mice bleeding challenge

[0448] FIX-deficient mice were administered a single intravenous injection of 100 IU/kg of BeneFIX® or rFIX-CTP₃. The tail vein was slightly clipped 48 hours post-dosing, and tail

vein bleeding time (TVBT) and bleeding intensity (hemoglobin OD) were evaluated. A second bleeding challenge was performed 15 minutes after reaching homeostasis, and the same parameters were measured. Following the first bleeding challenge, FIX-CTP₃-administered animals' bleeding was significantly less intense than BeneFIX® bleeding as demonstrated by the Hemoglobin OD values (Figure 17).

[0449] Since it was previously reported that during the first bleeding challenge in hemophilic mice, the bleeding time does not necessarily correlate with treatment efficacy, it is recommended to evaluate the homeostasis following additional bleeding. Once the first bleeding was spontaneously or manually stopped, a second bleeding challenge was performed 15 minutes following the first one, and the time and bleeding intensity were re-measured. During the second bleeding episode FIX-CTP₃-administered animals had reduced bleeding time and intensity, demonstrating that FIX-CTP₃ was potent at a later time points (Figure 18).

[0450] Finally, the animals were further observed for the 12 hours following the second bleeding challenge, and all recurring bleeding events were documented. FIX-CTP₃-administered animals were able to maintain blood homeostasis for the next 12 hours with no re-occurring bleeding events. In contrast, 50% of BeneFIX®-treated mice had spontaneous bleeding episodes from the tail (Table 29).

Table 29: Outcome 12 hours after tail transection

Mouse group	Delayed rebleeding	Death or Distress Requiring Euthanasia
FIX-CTP₃ (100IU/kg)	0/5 (0%)	0/5
BeneFIX® (100IU/kg)	3/6 (50%)	0/6
FIX-/- (untreated)	5/6 (100%)	1/6

[0451] Recombinant FIX-CTP₃, a fusion protein comprised of a single molecule of FIX fused to three CTP "cassettes" in tandem was developed to address the short half-life of currently available FIX products used to treat patients with hemophilia B. We have demonstrated that the elimination half-life of rFIX-CTP₃ was consistently 2.5- to 4-fold longer than rFIX in rats (as previously reported) and in FIX-deficient mice.

[0452] Without being bound by theory, the fusion protein reduces clearance of FIX and protects FIX from protease activity, degradation by masking and reduces the affinity of FIX for hepatic receptors. Taken together these characteristics of the CTP domain extend the half-life of FIX.

5 [0453] In addition to pharmacokinetic analysis of rFIX-CTP₃, we examined the pharmacodynamic properties of FIX-CTP₃ in FIX-deficient mice. rFIX-CTP₃ and rFIX, were administered at comparable doses (in units) to compensate for the clotting deficiency levels in FIX-deficient mice. However, the effect of rFIX-CTP₃ in FIX-deficient mice was significantly prolonged to at least 76 hr after dosing, reaching a higher activity peak. FIX-
10 CTP₃ clotting activity began after a 1-hour delay compared to BeneFIX®. FIX activation may be required since the addition of three tandem CTPs might theoretically mask the activation site and delay cascade onset. Following FIX-CTP₃ administration, a 100% peak activity was observed, while BeneFIX® activity was only 40%. The superior initial activity is a very important parameter and demonstrates that addition of 3 CTPs has the potential to improve
15 recovery.

[0454] Prophylactic FIX replacement therapy for patients with hemophilia B aims to maintain plasma levels of 1–2% normal clotting activity. The tail vein bleeding assay is a sensitive *in vivo* test that measures the ability to maintain bleeding homeostasis at low activity values mimicking human bleeding homeostasis model. In response to tail vein bleeding
20 challenge 48 hours post-dosing, rFIX-CTP₃-administered animals maintained blood homeostasis with shorter and less severe bleeding episodes, demonstrating sustained clotting activity.

[0455] FIX is a complex protein that contains a number of functional domains which undergo extensive post-translational modifications. One of the essential post-translational
25 modifications for FIX activity is gamma-carboxylation of the first 12 glutamic acids in the Gla domain by vitamin K-dependent γ -glutamyl carboxylase. This modification facilitates the binding of FIX to phospholipid membranes and, thus, is critical to its function. FIX that is not gamma-carboxylated is not functional, and hence gamma-carboxylation is a rate-limiting step.

[0456] This PK-PD study was conducted using transiently transfected cells. An extensive
30 analytical evaluation of post-translational modifications is performed on the stable FIX-CTP₃ protein produced and secreted from stable optimized clone.

[0457] Based on the presented data, FIX-CTP₃ coagulation factor has the potential to reduce the frequency of injections in patients receiving routine prophylactic doses of FIX replacement therapy. It is anticipated that rFIX-CTP₃ can confer prolonged protection from

bleeding following each dose of factor, decrease the overall units of factor needed to treat bleeding episodes, and/or maintain adequate hemostasis during surgical procedures with fewer injections.

EXAMPLE 4

5 *Generation and utilization of Coagulation Factor FVII*

[0458] A long-acting version of activated Factor VII (FVIIa) coagulation factor will be useful for the treatment of patients with hemophilia A and B. FVIIa-CTP₃ recombinant protein has the clinical potential to improve the treatment of hemophilia patients by reducing the frequency of infusions and even by reducing the drug load, enabling a prophylactic
10 treatment approach which can significantly improves a patient's quality of life, avoid spontaneous bleeding episodes and accumulated damage to the joint and other organs.

[0459] The generation of a recombinant FVIIa-CTP molecule with an extended half-life based on fusion of FVII to a human CTP is described herein. The recombinant FVIIa-CTP was expressed in mammalian cells and characterized *in vitro* and *in vivo*. It was demonstrated
15 that rFVII-CTP activity was comparable to rFVII. Pharmacokinetic and efficacy studies in rats demonstrated improved properties of rFVII-CTP. The results of this study demonstrated that it is feasible to develop a half-life extended rFVIIa molecule with very similar haemostatic properties to the wild-type enzyme.

[0460] **Cloning and expression of recombinant FVII molecule:** Several Factor VII clones
20 were constructed in our eukaryotic expression vector (pCI-dhfr) (Figure 19). Human MGC verified FL cDNA clone (IRCM) containing the sequence of *homo sapiens* coagulation Factor VII was ordered from "Open Biosystems" (OB-MHS4426). The following primers were synthesized by Sigma-Genosys in the following sequence: Primer 67: 5'CTCGAGGACATGGTCTCCCAGGCC3' (contains the 5' end of Factor VII DNA and the
25 restriction site of XhoI) (SEQ ID NO: 5); Primer 68^R: 5' TCTAGAATAGGTATTTTCCACATG3' (contains the restriction site of XbaI) (SEQ ID NO: 6); Primer 69: 5' TCTAGAAAAAAGAAATGCCAGC3' (contains the restriction site of XbaI) (SEQ ID NO: 7); and Primer 70^R: 5'GCGGCCGCATCCTCAGGGAAATGGGGCTCGCA3' (contains the 3' end of Factor VII
30 DNA and the restriction site of NotI) (SEQ ID NO: 8).

[0461] Cloning was performed in two sets of PCR reactions. The first reaction was conducted with primer 67 and primer 68^R using a cDNA plasmid with the Factor VII sequence (OB-MHS4426) as a template; as a result of the PCR amplification, a ~534 bp product was formed, isolated and ligated into a TA cloning vector (Invitrogen, Catalog No:

K2000-01). A XhoI –XbaI fragment containing the amino terminus of the Factor VII sequence was isolated. The second reaction was conducted with primer 69 and primer 70^R and again, a cDNA plasmid with the Factor VII sequence (OB-MHS4426) was used as a template. As a result of the PCR amplification, a ~813 bp product was formed and ligated into TA
5 cloning vector (Invitrogen, Catalog No: K2000-01). A XbaI-NotI fragment containing the carboxy terminus of Factor VII sequence was isolated. The two fragments were inserted into our eukaryotic expression vector pCI-dhfr (triple ligation) to yield the 501-0-p136-1 clone.

[0462] Plasmid 501-p136-1 (Factor VII in pCI-dhfr vector) was digested with restriction enzymes XhoI and KpnI. A fragment of ~1186 bp was isolated. A partial Factor VII clone
10 (1180 bp-1322 bp) followed by a CTP sequence, termination sequence and NotI sequence that was synthesized by GeneArt (0721543) was digested with restriction enzymes KpnI and NotI. A fragment of ~253 bp was isolated. The two fragments were inserted into our eukaryotic expression vector pCI-dhfr (triple ligation) to yield the 501-1-p137-2 clone. pCI-dhfr-701-2-p24-2 was digested with restriction enzymes XhoI and ApaI, and the large fragment (vector)
15 was isolated.

[0463] pCI-dhfr-501-2-p137-2 (Factor VII-ctp x1) was digested with restriction enzymes XhoI and ApaI, and a ~1200 bp insert was isolated. The vector and insert were ligated to yield 501-2-p139-2. Dg44 cells were plated in 100 mm tissue culture dishes and grown to confluence of 50-60%. A total of 2 µg of DNA was used for transfection of one 100 mm plate
20 using the FuGene reagent (Roche) in protein-free medium (Invitrogen CD Dg44). The medium was removed 48 hours post-transfection and replaced with a protein-free medium (Invitrogen CD Dg44) without nucleosides. After 14 days, the transfected cell population was transferred into T25 tissue culture flasks, and the selection was continued for 10-14 days until the cells began to grow well as a stable clone. High-expressing clones were selected and
25 approximately 2×10^7 cells were used to inoculate 300 ml of growth medium in a 1700cm² roller bottle (Corning, Corning NY) supplemented with 5 ng/ml of Vitamin K3 (menadione sodium bisulfate; Sigma). The production medium (harvest) was collected after a rapid decrease in the cell viability to around 70%. The production medium was first clarified and then concentrated approximately 20-fold and dialyzed to PBS using flow filtration cassette
30 (10KDaMWCO; Millipore Corp, Billerica, MA).

[0464] **Determination of FVII antigen level**

[0465] The cDNA coding the CTP peptide was fused to the 3' end of the cDNA coding human FVII. The corresponding rFVII construct was transfected into Dg44 cells. As a control, a human rFVII cDNA was utilized. The production medium (harvest) was collected,

concentrated and the secreted recombinant FVII was further evaluated. rFVII, rFVII-CTP and rFVII-CTP-CTP antigen levels were determined by AssayMax Human FVII ELISA kit (AssayPro) (Figure 20A). There was no significant difference in the secretion level of rFVII-CTP and rFVII-(CTP)₂ compared to native rFVII.

5 [0466] **SDS-PAGE blots**

[0467] SDS-PAGE analysis was done by loading 50 ng of either harvest, purified or activated rFVII protein. Samples were loaded on 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE analysis was done by performing a Western immunoblot using an anti-human FVII monoclonal antibody (Ab) (R&D systems) or
10 anti-CTP polyclonal antibody generated in Rabbit.

[0468] The level of rFVII antigen correlated with the detected protein level in a SDS-PAGE immunoblotted with anti-FVII Ab. rFVII-CTP migrated as a single band, while the corresponding molecular weight of the FVII control was approximately 52 KDa (data not shown). Both proteins reacted with antibodies specific for FVII on immunoblots. The rFVII-CTP also reacted with antibodies specific for CTP. rFVII was secreted in its zymogene form
15 with no trace of activated protein.

[0469] **FVII Chromogenic activity:**

[0470] rFVII, rFVII-CTP and rFVII-(CTP)₂ harvest activities were determined using a commercially available chromogenic test kit (AssaySense Human FVII Chromogenic Activity Assay Kit (AssayPro). For functional characterization of the rFVII-CTP and its ability to be
20 further activated (FVIIa), concentrated rFVII-CTP (harvests) were placed in a commercially available chromogenic test kit that measure the ability of TF/FVIIa to activate Factor X to Factor Xa that in the presence of FXa specific substrate releases a quantitated signal (AssayPro). The addition of the CTP peptide at the C-terminal of the rFVII protein did not
25 impair the FVII serine protease activity (Figure 20B, 20C).

[0471] **FVII clotting activity:**

[0472] Prothrombin time (PT) measures the extrinsic pathway of coagulation. The PT is the time (measured in seconds) it takes plasma to clot following the addition of an extrinsic pathway activator, phospholipid and calcium. It is used to determine the clotting tendency of
30 blood, specifically in the measure of warfarin dosage, liver damage, and vitamin K status. The reference range for prothrombin time is usually around 12–15 seconds. Specifically, the assay quantitated the ability of FVII-CTP and FVII-(CTP)₂ harvest to restore the clotting activity of FVII-depleted human plasma by the addition of rhFVII. 300 µl of FVII-deficient human plasma was mixed with 100 µl of FVII, FVII-CTP and FVII-(CTP)₂ harvests at specific

concentrations, or normal pool human plasma and were further diluted. Following a 60 second incubation at 37°C, Tissue Factor (TF), CaCl₂, and phospholipids were added to the mixture. The clotting time in seconds was determined. Potency was assessed by comparing a dose-response curve of FVII-CTP and FVII-(CTP)₂ harvests to a reference preparation consisting of rhFVII or human pool plasma. One unit of active FVII was defined as the amount of FVII which equals to the activity of one ml human normal plasma. The PT Clotting activity of rFVII and rFVII-CTP was measured on a coagulometer (Instrumentation Laboratory).

[0473] As previously shown, the addition of a CTP peptide at the C-terminal of the rFVII protein did not damage its serine protease activity and lead to the initiation and activation of a native Factor X and Factor IX in human plasma. Following the insertion of an additional CTP at the C terminal, there was a three-fold reduction in the serine protease activity (data not shown).

[0474] **Pharmacokinetics study:**

[0475] rFVII, rFVII-CTP, and rFVII-(CTP)₂ harvests were administered intravenously to Sprague-Dawley rats (six rats per substance) with a dose of 100µg/kg body weight. For all of the *in vivo* experiments, the amount of the respective protein was determined on the basis of FVII ELISA kit. For each FVII test substance, the injected amount was calculated by taking into account the differences in the molecular weight of rFVII versus rFVII-CTP, which leads to a different molar concentration.

[0476] Blood samples were drawn retro-orbitally using an altering sampling scheme to minimize interference of the sampling procedure levels to be quantified: from 3 rats at 30 and 90 min and at 2, 6, and 48 hrs, and from the remaining three rats at 15 and 60 min and at 1.5, 4, and 24 hrs alternately. Plasma was prepared immediately after sampling and stored at -20°C until analysis. FVII concentration was quantified by FVII ELISA specific assay. Half-life and area under the curve (AUC) were calculated using a linear trapezoidal rule. Comparison of these clearance parameters revealed that the *in vivo* half-life and rFVII-(CTP)₂ AUC are significantly higher than those of rFVII (Table 30).

Table 30: PK study parameters

FVII-CTP	IV	60	$\beta=51.06$	31353.9	0.287	73.7
FVII-CTP-CTP	IV	60	$\beta=13.66$	7626.8	1.18	15.4
FVII-CTP	IV	60	4.07	3314.7	6.195	6.2

[0477] **Characterization of recombinant FVIIa-CTP:**

[0478] During activation, FVII is cleaved at R152 resulting in heavy and light chain domains that are held together by a single disulfide bridge. rFVIIa-(CTP)₂ is purified and activated by an ion exchange column purification process. In order to fully evaluate rFVIIa-(CTP)₂, the protein is loaded on SDS-PAGE under reducing conditions to commercial FVIIa (NovoSeven®). The heavy and the light chain domains are separated and migrate as separated bands of molecular weights 55 and 25 KDa. Both proteins react with antibodies specific for FVII, but the heavy chain of the rFVIIa-CTP specifically reacts with anti-CTP-specific antibodies, indicating that this band represents the FVII heavy chain fused to CTP. The light chain reacts specifically with anti-gamma carboxylase Ab. The FVIIa protein concentration is determined by FVIIa-specific ELISA kit.

[0479] **FVIIa N-terminal sequencing:**

[0480] rFVII-CTP-CTP in activated or zymogene purified proteins is separated by SDS-PAGE (on 12% Tris-Glycine) and subsequently electroblotted to a PVDF membrane. The bands of interest are cut out and put on a purified Biobrene-treated glass fiber filter. The N-terminal sequence analysis is carried out by Edmann degradation using a pulsed liquid protein sequencer equipped with a 140C HPLC microgradient system. The identity of the recombinant protein and proper pro-peptide cleavage is further verified by N-terminal sequencing.

[0481] **FVIIa clotting activity:**

[0482] In order to evaluate FVII-(CTP)₂ coagulation activity, activated partial thromboplastin time assay (aPTT) is performed. FVII-deficient plasma sample is substituted with rFVIIa (NovoSeven®) or rFVIIa-(CTP)₂. 300 µl of FVII deficient human plasma is mixed with 100 µl of FVIIa or rFVIIa-(CTP)₂ at specific concentrations, or normal pooled human plasma which is further diluted. Following 60 seconds incubation at 37°C. Tissue Factor (TF), CaCl₂, and phospholipids are added to the mixture. Clotting time in seconds is determined. Potency is assessed by comparing a dose-response curve of rFVIIa-(CTP)₂ to a reference preparation consisting of rhFVIIa or human pool normal plasma. One unit of FVIIa is defined as the amount of FVIIa which equals to the activity of 1 ml human normal plasma. The aPTT clotting activity of rFVII and rFVIIa-(CTP)₂ is measured on a coagulometer (Instrumentation Laboratory). The aPTT clotting activity of rFVIIa and rFVIIa-(CTP)₂ is similar.

[0483] **Pharmacokinetics studies in rats:**

[0484] In order to characterize the influence of the CTP addition to the rFVIIa on its longevity potential, a comparative pharmacokinetic study in rats is performed. NovoSeven® (rFVIIa) and rFVIIa-(CTP)₂ in TBS are injected IV to 6 SD rats. The levels of FVIIa over time are detected using a FVIIa ELISA kit. The half-life and AUC are calculated for each protein. Comparison of these clearance parameters reveals that the *in vivo* measures of half-life, recovery, and AUC of the rFVIIa-(CTP)₂ are superior to those of NovoSeven®.

[0485] **FVIIa-CTP *in vivo* efficacy model (FVIII-deficient mouse model of hemophilia):**

[0486] In order to assess the *in vivo* activity model, FVIII knockout mice are obtained, and a breeding colony is established. 10 µg of either commercial recombinant hFVIIa (NovoSeven®) or rFVIIa-(CTP)₂ are injected into the tail vein of an anaesthetized FVIII knockout mouse (22-28g). The amount of injected protein equals to the required concentration of FVIII in normal plasma (5µg/ml). Blood samples are taken from the clipped tail into heparinized capillary tubes at specific time points. Plasma samples are assessed for FVIIa levels by ELISA, and efficacy is measured by a PTT coagulation assay.

[0487] In this study, a fusion construct of FVII with CTP is generated. This recombinant protein is the basis for a treatment that provides a prolonged half-life and retention of therapeutic potency.

[0488] These results suggest that rFVIIa-(CTP)₂ has a similar therapeutic efficacy to rFVIIa in hemophilia patients. Moreover, this technology requires less frequent dosing. It appears that a single injection of rFVIIa-(CTP)₂ is sufficient to control bleeding episodes and reduce the number of injections that are needed during surgical intervention. This recombinant protein may be used as a long term prophylactic treatment.

EXAMPLE 5

Comparative assessment of Purified FVII-CTP₃, FVII-CTP₄, and FVII-CTP₅

[0489] 5.1 Study objective

[0490] Comparative assessment of pharmacokinetic parameters and clotting activity of FVII-CTP₄ and FVII-CTP₅ versus FVII-CTP₃.

[0491] 5.2 Production of FVII-CTP₄ and FVII-CTP₅ harvests

[0492] FVII cDNA fused at the C-terminal to four or five tandem CTP sequences was expressed in Dg44 cells using the Excellgene expressing system in the presence of 20 µg/L of vitamin K3 (Sigma, Mennadion). The harvest was collected (300 ml), filtered and frozen.

[0493] 5.3 Production of FVII-CTP₃ harvest

[0494] FVII-CTP₃ was expressed in-house in mammalian expressing system, CHO cells, using pCI-DHFR vector. Stable transfected pool #71 was grown in shake flasks, in the presence of 25 ng/L of vitamin K3 (Sigma). The harvests were collected and filtered.

[0495] All FVII-CTP harvests (3, 4 and 5 CTPs) were concentrated and dialyzed against TBS (50 mM Tris, 150mM NaCl, pH 7.4) using Pellicon XL MWCO 10kDa.

[0496] 5.4 Determination of FVII antigen level

[0497] FVII antigen level was determined using Human FVII ELISA kit (Zymotest HyPhen) (Table 31). The calculated protein concentration is the average of two independent runs.

Table 31: FVII antigen level

	FVII-CTP ₃	FVII-CTP ₄	FVII-CTP ₅
Av. (ng/ml)	224357.3	87884.1	589423
SD	44789.5	3248.7	5309
%CV	20.0	3.7	9

[0498] 5.5 FVII-CTP immune-blot

[0499] FVII-CTP₃, FVII-CTP₄, and FVII-CTP₅ harvests were loaded on 12% Tris-Glycine gel (*expedion*) using Precision plus dual color protein marker (Bio-Rad). The SDS-PAGE analysis was performed by Western immune-blot using anti-CTP polyclonal Ab (Adar Biotech Production) or anti-Gla Ab (American Diagnostica).

[0500] FVII fused to three, four and five CTP migrated at 80, 90 and 100kDa, respectively. As expected, FVII-CTP₄ and FVII-CTP₅ harvests from Excellgene contain low gamma carboxylation content as compared to FVII-CTP₃ harvest which was produced at Prolor since the production process wasn't optimized (Figure 21).

[0501] 5.6 Comparative assessment of FVII *in vitro* potency

[0502] A comparative assessment of the *in vitro* potency of HA purified (highly gamma carboxylated fraction) FVII-CTP₃, FVII-CTP₄, and FVII-CTP₅ versus normal human pool plasma was performed using a commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221304). All samples were serially diluted, and the potency was assessed by comparing a dose-response curve to a reference preparation consisting of normal human plasma. FVII-CTP₃ and FVII-CTP₅ demonstrated chromogenic activity lower than pooled normal plasma (Figure 22). FVII-CTP₄ demonstrated higher activity as reflected by EC50 ratios, compared to FVII-CTP₃ and FVII-CTP₅ (Table 32).

Table 32: FVII *In Vitro* Clotting Activity

Sample	EC50 (ng/ml)	Sample /plasma EC50 ratio
Plasma	0.05	
FVII 3CTP	0.12	2.72
FVII 4CTP	0.03	0.71
FVII 5CTP	0.06	1.35

5

5.7 FVII *In Vitro* Clotting Activity:

[0504] Factor VII (FVII) activity assay, which was performed in Sheba Medical Center, the Israel National Coagulation Center, is a prothrombin (PT)-based assay using immuno-adsorbed plasma deficient in Factor VII (Siemens). The PT reagent is innovin, and the assay is performed in the Sysmex® CA 1500 instrument. FVII normal range is within 55-145%.

10

Table 33: FVII *In Vitro* Chromogenic Activity

Sample	FVII % of activity	Concentration tested (µg/ml)	Concentration in sample (µg/ml)
FVII 3CTP	36	0.5	224.2
	18	0.25	
	6	0.125	
FVII 4 CTP	334	0.5	87.9
	176	0.25	
	93	6.25	
FVII 5 CTP	38	0.5	58.9
	19	0.25	
	10	0.125	

[0505] Since the normal level of circulating FVII in the body is around 0.5 µg/ml, FVII-CTP₃ and FVII-CTP₅ harvests exhibit 3-fold reductions in their coagulation activity versus normal human pool plasma; this result correlates with the obtained chromogenic activity (Table 33).

15

[0506] The FVII-CTP₄ harvest exhibits a 3-fold increase in its potential coagulation activity versus normal human pool plasma as observed in the chromogenic activity assay (Table 33). The activity percentage of FVII-CTP₄ is much higher compared to activity percentage of

FVII-CTP₃ and FVII-CTP₅. Methodological limitations of the ELISA method may limit the accuracy of Ag level calculations of FVII-CTP₄.

[0507] 5.8 Pharmacokinetic study

[0508] Two pharmacokinetic studies were performed in order to determine the FVII-CTP₃,
 5 FVII-CTP₄, and FVII-CTP₅ pharmacokinetics (PK) parameters. During the first study, FVII-CTP₃, FVII-CTP₄, and FVII-CTP₅ (Group A, B and C, respectively) were administered in a single intravenous injection to Sprague Dawley rats (six rats per treatment) in a dose of 250 µg/kg body weight. Blood samples were drawn retro-orbitally from 3 rats alternately at 0.083, 0.5, 2, 5, 8, 24, 48, 72 and 96 hours post-dosing (Table 34). Citrated plasma (0.38%) was
 10 prepared immediately after sampling and stored at -20°C until analysis.

Table 34: Pharmacokinetic Study Design – Concentrated Harvest

Treatm ent Group	Test Article	No. of animals/ group/ time point	Dose Rou te	Dose Level (µg per animal)	Inje cted Vol. (µl)	Conc. (µg/ml)	Time-Points (hours post-dose)
A	FVII- CTP*3	6	IV	50	200	250	0 (Pre-dose) 0.083, 0.5, 2, 5, 8, 24, 48, 72, 96
B	FVII- CTP*4	6	IV	50	200	250	0 (Pre-dose) 0.083, 0.5, 2, 5, 8, 24, 48, 72, 96
C	FVII- CTP*5	6	IV	50	200	250	0 (Pre-dose) 0.083, 0.5, 2, 5, 8, 24, 48, 72, 96

[0509] FVII concentration in plasma samples were quantified using human FVII Elisa kits
 15 (Zymutest FVII-Biophen). The pharmacokinetic profile was calculated and is the mean of 3 animals at each time point. Terminal half-life values were calculated using PK Solutions 2.0 Software. Table 35 below summarizes the calculated FVII concentrations at the different sampling time points. The PK profile (Figures 23-24) and a summary of the PK parameters (Table 36) are also presented below. FVII-CTP₅ demonstrated a superior profile as compared
 20 to FVII-CTP₃ and FVII-CTP₄ (Table 36).

Table 35: First Pharmacokinetic Study - FVII Concentrations

Time (hr)	AVE-FVII- 3-CTP (ng/ml)	SD	AVE-FVII- 4-CTP (ng/ml)	SD	AVE-FVII- 5-CTP (ng/ml)	SD
--------------	-------------------------------	----	-------------------------------	----	-------------------------------	----

0.083	4214	583	3600	427	4888	504
0.5	3386	892	5213	1682	5384	2549
2	1138	219	3603	1338	3082	289
5	1390	374	2726	1127	2480	561
8	333	167	1349	44	2316	633
24	133	12	476	98	788	34
48	38	3	165	24	384	61
72	12	2	91	62	167	31
96	26	1	42	8	93	49

Table 36: Pharmacokinetic Analysis

	FVII-3-CTP	FVII-4-CTP	FVII-5CTP
half-life (0.083-8 hr) (hr)	2.5	4.9	6.6
half-life (8-72hr) (hr)	13.3	16.6	17.7
AUC (ng-hr/ml)(8-72hr)	18374.6	51224.4	72954.2
Vd (ml/kg)(8-72hr)	203.7	91.9	67.7
CL(ml/hr/kg) (8-72hr)	10.6	3.8	2.7

5 [0510] The addition of four or five CTPs significantly elongated FVII half-life as compared to 3 CTPs by 2- and 3-fold, respectively (Table 36). This superiority was more significant in the initial part of the study (0.083-8 hr), suggesting a potential improved protein recovery and reduced extra vascular clearance. AUC following FVII-CTP₄ and FVII-CTP₅ administration increased by 3- and 4-fold, respectively, versus FVII-CTP₃. Clearance was also reduced while adding 4 and 5 CTPs to FVII (Table 36).

[0511] As observed in the study, the addition of four and five CTPs significantly elongated FVII half-life as compared to 3 CTPs, both in the initial and terminal half-life. The half-life values in the first and second study are different due to a different analysis approach which was effected by the dose and study duration, nevertheless the overall trend was maintained.

15 The AUC following FVII-CTP₄ and FVII-CTP₅ administration increased by 2.5- and 7-fold, respectively, versus FVII-CTP₃.

[0512] **5.9 Conclusions:**

[0513] In this study, the PK parameters and potential clotting activity of FVII-CTP₃, FVII-CTP₄, and FVII-CTP₅ were assessed. Fusion of 4 and 5 CTPs to FVII provided a superior and improved half-life, exposure and reduced clearance as compared to FVII-CTP₃ while

maintaining a similar chromogenic and *in vitro* clotting activity. These results were observed at different concentrations of protein and were consistent for both harvest and purified protein. While evaluating the overall effect of fusion of CTP at the C terminus to FVII, fusion of 1-5 CTPs considerably increased the half-life and AUC of FVII in a CTP proportional manner, suggesting that as the CTP portion of the molecule increases, FVII longevity and stability is significantly improved while maintaining its initial *in vitro* clotting activity, as summarized in Table 37 hereinbelow.

Table 37:

Comparative assessment	T _{1/2} percent increase	AUC percent increase
FVII vs. FVII-CTP₂	268	200
FVII-CTP₂ vs. FVII-CTP₃	67	57.8
FVII-CTP₃ vs. FVII-CTP₄	24	178
FVII-CTP₄ vs. FVII-CTP₅	6	42

[0514] As previously reported, FVII half-life correlates with the half-life of the activated form of FVII (FVIIa) both in humans and animals. Therefore, it is anticipated that a similar improvement in half-life will be obtained for the activated versions following CTP fusion.

EXAMPLE 6***FVII-CTP₃ feasibility studies in FVIII-deficient hemophilic mice***

[0515] Studies described hereinabove testing FVII-CTP, FVII-CTP₂ and FVII-CTP₃ harvest PK profile and coagulation activity vs. a commercial FVII were conducted. FVII-CTP₃ exhibited an improved PK profile while maintaining its coagulation activity vs. FVII-CTP and FVII-CTP₂ harvests or rhFVII. In order to further characterize FVII-CTP₃ *in vitro* and *in vivo* properties, a mini stable pool expressing and secreting the protein was generated, and purification and activation processes were developed.

[0516] In the current study, the pharmacokinetic and pharmacodynamic properties of FVIIa-CTP₃ were tested in FVIII-deficient mice. The PK profile of the protein was evaluated. A FVIIa specific activity-based PK profile was established and compared to commercial product NovoSeven®. In addition, the long-lasting *in vivo* hemostatic capabilities of FVIIa-CTP₃ to induce coagulation in FVIII-deficient mice after a tail vein transection (survival study) were tested.

[0517] Study Objectives:

[0518] To evaluate the pharmacokinetic and pharmacodynamic parameters of FVIIa-CTP₃ vs. commercial rhFVIIa (NovoSeven®) in FVIII-deficient mice following a single IV administration at a similar activity dose.

[0519] To determine the *in vivo* ability of FVIIa-CTP₃ to maintain homeostasis in FVIII-deficient mice by a single IV administration of FVIIa-CTP₃ and NovoSeven® at a similar activity dose followed by a challenge of tail vein transection (survival study).

[0520] **Production of FVII-CTP₃ harvest:**

[0521] FVII-CTP₃ was expressed in-house in Dg44 cells using a pCI-DHFR vector. Stable transfected pool #71 was grown in shake flasks, in the presence of 25 ng/L of Vitamin K3 (Sigma). Cell suspension was cultured and harvested following viability decline to 60-80%. The harvest was filtered and frozen at -70°C.

[0522] **Determination of harvest FVII antigen level:**

[0523] FVII antigen level was determined using human FVII ELISA kit (Zymotest HyPhen) (Table 38). The antigen level was calculated per each pooled harvest batch.

Table 38: FVII-CTP₃ antigen level

	FVII antigen level		
	PK-PD study		Survival study
	harvest 31A	harvest 31B	harvest 38
Av (µg/ml)	16.0	15.9	16.6
STD	1.5	0.0	0.8
% CV	9.1	0.1	4.9

[0524] **FVII-CTP₃ purification process (Figure 25)**

[0525] **Process outline**

[0526] Following a short purification study, the following purification process using 2 columns was performed. VII-Select affinity column (GE) and Ceramic Hydroxyapatite type 1 (HA), 40 µm (Bio Rad), FVII-CTP₃ γ-carboxylated enriched protein was purified. Auto-activation was induced by incubation of purified FVII-CTP₃ in the presence of CaCl₂ overnight at 2-8°C. The purification process is in its final developmental stage and is being optimized, thus part of the purification steps are not identical in the two batches.

[0527] **Ultra-filtration/diafiltration (UFDF) using 10kDa hollow fiber or Pellicon cassette**

[0528] Clarified harvest was thawed at 4°C over the weekend (2-3 days).

[0529] In Batch 31, clarified harvest (12 liters) was concentrated 4-fold (in two successive runs) using a hollow fiber cartridge (GE Healthcare Catalog # UFP-10-C-4X2MA) with a 10 KDa molecular weight cut-off. Concentrated harvest was dia-filtrated against 1-2 volumes of TBS (50mM Tris 150mM NaCl pH 7.4).

[0530] In Batch 38, clarified harvest (8.5 liters) was concentrated 4-fold using a Pellicon 2 (Millipore) cassette with a 10 KDa molecular weight cut-off. Concentrated harvest was directly loaded on VII-Select column.

[0531] Both ultra-filtrations were performed on ice with ice cold buffers. UFDF samples
5 were filtered 0.22 μ m before loading.

[0532] **Capture on FVII-Select column**

[0533] The UFDF or concentrated harvest was loaded on VII-Select column (XK16/20, CV 18ml), pre-equilibrated with TBS pH 7.4. The column was washed with 50 mM Tris-HCl, 0.5M NaCl pH 7.5, and FVII-CTP₃ was eluted with 50 mM Tris-HCl, 1M NaCl 50% (v/v),
10 Propylene Glycol pH 7.5. The process was performed in two successive cycles utilizing the same column.

[0534] **Gamma carboxylation-based separation on a ceramic hydroxyapatite column**

[0535] The eluted product was diluted 1:10 with 10 mM sodium phosphate pH 6.8 and loaded on ceramic hydroxyapatite columns (XK16/20, CV 24ml). The column was washed
15 with 59 mM sodium phosphate pH 6.8 and the γ -carboxylated rich fraction of Factor VII was eluted with 500mM sodium phosphate pH 6.8. This process was performed in two successive cycles on the same column. At each batch, the eluates of the two cycles were pooled and concentrated to 1.7-2 mg/ml and dia-filtered with 20 mM Tris-HCl, 100 mM NaCl pH 8.2 to reduce volume and prepare the material for the activation step.

20 [0536] **FVII activation**

[0537] Purified FVII-CTP₃ was diluted to 1 mg/ml and incubated in 20 mM Tris-HCl, 100 mM NaCl and 1mM CaCl₂ pH 8.2 at 2-8°C for 24 hours. Activation was terminated by buffer exchange (UFDF) to preliminary formulation buffer (20 mM Citrate, 240 mM NaCl, 13.3 mM Glycine, pH 6.9).

25 [0538] **FVII-CTP₃ and FVIIa-CTP₃ analytical properties:**

[0539] **SDS-PAGE and Western blots**

[0540] Purified FVII-CTP₃, and FVIIa-CTP₃ were loaded on 12% Tris-Glycine gel using Precision Plus Dual Color Protein Marker (Bio-Rad). The SDS-PAGE Coomassie analysis was performed by staining the gel with Coomassie brilliant blue reagent (5 or 10 μ g of protein/lane). Western blot analysis was performed (1 μ g of protein/ lane) using anti-human
30 FVII polyclonal Ab (R&D systems; AF2338), anti-human gamma carboxylation monoclonal antibody (American Diagnostics Catalog #499, 3570), and anti-CTP polyclonal Ab. Under reduced conditions, FVII-CTP₃ migrated at 75KDa, and FVIIa-CTP₃ migrated as two main

bands: a heavy chain at 50 kDa, and a light chain at 25 kDa, represented in Figure 26 as Bands 2 and 3, respectively.

[0541] The purification procedure significantly enriched the FVII-CTP₃ portion while reducing impurities. The purification process yield was 25-30% FVII (according to ELISA).

5 Most of the protein lost during purification had low FVII chromogenic activity or no activity. Based on Coomassie-stained SDS-PAGE, the reduced FVIIa-CTP₃ contains more than the predicted bands. A band migrating to around ~75 kDa represents non-activated FVII (Figure 26, Band 1). This band consists of two bands with minor MW differences, which might reflect different γ -carboxylation content. Additional bands with MW lower than 20 kDa were
10 observed. This was previously reported to be degradation products of the heavy chain.

[0542] **FVII-CTP₃ chromogenic activity:**

[0543] A comparative assessment of the *in vitro* potency of FVII-CTP₃ harvest, in-process fractions, and purified FVII-CTP₃ versus human pool normal plasma was performed using a commercially available chromogenic activity test kit, BIOPHEN (Hyphen BioMed 221304).
15 FVII-CTP₃ harvest and protein were serially diluted and the potency was assessed by comparing a dose-response curve to a reference preparation of normal human plasma. Following FVII-CTP₃ purification, the chromogenic activity was significantly improved, and non-active fractions were separated mainly by HA column (Figure 27). A strong correlation between FVII chromogenic activity and detection of FVII with monoclonal anti-Gla
20 antibodies in Western blot was observed. The potency of FVII chromogenic activity as reflected by EC50 value in harvest is affected from both carboxylated and non-carboxylated FVII fractions. Following purification and enrichment of FVII-CTP₃ γ -carboxylated fraction, the activity was improved, demonstrating the important contribution of γ -carboxylation to FVII activity (Figure 27). This parameter is crucial for proper FVII *in vivo* activity and will be
25 further addressed in a clone development program.

[0544] **Protein determination by A280**

[0545] The theoretical extinction coefficient of FVIIa-CTP₃ and NovoSeven® was calculated using the ProtParam algorithm (<http://web.expasy.org/protparam>). The calculation is based on amino acid sequence. The calculated extinction coefficients for FVII-CTP₃ and
30 NovoSeven® is 1.186 and 1.406, respectively. These values represent the absorbance of 1 g/L at 280 nm.

[0546] The extinction coefficient difference between the two proteins derives solely from the increase in molecular weight of FVIIa-CTP₃ compared to NovoSeven®, since CTP lacks aromatic and cysteine residues, thus does not contribute to the absorbance.

[0547] Protein determination by A280 is used for final FVII, and for purified in-process samples, starting from the elution of VII-Select column.

[0548] **Determination of FVIIa antigen level**

[0549] FVIIa antigen level was determined using Human FVIIa ELISA kit (IMUBIND, American Diagnostica). The antigen level was calculated per each batch. However, this tool was not useful for the determination of the dose for injection, since it did not represent the amount of active product.

[0550] **Clotting assay of FVIIa- Staclot® VIIa-rTF**

[0551] FVIIa is derived from an intra-chain cleavage of the single-chain FVII. Native tissue factor (TF) is a cofactor of FVIIa. Upon binding to TF, FVII mediates the activation of Factor X to Xa, while itself is transformed to FVIIa. The soluble tissue factor is the extracellular part of native tissue factor. It can no longer activate FVII by auto-activation, but the FVIIa bound to tissue factor can activate FX to FXa.

[0552] The recombinant soluble tissue factor (rsTF) used in this assay utilizes the FVIIa specificity to construct a FVIIa clotting test. rsTF, in the presence of FVIIa, calcium and phospholipids leads to coagulation of plasma, without activating FVII to FVIIa.

[0553] The observed clotting time in this system has an inverse relationship with the FVIIa content in the tested sample, with no interference of FVII presence in the sample.

[0554] The assay was performed by Omri Laboratories (Nes-Ziona, Israel). FVIIa activity was evaluated for both NovoSeven® following reconstitution and FVIIa-CTP₃ prior to each study. NovoSeven® activity did not correlate with the anticipated activity as reported on the vial, but the discrepancy might be due to a different approach for activity evaluation. Table 39 summarizes the FVIIa clotting activity per volume without considering the protein concentration.

Table 39: FVIIa clotting activity of batch products

	PK study		Survival Study	
	FVIIa-3*CTP (FVIIa 31)	NovoSeven®	FVIIa-3*CTP (FVIIa 38)	NovoSeven®
Activity (U/ml)	1.3E+06	2.5E+05	1.3E+06	7.4E+05

[0555] **Specific activity of FVIIa-CTP₃**

[0556] FVIIa specific activity (which is calculated as the activity/ml divided by protein concentration) was calculated based on A280 and is presented in Table 40. When comparing the specific activity of the two molecules, which differ in MW, compensation must be made

in order to normalize the activity (i.e. because of the molecular weight difference, the number of active sites in 1 mg of NovoSeven® is 1.185-fold higher than in FVIIa-CTP₃). Calculation of the conversion factor is presented in the following equation:

$$\begin{aligned} \text{Normalized_SA} &= \frac{\text{SA(FVIIa-CTP}_3\text{)}}{\text{MW(FVII-CTP}_3\text{)}} \times \text{MW(Native_FVII)} = \\ &= \frac{\text{SA(FVIIa-CTP}_3\text{)}}{53419.5\text{Da}} \times 45079.1\text{Da} = \text{SA(FVIIa-CTP}_3\text{)} \times 1.185 \end{aligned}$$

Table 40: FVIIa-CTP₃ specific activity compared to NovoSeven®

Sample	Average A280	STDV (n=9)	% CV	Extinction coefficient	Prot conc. (mg/ml)	U/ml	Specific Activity		Fold decrease from NovoSeven®
							U/mg protein	U/mg FVIIa	
NovoSeven®	1.274	0.031	2.398	1.406	0.906	8.36E+05	9.23E+05	9.23E+05	1.0
FVIIa-CTP ₃	4.396	0.092	2.094	1.186	3.706	7.23E+05	1.95E+05	2.31E+05	4.0

[0557] **FVIIa-CTP₃ PK-PD study:**

[0558] **Study outline**

[0559] FVIIa-CTP₃ and rhFVIIa (NovoSeven®, NS) were administered in a single intravenous injection to C57B FVIII-deficient mice at a dose of 6.4E6 U/kg body weight (160,000 U/animal). Blood samples were drawn retro-orbitally from 4 mice alternately at 0.166, 0.5, 2, 4, 8, 12, 24, 34, 48, 58, and 72 hours post-dosing (Table 41). Citrated plasma (0.32%) was prepared immediately after sampling and stored at -20°C until analysis. FVIIa clotting activity level was evaluated, and a detailed PK analysis was performed. The study was performed by Omri Laboratories (Nes-Ziona, Israel).

Table 41: Study outline

Treated Groups	Test Article	No. of animals/ group/ timepoint	Dose Route	Amount of Units/ animal	Injected Vol. (μl)	Time-Points (hours post-dose)
A	rhFVIIa	4	IV	1.6e5	200	0 (Pre-dose) 0.166, 0.5, 2, 4, 8, 12, 24, 34, 48, 58, 72

B	FVIIa-CTP ₃	4	IV	1.6e5	200	0 (Pre-dose) 0.166, 0.5, 2, 4, 8, 12, 24, 34, 48, 58, 72
---	------------------------	---	----	-------	-----	--

[0560] **FVIIa-CTP₃ PK profile in FVIII-deficient mice**

[0561] FVIIa activity in blood samples was quantitated using a StacLOT® VIIa-rTF kit (Stago, Parsippany, NJ). The pharmacokinetic profile was calculated for each protein and represents the mean of 4 animals at each time point. Figure 28 presents the PK profile of FVIIa throughout the experiment. FVIIa recovery is presented in Table 42. A summary of the PK parameters is presented in Table 43.

[0562] Table 41 summarizes the clotting activity values following administration of either NovoSeven® or FVIIa-CTP₃. FVIIa-CTP₃ and NovoSeven® reached maximal activity half an hour post-dosing. NovoSeven®'s highest activity value reached only 43% of FVIIa-CTP₃'s maximal activity value. FVIIa-CTP₃ clotting activity was maintained for a longer period of time, demonstrating elongated activity. Clotting activity for the NovoSeven®-treated mice was undetectable at time points later than 12 hours, while FVII-CTP₃ treated mice continued to retain measurable activity at 48 hours post dosing (Table 41 and Figure 28).

[0563] The addition of three tandem CTP copies to FVIIa elevated recovery by 100% (Table 42), as measured by the highest activity post-dosing and compared to the anticipated activity based on *in vitro* analysis, and increased the half-life and mean resident time (MRT) 5-fold. The exposure time (AUC) was increased 3-fold (Table 43).

Table 41: FVIIa clotting activity following single IV injection

Time after administration (hours)	Average FVIIa Clotting Activity (U/ml)	
	FVIIa-CTP ₃	NovoSeven®
0.16	6.8E+07	3.2E+07
0.5	9.7E+07	4.3E+07
2	2.1E+07	3.9E+06
4	7.7E+06	7.3E+05
8	2.7E+06	4.2E+04
12	3.7E+05	6.2E+03
24	2.4E+04	BLQ
34	4.6E+03	BLQ
48	1.5E+03	BLQ

Table 42: FVIIa-CTP₃ recovery

Treated Groups	Test Article	Amount of Units/animal	Practical administered dose (U/ml)	*Anticipated Cmax (U/ml blood)	Cmax (U/ml)	% Recovery
A	rFVIIa	1.60E+05	1.20E+06	1.40E+05	4.25E+04	30
B	FVIIa-CTP ₃	1.60E+05	1.29E+06	1.50E+05	9.74E+04	64.6

*anticipated Cmax is derived from administered dose divided in blood volume

5

Table 43: PK parameters of FVIIa-CTP₃ vs. NovoSeven®

PK Parameters	NovoSeven®	FVIIa-CTP ₃
Half-life- α (0.5-12hr)	0.94	1.57
Half-life- β (12-48hr)	NA	4.62
AUC (mU*hr/ml)	5.80E+07	1.80E+08
Vd/Kg (ml/Kg)	1408	2375
CL/Kg (ml/hr/Kg)	1034	356
MRT (hr)	1.3	6.7

[0564] **Thrombin generation assay (TGA)**

[0565] The generation of thrombin is a fundamental part of the clotting cascade and as such an estimate of how well a particular individual can generate thrombin may correlate with either a risk of bleeding or thrombosis. Commonly measured variables when analyzing thrombin generation include: the lag time, the time to peak thrombin generation, the peak, the endogenous thrombin potential [ETP] (i.e., the area under the curve and the tail), the time course of the thrombogram ("TG"). After a lag time, a burst of thrombin is observed. However, clotting occurs at the end of the lag time, when more than 95% of all thrombin has not yet formed. The thrombin generation assay was performed at Omri Laboratories, using Thrombinoscope reagents supplemented with human hemophilic plasma. TGA reflects of the clotting ability in mice plasma, derived from injection of NovoSeven® and FVIIa-CTP₃. Figure 29 presents TGA parameter values for mice plasma following administration of either FVIIa-CTP₃ or NovoSeven®. Following FVIIa-CTP₃ administration, all three parameters (rate of thrombin generation, maximal amount of generated thrombin and KIIa) demonstrate an advantage of FVII-CTP₃ over NovoSeven® treatment. This further strengthens the notion of potential long-acting superiority of FVII-CTP₃ as compared to NovoSeven®.

[0566] **FVIIa-CTP₃ Tail Vain Transection (TVT) study:**

[0567] **Study outline**

[0568] The data obtained from the PK/PD test for FVIIa-CTP₃ provided insight into the functionality of FVIIa-CTP₃, and demonstrated that FVIIa-CTP₃ had a pharmacokinetic advantage when compared with NovoSeven®. However, the ability of the protein to induce a clot *in vivo*, after a traumatic event has not yet been demonstrated. In order to evaluate the ability of FVIIa-CTP₃ to stop bleeding, the same FVIII-deficient mice model was employed for a bleeding challenge.

[0569] FVIII-deficient mice were administered a single intravenous injection of FVIIa-CTP₃ or NovoSeven®. The mice were dosed with drug in amounts that provided equivalent FVIIa activity (1.6E05 units, 200 µl), calculated according to the potency of each drug evaluated in the FVIIa clot activity assay (Table 44). The administered doses were 9 mg/kg of NovoSeven®, and 40 mg/kg of FVII-CTP₃ due to the reduced activity of FVIIa-CTP₃. A control group was injected with 200 µl vehicle.

[0570] The tail vein was transected 2.7 cm from the tail tip 15 min (injection 1), 24 hours (injection 2) or 48 hours (injection 3) post-administration, and mice survival was recorded for 24 hours.

Table 44: Evaluation of injected samples

Injection No.	NovoSeven®			FVIIa-CTP ₃			
	protein conc. (mg/ml)	Activity (U/ml)	Specific Activity (U/mg)	protein conc. (mg/ml)	Activity (U/ml)	Specific Activity (U/mg)	Specific Activity (normalized)
1	0.91	8.0E+05	8.8E+05	3.63	6.6E+05	1.8E+05	2.2E+05
2	0.92	8.3E+05	9.0E+05	3.81	7.8E+05	2.0E+05	2.4E+05
3	0.89	8.8E+05	9.9E+05	3.68	7.3E+05	2.0E+05	2.3E+05

[0571] Protein concentration was determined by A280.

[0572] Results

[0573] Data from the vehicle-injected control groups for the three injections (5 animals x 3 injections), were summarized and are presented in Figure 30. 30% survival was observed 24 hours after tail vein transection.

[0574] NovoSeven® and FVIIa-CTP₃-treated mice demonstrated proper hemostatic activity after tail vein transection performed 15 min after FVIIa administration. A 100% survival rate was observed in FVIIa-CTP₃ and NovoSeven® treated animals (Figure 30).

[0575] The reduced clearance rate of FVII-CTP₃ which was demonstrated in the PK/PD study is most clearly appreciated after a tail vein transection performed 24 hours post-administration. A decline in the survival rate of NovoSeven® is observed. Similar to the control group, 50% death is observed within 10 hours. Meanwhile, 90% of FVIIa-CTP₃ treated mice survived (Figure 30). This result emphasizes the long-lasting efficacy of the FVIIa-CTP₃ treatment.

[0576] 48 hours after administration, a decline in survival rate is demonstrated in groups treated with either FVIIa-CTP₃ or NovoSeven® (Figure 30C). A slight improvement in FVIIa-CTP mice was observed, but the difference did not reach statistical significance.

[0577] **Discussion:**

[0578] CTP fusion to recombinant proteins extends the circulatory half-life of proteins while maintaining comparable activity. While the mechanism behind the reduced clearance of protein above a threshold size of 70 KDa is well understood with respect to renal clearance, additional protection is achieved following CTP fusion. CTP fusion is believed to sweep around the protein shield and protect it from proteolytic cleavage, to increase its radial molecular weight due to the highly negative charge and to reduce its affinity to hepatic clearance receptors.

[0579] The present study was aimed to provide specific insight on the impact of CTP fusion to FVII on protein half-life and clearance and also address the paradigm of its specific activity following this modification. FVIII-deficient mice were administered with a single IV injection of FVIIa-CTP₃ or recombinant commercial FVIIa (NovoSeven®) at similar dose (unit based) and a PK activity-based analysis was performed. FVIIa-CTP₃ demonstrated a superior longevity as reflected by 5- and 3.5-fold increase in its half-life and AUC, respectively. The specific activity (U/mg) of FVIIa-CTP as calculated by the Staclot® activity kit divided by the protein concentration measured by A280 was shown to be 4-5 times lower than the specific activity of NovoSeven®.

[0580] To build on the understanding of how CTP affects the haemostatic effects of FVIIa *in vivo*, the ability of FVIIa-CTP₃ to reduce bleeding was investigated. In the tail vein transection bleeding model in hemophilic mice model, rFVIIa administration can improve the survival rate of challenged animals and avoid their bleeding to death. In the study described herein, animals were administered with FVIIa-CTP₃ or NovoSeven®. Both molecules were

able to maintain homeostasis when the transection was performed 0.25 hours post-dosing. A significantly prolonged duration of activity was demonstrated for the FVIIa-CTP₃-treated group when the tail transection was performed 24 hr post dosing. The vehicle-treated group's survival rate was higher than anticipated and higher than that obtained in previous studies (50% vs. 20% in previous studies, data not shown). The percent survival of treated animals at is further evaluated at earlier time points, including at 36 hr post dosing.

[0581] In conclusion, it was demonstrated that FVIIa-CTP₃ has an increased duration of activity in hemophilic mice which translates into a longer duration of haemostatic effect when compared to NovoSeven®. The data gathered suggest that fusion of CTP to FVII is a technology with the potential to significantly improve prophylactic treatment in patients with hemophilia.

EXAMPLE 7: COMPARATIVE ASSESSMENT OF PURIFIED FVII-CTP₃ vs. FVII-CTP₅ PROFILE FOLLOWING SINGLE IV or SC INJECTION TO SD RATS

Study objective

[0582] Two studies were carried out:

[0583] The first study objective was to determine the pharmacokinetic parameters of rFVII-CTP₃ versus rFVII-CTP₅ following FVII select- and HA-column purification in male Sprague Dawley rats, after a single intravenous administration of 50µg/animal.

[0584] In the second study, rFVII-CTP₃-HA versus rFVII-CTP₅-HA pharmacokinetic parameters, were examined in male Sprague Dawley rats following a single intravenous or subcutaneous administration of 100 µg/animal.

RESULTS

[0585] Determination of FVII-CTP 3 and FVII-CTP 5 antigen level

[0586] FVII antigen level was determined using Human FVII ELISA kit (Zymotest HyPhen) (See table 45). T

[0587] **Table 45.** Summarizes the calculated protein concentration which is the average of three independent runs.

	FVII 3 CTP		FVII 5 CTP	
	<u>FVIIS 46 el.</u>	<u>FVII HA 46 el.</u>	<u>FVIIS el.</u>	<u>FVII HA 5 100%B</u>
	<u>Conc. Dial</u>	<u>Conc. Dial</u>	<u>Conc. Dial</u>	<u>Conc. Dial</u>
AVE (ng/ml)	3.78E+06	1.59E+06	1.88E+06	7.92E+05
SD	1.30E+06	6.03E+05	7.15E+05	3.57E+05
CV (%)	3.43E+01	3.80E+01	3.80E+01	4.51E+01

Western blot analysis of the examined samples

[0588] FVII-CTP_{3,5} samples were loaded on 4-12% bisTrisgel (*NuPage, invitrogene*) using Precision plus dual color protein marker (Bio-Rad). The SDS-PAGE analysis was performed by western immune-blot using polyclonal anti FVII Ab (R&D systems), anti CTP polyclonal Ab (Adar biotech production) or anti Gla Ab (American diagnostica). In summary, FVII fused to three and five CTP migrated at 80 and 100kDa, respectively (see Figure 31).

Comparative assessment of FVII *in vitro* potency

[0589] FVII activity assay, which was performed in Sheba medical center, the national coagulation center, is a PT based assay using immunoadsorbed plasma deficient in factor VII (Siemens). The PT reagent is innovin and the assay is performed in the Sysmex CA 1500 instrument. FVII normal range is within 55-145%. Sample activities are summarized in Table 46.

Table 46: Sample activity

Sample	Concentration (mg/ml) according to (NANODROP)	Concentration in tested sample (µg/ml)	Results (%)	Average-% of plasma
FVII-5CTP FVIIS el. Conc. Dial	2.19	2	87	16%
		1	30	
		0.5	10	
FVII-5CTP HA 5 100%B conc. Dial	1	2	97	21%
		1	36	
		0.5	13	
FVIIS 46 el. Conc. Dial	3.17	2	100	18%
		1	35	
		0.5	12	
FVII HA 46 el. Conc. Dial (1)	1.5	2	92	20%
		1	33	
		0.5	10	

[0590] The normal level of circulating FVII in the body is around 0.5µg/ml. Both, FVII-CTP₃ and FVII-CTP₅ exhibit about 5-fold reductions in their coagulation activity versus normal human pool plasma.

Pharmacokinetic study

[0591] Two pharmacokinetic studies were performed in order to determine the FVII-CTP₃ and FVII-CTP₅ (after FVII select and FVII HA column) pharmacokinetics (PK) profile and parameters. In the first study, FVII-CTP₃ and FVII-CTP₅ following FVII select/ HA purification were administered in a single intravenous injection to Sprague Dawley rats (six rats per substance) in a dose of 50 µg/rat.

[0592] Blood samples were drawn retro-orbital from 3 rats alternately at 0.083, 0.5, 2, 5, 8, 24, 48, 72, 96 and 120 hours post dosing. Citrated plasma (0.38%) was prepared immediately after sampling and stored at -20 until analysis.

[0593] In the second study, only samples after HA column were tested. These samples were administered in a single intravenous or subcutaneous injection to Sprague Dawley rats (six rats per substance) using a dose of 100 µg/rat. Blood samples were collected at the same time points and conditions as at the first study above.

[0594] **Table 47.** First study design (FVII select vs. FVII HA).

Treat ed Grou ps	Test Article	No. of animals/ group/	Dose Route	Dose Level (µg per animal)	Injected Vol. (µl)	Conc. (µg/ml)	Time- Points (hours post-dose)
A	FVII-CTP*3 batch 46 HA	6	IV	50	200	250	0 (Pre- dose) 0.083 0.5, 2, 5, 8, 24, 48, 72, 96, 120
B	FVII-CTP*3 batch 46 FVIIS	6	IV	50	200	250	0 (Pre- dose) 0.083 0.5, 2, 5, 8, 24, 48, 72, 96, 120
C	FVII- CTP*5batch 5 HA	6	IV	50	200	250	0 (Pre- dose) 0.083 0.5, 2, 5, 8, 24, 48, 72, 96, 120
D	FVII-CTP*5 batch 5 FVIIS	6	IV	50	200	250	0 (Pre- dose) 0.083 0.5, 2, 5, 8, 24, 48, 72, 96, 120

[0595] **Table 18.** Second study design (IV vs. SC)

Treated Groups	Test Article	No. of animals/ group/	Dose Route	Dose Level (µg per animal)	Injected Vol. (µl)	Conc. (µg/ml)	Time-Points (hours post-dose)
A	FVII-CTP*3 batch 46 HA	6	IV	100	200	500	0 (Pre-dose) 0.083 0.5, 2, 5, 8, 24,48,72, 96,120
B	FVII-CTP*3 batch 46 HA	6	SC	100	200	500	0 (Pre-dose) 0.083 0.5, 2, 5, 8, 24,48,72, 96,120
C	FVII-CTP*5batch 5 HA	6	IV	100	200	500	0 (Pre-dose) 0.083 0.5, 2, 5, 8, 24,48,72, 96,120
D	FVII-CTP*5 batch 5 HA	6	SC	100	200	500	0 (Pre-dose) 0.083 0.5, 2, 5, 8, 24,48,72, 96,120

[0596] The main differences between these two studies are the dosages and the route of administration. In the first study, rats were injected IV with 50 µg/rat, while in the second study, the rats were injected IV or SC with 100 µg/rat (total 500 µg/kg; rats weigh 200 g). The increase in the dosage is due to the change in the type of administration; SC administration requires higher amounts to achieve effects similar to IV administration.

5

Analysis of PK study

[0597] FVII concentration in plasma samples were quantified using human FVII Elisa kits (zymutest FVII-Biophen). Pharmacokinetic profiles were calculated and reflect the mean for 3 animals at each time point. Terminal half-live values were calculated using PK solutions 2.0 software. The table below summarizes the calculated FVII concentrations at the different sampling time points. PK profile and a summary of the PK parameters are presented in table below.

[0598] **Table 49.** First pharmacokinetic study (FVII select vs. FVII HA) -FVII concentrations (ng/ml).

Time (hour)	FVII CTP*3 BATCH 46 HA	FVII CTP*3 BATCH 46 FVII S	FVII CTP*5 BATCH 5 HA	FVII CTP*5 BATCH 5 FVII S
0.083	1816.3	1633.9	2064.3	1853.5
0.5	1523.7	1409.9	1351.4	1418.0
2	1284.9	1041.7	1389.7	834.4
5	607.9	531.6	722.7	737.2
8	524.2	430.0	712.2	614.6
24	115.5	132.9	272.5	201.8
48	21.1	31.6	62.3	90.4
72	9.5	15.8	29.1	31.8
96	BLQ	5.8	7.0	16.9
120	BLQ	BLQ	8.5	13.4

[0599] **Table 50.** Second pharmacokinetic study (IV vs. SC) -FVII concentrations (ng/ml).

Time (hour)	FVII CTP*3 BATCH 46 HA-IV	FVII CTP*5 BATCH 5 HA-IV	FVII CTP*3 BATCH 46 HA-SC	FVII CTP*5 BATCH 5 HA-SC
0.083	6452.6	6153.3	5.0	BLQ
0.5	3930.7	3660.6	14.5	14.6
2	1992.3	2176.2	113.6	96.2
5	1598.9	2087.3	106.6	70.5
8	781.6	1075.6	188.9	129.7
24	268.5	627.2	155.0	239.2
48	51.9	143.3	43.0	88.6
72	8.8	39.0	7.0	36.7
96	BLQ	10.8	BLQ	10.4
120	BLQ	8.2	BLQ	8.7

[0600] **Table 51.** PK Analysis- first pharmacokinetic study (FVII S vs. HA).

	FVII CTP*3 BATCH 46 HA	FVII CTP*3 BATCH 46 FVII S	FVII CTP*5 BATCH 5 HA	FVII CTP*5 BATCH 5 FVII S
half-life (0.083-8 hr) (hr)	4.3	4.0	5.51	5.59
half-life (8-72/96/120hr) (hr)	11.1	12.1	16.46	20.29
half life (8-72) (hr)	11.1	13.4	13.62	15.64
AUC(O-t)(obs area)(8-72/96/120hr)	14566.9	13686.4	21812.7	19307.9
AUC (∞) area(8-72/96/120hr)	14718.2	13788.1	22013.9	19701
Vd(area)/kg (ml/kg)(8-2/96/120hr)	271.1	316.1	269.7	371.5
CL(area)/kg(ml/hr/kg) (8-72/96/120hr)	17.0	18.1	11.356	12.69

5 [0601] The addition of five CTP elongated FVII half-life compared to 3 CTPs. Both forms of 5 CTP (i.e FVIIS and FVII HA) were detected at the long time points (96 and 120 hr), while FVII-3 CTP HA and FVIIS -3 CTP were detected until 72 hr and 96 hr, respectively. Based on this fact, the half-life of FVII-5 CTPs is longer than 3CTPs variants (see Fig. 32). Comparing half-life of all examined materials (3 and 5 CTPs) at the same time points (8-72

10 hr) showed that the half-life are similar, although 5 CTP are quite longer (Fig. 32).

[0602] **Table 52: PK analysis - second pharmacokinetic study-(IV vs. SC).**

	FVII CTP*3 BATCH 46 HA- IV	FVII CTP*5 BATCH 5 HA- IV	FVII CTP*3 BATCH 46 HA- SC	FVII CTP*5 BATCH H 5 HA-SC	Bioviability CTP*3	Bioviability CTP*5
half-life (0.083-8 hr) (hr)	3.0	3.9	-1.8	-3.18		
half-life (8- 72/96/120hr) (hr)	9.9	14.6	13.14	22.94		
half-life (8- 72) (hr)	9.9	13.0	13.14	29.47		
AUC(O- t)(obs area)(8- 72/96/120hr)	28866.8	43761.0	6600	9822.7	22.9	22.4
AUC (∞) area(8- 72/96/120hr)	28993.0	43934.4	6733	10110.8	23.22	23.01
Vd(area)/kg (ml/kg)(8- 72/96/120hr)	246.4	240.5	1407.6	1636.8		
CL(area)/kg(ml/hr/kg) (8- 72/96/120hr)	17.2	11.4	74.261	49.452		

[0603] Again, as observed in the first study, the addition of 5 CTPs elongated FVII half-life as compared to adding 3 CTP, both in the initial and terminal half-life and in both administration ways (IV and SC, see Fig. 33). As expected, following SC administration, FVII was first detected in the blood at a later time point as compared to when it was administered IV.

[0604] In the above, two PK studies were summarized. The main purpose of the first study was to check the difference between FVII-3CTP and FVII-5 CTP after 2 different columns: FVII select and FVII HA. In our previous studies, harvest vs. purified proteins were checked and it was found that the difference between 3 and 5 CTP versions of FVII was greater when harvest was injected to the rats.

[0605] There was no significant difference between the results of FVII 3\5 CTP after both columns, hence it was decided to inject FVII HA 3\5 CTP in the second study.

EXAMPLE 8: FVIIa-CTP₃ (MOD-5014) SURVIVAL STUDY IN FVIII DEFICIENT MICE FOLLOWING SUBCUTANEOUS INJECTION

Study Objective

[0606] To evaluate the efficacy of NovoSeven®, MOD-5014 (FVIIA-CTP₃) and MOD-5019 (FVIIA-CTP₅) in a tail vein transection study, following subcutaneous administration.

FVIIa-CTP₃(MOD-5014) and FVIIa-CTP₅(MOD 5019) analytical properties:

Protein determination by A280

[0607] Theoretical extinction coefficient of NovoSeven® was calculated using ProtParam algorithm (<http://web.expasy.org/protparam>). The calculation is based on amino acid sequence. The calculated extinction coefficient for NovoSeven® is 1.406, and for MOD-5019 is 1.075 (values represent the absorbance of 1 g/L at 280 nm). Extinction coefficient of MOD-5014 was determined by amino acid analysis at Mscan. The extinction coefficients for MOD-5014 is 1.27.

Clotting assay of FVIIa - STACLOT VIIa-rTF

[0608] FVIIa is derived from intra-chain cleavage of the single-chain FVII. Native tissue factor (TF) is a cofactor of FVIIa, upon binding to TF, FVII mediates the activation of Factor X to Xa, while itself is transformed to FVIIa. The soluble tissue factor is the extra cellular part of native tissue factor. It can no longer activate FVII by auto activation, but the FVIIa bound to tissue factor can activate FX to FXa.

[0609] The recombinant soluble tissue factor (rsTF) used in this assay is utilizing the FVIIa specificity to construct a FVIIa clotting test. Recombinant soluble tissue factor (rsTF), in the presence of FVIIa, calcium and phospholipids, produces coagulation of plasma without activating FVII to FVIIa.

[0610] The observed clotting time in this system has an inverse relationship with the FVIIa content in the tested sample, with no interference of FVII presence in the sample.

[0611] FVIIa activity was evaluated for reconstituted NovoSeven®, and for MOD-5014 and MOD-5019 prior to each study.

[0612] FVIIa specific activity (which is calculated as the activity/ ml divided by protein concentration) was calculated based on A280 and is presented in Table 53. When comparing specific activity of the two molecules, which differ in molecular weight, compensation must be made in order to normalize the activity (i.e. because of the molecular weight difference, the number of active sites in 1 mg of NovoSeven® is 1.185-fold higher than in MOD-5014 and 1.307-fold higher than MOD-5019). Hence, calculation of the conversion factor is presented in the following formula:

$$\text{Normalized_SA} = \frac{\text{SA(FVIIa-CTP}_3\text{)}}{\text{MW. (Native_FVII)}} \times \text{MW(FVII-CTP}_3\text{)} =$$

$$= \frac{\text{SA(FVIIa-CTP}_3\text{)}}{45079.1\text{Da}} \times 53419.5\text{Da} = \text{SA(FVIIa-CTP}_3\text{)} * 1.185$$

Table 53- MOD-5014 Specific activity compared to NovoSeven®

Sample	Protein conc. By A280 (mg/ml)	Specific Activity (U/mg FVIIa)	Fold decrease from @NovoSeven
@NovoSeven	0.93	52,487	1.0
MOD-5014 batch 73	1.4	25,490	2.05
MOD-5019 batch 9	3.0	11,698	4.48

Study outline

5 [0613] The most significant measurement is the ability of the protein to induce a clot *in vivo*, after a traumatic event. In order to evaluate the ability of MOD-5014 to stop bleeding, the same FVIII deficient mice model was employed for a bleeding challenge.

[0614] FVIII deficient mice were administrated with a single subcutaneous injection of MOD-5014, MOD-5019 or NovoSeven®. Group A and B were dosed with NovoSeven® and MOD-5014 respectively, in equivalent amounts as FVIIa activity. Group C was dosed with MOD-5019 in equivalent amount FVIIa protein as MOD-5014, in order to evaluate the critical factor (activity or amount of protein). The administrated doses were 4.2 mg/kg of NovoSeven®, and 8.6 mg/kg of MOD-5014 and MOD-5019. The tail vein was transected 2.7cm from tail tip 12 hours post administration, and mice survival was recorded for 24 hours.

15 [0615] **Table 54 – Group designation**

Group	Injection date	Test Article	Administered Dose		Injected Volume (µl)	No. of mice per group	Bleeding time, hours post dosing
			mg FVII /Kg	mU/Kg			
A	13.1.13	@NovoSeven	4.23	221,876	100	10	12
B	15.1.13	MOD-5014, batch 73	8.59	218,750	160	10	12
C	27.1.13	MOD-5019, batch 9	8.59	100,496	160	10	12

[0616] RESULTS

[0617] The experiment data is summarized in Table 55 and in Figure 34.

20 [0618] **Table 55. TVT study results**

Time	No. of surviving mice	% survival
------	-----------------------	------------

post TVT (h)	NovoSeven®	MOD- 5014	MOD- 5019	NovoSeven ®	MOD- 5014	MOD- 5019
0	9	10	10	100	100	100
1	9	10	10	100	100	100
2	9	10	10	100	100	100
3	8	10	8	89	100	80
4	6	9	8	67	90	80
5	5	9	7	56	90	70
6	4	8	5	44	80	50
7	3	8	5	33	80	50
8	2	7	5	22	70	50
9	1	6	5	11	60	50
10	1	5	5	11	50	50
11	1	3	5	11	30	50
12	1	3	5	11	30	50
24	1	3	4	11	30	40

[0619] 24 hours post TVT, only 11% of NovoSeven® injected mice have survived. 30% of MOD-5014 and 40% of MOD-5019 have survived to this time point. Subcutaneously injected MOD-5014 and MOD-5019 shows improved mice survival in comparison to NovoSeven®.

5 Nevertheless, the results are not optimal since more than 50% of the animals died during the experiment.

[0620] Factor VIIa, like other coagulation factors, is normally injected intravenously, in order to be directly available in the blood stream. However, the present invention shows that the compositions provided herein are surprisingly more effectively absorbed into the bloodstream after SC administration. To be able to administer FVIIa subcutaneously serves as an advantage as it can be used for prophylactic applications. Subcutaneous injections are also much easier for patients to self-inject, and are advantage when the patients are very young and their veins are small and difficult to find.

[0621] Hence, the subcutaneous application can be used for prophylactic treatment,

15

EXAMPLE 9: COMPARATIVE PK-PD STUDY OF RECOMBINANT MOD-5014 VS. NOVOSEVEN® FOLLOWING SUBCUTANEOUS ADMINISTRATION IN SD RATS

Study Objectives

[0622] To determine the pharmacokinetic and pharmacodynamic parameters of MOD-5014 versus commercial rFVIIa in SD rats following a single SC administration.

20

[0623] To compare two independent experiments (05010 & 05034) containing MOD-5014 products originated from two different clones (clone no. 28 vs. 61) by their pharmacokinetics parameters.

[0624] **Experimental Methods**

5 **Animals**

[0625] 24 males SD rats arrived from Harlan Laboratories Israel, Ltd, at least 4 days before the injections begin. The animals were healthy young adults, at ~200 gr at study initiation. The body weight variation of animals at the time of treatment initiation should not exceed $\pm 20\%$ of the mean weight of each sex. The health status of the animals used in this study is
10 examined on arrival. Only animals in good health are acclimatized to laboratory conditions and are used in the study.

Clotting assay of FVIIa - STACLOT VIIa-Rtf

[0626] The recombinant soluble tissue factor (rsTF) used in this assay is utilizing the FVIIa specificity to construct a FVIIa clotting test. rsTF, In the presence of FVIIa, calcium and
15 phospholipids produce coagulation of plasma, without activating FVII to FVIIa.

[0627] The observed clotting time in this system has an inverse relationship with the FVIIa content in the tested sample, with no interference of FVII presence in the sample.

[0628] FVIIa activity was evaluated for both NovoSeven® following reconstitution and MOD-5014 prior to each study. FVIIa specific activity was calculated based on A280. When
20 comparing specific activity of the two molecules, which differ in MW, compensation must be made in order to normalize the activity (i.e. because of the molecular weight difference, the number of active sites in 1 mg of NovoSeven® is 1.185-fold higher than in MOD-5014).

PK solver software

[0629] The pharmacokinetic parameters were calculated using PK solver software. The IV
25 administration curve analyzed as two compartmental CA bolus, and the SC administration as NCA Extravascular- Log linear trapezoidal analysis. Half-life, AUC, clearance and volume distribution specifications were calculated and the output parameters were studied in comparison between groups of experiments.

Experimental materials

30 [0630] Experiment no. 05010:

- A. NovoSeven® RT: (Lot # AU61553 prepared on 31.7.12*) FVIIa concentration by A280: 0.86 mg/ml. FVIIa Staclot activity assay: 56,867 U/mg. Injected dose: **946µg/kg**. *Pool of NovoSeven® aliquots, all from the same Lot no.

- B. Clone 28: MOD-5014 RS12-001: 0.77 mg/ml** based on A280. FVIIa Staclot activity assay: 34,162 U/mg. Injected dose: **850µg FVIIa/kg**.

[0631] Experiment no. 05034:

- A. NovoSeven® RT: (Lot #AU61347 prepared on 1.1.13) FVIIa concentration by A280: 0.82mg/ml, diluted to 0.4 mg/ml with sterile NS buffer. FVIIa Staclot activity assay: 55,688 U/mg. Injected dose: **360µg/kg** and **20,047.7 U/kg**.
- B. Clone 61: MOD-5014 Batch 75: 1.9 mg/ml** based on A280, diluted to 0.89 mg/ml with formulation buffer. Injected dose: **20,047.7 U/kg**. FVIIa clotting activity: 25,002* U/mg based on FVIIa Staclot activity assay.
- C. Clone 61: MOD-5014 Batch 81A: 2.36 mg/ml based on A280 (filtered on the morning of study day and re-measured at 280nm), diluted to 0.4 mg/ml with formulation buffer. Injected dose: **360µgFVIIa/kg**. FVIIa clotting activity: 24943U/mg based on FVIIa Staclot activity assay.
- D. Clone 61: MOD-5014 Batch 81A: 2.36 mg/ml based on A280, diluted to 0.89 mg/ml with formulation buffer. Injected dose: **20,047.7 U/kg**. FVIIa clotting activity: 24,943U/mg based on FVIIa Staclot activity assay.

[0632] **Study outlines**

[0633] **Experiment no. 05010**

[0634] MOD-5014 and NovoSeven® were administered in a single intravenous or subcutaneous injection to SD Rats in a dose of 0.9 mg/kg body weight. Blood samples were drawn from sinus orbital eye from 3 rats alternately at 0.5, 4, 8, 12, 24, 34, 48 and 58 hours post dosing. Citrated plasma (0.32%) was prepared immediately after sampling and stored at -20°C until analysis. The study was performed at "Science in Action", Nes-Ziona. FVIIa clotting activity level was evaluated and detailed PK analysis was performed at Prolor-Biotech.

Table 55: Study design 05010

Treated Groups	Test Article	No. of animals /group	No. of animals/ group/ Time point	Dose Route	Gender	Dose Level (µg/kg)	Injected Vol. (µl)	Time-Points (hours post-dose)
A	rFVIIa (Novo Seven®)	6	3	IV	M	946	220	0, 0.5, 4, 8, 12, 24, 34, 48, 58

B	rFVIIa RS12-001 (clone 28)	6	3	IV	M	850	220	0, 0.5, 4, 8, 12, 24, 34, 48, 58
C	rFVIIa (Novo Seven®)	6	3	SC	M	946	220	0, 0.5, 4, 8, 12, 24, 34, 48, 58
D	rFVIIa RS12-001 (clone 28)	6	3	SC	M	850	220	0, 0.5, 4, 8, 12, 24, 34, 48, 58

[0635] **Experiment no. 05034**

[0636] MOD-5014 and NovoSeven® were administered in a single subcutaneous injection to SD Rats in a dose of 0.9 mg/kg body weight. Blood samples were drawn from sinus orbital eye from 3 rats alternately at 0.5, 2, 4, 6, 8, 12, 24, 34, 48 and 72 hours post dosing. Citrated plasma (0.32%) was prepared immediately after sampling and stored at -20 °C until analysis. The study was performed at “Science in Action”, Nes-Ziona.

[0637] FVIIa clotting activity level was evaluated and detailed PK analysis was performed at Prolor-Biotech.

10 **Table 56: Study design 05034**

Treated Groups	Test Article	No. of animals/ group/ Time-point ***	Dose Route	Gender	Dose Level Per Animal (µg/kg)	Dose Level Per Animal (U/kg)	Injected Vol. (µl)	Time-Points (hours post-dose)
A	FVIIa (NovoSeven®)	3	SC	M	360	20047.7	207	0, 0.5, 2, 4, 6, 8, 12, 24, 34, 48, 72
B	FVIIa 75 (clone 61)	3	SC	M	801.84	20047.7	207	0, 0.5, 2, 4, 6, 8, 12, 24, 34, 48, 72
C	FVIIa 81A (clone	3	SC	M	360	8979.48	207	0, 0.5, 2, 4, 6, 8, 12, 24,

	61)							34, 48, 72
D	FVIIa 81A (clone 61)	3	SC	M	803.7 4	20047. 7	207	0, 0.5, 2, 4, 6, 8, 12, 24, 34, 48, 72

RESULTS

[0638] FVIIa activity in blood samples was quantitated using STACLOT VIIa-rTF kit (Stago). Pharmacokinetic profile was calculated for each protein and is the mean of 3 animals at each time point.

[0639] **Experiment no. 05010**

[0640] After background reduction: 15 mU/ml.

[0641] Figure 35 presents the PK profile of FVIIa following IV and SC administration of either NovoSeven® or MOD-5014. Summary of FVIIa activity values for each time point is presented in Table 57. IV and SC administration have different PK pattern (see Fig. 35; after background reduction: 15 mU/ml). The C_{max} following IV injection is higher than that obtained following SC injection, due to the presence of the drug immediately after administration in the blood (measured at 0.5hr, Table 57 and Table 58). However, after SC administration drug molecules transfer to intracellular matrix and tissues, thus C_{max} can be measured only after 2hr from injection. The total recovery of the drug after SC administration is lower than C_{max} value after IV injection.

[0642] 8hr after injection, NovoSeven® manifested an equal PK pattern when injected by either IV or SC, (After background reduction: 15mU/ml, Fig. 35). Moreover, clotting activity for the NovoSeven® treated mice was undetectable at time points later than 12 hours, while MOD-5014 treated mice continued to retain measurable activity at 58 hours post dosing (Table 57; after background reduction: 15 mU/ml; Fig. 35).

[0643] **Table 57.** FVIIa clotting activity of MOD-5014 vs. NovoSeven® following IV or SC administration

Time (hr)	NovoSeven® IV (A)		MOD-5014 IV (B)		NovoSeven® SC (C)		MOD-5014 SC (D)	
	mU/ml	%CV	mU/ml	%CV	mU/ml	%CV	mU/ml	%CV
0.5	304651.7	18.7	232818.3	5.0	11491.7	2.4	3691.7	19.0
4	40068.3	7.8	62085.0	9.5	21385.0	22.6	12018.3	15.8
8	5276.7	2.5	25931.7	6.1	5525.0	32.5	6445.0	2.2
12	255.0	13.8	5633.3	9.3	297.7	41.4	924.7	24.1
24	1.3	7.1	251.3	11.8	1.3	89.2	249.3	60.3
34	0.0		78.3	4.5	0.0		63.7	85.5
48			29.0	9.9	0.0		35.0	47.2
58			10.3	4.6	0.0		13.7	33.5

After background reduction: 15mU/ml.

[0644]

- 5 **Table 58: PK parameters of MOD-5014 vs. NovoSeven® following IV or SC administration**

A. IV

PK Parameters	NovoSeven® RT (A)	MOD-5014 (RS 12-001) (B)
Half-life- α (0.5-4hr)	0.24	1.04
Half-life- β (4-58hr)	1.31	3.17
AUC o-inf mU/ml*h	702467.95	820778.67
Vss [U/Kg/(mU/ml)]	0.13	0.13
CL [(U/Kg)/(mU/ml)/h]	0.08	0.04
MRT (hr)	1.74	3.62

B. SC

PK Parameters	NovoSeven® RT (B)	MOD-5014 (RS 12-001) (C)
Half-Life (hr)	1.40	7.78
Cmax (mU/ml)	21385.00	12018.33
AUC 0-inf (mU/ml*h)	115099.72	84158.87
MRT 0-inf (hr)	4.32	7.04
Vz/F (U/Kg)/(mU/ml)	0.95	3.88
Cl/F (U/Kg)/(mU/ml)/h	0.47	0.35

Experiment no. 05034

[0645] Figure 36 presents the PK profile of FVIIa following SC administration of either NovoSeven® or MOD-5017. Two different batches of clone no. 61 (#75 and #81) were examined in the same concentration or the same activity units, compared to NovoSeven®.

5 Summary of FVIIa activity values for each time point is presented in Table 59.

[0646] The results indicate a similar PK pattern after SC administration corresponding to previous experiments. Moreover, clotting activity for the NovoSeven® treated mice was undetectable at time points later than 12 hours, while MOD-5014 treated mice continued to retain measurable activity at 24 hours post dosing (Table 59 and Figure 36; and after
10 background reduction: 56 mU/ml (8, 12 hr) or 32 mU/ml (0.5, 2, 6, 14 hr)).

[0647] Clone no. 61 batch #81 (D) C_{max} (1,301mU/ml) was lower than the C_{max} values of clone no. 61 batch #75 (B) and NovoSeven® (A) (3,521mU/ml and 5,908mU/ml respectively), although they were all injected by the same unit activity (Table 6). However, batch #75 (B) and #81 (D) have the same activity units (559 mU/ml and 478 mU/ml
15 respectivaly) measured 8hr after injection (Figure 36 and Table 59; and after background reduction: 56 mU/ml (8, 12 hr) or 32 mU/ml (0.5, 2, 6, 14 hr)).

[0648] **Table 59:** FVIIa clotting activity of MOD-5014 (Clone 61 #75, #81) vs. NovoSeven® following single SC administration.

Time (hr)	NovoSeven® (A)		MOD-5014 Clone 61 Batch 75 (B) - equal U/kg		MOD-5014 Clone 61 Batch 81A (C) - equal conc.FVIIa µg/kg		MOD-5014 Clone 61 Batch 81A (D) - equal U/kg	
	mU/ml	%CV	mU/ml	%CV	mU/ml	%CV	mU/ml	%CV
0.5	3271.3	46.5	350.3	26.6	101.3	24.1	208.7	51.2
2	5908.0	18.1	3521.3	70.9	1294.7	7.0	1301.3	31.6
6	1411.7	23.6	1349.7	45.6	425.3	27.6	663.0	13.4
8	1029.0	12.4	559.3	52.7	152.7	19.5	478.0	25.4
12	121.3	9.9	563.0	17.4	148.7	36.3	712.7	16.2
24	1.0	25.0	117.0	41.9	21.3	36.4	99.0	36.7

After background reduction: 56mU/ml (8,12hr) or 32mU/ml (0.5, 2, 6, 14hr).

[0649] **Table 60:** PK parameters of MOD-5014 (Clone 61 #75, #81) vs. NovoSeven® following single SC administration.

PK Parameters	NovoSeven® RT (A)	MOD-5014 Clone 61 Batch 75 (B)- equal U/kg	MOD-5014 Clone 61 Batch 81A (C)- equal conc.FVIIa µg/kg	MOD-5014 Clone 61 Batch 81A (D)- equal U/kg
Half-Life (hr)	1.67	5.70	4.62	6.41
Cmax (mU/ml)	5908.00	3521.33	1294.67	1301.33
AUC 0-inf (mU/ml*h)	24688.18	20456.96	6260.23	13098.16
MRT 0-inf (hr)	3.73	7.86	6.40	10.59
Vz/F (U/Kg)/(mU/ml)	1.96	8.06	9.55	14.15
Cl/F (U/Kg)/(mU/ml)/h	0.81	0.98	1.43	1.53

[0650] This report summarized two PK studies; 05010 & 05034. We aim to provide specific insight on the impact of CTP fusion to FVII on protein half-life and clearance in subcutaneous administration and address the paradigm of its specific activity following this modification. In these studies, SD rats were administered with a single SC injection of MOD-5014 originated from two clones, and two different batches, compared to recombinant commercial FVIIa (NovoSeven®). The components were injected at similar FVIIa concentration (µg/Kg) or at the same activity level (U/Kg) and the PK activity based analysis was performed.

[0651] The main purpose of the first study was to verify the different PK parameters after IV and SC administration. Based on this study we can conclude that there is a difference between the PK pattern measured after IV or SC administration. A $t^{1/2}$ of 7.78 hr measured after MOD-5014 SC injection, and only 4.2 hr after IV injection. AUC values were the same (Table 58).

[0652] The second study however, focuses on the differences between two batches of MOD-5014 clone no. 61, which were injected by the same FVIIa concentration or at an equal activity unit, compare to NovoSeven®. At this study we showed that clone 61 batch #75 manifested better PK parameters than batch #81. Batch #81, which was injected by the same unit activity level, had lower Cmax from an unknown reason. Moreover, the same Cmax was

measured when injecting clone 61 batch #81 in two different doses (by FVIIa concentration or by unit activity), instead of 2.5-fold between the two activity values. Following analysis of both studies together, we can conclude that clone 28 manifested prolong $t^{1/2}$ parameter that clone 61 #75 (the better batch) after SC injection (7.78hr and 5.7hr respectively, Table 60).

5 We can also conclude that dissimilar time point samples create different PK pattern, which lead to variation in the PK curves. The patterns of the curves can teach us more about the drug behavior in the blood. Therefore, we decided to determine the time points similar to those detected by Baxter (0, 0.5, 2, 6, 8, 12, 24, 34, 48, 72hr). Moreover, the FVIIa concentration in 05010 experiment was too high, and was revised in the following SC experiment (05034). For
10 future PK studies, we decided to inject the component at 360 μ g FVIIa/kg for a dose.

[0653] Taken all together, we can learn more about our MOD-5014 product after SC administration in order to determine the most quality clone and batch and to decide the best method for MOD-5014 injection amount- by FVIIa concentration or activity units.

[0654] While certain features of the invention have been illustrated and described herein,
15 many modifications, substitutions, changes, and equivalents will now occur to those of ordinary skill in the art. It is, therefore, to be understood that the appended claims are intended to cover all such modifications and changes as fall within the true spirit of the invention.

1. A chorionic gonadotropin carboxy terminal peptide (CTP)-modified coagulation factor consisting of a coagulation factor and three CTPs attached to the carboxy terminus of said coagulation factor, wherein said coagulation factor is selected from:
 - a. a CTP-modified Factor IX (FIX) polypeptide having an amino acid sequence selected from the group consisting of: SEQ ID NO: 31 and amino acids 47-545 of SEQ ID NO: 31,
 - b. a CTP-modified Factor VII (FVII) polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 25 and amino acids 39-528 of SEQ ID NO: 25, and
 - c. a CTP-modified activated Factor VII (FVIIa) polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 25 and amino acids 39-528 of SEQ ID NO: 25.
2. The CTP-modified coagulation factor according to claim 1, wherein at least one CTP is encoded by the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 2.
3. The CTP-modified coagulation factor according to any one of claims 1 to 2, wherein at least one CTP is glycosylated.
4. The CTP-modified coagulation factor according to any one of claims 1 to 3, wherein at least one CTP is truncated.
5. The CTP-modified coagulation factor according to any one of claims 1 to 4, wherein at least one CTP is attached to the coagulation factor via a linker.
6. The CTP-modified coagulation factor according to claim 5, wherein the linker is a peptide bond.
7. The CTP-modified coagulation factor according to any one of claims 1-6, wherein said coagulation factor is a FIX polypeptide having an amino acid sequence selected from the group consisting of: SEQ ID NO: 31 and amino acids 47-545 of the sequence of SEQ ID NO: 31.

8. The CTP-modified coagulation factor according to any one of claims 1-6, wherein said coagulation factor is a CTP-modified FVII or FVIIa polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 25 and amino acids 39-528 of SEQ ID NO: 25.
9. A pharmaceutical composition comprising a CTP-modified coagulation factor as defined in any one of claims 1-8, and a pharmaceutically acceptable carrier.
10. Use of the CTP-modified coagulation factor according to any one of claims 1-8 or the pharmaceutical composition according to claim 9, in the preparation of a medicament for preventing or treating a blood clotting or a coagulation disorder in a subject.
11. The use according to claim 10, wherein the blood clotting or a coagulation disorder is hemophilia.
12. The use according to any one of claims 10-11, wherein said medicament is for subcutaneous administration.
13. The use according to any one of claims 10 to 12, wherein the subject is a child.
14. A method of extending the biological half-life, improving the area under the curve (AUC), reducing the dosing frequency, or reducing the clearance rate of a coagulation factor, the method comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of a FIX polypeptide, or comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of a FVII or FVIIa polypeptide, thereby producing a CTP-modified coagulation factor as defined in any one of claims 1-8, in which the coagulation factor has an extended biological half-life, an improved area under the curve (AUC), a reduced dosing frequency, or a reduced clearance rate.
15. A method of producing the CTP-modified coagulation factor of any one of claims 1-8, the method comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of a FIX polypeptide, or comprising the step of attaching three chorionic gonadotrophin carboxy terminal peptides (CTPs) to the carboxy terminus of a FVII or a FVIIa polypeptide, thereby producing the CTP-modified coagulation factor.
16. A polynucleotide encoding the CTP-modified coagulation factor according to any one of claims 1-8.

17. The polynucleotide of claim 16, wherein the nucleic acid sequence of said polynucleotide is as set forth in SEQ ID NO: 30.

18. The polynucleotide of claim 16, wherein the nucleic acid sequence of said polynucleotide is as set forth in SEQ ID NO: 24.