

AQUEOUS ANTI-PD-L1 ANTIBODY FORMULATION

The present invention relates to a novel anti-PD-L1 antibody formulation. In particular, the invention relates to an aqueous pharmaceutical formulation of the anti-PD-L1
5 antibody Avelumab.

Background of the invention

The programmed death 1 (PD-1) receptor and PD-1 ligands 1 and 2 (PD-L1, PD-L2)
10 play integral roles in immune regulation. Expressed on activated T cells, PD-1 is activated by PD-L1 and PD-L2 expressed by stromal cells, tumor cells, or both, initiating T-cell death and localized immune suppression (Dong H, Zhu G, Tamada K, Chen L. B7-H1, a third member of the B7 family, co-stimulates T-cell proliferation and interleukin-10 secretion. Nat Med 1999;5:1365-69; Freeman GJ, Long AJ, Iwai Y, et al.
15 Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. J Exp Med 2000;192:1027-34; Dong H, Strome SE, Salomao DR, et al. Tumor-associated B7-H1 promotes T-cell apoptosis: a potential mechanism of immune evasion. Nat Med 2002; 8:793-800. Erratum, Nat Med 2002;8:1039; Topalian SL, Drake CG, Pardoll DM.
20 Targeting the PD-1/B7-H1(PD-L1) pathway to activate anti-tumor immunity. Curr Opin Immunol 2012;24:207-12), potentially providing an immune-tolerant environment for tumor development and growth. Conversely, inhibition of this interaction can enhance local T-cell responses and mediate antitumor activity in nonclinical animal models (Dong H, Strome SE, Salomao DR, et al. Nat Med 2002; 8:793-800. Erratum, Nat Med
25 2002;8:1039; Iwai Y, Ishida M, Tanaka Y, et al. Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. Proc Natl Acad Sci USA 2002;99:12293-97). In the clinical setting, treatment with antibodies that block the PD-1 – PD-L1 interaction have been reported to produce objective response rates of 7% to 38% in patients with advanced or metastatic solid
30 tumors, with tolerable safety profiles (Hamid O, Robert C, Daud A, et al. Safety and tumor responses with lambrolizumab (Anti-PD-1) in melanoma. N Engl J Med 2013;369:134-44; Brahmer JR, Tykodi SS, Chow LQ, et al. Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. N Engl J Med 2012;366(26):2455-65;

Topalian SL, Hodi FS, Brahmer JR, et al. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. N Engl J Med 2012;366(26):2443-54; Herbst RS, Soria J-C, Kowanetz M, et al. Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. Nature 2014;515:563-67). Notably, responses
5 appeared prolonged, with durations of 1 year or more for the majority of patients.

Avelumab (also known as MSB0010718C) is a fully human monoclonal antibody of the immunoglobulin (Ig) G1 isotype. Avelumab selectively binds to PD-L1 and competitively blocks its interaction with PD-1.

10

Compared with anti-PD-1 antibodies that target T-cells, Avelumab targets tumor cells, and therefore is expected to have fewer side effects, including a lower risk of autoimmune-related safety issues, as blockade of PD-L1 leaves the PD-L2 – PD-1 pathway intact to promote peripheral self-tolerance (Latchman Y, Wood CR, Chernova
15 T, et al. PD-L1 is a second ligand for PD-1 and inhibits T cell activation. Nat Immunol 2001;2(3):261-68).

Avelumab is currently being tested in the clinic in a number of cancer types including non-small cell lung cancer, urothelial carcinoma, mesothelioma, Merkel cell carcinoma,
20 gastric or gastroesophageal junction cancer, ovarian cancer, and breast cancer.

The amino acid sequences of Avelumab and sequence variants and antigen binding fragments thereof, are disclosed in WO2013079174, where the antibody having the amino acid sequence of Avelumab is referred to as A09-246-2. Also disclosed are
25 methods of manufacturing and certain medical uses.

Further medical uses of Avelumab are described in WO2016137985, WO2016181348, WO2016205277, PCT/US2016/053939, U.S. patent application Ser. No. 62/423,358. WO2013079174 also describes in section 2.4 a human aqueous formulation of an antibody having the amino acid sequence of Avelumab. This formulation comprises the
30 antibody in a concentration of 10 mg/ml, methionine as an antioxidant and has a pH of 5.5. Avelumab formulations not comprising an antioxidant are described in PCT/EP2016/002040.

A formulation study for an aglycosylated anti-PD-L1 antibody of the IgG1 type is described in WO2015048520, where a formulation with a pH of 5.8 was selected for clinical studies.

5 Description of the invention

As Avelumab is generally delivered to a patient via intravenous infusion, and is thus provided in an aqueous form, the present invention relates to further aqueous formulations that are suitable to stabilize Avelumab with its post-translational
10 modifications, and at higher concentrations as disclosed in WO2013079174.

Figure 1a (SEQ ID NO:1) shows the full length heavy chain sequence of Avelumab, as expressed by the CHO cells used as the host organism.

It is frequently observed, however, that in the course of antibody production the C-
15 terminal lysine (K) of the heavy chain is cleaved off. Located in the Fc part, this modification has no influence on the antibody – antigen binding. Therefore, in some embodiments the C-terminal lysine (K) of the heavy chain sequence of Avelumab is absent. The heavy chain sequence of Avelumab without the C-terminal lysine is shown in Figure 1b (SEQ ID NO:2).

20 Figure 2 (SEQ ID NO:3) shows the full length light chain sequence of Avelumab.

A post-translational modification of high relevance is glycosylation.

Most of the soluble and membrane-bound proteins that are made in the endoplasmatic reticulum of eukaryotic cells undergo glycosylation, where enzymes called
25 glycosyltransferases attach one or more sugar units to specific glycosylation sites of the proteins. Most frequently, the points of attachment are NH₂ or OH groups, leading to N-linked or O-linked glycosylation.

This applies also to proteins, such as antibodies, which are recombinantly produced in eukaryotic host cells. Recombinant IgG antibodies contain a conserved N-linked
30 glycosylation site at a certain asparagine residue of the Fc region in the CH2 domain. There are many known physical functions of N-linked glycosylation in an antibody such as affecting its solubility and stability, protease resistance, binding to Fc receptors, cellular transport and circulatory half-life in vivo (Hamm M. et al.,

Pharmaceuticals 2013, 6, 393-406). IgG antibody N-glycan structures are predominantly biantennary complex-type structures, comprising b-D-N-acetylglucosamine (GlcNAc), mannose (Man) and frequently galactose (Gal) and fucose (Fuc) units.

- 5 In Avelumab the single glycosylation site is Asn300, located in the CH2 domain of both heavy chains. Details of the glycosylation are described in Example 1.

Since glycosylation affects the solubility and stability of an antibody, it is prudent to take this parameter into account when a stable, pharmaceutically suitable formulation of the
10 antibody is to be developed.

Surprisingly, it has been found by the inventors of the present patent application that it is possible to stabilize Avelumab, fully characterized by its amino acid sequence and its post-translational modifications, in a number of aqueous formulations without the
15 presence of an antioxidant, at pH values even below 5.2.

Figures

- Figure 1a: Heavy chain sequence of Avelumab (SEQ ID NO:1)
20 Figure 1b: Heavy chain sequence of Avelumab, lacking the C-terminal K (SEQ ID NO:2)
Figure 2: Light chain sequence of Avelumab (SEQ ID NO:3)
Figure 3: Secondary structure of Avelumab
Figure 4: 2AB HILIC-UPLC Chromatogram of Avelumab Glycans
25 Figure 5: Numbering of the peaks of Figure 4

Definitions

Unless otherwise stated, the following terms used in the specification and claims have
30 the following meanings set out below.

References herein to "Avelumab" include the anti-PD-L1 antibody of the IgG1 type as defined in WO2013079174 by its amino acid sequence, and as defined in the present patent application by its amino acid sequence and by its post-translational

modifications. References herein to “Avelumab” may include biosimilars which, for instance, may share at least 75%, suitably at least 80%, suitably at least 85%, suitably at least 90%, suitably at least 95%, suitably at least 96%, suitably at least 97%, suitably at least 98% or most suitably at least 99% amino acid sequence identity with the amino acid sequences disclosed in WO2013079174. Alternatively or additionally, references
5 herein to “Avelumab” may include biosimilars which differ in the post-translational modifications, especially in the glycosylation pattern, herein disclosed.

The term “biosimilar” (also known as follow-on biologics) is well known in the art, and
10 the skilled person would readily appreciate when a drug substance would be considered a biosimilar of Avelumab. The term “biosimilar” is generally used to describe subsequent versions (generally from a different source) of “innovator biopharmaceutical products” (“biologics” whose drug substance is made by a living organism or derived from a living organism or through recombinant DNA or controlled gene expression
15 methodologies) that have been previously officially granted marketing authorisation. Since biologics have a high degree of molecular complexity, and are generally sensitive to changes in manufacturing processes (e.g. if different cell lines are used in their production), and since subsequent follow-on manufacturers generally do not have access to the originator’s molecular clone, cell bank, know-how regarding the
20 fermentation and purification process, nor to the active drug substance itself (only the innovator’s commercialized drug product), any “biosimilar” is unlikely to be exactly the same as the innovator drug product.

Herein, the term “buffer” or “buffer solution” refers to a generally aqueous solution comprising a mixture of an acid (usually a weak acid, e.g. acetic acid, citric acid,
25 imidazolium form of histidine) and its conjugate base (e.g. an acetate or citrate salt, for example, sodium acetate, sodium citrate, or histidine) or alternatively a mixture of a base (usually a weak base, e.g. histidine) and its conjugate acid (e.g. protonated histidine salt). The pH of a “buffer solution” will change very only slightly upon addition of a small quantity of strong acid or base due to the “buffering effect” imparted by the
30 “buffering agent”.

Herein, a “buffer system” comprises one or more buffering agent(s) and/or an acid/base conjugate(s) thereof, and more suitably comprises one or more buffering agent(s) and an acid/base conjugate(s) thereof, and most suitably comprises one buffering agent

only and an acid/base conjugate thereof. Unless stated otherwise, any concentrations stipulated herein in relation to a “buffer system” (i.e. a buffer concentration) suitably refers to the combined concentration of the buffering agent(s) and/or acid/base conjugate(s) thereof. In other words, concentrations stipulated herein in relation to a
5 “buffer system” suitably refer to the combined concentration of all the relevant buffering species (i.e. the species in dynamic equilibrium with one another, e.g. citrate/citric acid). As such, a given concentration of a histidine buffer system generally relates to the combined concentration of histidine and the imidazolium form of histidine. However, in the case of histidine, such concentrations are usually straightforward to calculate by
10 reference to the input quantities of histidine or a salt thereof. The overall pH of the composition comprising the relevant buffer system is generally a reflection of the equilibrium concentration of each of the relevant buffering species (i.e. the balance of buffering agent(s) to acid/base conjugate(s) thereof).

15 Herein, the term “buffering agent” refers to an acid or base component (usually a weak acid or weak base) of a buffer or buffer solution. A buffering agent helps maintain the pH of a given solution at or near to a pre-determined value, and the buffering agents are generally chosen to complement the pre-determined value. A buffering agent is suitably a single compound which gives rise to a desired buffering effect, especially when said
20 buffering agent is mixed with (and suitably capable of proton exchange with) an appropriate amount (depending on the pre-determined pH desired) of its corresponding “acid/base conjugate”, or if the required amount of its corresponding “acid/base conjugate” is formed *in situ* – this may be achieved by adding strong acid or base until the required pH is reached. For example in the sodium acetate buffer system, it is
25 possible to start out with a solution of sodium acetate (basic) which is then acidified with, e.g., hydrochloric acid, or to a solution of acetic acid (acidic), sodium hydroxide or sodium acetate is added until the desired pH is reached.

Generally, a “stabiliser” refers to a component which facilitates maintenance of the
30 structural integrity of the biopharmaceutical drug, particularly during freezing and/or lyophilization and/or storage (especially when exposed to stress). This stabilising effect may arise for a variety of reasons, though typically such stabilisers may act as osmolytes which mitigate against protein denaturation. As used herein, stabilisers can be sugar alcohols (e.g. inositol, sorbitol), disaccharides (e.g. sucrose, maltose),

monosaccharides (e.g. dextrose (D- glucose)), or forms of the amino acid lysine (e.g. lysine monohydrochloride, acetate or monohydrate), or salts (e.g. sodium chloride).

Agents used as buffering agents, antioxidants or surfactants according to the invention,
5 are excluded from the meaning of the term “stabilisers” as used herein, even if they may exhibit, i.a. stabilising activity.

Herein, the term "surfactant" refers to a surface-active agent, preferably a nonionic surfactant. Examples of surfactants used herein include polysorbate, for example,
10 polysorbate 80 (polyoxyethylene (80) sorbitan monooleate, also known under the tradename Tween 80); polyoxyl castor oil, such as polyoxyl 35 castor oil, made by reacting castor oil with ethylene oxide in a molar ratio of 1 : 35, also known under the tradename Kolliphor ELP; or Kollidon 12PF or 17PF, which are low molecular weight povidones (polyvinylpyrrolidones), known under the CAS number 9003-39-8 and having
15 slightly different molecular weights (12PF: 2000-3000 g/mol, 17PF: 7000-11000 g/mol).

Agents used as buffering agents, antioxidants or stabilisers according to the invention, are excluded from the meaning of the term “surfactants” as used herein, even if they may exhibit, i.a. surfactant activity.
20

Herein, the term “stable” generally refers to the physical stability and/or chemical stability and/or biological stability of a component, typically an active or composition thereof, during preservation/storage.

25 Herein, the term “antioxidant” refers to an agent capable of preventing or decreasing oxidation of the biopharmaceutical drug to be stabilized in the formulation.

Antioxidants include radical scavengers (e.g. ascorbic acid, BHT, sodium sulfite, p-amino benzoic acid, glutathione or propyl gallate), chelating agents (e.g. EDTA or citric acid) or chain terminators (e.g. methionine or N-acetyl cysteine).

30 Agents used as buffering agents, stabilisers or surfactants according to the invention, are excluded from the meaning of the term “antioxidants” as used herein, even if they may exhibit, i.a. antioxidative activity.

A "diluent" is an agent that constitutes the balance of ingredients in any liquid pharmaceutical composition, for instance so that the weight percentages total 100%. Herein, the liquid pharmaceutical composition is an aqueous pharmaceutical composition, so that a "diluent" as used herein is water, preferably water for injection (WFI).

Herein, the term "particle size" or "pore size" refers respectively to the length of the longest dimension of a given particle or pore. Both sizes may be measured using a laser particle size analyser and/or electron microscopes (e.g. tunneling electron microscope, TEM, or scanning electron microscope, SEM). The particle count (for any given size) can be obtained using the protocols and equipment outlined in the Examples, which relates to the particle count of sub-visible particles.

Herein, the term "about" refers to the usual error range for the respective value readily known to the skilled person in this technical field. Reference to "about" a value or parameter herein includes (and describes) embodiments that are directed to that value or parameter per se. In case of doubt, or should there be no art recognized common understanding regarding the error range for a certain value or parameter, "about" means $\pm 5\%$ of this value or parameter.

Herein, the term "percent share" in connection with glycan species refers directly to the number of different species. For example the term "said FA2G1 has a share of 25% - 41% of all glycan species" means that in 50 antibody molecules analysed, having 100 heavy chains, 25-41 of the heavy chains will exhibit the FA2G1 glycosylation pattern.

It is to be appreciated that references to "treating" or "treatment" include prophylaxis as well as the alleviation of established symptoms of a condition. "Treating" or "treatment" of a state, disorder or condition therefore includes: (1) preventing or delaying the appearance of clinical symptoms of the state, disorder or condition developing in a human that may be afflicted with or predisposed to the state, disorder or condition but does not yet experience or display clinical or subclinical symptoms of the state, disorder or condition, (2) inhibiting the state, disorder or condition, i.e., arresting, reducing or delaying the development of the disease or a relapse thereof (in case of maintenance treatment) or at least one clinical or subclinical symptom thereof, or (3) relieving or

attenuating the disease, i.e., causing regression of the state, disorder or condition or at least one of its clinical or subclinical symptoms.

Aqueous anti-PD-L1 Antibody Formulation

5

In a first aspect, the invention provides a novel aqueous pharmaceutical antibody formulation, comprising:

- (i) Avelumab in a concentration of 1 mg/mL to 30 mg/mL as the antibody;
 - (ii) glycine, succinate, citrate phosphate or histidine in a concentration of 5 mM to 35 mM as the buffering agent;
 - (iii) lysine monohydrochloride, lysine monohydrate, lysine acetate, dextrose, sucrose, sorbitol or inositol in a concentration of 100 mM to 320 mM as the stabiliser;
 - (iv) povidone, polyoxyl castor oil or polysorbate in a concentration of 0.25 mg/mL to 0.75 mg/mL, as the surfactant;
- wherein the formulation does not comprise methionine, and further wherein the formulation has a pH of 3.8 to 5.2.

In a preferred embodiment the formulation does not comprise any antioxidant.

- In an embodiment the concentration of Avelumab in the said formulation is about 10 mg/mL to about 20 mg/mL.

In yet another embodiment the concentration of glycine, succinate, citrate phosphate or histidine in the said formulation is about 10 mM to about 20 mM.

- In further embodiments, in the said formulation, the concentration of lysine monochloride is about 140 mM to about 280 mM, or the concentration of said lysine monohydrate is about 280 mM, or the concentration of the said lysine acetate is about 140 mM.

In yet another embodiment the concentration of dextrose, sucrose, sorbitol or inositol in the said formulation is about 280 mM.

- In yet another embodiment the concentration of povidone, polyoxyl castor oil or polysorbate inositol in the said formulation is about 0.5 mg/mL.

In a preferred embodiment the said povidone in the said formulation is the low molecular weight polyvinylpyrrolidone Kollidon 12PF or 17PF of CAS number 9003-39-8.

In another preferred embodiment the said polyoxyl castor oil is Polyoxyl 35 Castor Oil.
In yet another preferred embodiment the said polysorbate is Polysorbate 80.

In a more preferred embodiment, the novel aqueous pharmaceutical antibody

5 formulation, comprises:

- (i) Avelumab in a concentration of 1 mg/mL to about 20 mg/mL as the antibody;
 - (ii) glycine in a concentration of 5 mM to 15 mM as the buffering agent, and not comprising any other buffering agent;
 - (iii) lysine monohydrochloride, dextrose, sucrose or sorbitol in a concentration of 100
10 mM to 320 mM as the stabiliser, and not comprising any other stabiliser;
 - (iv) Kollidon 12PF, polyoxyl 35 castor oil or Polysorbate 80 in a concentration of 0.25 mg/mL to 0.75 mg/mL, as the surfactant, and not comprising any other surfactant;
- wherein the formulation has a pH of 3.8 to 4.6, and does not comprise an antioxidant.

15 In an equally preferred embodiment, the novel aqueous pharmaceutical antibody formulation, comprises:

- (i) Avelumab in a concentration of 1 mg/mL to about 20 mg/mL as the antibody;
 - (ii) succinate in a concentration of 5 mM to 15 mM as the buffering agent, and not comprising any other buffering agent;
 - (iii) lysine monohydrochloride, dextrose, sucrose or sorbitol in a concentration of 100
20 mM to 320 mM as the stabiliser, and not comprising any other stabiliser;
 - (iv) Kollidon 12PF or polyoxyl 35 castor oil in a concentration of 0.25 mg/mL to 0.75 mg/mL, as the surfactant, and not comprising any other surfactant;
- wherein the formulation has a pH of 4.9 to 5.2, and does not comprise an antioxidant.

25

In an equally preferred embodiment, the novel aqueous pharmaceutical antibody formulation, comprises:

- (i) Avelumab in a concentration of 1 mg/mL to about 20 mg/mL as the antibody;
- (ii) citrate phosphate in a concentration of 10 mM to 20 mM as the buffering agent, and
30 not comprising any other buffering agent;
- (iii) lysine monohydrochloride, dextrose, sucrose or sorbitol in a concentration of 100 mM to 320 mM as the stabiliser, and not comprising any other stabiliser;
- (iv) Kollidon 12PF or polyoxyl 35 castor oil in a concentration of 0.25 mg/mL to 0.75 mg/mL, as the surfactant, and not comprising any other surfactant;

wherein the formulation has a pH of 3.8 to 4.7, and does not comprise an antioxidant.

In an equally preferred embodiment, the novel aqueous pharmaceutical antibody formulation, comprises:

- 5 (i) Avelumab in a concentration of 1 mg/mL to about 20 mg/mL as the antibody;
- (ii) glycine in a concentration of about 10 mM as the buffering agent, and not comprising any other buffering agent;
- (iii) lysine monohydrochloride in a concentration of about 140 mM as the stabiliser, and not comprising any other stabiliser;
- 10 (iv) polyoxyl 35 castor oil in a concentration of about 0.5 mg/mL as the surfactant, and not comprising any other surfactant;

wherein the formulation has a pH of 4.2 to 4.6, and does not comprise an antioxidant.

- In a more preferred embodiment, the novel aqueous pharmaceutical antibody formulation, comprises:

- 15 (i) Avelumab in a concentration of 1 mg/mL to about 20 mg/mL as the antibody;
- (ii) glycine in a concentration of about 10 mM as the buffering agent, and not comprising any other buffering agent;
- (iii) lysine acetate in a concentration of about 140 mM as the stabiliser, and not
- 20 comprising any other stabiliser;
- (iv) polyoxyl 35 castor oil in a concentration of about 0.5 mg/mL as the surfactant, and not comprising any other surfactant;

wherein the formulation has a pH of 4.2 to 4.6, and does not comprise an antioxidant.

- 25 In an equally preferred embodiment, the novel aqueous pharmaceutical antibody formulation, comprises:

- (i) Avelumab in a concentration of 1 mg/mL to about 20 mg/mL as the antibody;
- (ii) histidine in a concentration of about 10 mM as the buffering agent, and not comprising any other buffering agent;
- 30 (iii) sucrose in a concentration of about 280 mM as the stabiliser, and not comprising any other stabiliser;
- (iv) Kollidon 12PF in a concentration of about 0.5 mg/mL as the surfactant, and not comprising any other surfactant;

wherein the formulation has a pH of 4.8 to 5.2, and does not comprise an antioxidant.

In an equally preferred embodiment, the novel aqueous pharmaceutical antibody formulation, comprises:

- (i) Avelumab in a concentration of 1 mg/mL to about 20 mg/mL as the antibody;
 - 5 (ii) succinate in a concentration of about 10 mM as the buffering agent, and not comprising any other buffering agent;
 - (iii) lysine monohydrochloride in a concentration of about 140 mM as the stabiliser, and not comprising any other stabiliser;
 - (iv) polyoxyl 35 castor oil in a concentration of about 0.5 mg/mL as the surfactant, and
 - 10 not comprising any other surfactant;
- wherein the formulation has a pH of 4.8 to 5.2, and does not comprise an antioxidant.

In a more preferred embodiment of the above described embodiments, the concentration of Avelumab is about 20 mg/ml.

15

In an even more preferred embodiments the said formulation consists of:

- (i) Avelumab in a concentration of 20 mg/mL;
 - (ii) glycine in a concentration of 10 mM;
 - 20 (iii) lysine monohydrochloride in a concentration of 140 mM;
 - (iv) polyoxyl 35 castor oil in a concentration of 0.5 mg/mL;
 - (v) HCl or NaOH to adjust the pH;
 - (vi) water (for injection) as the solvent;
- and has a pH of 4.4 (± 0.1);

25

or

- (i) Avelumab in a concentration of 20 mg/mL;
 - (ii) glycine in a concentration of 10 mM;
 - 30 (iii) lysine acetate in a concentration of 140 mM;
 - (iv) polyoxyl 35 castor oil in a concentration of 0.5 mg/mL;
 - (v) HCl or NaOH to adjust the pH;
 - (vi) water (for injection) as the solvent;
- and has a pH of 4.4 (± 0.1);

or

- (i) Avelumab in a concentration of 20 mg/mL;
- 5 (ii) histidine in a concentration of 10 mM;
- (iii) sucrose in a concentration of 280 mM;
- (iv) Kollidon 12PF in a concentration of 0.5 mg/mL;
- (v) HCl or NaOH to adjust the pH;
- (vi) water (for injection) as the solvent;
- 10 and has a pH of 5.0 ($\pm$ 0.1);

or

- (i) Avelumab in a concentration of 20 mg/mL;
- 15 (ii) succinate in a concentration of 10 mM;
- (iii) lysine monohydrochloride in a concentration of 140 mM;
- (iv) polyoxyl 35 castor oil in a concentration of 0.5 mg/mL;
- (v) HCl or NaOH to adjust the pH;
- (vi) water (for injection) as the solvent;
- 20 and has a pH of 5.0 ($\pm$ 0.1).

In another preferred embodiment, the formulation has an osmolality between 270 and 330 mOsm/kg.

- 25 In an embodiment said Avelumab in the formulations as described above has the heavy chain sequence of either Fig. 1a (SEQ ID NO:1) or Fig. 1b (SEQ ID NO:2), the light chain sequence of Fig. 2 (SEQ ID NO:3), and carries a glycosylation on Asn300 comprising FA2 and FA2G1 as the main glycan species, having a joint share of > 70% of all glycan species.

30

In a preferred embodiment, in the Avelumab glycosylation the said FA2 has a share of 44% - 54% and said FA2G1 has a share of 25% - 41% of all glycan species.

In a preferred embodiment, in the Avelumab glycosylation the said FA2 has a share of 47% - 52% and said FA2G1 has a share of 29% - 37% of all glycan species.

In a preferred embodiment, in the Avelumab glycosylation the said FA2 has a share of about 49% and said FA2G1 has a share of about 30% - about 35% of all glycan species.

- 5 In a preferred embodiment the Avelumab glycosylation further comprises as minor glycan species A2 with a share of < 5%, A2G1 with a share of < 5%, A2G2 with a share of < 5% and FA2G2 with a share of < 7% of all glycan species.

- 10 In a preferred embodiment, in the Avelumab glycosylation said A2 has a share of 3%-5%, said A2G1 has a share of < 4%, said A2G2 has a share of < 3% and said FA2G2 has a share of 5%-6% of all glycan species.

- 15 In a preferred embodiment, in the Avelumab glycosylation said A2 has a share of about 3.5% - about 4.5%, said A2G1 has a share of about 0.5% - about 3.5%, said A2G2 has a share of < 2.5% and said FA2G2 has a share of about 5.5% of all glycan species.

In an embodiment the said Avelumab in the formulation as described above has the heavy chain sequence of Fig. 1b (SEQ ID NO:2).

- 20 In an embodiment the Avelumab formulation as described above is for intravenous (IV) administration.

Drug-delivery Device

- 25 In a second aspect the present invention provides a drug delivery device comprising a liquid pharmaceutical composition as defined herein. Suitably the drug delivery device comprises a chamber within which the pharmaceutical composition resides. Suitably the drug delivery device is sterile.

- 30 The drug delivery device may a vial, ampoule, syringe, injection pen (e.g. essentially incorporating a syringe), or i.v. (intravenous) bag.

The aqueous pharmaceutical formulations are parenterally administered, preferably via sub-cutaneous injection, intramuscular injection, i.v. injection or i.v. infusion. The most preferred way of administration is i.v. infusion.

In a preferred embodiment, the drug delivery device is a vial containing the formulation as described above.

In a more preferred embodiment the said vial contains 200 mg avelumab in 10 mL of solution for a concentration of 20 mg/mL.

- 5 In an even more preferred embodiment the vial is a glass vial.

Medical Treatment

In a third aspect, the invention provides a method of treating cancer comprising

- 10 administering the formulation as described above to a patient.

In an embodiment the cancer to be treated is selected from non-small cell lung cancer, urothelial carcinoma, bladder cancer, mesothelioma, Merkel cell carcinoma, gastric or gastroesophageal junction cancer, ovarian cancer, breast cancer, thymoma, adenocarcinoma of the stomach, adrenocortical carcinoma, head and neck squamous
15 cell carcinoma, renal cell carcinoma, melanoma, and/or classical Hodgkin's lymphoma.

Methods of manufacturing

The present invention also provides a method of manufacturing an aqueous
20 pharmaceutical formulation as defined herein. The method suitably comprises mixing together, in any particular order deemed appropriate, any relevant components required to form the aqueous pharmaceutical formulation. The skilled person may refer to the examples or techniques well known in the art for forming aqueous pharmaceutical formulations (especially those for injection via syringe, or i.v. infusion).

25

The method may involve first preparing a pre-mixture (or pre-solution) of some or all components (optionally with some or all of the diluent) excluding Avelumab, and Avelumab may then itself (optionally with or pre-dissolved in some of the diluent) be mixed with the pre-mixture (or pre-solution) to afford the aqueous pharmaceutical
30 formulation, or a composition to which final components are then added to furnish the final aqueous pharmaceutical formulation. Preferably, the method involves forming a buffer system, suitably a buffer system comprising a buffering agent as defined herein. The buffer system is suitably formed in a pre-mixture prior to the addition of Avelumab. The buffer system may be formed through simply mixing the buffering agent (supplied

ready-made) with its acid/base conjugate (suitably in appropriate relative quantities to provide the desired pH – this can be determined by the skilled person either theoretically or experimentally). In the case of an acetate buffer system, this means e.g. mixing sodium acetate with HCl, or mixing acetic acid with NaOH or acetate. The pH of
5 either the pre-mixture of final aqueous pharmaceutical formulation may be judiciously adjusted by adding the required quantity of base or acid, or a quantity of buffering agent or acid/base conjugate.

In certain embodiments, the buffering agent and/or buffer system is pre-formed as a
10 separate mixture, and the buffer system is transferred to a precursor of the aqueous pharmaceutical formulation (comprising some or all components save for the buffering agent and/or buffer system, suitably comprising Avelumab and potentially only Avelumab) via buffer exchange (e.g. using diafiltration until the relevant concentrations or osmolality is reached). Additional excipients may be added thereafter if necessary in
15 order to produce the final liquid pharmaceutical composition. The pH may be adjusted once or before all the components are present.

Any, some, or all components may be pre-dissolved or pre-mixed with a diluent prior to mixing with other components.
20 The final aqueous pharmaceutical formulation may be filtered, suitably to remove particulate matter. Suitably filtration is through filters sized at or below 1 μm , suitably at 0.22 μm . Suitably, filtration is through either PES filters or PVDF filters, suitably with 0.22 μm PES filters.

25 The person of skill in the art is well aware how an aqueous pharmaceutical formulation can be used to prepare an IV solution, so that the antibody drug substance can be administered intravenously.

The preparation of the IV solution typically consists of a certain amount of solution being withdrawn from saline bags (e.g. 0.9% or 0.45% saline) with a plastic syringe (PP) and a
30 needle and replaced with aqueous pharmaceutical formulation. The amount of solution replaced will depend on the body weight of the patients.

Abbreviations

	ANOVA	Analysis of variance
	CD	Circular dichroism
5	CE – SDS	Capillary electrophoresis sodium dodecyl sulfate
	cIEF	Capillary isoelectrofocusing
	DoE	Design of Experiment
	DP	Drug Product
	DS	Drug Substance
10	FT	Freeze - thawing
	HMW	Higher Molecular Weight
	LMW	Low Molecular Weight
	SE-HPLC	Size Exclusion High Performance Liquid Chromatography
	OD	Optical Density
15	PES	Polyethersulphone
	PVDF	Polyvinylidene fluoride
	RH	Relative Humidity
	SE – HPLC	Size – exclusion high performance chromatography
	UV	Ultraviolet
20	WFI	Water for Injection

Examples25 Example 1 – Structure of Avelumab

1.1 Primary Structure

30 Avelumab is an IgG with two heavy and two light chain molecules. The amino acid sequences of the two chains are shown in Figures 1a (SEQ ID NO:1) / 1b (SEQ ID NO:2) and 2 (SEQ ID NO:3), respectively.

1.2 Secondary Structure

LC-MS and MS/MS methods were used to confirm the intact chains of the molecule and the presence of post-translational modifications to the proteins. The secondary structure of the Avelumab molecule subunits are shown in Figure 3.

As confirmed by UPLC-Q-TOF mass spectrometry of peptides obtained by trypsin digestion, the disulfide bonds Cys21-Cys96, Cys21-Cys90, Cys147-Cys203, Cys138-Cys197, Cys215-Cys223, Cys229-Cys229, Cys232-Cys232, Cys264-Cys324 and Cys370-Cys428 are forming the nine typical IgG bonding pattern.



1.3 Glycosylation











The molecule contains one N-glycosylation site on Asn300 of the heavy chain. As determined by peptide mapping, the main structure identified by MALDI-TOF was a complex, biantennary type core fucosylated oligosaccharide with zero (G0F), one (G1F), or two galactose (G2F) residues. The main species are G0F and G1F.

Avelumab glycans fluorescence labeled by 2-aminobenzamide have been analysed by HILIC-UPLC-ESI-Q-TOF. Figure 4 shows the UPLC profile of the glycan species found.

20

Table 1: Peak identification of 2AB HILIC-UPLC chromatogram

Peak	RT (min)	Measured MW	Expected MW	Identification	Oxford nomenclature	Identification by
1a	5.99	1380.52 (M+H)	1380.54 (M+H)		FA1	Manually identified by MS
2	6.01	1437.54	1437.56		A2	Manually identified by MS
3	7.02	1583.74 (M+H)	1583.62 (M+H)		FA2	MS in source fragmentation by GlycoworkBench

5	4	7.77	1355.57 (M+H)	1355.51 (M+H)		M5	Manually identified by MS
	5	8.16	1599.77 (M+H)	1599.62 (M+H)		A2G1	Manually identified by MS
	6	9.82	1744.79	1744.67		FA2G1	MS in source fragmentation by GlycoworkBench
			1462.90	1462.54		FA2 freeEnd	GlycoworkBench identified by MS
	7	10.07	1744.80	1744.67		FA2G1	MS in source fragmentation by GlycoworkBench
			1462.91	1462.54		FA2 freeEnd	GlycoworkBench identified by MS
	8	10.44	1462.90	1462.54		FA2 freeEnd	GlycoworkBench identified by MS
			1744.79	1744.67		FA2G1	Manually identified by MS
	9	12.15	1177.50 (M+H)	1177.46 (M+H)		FM3	GlycoworkBench identified by MS
	10	16.66	No ionization	No ionization			
10	11	13.42	1906.33	1906.72		FA2G2	MS in source fragmentation by GlycoworkBench
			1624.71	1624.59		FA2G1 freeEnd	GlycoworkBench identified by MS
	12	13.71	954.40 (M+2H)/2	954.36 (M+2H)/2		FA2G2	Manually identified by MS
			1626.69	1626.61		FA2G1 redEnd	GlycoworkBench identified by MS
	13	17.46	1099.97 (M+2H)/2	1099.91 (M+2H)/2		FA2G2S	MS in source fragmentation by GlycoworkBench
	14	18.54	1079.91 (M+2H)/2	1079.86 (M+2H)/2		FA2G2S freeEnd+S (probable-small traces)	Manually identified by MS
15	15	21.04	2489.05	2488.91		FA2G2S2	Manually identified by MS

The geometric shapes representing the glycan building blocks correspond to the following molecular entities:

● Man △ Fuc ○ Gal □ GalNAc ▨ GlcNAc ◇ NANA

Man: mannose, Fuc: fucose, Gal: galactose, GalNAc: N-Acetylgalactosamine, NANA: sialic acid

5

The glycan nomenclature used follows the Oxford Notation as proposed by Harvey et al. (Proteomics 2009, 9, 3796-3801). In species containing fucose (FA2, FA2G1, FA2G2), the Fuc-GlcNAc connectivity is α 1-6. In species having a terminal GlcNAc, the GlcNAc-Man connectivity is β 1-2. In species containing galactose, the Gal-GlcNAc connectivity is β 1-4.

10

The reported chromatographic profile has been integrated and yielded the Glycan Species Distribution of Avelumab as shown in Table 2a.

15 Table 2a

A2	FA2	A2G1	FA2G1	A2G2	FA2G2	M5**
3.6	48.7	3.4	35.6	2.3	5.4	1.0

** Probably Mannose 5, coelution with biantennary mono-galactosylated species

The glycan mapping analysis confirmed the identification carried out by peptide mapping (that allowed to identify the two main glycan species), in addition secondary and minor species were also characterized by this method, specific for glycan analysis.

20

In another measurement the following Glycan Species Distribution was observed.

25 Table 2b:

A2	FA2	A2G1	FA2G1	A2G2	FA2G2
4.0	50.2	1.0	30.0	0.1	5.6

30

Example 2 – DoE screening

A Design of Experiment screening at 20 mg/mL Avelumab assessed the impact of several factors such as varying buffer type/pH, stabilisers, surfactant type and relevant concentration. The study, testing 80 different formulations, led to the selection of the suitable conditions that can maximize protein stability.

Four different buffers were examined in this DoE covering different buffer types and effective pH buffering range:

10 Amino acid buffers such as Glycine (effective pH 4.0 to 7.5) and Histidine (effective pH 5.0 to 6.6).

Chelating ionic buffer such as Citrate (effective pH 4.0 to 7.5).

Succinate (effective pH 5.0 to 6.0).

15 Seven stabilisers were selected in the DoE on the basis of their chemical structure.

Included in the DoE were sugars, polyols, salts, and amino acids. The breakdown is as follows:

Sugars: The disaccharides Sucrose and Maltose were selected as well as the monosaccharide Dextrose (D-Glucose).

20 Sugar alcohols: Two sugar alcohols / polyols were selected for the DoE - Sorbitol and Inositol.

Salt: Sodium chloride was investigated as a stand-alone stabiliser in this DoE.

Amino acid: Lysine, a positively charged amino acid was investigated.

25 Table 3 lists the samples and their respective compositions.

Table 3: DoE screening formulations

Sample ID	pH	Buffer	Buffer Strength (mM)	Stabiliser (280 mM)	Surfactant (0.5 mg/mL)
1	4	Citrate-phosphate	10	Sorbitol	Kollidon 12PF
2	4	Citrate-phosphate	50	Dextrose	Tween 40
3	4.5	Citrate-phosphate	20	Dextrose	Tween 40
4	4.5	Citrate-phosphate	30	Inositol	Kollidon 12PF
5	4.8	Citrate-phosphate	40	Maltose	Kolliphor ELP

6	4.8	Citrate-phosphate	40	Lysine	Tween 80
7	5.2	Citrate-phosphate	50	Dextrose	Kolliphor ELP
8	5.2	Citrate-phosphate	10	Sodium chloride	Tween 80
9	5.2	Citrate-phosphate	20	Lysine	Kolliphor ELP
10	5.5	Citrate-phosphate	20	Sucrose	Kollidon 12PF
11	5.5	Citrate-phosphate	30	Lysine	Tween 40
12	6	Citrate-phosphate	20	Maltose	Tween 40
13	6	Citrate-phosphate	30	Sodium chloride	Kolliphor ELP
14	6.5	Citrate-phosphate	30	Dextrose	Tween 80
15	6.5	Citrate-phosphate	30	Sorbitol	Tween 80
16	7	Citrate-phosphate	50	Sucrose	Tween 80
17	7	Citrate-phosphate	10	Lysine	Kollidon 12PF
18	7	Citrate-phosphate	10	Inositol	Tween 80
19	7	Citrate-phosphate	30	Sodium chloride	Tween 40
20	7.5	Citrate-phosphate	50	Inositol	Kolliphor ELP
21	7.5	Citrate-phosphate	50	Sorbitol	Tween 40
22	4	Glycine	10	Sodium chloride	Kollidon 12PF
23	4	Glycine	10	Dextrose	Tween 40
24	4	Glycine	30	Sorbitol	Tween 40
25	4.3	Glycine	50	Sorbitol	Kolliphor ELP
26	4.3	Glycine	50	Inositol	Tween 40
27	4.3	Glycine	50	Dextrose	Tween 80
28	4.5	Glycine	30	Sodium chloride	Tween 40
29	4.8	Glycine	40	Lysine	Tween 80
30	4.8	Glycine	40	Maltose	Tween 80
31	5.8	Glycine	50	Lysine	Kollidon 12PF
32	5.8	Glycine	30	Maltose	Kollidon 12PF
33	6	Glycine	30	Sucrose	Kolliphor ELP
34	6.5	Glycine	30	Sodium chloride	Tween 80
35	6.8	Glycine	40	Dextrose	Kolliphor ELP
36	6.8	Glycine	10	Inositol	Tween 80
37	6.8	Glycine	10	Sorbitol	Kollidon 12PF
38	7	Glycine	10	Lysine	Kolliphor ELP
39	7	Glycine	10	Inositol	Tween 40
40	7	Glycine	30	Sodium chloride	Tween 80
41	7.5	Glycine	30	Inositol	Kolliphor ELP
42	7.5	Glycine	50	Dextrose	Kollidon 12PF
43	7.5	Glycine	10	Sucrose	Kollidon 12PF
44	5	Histidine	10	Maltose	Kolliphor ELP
45	5	Histidine	10	Sorbitol	Kolliphor ELP
46	5	Histidine	20	Dextrose	Kollidon 12PF
47	5.2	Histidine	50	Inositol	Kolliphor ELP
48	5.2	Histidine	50	Maltose	Kolliphor ELP
49	5.2	Histidine	10	Maltose	Kollidon 12PF
50	5.5	Histidine	50	Maltose	Tween 40
51	5.5	Histidine	20	Sodium chloride	Tween 40
52	5.8	Histidine	10	Inositol	Kollidon 12PF
53	5.8	Histidine	10	Inositol	Tween 80
54	5.8	Histidine	50	Lysine	Kolliphor ELP
55	6	Histidine	50	Sodium chloride	Tween 80

56	6	Histidine	10	Sucrose	Kolliphor ELP
57	6	Histidine	10	Sorbitol	Tween 40
58	6	Histidine	30	Sodium chloride	Kollidon 12PF
59	6.5	Histidine	40	Sorbitol	Kollidon 12PF
60	6.5	Histidine	40	Maltose	Tween 80
61	6.5	Histidine	50	Sucrose	Kollidon 12PF
62	6.5	Histidine	50	Dextrose	Tween 40
63	6.6	Histidine	30	Lysine	Tween 80
64	5	Succinate	50	Inositol	Kollidon 12PF
65	5	Succinate	10	Maltose	Kollidon 12PF
66	5	Succinate	50	Sodium chloride	Tween 80
67	5.2	Succinate	30	Sodium chloride	Tween 40
68	5.2	Succinate	50	Lysine	Kolliphor ELP
69	5.2	Succinate	50	Dextrose	Kollidon 12PF
70	5.4	Succinate	10	Maltose	Tween 80
71	5.4	Succinate	30	Inositol	Tween 40
72	5.4	Succinate	10	Dextrose	Tween 40
73	5.5	Succinate	30	Sodium chloride	Kollidon 12PF
74	5.5	Succinate	30	Sucrose	Kollidon 12PF
75	5.5	Succinate	40	Dextrose	Tween 80
76	5.8	Succinate	10	Lysine	Tween 40
77	5.8	Succinate	20	Inositol	Kolliphor ELP
78	5.8	Succinate	50	Sucrose	Kolliphor ELP
79	6	Succinate	30	Sorbitol	Tween 80
80	6	Succinate	50	Sodium chloride	Tween 40

Table 4 lists the analytical tests conducted (short-term stability, mechanical stress, light exposure, F/T) in the framework of this DoE screening and presented herein.

5 Table 4 Panel of analyses conducted on DoE screening formulations

Analysis	Time zero	Stability Study (4 weeks at 40±2°C 75%RH)	Light Stress	Mechanical Stress	Freeze/Thaw Stress
Protein content by OD	X	-	-	-	-
Aggregation by Optical Density	X	X	X	X	X
Visual Inspection	X	X	X	X	X
LMW Fragments by Bioanalyzer ¹ (NR)	X	X	-	X	-
LMW and HMW by CE-SDS (NR)	X	-	X	-	-
HMW by SE-HPLC	X	X	X	X	X
Isoforms by cIEF	X	-	X	-	-

(1) 2100 Bioanalyzer (Agilent)

2.1 Methods used to determine stability

Thermal stability

5

The thermal stability of the formulations was examined after four weeks of storage at 40±2°C (75% R.H.) for the following:

- Aggregation index: calculated by optical density to track aggregation and formation of HMW impurities
- 10 • Visual inspection for presence of visual particles
- HMW content by SE – HPLC (to track aggregation)
- LMW content by bioanalyzer (to track fragmentation)

Light stress

15

The formulations was exposed to 7 hours of light at an intensity of 765 W/m² which satisfies ICHQ1B guideline requirements. The formulations was analyzed by the following techniques:

- Aggregation Index: calculated by OD, measures the extent of aggregate formation
- 20 which results from light stress
- Visual Inspection: for presence of visible particles resulting from aggregation
- CE-SDS: for production of LMW impurities, also indicative of HMW impurities
- SE-HPLC: quantitation of HMW impurities resulting from aggregation
- cIEF: provides insight into relative quantity of charge variants, can monitor oxidation
- 25 (by product of light stress)

Mechanical stress

- 30 Mechanical (shaking) stress is often associated with a production of aggregates due to protein self-association and interaction among hydrophobic regions of the protein in solution. The DoE formulations in this study was examined for resistance to shaking stress after 24 hours of stirring at 200 rpm at room temperature. The shaking stress formulations was analyzed as follows:

- Aggregation index: calculated by optical density to track aggregation and formation of HMW impurities
- Visual inspection for presence of visual particles
- HMW content by SE-HPLC (to track HMW impurity generation and hence monitor aggregations)
- LMW content by bioanalyzer (to track fragmentation)

Freeze/thaw stress

- As a protein formulation freezes, an interface is formed as micro-regions within the solution begin solidify. In these micro-environments there is a change in polarity as different component of the formulation buffer are excluded or included from the liquid matrix that is solidifying. What results is precipitation of protein as hydrophilic/hydrophobic interactions are forced upon the molecules in these changing micro-environments. To ascertain the effectiveness of the various stabilisers and surfactants in the DoE the samples were exposed to three cycles of freeze-thawing. The samples were then examined by the following analyses to determine their resistance to precipitation / aggregation / degradation by freeze-thawing:
- Aggregation index: calculated by optical density to track aggregation and formation of HMW impurities
 - Visual inspection for presence of visual particles
 - HMW content by SE-HPLC (to track HMW impurity generation and hence monitor aggregations)

25

2.2 Manufacturing

- A drug substance material of the composition: 20.6 mg/mL Avelumab, 51 mg/mL D-Mannitol, 0.6 mg/mL glacial acetic acid, pH 5.2 (surfactant – free) was equilibrated by tangential flow filtration (using a Pellicon XL Cassette Biomax cut – off 10 KDa in PES) in the three buffers:

- 10 mM Citrate – phosphate pH 5.2,
- 10 mM Glycine pH 5.2,

- 10 mM Histidine pH 5.2,
- 10 mM Succinate pH 5.2.

The buffer exchange was carried out with a 5-fold dilution of the above mentioned DS in one of the four relevant buffers and equilibrating/concentrating until the initial volume
5 was obtained. The operation was repeated three times. The four equilibrated drug substance materials were tested for protein content by OD prior to formulations manufacturing.

Formulations 1-21 (in citrate – phosphate buffer)

- 10 The exchanged DS material (26.4 mg/mL) was weighed in a glass beaker (30.30 grams). If needed, the strength of the buffer was adjusted (starting molarity of the exchanged DS: 10 mM; molarity range in the DoE formulas: 10 – 50 mM) by adding di-sodium hydrogen phosphate dihydrate and citric acid monohydrate. The solution was stirred until complete dissolution. The stabiliser was then added: Sorbitol (2.04 grams)
15 or Dextrose (2.02 g) or Inositol (2.02 g) or Maltose monohydrate (4.04 g) or Lysine monohydrochloride (2.02 g) or Sodium Chloride (0.327 g) or Sucrose (3.83 g). The solution was stirred until complete dissolution. The surfactant was then added: 0.4 mL of a 50 mg/mL Tween 40 stock or 0.4 mL of a 50 mg/mL Tween 80 stock or 0.4 mL of a 50 mg/mL Kolliphor ELP stock or 20 mg of Kollidon 12PF (no stock solution needed).
20 The solution was stirred until complete dissolution. The pH was measured and adjusted to target with diluted o-phosphoric acid or sodium hydroxide. The solution was brought to final weight (40 g) with the relevant buffer.

Formulations 22-31 (in glycine buffer)

- 25 The exchanged DS material (24.5 mg/mL) was weighed in a glass beaker (32.65 g). If needed, the strength of the buffer was adjusted (starting molarity of the exchanged DS: 10 mM; molarity range in the DoE formulas: 10 – 50 mM) by adding glycine. The solution was stirred until complete dissolution. The stabiliser was then added: Sorbitol (2.04 g) or Dextrose (2.02 g) or Inositol (2.02 g) or Maltose monohydrate (4.04 g) or
30 Lysine monohydrochloride (2.02 g) or Sodium Chloride (0.327 g) or Sucrose (3.83 g). The solution was stirred until complete dissolution. The surfactant was then added: 0.4 mL of a 50 mg/mL Tween 40 stock or 0.4 mL of a 50 mg/mL Tween 80 stock or 0.4 mL of a 50 mg/mL Kolliphor ELP stock or 20 mg of Kollidon 12PF (no stock solution needed). The solution was stirred until complete dissolution. The pH was measured and

adjusted to target with diluted hydrochloric acid or sodium hydroxide. The solution was brought to final weight (40 g) with the relevant buffer.

Formulations 32-43 (in glycine buffer)

- 5 The exchanged DS material (23.2 mg/mL) was weighed in a glass beaker (34.48 g). If needed, the strength of the buffer was adjusted (starting molarity of the exchanged DS: 10 mM; molarity range in the DoE formulas: 10 – 50 mM) by adding glycine. The solution was stirred until complete dissolution. The stabiliser was then added: Sorbitol (2.04 g) or Dextrose (2.02 g) or Inositol (2.02 g) or Maltose monohydrate (4.04 g) or
- 10 Lysine monohydrochloride (2.02 g) or Sodium Chloride (0.327 g) or Sucrose (3.83 g). The solution was stirred until complete dissolution. The surfactant was then added: 0.4 mL of a 50 mg/mL Tween 40 stock or 0.4 mL of a 50 mg/mL Tween 80 stock or 0.4 mL of a 50 mg/mL Kolliphor ELP stock or 20 mg of Kollidon 12PF (no stock solution needed). The solution was stirred until complete dissolution. The pH was measured and
- 15 adjusted to target with diluted hydrochloric acid or sodium hydroxide. The solution was brought to final weight (40 g) with the relevant buffer.

Formulations 64-80 (in succinic buffer)

- The exchanged DS material (22.5 mg/mL) was weighed in a glass beaker (35.55
- 20 grams). If needed, the strength of the buffer was adjusted (starting molarity of the exchanged DS: 10 mM; molarity range in the DoE formulas: 10 – 50 mM) by adding succinic acid. The solution was stirred until complete dissolution. The stabiliser was then added: Sorbitol (2.04 g) or Dextrose (2.02 g) or Inositol (2.02 g) or Maltose monohydrate (4.04 g) or Lysine monohydrochloride (2.02 g) or Sodium Chloride (0.327
- 25 g) or Sucrose (3.83 g). The solution was stirred until complete dissolution. The surfactant was then added: 0.4 mL of a 50 mg/mL Tween 40 stock or 0.4 mL of a 50 mg/mL Tween 80 stock or 0.4 mL of a 50 mg/mL Kolliphor ELP stock or 20 mg of Kollidon 12PF (no stock solution needed). The solution was stirred until complete dissolution. The pH was measured and adjusted to target with diluted hydrochloric acid
- 30 or sodium hydroxide. The solution was brought to final weight (40 grams) with the relevant buffer.

Formulations 44-63 (in histidine buffer)

The exchanged DS material (24.4 mg/mL) was weighed in a glass beaker (32.80 g). If needed, the strength of the buffer was adjusted (starting molarity of the exchanged DS: 10 mM; molarity range in the DoE formulas: 10 – 50 mM) by adding histidine. The solution was stirred until complete dissolution. The stabiliser was then added: Sorbitol (2.04 g) or Dextrose (2.02 g) or Inositol (2.02 g) or Maltose monohydrate (4.04 g) or Lysine monohydrochloride (2.02 g) or Sodium Chloride (0.327 g) or Sucrose (3.83 g). The solution was stirred until complete dissolution. The surfactant was then added: 0.4 mL of a 50 mg/mL Tween 40 stock or 0.4 mL of a 50 mg/mL Tween 80 stock or 0.4 mL of a 50 mg/mL Kolliphor ELP stock or 20 mg of Kollidon 12PF (no stock solution needed). The solution was stirred until complete dissolution. The pH was measured and adjusted to target with diluted hydrochloric acid or sodium hydroxide. The solution was brought to final weight (40 grams) with the relevant buffer.

15 Filtration and filling

Each formulation was filtered through a 0.22 micron filter assembled on a 50 mL syringe (Millex GP 0.22 μ m Express PES membrane or Millex GV 0.22 μ m Durapore PVDF membrane) were used. The filtered solution was then filled in the relevant container (2 mL/container).

20

2.3 RESULTS*Check of protein content by OD upon manufacturing*

The protein content was determined by OD at time 0 (upon manufacturing). Values in line with the expected target (20 mg/mL) were found.

25

2.3.1 Thermal stress*Aggregation index by OD*

The aggregation index was determined by OD. Additional information on aggregation index as a tool to detect sub-visible particles/larger aggregates not detectable by SE-HPLC are provided in the Annex section.

30

It was found that histidine buffer is generally associated to higher increases in aggregation index upon stress (i.e. larger increase in particles), most significantly when

the pH is increased from 5.0 to 6.6 (pH dependent effect).

In the other buffers, changes in aggregation index are generally lower, thus indicating lower increases in sub-visible particles.

The increases in aggregation index observed in some (few) samples formulated in

- 5 citrate-phosphate and glycine buffer are not directly attributable to a specific factor (e.g. stabiliser or surfactant type).

The data were statistically evaluated by ANOVA for Response Surface Linear Model, which provided the following outcome:

Statistically significant impact of buffer type, strength and pH (all have a p-value <

- 10 0.001): in order to minimize the aggregation index low buffer strengths should be targeted (10 mM), in association with low pH ranges in citrate-phosphate (4.0 – 5.0) and glycine (4.0 – 5.8) and succinate (5.0 – 5.5), while histidine generally determines a negative impact on sub-visible particles/larger aggregates formation.

15 *Total aggregates by SE - HPLC*

Total aggregates (HMWs) were determined by SE – HPLC at time 0 and upon thermal stress. Citrate – phosphate generally leads to higher aggregation than reference

formula (reference threshold highlighted as a red horizontal bar in the chart), most

particularly as pH increases. In glycine buffer, low pH ranges are to be preferred (lower

- 20 than 5.0), being higher pH values associated with higher aggregation (similarly to when citrate – buffer is used). Succinate generally leads to higher aggregation values than the reference at all conditions, while histidine buffer at low pH (5.0 – 5.5) seems to provide aggregation values comparable to the reference.

The data were also statistically evaluated by ANOVA for Response Surface Linear

- 25 Model and buffer type was confirmed to be a significant factor (p-value = 0.02).

Overall, in order to reduce aggregates upon thermal stress, citrate – phosphate (pH range 4.0 – 5.0), glycine (pH range 4.0 – 6.8) and histidine (pH range 5.0 – 5.8) should be preferred over succinate buffer.

- 30 Combinations like those present in formulations # 2 (Tween 40 + Dextrose in citrate – phosphate buffer pH 4.0), formulation # 22 (Kollidon 12PF + Sodium chloride in glycine buffer pH 4.0) and formulation # 28 (Tween 40 + sodium chloride in glycine buffer pH 4.5) seem to be unfavorable to protein stabilization (significant increase in aggregation despite the optimal pH/buffer conditions applied) possibly due to incompatibility of

Kollidon 12PF and Tween 40 with low pH (about 4.0 – 4.5)/interaction with specific stabilisers like sodium chloride.

Fragments by Bioanalyzer

5 Fragmentation levels were assessed by Bioanalyzer. Although no statistically significant results could be highlighted by ANOVA evaluation, conditions which were most effective in minimizing fragmentation providing LMWs percentages in line with reference composition could be highlighted:

- Citrate – phosphate buffer in the pH range of 4.5 – 7.0
- 10 - Glycine buffer in the pH range 4.0 – 5.8.

Considering the variability of the method (up to $\pm 2 - 3\%$ in LMWs is common when Bioanalyzer is applied), other conditions (like the remaining compositions in histidine and succinate buffers) were observed to maintain the LMWs % relatively low and are therefore worth investigating further.

15

Visible particles by visual inspection

The presence of visible particles was assessed by visual inspection before and after thermal stress. Varying conditions in citrate – phosphate buffer can generate the presence of visible particles (most typically particulate – like suspensions) following
20 thermal stress.

In glycine buffer, particles formation is most frequently associated to the presence of Tween species (Sampe ID # 23, 24, 26, 28 containing Tween 40) and formulation # 30 containing Tween 80. Other formulations in glycine buffer (Sampe ID from # 32 to # 39) showed presence of particles at time 0 which tended to decrease upon stress (possible
25 reversible clusters).

In histidine, Tween species are generally associated to visible particles formation upon stress (all formulations showing visible particles after stress contain one of the two Tween alternatives).

In succinate buffer, particles observed at time 0 in most formulations were found to
30 decrease upon thermal stress (possible disruption of reversible associations over time).

Summary: thermal stress

According to SE – HPLC, OD and Bioanalyzer upon thermal stress, conditions that can provide favorable performances include:

- Buffers: Citrate – phosphate or glycine (preferably at more acidic pH and most relevantly in the range 4.0 – 5.0 for citrate phosphate and 4.0 – 5.8 for glycine),
- Buffer strength: preferably low (as per aggregation index outcome),
- Stabiliser: no specific indication obtained,
- 5 - Surfactant: Kolliphor ELP observed to be effective in reducing sub-visible particles.

2.3.2 Light stress

Aggregation index by O.D.

- 10 Aggregation index in most DoE compositions in citrate – phosphate buffer was found to be higher than in reference formula (most significantly in the higher pH range). The pH effect was also confirmed in glycine buffer, which was however found to considerably lower the aggregation index with respect to citrate – phosphate buffer (in the pH range 4.0 – 4.5 values comparable with reference compositions or lower were highlighted).
- 15 Histidine can generally cause considerable increases in aggregation index as well as succinate buffer (histidine remarkably worse than succinate).

The statistical analysis by ANOVA confirmed the significant impact from buffer type, pH and strength (p – value < 0.0001), indicating that the best conditions to minimize particles formation include utilisation of citrate phosphate buffer (in the range 4.0 – 5.0 and at low buffer strength), glycine (in the range 4.0 – 5.8).

20 Surfactant was also observed to have some impact on stability, being Kolliphor ELP the best option to be taken into account when aiming at particles reduction.

Total aggregates by SE - HPLC

- 25 Total aggregates (HMWs) were determined by SE – HPLC at time 0 and upon light stress. Citrate – phosphate generally leads to higher aggregation than reference formula, most particularly as pH increases. In glycine buffer, low pH ranges are to be preferred (lower than 4.8), being higher pH values associated with higher aggregation (similarly to when citrate – buffer is used). Succinate generally leads to higher
- 30 aggregation values than the reference at all conditions, while histidine buffer (whole range aside from few exceptions) seems to provide aggregation values comparable to the reference.

The data were also statistically evaluated by ANOVA for Response Surface Linear Model and buffer type and pH were confirmed to be significant factors (p-value < 0.0001).

Overall, in order to reduce aggregates upon thermal stress, glycine (pH range 4.0 – 5.0) and histidine (pH range 5.0 – 6.0) should be preferred over succinate and citrate phosphate buffers.

Importantly, stabilisers like Lysine, Dextrose, Sorbitol and Sucrose provide better stabilization against light stress than sodium chloride, maltose and Inositol (p – value < 0.01).

Purity by CE - SDS

Purity as determined by CE – SDS carries the information of both HMWs and LMWs species as it is the results of the calculation: 100 - % HMWs by CE – SDS - % LMWs by CE – SDS.

Purity values were determined before and after light stress.

Most formulations show higher purity than reference compositions upon light stress.

Conditions that can impact negatively on stability are typically: citrate phosphate at high pH (>7.0) and glycine buffer at low pH (4.0); the latter is most probably to be explained with the negative impact from Tween 40 / Kollidon 12PF at low pH.

Histidine was found to positively impact on purity, maximising formulation performances against light exposure.

Statistical analysis by ANOVA confirmed superior behaviour associated to histidine utilisation as a buffer, with comparable performances obtained when using citrate – phosphate, glycine or succinate buffers.

Isoforms profile by cIEF

Isoforms profiles were determined at time 0 and after light exposure. Light exposure generally determines an increase in acidic isoforms due to photo-oxidation phenomena. Such increase was calculated for all DoE formulations.

Several conditions are favourable to protein stabilization (i.e. lower changes in isoforms profile), such as citrate – phosphate and glycine buffer (most typically in the lower pH range). Lower performances observed when histidine is used as formulation buffer. The data, evaluated by ANOVA for Response Surface Linear Model confirmed the above (buffer type statistically significant factor with p – value < 0.0001).

The statistical analysis also confirmed a positive impact (reduction in acidic isoforms change) when L-Lysine is used as stabiliser. The effect is quite clear when observing the changes found in formulations # 11, 29, 31, 38, considerably lower than those in the surrounding formulation space with alternative stabilisers.

5

Visible particles by visual inspection

The presence of visible particles was assessed by visual inspection before and after light stress. Most formulations are not impacted by light stress in terms of visible particles. No specific conditions related to particle formation upon light stress.

10

Summary: light exposure stress

According to SE – HPLC, OD, CE – SDS, cIEF and visual inspection upon light stress, conditions that can provide favorable performances include:

- Buffers: glycine buffer (preferably at more acidic pH and most relevantly in the range 4.0 – 4.5),
- Buffer strength: preferably low (as per aggregation index outcome),
- Stabiliser: Lysine (monohydrochloride), dextrose and sorbitol showed a positive impact on protein stability
- Surfactant: Kolliphor ELP observed to be effective in reducing sub-visible particles

20

2.3.3 Freeze – thawing

Aggregation index by optical density

After 3X freeze – thawing cycles ($-80^{\circ}\text{C} \rightarrow$ room temperature), once again, glycine buffer (low pH) is confirmed to provide the lowest values indicating lower particle formation. An increase in aggregation index is observed both in citrate – phosphate buffer and glycine buffer as pH increase (pH effect more critical in citrate – phosphate buffer). Generally higher aggregation index values than reference composition are observed in histidine and succinate buffers.

The statistical analysis by ANOVA highlighted a moderately significant impact from buffer type, pH and surfactant type ($0.01 < p - \text{value} < 0.05$), indicating that citrate – phosphate and glycine buffers at pH lower than 6.0 are the best option for protein stabilisation against particles formation induced by freeze – thawing, being succinate and histidine buffer slightly pejorative with respect to reference composition.

A comparison of the impact of the different surfactants shows comparable performances from Tween 80, Kollidon 12PF and Kolliphor ELP (slightly preferable), while Tween 40 is expected to increase aggregation index.

5 *Total aggregates by SE – HPLC*

All formulations show lower total aggregates than reference composition upon freeze – thawing stress (values comparable to time 0).

In citrate – phosphate buffer, aggregates tend to increase up to the level of reference composition as the primary effect of pH (2.0 – 2.5% HMWs) being increased up to the
10 range 7.0 - 7.5 with minor/negligible changes upon freeze – thawing, whilst at pH < 7.0 total aggregates typically amount to lower than 1.5% (before and after stress).

In glycine and histidine buffer all total aggregates values after stress amount to less than 1% (comparable with time 0 values). In succinate, freeze – thawing was not found to determine critical changes with respect to time 0, however total aggregates are
15 generally slightly higher than in glycine and histidine (still equal to or lower than 1.5%, i.e. considerably lower than reference after stress).

Statistical analysis confirmed the significant impact from buffer type and pH (p – value < 0.0001), being citrate – phosphate buffer (pH 4.0 – 6.0), glycine buffer (pH 4.0 – 7.0) and histidine (5.0 – 6.6) the best options for protein stabilisation against freeze –
20 thawing.

A significant impact (p – value < 0.01) was also highlighted for the stabiliser type factor: Lysine hydrochloride minimises time 0 aggregation and the effects related to freeze – thawing stress (cf. Sampe ID # 6-9-11-17 in citrate – buffer); sucrose and dextrose, similarly, show stabilising properties.

25

Visible particles by visual inspection

In the results of visual inspection upon freeze – thawing the general trends that can be highlighted:

- In citrate – phosphate, particle formation is more likely at higher pH,
- 30 - In glycine buffer at low pH (< 5), particle formation is primarily related to the presence of Tween 40 (destabilising surfactant),
- In histidine buffer, Tween species are generally related to particle formation,
- In succinate, no specific factors seem to be related to particle formation, which is however quite a frequent occurrence when this buffer is used.

Summary: freeze – thawing stress

According to SE – HPLC, OD and visual inspection upon 3X freeze – thawing cycles (-80°C → room temperature), conditions that can provide favorable improved

5 performances include:

- Buffers: glycine or citrate – phosphate buffers (preferably at more acidic pH and most relevantly in the range 4.0 – 6.0),
- Stabiliser: Lysine (monohydrochloride), dextrose and sucrose showed a positive impact on protein stability (reduction of total aggregates by SE – HPLC),
- 10 - Surfactant: incompatibilities of Tween species with glycine and histidine buffered formulations are to be taken into account and avoided to minimize visible particles formation.

2.3.4 Mechanical stress

15

Aggregation index by Optical density

As previously shown, the factors that allow aggregation index values most similar to reference (i.e. minimal or no increases with respect to time 0) are:

20 Citrate – phosphate generally leads to higher aggregation index values than reference, most particularly as pH increases and in presence of Tween species: Sampe ID # 2 (Tween 40), # 8 (Tween 80), # 11 (Tween 40), # 19 (Tween 40), # 21 (Tween 40).

Glycine provides a conspicuous stabilising effect in the low pH range (aggregation index values slightly lower than reference).

25 Histidine buffer is to be preferably used at pH values close to 5.0 and without Tween 40 and Tween 80, which appear to be related to the highest aggregation index values: Sampe ID # 50 (Tween 40), # 60 (Tween 80), # 62 (Tween 40).

Succinate generally leads to aggregation index values slightly higher than reference composition, regardless of the specific factors involved.

30 The above results were confirmed by ANOV, which indicated buffer type and pH as statistically significant factors (p – value < 0.01) and surfactant as moderately significant factor ($0.01 < p$ –value < 0.05).

Glycine buffer at low pH (4.0 – 5.5) is highlighted as the selection buffer to minimise the aggregation index. The tendency towards an increase in aggregation index given by Tween species (Tween 40 worse than Tween 80) is confirmed by the surface response models.

5

Total aggregates by SE – HPLC

Minimal increase with respect to time 0 were observed for most formulations indicating a minor impact from this type of stress. Differentiation in terms of total aggregates appears to be the primary effect of buffer type and pH, as already highlighted

10 Buffer type and pH confirmed to be statistically significant factors by ANOVA (p – value < 0.0001); as well as buffer strength (p – value < 0.01) and stabiliser type (0.01 < p – value < 0.05).

Preferable ranges and conditions to minimise aggregates to the level of reference composition (< 1%) include: citrate – phosphate buffer (pH < 5 and low ionic strength);

15 glycine buffer (whole pH and ionic strength range); histidine buffer (whole range) and succinate buffer (pH 5.0 – 5.5 and low ionic strength). Preferable stabilisers are L-Lysine monohydrochloride, Maltose, Sucrose and Dextrose.

Fragments by Bioanalyzer

20 Except for Sampe ID # 22-23-24 (in glycine buffer, pH 4.0, containing Tween 40 or Kollidon 12PF), the remaining formulations showed LMWs % comparable to or lower than reference composition upon mechanical stress, also taking into account the variability of this method ($\pm$ 2-3% in LMWs% results is characteristic). Therefore, it can be concluded that most conditions tested can help improve protein resistance against

25 fragmentation provided that combinations like glycine buffer (low pH) + Tween 40 are avoided.

The statistical elaboration highlighted the better performances of formulations in succinate and histidine buffers, to be however carefully considered and evaluated as substantially comparable to/slightly better than the other formulas in citrate – phosphate

30 and glycine buffer due to the above discussed method variability.

Visible particles by visual inspection

In the results of visual inspection upon freeze – thawing are the general trends that can be highlighted:

- In citrate – phosphate buffer (Sampe ID # 1 – 21), particle formation occurs at almost all conditions regardless of specific factors involved,
- In glycine buffer, particle formation is primarily related to the presence of Tween 40 (Sampe ID # 23, 26, 28 and Kollidon 12PF (Formulations # 22, 32, 37, 43)
- 5 - In histidine buffer, all formulas showing increase of visible particles upon mechanical shaking contain either Tween 40 or Tween 80,
- In succinate, no specific factors seem to be related to particle formation.

Summary: mechanical stress

- 10 According to SE – HPLC, OD, Bioanalyzer and visual inspection upon mechanical shaking, conditions that can provide favorable performances with respect to reference compositions include:
- Buffers: glycine (preferably at more acidic pH and most relevantly in the range 4.0 – 5.5), histidine and succinate at pH of about 5.0.
 - 15 - Stabiliser: Lysine (monohydrochloride), Sucrose, Maltose and Dextrose showed a positive impact on protein stability (reduction of total aggregates by SE – HPLC),
 - Surfactant: incompatibilities of Tween species with glycine, citrate-phosphate and histidine buffered formulations are to be taken into account and avoided to minimize visible particles formation.

20

Example 3 – Formulations optimisation

3.1 Formulation optimisation

- 25 The data shown in Example 2 were combined to identify the formulation space which could suitably stabilise Avelumab (factors evaluated: buffer type, pH and strength, stabiliser type and surfactant) against thermal, freeze – thaw, mechanical and light stress.
- 30 Using the following criteria
- Minimise HMWs (by SE – HPLC) after thermal stress, mechanical shaking, freeze – thawing and light stress,
 - Minimise LMWs (by Bioanalyser) after thermal stress and mechanical shaking,
 - Maximise purity (by CE-SDS) after light stress,

- Minimise acidic isoforms (by cIEF) change after light stress,
 - Target Aggregation index values (by OD) lower than 2 after thermal stress, mechanical shaking, freeze – thawing and light stress,
- for each buffer type the 10 most promising formulations were extrapolated as shown in
- 5 Table 5.

Table 5: Candidate formulations (DoE extrapolation)

Number	Buffer pH	Buffer strength (mM)	Surfactant (0.5 mg/mL)	Stabiliser (280 mM)	Buffer type
1	4,4	10	Kolliphor ELP	Lysine	Glycine
2	4,1	10	Tween 80	Lysine	Glycine
3	4,0	10	Tween 80	Lysine	Glycine
4	4,0	10	Kolliphor ELP	Dextrose	Glycine
5	4,0	10	Kolliphor ELP	Dextrose	Glycine
6	4,0	10	Kolliphor ELP	Dextrose	Glycine
7	4,0	10	Kollidon 12PF	Lysine	Glycine
8	4,0	10	Kolliphor ELP	Sorbitol	Glycine
9	4,0	10	Kolliphor ELP	Sucrose	Glycine
10	4,0	10	Kolliphor ELP	Sucrose	Glycine

Number	Buffer pH	Buffer strength (mM)	Surfactant (0.5 mg/mL)	Stabiliser (280 mM)	Buffer type
1	4,0	10	Kollidon 12PF	Sorbitol	Citrate-phosphate
2	4,2	15	Kollidon 12PF	Lysine	Citrate-phosphate
3	4,3	17	Kollidon 12PF	Sucrose	Citrate-phosphate
4	4,1	20	Kollidon 12PF	Lysine	Citrate-phosphate
5	4,1	15	Tween 80	Lysine	Citrate-phosphate
6	4,0	27	Kollidon 12PF	Sucrose	Citrate-phosphate
7	4,1	19	Kolliphor ELP	Sucrose	Citrate-phosphate
8	4,1	22	Kolliphor ELP	Dextrose	Citrate-phosphate
9	4,2	13	Tween 80	Sorbitol	Citrate-phosphate
10	4,2	17	Kolliphor ELP	Dextrose	Citrate-phosphate

Number	Buffer pH	Buffer strength (mM)	Surfactant (0.5 mg/mL)	Stabiliser (280 mM)	Buffer type
1	5,0	10	Kolliphor ELP	Dextrose	Histidine
2	5,0	10	Kolliphor ELP	Dextrose	Histidine
3	5,0	10	Kolliphor ELP	Sorbitol	Histidine
4	5,0	10	Kolliphor ELP	Sucrose	Histidine
5	5,0	11	Kolliphor ELP	Sucrose	Histidine
6	5,1	10	Kolliphor ELP	Sorbitol	Histidine
7	5,1	10	Kolliphor ELP	Sucrose	Histidine
8	5,0	10	Kolliphor ELP	Inositol	Histidine
9	5,0	15	Kolliphor ELP	Sorbitol	Histidine
10	5,0	10	Kolliphor ELP	Lysine	Histidine

Number	Buffer pH	Buffer strength (mM)	Surfactant (0.5 mg/mL)	Stabiliser (280 mM)	Buffer type
1	5,0	10	Kollidon 12PF	Sucrose	Succinate
2	5,0	10	Kolliphor ELP	Lysine	Succinate
3	5,0	10	Kolliphor ELP	Lysine	Succinate
4	5,0	12	Kolliphor ELP	Lysine	Succinate
5	5,1	10	Kolliphor ELP	Lysine	Succinate
6	5,1	10	Kollidon 12PF	Sucrose	Succinate
7	5,0	10	Kollidon 12PF	Sorbitol	Succinate
8	5,0	10	Kollidon 12PF	Lysine	Succinate
9	5,0	10	Kollidon 12PF	Dextrose	Succinate
10	5,0	14	Kollidon 12PF	Sucrose	Succinate

3.2 Lead formulations to be further assessed

5

Out of the formulations of Table 5, the eleven formulations listed in Table 6 appeared most promising. Hence, they were manufactured and evaluated upon thermal stress and repeated freeze-thawing cycles as per the analytical panel shown in Table 7.

- 10 Thermal stress was selected as the most relevant stress conditions to evaluate formulation performances and possibly predict stability at refrigerated conditions. Freeze – thawing was also considered in order to anticipate any issues related to temperature excursions/storage of pre-formulated DS materials.
- The results of the experiments carried out on these formulation are described in the
- 15 following paragraphs.

Table 6: Lead formulations resulting from DoE

DP	pH (± 0.1)	Buffer	Buffer Strength (mM)	Stabiliser	Surfactant (0.5 mg/mL)
1	4.4	Glycine	10	Lysine (monohydrochloride) 280 mM	Kolliphor ELP
2	4.4	Glycine	10	Lysine (monohydrochloride) 140 mM	Kolliphor ELP
3	4.4	Glycine	10	Lysine (monohydrate) 280 mM	Kolliphor ELP
4	4.4	Glycine	10	Lysine acetate 140 mM	Kolliphor ELP
5	4.1	Glycine	10	Lysine (monohydrate) 280 mM	Tween 80
6	5.0	Histidine	10	Dextrose 280 mM	Kolliphor ELP
7	5.0	Histidine	10	Sucrose 280 mM	Kolliphor ELP
8	4.2	Citrate- Phosphate	15	Lysine (monohydrochloride) 140 mM	Kollidon 17PF
9	4.3	Citrate- Phosphate	17	Sucrose 280 mM	Kollidon 17PF
10	5.0	Succinate	10	Lysine (monohydrochloride) 140 mM	Kolliphor ELP
11	5.0	Succinate	10	Sucrose 280 mM	Kollidon 17PF

5 Table 7: Panel of analyses conducted on lead formulations

Test	Time 0	Thermal Stress (4 weeks at 40 $\pm$ 2°C 75%RH)	Freeze-thaw 3X
Visible particles (visual)	X	X	X
pH	X	X	-
Turbidity (OD)	X	X	X
Sub-visible particles (PAMAS)	X	X	X
Protein content (OD)	X	X	-
HMWs by SE - HPLC	X	X	X
LMWs by Bioanalyzer	X	X	-
Isoforms profile by iCE	X	X	-
Tertiary structure by CD	X	X	-

3.3 Manufacturing of lead formulations resulting from DoE step

A drug substance material of the composition: 18.6 mg/mL avelumab, 51 mg/mL D-Mannitol, 0.6 mg/mL glacial acetic acid, pH 5.2 (surfactant – free) was equilibrated by tangential flow filtration (using a Pellicon XL Cassette Biomax cut – off 50 KDa in PES) in the three buffers:

- 10 mM Glycine pH 4.4,
- 10 mM histidine pH 5.0,
- 15 mM citrate-phosphate pH 4.2,
- 10 mM succinate pH 5.0.

The buffer exchange was carried out with a 5-fold dilution of the above mentioned DS in one of the four relevant buffers and equilibrating / concentrating until the initial volume was obtained. The operation was repeated three times. The four equilibrated drug substance materials were tested for protein content by OD prior to formulations manufacturing.

Formulations 1-5 (in glycine buffer)

The exchanged DS material (21.8 mg/mL) was weighed in a glass beaker (64.2 g). The stabiliser was then added: Lysine monohydrochloride (3.58 grams for DP1 or 1.79 g for DP2) or Lysine monohydrate (3.22 grams for DP3 and DP5) or Lysine Acetate (2.02 g for DP4). The solution was stirred until complete dissolution. The surfactant was then added: 0.7 mL of a 50 mg/mL Kolliphor ELP stock (in 10 mM glycine pH 4.4 for DP 1-2-3-4) or 0.7 mL of a 50 mg/mL Tween 80 (in 10 mM glycine pH 4.1 for DP5). The solution was stirred until complete dissolution. The pH was measured and adjusted to target with diluted hydrochloric acid or sodium hydroxide. The solution was brought to final weight (70 g) with the relevant buffer.

Formulations 6-7 (in histidine buffer)

The exchanged DS material (23.2 mg/mL) was weighed in a glass beaker (60.3 g). The stabiliser was then added: Dextrose (3.53 g for DP6) or Sucrose (6.71 g for DP7). The solution was stirred until complete dissolution. The surfactant was then added: 0.7 mL of a 50 mg/mL Kolliphor ELP stock (in 10 mM histidine buffer pH 5.0 for DP6 and 7). The solution was stirred until complete dissolution. The pH was measured and adjusted to target (pH 5.0) with diluted hydrochloric acid or sodium hydroxide. The solution was

brought to final weight (70 g) with relevant buffer (10 mM histidine buffer pH 5.0).

Formulations 8-9 (in citrate-phosphate buffer)

The exchanged DS material (23.4 mg/mL) was weighed in a glass beaker (59.8 g). If
5 needed (DP9), the strength of the buffer was adjusted by adding citric acid
(monohydrate) and di-sodium phosphate hydrogen (dihydrate). The stabiliser was then
added: Lysine monohydrochloride (1.79 g for DP8) or Sucrose (6.71 g for DP9). The
solution was stirred until complete dissolution. The surfactant was then added: 35 mg of
Kollidon 17PF (for both DP8 and 9). The solution was stirred until complete dissolution.
10 The pH was measured and adjusted to target (pH 4.2 for DP8 and 4.3 for DP9) with
diluted o-phosphoric acid or sodium hydroxide. The solution was brought to final weight
(70 g) with the relevant buffer.

Formulations 10-11 (in succinate buffer)

15 The exchanged DS material (24.5 mg/mL) was weighed in a glass beaker (57.1 g) (g
ms). The stabiliser was then added: Lysine monohydrochloride (1.79 g for DP10) or
Sucrose (6.71 g for DP11). The solution was stirred until complete dissolution. The
surfactant was then added: 0.7 mL of a 50 mg/mL Kolliphor ELP stock solution in 10
mM succinate buffer pH 5.0 (DP10) or 35 mg of Kollidon 17PF (DP11). The solution
20 was stirred until complete dissolution. The pH was measured and adjusted to target (pH
5.0 for DP10 and 11) with diluted hydrochloric acid or sodium hydroxide. The solution
was brought to final weight (70 g) with 10 mM succinate buffer pH 5.0.

3.4 Results

25

3.4.1 Thermal stress

Protein content by OD: No major changes observed with respect to time 0 after 4 weeks
at 40°C.

30

pH: The pH values at time 0 were in line with the target. No major changes were
observed with respect to time 0 after 4 weeks at 40°C.

Visible particles by visual inspection

All formulations were found to be free of visible particles at time 0. Upon stress, one formulation (DP6) showed the presence of particles (possibly formulation – related).

5 *Turbidity by Nephelometry*

Most formulations have turbidity values in the clear or slightly opalescent range with minimal changes after stress (DP 2-4-6-7-9-10-11). Other formulations show either higher turbidity changes from the slightly opalescent to the opalescent range (DP1) or values in the opalescent range already at time 0 with minor/negligible changes after stress (DP 3-8). Formulation DP5 shows a significant increase in turbidity (> 18 NTU) after stress.

15 *Sub-visible particles by light obscuration*

Particles ≥ 25 micron were well below the Pharmacopoeia limit of 600 particles / container (typically < 100 particles). Particles ≥ 10 micron had somewhat larger counts, but were still below the 6000 particles / container limit. DP8 and 9, in citrate-phosphate buffer, showed higher counts than the others (still below the above limit) at time 0, with significant reduction after stress.

20

Total aggregates by SE - HPLC

With respect to total aggregates by SE – HPLC at time 0 and after thermal stress, DP 1-2-3-4 (glycine buffer) varied for the stabiliser type and amount, but had the same buffer strength, surfactant and pH): reduction in Lysine monohydrochloride from 280 mM (DP1) to 140 mM (DP2) seems to favor protein stability. The higher aggregation rate was confirmed when Lysine monohydrate at 280 mM was used (DP3). Lysine acetate (140 mM) provided similar performances as Lysine monohydrochloride used at the same concentration (DP2).

DP5 (glycine buffer) showed significant increase in aggregates (probably due an unfavourable combination of Lysine monohydrate at 280 mM + Tween 80 instead of Kolliphor ELP).

DP6-7 (histidine buffer) showed no changes in aggregates.

DP8-9 (citrate-phosphate buffer): sucrose in DP9 seems to be the critical factor which can significantly improve formulation performance with respect to DP8 (Lysine

monohydrate) being the other ingredients/parameters pretty similar (same buffer type, same surfactant and similar pH: 4.2 vs. 4.3).

DP10-11 (succinate buffer): no significant changes in aggregation were observed (similar performances of Lysine monohydrate and Sucrose in this buffer).

5

Lower molecular weights by Bioanalyzer

Fragments by Bioanalyzer at time 0 and after thermal stress:

DP 1-2-3-4 (glycine buffer) varied for the stabiliser type and amount, but had the same buffer strength, surfactant and pH): similar increase in fragments (+3-5% after stress).

10 DP5 (glycine buffer) showed significant increase in lower molecular weight species (probably due an unfavourable combination of Lysine monohydrate at 280 mM + Tween 80 instead of Kolliphor ELP): +13% increase after stress.

DP6-7 (histidine buffer) showed no changes in fragments.

15 DP8-9 (citrate-phosphate buffer): sucrose in DP9 (+6% in fragments after stress) seems to be the critical factor which can significantly improve formulation performance with respect to DP8 (Lysine monohydrate; +11% in fragments) being the other ingredients/parameters pretty similar (same buffer type, same surfactant and similar pH: 4.2 vs. 4.3).

20 DP10-11 (succinate buffer): minimal changes for both (similar performances of Lysine monohydrate and Sucrose in this buffer): +1-3% in lower molecular weight species after stress.

Isoforms profile by cIEF

25 Isoforms profile at time 0 and after thermal stress: Upon thermal stress all samples generally tended to lose part of the main species with concurrent increase in acidic species and minor changes in the basic isoforms. More in detail: DP 1-2-3-4-5 (glycine buffer): similar changes were observed in isoforms profile. For the five samples, main species decreased by about 10-12% (increase in acidic isoforms of 14 – 17% and decrease in basic isoforms of -4/-6%).

30 DP 6-7 (histidine buffer): DP6 showed major changes in isoforms profile and the profiles obtained could not be elaborated due to likely instability from the components chosen and/or contamination of the sample prior to analysis. DP7 showed changes similar to samples in glycine buffer.

DP8-9 (citrate-phosphate buffer): significant changes in both formulations, higher than observed in the other buffers. Acidic species were found to increase up to 24 – 29% after stress.

- 5 DP10-11 (succinate buffer): DP10 showed minimal changes, even lower than the other samples in the other buffers: main species decreased by about 7% (increase in acidic isoforms of about 12% and decrease in basic isoforms of about -5%). DP11 showed higher changes (increase in acidic isoforms after stress was +20%).

Tertiary structure by Circular dichroism

- 10 Circular dichroism was run before and after stress on the lead formulations. The samples were diluted with WFI to 1.5 mg/mL and then tested in 1 cm – pathlength quartz cuvettes with a Jasco J-810 spectropolarimeter in the range 250 nm – 320 nm at a scanning speed of 20 nm/min (sensitivity: standard; bandwidth: 1 nm; data pitch 0.2 nm; D.I.T.: 8 seconds; 4 replicates) at room temperature.
- 15 Protein conformation in most formulations could be effectively retained, with only slight changes in the region 260 – 280 nm (tyrosine and phenylalanine signals). However, a few exceptions could be observed, where more significant changes could be found which may indicate partial disruption/unfolding and loss of structure following thermal stress: DP5 (possible effect of the surfactant type present), DP8 and 9 (formulations in
- 20 citrate – phosphate buffer; possible effect of the buffer type and combination with other ingredients present).

3.4.2 Freeze – thawing

- 25 *Visible particles by visual inspection*

Repeated FT cycles were not observed to cause significant increase in visible particles. Some formulations presented fibers-like particles upon stress (not particulate/precipitate or other forms typically formulation – related).

- 30 *Turbidity by Nephelometry*

Upon freeze – thawing, no significant changes occur in the formulations tested. Most formulations are clear or slightly opalescent at time 0 and after stress (exception: DP3, 5, 8, opalescent solution range at time 0, with negligible changes after stress).

Sub-visible particles by light obscuration method

Particles ≥ 25 micron were well below the Pharmacopoeia limit of 600 particles / container (typically ≤ 100 particles).

- Particles ≥ 10 micron had larger counts, but still below the 6000 particles / container limit. DP8 and 9, in citrate-phosphate buffer, show higher counts than the others (still below the above limit) at time 0, with no further increase upon FT stress.

Total aggregates by SE – HPLC

- In the total aggregates by SE – HPLC before and after FT stress, minimal changes were observed for all formulations (total aggregates increased by 0.2 – 0.5% after 3 FT cycles).

3.5 Conclusion

- 15 In glycine buffer, the most suitable conditions for antibody stabilisation include:
low ionic strength (10 mM),
low pH (4.0 – 4.4),
Lysine (monohydrochloride), Dextrose, Sucrose and Sorbitol as stabilisers,
Preferred surfactants: Kolliphor ELP and Kollidon 12PF (Tween 80 to be possibly
20 avoided to due visible particles concerns).

In succinate buffer, the most suitable conditions for antibody stabilisation include:
low ionic strength (10 mM),
pH 5.0 – 5.1

- 25 Lysine (monohydrochloride), Dextrose, Sucrose or Sorbitol as stabilisers,
Preferred surfactants: Kolliphor ELP and Kollidon 12PF (Tween 80 to be possibly
avoided to due visible particles concerns).

- In citrate – phosphate buffer, the most suitable conditions for antibody stabilisation
30 include:
low ionic strength (10 – 30 mM),
low pH (4.0 – 4.5),
Lysine (monohydrochloride), Dextrose, Sucrose or Sorbitol as stabilisers,

Preferred surfactants: Kolliphor ELP and Kollidon 12PF (Tween 80 to be possibly avoided to due visible particles concerns).

In histidine buffer, the most suitable conditions for antibody stabilisation include:

5 low ionic strength (10 - 15 mM),

pH 5.0 – 5.1,

Dextrose, Sucrose, Lysine (monohydrochloride), Inositol, Sorbitol as stabilisers,

Preferred surfactants: Kolliphor ELP and Kollidon 12PF (Tween 80 to be possibly avoided to due visible particles concerns).

10

The most favourable formulations of Table 6 were found to be DP 2, 4, 7, and 10.