

ANDROGEN RECEPTOR MODULATORS AND USES THEREOF

RELATED APPLICATIONS

[0001] This application claims the benefit of U.S provisional patent application no. 61/305,082 entitled "ANDROGEN RECEPTOR MODULATORS AND USES THEREOF" filed on February 16, 2010; U.S provisional patent application no. 61/329,023 entitled "ANDROGEN RECEPTOR MODULATORS AND USES THEREOF" filed on April 28, 2010; and U.S provisional patent application no. 61/388,457 entitled "ANDROGEN RECEPTOR MODULATORS AND USES THEREOF" filed on September 30, 2010; all of which are incorporated by reference in their entirety.

FIELD OF THE INVENTION

[0002] Described herein are compounds, including pharmaceutically acceptable salts, solvates, metabolites, prodrugs thereof, methods of making such compounds, pharmaceutical compositions comprising such compounds, and methods of using such compounds to treat, prevent or diagnose diseases or conditions that are androgen receptor dependent or androgen receptor mediated.

BACKGROUND OF THE INVENTION

[0003] The androgen receptor ("AR") is a ligand-activated transcriptional regulatory protein that mediates induction of a variety of biological effects through its interaction with endogenous androgens. Endogenous androgens include steroids such as testosterone and dihydrotestosterone. Testosterone is converted to dihydrotestosterone by the enzyme 5 alpha-reductase in many tissues.

[0004] The actions of androgens with androgen receptors have been implicated in a number of diseases or conditions, such as androgen dependent cancers, virilization in women, and acne, among others. Compounds that diminish the effects of androgens with androgen receptors and/or lower the the concentrations of androgen receptors find use in the treatment of diseases or conditions in which androgen receptors play a role.

SUMMARY OF THE INVENTION

[0005] In one aspect, presented herein are compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) that diminish the effects of androgens with androgen receptors and/or lower the the concentrations of androgen receptors, and therefore, are useful as agents for the treatment or prevention of diseases or conditions in which androgen receptors play a role, such as diseases or conditions in which androgen receptors participate, are

involved in the etiology or pathology of the disease or condition, or contribute to at least one symptom of the disease or condition.


[0006] In one aspect, compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are useful for the treatment of benign prostate hyperplasia,

hirsutism, acne, adenomas and neoplasies of the prostate, benign or malignant tumor cells containing the androgen receptor, hyperpilosity, seborrhea, endometriosis, polycystic ovary syndrome, androgenic alopecia, hypogonadism, osteoporosis, suppression of spermatogenesis, libido, cachexia, anorexia, androgen supplementation for age related decreased testosterone levels, prostate cancer, breast cancer, endometrial cancer, uterine cancer, hot flashes, and

Kennedy's disease muscle atrophy and weakness, skin atrophy, bone loss, anemia, arteriosclerosis, cardiovascular disease, loss of energy, loss of well-being, type 2 diabetes and abdominal fat accumulation. In some embodiments, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is used in the treatment of prostate cancer. In some embodiments, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) are used in the treatment of hormone-sensitive prostate cancer. In some embodiments, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is used in the treatment of hormone refractory prostate cancer.

[0007] In one aspect, described herein are compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X), pharmaceutically acceptable salts, solvates, metabolites and prodrugs thereof. In some embodiments, compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are androgen receptor modulators. In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is an androgen receptor antagonist, an androgen receptor inverse agonist, an androgen receptor degrader, an androgen receptor trafficking modulator and/or an androgen receptor DNA-binding inhibitor. In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is an androgen receptor antagonist. In some embodiments, the compound of Formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX) or (X) is an androgen receptor inverse agonist. In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is an androgen receptor degrader. In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is an androgen receptor trafficking modulator. In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is an androgen receptor DNA-binding inhibitor.

[0008] In one aspect, provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (I)

- 5 wherein,
- ring A is monocyclic heteroaryl, bicyclic heteroaryl, or naphthyl;
- m is 0, 1, 2, 3 or 4;
- each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl;
- 10 each R¹ is independently selected from H, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-C₆fluoroalkyl;
- or both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₁₀cycloalkyl or a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;
- 20 each R² is H; or both R² are taken together with the carbon to which they are attached to form -C(=S)- or -C(=O)-;
- each R³ is H; or both R³ are taken together with the carbon to which they are attached to form -C(=S)- or -C(=O)-; provided that each R² is not H if each R³ is H;
- 25 ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;
- n is 0, 1, 2, 3 or 4;
- each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy,
- 30

substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-C₆heteroalkyl;

R⁵ is substituted or unsubstituted C₂-C₁₀alkyl, substituted or unsubstituted C₂-C₁₀fluoroalkyl, substituted or unsubstituted C₂-C₁₀alkoxy, substituted or unsubstituted C₂-C₁₀fluoroalkoxy, substituted or unsubstituted C₂-C₁₀heteroalkyl, substituted or unsubstituted C₂-C₁₀heterofluoroalkyl, or -L¹-L²-R⁶;

L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-, or -S(=O)₂NH-;

L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene;

R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl;

each R⁹ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or

two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;

R¹⁰ is substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-

C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl);

R¹¹ is H or C₁-C₄alkyl.

5 [0009] For any and all of the embodiments, substituents are selected from among from a subset of the listed alternatives. For example, in some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₁₀cycloalkyl. In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted cyclopropyl, cyclobutyl, cyclopentyl, or
10 cyclohexyl. In some embodiments, each R¹ is independently selected from H, -OH, C₁-C₆alkyl, and C₁-C₆fluoroalkyl. In some embodiments, each R¹ is independently selected from H, -CH₃ and -CF₃.


[0010] In some embodiments, both R² are taken together with the carbon to which they are attached to form -C(=S)-; both R³ are taken together with the carbon to which they are attached
15 to form -C(=O)-.

[0011] In some embodiments, ring A is C-linked monocyclic heteroaryl, C-linked bicyclic heteroaryl, or naphthyl.

[0012] In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₁₀cycloalkyl; both R² are taken together
20 with the carbon to which they are attached to form -C(=S)-; both R³ are taken together with the carbon to which they are attached to form -C(=O)-; ring A is *N*-containing monocyclic heteroaryl, or *N*-containing bicyclic heteroaryl; each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or
25 unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl.

[0013] In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; ring A is *N*-
30 containing monocyclic heteroaryl selected from pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, triazinyl, pyrrolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, imidazolyl, pyrazolyl, isothiazolyl, triazolyl, and tetrazolyl.

[0014] In some embodiments, ring A is pyridinyl, pyrimidinyl, pyridazinyl, or pyrazinyl. In some embodiments, ring A is pyridinyl.



[0016] In some embodiments, ring A is pyridinyl; each R^A is independently selected from H, halogen, -CN, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl.

[0017] In some embodiments, both R^1 are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; ring A is *N*-containing bicyclic heteroaryl selected from quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, indolyl, indazolyl, benzoxazolyl, benzisoxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzimidazolyl, purinyl, cinnolinyl, phthalazinyl, pteridinyl, pyridopyrimidinyl, pyrazolopyrimidinyl, azaindolyl, pyrazolopyridinyl, thiazolopyrimidinyl, triazolopyridazinyl, thiazolopyridinyl, pyridothienyl, pyrimidothienyl and pyrrolopyrimidinyl; each R^A is independently selected from H, halogen, -CN, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl.

[0018] In some embodiments, ring A is *N*-containing bicyclic heteroaryl selected from quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, pyridopyrimidinyl, pyrazolopyrimidinyl, triazolopyridazinyl, and pyrrolopyrimidinyl.

[0019] In some embodiments, ring A is [1,2,4]triazolo[4,3-*b*]pyridazinyl.

[0020] In some embodiments, ring B is phenyl or monocyclic heteroaryl; each R^4 is independently selected from H, halogen, -CN, -NO₂, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-

C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-C₆heteroalkyl; R⁵ is substituted or unsubstituted C₂-C₁₀alkyl, substituted or unsubstituted C₂-C₁₀fluoroalkyl, substituted or unsubstituted C₂-C₁₀alkoxy, substituted or unsubstituted C₂-C₁₀fluoroalkoxy, substituted or unsubstituted C₂-C₁₀heteroalkyl, substituted or unsubstituted C₂-C₁₀heterofluoroalkyl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHS(=O)₂-, or -S(=O)₂NH-; L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene; R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.


[0021] In some embodiments, ring B is phenyl or monocyclic heteroaryl; each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, substituted or unsubstituted C₁-C₄alkyl, substituted or unsubstituted C₁-C₄fluoroalkyl, substituted or unsubstituted C₁-C₄fluoroalkoxy, and substituted or unsubstituted C₁-C₄alkoxy; R⁵ is substituted or unsubstituted C₂-C₁₀alkyl, substituted or unsubstituted C₂-C₁₀fluoroalkyl, substituted or unsubstituted C₂-C₁₀alkoxy, substituted or unsubstituted C₂-C₁₀fluoroalkoxy, substituted or unsubstituted C₂-C₁₀heteroalkyl, substituted or unsubstituted C₂-C₁₀heterofluoroalkyl, or -L¹-L²-R⁶; L¹ is absent, -O-, or -C(=O)NH-; L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

[0022] In some embodiments, L² is substituted or unsubstituted C₁-C₆alkylene; R⁶ is -CN, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

[0023] In some embodiments, R⁶ is substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

[0024] In some embodiments, L^2 is substituted or unsubstituted C_1 - C_6 alkylene; R^6 is -CN, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, or -C(=O)N(R⁹)₂; each R⁹ is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, - C_1 - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), - C_1 - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), - C_1 - C_4 alkylene-(substituted or unsubstituted aryl), and - C_1 - C_4 alkylene-(substituted or unsubstituted heteroaryl); or two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl; R¹⁰ is substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, - C_1 - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), - C_1 - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), - C_1 - C_4 alkylene-(substituted or unsubstituted aryl), or - C_1 - C_4 alkylene-(substituted or unsubstituted heteroaryl).

[0025] In some embodiments, provided is a compound of Formula (Ia), or a pharmaceutically acceptable salt, or *N*-oxide thereof:



Formula (Ia)

wherein,

ring A is monocyclic heteroaryl, bicyclic heteroaryl, or naphthyl;

m is 1, 2, 3 or 4;

each R^A is independently H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, C_1 - C_6 alkoxy, or C_1 - C_6 heteroalkyl;

both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C_3 - C_6 cycloalkyl or a substituted or unsubstituted C_2 - C_6 heterocycloalkyl;

or each R¹ is independently selected from H, -OH, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, and C_1 - C_6 fluoroalkyl;

X is S or O;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, 2, 3 or 4;

each R^4 is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -

CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -

NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, C₁-

C₆alkoxy, and C₁-C₆heteroalkyl;

R^5 is -L¹-L²-R⁶ or -L¹-R⁷;

L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -

NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -

NHS(=O)₂-, or -S(=O)₂NH-;

L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene;

R^6 is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -

S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -

-OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, C₁-

C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆heteroalkyl, substituted or unsubstituted C₃-

C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or

unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic

heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted

naphthyl;

R^7 is substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, or substituted or

unsubstituted monocyclic heteroaryl;

each R^9 is independently selected from H, C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-

C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted

monocyclic heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₆cycloalkyl), -

C₁-C₄alkylene-(substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl), -C₁-

C₄alkylene-(substituted or unsubstituted phenyl), and -C₁-C₄alkylene-(substituted or

unsubstituted monocyclic heteroaryl); or

two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted monocyclic C₂-

C₆heterocycloalkyl;

R^{10} is C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-

C₆cycloalkyl, a substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, a

substituted or unsubstituted phenyl, a substituted or unsubstituted benzyl, a substituted

or unsubstituted monocyclic heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-

C₆cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted monocyclic C₂-

C₆heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted phenyl), or -C₁-C₄alkylene-(substituted or unsubstituted monocyclic heteroaryl);

R¹¹ is H or C₁-C₄alkyl.

[0026] In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₆cycloalkyl; X is S; ring A is *N*-containing monocyclic heteroaryl; each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, and C₁-C₆alkoxy; ring B is phenyl or monocyclic heteroaryl.

[0027] In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; ring A is pyridinyl, pyrimidinyl, pyridazinyl, or pyrazinyl.

[0028] In some embodiments, ring A is pyridinyl; each R^A is independently selected from H, halogen, -CN, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, and C₁-C₆alkoxy.

[0029] In some embodiments, ring B is phenyl; R⁵ is -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHS(=O)₂-, or -S(=O)₂NH-; L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene; R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

[0030] In some embodiments, ring B is phenyl; each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, and C₁-C₄alkoxy; R⁵ is -L¹-L²-R⁶; L¹ is absent, -O-, or -C(=O)NH-; L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

[0031] In some embodiments, L² is C₁-C₆alkylene; R⁶ is -CN, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

[0032] In some embodiments, R^6 is substituted or unsubstituted C_3 - C_6 cycloalkyl, substituted or unsubstituted C_2 - C_6 heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.


[0033] In some embodiments, L^2 is C_1 - C_6 alkylene; R^6 is a substituted or unsubstituted C_2 -
5 C_6 heterocycloalkyl.

[0034] In some embodiments, ring B is phenyl; each R^4 is independently selected from H, halogen, $-CN$, $-NO_2$, $-OH$, C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, C_1 - C_4 fluoroalkoxy, and C_1 - C_4 alkoxy; R^5 is $-L^1-R^7$; L^1 is absent, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, $-NHC(=O)O-$, $-NHC(=O)NH-$, $-C(=O)O-$, $-OC(=O)-$, $-OC(=O)O-$, $-OC(=O)NH-$, $-NHS(=O)_2-$,
10 or $-S(=O)_2NH-$; R^7 is substituted or unsubstituted monocyclic C_2 - C_6 heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl;

[0035] In some embodiments, R^5 is $-L^1-R^7$; L^1 is $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$; R^7 is substituted or unsubstituted monocyclic C_2 -
 C_6 heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.


[0036] In some embodiments, R^5 is $-L^1-R^7$; L^1 is absent, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$; R^7 is substituted or unsubstituted monocyclic heteroaryl.

[0037] In some embodiments, the compound of Formula (I) or Formula (Ia) has the structure of Formula (II):




Formula (II).

[0038] In some embodiments, the compound of Formula (Ia) has the structure of Formula (II):



Formula (II)

wherein,



m is 2;

one R^A is -CN, -NO₂, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, or -

C(=O)N(R⁹)₂; and the other R^A is H, halogen, -OH, C₁-C₆alkyl, C₁-C₆fluoroalkyl,

C₁-C₆fluoroalkoxy, or C₁-C₆alkoxy;

n is 0 or 1;

each R^4 is independently selected from H, halogen, -CN, -NO₂, -OH, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, and C₁-C₆alkoxy;

R^5 is -L¹-L²-R⁶ or -L¹-R⁷;


L¹ is absent, -O-, or -C(=O)NH-;

L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene;

R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -

N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl;

R⁷ is substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.



[0039] In some embodiments, one R^A is -CN and the other R^A is

H, halogen, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, or C₁-C₄alkoxy; each R^4 is

independently selected from H, halogen, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy,

and C₁-C₄alkoxy; R^5 is -L¹-L²-R⁶ or -L¹-R⁷; L¹ is absent, -O-, or -C(=O)NH-; L² is C₁-

C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -


S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-

C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted

monocyclic heteroaryl, or substituted or unsubstituted phenyl; R⁷ is substituted or unsubstituted

monocyclic C₂-C₆heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.

[0040] In some embodiments, described herein is a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, N-oxide, metabolite or prodrug thereof.



Formula (I)

wherein,

ring A is bicyclic heteroaryl, or naphthyl;

m is 0, 1, 2, or 3;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -

CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or

unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy,

substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted C₁-C₆heteroalkyl,

substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-

C₁₀heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl;

each R^1 is independently selected from H, -OH, substituted or unsubstituted C₁-C₆alkyl,

substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-

C₆fluoroalkyl;

or both R^1 are taken together with the carbon atom to which they are attached to form a

substituted or unsubstituted C₃-C₁₀cycloalkyl or a substituted or unsubstituted C₂-

C₁₀heterocycloalkyl;

each R^2 is H; or both R^2 are taken together with the carbon to which they are attached to

form -C(=S)- or -C(=O)-;

each R^3 is H; or both R^3 are taken together with the carbon to which they are attached to

form -C(=S)- or -C(=O)-; provided that each R^2 is not H if each R^3 is H;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

each R^4 is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -

CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -

NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₆alkyl, substituted

or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy,

substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-

C₆heteroalkyl;

R^5 is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₁₀alkyl, substituted or unsubstituted C₁-C₁₀fluoroalkyl, substituted or unsubstituted C₁-C₁₀fluoroalkoxy, C₁-C₁₀alkoxy, substituted or unsubstituted C₁-C₁₀heteroalkyl, or -L¹-L²-R⁶;

L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-, or -S(=O)₂NH-;

L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene;

R^6 is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl;

each R⁹ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or

two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;

R^{10} is substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-

C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl);

R¹¹ is H or C₁-C₄alkyl.


[0041] In some embodiments, ring A is selected from quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, indolyl, indazolyl, benzoxazolyl, benzisoxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzimidazolyl, purinyl, cinnolinyl, phthalazinyl, pteridinyl, pyridopyrimidinyl, pyrazolopyrimidinyl, azaindolyl, pyrazolopyridinyl, thiazolopyrimidinyl, triazolopyridazinyl, thiazolopyridinyl, pyridothienyl, pyrimidothienyl, pyrrolopyrimidinyl and naphthyl.

[0042] In some embodiments, ring A is selected from quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, pyridopyrimidinyl, pyrazolopyrimidinyl, triazolopyridazinyl, pyrrolopyrimidinyl, and naphthyl.

[0043] In some embodiments, ring A is [1,2,4]triazolo[4,3-b]pyridazinyl.

[0044] In some embodiments, described herein is a compound of Formula (I), or a

pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (I)

wherein,

ring A is a 5-membered heteroaryl;

m is 0, 1, 2, or 3;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -

CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₄alkyl, substituted or

unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy,

substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted C₁-C₆heteroalkyl,

substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-

C₁₀heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl;

each R¹ is independently selected from H, -OH, substituted or unsubstituted C₁-C₆alkyl,

substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-

C₆fluoroalkyl;

or both R^1 are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C_3 - C_{10} cycloalkyl or a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

each R^2 is H; or both R^2 are taken together with the carbon to which they are attached to form $-C(=S)-$ or $-C(=O)-$;

each R^3 is H; or both R^3 are taken together with the carbon to which they are attached to form $-C(=S)-$ or $-C(=O)-$; provided that each R^2 is not H if each R^3 is H;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

each R^4 is independently selected from H, halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, -

$S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, -

CO_2R^9 , $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, -

$NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_1 - C_6 alkyl, substituted

or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy,

substituted or unsubstituted C_1 - C_6 alkoxy, and substituted or unsubstituted C_1 -

C_6 heteroalkyl;

R^5 is H, halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, -

$S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, -

$OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or

unsubstituted C_1 - C_{10} alkyl, substituted or unsubstituted C_1 - C_{10} fluoroalkyl, substituted or

unsubstituted C_1 - C_{10} fluoroalkoxy, substituted or unsubstituted C_1 - C_{10} alkoxy, substituted

or unsubstituted C_1 - C_{10} heteroalkyl, or $-L^1-L^2-R^6$;

L^1 is absent, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, -

$NHC(=O)O-$, $-NHC(=O)NH-$, $-C(=O)O-$, $-OC(=O)-$, $-OC(=O)O-$, $-OC(=O)NH-$, -

$NHS(=O)_2-$, or $-S(=O)_2NH-$;

L^2 is substituted or unsubstituted C_1 - C_6 alkylene, substituted or unsubstituted C_1 -

C_6 fluoroalkylene or substituted or unsubstituted C_1 - C_6 heteroalkylene;

R^6 is halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, -

$S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, -

$-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted

or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl,

substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_3 -

C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or

unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic

heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl;


each R⁹ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or

two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;


R¹⁰ is substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl);

R¹¹ is H or C₁-C₄alkyl.

[0045] In some embodiments, ring A is selected from pyrrolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, imidazolyl, pyrazolyl, isothiazolyl, triazolyl, and tetrazolyl.



[0046] In some embodiments, is selected from:




---^A , ---^A , and ---^A ; where X is O, S, or NR^A. In some embodiments, X is O. In some embodiments, X is S. In some embodiments, X is NR^A.

[0047] In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a C₃-C₁₀cycloalkyl; both R² are taken together with the carbon to which

they are attached to form $-C(=S)-$; both R^3 are taken together with the carbon to which they are attached to form $-C(=O)-$.

[0048] In some embodiments, the compound described herein has the structure of Formula (III):



Formula (III)

wherein,

p is 0, 1, 2, or 3.

[0049] In some embodiments, each R^A is independently selected from H, halogen, $-CN$, $-NO_2$, $-OH$, $-S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-CO_2R^9$, $-C(=O)N(R^9)_2$, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, substituted or unsubstituted C_1 - C_6 alkoxy, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl.


[0050] In some embodiments, each R^4 is independently selected from H, halogen, $-CN$, $-NO_2$, $-OH$, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, and substituted or unsubstituted C_1 - C_6 alkoxy; R^5 is substituted or unsubstituted C_2 - C_{10} alkyl, substituted or unsubstituted C_2 - C_{10} fluoroalkyl, substituted or unsubstituted C_2 - C_{10} alkoxy, substituted or unsubstituted C_2 - C_{10} fluoroalkoxy, substituted or unsubstituted C_2 - C_{10} heteroalkyl, substituted or unsubstituted C_2 -

C_{10} heterofluoroalkyl, or $-L^1-L^2-R^6$; L^1 is absent, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, $-NHS(=O)_2-$, or $-S(=O)_2NH-$; L^2 is substituted or unsubstituted C_1 - C_6 alkylene, substituted or unsubstituted C_1 - C_6 fluoroalkylene or substituted or unsubstituted C_1 - C_6 heteroalkylene; R^6 is halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-CO_2R^9$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

[0051] In some embodiments, each R^4 is independently selected from H, halogen, $-CN$, $-NO_2$, $-OH$, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, and substituted or unsubstituted C_1 - C_6 alkoxy;

R^5 is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₁₀alkyl, substituted or unsubstituted C₁-C₁₀fluoroalkyl, substituted or unsubstituted C₁-C₁₀fluoroalkoxy, substituted or unsubstituted C₁-C₁₀alkoxy, substituted or unsubstituted C₁-C₁₀heteroalkyl, or -L¹-L²-R⁶; L¹ is absent, -O-, or -C(=O)NH-; L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

[0052] In some embodiments, described herein is a compound of Formula (IV), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (IV)

wherein,

each R¹ is independently selected from H, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-C₆fluoroalkyl;

or both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₁₀cycloalkyl or a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;

X¹ is CR^A or N;

X² is CR^A or N;

X³ is CR^A or N; provided that at least two of X¹, X², and X³ is CR^A;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or

unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

R⁴ is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, or substituted or unsubstituted C₁-C₆heteroalkyl;

R⁵ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₂-C₁₀alkyl, substituted or unsubstituted C₂-C₁₀fluoroalkyl, substituted or unsubstituted C₁-C₁₀fluoroalkoxy, substituted or unsubstituted C₁-C₁₀alkoxy, substituted or unsubstituted C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶;

L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-, or -S(=O)₂NH-;

L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene;





R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl;


each R⁹ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-





(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;






R¹⁰ is substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl);


R¹¹ is H or C₁-C₄alkyl.

[0053] In some embodiments,  is , , ,


or ; m is 0, 1, 2, 3, or 4.

[0054] In some embodiments,  is , , ,

, , , , , or

.

[0055] In some embodiments, the compound of Formula (IV) has the structure of Formula (V):




Formula (V).

- [0056] In some embodiments, ring B is phenyl; R^4 is H, halogen, -CN, -OH, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, or substituted or unsubstituted C_1 - C_6 alkoxy; R^5 is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, substituted or unsubstituted C_2 - C_{10} alkyl, substituted or unsubstituted C_2 - C_{10} fluoroalkyl, substituted or unsubstituted C_1 - C_{10} fluoroalkoxy, substituted or unsubstituted C_1 - C_{10} alkoxy, substituted or unsubstituted C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, or -S(=O)₂NH-; L² is substituted or unsubstituted C_1 - C_6 alkylene, substituted or unsubstituted C_1 - C_6 fluoroalkylene or substituted or unsubstituted C_1 - C_6 heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl.


- [0057] In some embodiments, each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, or substituted or unsubstituted C_1 - C_6 alkoxy.

[0058] In some embodiments, the compound of Formula (I), (Ia), (II), (IV) or (V) has the structure of Formula (VI):



Formula (VI).

- [0059] In some embodiments, the compound of Formula (I), (Ia), (II), (IV) or (V) has the structure of Formula (VI):



Formula (VI)

wherein,

one R^A is $-\text{CN}$, $-\text{NO}_2$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{R}^{10}$, $-\text{CO}_2\text{R}^9$, or -

$\text{C}(=\text{O})\text{N}(\text{R}^9)_2$; and the other R^A is H, halogen, $-\text{OH}$, $\text{C}_1\text{-C}_6\text{alkyl}$, $\text{C}_1\text{-C}_6\text{fluoroalkyl}$,

$\text{C}_1\text{-C}_6\text{fluoroalkoxy}$, or $\text{C}_1\text{-C}_6\text{alkoxy}$;

both R^1 are taken together with the carbon atom to which they are attached to form a $\text{C}_3\text{-C}_6\text{cycloalkyl}$;


or each R^1 is independently $\text{C}_1\text{-C}_4\text{alkyl}$;

R^4 is H, halogen, $-\text{OH}$, $\text{C}_1\text{-C}_4\text{alkyl}$, $\text{C}_1\text{-C}_4\text{fluoroalkyl}$, $\text{C}_1\text{-C}_4\text{fluoroalkoxy}$, or $\text{C}_1\text{-C}_4\text{alkoxy}$.

[0060] In some embodiments, R^5 is $-\text{L}^1\text{-L}^2\text{-R}^6$ or $-\text{L}^1\text{-R}^7$; L^1 is absent, $-\text{O}-$, or $-\text{C}(=\text{O})\text{NH}-$; L^2 is $\text{C}_1\text{-C}_6\text{alkylene}$, $\text{C}_1\text{-C}_6\text{fluoroalkylene}$ or $\text{C}_1\text{-C}_6\text{heteroalkylene}$; R^6 is $-\text{CN}$, $-\text{NO}_2$, $-\text{OH}$, $-\text{OR}^9$, $-\text{SR}^9$, $-\text{S}(=\text{O})\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{C}(=\text{O})\text{R}^{10}$, $-\text{CO}_2\text{R}^9$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, substituted or unsubstituted $\text{C}_3\text{-C}_6\text{cycloalkyl}$, substituted or unsubstituted $\text{C}_2\text{-C}_6\text{heterocycloalkyl}$, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl; R^7 is substituted or unsubstituted monocyclic $\text{C}_2\text{-C}_6\text{heterocycloalkyl}$, or substituted or unsubstituted monocyclic heteroaryl.

[0061] In some embodiments, one R^A is $-\text{CN}$, $-\text{NO}_2$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{R}^{10}$, $-\text{CO}_2\text{R}^9$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl; and the other R^A is H, halogen, $-\text{OH}$, substituted or unsubstituted $\text{C}_1\text{-C}_6\text{alkyl}$, substituted or unsubstituted $\text{C}_1\text{-C}_6\text{fluoroalkyl}$, substituted or unsubstituted $\text{C}_1\text{-C}_6\text{fluoroalkoxy}$, or substituted or unsubstituted $\text{C}_1\text{-C}_6\text{alkoxy}$; R^4 is H, halogen, $-\text{CN}$, $-\text{OH}$, substituted or unsubstituted $\text{C}_1\text{-C}_6\text{alkyl}$, substituted or unsubstituted $\text{C}_1\text{-C}_6\text{fluoroalkyl}$, substituted or unsubstituted $\text{C}_1\text{-C}_6\text{fluoroalkoxy}$, or substituted or unsubstituted $\text{C}_1\text{-C}_6\text{alkoxy}$; R^5 is halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{OH}$, $-\text{OR}^9$, $-\text{SR}^9$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{R}^{10}$, $-\text{CO}_2\text{R}^9$, substituted or unsubstituted $\text{C}_2\text{-C}_{10}\text{alkyl}$, substituted or unsubstituted $\text{C}_2\text{-C}_{10}\text{fluoroalkyl}$, substituted or unsubstituted $\text{C}_1\text{-C}_{10}\text{fluoroalkoxy}$, substituted or unsubstituted $\text{C}_1\text{-C}_{10}\text{alkoxy}$, substituted or unsubstituted $\text{C}_1\text{-C}_{10}\text{heteroalkyl}$, substituted or unsubstituted $\text{C}_3\text{-C}_{10}\text{cycloalkyl}$, substituted or unsubstituted $\text{C}_2\text{-C}_{10}\text{heterocycloalkyl}$, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or $-\text{L}^1\text{-L}^2\text{-R}^6$; L^1 is absent, $-\text{O}-$, or $-\text{C}(=\text{O})\text{NH}-$; L^2 is substituted or unsubstituted $\text{C}_1\text{-C}_6\text{alkylene}$, substituted or unsubstituted $\text{C}_1\text{-C}_6\text{fluoroalkylene}$ or substituted or unsubstituted $\text{C}_1\text{-C}_6\text{heteroalkylene}$; R^6 is $-\text{CN}$, $-\text{NO}_2$, $-\text{OH}$, $-\text{OR}^9$, $-\text{SR}^9$, $-\text{S}(=\text{O})\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{CO}_2\text{R}^9$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

[0062] In some embodiments, described herein is a compound of Formula (VII), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (VII)

wherein,

each R^1 is independently selected from H, -OH, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 alkoxy, and substituted or unsubstituted C_1 - C_6 fluoroalkyl;

or both R^1 are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C_3 - C_{10} cycloalkyl or a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, substituted or unsubstituted C_1 - C_6 alkoxy, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl;

ring B is 5-membered heteroaryl, bicyclic heteroaryl or naphthyl;

n is 0, 1, or 2;

R^4 is H, halogen, -CN, -NO₂, -OH, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, or substituted or unsubstituted C_1 - C_6 alkoxy;

R^5 is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C_2 - C_{10} alkyl, substituted or unsubstituted C_2 - C_{10} fluoroalkyl, substituted or unsubstituted C_1 - C_{10} fluoroalkoxy, substituted or unsubstituted C_1 - C_{10} alkoxy, substituted or unsubstituted C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶;

L^1 is absent, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, $-NHC(=O)O-$, $-NHC(=O)NH-$, $-C(=O)O-$, $-OC(=O)-$, $-OC(=O)O-$, $-OC(=O)NH-$, $-NHS(=O)_2-$, or $-S(=O)_2NH-$;

L^2 is substituted or unsubstituted C_1 - C_6 alkylene, substituted or unsubstituted C_1 - C_6 fluoroalkylene or substituted or unsubstituted C_1 - C_6 heteroalkylene;


R^6 is halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl;

each R^9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), and $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl); or two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

R^{10} is substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), or $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl);

R^{11} is H or C_1 - C_4 alkyl.

[0063] In some embodiments, described herein is a compound of Formula (VIII), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (VIII)

wherein,

each R^1 is independently selected from H, -OH, substituted or unsubstituted C_1 - C_6 alkyl,
 5 substituted or unsubstituted C_1 - C_6 alkoxy, and substituted or unsubstituted C_1 - C_6 fluoroalkyl;

or both R^1 are taken together with the carbon atom to which they are attached to form a
 substituted or unsubstituted C_3 - C_6 cycloalkyl or a substituted or unsubstituted C_2 - C_6 heterocycloalkyl;

10 X is O or S;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -
 S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -
 CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 heteroalkyl, substituted
 or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl,
 15 substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic
 heteroaryl;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

R^4 is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, substituted or
 20 unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted
 or unsubstituted C_1 - C_6 fluoroalkoxy, substituted or unsubstituted C_1 - C_6 alkoxy, or
 substituted or unsubstituted C_1 - C_6 heteroalkyl;

R^5 is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰,
 -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -
 25 C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -
 NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C_1 - C_{10} alkyl, substituted or
 unsubstituted C_1 - C_{10} fluoroalkyl, substituted or unsubstituted C_1 - C_{10} fluoroalkoxy,
 substituted or unsubstituted C_1 - C_{10} alkoxy, substituted or unsubstituted C_1 -
 C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or
 30 unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl,
 substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶;

L^1 is absent, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, $-NHC(=O)O-$, $-NHC(=O)NH-$, $-C(=O)O-$, $-OC(=O)-$, $-OC(=O)O-$, $-OC(=O)NH-$, $-NHS(=O)_2-$, or $-S(=O)_2NH-$;

L^2 is substituted or unsubstituted C_1 - C_6 alkylene, substituted or unsubstituted C_1 - C_6 fluoroalkylene or substituted or unsubstituted C_1 - C_6 heteroalkylene;

R^6 is $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl;

each R^9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), and $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl); or two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

R^{10} is substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), or $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl);


R^{11} is H or C_1 - C_4 alkyl.

[0064] In some embodiments, X is S; ring B is phenyl; R^4 is H, halogen, $-CN$, $-OH$, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, or C_1 - C_6 alkoxy; R^5 is halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-CO_2R^9$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, substituted or unsubstituted C_1 - C_{10} alkyl, substituted or unsubstituted C_1 - C_{10} fluoroalkyl,

substituted or unsubstituted C₁-C₁₀fluoroalkoxy, substituted or unsubstituted C₁-C₁₀alkoxy, substituted or unsubstituted C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, or -S(=O)₂NH-; L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl.

[0065] In some embodiments, each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, C₁-C₆alkyl, and C₁-C₆alkoxy.

[0066] In some embodiments, described herein is a compound of Formula (VIIIa), or a pharmaceutically acceptable salt, or *N*-oxide thereof:



Formula (VIIIa)

wherein,

both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₆cycloalkyl or a substituted or unsubstituted C₂-C₆heterocycloalkyl;

or

each R¹ is independently H, -OH, C₁-C₆alkyl, C₁-C₆alkoxy, or C₁-C₆fluoroalkyl;

X is O or S;

R^A is C₁-C₆alkyl;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

R⁴ is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, C₁-C₆alkoxy, or C₁-C₆heteroalkyl;

R⁵ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -

$C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, C_1 - C_{10} alkyl, C_1 - C_{10} fluoroalkyl, C_1 - C_{10} fluoroalkoxy, C_1 - C_{10} alkoxy, C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or $-L^1-L^2-R^6$;

L^1 is absent, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, $-NHC(=O)O-$, $-NHC(=O)NH-$, $-C(=O)O-$, $-OC(=O)-$, $-OC(=O)O-$, $-OC(=O)NH-$, $-NHS(=O)_2-$, or $-S(=O)_2NH-$;

L^2 is C_1 - C_6 alkylene, C_1 - C_6 fluoroalkylene or C_1 - C_6 heteroalkylene;

R^6 is $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl;

each R^9 is independently selected from H, C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), and $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl); or

two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

R^{10} is C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), or $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl);


R^{11} is H or C_1 - C_4 alkyl.

[0067] In some embodiments, X is S; ring B is phenyl; R^4 is H, halogen, $-CN$, $-OH$, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, or C_1 - C_6 alkoxy; R^5 is halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, -

SR^9 , $-\text{S}(=\text{O})\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{R}^{10}$, $-\text{CO}_2\text{R}^9$, $-\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, C_1 - C_{10} alkyl, C_1 - C_{10} fluoroalkyl, C_1 - C_{10} fluoroalkoxy, C_1 - C_{10} alkoxy, C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or $-\text{L}^1$ - L^2 - R^6 ; L^1 is absent, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$, $-\text{NH}-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{O})\text{NH}-$, or $-\text{S}(=\text{O})_2\text{NH}-$; L^2 is C_1 - C_6 alkylene, C_1 - C_6 fluoroalkylene or C_1 - C_6 heteroalkylene; R^6 is $-\text{CN}$, $-\text{NO}_2$, $-\text{OH}$, $-\text{OR}^9$, $-\text{SR}^9$, $-\text{S}(=\text{O})\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{CO}_2\text{R}^9$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl.

[0068] In some embodiments, R^A is C_1 - C_6 alkyl; both R^1 are taken together with the carbon atom to which they are attached to form a C_3 - C_6 cycloalkyl; or each R^1 is independently C_1 - C_4 alkyl.


[0069] In some embodiments, the compound of Formula (VIII) has the structure of Formula (IX):



Formula (IX).

[0070] In some embodiments, each R^A is independently selected from H, halogen, $-\text{OH}$, C_1 - C_6 alkyl, and C_1 - C_6 alkoxy.

[0071] In some embodiments, the compound of Formula (VIII) or Formula (VIIIa) has the structure of Formula (IXa):



Formula (IXa).


[0072] In some embodiments, each R^9 is independently selected from H, C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_6 cycloalkyl, a substituted or unsubstituted C_2 - C_6 heterocycloalkyl, a substituted or unsubstituted phenyl, a substituted or unsubstituted benzyl, and a substituted or unsubstituted monocyclic heteroaryl; or two R^9 groups attached to the same N atom are taken together with the N atom to which they are

attached to form a substituted or unsubstituted C₂-C₆heterocycloalkyl; R¹⁰ is C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₆cycloalkyl, a substituted or unsubstituted C₂-C₆heterocycloalkyl, a substituted or unsubstituted phenyl, or a substituted or unsubstituted monocyclic heteroaryl.

- 5 [0073] In some embodiments, R⁵ is halogen, -CN, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)NH(R⁹), C₁-C₁₀alkyl, C₁-C₁₀fluoroalkyl, C₁-C₁₀fluoroalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, or -S(=O)₂NH-; L² is C₁-C₆alkylene; R⁶ is -CN, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

- 15 [0074] In some embodiments, R^A is -CH₃; both R¹ are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; or each R¹ is -CH₃; R⁵ is halogen, -CN, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)NH(R⁹), C₁-C₁₀alkyl, C₁-C₁₀fluoroalkyl, C₁-C₁₀fluoroalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, or -S(=O)₂NH-; L² is C₁-C₆alkylene; R⁶ is substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

- 25 [0075] In some embodiments, described herein is a compound of Formula (X), or a pharmaceutically acceptable salt, solvate, N-oxide, metabolite or prodrug thereof:



wherein,

each R¹ is independently selected from H and -CH₃;

- 30 or both R¹ are taken together with the carbon atom to which they are attached to form a cyclobutyl;

X is O or S;


X^1 is CH or N;

R^A is $-\text{CN}$ or $-\text{C}(=\text{O})\text{NH}_2$;

R^5 is $-\text{CO}_2\text{H}$, or $-\text{C}(=\text{O})\text{NH}_2$.

[0076] In some embodiments, described herein is a compound of Formula (X), or a

5 pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



wherein,

each R^1 is independently selected from H and $-\text{CH}_3$;

10 or both R^1 are taken together with the carbon atom to which they are attached to form a cyclobutyl;

X is O;


X^1 is CH or N;

R^A is $-\text{CN}$ or $-\text{C}(=\text{O})\text{NH}_2$;

15 R^5 is $-\text{CO}_2\text{H}$, $-\text{C}(=\text{O})\text{NH}_2$ or $-\text{C}(=\text{O})\text{NH}(\text{CH}_3)$.

[0077] In some embodiments, described herein is a compound of Formula (X), or a

pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



20 wherein,

each R^1 is independently selected from H and $-\text{CH}_3$;

or both R^1 are taken together with the carbon atom to which they are attached to form a cyclobutyl;

X is O or S;

25 X^1 is CH or N;

R^A is $-\text{C}(=\text{O})\text{NH}_2$;

R^5 is $-\text{CO}_2\text{H}$, $-\text{C}(=\text{O})\text{NH}_2$ or $-\text{C}(=\text{O})\text{NH}(\text{CH}_3)$.

[0078] Throughout the specification, groups and substituents thereof can be chosen by one skilled in the field to provide stable moieties and compounds.

[0079] Compounds disclosed herein are selective androgen receptor modulators. In some embodiments, compounds disclosed herein have high specificity for the androgen receptor and have desirable, tissue-selective pharmacological activities. Desirable, tissue-selective pharmacological activities include, but are not limited to, AR antagonist activity in prostate cells and no AR antagonist activity in non-prostate cells. In some embodiments, compounds disclosed herein are antiandrogens that display negligible or no AR agonist activity.

[0080] In some embodiments, presented herein are compounds selected from active metabolites, tautomers, pharmaceutically acceptable solvates, pharmaceutically acceptable salts or prodrugs of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X).

[0081] In some embodiments, provided is a pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X). In some embodiments, the pharmaceutical composition also contains at least one pharmaceutically acceptable inactive ingredient. In some embodiments, the pharmaceutical composition is formulated for intravenous injection, subcutaneous injection, oral administration, or topical administration. In some embodiments, the pharmaceutical composition is a tablet, a pill, a capsule, a liquid, a suspension, a gel, a colloid, a dispersion, a suspension, a solution, an emulsion, an ointment, or a lotion.

[0082] In some embodiments, the pharmaceutical composition further comprises one or more additional therapeutically active agents selected from: corticosteroids, anti-emetic agents, analgesics, anti-cancer agents, anti-inflammatories, kinase inhibitors, HSP90 inhibitors, histone deacetylase (HDAC) inhibitors.

[0083] In some embodiments, provided is a method comprising administering a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) to a human with a disease or condition that is androgen receptor mediated or androgen receptor dependent. In some embodiments, the human is already being administered one or more additional therapeutically active agents other than a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X). In some embodiments, the method further comprises administering one or more additional therapeutically active agents other than a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X).

[0084] In some embodiments, the one or more additional therapeutically active agents other than a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) are selected from: corticosteroids, anti-emetic agents, analgesics, anti-cancer

agents, anti-inflammatories, kinase inhibitors, HSP90 inhibitors, histone deacetylase (HDAC) inhibitors.

[0085] In some embodiments, described herein is a method of treating an androgen receptor dependent or androgen receptor mediated disease or condition in mammal comprising administering to the mammal a therapeutically effective amount of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X), or a pharmaceutically acceptable salt thereof or N-oxide thereof. In some embodiments, the androgen receptor dependent or androgen receptor mediated disease or condition is benign prostate hyperplasia, hirsutism, acne, adenomas and neoplasies of the prostate, benign or malignant tumor cells containing the androgen receptor, hyperpilosity, seborrhea, endometriosis, polycystic ovary syndrome, androgenic alopecia, hypogonadism, osteoporosis, suppression of spermatogenesis, libido, cachexia, anorexia, androgen supplementation for age related decreased testosterone levels, prostate cancer, breast cancer, endometrial cancer, uterine cancer, hot flashes, and Kennedy's disease muscle atrophy and weakness, skin atrophy, bone loss, anemia, arteriosclerosis, cardiovascular disease, loss of energy, loss of well-being, type 2 diabetes or abdominal fat accumulation.

[0086] In some embodiments, described herein is a method of treating cancer in a mammal comprising administering to the mammal a therapeutically effective amount of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X), or a pharmaceutically acceptable salt thereof or N-oxide thereof. In some embodiments, the cancer is a hormone dependent cancer. In some embodiments, the hormone dependent cancer is an androgen receptor dependent cancer. In some embodiments, the cancer is prostate cancer. In some embodiments, the cancer is hormone refractory prostate cancer. In some embodiments, the method of treating cancer further comprises administering to the mammal at least one additional anti-cancer agent.

[0087] In some embodiments, described herein is a method of treating cancer in a mammal comprising administering to the mammal a therapeutically effective amount of a compound, wherein the compound is: an androgen receptor inverse agonist; androgen receptor antagonist; androgen receptor degrader; androgen receptor trafficking modulator; androgen receptor degrader; androgen receptor DNA-binding inhibitor; or combinations thereof. In some embodiments, the cancer is prostate cancer. In some embodiments, the cancer is hormone refractory prostate cancer. In some embodiments, the compound is a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X), or a pharmaceutically acceptable salt thereof or N-oxide thereof.

[0088] Pharmaceutical formulations described herein are administerable to a subject in a variety of ways by multiple administration routes, including but not limited to, oral, parenteral (e.g., intravenous, subcutaneous, intramuscular), buccal, topical or transdermal administration routes. The pharmaceutical formulations described herein include, but are not limited to, aqueous liquid dispersions, self-emulsifying dispersions, solid solutions, liposomal dispersions, solid dosage forms, powders, immediate release formulations, controlled release formulations, fast melt formulations, tablets, capsules, pills, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate and controlled release formulations.

[0089] In some embodiments, the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are administered orally.

[0090] In some embodiments, the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are administered topically. In such embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is formulated into a variety of topically administrable compositions, such as solutions, suspensions, lotions, gels, pastes, shampoos, scrubs, rubs, smears, medicated sticks, medicated bandages, balms, creams or ointments. Such pharmaceutical compounds can contain solubilizers, stabilizers, tonicity enhancing agents, buffers and preservatives. In one aspect, the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are administered topically to the skin.

[0091] In another aspect is the use of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) in the manufacture of a medicament for treating a disease, disorder or conditions in which the activity of androgen receptors contributes to the pathology and/or symptoms of the disease or condition. In one aspect, the disease or condition is any of the diseases or conditions specified herein.

[0092] In any of the aforementioned aspects are further embodiments in which: (a) the effective amount of the compound of Formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX) or (X) is systemically administered to the mammal; and/or (b) the effective amount of the compound is administered orally to the mammal; and/or (c) the effective amount of the compound is intravenously administered to the mammal; and/or (d) the effective amount of the compound is administered by injection to the mammal; and/or (e) the effective amount of the compound is administered topically to the mammal; and/or (f) the effective amount is administered non-systemically or locally to the mammal.

[0093] In any of the aforementioned aspects are further embodiments comprising single administrations of the effective amount of the compound, including further embodiments in

which (i) the compound is administered once; (ii) the compound is administered to the mammal multiple times over the span of one day; (iii) continually; or (iv) continuously.

[0094] In any of the aforementioned aspects are further embodiments comprising multiple administrations of the effective amount of the compound, including further embodiments in

5 which (i) the compound is administered continuously or intermittently: as in a single dose; (ii) the time between multiple administrations is every 6 hours; (iii) the compound is administered to the mammal every 8 hours; (iv) the compound is administered to the mammal every 12 hours; (v) the compound is administered to the mammal every 24 hours. In further or alternative embodiments, the method comprises a drug holiday, wherein the administration of the
10 compound is temporarily suspended or the dose of the compound being administered is temporarily reduced; at the end of the drug holiday, dosing of the compound is resumed. In one embodiment, the length of the drug holiday varies from 2 days to 1 year.

[0095] Also provided is a method of reducing AR activation in a mammal comprising administering to the mammal at least one compound having the structure of Formula (I), (Ia),
15 (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X). In some embodiments, the method comprises reducing AR activation in prostate cells in the mammal. In some embodiments, the method comprises reducing AR activation in prostate cells in the mammal and not in non-prostate cells. In some embodiments, the method of reducing AR activation comprises reducing the binding of androgens to the androgen receptor. In some embodiments,
20 the method of reducing AR activation comprises reducing AR concentrations.

[0096] In some cases disclosed herein is the use of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) in the manufacture of a medicament for the treatment of diseases or conditions that are androgen receptor dependent or androgen receptor mediated. In some embodiments, the disease or condition is prostate cancer. In some
25 embodiments, the androgen receptor dependent or androgen receptor mediated disease or condition is described herein.

[0097] In some cases disclosed herein is the use of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) in the treatment or prevention of diseases or conditions that are androgen receptor dependent or androgen receptor mediated. In
30 some embodiments, the androgen receptor dependent or androgen receptor mediated disease or condition is described herein.

[0098] In any of the embodiments disclosed herein, the mammal is a human.

[0099] In some embodiments, compounds provided herein are administered to a human.

[00100] In some embodiments, compounds provided herein are orally administered.

[00101] In some embodiments, compounds provided herein are used to diminish, reduce, or eliminate the activity of androgen receptors.

[00102] Articles of manufacture, which include packaging material, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) within the packaging material, and a label that indicates that the compound or composition, or pharmaceutically acceptable salt, tautomers, pharmaceutically acceptable N-oxide, pharmaceutically active metabolite, pharmaceutically acceptable prodrug, or pharmaceutically acceptable solvate thereof, is used for reducing, diminishing or eliminating the effects of androgen receptors, or for the treatment, prevention or amelioration of one or more symptoms of a disease or condition that would benefit from a reduction or elimination of androgen receptor activity, are provided.

[00103] Other objects, features and advantages of the compounds, methods and compositions described herein will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments, are given by way of illustration only, since various changes and modifications within the spirit and scope of the instant disclosure will become apparent to those skilled in the art from this detailed description

DETAILED DESCRIPTION OF THE INVENTION

[00104] Androgen receptor (AR) is a member of the steroid and nuclear receptor superfamily.

Among this large family of proteins, only five vertebrate steroid receptors are known and include the androgen receptor, estrogen receptor, progesterone receptor, glucocorticoid receptor, and mineralocorticoid receptor. AR is a soluble protein that functions as an intracellular transcriptional factor. AR function is regulated by the binding of androgens, which initiates sequential conformational changes of the receptor that affect receptor–protein interactions and receptor–DNA interactions.

[00105] AR is mainly expressed in androgen target tissues, such as the prostate, skeletal muscle, liver, and central nervous system (CNS), with the highest expression level observed in the prostate, adrenal gland, and epididymis. AR can be activated by the binding of endogenous androgens, including testosterone and 5 α -dihydrotestosterone (5 α -DHT).

[00106] The androgen receptor (AR), located on Xq11-12, is a 110 kD nuclear receptor that, upon activation by androgens, mediates transcription of target genes that modulate growth and differentiation of prostate epithelial cells. Similar to the other steroid receptors, unbound AR is mainly located in the cytoplasm and associated with a complex of heat shock proteins (HSPs) through interactions with the ligand-binding domain. Upon agonist binding, AR goes through a series of conformational changes: the heat shock proteins dissociate from AR, and the

transformed AR undergoes dimerization, phosphorylation, and translocation to the nucleus, which is mediated by the nuclear localization signal. Translocated receptor then binds to the androgen response element (ARE), which is characterized by the six-nucleotide half-site consensus sequence 5'-TGTTCT-3' spaced by three random nucleotides and is located in the

5 promoter or enhancer region of AR gene targets. Recruitment of other transcription co-regulators (including co-activators and co-repressors) and transcriptional machinery further ensures the transactivation of AR-regulated gene expression. All of these processes are initiated by the ligand-induced conformational changes in the ligand-binding domain.

[00107] AR signaling is crucial for the development and maintenance of male reproductive
10 organs including the prostate gland, as genetic males harboring loss of function AR mutations and mice engineered with AR defects do not develop prostates or prostate cancer. This dependence of prostate cells on AR signaling continues even upon neoplastic transformation. Androgen depletion (now using GnRH agonists) continues to be the mainstay of prostate cancer treatment. However androgen depletion is usually effective for a limited duration and prostate
15 cancer evolves to regain the ability to grow despite low levels of circulating androgens.

Treatment options for castration resistant prostate cancer (CRPC) are an unmet need with docetaxel being the only agent that has been shown to prolong survival. Interestingly, while a small minority of CRPC does bypass the requirement for AR signaling, the vast majority of CRPC, though frequently termed "androgen independent prostate cancer" or "hormone
20 refractory prostate cancer," retains its lineage dependence on AR signaling.

[00108] Prostate cancer is the second most common cause of cancer death in men in the US, and approximately one in every six American men will be diagnosed with the disease during his lifetime. Treatment aimed at eradicating the tumor is unsuccessful in 30% of men, who develop recurrent disease that is usually manifest first as a rise in plasma prostate-specific antigen (PSA)
25 followed by spread to distant sites. Given that prostate cancer cells depend on androgen receptor (AR) for their proliferation and survival, these men are treated with agents that block production of testosterone (e.g. GnRH agonists), alone or in combination with anti-androgens (e.g. bicalutamide), which antagonize the effect of any residual testosterone. The approach is effective as evidenced by a drop in PSA and regression of visible tumor (if present); however,
30 this is followed by regrowth as a "castration resistant" prostate cancer (CRPC) to which most patients eventually succumb. Recent studies on the molecular basis of CRPC have demonstrated that CRPC continues to depend on AR signaling and that a key mechanism of acquired resistance is an elevated level of AR protein (*Nat. Med.*, 2004, 10, 33-39). AR targeting agents with activity in hormone sensitive and castration resistant prostate cancer have great
35 promise in treating this lethal disease.

[00109] Anti-androgens are useful for the treatment of prostate cancer during its early stages. However, prostate cancer often advances to a hormone-refractory state in which the disease progresses in the presence of continued androgen ablation or anti-androgen therapy. Instances of antiandrogen withdrawal syndrome have also been reported after prolonged treatment with anti-androgens. Antiandrogen withdrawal syndrome is commonly observed clinically and is defined in terms of the tumor regression or symptomatic relief observed upon cessation of antiandrogen therapy. AR mutations that result in receptor promiscuity and the ability of these anti-androgens to exhibit agonist activity might at least partially account for this phenomenon. For example, hydroxyflutamide and bicalutamide act as AR agonists in T877A and W741L/W741C AR mutants, respectively.

[00110] In the setting of prostate cancer cells that were rendered “castration resistant” via overexpression of AR, it has been demonstrated that certain anti-androgen compounds, such as bicalutamide, have no antagonist activity, but instead have modest agonist activity (Science, 2009 May 8;324(5928): 787-90). This agonist activity helps to explain a clinical observation, called the anti-androgen withdrawal syndrome, whereby about 30% of men who progress on AR antagonists experience a decrease in serum PSA when therapy is discontinued (*J Clin Oncol*, 1993. 11(8): p. 1566-72).

[00111] Given the central role of AR in prostate cancer development and progression, compounds disclosed herein are useful in the treatment of prostate cancer, either alone or in combination with other agent agents that can modulate other critical pathways in prostate cancer, including but not limited to those that target IGF1R, the PI3K/AKT/mTOR axis, HSP90, or histone deacetylases.

[00112] AR-related diseases or conditions include, but are not limited to, benign prostate hyperplasia, hirsutism, acne, adenomas and neoplasies of the prostate, benign or malignant tumor cells containing the androgen receptor, hyperpilosity, seborrhea, endometriosis, polycystic ovary syndrome, androgenic alopecia, hypogonadism, osteoporosis, suppression of spermatogenesis, libido, cachexia, anorexia, androgen supplementation for age related decreased testosterone levels, prostate cancer, breast cancer, endometrial cancer, uterine cancer, hot flashes, and Kennedy’s disease muscle atrophy and weakness, skin atrophy, bone loss, anemia, arteriosclerosis, cardiovascular disease, loss of energy, loss of well-being, type 2 diabetes and abdominal fat accumulation.

[00113] In some embodiments, compounds disclosed herein inhibit AR nuclear translocation, DNA binding to androgen response elements, and coactivator recruitment. In some embodiments, compounds disclosed herein exhibit no agonist activity in AR-overexpressing prostate cancer cells.


[00114] In some embodiments, compounds disclosed herein are used to treat prostate cancer in a mammal, wherein the mammal is chemotherapy-naïve.

[00115] In some embodiments, compounds disclosed herein are used to treat prostate cancer in a mammal, wherein the mammal is being treated for prostate cancer with at least one anti-cancer agent. In one embodiment, the prostate cancer is hormone refractory prostate cancer. In one embodiment, the prostate cancer is bicalutamide-resistant prostate cancer.

Compounds

[00116] Compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X), including pharmaceutically acceptable salts, prodrugs, and pharmaceutically acceptable solvates thereof, are androgen receptor modulators, such as, for example, AR inverse agonists, AR antagonists, AR degraders, AR trafficking modulators and/or AR DNA-binding inhibitors, and are useful in the treatment of androgen receptor-dependent or androgen receptor-mediated conditions or diseases or conditions.

[00117] In one aspect, provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (I)

wherein,

ring A is monocyclic heteroaryl, bicyclic heteroaryl, or naphthyl;

m is 0, 1, 2, 3 or 4;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl;

each R¹ is independently selected from H, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-C₆fluoroalkyl;

or both R^1 are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C_3 - C_{10} cycloalkyl or a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

each R^2 is H; or both R^2 are taken together with the carbon to which they are attached to form $-C(=S)-$ or $-C(=O)-$;

each R^3 is H; or both R^3 are taken together with the carbon to which they are attached to form $-C(=S)-$ or $-C(=O)-$; provided that each R^2 is not H if each R^3 is H;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, 2, 3 or 4;

each R^4 is independently selected from H, halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, substituted or unsubstituted C_1 - C_6 alkoxy, and substituted or unsubstituted C_1 - C_6 heteroalkyl;

R^5 is substituted or unsubstituted C_2 - C_{10} alkyl, substituted or unsubstituted C_2 - C_{10} fluoroalkyl, substituted or unsubstituted C_2 - C_{10} alkoxy, substituted or unsubstituted C_2 - C_{10} fluoroalkoxy, substituted or unsubstituted C_2 - C_{10} heteroalkyl, substituted or unsubstituted C_2 - C_{10} heterofluoroalkyl, or $-L^1-L^2-R^6$;

L^1 is absent, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, $-NHC(=O)O-$, $-NHC(=O)NH-$, $-C(=O)O-$, $-OC(=O)-$, $-OC(=O)O-$, $-OC(=O)NH-$, $-NHS(=O)_2-$, or $-S(=O)_2NH-$;

L^2 is substituted or unsubstituted C_1 - C_6 alkylene, substituted or unsubstituted C_1 - C_6 fluoroalkylene or substituted or unsubstituted C_1 - C_6 heteroalkylene;

R^6 is halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl;

each R⁹ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or

two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;

R¹⁰ is substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl);

R¹¹ is H or C₁-C₄alkyl.

[00118] For any and all of the embodiments, substituents are selected from among from a subset of the listed alternatives. For example, in some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₁₀cycloalkyl. In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a C₃-C₆cycloalkyl. In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a cyclopropyl. In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a cyclobutyl. In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a cyclopentyl. In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a cyclohexyl. In some embodiments, each R¹ is independently selected from H, -OH, C₁-C₆alkyl, and C₁-C₆fluoroalkyl. In some embodiments, each R¹ is independently selected from H, -CH₃ and -CF₃. In some embodiments, each R¹ is -CH₃. In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a cyclobutyl or each R¹ is -CH₃.

[00119] In some embodiments, both R^2 are taken together with the carbon to which they are attached to form $-C(=S)-$ or $-C(=O)-$; both R^3 are taken together with the carbon to which they are attached to form $-C(=O)-$. In some embodiments, both R^2 are taken together with the carbon to which they are attached to form $-C(=S)-$; both R^3 are taken together with the carbon to which they are attached to form $-C(=O)-$.


[00120] In some embodiments, ring A is C-linked monocyclic heteroaryl, C-linked bicyclic heteroaryl, or naphthyl. In some embodiments, ring A is C-linked monocyclic heteroaryl. In some embodiments, ring A is C-linked monocyclic 5-membered or 6-membered heteroaryl. In some embodiments, ring A is C-linked monocyclic 6-membered heteroaryl. In some

embodiments, ring A is an *N*-containing heteroaryl.

[00121] In some embodiments, both R^1 are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C_3-C_{10} cycloalkyl; both R^2 are taken together with the carbon to which they are attached to form $-C(=S)-$; both R^3 are taken together with the carbon to which they are attached to form $-C(=O)-$; ring A is *N*-containing monocyclic heteroaryl, or *N*-containing bicyclic heteroaryl; each R^A is independently selected from H, halogen, $-CN$, $-NO_2$, $-OH$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-C(=O)N(R^9)_2$, substituted or unsubstituted C_1-C_6 alkyl, substituted or unsubstituted C_1-C_6 fluoroalkyl, substituted or unsubstituted C_1-C_6 fluoroalkoxy, substituted or unsubstituted C_1-C_6 alkoxy, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl.

[00122] In some embodiments, both R^1 are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; ring A is *N*-containing monocyclic heteroaryl selected from pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, triazinyl, pyrrolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, imidazolyl, pyrazolyl, isothiazolyl, triazolyl, and tetrazolyl.

[00123] In some embodiments, ring A is pyridinyl, pyrimidinyl, pyridazinyl, or pyrazinyl. In some embodiments, ring A is pyridinyl.



[00125] In some embodiments, ring A is monosubstituted with R^A . In some embodiments, ring A is disubstituted with R^A . In some embodiments, ring A is trisubstituted with R^A . In some embodiments, ring A is substituted with $-\text{CN}$ and at least one additional R^A . In some embodiments, ring A is substituted with $-\text{CN}$ and zero, one or two additional R^A . In some embodiments, ring A is substituted with $-\text{CN}$ and one or two additional R^A . In some embodiments, ring A is substituted with $-\text{CN}$ and one additional R^A .

[00126] In some embodiments, ring A is pyridinyl; each R^A is independently selected from H, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, substituted or unsubstituted C_1 - C_6 alkoxy, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl.

[00127] In some embodiments, both R^1 are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; ring A is *N*-containing bicyclic heteroaryl selected from quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, indolyl, indazolyl, benzoxazolyl, benzisoxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzimidazolyl, purinyl, cinnolinyl, phthalazinyl, pteridinyl, pyridopyrimidinyl, pyrazolopyrimidinyl, azaindolyl, pyrazolopyridinyl, thiazolopyrimidinyl, triazolopyridazinyl, thiazolopyridinyl, pyridothienyl, pyrimidithienyl and pyrrolopyrimidinyl; each R^A is independently selected from H, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, substituted or unsubstituted C_1 - C_6 alkoxy, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl.

[00128] In some embodiments, ring A is *N*-containing bicyclic heteroaryl selected from quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, pyridopyrimidinyl, pyrazolopyrimidinyl, triazolopyridazinyl, and pyrrolopyrimidinyl.

[00129] In some embodiments, ring A is [1,2,4]triazolo[4,3-*b*]pyridazinyl.

5 [00130] In some embodiments, ring B is phenyl or monocyclic heteroaryl; each R^4 is independently selected from H, halogen, -CN, -NO₂, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-C₆heteroalkyl; R^5 is substituted or unsubstituted C₂-C₁₀alkyl, substituted or unsubstituted C₂-C₁₀fluoroalkyl, substituted or unsubstituted C₂-C₁₀alkoxy, substituted or unsubstituted C₂-C₁₀fluoroalkoxy, substituted or unsubstituted C₂-C₁₀heteroalkyl, substituted or unsubstituted C₂-C₁₀heterofluoroalkyl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHS(=O)₂-, or -S(=O)₂NH-; L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene; R^6 is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

20 [00131] In some embodiments, ring B is phenyl or monocyclic heteroaryl; each R^4 is independently selected from H, halogen, -CN, -NO₂, -OH, substituted or unsubstituted C₁-C₄alkyl, substituted or unsubstituted C₁-C₄fluoroalkyl, substituted or unsubstituted C₁-C₄fluoroalkoxy, and substituted or unsubstituted C₁-C₄alkoxy; R^5 is substituted or unsubstituted C₂-C₁₀alkyl, substituted or unsubstituted C₂-C₁₀fluoroalkyl, substituted or unsubstituted C₂-C₁₀alkoxy, substituted or unsubstituted C₂-C₁₀fluoroalkoxy, substituted or unsubstituted C₂-C₁₀heteroalkyl, substituted or unsubstituted C₂-C₁₀heterofluoroalkyl, or -L¹-L²-R⁶; L¹ is absent, -O-, or -C(=O)NH-; L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene; R^6 is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.


35 [00132] In some embodiments, L² is substituted or unsubstituted C₁-C₆alkylene; R^6 is -CN, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-

C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

[00133] In some embodiments, R⁶ is substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

[00134] In some embodiments, L² is substituted or unsubstituted C₁-C₆alkylene; R⁶ is -CN, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, or -C(=O)N(R⁹)₂; each R⁹ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C₂-C₁₀heterocycloalkyl; R¹⁰ is substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl).

[00135] In some embodiments, provided is a compound of Formula (Ia), or a pharmaceutically acceptable salt, or *N*-oxide thereof:



Formula (Ia)

wherein,

ring A is monocyclic heteroaryl, bicyclic heteroaryl, or naphthyl;

m is 1, 2, 3 or 4;

each R^A is independently H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, C₁-C₆alkoxy, or C₁-C₆heteroalkyl;

both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₆cycloalkyl or a substituted or unsubstituted C₂-C₆heterocycloalkyl;

or each R¹ is independently selected from H, -OH, C₁-C₆alkyl, C₁-C₆alkoxy, and C₁-C₆fluoroalkyl;

X is S or O;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, 2, 3 or 4;

each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, C₁-C₆alkoxy, and C₁-C₆heteroalkyl;

R⁵ is -L¹-L²-R⁶ or -L¹-R⁷;

L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-, or -S(=O)₂NH-;

L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene;

R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl;

R⁷ is substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl;

each R⁹ is independently selected from H, C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₆cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted phenyl), and -C₁-C₄alkylene-(substituted or unsubstituted monocyclic heteroaryl); or

two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl;

R¹⁰ is C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₆cycloalkyl, a substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, a substituted or unsubstituted phenyl, a substituted or unsubstituted benzyl, a substituted or unsubstituted monocyclic heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₆cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted phenyl), or -C₁-C₄alkylene-(substituted or unsubstituted monocyclic heteroaryl);

R¹¹ is H or C₁-C₄alkyl.

[00136] In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₆cycloalkyl; X is S; ring A is N-containing monocyclic heteroaryl; each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, and C₁-C₆alkoxy; ring B is phenyl or monocyclic heteroaryl.

[00137] In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; ring A is pyridinyl, pyrimidinyl, pyridazinyl, or pyrazinyl.

[00138] In some embodiments, ring A is pyridinyl; each R^A is independently selected from H, halogen, -CN, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, and C₁-C₆alkoxy.

[00139] In some embodiments, each R^A is independently selected from H, halogen, -CN, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, and C₁-C₆alkoxy. In some embodiments, each R^A is independently selected from H, halogen, -CN, -OH, -C(=O)N(R⁹)₂, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, and C₁-C₄alkoxy. In some embodiments, each R^A is independently selected from H, F, Cl, -CN, -OH, -C(=O)NH₂, -CH₃, -CH₂H₃, -CF₃, -OCF₃, -OCH₃, -CH₂CH₃.

[00140] In some embodiments, ring B is phenyl; R⁵ is -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHS(=O)₂-, or -S(=O)₂NH-; L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene; R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-

C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

[00141] In some embodiments, ring B is phenyl; each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, and C₁-C₄alkoxy; R⁵ is -L¹-L²-R⁶; L¹ is absent, -O-, or -C(=O)NH-; L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

[00142] In some embodiments, L² is C₁-C₆alkylene; R⁶ is -CN, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

[00143] In some embodiments, R⁶ is substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

[00144] In some embodiments, if R⁶ is substituted, then R⁶ is substituted with 1 or 2 groups independently selected from halogen, -CN, -NH₂, -NH(CH₃), -N(CH₃)₂, -OH, -CO₂H, -CO₂(C₁-C₄alkyl), -C(=O)C₁-C₄alkyl, -C(=O)NH₂, -C(=O)NH(C₁-C₄alkyl), -C(=O)N(C₁-C₄alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(C₁-C₄alkyl), -S(=O)₂N(C₁-C₄alkyl)₂, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄fluoroalkyl, C₁-C₄heteroalkyl, C₁-C₄alkoxy, C₁-C₄fluoroalkoxy, -S-C₁-C₄alkyl, or -S(=O)₂C₁-C₄alkyl. In some embodiments, if R⁶ is substituted, then R⁶ is substituted with 1 or 2 groups independently selected from halogen, -CN, -OH, -CO₂(C₁-C₄alkyl), -C(=O)C₁-C₄alkyl, -C(=O)NH₂, -C(=O)NH(C₁-C₄alkyl), -C(=O)N(C₁-C₄alkyl)₂, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄fluoroalkyl, C₁-C₄heteroalkyl, C₁-C₄alkoxy, and C₁-C₄fluoroalkoxy.

[00145] In some embodiments, L² is C₁-C₆alkylene; R⁶ is a substituted or unsubstituted C₂-C₆heterocycloalkyl. In some embodiments, L¹ is absent, -O-, or -C(=O)NH-; L² is C₁-C₆alkylene; R⁶ is a substituted or unsubstituted C₂-C₆heterocycloalkyl. In some embodiments, L¹ is -C(=O)NH-. In some embodiments, R⁶ is a substituted or unsubstituted C₃-C₆heterocycloalkyl. In some embodiments, R⁶ is a substituted or unsubstituted N-containing C₂-C₆heterocycloalkyl.

[00146] In some embodiments, ring B is phenyl; each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, and C₁-C₄alkoxy; R⁵ is -L¹-R⁷; L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-,

or $-S(=O)_2NH-$; R^7 is substituted or unsubstituted monocyclic C_2 - C_6 heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.

[00147] In some embodiments, L^2 is C_1 - C_6 alkylene. In some embodiments, L^2 is C_1 - C_4 alkylene.

In some embodiments, L^2 is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2CH_2CH_2CH_2-$, -

5 $CH_2CH_2CH_2CH_2CH_2-$, or $-CH_2CH_2CH_2CH_2CH_2CH_2-$. In some embodiments, L^2 is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, or $-CH_2CH_2CH_2CH_2-$.

[00148] In some embodiments, each R^4 is independently selected from H, halogen, $-OH$, C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, C_1 - C_4 fluoroalkoxy, and C_1 - C_4 alkoxy.

[00149] In some embodiments, R^5 is $-L^1-R^7$.

10 [00150] In some embodiments, L^1 is absent, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$. In some embodiments, L^1 is absent. In some embodiments, L^1 is $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$. In some embodiments, L^1 is absent. In some embodiments, L^1 is $-O-$. In some embodiments, L^1 is absent. In some embodiments, L^1 is $-C(=O)NH-$.

15 [00151] In some embodiments, R^5 is $-L^1-R^7$; L^1 is $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$; R^7 is substituted or unsubstituted monocyclic C_2 - C_6 heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.

[00152] In some embodiments, if R^7 is substituted, then R^7 is substituted with 1 or 2 groups independently selected from halogen, $-CN$, $-NH_2$, $-NH(CH_3)$, $-N(CH_3)_2$, $-OH$, $-CO_2H$, $-CO_2(C_1$ -


20 C_4 alkyl), $-C(=O)C_1$ - C_4 alkyl, $-C(=O)NH_2$, $-C(=O)NH(C_1$ - C_4 alkyl), $-C(=O)N(C_1$ - C_4 alkyl) $_2$, $-S(=O)_2NH_2$, $-S(=O)_2NH(C_1$ - C_4 alkyl), $-S(=O)_2N(C_1$ - C_4 alkyl) $_2$, C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, C_1 - C_4 fluoroalkyl, C_1 - C_4 heteroalkyl, C_1 - C_4 alkoxy, C_1 - C_4 fluoroalkoxy, $-S-C_1$ - C_4 alkyl, or $-S(=O)_2C_1$ - C_4 alkyl. In some embodiments, if R^7 is substituted, then R^7 is substituted with 1 or 2 groups independently selected from halogen, $-CN$, $-OH$, $-CO_2(C_1$ - C_4 alkyl), $-C(=O)C_1$ - C_4 alkyl, -

25 $C(=O)NH_2$, $-C(=O)NH(C_1$ - C_4 alkyl), $-C(=O)N(C_1$ - C_4 alkyl) $_2$, C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, C_1 - C_4 fluoroalkyl, C_1 - C_4 heteroalkyl, C_1 - C_4 alkoxy, and C_1 - C_4 fluoroalkoxy.

[00153] In some embodiments, R^5 is $-L^1-R^7$; L^1 is absent, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$; R^7 is substituted or unsubstituted monocyclic heteroaryl.


[00154] In some embodiments, the compound of Formula (I) or Formula (Ia) has the structure of

30 Formula (II):




Formula (II).

[00155] In some embodiments, the compound of Formula (Ia) has the structure of Formula (II):



Formula (II)

wherein,




m is 2;

one R^A is $-\text{CN}$, $-\text{NO}_2$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{R}^{10}$, $-\text{CO}_2\text{R}^9$, or - $\text{C}(=\text{O})\text{N}(\text{R}^9)_2$; and the other R^A is H, halogen, $-\text{OH}$, $\text{C}_1\text{-C}_6\text{alkyl}$, $\text{C}_1\text{-C}_6\text{fluoroalkyl}$, $\text{C}_1\text{-C}_6\text{fluoroalkoxy}$, or $\text{C}_1\text{-C}_6\text{alkoxy}$;

n is 0 or 1;


each R^4 is independently selected from H, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{OH}$, $\text{C}_1\text{-C}_6\text{alkyl}$, $\text{C}_1\text{-C}_6\text{fluoroalkyl}$, $\text{C}_1\text{-C}_6\text{fluoroalkoxy}$, and $\text{C}_1\text{-C}_6\text{alkoxy}$; R^5 is $-\text{L}^1\text{-L}^2\text{-R}^6$ or $-\text{L}^1\text{-R}^7$; L^1 is absent, $-\text{O}-$, or $-\text{C}(=\text{O})\text{NH}-$; L^2 is $\text{C}_1\text{-C}_6\text{alkylene}$, $\text{C}_1\text{-C}_6\text{fluoroalkylene}$ or $\text{C}_1\text{-C}_6\text{heteroalkylene}$; R^6 is $-\text{CN}$, $-\text{NO}_2$, $-\text{OH}$, $-\text{OR}^9$, $-\text{SR}^9$, $-\text{S}(=\text{O})\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{C}(=\text{O})\text{R}^{10}$, $-\text{CO}_2\text{R}^9$, - $\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, substituted or unsubstituted $\text{C}_3\text{-C}_6\text{cycloalkyl}$, substitutedor unsubstituted $\text{C}_2\text{-C}_6\text{heterocycloalkyl}$, substituted or unsubstituted monocyclic

heteroaryl, or substituted or unsubstituted phenyl;

 R^7 is substituted or unsubstituted monocyclic $\text{C}_2\text{-C}_6\text{heterocycloalkyl}$, or substituted or unsubstituted monocyclic heteroaryl.[00156] In some embodiments, one R^A is $-\text{CN}$ and the other R^A isH, halogen, $-\text{OH}$, $\text{C}_1\text{-C}_4\text{alkyl}$, $\text{C}_1\text{-C}_4\text{fluoroalkyl}$, $\text{C}_1\text{-C}_4\text{fluoroalkoxy}$, or $\text{C}_1\text{-C}_4\text{alkoxy}$; each R^4 isindependently selected from H, halogen, $-\text{OH}$, $\text{C}_1\text{-C}_4\text{alkyl}$, $\text{C}_1\text{-C}_4\text{fluoroalkyl}$, $\text{C}_1\text{-C}_4\text{fluoroalkoxy}$,and $\text{C}_1\text{-C}_4\text{alkoxy}$; R^5 is $-\text{L}^1\text{-L}^2\text{-R}^6$ or $-\text{L}^1\text{-R}^7$; L^1 is absent, $-\text{O}-$, or $-\text{C}(=\text{O})\text{NH}-$; L^2 is $\text{C}_1\text{-}$

C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl; R⁷ is substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.

[00157] In some embodiments, described herein is a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (I)

wherein,

ring A is bicyclic heteroaryl, or naphthyl;

m is 0, 1, 2, or 3;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -

CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or

unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy,

substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted C₁-C₆heteroalkyl,

substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-

C₁₀heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted

monocyclic heteroaryl;

each R¹ is independently selected from H, -OH, substituted or unsubstituted C₁-C₆alkyl,

substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-

C₆fluoroalkyl;

or both R¹ are taken together with the carbon atom to which they are attached to form a

substituted or unsubstituted C₃-C₁₀cycloalkyl or a substituted or unsubstituted C₂-

C₁₀heterocycloalkyl;

each R² is H; or both R² are taken together with the carbon to which they are attached to form -C(=S)- or -C(=O)-;

each R³ is H; or both R³ are taken together with the carbon to which they are attached to

form -C(=S)- or -C(=O)-; provided that each R² is not H if each R³ is H;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

each R^4 is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -

CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -

NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₆alkyl, substituted

or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy,

substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-

C₆heteroalkyl;

R^5 is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -

S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -

OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or

unsubstituted C₁-C₁₀alkyl, substituted or unsubstituted C₁-C₁₀fluoroalkyl, substituted or

unsubstituted C₁-C₁₀fluoroalkoxy, C₁-C₁₀alkoxy, substituted or unsubstituted C₁-

C₁₀heteroalkyl, or -L¹-L²-R⁶;

L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -

NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -

NHS(=O)₂-, or -S(=O)₂NH-;

L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-

C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene;

R^6 is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -

S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂,

-OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted

or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl,

substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₃-

C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or

unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic

heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted

naphthyl;

each R^9 is independently selected from H, substituted or unsubstituted C₁-C₆alkyl,

substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-

C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or

unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted

or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-

C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -

C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or

unsubstituted heteroaryl); or

two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

R^{10} is substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), or $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl);


R^{11} is H or C_1 - C_4 alkyl.

[00158] In some embodiments, ring A is selected from quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, indolyl, indazolyl, benzoxazolyl, benzisoxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzimidazolyl, purinyl, cinnolinyl, phthalazinyl, pteridinyl, pyridopyrimidinyl, pyrazolopyrimidinyl, azaindolyl, pyrazolopyridinyl, thiazolopyrimidinyl, triazolopyridazinyl, thiazolopyridinyl, pyridothienyl, pyrimidothienyl, pyrrolopyrimidinyl and naphthyl.

[00159] In some embodiments, ring A is selected from quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, pyridopyrimidinyl, pyrazolopyrimidinyl, triazolopyridazinyl, pyrrolopyrimidinyl, and naphthyl.

[00160] In some embodiments, ring A is [1,2,4]triazolo[4,3-b]pyridazinyl.

[00161] In some embodiments, described herein is a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (I)

wherein,

ring A is a 5-membered heteroaryl;

m is 0, 1, 2, or 3;

each R^A is independently selected from H, halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, substituted or unsubstituted C_1 - C_4 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, substituted or unsubstituted C_1 - C_6 alkoxy, substituted or unsubstituted C_1 - C_6 heteroalkyl,

substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl;

each R¹ is independently selected from H, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-C₆fluoroalkyl;

or both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₁₀cycloalkyl or a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;

each R² is H; or both R² are taken together with the carbon to which they are attached to form -C(=S)- or -C(=O)-;

each R³ is H; or both R³ are taken together with the carbon to which they are attached to form -C(=S)- or -C(=O)-; provided that each R² is not H if each R³ is H;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₆alkyl, substituted

or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-C₆heteroalkyl;

R⁵ is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -

S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or

unsubstituted C₁-C₁₀alkyl, substituted or unsubstituted C₁-C₁₀fluoroalkyl, substituted or unsubstituted C₁-C₁₀fluoroalkoxy, substituted or unsubstituted C₁-C₁₀alkoxy, substituted or unsubstituted C₁-C₁₀heteroalkyl, or -L¹-L²-R⁶;

L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -

NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-, or -S(=O)₂NH-;

L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene;

R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -

S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂,

-OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl;


each R⁹ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or

two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;


R¹⁰ is substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl);

R¹¹ is H or C₁-C₄alkyl.

[00162] In some embodiments, ring A is selected from pyrrolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, imidazolyl, pyrazolyl, isothiazolyl, triazolyl, and tetrazolyl.




[00163] In some embodiments, is selected from:



O. In some embodiments, X is S. In some embodiments, X is NR^A.

[00164] In some embodiments, both R¹ are taken together with the carbon atom to which they are attached to form a C₃-C₁₀cycloalkyl; both R² are taken together with the carbon to which they are attached to form -C(=S)-; both R³ are taken together with the carbon to which they are attached to form -C(=O)-.

[00165] In some embodiments, the compound described herein has the structure of Formula (III):



Formula (III)

wherein,

p is 0, 1, 2, or 3.

[00166] In some embodiments, each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl.

[00167] In some embodiments, each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, and substituted or unsubstituted C₁-C₆alkoxy; R⁵ is substituted or unsubstituted C₂-C₁₀alkyl, substituted or unsubstituted C₂-C₁₀fluoroalkyl, substituted or unsubstituted C₂-C₁₀alkoxy, substituted or unsubstituted C₂-C₁₀fluoroalkoxy, substituted or unsubstituted C₂-C₁₀heteroalkyl, substituted or unsubstituted C₂-


C₁₀heterofluoroalkyl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHS(=O)₂-, or -S(=O)₂NH-; L² is substituted or unsubstituted C₁-

C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene; R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

[00168] In some embodiments, each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, and substituted or unsubstituted C₁-C₆alkoxy;

R⁵ is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₁-C₁₀alkyl, substituted or unsubstituted C₁-C₁₀fluoroalkyl, substituted or unsubstituted C₁-C₁₀fluoroalkoxy, substituted or unsubstituted C₁-C₁₀alkoxy, substituted or unsubstituted C₁-C₁₀heteroalkyl, or -L¹-L²-R⁶; L¹ is absent, -O-, or -C(=O)NH-; L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl.

[00169] In some embodiments, the compound described herein has the structure of Formula (III):



Formula (III)

wherein,

ring A is monocyclic heteroaryl, bicyclic heteroaryl, or naphthyl;

m is 1 or 2;

each R^A is independently H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, C₁-C₆alkoxy, or C₁-C₆heteroalkyl;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1 or 2;

each R^4 is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -

CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-

C₆fluoroalkoxy, C₁-C₆alkoxy, and C₁-C₆heteroalkyl;

R⁵ is substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-

C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted

monocyclic heteroaryl, -L¹-L²-R⁶ or -L¹-R⁷;

L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -

NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -

NHS(=O)₂-, or -S(=O)₂NH-;

L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene;

R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -

S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -

-OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, C₁-

C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆heteroalkyl, substituted or unsubstituted C₃-

C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or

unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic

heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted

naphthyl;

R⁷ is substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted

monocyclic C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic phenyl,

or substituted or unsubstituted monocyclic heteroaryl;

each R⁹ is independently selected from H, C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-

C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or

unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a

substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, -C₁-

C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-

(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or

unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or

two R⁹ groups attached to the same N atom are taken together with the N atom to which

they are attached to form a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;

R¹⁰ is C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-

C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or


unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl);

R¹¹ is H or C₁-C₄alkyl;

p is 0, 1, 2, or 3.

[00170] In some embodiments, R⁵ is C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic 5-membered or 6-membered heteroaryl, -L¹-L²-R⁶ or -L¹-R⁷; L¹ is absent, -O-, or -C(=O)NH-; L² is C₁-C₆alkylene; R⁶ is C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic 5-membered or 6-membered heteroaryl, or substituted or unsubstituted phenyl; R⁷ is C₃-C₆cycloalkyl, substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic phenyl, or substituted or unsubstituted monocyclic 5-membered or 6-membered heteroaryl.

[00171] In some embodiments, described herein is a compound of Formula (IV), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (IV)

wherein,

each R¹ is independently selected from H, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆alkoxy, and substituted or unsubstituted C₁-C₆fluoroalkyl;

or both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₁₀cycloalkyl or a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;

X¹ is CR^A or N;

X² is CR^A or N;

X³ is CR^A or N; provided that at least two of X¹, X², and X³ is CR^A;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰,

-CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

R⁴ is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, substituted or unsubstituted C₁-C₆alkoxy, or substituted or unsubstituted C₁-C₆heteroalkyl;

R⁵ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₂-C₁₀alkyl, substituted or unsubstituted C₂-C₁₀fluoroalkyl, substituted or unsubstituted C₁-C₁₀fluoroalkoxy, substituted or unsubstituted C₁-C₁₀alkoxy, substituted or unsubstituted C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶;

L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-, or -S(=O)₂NH-;

L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene;



R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl;






each R⁹ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-


C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C₂-C₁₀heterocycloalkyl;

R¹⁰ is substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆heteroalkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₁₀cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted C₂-C₁₀heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl);


R¹¹ is H or C₁-C₄alkyl.

[00172] In some embodiments,  is ; where X¹ is CR^A or N; X² is CR^A or N; X³ is CR^A or N; provided that at least two of X¹, X², and X³ is CR^A.

[00173] In some embodiments,  is , , , or ; m is 0, 1, 2, 3, or 4.



[00175] In some embodiments, the compound of Formula (IV) has the structure of Formula (V):




Formula (V).

[00176] In some embodiments, ring B is phenyl; R^4 is H, halogen, -CN, -OH, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, or substituted or unsubstituted C_1 - C_6 alkoxy; R^5 is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, substituted or unsubstituted C_2 - C_{10} alkyl, substituted or unsubstituted C_2 - C_{10} fluoroalkyl, substituted or unsubstituted C_1 - C_{10} fluoroalkoxy, substituted or unsubstituted C_1 - C_{10} alkoxy, substituted or unsubstituted C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, or -S(=O)₂NH-; L² is substituted or unsubstituted C_1 - C_6 alkylene, substituted or unsubstituted C_1 - C_6 fluoroalkylene or substituted or unsubstituted C_1 - C_6 heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl.

[00177] In some embodiments, each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or


unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, or substituted or unsubstituted C₁-C₆alkoxy.

[00178] In some embodiments, the compound of Formula (I), (Ia), (II), (IV) or (V) has the structure of Formula (VI):



Formula (VI).

[00179] In some embodiments, the compound of Formula (I), (Ia), (II), (IV) or (V) has the structure of Formula (VI):



Formula (VI)

wherein,

one R^A is -CN, -NO₂, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, or -

C(=O)N(R⁹)₂; and the other R^A is H, halogen, -OH, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, or C₁-C₆alkoxy;

both R¹ are taken together with the carbon atom to which they are attached to form a C₃-C₆cycloalkyl;

or each R¹ is independently C₁-C₄alkyl;

R⁴ is H, halogen, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, or C₁-C₄alkoxy.

[00180] In some embodiments, R⁵ is -L¹-L²-R⁶ or -L¹-R⁷; L¹ is absent, -O-, or -C(=O)NH-; L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl; R⁷ is substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.

[00181] In some embodiments, one R^A is -CN; and the other R^A is H, F, Cl, -OH, -CH₃, -CF₃, -OCF₃, -OCH₃ or -OCH₂CH₃. In some embodiments, one R^A is -CN; and the other R^A is -CH₃, or -CF₃.

[00182] In some embodiments, both R^1 are taken together with the carbon atom to which they are attached to form a cyclobutyl; or each R^1 is -CH₃.


[00183] In some embodiments, R^5 is -L¹-L²-R⁶ or -L¹-R⁷; L¹ is absent, -O-, or -C(=O)NH-; L² is C₁-C₆alkylene; R⁶ is substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl; R⁷ is substituted or unsubstituted monocyclic C₂-

C₆heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl. In some embodiments, R^5 is -L¹-L²-R⁶; L¹ is -C(=O)NH-; L² is C₁-C₄alkylene; substituted or unsubstituted N-containing 5-membered or 6-membered C₂-C₆heterocycloalkyl. In some embodiments, R^5 is -L¹-R⁷; L¹ is absent, -O-, or -C(=O)NH-; R⁷ is substituted or unsubstituted monocyclic 5-membered or 6-membered C₂-C₆heterocycloalkyl, or substituted or unsubstituted monocyclic 5-membered or 6-membered heteroaryl.

[00184] In some embodiments, one R^A is -CN, -NO₂, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl; and the other R^A is H, halogen, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-

C₆fluoroalkoxy, or substituted or unsubstituted C₁-C₆alkoxy; R⁴ is H, halogen, -CN, -OH, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₁-C₆fluoroalkoxy, or substituted or unsubstituted C₁-C₆alkoxy; R⁵ is halogen, -CN, -NO₂, -OH, -OR⁹, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, substituted or unsubstituted C₂-C₁₀alkyl, substituted or unsubstituted C₂-C₁₀fluoroalkyl, substituted or unsubstituted C₁-C₁₀fluoroalkoxy, substituted or unsubstituted C₁-C₁₀alkoxy, substituted or unsubstituted C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶; L¹ is absent, -O-, or -C(=O)NH-; L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

[00185] In some embodiments, described herein is a compound of Formula (VII), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (VII)

wherein,

each R^1 is independently selected from H, -OH, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 alkoxy, and substituted or unsubstituted C_1 - C_6 fluoroalkyl;

or both R^1 are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C_3 - C_{10} cycloalkyl or a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, substituted or unsubstituted C_1 - C_6 alkoxy, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic heteroaryl;

ring B is 5-membered heteroaryl, bicyclic heteroaryl or naphthyl;

n is 0, 1, or 2;

R^4 is H, halogen, -CN, -NO₂, -OH, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkoxy, or substituted or unsubstituted C_1 - C_6 alkoxy;

R^5 is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C_2 - C_{10} alkyl, substituted or unsubstituted C_2 - C_{10} fluoroalkyl, substituted or unsubstituted C_1 - C_{10} fluoroalkoxy, substituted or unsubstituted C_1 - C_{10} alkoxy, substituted or unsubstituted C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶;

L^1 is absent, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, $-NHC(=O)O-$, $-NHC(=O)NH-$, $-C(=O)O-$, $-OC(=O)-$, $-OC(=O)O-$, $-OC(=O)NH-$, $-NHS(=O)_2-$, or $-S(=O)_2NH-$;

L^2 is substituted or unsubstituted C_1 - C_6 alkylene, substituted or unsubstituted C_1 - C_6 fluoroalkylene or substituted or unsubstituted C_1 - C_6 heteroalkylene;


R^6 is halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl;

each R^9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), and $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl); or two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

R^{10} is substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), or $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl);

R^{11} is H or C_1 - C_4 alkyl.

[00186] In some embodiments, described herein is a compound of Formula (VIII), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



Formula (VIII)

wherein,

each R^1 is independently selected from H, -OH, substituted or unsubstituted C_1 - C_6 alkyl,
 5 substituted or unsubstituted C_1 - C_6 alkoxy, and substituted or unsubstituted C_1 - C_6 fluoroalkyl;

or both R^1 are taken together with the carbon atom to which they are attached to form a
 substituted or unsubstituted C_3 - C_6 cycloalkyl or a substituted or unsubstituted C_2 - C_6 heterocycloalkyl;

10 X is O or S;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -
 S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -
 CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 heteroalkyl, substituted
 or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl,
 15 substituted or unsubstituted phenyl or substituted or unsubstituted monocyclic
 heteroaryl;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

R^4 is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, substituted or
 20 unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted
 or unsubstituted C_1 - C_6 fluoroalkoxy, substituted or unsubstituted C_1 - C_6 alkoxy, or
 substituted or unsubstituted C_1 - C_6 heteroalkyl;

R^5 is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰,
 -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -
 25 C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -
 NR¹¹C(=O)OR¹⁰, substituted or unsubstituted C_1 - C_{10} alkyl, substituted or
 unsubstituted C_1 - C_{10} fluoroalkyl, substituted or unsubstituted C_1 - C_{10} fluoroalkoxy,
 substituted or unsubstituted C_1 - C_{10} alkoxy, substituted or unsubstituted C_1 -
 C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or
 30 unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl,
 substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶;

L^1 is absent, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, $-NHC(=O)O-$, $-NHC(=O)NH-$, $-C(=O)O-$, $-OC(=O)-$, $-OC(=O)O-$, $-OC(=O)NH-$, $-NHS(=O)_2-$, or $-S(=O)_2NH-$;

L^2 is substituted or unsubstituted C_1 - C_6 alkylene, substituted or unsubstituted C_1 - C_6 fluoroalkylene or substituted or unsubstituted C_1 - C_6 heteroalkylene;

R^6 is $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl;

each R^9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), and $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl); or two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

R^{10} is substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 heteroalkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), or $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl);


R^{11} is H or C_1 - C_4 alkyl.

[00187] In some embodiments, X is S; ring B is phenyl; R^4 is H, halogen, $-CN$, $-OH$, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, or C_1 - C_6 alkoxy; R^5 is halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-CO_2R^9$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, substituted or unsubstituted C_1 - C_{10} alkyl, substituted or unsubstituted C_1 - C_{10} fluoroalkyl,

substituted or unsubstituted C₁-C₁₀fluoroalkoxy, substituted or unsubstituted C₁-C₁₀alkoxy, substituted or unsubstituted C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, or -S(=O)₂NH-; L² is substituted or unsubstituted C₁-C₆alkylene, substituted or unsubstituted C₁-C₆fluoroalkylene or substituted or unsubstituted C₁-C₆heteroalkylene; R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl.

[00188] In some embodiments, each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, C₁-C₆alkyl, and C₁-C₆alkoxy.

[00189] In some embodiments, described herein is a compound of Formula (VIIIa), or a pharmaceutically acceptable salt, or *N*-oxide thereof:



Formula (VIIIa)

wherein,

both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₆cycloalkyl or a substituted or unsubstituted C₂-C₆heterocycloalkyl;

or

each R¹ is independently H, -OH, C₁-C₆alkyl, C₁-C₆alkoxy, or C₁-C₆fluoroalkyl;

X is O or S;

R^A is C₁-C₆alkyl;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

R⁴ is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, C₁-C₆alkoxy, or C₁-C₆heteroalkyl;

R⁵ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -

$C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, C_1 - C_{10} alkyl, C_1 - C_{10} fluoroalkyl, C_1 - C_{10} fluoroalkoxy, C_1 - C_{10} alkoxy, C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or $-L^1-L^2-R^6$;

L^1 is absent, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, $-NHC(=O)-$, $-NHC(=O)O-$, $-NHC(=O)NH-$, $-C(=O)O-$, $-OC(=O)-$, $-OC(=O)O-$, $-OC(=O)NH-$, $-NHS(=O)_2-$, or $-S(=O)_2NH-$;

L^2 is C_1 - C_6 alkylene, C_1 - C_6 fluoroalkylene or C_1 - C_6 heteroalkylene;

R^6 is $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-N(R^{11})S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-OC(=O)R^{10}$, $-CO_2R^9$, $-OCO_2R^{10}$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl;

each R^9 is independently selected from H, C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), and $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl); or

two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

R^{10} is C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), or $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl);


R^{11} is H or C_1 - C_4 alkyl.

[00190] In some embodiments, X is S; ring B is phenyl; R^4 is H, halogen, $-CN$, $-OH$, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, or C_1 - C_6 alkoxy; R^5 is halogen, $-CN$, $-NO_2$, $-OH$, $-OR^9$, -

SR^9 , $-\text{S}(=\text{O})\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{R}^{10}$, $-\text{CO}_2\text{R}^9$, $-\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, C_1 - C_{10} alkyl, C_1 - C_{10} fluoroalkyl, C_1 - C_{10} fluoroalkoxy, C_1 - C_{10} alkoxy, C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or $-\text{L}^1$ - L^2 - R^6 ; L^1 is absent, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$, $-\text{NH}-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{O})\text{NH}-$, or $-\text{S}(=\text{O})_2\text{NH}-$; L^2 is C_1 - C_6 alkylene, C_1 - C_6 fluoroalkylene or C_1 - C_6 heteroalkylene; R^6 is $-\text{CN}$, $-\text{NO}_2$, $-\text{OH}$, $-\text{OR}^9$, $-\text{SR}^9$, $-\text{S}(=\text{O})\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{CO}_2\text{R}^9$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl.

[00191] In some embodiments, R^A is C_1 - C_6 alkyl; both R^1 are taken together with the carbon atom to which they are attached to form a C_3 - C_6 cycloalkyl; or each R^1 is independently C_1 - C_4 alkyl.


[00192] In some embodiments, the compound of Formula (VIII) has the structure of Formula (IX):



Formula (IX).

[00193] In some embodiments, each R^A is independently selected from H, halogen, $-\text{OH}$, C_1 - C_6 alkyl, and C_1 - C_6 alkoxy.

[00194] In some embodiments, the compound of Formula (VIIIa) has the structure of Formula (IXa):



Formula (IXa).


[00195] In some embodiments, each R^9 is independently selected from H, C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_6 cycloalkyl, a substituted or unsubstituted C_2 - C_6 heterocycloalkyl, a substituted or unsubstituted phenyl, a substituted or unsubstituted benzyl, and a substituted or unsubstituted monocyclic heteroaryl; or two R^9 groups attached to the same N atom are taken together with the N atom to which they are

attached to form a substituted or unsubstituted C₂-C₆heterocycloalkyl; R¹⁰ is C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₆cycloalkyl, a substituted or unsubstituted C₂-C₆heterocycloalkyl, a substituted or unsubstituted phenyl, or a substituted or unsubstituted monocyclic heteroaryl.

- 5 [00196] In some embodiments, R⁵ is halogen, -CN, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)NH(R⁹), C₁-C₁₀alkyl, C₁-C₁₀fluoroalkyl, C₁-C₁₀fluoroalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, or -S(=O)₂NH-; L² is C₁-C₄alkylene; R⁶ is -CN, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

- 15 [00197] In some embodiments, R^A is -CH₃; both R¹ are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; or each R¹ is -CH₃; R⁵ is halogen, -CN, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)NH(R⁹), C₁-C₁₀alkyl, C₁-C₁₀fluoroalkyl, C₁-C₁₀fluoroalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶; L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, or -S(=O)₂NH-; L² is C₁-C₄alkylene; R⁶ is substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

- 25 [00198] In some embodiments, the compound of Formula (VIII) or Formula (VIIIa) has the following structure:




[00199] In some embodiments, R^A is H, halogen, -OH, C₁-C₆alkyl, or C₁-C₆alkoxy. In some embodiments, R^A is C₁-C₆alkyl. In some embodiments, R^A is -CH₃.

- 30 [00200] In some embodiments, R⁴ is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, C₁-C₆alkoxy, or C₁-C₆heteroalkyl. In some embodiments, R⁴ is halogen. In some embodiments, R⁴ is F.

[00201] In some embodiments, R^5 is halogen, $-\text{CN}$, $-\text{OH}$, $-\text{OR}^9$, $-\text{SR}^9$, $-\text{S}(=\text{O})\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{R}^{10}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^9)_2$, $-\text{C}(=\text{O})\text{R}^{10}$, $-\text{CO}_2\text{R}^9$, $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$, C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, C_1 - C_4 fluoroalkoxy, C_1 - C_4 alkoxy. In some embodiments, R^5 is $-\text{CO}_2\text{R}^9$ or $-\text{C}(=\text{O})\text{N}(\text{R}^9)_2$. In some embodiments, R^5 is $-\text{CO}_2\text{H}$, $-\text{C}(=\text{O})\text{NH}_2$ or $-\text{C}(=\text{O})\text{NH}(\text{CH}_3)$.

[00202] In some embodiments, described herein is a compound of Formula (X), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



wherein,

each R^1 is independently selected from H and $-\text{CH}_3$;

or both R^1 are taken together with the carbon atom to which they are attached to form a cyclobutyl;


X is O or S;

X^1 is CH or N;

R^A is $-\text{CN}$ or $-\text{C}(=\text{O})\text{NH}_2$;

R^5 is $-\text{CO}_2\text{H}$, or $-\text{C}(=\text{O})\text{NH}_2$.

[00203] In some embodiments, described herein is a compound of Formula (X), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



wherein,

each R^1 is independently selected from H and $-\text{CH}_3$;

or both R^1 are taken together with the carbon atom to which they are attached to form a cyclobutyl;


X is O;

X^1 is CH or N;

R^A is $-\text{CN}$ or $-\text{C}(=\text{O})\text{NH}_2$;

R^5 is $-\text{CO}_2\text{H}$, $-\text{C}(=\text{O})\text{NH}_2$ or $-\text{C}(=\text{O})\text{NH}(\text{CH}_3)$.

[00204] In some embodiments, described herein is a compound of Formula (X), or a pharmaceutically acceptable salt, solvate, *N*-oxide, metabolite or prodrug thereof:



wherein,

each R^1 is independently selected from H and $-CH_3$;

or both R^1 are taken together with the carbon atom to which they are attached to form a cyclobutyl;

X is O or S;

X^1 is CH or N;

R^A is $-C(=O)NH_2$;

R^5 is $-CO_2H$, $-C(=O)NH_2$ or $-C(=O)NH(CH_3)$.








[00205] Throughout the specification, groups and substituents thereof can be chosen by one skilled in the field to provide stable moieties and compounds.







[00206] In one aspect, compounds described herein include compounds in Table 1, or a pharmaceutically acceptable salt thereof, or *N*-oxide thereof:

Table 1.

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
1	5-(5-(3-Fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		
2	5-(4-Hydroxyphenyl)-6-thioxo-7-(3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-yl)-5,7-diazaspiro[3.4]octan-8-one		435.4








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
3	Ethyl 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoate		
4	5-(5-(4-Bromo-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		500.3
5	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		424.0
6	4-(3-(6-Cyano-5-methylpyridin-3-yl)-4-oxo-2-thioxo-1,3-diazaspiro[4.4]nonan-1-yl)-2-fluoro-N-methylbenzamide		438.1
7	3-Methyl-5-(5-(4-(5-methylfuran-2-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		429.1
8	Ethyl 5-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoate		493.4
9	5-(5-(Naphthalen-2-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		453.0

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
10	5-(5-(2-Cyanophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		428.0
11	5-(5-(3-Cyanophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		428.4
12	5-(5-(4-Cyanophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		428.4
13	5-(5-([1,1'-Biphenyl]-4-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		479.5
14	5-(5-([1,1'-Biphenyl]-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		479.0
15	5-(5-([1,1'-Biphenyl]-2-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		479.0
16	5-(8-Oxo-5-(4-(pyridin-2-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		480.1








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
17	5-(8-Oxo-5-(4-(pyridin-3-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		480.1
18	5-(8-Oxo-5-(4-(pyridin-4-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		480.1
19	5-(8-Oxo-5-(pyridin-3-yl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		404.4
20	5-(8-Oxo-6-thioxo-5-(o-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		417.5
21	5-(8-Oxo-6-thioxo-5-(m-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		417.5
22	5-(8-Oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		417.5

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
23 ^a	5-(8-Oxo-5-(2-phenoxyphenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		495.5
24	5-(8-Oxo-5-(3-phenoxyphenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		495.5
25	5-(8-Oxo-5-(4-phenoxyphenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		495.5
26	5-(5-(2-Fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		421.5
27	5-(5-(3-Fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		421.5
28	5-(5-(4-Fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		421.5








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
29	5-(5-(4-Fluoro-2-methoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		451.5
30	6-Thioxo-5-(p-tolyl)-7-(3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-yl)-5,7-diazaspiro[3.4]octan-8-one		433.5
31	3-Methyl-5-(8-oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		363.5
32	5-(5-(4-Fluoro-3-methoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		451.0
33	5-(4,4-Dimethyl-5-oxo-2-thioxo-3-(p-tolyl)imidazolidin-1-yl)-3-(trifluoromethyl)picolinonitrile		405.5
34	6-(8-Oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)nicotinonitrile		417.5
35	4-(7-(5-Cyano-6-(trifluoromethyl)pyridin-2-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		478.5

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
36	5-(8-Oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)quinoline-8-carbonitrile		399.6
37	4-(7-(8-Cyanoquinolin-5-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		460.6
38	5-(5-(4-Fluoro-2-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		437.5
39	5-(8-Oxo-5-(4-(pyridin-3-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		496.1
40	5-(8-Oxo-5-(4-(pyridin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		496.4
41	5-(8-Oxo-5-(4-(pyridin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		496.6
42	5-(8-Oxo-5-(4-(pyrimidin-5-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		496.9

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
43	4-(3-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-5,5-dimethyl-4-oxo-2-thioxoimidazolidin-1-yl)-2-fluoro-N-methylbenzamide		466.5
44	5-(8-Oxo-6-thioxo-5-(4-(trifluoromethoxy)phenyl)-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		487.5
45	5-(8-Oxo-6-thioxo-5-(3-(trifluoromethoxy)phenyl)-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		487.5
46	5-(8-Oxo-6-thioxo-5-(4-(trifluoromethyl)phenyl)-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		471.5
47	4-(3-(6-Cyano-5-methylpyridin-3-yl)-5,5-dimethyl-4-oxo-2-thioxoimidazolidin-1-yl)-2-fluoro-N-methylbenzamide		412.5
48	5-(8-Oxo-5-phenyl-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		403.5
49	5-(5-(3-Fluoro-4-methylphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		435.5

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
50	5-(5-(2-Fluoro-4-methylphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		435.5
51	5-(5-(Isoquinolin-6-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		454.6
52	5-(5-(Isoquinolin-7-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		454.6
53	5-(5-(Cyclohexyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		409.6
54	5-(5-(3-Fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		383.0
55	5-(5-(3-Fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)quinoline-8-carbonitrile		419.0
56	5-(5-(4-Cyano-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		446.6








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
57	3-Chloro-5-(8-oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		383.0
58	4-(7-(5-Chloro-6-cyanopyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		444.0
59	3-Methoxy-5-(8-oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		379.0
60	4-(7-(6-Cyano-5-methoxypyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		440.0
61	6-Thioxo-5-(p-tolyl)-7-(3-(trifluoromethyl)-[2,3'-bipyridin]-5-yl)-5,7-diazaspiro[3.4]octan-8-one		469.0
62	7-(Imidazo[1,2-a]pyridin-6-yl)-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-8-one		363.0
63	5-(5-(4-Hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		419.9

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
64	5-(5-(2-Fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		437.8
65	4-(8-Oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)isoquinoline-1-carbonitrile		399.0
66	4-(7-(1-Cyanoisoquinolin-4-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		460.0
67	5-(5-(3-(Hydroxymethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		433.0
68	5-(5-(2-(Hydroxymethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		433.0
69	5-(5-(4-(Hydroxymethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		433.1
70	5-(5-(4-Cyano-2-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		445.9

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
71	Methyl 4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-3-fluorobenzoate		425.0
72	5-(5-(2,3-Difluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		438.9
73	5-(5-(2,6-Difluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		438.9
74	5-(5-(2,5-Difluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		439.0
75	5-(8-Oxo-6-thioxo-5-(2,3,6-trifluorophenyl)-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		456.9
76	5-(5-(2,4-Difluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		438.9
77	5-(5-(4-Hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		365.0

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
78	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-methylbenzamide		406.1
79	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-methylbenzamide		460.0
80	5-(5-(4-(Methylsulfonyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		481.0
81	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)benzenesulfonamide		482.0
82	4-(8-Oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)pyrazolo[1,5-a]pyridine-7-carbonitrile		388.0
83	4-(7-(7-Cyanopyrazolo[1,5-a]pyridin-4-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		449.0
84	5-(5-(4-Methylbenzyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		431.0

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
85	5-(5-(4-Methylphenethyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		445.0
86	5-(5-(4-(3-Hydroxypropyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		461.1
87	5-(5-(1H-Indazol-5-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		443.7
88	5-(5-(1H-Indazol-6-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		443.7
89	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-3-fluoro-N-methylbenzamide		478.3
90	5-(5-(4-(4-Methylpiperazin-1-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		501.1
91	5-(8-Oxo-5-(4-((2-(pyridin-4-yl)ethyl)sulfonyl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		572.1

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
92	5-(5-(4-((Methyl(pyridin-4-ylmethyl)amino)methyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		537.1
93	5-(5-(4-((4-Methylpiperazin-1-yl)methyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		515.1
94	4-(8-Oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)-1-naphthonitrile		398.6
95	4-(7-(4-Cyanonaphthalen-1-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		459.5
96	3-Methyl-5-(5-(4-(6-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		440.0
97	3-Methyl-5-(5-(4-(4-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		440.1
98	3-Methyl-5-(5-(4-(5-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		440.1








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
99	3-Methyl-5-(5-(4-(2-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		440.0
100	5-(3-(4-Hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-3-(trifluoromethyl)picolinonitrile		407.0
101	5-(3-(4-Hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-3-methylpicolinonitrile		353.0
102	3-Methyl-5-(5-(4-(methyl(1-methylpiperidin-4-yl)amino)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		475.1
103	3-Methyl-5-(8-oxo-5-(4-(pyrimidin-5-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		427.0
104	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-3-fluoro-N-methylbenzamide		424.0
105	5-(5-(4-(5-Fluoropyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		444.0








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
106	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N,2-dimethylbenzamide		419.9
107	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-methoxy-N-methylbenzamide		436.0
108	2-Chloro-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-methylbenzamide		440.0
109	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-methyl-2-(trifluoromethyl)benzamide		474.1
110	5-(5-(Naphthalen-1-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		453.0
111	3-Methyl-5-(8-oxo-5-(1-oxoisindolin-5-yl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		404.1
112	3-Methyl-5-(8-oxo-5-(4-(tetrahydro-2H-pyran-4-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		433.0








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
113	4-(7-(6-Cyano-5-(difluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		460.0
114	1-Methyl-4-(8-oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)-1H-pyrrole-2-carbonitrile		352.1
115	5-(8-Oxo-5-(4-(tetrahydro-2H-pyran-4-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		487.1
116	5-(5-(4-(Furan-2-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		415.1
117	5-(3-Fluoro-4-hydroxyphenyl)-6-thioxo-7-(3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-yl)-5,7-diazaspiro[3.4]octan-8-one		453.5
118	5-(8-Oxo-5-(4-(piperazin-1-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		487.1
119	tert-Butyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)piperazine-1-carboxylate		587.2

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
120	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-methylbenzenesulfonamide		518.0
121	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N,N-dimethylbenzenesulfonamide		532.0
122	Methyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)butanoate		503.0
123	5-(5-(4-Fluoro-3-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		437.5
124	5-(5-(4-(3-(4-Methylpiperazin-1-yl)propyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		543.1
125	5-(5-(Benzo[d]oxazol-6-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		444.0
126	5-(5-(3-Fluoro-4-(2-(1-methyl-1H-pyrazol-5-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		545.5







Cmpd.	Name	Structure	LCMS * [M+1] ⁺
127	5-(5-(3-Fluoro-4-(2-(pyrrolidin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		534.9
128	5-(5-(4-(Benzyloxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		527.6
129	5-(5-(4-((1-Methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		516.0
130	5-(5-(3-Fluoro-4-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		542.6
131	5-(5-(3-Fluoro-4-(2-methoxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		495.5
132	Ethyl 2-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorophenoxy)acetate		523.5
133	5-(5-(3-Fluoro-4-(4-methoxybutoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		523.6








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
134	5-(5-(3-Fluoro-4-((4,4,5,5,5-pentafluoropentyl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		597.7
135	5-(5-(4-(3-(Benzyloxy)propoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		585.5
136	5-(5-(4-(4-(Benzyloxy)butoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		599.8
137	5-(5-(3-Fluoro-4-(2-methylphenethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		555.8
138	5-(5-(3-Fluoro-4-(3-phenylpropoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		555.8
139	5-(5-(3-Fluoro-4-(3,3,4,4,4-pentafluorobutoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		583.4
140	5-(5-(3-Fluoro-4-(2-(naphthalen-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		591.2








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
141	5-(5-(3-Fluoro-4-(2-(naphthalen-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		591.2
142	5-(5-(3-Fluoro-4-(2-(pyridin-4-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		542.7
143	5-(5-(3-Fluoro-4-(2-morpholinoethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		550.6
144	5-(5-(3-Fluoro-4-(2-(pyridin-3-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		542.5
145	5-(5-(3-Fluoro-4-(3-(pyridin-3-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		556.5
146	5-(5-(4-(2-(1H-Pyrrol-1-yl)ethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		530.5
147	5-(5-(3-Fluoro-4-(pyridin-2-ylmethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		528.8








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
148	5-(5-(3-Fluoro-4-(pyridin-3-ylmethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		528.5
149	5-(5-(3-Fluoro-4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		563.8
150	5-(5-(3-Fluoro-4-(3-morpholinopropoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		564.7
151	5-(5-(3-Fluoro-4-(4-((4,4,5,5,5-pentafluoropentyl)thio)butoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		685.5
152	5-(5-(3-Fluoro-4-(4-((4,4,5,5,5-pentafluoropentyl)sulfinyl)butoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		701.5
153	5-(5-(3-Fluoro-4-(3-((4,4,5,5,5-pentafluoropentyl)thio)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		671.8
154	5-(5-(3-Fluoro-4-(3-((4,4,5,5,5-pentafluoropentyl)sulfinyl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		687.5








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
155	5-(4-(2-(Pyridin-2-yl)ethoxy)phenyl)-6-thioxo-7-(3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-yl)-5,7-diazaspiro[3.4]octan-8-one		540.6
156	5-(5-(3-Fluoro-4-(pyridin-4-ylmethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		528.5
157	5-(5-(3-Fluoro-4-(3-(4-methylpiperazin-1-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		577.5
158	5-(5-(4-Fluoro-3-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		542.1
159	5-(5-(4-Fluoro-3-(2-(pyridin-3-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		542.0
160	5-(5-(4-Fluoro-3-(2-(pyridin-4-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		542.5
161	5-(5-(3-(Benzyloxy)-4-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		527.5

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
162	5-(5-(4-Fluoro-3-phenethoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		541.5
163	5-(5-(4-Fluoro-3-(3-phenylpropoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		555.5
164	5-(5-(4-Fluoro-3-(2-morpholinoethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		550.1
165	5-(5-(4-Fluoro-3-(3-morpholinopropoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		564.7
166	5-(5-(4-Fluoro-3-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		563.5
167	5-(5-(4-Fluoro-3-(3-(4-methylpiperazin-1-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		577.6








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
168	5-(5-(3-Fluoro-4-(2-methoxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		441.6
169	5-(5-(3-Fluoro-4-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		488.7
170	5-(5-(3-Fluoro-4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		509.8
171	5-(5-(3-Fluoro-4-(2-methoxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)quinoline-8-carbonitrile		477.6
172	5-(5-(3-Fluoro-4-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)quinoline-8-carbonitrile		524.6
173	5-(5-(3-Fluoro-4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)quinoline-8-carbonitrile		545.6
174	5-(5-(3-Fluoro-4-(2-(piperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		549.9

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
175	5-(5-(3-Fluoro-4-(2-thiomorpholinoethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		566.8
176	5-(5-(3-Fluoro-4-(2-(pyrazin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		543.8
177	5-(5-(3-Fluoro-4-(2-(piperidin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		548.8
178	5-(5-(4-(2-(4-Methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		545.8
179	5-(5-(4-(2-Cyclohexylethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		547.9
180	5-(5-(3-Fluoro-4-(2-(tetrahydro-2H-pyran-4-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		549.8
181	5-(5-(2-Fluoro-4-(2-methoxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		495.7








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
182	5-(5-(2-Fluoro-4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		563.9
183	5-(5-(2-Fluoro-4-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		542.0
184	5-(5-(2-Fluoro-4-(3-(4-methylpiperazin-1-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		577.1
185	5-(5-(3-Fluoro-4-(3-(pyrrolidin-1-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		548.1
186	5-(5-(4-(2-(1,1-Dioxidothiomorpholino)ethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		598.0
187	5-(5-(2-Fluoro-4-(2-(piperidin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		548.1
188	5-(5-(3-Fluoro-4-phenethoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		541.1








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
189	5-(5-(3-Fluoro-4-(2-(pyrimidin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		543.1
190	5-(5-(2-Fluoro-4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		534.0
191	3-Methyl-5-(5-(4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		462.1
192	5-(5-(3-Fluoro-4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		480.1
193	5-(5-(3-Fluoro-4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		534.2
194	3-Methyl-5-(8-oxo-5-(4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		449.0
195	3-Methyl-5-(8-oxo-5-(4-((tetrahydro-2H-thiopyran-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		465.0

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
196	5-(4,4-Dimethyl-3-(4-((1-methylpiperidin-4-yl)oxy)phenyl)-5-oxo-2-thioxoimidazolidin-1-yl)-3-(trifluoromethyl)picolinonitrile		504.1
197	5-(4,4-Dimethyl-3-(4-((1-methylpiperidin-4-yl)oxy)phenyl)-5-oxo-2-thioxoimidazolidin-1-yl)-3-methylpicolinonitrile		450.1
198	5-(8-Oxo-5-(4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		503.0
199	3-Methyl-5-(8-oxo-6-thioxo-5-(4-(2-(4-(2,2,2-trifluoroethyl)piperazin-1-yl)ethoxy)phenyl)-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		559.1
200	3-Methyl-5-(5-(4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		491.2
201	5-(5-(4-((1,1-Dioxidotetrahydro-2H-thiopyran-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		497.0
202	5-(5-(4-(2-(4-Acetylpiperazin-1-yl)ethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		591.9

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
203	5-(5-(3-Fluoro-4-methoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		451.4
204	5-(5-(4-Methoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		433.8
205	5-(5-(4-Methoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		379.0
206	5-(8-Oxo-5-(4-(pyrimidin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		497.8
207	5-(8-Oxo-5-(4-(pyrazin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		496.9
208	5-(8-Oxo-5-(4-(pyrimidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		497.1
209	3-Methyl-5-(8-oxo-5-(4-(pyrimidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		443.0

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
210	3-Methyl-5-(8-oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile		448.1
211	5-(8-Oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		502.2
212	5-(8-Oxo-5-(4-((1-propionylpiperidin-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		558.1
213	5-(5-(4-((1-isobutyrylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		572.1
214	ethyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidine-1-carboxylate		574.0
215	5-(5-(4-((1-acetylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		490.0

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
216	5-(5-(4-((1-Ethylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		476.1
217	5-(5-(4-((1-(2-Hydroxyethyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		492.1
218	5-(5-(4-((1-(2-hydroxyethyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		546.1
219	5-(5-(4-((1-(methylsulfonyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		580.1
220	5-(5-(4-((1-Isopropylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		544.0
221	Ethyl 2-(4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidin-1-yl)acetate		588.1
222	4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidine-1-carboxamide		545.2








Cmpd.	Name	Structure	LCMS * [M+1] ⁺
223	5-(5-(4-(2-Hydroxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile		409.0
224	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid		
225	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid		411.0
226	5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid		465.4
227	4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)butanoic acid		489.0
228	2-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorophenoxy)acetic acid		495.5
229	5-(7-(6-Carbamoyl-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid		483.4

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
230	4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)-N-methylbutanamide		502.0
231	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		464.0
232	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		410.0
233	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-methylbenzamide		406.1
234	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyrrolidin-1-yl)ethyl)benzamide		561.1
235	N-Benzyl-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		554.5
236	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-2-yl)ethyl)benzamide		569.5

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
237	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(pyrrolidin-1-yl)propyl)benzamide		575.1
238	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-4-yl)ethyl)benzamide		569.5
239	5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-2-yl)ethyl)benzamide		569.5
240	5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-3-yl)ethyl)benzamide		569.5
241	5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-4-yl)ethyl)benzamide		569.5
242	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-morpholinoethyl)benzamide		577.5
243	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(4-methylpiperazin-1-yl)ethyl)benzamide		590.5

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
244	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-3-yl)ethyl)benzamide		569.5
245	Ethyl 2-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamido)acetate		550.5
246	Ethyl 3-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamido)propanoate		564.5
247	Ethyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamido)butanoate		578.5
248	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-methoxyethyl)benzamide		522.5
249	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-methoxypropyl)benzamide		536.5
250	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-methoxyethyl)-N-methylbenzamide		536.5

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
251	5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyrrolidin-1-yl)ethyl)benzamide		561.6
252	5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(pyrrolidin-1-yl)propyl)benzamide		575.1
253	5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-morpholinoethyl)benzamide		577.6
254	5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(4-methylpiperazin-1-yl)ethyl)benzamide		590.6
255	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-phenylbenzamide		540.5
256	N-(2-(4-Bromo-1-methyl-1H-pyrazol-5-yl)ethyl)-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		652.2
257	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-phenethylbenzamide		568.5

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
258	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-phenylpropyl)benzamide		582.5
259	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-4-ylmethyl)benzamide		555.5
260	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(thiophen-2-ylmethyl)benzamide		560.5
261	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-2-yl)benzamide		541.5
262	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-3-yl)benzamide		541.4
263	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-4-yl)benzamide		541.4
264	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-2-ylmethyl)benzamide		555.9

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
265	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-3-ylmethyl)benzamide		556.1
266	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(furan-2-ylmethyl)benzamide		544.4
267	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-cyclopentyl-2-fluorobenzamide		532.5
268	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-morpholinopropyl)benzamide		591.0
269	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(4-methylpiperazin-1-yl)propyl)benzamide		604.0
270	5-(5-(3-Fluoro-4-(pyrrolidine-1-carbonyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		518.0
271	5-(5-(3-Fluoro-4-(morpholine-4-carbonyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		534.0

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
272	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-phenylbenzamide		486.8
273	N-Benzyl-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		500.8
274	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(thiophen-2-ylmethyl)benzamide		506.8
275	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(pyrrolidin-1-yl)propyl)benzamide		521.9
276	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(1-methylpiperidin-4-yl)benzamide		561.1
277	N-Butyl-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		520.0
278	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-propylbenzamide		506.1

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
279	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-isopropylbenzamide		506.1
280	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-isobutylbenzamide		520.0
281	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(prop-2-yn-1-yl)benzamide		502.0
282	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-isopentylbenzamide		532.1
283	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-(cyclopentylmethyl)-2-fluorobenzamide		546.0
284	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-cyclopropyl-2-fluorobenzamide		504.0
285	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-hydroxy-2-methylpropyl)benzamide		519.0 (-OH)

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
286	N-(tert-Butyl)-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		520.0
287	N-(2-Chlorobenzyl)-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		588.4
288	N-(3-Chlorobenzyl)-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		588.4
289	N-(4-Chlorobenzyl)-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		588.4
290	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyrazin-2-ylmethyl)benzamide		556.1
291	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-((5-methylfuran-2-yl)methyl)benzamide		558.2
292	4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)-N-phenylbutanamide		564.0

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
293	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(4-fluorobenzyl)benzamide		518.1
294	N-(2-Chlorophenyl)-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		520.1
295	N-(3-Chlorophenyl)-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		520.1
296	N-(4-Chlorophenyl)-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide		520.1
297	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-fluorophenyl)benzamide		504.2
298	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-fluorophenyl)benzamide		504.1
299	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(4-fluorophenyl)benzamide		503.1

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
300	4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(oxazol-2-yl)benzamide		477.1
301	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(thiazol-2-yl)benzamide		493.1
302	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(1-methyl-1H-pyrazol-5-yl)benzamide		490.9
303	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(1-methyl-1H-pyrazol-3-yl)benzamide		490.2
304	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(6-methylpyridin-3-yl)benzamide		501.2
305	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(5-fluoropyridin-3-yl)benzamide		505.1
306	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-hydroxypropyl)benzamide		521.9

Cmpd.	Name	Structure	LCMS * [M+1] ⁺
307	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-(3-(cyclopentyl(methyl)amino)propyl)-2-fluorobenzamide		603.2
308	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(2-oxopyrrolidin-1-yl)propyl)benzamide		589.0
309	5-(5-(3-Fluoro-4-(2-(4-methylpiperazin-1-yl)-2-oxoethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile		577.1
310	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(tetrahydro-2H-pyran-4-yl)benzamide		494.1
311	4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-((tetrahydro-2H-pyran-4-yl)methyl)benzamide		508.0
312	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-6,8-dioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide		462.6
313	4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-(3-(3,3-difluoropyrrolidin-1-yl)propyl)-2-fluorobenzamide		611.1

* mass spectrometric data

[00207] Compounds disclosed herein are AR modulators. In specific embodiments, compounds disclosed herein are AR inverse agonists, AR antagonists, AR degraders, AR trafficking modulators and/or AR DNA-binding inhibitors. In some embodiments, compounds disclosed herein are AR inverse agonists. In some embodiments, compounds disclosed herein are AR antagonists. In some embodiments, compounds disclosed herein are AR degraders. In some embodiments, compounds disclosed herein are AR trafficking modulators. In some embodiments, compounds disclosed herein are AR DNA-binding inhibitors. The overall profile of an AR modulator for the treatment of prostate cancer includes one or more of the foregoing profiles of an AR modulator.

[00208] In some embodiments, compounds disclosed herein have the following properties: full AR antagonist in prostate cancer cells, inverse AR agonist in prostate cancer cells, no AR agonist activity in prostate cancer cells, AR Degradation activity in prostate cancer cells, no antagonist or agonist or degradation activity in non- prostate cancer cells, inhibition of prostate cancer growth.

[00209] In some embodiments, compounds disclosed herein have the following properties: full AR antagonist in prostate cancer cells, inverse AR agonist in prostate cancer cells, no AR agonist activity in prostate cancer cells, AR Degradation activity in prostate cancer cells, and inhibition of prostate cancer growth.

[00210] In some embodiments, compounds disclosed herein have the following properties: full AR antagonist in prostate cancer cells, inverse AR agonist in prostate cancer cells, no AR agonist activity in prostate cancer cells, inhibition of prostate cancer growth

[00211] In some embodiments, compounds disclosed herein have the following properties: full AR antagonist in prostate cancer cells, no AR agonist activity in prostate cancer cells, inhibition of prostate cancer growth

[00212] In some embodiments, compounds disclosed herein have the following properties: AR Trafficking Modulator, no AR agonist activity in prostate cancer cells, no antagonist or agonist or degradation activity in non- prostate cancer cells, inhibition of prostate cancer growth

[00213] In some embodiments, compounds disclosed herein have the following properties: AR Trafficking Modulator, no AR agonist activity in prostate cancer cells, inhibition of prostate cancer growth.

[00214] In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) has minimal pro-convulsant activity and/or minimal impact on seizure threshold.

[00215] In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) displays minimal modulation of the GABA-gated chloride channel.

[00216] In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) displays minimal binding to the GABA-gated chloride channel.

[00217] In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) has minimal antagonism of the GABA-gated chloride channel.

[00218] In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is an androgen receptor modulator with minimal interaction with a GABA-gated chloride channel.

[00219] In some embodiments, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is an androgen receptor modulator with minimal interaction with the GABA_A-gated chloride channel.

[00220] GABA assays are known and include, but are not limited to, those described in Ashok K. Mehta and Maharaj K. Ticku "Characterization of the Picrotoxin Site of GABA_A Receptors" *Current Protocols in Pharmacology* (2000) 1.18.1-1.18.17; Copyright © 2000 by John Wiley & Sons, Inc., which is herein incorporated by reference.

[00221] In some embodiments, described herein is a method of treating cancer in a mammal comprising administering to the mammal a therapeutically effective amount of a compound, wherein the compound: a) is an androgen receptor inverse agonist; androgen receptor antagonist; is an androgen receptor degrader; is an androgen receptor trafficking modulator; is an androgen receptor DNA-binding inhibitor; or combinations thereof; and b) has minimal pro-convulsant activity and/or minimal impact on seizure threshold; displays minimal modulation of the GABA-gated chloride channel; displays minimal binding to the GABA-gated chloride channel; has minimal antagonism of the GABA-gated chloride channel; has minimal interaction with a GABA-gated chloride channel; or combinations thereof.

[00222] In some embodiments, the compound has minimal interaction with the GABA_A-gated chloride channel. In some embodiments, the cancer is prostate cancer. In some embodiments, the cancer is hormone refractory prostate cancer. In some embodiments, the compound is a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X), or a pharmaceutically acceptable salt thereof or N-oxide thereof.

[00223] In some embodiments, described herein is a method of identifying an androgen receptor modulator comprising: 1) testing a compound for androgen receptor modulatory activity in an appropriate assay; and 2) testing the same compound for activity on the GABA-gated chloride channel in an appropriate in vitro or in vivo assay; wherein the compound is an androgen receptor modulator if it exhibits activity in 1) and exhibits any one of the following in 2): displays minimal modulation of the GABA-gated chloride channel; displays minimal binding to the GABA-gated chloride channel; has minimal antagonism of the GABA-gated chloride channel; or has minimal interaction with the GABA-gated chloride channel.


Synthesis of Compounds

[00224] Compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) described herein are synthesized using standard synthetic techniques or using methods known in the art in combination with methods described herein. In additions, solvents, temperatures and other reaction conditions presented herein may vary.

[00225] The starting material used for the synthesis of the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are either synthesized or obtained from commercial sources, such as, but not limited to, Sigma-Aldrich, Fluka, Acros Organics, Alfa Aesar, and the like. The compounds described herein, and other related compounds having different substituents are synthesized using techniques and materials described herein or otherwise known, including those found in March, ADVANCED ORGANIC CHEMISTRY 4th Ed., (Wiley 1992); Carey and Sundberg, ADVANCED ORGANIC CHEMISTRY 4th Ed., Vols. A and B (Plenum 2000, 2001), and Green and Wuts, PROTECTIVE GROUPS IN ORGANIC SYNTHESIS 3rd Ed., (Wiley 1999). General methods for the preparation of compounds can be modified by the use of appropriate reagents and conditions for the introduction of the various moieties found in the formulae as provided herein.

[00226] In some embodiments, the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are prepared as outlined in the following Scheme.

Scheme 1:




[00227] In some embodiments, amines of structure I are treated with ketones or aldehydes $R^1-C(=O)-R^1$ in the presence of NaCN, in a suitable solvent to provide compounds of structure II. In some embodiments, the suitable solvent is acetic acid. In some embodiments, the reaction is performed at a temperature from about 25°C to about 80°C.

[00228] In some embodiments, compounds of structure II are treated with thiophosgene and compounds of structure III in a suitable solvent, followed by treated with an acid to provide thiohydantions of structure IV. In some embodiments, the suitable solvent is dimethylacetamide. In some embodiments, the reaction is heated to about 60°C. In some embodiments, treatment with an acid encompasses treatment with hydrochloric acid. In some embodiments, treatment with an acid encompasses treatment with HCl, MeOH, at reflux for 2h.

[00229] In some embodiments, compounds disclosed herein are prepared as outlined in Scheme 2.


Scheme 2



[00230] In some embodiments, compounds of structure V are elaborated into compounds of structure VI by reacting compounds of structure V with electrophiles or nucleophiles in the presence of a coupling agent. For example, in some embodiments, compounds of structure V are treated with compounds such as R^6-L^2-OH in the presence of DIAD and PPh_3 in a suitable solvent to provide compounds of structure VI. In some embodiments, the suitable solvent is tetrahydrofuran.

[00231] In some embodiments, compounds disclosed herein are prepared as outlined in Scheme 3.

Scheme 3



[00232] Treatment of esters of structure **VII** with a suitable base in a suitable solvent provides carboxylic acids of structure **VIII**. In some embodiments, the suitable base is lithium hydroxide. In some embodiments, the suitable solvent is tetrahydrofuran. Carboxylic acids of structure **VIII** are then coupled with a variety of agents to provide compounds disclosed herein.

5 In some embodiments, carboxylic acids of structure **VIII** are reacted with amines $R^6-L^2-NH_2$ in the presence of a coupling reagent to provide amides of structure IX. In some embodiments, carboxylic acids of structure **VIII** are reacted with alcohols R^6-L^2-OH in the presence of a coupling reagent to provide esters as described herein. In some embodiments, the coupling reagent is EDC, DCC, BOP, HATU or the like. In some embodiments, the coupling reaction is

10 performed in a solvent selected from dichloromethane, dichloroethane, tetrahydrofuran, dimethoxyethane, dimethylformamide or the like in the presence of a base. Suitable bases include, but are not limited to, triethylamine, diisopropylethylamine, N-methylmorpholine, pyridine or the like.

[00233] In one aspect, compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are synthesized as outlined in the Examples.

[00234] Throughout the specification, groups and substituents thereof are chosen by one skilled in the field to provide stable moieties and compounds.

[00235] A detailed description of techniques applicable to the creation of protecting groups and their removal are described in Greene and Wuts, Protective Groups in Organic Synthesis, 3rd

20 Ed., John Wiley & Sons, New York, NY, 1999, and Kocienski, Protective Groups, Thieme Verlag, New York, NY, 1994, which are incorporated herein by reference for such disclosure.

Further Forms of Compounds

[00236] In one aspect, compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) possess one or more stereocenters and each stereocenter exists

25 independently in either the R or S configuration. The compounds presented herein include all diastereomeric, enantiomeric, and epimeric forms as well as the appropriate mixtures thereof. The compounds and methods provided herein include all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers as well as the appropriate mixtures thereof. In certain embodiments, compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and

30 (X) are prepared as their individual stereoisomers by reacting a racemic mixture of the compound with an optically active resolving agent to form a pair of diastereoisomeric compounds/salts, separating the diastereomers and recovering the optically pure enantiomers. In some embodiments, resolution of enantiomers is carried out using covalent diastereomeric derivatives of the compounds described herein. In another embodiment, diastereomers are

35 separated by separation/resolution techniques based upon differences in solubility. In other

embodiments, separation of stereoisomers is performed by chromatography or by the forming diastereomeric salts and separation by recrystallization, or chromatography, or any combination thereof. Jean Jacques, Andre Collet, Samuel H. Wilen, "Enantiomers, Racemates and Resolutions", John Wiley And Sons, Inc., 1981. In some embodiments, stereoisomers are
5 obtained by stereoselective synthesis.

[00237] The methods and compositions described herein include the use of amorphous forms as well as crystalline forms (also known as polymorphs). In one aspect, compounds described herein are in the form of pharmaceutically acceptable salts. As well, active metabolites of these compounds having the same type of activity are included in the scope of the present disclosure.

10 In addition, the compounds described herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein.

[00238] In some embodiments, compounds described herein are prepared as prodrugs. A "prodrug" refers to an agent that is converted into the parent drug *in vivo*. Prodrugs are often
15 useful because, in some situations, they may be easier to administer than the parent drug. They may, for instance, be bioavailable by oral administration whereas the parent is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. In some embodiments, the design of a prodrug increases the effective water solubility. An example, without limitation, of a prodrug is a compound described herein, which is
20 administered as an ester (the "prodrug") to facilitate transmittal across a cell membrane where water solubility is detrimental to mobility but which then is metabolically hydrolyzed to the carboxylic acid, the active entity, once inside the cell where water-solubility is beneficial. A further example of a prodrug might be a short peptide (polyaminoacid) bonded to an acid group where the peptide is metabolized to reveal the active moiety. In certain embodiments, upon *in*
25 *vivo* administration, a prodrug is chemically converted to the biologically, pharmaceutically or therapeutically active form of the compound. In certain embodiments, a prodrug is enzymatically metabolized by one or more steps or processes to the biologically, pharmaceutically or therapeutically active form of the compound.

[00239] In one aspect, prodrugs are designed to alter the metabolic stability or the transport
30 characteristics of a drug, to mask side effects or toxicity, to improve the flavor of a drug or to alter other characteristics or properties of a drug. By virtue of knowledge of pharmacokinetic, pharmacodynamic processes and drug metabolism *in vivo*, once a pharmaceutically active compound is known, the design prodrugs of the compound is possible. (see, for example, Nogrady (1985) *Medicinal Chemistry A Biochemical Approach*, Oxford University Press, New
35 York, pages 388-392; Silverman (1992), *The Organic Chemistry of Drug Design and Drug*

Action, Academic Press, Inc., San Diego, pages 352-401, Rooseboom *et al.*, *Pharmacological Reviews*, 56:53-102, 2004; Aesop Cho, "Recent Advances in Oral Prodrug Discovery", *Annual Reports in Medicinal Chemistry*, Vol. 41, 395-407, 2006; T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems*, Vol. 14 of the A.C.S. Symposium Series).

5 [00240] Prodrug forms of the herein described compounds, wherein the prodrug is metabolized *in vivo* to produce a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) as set forth herein are included within the scope of the claims. In some cases, some of the herein-described compounds may be a prodrug for another derivative or active compound.

10 [00241] In some embodiments, sites on the aromatic ring portion of compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are susceptible to various metabolic reactions. Therefore incorporation of appropriate substituents on the aromatic ring structures will reduce, minimize or eliminate this metabolic pathway. In specific embodiments, the appropriate substituent to decrease or eliminate the susceptibility of the aromatic ring to metabolic reactions is, by way of example only, a halogen, or an alkyl group.

15 [00242] In another embodiment, the compounds described herein are labeled isotopically (e.g. with a radioisotope) or by another other means, including, but not limited to, the use of chromophores or fluorescent moieties, bioluminescent labels, or chemiluminescent labels.

[00243] Compounds described herein include isotopically-labeled compounds, which are
20 identical to those recited in the various formulae and structures presented herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into the present compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, fluorine and chlorine, such as, for example, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O ,
25 ^{35}S , ^{18}F , ^{36}Cl . In one aspect, isotopically-labeled compounds described herein, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. In one aspect, substitution with isotopes such as deuterium affords certain therapeutic advantages resulting from greater metabolic stability, such as, for example, increased *in vivo* half-life or reduced dosage requirements.

30 [00244] In additional or further embodiments, the compounds described herein are metabolized upon administration to an organism in need to produce a metabolite that is then used to produce a desired effect, including a desired therapeutic effect.

[00245] "Pharmaceutically acceptable," as used herein, refers a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compound, and is
35 relatively nontoxic, i.e., the material may be administered to an individual without causing

undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

[00246] The term “pharmaceutically acceptable salt” refers to a formulation of a compound that does not cause significant irritation to an organism to which it is administered and does not abrogate the biological activity and properties of the compound. In some embodiments, pharmaceutically acceptable salts are obtained by reacting a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) with acids. Pharmaceutically acceptable salts are also obtained by reacting a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) with a base to form a salt.

[00247] Compounds described herein may be formed as, and/or used as, pharmaceutically acceptable salts. The type of pharmaceutical acceptable salts, include, but are not limited to: (1) acid addition salts, formed by reacting the free base form of the compound with a pharmaceutically acceptable: inorganic acid, such as, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, metaphosphoric acid, and the like; or with an organic acid, such as, for example, acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, trifluoroacetic acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, toluenesulfonic acid, 2-naphthalenesulfonic acid, 4-methylbicyclo-[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, butyric acid, phenylacetic acid, phenylbutyric acid, valproic acid, and the like; (2) salts formed when an acidic proton present in the parent compound is replaced by a metal ion, e.g., an alkali metal ion (e.g. lithium, sodium, potassium), an alkaline earth ion (e.g. magnesium, or calcium), or an aluminum ion. In some cases, compounds described herein may coordinate with an organic base, such as, but not limited to, ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine, dicyclohexylamine, tris(hydroxymethyl)methylamine. In other cases, compounds described herein may form salts with amino acids such as, but not limited to, arginine, lysine, and the like. Acceptable inorganic bases used to form salts with compounds that include an acidic proton, include, but are not limited to, aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate, sodium hydroxide, and the like.

[00248] It should be understood that a reference to a pharmaceutically acceptable salt includes the solvent addition forms or crystal forms thereof, particularly solvates or polymorphs.

Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and may be formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, and the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Solvates of compounds described herein can be

5 conveniently prepared or formed during the processes described herein. In addition, the compounds provided herein can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein.

[00249] Compounds described herein, such as compounds of Formula (I), (Ia), (II), (III), (IV),
10 (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X), may be in various forms, including but not limited to, amorphous forms, milled forms and nano-particulate forms. In addition, compounds described herein include crystalline forms, also known as polymorphs. Polymorphs include the different crystal packing arrangements of the same elemental composition of a compound. Polymorphs usually have different X-ray diffraction patterns, melting points, density, hardness,
15 crystal shape, optical properties, stability, and solubility. Various factors such as the recrystallization solvent, rate of crystallization, and storage temperature may cause a single crystal form to dominate.

Certain Terminology

[00250] Unless otherwise stated, the following terms used in this application, including the
20 specification and claims, have the definitions given below. It must be noted that, as used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. Unless otherwise indicated, conventional methods of mass spectroscopy, NMR, HPLC, protein chemistry, biochemistry, recombinant DNA techniques and pharmacology are employed. In this application, the use of “or” or “and”
25 means “and/or” unless stated otherwise. Furthermore, use of the term “including” as well as other forms, such as “include”, “includes,” and “included,” is not limiting. The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

[00251] An “alkyl” group refers to an aliphatic hydrocarbon group. The alkyl group may be a
30 saturated alkyl group or the the alkyl group may be an unsaturated alkyl group. The alkyl moiety, whether saturated or unsaturated, may be branched or straight chain. The “alkyl” group may have 1 to 10 carbon atoms (whenever it appears herein, a numerical range such as “1 to 10” refers to each integer in the given range; *e.g.*, “1 to 10 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 10
35 carbon atoms, although the present definition also covers the occurrence of the term “alkyl”

where no numerical range is designated). In one aspect the alkyl is selected from the group consisting of methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and t-butyl. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tertiary butyl, pentyl, neopentyl, hexyl, allyl, but-2-enyl, but-3-enyl, and the like. In some embodiments, an alkyl is a C₁-C₆alkyl.

[00252] The term “alkylene” refers to a divalent alkyl radical. Any of the above mentioned monovalent alkyl groups may be an alkylene by abstraction of a second hydrogen atom from the alkyl. In one aspect, an alkylene is a C₁-C₁₀alkylene. In another aspect, an alkylene is a C₁-C₆alkylene. Typical alkylene groups include, but are not limited to, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH₂CH₂-, -CH₂CH(CH₃)-, -CH₂C(CH₃)₂-, -CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂CH₂-, and the like.

[00253] An “alkoxy” group refers to a (alkyl)O- group, where alkyl is as defined herein.

[00254] The term “alkylamine” refers to the -N(alkyl)_xH_y group, where x and y are selected from the group x=1, y=1 and x=2, y=0.

[00255] The term “aromatic” refers to a planar ring having a delocalized π -electron system containing $4n+2$ π electrons, where n is an integer. Aromatic rings can be formed from five, six, seven, eight, nine, ten, or more than ten atoms. Aromatics are optionally substituted. The term “aromatic” includes both carbocyclic aryl (“aryl”, *e.g.*, phenyl) and heterocyclic aryl (or “heteroaryl” or “heteroaromatic”) groups (*e.g.*, pyridine). The term includes monocyclic or fused-ring polycyclic (*i.e.*, rings which share adjacent pairs of carbon atoms) groups.

[00256] The term “carbocyclic” or “carbocycle” refers to a ring or ring system where the atoms forming the backbone of the ring are all carbon atoms. The term thus distinguishes carbocyclic from heterocyclic rings in which the ring backbone contains at least one atom which is different from carbon.

[00257] As used herein, the term “aryl” refers to an aromatic ring wherein each of the atoms forming the ring is a carbon atom. Aryl rings are formed by five, six, seven, eight, nine, or more than nine carbon atoms. Aryl groups are optionally substituted. In one aspect, an aryl is a phenyl or a naphthalenyl. In one aspect, an aryl is a phenyl. In one aspect, an aryl is a C₆-C₁₀aryl. Depending on the structure, an aryl group can be a monoradical or a diradical (*i.e.*, an arylene group). Exemplary arylenes include, but are not limited to, phenyl-1,2-ene, phenyl-1,3-ene, and phenyl-1,4-ene.

[00258] The term “cycloalkyl” refers to a monocyclic or polycyclic aliphatic, non-aromatic radical, wherein each of the atoms forming the ring (*i.e.* skeletal atoms) is a carbon atom. Cycloalkyls may be saturated, or partially unsaturated. Cycloalkyls may be fused with an aromatic ring, and the point of attachment is at a carbon that is not an aromatic ring carbon

atom. Cycloalkyl groups include groups having from 3 to 10 ring atoms. In some embodiments, cycloalkyl groups are selected from among cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, and cyclooctyl. Cycloalkyl groups may be substituted or unsubstituted. Depending on the structure, a cycloalkyl group can be a monoradical or a

5 diradical (i.e., an cycloalkylene group, such as, but not limited to, cyclopropan-1,1-diyl, cyclobutan-1,1-diyl, cyclopentan-1,1-diyl, cyclohexan-1,1-diyl, cyclohexan-1,4-diyl, cycloheptan-1,1-diyl, and the like). In one aspect, a cycloalkyl is a C₃-C₆cycloalkyl.

[00259] The term “halo” or, alternatively, “halogen” or “halide” means fluoro, chloro, bromo or iodo.

10 [00260] The term “haloalkyl” refers to an alkyl group in which one or more hydrogen atoms are replaced by one or more halide atoms. In one aspect, a haloalkyl is a C₁-C₆haloalkyl.

[00261] The term “haloalkylene” refers to an alkylene group in which one or more hydrogen atoms are replaced by one or more halide atoms. In one aspect, a haloalkylene is a C₁-C₆haloalkylene.

15 [00262] The term “fluoroalkyl” refers to an alkyl in which one or more hydrogen atoms are replaced by a fluorine atom. In one aspect, a fluoroalkyl is a C₁-C₆fluoroalkyl.

[00263] The term “fluoroalkylene” refers to an alkylene in which one or more hydrogen atoms are replaced by a fluorine atom. In one aspect, a fluoroalkylene is a C₁-C₆fluoroalkylene.

[00264] The term “heteroalkyl” refers to an alkyl group in which one or more skeletal atoms of the alkyl are selected from an atom other than carbon, *e.g.*, oxygen, nitrogen (*e.g.* -NH-, -N(alkyl)-, sulfur, or combinations thereof. In one aspect, a heteroalkyl is a C₁-C₆heteroalkyl.


[00265] The term “heteroalkylene” refers to an alkylene group in which one or more skeletal atoms of the alkyl are selected from an atom other than carbon, *e.g.*, oxygen, nitrogen, sulfur, phosphorus or combinations thereof. In one aspect, a heteroalkylene is a C₁-C₆heteroalkylene.


25 Exemplary heteroalkylenes include, but are not limited to, -OCH₂-, -OCH(CH₃)-, -OC(CH₃)₂-, -OCH₂CH₂-, -OCH₂CH₂CH₂-, -CH₂O-, -CH(CH₃)O-, -C(CH₃)₂O-, -CH₂CH₂O-, -CH₂OCH₂-, -CH₂OCH₂CH₂-, -CH₂CH₂OCH₂-, -SCH₂-, -SCH(CH₃)-, -SC(CH₃)₂-, -SCH₂CH₂-, -CH₂S-, -CH(CH₃)S-, -C(CH₃)₂S-, -CH₂CH₂S-, -CH₂SCH₂-, -CH₂SCH₂CH₂-, -CH₂CH₂SCH₂-, -SO₂CH₂-, -SO₂CH(CH₃)-, -SO₂C(CH₃)₂-, -SO₂CH₂CH₂-, -CH₂SO₂-, -CH(CH₃)SO₂-, -C(CH₃)₂SO₂-, -CH₂CH₂SO₂-, -CH₂SO₂CH₂-, -CH₂SO₂CH₂CH₂-, -CH₂CH₂SO₂CH₂-, -NHCH₂-, -NHCH(CH₃)-, -NHC(CH₃)₂-, -NHCH₂CH₂-, -CH₂NH-, -CH(CH₃)NH-, -C(CH₃)₂NH-, -CH₂CH₂NH-, -CH₂NHCH₂-, -CH₂NHCH₂CH₂-, -CH₂CH₂NHCH₂-, and the like.

[00266] The term “heterocycle” or “heterocyclic” refers to heteroaromatic rings (also known as heteroaryls) and heterocycloalkyl rings (also known as heteroalicyclic groups) containing one to four heteroatoms in the ring(s), where each heteroatom in the ring(s) is selected from O, S and

N, wherein each heterocyclic group has from 4 to 10 atoms in its ring system, and with the proviso that the any ring does not contain two adjacent O or S atoms. Non-aromatic heterocyclic groups (also known as heterocycloalkyls) include groups having only 3 atoms in their ring system, but aromatic heterocyclic groups must have at least 5 atoms in their ring system. The heterocyclic groups include benzo-fused ring systems. An example of a 3-membered heterocyclic group is aziridinyl. An example of a 4-membered heterocyclic group is azetidiny. An example of a 5-membered heterocyclic group is thiazolyl. An example of a 6-membered heterocyclic group is pyridyl, and an example of a 10-membered heterocyclic group is quinolinyl. Examples of non-aromatic heterocyclic groups are pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, oxazolidinonyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, thioxanyl, piperazinyl, aziridinyl, azetidiny, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepinyl, diazepinyl, thiazepinyl, 1,2,3,6-tetrahydropyridinyl, pyrrolin-2-yl, pyrrolin-3-yl, indolinyl, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazolinyl, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl, 3H-indolyl and quinoliziny. Examples of aromatic heterocyclic groups are pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnoliny, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridinyl, and furopyridinyl. The foregoing groups may be C-attached (or C-linked) or N-attached where such is possible. For instance, a group derived from pyrrole may be pyrrol-1-yl (*N*-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole may be imidazol-1-yl or imidazol-3-yl (both *N*-attached) or imidazol-2-yl, imidazol-4-yl or imidazol-5-yl (all C-attached). The heterocyclic groups include benzo-fused ring systems. Non-aromatic heterocycles may ber substituted with one or two oxo (=O) moieties, such as pyrrolidin-2-one.


[00267] The terms “heteroaryl” or, alternatively, “heteroaromatic” refers to an aryl group that includes one or more ring heteroatoms selected from nitrogen, oxygen and sulfur. Illustrative examples of heteroaryl groups include the following moieties:





and the like. Monocyclic

- 5 heteroaryls include pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, pyridazinyl, triazinyl, oxadiazolyl, thiadiazolyl, and furazanyl. In some embodiments, a heteroaryl contains 0-3 N atoms in the ring. In some embodiments, a heteroaryl contains 1-3 N atoms in the ring. In some embodiments, a heteroaryl contains 0-3 N atoms, 0-1 O atoms, and 0-1 S atoms in the ring. In
- 10 some embodiments, a heteroaryl is a monocyclic or bicyclic heteroaryl. In some embodiments, heteroaryl is a C₁-C₉heteroaryl. In some embodiments, monocyclic heteroaryl is a C₁-C₅heteroaryl. In some embodiments, monocyclic heteroaryl is a 5-membered or 6-membered heteroaryl. In some embodiments, bicyclic heteroaryl is a C₆-C₉heteroaryl. Depending on the structure, a heteroaryl group can be a monoradical or a diradical (i.e., a heteroarylene group).
- 15 **[00268]** A “heterocycloalkyl” or “heteroalicyclic” group refers to a cycloalkyl group that includes at least one heteroatom selected from nitrogen, oxygen and sulfur. The radicals may be fused with an aryl or heteroaryl. Illustrative examples of heterocycloalkyl groups, also referred to as non-aromatic heterocycles, include:




and the like. In some embodiments, the heterocycloalkyl is selected from oxazolidinonyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, and indolinyl. In some embodiments,

the heterocycloalkyl is pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, or piperazinyl. The term heteroalicyclic also includes all ring forms of the carbohydrates, including but not limited to the monosaccharides, the disaccharides and the oligosaccharides. In one aspect, a heterocycloalkyl is a C₂-C₁₀heterocycloalkyl. In another aspect, a heterocycloalkyl is a C₄-C₁₀heterocycloalkyl. In some embodiments, a heterocycloalkyl contains 0-2 N atoms in the ring. In some embodiments, a heterocycloalkyl contains 0-2 N atoms, 0-2 O atoms and 0-1 S atoms in the ring.

[00269] The term “bond” or “single bond” refers to a chemical bond between two atoms, or two moieties when the atoms joined by the bond are considered to be part of larger substructure. In one aspect, when a group described herein is a bond, the referenced group is absent thereby allowing a bond to be formed between the remaining identified groups.

[00270] The term “moiety” refers to a specific segment or functional group of a molecule. Chemical moieties are often recognized chemical entities embedded in or appended to a molecule.

[00271] As used herein, “carboxylic acid bioisostere” refers to a functional group or moiety that exhibits similar physical, biological and/or chemical properties as a carboxylic acid moiety. Examples of carboxylic acid bioisosteres include, but are not limited to,



[00272] The term “optionally substituted” or “substituted” means that the referenced group may be substituted with one or more additional group(s) individually and independently selected from alkyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, alkylthio, arylthio, alkylsulfoxide, arylsulfoxide, alkylsulfone, arylsulfone, cyano, halo, nitro, haloalkyl, fluoroalkyl, fluoroalkoxy, and amino, including mono- and di-substituted amino groups, and the protected derivatives thereof. By way of example optional substituents are independently selected from halide, -CN, -NO₂, and -LR, wherein each L is independently selected from a bond, -O-, -C(=O)-, -C(=O)O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -NHC(=O)-, -C(=O)NH-, S(=O)₂NH-, -NHS(=O)₂-, -OC(=O)NH-, -NHC(=O)O-, or -(C₁-C₆alkylene)-; and each R is selected from H, alkyl, fluoroalkyl, heteroalkyl, cycloalkyl, aryl, heteroaryl, or heterocycloalkyl. In some embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -NH(CH₃), -N(CH₃)₂, -OH, alkyl, fluoroalkyl, heteroalkyl, cycloalkyl, heterocycloalkyl,

aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, alkylsulfoxide, arylsulfoxide, alkylsulfone, and arylsulfone. In some embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -NH(CH₃), -N(CH₃)₂, -OH, -CO₂H, -CO₂alkyl, -C(=O)alkyl, -C(=O)NH₂, -C(=O)NH(alkyl), -C(=O)N(alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(alkyl), -S(=O)₂N(alkyl)₂, alkyl, cycloalkyl, fluoroalkyl, heteroalkyl, alkoxy, fluoroalkoxy, -S-alkyl, or -S(=O)₂alkyl. In some
5 embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -NH(CH₃), -N(CH₃)₂, -OH, -CO₂H, -CO₂alkyl, -C(=O)alkyl, -C(=O)NH₂, -C(=O)NH(alkyl), -C(=O)N(alkyl)₂, alkyl, cycloalkyl, fluoroalkyl, heteroalkyl, alkoxy, and fluoroalkoxy. In some
embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -OH, -
10 NH(CH₃), -N(CH₃)₂, -CH₃, -CH₂CH₃, -CF₃, -CH₂CF₃, -OCH₃, -OCH₂CH₃, and -OCF₃. In some
embodiments, substituted groups are substituted with one or two of the preceding groups. In
some embodiments, an optional substituent on an aliphatic carbon atom (acyclic or cyclic,
saturated or unsaturated carbon atoms, excluding aromatic carbon atoms) includes oxo (=O).

[00273] In certain embodiments, the compounds presented herein possess one or more
15 stereocenters and each center independently exists in either the R or S configuration. The
compounds presented herein include all diastereomeric, enantiomeric, and epimeric forms as
well as the appropriate mixtures thereof. Stereoisomers are obtained, if desired, by methods
such as, stereoselective synthesis and/or the separation of stereoisomers by chiral
chromatographic columns.

[00274] The methods and formulations described herein include the use of *N*-oxides (if
20 appropriate), crystalline forms (also known as polymorphs), or pharmaceutically acceptable
salts of compounds having the structure of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII),
(VIII), (VIIIa), (IX), (IXa) or (X), as well as active metabolites of these compounds having the
same type of activity. In some situations, compounds may exist as tautomers. All tautomers are
25 included within the scope of the compounds presented herein. In specific embodiments, the
compounds described herein exist in solvated forms with pharmaceutically acceptable solvents
such as water, ethanol, and the like. In other embodiments, the compounds described herein
exist in unsolvated form.

[00275] The term “acceptable” with respect to a formulation, composition or ingredient, as used
30 herein, means having no persistent detrimental effect on the general health of the subject being
treated.

[00276] The term “modulate” as used herein, means to interact with a target either directly or
indirectly so as to alter the activity of the target, including, by way of example only, to enhance
the activity of the target, to inhibit the activity of the target, to limit the activity of the target, or
35 to extend the activity of the target.

[00277] The term “modulator” as used herein, refers to a molecule that interacts with a target either directly or indirectly. The interactions include, but are not limited to, the interactions of an agonist, partial agonist, an inverse agonist, antagonist, degrader, AR trafficking modulator, AR DNA-binding inhibitor. In some embodiments, a modulator is an antagonist. In some
5 embodiments, a modulator is an inverse agonist, antagonist, degrader, AR trafficking modulator and/or a DNA binding inhibitor.

[00278] The term “antagonist” as used herein, refers to a small -molecule agent that binds to a nuclear hormone receptor and subsequently decreases the agonist induced transcriptional activity of the nuclear hormone receptor.

10 [00279] The term “agonist” as used herein, refers to a small-molecule agent that binds to a nuclear hormone receptor and subsequently increases nuclear hormone receptor transcriptional activity in the absence of a known agonist.

[00280] The term “inverse agonist” as used herein, refers to a small-molecule agent that binds to a nuclear hormone receptor and subsequently decreases the basal level of nuclear hormone
15 receptor transcriptional activity that is present in the absence of a known agonist.

[00281] The term “degrader” as used herein, refers to a small molecule agent that binds to a nuclear hormone receptor and subsequently lowers the steady state protein levels of said receptor.

[00282] The term “AR trafficking modulator” as used herein, refers to a small-molecule agent
20 that binds to a nuclear hormone receptor and subsequently alters the normal subcellular location of the receptor thereby interfering with its function and signaling.

[00283] The term “DNA-binding inhibitor” as used herein, refers to a small-molecule agent that binds to a nuclear hormone receptor and subsequently prevents DNA binding of the receptor thereby interfering with its function and signaling.

25 [00284] The term “AR-dependent”, as used herein, refers to diseases or conditions that would not occur, or would not occur to the same extent, in the absence of androgen receptors.

[00285] The term “AR-mediated”, as used herein, refers to diseases or conditions that might occur in the absence of androgen receptors but can occur in the presence of androgen receptors.

[00286] “Selective” with respect to androgen receptors means that the compound preferentially
30 binds to androgen receptors versus other nuclear receptors. In some embodiments, a selective androgen receptor modulator preferentially binds to androgen receptors and displays little, if any, affinity to other nuclear receptors.

[00287] The term “cancer” as used herein refers to an abnormal growth of cells which tend to proliferate in an uncontrolled way and, in some cases, to metastasize (spread).

[00288] The terms “co-administration” or the like, as used herein, are meant to encompass administration of the selected therapeutic agents to a single patient, and are intended to include treatment regimens in which the agents are administered by the same or different route of administration or at the same or different time.

5 [00289] The terms “effective amount” or “therapeutically effective amount,” as used herein, refer to a sufficient amount of an agent or a compound being administered which will relieve to some extent one or more of the symptoms of the disease or condition being treated. The result can be reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an “effective amount” for therapeutic
10 uses is the amount of the composition comprising a compound as disclosed herein required to provide a clinically significant decrease in disease symptoms. An appropriate “effective” amount in any individual case may be determined using techniques, such as a dose escalation study.

[00290] The terms “enhance” or “enhancing,” as used herein, means to increase or prolong
15 either in potency or duration a desired effect. Thus, in regard to enhancing the effect of therapeutic agents, the term “enhancing” refers to the ability to increase or prolong, either in potency or duration, the effect of other therapeutic agents on a system. An “enhancing-effective amount,” as used herein, refers to an amount adequate to enhance the effect of another therapeutic agent in a desired system.

20 [00291] The term “pharmaceutical combination” as used herein, means a product that results from the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. The term “fixed combination” means that the active ingredients, e.g. a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) and a co-agent, are both administered to a patient simultaneously in
25 the form of a single entity or dosage. The term “non-fixed combination” means that the active ingredients, e.g. a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) and a co-agent, are administered to a patient as separate entities either simultaneously, concurrently or sequentially with no specific intervening time limits, wherein such administration provides effective levels of the two compounds in the body of the patient.
30 The latter also applies to cocktail therapy, e.g. the administration of three or more active ingredients.

[00292] The terms “kit” and “article of manufacture” are used as synonyms.

[00293] A “metabolite” of a compound disclosed herein is a derivative of that compound that is formed when the compound is metabolized. The term “active metabolite” refers to a
35 biologically active derivative of a compound that is formed when the compound is metabolized.

The term “metabolized,” as used herein, refers to the sum of the processes (including, but not limited to, hydrolysis reactions and reactions catalyzed by enzymes) by which a particular substance is changed by an organism. Thus, enzymes may produce specific structural alterations to a compound. For example, cytochrome P450 catalyzes a variety of oxidative and reductive reactions while uridine diphosphate glucuronyltransferases catalyze the transfer of an activated glucuronic-acid molecule to aromatic alcohols, aliphatic alcohols, carboxylic acids, amines and free sulphydryl groups. Metabolites of the compounds disclosed herein are optionally identified either by administration of compounds to a host and analysis of tissue samples from the host, or by incubation of compounds with hepatic cells in vitro and analysis of the resulting compounds.

[00294] The term “subject” or “patient” encompasses mammals. Examples of mammals include, but are not limited to, any member of the Mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. In one embodiment, the mammal is a human.

[00295] The terms “treat,” “treating” or “treatment,” as used herein, include alleviating, abating or ameliorating at least one symptom of a disease or condition, preventing additional symptoms, inhibiting the disease or condition, e.g., arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or stopping the symptoms of the disease or condition either prophylactically and/or therapeutically.

Routes of Administration

[00296] Suitable routes of administration include, but are not limited to, oral, intravenous, rectal, aerosol, parenteral, ophthalmic, pulmonary, transmucosal, transdermal, vaginal, otic, nasal, and topical administration. In addition, by way of example only, parenteral delivery includes intramuscular, subcutaneous, intravenous, intramedullary injections, as well as intrathecal, direct intraventricular, intraperitoneal, intralymphatic, and intranasal injections.

[00297] In certain embodiments, a compound as described herein is administered in a local rather than systemic manner, for example, via injection of the compound directly into an organ, often in a depot preparation or sustained release formulation. In specific embodiments, long acting formulations are administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Furthermore, in other embodiments, the drug is delivered in a targeted drug delivery system, for example, in a liposome coated with organ-specific antibody. In such embodiments, the liposomes are targeted to and taken up selectively by the organ. In yet other embodiments, the compound as described herein is

provided in the form of a rapid release formulation, in the form of an extended release formulation, or in the form of an intermediate release formulation. In yet other embodiments, the compound described herein is administered topically.

Pharmaceutical Compositions/Formulations

5 [00298] In some embodiments, the compounds described herein are formulated into pharmaceutical compositions. Pharmaceutical compositions are formulated in a conventional manner using one or more pharmaceutically acceptable inactive ingredients that facilitate processing of the active compounds into preparations that can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. A summary of
10 pharmaceutical compositions described herein can be found, for example, in Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery
15 Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999), herein incorporated by reference for such disclosure.

[00299] Provided herein are pharmaceutical compositions that include a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) and at least one pharmaceutically acceptable inactive ingredient. In some embodiments, the compounds
20 described herein are administered as pharmaceutical compositions in which a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is mixed with other active ingredients, as in combination therapy. In other embodiments, the pharmaceutical compositions include other medicinal or pharmaceutical agents, carriers, adjuvants, preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for
25 regulating the osmotic pressure, and/or buffers. In yet other embodiments, the pharmaceutical compositions include other therapeutically valuable substances.

[00300] A pharmaceutical composition, as used herein, refers to a mixture of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) with other chemical components (i.e. pharmaceutically acceptable inactive ingredients), such as carriers,
30 excipients, binders, filling agents, suspending agents, flavoring agents, sweetening agents, disintegrating agents, dispersing agents, surfactants, lubricants, colorants, diluents, solubilizers, moistening agents, plasticizers, stabilizers, penetration enhancers, wetting agents, anti-foaming agents, antioxidants, preservatives, or one or more combination thereof. The pharmaceutical composition facilitates administration of the compound to a mammal.

[00301] A therapeutically effective amount can vary widely depending on the severity of the disease, the age and relative health of the subject, the potency of the compound used and other factors. The compounds can be used singly or in combination with one or more therapeutic agents as components of mixtures.

5 [00302] The pharmaceutical formulations described herein are administered to a subject by appropriate administration routes, including but not limited to, oral, parenteral (e.g., intravenous, subcutaneous, intramuscular), intranasal, buccal, topical, rectal, or transdermal administration routes. The pharmaceutical formulations described herein include, but are not limited to, aqueous liquid dispersions, self-emulsifying dispersions, solid solutions, liposomal
10 dispersions, aerosols, solid dosage forms, powders, immediate release formulations, controlled release formulations, fast melt formulations, tablets, capsules, pills, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate and controlled release formulations.

[00303] Pharmaceutical compositions including a compound of Formula (I), (Ia), (II), (III), (IV),
15 (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) are manufactured in a conventional manner, such as, by way of example only, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

[00304] The pharmaceutical compositions will include at least one compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) as an active ingredient in
20 free-acid or free-base form, or in a pharmaceutically acceptable salt form. In addition, the methods and pharmaceutical compositions described herein include the use of *N*-oxides (if appropriate), crystalline forms, amorphous phases, as well as active metabolites of these compounds having the same type of activity. In some embodiments, compounds described herein exist in unsolvated form or in solvated forms with pharmaceutically acceptable solvents
25 such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein.

[00305] The pharmaceutical compositions described herein, which include a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) are formulated into any suitable dosage form, including but not limited to, aqueous oral dispersions,
30 liquids, gels, syrups, elixirs, slurries, suspensions, solid oral dosage forms, controlled release formulations, fast melt formulations, effervescent formulations, lyophilized formulations, tablets, powders, pills, dragees, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate release and controlled release formulations.

[00306] Pharmaceutical preparations that are administered orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In some embodiments, the push-fit capsules do not include any other ingredient besides the capsule shell and the active ingredient. In soft capsules, the active compounds are dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In some embodiments, stabilizers are added.

[00307] All formulations for oral administration are in dosages suitable for such administration.

[00308] In one aspect, solid oral dosage forms are prepared by mixing a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) with one or more of the following: antioxidants, flavoring agents, and carrier materials such as binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, and diluents.

[00309] In some embodiments, the solid dosage forms disclosed herein are in the form of a tablet, (including a suspension tablet, a fast-melt tablet, a bite-disintegration tablet, a rapid-disintegration tablet, an effervescent tablet, or a caplet), a pill, a powder, a capsule, solid dispersion, solid solution, bioerodible dosage form, controlled release formulations, pulsatile release dosage forms, multiparticulate dosage forms, beads, pellets, granules. In other embodiments, the pharmaceutical formulation is in the form of a powder. In still other embodiments, the pharmaceutical formulation is in the form of a tablet. In other embodiments, pharmaceutical formulations of the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is in the form of a capsule.

[00310] In some embodiments, solid dosage forms, e.g., tablets, effervescent tablets, and capsules, are prepared by mixing particles of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) with one or more pharmaceutical excipients to form a bulk blend composition. The bulk blend is readily subdivided into equally effective unit dosage forms, such as tablets, pills, and capsules. In some embodiments, the individual unit dosages include film coatings. These formulations are manufactured by conventional formulation techniques.

[00311] Conventional formulation techniques include, e.g., one or a combination of methods: (1) dry mixing, (2) direct compression, (3) milling, (4) dry or non-aqueous granulation, (5) wet granulation, or (6) fusion. Other methods include, e.g., spray drying, pan coating, melt granulation, granulation, fluidized bed spray drying or coating (e.g., wurster coating), tangential coating, top spraying, tableting, extruding and the like.

[00312] In some embodiments, tablets will include a film surrounding the final compressed tablet. In some embodiments, the film coating can provide a delayed release of the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) from the formulation. In other embodiments, the film coating aids in patient compliance (e.g., Opadry[®] coatings or sugar coating). Film coatings including Opadry[®] typically range from about 1% to about 3% of the tablet weight.

[00313] A capsule may be prepared, for example, by placing the bulk blend of the formulation of the compound described above, inside of a capsule. In some embodiments, the formulations (non-aqueous suspensions and solutions) are placed in a soft gelatin capsule. In other embodiments, the formulations are placed in standard gelatin capsules or non-gelatin capsules such as capsules comprising HPMC. In other embodiments, the formulation is placed in a sprinkle capsule, wherein the capsule is swallowed whole or the capsule is opened and the contents sprinkled on food prior to eating.

[00314] In various embodiments, the particles of the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) and one or more excipients are dry blended and compressed into a mass, such as a tablet, having a hardness sufficient to provide a pharmaceutical composition that substantially disintegrates within less than about 30 minutes, less than about 35 minutes, less than about 40 minutes, less than about 45 minutes, less than about 50 minutes, less than about 55 minutes, or less than about 60 minutes, after oral administration, thereby releasing the formulation into the gastrointestinal fluid.

[00315] In other embodiments, a powder including a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is formulated to include one or more pharmaceutical excipients and flavors. Such a powder is prepared, for example, by mixing the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) and optional pharmaceutical excipients to form a bulk blend composition. Additional embodiments also include a suspending agent and/or a wetting agent. This bulk blend is uniformly subdivided into unit dosage packaging or multi-dosage packaging units.

[00316] In still other embodiments, effervescent powders are also prepared. Effervescent salts have been used to disperse medicines in water for oral administration.

[00317] In some embodiments, the pharmaceutical solid oral dosage forms are formulated to provide a controlled release of the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X). Controlled release refers to the release of the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) from a dosage form in which it is incorporated according to a desired profile over an extended period of time. Controlled release profiles include, for example, sustained release,

prolonged release, pulsatile release, and delayed release profiles. In contrast to immediate release compositions, controlled release compositions allow delivery of an agent to a subject over an extended period of time according to a predetermined profile. Such release rates can provide therapeutically effective levels of agent for an extended period of time and thereby provide a longer period of pharmacologic response while minimizing side effects as compared to conventional rapid release dosage forms. Such longer periods of response provide for many inherent benefits that are not achieved with the corresponding short acting, immediate release preparations.

[00318] In some embodiments, the solid dosage forms described herein are formulated as enteric coated delayed release oral dosage forms, i.e., as an oral dosage form of a pharmaceutical composition as described herein which utilizes an enteric coating to affect release in the small intestine or large intestine. In one aspect, the enteric coated dosage form is a compressed or molded or extruded tablet/mold (coated or uncoated) containing granules, powder, pellets, beads or particles of the active ingredient and/or other composition components, which are themselves coated or uncoated. In one aspect, the enteric coated oral dosage form is in the form of a capsule containing pellets, beads or granules, which include a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) that are coated or uncoated.

[00319] Conventional coating techniques such as spray or pan coating are employed to apply coatings. The coating thickness must be sufficient to ensure that the oral dosage form remains intact until the desired site of topical delivery in the intestinal tract is reached.

[00320] In other embodiments, the formulations described herein are delivered using a pulsatile dosage form. A pulsatile dosage form is capable of providing one or more immediate release pulses at predetermined time points after a controlled lag time or at specific sites. Exemplary pulsatile dosage forms and methods of their manufacture are disclosed in U.S. Pat. Nos.

5,011,692, 5,017,381, 5,229,135, 5,840,329 and 5,837,284. In one embodiment, the pulsatile dosage form includes at least two groups of particles, (i.e. multiparticulate) each containing the formulation described herein. The first group of particles provides a substantially immediate dose of the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) upon ingestion by a mammal. The first group of particles can be either uncoated or include a coating and/or sealant. In one aspect, the second group of particles comprises coated particles. The coating on the second group of particles provides a delay of from about 2 hours to about 7 hours following ingestion before release of the second dose. Suitable coatings for pharmaceutical compositions are described herein or known in the art.

[00321] In some embodiments, pharmaceutical formulations are provided that include particles of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa)

or (X) and at least one dispersing agent or suspending agent for oral administration to a subject. The formulations may be a powder and/or granules for suspension, and upon admixture with water, a substantially uniform suspension is obtained.

[00322] In one aspect, liquid formulation dosage forms for oral administration are in the form of aqueous suspensions selected from the group including, but not limited to, pharmaceutically acceptable aqueous oral dispersions, emulsions, solutions, elixirs, gels, and syrups. See, e.g., Singh *et al.*, Encyclopedia of Pharmaceutical Technology, 2nd Ed., pp. 754-757 (2002). In addition to the particles of the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X), the liquid dosage forms include additives, such as: (a) disintegrating agents; (b) dispersing agents; (c) wetting agents; (d) at least one preservative, (e) viscosity enhancing agents, (f) at least one sweetening agent, and (g) at least one flavoring agent. In some embodiments, the aqueous dispersions can further include a crystalline inhibitor.

[00323] Buccal formulations that include a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) are administered using a variety of formulations known in the art. For example, such formulations include, but are not limited to, U.S. Pat. Nos. 4,229,447, 4,596,795, 4,755,386, and 5,739,136. In addition, the buccal dosage forms described herein can further include a bioerodible (hydrolysable) polymeric carrier that also serves to adhere the dosage form to the buccal mucosa. For buccal or sublingual administration, the compositions may take the form of tablets, lozenges, or gels formulated in a conventional manner.

[00324] In some embodiments, compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) are prepared as transdermal dosage forms. In one embodiment, the transdermal formulations described herein include at least three components: (1) a formulation of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X); (2) a penetration enhancer; and (3) an aqueous adjuvant. In some embodiments the transdermal formulations include additional components such as, but not limited to, gelling agents, creams and ointment bases, and the like. In some embodiments, the transdermal formulation further include a woven or non-woven backing material to enhance absorption and prevent the removal of the transdermal formulation from the skin. In other embodiments, the transdermal formulations described herein can maintain a saturated or supersaturated state to promote diffusion into the skin.

[00325] In one aspect, formulations suitable for transdermal administration of compounds described herein employ transdermal delivery devices and transdermal delivery patches and can be lipophilic emulsions or buffered, aqueous solutions, dissolved and/or dispersed in a polymer or an adhesive. In one aspect, such patches are constructed for continuous, pulsatile, or on

demand delivery of pharmaceutical agents. Still further, transdermal delivery of the compounds described herein can be accomplished by means of iontophoretic patches and the like. In one aspect, transdermal patches provide controlled delivery of the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X). In one aspect, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a rate controlling barrier to deliver the compound to the skin of the host at a controlled and predetermined rate over a prolonged period of time, and means to secure the device to the skin.

[00326] In one aspect, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is formulated into a pharmaceutical composition suitable for intramuscular, subcutaneous, or intravenous injection. In one aspect, formulations suitable for intramuscular, subcutaneous, or intravenous injection include physiologically acceptable sterile aqueous or non-aqueous solutions, dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and non-aqueous carriers, diluents, solvents, or vehicles include water, ethanol, polyols (propyleneglycol, polyethylene-glycol, glycerol, cremophor and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants. In some embodiments, formulations suitable for subcutaneous injection also contain additives such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the growth of microorganisms can be ensured by various antibacterial and antifungal agents, such as parabens, chlorobutanol, phenol, sorbic acid, and the like. In some cases it is desirable to include isotonic agents, such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, such as aluminum monostearate and gelatin.

[00327] For intravenous injections, compounds described herein are formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. For other parenteral injections, appropriate formulations include aqueous or nonaqueous solutions, preferably with physiologically compatible buffers or excipients. Such excipients are known.

[00328] Parenteral injections may involve bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose

containers, with an added preservative. The pharmaceutical composition described herein may be in a form suitable for parenteral injection as a sterile suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. In one aspect, the active ingredient is in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

[00329] In certain embodiments, delivery systems for pharmaceutical compounds may be employed, such as, for example, liposomes and emulsions. In certain embodiments, compositions provided herein can also include an mucoadhesive polymer, selected from among, for example, carboxymethylcellulose, carbomer (acrylic acid polymer), poly(methylmethacrylate), polyacrylamide, polycarbophil, acrylic acid/butyl acrylate copolymer, sodium alginate and dextran.

[00330] In some embodiments, the compounds described herein may be administered topically and can be formulated into a variety of topically administrable compositions, such as solutions, suspensions, lotions, gels, pastes, medicated sticks, balms, creams or ointments. Such pharmaceutical compounds can contain solubilizers, stabilizers, tonicity enhancing agents, buffers and preservatives.

Methods of Dosing and Treatment Regimens

[00331] In one embodiment, the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are used in the preparation of medicaments for the treatment of diseases or conditions in a mammal that would benefit from a reduction of androgen receptor activity. Methods for treating any of the diseases or conditions described herein in a mammal in need of such treatment, involves administration of pharmaceutical compositions that include at least one compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) or a pharmaceutically acceptable salt, active metabolite, prodrug, or pharmaceutically acceptable solvate thereof, in therapeutically effective amounts to said subject.

[00332] In certain embodiments, the compositions containing the compound(s) described herein are administered for prophylactic and/or therapeutic treatments. In certain therapeutic applications, the compositions are administered to a patient already suffering from a disease or condition, in an amount sufficient to cure or at least partially arrest at least one of the symptoms of the disease or condition. Amounts effective for this use depend on the severity and course of the disease or condition, previous therapy, the patient's health status, weight, and response to the drugs, and the judgment of the treating physician. Therapeutically effective amounts are optionally determined by methods including, but not limited to, a dose escalation clinical trial.

[00333] In prophylactic applications, compositions containing the compounds described herein are administered to a patient susceptible to or otherwise at risk of a particular disease, disorder

or condition. Such an amount is defined to be a "prophylactically effective amount or dose." In this use, the precise amounts also depend on the patient's state of health, weight, and the like. When used in a patient, effective amounts for this use will depend on the severity and course of the disease, disorder or condition, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician. In one aspect, prophylactic treatments include administering to a mammal, who previously experienced at least one symptom of the disease being treated and is currently in remission, a pharmaceutical composition comprising a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) in order to prevent a return of the symptoms of the disease or condition.

[00334] In certain embodiments wherein the patient's condition does not improve, upon the doctor's discretion the administration of the compounds are administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disease or condition.

[00335] In certain embodiments wherein a patient's status does improve, the dose of drug being administered may be temporarily reduced or temporarily suspended for a certain length of time (*i.e.*, a "drug holiday"). In specific embodiments, the length of the drug holiday is between 2 days and 1 year, including by way of example only, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, or more than 28 days. The dose reduction during a drug holiday is, by way of example only, by 10%-100%, including by way of example only 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, and 100%.

[00336] Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, in specific embodiments, the dosage or the frequency of administration, or both, is reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. In certain embodiments, however, the patient requires intermittent treatment on a long-term basis upon any recurrence of symptoms.

[00337] The amount of a given agent that corresponds to such an amount varies depending upon factors such as the particular compound, disease condition and its severity, the identity (*e.g.*, weight, sex) of the subject or host in need of treatment, but can nevertheless be determined according to the particular circumstances surrounding the case, including, *e.g.*, the specific agent being administered, the route of administration, the condition being treated, and the subject or host being treated.

[00338] In general, however, doses employed for adult human treatment are typically in the range of 0.01 mg-5000 mg per day. In one aspect, doses employed for adult human treatment are from about 1 mg to about 1000 mg per day. In one embodiment, the desired dose is

conveniently presented in a single dose or in divided doses administered simultaneously (or over a short period of time) or at appropriate intervals, for example as two, three, four or more sub-doses per day.

[00339] In one embodiment, the daily dosages appropriate for the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) described herein are from about 0.01 to about 10 mg/kg per body weight. In specific embodiments, an indicated daily dosage in a large mammal, including, but not limited to, humans, is in the range from about 0.5 mg to about 1000 mg, conveniently administered in divided doses, including, but not limited to, up to four times a day. In one embodiment, the daily dosage is administered in extended release form. In certain embodiments, suitable unit dosage forms for oral administration comprise from about 1 to 500 mg active ingredient. In other embodiments, the daily dosage or the amount of active in the dosage form are lower or higher than the ranges indicated herein, based on a number of variables in regard to an individual treatment regime. In various embodiments, the daily and unit dosages are altered depending on a number of variables including, but not limited to, the activity of the compound used, the disease or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disease or condition being treated, and the judgment of the practitioner.

[00340] Toxicity and therapeutic efficacy of such therapeutic regimens are determined by standard pharmaceutical procedures in cell cultures or experimental animals, including, but not limited to, the determination of the LD₅₀ and the ED₅₀. The dose ratio between the toxic and therapeutic effects is the therapeutic index and it is expressed as the ratio between LD₅₀ and ED₅₀. In certain embodiments, the data obtained from cell culture assays and animal studies are used in formulating the therapeutically effective daily dosage range and/or the therapeutically effective unit dosage amount for use in mammals, including humans. In some embodiments, the daily dosage amount of the compounds described herein lies within a range of circulating concentrations that include the ED₅₀ with minimal toxicity. In certain embodiments, the daily dosage range and/or the unit dosage amount varies within this range depending upon the dosage form employed and the route of administration utilized.

Combination Treatments

[00341] In certain instances, it is appropriate to administer at least one compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) in combination with another therapeutic agent.

[00342] In one embodiment, the therapeutic effectiveness of one of the compounds described herein is enhanced by administration of an adjuvant (*i.e.*, by itself the adjuvant may have minimal therapeutic benefit, but in combination with another therapeutic agent, the overall

therapeutic benefit to the patient is enhanced). Or, in some embodiments, the benefit experienced by a patient is increased by administering one of the compounds described herein with another therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit.

5 [00343] In one specific embodiment, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is co-administered with a second therapeutic agent, wherein the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) and the second therapeutic agent modulate different aspects of the disease, disorder or condition being treated, thereby providing a greater overall benefit than

10 administration of either therapeutic agent alone.

[00344] In any case, regardless of the disease, disorder or condition being treated, the overall benefit experienced by the patient may simply be additive of the two therapeutic agents or the patient may experience a synergistic benefit.

15 [00345] In certain embodiments, different therapeutically-effective dosages of the compounds disclosed herein will be utilized in formulating pharmaceutical composition and/or in treatment regimens when the compounds disclosed herein are administered in combination with one or more additional agent, such as an additional therapeutically effective drug, an adjuvant or the like. Therapeutically-effective dosages of drugs and other agents for use in combination treatment regimens can be determined by means similar to those set forth hereinabove for the actives themselves. Furthermore, the methods of prevention/treatment described herein encompasses the use of metronomic dosing, i.e., providing more frequent, lower doses in order to minimize toxic side effects. In some embodiments, a combination treatment regimen encompasses treatment regimens in which administration of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is initiated prior to, during, or

20 after treatment with a second agent described herein, and continues until any time during treatment with the second agent or after termination of treatment with the second agent. It also includes treatments in which a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) and the second agent being used in combination are administered simultaneously or at different times and/or at decreasing or increasing intervals

25 during the treatment period. Combination treatment further includes periodic treatments that start and stop at various times to assist with the clinical management of the patient.

[00346] It is understood that the dosage regimen to treat, prevent, or ameliorate the condition(s) for which relief is sought, is modified in accordance with a variety of factors. These factors include the disease, disorder or condition from which the subject suffers, as well as the age,

30 weight, sex, diet, and medical condition of the subject. Thus, in some instances, the dosage

regimen actually employed varies and, in some embodiments, deviates from the dosage regimens set forth herein.

[00347] For combination therapies described herein, dosages of the co-administered compounds vary depending on the type of co-drug employed, on the specific drug employed, on the disease or condition being treated and so forth. In additional embodiments, when co-administered with one or more other therapeutic agents, the compound provided herein is administered either simultaneously with the one or more other therapeutic agents, or sequentially.

[00348] In combination therapies, the multiple therapeutic agents (one of which is one of the compounds described herein) are administered in any order or even simultaneously. If administration is simultaneous, the multiple therapeutic agents are, by way of example only, provided in a single, unified form, or in multiple forms (e.g., as a single pill or as two separate pills). In one embodiment, one of the therapeutic agents is given in multiple doses, and in another, two (or more if present) are given as multiple doses. In some embodiments of non-simultaneous administration, the timing between the multiple doses vary from more than zero weeks to less than four weeks. In addition, the combination methods, compositions and formulations are not to be limited to the use of only two agents; the use of multiple therapeutic combinations is also envisioned.

[00349] The compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X), as well as combination therapies that include compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X), are administered before, during or after the occurrence of a disease or condition, and the timing of administering the composition containing a compound varies. Thus, in one embodiment, the compounds described herein are used as a prophylactic and are administered continuously to subjects with a propensity to develop conditions or diseases in order to prevent the occurrence of the disease or condition. In another embodiment, the compounds and compositions are administered to a subject during or as soon as possible after the onset of the symptoms. In specific embodiments, a compound described herein is administered as soon as is practicable after the onset of a disease or condition is detected or suspected, and for a length of time necessary for the treatment of the disease. In some embodiments, the length required for treatment varies, and the treatment length is adjusted to suit the specific needs of each subject. For example, in specific embodiments, a compound described herein or a formulation containing the compound is administered for at least 2 weeks, about 1 month to about 5 years.

Exemplary Agent for use in Combination Therapy

[00350] In some embodiments, methods for treatment of androgen receptor-dependent or androgen receptor-mediated conditions or diseases, such as proliferative disorders, including

cancer, comprises administration to a mammal a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) in combination with at least one additional agent selected, by way of example only, alemtuzumab, arsenic trioxide, asparaginase (pegylated or non-), bevacizumab, cetuximab, platinum-based compounds such as cisplatin, cladribine, daunorubicin/doxorubicin/idarubicin, irinotecan, fludarabine, 5-fluorouracil, gemtuzumab, methotrexate, taxol, temozolomide, thioguanine, or classes of drugs including hormones (an antiestrogen, an antiandrogen, or gonadotropin releasing hormone analogues, interferons such as alpha interferon, nitrogen mustards such as busulfan or melphalan or mechlorethamine, retinoids such as tretinoin, topoisomerase inhibitors such as irinotecan or topotecan, tyrosine kinase inhibitors such as gefinitinib or imatinib, or agents to treat signs or symptoms induced by such therapy including allopurinol, filgrastim, granisetron/ondansetron/palonosetron, dronabinol.

[00351] In one aspect, the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is administered or formulated in combination with one or more anti-cancer agents. In some embodiments, one or more of the anti-cancer agents are proapoptotic agents. Examples of anti-cancer agents include, but are not limited to, any of the following: gossypol, genasense, polyphenol E, Chlorofusin, all trans-retinoic acid (ATRA), bryostatin, tumor necrosis factor-related apoptosis-inducing ligand (TRAIL), 5-aza-2'-deoxycytidine, all trans retinoic acid, doxorubicin, vincristine, etoposide, gemcitabine, imatinib, geldanamycin, 17-N-Allylamino-17-Demethoxygeldanamycin (17-AAG), flavopiridol, LY294002, bortezomib, trastuzumab, BAY 11-7082, PKC412, or PD184352, paclitaxel, and analogs of paclitaxel. Compounds that have the basic taxane skeleton as a common structure feature, have also been shown to have the ability to arrest cells in the G2-M phases due to stabilized microtubules and may be useful for treating cancer in combination with the compounds described herein.

[00352] Further examples of anti-cancer agents for use in combination with the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) include inhibitors of mitogen-activated protein kinase signaling, e.g., U0126, PD98059, PD184352, PD0325901, ARRY-142886, SB239063, SP600125, BAY 43-9006, wortmannin, or LY294002; Syk inhibitors; mTOR inhibitors; and antibodies (e.g., rituxan).

[00353] Other anti-cancer agents for use in combination with the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) include one or more of the following: abiraterone, adriamycin, dactinomycin, bleomycin, vinblastine, cisplatin, acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin;

asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide;
 bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium;
 bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin;
 carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin;
 5 cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; daunorubicin
 hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone;
 doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone
 propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitrucin; enloplatin;
 enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride;
 10 estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate;
 etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate;
 fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine
 hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; iimofosine; interleukin II
 (including recombinant interleukin II, or rIL2), interferon alfa-2a; interferon alfa-2b; interferon
 15 alfa-n1; interferon alfa-n3; interferon beta-1 a; interferon gamma-1 b; iproplatin; irinotecan
 hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride;
 lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine;
 mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril;
 mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedpa; mitindomide;
 20 mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone
 hydrochloride; mycophenolic acid; nocodazoie; nogalamycin; ormaplatin; oxisuran;
 pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman;
 piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium;
 porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin
 25 hydrochloride; pyrazofurin; riboprine; rogletimide; safingol; safingol hydrochloride; semustine;
 simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine;
 spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur;
 teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine;
 thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; triciribine
 30 phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil
 mustard; uredepa; vaporeotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine;
 vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine
 tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin
 hydrochloride.

[00354] Yet other anticancer agents for use in combination with the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) include alkylating agents, antimetabolites, natural products, or hormones, e.g., nitrogen mustards (e.g., mechloroethamine, cyclophosphamide, chlorambucil, etc.), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine, lomusitne, etc.), or triazenes (decarbazine, etc.). Examples of antimetabolites include but are not limited to folic acid analog (e.g., methotrexate), or pyrimidine analogs (e.g., Cytarabine), purine analogs (e.g., mercaptopurine, thioguanine, pentostatin).

[00355] Examples of natural products for use in combination with the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) include but are not limited to vinca alkaloids (e.g., vinblastin, vincristine), epipodophyllotoxins (e.g., etoposide), antibiotics (e.g., daunorubicin, doxorubicin, bleomycin), enzymes (e.g., L-asparaginase), or biological response modifiers (e.g., interferon alpha).

[00356] Examples of alkylating agents for use in combination with the compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) include, but are not limited to, nitrogen mustards (e.g., mechloroethamine, cyclophosphamide, chlorambucil, meiphalan, etc.), ethylenimine and methylmelamines (e.g., hexamethylmelamine, thiotepa), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine, lomusitne, semustine, streptozocin, etc.), or triazenes (decarbazine, etc.).

[00357] In some embodiments, compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are used to treat cancer in combination with: an antiestrogen (e.g., tamoxifen), an antiandrogen (e.g., bicalutamide, flutamide), gonadotropin releasing hormone analog (e.g., leuprolide).

[00358] Other agents that can be used in the methods and compositions described herein for the treatment or prevention of cancer include platinum coordination complexes (e.g., cisplatin, carboplatin), anthracenedione (e.g., mitoxantrone), substituted urea (e.g., hydroxyurea), methyl hydrazine derivative (e.g., procarbazine), adrenocortical suppressant (e.g., mitotane, aminoglutethimide).

[00359] Examples of anti-cancer agents which act by arresting cells in the G2-M phases due to stabilized microtubules include without limitation the following marketed drugs and drugs in development: Erbulozole, Dolastatin 10, Mivobulin isethionate, Vincristine, NSC-639829, Discodermolide, ABT-751, Altorhyrtins (such as Altorhyrtin A and Altorhyrtin C), Spongistatins (such as Spongistatin 1, Spongistatin 2, Spongistatin 3, Spongistatin 4, Spongistatin 5, Spongistatin 6, Spongistatin 7, Spongistatin 8, and Spongistatin 9), Cemadotin hydrochloride, Epothilones (such as Epothilone A, Epothilone B, Epothilone C, Epothilone D,

Epothilone E, Epothilone F, Epothilone B N-oxide, Epothilone A N-oxide, 16-aza-epothilone B, 21-aminoepothilone B, 21-hydroxyepothilone D, 26-fluoroepothilone, Auristatin PE, Soblidotin, Vincristine sulfate, Cryptophycin 52, Vitilevuamide, Tubulysin A, Canadensol, Centaureidin, Oncocidin A1 Fijianolide B, Laulimalide, Narcosine, Nascapine, Hemiasterlin, Vanadocene acetylacetonate, Indanocine Eleutherobins (such as Desmethyleleutherobin, Desacetyeleutherobin, Isoeleutherobin A, and Z-Eleutherobin), Caribaeoside, Caribaeolin, Halichondrin B, Diazonamide A, Taccalonolide A, Diozostatin, (-)-Phenylahistin, Myoseverin B, Resverastatin phosphate sodium.

[00360] In one aspect, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is co-administered with thrombolytic agents (e.g., alteplase, anistreplase, streptokinase, urokinase, or tissue plasminogen activator), heparin, tinzaparin, warfarin, dabigatran (e.g., dabigatran etexilate), factor Xa inhibitors (e.g., fondaparinux, draparinux, rivaroxaban, DX-9065a, otamixaban, LY517717, or YM150), ticlopidine, clopidogrel, CS-747 (prasugrel, LY640315), ximelagatran, or BIBR 1048.

[00361] In some embodiments, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is used in combination with anti-emetic agents to treat nausea or emesis, which may result from the use of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X), anti-cancer agent(s) and/or radiation therapy.

[00362] Anti-emetic agents include, but are not limited to: neurokinin-1 receptor antagonists, 5HT₃ receptor antagonists (such as ondansetron, granisetron, tropisetron, Palonosetron, and zatisetron), GABA_B receptor agonists (such as baclofen), corticosteroids (such as dexamethasone, prednisone, prednisolone, or others), dopamine antagonists (such as, but not limited to, domperidone, droperidol, haloperidol, chlorpromazine, promethazine, prochlorperazine, metoclopramide), antihistamines (H₁ histamine receptor antagonists, such as but not limited to, cyclizine, diphenhydramine, dimenhydrinate, meclizine, promethazine, hydroxyzine), cannabinoids (such as but not limited to, cannabis, marinol, dronabinol), and others (such as, but not limited to, trimethobenzamide; ginger, emetrol, propofol).

[00363] In some embodiments, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is used in combination with an agent useful in the treatment of anemia. Such an anemia treatment agent is, for example, a continuous erythropoiesis receptor activator (such as epoetin- α).

[00364] In some embodiments, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is used in combination with an agent useful in the treatment of neutropenia. Examples of agents useful in the treatment of neutropenia include, but are not

limited to, a hematopoietic growth factor which regulates the production and function of neutrophils such as a human granulocyte colony stimulating factor, (G-CSF). Examples of a G-CSF include filgrastim.

[00365] In some embodiments, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is administered with corticosteroids. Corticosteroids, include, but are not limited to: betamethasone, prednisone, alclometasone, aldosterone, amcinonide, beclometasone, betamethasone, budesonide, ciclesonide, clobetasol, clobetasone, clocortolone, cloprednol, cortisone, cortivazol, deflazacort, deoxycorticosterone, desonide, desoximetasone, desoxycortone, dexamethasone, diflorasone, diflucortolone, difluprednate, flucolorolone, fludrocortisone, fludroxycortide, flumetasone, flunisolide, fluocinolone acetonide, fluocinonide, fluocortin, fluocortolone, fluorometholone, fluperolone, fluprednidene, fluticasone, formocortol, halcinonide, halometasone, hydrocortisone/cortisol, hydrocortisone aceponate, hydrocortisone buteprate, hydrocortisone butyrate, loteprednol, medrysone, meprednisone, methylprednisolone, methylprednisolone aceponate, mometasone furoate, paramethasone, prednicarbate, prednisone/prednisolone, rimexolone, tixocortol, triamcinolone, and ulobetasol.

[00366] In one embodiment, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is administered to a mammal in combination with a non-steroidal anti-inflammatory drug (NSAID). NSAIDs include, but are not limited to: aspirin, salicylic acid, gentisic acid, choline magnesium salicylate, choline salicylate, choline magnesium salicylate, choline salicylate, magnesium salicylate, sodium salicylate, diflunisal, carprofen, fenoprofen, fenoprofen calcium, flurobiprofen, ibuprofen, ketoprofen, nabutone, ketolorac, ketolorac tromethamine, naproxen, oxaprozin, diclofenac, etodolac, indomethacin, sulindac, tolmetin, meclofenamate, meclofenamate sodium, mefenamic acid, piroxicam, meloxicam, COX-2 specific inhibitors (such as, but not limited to, celecoxib, rofecoxib, valdecoxib, parecoxib, etoricoxib, lumiracoxib, CS-502, JTE-522, L-745,337 and NS398).

[00367] In some embodiments, compounds of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) and (X) are coadministered with analgesics.

[00368] In some embodiments, a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is used in combination with radiation therapy (or radiotherapy). Radiation therapy is the treatment of cancer and other diseases with ionizing radiation. Radiation therapy can be used to treat localized solid tumors, such as cancers of the skin, tongue, larynx, brain, breast, prostate, colon, uterus and/or cervix. It can also be used to treat leukemia and lymphoma (cancers of the blood-forming cells and lymphatic system, respectively).

[00369] A technique for delivering radiation to cancer cells is to place radioactive implants directly in a tumor or body cavity. This is called internal radiotherapy (brachytherapy, interstitial irradiation, and intracavitary irradiation are types of internal radiotherapy.) Using internal radiotherapy, the radiation dose is concentrated in a small area, and the patient stays in the hospital for a few days. Internal radiotherapy is frequently used for cancers of the tongue, uterus, prostate, colon, and cervix.

[00370] The term “radiotherapy” or “ionizing radiation” include all forms of radiation, including but not limited to α , β , and γ radiation and ultraviolet light.

Kits/Articles of Manufacture

[00371] For use in the therapeutic applications described herein, kits and articles of manufacture are also described herein. Such kits can comprise a carrier, package, or container that is compartmentalized to receive one or more containers such as vials, tubes, and the like, each of the container(s) comprising one of the separate elements to be used in a method described herein. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers are formed from any acceptable material including, e.g., glass or plastic.

[00372] For example, the container(s) can comprise one or more compounds described herein, optionally in a composition or in combination with another agent as disclosed herein. The container(s) optionally have a sterile access port (for example the container can be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). Such kits optionally comprising a compound with an identifying description or label or instructions relating to its use in the methods described herein.

[00373] A kit will typically comprise one or more additional containers, each with one or more of various materials (such as reagents, optionally in concentrated form, and/or devices) desirable from a commercial and user standpoint for use of a compound described herein. Non-limiting examples of such materials include, but not limited to, buffers, diluents, filters, needles, syringes; carrier, package, container, vial and/or tube labels listing contents and/or instructions for use, and package inserts with instructions for use. A set of instructions will also typically be included.

[00374] A label can be on or associated with the container. A label can be on a container when letters, numbers or other characters forming the label are attached, molded or etched into the container itself; a label can be associated with a container when it is present within a receptacle or carrier that also holds the container, e.g., as a package insert. A label can be used to indicate that the contents are to be used for a specific therapeutic application. The label can also indicate directions for use of the contents, such as in the methods described herein.

EXAMPLES

[00375] These examples are provided for illustrative purposes only and not to limit the scope of the claims provided herein.


Synthesis of Compounds

5

INTERMEDIATE 1


5-Amino-3-(trifluoromethyl)picolinonitrile

Step 1: 5-Iodo-3-(trifluoromethyl)pyridin-2-ol




[00376] A mixture of 5-iodo-3-(trifluoromethyl)pyridin-2-ol (21.4g, 0.13 mmol) and NIS (29.5 g, 0.13 mmol) in acetonitrile (300 mL) was heated to 80°C for 6h. The reaction mixture was cooled to room temperature and a saturated solution of sodium bicarbonate was added. The solid that precipitated was filtered off and acetonitrile was removed in vacuo. The aqueous layer was extracted with EtOAc (4x), the organics were combined, washed with an aqueous saturated solution of sodium thiosulfate, dried over sodium sulfate, and concentrated to dryness to afford 24 g of 5-iodo-3-(trifluoromethyl)pyridin-2-ol as a yellow solid. ¹H NMR (300 MHz, DMSO-d₆) δ 12.57 (s, 1H), 8.01 (s, 1H), 7.99 (s, 1H).

Step 2: 2-Chloro-5-iodo-3-(trifluoromethyl)pyridine



[00377] A mixture of 5-iodo-3-(trifluoromethyl)pyridin-2-ol (53 g, 0.18 mmol) and POCl₃ (67 mL, 0.73 mmol) in DMF (50 mL) was heated to 110°C for 4h. The reaction mixture was cooled to room temperature and then slowly poured into an ice/water bath. The brown solid was filtered and washed with water. It was then dissolved in DCM, washed with an aqueous saturated solution of sodium thiosulfate (2x), dried over sodium sulfate, and concentrated to dryness to afford 47 g of 2-chloro-5-iodo-3-(trifluoromethyl)pyridine as a pale yellow solid. ¹H NMR (300 MHz, DMSO-d₆) δ 8.79 (d, 1H), 8.29 (d, 1H).

Step 3: 6-Chloro-N-(4-methoxybenzyl)-5-(trifluoromethyl)pyridin-3-amine




[00378] An oil bath was preheated to 130°C. A mixture of 2-chloro-5-iodo-3-(trifluoromethyl)pyridine (47 g, 152.8 mmol), (4-methoxyphenyl)methanamine (23.8 mL, 183.4

mmol), Xantphos (2.6 g, 4.58 mmol) and NaOtBu (22 g, 229.2 mmol) in toluene (500 mL) was stirred at room temperature while bubbling nitrogen for 5 min. Pd₂(dba)₃ (2.8 g, 3.05 mmol) was then added and the reaction mixture was refluxed for 2h. The mixture was cooled to room temperature and it was filtered through a pad of celite. The pad of celite was washed with toluene and the filtrate was concentrated. The residue obtained was dissolved in EtOAc, the organic layer was washed with water (4x), and it was then evaporated to dryness. The dark oil obtained was dissolved in DCM and hexanes was added with swirling until a brown solid precipitated. The brown solid was filtered, washed with hexanes, and dried to afford 38.8g (80%) of 6-chloro-N-(4-methoxybenzyl)-5-(trifluoromethyl)pyridin-3-amine as a brown solid.

¹H NMR (300 MHz, DMSO-d₆) δ 7.96 (d, 1H), 7.38 (d, 1H), 7.29 (d, 2H), 7.07 (t, 1H), 6.92 (d, 2H), 4.29 (d, 2H), 3.73 (s, 3H). The filtrate was purified by column chromatography on Silica Gel eluting with 0 to 50% EtOAc/Hexanes to afford an additional 4 g of 6-chloro-N-(4-methoxybenzyl)-5-(trifluoromethyl)pyridin-3-amine as an orange solid (Total yield: 42.8 g).

Step 4: 5-((4-Methoxybenzyl)amino)-3-(trifluoromethyl)picolinonitrile




[00379] An oil bath was preheated to 180°C. A mixture of 6-chloro-N-(4-methoxybenzyl)-5-(trifluoromethyl)pyridin-3-amine (19.45 g, 61.55 mmol), Zn(CN)₂ (8.7 g, 73.86 mmol), and dppf (6.8 g, 12.31 mmol) in DMF (250 mL) was stirred at room temperature while bubbling nitrogen for 5 min. Pd₂(dba)₃ (2.8 g, 3.07 mmol, 0.05 eq.) was then added and the reaction mixture was placed inside the pre-heated bath. The bath temperature dropped to 160°C. The mixture started refluxing when the bath temperature reached 170°C (takes 20 min). Refluxed for 10 min. The reaction mixture was cooled to room temperature and it was filtered through a pad of celite. The pad of celite was washed with EtOAc. The solvent was removed using a rotovap hooked up to a high vac. pump. The black residue was partitioned between EtOAc and water.

An orange solid (some dppf by-product) precipitated and was filtered and washed with EtOAc. The organic layer was washed with brine (2x), dried over sodium sulfate, and concentrated to dryness. The black residue was dissolved in DCM and loaded onto a silica gel plug. The plug was washed with DCM (to get rid of some of the fast eluting impurities) followed by 1:1 EtOAc/hexanes (to move the desired). The fractions containing the desired were combined, evaporated to dryness and the residue obtained was purified by column chromatography on silica gel eluting with 0 to 50% EtOAc/hexanes to afford 15 g of 5-((4-methoxybenzyl)amino)-

3-(trifluoromethyl)picolinonitrile as reddish thick oily residue. ¹H NMR (300 MHz, DMSO-d₆) δ 8.28 (s, 1H), 8.01 (t, 1H), 7.31 (m, 3H), 6.92 (d, 2H), 4.41 (d, 2H), 3.73 (s, 3H).

Step 5: 5-Amino-3-(trifluoromethyl)picolinonitrile




5 [00380] 5-((4-Methoxybenzyl)amino)-3-(trifluoromethyl)picolinonitrile (15 g, 49.02 mmol) was taken up in DCM (30 mL) and TFA (50 mL) was added. The resulting reaction mixture was stirred at room temperature for 3h (until done by LC-MS). The solvent and TFA were removed in vacuo and MeOH was added to the residue. The beige solid that was not soluble in MeOH was filtered and washed with MeOH. The filtrate was evaporated to dryness and the residue
10 was partitioned between EtOAc and a saturated solution of sodium bicarbonate. The organic layer was washed two more times with a saturated aqueous sodium bicarbonate, dried over sodium sulfate, and evaporated to dryness. The residue was dissolved in DCM and hexanes with swirling until a yellow solid precipitated. This solid was filtered and washed with hexanes to afford 7.2 g of 5-amino-3-(trifluoromethyl)picolinonitrile as a pale yellow solid. ¹H NMR
15 (300 MHz, DMSO-d₆) δ 8.17 (d, 1H), 7.28 (d, 1H), 6.99 (bs, 2H).

INTERMEDIATE 2


5-Amino-3-methylpicolinonitrile

Step 1: 3-Methyl-5-nitropicolinonitrile



20 [00381] A mixture of tetrabutylammonium nitrate (17 g, 55.1 mmol) and trifluoroacetic anhydride (7.6 mL, 55.1 mmol) in DCM (50 mL) was slowly added to a cooled (0°C) solution of 3-methylpicolinonitrile (5 g, 42.4 mmol) in DCM (100 mL) and the resulting mixture was stirred for 3 days (temperature allowed to warm to room temperature). A saturated solution of
25 sodium bicarbonate (80 mL) was added and the mixture was stirred at room temperature for 1h. The two layers were separated and the aqueous was extracted with DCM (2x). The organics were combined, dried over sodium sulfate, and concentrated to dryness. The residue was purified by column chromatography on silica gel eluting with 0 to 50% EtOAc/hexanes to afford 2.9 g of 3-methyl-5-nitropicolinonitrile as a pale yellow solid. ¹H NMR (300 MHz,
30 DMSO-d₆) δ 9.32 (s, 1H), 8.83 (s, 1H), 2.64 (s, 3H).

Step 2: 5-Amino-3-methylpicolinonitrile




[00382] A mixture of 3-methyl-5-nitropicolinonitrile (2.9 g, 17.8 mmol) and Pd/C (cat.) in MeOH (60 mL) was hydrogenated overnight with a balloon of hydrogen. Pd/C was removed through a pad of celite and the filtrate was evaporated to dryness to afford 2.2 g of 5-amino-3-methylpicolinonitrile as a greenish solid. ¹H NMR (300 MHz, DMSO-d₆) δ 7.83 (s, 1H), 6.79 (s, 1H), 6.32 (s, 2H), 2.29 (s, 3H).

INTERMEDIATE 3


3-(Trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-amine

Step 1: 6-Chloro-3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazine




[00383] A mixture of 3-chloro-6-hydrazinylpyridazine (3.3 g, 22.8 mmol) in TFA (60 mL) was heated to 85°C for 18h. The reaction mixture was cooled to room temperature and TFA was removed in vacuo. The residue was partitioned between DCM and saturated aqueous sodium bicarbonate and the organic layer was washed with a saturated solution of sodium bicarbonate (3x), dried over sodium sulfate, and evaporated to dryness to afford 2.8 g of 6-chloro-3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazine as a beige solid. ¹H NMR (300 MHz, DMSO-d₆) δ 8.72 (d, 1H), 7.78 (d, 1H).

Step 2: 3-(Trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-amine




[00384] A mixture of 6-chloro-3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazine (1.14 g, 5.1 mmol) and ammonium hydroxide (10 mL) in THF (10 mL) was heated to 60°C for 18h. The reaction mixture was cooled to room temperature and water and EtOAc were added. The aqueous layer was extracted with EtOAc (4x), the organics were combined, dried over sodium sulfate, and evaporated to dryness to afford 1 g of 3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-amine as beige solid. ¹H NMR (300 MHz, DMSO-d₆) δ 8.12 (d, 1H), 7.23 (s, 2H), 6.97 (d, 1H).


INTERMEDIATE 4

6-Amino-2-(trifluoromethyl)nicotinonitrile

[00385] A mixture of 5-bromo-6-(trifluoromethyl)pyridin-2-amine (3.0 g, 12.3 mmol), $\text{Zn}(\text{CN})_2$ (0.81 g, 6.9 mmol), $\text{Pd}_2(\text{dba})_3$ (0.57 g, 0.6 mmol), and Xantphos (0.72 g, 1.2 mmol) in DMA (12 mL) was placed in a sealed tube. The mixture was degassed with argon and stirred at 160°C for 20h. The mixture was poured into water and extracted with EtOAc (2x). The combined organic layers were dried over magnesium sulfate and concentrated to dryness. The residue was purified by column chromatography on silica gel eluting with 3:1 EtOAc/hexanes to afford 1.9 g of 6-amino-2-(trifluoromethyl)nicotinonitrile as a solid. ^1H NMR (400 MHz, CDCl_3) δ 7.74 (d, 1H), 6.67 (d, 1H).

INTERMEDIATE 5**3-(Trifluoromethyl)-[2,3'-bipyridin]-5-amine****Step1: N-(4-Methoxybenzyl)-3-(trifluoromethyl)-[2,3'-bipyridin]-5-amine**

[00386] To a solution of 6-chloro-N-(4-methoxybenzyl)-5-(trifluoromethyl)pyridin-3-amine (Intermediate 1, Step 3, 1.2 g, 3.8 mmol) and 3-(diethylboryl)pyridine (0.613 g, 4.17 mmol) in 1,4-dioxane (100 mL) was added K_3PO_4 (2.52 g, 9.48 mmol). The mixture was degassed then flushed with nitrogen before $\text{Pd}_2(\text{dba})_3$ (173 mg, 0.02 mmol) and Xantphos (0.218 mg, 0.04 mmol) were added. The mixture was heated at 105°C overnight. The reaction mixture was cooled to room temperature and filtered. The filtrate was concentrated in vacuo and the residue was purified by column chromatography on silica gel using EtOAc/PE = 1/ to afford 1 g of N-(4-methoxybenzyl)-3-(trifluoromethyl)-[2,3'-bipyridin]-5-amine.


Step 2: 3-(Trifluoromethyl)-[2,3'-bipyridin]-5-amine

[00387] The title compound was synthesized as described for Intermediate 1, Step 5 using N-(4-methoxybenzyl)-3-(trifluoromethyl)-[2,3'-bipyridin]-5-amine as the starting material. ^1H NMR

(300MHz, CDCl₃) δ 8.71 (dd, 1H), 8.64 (dd, 1H), 8.30 (dd, 1H), 7.78 (m, 1H), 7.35 (m, 1H), 7.32 (d, 1H), 4.05 (s, 2H).

INTERMEDIATE 6 (Isothiocyanate method A)


5-Isothiocyanto-3-methylpicolinonitrile



[00388] Thiophosgene (0.24 mL, 3.1 mmol) was added to a stirred biphasic solution of 5-amino-3-methylpicolinonitrile (Intermediate 2, 275 mg, 2.07 mmol) in CHCl₃ (6 mL)/water (10 mL) and the resulting orange mixture was stirred at room temperature overnight. The two layers were separated and the aqueous layer was extracted with DCM (3x). The organics were combined, washed with a saturated solution of sodium bicarbonate (2x), dried over sodium sulfate, and evaporated to dryness to afford 300 mg of 5-isothiocyanto-3-methylpicolinonitrile as an orange solid. ¹H NMR (300 MHz, DMSO-d₆) δ 8.66 (s, 1H), 8.06 (s, 1H), 2.5 (s, 3H).

INTERMEDIATE 7


6-Isothiocyanto-2-(trifluoromethyl)nicotinonitrile



[00389] A mixture of 6-amino-2-(trifluoromethyl)nicotinonitrile (648 mg, 3.5 mmol) and thiophosgene (0.4 mL, 5.2 mmol) in DCE (8 mL) was heated to 80°C overnight. The mixture was cooled to room temperature and diluted with DCM/water. The two layers were separated and the aqueous layer was extracted with DCM (3x). The organics were combined, washed with a saturated solution of sodium bicarbonate (2x), dried over sodium sulfate, and evaporated to dryness to afford 521 mg of 6-isothiocyanto-2-(trifluoromethyl)nicotinonitrile (70% pure) as a brown oil. ¹H NMR (300 MHz, DMSO-d₆) δ 8.73 (d, 1H), 7.93 (d, 1H).

INTERMEDIATE 8

6-Isothiocyanto-3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazine




[00390] Thiophosgene (1 mL, 12.7 mmol) was added to a solution of 3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-amine (860 mg, 4.23 mmol) in pyridine (10 mL) and THF (15

mL) and the resulting dark mixture was stirred at room temperature for 2h. The dark mixture was partitioned between EtOAc and 1M HCl. The organic layer was washed with 1M HCl (2x), dried over sodium sulfate, and evaporated to dryness to afford 545 mg (60% pure) of 6-isothiocyanato-3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazine as a dark purple solid.

5 LCMS $[M + 1]^+$ 278.5 (OMe adduct).

INTERMEDIATE 9 (Isothiocyanate method B)

4-Isothiocyanatoisoquinoline-1-carbonitrile







10 **[00391]** A mixture of 4-aminoisoquinoline-1-carbonitrile (500 mg, 2.95 mmol) and thiophosgene (0.34 mL, 4.44 mmol) in DMA (5 mL) was stirred at room temperature overnight. The mixture was diluted with EtOAc/water. The two layers were separated and the aqueous layer was extracted with EtOAc (3x). The organics were combined, washed with a saturated solution of sodium bicarbonate (2x), dried over sodium sulfate, and evaporated to dryness to afford 553 mg of 4-isothiocyanatoisoquinoline-1-carbonitrile as an orange oil. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.88 (s, 1H), 8.32 (d, 1H), 8.23 (d, 1H), 8.12 (td, 1H), 8.05 (td, 1H).

15

[00392] ISOTHIOCYANATE INTERMEDIATES 10-16 were synthesized from the appropriate amines (amines were either commercially available, synthesized in the Intermediates section, or synthesized from published procedures) according to the conditions indicated (see footnotes).

20

Intermediate #	Name	Structure
10^a	5-Isothiocyanato-3-(trifluoromethyl)picolinonitrile	
11^a	3-Chloro-5-isothiocyanatopicolinonitrile	
12^a	5-Isothiocyanato-3-methoxypicolinonitrile	


Intermediate #	Name	Structure
13 ^a	5-Isothiocyanatoquinoline-8-carbonitrile	
14 ^a	6-Isothiocyanatoimidazo[1,2-a]pyridine	
15 ^a	5-Isothiocyanato-3-(trifluoromethyl)-2,3'-bipyridine	
16 ^b	4-Isothiocyanatopyrazolo[1,5-a]pyridine-7-carbonitrile	

^a Isothiocyanate method A, ^b Isothiocyanate method B

INTERMEDIATE 17


4-((2-(Pyridin-4-yl)ethyl)sulfonyl)aniline

5 Step 1: 4-(2-((4-Nitrophenyl)thio)ethyl)pyridine




[00393] A mixture of 2-(pyridin-4-yl)ethanethiol hydrochloride (3.84 g, 21.8 mmol, 4-fluoro-1-nitrobenzene (6.12 g, 43.3 mmol), and K₂CO₃ (9.0 g, 65.1 mmol) in acetonitrile (150 mL) was refluxed overnight. The reaction mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by column chromatography on silica gel (EtOAc/PE = 1/1) to afford 2.0 g of 4-(2-((4-nitrophenyl)thio)ethyl)pyridine as a light yellow solid. ¹H NMR (300 MHz, CDCl₃) δ 8.55 (d, 2H), 8.14 (d, 2H), 7.34 (d, 2H), 7.15 (d, 2H), 3.30 (t, 2H), 3.01 (t, 2H).

Step 2: 4-(2-((4-Nitrophenyl)sulfonyl)ethyl)pyridine



[00394] H₂O₂ (30%, 5 mL) was added dropwise to a solution of 4-(2-((4-nitrophenyl)thio)ethyl)pyridine (338 mg, 1.30 mmol) in acetic acid (20 mL) heated at 60°C and the resulting mixture was stirred at 60°C for 30min. The mixture was poured into a sat. Na₂CO₃ (100 mL) and extracted with DCM. The organic layer was dried over sodium sulfate and concentrated to dryness. The residue was purified by column chromatography on silica gel (EA/PE = 1/3) to afford 270 mg of 4-(2-((4-nitrophenyl)sulfonyl)ethyl)pyridine as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 8.51 (d, 2H), 8.42 (dd, 2H), 7.13 (dd, 2H), 7.08 (d, 2H), 3.45-3.40 (m, 2H), 3.14-3.08 (m, 2H).

Step 3: 4-((2-(Pyridin-4-yl)ethyl)sulfonyl)aniline




[00395] A mixture of 4-(2-((4-nitrophenyl)sulfonyl)ethyl)pyridine (210 mg, 0.76 mmol) and 10%Pd/C (100 mg) in EtOH (100 mL) was stirred at room temperature for 3 h under H₂ atmosphere (balloon). The reaction mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by column chromatography on silica gel (MeOH/DCM = 1/50) to afford 100 mg of 4-((2-(pyridin-4-yl)ethyl)sulfonyl)aniline as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 8.48 (m, 2H), 7.67 (m, 2H), 7.05 (m, 2H), 6.70 (m, 2H), 4.23 (bs, 2H), 3.34-3.28 (m, 2H), 3.07-3.01 (m, 2H).

INTERMEDIATE 18


4-((Methyl(pyridin-4-ylmethyl)amino)methyl)aniline

Step 1: N-(4-Nitrobenzyl)-1-(pyridin-4-yl)methanamine




[00396] A mixture of 4-nitrobenzaldehyde (3.0 g, 19.8 mmol), 4-pyridylmethylamine (2.36 g, 21.8 mmol), and NaBH₃CN (1.75 g, 27.7 mmol) in 1,2-dichloroethane (40.0 mL) was stirred at room temperature for 36h and then concentrated to dryness. The residue was partitioned between water and EtOAc. The aqueous layer was extracted with EtOAc (3x). The combined organics were washed with brine (2x), dried over sodium sulfate, and concentrated. The residue was purified by column chromatography on silica gel (MeOH/DCM = 1/400) to afford 1.3 g of N-(4-nitrobenzyl)-1-(pyridin-4-yl)methanamine as brown oil.

Step 2: N-Methyl-N-(4-nitrobenzyl)-1-(pyridin-4-yl)methanamine



[00397] The title compound was synthesized as described in the previous step using N-(4-nitrobenzyl)-1-(pyridin-4-yl)methanamine and (HCHO)_n as the starting materials.


Step 3: 4-((Methyl(pyridin-4-ylmethyl)amino)methyl)aniline



[00398] The title compound was synthesized as described for Intermediate 17, Step 3 using N-methyl-N-(4-nitrobenzyl)-1-(pyridin-4-yl)methanamine as the starting material.

INTERMEDIATE 19 (Aminonitrile synthesis method A)


1-((3-Fluoro-4-hydroxyphenyl)amino)cyclobutanecarbonitrile



[00399] Sodium cyanide (2.5 g, 50.8 mmol) was added to a mixture of 4-amino-2-fluorophenol (4.3 g, 33.9 mmol) and cyclobutanone (3.8 mL, 50.8 mmol) in acetic acid (99%, 30 mL) hooked up to a NaOH scrubber and the resulting mixture was stirred at room temperature for 6h. The mixture was poured in water and the aqueous layer was extracted with EtOAc (3x). The organics were combined, washed with a saturated solution of sodium bicarbonate (4x), dried over sodium sulfate, and evaporated to dryness. The residue was purified by column chromatography on silica gel eluting with 0 to 50% EtOAc/hexanes to afford 4.8 g of 1-((3-fluoro-4-hydroxyphenyl)amino)cyclobutanecarbonitrile as a beige solid. ¹H NMR (300 MHz, DMSO-d₆) δ 8.98 (s, 1H), 6.80 (t, 1H), 6.36 (dd, 1H), 6.28 (m, 2H), 2.71-2.62 (m, 2H), 2.35-2.25 (m, 2H), 2.12-2.00 (m, 2H).

INTERMEDIATE 20 (Aminonitrile synthesis method B)

Ethyl 4-((1-cyanocyclobutyl)amino)-2-fluorobenzoate




[00400] Sodium cyanide (1.3 g, 26.6 mmol) was added to a mixture of ethyl 4-amino-2-fluorobenzoate (3.25 g, 17.8 mmol) and cyclobutanone (2 mL, 26.6 mmol) in acetic acid (99%, 40 mL) hooked up to a NaOH scrubber and the resulting mixture was heated to 80°C overnight.

The mixture was cooled to room temperature and poured in water. The pink precipitate was filtered, washed with water, and redissolved in DCM (100 mL). The organic layer was washed with a saturated solution of sodium bicarbonate (4x), dried over sodium sulfate, and evaporated to dryness to afford 3.9 g of ethyl 4-((1-cyanocyclobutyl)amino)-2-fluorobenzoate as an orange solid. ¹H NMR (300 MHz, CDCl₃) δ 7.87 (t, 1H), 6.43 (dd, 1H), 6.34 (dd, 1H), 4.55 (s, 1H), 4.36 (q, 2H), 2.91-2.82 (m, 2H), 2.46-2.30 (m, 2H), 2.28-2.13 (m, 2H), 1.39 (t, 3H).

INTERMEDIATE 21 ((Aminonitrile synthesis method C)


4-((1-Cyanocyclobutyl)amino)-2-fluoro-N-methylbenzamide




[00401] Sodium cyanide (5.8 g, 119 mmol) was added to a mixture of 4-amino-2-fluoro-N-methylbenzamide (5 g, 29.8 mmol) and cyclobutanone (4.4 mL, 59.5 mmol) in AcOH (90%, 25 mL) and EtOH (25 mL) hooked up to a NaOH scrubber and the resulting mixture was heated to 80°C overnight. The mixture was cooled to room temperature and poured in water. The yellow precipitate was filtered, washed with water, and dried to afford 6.5 g of 4-((1-cyanocyclobutyl)amino)-2-fluoro-N-methylbenzamide as a pale yellow solid. ¹H NMR (300 MHz, DMSO-d₆) δ 7.81 (t, 1H), 7.56 (t, 1H), 7.37 (s, 1H), 6.46 (dd, 1H), 6.30 (dd, 1H), 2.80-2.71 (m, 5H), 2.41-2.31 (m, 2H), 2.13-2.00 (m, 2H)

INTERMEDIATE 22 (Aminonitrile synthesis method D)


1-((2,6-Difluorophenyl)amino)cyclobutanecarbonitrile



[00402] A mixture of 2,6-difluoroaniline (0.5 g, 3.9 mmol), TMS-CN (0.72 mL, 5.8 mmol), and cyclobutanone (0.43 mL, 5.8 mmol) was heated to 90°C overnight. The mixture was poured in water and the aqueous layer was extracted with EtOAc (3x). The organics were combined, washed with brine. (2x), dried over sodium sulfate, and evaporated to dryness. The residue was purified by column chromatography on silica gel eluting with 0 to 20% EtOAc/hexanes to afford 0.7 g of 1-((2,6-difluorophenyl)amino)cyclobutanecarbonitrile as a colorless oil. ¹H NMR (300 MHz, DMSO-d₆) δ 7.07-7.01 (m, 2H), 6.91-6.81 (m, 1H), 6.15 (s, 1H), 2.65-2.56 (m, 2H), 2.52-2.42 (m, 2H), 2.01-1.91 (m, 2H).


INTERMEDIATE 23 (Aminonitrile synthesis method E)**4-((2-Cyanopropan-2-yl)amino)-2-fluoro-N-methylbenzamide**

- 5 [00403] A mixture of 4-amino-2-fluoro-N-methylbenzamide (2 g, 11.9 mmol), TMSCN (2.2 mL, 17.9 mmol), TMSOTf (0.1 mL, 0.6 mmol), and acetone (10 mL, 140 mmol) in DCM (20 mL) was stirred at room temperature overnight. The white solid was filtered, washed with a small amount of DCM, and dried to afford 1.12 g of 4-((2-cyanopropan-2-yl)amino)-2-fluoro-N-methylbenzamide as a white solid. ¹H NMR (300 MHz, DMSO-d₆) δ 7.79 (t, 1H), 7.56 (t, 10 1H), 6.88 (s, 1H), 6.67 (dd, 1H), 6.54 (dd, 1H), 2.75 (d, 3H), 1.67 (s, 6H).

INTERMEDIATE 24**1-(4-(Pyridin-4-yloxy)phenylamino)cyclobutanecarbonitrile**

- 15 [00404] Sodium cyanide (395 mg, 8.06 mmol) was added to a mixture of 4-(pyridin-4-yloxy)aniline (500 mg, 2.69 mmol) and cyclobutanone (376 mg, 5.37 mmol) in AcOH (99%, 6 mL) in a microwave vial. The reaction mixture was irradiated with microwaves (Biotage) at 120°C for 30 min. The slightly yellowish solution was concentrated to dryness. The residue was diluted with water and extracted with EtOAc (2x). The combined organic layers were dried 20 over MgSO₄ and concentrated to dryness to afford 463 mg (65%) of 1-(4-(pyridin-4-yloxy)phenylamino)-cyclobutanecarbonitrile as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 8.46-8.38 (m, 2H), 7.02-6.94 (m, 2H), 6.79 (m, 2H), 6.72-6.64 (m, 2H), 2.86-2.74 (m, 2H), 2.46-2.34 (m, 2H), 2.22 (m, 2H)

25










INTERMEDIATE 25**2-((4-Hydroxyphenyl)amino)-2-methylpropanenitrile**

- [00405] A mixture of 4-aminophenol (1 g, 9.17 mmol) and magnesium sulfate (4 g) in 2-hydroxy-2-methylpropanenitrile (10 mL) was heated to 80°C for 2h. Ice/water was added to a










cooled mixture and it was stirred for 30 min. The white solid that precipitated was filtered, washed with water, and dried to afford 1.2g (74%) of 2-((4-hydroxyphenyl)amino)-2-methylpropanenitrile as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 8.89 (s, 1H), 6.78 (d, 2H), 6.65 (d, 2H), 5.21 (s, 1H), 1.52 (s, 6H).










- 5 [00406] **INTERMEDIATES 26-107** were synthesized from the appropriate amines (amines were either commercially available, synthesized in the Intermediates section, or synthesized from published procedures) and ketones, according to the conditions indicated (see footnotes).

Intermediate #	Name	Structure
26 ^a	1-((2-Fluoro-4-hydroxyphenyl)amino)cyclobutanecarbonitrile	
27 ^b	1-((4-Fluoro-3-methoxyphenyl)amino)cyclobutanecarbonitrile	
28 ^b	1-([1,1'-Biphenyl]-2-ylamino)cyclobutanecarbonitrile	
29 ^b	1-([1,1'-Biphenyl]-3-ylamino)cyclobutanecarbonitrile	
30 ^b	1-([1,1'-Biphenyl]-4-ylamino)cyclobutanecarbonitrile	
31 ^b	1-(Naphthalen-1-ylamino)cyclobutanecarbonitrile	
32 ^b	1-(Naphthalen-2-ylamino)cyclobutanecarbonitrile	










Intermediate #	Name	Structure
33 ^b	Ethyl 5-((1-cyanocyclobutyl)amino)-2-fluorobenzoate	
34 ^b	2-((1-Cyanocyclobutyl)amino)benzonitrile	
35 ^b	3-((1-Cyanocyclobutyl)amino)benzonitrile	
36 ^b	4-((1-Cyanocyclobutyl)amino)benzonitrile	
37 ^b	1-(Pyridin-3-ylamino)cyclobutanecarbonitrile	
38 ^c	1-(o-Tolylamino)cyclobutanecarbonitrile	
39 ^c	1-(m-Tolylamino)cyclobutanecarbonitrile	
40 ^c	1-(p-Tolylamino)cyclobutanecarbonitrile	
41 ^b	1-((4-Fluoro-2-methoxyphenyl)amino)cyclobutanecarbonitrile	

Intermediate #	Name	Structure
42 ^b	1-((4-Fluoro-2-hydroxyphenyl)amino)cyclobutanecarbonitrile	
43 ^d	1-((2-Phenoxyphenyl)amino)cyclobutanecarbonitrile	
44 ^b	1-((3-Phenoxyphenyl)amino)cyclobutanecarbonitrile	
45 ^b	1-((4-Phenoxyphenyl)amino)cyclobutanecarbonitrile	
46 ^d	1-((4-Methylbenzyl)amino)cyclobutanecarbonitrile	
47 ^d	1-((4-Methylphenethyl)amino)cyclobutanecarbonitril e	
48 ^d	1-((4-(Pyridin-3-yloxy)phenyl)amino)cyclobutanecarbonitrile	
49 ^b	1-((4-(Pyridin-2-yloxy)phenyl)amino)cyclobutanecarbonitrile	
50 ^b	1-((4-(Pyrimidin-5-yloxy)phenyl)amino)cyclobutanecarbonitrile	

Intermediate #	Name	Structure
51 ^b	1-((3-(Trifluoromethoxy)phenyl)amino)cyclobutanecarbonitrile	
52 ^b	1-((4-(Trifluoromethoxy)phenyl)amino)cyclobutanecarbonitrile	
53 ^b	1-((2-Fluorophenyl)amino)cyclobutanecarbonitrile	
54 ^b	1-((3-Fluorophenyl)amino)cyclobutanecarbonitrile	
55 ^b	1-((4-Fluorophenyl)amino)cyclobutanecarbonitrile	
56 ^c	1-((4-(Trifluoromethyl)phenyl)amino)cyclobutanecarbonitrile	
57 ^b	1-(Phenylamino)cyclobutanecarbonitrile	
58 ^b	1-((3-Fluoro-4-methylphenyl)amino)cyclobutanecarbonitrile	
59 ^b	1-((2-Fluoro-4-methylphenyl)amino)cyclobutanecarbonitrile	




Intermediate #	Name	Structure
60^b	1-(Cyclohexylamino)cyclobutanecarbonitrile	
61^b	1-((4-(Methylsulfonyl)phenyl)amino)cyclobutanecarbonitrile	
62^b	4-((1-Cyanocyclobutyl)amino)benzenesulfonamide	
63^b	1-(Isoquinolin-6-ylamino)cyclobutanecarbonitrile	
64^b	1-(Isoquinolin-7-ylamino)cyclobutanecarbonitrile	
65^b	4-((1-Cyanocyclobutyl)amino)-2-fluorobenzonitrile	
66^b	4-((1-Cyanocyclobutyl)amino)-3-fluorobenzonitrile	
67^c	1-((2-(Hydroxymethyl)phenyl)amino)cyclobutanecarbonitrile	
68^c	1-((3-(Hydroxymethyl)phenyl)amino)cyclobutanecarbonitrile	

Intermediate #	Name	Structure
69 ^c	1-((4-(Hydroxymethyl)phenyl)amino)cyclobutanecarbonitrile	
70 ^b	4-((1-Cyanocyclobutyl)amino)-3-fluoro-N-methylbenzamide	
71 ^e	Methyl 4-((1-cyanocyclobutyl)amino)-3-fluorobenzoate	
72 ^d	1-((2,3-Difluorophenyl)amino)cyclobutanecarbonitrile	
73 ^d	1-((2,5-Difluorophenyl)amino)cyclobutanecarbonitrile	
74 ^d	1-((2,3,6-Trifluorophenyl)amino)cyclobutanecarbonitrile	
75 ^d	1-((2,4-Difluorophenyl)amino)cyclobutanecarbonitrile	
76 ^c	2-Methyl-2-(p-tolylamino)propanenitrile	
77 ^c	1-((1H-Indazol-5-yl)amino)cyclobutanecarbonitrile	

Intermediate #	Name	Structure
78 ^c	1-((1H-Indazol-6-yl)amino)cyclobutanecarbonitrile	
79 ^e	1-(Benzo[d]oxazol-6-ylamino)cyclobutanecarbonitrile	
80 ^b	4-(4-((1-Cyanocyclobutyl)amino)phenyl)butanoic acid	
81 ^b	1-((4-(3-Hydroxypropyl)phenyl)amino)cyclobutanecarbonitrile	
82 ^e	1-((4-(4-Methylpiperazin-1-yl)phenyl)amino)cyclobutanecarbonitrile	
83 ^e	1-((4-(Pyridin-4-yl)phenyl)amino)cyclobutanecarbonitrile	
84 ^e	1-((4-(Pyridin-3-yl)phenyl)amino)cyclobutanecarbonitrile	
85 ^e	1-((4-(pyridin-2-yl)phenyl)amino)cyclobutanecarbonitrile	
86 ^e	tert-Butyl 4-(4-((1-cyanocyclobutyl)amino)phenyl)piperazine-1-carboxylate	

Intermediate #	Name	Structure
87 ^e	1-((4-((2-(Pyridin-4-yl)ethyl)sulfonyl)phenyl)amino)cyclobutanecarbonitrile	
88 ^e	1-((4-((Methyl(pyridin-4-ylmethyl)amino)methyl)phenyl)amino)cyclobutanecarbonitrile	
89 ^e	1-((4-((4-Methylpiperazin-1-yl)methyl)phenyl)amino)cyclobutanecarbonitrile	
90 ^b	1-((4-Bromo-3-fluorophenyl)amino)cyclobutanecarbonitrile	
91 ^c	1-((4-(6-Methylpyridin-3-yl)phenyl)amino)cyclobutanecarbonitrile	
92 ^c	1-((4-(4-Methylpyridin-3-yl)phenyl)amino)cyclobutanecarbonitrile	
93 ^c	1-((4-(5-Methylpyridin-3-yl)phenyl)amino)cyclobutanecarbonitrile	
94 ^c	1-((4-(2-Methylpyridin-3-yl)phenyl)amino)cyclobutanecarbonitrile	
95 ^a	1-((4-(Methyl(1-methylpiperidin-4-yl)amino)phenyl)amino)cyclobutanecarbonitrile	

Intermediate #	Name	Structure
96 ^c	1-((4-(Pyrimidin-5-yl)phenyl)amino)cyclobutanecarbonitrile	
97 ^c	1-((4-(5-Fluoropyridin-3-yl)phenyl)amino)cyclobutanecarbonitrile	
98 ^c	4-((1-Cyanocyclobutyl)amino)-N,2-dimethylbenzamide	
99 ^c	4-((1-Cyanocyclobutyl)amino)-2-methoxy-N-methylbenzamide	
100 ^c	2-Chloro-4-((1-cyanocyclobutyl)amino)-N-methylbenzamide	
101 ^c	1-((4-(Tetrahydro-2H-pyran-4-yl)phenyl)amino)cyclobutanecarbonitrile	
102 ^c	4-((1-Cyanocyclobutyl)amino)-N-methyl-2-(trifluoromethyl)benzamide	
103 ^c	4-((1-Cyanocyclopentyl)amino)-2-fluoro-N-methylbenzamide	
104 ^c	1-((1-Oxoisindolin-5-yl)amino)cyclobutanecarbonitrile	


Intermediate #	Name	Structure
105 ^c	1-((4-(Furan-2-yl)phenyl)amino)cyclobutanecarbonitrile	
106 ^c	1-((4-(5-Methylfuran-2-yl)phenyl)amino)cyclobutanecarbonitrile	
107 ^a	1-((4-Hydroxyphenyl)amino)cyclobutanecarbonitrile	

^a Aminonitrile synthesis method A, ^b Aminonitrile synthesis method B, ^c Aminonitrile synthesis method C, ^d Aminonitrile synthesis method D, ^e Aminonitrile synthesis method E.

INTERMEDIATE 108


4-((4,4,5,5,5-Pentafluoropentyl)thio)butan-1-ol

Step 1: 4-(Benzyloxy)butyl methanesulfonate



[00407] Methanesulfonyl chloride (1 mL, 13.3 mmol) was added to a cooled (0°C) solution of 4-(benzyloxy)butan-1-ol (2 g, 11.1 mmol) and triethylamine (1.9 mL, 13.3 mmol) in DCM (20 mL) and it was stirred at 0°C for 1h. HCl (1M) was added to the reaction mixture and the aqueous layer was extracted with DCM (2x). The organics were combined, dried over sodium sulfate, and evaporated to dryness to afford 4-(benzyloxy)butyl methanesulfonate as a yellow oil. LCMS [M + 1]⁺ 259.5.


Step 2: 4,4,5,5,5-Pentafluoropentyl 4-methylbenzenesulfonate



[00408] A mixture of 4,4,5,5,5-pentafluoropentan-1-ol (1 g, 5.6 mmol), p-toluenesulfonylchloride (1.3 g, 6.8 mmol), and triethylamine (1.2 mL, 8.4 mmol) in DCM (20 mL) was stirred at room temperature for 18h. Water was added to the reaction mixture, the two layers were separated, and the aqueous layer was extracted with DCM (2x). The organics were combined, dried over sodium sulfate, and evaporated to dryness. The residue was purified by


column chromatography on silica gel eluting with 0 to 20% EtOAc/hexanes to afford 1.7 g of 4,4,5,5,5-pentafluoropentyl 4-methylbenzenesulfonate as a colorless oil.

Step 3: 2-(4,4,5,5,5-Pentafluoropentyl)isothiuronium 4-methylbenzenesulfonate




- 5 [00409] A mixture of 4,4,5,5,5-pentafluoropentyl 4-methylbenzenesulfonate (1.7 g, 5.1 mmol) and thiourea (430 mg, 5.6 mmol) in ethanol (20 mL) was refluxed for 18h. The mixture was cooled to room temperature and most of the ethanol was removed in vacuo. Hexanes and Et₂O were added until a white precipitate started to form and the mixture was allowed to sit at room temperature for 30 min. The white solid was filtered, washed with hexanes, and dried to afford
- 10 1.7 g of 2-(4,4,5,5,5-pentafluoropentyl)isothiuronium 4-methylbenzenesulfonate as a white solid. LCMS [M + 1]⁺ 237.4.

Step 4: (4-(Benzyloxy)butyl)(4,4,5,5,5-pentafluoropentyl)sulfane




- 15 [00410] A mixture of 4-(benzyloxy)butyl methanesulfonate (759 mg, 2.9 mmol), 2-(4,4,5,5,5-pentafluoropentyl)isothiuronium 4-methylbenzenesulfonate (1.2 g, 2.9 mmol), NaOH (5M, 5 mL) in DMF (20 mL) was stirred at room temperature for 4h. EtOAc was added to the reaction mixture and the organic layer was washed with brine (2x), dried over sodium sulfate, and evaporated to dryness. The residue was purified by column chromatography on silica gel eluting with 0 to 20% EtOAc/hexanes to afford 960 mg of (4-(benzyloxy)butyl)(4,4,5,5,5-
- 20 pentafluoropentyl)sulfane as a yellow oil. LCMS [M + 1]⁺ 357.5.

Step 5: 4-((4,4,5,5,5-Pentafluoropentyl)thio)butan-1-ol




- [00411] BBr₃ (1M in DCM, 2.9 mL, 2.9 mmol) was added dropwise to a solution of (4-(benzyloxy)butyl)(4,4,5,5,5-pentafluoropentyl)sulfane (340 mg, 0.95 mmol) in DCM (5 mL)
- 25 cooled to -78°C. The temperature was allowed to warm up to room temperature and the mixture was stirred for 18h. Water was slowly added and the reaction mixture was stirred for 10 min. The two layers were separated and the aqueous layer was extracted with DCM (3x). The organics were combined, dried over sodium sulfate, and evaporated to dryness. The residue was purified by column chromatography on silica gel eluting with 50 to 100% EtOAc/hexanes
- 30 to afford 171 mg of 4-((4,4,5,5,5-pentafluoropentyl)thio)butan-1-ol as a colorless oil.

INTERMEDIATE 109**4-((4,4,5,5,5-Pentafluoropentyl)sulfinyl)butan-1-ol**


5 [00412] m-CPBA (64 mg, 0.37 mmol) was added to a cooled (0°C) solution of 4-((4,4,5,5,5-pentafluoropentyl)thio)butan-1-ol (76 mg, 0.28 mmol) in DCM (3 mL) and it was stirred at 0°C for 30 min. The reaction mixture was warmed to room temperature and was partitioned between a saturated solution of sodium bicarbonate and DCM. The organic layer was dried over sodium sulfate, and evaporated to dryness. The residue was purified by column chromatography on silica gel eluting with 0 to 10% MeOH/DCM to afford 52 mg of 4-((4,4,5,5,5-pentafluoropentyl)sulfinyl)butan-1-ol as a white solid.

INTERMEDIATE 110**5-Amino-3-(difluoromethyl)picolinonitrile**

15

Step 1: (5-Bromo-2-chloropyridin-3-yl)methanol


[00413] To a mixture of NaBH₄ (5.8 g, 152.6 mmol, 4 eq) and anhydrous CaCl₂ (16.9 g, 152.6 mmol) in dry DCM (100 mL) at 0°C, was added slowly methyl 5-bromo-2-chloronicotinate (9.5 g, 38.15 mmol). The resulting mixture was stirred at room temperature for 12h. Water was added to the reaction mixture at 0°C. The organic layer was separated, dried over sodium sulfate, and concentrated in vacuo, to afford 6.8 g (crude) of (5-bromo-2-chloropyridin-3-yl)methanol, which was used in the next step without further purification. LCMS [M+H]⁺ 223.1.

Step 2: 5-Bromo-2-chloronicotinaldehyde

25


[00414] To a solution of (5-bromo-2-chloropyridin-3-yl)methanol (6.8 g, 30.6 mmol) in DCM (200 mL), was added Dess-Martin periodinane (32.4 g, 76.6 mmol) at 0°C. The resulting mixture was stirred at room temperature for 12h. The reaction mixture was concentrated in vacuo and the residue was purified by column chromatography on silica gel (0 to 30% EtOAc in petroleum ether) to afford 5.05 g of 5-bromo-2-chloronicotinaldehyde. LCMS [M+H]⁺ 221.1.

Step 3: 5-Bromo-2-chloro-3-(difluoromethyl)pyridine



[00415] To a solution of 5-bromo-2-chloronicotinaldehyde (5.05 g, 23.2 mmol) in DCM (30 mL), were added ethanol (0.13 mL, 2.32 mmol) and DAST (7.51 g, 46.4 mmol) at room temperature. The resulting mixture was stirred for 2h at that temperature. To the reaction mixture was slowly added saturated aqueous sodium bicarbonate (100 mL). The organic layer was separated, dried over sodium sulfate, and concentrated in vacuo. The residue was purified by column chromatography on silica gel (0 to 50% EtOAc in petroleum ether) to afford 3.80 g of 5-bromo-2-chloro-3-(difluoromethyl)pyridine. LCMS $[M+H]^+$ 243.1.


Step 4: 5-Amino-3-(difluoromethyl)picolinonitrile



[00416] The title compound was synthesized as described in Intermediate 1 (Steps 3-5) using 5-bromo-2-chloro-3-(difluoromethyl)pyridine as the starting material. ^1H NMR (400 MHz, DMSO- d_6) δ 8.09 (d, 1H), 7.28-7.01 (m, 2H), 6.80 (s, 2H). LCMS $[M+H]^+$ 170.1.

EXAMPLE 1: Synthesis of Compound 1 (Thiohydantoin synthesis method A)

5-(5-(3-Fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile




[00417] Thiophosgene (1.3 mL, 16.04 mmol) was added to a mixture of 5-amino-3-(trifluoromethyl) picolinonitrile (Intermediate 1, 3 g, 16.0 mmol) and 1-((3-fluoro-4-hydroxyphenyl)amino)cyclobutane-carbonitrile (Intermediate 19, 3.3 g, 16.0 mmol) in DMA (40 mL) and the mixture was heated to 60°C for 18h. MeOH (60 mL) and HCl (2M, 40 mL) were added and the mixture was refluxed for 2h. The mixture was cooled to room temperature and slowly poured into an ice/water bath. The brown solid was filtered, washed with water, dried, and purified by column chromatography on silica gel eluting with 0 to 50% EtOAc/hexanes to afford 3 g of 5-(5-(3-fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a beige solid. ^1H NMR (300

MHz, DMSO- d_6) δ 10.44 (s, 1H), 9.21 (d, 1H), 8.74 (d, 1H), 7.27-7.05 (m, 3H), 2.64-2.42 (m, 4H), 2.01-1.91 (m, 1H), 1.61-1.54 (m, 1H).

EXAMPLE 2: Synthesis of Compound 2 (Thiohydantoin synthesis method B)


5 5-(4-Hydroxyphenyl)-6-thioxo-7-(3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-yl)-5,7-diazaspiro[3.4]octan-8-one



[00418] A mixture of 6-isothiocyanato-3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazine (Intermediate 8, 545 mg, 2.22 mmol) and 1-((4-hydroxyphenyl)amino)cyclobutanecarbonitrile (Intermediate 25, 418 mg, 2.22 mmol) in DMA (15 mL) was heated to 60°C for 2h. MeOH (10 mL) and HCl (2M, 10 mL) were added and the mixture was refluxed for 1h. The mixture was cooled to room temperature and slowly poured into an ice/water bath. The dark brown solid was filtered, washed with water, dried, and purified by reverse phase HPLC (Acetonitrile/water:TFA) to afford 5-(4-hydroxyphenyl)-6-thioxo-7-(3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-yl)-5,7-diazaspiro[3.4]octan-8-one as a light brown solid. LCMS $[M + 1]^+$ 435.4.

EXAMPLE 3: Synthesis of Compound 3

20 Ethyl 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoate




[00419] A mixture of thiophosgene (0.55 mL, 5.35 mmol), 5-amino-3-(trifluoromethyl)picolinonitrile (Intermediate 1, 1g, 5.35 mmol), and ethyl 4-((1-cyanocyclobutyl)amino)-2-fluorobenzoate (Intermediate 20, 1.4g, 5.35 mmol) in DMA (25 mL) was heated to 70°C overnight. MeOH (20 mL) and HCl (2M, 20 mL) were added and the mixture was refluxed for 2h then cooled to room temperature. Water/ice was added and the aqueous layer was extracted with DCM (4x). The organics were combined, dried over sodium sulfate, and evaporated to dryness. The residue was purified by column chromatography on silica gel eluting with 0 to 50% EtOAc/hexanes to afford 1g of ethyl 4-(7-(6-cyano-5-

(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoate as a yellow foam. ^1H NMR (300 MHz, DMSO- d_6) δ 9.21 (s, 1H), 8.74 (s, 1H), 8.13 (m, 1H), 7.53 (d, 1H), 7.45 (d, 1H), 4.37 (q, 2H), 2.65-2.62 (m, 2H), 2.56-2.45 (m, 2H), 2.05-1.91 (m, 1H), 1.61-1.57 (m, 1H), 1.34 (t, 3H).

5

EXAMPLE 4: Synthesis of Compound 4

5-(5-(4-Bromo-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile




- 10 [00420] Thiophosgene (0.36 mL, 4.67 mmol) was added dropwise to a solution of 5-amino-3-(trifluoromethyl)picolinonitrile (Intermediate 1, 795 mg, 4.25 mmol) and 1-((4-bromo-3-fluorophenyl)amino)cyclobutanecarbonitrile (Intermediate 90, 1.14g, 4.25 mmol) in DMA (30 mL). The resulting mixture was heated to 60°C overnight. MeOH (8 mL) and HCl (2M, 8 mL) were added and the mixture was refluxed for 2h then cooled to room temperature. Water was
- 15 slowly added until a light brown solid precipitated. The solid was filtered, washed with water, dried, dissolved in MeOH, and absorbed on silica gel. Purification by column chromatography on silica gel eluting with 10% EtOAc/hexanes afforded 616 mg of 5-(5-(4-bromo-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a white solid. ^1H NMR (300 MHz, DMSO- d_6) δ 9.21 (d, 1H), 8.74 (d, 1H), 8.00 (t, 1H), 7.55
- 20 (dd, 1H), 7.28 (dd, 1H), 2.67-2.59 (m, 2H), 2.55-2.45 (m, 2H), 1.99-1.95 (m, 1H), 1.61-1.56 (m, 1H).

EXAMPLE 5: Synthesis of Compound 5

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide

25




[00421] A mixture of 5-isothiocyanato-3-methylpicolinonitrile (Intermediate 6, 2.9g, 16.57 mmol) and 4-((1-cyanocyclobutyl)amino)-2-fluoro-N-methylbenzamide (Intermediate 21, 2.7g, 11.05 mmol) in DMA was heated to 65°C overnight. MeOH (50 mL) and HCl (2M, 70 mL)

were added and the mixture was heated to 105°C for 1h then cooled to room temperature.

Water was slowly added until a yellow solid precipitated and the mixture was stirred at room temperature for 3h. The solid was filtered, washed with water, dissolved in DCM, dried over sodium sulfate, and absorbed on silica gel. Purification by column chromatography on silica gel eluting with 0 to 100% EtOAc/hexanes afforded a pale yellow solid. This solid was triturated in MeOH, filtered, and dried to afford 1.39g (29%) of 4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide as a white solid. ¹H NMR (300 MHz, DMSO-d₆) δ 8.64 (d, 1H), 8.40 (m, 1H), 8.06 (d, 1H), 7.75 (t, 1H), 7.43 (dd, 1H), 7.32 (dd, 1H), 2.73 (d, 3H), 2.59-2.53 (m, 2H), 2.52 (s, 3H), 2.44-2.23 (m, 2H), 1.91-1.88 (m, 1H), 1.52-1.48 (m, 1H). LCMS [M + 1]⁺ 424.0.

EXAMPLE 6: Synthesis of Compound 6


4-(3-(6-Cyano-5-methylpyridin-3-yl)-4-oxo-2-thioxo-1,3-diazaspiro[4.4]nonan-1-yl)-2-fluoro-N-methylbenzamide



[00422] Thiophosgene (0.35 mL) was added dropwise to a solution of 4-((1-cyanocyclopentyl)amino)-2-fluoro-N-methylbenzamide (Intermediate 103, 400 mg, 1.53 mmol) and 5-amino-3-methylpicolinonitrile (Intermediate 2, 303 mg, 2.28 mmol) in DMA (30 mL) under N₂ atmosphere. The resulting mixture was stirred at 60°C overnight. MeOH (22 mL) and HCl (2M, 11 mL) were added and the mixture was refluxed for 2h then cooled to room temperature. The reaction mixture was poured into water (75 mL) and extracted with EtOAc. The organic layer was dried over sodium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica gel (1:5 EtOAc/petroleum ether), to afford 300 mg of 4-(3-(6-cyano-5-methylpyridin-3-yl)-4-oxo-2-thioxo-1,3-diazaspiro[4.4]nonan-1-yl)-2-fluoro-N-methylbenzamide as a light yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.74 (s, 1H), 8.47 (m, 1H), 8.17 (s, 1H), 7.78 (m, 1H), 7.55 (d, 1H), 7.41 (d, 1H), 2.81 (d, 3H), 2.60 (s, 3H), 2.29-2.21 (m, 4H), 1.72 (m, 2H), 1.41 (m, 2H). LCMS [M+1]⁺ 438.1.

EXAMPLE 7: Synthesis of Compound 7

3-Methyl-5-(5-(4-(5-methylfuran-2-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile




[00423] Thiophosgene (0.1 mL) was added dropwise to a solution of 1-((4-(5-methylfuran-2-yl)phenyl)amino)cyclobutanecarbonitrile (Intermediate 106, 280 mg, 1.11 mmol) and 5-amino-3-methylpicolinonitrile (Intermediate 2, 177 mg, 1.33 mmol) in DMA (20 mL). The resulting mixture was stirred at 60°C overnight. MeOH (16 mL) and HCl (2M, 8 mL) were added and the mixture was refluxed for 2h then cooled to room temperature. The reaction mixture was poured into water (200 mL) and extracted with EtOAc. The organic layer was washed with brine, dried over Sodium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica gel (1:3 EtOAc/petroleum ether), to afford 140 mg of 3-methyl-5-(5-(4-(5-methylfuran-2-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile. ¹HNMR (400 MHz, DMSO-d₆) δ 8.74 (s, 1H), 8.16 (s, 1H), 7.86 (d, 2H), 7.46 (d, 2H), 6.98 (d, 1H), 6.28 (d, 1H), 2.63 (m, 2H), 2.60 (s, 3H), 2.51-2.43 (m, 2H), 2.38 (s, 3H), 1.97 (m, 1H), 1.57 (m, 1H). LCMS [M+1]⁺ 429.1.

[00424] **Compounds 8 to 117** were synthesized from the appropriate amines (amines were either commercially available, synthesized in the Intermediates section, or synthesized from literature procedures) and amino cyanohydrins, according to the conditions indicated (see footnotes): 8^a, 9^a, 10^a, 11^a, 12^a, 13^a, 14^a, 15^a, 16^b, 17^b, 18^b, 19^a, 20^a, 21^a, 22^a, 23^a, 24^b, 25^b, 26^b, 27^b, 28^b, 29^a, 30^b, 31^b, 32^a, 33^a, 34^b, 35^b, 36^b, 37^b, 38^a, 39^b, 40^b, 41^b, 42^a, 43^a, 44^b, 45^b, 46^a, 47^b, 48^b, 49^b, 50^b, 51^b, 52^b, 53^b, 54^b, 55^b, 56^a, 57^b, 58^b, 59^b, 60^b, 61^b, 62^b, 63^a, 64^a, 65^b, 66^b, 67^a, 68^a, 69^a, 70^a, 71^a, 72^a, 73^a, 74^a, 75^a, 76^a, 77^a, 78^a, 79^a, 80^b, 81^b, 82^b, 83^b, 84^b, 85^b, 86^a, 87^a, 88^a, 89^b, 90^b, 91^b, 92^b, 93^b, 94^a, 95^a, 96^a, 97^a, 98^a, 99^a, 100^a, 101^a, 102^a, 103^a, 104^a, 105^a, 106^a, 107^a, 108^a, 109^b, 110^a, 111^b, 112^a, 113^a, 114^a, 115^a, 116^b, 117^b. ^a Thiohydantoin synthesis method A. ^b Thiohydantoin synthesis method B.

EXAMPLE 8: Synthesis of Compound 118

5-(8-Oxo-5-(4-(piperazin-1-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile




[00425] A mixture of *tert*-butyl 4-(4-(1-cyanocyclobutylamino)phenyl)piperazine-1-carboxylate (Intermediate 86, 0.25 g, 0.70 mmol) and 5-isothiocyanato-3-(trifluoromethyl)picolinonitrile (Intermediate 10, 241 mg, 1.05 mmol) in DMA (5 mL) was heated to 80°C for 12 h. 2M

HCl/MeOH = 1/1 (10 mL) was added then and the resulting mixture was heated at 100 °C for 2h. The reaction mixture was cooled to room temperature and diluted with EtOAc. The organic layer was washed with water (3x), dried over sodium sulfate, and concentrated in vacuo. The residue was purified by Prep-HPLC (20% acetonitrile in water) to afford 75 mg of 5-(8-oxo-5-(4-(piperazin-1-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl) -3-

(trifluoromethyl)picolinonitrile. ¹H NMR (DMSO-d₆, 300 MHz) δ 9.26 (s, 1H), 9.19 (m, 1H), 8.74 (m, 1H), 7.25 (d, 2H), 7.15 (d, 2H), 3.48 (m, 4H), 3.22 (m, 4H), 2.60-2.49 (m, 2H), 2.48-2.40 (m, 2H), 1.95 (m, 1H), 1.63 (m, 1H). LCMS [M+H]⁺ 487.1.

EXAMPLE 9: Synthesis of Compound 119


[0001] *tert*-Butyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)piperazine-1-carboxylate



[00426] To a solution of 5-(8-oxo-5-(4-(piperazin-1-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl) -3-(trifluoromethyl)picolinonitrile (15 mg, 0.031 mmol) in dry DCM (2 mL) was added TEA (6 µl, 0.062 mmol) and (Boc)₂O (10 mg, 0.046 mmol). The mixture was stirred at room temperature for 3h and then concentrated. The residue was purified by column chromatography on silica gel (0 to 20% EtOAc in petroleum ether) to afford 10 mg of *tert*-butyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)piperazine-1-carboxylate. ¹H NMR (CDCl₃, 300 MHz) δ 9.10 (d, 1H), 8.36 (d, 1H), 7.18 (d, 2H), 7.04 (d, 2H), 3.60 (m, 4H), 3.27 (m, 4H), 2.63-2.60 (m, 4H), 2.24-1.97 (m, 1H), 1.71-1.63 (m, 1H), 1.49 (s, 9H).

EXAMPLE 10: Synthesis of Compound 120


4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-methylbenzenesulfonamide



[00427] A mixture of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro-[3.4]octan-5-yl)benzenesulfonamide (Compound 81, 97 mg, 0.20 mmol), methyl tosylate (37 mg, 0.20 mmol), and Cs₂CO₃ (65 mg, 0.20 mmol) in acetonitrile (1 mL) was heated at 50°C for 20h in a sealed tube. After being cooled room temperature, the reaction mixture was diluted with water and extracted with EtOAc (3x). The organic layer was dried over magnesium sulfate and concentrated to give a residue that was purified by column chromatography on silica gel (hexanes:DCM:acetone = 5:9:1) to afford 30 mg of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]-octan-5-yl)-N-methylbenzenesulfonamide as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.11 (s, 1H), 8.37 (s, 1H), 8.10 (dd, 2H), 7.58-7.49 (m, 2H), 4.52 (d, 1H), 2.80 (m, 3H), 2.78-2.69 (m, 2H), 2.63-2.47 (m, 2H), 2.29 (m, 1H), 1.74 (m, 1H). LCMS [M + 1 + Na]⁺ 518.0.

EXAMPLE 11: Synthesis of Compound 121


4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N,N-dimethylbenzenesulfonamide



[00428] The title compound was obtained as a side product from the procedure described for Example 10. ¹H NMR (400 MHz, CDCl₃) δ 9.11 (s, 1H), 8.37 (s, 1H), 8.02 (dd, 2H), 7.52 (dd, 2H), 2.84 (s, 6H), 2.83-2.68 (m, 2H), 2.55 (m, 2H), 2.37-2.21 (m, 1H), 1.80-1.64 (m, 1H). LCMS [M + Na]⁺ 532.0.

EXAMPLE 12: Synthesis of Compound 122


Methyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)butanoate




[00429] A mixture of 5-isothiocyanato-3-trifluoromethylpyridine-2-carbonitrile (Intermediate 10, 582 mg, 2.54 mmol) and 4-(4-(1-cyanocyclobutylamino)phenyl)butanoic acid (Intermediate 80, 410 mg, 1.59 mmol) in DMF (3 mL) was stirred at room temperature overnight. To this mixture were added methanol (4 mL) and HCl (2M, 2 mL) and the resulting mixture was refluxed for 3h. After being cooled room temperature, the reaction mixture was poured into cold water and extracted with EtOAc (3x). The organic layers were dried over magnesium sulfate and concentrated to dryness. Column chromatography of the residue on silica gel (Hexane:EtOAc = 4:1) afforded 256 mg of methyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)-pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)butanoate as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.11 (d, 1H), 8.38 (d, 1H), 7.42 (d, 2H), 7.36-7.19 (m, 2H), 3.70 (s, 3H), 2.87-2.75 (m, 2H), 2.75-2.53 (m, 4H), 2.42 (t, 2H), 2.26 (m, 1H), 2.05 (m, 2H), 1.78-1.61 (m, 1H). LCMS [M + 1]⁺ 503.0.

EXAMPLE 13: Synthesis of Compound 123

5-(5-(4-Fluoro-3-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile



[00430] Trimethylsilyl iodide (7.0 ml, 51.1 mmol) was added to a solution of 5-(5-(4-fluoro-3-methoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Compound 32, 3 g, 5.1 mmol) in acetonitrile (50 mL) and the mixture was refluxed for 24h. The mixture was cooled to room temperature, poured into water, and extracted with EtOAc (3x). The organics were combined, washed with sat'd solution of NaHSO₃, dried over MgSO₄, and evaporated to dryness. The residue was purified by column chromatography on silica gel eluting with EtOAc/Hexane = 1/5 to afford 1.1 g of 5-(5-(4-fluoro-3-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile. ¹H NMR (400 MHz, CDCl₃) δ 9.10 (d, 1H), 8.38 (d, 1H), 7.27-7.22 (m, 1H), 6.93 (dd, 1H), 6.78 (m, 1H), 2.74-2.55 (m, 1H), 2.23 (m, 1H), 1.80-1.59 (m, 4H).


EXAMPLE 14: Synthesis of Compound 124**5-(5-(4-(3-(4-Methylpiperazin-1-yl)propyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile**

- 5 [00431] A solution of 5-(5-(4-(3-hydroxypropyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Compound 86, 66 mg, 0.14 mmol) and methanesulfonyl chloride (0.033 mL, 0.42 mmol) in DCM (2 mL) was stirred at room temperature for 2h. The solution was washed with water (2x) and brine, dried over magnesium sulfate and concentrated to give 80 mg of the corresponding mesylate without purification. The
- 10 mesylate (45 mg, 0.084 mmol), methyl piperazine (0.02 mL, 0.17 mmol) and triethylamine (0.04 mL, 0.25 mmol) in dichloromethane was stirred at room temperature overnight, then at reflux for 16h. The mixture was partitioned between water and DCM and the aqueous was further extracted with DCM (2x). The combined organics were washed with brine, dried (MgSO₄) and concentrated. Purification by silica gel chromatography (0 to 80% EtOAc/hexanes) afforded 20 mg of 5-(5-(4-(3-(4-methylpiperazin-1-yl)propyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a beige solid. ¹H NMR (300 MHz, DMSO-d₆) δ 9.23 (d, 1H), 8.77 (d, 1H), 7.45 (d, 2H), 7.31 (d, 2H), 2.76-2.57 (m, 4H), 2.49-2.23 (m, 5H), 2.14 (s, 3H), 2.02-1.89 (m, 1H), 1.85-1.73 (m, 3H), 1.59-1.46 (m, 1H). LCMS [M + 1]⁺ 543.1.

20

EXAMPLE 15: Synthesis of Compound 125**5-(5-(Benzo[d]oxazol-6-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile****Step 1: 5-(5-(4-Amino-3-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile**


25



[00432] A mixture of 5-amino-3-(trifluoromethyl)picolinonitrile (Intermediate 1, 110 mg, 0.59 mmol), benzo[d]oxazol-6-amine (Intermediate 79, 125 mg, 0.59 mmol), and thiophosgene (0.05

mL, 0.60 mmol) in anhydrous DMA (3 mL) was stirred at 60°C for 16h. Methanol (3 mL) and 2M HCl (1.5 mL) were then added and the mixture was heated at 90°C for 2h. The reaction mixture was cooled and poured into ice/water (40 mL) and the resultant solid was filtered to afford 150 mg of 5-(5-(4-amino-3-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a brown powder. LCMS $[M + 1]^+$ 434.0.


Step 2: 5-(5-(Benzo[d]oxazol-6-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile



[00433] A mixture of 5-(5-(4-amino-3-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (60 mg, 0.14 mmol) and triethylorthoformate (0.050 mL, 0.28 mmol) in anhydrous DMF (1 mL) was heated at 100°C, 1h. The mixture was diluted with water, extracted with EtOAc (3x), the organics combined and washed with water (2x), brine, dried ($MgSO_4$) and concentrated. Purification by column chromatography, eluting with 30% EtOAc/hexanes provided 23 mg of 5-(5-(benzo[d]oxazol-6-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a pale yellow solid. ^1H NMR (300 MHz, $DMSO-d_6$) δ 9.24 (d, 1H), 8.95 (s, 1H), 8.78 (d, 1H), 8.05 (d, 1H), 7.94 (d, 1H), 7.45 (dd, 1H), 2.74-2.61 (m, 2H), 2.59-2.47 (m, 2H), 2.03-1.90 (m, 1H), 1.61-1.48 (m, 1H). LCMS $[M + 1]^+$ 444.0.

EXAMPLE 16: Synthesis of Compound 126

5-(5-(3-Fluoro-4-(2-(1-methyl-1H-pyrazol-5-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile




[00434] A mixture of 5-(5-(3-fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Compound 1, 100 mg, 0.23 mmol), 2-(1-methyl-1H-pyrazol-5-yl)ethanol (43.5 mg, 0.34 mmol), triphenylphosphine (89 mg, 0.34 mmol), and diisopropyl azodicarboxylate (0.07 mL, 0.34 mmol) in THF (5 mL) was stirred at room temperature for 18h. The reaction mixture was absorbed on silica gel and purified by column chromatography on silica gel eluting with EtOAc/hexanes to afford impure desired product that

was repurified by reverse phase HPLC (acetonitrile/water:TFA). The fractions containing the desired compound were combined, acetonitrile was removed in vacuo, and the remaining aqueous layer was treated with a saturated solution of sodium bicarbonate. The aqueous layer was extracted with DCM (3x), the organics were combined, dried over sodium sulfate, and evaporated to dryness to afford 29 mg of 5-(5-(3-fluoro-4-(2-(1-methyl-1H-pyrazol-5-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a beige solid. LCMS $[M + 1]^+$ 545.5.

EXAMPLE 17: Synthesis of Compound 127


5-(5-(3-Fluoro-4-(2-(pyrrolidin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile



[00435] A mixture of 5-(5-(3-fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Compound 1, 100 mg, 0.23 mmol), 2-(pyrrolidin-1-yl)ethanol (35 μ L, 0.30 mmol), triphenylphosphine (79 mg, 0.30 mmol), and diisopropyl azodicarboxylate (60 μ L, 0.30 mmol) in THF (5 mL) was stirred at room temperature for 2 days. The reaction mixture was absorbed on silica gel and purified by column chromatography on silica gel eluting with 0 to 10% MeOH/DCM to afford impure desired product that was repurified by reverse phase HPLC (acetonitrile/water, 0.1% TFA). The fractions containing the desired compound were combined, acetonitrile was removed in vacuo, and the remaining aqueous layer was treated with a saturated solution of sodium bicarbonate. The aqueous layer was extracted with DCM (3x), the organics were combined, dried over sodium sulfate, and evaporated to dryness to afford 30 mg of 5-(5-(3-fluoro-4-(2-(pyrrolidin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a white solid. ^1H NMR (300 MHz, DMSO- d_6) δ 9.21 (d, 1H), 8.74 (d, 1H), 7.40 (t, 1H), 7.35 (dd, 1H), 7.22 (d, 1H), 4.24 (t, 2H), 2.86 (t, 2H), 2.65-2.42 (m, 8H), 1.98-1.95 (m, 1H), 1.70 (m, 4H), 1.60-1.54 (m, 1H). LCMS $[M + 1]^+$ 534.9.

EXAMPLE 18: Synthesis of Compound 128


5-(5-(4-(Benzyloxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile



[00436] A mixture of 5-(5-(3-fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Example 1, 100 mg, 0.23 mmol), benzyl alcohol (35 μ L, 0.30 mmol) triphenylphosphine (79 mg, 0.30 mmol), and diisopropyl azodicarboxylate (60 μ L, 0.30 mmol) in THF (5 mL) was stirred at room temperature for 2 days then heated to 60°C overnight. The reaction mixture was absorbed on silica gel and purified by column chromatography on silica gel eluting with 0 to 50% EtOAc/Hexanes to afford impure desired product. The solid was triturated in MeOH, filtered, and dried to afford 36 mg of 5-(5-(4-(benzyloxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a white solid. ^1H NMR (300 MHz, DMSO- d_6) δ 9.21 (d, 1H), 8.75 (d, 1H), 7.53-7.36 (m, 7H), 7.23 (d, 1H), 5.28 (s, 2H), 2.66-2.60 (m, 2H), 2.53-2.43 (m, 2H), 2.02-1.92 (m, 1H), 1.61-1.55 (m, 1H).

EXAMPLE 19: Synthesis of Compound 129

5-(5-(4-((1-Methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile




[00437] To a solution of 5-(5-(4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Compound 63, 150 mg, 0.36 mmol), 1-methylpiperidin-4-ol (50 μ L, 0.39 mmol), and triphenylphosphine (125 mg, 0.47 mmol) in anhydrous THF (2 mL) was added diisopropyl azodicarboxylate (0.1 mL, 0.47 mmol) and the reaction mixture was stirred at room temperature overnight. Additional 1-methylpiperidin-4-ol (50 μ L, 0.39 mmol), triphenylphosphine (125 mg, 0.47 mmol) and diisopropyl azodicarboxylate (0.1 mL, 0.47 mmol) was added and the reaction was allowed to stir an additional 16h. The mixture was partitioned between EtOAc and water and the aqueous layer was extracted with EtOAc (2x). The combined organics were washed with water and brine, then dried over magnesium sulfate and concentrated in vacuo. Purification by silica gel chromatography (0 to 80% EtOAc/hexanes, then 0 to 5% MeOH/dichloromethane) gave impure product which was further

purified by preparative HPLC (40 to 75% acetonitrile/water, 0.1% TFA) to afford 45 mg of 5-(5-(4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a pale yellow solid. ¹H NMR (300 MHz, DMSO-d₆) δ 9.22 (d, 1H), 8.76 (d, 1H), 7.30 (d, 2H), 7.15 (d, 2H), 4.52-4.39 (m, 1H), 2.72-2.56 (m, 4H), 2.49-2.38 (m, 2H), 2.25-2.14 (m, 5H), 2.03-1.90 (m, 3H), 1.77-1.49 (m, 3H). LCMS [M + 1]⁺ 516.0.

[00438] **Compounds 130 to 201** were synthesized following the procedure described in Example 16 from the appropriate phenols and alcohols (phenols and alcohols were either commercially available, synthesized in the Intermediates section, or synthesized from published procedures).

EXAMPLE 20: Synthesis of Compound 202


5-(5-(4-(2-(4-Acetylpiperazin-1-yl)ethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile



[00439] A mixture of 5-(5-(3-fluoro-4-(2-(piperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Compound 174, 50 mg, 0.1 mmol), acetic anhydride (13 μL, 0.14 mmol), and triethylamine (40 μL, 0.27 mmol) in DCM (2 mL) was stirred at room temperature for 2 days. The reaction mixture was absorbed on silica gel and purified by column chromatography on silica gel eluting with 0 to 10% MeOH/DCM to afford 34 mg of 5-(5-(4-(2-(4-acetylpiperazin-1-yl)ethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a white solid. ¹H NMR (300 MHz, DMSO-d₆) δ 9.21 (s, 1H), 8.74 (s, 1H), 7.42 (t, 1H), 7.35 (dd, 1H), 7.22 (m, 1H), 4.28 (t, 2H), 3.42 (m, 4H), 2.80 (t, 2H), 2.65-2.59 (m, 2H), 2.54-2.44 (m, 6H), 1.98 (m, 4H), 1.57 (m, 1H). LCMS [M + 1]⁺ 591.9.

EXAMPLE 21: Synthesis of Compound 203

5-(5-(3-Fluoro-4-methoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile




[00440] A mixture of 5-(5-(3-fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Compound 1, 100 mg, 0.23 mmol), methyl iodide (0.14 mL, 0.23 mmol), and potassium carbonate (31 mg, 0.23 mmol) in acetone (4 mL) was stirred at room temperature for 4h. Excess potassium carbonate was removed and the reaction mixture was absorbed on silica gel and purified by column chromatography on silica gel eluting with 0 to 30% EtOAc/hexanes to afford 74 mg of 5-(5-(3-fluoro-4-methoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a white solid. LCMS $[M + 1]^+$ 451.4

[00441] **Compounds 204 and 205** were synthesized following the procedure described in Example 21 from the appropriate phenols.

EXAMPLE 22: Synthesis of Compound 206


5-(8-Oxo-5-(4-(pyrimidin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile



[00442] A mixture of 5-(5-(4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Compound 63, 50 mg, 0.12 mmol), 2-chloropyrimidine (17 mg, 0.15 mmol), and cesium carbonate (60 mg, 0.18 mmol) in anhydrous THF (1.2 mL) was heated at reflux overnight. The reaction mixture was cooled to room temperature and water was added. The aqueous was extracted with EtOAc (3x), the organics combined and washed with brine, dried ($MgSO_4$) and concentrated. Purification by column chromatography on silica gel eluting with 0 to 60% EtOAc/hexanes provided 30 mg of 5-(8-oxo-5-(4-(pyrimidin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as an off-white solid. ^1H NMR (300 MHz, $DMSO-d_6$) δ 9.24 (d, 1H), 8.78 (d, 1H), 8.73 (d, 2H), 7.48 (dd, 4H), 7.35 (t, 1H), 2.72-2.61 (m, 2H), 2.54-2.42 (m, 2H), 2.10-1.93 (m, 1H), 1.69-1.52 (m, 1H). LCMS $[M + 1]^+$ 497.8.

EXAMPLE 23: Synthesis of Compound 207

5-(8-Oxo-5-(4-(pyrazin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile




[00443] A mixture of 5-(5-(4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (Compound 63, 100 mg, 0.24 mmol), NaH (14 mg, 0.36 mmol, 60% dispersion in oil) and chloropyrazine (0.025 mL, 0.29 mmol) in anhydrous DMF (1.4 mL) was heated at 90°C for 16h. Additional chloropyrazine (0.025 mL, 0.29 mmol) was added and heating was continued for 16 h. The reaction mixture was cooled to room temperature and water was added. The aqueous was extracted with EtOAc (3x), the organics combined and washed with brine, dried (MgSO₄) and concentrated. Purification by column chromatography on silica gel eluting with 0 to 50% EtOAc/hexanes afforded 50 mg of impure product. Further purification by preparative HPLC (30 to 100% acetonitrile/water, 10 min) provided 10 mg of 5-(8-oxo-5-(4-(pyrazin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as an off-white solid. ¹H NMR (300 MHz, DMSO-d₆) δ 9.23 (d, 1H), 8.78 (d, 1H), 8.65 (d, 1H), 8.46 (d, 1H), 8.31 (dd, 1H), 7.48 (dd, 4H), 2.75-2.60 (m, 2H), 2.57-2.42 (m, 2H), 2.06-1.94 (m, 1H), 1.68-1.53 (m, 1H). LCMS [M + 1]⁺ 496.9.

[00444] **Compounds 208 and 209** were synthesized following the procedure described in Example 23 from the appropriate phenols.

EXAMPLE 24: Synthesis of Compound 210

3-Methyl-5-(8-oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile




[00445] Tert-butyl 4-(4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidine-1-carboxylate (prepared by reaction of 5-(5-(4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile (Compound 77) with tert-butyl 4-hydroxypiperidine-1-carboxylate, according to Example 16, 300 mg, 0.55 mmol) was stirred in 2M HCl/methanol (1.5 mL) at room temperature overnight. The mixture was concentrated and purified by column chromatography, eluting with 10%

MeOH/DCM to provide 230 mg of 3-methyl-5-(8-oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile as a white solid. ¹H NMR (300 MHz, DMSO-d₆) δ 8.72 (d, 1H), 8.14 (d, 1H), 7.30 (d, 2H), 7.13 (d, 2H), 4.53-4.42 (m, 1H), 3.94-3.86 (m, 1H), 3.02-2.83 (m, 3H), 2.66-2.50 (m, 5H), 2.49-2.33 (m, 3H), 2.11-1.80 (m, 3H), 1.60-1.33 (m, 3H). LCMS [M + 1]⁺ 448.1.

EXAMPLE 25: Synthesis of Compound 211


5-(8-Oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile



[00446] The title compound was synthesized as described in Example 24 using Compound 63 as the starting material. ¹H NMR (300 MHz, DMSO-d₆) δ 9.22 (d, 1H), 8.75 (d, 1H), 7.29 (d, 2H), 7.16 (d, 2H), 4.48 (m, 1H), 3.32 (s, 1H), 3.01-2.94 (dt, 2H), 2.64-2.57 (m, 4H), 2.51-2.38 (m, 2H), 1.98-1.95 (m, 3H), 1.58-1.43 (m, 3H). LCMS [M + 1]⁺ 502.2.

EXAMPLE 26: Synthesis of Compound 212

5-(8-Oxo-5-(4-((1-propionylpiperidin-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile



[00447] Propionyl chloride (17 μL, 0.2 mmol) was added to a mixture of 5-(8-oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (50 mg, 0.1 mmol) and triethylamine (28 μL, 0.2 mmol) in THF (2mL) and the resulting milky mixture was stirred at room temperature overnight. Methanol was added to quench the reaction and the mixture was absorbed on silica gel and purified by column chromatography on silica gel eluting with 50 to 100% EtOAc/hexanes to afford 21 mg of 5-(8-oxo-5-(4-((1-propionylpiperidin-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as a white solid. ¹H NMR (300 MHz, DMSO-d₆) δ 9.15


(d, 1H), 8.69 (s, 1H), 7.25 (d, 2H), 7.13 (d, 2H), 4.62 (m, 1H), 3.85 (m, 1H), 3.62 (m, 1H), 3.29 (m, 1H), 3.15 (m, 1H), 2.55-2.52 (m, 2H), 2.42 (m, 2H), 2.39-2.25 (m, 2H), 1.92-1.88 (m, 3H), 1.50-1.45 (m, 3H), 0.93 (t, 3H). LCMS $[M + 1]^+$ 558.1.

[00448] **Compounds 213 to 219** were synthesized following the procedure described in

- 5 Example 26 from the appropriate piperidines and the appropriate acid chlorides, chloroformates, alkyl bromides, or sulfonyl chlorides.

EXAMPLE 27: Synthesis of Compound 220


5-(5-(4-((1-Isopropylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile



[00449] To a solution of 5-(8-oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (50 mg, 0.10 mmol) in anhydrous THF (1 mL) was added 2-bromopropane (20 μ L, 0.20 mmol) followed by cesium carbonate (100 mg, 0.30 mmol). The reaction mixture was heated at 70°C overnight. DMF (1 mL) was added to improve solubility and the reaction was heated at 85°C for 3 h. The mixture was cooled to room temperature and partitioned between water and EtOAc. The aqueous was extracted further with EtOAc (2x) and the combined organics were washed with water, then brine. The organics were dried over magnesium sulfate, filtered and concentrated. Purification by preparative HPLC provided 5 mg of 5-(5-(4-((1-isopropylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile as an off-white solid. ^1H NMR (300 MHz, DMSO- d_6) δ 9.04 (d, 1H), 8.30 (d, 1H), 7.13 (d, 2H), 6.99 (d, 2H), 4.37-4.26 (m, 1H), 2.81-2.33 (m, 10H), 2.25-1.74 (m, 5H), 1.00 (d, 6H). LCMS $[M + 1]^+$ 544.0.

EXAMPLE 28: Synthesis of Compound 221


Ethyl 2-(4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidin-1-yl)acetate



[00450] The title compound was synthesized as described in Example 27 using 5-(8-oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile and ethyl 2-bromoacetate as starting materials. LCMS $[M + 1]^+$ 588.1.

EXAMPLE 29: Synthesis of Compound 222


4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidine-1-carboxamide



[00451] A mixture of 5-(8-oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile (70 mg, 0.14 mmol) and isocyanatotrimethylsilane (23 μ L, 0.17 mmol) in DCM (3 mL) was stirred at room temperature overnight. Water was added and the mixture was stirred vigorously for 1h. The aqueous layer was extracted with DCM (3x), the organics were combined, dried over sodium sulfate, and evaporated to dryness. Column chromatography on silica gel (0 to 20% MeOH/DCM) afforded 55 mg of 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidine-1-carboxamide as a white solid. ^1H NMR (300 MHz, DMSO- d_6) δ 9.15 (s, 1H), 8.69 (s, 1H), 7.23 (d, 2H), 7.12 (d, 2H), 5.92 (s, 2H), 4.53 (m, 1H), 3.67-3.63 (m, 2H), 3.08-3.04 (m, 2H), 2.44 (m, 2H), 2.39 (m, 2H), 1.88 (m, 3H), 1.46 (m, 3H). LCMS $[M + 1]^+$ 545.2

EXAMPLE 30: Synthesis of Compound 223


5-(5-(4-(2-Hydroxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile



[00452] Dihydrofuran-2,5-dione (604 mg, 6.87 mmol) in DMF (5mL) was added to a mixture of 5-(5-(4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile (500 mg, 1.37 mmol) and potassium carbonate (378 mg, 2.74 mmol) in DMF (10 mL) and the resulting mixture was heated to 85°C overnight. The mixture was cooled to room temperature and partitioned between water and EtOAc. The aqueous was extracted further with EtOAc (2x) and the combined organics were washed with water, then brine. The organics were dried over sodium sulfate, filtered, and concentrated. Column chromatography on silica gel (50 to 100% EtOAc/hexanes) afforded 396 mg of 5-(5-(4-(2-hydroxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile as a white solid. ¹H NMR (300 MHz, DMSO-d₆) δ 8.71 (d, 1H), 8.14 (d, 1H), 7.33 (d, 2H), 7.13 (d, 2H), 4.92 (t, 1H), 4.11-4.05 (m, 2H), 3.78-3.73 (m, 2H), 2.63-2.5 (m, 5H), 2.49-2.37 (m, 2H), 2.00-1.90 (m, 1H), 1.56-1.51 (m, 1H). LCMS [M + 1]⁺ 409.0.


EXAMPLE 31: Synthesis of Compound 224

4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid



[00453] A mixture of ethyl 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoate (Compound 3, 1 g, 2.03 mmol) and NaOH (3M, 10 mL) in MeOH (10 mL) was stirred at room temperature for 18h. Aqueous HCl (2M) was added to the reaction mixture until the pH =2 and the aqueous layer was extracted with DCM (5x), the organics were combined, dried over sodium sulfate, and evaporated to dryness to afford 900 mg of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid as a light yellow solid. ¹H NMR (300 MHz, DMSO-d₆) δ 13.58 (s, 1H), 9.22 (s, 1H), 8.75 (s, 1H), 8.13 (t, 1H), 7.51 (dd, 1H), 7.43 (dd, 1H), 2.69-2.49 (m, 4H), 2.03-1.93 (m, 1H), 1.63-1.57 (m, 1H).


[00454] **Compounds 225 to 228** were synthesized following the procedure described in Example 31 from the corresponding ester.

EXAMPLE 32: Synthesis of Compound 229**5-(7-(6-Carbamoyl-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid**

[00455] The title compound was synthesized as a by-product in the synthesis of Compound 226.

¹H NMR (300 MHz, DMSO-d₆) δ 13.63 (s, 1H), 8.95 (s, 1H), 8.50 (s, 1H), 8.27 (s, 1H), 7.95 (m, 2H), 7.72 (m, 1H), 7.58 (t, 1H), 2.65 (m, 2H), 2.45 (m, 2H), 1.97 (m, 1H), 1.56 (m, 1H).

LCMS [M + 1]⁺ 483.4.


EXAMPLE 33: Synthesis of Compound 230**4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)-N-methylbutanamide**

[00456] To a solution of 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)butanoic acid (Compound 227, 85 mg, 0.17 mmol) in DCM (2 mL) were added triethylamine (0.013 mL, 0.09 mmol) and 4-nitrophenyl chloroformate (15 mg, 0.07 mmol). The mixture was stirred at room temperature for 1h and methylamine (0.3 mL, 2M in THF) was added. After 30 min, the mixture was diluted with water and extracted with

EtOAc (2x). The organic layers were dried over magnesium sulfate and concentrated. Column chromatography on silica gel (Hexane:EtOAc = 1:1) afforded 35 mg (40%) of 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)-N-methylbutanamide as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.11 (d, 1H), 8.38 (d, 1H), 7.42 (d, 2H), 7.22 (d, 2H), 2.86-2.81 (m, 3H), 2.81-2.73 (t, 2H), 2.73-2.64 (m, 2H), 2.64-2.53 (m, 2H), 2.26 (t, 2H), 2.24-2.17 (m, 1H), 2.11-2.01 (m, 2H), 1.69 (m, 1H). LCMS [M + 1]⁺ 502.0.

EXAMPLE 34: Synthesis of Compound 231

4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide




[00457] To a suspension of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid (Compound 224, 500 mg, 1.08 mmol) in DCM was added DMF (cat., 0.1 mL), followed by oxalyl chloride (0.14 mL, 1.61 mmol). The mixture was stirred at room temperature for 4h then concentrated in vacuo to produce a yellow residue that was further dried on a high vacuum pump. Ammonia (0.5M in dioxane, 40 mL, 20 mmol) was directly added to the residue and the mixture was stirred at room temperature overnight. MeOH was added and the mixture was absorbed onto silica gel and purified by flash chromatography (50 to 100% EtOAc/Hexanes) to afford impure desired product that was repurified by reverse phase HPLC (acetonitrile/water:TFA). The fractions containing the desired compound were combined, acetonitrile was removed in vacuo, and the remaining aqueous layer was treated with a saturated solution of sodium bicarbonate. The aqueous layer was extracted with DCM (3x), the organics were combined, dried over sodium sulfate, and evaporated to dryness to afford 100 mg of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide as a white solid. ¹H NMR (300 MHz, DMSO-d₆) δ 9.21 (s, 1H), 8.75 (s, 1H), 7.95 (s, 1H), 7.87 (t, 1H), 7.80 (s, 1H), 7.46 (dd, 1H), 7.37 (dd, 1H), 2.69-2.62 (m, 2H), 2.55-2.47 (m, 2H), 2.00 (m, 1H), 1.58 (m, 1H).

[00458] **Compounds 232 and 233** were synthesized following the procedure described in Example 34 from the appropriate carboxylic acids.

EXAMPLE 35: Synthesis of Compound 234

4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyrrolidin-1-yl)ethyl)benzamide




[00459] A mixture of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid (Compound 224, 100 mg, 0.21 mmol), 2-(pyrrolidin-1-yl)ethanamine (32 mg, 0.28 mmol), HATU (106 mg, 0.28 mmol), and DIEA (0.1

mL, 0.63 mmol) in DCM (3 mL) and DMF (1.5 mL) was stirred at room temperature for 18h. Brine and EtOAc were added and the aqueous layer was extracted with EtOAc (4x), the organics were combined, dried over sodium sulfate, and evaporated to dryness. The residue was purified by column chromatography on silica gel eluting with 0 to 10% MeOH/DCM to afford impure desired product that was repurified by reverse phase HPLC (acetonitrile/water:TFA). The fractions containing the desired compound were combined, acetonitrile was removed in vacuo, and the remaining aqueous layer was treated with a saturated solution of sodium bicarbonate. The aqueous layer was extracted with DCM (3x), the organics were combined, dried over sodium sulfate, and evaporated to dryness to afford 88 mg of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyrrolidin-1-yl)ethyl)benzamide as an off-white solid. LCMS $[M + 1]^+$ 561.1.

EXAMPLE 36: Synthesis of Compound 235


N-Benzyl-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide



[00460] A mixture of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid (Compound 224, 68 mg, 0.14 mmol), benzylamine (22 mg, 0.21 mmol), HATU (70 mg, 0.18 mmol), and DIEA (40 μ L, 0.21 mmol) in DMF (3 mL) was stirred at room temperature for 3h. The crude reaction mixture was purified by reverse phase HPLC (acetonitrile/water:TFA) to afford 51 mg of N-benzyl-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide as a white solid. ^1H NMR (300 MHz, DMSO- d_6) δ 9.22 (d, 1H), 9.12 (t, 1H), 8.75 (d, 1H), 7.87 (t, 1H), 7.49 (dd, 1H), 7.40 (dd, 1H), 7.37-7.31 (m, 4H), 7.29-7.24 (m, 1H), 4.51 (d, 1H), 2.69-2.63 (m, 2H), 2.55-2.44 (m, 2H), 2.03-1.94 (m, 1H), 1.62-1.56 (m, 1H). LCMS $[M + 1]^+$ 554.5.

EXAMPLE 37: Synthesis of Compound 236


4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-2-yl)ethyl)benzamide



[00461] A mixture of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid (Compound 224, 70 mg, 0.15 mmol), 2-(pyridin-2-yl)ethanamine (24 mg, 0.20 mmol), HATU (74 mg, 0.20 mmol), and DIEA (80 μ L, 0.45 mmol) in DMF (3 mL) was stirred at room temperature overnight. The crude reaction mixture was purified by reverse phase HPLC (acetonitrile/water, 0.1% TFA). The fractions containing the desired compound were combined, acetonitrile was removed in vacuo, and the remaining aqueous layer was treated with a saturated solution of sodium bicarbonate. The aqueous layer was extracted with DCM (3x), the organics were combined, dried over sodium sulfate, and evaporated to dryness to afford 51 mg of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-2-yl)ethyl)benzamide as a white solid. ^1H NMR (300 MHz, DMSO- d_6) δ 9.22 (d, 1H), 8.75 (d, 1H), 8.66 (t, 1H), 8.52 (d, 1H), 7.82-7.70 (m, 2H), 7.47-7.22 (m, 4H), 3.65 (q, 2H), 3.01 (t, 2H), 2.68-2.62 (m, 2H), 2.54-2.43 (m, 2H), 2.03-1.93 (m, 1H), 1.60-1.57 (m, 1H). LCMS $[M + 1]^+$ 569.5.

EXAMPLE 38: Synthesis of Compound 237

4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(pyrrolidin-1-yl)propyl)benzamide



[00462] A mixture of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid (Compound 224, 550 mg, 1.18 mmol), 3-(pyrrolidin-1-yl)propan-1-amine (303 mg, 2.37 mmol), HATU (673 mg, 1.77 mmol), and DIEA (0.6 mL, 3.54 mmol) in DMF (25 mL) was stirred at room temperature overnight. Brine and EtOAc were added and the aqueous layer was extracted with EtOAc (4x), the organics were combined, dried over sodium sulfate, and evaporated to dryness. The residue was purified by column chromatography on silica gel eluting with 0 to 20% MeOH/DCM to afford 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-


N-(3-(pyrrolidin-1-yl)propyl)benzamide as a pale orange solid. ^1H NMR (300 MHz, DMSO-d_6) δ 9.22 (d, 1H), 8.75 (d, 1H), 8.64 (t, 1H), 7.83 (t, 1H), 7.47 (dd, 1H), 7.38 (dd, 1H), 3.35 (m, 4H), 2.65-2.54 (m, 6H), 2.53-2.43 (m, 2H), 2.03-1.93 (m, 1H), 1.74 (m, 6H), 1.64-1.53 (m, 1H). LCMS $[\text{M} + 1]^+$ 575.1.

5 [00463] The trifluoroacetic acid salt was prepared according to the following procedure: A mixture of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid (Compound 224, 70 mg, 0.15 mmol), 3-(pyrrolidin-1-yl)propan-1-amine (25 mg, 0.20 mmol), HATU (74 mg, 0.20 mmol), and DIEA (80 μL , 0.45 mmol) in DMF (3 mL) was stirred at room temperature overnight. The crude
10 reaction mixture was purified by reverse phase HPLC (acetonitrile/water, 0.1% TFA) to afford 60 mg of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(pyrrolidin-1-yl)propyl)benzamide 2,2,2-trifluoroacetate as pale yellow solid. ^1H NMR (300 MHz, DMSO-d_6) δ 9.47 (s, 1H), 9.22 (d, 1H), 8.75 (d, 1H), 8.68 (t, 1H), 7.86 (t, 1H), 7.49 (dd, 1H), 7.41 (dd, 1H), 3.56 (m, 2H), 3.40-
15 3.34 (m, 2H), 3.23-3.16 (m, 2H), 3.07-3.96 (m, 2H), 2.70-2.64 (m, 2H), 2.54-2.43 (m, 2H), 2.08-1.85 (m, 7H), 1.60-1.55 (m, 1H). LCMS $[\text{M} + 1]^+$ 575.6.

[00464] **Compounds 238 to 311** were synthesized following the procedure described in Example 35 from the appropriate acids and amines (amines were either commercially available or synthesized from literature procedures).

EXAMPLE 39: Synthesis of Compound 312

4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-6,8-dioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide




25 [00465] A mixture of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide (synthesized as described for Example 1 using 5-amino-3-(trifluoromethyl) picolinonitrile (Intermediate 1) and of 4-((1-cyanocyclobutyl)amino)-2-fluoro-N-methylbenzamide (Intermediate 21) as starting materials) (90 mg, 0.19 mmol) and H_2O_2 (30%, 0.72 mL) in MeOH (8 mL) was stirred at room
30 temperature for 5 days. The reaction mixture was diluted with EtOAc (10 mL) and the organic layer was washed with water, brine, and dried over magnesium sulfate. The residue obtained

was purified by reverse phase HPLC (acetonitrile/water:TFA) to obtain 10 mg of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-6,8-dioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide as a white solid. ¹H NMR (500 MHz, CDCl₃) δ 9.28 (d, 1H), 8.54 (d, 1H), 8.23 (t, 1H), 7.33 (m, 1H), 7.27 (m, 1H), 6.93 (m, 1H), 3.07 (d, 3H), 2.71 (m, 2H), 2.57 (m, 2H), 2.28 (m, 1H), 1.82 (m, 1H).

EXAMPLE 40: Synthesis of Compound 313

4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-(3-(3,3-difluoropyrrolidin-1-yl)propyl)-2-fluorobenzamide


Step 1: 3-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamido)propyl methanesulfonate



[00466] A mixture of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-hydroxypropyl)benzamide (Compound 306, 80 mg, 0.15 mmol) and triethylamine (22 μL, 0.16 mmol) in THF (5 mL) was cooled to 0°C.

Methanesulfonyl chloride (23 μL, 0.3 mmol) was added dropwise and the mixture was stirred at room temperature overnight. Brine and EtOAc were added and the aqueous layer was extracted with EtOAc (2x), the organics were combined, dried over sodium sulfate, and evaporated to dryness to afford crude 3-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamido)propyl methanesulfonate that was used as is.

Step 2: 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-(3-(3,3-difluoropyrrolidin-1-yl)propyl)-2-fluorobenzamide



[00467] A mixture of 3-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamido)propyl methanesulfonate (45 mg, 0.075 mmol), 3,3-difluoropyrrolidine hydrochloride (54 mg, 0.37 mmol), and triethylamine (0.14 mL, 1.05 mmol) in THF (2mL) was heated to 70°C overnight. Brine and EtOAc were added to a cooled reaction mixture and the aqueous layer was extracted with EtOAc (2x), the organics were

combined, dried over sodium sulfate, and evaporated to dryness. The residue obtained was purified by reverse phase HPLC (acetonitrile/water:TFA) to obtain 4.2 mg of 4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-(3-(3,3-difluoropyrrolidin-1-yl)propyl)-2-fluorobenzamide. ¹H NMR (300 MHz, DMSO-d₆) δ 9.15 (s, 1H), 8.68 (d, 1H), 8.51 (t, 1H), 7.72 (t, 1H), 7.39 (dd, 1H), 7.30 (dd, 1H), 3.25 (m, 4H), 2.80 (t, 2H), 2.64-2.58 (m, 4H), 2.44-2.40 (m, 2H), 2.16 (m, 2H), 1.91 (m, 1H), 1.64-1.45 (m, 3H). LCMS [M + 1]⁺ 611.1.

ASSAYS

EXAMPLE 41 - AR In Cell Western Assay

[00468] LNCaP cells (8,000/well) were plated in RPMI media containing 10% Charcoal Dextran Stripped Serum into plates coated with poly-d-lysine. After 24 hours cells were treated with compound from 30μM to 0.0003μM. At 20 hours post compound addition the cells were fixed (30% formaldehyde in PBS) for 20'. Cells were permeabilized in PBS 0.1% Triton (50μl/well, three times for 5' each) and blocked with LiCor blocking buffer (50μl/well, 90'). The wells were then incubated overnight at 4°C with the rabbit IgG androgen receptor antibody (AR-N20, Santa Cruz antibody) diluted 1:1000 in LiCor blocking buffer/0.1% Tween-20. Wells were washed with 0.1% Tween-20/PBS (50μl/well, 5' each) and then incubated in goat anti-rabbit IRDyeTM 800CW (1:1000) and DRAQ5 DNA dye (1:10,000 for 5mM stock) diluted in 0.2%Tween-20/0.01%SDS/LiCor blocking buffer in the dark (90'). Cells were washed (50μl/well, 5' each) in 0.1%Tween-20/PBS. Wash buffer was removed and plates were read using the LiCor Odyssey.

EXAMPLE 42 - AR Fluorescent Polarization Assay (Invitrogen)

[00469] Compounds to be tested were diluted to 2X the final desired concentration in AR Green Assay Buffer (final DMSO: 0.6%). Fluormone AL Green and the rat AR Ligand Binding Domain were diluted to 2X the final desired concentration (Fluormone: 2nM, AR LBD: 50nM) in AR Green Assay Buffer containing 2mM DTT. The AR LBD/Fluormone solution was added to all the wells of a 384 well black plate (10μl/well). The compounds were added to the AR LBD/Fluormone solution (10μl/well). The plate was incubated for 4 hours in the dark. The fluorescence polarization of each well was measured using an excitation wavelength of 485nm and emission wavelength of 530nm.

EXAMPLE 43 - Prostate Cancer Cell Viability Assays

[00470] LNCaP, LNCaP AR, PC3, and VCaP cells were adjusted to a concentration of 30,000 cells per mL in RPMI containing 10% FBS and 20 mM HEPES. 16 microliters of the cell suspension was added to each well of a 384 well plate, and the cells were incubated overnight to allow the cells to adhere. The following day a seven point, serial semilog dilution of each

compound was added to the cells in 16 μ L at a final concentration ranging from 30-0.003 μ M. After 5 days' compound exposure, 16 μ L of CellTiter-Glo (Promega, Madison WI) was added to the cells the the relative luminescence units (RLUs) of each well was determined. CellTiter-Glo added to 32 μ L of medium without cells was used to obtain a background value. The

Percent viability of each sample was determined as follows:

[00471] $(RLU \text{ sample} - RLU \text{ background} / RLU \text{ untreated cells} - RLU \text{ background}) \times 100 = \% \text{viability}$

EXAMPLE 44 - LNCaP-AR Luciferase Transcriptional Reporter Assays

[00472] LNCaP-AR-Luc were maintained in RPMI 1640 supplemented with 10% FCS (Cancer Res 2006; 66: (21). November 1, 2006). Transcriptional assays were performed by seeding 100 μ L of cells at a density of 25,000 cells/mL into 96-well cell culture plates in RPMI 1640 supplemented with 10% charcoal stripped serum and allowed to attach overnight. For AR agonist assays, the compounds were serially diluted and 50 μ L of compound plus RPMI 1640 supplemented with charcoal stripped serum was added to the cells. For AR antagonist assays, the compounds were serially diluted and 50 μ L of compound with RPMI plus R1881 supplemented with charcoal stripped serum were added to the cells. The final R1881 concentration used in the antagonist assays was 0.1 nM. Following 40 hour incubation the medium was removed and the cells were lysed in 40 μ L of lysis buffer (25mM Tris Phosphate, 2mM CDTA, 10% Glycerol, 0.5% Triton X-100, 2 mM DTT). Firefly luciferase activity was measured immediately following the addition of 50 μ L luciferase buffer (20mM tricine, 0.1 mM EDTA, 1.07 mM $(MgCo_3)_4$ $Mg(OH)_2 \cdot 5H_2O$, 2.67 mM $MgSO_4$, 33.3 mM DTT, 270 μ M Coenzyme A, 470 μ M luciferin, 530 μ M ATP).

EXAMPLE 45 – AR-VP16 DNA Binding Assays

[00473] HepG2 cells were maintained in RPMI 1640 supplemented with 10% FCS. AR-VP16 DNA binding assays were performed by seeding 100 μ L of cells at a density of 250,000 cells/mL into 96-well cell culture plates in RPMI 1640 supplemented with 10% charcoal stripped serum and allowed to attach overnight. Cells were transiently transfected using Lipofectin (Life Technologies) according to the manufacturer's protocol. Triplicate transfections were performed using 33.3 ng AR-VP16 pCDNA3 (expression vector), 66.7 ng 4X ARE-Luciferase (reporter vector), 16.7 ng CMVpRL (normalization vector), and 43.3 ng pCMX (filler DNA). Transfected cells were incubated overnight then treated with ligand. For agonist assays, compounds were serially diluted and 50 μ L of compound plus RPMI 1640 supplemented with charcoal stripped serum was added to the cells. For antagonist assays, the compounds were serially diluted and 50 μ L of compound with RPMI supplemented with

charcoal stripped serum plus methyltrienolone (R1881) were added to the cells. The final R1881 concentration used in the antagonist assays was 0.1 nM. Following 48 hour incubation the medium was removed and the cells were lysed in 40 μ L of lysis buffer (25mM Tris Phosphate, 2mM CDTA, 10% Glycerol, 0.5% Triton X-100, 2 mM DTT). Firefly luciferase activity was measured immediately following the addition of 40 μ L luciferase buffer (20mM

5 tricine, 0.1 mM EDTA, 1.07 mM $(\text{MgCo}_3)_4$ $\text{Mg}(\text{OH})_2 \cdot 5\text{H}_2\text{O}$, 2.67 mM MgSO_4 , 33.3 mM DTT, 270 μ M Coenzyme A, 470 μ M luciferin, 530 μ M ATP). Renilla luciferase was measured following the addition of 40 μ L coelenterazine buffer (1.1 M NaCl, 2.2 mM Na_2EDTA , 0.22 M K_2PO_4 (pH 5.1), 0.44 mg/mL BSA, 1.3 mM NaN_3 , 1.43 μ M coelenterazine, final pH adjusted to

10 5.0).

[00474] Illustrative biological data for representative compounds disclosed herein is presented in the following table:

Compound #	ARVP16 DNA Binding Assay (Agonist mode): Agonist? ¹	ARVP16 DNA Binding Assay (Antagonist mode): $\text{IC}_{50} < 1.0 \mu\text{M}$
1	No	Yes
2	No	No
3	No	No
4	No	Yes
5	No	Yes
6	No	Yes
7	No	Yes
8	No	No
9	No	Yes
10	No	No
11	No	Yes
12	No	Yes
13	No	Yes
14	No	No
15	No	No
16	No	Yes
17	Yes	Yes
18	Yes	Yes
19	No	No
20	No	Yes
21	No	Yes
22	No	Yes
23	No	No
24	No	No
25	No	Yes
26	Yes	Yes
27	No	Yes
28	No	Yes
29	No	Yes
30	No	Yes

Compound #	ARVP16 DNA Binding Assay (Agonist mode): Agonist? ¹	ARVP16 DNA Binding Assay (Antagonist mode): IC ₅₀ < 1.0 μ M
31	No	Yes
32	No	Yes
33	No	Yes
34	No	No
35	No	No
36	No	Yes
37	No	Yes
38	No	No
39	Yes	Yes
40	No	Yes
41	No	Yes
42	No	Yes
43	No	Yes
44	No	Yes
45	No	Yes
46	No	Yes
47	No	Yes
48	No	Yes
49	No	Yes
50	No	Yes
51	No	Yes
52	No	Yes
53	Yes	Yes
54	No	No
55	Yes	Yes
56	No	Yes
57	No	Yes
58	Yes	Yes
59	No	Yes
60	Yes	Yes
61	No	No
62	No	No
63	No	Yes
64	No	Yes
65	Yes	Yes
66	Yes	Yes
67	Yes	Yes
68	No	No
69	No	Yes
70	No	Yes
71	No	No
72	Yes	Yes
73	Yes	Yes
74	Yes	Yes
75	Yes	Yes
76	Yes	Yes
77	No	No
78	No	No

Compound #	ARVP16 DNA Binding Assay (Agonist mode): Agonist? ¹	ARVP16 DNA Binding Assay (Antagonist mode): IC ₅₀ < 1.0 μ M
79	No	Yes
80	No	Yes
81	No	No
82	No	Yes
83	No	No
84	Yes	Yes
85	Yes	Yes
86	No	Yes
87	No	Yes
88	No	Yes
89	No	Yes
90	No	No
91	No	No
92	No	Yes
93	No	No
94	No	Yes
95	Yes	Yes
96	No	Yes
97	Yes	Yes
98	No	Yes
99	Yes	Yes
100	No	Yes
101	No	No
102	No	No
103	No	Yes
104	No	Yes
105	No	Yes
106	No	Yes
107	No	Yes
108	Yes	Yes
109	No	Yes
110	No	Yes
111	No	No
112	No	Yes
113	No	Yes
114	No	No
115	No	Yes
116	No	Yes
117	No	No
118	No	No
119	No	Yes
120	No	No
121	No	No
122	No	No
123	No	No
124	No	Yes
125	No	Yes
126	Yes	Yes

Compound #	ARVP16 DNA Binding Assay (Agonist mode): Agonist? ¹	ARVP16 DNA Binding Assay (Antagonist mode): IC ₅₀ < 1.0 µM
127	No	Yes
128	No	Yes
129	No	Yes
130	No	Yes
131	No	Yes
132	No	No
133	No	Yes
134	No	Yes
135	No	Yes
136	No	Yes
137	No	Yes
138	No	No
139	No	Yes
140	No	No
141	Yes	No
142	Yes	Yes
143	Yes	Yes
144	Yes	Yes
145	Yes	Yes
146	No	Yes
147	No	Yes
148	Yes	Yes
149	No	Yes
150	Yes	Yes
151	No	No
152	No	Yes
153	No	Yes
154	No	Yes
155	No	No
156	Yes	Yes
157	Yes	Yes
158	No	No
159	No	No
160	No	No
161	No	No
162	No	No
163	No	No
164	No	No
165	No	No
166	No	No
167	No	No
168	No	Yes
169	No	Yes
170	Yes	Yes
171	No	Yes
172	No	Yes
173	No	Yes
174	No	Yes

Compound #	ARVP16 DNA Binding Assay (Agonist mode): Agonist? ¹	ARVP16 DNA Binding Assay (Antagonist mode): IC ₅₀ < 1.0 μ M
175	No	Yes
176	No	Yes
177	No	Yes
178	No	Yes
179	Yes	Yes
180	Yes	Yes
181	Yes	Yes
182	Yes	Yes
183	No	Yes
184	Yes	Yes
185	No	Yes
186	Yes	Yes
187	No	Yes
188	No	Yes
189	No	Yes
190	No	Yes
191	No	No
192	No	No
193	No	Yes
194	No	Yes
195	No	Yes
196	No	Yes
197	No	No
198	No	Yes
199	No	Yes
200	Yes	Yes
201	No	Yes
202	Yes	Yes
203	No	Yes
204	No	Yes
205	No	Yes
206	No	Yes
207	No	Yes
208	No	Yes
209	No	Yes
210	No	No
211	No	Yes
212	Yes	Yes
213	Yes	Yes
214	Yes	Yes
215	Yes	Yes
216	No	No
217	No	Yes
218	No	Yes
219	Yes	Yes
220	No	Yes
221	No	No
222	Yes	Yes

Compound #	ARVP16 DNA Binding Assay (Agonist mode): Agonist? ¹	ARVP16 DNA Binding Assay (Antagonist mode): IC ₅₀ < 1.0 μ M
223	No	Yes
224	No	No
225	No	Yes
226	No	No
227	No	No
228	No	No
229	No	Yes
230	No	Yes
231	No	Yes
232	No	Yes
233	No	No
234	No	No
235	No	Yes
236	No	Yes
237	No	Yes
238	No	No
239	No	No
240	No	No
241	No	No
242	No	Yes
243	No	No
244	No	Yes
245	No	No
246	No	No
247	No	No
248	No	No
249	No	Yes
250	No	No
251	No	No
252	No	Yes
253	No	No
254	No	No
255	No	Yes
256	No	Yes
257	No	Yes
258	No	Yes
259	No	Yes
260	No	Yes
261	No	Yes
262	No	Yes
263	Yes	Yes
264	No	Yes
265	No	Yes
266	No	Yes
267	No	Yes
268	No	Yes
269	No	Yes
270	No	No

Compound #	ARVP16 DNA Binding Assay (Agonist mode): Agonist? ¹	ARVP16 DNA Binding Assay (Antagonist mode): IC ₅₀ < 1.0 μ M
271	No	No
272	No	Yes
273	No	Yes
274	No	Yes
275	No	No
276	No	No
277	No	Yes
278	No	Yes
279	No	Yes
280	No	Yes
281	No	Yes
282	No	Yes
283	No	Yes
284	No	Yes
285	No	No
286	No	Yes
287	No	Yes
288	No	Yes
289	No	Yes
290	No	Yes
291	No	Yes
292	No	No
293	No	Yes
294	No	Yes
295	No	Yes
296	No	Yes
297	No	Yes
298	No	Yes
299	No	Yes
300	No	Yes
301	No	Yes
302	No	Yes
303	No	No
304	No	Yes
305	No	Yes
306	No	Yes
307	No	Yes
308	No	Yes
309	No	Yes
310	No	No
311	No	No
312	No	No
313	No	Yes

¹: Agonist defined as a compound whose Emax in the ARVP16 DNA Binding Assay (Agonist mode) is >20x DMSO control

EXAMPLE 46 – GABA-gated Cl Channel Antagonist Radioligand Binding Assay

[00475] Membrane homogenates of cerebral cortex (120 µg protein) are incubated for 120 min at 22°C with 3 nM [³⁵S]-TBPS in the absence or presence of the test compound in a buffer containing 50 mM Na₂HPO₄/KH₂PO₄ (pH 7.4) and 500 mM NaCl. Nonspecific binding is determined in the presence of 20 µM picrotoxinin. Following incubation, the samples are filtered rapidly under vacuum through glass fiber filters (GF/B, Packard) presoaked with 0.3% PEI and rinsed several times with ice-cold 50 mM Tris-HCl using a 96-sample cell harvester (Unifilter, Packard). The filters are dried then counted for radioactivity in a scintillation counter (Topcount, Packard) using a scintillation cocktail (Microscint 0, Packard). The results are expressed as a percent inhibition of the control radioligand specific binding. The standard reference compound is picrotoxinin, which is tested in each experiment at several concentrations to obtain a competition curve from which its IC₅₀ is calculated.

[00476] In this assay, the following representative compounds disclosed herein demonstrated an inhibition of less than 65% at 10 µM in the GABA-gated Cl⁻ Channel Binding Assay:

Compound 5, Compound 13, Compound 124, Compound 168, Compound 171, Compound 208, Compound 218, Compound 237, Compound 268, Compound 291.

***In vivo* Assay(s)**

Example 47: Castrate Resistant Prostate Cancer Xenograft Studies

[00477] Six to Seven week old male SCID Hairless Outbred mice (SHO, Charles Rivers Laboratories) underwent bilateral orchiectomy under isoflurane anesthesia. LNCaP/AR cells were grown in RPMI at 5% CO₂, 37°C. Cells were spun down and re-suspended in 50% serum-free RPMI and 50% Matrigel at 1X 10⁷ cells/ml. LNCaP/AR cells were subcutaneously injected (100µl/animal) on the right flank 3-5 days post castration. Tumor volume (length x width²/2) was monitored weekly. When tumors reached an average volume of ~200mm³ animals were randomized into treatment groups. During the treatment period tumor volume was monitored bi-weekly. At the termination of study tumors were collected and stored for further analyses.

Pharmaceutical Compositions

Example 48: Parenteral Composition

[00478] To prepare a parenteral pharmaceutical composition suitable for administration by injection (subcutaneous, intravenous, and the like), 100 mg of a water-soluble salt of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is dissolved in sterile water and then mixed with 10 mL of 0.9% sterile saline. The mixture is incorporated into a dosage unit form suitable for administration by injection

[00479] In another embodiment, the following ingredients are mixed to form an injectable formulation: 1.2 g of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X), 2.0 mL of sodium acetate buffer solution (0.4 M), HCl (1 N) or

NaOH (1 M) (q.s. to suitable pH), water (distilled, sterile) (q.s.to 20 mL). All of the above ingredients, except water, are combined and stirred and if necessary, with slight heating if necessary. A sufficient quantity of water is then added.

Example 49: Oral Solution

5 [00480] To prepare a pharmaceutical composition for oral delivery, an aqueous 20% propylene glycol solution is prepared. To this is added a sufficient amount of compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) to provide a 20 mg/mL solution of the compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X).

10 **Example 50: Oral Capsule**

[00481] To prepare a pharmaceutical composition for oral delivery, 100 mg of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is mixed with 750 mg of starch. The mixture is incorporated into an oral dosage unit such as a hard gelatin capsule, which is suitable for oral administration.

15 [00482] In another embodiment, 100 mg of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is placed into Size 4 capsule, or size 1 capsule (hypromellose or hard gelatin) and the capsule is closed.

Example 51: Oral Tablet

20 [00483] A tablet is prepared by mixing 48% by weigh of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X), 45% by weight of microcrystalline cellulose, 5% by weight of low-substituted hydroxypropyl cellulose, and 2% by weight of magnesium stearate. Tablets are prepared by direct compression. The total weight of the compressed tablets is maintained at 250 mg.


Example 52: Topical Gel Composition

25 [00484] To prepare a pharmaceutical topical gel composition, 100 mg of a compound of Formula (I), (Ia), (II), (III), (IV), (V), (VI), (VII), (VIII), (VIIIa), (IX), (IXa) or (X) is mixed with 1.75 g of hydroxypropyl cellulose, 10 mL of propylene glycol, 10 mL of isopropyl myristate and 100 mL of purified alcohol USP. The resulting gel mixture is then incorporated into containers, such as tubes, which are suitable for topical administration.

30 [00485] The examples and embodiments described herein are for illustrative purposes only and various modifications or changes suggested to persons skilled in the art are to be included within the spirit and purview of this application and scope of the appended claims.

WHAT IS CLAIMED IS:

1. A compound of Formula (VIIIa), or a pharmaceutically acceptable salt, or *N*-oxide thereof:



Formula (VIIIa)

wherein,

both R^1 are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C_3 - C_6 cycloalkyl or a substituted or unsubstituted C_2 - C_6 heterocycloalkyl;

or

each R^1 is independently H, -OH, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, or C_1 - C_6 fluoroalkyl;

X is O or S;

R^A is C_1 - C_6 alkyl;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, or 2;

R^4 is H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, C_1 - C_6 alkoxy, or C_1 - C_6 heteroalkyl;

R^5 is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, C_1 - C_{10} alkyl, C_1 - C_{10} fluoroalkyl, C_1 - C_{10} fluoroalkoxy, C_1 - C_{10} alkoxy, C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl,

substituted or unsubstituted monocyclic heteroaryl, or -L¹-L²-R⁶;

L¹ is absent, -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-, or -S(=O)₂NH-;

L² is C_1 - C_6 alkylene, C_1 - C_6 fluoroalkylene or C_1 - C_6 heteroalkylene;

R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -

$C(=O)N(R^9)_2$, $-OC(=O)N(R^9)_2$, $-NR^{11}C(=O)N(R^9)_2$, $-NR^{11}C(=O)R^{10}$, $-NR^{11}C(=O)OR^{10}$, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl;

each R^9 is independently selected from H, C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), and $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl); or two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl;

R^{10} is C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 fluoroalkyl, a substituted or unsubstituted C_3 - C_{10} cycloalkyl, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted benzyl, a substituted or unsubstituted heteroaryl, $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_3 - C_{10} cycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted C_2 - C_{10} heterocycloalkyl), $-C_1$ - C_4 alkylene-(substituted or unsubstituted aryl), or $-C_1$ - C_4 alkylene-(substituted or unsubstituted heteroaryl);

R^{11} is H or C_1 - C_4 alkyl.

2. The compound of claim 1, wherein:

X is S;

ring B is phenyl;

R^4 is H, halogen, -CN, -OH, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, or C_1 - C_6 alkoxy;

R^5 is halogen, -CN, $-NO_2$, -OH, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-CO_2R^9$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, C_1 - C_{10} alkyl, C_1 - C_{10} fluoroalkyl, C_1 - C_{10} fluoroalkoxy, C_1 - C_{10} alkoxy, C_1 - C_{10} heteroalkyl, substituted or unsubstituted C_3 - C_{10} cycloalkyl, substituted or unsubstituted C_2 - C_{10} heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or $-L^1-L^2-R^6$;

L^1 is absent, -O-, -S-, -S(O)-, $-S(O)_2$ -, -NH-, $-C(=O)$ -, $-C(=O)NH$ -, or $-S(=O)_2NH$ -;

L^2 is C_1 - C_6 alkylene, C_1 - C_6 fluoroalkylene or C_1 - C_6 heteroalkylene;

R^6 is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, or substituted or unsubstituted aryl.

3. The compound of claim 2, wherein:


R^A is C₁-C₆alkyl;

both R^1 are taken together with the carbon atom to which they are attached to form a C₃-C₆cycloalkyl;

or

each R^1 is independently C₁-C₄alkyl.

4. The compound of claim 3, wherein the compound of Formula (VIIIa) has the structure of Formula (IXa):



Formula (IXa).

5. The compound of any one of claims 1-4, wherein:

each R^9 is independently selected from H, C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₆cycloalkyl, a substituted or unsubstituted C₂-C₆heterocycloalkyl, a substituted or unsubstituted phenyl, a substituted or unsubstituted benzyl, and a substituted or unsubstituted monocyclic heteroaryl; or

two R^9 groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted C₂-C₆heterocycloalkyl;

R^{10} is C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₆cycloalkyl, a substituted or unsubstituted C₂-C₆heterocycloalkyl, a substituted or unsubstituted phenyl, or a substituted or unsubstituted monocyclic heteroaryl.

6. The compound of claim 5, wherein:

R^5 is halogen, -CN, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -

C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)NH(R⁹), C₁-C₁₀alkyl, C₁-C₁₀fluoroalkyl, C₁-

C₁₀fluoroalkoxy, C₁-C₁₀alkoxy, C₁-C₁₀heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or

unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or $-L^1-L^2-R^6$;

L^1 is absent, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$;

L^2 is C_1-C_6 alkylene;

R^6 is $-CN$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-CO_2R^9$, $-C(=O)N(R^9)_2$, substituted or unsubstituted C_3-C_6 cycloalkyl, substituted or unsubstituted C_2-C_6 heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

7. The compound of claim 6, wherein:

R^A is $-CH_3$;

both R^1 are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl;

or

each R^1 is $-CH_3$;

R^5 is halogen, $-CN$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-CO_2R^9$, $-N(R^9)_2$, $-C(=O)NH(R^9)$, C_1-C_{10} alkyl, C_1-C_{10} fluoroalkyl, C_1-C_{10} fluoroalkoxy, C_1-C_{10} alkoxy, C_1-C_{10} heteroalkyl, substituted or unsubstituted C_3-C_6 cycloalkyl, substituted or unsubstituted C_2-C_6 heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, or $-L^1-L^2-R^6$;

L^1 is absent, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$;

L^2 is C_1-C_6 alkylene;

R^6 is substituted or unsubstituted C_2-C_6 heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

8. The compound of claim 1, wherein the compound is:

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-methylbenzamide;

4-(3-(6-Cyano-5-methylpyridin-3-yl)-4-oxo-2-thioxo-1,3-diazaspiro[4.4]nonan-1-yl)-2-fluoro-N-methylbenzamide;

3-Methyl-5-(5-(4-(5-methylfuran-2-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

3-Methyl-5-(8-oxo-6-thioxo-5-(p-tolyl)-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

4-(3-(6-Cyano-5-methylpyridin-3-yl)-5,5-dimethyl-4-oxo-2-thioxoimidazolidin-1-yl)-2-fluoro-N-methylbenzamide;

5-(5-(3-Fluoro-4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

Methyl 4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-3-fluorobenzoate;

5 5-(5-(4-Hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile; 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-methylbenzamide;

3-Methyl-5-(5-(4-(6-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

10 3-Methyl-5-(5-(4-(4-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

3-Methyl-5-(5-(4-(5-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

15 3-Methyl-5-(5-(4-(2-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

5-(3-(4-Hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-3-methylpicolinonitrile;

3-Methyl-5-(5-(4-(methyl(1-methylpiperidin-4-yl)amino)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

20 3-Methyl-5-(8-oxo-5-(4-(pyrimidin-5-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-3-fluoro-N-methylbenzamide;

25 5-(5-(4-(5-Fluoropyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N,2-dimethylbenzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-methoxy-N-methylbenzamide;

30 2-Chloro-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-methylbenzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-methyl-2-(trifluoromethyl)benzamide;

35 3-Methyl-5-(8-oxo-5-(1-oxoisindolin-5-yl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

3-Methyl-5-(8-oxo-5-(4-(tetrahydro-2H-pyran-4-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

5-(5-(4-(Furan-2-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5 5-(5-(3-Fluoro-4-(2-methoxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5-(5-(3-Fluoro-4-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

10 5-(5-(3-Fluoro-4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

3-Methyl-5-(5-(4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

5-(5-(3-Fluoro-4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

15 3-Methyl-5-(8-oxo-5-(4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

3-Methyl-5-(8-oxo-5-(4-((tetrahydro-2H-thiopyran-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

20 5-(4,4-Dimethyl-3-(4-((1-methylpiperidin-4-yl)oxy)phenyl)-5-oxo-2-thioxoimidazolidin-1-yl)-3-methylpicolinonitrile;

3-Methyl-5-(8-oxo-6-thioxo-5-(4-(2-(4-(2,2,2-trifluoroethyl)piperazin-1-yl)ethoxy)phenyl)-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

3-Methyl-5-(5-(4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

25 5-(5-(4-((1,1-Dioxidotetrahydro-2H-thiopyran-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5-(5-(4-Methoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

3-Methyl-5-(8-oxo-5-(4-(pyrimidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

30 3-Methyl-5-(8-oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

5-(5-(4-((1-acetylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5-(5-(4-((1-Ethylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5-(5-(4-((1-(2-Hydroxyethyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5 5-(5-(4-(2-Hydroxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzoic acid;

10 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-methylbenzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-phenylbenzamide;

15 N-Benzyl-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(thiophen-2-ylmethyl)benzamide;

20 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(pyrrolidin-1-yl)propyl)benzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(4-fluorobenzyl)benzamide;

N-(2-Chlorophenyl)-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;

25 N-(3-Chlorophenyl)-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;

N-(4-Chlorophenyl)-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;

30 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-fluorophenyl)benzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-fluorophenyl)benzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(4-fluorophenyl)benzamide;

4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(oxazol-2-yl)benzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(thiazol-2-yl)benzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(1-methyl-1H-pyrazol-5-yl)benzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(6-methylpyridin-3-yl)benzamide;


4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(5-fluoropyridin-3-yl)benzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(tetrahydro-2H-pyran-4-yl)benzamide;

4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-((tetrahydro-2H-pyran-4-yl)methyl)benzamide;

or a pharmaceutically acceptable salt thereof or *N*-oxide thereof.

9. A compound of Formula (Ia), or a pharmaceutically acceptable salt, or *N*-oxide thereof:



Formula (Ia)

wherein,

ring A is monocyclic heteroaryl, bicyclic heteroaryl, or naphthyl;

m is 1, 2, 3 or 4;

each R^A is independently H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, C₁-C₆alkoxy, or C₁-C₆heteroalkyl;

both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₆cycloalkyl or a substituted or unsubstituted C₂-C₆heterocycloalkyl;

or each R¹ is independently selected from H, -OH, C₁-C₆alkyl, C₁-C₆alkoxy, and C₁-C₆fluoroalkyl;

X is S or O;

ring B is phenyl, naphthyl, monocyclic heteroaryl, or bicyclic heteroaryl;

n is 0, 1, 2, 3 or 4;

each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -

5 S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, C₁-C₆alkoxy, and C₁-C₆heteroalkyl;

R⁵ is -L¹-L²-R⁶ or -L¹-R⁷;

10 L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-, or -S(=O)₂NH-;

L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene;

15 R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR¹¹C(=O)N(R⁹)₂, -NR¹¹C(=O)R¹⁰, -NR¹¹C(=O)OR¹⁰, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆heteroalkyl, substituted or unsubstituted C₃-C₁₀cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, substituted or unsubstituted bicyclic heteroaryl, substituted or unsubstituted phenyl, or substituted or unsubstituted naphthyl;

20 R⁷ is substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl;

25 each R⁹ is independently selected from H, C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₁₀heterocycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted monocyclic heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₆cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted phenyl), and -C₁-C₄alkylene-(substituted or unsubstituted monocyclic heteroaryl); or

30 two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl;

35 R¹⁰ is C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted C₃-C₆cycloalkyl, a substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, a

substituted or unsubstituted phenyl, a substituted or unsubstituted benzyl, a substituted or unsubstituted monocyclic heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted C₃-C₆cycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted phenyl), or -C₁-C₄alkylene-(substituted or unsubstituted monocyclic heteroaryl);

R¹¹ is H or C₁-C₄alkyl.

10. The compound of claim 9, wherein:

both R¹ are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted C₃-C₆cycloalkyl;

X is S;

ring A is *N*-containing monocyclic heteroaryl;

each R^A is independently selected from H, halogen, -CN, -NO₂, -OH, -S(=O)₂R¹⁰, -N(R¹¹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, and C₁-C₆alkoxy;

ring B is phenyl or monocyclic heteroaryl.

11. The compound of claim 10, wherein:

both R¹ are taken together with the carbon atom to which they are attached to form a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl;

ring A is pyridinyl, pyrimidinyl, pyridazinyl, or pyrazinyl.

12. The compound of claim 11, wherein:

ring A is pyridinyl;

each R^A is independently selected from H, halogen, -CN, -OH, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)N(R⁹)₂, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₁-C₆fluoroalkoxy, and C₁-C₆alkoxy.

13. The compound of any one of claims 9-12, wherein:

ring B is phenyl;

R⁵ is -L¹-L²-R⁶;

L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHS(=O)₂-, or -S(=O)₂NH-;

L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene;

R⁶ is halogen, -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -CO₂R⁹, -N(R⁹)₂, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

14. The compound of claim 13, wherein:

ring B is phenyl;

each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, and C₁-C₄alkoxy;

R⁵ is -L¹-L²-R⁶;

L¹ is absent, -O-, or -C(=O)NH-;

L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene;

R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

15. The compound of claim 14, wherein:

L² is C₁-C₆alkylene;

R⁶ is -CN, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

16. The compound of claim 15, wherein:

R⁶ is substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl.

17. The compound of claim 16, wherein:

L² is C₁-C₆alkylene;

R⁶ is a substituted or unsubstituted C₂-C₆heterocycloalkyl.

18. The compound of any one of claims 9-12, wherein:

ring B is phenyl;

each R⁴ is independently selected from H, halogen, -CN, -NO₂, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, and C₁-C₄alkoxy;

R⁵ is -L¹-R⁷;

L¹ is absent, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -C(=O)-, -C(=O)NH-, -NHC(=O)-, -NHC(=O)O-, -NHC(=O)NH-, -C(=O)O-, -OC(=O)-, -OC(=O)O-, -OC(=O)NH-, -NHS(=O)₂-, or -S(=O)₂NH-;

R⁷ is substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl;

19. The compound of claim 18, wherein:

R⁵ is -L¹-R⁷;

L^1 is $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$;

R^7 is substituted or unsubstituted monocyclic C_2 - C_6 heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.


20. The compound of claim 18, wherein:

5 R^5 is $-L^1-R^7$;

L^1 is absent, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(=O)-$, $-C(=O)NH-$, or $-S(=O)_2NH-$;


R^7 is substituted or unsubstituted monocyclic heteroaryl.

21. The compound of claim 9, wherein the compound of Formula (I) has the structure of Formula (II):



Formula (II)

wherein,



15 m is 2;

one R^A is $-CN$, $-NO_2$, $-S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-CO_2R^9$, or $-$

$C(=O)N(R^9)_2$; and the other R^A is H , halogen, $-OH$, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, or C_1 - C_6 alkoxy;

n is 0 or 1;

20 each R^4 is independently selected from H , halogen, $-CN$, $-NO_2$, $-OH$, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, and C_1 - C_6 alkoxy;

R^5 is $-L^1-L^2-R^6$ or $-L^1-R^7$;


L^1 is absent, $-O-$, or $-C(=O)NH-$;

L^2 is C_1 - C_6 alkylene, C_1 - C_6 fluoroalkylene or C_1 - C_6 heteroalkylene;

25 R^6 is $-CN$, $-NO_2$, $-OH$, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-C(=O)R^{10}$, $-CO_2R^9$, $-N(R^9)_2$, $-C(=O)N(R^9)_2$, substituted or unsubstituted C_3 - C_6 cycloalkyl, substituted or unsubstituted C_2 - C_6 heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl;

R^7 is substituted or unsubstituted monocyclic C_2 - C_6 heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.

22. The compound of claim 21, wherein:



one R^A is -CN and the other R^A is H, halogen, -OH, C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, C_1 - C_4 fluoroalkoxy, or C_1 - C_4 alkoxy;

each R^4 is independently selected from H, halogen, -OH, C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, C_1 - C_4 fluoroalkoxy, and C_1 - C_4 alkoxy;

R^5 is $-L^1-L^2-R^6$ or $-L^1-R^7$;


L^1 is absent, -O-, or $-C(=O)NH-$;

L^2 is C_1 - C_6 alkylene, C_1 - C_6 fluoroalkylene or C_1 - C_6 heteroalkylene;

R^6 is -CN, $-NO_2$, -OH, $-OR^9$, $-SR^9$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-C(=O)R^{10}$, $-CO_2R^9$, $-C(=O)N(R^9)_2$, substituted or unsubstituted C_3 - C_6 cycloalkyl, substituted or unsubstituted C_2 - C_6 heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl;

R^7 is substituted or unsubstituted monocyclic C_2 - C_6 heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.

23. The compound of claim 9, wherein the compound of Formula (Ia) has the structure of Formula (VI):



Formula (VI)

wherein,

one R^A is -CN, $-NO_2$, $-S(=O)_2R^{10}$, $-S(=O)_2N(R^9)_2$, $-C(=O)R^{10}$, $-CO_2R^9$, or -

$C(=O)N(R^9)_2$; and the other R^A is H, halogen, -OH, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_1 - C_6 fluoroalkoxy, or C_1 - C_6 alkoxy;

both R^1 are taken together with the carbon atom to which they are attached to form a C_3 - C_6 cycloalkyl;

or each R^1 is independently C_1 - C_4 alkyl;

R⁴ is H, halogen, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, or C₁-C₄alkoxy.

24. The compound of claim 23, wherein:

R⁵ is -L¹-L²-R⁶ or -L¹-R⁷;

L¹ is absent, -O-, or -C(=O)NH-;

L² is C₁-C₆alkylene, C₁-C₆fluoroalkylene or C₁-C₆heteroalkylene;

R⁶ is -CN, -NO₂, -OH, -OR⁹, -SR⁹, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -C(=O)R¹⁰, -CO₂R⁹, -C(=O)N(R⁹)₂, substituted or unsubstituted C₃-C₆cycloalkyl, substituted or unsubstituted C₂-C₆heterocycloalkyl, substituted or unsubstituted monocyclic heteroaryl, or substituted or unsubstituted phenyl;

R⁷ is substituted or unsubstituted monocyclic C₂-C₆heterocycloalkyl, or substituted or unsubstituted monocyclic heteroaryl.

25. The compound of claim 9, wherein the compound is:

3-Methyl-5-(5-(4-(5-methylfuran-2-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

5-(8-Oxo-5-(4-(pyridin-2-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(8-Oxo-5-(4-(pyridin-3-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(8-Oxo-5-(4-(pyridin-4-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(8-Oxo-5-(4-(pyridin-3-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(8-Oxo-5-(4-(pyridin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(8-Oxo-5-(4-(pyridin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(8-Oxo-5-(4-(pyrimidin-5-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(5-(3-(Hydroxymethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(5-(2-(Hydroxymethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(5-(4-(Hydroxymethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

- 5-(5-(4-(3-Hydroxypropyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-(4-Methylpiperazin-1-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5 (5-(8-Oxo-5-(4-((2-(pyridin-4-yl)ethyl)sulfonyl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-((Methyl(pyridin-4-ylmethyl)amino)methyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-((4-Methylpiperazin-1-yl)methyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 10 3-Methyl-5-(5-(4-(6-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;
- 3-Methyl-5-(5-(4-(4-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;
- 15 3-Methyl-5-(5-(4-(5-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;
- 3-Methyl-5-(5-(4-(2-methylpyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;
- 3-Methyl-5-(5-(4-(methyl(1-methylpiperidin-4-yl)amino)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;
- 20 3-Methyl-5-(8-oxo-5-(4-(pyrimidin-5-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile; 5-(5-(4-(5-Fluoropyridin-3-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;
- 3-Methyl-5-(8-oxo-5-(4-(tetrahydro-2H-pyran-4-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;
- 25 5-(8-Oxo-5-(4-(tetrahydro-2H-pyran-4-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-(Furan-2-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;
- 30 5-(8-Oxo-5-(4-(piperazin-1-yl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- tert-Butyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)piperazine-1-carboxylate;
- Methyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)butanoate;
- 35

- 5-(5-(4-(3-(4-Methylpiperazin-1-yl)propyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-(1-methyl-1H-pyrazol-5-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5 5-(5-(3-Fluoro-4-(2-(pyrrolidin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-(Benzyloxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-((1-Methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-10 7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-methoxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 15 Ethyl 2-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorophenoxy)acetate;
- 5-(5-(3-Fluoro-4-(4-methoxybutoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-((4,4,5,5,5-pentafluoropentyl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 20 5-(5-(4-(3-(Benzyloxy)propoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-(4-(Benzyloxy)butoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 25 5-(5-(3-Fluoro-4-(2-methylphenethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(3-phenylpropoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(3,3,4,4,4-pentafluorobutoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 30 5-(5-(3-Fluoro-4-(2-(naphthalen-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-(naphthalen-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

- 5-(5-(3-Fluoro-4-(2-(pyridin-4-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-morpholinoethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5 5-(5-(3-Fluoro-4-(2-(pyridin-3-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(3-(pyridin-3-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-(2-(1H-Pyrrol-1-yl)ethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 10 5-(5-(3-Fluoro-4-(pyridin-2-ylmethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(pyridin-3-ylmethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 15 5-(5-(3-Fluoro-4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(3-morpholinopropoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(4-((4,4,5,5,5-pentafluoropentyl)thio)butoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 20 5-(5-(3-Fluoro-4-(4-((4,4,5,5,5-penta fluoropentyl)sulfinyl)butoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(3-((4,4,5,5,5-pentafluoropentyl)thio)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 25 5-(5-(3-Fluoro-4-(3-((4,4,5,5,5-pentafluoropentyl)sulfinyl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(4-(2-(Pyridin-2-yl)ethoxy)phenyl)-6-thioxo-7-(3-(trifluoromethyl)-[1,2,4]triazolo[4,3-b]pyridazin-6-yl)-5,7-diazaspiro[3.4]octan-8-one;
- 5-(5-(3-Fluoro-4-(pyridin-4-ylmethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 30 5-(5-(3-Fluoro-4-(3-(4-methylpiperazin-1-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-Fluoro-3-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

- 5-(5-(4-Fluoro-3-(2-(pyridin-3-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-Fluoro-3-(2-(pyridin-4-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5 5-(5-(3-(Benzyloxy)-4-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-Fluoro-3-phenethoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-Fluoro-3-(3-phenylpropoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 10 5-(5-(4-Fluoro-3-(2-morpholinoethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-Fluoro-3-(3-morpholinopropoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 15 5-(5-(4-Fluoro-3-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-Fluoro-3-(3-(4-methylpiperazin-1-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-methoxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;
- 20 5-(5-(3-Fluoro-4-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;
- 5-(5-(3-Fluoro-4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;
- 25 5-(5-(3-Fluoro-4-(2-methoxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)quinoline-8-carbonitrile;
- 5-(5-(3-Fluoro-4-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)quinoline-8-carbonitrile;
- 5-(5-(3-Fluoro-4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)quinoline-8-carbonitrile;
- 30 5-(5-(3-Fluoro-4-(2-(piperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-thiomorpholinoethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

- 5-(5-(3-Fluoro-4-(2-(pyrazin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-(piperidin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5 5-(5-(4-(2-(4-Methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(4-(2-Cyclohexylethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-(tetrahydro-2H-pyran-4-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 10 5-(5-(2-Fluoro-4-(2-methoxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile; 5-(5-(2-Fluoro-4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 15 5-(5-(2-Fluoro-4-(2-(pyridin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(2-Fluoro-4-(3-(4-methylpiperazin-1-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(3-(pyrrolidin-1-yl)propoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 20 5-(5-(4-(2-(1,1-Dioxidothiomorpholino)ethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(2-Fluoro-4-(2-(piperidin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 25 5-(5-(3-Fluoro-4-phenethoxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(3-Fluoro-4-(2-(pyrimidin-2-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 5-(5-(2-Fluoro-4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
- 30 3-Methyl-5-(5-(4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;
- 5-(5-(3-Fluoro-4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5-(5-(3-Fluoro-4-((1-methylpiperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

3-Methyl-5-(8-oxo-5-(4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

5 3-Methyl-5-(8-oxo-5-(4-((tetrahydro-2H-thiopyran-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

5-(4,4-Dimethyl-3-(4-((1-methylpiperidin-4-yl)oxy)phenyl)-5-oxo-2-thioxoimidazolidin-1-yl)-3-(trifluoromethyl)picolinonitrile;

5-(4,4-Dimethyl-3-(4-((1-methylpiperidin-4-yl)oxy)phenyl)-5-oxo-2-thioxoimidazolidin-1-yl)-3-methylpicolinonitrile;

10 5-(8-Oxo-5-(4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

3-Methyl-5-(8-oxo-6-thioxo-5-(4-(2-(4-(2,2,2-trifluoroethyl)piperazin-1-yl)ethoxy)phenyl)-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

15 3-Methyl-5-(5-(4-(2-(4-methylpiperazin-1-yl)ethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

5-(5-(4-((1,1-Dioxidotetrahydro-2H-thiopyran-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5-(5-(4-(2-(4-Acetylpiperazin-1-yl)ethoxy)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

20 5-(8-Oxo-5-(4-(pyrimidin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(8-Oxo-5-(4-(pyrazin-2-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

25 5-(8-Oxo-5-(4-(pyrimidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

3-Methyl-5-(8-oxo-5-(4-(pyrimidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

3-Methyl-5-(8-oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile;

30 5-(8-Oxo-5-(4-(piperidin-4-yloxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(8-Oxo-5-(4-((1-propionylpiperidin-4-yl)oxy)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(5-(4-((1-isobutyryl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

ethyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidine-1-carboxylate;

5-(5-(4-((1-acetyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5-(5-(4-((1-Ethyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5-(5-(4-((1-(2-Hydroxyethyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

5-(5-(4-((1-(2-hydroxyethyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(5-(4-((1-(methylsulfonyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

5-(5-(4-((1-Isopropyl)piperidin-4-yl)oxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;

Ethyl 2-(4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidin-1-yl)acetate;

4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenoxy)piperidine-1-carboxamide;

5-(5-(4-(2-Hydroxyethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-methylpicolinonitrile;

4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)butanoic acid;

2-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorophenoxy)acetic acid;

4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)-N-methylbutanamide;

4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-

diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyrrolidin-1-yl)ethyl)benzamide;

N-Benzyl-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;

4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-2-yl)ethyl)benzamide;

- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(pyrrolidin-1-yl)propyl)benzamide;
- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-4-yl)ethyl)benzamide;
- 5 5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-2-yl)ethyl)benzamide;
- 5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-3-yl)ethyl)benzamide;
- 10 5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-4-yl)ethyl)benzamide;
- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-morpholinoethyl)benzamide;
- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(4-methylpiperazin-1-yl)ethyl)benzamide;
- 15 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyridin-3-yl)ethyl)benzamide;
- Ethyl 2-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamido)acetate;
- Ethyl 3-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamido)propanoate;
- 20 Ethyl 4-(4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamido)butanoate;
- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-methoxyethyl)benzamide;
- 25 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-methoxypropyl)benzamide;
- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-methoxyethyl)-N-methylbenzamide;
- 5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(pyrrolidin-1-yl)ethyl)benzamide;
- 30 5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(pyrrolidin-1-yl)propyl)benzamide;
- 5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-morpholinoethyl)benzamide;

- 5-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-(4-methylpiperazin-1-yl)ethyl)benzamide;
N-(2-(4-Bromo-1-methyl-1H-pyrazol-5-yl)ethyl)-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;
- 5 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-phenethylbenzamide;
4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-phenylpropyl)benzamide;
- 10 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-4-ylmethyl)benzamide;
4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(thiophen-2-ylmethyl)benzamide;
- 15 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-2-yl)benzamide;
4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-3-yl)benzamide;
- 20 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-4-yl)benzamide;
4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-2-ylmethyl)benzamide;
- 25 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyridin-3-ylmethyl)benzamide;
4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(furan-2-ylmethyl)benzamide;
- 30 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-morpholinopropyl)benzamide;
4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(4-methylpiperazin-1-yl)propyl)benzamide;
N-Benzyl-4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;
- 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(thiophen-2-ylmethyl)benzamide;
4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(pyrrolidin-1-yl)propyl)benzamide;

- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(1-methylpiperidin-4-yl)benzamide;
- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-(cyclopentylmethyl)-2-fluorobenzamide;
- 5 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(2-hydroxy-2-methylpropyl)benzamide;
- N-(2-Chlorobenzyl)-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;
- N-(3-Chlorobenzyl)-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;
- 10 N-(4-Chlorobenzyl)-4-(7-(6-cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluorobenzamide;
- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(pyrazin-2-ylmethyl)benzamide;
- 15 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-((5-methylfuran-2-yl)methyl)benzamide;
- 4-(4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)phenyl)-N-phenylbutanamide;
- 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(4-fluorobenzyl)benzamide;
- 20 4-(7-(6-cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(oxazol-2-yl)benzamide;
- 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(thiazol-2-yl)benzamide;
- 25 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(1-methyl-1H-pyrazol-5-yl)benzamide;
- 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(6-methylpyridin-3-yl)benzamide;
- 30 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(5-fluoropyridin-3-yl)benzamide;
- 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-hydroxypropyl)benzamide;

4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-(3-(cyclopentyl(methyl)amino)propyl)-2-fluorobenzamide;
 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(3-(2-oxopyrrolidin-1-yl)propyl)benzamide;
 5 5-(5-(3-Fluoro-4-(2-(4-methylpiperazin-1-yl)-2-oxoethoxy)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile;
 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-(tetrahydro-2H-pyran-4-yl)benzamide;
 4-(7-(6-Cyano-5-methylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-2-fluoro-N-((tetrahydro-2H-pyran-4-yl)methyl)benzamide;
 10 4-(7-(6-Cyano-5-(trifluoromethyl)pyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-5-yl)-N-(3-(3,3-difluoropyrrolidin-1-yl)propyl)-2-fluorobenzamide;

or a pharmaceutically acceptable salt thereof or *N*-oxide thereof.

- 15 26. A pharmaceutical composition comprising a compound of any one of claims 1-25, or a pharmaceutically acceptable salt thereof.
27. A method of treating an androgen receptor dependent or androgen receptor mediated disease or condition in mammal comprising administering to the mammal a therapeutically effective amount of a compound according to any one of claims 1-25.
- 20 28. The method of claim 27, wherein androgen receptor dependent or androgen receptor mediated disease or condition is selected from benign prostate hyperplasia, hirsutism, acne, adenomas and neoplasies of the prostate, benign or malignant tumor cells containing the androgen receptor, hyperpilosity, seborrhea, endometriosis, polycystic ovary syndrome, androgenic alopecia, hypogonadism, osteoporosis, suppression of
 25 spermatogenesis, libido, cachexia, anorexia, androgen supplementation for age related decreased testosterone levels, prostate cancer, breast cancer, endometrial cancer, uterine cancer, hot flashes, and Kennedy's disease muscle atrophy and weakness, skin atrophy, bone loss, anemia, arteriosclerosis, cardiovascular disease, loss of energy, loss of well-being, type 2 diabetes and abdominal fat accumulation.
- 30 29. A method of treating cancer in a mammal comprising administering to the mammal a therapeutically effective amount of a compound of any one of claims 1-25.
30. The method of claim 29, wherein the cancer is a hormone dependent cancer.
31. The method of claim 29, wherein the cancer is prostate cancer.
32. The method of claim 29, further comprising administering to the mammal at least one
 35 additional anti-cancer agent.

33. A method of treating cancer in a mammal comprising administering to the mammal a thereapeutically effective amount of a compound, wherein the compound:
- 5 a) is an androgen receptor inverse agonist; androgen receptor antagonist; is an androgen receptor degrader; is an androgen receptor trafficking modulator; is an androgen receptor DNA-binding inhibitor; or combinations thereof; and
- b) has minimal pro-convulsant activity and/or minimal impact on seizure threshold; displays minimal modulation of the GABA-gated chloride channel; displays minimal binding to the GABA-gated chloride channel; has minimal antagonism of the GABA-gated chloride channel; has minimal interaction with a GABA-gated chloride channel;
- 10 has minimal interaction with the GABA-gated chloride channel; or combinations thereof.
34. The method of claim 33, wherein the cancer is prostate cancer and the compound is a compound of any one of claims 1-25.

15