


Ref.: S 949 EP/I
Case: 5203
Scripps Clinic and Research
Foundation

1108094-0 1985

17. Mai 1993
VOSSIUS & PARTNER
PATENTANWÄLTE
SIEBERTSTR. 4
8000 MÜNCHEN 80

MONOCLONAL ANTIBODIES TO NEW FACTOR VIII
COAGULANT POLYPEPTIDES

This invention relates to a monoclonal antibody of class IgG produced by hybridoma formed by fusion of cells from a mouse myeloma and spleen cells from a mouse previously immunized with human factor VIII:C, which antibody reacts with human factor VIII:C polypeptides. The invention has utility in the therapy for classic hemophilia, and in the further study and characterization of the polypeptide or polypeptide complexes which provide desired clotting behavior to the blood of humans and other mammals.



It has long been known that plasma factor VIII plays a crucial role in blood coagulation, and that thrombin activates the coagulant effect of factor VIII. Recent attempts to characterize factor VIII have postulated that factor VIII is a complex of at least two polypeptides, which are known as VIII:C and VIII:R, and have found coagulant activity to reside in the VIII:C portion. Studies of the effect of thrombin on factor VIII:C have led to the conclusion that thrombin activates this factor by breaking it down into several smaller polypeptides. However, no prior studies have been able to associate thrombin-induced factor VIII activation in humans with any defined polypeptides formed from human factor VIII:C.

This Sheet Has Been Replaced



0-1000000000000

1 For instance, Hoyer and Trabold, in "The effect of
thrombin on human factor VIII", J. Lab. Clin. Med. 97:50-64
(1981), sought to purify human factor VIII:C by immunoadsorbent
5 chromatography using a polyclonal antibody to factor VIII:R
raised in the rabbit. They then incubated the factor VIII:C
with purified human α -thrombin, and determined that small
10 amounts of the thrombin activated the factor VIII:C whereas
larger amounts activated it less or not at all. They also
concluded that thrombin activation is accompanied by a decrease
in the size of the protein, and they proposed a molecular
15 weight of about 116,000 for the activated factor VIII:C. Most
significantly, they could not identify specific polypeptides
that retained factor VIII:C activity and VIII:C antigen
determinants.

20 Fulcher, C.A. and Zimmerman, T.S. in "Characterization
of the human factor VIII procoagulant protein with a
heterologous precipitating antibody", Proc. Natl. Acad. of Sci.
25 USA, 79:1648-1652 (1982) obtained highly purified human factor
VIII:C from plasma concentrate by passing the concentrate
through a column containing a monoclonal antibody to factor
VIII:R, eluting the VIII:C from the adsorbed VIII:C/VIII:R
30 complex, and concentrating the factor VIII:C on a second
column. The purified factor VIII:C was then analyzed by sodium
dodecyl sulfate/polyacrylamide gel electrophoresis (hereafter,
"SDS-PAGE"), both before and after addition of thrombin to the
35



1 purified material. The purified factor VIII:C prior to
thrombin addition showed a wide array of bands on SDS-PAGE
corresponding to polypeptides of various molecular weights
5 (M_r), including a relatively strong doublet at M_r of 80,000
and 79,000, and at least six additional faintly staining
polypeptides with larger M_r including one at M_r about
10 92,000. Addition of thrombin to the purified factor VIII:C
caused the diminution or disappearance of all of the
polypeptides shown prior to thrombin addition.

The coagulant activity of the plasma concentrate rose
15 following thrombin addition to a maximum of three times that of
the material prior to thrombin addition, and then diminished.
The coagulant activity of the purified factor VIII:C also rose
to a maximum of three times that of the pre-activated
20 material. That is, the thrombin had essentially the same
activating effect on the factor VIII:C in each case. Thus,
whereas the purified factor VIII:C possessed a reported
specific activity some 3280 times that of the starting
25 material, one skilled in this art would conclude that the
increase in specific activity was due to the high degree of
purification achieved. There is no basis in this article for
30 ascribing activated coagulant activity to any specific one or
more of the large number of polypeptides associated with the
bands observed in the purified factor VIII:C prior to
activation with thrombin.

35

This Sheet Has Been Replaced



1 The present invention relates to a
 monoclonal antibody of class IgG which reacts
with human Factor VIII:C polypeptides selected from the group
5 consisting of:

(A) polypeptides of $M_r = 44,000$; $92,000$; $108,000$; and
undigested Factor VIII:C; and not polypeptides of $M_r = 54,000$;
 $71,000$; $72,000$; $79,000$; or $80,000$;

10 (B) polypeptides of $M_r = 54,000$; $92,000$; $108,000$; and
undigested Factor VIII:C; and not with polypeptides of $M_r =$
 $44,000$; $71,000$; $72,000$; $79,000$; or $80,000$; and

(C) polypeptides of $M_r = 79,000$ and $80,000$; $108,000$;
15 and undigested Factor VIII:C; and not with polypeptides of $M_r =$
 $44,000$; $54,000$; $71,000$; $72,000$; or $92,000$ wherein the M_r values
are determined by subjecting the polypeptides to sodium dodecyl
sulfate-polyacrylamide gel electrophoresis.

20 Other aspects of the invention include biological preparations
containing antibodies exhibiting different reaction profiles.

The polypeptide coagulants exhibit factor VIII:C coagulant
activity and factor VIII:C immunological behavior, and
25 show characteristic M_r value(s) when analyzed by sodium
dodecyl sulfate-polyacrylamide gel electrophoresis (hereafter,
"SDS-PAGE"). By "polypeptide complex" hereinafter is meant a
coagulant comprising combinations of two or more polypeptides
known to be physically distinct, but also preparations contain-
30 ing only one discernible polypeptide, i.e., that which exhibits



1 a band at an M_r of 92,000.

2 The description below showing the preparation of the
3 coagulant utilizes human factor VIII:C which has been
4 highly purified, in order to free the identification and
5 characterization of the novel product from the effects of
6 extraneous polypeptides. It should be recognized, though, that
7 the invention itself is not dependent on how the starting
8 material has previously been treated, except where specifically
9 indicated. The active factor VIII:C polypeptides of the
10 present invention can be prepared not only by digestion with
11 thrombin as described in greater detail below, or with other
12 similarly acting proteases such as Factor Xa, Factor IXa, or
13 Russell's viper venom-V, but also by recombinant DNA techniques
14 in which the polypeptides of interest are produced by bacteria,
15 yeast, or other cells into which one or more genes for
16 producing the polypeptides of interest are inserted by
17 techniques known to those of ordinary skill in the art. Either
18 process for producing the complex of interest can be expected
19 to produce the complex in a mixture with one or more other
20 polypeptides.
21
22
23
24
25

26 Any plasma or plasma concentrate containing human
27 factor VIII:C can be employed to advantage. The novel
28 coagulant can be prepared from human factor
29 VIII:C which has been ultrapurified in accordance with the
30 process described in U.S. Patent No. 4,361,509, issued November
31 30, 1982, the disclosure of which is hereby incorporated herein
32
33
34
35

This Sheet Has Been Replaced



1 by reference. In that process, a source of factor VIII:C such
as plasma or a plasma concentrate is passed through an
immunoabsorbent column to which monoclonal antibodies to factor
5 VIII:R have been attached. The factor VIII:R/VIII:C complex is
adsorbed on the column, and then the factor VIII:C is eluted
and passed through a second column such as aminohexyl agarose.
It should be noted that the second column can also be an
10 immunoabsorbent containing antibodies to factor VIII:C. The
factor VIII:C should be kept free of α -thrombin and other
proteases. The factor VIII:C is conveniently stored in a
15 saline, buffered solution containing, e.g., 0.3M calcium
chloride, at a pH of 6.8 to 7.4.

The human factor VIII:C is then digested with
 α -thrombin under conditions effective to form the polypeptide
20 complex described below. The purified α -thrombin can be
prepared by the procedure described by Fenton, J.W. II; Fasco,
M.J.; Stackrow, A.B.; Aronson, D.L.; Young, A.M.; and
25 Finlayson, J.S. "Human Thrombin. Production, Evaluation, and
Properties of α -Thrombin". J. Biol. Chem. 252: 3587-3598
(1977).

The α -thrombin and the factor VIII:C are combined in
30 an aqueous system, preferably buffered at a pH of about 6.8 to
about 7.4. The thrombin should be present in an amount
relative to the factor VIII:C that is sufficient to permit
reaction with the factor VIII:C, but not so high that the
35



1 factor VIII:C degrades to inactive polypeptides before the
desired active polypeptide complex can be recovered. As an
illustration, a preparation containing 200-400 units (0.2
5 mg/ml) of factor VIII:C per ml should be digested with about
0.1 to about 0.5 units/ml of α -thrombin. The digestion can
proceed at room temperature; too high a temperature can
denature the polypeptides, and too low a temperature can retard.
10 the progress of the digestion.

The digestion is allowed to proceed for a time long
enough to permit the formation of the desired polypeptide
15 complex. The optimum time will be from 0.1 to 60
minutes, with times from 0.1 to 30 minutes preferred. Times
from 1 to 10 minutes have been found highly satisfactory,
although it will be recognized that optimum times can be
20 identified with minimal experimentation using aliquots of the
factor VIII:C starting material being treated. An optimum time
is that which forms the maximal amount of the polypeptide
complex exhibiting an M_r of about 92,000, accompanied by the
25 formation of a protein complex exhibiting an M_r doublet of
about 79,000 and about 80,000, without degrading the M_r
92,000 polypeptide significantly. The M_r 79,000-80,000
30 doublet may possibly function after it has undergone
degradation to a complex exhibiting a doublet at M_r values of
71,000-72,000. The 71,000-72,000 doublet alone does not
exhibit factor VIII:C activity.

35

This Sheet Has Been Replaced



1 The digestion is then discontinued by adding to the
reaction mixture an effective amount of (p-amidinophenyl)
methanesulfonyl fluoride (hereafter, "p-APMSF") or another
5 thrombin inhibitor. The p-APMSF prevents the α -thrombin from
reacting further with the factor VIII:C proteins, without
itself degrading those proteins. The amount of p-APMSF to add
should comprise about 1.5 to about 2.5 millimoles per unit of
10 α -thrombin activity initially present in the reaction mixture.

p-APMSF can be obtained through California Medicinal
Chemistry Company, San Francisco, California, and its
15 preparation is described in Laura, R.; Robinson, D.J.; and
Bing, D.H. "(p-Amidinophenyl)methanesulfonyl Fluoride, an
Irreversible Inhibitor of Serine Proteases", Biochemistry
(1980) 19, 4859-4864, at 4861.

20 The reaction mixture can then be treated to
concentrate the polypeptide complex comprising the present
invention. Preferably, the polypeptide complex is concentrated
with respect to other factor VIII and non-factor VIII
25 proteinaceous material, to provide the complex in a form which
affords the very high activity possessed by the complex in the
purified form. Purification techniques include, for example,
30 ultrafiltration, ultracentrifugation, ion exchange, gel
permeation chromatography, preparative electrophoresis,
isoelectric focusing, and gel and affinity chromatography.

35 The desired complex can also be concentrated and/or
recovered by passing the reaction mixture, which can already



1 have been concentrated by another technique, through an
immunoabsorbent column containing anti-human VIII:C antibodies,
or equivalent antibodies against VIII:C other than human, that
5 react with the polypeptide(s) of the complex. The antibodies
are attached to agarose (see Example I below). Several
antibodies which can be used to concentrate the complex and/or
10 isolate certain of the polypeptides thereof are described
below. The active VIII:C complex adsorbs preferentially to the
column, and is then eluted from the column with a solution of
calcium ions (e.g., CaCl_2) which can optionally also contain
15 a non-ionic surfactant. Suitable non-ionic surfactants include
alkyl phenyl polyoxyethylenes such as Triton-X-100[®], -N-101[®], or
-X-405[®] (Eastman Chemical Co.); Tween-20[®], -60[®] or -80[®] (Sigma
Chemical Co.); and Nonidet P-40[®] (Sigma Chemical Co.); all of
20 these are well-known articles of commerce having known chemical
formulas.

The amounts of calcium ion and surfactant to use
25 should be high enough to desorb the polypeptide complex, but
not so high that the eluant inactivates the polypeptide. A
calcium ion concentration of up to 0.5 M or even up to
1.0 M is satisfactory, and 0.25 M is preferred. A surfactant
30 concentration of up to 1 wt. % is satisfactory, and about
0.1 wt. % is preferred. The eluant is applied to the
immunoabsorbent column at 1 to 8 bed volumes per
hour, and preferably about 3 to 4 per hour. Too high a
35



1 flow rate risks disruption of the column and non-absorption of
the polypeptide complex. The skilled practitioner will readily
adapt these guidelines to immunoadsorbent processes other than
5 fixed-bed columns.

The VIII:C polypeptide complex is recovered in a
suitable buffer, at a pH of 6.8 to 7.4, which also
contains calcium ion and surfactant from the eluant solution.
10 The calcium and surfactant concentrations can be lowered, and
the surfactant preferably removed, such as by dialyzing the
solution against a buffer such as the VIII:C buffer used in
15 Example 1 which contains a lower amount of calcium ion. The
complex can be stored in this solution, or
lyophilized. The complex can be administered to patients with
hemophilic clotting disorders by adjusting the calcium content
20 to be physiologically compatible and injecting a sterile saline
solution thereof.

When analyzed by SDS-PAGE as shown below, the coagulant
exhibits an M_r of about 92,000, and can
25 normally contain material exhibiting an M_r doublet of about
79,000 and about 80,000, some of which can have undergone
degradation to exhibit an M_r doublet of about 71,000 and
30 about 72,000. In its purified form, this band or the group of
these bands are essentially the only bands that appear.
However,

less than 100%,

35

This Sheet Has Been Replaced



1 i.e., 95%, 90%, or even 80%, 70% or 60%, or even as little
as 30%, 20%, 10%, or 1% of the proteinaceous matter present
in the biological preparation may comprise the coagulant.

5

The protein complex possesses
specific VIII:C coagulant activity (that is, activity per
milligram of all protein present) higher than that exhibited by
10 purified human factor VIII:C, for instance human factor VIII:C
purified by the process disclosed and claimed in the
aforementioned U.S. Patent No. 4,361,509. Indeed, the activity
of the purified complex should be several times, e.g., 3 to 5
15 times, that of purified human factor VIII:C, and is
advantageously at least 10 times or even 50 times as active.
Likewise, biological preparations comprising the complex
20 in association with one or more other proteins
can be prepared which exhibit higher specific activity than
that afforded in previously known coagulant preparations. The
specific activity of the complex, and of biological
25 preparations containing it, is higher than 1800 Units/mg,
advantageously higher than 5400 Units/mg, and in more
advantageous embodiments exceeds 7,500 and even 10,000
Units/mg. Preferably, the specific activity of the
30 preparations exceeds that of the purified human factor VIII:C
used herein by a factor of 3 to 5, and more advantageously by
at least 10 times, by 50, or even by 100 times.

35

This Sheet Has Been Replaced



The protein complex and biological preparations thereof, are characterized in that the enhanced activity described above remains present over a continuous period of at least 10 minutes and preferably at least 30 minutes. Of course, activity will generally be stable for much longer. The complex also possesses the immunological characteristics of a factor VIII:C protein, i.e., it binds to an antibody for human factor VIII:C. This can be ascertained, for instance, by growing a monoclonal antibody to factor VIII:C as described below, attaching the antibody to an agarose column, passing an aqueous solution of the complex through the column, and assaying the resultant solution for factor VIII:C activity.

20 The polypeptides of $M_r=92,000$ and $M_r=79,000-80,000$ are stable, relative to native human factor VIII:C which is notoriously susceptible to proteolysis, degradation, and resultant loss of coagulant activity. This stability is
25 demonstrated by the polypeptides' ability to survive the treatment steps described herein.

30 EXAMPLE I

This Example shows how factor VIII:C was purified from commercial concentrate and digested with purified α -thrombin.



1 A monoclonal antibody to factor VIII:C was produced and used to
identify VIII:C polypeptides. At several selected points
during the digestion, portions of the digestion mixture were
5 assayed for VIII:C (coagulant) activity, and for protein bands
using SDS-PAGE.

Purification of VIII:C. All steps were at room
temperature. Chemicals were reagent grade. 40 bottles of
10 commercial factor VIII concentrate (provided by Armour
Pharmaceutical) were reconstituted in 1000 ml of VIII:C buffer
(0.02 M imidazole/0.15M sodium chloride/0.1 M L-lysine
15 HCl/0.02% sodium azide, pH 6.8). This sample, which contained
a total of 17,000 units of VIII:C activity, was applied to a
2.5 - 3.0 liter bed volume immunoabsorbent column. The column
was cyanogen bromide-activated agarose (Sephacrose 4B,
20 Pharmacia, Piscataway, New Jersey), to which monoclonal
antibodies to VIII:C had been covalently bonded. The
antibodies were raised and attached to the column as described
in the aforementioned U.S. Patent No. 4,361,509. The
25 antibodies were precipitated from ascites fluid using 50%
ammonium sulfate, reprecipitated two more times, and then
attached to the column at a density of 2-4 mg/ml of Sepharose.
30 The immunoabsorbent was pre-eluted with 3M sodium thiocyanate,
washed with VIII:C buffer (0.02M imidazole HCl pH 7.0, 0.15 M
NaCl, 0.1 M L-lysine-HCl, 0.02% sodium azide), treated twice
with 2mM di-isopropyl fluorophosphate, and then the concentrate
35 was added.



1 The column was washed with 20 liters of VIII:C buffer
containing 0.15 M sodium chloride, and VIII:C was then eluted
from the VIII:R with VIII:C buffer containing 0.35 M calcium
5 chloride. Active fractions were pooled and concentrated under
nitrogen pressure 100-fold in an Amicon stirred cell with a
YM10 membrane. The concentrate was then diluted 1:10 in VIII:C
buffer and applied to a 4 ml column of aminohexyl-Sepharose
10 (Pharmacia) equilibrated in VIII:C buffer containing 0.025 M
calcium chloride, VIII:C was eluted in high concentration with
VIII:C buffer containing 0.3 M calcium chloride at a flow rate
15 of 10 ml/hr. The concentrated immunoabsorbent pool was
adjusted to 0.25 M calcium chloride and adsorbed twice for 1 h
each time with 1/10 vol/vol of a mixture of monoclonal
anti-fibrinogen, anti-fibronectin and anti-vWF antibodies which
20 had been coupled to cyanogen bromide-activated Sepharose.

Production of monoclonal antibody against VIII:C.

Monoclonal antibodies were produced as described in U.S. Patent
No. 4,361,509 using purified VIII:C as immunogen. The
25 antibodies were selected with a solid-phase assay in
Linbro-Titertek[®] (Flow Laboratories, Inglewood, CA) plates and
an enzyme-linked immunoabsorbent (ELISA) detection system
30 described in Engvall, E. and Perlmann, P. "Enzyme-linked
immunoabsorbent assay (ELISA), Quantitative assay of
immunoglobulin G" Immunochemistry 8:871-874 (1971)) using a
peroxidase-antibody conjugate (Zymed Laboratories, Burlingame,
35

This Sheet Has Been Replaced



1 CA). The plates were coated with 100 ng of purified VIII:C per
well. The ELISA-positive culture supernatant of the clone
selected for use in this study also inhibited plasma VIII:C
5 activity.

Thrombin activation time course analysis of purified
VIII:C. Purified human α -thrombin(sp. act. 2534 U/mg, final
concentration 0.5 U/ml), was added to the purified VIII:C
10 (final concentration 167 μ g/ml) in imidazole saline buffer
containing 0.04 M CaCl_2 . Buffer alone was added to a control
aliquot. The solutions were incubated at room temperature and
15 at various time intervals samples of the VIII:C-thrombin
mixture were added to tubes containing p-APMSF (California
Medicinal Chemistry Co.) to inactivate the thrombin rapidly and
irreversibly. In order to minimize hydrolysis of p-APMSF, it
20 was diluted 1:10 from a stock solution (100 mM in methanol)
into imidazole saline buffer 60 seconds before reaction with
the VIII:C-thrombin samples. The final p-APMSF concentration
was 1 mM. The control aliquot was treated similarly with
25 p-APMSF at the start of the experiment. At the end of the 60
minute time course, all VIII:C samples were assayed for VIII:C
activity using an activated partial thromboplastin time assay
30 described in the literature and then prepared for SDS-PAGE.

SDS-PAGE "Procedure A":

Discontinuous SDS polyacrylamide slab gel
electrophoresis was performed based on the procedure of
35

This Sheet Has Been Replaced



1 Laemmli, U.K., Nature 227, 680-685, 1970. The "Procedure A" followed is:

I. Sample Preparation

- 5 1. Dialyze the protein sample (ideally 50-100 microliters containing 5-60 micrograms of protein) against sample buffer overnight at room temperature. If the sample contains calcium ion, include 10 millimolar ethylenediamine tetracetic acid (EDTA) in the sample buffer.
- 10 2. Place the dialyzed sample in a tube and add 1/10 volume of 10% SDS. Cover the tube with aluminum foil. Heat the sample in a boiling waterbath for 10 minutes.
- 15 3. Remove the sample from the waterbath and add to it 1/10 volume of 500 millimolar dithiothreitol. Incubate it at 56°C for 4 hours.
4. Allow the sample to cool to room temperature and prepare it for layering onto the gel by adding stock glycerol solution to 10% final concentration and stock bromophenol blue dye solution to 0.05% final concentration.

20 II. Preparation of gel solutions (use deionized, distilled water)

1. Stock glycerol solution: 50% glycerol
- 25 2. Stock bromophenol blue dye solution: 0.5% bromophenol blue
3. Lower gel stock solution:
- 18.2 grams of Tris base
4 ml of 10% SDS
final volume 100 ml.
Adjust pH to 8.8 with concentrated hydrochloric acid. Filter.
- 30 4. Upper gel stock solution:
- 6.1 grams of Tris base
4 ml of 10% SDS
Final volume 100 ml
Adjust pH to 6.8 with concentrated hydrochloric acid. Filter.
- 35



- 1 5. Sample buffer
0.01M sodium phosphate
1.0% SDS
10 millimolar disodium EDTA
Final volume 1 liter
5 pH adjusted to 7.0 with sodium hydroxide or
phosphoric acid.
- 10 6. Acrylamide stock solution:
Dissolve 30 g of acrylamide in 50 ml of water and
add 0.8 g of bisacrylamide and dissolve. Bring
to 100 ml final volume. Filter the solution and
store in the dark at 4°C.
- 15 7. Stock electrode buffer solution;
30.3 g Tris base
144.1 g glycine
final volume 1 liter
- 20 8. Electrode buffer
100 ml of stock electrode buffer solution
890 ml water
10 ml of 10% SDS
- 25 9. Stock Coomassie blue dye solution
1% Coomassie blue R 250 in water
Dissolve with stirring for at least 30 minutes at
room temperature and filter.
- 30 10. Ammonium persulfate solution
10% ammonium persulfate.
Stored in dark at 4°C and made fresh every week.
- III. Gel Preparation and Running: Final acrylamide
concentration = 7.5%
- 35 1. Lower gel solution
20 ml of lower gel
Stock lower gel solution 5 ml
Stock acrylamide solution 5 ml
Water 10 ml
N,N,N',N'-tetramethylethylenediamine .005 ml
(TEMED)
10% ammonium persulfate 0.1 ml



1 2. Upper gel solution 10 ml of upper gel

	Stock upper gel solution	2.5 ml
	Stock acrylamide solution	1.0 ml
5	Water	6.5 ml
	N,N,N',N'-tetramethylethylenediamine (TEMED)	0.01 ml
	10% ammonium persulfate	0.03 ml

3. Procedure:

10 a. Prepare the slab gel apparatus for a 14.5 cm x 9.0 cm x 0.8 mm slab gel. The apparatus is a standard gel electrophoresis apparatus, available, for instance, from Hoeffer Scientific Instruments, San Francisco, California.

15 b. Mix all lower gel ingredients except the TEMED and ammonium persulfate in a 50 ml vacuum flask and de-aerate. Then add the TEMED and ammonium persulfate, mix gently and pour the lower gel immediately. Layer the lower gel with water-saturated butanol and allow it to polymerize undisturbed for at least 1 hour, preferably 2-6 hours.

20 c. Pour off the butanol layer and rinse the top of the lower gel with the complete upper gel mixture. (The upper gel mixture is prepared, de-aerated and TEMED and ammonium persulfate are added as for the lower gel above).

25 d. Pour the upper gel and insert the comb into the upper gel allowing at least 1.0 cm between the bottom of the comb teeth and the upper gel-lower gel interface. Fill with upper gel solution as full as possible. Allow upper gel to polymerize at least 1 hour before running the gel.

30 e. To remove the comb, pipet electrode buffer over the top of the upper gel and gently remove the comb. Rinse the upper gel wells with electrode buffer several times.

35 f. Assemble the apparatus for running and add electrode buffer. Apply the sample(s) by layering it into the upper gel wells underneath the buffer layer.



- 1 g. Run the gel using constant current:
 8 milliamperes while samples are in the upper gel
 and 15 milliamperes while samples are in the
 lower gel. Stop the electrophoresis when the
 bromophenol blue dye front is 1.0 cm from the
 5 bottom of the lower gel.

IV. Fixing and Staining the gel

Reference: Fairbanks, G., Steck, T.L., and Wallach, D.F.N., Biochemistry 10, 2606-2617, 1971.

- 10 1. Fix the gel at least overnight in a sealed chamber in a solution containing 25% isopropanol, 10% acetic acid, 10 ml of 1% Coomassie blue stock solution, final volume 400 ml.
- 15 2. Next, soak the gel at least 1 hour in 10% isopropanol, 10% acetic acid and 1.0 ml of 1% Coomassie blue stock solution, final volume 400 ml.
3. Soak the gel about 4 hours in 10% acetic acid with changes or until completely destained.
- 20 4. The destained gel can be dried onto filter paper using a gel dryer to increase contrast.

Approximately 5-20 g of protein was applied to the gels. VIII:C M_r s were calculated for reduced samples by semilogarithmic plots of M_r versus migration distance, using reduced fibronectin (M_r 200,000), phosphorylase b (M_r 95,000), bovine serum albumin (M_r 68,000), and IgG heavy chain (M_r 50,000), and ovalbumin (M_r 43,000) as standards.

25 Scanning and integration of a photographic print of the finished gel was done using a Zeineh soft laser scanning densitometer.

Results. The specific activity of the purified factor VIII:C was 2000 units/mg. Thrombin activation of purified VIII:C activity was analyzed over a 60 minute time course.

Before thrombin exposure the untreated VIII:C sample showed the characteristic array of VIII:C forms ranging from a doublet at

35



1 $M_r = 79,000-80,000$ to a band at $M_r = 188,000$. A band above
 $M_r = 188,000$ and two bands below $M_r = 79,000$ did not bind
to the monoclonal anti-VIII:C antibody immunoabsorbent. The
5 bands between $M_r=79,000$ and $M_r=188,000$ did bind to the
anti-VIII:C antibody.

During the first 5 minutes of the thrombin activation time
course, all but one of the monoclonal anti-VIII:C antibody
10 reactive bands with an M_r greater than 92,000 gradually
disappeared and were undetectable when VIII:C activity reached
its peak at 5 minutes.

15 A band at $M_r = 122,000$ appeared thrombin-resistant in only
some experiments, but after extensive thrombin treatment
neither this band nor any other band was reactive with the
immobilized monoclonal anti-VIII:C antibody.

20 A band at $M_r = 92,000$ increased in intensity as
VIII:C activity increased. A doublet at $M_r = 79,000$ and
80,000 appeared to be converted to a doublet at $M_r =$
25 71,000-72,000, with the latter form predominant from 5 to 60
min. as VIII:C activity decreased. Two bands, at $M_r = 54,000$
and $M_r = 44,000$, became clearly visible from 5 to 60 min.
The $M_r = 44,000$ band also appeared as a doublet in some
30 experiments. The $M_r = 71,000-72,000$ doublet, the $M_r =$
54,000 band and the $M_r = 44,000$ band were not removed to a
significant degree by the immobilized monoclonal anti-VIII:C
antibody.
35



1 Scanning and integration of the gel discussed above
allowed correlation of changes in polypeptide concentration
with changes in VIII:C activity. The results are shown in the
5 Table. As shown, the $M_r = 92,000$ band increased and then
decreased in concentration in parallel with VIII:C activity.
This suggests that the $M_r = 92,000$ band is an active form of
VIII:C whose concentration increased with thrombin activation.
10 The $M_r = 54,000$ and $M_r = 44,000$ bands increased steadily in
concentration between 1 and 40 min., even after activity of the
mixture declined.

15 Most of the $M_r = 79,000-80,000$ doublet was lost
during the first 0.1 to 10 min. as VIII:C activity peaked,
while most of the $M_r = 71,000-72,000$ doublet appeared during
this time and predominated even as VIII:C activity decreased.
20 These data suggest that the $M_r = 71,000-72,000$ doublet was
derived from the $M_r = 79,000-80,000$ doublet and that the M_r
 $= 71,000-72,000$ doublet is inactive by itself. These data are
consistent with the retention by the $M_r = 71,000-72,000$
25 doublet of the ability to complex with the $M_r = 92,000$
polypeptide; this complex would also have activity.

Direct evidence that the $M_r = 92,000$ polypeptide is
30 complexed with the $M_r = 79,000-80,000$ doublet derives from
experiments utilizing the anti-VIII:C monoclonal
immunoabsorbent. It was first shown that the monoclonal
antibody reacts predominantly with the $M_r = 79,000-80,000$
35 doublet and not with the $M_r = 92,000$ polypeptide. This was



1 shown by electrophoretic transfer experiments. Then it was
shown that the monoclonal anti-immunoabsorbent removed both the
 $M_r = 79,000-80,000$ doublet and the $M_r=92,000$ polypeptide
5 from the solution. The $M_r=92,000$ polypeptide could be eluted
from the anti-VIII:C immunoabsorbent column with 10 mM EDTA,
whereas the $M_r=79,000-80,000$ doublet remained bound. The
doublet could be subsequently eluted with 3M sodium
10 thiocyanate. These experiments demonstrated that the
 $M_r=92,000$ polypeptide bound to the immunoabsorbent because it
was complexed with the $M_r=79,000-80,000$ doublet and that it
15 was this doublet which bound directly to the immunoabsorbent.

20

25

30

35

This Sheet Has Been Replaced



	35	30	25	20	15	10	5	1		
Time (min) from addition of thrombin to addition of p-APMSF:	0*	0.1	1	2	5	10	20	30	40	60
Activity of digestion mixture	300	900	1300	1350	1400	1250	800	375	250	100
Amount of each poly- peptide (M_r) present, as % of total pro- tein present:										
$M_r = 92,000$	7.3	9.5	11.1	11.5	11.8	9.6	7.3	6.7	4.9	4.4
$M_r = 79-80,000$	21.9	18.8	16.1	12.9	11.0	8.4	9.0	8.6	7.4	7.4
$M_r = 71-72,000$	0.0	9.0	11.2	12.9	16.5	15.9	15.7	18.8	18.2	20.4
$M_r = 54,000$	4.9	5.9	5.1	5.7	7.1	9.4	10.2	11.8	13.7	13.5
$M_r = 44,000$	0.0	0.0	0.0	3.1	3.7	5.9	6.1	7.2	7.8	7.6

* Measurements just prior to addition of α -thrombin

Activity is in Factor VIII:C units/ml

EXAMPLE II

1 This Example describes the results of treating
5 purified human Factor VIII:C with purified human activated
protein C (hereafter, "APC"), a known anticoagulant enzyme.
Human Factor VIII:C was purified as described above. APC was
purified by the procedure described in Marlar, R.A. et al.,
10 "Mechanism of action of human activated protein C, a
thrombin-dependent anticoagulant enzyme." Blood, Vol. 59, 1067
(1982), except that a mono S column of a Pharmacia FPLC system
was used to separate APC from thrombin.
15

Assays: Samples were assayed for VIII:C activity as
described above using an activated partial thromboplastin time
assay with hemophilia A plasma substrate.

20 Preparation of samples for electrophoresis: Since
calcium ions are required for APC activity, APC was stopped at
various times by addition of 1/10 volume of 100 mM EDTA containing 10 μ M
dansylarginine N-(3-Ethyl-1,5-pentanediyI)amide (DAPA) to the VIII:C + APC
25 aliquots as well as to the control VIII:C and APC aliquots. A 1/10-volume
of 10 %
sodium dodecyl sulfate was then added to these aliquots. They
were then heated in a boiling waterbath for 5 minutes and
subsequently dialyzed for SDS-PAGE as described above.
30

Discontinuous sodium dodecyl sulfate 7.5%
polyacrylamide gel electrophoresis (PAGE) of reduced VIII:C,
staining with Coomassie blue R250 and scanning and integration
35 of the gel were as described above.



1 Sample preparation: A 339 μ g sample of VIII:C in 0.3
ml of VIII:C buffer containing 0.3 M calcium chloride was
dialyzed overnight against buffer (50 mM Tris-chloride, 0.15 M
5 sodium chloride, 5 mM calcium chloride, 0.02% sodium azide, pH
7.4). To the dialyzed VIII:C sample was added 1.095 ml of
buffer, 90 μ l of rabbit brain cephalin (Sigma Chemical Co., St.
Louis, MD, reconstituted, stored and thawed according to
10 manufacturer's instructions) and 15 μ l of 1 mM
DAPA , to give a final DAPA
concentration of 10 μ M. The DAPA was included to inhibit any
15 trace amounts of thrombin present in the APC since 10 μ M DAPA
does not significantly inhibit APC. The final volume of the
sample was 1.5 ml and the final VIII:C concentration was 226
 μ g/ml. Four hundred microliters of the 1.5 ml VIII:C sample
20 were withdrawn and set aside as a control (designated VIII:C).
To the remaining 1.1 ml was added 20 μ l (10 ug) of APC giving a
final APC concentration of 9 μ g/ml (designated VIII:C + APC).
A second control sample was prepared containing all components
25 at similar concentrations except that VIII:C was omitted
(designated APC).

Timepoints: VIII:C alone; the mixture of VIII:C and
30 APC, and APC alone were placed in a 37⁰C waterbath and at
given timepoints aliquots were withdrawn for SDS-PAGE and/or
VIII:C activity assay. It was also determined in control that
APC remained active in the hydrolysis of the synthetic
35 substrate S-2238 after prolonged incubation at 37⁰C.



1 Results: Digestion of purified VIII:C with APC was
associated with loss of approximately 85% of the control VIII:C
activity. APC inactivation of VIII:C activity resulted in the
5 diminution of all VIII:C polypeptides of M_r between 92,000
and 188,000 and generation of a polypeptide of $M_r=45,000$,
while leaving the doublet of $M_r = 79-80,000$ intact.

10 A time course inactivation of purified VIII:C by APC
showed the progressive disappearance of specific VIII:C
polypeptides as VIII:C activity decreased over a 360 minute
time course. Scanning and integration of the gel showed that
15 the polypeptide $M_r = 188,000$ and the polypeptide of $M_r =$
92,000 decreased in parallel with VIII:C activity.

 One other polypeptide of intermediate M_r was cleaved
by APC but it was not easily quantitated by gel scanning. In
20 this experiment unlike the digestion with APC, some VIII:C
polypeptides of $M_r = 92,000-188,000$ were resistant to the APC
digestion. However, as in the digestion with APC, the doublet
polypeptide at $M_r = 79-80,000$ was not proteolyzed by APC.

25 A polypeptide of $M_r = 45,000$ appeared to increase in
concentration as the polypeptides of $M_r = 188,000$ and 92,000
decreased suggesting that it is a proteolytic fragment derived
30 from them. No other digestion products were visualized with
Coomassie blue.

 As shown in Example I, during thrombin activation of
VIII:C the $M_r = 92,000$ polypeptide increased and decreased in
35



1 parallel with VIII:C activity. In order to determine whether a
linear relationship existed between proteolysis of the
polypeptide of $M_r = 92,000$ and the loss of VIII:C activity,
5 the data of this time course inactivation run were re-plotted
to examine percent VIII:C activity versus percent of
polypeptide of $M_r = 92,000$. The amount of VIII:C activity
appeared to be proportional to the amount of the polypeptide of
10 $M_r = 92,000$.

The present invention relates to
monoclonal antibodies which are specific in an unforeseeable
15 way to the various polypeptides which are formed by the
reaction of factor VIII:C with a protease such as
alpha-thrombin. Each antibody reacts with human factor VIII:C
which has not been activated nor digested with thrombin or
20 equivalent proteases. The antibodies are further characterized
by their individual properties, as follows:

(A) One reacts with the $M_r=92,000$ polypeptide
described herein, with polypeptides of $M_r=108,000$ and larger,
25 and with the polypeptide of $M_r = 44,000$ which is present in
terminal-thrombin digests. It does not react with the
polypeptide described herein which exhibits a doublet of $M_r =$
30 $79,000-80,000$, nor with the polypeptide described herein which
exhibits a doublet of $M_r = 71,000-72,000$.

(B) One reacts with the $M_r=92,000$ polypeptide
described herein, with polypeptides of $M_r=108,000$ and larger,
35



1 and with the polypeptide of $M_r = 54,000$ which is present in
terminal thrombin digests. It does not react with the
polypeptide described herein which exhibits a doublet of $M_r =$
5 79,000-80,000, nor with the polypeptide described herein which
exhibits a doublet of $M_r = 71,000-72,000$.

(C) One reacts with the doublet of $M_r=79,000-80,000$
and polypeptides of $M_r=108,000$ and greater, but not with the
10 polypeptide of $M_r=92,000$, nor with the polypeptides of $M_r =$
71,000-72,000, $M_r=54,000$, nor $M_r=44,000$.

(D) One reacts only with polypeptides of $M_r=108,000$
15 and greater.

Each of these antibodies can be used to concentrate
the complex described above, from mixtures which also contain
other polypeptides. One such mixture is produced by partial
20 digestion of human factor VIII:C with α' -thrombin or an
equivalent protease. Another such mixture is that produced by
recombinant DNA techniques in which a desired polypeptide or
(complex is expressed by a microorganism and must be recovered
25 from a mixture with other proteinaceous compounds. Antibody
(A), (B), (C) or (D), or a combination of two, three, or all of
them, can be attached to an immunoadsorbent column in the
30 manner described in Example 1, and a feed solution of the
mixture containing the polypeptide(s) comprising the complex is
poured through the column. Those polypeptides having M_r of
92,000, 79,000-80,000, and 71,000-72,000 which are present in
35



1 the feed solution adsorb onto the column, from which they can
be eluted as taught hereinabove after the source solution has
been washed through the column. The resulting eluted solution
5 is thereby concentrated in the desired activated VIII:C complex
compared to the feed solution.

The new antibodies are also useful for analytical
10 purposes, to detect the occurrence of the reaction of human
VIII:C with thrombin or other proteases, because of their
ability to react with products of that reaction. An antibody
having profile (B) and an antibody having profile (C) have been
15 found which neutralize VIII:C coagulant activity when either is
bound to factor VIII:C. This additional property can be of
value in the diagnosis of hemophilia-related disorders.

The discovery of these antibodies also permits further
20 characterization of the components of the polypeptide complex
described herein. Thus, the polypeptide which exhibits a band
of $M_r=92,000$ contains (at least) two epitopes (i.e. antibody
binding sites) that are not destroyed by thrombin digestion and
25 are not present on the polypeptides which exhibit the doublet
of $M_r=79,000-80,000$ nor the doublet of $M_r=71,000-72,000$.
One of these epitopes is also present on the polypeptide of
30 $M_r=44,000$, and the other on the polypeptide of $M_r=54,000$.
Thus, the $M_r=54,000$ and $M_r=44,000$ polypeptides derive from
the $M_r=92,000$ polypeptide. Also, the $M_r=92,000$ and
 $M_r=79,000-80,000$ polypeptides are derived from a common
35



1 precursor or precursors. The discovery of antibodies having
profile (B) and profile (C) each of which neutralizes factor
VIII:C coagulant activity is further support that the
5 $M_r=92,000$ and $M_r=79,000-80,000$ polypeptides are important
to the coagulant function. The polypeptides exhibiting the
doublet of $M_r=79,000-80,000$ contain an epitope which is
destroyed by thrombin digestion and which is not present on the
10 polypeptide of $M_r=92,000$.

These monoclonal antibodies can be prepared by the
general steps of purification of human factor VIII:C; raising
15 monoclonal antibodies to the purified VIII:C; partial digestion
and activation of purified VIII:C to produce the polypeptide
complex described above, including identification of the
specific polypeptides; reaction of the anti-VIII:C antibodies
20 with the products of the activation; and characterization of an
antibody by identification of the polypeptide(s) with which it
reacted. This sequence is set forth in more detail in Example
III below. Alternatively, one can prepare an antibody by
25 isolating the particular polypeptide of interest from the
partial thrombin digestion products, for instance by
immunoabsorption onto and then elution off of a column
30 comprising agarose to which is coupled a monoclonal antibody
known to react with the polypeptide of interest, and then
raising a monoclonal antibody against that polypeptide using
the procedure described in Example III.

35



1

EXAMPLE III

Monoclonal antibodies to human Factor VIII:C were
5 raised using the following procedure, starting from highly
purified VIII:C which had been prepared by the process of U.S.
Patent No. 4,361,509.

10 Mice were injected with highly purified factor VIII:C
according to the following procedure. On day zero, the mice
were injected intraperitoneally with a composition prepared by
dissolving (or suspending) 10 μ g of the protein in 0.1 ml of
15 buffer containing 0.05 M Tris, 0.15 M sodium chloride, 0.02%
sodium azide, 1 mM phenyl methyl sulfonyl fluoride, trasylol 10
units/ml at pH 7.3, and shaking with an equal volume of
complete Freund's adjuvant. On day 14, the mice were again
20 injected with the same material except that incomplete Freund's
adjuvant was substituted for complete Freund's adjuvant. On
day 21, the injection of day 14 was repeated. On day 38, the
mice were injected with purified VIII:C only. On day 42, the
25 spleens of the mice were removed and fused according to a
standard procedure, of the type described by J. P. Brown et al
"Protein Antigens of Normal and Malignant Human Cells
30 Identified by Immunoprecipitation with Monoclonal Antibodies",
JOURNAL OF BIOLOGICAL CHEMISTRY, Vol. 225, pp. 4980-4983
(1980). The standard technique was varied only to the extent
that 35% polyethylene glycol 1000 was substituted for 50%
35 polyethylene glycol.



1 The antibodies were selected using the assay procedure
described above in Example I, in the paragraph with the heading
"Production of monoclonal antibody against VIII:C", except that
5 antibodies which did not neutralize VIII:C activity were also
subcloned and treated as described below.

 The clones which were positive were subcloned twice
and stable clones producing antibody to VIII:C were then
10 injected into the peritoneal cavities of Balb/C mice which had
been pretreated intraperitoneally with 0.5 ml of pristane at
least four days prior to injection of cells. Hybridoma cells
15 were injected at concentrations of approximately 5×10^6
cells per mouse in 0.5 ml of Delbecco's modified Eagle's medium
without fetal bovine serum. The mice were tapped when bloated
and ascites fluid was collected in heparin at approximately 10
20 units/ml. Ascites fluid from multiple mice was pooled to
provide a convenient volume for subsequent isolation of the
monoclonal IgG. The antibodies were precipitated from ascites
fluid using 50% ammonium sulfate, and then reprecipitated two
25 more times. The anti-VIII:C antibodies so raised corresponding
to (B), (C) and (D) above were bound to agarose beads and shown
to bind purified VIII:C from solution.

30 Separate batches of VIII:C, one untreated and one
subjected to thrombin proteolysis, were then analyzed by the
SDS-PAGE steps described in Example 1. Then, each band was
transferred electrophoretically (Western transfers) from the
35 gel onto a nitrocellulose sheet. The apparatus used was



1 Bio-Rad "Trans-Blot" cell, and Bio-Rad Model 160.1.6 power
supply (Bio-Rad Laboratories, Richmond, California). The
transfer buffer was 25 mM Tris with glycine added to pH 8.3,
5 20% methanol. Transfers were carried out over 16-24 hours at
90 volts and 100 milliamperes.

The specific polypeptides with which each of the
antibodies reacted were determined using a procedure adapted
10 from W. M. Burnette, "Western Blotting: Electrophoretic
Transfer of Proteins from Sodium Dodecyl Sulfate Polyacrylamide
Gels to Unmodified Nitrocellulose and Radiographic Detection
15 with Antibodies and Radioiodinated Protein A", Analytical
Biochemistry, Vol. 112, pp. 195-203 (1981).

"Buffer D" was 10 mM Tris chloride, 0.15 M NaCl, 0.02%
sodium azide, pH 7.4. All reactions were carried out at room
20 temperature.

1. Place the nitrocellulose sheet with the
transferred protein in a tray containing 100 ml of Buffer D and
0.25% gelatin. Place the tray on a rotary shaker and shake
25 slowly for 30 minutes.

2. Then add the monoclonal antibody to the tray
(either 0.1% - 1% of ascites fluid or 1 mg of purified IgG).
30 Shake for 120 minutes.

3. Wash the nitrocellulose sheet as follows:
(a) 10 minutes with 100 ml of Buffer D.
(b) 30 minutes with 100 ml of Buffer D and 0.05 % of
35 Nonidet-P-40, with changes at 10 and 20 minutes.



1 (c) 10 minutes with 100 ml of Buffer D.

4. Soak the nitrocellulose sheet in Buffer D plus
0.25% gelatin and I^{125} -labelled purified rabbit anti-mouse
5 IgG for 30 minutes.

5. Wash the nitrocellulose sheet as follows:

(a) 10 minutes with 100 ml of Buffer D.

10 (b) 16-24 hours with 100 ml of Buffer D plus 0.1%
Nonidet P-40 and 0.5 M NaCl.

(c) 10 minutes with 100 ml of Buffer D.

6. Blot the nitrocellulose sheet between two sheets
15 of filter paper and then store the nitrocellulose sheet in a
sealed plastic bag.

7. To determine which antibodies and polypeptides
had reacted,

20 (a) Prepare an autoradiograph of the nitrocellulose
sheet, using standard procedures known in the literature.

(b) Compare the autoradiograph with a nitrocellulose
25 sheet onto which VIII:C was transferred but which had been
stained with Coomassie blue R250 rather than reacted with
monoclonal antibody.

These steps demonstrated that the following distinct
30 antibodies had been raised.

Four antibodies reacted with the polypeptide of
 $M_r=92,000$, the $M_r=108,000$ and larger polypeptides, one or
the other of the polypeptides of $M_r=54,000$ or $44,000$ present
35 in terminal thrombin digests, and no other polypeptides. This

This Sheet Has Been Replaced



1 demonstrates the origin of these two latter polypeptides from
the polypeptide of $M_r=92,000$ as the result of thrombin
cleavage. It also shows that two epitopes on the $M_r=92,000$
5 polypeptide survive that cleavage, and that these epitopes are
not present on the $M_r = 79,000-80,000$ doublet.

A fifth antibody reacted with the doublet of $M_r =$
79,000-80,000, and with polypeptides of $M_r=108,000$ and
10 greater, but not with the polypeptide of $M_r=92,000$ nor with
any of the polypeptides present in terminal thrombin digests.
This demonstrates that the $M_r=79,000-80,000$ doublet possesses
15 an epitope which is not present on the $M_r=92,000$ polypeptide;
and that the epitope is destroyed by thrombin digestion of the
 $M_r = 79,000-80,000$ doublet.

The reaction profiles of these five antibodies
20 indicate that the $M_r=92,000$ and $M_r = 79,000-80,000$
polypeptides derive from a common precursor or precursors.

A sixth antibody reacted only with polypeptides of
25 $M_r=108,000$ and greater. Since it did not react with any
polypeptides present in terminal thrombin digests, the epitope
with which it reacted is destroyed by thrombin digestion.

To prepare and store a biological preparation of one
30 or more of these antibodies, the corresponding monoclonal IgG
may be isolated from heparinized pooled ascites fluid
immediately after collection or a frozen portion of the stored
solution may be thawed. Regardless of whether fresh or frozen
35



1 material is used, the solution is brought to 4°C and treated
with an equal volume of phosphate buffered saline solution
(PBS) (PBS: 1.6 g sodium phosphate, monobasic monohydrate; 8.4
5 g sodium phosphate, dibasic anhydrous; 61.4 g sodium chloride;
water to 7 liters; pH=7.2). The diluted ascites is
precipitated by dropwise addition with stirring at 4°C.
Centrifugations are preferably carried out at 14,000 rpm for 60
10 minutes (30,000 x g). The supernatant solution of ascites is
precipitated twice more with SAS and the mixture of precipitate
and supernatant liquid stirred and centrifuged in the same
15 manner as in the first cycle. The pellets resulting from the
third precipitation are resuspended in a volume of PBS equal to
that of the diluted ascites fluid and then dialyzed
exhaustively against PBS. Clots appearing in the dialysis bags
20 are removed by centrifugation at 20°C. The dialyzed IgG is
adsorbed by stirring it with a 5% aqueous solution of aluminum
hydroxide at room temperature and centrifuging at 20°C after
(adsorption. The adsorption treatment is repeated at least
25 three more times using 2.5% aluminum hydroxide solution for
each treatment after the first. The adsorbed IgG is brought to
4°C and reprecipitated once with SAS as described above. The
30 precipitated pellets may be stored at -20°C until used.

Two preferred procedures for purifying the monoclonal
antibodies and maintaining biological preparations containing
them are described in P. L. Ey et al., "Isolation of Pure



- 1 IgG₁, IgG_{2a}, and IgG_{2b} Immunoglobulins from Mouse Serum
Using Protein A-Sepharose." Immunochemistry, Vol. 15, pp.
429-436; and in C. Bruck et al., "One-Step Purification of
5 Mouse Monoclonal Antibodies from Ascitic Fluid by DEAE Affi-Gel
Blue Chromatography." J. Immunological Methods, Vol. 53, pp.
313-319 (1982).

10

15

20

25

30

35

This Sheet Has Been Replaced



Claims

1. A monoclonal antibody of class IgG which reacts with human Factor VIII:C polypeptides selected from the group consisting of:

(A) polypeptides of $M_r = 44,000$; $92,000$; $108,000$; and undigested Factor VIII:C; and not polypeptides of $M_r = 54,000$; $71,000$; $72,000$; $79,000$; or $80,000$;

(B) polypeptides of $M_r = 54,000$; $92,000$; $108,000$; and undigested Factor VIII:C; and not with polypeptides of $M_r = 44,000$; $71,000$; $72,000$; $79,000$; or $80,000$; and

(C) polypeptides of $M_r = 79,000$ and $80,000$; $108,000$; and undigested Factor VIII:C; and not with polypeptides of $M_r = 44,000$; $54,000$; $71,000$; $72,000$; or $92,000$.

2. A biological preparation comprising the antibody according to claim 1 which exhibits reaction profile A, said preparation is essentially free of antibodies which exhibit reaction profiles B and C.

3. A biological preparation comprising the antibody according to claim 1 which exhibits reaction profile B, said preparation is essentially free of antibodies which exhibit reaction profiles A and C.





4. A biological preparation comprising the antibody according to claim 1 which exhibits reaction profile C, said preparation is essentially free of antibodies which exhibit reaction profiles A and B.

5. A biological preparation comprising a mixture of antibodies according to claim 1 which exhibit reaction profiles A and B and is essentially free of antibodies which exhibit reaction profile C.

6. A biological preparation comprising a mixture of antibodies according to claim 1 which exhibit reaction profiles A, B and C.

7. A monoclonal antibody according to claim 1 having reaction profile B or C, further characterized in that said antibody neutralizes human Factor VIII:C coagulant activity when said antibody is bound to Factor VIII:C.

8. A hybridoma that secretes a monoclonal antibody according to claim 1(A).

9. A hybridoma that secretes a monoclonal antibody according to claim 1(B).

10. A hybridoma that secretes a monoclonal antibody according to claim 1(C).





11. A human Factor VIII:C monoclonal antibody produced in response to immunization with a Factor VIII:C protein composition having a Factor VIII:C protein doublet with a molecular weight of 80,000 and 79,000 complexed with a 92,000 protein, said composition exhibiting a biological activity of Factor VIII:C, and wherein said antibody specifically binds to at least one polypeptide derived from said composition.

12. Monoclonal antibodies prepared in response to an immunogen and specifically binding to said immunogen, said immunogen consisting of a protein composition substantially free of other proteins and consisting of a complex of an 80,000 and 79,000 protein calcium bridged with a 92,000 protein exhibiting a biological activity of Factor VIII:C.

13. A monoclonal antibody of class IgG which reacts with human Factor VIII:C polypeptides of $M_r = 108,000$ and undigested Factor VIII:C; and not polypeptides of $M_r =$ less than 108,000.

14. A hybridoma that secretes a monoclonal antibody according to claim 13.





15. An immunoabsorbent column comprising particles of solid immunologically inert adsorbent material bound to an antibody according to claim 1 or 13.

16. A column according to claim 15 wherein the adsorbent material is agarose.

