Syndrome (AIDS), adult respiratory distress syndrome, and ataxia telengiectasia, said composition comprising an excipient and a therapeutically effective amount of a compound of claim 1, or pharmaceutically acceptable salts thereof.

17. ~~A method of treating hypertension, chronic and congestive heart failure, cardiac hypertrophy, restenosis, chronic renal failure, cerebral vasospasm after subarachnoid bleeding, pulmonary hypertension, atherosclerosis, asthma, male erectile dysfunctions, female sexual dysfunction, over-active bladder syndrome, spinal-cord injury, traumatic brain injury, stroke, inflammatory and demyelinating diseases, Alzheimer's disease, pain, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, inflammatory pain, rheumatoid arthritis, osteoarthritis, irritable bowel syndrome, Crohn's disease, psoriasis, ulcerative colitis, lupus, inflammatory bowel disease, cancer, tumor metastasis, viral infections, bacterial infections, insulin resistance, diabetes, ocular hypertension, preterm labor, atherosclerosis, spinal muscular atrophy, encephalomyelitis, osteoporosis, HIV-1, encephalitis, ischemic CNS disorders, vascular or AD type dementia, glaucoma, retinopathy, benign prostatic hypertrophy, psychiatric disorders including depression, schizophrenia, obsessive compulsive disorder and bipolar disorder, epilepsy and seizure disorders, for decreasing ischemia-reperfusion injury, myocardial infarct size and myocardial fibrosis, for the prevention of graft failure, and cystic fibrosis, in a patient, said method comprising administering to the patient~~ Use of a therapeutically effective amount of a compound of claim 1, or pharmaceutically acceptable salts thereof, in the manufacture of a medicament for treating hypertension, chronic and congestive heart failure, cardiac hypertrophy, restenosis, chronic renal failure, cerebral vasospasm after subarachnoid bleeding, pulmonary hypertension, atherosclerosis, asthma, male erectile dysfunctions, female sexual dysfunction, over-active bladder syndrome, spinal-cord injury, traumatic brain injury, stroke, inflammatory and demyelinating diseases, Alzheimer's disease, pain, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, inflammatory pain, rheumatoid arthritis, osteoarthritis, irritable bowel syndrome, Crohn's disease, psoriasis, ulcerative colitis, lupus, inflammatory bowel disease, cancer, tumor metastasis, viral infections, bacterial infections, insulin resistance, diabetes, ocular hypertension, preterm labor, atherosclerosis, spinal muscular atrophy, encephalomyelitis, osteoporosis, HIV-1, encephalitis, ischemic CNS disorders, vascular or AD type dementia, glaucoma, retinopathy, benign prostatic hypertrophy, psychiatric disorders including depression, schizophrenia, obsessive compulsive disorder and bipolar disorder, epilepsy and seizure disorders, for decreasing ischemia-reperfusion injury, myocardial infarct size and myocardial fibrosis, for the prevention of graft failure, and cystic fibrosis, in a patient.

18. ~~A method of treating inflammatory and tissue repair disorders including rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease), osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultra-violet induced skin damage; autoimmune diseases including systemic lupus erythematosis, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, athersclerosis, restenosis, diabetes, glomerulonephritis, cancer, including breast cancer, prostate cancer, lung cancer, colon cancer, cervix cancer, ovarian cancer, skin cancer, CNS cancer, bladder cancer, pancreatic cancer, leukemia, lymphoma or Hodgkin's disease, cachexia, inflammation associated with infection and viral infections, including Acquired Immune Deficiency Syndrome (AIDS), adult respiratory distress syndrome, and ataxia telengiectasia in a patient, said method comprising administering to the patient~~ Use of a therapeutically effective amount of a compound of claim 1, or pharmaceutically acceptable salts thereof, in the manufacture of a medicament for treating inflammatory and tissue repair disorders including rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease), osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultra-violet induced skin damage; autoimmune diseases including systemic lupus erythematosis, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, athersclerosis, restenosis, diabetes, glomerulonephritis, cancer, including breast cancer, prostate cancer, lung cancer, colon cancer, cervix cancer, ovarian cancer, skin cancer, CNS cancer, bladder cancer, pancreatic cancer, leukemia, lymphoma or Hodgkin's disease, cachexia, inflammation associated with infection and viral infections, including Acquired Immune Deficiency Syndrome (AIDS), adult respiratory distress syndrome, and ataxia telengiectasia in a patient.

19. ~~A method of treating hypertension, chronic and congestive heart failure, cardiac hypertrophy, restenosis, chronic renal failure, cerebral vasospasm after subarachnoid bleeding, pulmonary hypertension, atherosclerosis, asthma, male erectile dysfunctions, female sexual dysfunction, over-active bladder syndrome, spinal-cord injury, traumatic brain injury, stroke, inflammatory and demyelinating diseases, Alzheimer's disease, pain, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, inflammatory pain, rheumatoid arthritis, osteoarthritis, irritable bowel syndrome, Crohn's disease, psoriasis, ulcerative colitis, lupus, inflammatory bowel disease, cancer, tumor metastasis, viral infections, bacterial infections, insulin resistance, diabetes, ocular hypertension, preterm labor, atherosclerosis, spinal muscular atrophy, encephalomyelitis, osteoporosis, HIV-1, encephalitis, ischemic CNS disorders, vascular or AD type dementia, glaucoma, retinopathy, benign prostatic hypertrophy, psychiatric disorders, including depression, schizophrenia, obsessive compulsive disorder and bipolar disorder, epilepsy and seizure disorders, for decreasing ischemia-reperfusion injury, myocardial infarct size and myocardial fibrosis, for the prevention of graft failure, and cystic fibrosis in a patient, said method comprising administering to the patient therapeutically effective amount of the compound of claim 1, or pharmaceutically acceptable salts thereof; and~~ Use of a therapeutically effective amount of one additional therapeutic agent or more than one additional therapeutic agent, in the manufacture of a medicament for treating hypertension, chronic and congestive heart failure, cardiac hypertrophy, restenosis, chronic renal failure, cerebral vasospasm after subarachnoid bleeding, pulmonary hypertension, atherosclerosis, asthma, male erectile dysfunctions, female sexual dysfunction, over-active bladder syndrome, spinal-cord injury, traumatic brain injury, stroke, inflammatory and demyelinating diseases, Alzheimer's disease, pain, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, inflammatory pain, rheumatoid arthritis, osteoarthritis, irritable bowel syndrome, Crohn's disease, psoriasis, ulcerative colitis, lupus, inflammatory bowel disease, cancer, tumor metastasis, viral infections, bacterial infections, insulin resistance, diabetes, ocular hypertension, preterm labor, atherosclerosis, spinal muscular atrophy, encephalomyelitis, osteoporosis, HIV-1, encephalitis, ischemic CNS disorders, vascular or AD type dementia, glaucoma, retinopathy, benign prostatic hypertrophy, psychiatric disorders, including depression, schizophrenia, obsessive compulsive disorder and bipolar disorder, epilepsy and seizure disorders, for decreasing ischemia-reperfusion injury, myocardial infarct size and myocardial fibrosis, for the prevention of graft failure, and cystic fibrosis in a patient.

20. ~~A method of treating inflammatory and tissue repair disorders including rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease), osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultra-violet induced skin damage; autoimmune diseases including systemic lupus erythematosis, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherselerosis, restenosis, diabetes, glomerulonephritis, cancer, including breast cancer, prostate cancer, lung cancer, colon cancer, cervix cancer, ovarian cancer, skin cancer, CNS cancer, bladder cancer, pancreatic cancer, leukemia, lymphoma or Hodgkin's disease, cachexia, inflammation associated with infection and certain viral infections, including Acquired Immune Deficiency Syndrome (AIDS), adult respiratory distress syndrome, and ataxia telengiectasia or spleen cancer in a patient, said method comprising administering to the patient~~ Use of a therapeutically effective amount of the compound of claim 1, or pharmaceutically acceptable salts thereof; and a therapeutically effective amount of one additional therapeutic agent or more than one additional therapeutic agent, in the manufacture of a medicament for treating inflammatory and tissue repair disorders including rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease), osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultra-violet induced skin damage; autoimmune diseases including systemic lupus erythematosis, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherselerosis, restenosis, diabetes, glomerulonephritis, cancer, including breast cancer, prostate cancer, lung cancer, colon cancer, cervix cancer, ovarian cancer, skin cancer, CNS cancer, bladder cancer, pancreatic cancer, leukemia, lymphoma or Hodgkin's disease, cachexia, inflammation associated with infection and certain viral infections, including Acquired Immune Deficiency Syndrome (AIDS), adult respiratory distress syndrome, and ataxia telengiectasia or spleen cancer in a patient.