

COMPOSITIONS OF
1-(4-BROMO-5-(1-ETHYL-7-(METHYLAMINO)-2-OXO-1,2-DIHYDRO-1,6-NAPHTHYRIDIN-3-YL)-2-FLUOROPHENYL)-3-PHENYLUREA

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S.S.N. 62/955,073 filed December 30, 2019,
5 U.S.S.N. 62/955,062 filed December 30, 2019, U.S.S.N. 62/968,695 filed January 31, 2020,
and U.S.S.N. 62/968,724 filed January 31, 2020, the contents of each of which are
incorporated herein by reference.

BACKGROUND

[0002] c-KIT (also known as KIT, CD117, and stem cell factor receptor) is a 145 kDa
10 transmembrane tyrosine kinase protein that acts as a type-III receptor. The c-KIT proto-
oncogene, located on chromosome 4q11-21, encodes the c-KIT receptor, whose ligand is the
stem cell factor (SCF), steel factor, kit ligand, and mast cell growth factor. The receptor has
tyrosine-protein kinase activity and binding of the ligand SCF leads to the
autophosphorylation of c-KIT and its association with substrates such as phosphatidylinositol
15 3-kinase (PI3K). Tyrosine phosphorylation by protein tyrosine kinases is of particular
importance in cellular signaling and can mediate signals for major cellular processes, such as
proliferation, survival, differentiation, apoptosis, attachment, invasiveness and migration.
Defects in c-KIT are a cause of piebaldism, an autosomal dominant genetic developmental
abnormality of pigmentation characterized by congenital patches of white skin and hair that
20 lack melanocytes. Gain-of-function mutations of the c-KIT gene and the expression of
constitutively phosphorylated c-KIT are found in most gastrointestinal stromal tumors (GIST)
and mastocytosis. Further, almost all gonadal seminomas/dysgerminomas exhibit c-KIT
membranous staining, and several reports have clarified that some (10-25%) have a c-KIT
gene mutation. c-KIT defects have also been associated with testicular tumors including germ
25 cell tumors (GCT) and testicular germ cell tumors (TGCT). C-KIT mutations also have been
associated with a subset of cutaneous or acral melanoma.

[0003] Oncogenic genomic alterations of PDGFR α kinase or overexpression of PDGFR α
kinase have been shown to be causative of human cancers. Missense mutations of PDGFR α
kinase have been shown to be causative of a subset of GISTs. PDGFR α mutations are
30 oncogenic drivers in approximately 8-10% of GISTs. The predominant PDGFR α mutation is
exon 18 D842V, although other exon 18 mutations including D846Y, N848K, and Y849K,
and exon 18 insertion-deletion mutations (INDELs) including RD841-842KI, DI842-843-IM,

and HDSN845-848P have also been reported. Furthermore, rare mutations in PDGFR α exons 12 and 14 have also been reported. The PDGFR α exon 18 deletion mutations Δ D842-H845 and Δ I843-D846 have been reported in GIST. Amplification or mutations of PDGFR α have been described in human tissues of malignant peripheral nerve sheath tumors (MPNST).

5 Amplification of PDGFR α has been described in multiple skin lesions of undifferentiated pleomorphic sarcoma and in intimal sarcoma. Amplification of PDGFR α has been linked to a subset of lung cancer patients. 4q12, containing the PDGFR α gene locus, is amplified in 3–7% of lung adenocarcinomas and 8–10% of lung squamous cell carcinomas. PDGFR α amplification is common in pediatric and adult high- grade astrocytomas and identified a
10 poor prognostic group in IDH1 mutant glioblastoma. PDGFR α amplification was frequent in pediatric (29.3%) and adult (20.9%) tumors. PDGFR α amplification was reported to increase with grade and in particular to be associated with a less favorable prognosis in IDH1 mutant de novo GBMs. The PDGFR α locus in PDGFR α -amplified gliomas has been demonstrated to present a PDGFR α exon 8,9 intragenic deletion rearrangement. This intragenic deletion
15 was common, being present in 40% of the glioblastoma multiformes (GBMs) presenting with PDGFR α amplification. Tumors with this rearrangement displayed histologic features of oligodendroglioma, and the PDGFR α exon 8,9 intragenic deletion showed constitutively elevated tyrosine kinase activity. The FIP1L1-PDGFR α fusion protein is oncogenic in a subset of patients with hypereosinophilic syndrome. FIP1L1- PDGFR α fusion has also been
20 identified in eosinophilia-associated acute myeloid leukemia and lymphoblastic T-cell lymphoma.


[0004] Such a broad-spectrum c-KIT inhibitor, and formulations thereof, would be of high therapeutic value in the treatment of refractory GIST patients and those suffering from other disorders. There is a need for oral formulations that provide significantly stable
25 products to patients. Mutations, deletions, rearrangements, and amplification of the PDGFR α gene are linked to a number of solid and hematological cancers. Given the complex function of the PDGFR α gene and the potential utility for PDGFR α inhibitors in the treatment of various solid and hematological cancers, there is a need for oral formulations of inhibitors with good therapeutic properties.

30

SUMMARY


[0005] Provided herein, in part, are compositions comprising a compound of Formula (I) with the purity and safety aspects to be considered for pharmaceutical preparations. A

compound of Formula (I) as described herein has the following structure:



Formula (I)


[0006] Provided herein, in part, are pharmaceutical compositions comprising a compound
5 of Formula (I):



Formula (I)

and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the
pharmaceutical composition comprises one or more anilinic substances, each in an amount
10 equal to or less than 3.0% by weight based on the weight of the compound of Formula (I).

[0007] Provided herein, in part, are pharmaceutical compositions comprising a compound
of Formula (I):




Formula (I)

15

and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the

pharmaceutical composition comprises one or more anilinic substances and a compound represented by Formula (III):




Formula (III)

- 5 each in an amount equal to or less than 3.0% by weight based on the weight of the compound of Formula (I).

[0008] In some embodiments, provided herein is the solid dispersion comprising a compound of Formula (I) and a polymer, wherein the pharmaceutical composition comprises one or more anilinic substances, each in an amount equal to or less than 3.0% by weight
10 based on the weight of the compound of Formula (I).

[0009] In some embodiments, provided herein is a pharmaceutical composition comprising the compound of Formula (I) and a pharmaceutically acceptable carrier, wherein the composition has less than 3% w/w of each of: 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methyamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)), aniline
15 and diphenyl urea.

[0010] In an embodiment, provided herein is a pharmaceutical composition comprising the compound of Formula (I) and a pharmaceutically acceptable carrier, wherein the composition has less than 3% w/w of each of: 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methyamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)), a compound
20 represented by Formula (III):



Formula (III)

aniline and diphenyl urea.

[0011] In an embodiment, described herein is a substantially purified compound represented by Formula (I) having less than about 3.0% by weight of an impurity selected from the group consisting of the compound of Formula (II), aniline, and combinations thereof. In some embodiments, the compound comprises less than 0.5% of the impurity.

5 [0012] In an embodiment, described herein is a substantially purified compound represented by Formula (I) having less than about 3.0% by weight of an impurity selected from the group consisting of the compound of Formula (II), the compound of Formula (III), and combinations thereof. In some embodiments, the compound comprises less than 0.5% of the impurity.


10 [0013] In an embodiment, described herein is a high purity compound represented by the compound of Formula (I) having less than about 3.0% of anilinic substance impurities. In some embodiments, the compound comprises less than 0.5% of each of one or more anilinic substances.

[0014] In another embodiment, provided herein is a high purity compound represented by
15 the compound of Formula (I) having less than about 0.05% of a diphenyl urea impurity.

[0015] Described herein, in an embodiment, is a pharmaceutical composition comprising the of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition has less than about 0.05% of diphenyl urea.

[0016] Described herein, in an embodiment, is a pharmaceutical composition comprising the
20 of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition has less than about 0.05% of diphenyl urea as detected by HPLC.


[0017] Described herein, in an embodiment, is a pharmaceutical composition comprising: a)
a solid dispersion comprising: a compound represented by Formula (I) or pharmaceutically
25 acceptable salt thereof,



Formula (I)


and a pharmaceutically acceptable polymer; and b) one or more pharmaceutically acceptable carriers, wherein no single anilinic substance is present in an amount equal to or less than about 3.0% by weight.

- [0018] Described herein, in an embodiment, pharmaceutical composition comprising: a) a
5 solid dispersion comprising: an amorphous form of a compound represented by Formula (I)



Formula (I)


and a pharmaceutically acceptable polymer; and b) one or more pharmaceutically acceptable carriers; wherein each of: a compound represented by Formula (II):



Formula (II)

aniline and diphenyl urea are present in amount equal to or less than about 1.5% by weight based on the total weight of the composition.


- [0019] Described herein, in an embodiment, is a pharmaceutical composition comprising: a)
15 a solid dispersion comprising: an amorphous form of a compound represented by Formula (I)



Formula (I)


- and a pharmaceutically acceptable polymer; and b) one or more pharmaceutically acceptable carriers; wherein diphenyl urea is present in amount equal to or less than about 0.5% by
20 weight based on the total weight of the composition.

[0020] Described herein, in an embodiment, is a pharmaceutically acceptable composition for a 50 mg single dosage unit of a compound represented by:




comprising: an intragranular blend, wherein the intragranular blend comprises: a solid dispersion having 50 mg of the compound wherein the compound is present in amorphous form, hydroxypropyl methyl cellulose acetate succinate; a bulking agent and/or filler; and a lubricant and/or a glidant; and an extragranular blend comprising a glidant and/or a lubricant; wherein diphenyl urea is present in amount equal to or less than about 0.5% by weight based on the total weight of the composition.

10 [0021] Described herein, in an embodiment, is a pharmaceutically acceptable composition for orally delivering to a patient 50 mg of a compound represented by:



comprising: a solid dispersion having: 50 mg of the compound wherein the compound is present in amorphous form; hydroxypropyl methyl cellulose acetate succinate; a bulking agent, a filler, and a lubricant and/or a glidant; wherein diphenyl urea is present in amount equal to or less than about 0.5% by weight based on the total weight of the composition.

[0022] Described herein, in an embodiment, is a pharmaceutically acceptable composition for orally delivering 50 mg of a compound represented by:




comprising:

a solid dispersion having 50 mg of the compound wherein the compound is present in amorphous form, and a polymer; about 25-45% by weight of a bulking agent based on the total amount of the pharmaceutical composition; about 25-45% by weight of a filler based on the total amount of the pharmaceutical composition; and a glidant and/or a lubricant; wherein the composition has not more than about 1.5% (w/w) of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one, when exposed to 60% relative humidity at 25 °C for 1 month, 3 months or 6 months.

[0023] Described herein, in an embodiment, is a pharmaceutically acceptable tablet

having 50 mg of a compound represented by:



wherein the tablet comprises: a solid dispersion having 50 mg of the compound wherein the compound is present in amorphous form, and hydroxypropyl methyl cellulose acetate succinate; about 25-35% by weight microcrystalline cellulose based on the total weight of the tablet; and about 25-35% by weight of lactose or a hydrate thereof based on the total amount of the pharmaceutical composition; wherein diphenyl urea is present in amount equal to or less than about 0.5% by weight based on the total weight of the composition.

[0024] In an embodiment, provided herein is a pharmaceutical composition comprising:

(a) an intragranular blend comprising: (i) a solid dispersion comprising a compound represented by Formula (I), wherein the pharmaceutical composition is comprises one or more anilinic substance, each in an amount equal to or less than 3% by weight based on the weight of the compound of Formula (I), and a pharmaceutically acceptable polymer; (ii) one or more fillers; (iii) a disintegrant; (iv) a glidant; and (v) a lubricant; and (b) an extragranular blend comprising: (i) a glidant; and (ii) a lubricant. In some embodiments, each anilinic substance is present in an amount equal to or less than about 5.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 4.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 2.0% by weight based on the weight of the compound of

Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 1.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.7% by weight based on the weight of the compound of Formula (I). In some
5 embodiments, each anilinic substance is present in an amount equal to or less than about 0.5% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.3% by weight based on the weight of the compound of Formula (I).

[0025] In an embodiment, the disclosure provides a pharmaceutical composition
10 comprising: (a) an intragranular blend comprising: (i) about 33% by weight of a solid dispersion based on the total weight of the composition, the solid dispersion comprising a compound represented by Formula (I) and hydroxypropyl methyl cellulose acetate succinate, wherein the solid dispersion comprises about 25% by weight of the compound represented by Formula (I) based on the total weight of the solid dispersion; (ii) about 30% by weight of
15 microcrystalline cellulose based on the total amount of the of the composition; (iii) about 30% by weight of lactose monohydrate based on the total amount of the of the composition; (iv) about 5% by weight of crospovidone based on the total amount of the of the composition; (v) about 0.5% by weight of silicon dioxide based on the total amount of the of the composition; and (vi) about 0.5% by weight of magnesium stearate based on the total amount
20 of the of the composition; and (b) an extragranular blend comprising: (i) about 0.5% by weight of silicon dioxide based on the total amount of the of the composition; and (ii) about 0.5% by weight of magnesium stearate based on the total amount of the composition.

[0026] In an embodiment, provided herein is a pharmaceutical composition comprising:
25 (a) an intragranular blend comprising: (i) about 200 mg of a solid dispersion comprising a compound represented by Formula (I), wherein the pharmaceutical composition comprises one or more anilinic substance, each in an amount equal to or less than 3% by weight based on the weight of the compound of Formula (I) and hydroxypropyl methyl cellulose acetate succinate, wherein the solid dispersion comprises about 50 mg of the compound represented by Formula (I); (ii) about 179 mg of microcrystalline cellulose; (iii)
30 about 179 mg of lactose monohydrate; (iv) about 30 mg of crospovidone; (v) about 3 mg of silicon dioxide; and (vi) about 3 mg of magnesium stearate; and (b) an extragranular blend comprising: (i) about 3 mg of silicon dioxide; and (ii) about 3 mg of magnesium stearate. In some embodiments, each anilinic substance is present in an amount equal to or less than about 5.0% by weight based on the weight of the compound of Formula (I). In some

embodiments, each anilinic substance is present in an amount equal to or less than about 4.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 2.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 1.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.7% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.5% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.3% by weight based on the weight of the compound of Formula (I).

[0027] In an embodiment, provided herein is a tablet providing about 50 mg of a compound represented by Formula (I), wherein the tablet comprises one or more anilinic substance impurities, each in an amount equal to or less than 3% by weight based on the weight of the compound of Formula (I), wherein the tablet comprises: (a) an intragranular blend comprising: (i) about 195 mg to about 205 mg of a solid dispersion that comprises about 50 mg of the compound and hydroxypropyl methyl cellulose acetate succinate; (ii) about 177 mg to about 181 mg of microcrystalline cellulose; (iii) about 177 mg to about 181 mg of lactose monohydrate; and (iv) about 28 mg to about 32 mg of crospovidone; and (b) an extragranular blend comprising: (i) about 2 mg to about 4 mg of silicon dioxide; and (ii) about 2 mg to about 4 mg of magnesium stearate. In some embodiments, each anilinic substance is present in an amount equal to or less than about 5.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 4.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 2.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 1.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.7% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.5% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.3% by weight based on the weight of the compound of Formula (I).

[0028] In an embodiment, provided herein are methods for treating a disease caused by the kinase activity of c-KIT or PDGFRA, and oncogenic forms thereof, wherein the disease is gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma (e.g., cutaneous melanoma, noncutaneous melanoma, KIT driven melanoma or PDGFRA driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, or non-small cell lung cancer. In some embodiments, melanoma is cutaneous melanoma or noncutaneous melanoma. In some embodiments, melanoma is cutaneous melanoma. In some embodiments, cutaneous melanoma is superficial spreading melanoma, nodular melanoma, acral-lentiginous melanoma, or amelanotic and desmoplastic melanoma. In some embodiments, melanoma is noncutaneous (non-skin) melanoma. In some embodiments, noncutaneous melanoma is ocular melanoma or mucosal melanoma.

[0029] Also provided herein, in another embodiment, is a method of treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), NF-1-deficient gastrointestinal stromal tumors, succinate dehydrogenase (SDH)-deficient gastrointestinal stromal tumors, KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma, acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein.

[0030] Also provided herein, in another embodiment, is a method of treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, lung cancer, glioblastoma, a glioma, malignant peripheral nerve sheath sarcoma, and hypereosinophilic syndrome in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein.

[0031] Also provided herein, in another embodiment, is a method of treating a disease selected from the group consisting of KIT driven germ cell tumor (e.g., testicular germ cell), KIT driven skin cancer, or KIT driven renal cell carcinoma in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein.

[0032] Also provided herein, in another embodiment, is a method of treating a disease selected from the group consisting of penile cancer, PDGFRA driven penile cancer, prostate cancer, PDGFRA driven prostate cancer, PDGFRA driven non-melanoma skin cancer, PDGFRA driven glioma, PDGFRA driven sarcoma, PDGFRA driven glioblastoma, or PDGFRA driven pancreatic cancer in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition or one or more tablets described herein.

[0033] Also provided herein, in another embodiment, is a method of treating a disease comprising a PDGFRB mutation selected from the group consisting of vaginal cancer, prostate cancer, penile cancer, non-melanoma skin cancer, melanoma, or breast sarcoma in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein.

[0034] In some embodiments, provided herein is a method for treating diseases driven by KIT mutations or PDGFRA mutations in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition or one or more tablets described herein. In some embodiments, provided herein is a method for treating diseases driven by KIT mutations and PDGFRA mutations in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition or one or more tablets described herein. In some embodiments, provided herein is a method for treating diseases driven by KIT mutations or PDGFRA mutations, comprising passenger PDGFRB mutations in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition or one or more tablets described herein. In some embodiments, provided herein is a method for treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma (e.g., KIT driven melanoma or PDGFRA driven melanoma or PDGFR driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric

glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer in a patient in need thereof, comprising

5 administering to the patient a therapeutically effective amount of a composition or one or more tablets described herein. In some embodiments, the melanoma is cutaneous melanoma or noncutaneous melanoma. In some embodiments, the melanoma is cutaneous melanoma. In some embodiments, the cutaneous melanoma is superficial spreading melanoma, nodular melanoma, acral-lentiginous melanoma, or amelanotic and desmoplastic melanoma. In some

10 embodiments, the melanoma is noncutaneous (non-skin) melanoma. In some embodiments, the noncutaneous melanoma is ocular melanoma or mucosal melanoma. In some embodiments, the disease is caused by the kinase activity of c-KIT and/or PDGFRA, and/or oncogenic forms thereof. In some embodiments, the disease is selected from the group consisting of KIT driven germ cell tumor (e.g., testicular germ cell), KIT driven skin cancer

15 (e.g., KIT driven cutaneous squamous cell carcinoma, KIT driven Merkel cell carcinoma, uveal melanoma, non-melanoma skin cancer), or KIT driven renal cell carcinoma (e.g., renal cell carcinoma, chromophobe renal cell carcinoma). In some embodiments, the disease is selected from the group consisting of penile cancer, PDGFRA driven penile cancer, prostate cancer, PDGFRA driven prostate cancer, PDGFRA driven non-melanoma skin cancer,

20 PDGFRA driven glioma, PDGFRA driven sarcoma, PDGFRA driven glioblastoma, or PDGFRA driven pancreatic cancer. In some embodiments, the disease comprising a PDGFRB mutation is selected from the group consisting of vaginal cancer, prostate cancer, penile cancer, non-melanoma skin cancer, melanoma, or breast sarcoma.


[0035] Also provided herein, in another embodiment, is a use of a composition or tablets

25 described herein for the preparation of a medicament for the treatment of a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma, acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma,

30 pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer.


[0036] In another embodiment, described herein is a process for the preparation of the

solid dispersion comprising a compound of Formula (I)



Formula (I)

e.g., a solid dispersion described herein, the process comprising: (a) mixing the compound of
 5 Formula (I), a solvent, the polymer and water to obtain a suspension; (b) optionally agitating
 and/or mixing the suspension while maintaining a temperature of about 10 to about 25 °C; (c)
 heating the suspension to dissolve the suspended particles prior to introduction into a spray-
 dryer; and (d) spray-drying the suspension to obtain a spray-dried dispersion; (e) drying the
 spray-dried dispersion; thereby obtaining the solid dispersion. In some embodiments, a solid
 10 dispersion comprising a compound of Formula (I)



Formula (I)

e.g., a solid dispersion described herein, is produced by said process.


DETAILED DESCRIPTION

15 **[0037]** The features and other details of the disclosure will now be more particularly
 described. Certain terms employed in the specification, examples and appended claims are
 collected here. These definitions should be read in light of the remainder of the disclosure and
 as understood by a person of skill in the art. Unless defined otherwise, all technical and
 scientific terms used herein have the same meaning as commonly understood by a person of
 20 ordinary skill in the art.

Definitions

[0038] As used herein, the term “excipient” refers to a substance that may be beneficial to include in a composition with an active agent. The term “excipient” includes inert substances as well as functional excipients that may result in beneficial properties of the composition. Exemplary excipients include but are not limited to polymers, glidants, sugars, lubricant, salts, buffers, fats, fillers, disintegrating agents, binders, surfactants, high surface area substrates, flavorants, carriers, matrix materials, and so forth.

[0039] As used herein, the terms “Anilinic impurity A,” “Impurity A,” and Compound 2 each refer to the compound 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one, the structure of which is the compound of Formula (II):



Formula (II)

In some embodiments, an anilinic substance may be Impurity A.

[0040] As used herein, the terms “Anilinic impurity B,” “Impurity B” refer to aniline. In some embodiments, an anilinic substance may be Impurity B.

[0041] As used herein, the terms “Anilinic substances,” “anilinic substance impurity,” “anilinic substance impurities” are alone or together and may include but are not limited to 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (compound of Formula (II)) or aniline.

[0042] As used herein, the terms “Anilinic impurities,” “anilinic impurity,” “anilinic substance impurity,” “anilinic substance impurities” are alone or together and may include but are not limited to 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (compound of Formula (II)) or aniline.

[0043] As used herein, the terms “Individual,” “patient,” or “subject” are used interchangeably and include any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans. The compounds described herein can be administered to a mammal, such as a human, but can also be administered to other mammals such as an animal in need of

veterinary treatment, *e.g.*, domestic animals (*e.g.*, dogs, cats, and the like), farm animals (*e.g.*, cows, sheep, pigs, horses, and the like) and laboratory animals (*e.g.*, rats, mice, guinea pigs, and the like).

5 [0044] As used herein, the terms “Pharmaceutically acceptable” or “pharmacologically acceptable” includes molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate. For human administration, preparations should meet sterility, pyrogenicity, and general safety and purity standards as required by FDA Office of Biologics standards.

10 [0045] As used herein, the term “pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” as used herein refers to any and all solvents, dispersion media, coatings, isotonic and absorption delaying agents, and the like, that are compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. The compositions may also contain other active compounds providing supplemental, additional, or enhanced therapeutic
15 functions.

[0046] As used herein, the term “pharmaceutical composition” as used herein refers to a composition comprising at least one compound as disclosed herein formulated together with one or more pharmaceutically acceptable carriers, excipients or diluents.

20 [0047] As used herein, the term “therapeutically effective amount” means the amount of the subject compound that will elicit the biological or medical response of a tissue, system or animal, (*e.g.*, mammal or human) that is being sought by the researcher, veterinarian, medical doctor or other clinician. The compounds described herein are administered in therapeutically effective amounts to treat a disorder.

25 [0048] As used herein, the term “treating” includes any effect, *e.g.*, lessening, reducing, modulating, or eliminating, that results in the improvement of the condition, disease, disorder and the like.

[0049] As used herein, the term “active agent” means a drug, medicament, pharmaceutical, therapeutic agent, for example, a compound of Formula (I) as described herein.

30 [0050] As used herein, the term “oral formulation” as used herein, refers to a composition or medium used to administer a compound as disclosed herein (*e.g.*, a compound of Formula (I)) to a subject in need thereof by oral administration. Typically, an oral formulation is


administered via the mouth, however, "oral formulation" as used herein is intended to cover any substance which is administered to a subject and is absorbed across a membrane, e.g., a mucosal membrane, of the gastrointestinal tract, including, e.g., the mouth, esophagus, stomach, small intestine, large intestine, and colon. In one embodiment, the oral formulation is a solid oral formulation. In one embodiment, the oral formulation is a solid oral formulation administered to a subject in need thereof via the mouth.

[0051] A "combination therapy" is a treatment that includes the administration of two or more therapeutic agents, e.g., a compound of Formula I and a MAPKAP pathway inhibitor, to a patient. The two or more therapeutic agents may be delivered at the same time, e.g., in separate pharmaceutical compositions or in the same pharmaceutical composition, or they may be delivered at different times. For example, they may be delivered concurrently or during overlapping time periods, and/or one therapeutic agent may be delivered before or after the other therapeutic agent(s). Treatment with a combination therapy optionally includes treatment with either single agent, preceded or followed by a period of concurrent treatment with both agents. However, it is contemplated that during some time period, effective amounts of the two or more therapeutic agents are present within the patient.

[0052] All ranges recited herein include the endpoints, including those that recite a range "between" two values. Term "substantially" and "about" is to be construed as modifying a term or value such that it is not an absolute. This includes, at very least, the degree of expected experimental error, technique error and instrument error for a given technique used to measure a value.

Process


[0053] In one aspect, provided herein, is a process of preparing a composition comprising a compound of Formula (I)



Formula (I),


having one or more anilinic substances, each in an amount equal to or less than 3% by weight of the compound of Formula (I) comprising:

- (a) weighing and dispensing the compound of Formula (I), a solvent, polymer and water;
- (b) charging and suspending the compound of Formula (I);
- (c) optionally agitating and mixing the final suspension while maintaining a temperature of about 10-25 °C; and
- (d) passing the resulting suspension through an in-line heat exchanger to dissolve the suspended particles prior to introduction into the spray dryer; and
- (e) optionally drying the spray dried compound of Formula (I).
- (f) In another embodiment, described herein is a process for the preparation of the solid dispersion comprising a compound of Formula (I)



Formula (I)

- e.g., a solid dispersion described herein, the process comprising: (a) mixing the compound of Formula (I), a solvent, the polymer and water to obtain a suspension; (b) optionally agitating and/or mixing the suspension while maintaining a temperature of about 10 to about 25 °C; (c) heating the suspension to dissolve the suspended particles prior to introduction into a spray-dryer; and (d) spray-drying the suspension to obtain a spray-dried dispersion; (e) drying the spray-dried dispersion; thereby obtaining the solid dispersion. In some embodiments, heating comprises passing the suspension through an in-line heat exchanger. In some embodiments, a solid dispersion comprising a compound of Formula (I)



Formula (I)

e.g., a solid dispersion described herein, is produced by said process.

[0054] For purposes described herein, one of ordinary skill in the art would understand that anilinic substances are considered impurities in the compositions, pharmaceutical compositions, and solid dispersions as described herein. The concentration of the impurities in the composition, pharmaceutical composition or solid dispersions described herein depend
5 on the concentration of the compound of Formula (I). For example, the concentration of anilinic substances in the composition, pharmaceutical composition or solid dispersion of the inventions described herein would be expected in some embodiments, each anilinic substance is present in an amount equal to or less than about 5.0% by weight based on the weight of the compound of Formula (I) present in the composition, pharmaceutical composition or solid
10 dispersion as described herein. In some embodiments, each anilinic substance is present in an amount equal to or less than about 4.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 2.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than
15 about 1.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.7% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.5% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic
20 substance is present in an amount equal to or less than about 0.3% by weight based on the weight of the compound of Formula (I).

[0055] In some embodiments, the compound of Formula (I), solvent, polymer and water are combined, and the mixture is agitated and mixed to a suspension. In some embodiments, the solvent, water and polymer are combined and agitated prior to the addition of the
25 compound of Formula (I). In some embodiments, the solvent and water are combined and agitated prior to the addition of the polymer and the compound of Formula (I). In some embodiments, the solvent and water are combined and agitated followed by addition of the polymer, followed by addition of the compound of Formula (I).

[0056] In some embodiments, the solvent:water ratio may be about 95:5, followed by the
30 addition and dissolution of the polymer. In some embodiments, the solvent:water ratio may be about 90:10, followed by the addition and dissolution of the polymer. In some embodiments, the solvent:water ratio may be about 85:15, followed by the addition and dissolution of the polymer. In some embodiments, the solvent:water ratio may be about 80:20, followed by the addition and dissolution of the polymer. In some embodiments, the

solvent:water ratio may be about 75:25, followed by the addition and dissolution of the polymer. In some embodiments, the solvent:water ratio may be about 70:30, followed by the addition and dissolution of the polymer. In some embodiments, the solvent:water ratio may be about 65:35, followed by the addition and dissolution of the polymer. In some
5 embodiments, the solvent:water ratio may be about 60:40, followed by the addition and dissolution of the polymer. In some embodiments, the solvent:water ratio may be about 55:45, followed by the addition and dissolution of the polymer. In some embodiments, the solvent:water ratio may be about 50:50, followed by the addition and dissolution of the polymer.

- 10 **[0057]** In some embodiments, the solvent is an organic compound in which the active agent and polymer are mutually soluble. In some embodiments, the solvent is an alcohol, ketone, ether, ester, halogenated alkane, amide, sulfone, acid, or a nitro compound. In some embodiments, the solvent is methanol, ethanol, n-propanol, iso-propanol, or butanol. In some
15 embodiments, the solvent is acetone, methyl ethyl ketone (MEK), or methyl isobutyl ketone (MIBK). In some embodiments, the solvent is methyl acetate, ethyl acetate, or propylacetate. In some embodiments, the solvent is diethylether, tetrahydrofuran (THF), 2-methyl THF, 2,5-dimethyl THF, or 2,2,5,5-tetramethyl THF. In some embodiments, the solvent is acetonitrile, methylene chloride, toluene, 1,1,1-trichloroethane, dimethyl acetamide (DMA), nitromethane, acetic acid, or dimethylsulfoxide (DMSO). Mixtures of solvent and water are
20 suitable as long as the polymer and the Compound of Formula (I) are sufficiently soluble to make the spray-drying process practicable. In some embodiments, the water:solvent mixture is water:acetone. In some embodiments, the water:solvent mixture is water:THF. In some embodiments, the water:solvent mixture is water:methanol. In some embodiments, the water:solvent mixture is water:ethanol. In some embodiments, the water:solvent mixture is water:methyl ethyl ketone. In some embodiments, the water:solvent mixture is water:ethyl
25 acetate. In some embodiments, the water:solvent mixture is water:methylene chloride. In some embodiments, mixtures of solvents are suitable as long as the polymer and the Compound of Formula (I) are sufficiently soluble to make the spray-drying process practicable. In some embodiments, the solvent:solvent mixture is is methanol:ethylacetate. In
30 some embodiments, the solvent:solvent mixture ethanol:ethylacetate. In some embodiments, the solvent:solvent mixture is methanol:dichloromethane. In some embodiments, the solvent:solvent mixture ethanol: dichloromethane.

[0058] In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 0-25 °C. In some embodiments, the temperature range for the

agitating and mixing of the final suspension is about 5-25 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 10-25 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 15-25 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 15-24 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 15-23 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 15-22 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 15-21 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 15-20 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 17-25 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 17-24 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 17-23 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 17-22 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 17-21 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 17-20 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 18-25 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 18-24 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 18-23 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 18-22 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 18-21 °C. In some embodiments, the temperature range for the agitating and mixing of the final suspension is about 18-20 °C

[0059] In some embodiments, the suspension flow rate through the inline heat exchanger operating range may be at about 5-100 kg/hr. In some embodiments, the suspension flow rate through the inline heat exchanger operating range may be at about 5-30 kg/hr. In some embodiments, the suspension flow rate through the inline heat exchanger operating range may be at about 5-25 kg/hr. In some embodiments, the suspension flow rate through the inline heat exchanger operating range may be at about 5-20 kg/hr. In some embodiments, the suspension flow rate through the inline heat exchanger operating range may be at about 5-15 kg/hr. In some embodiments, the suspension flow rate through the inline heat exchanger

operating range may be at about 5-10 kg/hr. In some embodiments, the suspension flow rate through the inline heat exchanger operating range may be at about 30-50 kg/hr. In some embodiments, the suspension flow rate is about 35-45 kg/hr. In some embodiments, the suspension flow rate is about 35-40 kg/hr. In some embodiments, the suspension flow rate is about 40-45 kg/hr. In some embodiments, the suspension flow rate is about 42-48 kg/hr. In some embodiments, the suspension flow rate is about 45-50 kg/hr. In some embodiments, the suspension flow rate through the inline heat exchanger operating range may be at about 50-100 kg/hr. In some embodiments, the suspension flow rate is about 50-90 kg/hr. In some embodiments, the suspension flow rate is about 50-80 kg/hr. In some embodiments, the suspension flow rate is about 50-70 kg/hr. In some embodiments, the suspension flow rate is about 50-60 kg/hr.

[0060] In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 110-130 °C, preferably about 115-125 °C, most preferably about 116 °C, about 117 °C, about 118 °C, about 119 °C, about 120 °C, about 121 °C, about 122 °C, about 123 °C, about 124 °C, about 125 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 15-25 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 20-25 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 10-100 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 20-90 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 20-80 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 20-70 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 20-60 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 20-50 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 20-40 °C. In some embodiments, the solution temperature near or at the nozzle of the spray dryer may be at about 20-30 °C.

[0061] In some embodiments, the spray drying nozzle sheath gas pressure may be at about 50-100 psig. In some embodiments, the spray dryer bulk drying gas flow rate may be about 400-500 kg/hr. In some embodiments, the spray dryer chamber outlet temperature may be about 45-75 °C. In some embodiments, the spray dryer condenser temperature may be about -5 to about -20 °C.

[0062] After completion of the spray drying, the spray dried intermediate undergoes an optional secondary spray drying in an agitated vacuum dryer. In some embodiments, the drying temperature may be at about 30-60 °C, preferably about 35-55 °C, most preferably about 40-50 °C. In some embodiments, the drying duration time may not be less than about 3 hours, preferably not less than about 6 hours, not less than about 7 hours, not less than about 8 hours, not less than about 9 hours. In some embodiments, the chamber pressure may be about 30-60 mbar, preferably about 35-55 mbar, most preferably about 40-50 mbar.

[0063] In some embodiments, the polymer may be ionic. In some embodiments, the polymer may be non-ionic. In some embodiments, the pharmaceutically acceptable polymer is selected from the group consisting of: polyvinyl pyrrolidone, polyethyleneoxide, polyethylene glycol, poly(vinyl pyrrolidone-co-vinyl acetate), polyoxyethylene-polyoxypropylene block copolymers, graft copolymers comprised of polyethylene glycol, polyvinyl caprolactam and polyvinyl acetate, polymethacrylates, polyoxyethylene alkyl ethers, polyoxyethylene castor oils, polycaprolactam, polylactic acid, polyglycolic acid, poly(lactic-glycolic)acid, lipids, cellulose, pullulan, dextran, maltodextrin, hyaluronic acid, polysialic acid, chondroitin sulfate, heparin, fucoidan, pentosan polysulfate, spirulan, hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose propionate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate, cellulose acetate isophthalate, carboxymethyl ethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose acetate phthalate, hydroxypropyl methylcellulose propionate phthalate, hydroxypropyl methylcellulose acetate trimellitate, hydroxypropyl methylcellulose propionate trimellitate, cellulose acetate succinate, methyl cellulose acetate succinate, dextran, dextran acetate, dextran propionate, dextran succinate, dextran acetate propionate, dextran acetate succinate, dextran propionate succinate, dextran acetate propionate succinate, poly(methacrylic acid-co-methyl methacrylate) 1:1, poly(methacrylic acid-co-methyl methacrylate) 1:2, poly(methacrylic acid-co-ethyl acrylate) 1:1, hydroxyethyl cellulose, methyl cellulose and hydroxy propyl cellulose, poly methacrylic acid-ethyl acrylate, poly methacrylic acid-methyl methacrylate, poly methyl methacrylate-ethyl acrylate, poly trimethylammonioethyl methacrylate chloride-methyl methacrylate-ethyl acrylate and poly(butylmethacrylate-co-(2-dimethylaminoethyl)methacrylate-co-methyl methacrylate), and mixtures thereof. In some embodiments, the pharmaceutically acceptable polymer is selected from the group consisting of polyvinyl pyrrolidone, polyethyleneoxide, polyethylene glycol, poly(vinyl pyrrolidone-co-

vinyl acetate), polyoxyethylene-polyoxypropylene block copolymers, graft copolymers comprised of polyethylene glycol, polyvinyl caprolactam and polyvinyl acetate, polymethacrylates, polyoxyethylene alkyl ethers, polyoxyethylene castor oils, polycaprolactam, polylactic acid, polyglycolic acid, poly(lactic-glycolic)acid, lipids,

5 cellulose, pullulan, dextran, maltodextrin, hyaluronic acid, polysialic acid, chondroitin sulfate, heparin, fucoidan, pentosan polysulfate, spirulan, hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose propionate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate, cellulose acetate isophthalate, carboxymethyl ethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose acetate phthalate, hydroxypropyl methylcellulose propionate phthalate, hydroxypropyl methylcellulose acetate trimellitate, hydroxypropyl methylcellulose propionate trimellitate, cellulose acetate succinate, methyl cellulose acetate succinate, dextran, dextran acetate, dextran propionate, dextran succinate,

15 dextran acetate propionate, dextran acetate succinate, dextran propionate succinate, dextran acetate propionate succinate, poly(methacrylic acid-co-methyl methacrylate) 1:1, poly(methacrylic acid-co-methyl methacrylate) 1:2, poly(methacrylic acid-co-ethyl acrylate) 1:1, and mixtures thereof. In some embodiments, the polymer is hydroxypropyl methyl cellulose, hydroxypropyl cellulose, carboxymethyl ethyl cellulose, hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, polyvinyl alcohols that have at least a portion of their repeat units in hydrolyzed form, polyvinyl pyrrolidone, poloxamers, or blends thereof. In some embodiments, the pharmaceutically acceptable polymer is hydroxypropyl methyl cellulose acetate succinate.

25 **[0064]** In some embodiment, the resulting composition comprising the compound of Formula (I) comprises one or more anilinic substance, each in an amount equal to or less than 3.0% by weight based on the weight of the compound of Formula (I). Other impurities, which may include diphenyl urea are equal to or less than 0.3% by weight based on the weight of Formula (I).

30 *Identifying the Impurities*

[0065] The purity of the Compound of Formula (I) may be analyzed, generally by methods such as high-performance liquid chromatography (HPLC), gas chromatography (GC) or thin layer chromatography (TLC), to determine whether the impurities are present at

levels suitable for pharmaceutical use. Generally, impurities are identified spectroscopically and provide a chromatographic peak on a chromatogram or as a spot on a TLC panel.

[0066] Once a peak position has been associated with a particular impurity, the impurity can be identified in a sample based on its position in the chromatogram, where the position in the chromatogram is measured in minutes between the injection of the sample in a column and elution of the impurity through the detector. The position in the chromatogram is known as the retention time and the ratio between the retention times is known as the relative retention time.

[0067] A relatively pure compound may be used as a reference standard. A reference standard is similar to a reference marker, except that the latter can be used not only for detecting impurities, but also for quantifying the amount of impurities present in the sample.

[0068] In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 5% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 4% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 3% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 2% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 1% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.75% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 1% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.75% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.70% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.65% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic

impurities each together or separately are present in an amount equal to or less than about 0.60% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.55% by weight based on the weight of the compound of Formula (I). In some
5 embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.50% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.45% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are
10 present in an amount equal to or less than about 0.40% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.35% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.30% by weight
15 based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.25% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.20% by weight based on the weight of the compound of Formula (I). In some
20 embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.15% by weight based on the weight of the compound of Formula (I). In some embodiments, the anilinic impurities each together or separately are present in an amount equal to or less than about 0.1% by weight based on the weight of the compound of Formula (I). In some embodiments the anilinic impurities are represented by one or more
25 anilinic impurities selected from 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one, aniline, and a combination thereof.

[0069] In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.30% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.30%. In some embodiments, the
30 diphenyl urea impurity is present in an amount equal to or less than about 0.20% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.20%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.10% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.10%. In some

embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.075% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.075%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.05% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.05%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.04% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.04%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.03% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.03%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.02% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.02%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.01% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.01%.

[0070] In another general aspect, provided herein is a pharmaceutical composition comprising compound of Formula (I) having purity by HPLC of greater than about 95%. In some embodiments, the purity by HPLC is greater than about 96%. In some embodiments, the purity by HPLC is greater than about 97%. In some embodiments, the purity by HPLC is greater than about 98%. In some embodiments, the purity by HPLC is greater than about 99%. In some embodiments, the purity by HPLC is greater than about 99.5%. In some embodiments, the purity by HPLC is greater than about 99.8%. In some embodiments, the purity by HPLC is greater than about 99.9%. In some embodiments, the purity by HPLC is greater than about 90%. In some embodiments, the purity by HPLC is greater than about 92%. In some embodiments, the purity by HPLC is greater than about 94%.

[0071] Dispersions of the active agent and pharmaceutically acceptable polymer as described herein may be made by a spray-drying process. As used herein, the term “spray-dried dispersion” or “spray-dried powdered dispersion” means a product of a spray-drying process wherein the product comprises a dispersion of at least one active agent and at least one excipient, such as a polymer.

[0072] In the spray-drying process, the active agent and one or more polymers are dissolved in a common solvent. “Common” here means that the solvent, which can be a

mixture of compounds, will dissolve both the active agent and the polymer(s). After both active agent and polymer have been dissolved, the solvent is rapidly removed by evaporation in the spray-drying apparatus, resulting in the formation of a substantially homogeneous solid dispersion. In such dispersions, the active agent is dispersed as homogeneously as possible throughout the polymer and can be thought of as a solid solution of active agent dispersed in the polymer(s).

[0073] The solvent is removed by the spray-drying process. The term “spray-drying” is used conventionally and broadly refers to processes involving breaking up liquid mixtures into small droplets (atomization) and rapidly removing solvent from the mixture in a spray-drying apparatus where there is a strong driving force for evaporation of solvent from the droplets. Spray-drying processes and spray-drying equipment are described generally in Perry's Chemical Engineers' Handbook, pages 20-54 to 20-57 (Sixth Edition 1984). More details on spray-drying processes and equipment are reviewed by Marshall, “Atomization and Spray-Drying,” 50 Chem. Eng. Prog. Monogr. Series 2 (1954), and Masters, Spray Drying Handbook (Fourth Edition 1985). Further, additional process and spray-drying techniques and equipment are described generally in US 8,343,550 and US 7,780,988. The strong driving force for solvent evaporation is generally provided by maintaining the partial pressure of solvent in the spray-drying apparatus well below the vapor pressure of the solvent at the temperature of the drying droplets. This is accomplished by (1) maintaining the pressure in the spray-drying apparatus at a partial vacuum (e.g., 0.01 to 0.50 atm); or (2) mixing the liquid droplets with a warm drying gas; or (3) both (1) and (2). In addition, a portion of the heat required for evaporation of solvent may be provided by heating the spray solution.

[0074] The drying gas may be virtually any gas, but to minimize the risk of fire or explosions due to ignition of flammable vapors, and to minimize undesirable oxidation of the active agent, concentration-enhancing polymer, or other materials in the dispersion, an inert gas such as nitrogen, nitrogen-enriched air, or argon is utilized. The temperature of the drying gas at the gas inlet of apparatus is typically from about 60 °C to about 300 °C. The temperature of the product particles, drying gas, and evaporated solvent at the outlet or distal end of collection cone typically ranges from about 0 °C. to about 100 °C.

[0075] Solvents suitable for spray-drying process can be any organic compound in which the active agent and polymer are mutually soluble. The solvent should have relatively low toxicity and be removed from the dispersion to a level that is acceptable according to The International Committee on Harmonization (ICH) guidelines. Removal of solvent to this level

may require a subsequent processing step such as tray-drying or secondary drying. In some embodiments, the solvent is an alcohol, ketone, ether, ester, halogenated alkane, amide, sulfone, acid, or a nitro compound. In some embodiments, the solvent is methanol, ethanol, n-propanol, iso-propanol, or butanol. In some embodiments, the solvent is acetone, methyl ethyl ketone (MEK), or methyl isobutyl ketone (MIBK). In some embodiments, the solvent is methyl acetate, ethyl acetate, or propylacetate. In some embodiments, the solvent is diethylether, tetrahydrofuran (THF), 2-methyl THF, 2,5-dimethyl THF, or 2,2,5,5-tetramethyl THF. In some embodiments, the solvent is acetonitrile, methylene chloride, toluene, 1,1,1-trichloroethane, dimethyl acetamide (DMA), nitromethane, acetic acid, or dimethylsulfoxide (DMSO). Mixtures of solvent and water are suitable as long as the polymer and the compound of Formula (I) are sufficiently soluble to make the spray-drying process practicable. In some embodiments, the water:solvent mixture is water:acetone. In some embodiments, the water:solvent mixture is water:THF. In some embodiments, the water:solvent mixture is water:methanol. In some embodiments, the water:solvent mixture is water:ethanol. In some embodiments, the water:solvent mixture is water:methyl ethyl ketone. In some embodiments, the water:solvent mixture is water:ethyl acetate. In some embodiments, the water:solvent mixture is water:methylene chloride. In some embodiments, mixtures of solvents are suitable as long as the polymer and the Compound of Formula (I) are sufficiently soluble to make the spray-drying process practicable. In some embodiments, the solvent:solvent mixture is methanol:ethylacetate. In some embodiments, the solvent:solvent mixture ethanol:ethylacetate. In some embodiments, the solvent:solvent mixture is methanol:dichloromethane. In some embodiments, the solvent:solvent mixture ethanol:dichloromethane.

[0076] The composition of the solvent-bearing feed will depend on the desired ratio of active agent-to-polymer in the dispersion and the solubility of the active agent and polymer in the solvent. Generally, it is desirable to use as high a combined active agent and polymer concentration in the solvent-bearing feed as possible, provided the active agent and polymer are dissolved in the solvent at the temperature range of the process, to reduce the total amount of solvent that must be removed to form the solid amorphous dispersion. In some embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 0.01 wt % to at least about 20 wt %. In some embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 0.01 wt %. In some embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 0.1 wt %. In some embodiments, the solvent-bearing

feed has a combined active agent and polymer concentration of at least about 0.5 wt %. In some embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 1.0 wt %. In some embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 2.0 wt %. In some
5 embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 3.0 wt %. In some embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 4.0 wt %. In some
10 embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 5.0 wt %. In some embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 6.0 wt %. In some
embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 7.0 wt %. In some embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 8.0 wt %. In some
15 embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 9.0 wt %. In some embodiments, the solvent-bearing feed has a combined active agent and polymer concentration of at least about 10.0 wt %.

[0077] The average residence time of particles in the drying chamber should be at least 10 seconds, preferably at least 20 seconds. Typically, following solidification, the powder formed stays in the spray-drying chamber for about 5 to 60 seconds, causing further
20 evaporation of solvent. The final solvent content of the solid dispersion as it exits the dryer should be low, since this reduces the mobility of active agent molecules in the dispersion, thereby improving its stability. Generally, the solvent content of the dispersion as it leaves the spray-drying chamber should be less than about 10 wt %. In some embodiments, the solvent content of the dispersion as it leaves the spray-drying chamber is less than about 9 wt%. In
25 some embodiments, the solvent content of the dispersion as it leaves the spray-drying chamber is less than about 8 wt %. In some embodiments, the solvent content of the dispersion as it leaves the spray-drying chamber is less than about 7 wt %. In some
embodiments, the solvent content of the dispersion as it leaves the spray-drying chamber is less than about 6 wt %. In some embodiments, the solvent content of the dispersion as it
30 leaves the spray-drying chamber is less than about 5 wt %. In some embodiments, the solvent content of the dispersion as it leaves the spray-drying chamber is less than about 4 wt
%. In some embodiments, the solvent content of the dispersion as it leaves the spray-drying chamber is less than about 3 wt %. In some embodiments, the solvent content of the
dispersion as it leaves the spray-drying chamber is less than about 2 wt %. In some

embodiments, the solvent content of the dispersion as it leaves the spray-drying chamber is less than about 1 wt %. In some embodiments, the acetone content of the dispersion as it leaves the spray-drying chamber is less than about 0.5 wt %. In some embodiments, the acetone content of the dispersion as it leaves the spray-drying chamber is less than about 0.3 wt %. In some embodiments, the acetone content of the dispersion as it leaves the spray-drying chamber is less than about 0.1 wt. A subsequent processing step, such as tray-drying, may be used to remove the solvent to this level.

Pharmaceutical Compositions and Formulations

[0078] Such compositions or pharmaceutical compositions, for example, can be in a form such as a tablet, capsule, pill, powder, liquids, suspensions, emulsions, granules, sustained release formulations, solution, and suspension. The pharmaceutical composition may be in an oral formulation suitable for single administration of precise dosages.

[0079] The compound of Formula (I) as described herein may be formed into a finished dosage form. The finished dosage form comprises one of more of a liquid, solid or semi-solid dosage forms depending on the route of administration.

[0080] The excipients employed in the pharmaceutical compositions can impart good powder flow and compression characteristics to the material being compressed. Desirable characteristics of excipients can include high-compressibilities as to allow for strong tablets to be made at low compression forces; good powder flow properties that can improve the powder flow of other excipients in the composition; and cohesiveness, for example to prevent a tablet from crumbling during processing, shipping, and handling. Such properties are imparted to these excipients through pretreatment steps, such as dry granulation (e.g., by roller compaction, slugging), wet granulation, spray drying spheronization (e.g., spray dried dispersion, solid nanodispersions) or crystallization (e.g., salt forms) of a pharmaceutical composition. They may be classified according to the role that they play in the final tablet. Other excipients which give physical characteristics to a finished tablet are coloring and flavoring agents (e.g., in the case of chewable tablets). Examples of excipients are described, for example, in the Handbook of Pharmaceutical Excipients (5th edition), edited by Raymond C. Rowe, Paul J. Sheskey, and Sian C. Owen; Publisher: Pharmaceutical Press.

[0081] As described herein, pharmaceutical compositions can also comprise a pharmaceutically acceptable polymer. The pharmaceutically acceptable polymers may be ionic or non-ionic. Exemplary pharmaceutically acceptable polymers include polyvinyl pyrrolidone, polyethyleneoxide, polyethylene glycol, poly(vinyl pyrrolidone-co-vinyl

acetate), polyoxyethylene-polyoxypropylene block copolymers, graft copolymers comprised of polyethylene glycol, polyvinyl caprolactam and polyvinyl acetate, polymethacrylates, polyoxyethylene alkyl ethers, polyoxyethylene castor oils, polycaprolactam, polylactic acid, polyglycolic acid, poly(lactic-glycolic)acid, lipids, cellulose, pullulan, dextran, maltodextrin,

5 hyaluronic acid, polysialic acid, chondroitin sulfate, heparin, fucoidan, pentosan polysulfate, spirulan, hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose propionate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate, cellulose acetate isophthalate,

10 carboxymethyl ethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose acetate phthalate, hydroxypropyl methylcellulose propionate phthalate, hydroxypropyl methylcellulose acetate trimellitate, hydroxypropyl methylcellulose propionate trimellitate, cellulose acetate succinate, methyl cellulose acetate succinate, dextran, dextran acetate, dextran propionate, dextran succinate, dextran acetate propionate,

15 dextran acetate succinate, dextran propionate succinate, dextran acetate propionate succinate, poly(methacrylic acid-co-methyl methacrylate) 1:1, poly(methacrylic acid-co-methyl methacrylate) 1:2, poly(methacrylic acid-co-ethyl acrylate) 1:1, hydroxyethyl cellulose, methyl cellulose and hydroxy propyl cellulose, poly methacrylic acid-ethyl acrylate, poly methacrylic acid-methyl methacrylate, poly methyl methacrylate-ethyl acrylate, poly

20 trimethylammonioethyl methacrylate chloride-methyl methacrylate-ethyl acrylate and poly(butylmethacrylate-co-(2-dimethylaminoethyl)methacrylate-co-methyl methacrylate), or mixtures thereof. In some embodiments, the pharmaceutically acceptable polymers is selected from the group consisting of polyvinyl pyrrolidone, polyethyleneoxide, polyethylene glycol, poly(vinyl pyrrolidone-co-vinyl acetate), polyoxyethylene-polyoxypropylene block

25 copolymers, graft copolymers comprised of polyethylene glycol, polyvinyl caprolactam and polyvinyl acetate, polymethacrylates, polyoxyethylene alkyl ethers, polyoxyethylene castor oils, polycaprolactam, polylactic acid, polyglycolic acid, poly(lactic-glycolic)acid, lipids, cellulose, pullulan, dextran, maltodextrin, hyaluronic acid, polysialic acid, chondroitin sulfate, heparin, fucoidan, pentosan polysulfate, spirulan, hydroxypropyl methyl cellulose

30 acetate succinate, hydroxypropyl methyl cellulose propionate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate, cellulose acetate isophthalate, carboxymethyl ethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose acetate phthalate, hydroxypropyl

methylcellulose propionate phthalate, hydroxypropyl methylcellulose acetate trimellitate, hydroxypropyl methylcellulose propionate trimellitate, cellulose acetate succinate, methyl cellulose acetate succinate, dextran, dextran acetate, dextran propionate, dextran succinate, dextran acetate propionate, dextran acetate succinate, dextran propionate succinate, dextran acetate propionate succinate, poly(methacrylic acid-co-methyl methacrylate) 1:1, poly(methacrylic acid-co-methyl methacrylate) 1:2, poly(methacrylic acid-co-ethyl acrylate) 1:1, and mixtures thereof. In some embodiments, the pharmaceutically acceptable polymer is hydroxypropyl methyl cellulose acetate succinate.

[0082] The pharmaceutical composition provided herein can contain one or more fillers, which are added, for example, to increase the bulk weight of the blend resulting in a practical size for compression. Fillers that may be used include one or more of calcium salts such as calcium phosphate dibasic and sugars such as lactose, sucrose, dextrose, microcrystalline cellulose, mannitol, and maltodextrin. Examples of pharmaceutically acceptable fillers and pharmaceutically acceptable diluents include, but are not limited to, confectioner's sugar, compressible sugar, dextrates, dextrin, dextrose, lactose, mannitol, microcrystalline cellulose, powdered cellulose, sorbitol, sucrose and talc. In some embodiments, the filler is microcrystalline cellulose, which can be manufactured by the controlled hydrolysis of alpha-cellulose. Suitable microcrystalline cellulose will have an average particle size of from about 20 nm to about 200 nm. Suitable microcrystalline cellulose includes Avicel PH 101, Avicel PH 102, Avicel PH 103, Avicel PH 105 and Avicel PH 200, e.g., manufactured by FMC Corporation. In some embodiments, the filler is lactose.

[0083] The pharmaceutical composition can also include a lubricant. The term "lubricant" as used herein is typically added to prevent the tableting materials from sticking to punches, minimize friction during tablet compression, and to allow for removal of the compressed tablet from the die. Examples of lubricants include, but are not limited to, colloidal silica, magnesium trisilicate, talc, magnesium carbonate, magnesium oxide, glyceryl behaptate, polyethylene glycol, ethylene oxide polymers (e.g., Carowax), sodium lauryl sulfate, magnesium stearate, aluminum stearate, calcium stearate, sodium stearyl fumarate, stearic acid, magnesium lauryl stearate, and mixtures of magnesium stearate with sodium lauryl sulfate. Exemplary lubricants include calcium stearate, magnesium stearate and sodium stearyl fumarate. In some embodiments, the lubricant is magnesium stearate.

[0084] The pharmaceutical composition provided herein can also contain a glidant. The term "glidant" as used herein is a substance added to a powder that can improve its

flowability, such as by reducing inter-particle friction. Exemplary glidants include but are not limited to colloidal silicas, colloidal silicon dioxide, fumed silica, CAB-O-SIL[®] M-5P, AEROSIL[®], talc, Syloid[®], starch, and magnesium aluminum silicates. In some embodiments, the glidant is silicon dioxide. It should be noted that excipients may serve multiple functions.


5 In some embodiments, the lubricant, for example magnesium stearate, may also function as a glidant.

[0085] A disintegrant may be present in an amount necessary to expedite dissolution (e.g., increase the rate of tablet disintegration). The term “disintegrant” as used herein refers to an excipient which can oppose the physical forces of particle bonding in a tablet or capsule
10 when the oral formulation is placed in an aqueous environment. Disintegrants include starch derivatives and salts of carboxymethylcellulose. Examples of pharmaceutically acceptable disintegrants include, but are not limited to, starches, e.g., sodium starch glycolate, pregelatinized starch; clays; celluloses; alginates; gums; cross-linked polymers, e.g., cross-linked polyvinyl pyrrolidone (e.g., polyplasdoneTM, polyvinyl polypyrrolidone,
15 crospovidone), cross-linked calcium carboxymethylcellulose and cross-linked sodium carboxymethylcellulose (sodium croscarmellose); and soy polysaccharides. In some embodiments, the disintegrant is crospovidone (e.g, PVP-XL).

[0086] In some embodiments, described herein is a pharmaceutical composition comprising a compound represented by Formula (I) and one or more pharmaceutically
20 acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises one or more anilinic substances each in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I). In some embodiments, the one or more anilinic substances is the compound of Formula (II) and is present in the composition in an amount of less than about 1 % by weight based on the weight of the compound of Formula
25 (I). In some embodiments, the one or more anilinic substances is the compound of Formula (II) and is present in the composition in an amount of less than about 0.1 % by weight to about 0.5% by weight based on the weight of the compound of Formula (I). In some embodiments, the one or more anilinic substances is the compound of Formula (II) and is present in the composition in an amount of less than about 0.01 % by weight to about 0.1%
30 by weight based on the weight of the compound of Formula (I). In some embodiments, the composition comprises less than about 1 % (w/w) of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one. In some embodiments, the composition comprises about 0.1 % (w/w) to about 0.5% (w/w) of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one. In some embodiments, the

composition comprises about 0.01 % (w/w) to about 0.1% (w/w) of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one.

[0087] In some embodiments, the composition further comprises less than 10% by weight of a compound represented by Formula (III):




5

Formula (III)

based on the weight of the compound of Formula (I).

[0088] In some embodiments, the composition further comprises less than 3% by weight of a compound represented by Formula (III):




10

Formula (III)

based on the weight of the compound of Formula (I).

[0089] In some embodiments, the composition further comprises less than 1% by weight of a compound represented by Formula (III):




15

Formula (III)

based on the weight of the compound of Formula (I).


[0090] In some embodiments, the composition further comprises less than about 0.1 % by weight to about 0.5% by weight of a compound represented by Formula (III):



Formula (III)

5 based on the weight of the compound of Formula (I).

[0091] In some embodiments, the composition further comprises about 0.01 % by weight to about 0.1% by weight of a compound represented by Formula (III):



Formula (III)

10 based on the weight of the compound of Formula (I).

[0092] In some embodiments, described herein is a pharmaceutical composition comprising a compound represented by Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises less than about 10% by weight of a compound represented by Formula (III) based on the weight of the compound of Formula (I). In some embodiments, described herein is a pharmaceutical composition comprising a compound represented by Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises less than about 7% by weight of a compound represented by Formula (III) based on the weight of the compound of Formula (I). In some embodiments, described herein is a pharmaceutical composition comprising a compound represented by Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises less than about 5% by weight of a compound represented by Formula (III) based on the weight of the compound of Formula (I). In some embodiments, described herein is a pharmaceutical composition comprising a compound

represented by Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises less than about 3% by weight of a compound represented by Formula (III) based on the weight of the compound of Formula (I). In some embodiments, described herein is a pharmaceutical composition


5 comprising a compound represented by Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises less than about 1% by weight of a compound represented by Formula (III) based on the weight of the compound of Formula (I). In some embodiments, described herein is a pharmaceutical composition comprising a compound represented by Formula (I) and one or

10 more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises less than about 0.1% by weight of a compound represented by Formula (III) based on the weight of the compound of Formula (I). In some embodiments, described herein is a pharmaceutical composition comprising a compound represented by Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents,

15 wherein the pharmaceutical composition comprises less than about 0.01% by weight of a compound represented by Formula (III) based on the weight of the compound of Formula (I).

[0093] In another embodiment, described herein is a pharmaceutical composition comprising the compound of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises one or

20 more anilinic substances and a compound represented by Formula (III):




Formula (III)

each in an amount equal to or less than 3.0% by weight based on the weight of the compound of Formula (I).


25 **[0094]** In some embodiments, the composition has equal to or less than about 0.5% by weight one or more anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the composition equal to or less than about 0.3% by weight anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the

one or more anilinic substances is selected from the group consisting of a compound represented by Formula (II):




Formula (II)

- 5 aniline, and a combination thereof. In some embodiments, the one or more anilinic substances is a compound represented by Formula (II):




Formula (II)

- and is present in the composition in an amount of less than about 1 % by weight based on the weight of the compound of Formula (I). In some embodiments, the one or more anilinic substances is a compound represented by Formula (II):



Formula (II)

- and is present in the composition in an amount of less than about 0.1 % by weight to about 0.5% by weight based on the weight of the compound of Formula (I). In some embodiments, the one or more anilinic substances is a compound represented by Formula (II):


**Formula (II)**

and is present in the composition in an amount of less than about 0.01 % by weight to about 0.1% by weight based on the weight of the compound of Formula (I). In some embodiments, the compound of Formula (III) is present in the composition in an amount of less than about 1% by weight based on the weight of the compound of Formula (I). In some embodiments, the compound of Formula (III) is present in the composition in an amount of less than about 0.1 % by weight to about 0.5% by weight by weight based on the weight of the compound of Formula (I). In some embodiments, the compound of Formula (III) is present in the composition in an amount of less than about 0.01 % by weight to about 0.1 % by weight by weight based on the weight of the compound of Formula (I).

[0095] In some embodiments, described herein is a solid dispersion comprising a compound represented by Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the solid dispersion comprises one or more anilinic substances each in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I). In some embodiments, the one or more anilinic substances is the compound of Formula (II) and is present in the composition in an amount of less than about 1 % by weight based on the weight of the compound of Formula (I). In some embodiments, the one or more anilinic substances is the compound of Formula (II) and is present in the composition in an amount of less than about 0.1 % by weight to about 0.5% by weight based on the weight of the compound of Formula (I). In some embodiments, the one or more anilinic substances is the compound of Formula (II) and is present in the composition in an amount of less than about 0.01 % by weight to about 0.1% by weight based on the weight of the compound of Formula (I). In some embodiments, the solid dispersion comprises less than about 1 % (w/w) of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)). In some embodiments, the solid dispersion comprises about 0.1 % (w/w) to about 0.5% (w/w) of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)). In some embodiments, the solid dispersion comprises about 0.01 % (w/w) to

about 0.1% (w/w) of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)).

[0096] In some embodiments, the solid dispersion further comprises less than 10% by weight of a compound represented by Formula (III):




5

Formula (III)

based on the weight of the compound of Formula (I).

[0097] In some embodiments, the solid dispersion further comprises less than 3% by weight of a compound represented by Formula (III):




10

Formula (III)

based on the weight of the compound of Formula (I).

[0098] In some embodiments, the solid dispersion comprises less than 1% by weight of a compound represented by Formula (III):




15

Formula (III)

based on the weight of the compound of Formula (I).


[0099] In some embodiments, the solid dispersion comprises less than about 0.1 % by weight to about 0.5% by weight of a compound represented by Formula (III):



Formula (III)

5 based on the weight of the compound of Formula (I).

[0100] In some embodiments, the solid dispersion comprises about 0.01 % by weight to about 0.1% by weight of a compound represented by Formula (III):




Formula (III)

10 based on the weight of the compound of Formula (I).

[0101] In some embodiments, described herein is a solid dispersion comprising a polymer and a compound represented by Formula (I) and a pharmaceutically acceptable polymer, wherein the solid dispersion comprises one or more anilinic substance, each in an amount equal to or less than about 0.50% by weight based on the weight of the compound of Formula (I). In some embodiments, the one or more anilinic substances is the compound of Formula (II) and is present in the composition in an amount of less than about 0.1 % based on the weight of the compound of Formula (I). In some embodiments, the one or more anilinic substances is the compound of Formula (II) and is present in the composition in an amount of less than about 0.01 % by weight to about 0.1% by weight based on the weight of the compound of Formula (I). In some embodiments, the solid dispersion comprises about 0.1 % (w/w) to about 0.5% (w/w) of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)). In some embodiments, the solid dispersion comprises about 0.01 % (w/w) to about 0.1% (w/w) of 3-

(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methyamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)).

[0102] In some embodiments, the solid dispersion further comprises less than about 0.1 % by weight to about 0.5% by weight of a compound represented by Formula (III):




5

Formula (III)

based on the weight of the compound of Formula (I).

[0103] In some embodiments, the solid dispersion further comprises about 0.01 % by weight to about 0.1% by weight of a compound represented by Formula (III):



10

Formula (III)

based on the weight of the compound of Formula (I).

[0104] In some embodiments, the solid dispersion comprises from about 10% to about 50%, or from about 10% to about 30%, or from about 20% to about 30%, by weight of the compound represented by Formula (I) based on the total weight of the solid dispersion. In some embodiments, the pharmaceutical compositions may comprise about 25% by weight of the compound represented by Formula (I) based on the total weight of the solid dispersion.

[0105] The solid dispersion provided herein comprises, in some embodiments, a pharmaceutically acceptable polymer selected from the group consisting of polyvinyl pyrrolidone, polyethyleneoxide, polyethylene glycol, poly(vinyl pyrrolidone-co-vinyl acetate), polyoxyethylene-polyoxypropylene block copolymers, graft copolymers comprised of polyethylene glycol, polyvinyl caprolactam and polyvinyl acetate, polymethacrylates, polyoxyethylene alkyl ethers, polyoxyethylene castor oils, polycaprolactam, polylactic acid,

polyglycolic acid, poly(lactic-glycolic)acid, lipids, cellulose, pullulan, dextran, maltodextrin, hyaluronic acid, polysialic acid, chondroitin sulfate, heparin, fucoidan, pentosan polysulfate, spirulan, hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose propionate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate, cellulose acetate isophthalate, carboxymethyl ethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose acetate phthalate, hydroxypropyl methylcellulose propionate phthalate, hydroxypropyl methylcellulose acetate trimellitate, hydroxypropyl methylcellulose propionate trimellitate, cellulose acetate succinate, methyl cellulose acetate succinate, dextran, dextran acetate, dextran propionate, dextran succinate, dextran acetate propionate, dextran acetate succinate, dextran propionate succinate, dextran acetate propionate succinate, poly(methacrylic acid-co-methyl methacrylate) 1:1, poly(methacrylic acid-co-methyl methacrylate) 1:2, poly(methacrylic acid-co-ethyl acrylate) 1:1, hydroxyethyl cellulose, methyl cellulose and hydroxy propyl cellulose, poly methacrylic acid-ethyl acrylate, poly methacrylic acid-methyl methacrylate, poly methyl methacrylate-ethyl acrylate, poly trimethylammonioethyl methacrylate chloride-methyl methacrylate-ethyl acrylate and poly(butylmethacrylate-co-(2-dimethylaminoethyl)methacrylate-co-methyl methacrylate), and mixtures thereof. The solid dispersion provided herein comprises, in some embodiments, a pharmaceutically acceptable polymer selected from the group consisting of polyvinyl pyrrolidone, polyethyleneoxide, polyethylene glycol, poly(vinyl pyrrolidone-co-vinyl acetate), polyoxyethylene-polyoxypropylene block copolymers, graft copolymers comprised of polyethylene glycol, polyvinyl caprolactam and polyvinyl acetate, polymethacrylates, polyoxyethylene alkyl ethers, polyoxyethylene castor oils, polycaprolactam, polylactic acid, polyglycolic acid, poly(lactic-glycolic)acid, lipids, cellulose, pullulan, dextran, maltodextrin, hyaluronic acid, polysialic acid, chondroitin sulfate, heparin, fucoidan, pentosan polysulfate, spirulan, hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose propionate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate, cellulose acetate isophthalate, carboxymethyl ethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose acetate phthalate, hydroxypropyl methylcellulose propionate phthalate, hydroxypropyl methylcellulose acetate trimellitate, hydroxypropyl methylcellulose propionate trimellitate, cellulose acetate succinate, methyl cellulose acetate succinate,

dextran, dextran acetate, dextran propionate, dextran succinate, dextran acetate propionate, dextran acetate succinate, dextran propionate succinate, dextran acetate propionate succinate, poly(methacrylic acid-co-methyl methacrylate) 1:1, poly(methacrylic acid-co-methyl methacrylate) 1:2, poly(methacrylic acid-co-ethyl acrylate) 1:1, and mixtures thereof. For example, the pharmaceutically acceptable polymer in the formulation provided herein is hydroxypropyl methyl cellulose acetate succinate.

[0106] In some embodiments, the solid dispersion comprises the compound represented by Formula (I) and the pharmaceutically acceptable polymer in a ratio from about 40:60 to about 10:90 or from about 30:70 to about 20:80. In some embodiments, the compound represented by Formula (I) and the pharmaceutically acceptable polymer may be in a ratio of about 25:75.

[0107] Also provided herein is a pharmaceutical composition comprising: (a) an intragranular blend comprising: (i) a solid spray-dried dispersion comprising a compound represented by Formula (I), and a pharmaceutically acceptable polymer; (ii) one or more fillers; (iii) a disintegrant; (iv) a glidant; and (v) a lubricant; and (b) an extragranular blend comprising: (i) a glidant; and (ii) a lubricant.

[0108] In some embodiments, the blend of the internal and extragranular blends is in a ratio of from about 90:10 to about 99.5:0.5. For example, the blend of the internal and extragranular blends may be in a ratio of about 99:1.

[0109] In some embodiments, the solid dispersion of the intragranular blend comprises from about 10% to about 50%, or from about 10% to about 30%, or from about 20% to about 30% by weight of the compound represented by Formula (I) based on the total weight of the solid spray-dried dispersion. In some embodiments, the solid spray-dried dispersion may comprise about 25% by weight of the compound represented by Formula (I) based on the total weight of the solid spray-dried dispersion.

[0110] In some embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable polymer selected from the group consisting of polyvinyl pyrrolidone, polyethyleneoxide, polyethylene glycol, poly(vinyl pyrrolidone-co-vinyl acetate), polyoxyethylene-polyoxypropylene block copolymers, graft copolymers comprised of polyethylene glycol, polyvinyl caprolactam and polyvinyl acetate, polymethacrylates, polyoxyethylene alkyl ethers, polyoxyethylene castor oils, polycaprolactam, polylactic acid, polyglycolic acid, poly(lactic-glycolic)acid, lipids, cellulose, pullulan, dextran, maltodextrin, hyaluronic acid, polysialic acid, chondroitin sulfate, heparin, fucoidan, pentosan polysulfate,

spirulan, hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose propionate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate, cellulose acetate isophthalate,

5 carboxymethyl ethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose acetate phthalate, hydroxypropyl methylcellulose propionate phthalate, hydroxypropyl methylcellulose acetate trimellitate, hydroxypropyl methylcellulose propionate trimellitate, cellulose acetate succinate, methyl cellulose acetate succinate, dextran, dextran acetate, dextran propionate, dextran succinate, dextran acetate propionate,

10 dextran acetate succinate, dextran propionate succinate, dextran acetate propionate succinate, poly(methacrylic acid-co-methyl methacrylate) 1:1, poly(methacrylic acid-co-methyl methacrylate) 1:2, poly(methacrylic acid-co-ethyl acrylate) 1:1, hydroxyethyl cellulose, methyl cellulose and hydroxy propyl cellulose, poly methacrylic acid-ethyl acrylate, poly methacrylic acid-methyl methacrylate, poly methyl methacrylate-ethyl acrylate, poly

15 trimethylammonioethyl methacrylate chloride-methyl methacrylate-ethyl acrylate and poly(butylmethacrylate-co-(2-dimethylaminoethyl)methacrylate-co-methyl methacrylate), and mixtures thereof. In some embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable polymer selected from the group consisting of polyvinyl pyrrolidone, polyethyleneoxide, polyethylene glycol, poly(vinyl pyrrolidone-co-vinyl

20 acetate), polyoxyethylene-polyoxypropylene block copolymers, graft copolymers comprised of polyethylene glycol, polyvinyl caprolactam and polyvinyl acetate, polymethacrylates, polyoxyethylene alkyl ethers, polyoxyethylene castor oils, polycaprolactam, polylactic acid, polyglycolic acid, poly(lactic-glycolic)acid, lipids, cellulose, pullulan, dextran, maltodextrin, hyaluronic acid, polysialic acid, chondroitin sulfate, heparin, fucoidan, pentosan polysulfate,

25 spirulan, hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose propionate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate, cellulose acetate isophthalate, carboxymethyl ethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl

30 methylcellulose acetate phthalate, hydroxypropyl methylcellulose propionate phthalate, hydroxypropyl methylcellulose acetate trimellitate, hydroxypropyl methylcellulose propionate trimellitate, cellulose acetate succinate, methyl cellulose acetate succinate, dextran, dextran acetate, dextran propionate, dextran succinate, dextran acetate propionate, dextran acetate succinate, dextran propionate succinate, dextran acetate propionate succinate,

poly(methacrylic acid-co-methyl methacrylate) 1:1, poly(methacrylic acid-co-methyl methacrylate) 1:2, poly(methacrylic acid-co-ethyl acrylate) 1:1, and mixtures thereof. For example, the pharmaceutically acceptable polymer is hydroxypropyl methyl cellulose acetate succinate.

- 5 **[0111]** In some embodiments, the pharmaceutical composition comprises the compound represented by Formula (I) and the pharmaceutically acceptable polymer in a ratio from about 40:60 to about 10:90, or from about 30:70 to about 20:80. In some embodiments, the compound represented by Formula (I) and the pharmaceutically acceptable polymer may be in a ratio of about 25:75.
- 10 **[0112]** In some embodiments, the intragranular blend of the pharmaceutical composition comprises one or more fillers, wherein the total amount of the one or more fillers is from about 40% to about 80% by weight based on the total weight of the pharmaceutical composition. One or more fillers are lactose, maltodextrin, mannitol, microcrystalline cellulose, pregelatinized starch, sucrose esters, or hydrates thereof. In some embodiments,
- 15 the intragranular blend comprises two fillers. When the intragranular blend comprises two fillers, each filler may independently be present in an amount from about 20% to about 40%, e.g., about 33%, by weight based on the total weight of the pharmaceutical composition. In some embodiments, one filler may be microcrystalline cellulose and the other filler may be lactose monohydrate.
- 20 **[0113]** In some embodiments, the intragranular blend of the pharmaceutical composition comprises from about 1% to about 10% by weight, e.g., about 5%, of the disintegrant based on the total weight of the pharmaceutical composition. The disintegrant is crospovidone, croscarmellose sodium, sodium starch glycolate, microcrystalline cellulose, or pregelatinized starch. In some embodiments, the disintegrant in the intragranular blend may be
- 25 crospovidone.
- [0114]** In some embodiments, the glidant of the intragranular blend is present in an amount from about 0.1% to about 1%, e.g., about 0.5%, based on the total weight of the pharmaceutical composition. For example, the glidant of the intragranular blend may be silicon dioxide.
- 30 **[0115]** In some embodiments, the glidant of the extragranular blend is present in an amount from about 0.1% to about 1%, e.g., about 0.5%, based on the total weight of the pharmaceutical composition. In some embodiments, the glidant of the extragranular blend may be silicon dioxide.


[0116] In some embodiments, the lubricant of the intragranular blend is present in an amount from about 0.1% to about 1%, e.g., about 0.5%, based on the total weight of the pharmaceutical composition. In some embodiments, the lubricant of the intragranular blend is magnesium stearate, calcium stearate, glyceryl monostearate, hydrogenated castor oil, sodium lauryl sulfate, sodium stearyl fumarate, stearic acid, zinc stearate, talc, microcrystalline cellulose, or sucrose esters. For example, the lubricant of the intragranular blend may be magnesium stearate.

[0117] In some embodiments, the lubricant of the extragranular blend is present in an amount from about 0.1% to about 1%, e.g., about 0.5%, based on the total weight of the pharmaceutical composition. In some embodiments, the lubricant of the extragranular blend is magnesium stearate, calcium stearate, glyceryl monostearate, hydrogenated castor oil, sodium lauryl sulfate, sodium stearyl fumarate, stearic acid, zinc stearate, talc, microcrystalline cellulose, or sucrose esters. As an example, the lubricant of the extragranular blend may be magnesium stearate.

[0118] In some embodiments, the composition comprises less than or equal to about 10 % by weight of one or more anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the composition comprises less than or equal to about 7 % by weight of one or more anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the composition comprises less than or equal to about 5 % by weight of one or more anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the composition comprises less than or equal to about 3 % by weight of one or more anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the composition comprises less than about 10 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the composition comprises less than about 7 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the composition comprises less than about 5 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the composition comprises less than about 3 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the

composition comprises less than about 2 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the composition comprises less than about 1 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the composition comprises about 0.1 % by weight to about 0.5% by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the composition comprises about 0.01 % by weight to about 0.1% by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I).


[0119] In some embodiments, the composition comprises less than 10% by weight of a compound represented by Formula (III):



Formula (III)

based on the weight of the compound of Formula (I).


[0120] In some embodiments, the composition comprises less than 7% by weight of a compound represented by Formula (III):



Formula (III)

based on the weight of the compound of Formula (I).


[0121] In some embodiments, the composition comprises less than 5% by weight of a compound represented by Formula (III):



Formula (III)

5 based on the weight of the compound of Formula (I).


[0122] In some embodiments, the composition comprises less than 3% by weight of a compound represented by Formula (III):



Formula (III)

10 based on the weight of the compound of Formula (I).


[0123] In some embodiments, the composition comprises less than 1% by weight of a compound represented by Formula (III):



Formula (III)

15 based on the weight of the compound of Formula (I).


[0124] In some embodiments, the composition comprises less than about 0.1 % by weight to about 0.5% by weight of a compound represented by Formula (III):



Formula (III)

5 based on the weight of the compound of Formula (I).

[0125] In some embodiments, the composition comprises about 0.01 % by weight to about 0.1% by weight of a compound represented by Formula (III):



Formula (III)


10 based on the weight of the compound of Formula (I).

[0126] In some embodiments, provided herein is a pharmaceutical composition comprising: (a) an intragranular blend comprising: (i) about 33% by weight of a solid spray-dried dispersion based on the total weight of the pharmaceutical composition, the solid spray-dried dispersion comprising a compound represented by Formula (I) having a purity by HPLC of greater than 95% and hydroxypropyl methyl cellulose acetate succinate, wherein the solid spray-dried dispersion comprises about 25% by weight of the compound represented by Formula (I) based on the total weight of the solid spray-dried dispersion; (ii) about 30% by weight of microcrystalline cellulose based on the total amount of the of the pharmaceutical composition; (iii) about 30% by weight of lactose monohydrate based on the total amount of the of the pharmaceutical composition; (iv) about 5% by weight of crospovidone based on the total amount of the of the pharmaceutical composition; (v) about 0.5% by weight of silicon dioxide based on the total amount of the of the pharmaceutical composition; and (vi) about 0.5% by weight of magnesium stearate based on the total amount of the of the pharmaceutical composition; and

(b) an extragranular blend comprising: (i) about 0.5% by weight of silicon dioxide based on the total amount of the of the pharmaceutical composition; and (ii) about 0.5% by weight of magnesium stearate based on the total amount of the of the pharmaceutical composition.

[0127] In some embodiments, provided herein is a pharmaceutical composition comprising: (a) an intragranular blend comprising: (i) about 200 mg of a solid spray-dried dispersion comprising a compound represented by Formula (I) and hydroxypropyl methyl cellulose acetate succinate, wherein the solid spray-dried dispersion comprises about 50 mg of the compound represented by Formula (I); (ii) about 179 mg of microcrystalline cellulose; (iii) about 179 mg of lactose monohydrate; (iv) about 30 mg of croscopovidone; (v) about 3 mg of silicon dioxide; and (vi) about 3 mg of magnesium stearate; and (b) an extragranular blend comprising: (i) about 3 mg of silicon dioxide; and (ii) about 3 mg of magnesium stearate.

[0128] Described herein, in an embodiment, is a pharmaceutical composition comprising: a) a solid dispersion comprising: a compound represented by Formula (I) or pharmaceutically acceptable salt thereof,




Formula (I)

and a pharmaceutically acceptable polymer; and b) one or more pharmaceutically acceptable carriers, wherein no single anilinic substance is present in an amount equal to or less than about 3.0% by weight.

[0129] In some embodiments, the anilinic substance is present in amount equal to or less than about 0.5% by weight based on the total weight of the compound of Formula (I) present in the composition.

[0130] In some embodiments, the anilinic substance is present in an amount equal to or less than about 0.3% by weight based on the total weight of the compound of Formula (I) present in the composition.

[0131] In some embodiments, the anilinic substance is one of : the compound of Formula (II)




Formula (II)

5 aniline, or a combination thereof.

[0132] In some embodiments, the compound is present in the solid dispersion as a free base in substantially amorphous form.


[0133] Described herein, in an embodiment, pharmaceutical composition comprising: a) a solid dispersion comprising: an amorphous form of a compound represented by Formula (I)



10

Formula (I)

and a pharmaceutically acceptable polymer; and b) one or more pharmaceutically acceptable carriers; wherein each of: a compound represented by Formula (II):




15

Formula (II)


aniline and diphenyl urea are present in amount equal to or less than about 1.5% by weight based on the total weight of the composition.

[0134] Described herein, in an embodiment, is a pharmaceutical composition comprising: a) a solid dispersion comprising: an amorphous form of a compound represented by Formula (I)

**Formula (I)**


- and a pharmaceutically acceptable polymer; and b) one or more pharmaceutically acceptable carriers; wherein diphenyl urea is present in amount equal to or less than about 0.5% by weight based on the total weight of the composition.

[0135] Described herein, in an embodiment, is a pharmaceutically acceptable composition for a 50 mg single dosage unit of a compound represented by:



- comprising: an intragranular blend, wherein the intragranular blend comprises: a solid dispersion having 50 mg of the compound wherein the compound is present in amorphous form, hydroxypropyl methyl cellulose acetate succinate; a bulking agent and/or filler; and a lubricant and/or a glidant; and an extragranular blend comprising a glidant and/or a lubricant; wherein diphenyl urea is present in amount equal to or less than about 0.5% by weight based on the total weight of the composition.


- [0136]** Described herein, in an embodiment, is a pharmaceutically acceptable composition for orally delivering to a patient 50 mg of a compound represented by:



comprising: a solid dispersion having: 50 mg of the compound wherein the compound is present in amorphous form; hydroxypropyl methyl cellulose acetate succinate; a bulking

agent, a filler, and a lubricant and/or a glidant; wherein diphenyl urea is present in amount equal to or less than about 0.5% by weight based on the total weight of the composition.

[0137] Described herein, in an embodiment, is a pharmaceutically acceptable composition for orally delivering 50 mg of a compound represented by:



5

comprising:

a solid dispersion having 50 mg of the compound wherein the compound is present in amorphous form, and a polymer; about 25-45% by weight of a bulking agent based on the total amount of the pharmaceutical composition; about 25-45% by weight of a filler based on the total amount of the pharmaceutical composition; and a glidant and/or a lubricant; wherein the composition has not more than about 1.5% (w/w) of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one, when exposed to 60% relative humidity at 25 °C for 1 month, 3 months or 6 months.

[0138] In some embodiments, the polymer is hydroxypropyl methyl cellulose acetate succinate.

15


[0139] In some embodiments, the bulking agent is microcrystalline cellulose.

[0140] In some embodiments, the filler is lactose or a hydrate thereof.

[0141] In some embodiments, the composition is in the form of a tablet.

[0142] In some embodiments, the composition comprises less than 10% by weight of a compound represented by Formula (III):


20



Formula (III)

based on the weight of the compound of Formula (I).


[0143] In some embodiments, the composition comprises less than 7% by weight of a compound represented by Formula (III):



Formula (III)

5 based on the weight of the compound of Formula (I).


[0144] In some embodiments, the composition comprises less than 5% by weight of a compound represented by Formula (III):



Formula (III)

10 based on the weight of the compound of Formula (I).


[0145] In some embodiments, the composition comprises less than 3% by weight of a compound represented by Formula (III):



Formula (III)

15 based on the weight of the compound of Formula (I).


[0146] In some embodiments, the composition comprises less than 1% by weight of a compound represented by Formula (III):



Formula (III)

5 based on the weight of the compound of Formula (I).


[0147] In some embodiments, the composition comprises less than about 0.1 % by weight to about 0.5% by weight of a compound represented by Formula (III):



Formula (III)

10 based on the weight of the compound of Formula (I).

[0148] In some embodiments, the composition comprises about 0.01 % by weight to about 0.1% by weight of a compound represented by Formula (III):




Formula (III)

15 based on the weight of the compound of Formula (I).

[0149] The pharmaceutical compositions may also be provided as tablets. Tablets may be uncoated, film, or sugar coated bisected, embossed, plain, layered, or sustained-release. They can be made in a variety of sizes, shapes, and colors. Tablets may be swallowed, chewed, or dissolved in the buccal cavity or beneath the tongue.

[0150] In some embodiments, provided herein is a tablet providing about 50 mg of a compound represented by Formula (I), wherein the tablet comprises: (a) an intragranular blend comprising: (i) about 195 mg to about 205 mg of a solid spray-dried dispersion that comprises about 50 mg of the compound and hydroxypropyl methyl cellulose acetate succinate; (ii) about 177 mg to about 181 mg of microcrystalline cellulose; (iii) about 177 mg to about 181 mg of lactose monohydrate; and (iv) about 28 mg to about 32 mg of crospovidone; and (b) an extragranular blend comprising: (i) about 2 mg to about 4 mg of silicon dioxide; and (ii) about 2 mg to about 4 mg of magnesium stearate.

[0151] Described herein, in an embodiment, is a pharmaceutically acceptable tablet having 50 mg of a compound represented by:




wherein the tablet comprises: a solid dispersion having 50 mg of the compound wherein the compound is present in amorphous form, and hydroxypropyl methyl cellulose acetate succinate; about 25-35% by weight microcrystalline cellulose based on the total weight of the tablet; and about 25-35% by weight of lactose or a hydrate thereof based on the total amount of the pharmaceutical composition; wherein diphenyl urea is present in amount equal to or less than about 0.5% by weight based on the total weight of the composition. In some embodiments, the tablet further comprises at least one of: magnesium stearate, crospovidone and silicon dioxide.

[0152] In some embodiments, the tablet comprises less than or equal to about 10 % by weight of one or more anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises less than or equal to about 7 % by weight of one or more anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises less than or equal to about 5 % by weight of one or more anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises less than or equal to about 3 % by weight of one or more anilinic substances based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises less than about 10 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of

Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises less than about 7 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises less than about 5 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises less than about 3 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises less than about 2 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises less than about 1 % by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises about 0.1 % by weight to about 0.5% by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I). In some embodiments, the tablet comprises about 0.01 % by weight to about 0.1% by weight of 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (the compound of Formula (II)) based on the weight of the compound of Formula (I).


[0153] In some embodiments, the tablet comprises less than 10% by weight of a compound represented by Formula (III):



Formula (III)


based on the weight of the compound of Formula (I).

[0154] In some embodiments, the tablet comprises less than 7% by weight of a compound represented by Formula (III):

**Formula (III)**


based on the weight of the compound of Formula (I).

- [0155] In some embodiments, the tablet comprises less than 5% by weight of a compound
5 represented by Formula (III):

**Formula (III)**


based on the weight of the compound of Formula (I).

- [0156] In some embodiments, the tablet comprises less than 3% by weight of a compound
10 represented by Formula (III):

**Formula (III)**

based on the weight of the compound of Formula (I).


[0157] In some embodiments, the tablet comprises less than 1% by weight of a compound represented by Formula (III):



Formula (III)

5 based on the weight of the compound of Formula (I).


[0158] In some embodiments, the tablet comprises less than about 0.1 % by weight to about 0.5% by weight of a compound represented by Formula (III):



Formula (III)

10 based on the weight of the compound of Formula (I).

[0159] In some embodiments, the tablet comprises about 0.01 % by weight to about 0.1% by weight of a compound represented by Formula (III):



Formula (III)

15 based on the weight of the compound of Formula (I).

Methods of Treatment

[0160] A pharmaceutical composition comprising a compound of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical

composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) as described herein is a broad-spectrum inhibitor of c-KIT.

[0161] Disorders that can be treated by the compound of Formula (I) include, but are not limited to: gastrointestinal stromal tumors (GIST), NF-1-deficient gastrointestinal stromal tumors, succinate dehydrogenase (SDH)-deficient gastrointestinal stromal tumors, KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma, acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, non-small cell lung cancer, lung cancer, glioblastoma, a glioma, malignant peripheral nerve sheath sarcoma, hypereosinophilic syndrome, KIT driven germ cell tumor (e.g., testicular germ cell), KIT driven skin cancer, KIT driven renal cell carcinoma, penile cancer, PDGFRA driven penile cancer, prostate cancer, PDGFRA driven prostate cancer, PDGFRA driven non-melanoma skin cancer, PDGFRA driven glioma, PDGFRA driven sarcoma, PDGFRA driven glioblastoma, PDGFRA driven pancreatic cancer, or a disease vaginal cancer, prostate cancer, penile cancer, non-melanoma skin cancer, melanoma, or breast sarcoma (e.g., a vaginal cancer, prostate cancer, penile cancer, non-melanoma skin cancer, melanoma, or breast sarcoma comprising a PDGFRB mutation).

[0162] Accordingly, provided herein, in another embodiment, is a method of treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma, acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein.

[0163] Also provided herein, in another embodiment, is a method of treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, lung cancer, glioblastoma, a glioma, malignant peripheral nerve sheath sarcoma, and
5 hypereosinophilic syndrome in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein. In some embodiments, the disease is gastrointestinal stromal tumors (GIST).

[0164] Also provided herein, in another embodiment, is a method of treating a disease selected from the group consisting of KIT driven germ cell tumor (e.g., testicular germ cell),
10 KIT driven skin cancer, or KIT driven renal cell carcinoma in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein.

[0165] Also provided herein, in another embodiment, is a method of treating a disease selected from the group consisting of penile cancer, PDGFRA driven penile cancer, prostate
15 cancer, PDGFRA driven prostate cancer, PDGFRA driven non-melanoma skin cancer, PDGFRA driven glioma, PDGFRA driven sarcoma, PDGFRA driven glioblastoma, or PDGFRA driven pancreatic cancer in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein.

[0166] Also provided herein, in another embodiment, is a method of treating a disease
20 comprising a PDGFRB mutation selected from the group consisting of vaginal cancer, prostate cancer, penile cancer, non-melanoma skin cancer, melanoma, or breast sarcoma in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein.

[0167] In some embodiments, provided herein is a method for treating diseases driven by
25 KIT mutations or PDGFRA mutations in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition or one or more tablets described herein. In some embodiments, provided herein is a method for treating diseases driven by KIT mutations and PDGFRA mutations in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition or one or
30 more tablets described herein. In some embodiments, provided herein is a method for treating diseases driven by KIT mutations or PDGFRA mutations, comprising passenger PDGFRB mutations in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition described herein.

- [0168] In some embodiments, provided herein is a method for treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma (e.g., KIT driven melanoma or PDGFRA driven melanoma or PDGFR driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a composition or one or more tablets described herein. In some embodiments, the melanoma is cutaneous melanoma or noncutaneous melanoma. In some embodiments, the melanoma is cutaneous melanoma. In some embodiments, the cutaneous melanoma is superficial spreading melanoma, nodular melanoma, acral-lentiginous melanoma, or amelanotic and desmoplastic melanoma. In some embodiments, the melanoma is noncutaneous (non-skin) melanoma. In some embodiments, the noncutaneous melanoma is ocular melanoma or mucosal melanoma. In some embodiments, the disease is caused by the kinase activity of c-KIT and/or PDGFRA, and/or oncogenic forms thereof. In some embodiments, the disease is selected from the group consisting of KIT driven germ cell tumor (e.g., testicular germ cell), KIT driven skin cancer (e.g., KIT driven cutaneous squamous cell carcinoma, KIT driven Merkel cell carcinoma, uveal melanoma, non-melanoma skin cancer), or KIT driven renal cell carcinoma (e.g., renal cell carcinoma, chromophobe renal cell carcinoma). In some embodiments, the disease is selected from the group consisting of penile cancer, PDGFRA driven penile cancer, prostate cancer, PDGFRA driven prostate cancer, PDGFRA driven non-melanoma skin cancer, PDGFRA driven glioma, PDGFRA driven sarcoma, PDGFRA driven glioblastoma, or PDGFRA driven pancreatic cancer. In some embodiments, the disease comprising a PDGFRB mutation is selected from the group consisting of vaginal cancer, prostate cancer, penile cancer, non-melanoma skin cancer, melanoma, or breast sarcoma.
- [0169] Also provided herein, in another embodiment, is a use of a composition described herein for the preparation of a medicament for the treatment of a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma, acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell

- leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell
- 5 lymphoma, and non-small cell lung cancer. In some embodiments, the preparation of a medicament for the treatment of a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, lung cancer, glioblastoma, a glioma, malignant peripheral nerve sheath sarcoma, and hypereosinophilic syndrome.
- 10 **[0170]** In some embodiments, provided herein is a method of treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma (e.g., cutaneous melanoma, noncutaneous melanoma, KIT driven melanoma or PDGFRA driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or
- 15 dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer,
- 20 comprising administering to a patient in need thereof a therapeutically effective amount of a pharmaceutical composition comprising a compound of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I). In
- 25 some embodiments, the disease is caused by the kinase activity of: c-KIT and/or PDGFRA, and/or oncogenic forms thereof. In some embodiments, the disease is gastrointestinal stromal tumors (GIST). In some embodiments, the disease is KIT driven gastrointestinal stromal tumors. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors. In some embodiments, the disease is lung cancer. In some embodiments, the disease
- 30 is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, provided herein, is a method of treating or preventing a PDGFR kinase-mediated tumor growth of tumor progression comprising administering to a patient in need thereof a therapeutically effective amount of the

pharmaceutical composition comprising a compound of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I). In some
5 embodiments, the tumor growth or tumor progression is caused by PDGFR α kinase overexpression, oncogenic PDGFR α missense mutations, oncogenic deletion PDGFR α mutations, oncogenic PDGFR α gene rearrangements leading to PDGFR α fusion proteins, PDGFR α intragenic in-frame deletions, and/or oncogenic PDGFR α gene amplification. In some embodiments, a pharmaceutical
10 composition comprising a compound of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is administered to a cancer patient wherein the cancer is PDGFRA driven gastrointestinal stromal tumors, lung
15 adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, gastrointestinal stromal tumors, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, or lymphoblastic T-cell lymphoma. In some embodiments, the disease is PDGFRA driven
20 gastrointestinal stromal tumors (GIST). In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, a pharmaceutical composition comprising a compound of Formula (I) and one or more
25 pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is administered as a single agent or in combination with other cancer targeted therapeutic agents, cancer-targeted biologicals, immune checkpoint inhibitors, or chemotherapeutic
30 agents.

[0171] In some embodiments, the methods of treatment described herein comprise administering a composition of the compound of Formula (I) described herein, to a subject in need thereof prior to surgery (as a neoadjuvant therapy). In some embodiments, the methods of treatment described herein comprise administering a composition of the compound of

Formula (I) described herein, to a subject in need thereof after to surgery (as an adjuvant therapy).

[0172] A solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) as described herein is a broad-spectrum inhibitor of c-KIT. In some embodiments, provided herein is a method of treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma (e.g., cutaneous melanoma, noncutaneous melanoma, KIT driven melanoma or PDGFRA driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer, comprising administering to a patient in need thereof a therapeutically effective amount of the solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I). In some embodiments, the disease is caused by the kinase activity of c-KIT and/or PDGFRA, and/or oncogenic forms thereof. In some embodiments, the disease is gastrointestinal stromal tumors (GIST). In some embodiments, the disease is KIT driven gastrointestinal stromal tumors. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors. In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, provided herein, is a method of treating or preventing a PDGFR kinase-mediated tumor growth or tumor progression comprising administering to a patient in need thereof a therapeutically effective amount of a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I). In some embodiments, the tumor growth or tumor progression

- is caused by PDGFR α kinase overexpression, oncogenic *PDGFR α* missense mutations, oncogenic deletion *PDGFR α* mutations, oncogenic *PDGFR α* gene rearrangements leading to PDGFR α fusion proteins, *PDGFR α* intragenic in-frame deletions, and/or oncogenic *PDGFR α* gene amplification. In some embodiments, a solid dispersion
- 5 comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is administered to a cancer patient wherein the cancer is PDGFRA driven gastrointestinal stromal tumors, lung adenocarcinoma, squamous cell lung cancer,
- 10 glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, gastrointestinal stromal tumors, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, or lymphoblastic T-cell lymphoma. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors (GIST). In
- 15 some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, wherein the
- 20 pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is administered as a single agent or in combination with other cancer targeted therapeutic agents, cancer-targeted biologicals, immune checkpoint inhibitors, or chemotherapeutic agents.
- 25 **[0173]** A solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, having a purity by HPLC of greater than about 95% as described herein is a broad-spectrum inhibitor of c-KIT. In some embodiments, provided herein is a method of treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal
- 30 tumors, melanoma (e.g., cutaneous melanoma, noncutaneous melanoma, KIT driven melanoma or PDGFRA driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome,

idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer, comprising administering to a patient in need thereof a therapeutically effective amount of a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, having a purity by HPLC of greater than about 95%. In some embodiments, the disease is caused by the kinase activity of: c-KIT and/or PDGFRA, and/or oncogenic forms thereof. In some embodiments, the disease is gastrointestinal stromal tumors (GIST). In some embodiments, the disease is KIT driven gastrointestinal stromal tumors. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors. In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, provided herein, is a method of treating or preventing a PDGFR kinase-mediated tumor growth or tumor progression comprising administering to a patient in need thereof a therapeutically effective amount of a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, having a purity by HPLC of greater than about 95%. In some embodiments, the tumor growth or tumor progression is caused by PDGFR α kinase overexpression, oncogenic PDGFR α missense mutations, oncogenic deletion PDGFR α mutations, oncogenic PDGFR α gene rearrangements leading to PDGFR α fusion proteins, PDGFR α intragenic in-frame deletions, and/or oncogenic PDGFR α gene amplification. In some embodiments, a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, having a purity by HPLC of greater than about 95% is administered to a cancer patient wherein the cancer is PDGFRA driven gastrointestinal stromal tumors, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, gastrointestinal stromal tumors, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, or lymphoblastic T-cell lymphoma. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors (GIST). In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, a solid dispersion comprising a

compound of Formula (I) and a pharmaceutically acceptable polymer, having a purity by HPLC of greater than about 95% is administered as a single agent or in combination with other cancer targeted therapeutic agents, cancer-targeted biologicals, immune checkpoint inhibitors, or chemotherapeutic agents.

- 5 [0174] A compound of Formula (I), wherein the compound comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) as described herein is a broad-spectrum inhibitor of c-KIT. In some embodiments, provided herein is a method of treating a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven
- 10 gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma (e.g., cutaneous melanoma, noncutaneous melanoma, KIT driven melanoma or PDGFRA driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral
- 15 nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of Formula (I), wherein the compound comprises one or more anilinic substances
- 20 each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I). In some embodiments, the disease is caused by the kinase activity of: c-KIT and/or PDGFRA, and/or oncogenic forms thereof. In some embodiments, the disease is gastrointestinal stromal tumors (GIST). In some embodiments, the disease is KIT driven gastrointestinal stromal tumors. In some embodiments, the disease is PDGFRA
- 25 driven gastrointestinal stromal tumors. In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, provided herein, is a method of treating or preventing a PDGFR kinase-mediated tumor
- 30 growth of tumor progression comprising administering to a patient in need thereof a therapeutically effective amount of a compound of Formula (I), wherein the compound comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I). In some embodiments, the tumor growth or tumor progression is caused by PDGFR α kinase

overexpression, oncogenic PDGFR α missense mutations, oncogenic deletion PDGFR α mutations, oncogenic PDGFR α gene rearrangements leading to PDGFR α fusion proteins, PDGFR α intragenic in-frame deletions, and/or oncogenic PDGFR α gene amplification. In some embodiments, a compound of Formula (I),
5 wherein the compound comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is administered to a cancer patient wherein the cancer is PDGFRA driven gastrointestinal stromal tumors, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, gastrointestinal stromal tumors, malignant
10 peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, or lymphoblastic T-cell lymphoma. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors (GIST). In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the
15 disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, a compound of Formula (I), wherein the compound comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is administered as a single agent or in
20 combination with other cancer targeted therapeutic agents, cancer-targeted biologicals, immune checkpoint inhibitors, or chemotherapeutic agents.

[0175] In some embodiments, a pharmaceutical composition comprising a compound of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises one or more anilinic substances each
25 present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) as described herein is used in the preparation of a medicament for the treatment of a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma (e.g., cutaneous melanoma, noncutaneous melanoma, KIT driven
30 melanoma or PDGFRA driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-

associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer. In some embodiments, the disease is caused by the kinase activity of: c-KIT and/or PDGFRA, and/or oncogenic forms thereof. In some embodiments, the disease is gastrointestinal stromal tumors (GIST). In some embodiments, the disease is KIT driven

5 gastrointestinal stromal tumors. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors. In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some

10 embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, a pharmaceutical composition comprising a compound of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is used in the preparation of a medicament for treating or preventing a PDGFR kinase-mediated

15 tumor growth of tumor. In some embodiments, the tumor growth or tumor progression is caused by PDGFR α kinase overexpression, oncogenic PDGFR α missense mutations, oncogenic deletion PDGFR α mutations, oncogenic PDGFR α gene rearrangements leading to PDGFR α fusion proteins, PDGFR α intragenic in-frame deletions, and/or oncogenic PDGFR α gene amplification. In some embodiments, a pharmaceutical

20 composition comprising a compound of Formula (I) and one or more pharmaceutically acceptable carriers, excipients or diluents, wherein the pharmaceutical composition comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is used in the preparation of a medicament for the treatment of a disease wherein the disease is PDGFRA driven

25 gastrointestinal stromal tumors, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, gastrointestinal stromal tumors, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, or lymphoblastic T-cell lymphoma. In some

30 embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors (GIST). In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome.

[0176] In some embodiments, a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, wherein the solid dispersion comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) as described herein is used in the preparation of a medicament for the treatment of a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma (e.g., cutaneous melanoma, noncutaneous melanoma, KIT driven melanoma or PDGFRA driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer. In some embodiments, the disease is caused by the kinase activity of: c-KIT and/or PDGFRA, and/or oncogenic forms thereof. In some embodiments, the disease is gastrointestinal stromal tumors (GIST). In some embodiments, the disease is KIT driven gastrointestinal stromal tumors. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors. In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, wherein the solid dispersion comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is used in the preparation of a medicament for treating or preventing a PDGFR kinase-mediated tumor growth of tumor. In some embodiments, the tumor growth or tumor progression is caused by PDGFR α kinase overexpression, oncogenic PDGFR α missense mutations, oncogenic deletion PDGFR α mutations, oncogenic PDGFR α gene rearrangements leading to PDGFR α fusion proteins, PDGFR α intragenic in-frame deletions, and/or oncogenic PDGFR α gene amplification. In some embodiments, a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, wherein the solid dispersion comprises one or more anilinic substances each present in an amount equal to or less than about 3.0% by weight based on the weight of the compound of Formula (I) is used in the preparation of a

medicament for the treatment of a disease wherein the disease is PDGFRA driven gastrointestinal stromal tumors, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, gastrointestinal stromal tumors, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, or lymphoblastic T-cell lymphoma. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors (GIST). In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome.

[0177] In some embodiments, a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, having a purity by HPLC of greater than about 95% is used in the preparation of a medicament for the treatment of a disease selected from the group consisting of gastrointestinal stromal tumors (GIST), KIT driven gastrointestinal stromal tumors, PDGFRA driven gastrointestinal stromal tumors, melanoma (e.g., cutaneous melanoma, noncutaneous melanoma, KIT driven melanoma or PDGFRA driven melanoma), acute myeloid leukemia, germ cell tumors of the seminoma or dysgerminoma, mastocytosis, mast cell leukemia, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, lymphoblastic T-cell lymphoma, and non-small cell lung cancer. In some embodiments, the disease is caused by the kinase activity of: c-KIT and/or PDGFRA, and/or oncogenic forms thereof. In some embodiments, the disease is gastrointestinal stromal tumors (GIST). In some embodiments, the disease is KIT driven gastrointestinal stromal tumors. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors. In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome. In some embodiments, a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, having a purity by HPLC of greater than about 95% is used in the preparation of a medicament for treating or preventing a PDGFR kinase-mediated tumor growth of tumor. In some embodiments, the tumor growth or tumor progression is caused by PDGFR α kinase

overexpression, oncogenic PDGFR α missense mutations, oncogenic deletion PDGFR α mutations, oncogenic PDGFR α gene rearrangements leading to PDGFR α fusion proteins, PDGFR α intragenic in-frame deletions, and/or oncogenic PDGFR α gene amplification. In some embodiments, a solid dispersion comprising a compound of Formula (I) and a pharmaceutically acceptable polymer, having a purity by HPLC of greater than about 95% is used in the preparation of a medicament for the treatment of a disease wherein the disease is PDGFRA driven gastrointestinal stromal tumors, lung adenocarcinoma, squamous cell lung cancer, glioblastoma, glioma, pediatric glioma, astrocytomas, sarcomas, gastrointestinal stromal tumors, malignant peripheral nerve sheath sarcoma, intimal sarcomas, hypereosinophilic syndrome, idiopathic hypereosinophilic syndrome, chronic eosinophilic leukemia, eosinophilia-associated acute myeloid leukemia, or lymphoblastic T-cell lymphoma. In some embodiments, the disease is PDGFRA driven gastrointestinal stromal tumors (GIST). In some embodiments, the disease is lung cancer. In some embodiments, the disease is glioblastoma. In some embodiments, the disease is a glioma. In some embodiments, the disease is malignant peripheral nerve sheath sarcoma. In some embodiments, the disease is a hypereosinophilic syndrome.

[0178] In some embodiments, each anilinic substance is present in an amount equal to or less than about 5.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 4.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 2.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 1.0% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.7% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.5% by weight based on the weight of the compound of Formula (I). In some embodiments, each anilinic substance is present in an amount equal to or less than about 0.3% by weight based on the weight of the compound of Formula (I).

[0179] In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.30% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.30%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.20% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a

maximum of about 0.20%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.10% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.10%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.075% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.075%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.05% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.05%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.04% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.04%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.03% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.03%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.02% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.02%. In some embodiments, the diphenyl urea impurity is present in an amount equal to or less than about 0.01% by weight based on the weight of the compound of Formula (I) which means from about 0.0001% to a maximum of about 0.01%.

[0180] The pharmaceutical compositions described herein may be administered to patients (animals and humans) in need of such treatment in dosages that will provide optimal pharmaceutical efficacy. It will be appreciated that the dose required for use in any particular application will vary from patient to patient, not only with the particular compound or composition selected, but also with the route of administration, the nature of the condition being treated, the age and condition of the patient, concurrent medication or special diets then being followed by the patient, and other factors which those skilled in the art will recognize, with the appropriate dosage ultimately being at the discretion of the attendant physician.

[0181] Treatment can be continued for as long or as short a period as desired. The compositions may be administered on a regimen of, for example, one to four or more times per day. A suitable treatment period can be, for example, at least about one week, at least about two weeks, at least about one month, at least about six months, at least about 1 year, or indefinitely. A treatment period can terminate when a desired result is achieved.

Combination Therapy

[0182] The present disclosure describes combination therapies that involve the administration of the compound of Formula (I) or a pharmaceutically acceptable salt thereof, or a composition comprising the compound of Formula (I) or a pharmaceutically acceptable salt thereof described herein, and one or more therapeutic agents. The combination therapies described herein can be used by themselves, or in further combination with one or more additional therapeutic agents (e.g., one or more additional therapeutic agents described below). For example, the compound of Formula (I) or a pharmaceutically acceptable salt thereof, or a composition comprising the compound of Formula (I) or a pharmaceutically acceptable salt thereof described herein can be administered together with a cancer targeted therapeutic agent, a cancer-targeted biological, an immune checkpoint inhibitor, or a chemotherapeutic agent. The therapeutic agents can be administered together with or sequentially with another therapeutic agent described herein in a combination therapy.

[0183] Combination therapy can be achieved by administering two or more therapeutic agents, each of which is formulated and administered separately. In one embodiment, a composition comprising the compound of Formula (I) or a pharmaceutically acceptable salt thereof described herein is administered in a separate formulation than a formulation comprising the one or more additional therapeutic agents, e.g., one or more additional therapeutic agents described herein. Alternatively, combination therapy can be achieved by administering two or more therapeutic agents in a single formulation.

[0184] Other combinations are also encompassed by combination therapy. While the two or more agents in the combination therapy can be administered simultaneously, they need not be. For example, administration of a first agent (or combination of agents) can precede administration of a second agent (or combination of agents) by minutes, hours, days, or weeks. Thus, the two or more agents can be administered within minutes of each other or within 1, 2, 3, 6, 9, 12, 15, 18, or 24 hours of each other or within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14 days of each other or within 2, 3, 4, 5, 6, 7, 8, 9, or weeks of each other. In some cases even longer intervals are possible. While in many cases it is desirable that the two or more agents used in a combination therapy be present in within the patient's body at the same time, this need not be so.

[0185] Combination therapy can also include two or more administrations of one or more of the agents used in the combination using different sequencing of the component agents. For example, if agent X and agent Y are used in a combination, one could administer them

sequentially in any combination one or more times, e.g., in the order X-Y-X, X-X-Y, Y-X-Y, Y-Y-X, X-X-Y-Y, etc.

[0186] In some embodiments, the additional therapeutic agent that may be administered according to the present disclosure include, but are not limited to, cytotoxic agents, cisplatin, doxorubicin, etoposide, irinotecan, topotecan, paclitaxel, docetaxel, the epothilones, tamoxifen, 5-fluorouracil, methotrexate, temozolomide, cyclophosphamide, lonafarib, tipifarnib, 4-((5-((4-(3-chlorophenyl)-3-oxopiperazin-1-yl)methyl)-1H-imidazol-1-yl)methyl)benzonitrile hydrochloride, (R)-1-((1H-imidazol-5-yl)methyl)-3-benzyl-4-(thiophen-2-ylsulfonyl)-2,3,4,5-tetrahydro-1H-benzo diazepine-7-carbonitrile, cetuximab, imatinib, interferon alfa-2b, pegylated interferon alfa-2b, aromatase combinations, gemcitabine, uracil mustard, chlormethine, ifosfamide, melphalan, chlorambucil, pipobroman, triethylenemelamine, triethylenethiophosphoramine, busulfan, carmustine, lomustine, streptozocin, dacarbazine, floxuridine, cytarabine, 6-mercaptopurine, 6-thioguanine, fludarabine phosphate, leucovorin, oxaliplatin, pentostatine, vinblastine, vincristine, vindesine, bleomycin, dactinomycin, daunorubicin, epirubicin, idarubicin, mithramycin, deoxycoformycin, mitomycin-C, L-asparaginase, teniposide 17 α -ethinyl estradiol, diethylstilbestrol, testosterone, prednisone, fluoxymesterone, dromostanolone propionate, testolactone, megestrol acetate, methylprednisolone, methyltestosterone, prednisolone, triamcinolone, chlorotrianisene, 17 α -hydroxyprogesterone, aminoglutethimide, estramustine, medroxyprogesterone acetate, leuprolide acetate, flutamide, toremifene citrate, goserelin acetate, carboplatin, hydroxyurea, amsacrine, procarbazine, mitotane, mitoxantrone, levamisole, vinorelbine, anastrozole, letrozole, capecitabine, raloxifene, droloxafine, hexamethylmelamine, bevacizumab, trastuzumab, tositumomab, bortezomib, ibritumomab tiuxetan, arsenic trioxide, porfimer sodium, cetuximab, thioTEPA, altretamine, fulvestrant, exemestane, rituximab, alemtuzumab, dexamethasone, bicalutamide, chlorambucil, and valrubicin.

[0187] In some embodiments, the additional therapeutic agent that can be administered may include, without limitation, an AKT inhibitor, alkylating agent, all-trans retinoic acid, antiandrogen, azacitidine, BCL2 inhibitor, BCL-XL inhibitor, BCR-ABL inhibitor, BTK inhibitor, BTK/LCK/LYN inhibitor, CDK1/2/4/6/7/9 inhibitor, CDK4/6 inhibitor, CDK9 inhibitor, CBP/p300 inhibitor, EGFR inhibitor, endothelin receptor antagonist, RAF inhibitor, MEK inhibitor, ERK inhibitor, farnesyltransferase inhibitor, FLT3 inhibitor, glucocorticoid receptor agonist, HDM2 inhibitor, histone deacetylase inhibitor, IKK β inhibitor, immunomodulatory drug (IMiD), ingenol, ITK inhibitor, JAK1/JAK2/JAK3/TYK2 inhibitor,

MTOR inhibitor, PI3 kinase inhibitor, dual PI3 kinase/MTOR inhibitor, proteasome inhibitor, protein kinase C agonist, SUV39H1 inhibitor, TRAIL, VEGFR2 inhibitor, Wnt/ β -catenin signaling inhibitor, decitabine, and anti-CD20 monoclonal antibody.

- [0188]** In some embodiments, the additional therapeutic agent is an immunomodulatory agent selected from the group consisting of CTLA4 inhibitors such as, but not limited to ipilimumab and tremelimumab; PD1 inhibitors such as, but not limited to pembrolizumab, and nivolumab; PDL1 inhibitors such as, but not limited to atezolizumab (formerly MPDL3280A), durvalumab (formerly MEDI4736), avelumab, PDR001; 4 1BB or 4 1BB ligand inhibitors such as, but not limited to urelumab and PF-05082566; OX40 ligand agonists such as, but not limited to MEDI6469; GITR agents such as, but not limited to TRX518; CD27 inhibitors such as, but not limited to varlilumab; TNFRSF25 or TL1A inhibitors; CD40 agonists such as, but not limited to CP-870893; HVEM or LIGHT or LTA or BTLA or CD160 inhibitors; LAG3 inhibitors such as, but not limited to BMS-986016; TIM3 inhibitors; Siglecs inhibitors; ICOS or ICOS ligand agonists; B7 H3 inhibitors such as, but not limited to MGA271; B7 H4 inhibitors; VISTA inhibitors; HHLA2 or TMIGD2 inhibitors; inhibitors of Butyrophilins, including BTNL2 inhibitors; CD244 or CD48 inhibitors; inhibitors of TIGIT and PVR family members; KIRs inhibitors such as, but not limited to lirilumab; inhibitors of ILTs and LIRs; NKG2D and NKG2A inhibitors such as, but not limited to IPH2201; inhibitors of MICA and MICB; CD244 inhibitors; CSF1R inhibitors such as, but not limited to emactuzumab, cabiralizumab, pexidartinib, ARRY382, BLZ945; IDO inhibitors such as, but not limited to INCB024360; thalidomide, lenalidomide, TGF β inhibitors such as, but not limited to galunisertib; adenosine or CD39 or CD73 inhibitors; CXCR4 or CXCL12 inhibitors such as, but not limited to ulocuplumab and (3S,6S,9S,12R,17R,20S,23S,26S,29S,34aS)-N-((S)-1-amino-5-guanidino-1-oxopentan-2-yl)-26,29-bis(4-aminobutyl)-17-((S)-2-((S)-2-((S)-2-(4-fluorobenzamido)-5-guanidinopentanamido)-5-guanidinopentanamido)-3-(naphthalen-2-yl)propanamido)-6-(3-guanidinopropyl)-3,20-bis(4-hydroxybenzyl)-1,4,7,10,18,21,24,27,30-nonaaxo-9,23-bis(3-ureidopropyl)triacontahydro-1H,16H-pyrrolo[2,1-p][1,2]dithia[5,8,11,14,17,20,23,26,29]nonaazacyclodotriacontine-12-carboxamide BKT140; phosphatidylserine inhibitors such as, but not limited to bavituximab; SIRPA or CD47 inhibitors such as, but not limited to CC-90002; VEGF inhibitors such as, but not limited to bevacizumab; and neuropilin inhibitors such as, but not limited to MNRP1685A.

[0189] In some embodiments, the additional therapeutic agent is a chemotherapeutic agent selected from the group consisting of chemotherapeutic agents including but not limited

to anti-tubulin agents (paclitaxel, paclitaxel protein-bound particles for injectable suspension such as nab-paclitaxel, eribulin, docetaxel, ixabepilone, vincristine), vinorelbine, DNA-alkylating agents (including cisplatin, carboplatin, oxaliplatin, cyclophosphamide, ifosfamide, temozolomide), DNA intercalating agents (including doxorubicin, pegylated liposomal
5 doxorubicin, daunorubicin, idarubicin, and epirubicin), 5-fluorouracil, capecitabine, cytarabine, decitabine, 5-aza cytidine, gemcitabine and methotrexate.

[0190] In some embodiments, the additional therapeutic agent is selected from the group consisting of paclitaxel, paclitaxel protein-bound particles for injectable suspension, eribulin, docetaxel, ixabepilone, vincristine, vinorelbine, cisplatin, carboplatin, oxaliplatin,
10 cyclophosphamide, ifosfamide, temozolomide, doxorubicin, pegylated liposomal doxorubicin, daunorubicin, idarubicin, epirubicin, 5-fluorouracil, capecitabine, cytarabine, decitabine, 5-azacytidine, gemcitabine, methotrexate, erlotinib, gefitinib, lapatinib, everolimus, temsirolimus, LY2835219, LEE011, PD 0332991, crizotinib, cabozantinib, sunitinib, pazopanib, sorafenib, regorafenib, axitinib, dasatinib, imatinib, nilotinib,
15 vemurafenib, dabrafenib, trametinib, idelasib, quizartinib, tamoxifen, fulvestrant, anastrozole, letrozole, exemestane, abiraterone acetate, enzalutamide, nilutamide, bicalutamide, flutamide, cyproterone acetate, prednisone, dexamethasone, irinotecan, camptothecin, topotecan, etoposide, etoposide phosphate, mitoxantrone, vorinostat, romidepsin, panobinostat, valproic acid, belinostat, DZNep 5-aza-2'-deoxycytidine, bortezomib, carfilzomib, thalidomide,
20 lenalidomide, pomalidomide, trastuzumab, pertuzumab, cetuximab, panitumumab, ipilimumab, labrolizumab, nivolumab, MPDL3280A, bevacizumab, aflibercept, brentuximab vedotin, ado-trastuzumab emtansine, radiotherapy, and sipuleucel T.

[0191] In some embodiments, the additional therapeutic agent is a kinase inhibitor selected from the group consisting of erlotinib, gefitinib, lapatinib, everolimus, temsirolimus,
25 LY2835219, LEE011, PD 0332991, crizotinib, cabozantinib, sunitinib, pazopanib, sorafenib, regorafenib, axitinib, dasatinib, imatinib, nilotinib, vemurafenib, dabrafenib, trametinib, idelalisib, and quizartinib.

[0192] In some embodiments, the additional therapeutic agent is an anti-PD1 therapeutic. Examples of anti-PD1 therapeutics that may be administered in combination with the
30 compound of Formula (I) or pharmaceutically acceptable salt thereof or a composition comprising the compound of Formula (I) or pharmaceutically acceptable salt thereof described herein include, but are not limited to, nivolumab, pidilizumab, cemiplimab, tislelizumab, AMP-224, AMP-514, and pembrolizumab.

[0193] In some embodiments, the additional therapeutic agent is selected from the group consisting of immunomodulatory agents including but not limited to anti-PD-L1 therapeutics including atezolizumab, durvalumab, BMS-936559, and avelumab, anti-TIM3 therapeutics including TSR-022 and MBG453, anti-LAG3 therapeutics including relatlimab, LAG525, and TSR-033, CD40 agonist therapeutics including SGN-40, CP-870,893 and RO7009789, anti-CD47 therapeutics including Hu5F9-G4, anti-CD20 therapeutics, anti-CD38 therapeutics, and other immunomodulatory therapeutics including thalidomide, lenalidomide, pomalidomide, prednisone, and dexamethasone. In some embodiments, the additional therapeutic agent is avelumab.

[0194] In some embodiments, the additional therapeutic agent is a chemotherapeutic agent selected from the group consisting of anti-tubulin agents (e.g., paclitaxel, paclitaxel protein-bound particles for injectable suspension, eribulin, abraxane, docetaxel, ixabepilone, taxiterem, vincristine or vinorelbine), LHRH antagonists including but not limited to leuprolide, goserelin, triptorelin, or histrelin, anti-androgen agents including but not limited to abiraterone, flutamide, bicalutamide, nilutamide, cyproterone acetate, enzalutamide, and apalutamide, , anti-estrogen agents including but not limited to tamoxifen, fulvestrant, anastrozole, letrozole, and exemestane, DNA-alkylating agents (including cisplatin, carboplatin, oxaliplatin, cyclophosphamide, ifosfamide, and temozolomide), DNA intercalating agents (including doxorubicin, pegylated liposomal doxorubicin, daunorubicin, idarubicin, and epirubicin), 5-fluorouracil, capecitabine, cytarabine, decitabine, 5-aza cytidine, gemcitabine methotrexate, bortezomib, and carfilzomib.

[0195] In some embodiments, the additional therapeutic agent is selected from the group consisting of targeted therapeutics including kinase inhibitors erlotinib, gefitinib, lapatanib, everolimus, temsirolimus, abemaciclib, LEE011, palbociclib, crizotinib, cabozantinib, sunitinib, pazopanib, sorafenib, regorafenib, axitinib, dasatinib, imatinib, nilotinib, vemurafenib, dabrafenib, trametinib, cobimetinib, binimetinib, idelalisib, quizartinib, avapritinib, BLU-667, BLU-263, Loxo 292, larotrectinib, and quizartinib, anti-estrogen agents including but not limited to tamoxifen, fulvestrant, anastrozole, letrozole, and exemestane, anti-androgen agents including but not limited to abiraterone acetate, enzalutamide, nilutamide, bicalutamide, flutamide, cyproterone acetate, steroid agents including but not limited to prednisone and dexamethasone, PARP inhibitors including but not limited to neraparib, olaparib, and rucaparib, topoisomerase I inhibitors including but not limited to irinotecan, camptothecin, and topotecan, topoisomerase II inhibitors including but not limited to etoposide, etoposide phosphate, and mitoxantrone, Histone Deacetylase

(HDAC) inhibitors including but not limited to vorinostat, romidepsin, panobinostat, valproic acid, and belinostat, DNA methylation inhibitors including but not limited to DZNep and 5-aza-2'-deoxycytidine, proteasome inhibitors including but not limited to bortezomib and carfilzomib, thalidomide, lenalidomide, pomalidomide, biological agents including but not limited to trastuzumab, ado-trastuzumab, pertuzumab, cetuximab, panitumumab, ipilimumab, tremelimumab, vaccines including but not limited to sipuleucel-T, and radiotherapy.

[0196] In some embodiments, the additional therapeutic agent is selected from the group consisting of an inhibitor of the TIE2 immunokinase including rebastinib or ARRY-614.

[0197] In some embodiments, the additional therapeutic agent is selected from the group consisting of an inhibitor of the TIE2 immunokinase including rebastinib or ARRY-614, and an anti-PD1 therapeutic.

[0198] In some embodiments, the additional therapeutic agent is selected from the group consisting of anti-angiogenic agents including AMG386, bevacizumab and aflibercept, and antibody-drug-conjugates (ADCs) including brentuximab vedotin, trastuzumab emtansine, and ADCs containing a payload such as a derivative of camptothecin, a pyrrolobenzodiazepine dimer (PBD), an indolinobenzodiazepine dimer (IGN), DM1, DM4, MMAE, or MMAF.

[0199] In some embodiments, the additional therapeutic agent is selected from a luteinizing hormone-releasing hormone (LHRH) analog, including goserelin and leuprolide.

[0200] In some embodiments, the additional therapeutic agent is selected from the group consisting of selected from the group consisting of everolimus, trabectedin, abraxane, TLK 286, AV-299, DN-101, pazopanib, GSK690693, RTA 744, ON 0910.Na, AZD 6244 (ARRY-142886), AMN-107, TKI-258, GSK461364, AZD 1152, enzastaurin, vandetanib, ARQ-197, MK-0457, MLN8054, PHA-739358, R-763, AT-9263, pemetrexed, erlotinib, dasatinib, nilotinib, decatanib, panitumumab, amrubicin, oregovomab, Lep-et, nolatrexed, azd2171, batabulin, ofatumumab, zanolimumab, edotecarin, tetrandrine, rubitecan, tesmilifene, oblimersen, ticilimumab, ipilimumab, gossypol, Bio 111, 131-I-TM-601, ALT-110, BIO 140, CC 8490, cilengitide, gimatecan, IL13-PE38QQR, INO 1001, IPdR1 KRX-0402, lucanthone, LY 317615, neuradiab, vitespan, Rta 744, Sdx 102, talampanel, atrasentan, Xr 311, romidepsin, ADS-100380, sunitinib, 5-fluorouracil, vorinostat, etoposide, gemcitabine, doxorubicin, irinotecan, liposomal doxorubicin, 5'-deoxy-5-fluorouridine, vincristine, temozolomide, ZK-304709, seliciclib; PD0325901, AZD-6244, capecitabine, L-Glutamic acid, N-[4-[2-(2-amino-4,7-dihydro-4-oxo-1H-pyrrolo[2,3-d]pyrimidin-5-yl)-ethyl]benzoyl]-

, disodium salt, heptahydrate, camptothecin, PEG-labeled irinotecan, tamoxifen, toremifene citrate, anastrozole, exemestane, letrozole, DES(diethylstilbestrol), estradiol, estrogen, conjugated estrogen, bevacizumab, IMC-1C11, CHIR-258,); 3-[5-(methylsulfonylpiperadinemethyl)-indolyl]-quinolone, vatalanib, AG-013736, AVE-0005, the

5 acetate salt of [D-Ser(Bu t) 6, Azgly 10] (pyro-Glu-His-Trp-Ser-Tyr-D-Ser(Bu t)-Leu-Arg-Pro-Azgly-NH₂ acetate [C₅₉H₈₄N₁₈O_{i4}-(C₂H₄O₂)_x where x=1 to 2.4], goserelin acetate, leuprolide acetate, triptorelin pamoate, medroxyprogesterone acetate, hydroxyprogesterone caproate, megestrol acetate, raloxifene, bicalutamide, flutamide, nilutamide, megestrol acetate, CP-724714; TAK-165, HKI-272, erlotinib, lapatanib, canertinib, ABX-EGF

10 antibody, erbitux, EKB-569, PKI-166, GW-572016, Ionafarnib, BMS-214662, tipifarnib; amifostine, NVP-LAQ824, suberoyl analide hydroxamic acid, valproic acid, trichostatin A, FK-228, SU11248, sorafenib, KRN951, aminoglutethimide, arnsacrine, anagrelide, L-asparaginase, Bacillus Calmette-Guerin (BCG) vaccine, bleomycin, buserelin, busulfan, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, cyproterone,

15 cytarabine, dacarbazine, dactinomycin, daunorubicin, diethylstilbestrol, epirubicin, fludarabine, fludrocortisone, fluoxymesterone, flutamide, gemcitabine, gleevac, hydroxyurea, idarubicin, ifosfamide, imatinib, leuprolide, levamisole, lomustine, mechlorethamine, melphalan, 6-mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, octreotide, oxaliplatin, pamidronate, pentostatin, plicamycin, porfimer,

20 procarbazine, raltitrexed, rituximab, streptozocin, teniposide, testosterone, thalidomide, thioguanine, thiotepa, tretinoin, vindesine, 13-cis-retinoic acid, phenylalanine mustard, uracil mustard, estramustine, altretamine, floxuridine, 5-deoxyuridine, cytosine arabinoside, 6-mercaptopurine, deoxycytosine, calcitriol, valrubicin, mithramycin, vinblastine, vinorelbine, topotecan, razoxin, marimastat, COL-3, neovastat, BMS-275291, squalamine,

25 endostatin, SU5416, SU6668, EMD121974, interleukin-12, IM862, angiostatin, vitaxin, droloxifene, idoxifene, spironolactone, finasteride, cimitidine, trastuzumab, denileukin diftitox, gefitinib, bortezomib, irinotecan, topotecan, doxorubicin, docetaxel, vinorelbine, bevacizumab (monoclonal antibody) and erbitux, cremophor-free paclitaxel, epithilone B, BMS-247550, BMS-310705, droloxifene, 4-hydroxytamoxifen, pipendoxifene, ERA-923,

30 arzoxifene, fulvestrant, acolbifene, lasofoxifene, idoxifene, TSE-424, HMR-3339, ZK186619, PTK787/ZK 222584, VX-745, PD 184352, rapamycin, 40-O-(2-hydroxyethyl)-rapamycin, temsirolimus, AP-23573, RAD001, ABT-578, BC-210, LY294002, LY292223, LY292696, LY293684, LY293646, wortmannin, ZM336372, L-779,450, PEG-filgrastim, darbepoetin, erythropoietin, granulocyte colony-stimulating factor, zolendronate, prednisone,

cetuximab, granulocyte macrophage colony-stimulating factor, histrelin, pegylated interferon alfa-2a, interferon alfa-2a, pegylated interferon alfa-2b, interferon alfa-2b, azacitidine, PEG-L-asparaginase, lenalidomide, gemtuzumab, hydrocortisone, interleukin-11, dexrazoxane, alemtuzumab, all-transretinoic acid, ketoconazole, interleukin-2, megestrol, immune
 5 globulin, nitrogen mustard, methylprednisolone, ibritumomab tiuxetan, androgens, decitabine, hexamethylmelamine, bexarotene, tositumomab, arsenic trioxide, cortisone, editronate, mitotane, cyclosporine, liposomal daunorubicin, Edwina-asparaginase, strontium
 10 89, casopitant, netupitant, an NK-1 receptor antagonists, palonosetron, aprepitant, diphenhydramine, hydroxyzine, metoclopramide, lorazepam, alprazolam, haloperidol, droperidol, dronabinol, dexamethasone, methylprednisolone, prochlorperazine, granisetron, ondansetron, dolasetron, tropisetron, pegfilgrastim, erythropoietin, epoetin alfa and darbepoetin alfa, ipilimumab, vemurafenib, and mixtures thereof.


[0201] In some embodiments, the additional therapeutic agent is an HSP90 inhibitor (e.g., AT13387). In some embodiments, the additional therapeutic agent is cyclophosphamide. In
 15 some embodiments, the additional therapeutic agent is an AKT inhibitor (e.g., perifosine). In some embodiments, the additional therapeutic agent is a BCR-ABL inhibitor (e.g., nilotinib). In some embodiments, the additional therapeutic agent is an mTOR inhibitor (e.g., RAD001). In some embodiments, the additional therapeutic agent is an FGFR inhibitor (e.g., erdafitinib, KO947, or BGJ398). In some embodiments, the additional therapeutic agent is an anti-PDL1
 20 therapeutic. In some embodiments, the additional therapeutic agent is a Bcl2 inhibitor (e.g., venetoclax). In some embodiments, the additional therapeutic agent is an autophagy inhibitor (e.g., hydroxychloroquine). In some embodiments, the additional therapeutic agent is a MET inhibitor.

EXAMPLES

25 **[0202]** The present disclosure is not to be limited in scope by the specific embodiments disclosed in the examples which are intended as illustrations of a few aspects of the disclosure and any embodiments that are functionally equivalent are within the scope of this disclosure. Indeed, various modifications in addition to those shown and described herein
 30 will become apparent to those skilled in the art and are intended to fall within the scope of the appended claims.

[0203] In the Examples provided below, the following abbreviations are used: “HPMCAS-HG” refers to hydroxymethylpropyl cellulose acetate succinate (high PH solubility grade); “SDD” refers to spray-dried dispersion; and “PVP-XL” refers to cross-

linked polyvinylpyrrolidone. "Compound 1" refers to the compound of Formula (I) described herein. "Compound 2" refers to the compound 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one, which has the structure:



5 [0204] Unless otherwise stated, percentages amounts of the compound of Formula (I) in the solid dispersions described below indicate weight percentages of the compound of Formula (I) with respect to the total weight of the solid dispersion.

[0205] As used below in Example 1, the "w/w suspension fraction" is the fraction of a component, as a weight percentage, of the suspension used to prepare the spray-dried
10 dispersion based on the amount of Compound 1 in the suspension.

[0206] Example 1. Preparation of a Spray-Dried Dispersion Comprising Compound 1 and HPMCAS-HG.

[0207] Suspension preparation. HPMCAS-HG is added to a purified water and acetone solution and mixed to ensure dissolution of the polymer. Compound 1 is added to the
15 solution, and the suspension is mixed at a temperature of 15-25 °C. The mixing remains on for the remainder of spray drying process.

[0208] Startup/shutdown solvent preparation and use. Purified water and acetone are mixed. The startup and shutdown solvent are sprayed at the beginning and end of the spray drying cycle.

20 [0209] Spray Drying. The suspension is passed through an inline heat exchanger (flow rate of 38-51 kg/hr) which heats the suspension to a temperature range of 112-124 °C to dissolve the suspended particles prior to spray drying. The solution is then spray dried in a pharmaceutical spray dryer (PSD-2 or equivalent) equipped with a capillary nozzle assisted with nitrogen sheath gas pressure of 65-85 psig using 400-500 kg/hr bulk drying gas, 50-70°C
25 chamber outlet temperature and -10 °C condenser temperature.

[0210] Secondary drying. The partially wet spray-dried intermediate resulting from the preparation described above is dried to provide an SDD comprising Compound 1 and HPMCAS-HG using agitated vacuum dryer at temperature range of 40-50 °C and chamber pressure of 40-50 mbar.

**Analytical Method for Determining Amount of Impurities in Compound of Formula (I)
(Compound 1).****Equipment, Reagents, and Impurity Marker Solutions**

Equipment	
HPLC system	Suitable RP-HPLC equipped with a photodiode array UV detector and data system.
Column	Zorbax Bonus RP, 4.6 x 150 mm, 3.5 μ m (Agilent) equipped with HPLC column filter (0.5 μ m), or equivalent

PLC Instrument Parameters for Identification, Assay, Degradation Products, and Uniformity of Dosage Units

Parameter	Value		
Column temperature	40°C		
Autosampler temperature	20°C		
Flow rate	1.0 mL/min		
Injection volume	3 µL		
Detection	UV at 240 nm Spectra collection 200 – 400 nm		
Acquisition run time	20 minutes		
Mobile phase	Mobile Phase A: purified water Mobile Phase B: acetonitrile		
Seal and needle wash	THF:HPLC-grade water:FA, 75:25:0.1 (v/v/v)		
Diluent	THF:purified water:FA, 75:25:0.1 (v/v/v)		
Gradient	Minutes	% Mobile Phase A	% Mobile Phase B
	0	80	20
	6.0	20	80
	16.0	0	100
	16.1	80	20
	20.0	80	20
Post analysis column wash	Post analysis column wash shall be performed in accordance with testing site procedure		

Abbreviations: FA: formic acid; THF: tetrahydrofuran; UV: ultraviolet; v: volume.

Characterization of Impurities from batches prepared according to Example 1.

Attribute	Impurities	Lot 1 made by the process of example 1	Lot 2 made by the process of example 1	Lot 3 made by the process of example 1	Lot 4 made by the process of example 1
Related Substances (% w/w with respect to the weight of Compound 1)	Impurity A	0.14%	0.13%	0.16%	0.15%
	Impurity B	< 0.05%	< 0.05%	< 0.05%	< 0.05%
	Diphenyl urea	< 0.05%	< 0.05%	< 0.05%	< 0.05%

5

Example 2. Preparation of a Spray-Dried Dispersion Comprising Compound 1 and HPMCAS-HG.

[0211] *Solution preparation.* Compound 1 is added to a purified water and THF solution and mixed to ensure dissolution of the compound. HPMCAS-HG is added to the solution and mixed at ambient temperature until the polymer dissolves. *Startup/shutdown solvent*

10

preparation and use. Purified water and THF are mixed. The startup and shutdown solvent are sprayed at the beginning and end of the spray drying cycle.

- [0212] *Spray Drying.* The solution is then spray dried in a pharmaceutical spray dryer at 175-205 g/min spray rate using 1550-2150 g/min bulk drying gas flow rate and 40-50 °C chamber outlet temperature.

[0213] *Secondary drying.* The partially wet spray-dried intermediate resulting from the preparation described above is dried to provide an SDD comprising Compound 1 and HPMCAS-HG using tray dryer at temperature range of 15-45 °C.

10 **Example 3. Purity studies of a solid dispersion of the compound of Formula (I).**

[0214] Purity studies on samples (Lot 1, Lot 2, Lot 3, and Lot 4) of a solid dispersion of the compound of Formula (I) were conducted using HPLC. Each lot was prepared according to the process outlined in Example 1. Results of the study are shown in Table 1 below.

15 **Table 1: Purity by HPLC for a solid dispersion of the compound of Formula (I).**

Lot Number	Lot 1	Lot 2	Lot 3	Lot 4
Appearance	White powder	White powder	White powder	White powder
Identification (HPLC)	Rep 1: 100.0% Rep 2: 99.9%	Rep 1: 100.0% Rep 2: 100.0%	Rep 1: 100.0% Rep 2: 100.0%	Rep 1: 100.0% Rep 2: 100.0%
Compound 1 Assay (% w/w)	24.8	24.9	24.9	24.9
Compound 2 (% w/w with respect to the weight of Compound 1)	0.14	0.13	0.13	0.13

Legend: LOD: Levels of Detection *Solid Dispersions*.

Example 4. Preparation of compound of Formula (III) reference standard.

- [0215] 3-(5-amino-2-bromo-4-fluorophenyl)-1-ethyl-7-(methylamino)-1,6-naphthyridin-2(1H)-one (40 g), phenyl isocyanate (30 g, 2.7 equiv.), pyridine (3 eq) and methanesulfonic acid (1 eq) were combined in a solvent comprised of 1-methyl-2-pyrrolidinone (10 vol) and tetrahydrofuran (5 vol). The mixture was stirred at 50 °C for 7 days with occasional addition of additional 0.1 - 0.2 eq of phenyl isocyanate (0.1-0.2 eq) to obtain crude 1-(3-(2-bromo-4-fluoro-5-(3-phenylureido)phenyl)-1-ethyl-2-oxo-1,2-dihydro-1,6-naphthyridin-7-yl)-1-methyl-3-phenylurea wet cake. The crude wetcake was crystallized from 1-methyl-2-pyrrolidinone (4 vol) and methanol (8 vol) to obtain 57 g of 1-(3-(2-bromo-4-fluoro-5-(3-

phenylureido)phenyl)-1-ethyl-2-oxo-1,2-dihydro-1,6-naphthyridin-7-yl)-1-methyl-3-phenylurea. MS m/z: 629 (M+1). ¹H NMR (400 MHz, DMSO-d₆): δ 11.44 (s, 1 H), 9.12 (s, 1 H), 8.84 (s, 1 H), 8.74 (s, 1 H), 8.29 and 8.27 (d, 1 H), 8.02 (s, 1 H), 7.72 and 7.70 (d, 1 H), 7.59 and 7.57 (d, 2 H), 7.45 and 7.43 (d, 2 H), 7.34-7.26 (m, 4 H), 7.24 (s, 1 H), 7.06-6.97 (m, 2 H), 4.35-4.28 (m, 2 H), 3.53 (s, 3 H), 1.27-1.23 (t, 3 H).

EQUIVALENTS

[0216] Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific embodiments described specifically herein. Such equivalents are intended to be encompassed in the scope of the following claims.