

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



1806264-8
1803 2017

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07C 403/16	A1	(11) International Publication Number: WO 98/49139 (43) International Publication Date: 5 November 1998 (05.11.98)
(21) International Application Number: PCT/EP98/02407 (22) International Filing Date: 23 April 1998 (23.04.98) (30) Priority Data: 97107175.8 30 April 1997 (30.04.97) EP (34) Countries for which the regional or international application was filed: DE et al. (71) Applicant (for all designated States except US): GIVAUDAN-ROURE (INTERNATIONAL) S.A. [CH/CH]; Vernier, CH-1412 Genève (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): COURTOIS, Didier [FR/FR]; 87 bis, rue de Coulmiers, F-45000 Orléans (FR). EHRET, Charles [FR/FR]; Usterstrasse 65, CH-8620 Wetzikon (CH). FIRMIN, Laurence, Marthe, Marie [FR/FR]; 23, rue Joseph Malleville, F-76290 Fontaine-la-Mallet (FR). (74) Common Representative: GIVAUDAN-ROURE (INTERNATIONAL) S.A.; P.O. Box 3255, CH-4002 Basle (CH).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PROCESS FOR THE PRODUCTION OF IRONES (57) Abstract A process for the preparation of irones by a chemical oxidation of iron precursors, comprising treating an iris rhizome substrate selected from the group consisting of iris rhizomes, parts of such rhizomes, iris extracts, iris extraction wastes, any plant tissues containing precursors of irones and mixtures thereof, is described.		

Process for the production of irones

Today, irone extracts are produced by solvent extraction or steam distillation of the rhizomes of any iris species, preferably *Iris pallida* or *Iris germanica*. After a cultivation period of 2-3 years the rhizomes are harvested and stored for 3-5 years at room temperature. During this maturation period the irones are released from their precursors (W. Krick, F.-J. Marner, and L. Jaenicke, Z. Naturforsch. 1983, 38c 179-184). Then the irones are recovered by steam distillation (as iris butter) or solvent extraction (leading to a resinoid) of the powdered rhizomes leading to valuable iris materials for the flavour and fragrance industry.

This long period of maturation is cumbersome, expensive, needs extensive storage capacity and leads to the loss of part of the isomers in uncontrolled degradation reactions.

Several alternatives to this non-effective and time consuming maturation period have been proposed. In a first proposal (FR-2 620 702, 24/3/89) by Buono et al., a lipophilic extract of the iris rhizomes containing the irone precursors is oxidized with KMnO_4 in an organic solvent; in this procedure the extraction of the precursors is laborious and due to this unusual downstream processing (working-up), namely the extraction of the precursors before the oxidation, the quality of the resulting product is quite different from the traditionally preferred Iris Butter or Absolute. A reduction of the maturation period to two months has been disclosed in FR-2 653 637, 3/5/91, by Baccou et al., with a preliminary treatment of the fresh, i.e. not matured iris rhizomes with ionizing radiation; but the claimed yields hardly reach the irone contents (< 400 mg / kg dry rhizomes) obtained during maturation by a 2-3 years period of storage. Bioconversions of lipophilic extracts of the rhizomes with fungi of the genus *Botryonia* (EP 0 443 925; 23/8/91) by Gil et al. or with peroxidizing enzymes (EP 0 443 926; 28/8/91) by Gil et al., have been disclosed; they lead to a good conversion of the precursors but again present the disadvantage of the preliminary solvent extraction and consequently lead to a product whose quality differs from the preferred iris butter. The bioconversion of powdered iris rhizomes by

bacteria of the family *Enterobacteriaceae* in the presence of a plant cell medium (EP 0 353 683, US 4,963,480) by Belcour et al., releases the irones from their precursors, and, after steam distillation produces an iris butter of excellent quality characterized by a high content of irones. A drawback is that the long maturation period has been replaced by a rather expensive microbiological treatment.

The present invention relates to a new process for the preparation of irones. In particular the invention relates to the treatment of iris rhizomes, parts of such rhizomes, iris extracts, plant tissues or any source of iron precursors, with nitrite salts in order to release, i.e. to form, to produce the irones.

Under irones there is to be understood a mixture of the isomers gamma-irone, cis-alpha-irone, trans-alpha-irone and beta-irone as usually found in extracts of iris species.

Under iron precursors there are to be understood compounds or any mixture of such compounds which, when treated with the nitrite, lead to irones. The use of selected clones of iris species characterized by a high content of iron precursors is preferred.

Surprisingly it has now been found that the treatment, e.g. at mild temperatures, of rhizomes, e.g. powdered fresh iris rhizomes with an aqueous solution of a nitrite salt liberates, i.e. forms the irones, under mild conditions and with excellent yields. The isomeric and enantiomeric distribution of the resulting irones is identical to that obtained after a traditional maturation.

Various cations may be associated to the nitrite anion. Good results were obtained with alkali metal, alkaline earth or ammonium nitrites, e.g. with sodium nitrite, potassium nitrite, calcium nitrite, ammonium nitrite, etc. preferably sodium or potassium nitrite.

A suitable concentration of nitrite salts is from about 0.1g/L to about 5 g/L, and, preferably from about 0.5 to about 1 g/L.

A suitable range of the ratio by weight of iris rhizome substrate to medium, i.e. the aqueous solution, is from about 1:20 to about 2:1, with about 1:5 to about 1:20 being preferred.

A suitable pH range for the treatment with aqueous nitrite solutions is from 1 to about 6, more preferably from about 2 to about 5. The preferred pH is between about 2 and about 3.

- 5 Any suitable acid, e.g. H_3PO_4 , or other mineral acids may be used to adjust the desired pH value.

The treatment is effective at mild temperatures. Best results are obtained between ca. 10 and ca. 50°C and preferably at ambient
10 temperatures, e.g. between 20 and 30°C.

The time necessary to complete the generation of the irones is dependant upon the conditions of the treatment. A suitable reaction time for the process is from about 1 hour to about 5 days, with about 5 to about 36
15 hours being preferred.

Fig. 1 shows the results of a kinetic study of the production of irones by treatment of iris rhizomes with a solution of sodium nitrate at ambient temperature.

20

In the example, the treatment of freshly harvested rhizomes of *Iris pallida* with an initial irone content of 12 mg/ kg dry rhizomes, in suspension at 100 g/L in a 0.1 % solution of sodium nitrite at pH 2.5 and at a temperature of 25°C liberates between 1200- 1400 mg irones / kg dry rhizomes
25 after 30 hours.

Rhizomes of the genera *Iris* (*Iridacea* family), parts of such rhizomes, iris wastes or extracts and more generally all plant tissues containing precursors of irones can be used. The substrate of the present invention is
30 preferably obtained from an iris plant selected from the group consisting of *I. pallida*, *I. germanica*, *I. florentina*, preferably *I. pallida*.

The iris substrate, which is preferably taken from the plant in a vegetative, i.e. growing state, may be used fresh or shortly after harvest, e.g.
35 in the form of :
-iris rhizomes, e.g. crushed or powdered,

Example 1

Production of irones by treatment of rhizomes of *Iris pallida*

5

Fresh rhizomes are first cleared from excess earth and ground in a Warring Blender.

To a solution of 5 g of sodium nitrite in 5 liters of water, ground fresh iris rhizomes (*Iris pallida*, 500 g, initial irone content: 12 mg irones / kg dry
10 rhizomes) are charged in a reaction vessel equipped with a mechanical stirrer. After the pH has been adjusted to 2.5 with 5 molar H_3PO_4 , the mixture is gently stirred at room temperature for 48 hours.

[For the determination of the irone content in the reaction mixture
15 after the "maturation", 50 ml of the reaction mixture are mixed with 20 ml of 1 molar sodium hydroxyde solution and extracted with a 80/20 v/v mixture of MTBE (methyl-t-butyl ether) and ethanol in the presence of an internal standard (anethole). After centrifugation the organic layer is analyzed by GC (Carbowax 20M column, 30 m, diameter 0.32 mm, program 60° C 4 min, 12°
20 C/min until 150° C, 3° C/min until 240° C). The matured suspension contains 1840 mg irones / kg dry rhizomes.]

The resulting matured mixture is hydrodistilled in a distillation apparatus equipped with a short path column, a condenser (phase separator) for the total distillate and a set-up for recycling of the aqueous
25 phase. The resulting Iris butter (1.85 g, yield 0.37 %/ fresh rhizomes or 0.92%/ dry rhizomes) is recovered at the surface of the receptor vessel. The Iris butter is analyzed by GC in the presence of an internal standard: ca. 50 mg of Iris butter are exactly weighed into a 50 ml flask; after addition of the internal standard and 5 ml of 1 molar sodium hydroxyde solution, the
30 heterogeneous mixture is stirred vigorously during 15 minutes; 5 ml of the resulting upper phase are mixed with 1 ml of methanol and analyzed by GC as described above. The resulting Iris butter has an irone content of 16.2 %.

Example 2

35 Production of irones by treatment of rhizomes of *Iris pallida* selected for their high irone precursor content

For these experiments, 2 types of *Iris pallida* rhizomes have been used:

Batch I: Fresh rhizomes from 56 different plants (3 years old) in a vegetative state are first cleared from excess earth. Three aliquots of each plant sample (3x5 g) are cut into small pieces and ground in a Warring Blender. Then the rhizomes are put in Erlenmeyer flasks (8g in 50ml of aqueous NaNO_2 1g/liter at pH 2.5 in each flask). The 3 flasks are placed on a gyratory shaker (100 revolutions/minute) at 25°C during 24 hours. A control experiment is performed following the same procedure without the NaNO_2 solution.

At the end of the 24 hours period, both the rhizomes and the aqueous phase are extracted with MTBE, together or separately, (i.e. by separating the rhizome parts by filtration or centrifugation). The combined extracts (extracted rhizomes and aqueous phase), are analyzed by gas chromatography (GC) as described in example 1.

Batch II: Two clones (GRN1 and GRN2) which previously had been selected for their growth and high content in irone precursors by the technique described in "C.R. Acad. Sci. Paris, Sciences de la vie/Life science, (317) 424-429 (1994)", from the 56 clones of various origins after 3 years of cultivation. The fresh rhizomes were treated as above (Batch I).

The results (expressed in mg of irones/kg of dry-weighed iris rhizome) are shown in Table 1. Standard deviations correspond to 3 replications in fields.

As currently demonstrated, the control sample (fresh harvested rhizome without any treatment) contains very few quantities of irones (less than 100mg/kg dry weight).

Table 1

	Batch I (*)	Batch II (GRN1)	Batch II (GRN2)
Treatment with NaNO_2	1339 (+208) (* mean of 56 clones of <i>Iris pallida</i> of various origins	1716 (+115)	1659 (+ 134)

Example 3

Production of irones by the treatment of an *Iris pallida* rhizome extract

A crude extract of iris rhizomes (Batch I, see example 1) is obtained of
5 50g of fresh material ground in a Warring Blendor, then follows an
extraction with 3 times 250ml CHCl_3 / MeOH (1/2) under agitation during 1
hour. After filtration, the combined fractions are concentrated. The
resulting crude extract is dissolved in 10 ml of MeOH. The solution is added
to a NaNO_2 solution (500 ml, 1 g/liter) at pH 2.5, 25°C, and stirred (100rpm)
10 during 24 hours. The total solution is extracted and analyzed as described in
Example 1. An aliquot of fresh rhizomes (before extraction) is treated as in
Example 1. The results (expressed in mg of irones/kg of dry-weighed iris
rhizome) are listed in Table 2

Table 2


15

	Fresh rhizomes (before extraction)	extract
Treatment with NaNO_2	1025 (+130)	1266 (+25)

Claims

1. A process for the preparation of irones, comprising treating iris rhizomes, parts of such rhizomes, iris extracts, iris extraction wastes, or
5 any iris plant tissue containing precursors of irones and mixtures thereof, with an aqueous solution of a nitrite, and recovering the irones formed.
2. The process according to claim 1 wherein the iris rhizome substrate is derived from an iris plant selected from the group consisting of *Iris*
10 *pallida*, *Iris germanica*, *Iris florentina*, *Iris* of Verona, and mixtures thereof.
3. The process according to claim 1 or 2, wherein the rhizomes are selected for their high content of irone precursors, e.g. are clones of *Iris*
15 *pallida*, e.g. are the clones GRN1 and GRN2.
4. The process according to claim 1, 2 or 3, wherein the reaction medium is the aqueous solution of a nitrite salt.
- 20 5. The process according to any one of claims 1 to 4 wherein an aqueous solution of an alkali, e.g. sodium, alkaline earth or ammonium nitrite is used.
6. The process according to any one of claims 1 to 5, wherein the iris
25 rhizome substrate is fresh, i.e. not matured.
7. The process according to any one of claims 1 to 6 wherein the iris rhizome substrate is an iris rhizome extract.
- 30 8. The process according to any one of claims 1 to 7 wherein the iris rhizome substrate is an iris rhizome extraction waste.

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED



9. The process according to any one of claims 1 to 8 wherein the iris rhizome substrate is a plant cell culture of iris.

10. The process according to any one of claims 1 to 9 wherein gamma-
5 irone, cis-alpha-irone, trans-alpha-irone and/or beta-irone are prepared.

11. The process according to any one of claims 1 to 10 wherein the process is carried out at about ambient temperatures, e.g. between ca. 20°C and ca. 30°C.

10

12. The process according to any one of claims 1 to 11 wherein the process is carried out for a period of about 1 to about 120 hours, preferably about 5 to about 36 hours.

15 13. The process according to any one of claims 5 to 12 wherein the concentration of the nitrite salt is ca. 0.1 to ca. 5g/liter of the reaction medium.

20 14. The process according to any one of claims 1 to 13 wherein the process is carried out in a pH range of about 1 to about 6, preferably, between about 2 to about 5.

25 15. The process according to any one of claims 1 to 14 wherein the range of iris rhizome substrate to medium is from about 1:20 to about 2:1 (weight/weight).

16. The process according to any one of claims 1 to 15 wherein the irones are isolated by extraction or distillation.

This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED



- 8 -

Claims

REPLACEMENT SHEET

REPLACEMENT SHEET

29 NOV 1999

1. A process for the preparation of irones, comprising treating a
substrate of iris rhizomes, parts of such rhizomes, iris extracts, iris extraction
5 wastes, or any iris plant tissue containing precursors of irones and mixtures
thereof, with a medium consisting of an aqueous solution of an inorganic
nitrite and recovering the irones formed, whereby the treatment is carried out
at a pH in the range of about 1 to about 6, preferably in the range of about 2
to about 5, especially between about 2 and about 3.
- 10
2. The process according to claim 1 wherein the iris rhizome substrate is
derived from an iris plant selected from the group consisting of *Iris pallida*,
Iris germanica, *Iris florentina*, Iris of Verona, and mixtures thereof.
- 15
3. The process according to claim 1 or 2, wherein the rhizomes are
selected for their high content of iron precursor, especially are clones of *Iris*
pallida, specifically are the clones GRN1 and GRN2.
4. The process according to claim 1, 2 or 3, wherein the reaction medium
20 is the aqueous solution of a nitrite salt.
5. The process according to any one of claims 1 to 4 wherein an aqueous
solution of an alkali nitrite, especially of sodium or alkaline earth nitrite, or
ammonium nitrite is used.
- 25
6. The process according to any one of claims 1 to 5, wherein the iris
rhizome substrate is fresh and not matured.
7. The process according to any one of claims 1 to 6 wherein the iris
30 rhizome substrate is an iris rhizome extract.
8. The process according to any one of claims 1 to 7 wherein the iris
rhizome substrate is an iris rhizome extraction waste.

9806269-8
29/11/99



REPLACEMENT SHEET

REPLACEMENT SHEET

29 NOV 1999

- 9 -

9. The process according to any one of claims 1 to 8 wherein the iris rhizome substrate is a plant cell culture of iris.

5 10. The process according to any one of claims 1 to 9 wherein gamma-irone, cis-alpha-irone, trans-alpha-irone and/or beta-irone are prepared.

10 11. The process according to any one of claims 1 to 10 wherein the process is carried out at about ambient temperatures, especially at a temperature of between about 20°C and about 30°C.

12. The process according to any one of claims 1 to 11 wherein the process is carried out for a period of about 1 to about 120 hours, preferably of about 5 to about 36 hours.

15

13. The process according to any one of claims 5 to 12 wherein the concentration of the nitrite salt is about 0.1 to about 5g/liter of the reaction medium.

20 14. The process according to any one of claims 1 to 13 wherein the range of iris rhizome substrate to medium is from about 1:20 to about 2:1 (weight/weight).

25 15. The process according to any one of claims 1 to 14 wherein the irones are isolated by extraction or distillation.
