

This Sheet Has Been Replaced




Ref. 20'007

THIS SHEET HAS BEEN
REPLACED

9604381-3


The invention relates to D-prolines of the formula



wherein

- 5 R is SH, benzyl or phenyl, optionally substituted by hydroxy or lower alkoxy

or



the group ;

- R¹ is hydrogen or halogen;
- 10 X is -(CH₂)_n-; -CH(R²)(CH₂)_n-; -CH₂O(CH₂)_n-; -CH₂NH-; benzyl, -C(R²)=CH-; -CH₂CH(OH)-; or thiazol-2,5-diyl;
- Y is -S-S-; -(CH₂)_n-; -O-; -NH-; -N(R²)-; -CH=CH-; -NHC(O)NH-; -N(R²)C(O)N(R²)-; -N[CH₂C₆H₅(OCH₃)₂]-; -N(CH₂C₆H₅)-; -N(CH₂C₆H₅)C(O)N(CH₂C₆H₅)-; -N(alkoxyalkyl)-; -N(cycloalkyl-methyl)-;
- 15 2,6-pyridyl; 2,5-furanyl; 2,5-thienyl; 1,2-cyclohexyl; 1,3-cyclohexyl; 1,4-cyclohexyl; 1,2-naphthyl; 1,4-naphthyl; 1,5-naphthyl; 1,6-naphthyl; biphenylen; or 1,2-phenylen, 1,3-phenylen and 1,4-phenylen, wherein the phenylen groups are optionally substituted by 1 - 4 substituents,

selected from halogen, lower alkyl, lower alkoxy, hydroxy, carboxy, -
COO-lower alkyl, nitrilo, 5-tetrazol, (2-carboxylic acid-pyrrolidin-1-yl)-
2-oxo-ethoxy, N-hydroxycarbamimidoyl, 5-oxo-[1,2,4]oxadiazolyl, 2-oxo-
[1,2,3,5]oxathiadiazolyl, 5-thioxo-[1,2,4]oxadiazolyl and 5-tert-
5 butylsulfanyl-[1,2,4]oxadiazolyl;

X' is $-(CH_2)_n-$; $-(CH_2)_nCH(R^2)-$; $-(CH_2)_nOCH_2-$; $-NHCH_2-$; benzyl,
 $-CH=C(R^3)-$; $-CH(OH)CH_2-$; or thiazol-2,5-diyl;

R² is lower alkyl, lower alkoxy or benzyl and

n is 0-3,

10 and to pharmaceutically acceptable salts and mono- and diesters thereof.

Compounds of formula I-A and I-B are novel compounds with the
exception of (R)-1-[(R)- and (R)-1-[(S)-3-mercapto-2-methyl-propionyl]-
pyrrolidine-2-carboxylic acid. These compounds are described in WO 97/10225,
having antibacterial activity against *B. fragilis*. Furthermore, they are
15 described in J. Comput.-Aided Mol. Des. (1987), 1(2) 133-42 in a theoretical
study of angiotensin-converting enzyme inhibitors.

The compounds of formulae I-A or I-B may contain 4 or 2 asymmetric
carbon atoms. Accordingly, the present invention includes all stereoisomeric
forms of the compounds of formula I-A or I-B, including each of the individual
20 enantiomers and mixtures thereof.

It has surprisingly been found that the D-prolines of formula I-A and I-B
can be used in the treatment or prevention of all forms of central and systemic
amyloidosis, which is a disorder of protein metabolism in which normally
soluble autologous proteins are deposited in the tissues as abnormal insoluble
25 fibrils, which cause structural and functional disruption. The most common
disorders associated with amyloidosis are Alzheimer's disease (AD), maturity
onset diabetes mellitus, or amyloidosis

- as a significant cause of non-ischaemic heart failure,
- as complication of long term haemodialysis in renal failure,
- 30 - as complication of monoclonal gammopathies,
- from chronic inflammatory disorders,

- from chronic infections or
- from certain types of cancer.

Furthermore, amyloidosis comprises many different diseases such as forms of hereditary amyloidosis most common familial amyloid polyneuropathy (FAP),
5 scrapie and Kreuzfeld-Jakob disease.

The common pathological feature is extracellular deposition of so called amyloid proteins in β -structured fibers and the same staining characteristics.

Serum amyloid P component (SAP) is a normal plasma protein and the precursor of amyloid component, a universal constituent of the abnormal
10 tissue deposits in amyloidosis. It is resistant to proteases and therefore plays a key role in the persistence of amyloid in vivo. For therapy pharmaceutically active compounds have to be found which would prevent the interaction of SAP with amyloid fibrils. This interaction has been demonstrated to be a protein fiber interaction, rather than an interaction with more general fiber
15 components such as glycosaminoglycans.

SAP consists as a pentamer of 5 identical non-covalently associated subunits. Two pentamers can non-covalently associate to a decamer with the two pentameric disk-like rings interacting face to face. SAP is a calcium-dependent ligand binding protein. It is produced and degraded exclusively in
20 hepatocytes and extremely stable outside the liver.

The participation of SAP in the pathogenesis of amyloidosis in vivo confirms that inhibition of binding to amyloid fibrils is an attractive therapeutic target in a range of serious human diseases.

Objects of the present invention are the aforementioned compounds of
25 formula I-A and I-B and salts and esters thereof per se and as therapeutically active substances, their manufacture and their use for therapeutic purposes and, respectively, for the production of corresponding medicaments as well as medicaments containing a compound of formula I-A and I-B or a salt thereof and the production of such medicaments for the said purpose.

The term "lower alkyl" denotes straight-chain or branched-chain saturated hydrocarbon residues, preferably with 1-4 C atoms, such as methyl, ethyl, propyl, isopropyl, n-butyl, 2-butyl, isobutyl and t-butyl.

"Halogen" denotes chlorine, iodine, fluorine and bromine.

- 5 Compounds of formula I-A and I-B can form salts with metals, e.g. alkali metal salts such as sodium or potassium salts or alkaline earth metal salts such as calcium or magnesium salts, with organic bases, e.g. salts with amines such as N-ethylpiperidine, procaine or dibenzylamine, or salts with basic amino acids such as salts with arginine or lysine. These salts can be formed
10 and isolated by methods well known in the art.

The compounds can also be used in the ester form, such esters being aliphatic or aromatic, such as, for example alkyl and phenolic esters. The most preferred esters are alkyl esters derived from C₁₋₄ alkanols, especially methyl and ethyl esters.

- 15 The compounds of formulae I-A and I-B can also be used in form of their prodrugs at either one or both carbonyl functions. Examples are esters, intramolecular esters, phosphate esters, double esters, glycolamide esters, glyceride conjugates, dihydropyridine derivatives or 8-(hydroxymethyl)-1-naphthylmethyldisulfide esters. The prodrugs may add to the value of the
20 present compounds advantages in absorption, pharmacokinetics in distribution and transport to the brain. (WO 9514705; H. Bundgaard et al., Drugs of the Future, 16, 443, 1991; A.N. Saab et al., Pharmaceutical Science, 79, 802, 1990; D.M. Lambert et al., Current Medical Chemistry 1, 376, 1995;

- Preferred are compounds of formula I-A. Especially preferred compounds
25 of formula I-A in the scope of the present invention are those in which X is CH(R²)(CH₂)_n- and wherein R² is methyl or methoxy and n is 0 or 1. The following are examples of such compounds:

(R)-1-[(S)-3-[(S)-3-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-methyl-3-oxopropyl-disulfanyl]-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid,

- (R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,7-dimethyl-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid,
(R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,7-dimethoxy-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid and
5 (R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

Especially preferred are also compounds, in which X is $-(CH_2)_n-$ and n is 0 or 1.

Such compounds are:

10

- (R)-1-[7-[(R)-2-Carboxy-pyrrolidin-1-yl]-7-oxo-heptanoyl]-pyrrolidine-2-carboxylic acid,
(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid,
15 (R)-1-[5-[(R)-2-Carboxy-pyrrolidin-1-yl]-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid,
(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]acetyl]-pyrrolidine-2-carboxylic acid,
(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-ureido]-pyrrolidine-2-
20 carboxylic acid,
(R)-1-[[Benzyl-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-amino]-acetyl]-pyrrolidine-2-carboxylic acid,
(R)-1-[cis-4-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid and
25 (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid.

Preferred are further compounds of formula I-A, wherein X is $-CH_2O-$.

Examples of such compounds are the following:

- 30 (R)-1-[[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,
(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,

5 (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid and

(R)-1-[[5-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid.

10 Compounds in which X is $-\text{CH}_2\text{NH}$ are further preferred.

An Example of such a compound is

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid.


15 Compounds, in which X is $-\text{CH}_2\text{CH}(\text{OH})-$ are further preferred.

Such a compound is, for example,


(2E,4E)-(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexa-2,4-dienoyl]-pyrrolidine-2-carboxylic acid.

20 The aforementioned compounds of formula I-A and I-B can be manufactured in accordance with the invention by

a) converting a compound of formula




into a compound of formula



25


and then into a compound of formula




wherein R^1 , X and X' have the significances given above and R^2 is lower alkyl,

or

- 5 b) treating a compound of formula



with a compound of formula




- 10 to a compound of formula I-A by cleaving off the protecting group, wherein X, Y and X' have the significances given above and R^4 is hydroxy or halogen,

or

- c) reacting a compound of formula


15



with an amine of formula



and cleaving off the protecting group from a compound of formula




20


wherein R^1 and R^3 are described as above and R^5 is hydrogen, lower alkyl, lower alkoxy, benzyl, lower alkoxyalkyl, cycloalkyl-methyl or $-\text{CH}_2\text{C}_6\text{H}_3(\text{OCH}_3)_2$,

or


- 5 d) reacting a compound of formula



with a compound of formula




- 10 and cleaving off the protecting group of a compound of formula



wherein R^1 , R^3 and X have the significances given above,


or

- e) reacting a compound of formula




15

with a compound of formula



and cleaving off the protecting group of compounds of formula




wherein R^1 , R^3 and X have the significances given above and R^7 is halogen, lower alkyl, lower alkoxy, hydroxy, carboxy, -COO-lower alkyl, nitrilo, 5-tetrazol, (2-carboxylic acid-pyrrolidin-1-yl)-2-oxo-ethoxy, N-

- 5 hydroxycarbamimidoyl, 5-oxo-[1,2,4]oxadiazolyl, 2-oxo-[1,2,3,5]oxathiadiazolyl, 5-thioxo-[1,2,4]oxadiazolyl and 5-tert-butylsulfanyl-[1,2,4]oxadiazolyl, and m is 0 - 4,

or

f) cleaving off a protecting group from a compound of formulae

10



wherein R, R^1 , X, Y and X' is as described as above and R^3 is a protecting group,

to give a compound of formula I-A or I-B, and, if desired,

15

converting a compound of general formulae I-A and I-B into a pharmaceutically usable salt or into a mono- and diester.

- In accordance with process variant a) a compound of formula I-A-1 is obtained by converting a compound of formula II, for example 1-[(S)-3-acetyl-sulfanyl-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid, into a
20 compound of formula I-B-1 and then into a compound of formula I-A-1. The reaction is conveniently effected under inert atmosphere at room temperature in the presence of ammonia in a solvent, such as methanol. After stirring for about 2 hours the compound is separated and subsequently the reaction
25 product can be worked-up to the desired pure product according to generally usual methods.

The compounds of formula I-A-1 are obtained by stirring the above compound in a solution of CuSO_4 in water at room temperature.

The precise reaction conditions are described in more detail in the working Examples.

- 5 In accordance with reaction step b) a protected D-proline is treated with a corresponding dicarboxylic acid or with a corresponding acetyl halide at 0°C . The following dicarboxylic acids are preferred:

- 2,4-dimethylglutaric acid, 2,3-dimethylsuccinic acid, cyclohexane-1,4-dicarboxylic acid, cyclohexane-1,3-dicarboxylic acid, cyclohexane-1,2-dicarboxylic acid, 1,4-phenylenediacetic acid, 1,3-phenylenediacetic acid, benzene-1,4-dioic acid, benzene-1,3-dioic acid, pyridine-2,6-dicarboxylic acid, thiophene-2,5-dicarboxylic acid, furan-2,5-dicarboxylic acid, adipic acid, 1,4-phenylenediacetic acid, 1,2-phenylenediacetic acid, (4-carboxymethyl-naphthyl-1-yl)acetic acid, (6-carboxymethyl-pyridin-2-yl)acetic acid, (5-carboxymethyl-thiophen-2-yl)acetic acid, 2,5-dimethoxy-hexanedioic acid, 2,5-dibenzyl-hex-3-enedioic acid or 2,5-diisopropyl-hex-3-enedioic acid. A detailed procedure is described in the Examples in the General Procedure A.
- 10
15

- The reaction step c) describes the treatment of an amine, for example propylamine, cyclopropylmethylamine, methoxyethylamine, benzylamine or veratrylamine with a compound of formula IX. This reaction is carried out at a temperature between 20 and 80°C in a solvent, such as acetonitrile.
- 20

- In accordance with variant d) a compound of formula I-B is prepared. To a compound of formula XV in dichloromethane at 0°C is added a corresponding bromoacetyl derivative, such as bromoacetyl bromide, and a compound of formula V. The deprotection is then carried out by methods known in the art.
- 25

- Compounds, in which Y is an optionally substituted 1,2-, 1,3- or 1,4-phenylene group, can be prepared in accordance with reaction variant e). To a compound of formula XV a corresponding dihydroxy-derivative of formula XVII is added. The reaction is carried out in dimethylformamide at room temperature. Preferred are the following dihydroxy-derivatives: hydroquinone,
- 30

tetrafluorohydroquinone, chorohydroquinone, methoxyhydroquinone, resorcinol, 2,6-dihydroxytoluene, 5-methoxyresorcinol, 3,5-dihydroxybenzoate, 3,5-dihydroxybenzonitrile, phloroglucinol, pyrogallol-1-methyl ether, 3-methylcatechol, tetrachlorocatechol, 2,6-dihydroxynaphthalene, 1,5-
5 dihydroxynaphthalene, 2,3-dihydroxynaphthalene, 2,2'-dihydroxybiphenyl, 1,4-naphthoquinone or 2,7-dihydroxynaphthalene,


In accordance with process variant f) a compound of formulae III or IV is deprotected to a compound of general formula I-A or I-B. Suitable protecting groups and methods for their cleavage will be familiar to any person skilled in
10 the art, although of course there can be used only those protecting groups which can be cleaved off by methods under the conditions of which other structural elements are not affected. The tert-butyl group and the benzyl group are preferred O-protecting groups. The process is carried out in conventional manner. For example, a compound of formula III is dissolved in a
15 suitable solvent or mixture of solvents such as ethanol and ethylacetate, and hydrogenated in the presence of Pd on carbon at room temperature and atmospheric pressure.

Pharmaceutically acceptable salts and esters can be manufactured according to methods which are known per se and familiar to any person
20 skilled in the art.

In schemes 1-9 are described processes for preparation of compounds of formulae I-A and I-B, starting from known compounds or from compounds, which can be prepared in conventional manner.


The starting materials of formulae V, VI, VIII, IX, X, XII, XIV, XVII, XX
25 and XXIV are commercial products or can be prepared according to methods known per se.

The preparation of compounds of formulae I-A and I-B are described in more detail in working Examples 1-104.



wherein R^1 , X and X' have the significances given above, R^2 is lower alkyl and R^3 is a protecting group.

Scheme 2




5

wherein X, Y and X' have the significances given above and R⁴ is hydroxy or halogen.


10

Scheme 3



wherein R^1 and R^3 are described as above and R^4 is hydrogen, lower alkyl, lower alkoxy, benzyl, lower alkoxyalkyl, cycloalkyl-methyl or - $CH_2C_6H_3(OCH_3)_2$.


Scheme 4



5

wherein R^1 , R^3 , X and X' have the significances given above and R^6 is hydrogen, lower alkyl, lower alkoxy, or benzyl.


Scheme 5



10


w wherein R^1 , R^3 and X have the significances given above.

Scheme 6




wherein R^1 , R^3 and X have the significances given above and R^7 is halogen, lower alkyl, lower alkoxy, hydroxy, carboxy, -COO-lower alkyl, nitrilo, 5-tetrazol, (2-carboxylic acid-pyrrolidin-1-yl)-2-oxo-ethoxy, N-hydroxycarbamimidoyl, 5-oxo-[1,2,4]oxadiazolyl, 2-oxo-[1,2,3,5]oxathiadiazolyl, 5-thioxo-[1,2,4]oxadiazolyl and 5-tert-butylsulfanyl-[1,2,4]oxadiazolyl, and m is 0 - 4.

Scheme 7




wherein R^1 , R^3 , R^5 , X and X' have the significances given above.

Scheme 8




wherein R^1 and R^2 have the significances given above.

Scheme 9

























wherein R^3 has the significance given above.






The preparation of the following examples is described in more detail:















X	Y	X'	R	R ¹	Expl.
$\begin{array}{c} \vdots \\ \\ -\text{CH}-\text{CH}_2- \end{array}$	-S-S-	$\begin{array}{c} \vdots \\ \\ -\text{CH}_2-\text{CH}- \end{array}$	-	H	1d
$\begin{array}{c} \\ -\text{CH}-\text{CH}_2- \end{array}$	-	-	SH	H	1c
$\begin{array}{c} \\ -\text{CH}-\text{CH}_2- \end{array}$	-S-S-	$\begin{array}{c} \\ -\text{CH}_2-\text{CH}- \end{array}$	-	H	2c
$\begin{array}{c} \vdots \\ \\ -\text{CH}-\text{CH}_2- \end{array}$	-	-	SH	H	2b
-(CH ₂) ₂ -	-S-S-	-(CH ₂) ₂ -	-	H	3
-(CH ₂) ₃ -	-CH ₂ -	-(CH ₂) ₃ -	-	H	4b
-CH(CH ₃)CH ₂ -	-(CH ₂) ₂ -	-CH ₂ CH(CH ₃)-	-	H	5
CH(OCH ₃)CH ₂ -	-(CH ₂) ₂ -	-CH ₂ CH(OCH ₃)-	-	H	6b
-(CH ₂) ₂ -	-CH ₂ -	-(CH ₂) ₂ -	-	H	7
-CH ₂ -	-(CH ₂) ₂ -	-CH ₂ -	-	H	8b (R),(R)
-CH ₂ -	-CH ₂ -	-CH ₂ -	-	H	9b
-CH ₂ -	a bond	-CH ₂ -	-	H	10b
-CH ₂ O(CH ₂) ₂ -	-O-	-(CH ₂) ₂ OCH ₂ -	-	H	11
-(CH ₂) ₂ -		-(CH ₂) ₂ -	-	H	12
-CH ₂ -		-CH ₂ -	-	H	13 (R),(R)
-CH ₂ O-		-OCH ₃ -	-	H	14b (R),(R)
-(CH ₂) ₂ -		-(CH ₂) ₂ -	-	H	15c
-(CH ₂) ₂ -	-N[(CH ₂) ₂ CH ₃]-	-(CH ₂) ₂ -	-	H	16c
-CH ₂ -	-NHC(O)NH-	-CH ₂ -	-	H	17c
-(CH ₂) ₃ -	-(CH ₂) ₂ -	-(CH ₂) ₃ -	-	H	18

$-(\text{CH}_2)_2-$	$-(\text{CH}_2)_2-$	$-(\text{CH}_2)_2-$	-	H	19
	a bond		-	H	20b
$-\text{CH}_2-$	-	-		H	21d
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	22b
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	23b
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	24b
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	25b
$-\text{CH}_2\text{O}-$	-	-		H	26b
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	27b
$-\text{CH}_2\text{O}-$	-	-		H	28b
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	29b
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	30b
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	31b
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	32
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	33b
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H	34

-CH ₂ O-		-OCH ₂ -	-	H	35b
-CH ₂ O-		-OCH ₂ -	-	H	36b
-CH ₂ O-		-OCH ₂ -	-	H	37b
-CH ₂ O-		-OCH ₂ -	-	H	38b
-CH ₂ O-		-OCH ₂ -	-	H	39b
-CH ₂ O-		-OCH ₂ -	-	H	40b
-CH ₂ O-		-OCH ₂ -	-	H	41b
-CH ₂ O-		-OCH ₂ -	-	H	42b
-CH ₂ O-		-OCH ₂ -	-	H	43b
-CH ₂ O-		-OCH ₂ -	-	H	44b
-CH ₂ O-		-OCH ₂ -	-	H	45b
-CH ₂ O-		-OCH ₂ -	-	H	46b

-CH ₂ O-		-OCH ₂ -	-	H	47b
-CH ₂ O-		-OCH ₂ -	-	H	48b
-CH ₂ O-		-OCH ₂ -	-	H	49b
-CH ₂ NH-		-NHCH ₂ -	-	H	50b
-CH ₂ NH-		-NHCH ₂ -	-	H	51b
-CH ₂ -	-N[(CH ₂) ₃ CH ₃]-	-CH ₂ -	-	H	52b
-CH ₂ -	-N[(CH ₂) ₂ OCH ₃]-	-CH ₂ -	-	H	53b
-CH ₂ -	-N(CH ₂ C ₆ H ₅)-	-CH ₂ -	-	H	54b
-CH ₂ -	-N[(CH ₂) ₃ CH ₃]- CO-N[(CH ₂) ₃ CH ₃]-	-CH ₂ -	-	H	55c
-CH ₂ -	-N[CH ₂ C ₆ H ₅]- CO-N[CH ₂ C ₆ H ₅]-	-CH ₂ -	-	H	56c
-CH ₂ NH-	-	-	benzyl	H	57
-CH(CH ₃)-	-CH ₂ -	-CH(CH ₃)-	-	H	58b
-CH(CH ₃)-	a bond	-CH(CH ₃)-	-	H	59b
a bond		a bond	-	H	60b
a bond		a bond	-	H	61b
a bond		a bond	-	H	62b
a bond		a bond	-	H	63b
-CH ₂ -		-CH ₂ -	-	H	64b

-CH ₂ -		-CH ₂ -	-	H	65b
a bond		a bond	-	H	66b
a bond		a bond	-	H	67b
a bond		a bond	-	H	68b
a bond		a bond	-	H	69b
a bond		a bond	-	H	70b
-CH ₂ -	-CH ₂ -	-CH ₂ -	-	H	71b (S),(S)
-CH ₂ -		-CH ₂ -	-	H	72b (S),(S)
-CH ₂ O-		-OCH ₂ -	-	H	73b (S),(S)
-CH ₂ -		-CH ₂ -	-	H	74d
-CH ₂ -		-CH ₂ -	-	H	75d
-CH ₂ -		-CH ₂ -	-	H	76e
	-(CH ₂) ₂ -		-	H	77d
	-(CH ₂) ₂ -		-	H	78f
	-(CH ₂) ₂ -		-	H	79 2 dia- stereomers
CH(CH ₂ C ₆ H ₅)	-(CH ₂) ₂ -	-CH(CH ₂ C ₆ H ₅)-	-	H	80d
-CH[(CH ₂) ₄]-	-(CH ₂) ₂ -	-CH[(CH ₂) ₄]-	-	H	81d
-CH[(CH ₂) ₄]-	-(CH ₂) ₂ -	-CH[(CH ₂) ₄]-	-	H	82 2 dia- stereomers
-CH(i-prop.)-	-(CH ₂) ₂ -	-CH(i-prop.)-	-	H	83d

$-\text{CH}[(\text{CH}_2)_2\text{OCH}_3]-$	$-(\text{CH}_2)_2-$	$-\text{CH}[(\text{CH}_2)_2\text{OCH}_3]-$	-	H	84d
$-\text{CH}(\text{CH}_3)-$		$-\text{CH}(\text{CH}_3)-$	-	H	85e 3 dia- stereomers
$-\text{C}(\text{CH}_3)=\text{CH}-$	a bond	$-\text{CH}=\text{C}(\text{CH}_3)-$	-	H	86b
$-\text{CH}(\text{CH}_3)-$	$-(\text{CH}_2)_2-$	$-\text{CH}(\text{CH}_3)-$	-	H	87
$-\text{CH}_2\text{CH}(\text{OH})-$	a bond	$-\text{CH}(\text{OH})\text{CH}_2-$	-	H	88c
$-\text{CH}_2-$	$-\text{CH}=\text{CH}-$	$-\text{CH}_2-$	-	H	89b
$-(\text{CH}_2)_2-$	$-\text{N}[(\text{CH}_2)_2\text{CH}_3]-$	$-(\text{CH}_2)_2-$	-	H	90c
$-(\text{CH}_2)_2-$	$-\text{N}(\text{CH}_2\text{cyclopropyl})-$	$-(\text{CH}_2)_2-$	-	H	91b
$-(\text{CH}_2)_2-$	$-\text{N}[\text{CH}_2\text{C}_6\text{H}_3(\text{OCH}_3)_2]-$	$-(\text{CH}_2)_2-$	-	H	92c
$-(\text{CH}_2)_2-$	$-\text{N}[(\text{CH}_2)_2\text{OCH}_3]-$	$-(\text{CH}_2)_2-$	-	H	93b
$-(\text{CH}_2)_2-$	$-\text{N}(\text{CH}_2\text{C}_6\text{H}_5)-$	$-(\text{CH}_2)_2-$	-	H	94b
$-(\text{CH}_2)_2-$	$-\text{NH}-$	$-(\text{CH}_2)_2-$	-	H	95c
$-(\text{CH}_2)_2-$	$-\text{N}(\text{CH}_2\text{CH}_3)-$	$-(\text{CH}_2)_2-$	-	H	96
$-(\text{CH}_2)_2-$	$-\text{N}[\text{CH}_2\text{C}_6\text{H}_4\text{CF}_3]-$	$-(\text{CH}_2)_2-$	-	H	97
$-\text{CH}_2-$	$-(\text{CH}_2)_2-$	$-\text{CH}_2-$	-	H	98c (R),(S)
$-\text{CH}_2-$	$-(\text{CH}_2)_2-$	$-\text{CH}_2-$	-	H	99c
$-\text{CH}_2-$	$-(\text{CH}_2)_2-$	$-\text{CH}_2-$	-	F	100e
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	F	101b
$-\text{CH}_2-$		$-\text{CH}_2-$	-	F	102b
$-\text{CH}_2-$	$-(\text{CH}_2)_2-$	$-\text{CH}_2-$	-	H, = bond	103f
$-\text{CH}_2\text{O}-$		$-\text{OCH}_2-$	-	H, = bond	104b

As mentioned earlier, the compounds of general formula I-A and I-B in accordance with the invention have valuable pharmacological properties. They can be used against all forms of central and systemic amyloidosis, which is a disorder of protein metabolism in which normally soluble autologous proteins are deposited in the tissues as abnormal insoluble fibrils, which cause structure and functional disruption.

Compounds of formula I-A and I-B have been tested by the following method:

10

Test method

Binding of SAP (serum amyloid P) to human amyloid A β (1-42) fibrils

Nunc Flouro Polysorp 96 well plates were coated with 0.5 μ g/well of A β 1-42, which had been aged for 7 days at 37 °C. Plates were dried for 3 days at 37 °C, washed with 2 x 150 μ l of TC (10 mM tris, 138 mM NaCl, 6 mM CaCl₂, 0.05% NaN₃ pH 8.0) with 1% bovine serum albumin. Then 50 μ l TC containing 8% bovine serum albumin, 25 μ l compound in TC and 25 μ l 40 nM [125I]serum amyloid protein in TE (10 mM EGTA instead of Ca) were added per well. Incubation was performed over night at room temperature and wells were washed twice with 180 μ l of TC containing 1% bovine serum albumin. To determine radioactivity 100 μ l Microscint 40 were added per well and radioactivity was measured in a TopCount (Packard). The IC₅₀ (μ M) of preferred compounds of formula I-A and I-B are in the range of about 0.2 - 2.0.

The compounds of formula I-A and I-B and their pharmaceutically acceptable acid addition salts, their mono- and diesters and cyclic imides thereof can be used as medicaments, e.g. in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. The administration can, however, also be effected rectally, e.g. in the form of suppositories, parenterally, e.g. in the form of injection solutions, or nasally.

For the manufacture of pharmaceutical preparations the compounds of formulae I-A and I-B and the pharmaceutically acceptable acid addition salts and esters thereof can be processed with pharmaceutically inert, inorganic or organic carriers. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts and the like can be used, for example, as such carriers for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carriers for soft gelatine capsules are, for example, vegetable oil, waxes, fats, semi-solid and liquid polyols and the like. Depending on the nature of the active substance no carriers are, however, usually required in the case of soft gelatine capsules.

5 Suitable carriers for the manufacture of solutions and syrups are, for example, water, polyols, glycerol, vegetable oils and the like. Suitable carriers for suppositories are, for example, natural or hardened oils, waxes, fats, semi-liquid or liquid polyols and the like.

10 Suitable carriers for the manufacture of solutions and syrups are, for example, water, polyols, glycerol, vegetable oils and the like. Suitable carriers for suppositories are, for example, natural or hardened oils, waxes, fats, semi-liquid or liquid polyols and the like.

The pharmaceutical preparations can, moreover, contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, coating agents or antioxidants. They can also contain still other therapeutically valuable substances.

15 The pharmaceutical preparations can, moreover, contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, coating agents or antioxidants. They can also contain still other therapeutically valuable substances.

Medicaments containing a compound of formulae I-A or I-B or a pharmaceutically acceptable acid addition salt or mono- and diesters thereof and a therapeutically inert carrier are also an object of the present invention, as is a process for their manufacture which comprises bringing one or more compounds of formula I-A and I-B and/or pharmaceutically acceptable acid addition salts and mono- and diesters thereof into a galenic administration form together with one or more therapeutically inert carriers.

20 Medicaments containing a compound of formulae I-A or I-B or a pharmaceutically acceptable acid addition salt or mono- and diesters thereof and a therapeutically inert carrier are also an object of the present invention, as is a process for their manufacture which comprises bringing one or more compounds of formula I-A and I-B and/or pharmaceutically acceptable acid addition salts and mono- and diesters thereof into a galenic administration form together with one or more therapeutically inert carriers.

25 Medicaments containing a compound of formulae I-A or I-B or a pharmaceutically acceptable acid addition salt or mono- and diesters thereof and a therapeutically inert carrier are also an object of the present invention, as is a process for their manufacture which comprises bringing one or more compounds of formula I-A and I-B and/or pharmaceutically acceptable acid addition salts and mono- and diesters thereof into a galenic administration form together with one or more therapeutically inert carriers.

In accordance with the invention compounds of general formula I-A and I-B as well as their pharmaceutically acceptable acid addition salts and mono- and diesters thereof can be used in the treatment or prevention of central and systemic amyloidosis. The most common disorders associated with amyloidosis are Alzheimer's disease (AD), maturity onset diabetes mellitus, or amyloidosis.

30 In accordance with the invention compounds of general formula I-A and I-B as well as their pharmaceutically acceptable acid addition salts and mono- and diesters thereof can be used in the treatment or prevention of central and systemic amyloidosis. The most common disorders associated with amyloidosis are Alzheimer's disease (AD), maturity onset diabetes mellitus, or amyloidosis.

- as a significant cause of non-ischaemic heart failure,
- as complication of long term haemodialysis in renal failure,
- as complication of monoclonal gammopathies,
- from chronic inflammatory disorders,
- 5 - from chronic infections and
- from certain types of cancer.

Furthermore, amyloidosis comprises many different diseases such as forms of hereditary amyloidosis most common familial amyloid polyneuropathy (FAP), scrapie and Kreuzfeld-Jakob disease.

10

Furthermore, the present compounds can be used for the manufacture of corresponding medicaments. The dosage can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In the case of oral administration the dosage lies in a range of about 0.1 mg per
15 dosage to about 5000 mg per day of a compound of general formulae I-A or I-B or the corresponding amount of a pharmaceutically acceptable acid addition salt or mono- and diesters thereof, although the upper limit can also be exceeded when this is shown to be indicated.

The following Examples illustrate the present invention in more detail.
20 However, they are not intended limit its scope in any manner. All temperatures are given in degrees Celsius.

Example 1

(R)-1-[(S)-3-[(S)-3-[(R)-2-carboxy-pyrrolidin-1-yl]-2-methyl-3-oxopropyl-
disulfanyll-2-methyl-propionyll-pyrrolidine-2-carboxylic acid

- 25 a) 1-[(S)-3-(Acetylsulfanyl)-20-methyl-propionyll-(R)-pyrrolidine-2-carboxylic
acid tert-butylester and
1-[(R)-3-(Acetylsulfanyl)-20-methyl-propionyll-(R)-pyrrolidine-2-carboxylic acid
tert-butylester

18.6 ml Triethylamine were given at 0-5 °C to a solution of 23.2 g (135 mmol)
30 D-proline-tert-butylester in 230 ml dry dichloromethane. A solution of 24.5 g (135 mmol) S-(3-chloro-2-methyl-3-oxopropyl)ethanethioic acid ester in 116 ml

dichloromethane were added at this temperature over a period of 1 hour and stirring was continued at room temperature for 2 hours. The precipitate was removed by filtration. The solution was washed with water and dried with sodium sulfate. Evaporation of the solvent at reduced pressure gave 41.4 g

5 colourless oil which was chromatographed on 4 kg silicagel with ether/cyclohexane 2/1 yielding 19.6 g (46%) 1-[(R)-3-(acetylsulfanyl)-20-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid tert-butylester and 18.2 g (43%) 1-[(S)-3-(Acetylsulfanyl)-20-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid tert-butylester and 1.6 g mixture of epimers.

10 MS m/e(%) = 315 (M^+ , 3), 259(10), 242(10), 214(100), 172(10), 145(32), 70(22);
[α]_D = -0.7 ° (1% EtOH).

MS m/e(%) = 315 (M^+ , 4), 259(7), 242(9), 214(100), 172(9), 145(33), 70(33);
[α]_D = +156.7 ° (1% EtOH).

b) 1-[(S)3-Acetylsulfanyl-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid

15 15.45 g (48.9 mmol) 1-[(S)-3-(Acetylsulfanyl)-20-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid tert-butylester were stirred with 99 ml trifluoric acid and 55 ml anisole under argon for three hours. The mixture was evaporated under vacuum. The residue was dissolved in about 100 ml icecold ethylacetate and washed with about 200 ml of an icecold aqueous solution of
20 sodiumbicarbonate. Concentrated hydrochloric acid was added under icecooling until ph 1-2. The aqueous phase was extracted four times with icecold ethylacetate, dried with sodium sulfate and evaporated. The yield was 11.6 g (91%) 1-[(S)3-acetylsulfanyl-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid that was used without further purification.

25 [α]_D = -11.8 ° (0.6% EtOH).

c) 1-[(S)3-Mercapto-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid

11.59 g (44.69 mmol) 1-[(S)3-acetylsulfanyl-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid were dissolved at room temperature under argon in 70 ml argon washed methanol. After addition of 70 ml 10N ammonia in
30 methanol stirring was continued for two hours at room temperature. Then the solvent was distilled off under vacuum. The residue was taken up with 5% aqueous KHSO₄ solution and extracted six times with dichloromethane.

The organic layers were washed twice with 5% aqueous KHSO₄ solution, three times with 1 N hydrochloric acid and dried over sodiumsulfate.

Evaporation of the solvent and crystallization from ethylacetate/hexane yielded 6.25 (64%) 1-[(S)3-mercapto-2-methyl-propionyl]-(R)-pyrrolidine-2-

5 carboxylic acid with melting point 99-101 °C.

$\alpha]_D = +40.7^\circ$ (1% EtOH).

d) (R)-1-[(S)-3-[(S)-3-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-methyl-3-oxopropyl]disulfanyl]-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid

A solution of 749 mg (3.0 mmol) CuSO₄ x 5 H₂O in 90 ml water was added at
10 room temperature to a solution of 651.85 mg (3.0 mmol) 1-[(S)3-mercapto-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid in 90 ml dichloromethane. The mixture was vigorously stirred for 10 minutes and filtered. The aqueous phase was washed 5 times with dichloromethane, the organic phases were washed with brine and dried with magnesiumsulfate and the solvent was
15 removed under vakuum. Crystallization from dichloromethane/hexane gave 275.3 mg (43%) (R)-1-[(S)-3-[(S)-3-[(R)-2-carboxy-pyrrolidin-1-yl]-2-methyl-3-oxopropyl]disulfanyl]-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid with melting point 142-144 °C.

$[\alpha]_D = +42.8^\circ$ (1% CHCl₃).

20

Example 2

(R)-1-[(R)-3-[(R)-3-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-methyl-3-oxopropyl]disulfanyl]-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid

a) 1-[(R)3-Acetylsulfanyl-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid

25 18.9 g (60.0 mmol) 1-(3-acetylsulfanyl-2-methyl-propionyl)-pyrrolidine-2-carboxylic acid tert-butyl ester were stirred with 120 ml trifluoric acid and 75 ml anisole under argon for three hours. The mixture was evaporated under vakuum. The residue was dissolved in icecold ethylacetate and washed with an icecold aqueous solution of sodiumbicarbonate. Concentrated hydrochloric
30 acid was added unter icecooling until ph 2-3. The aqueous phase was extracted three times with icecold ethylacetate, dried with sodium sulfate and evaporated. The yield was 15.3 g (98%) 1-[(R)3-acetylsulfanyl-2-methyl-

propionyl]-(R)-pyrrolidine-2-carboxylic acid that was used without further purification.

$[\alpha]_D^{+127.8}$ ° (1% EtOH).

b) 1-[(R)3-Mercapto-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid

- 5 2.98 g (11.5 mmol) 1-[(R)3-Acetylsulfanyl-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic were dissolved at room temperature under argon in 15 ml argon washed methanol. After addition of 15 ml 10N ammonia in methanol stirring was continued for two hours at room temperature. Then the solvent was distilled off under vacuum at room temperature. The residue was taken up
- 10 with 5% aqueous KHSO_4 solution and extracted six times with with dichloromethane and three times with ethylacetate. The organic layers were washed twice with 5% aqueous KHSO_4 solution, three times with 1 N hydrochloric acid and dried over sodiumsulfate. Evaporation of the solvent and crystallization from ethylacetate/hexane yielded 1.59 g (64%) 1-[(R)3-
- 15 mercapto-2-methyl-propionyl]-(R)-pyrrolidine-2-carboxylic acid with melting point 98-100 °C.

$[\alpha]_D^{+128.8}$ ° (1% EtOH).

c) (R)-1-[(R)-3-[(R)-3-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-methyl-3-oxopropyl]disulfanyl)-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid

- 20 Analogous to example 1d):

MS m/s (%): 432(M^+ ,2) 217(100), 184(76), 172(67), 142(13), 70(79), 41(21).

Example 3

- (R)-1-[3-[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]disulfanyl]-propionyl]-pyrrolidine-2-carboxylic acid
- 25

- 0.9 g (4 mmol) 1-(3-mercapto-propionyl)-(R)-pyrrolidine-2-carboxylic acid (raw material) were dissolved in dichloromethane and extracted with 50 ml of an saturated aqueous solution of CuSO_4 . The aqueous phase was extracted twice with dichloromethane, the combined organic phases filtered, dried with
- 30 magnesiumsulfate and evaporated. Chromatography with dichloromethane/acetone/ formic acid 80/20/1 gave 70 mg (R)-1-[3-[3-[(R)-2-carboxy-pyrrolidin-1-yl]-3-oxo-propyl]disulfanyl]-propionyl]-pyrrolidine-2-carboxylic acid as a

colourless oil.

ISP-MS: 403 (M-H)⁺.

Example 4

5 (R)-1-[9-[(R)-2-Carboxy-pyrrolidin-1-yl]-9-oxo-nonanoyl]pyrrolidine-2-carboxylic acid

a) (R)-1-[9-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-9-oxo-nonanoyl]pyrrolidine-2-carboxylic acid benzyl ester

0.97g (4 mmol) D-proline-benzylester hydrochloride in 25 ml dichloromethane
10 were stirred with 450 mg (2 mmol) azelaoyl chloride and 1.12 ml (8 mmol) triethylamine for 20 hours under argon at room temperature. Extraction with 2N hydrochloric acid and brine, drying with sodium sulfate and evaporation gave 1.2 g oil which was chromatographed over silicagel with acetoacetate to yield 0.9 g (80%) (R)-1-[9-[(R)-2-benzoyloxycarbonyl-pyrrolidin-1-yl]-9-oxo-
15 nonanoyl]pyrrolidine-2-carboxylic acid benzyl ester as colourless oil.

¹H-NMR (CDCl₃, ppm): 1.1-2.4 (m, 22H), 3.4-3.7 (m, 4H), 4.4-4.6 (m, 2H), 5.1-5.3 (2xAB, 4H), 7.34 (m, 10H).

b) (R)-1-[9-[(R)-2-Carboxy-pyrrolidin-1-yl]-9-oxo-nonanoyl]pyrrolidine-2-carboxylic acid.

20 100 mg (0.18 mmol) (R)-1-[9-[(R)-2-benzoyloxycarbonyl-pyrrolidin-1-yl]-9-oxo-nonanoyl]pyrrolidine-2-carboxylic acid benzyl ester in 20 ml ethanol were hydrogenated in the presence of 20 mg 5% Pd on carbon for two hours at room temperature. Filtration and evaporation gave 60 mg (R)-1-[9-[(R)-2-carboxy-pyrrolidin-1-yl]-9-oxo-nonanoyl]pyrrolidine-2-carboxylic acid, as a colourless oil.

25 ISP-MS: 383 (MH⁺).

Example 5

(R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,7-dimethyl-8-oxo-octanoyl]pyrrolidine-2-carboxylic acid

30 1.2 g (5.0 mmol) 2,7-dimethyl-octanedioyl chloride were dissolved in 100 ml dimethylformamide, 1.15 g (10 mmol) D-proline and 1.4 ml (10 mmol) triethylamine were added and the mixture warmed to 50 °C for five minutes.

Stirring was continued at room temperature over night. The solvent was distilled off and the residue taken up in 30 ml 2N hydrochloric acid. Extraction with ethylacetate, drying with sodiumsulfate, evaporation and chromatography over silicagel with chloroform/acetone/formic acid 80/15/5 gave 0.11 g (R)-1-[8-[(R)-2-carboxy-pyrrolidin-1-yl]-2,7-dimethyl-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid as colourless oil.

ISP-MS: 397 (MH)⁺.

Example 6

10 (R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,7-dimethoxy-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[8-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-2,7-dimethoxy-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid benzyl ester

To 0.25 g (1.1 mmol) 2,7-dimethoxy-octanedioic acid in a mixture of 25 ml tetrahydrofuran and 20 ml dichloromethane was added a solution of 0.35 g (2.1 mmol) carbonyldiimidazole in 15 ml tetrahydrofuran. After stirring at room temperature for two hours 0.52 g (2.16 mmol) D-proline-benzylester hydrochloride in 10 ml dichloromethane and 0.54 g triethylamine were added and stirring was continued for 18 hours.

20 After filtration the solvent was distilled off and the residue dissolved in ethylacetate and extracted with 2N hydrochloric acid and water. Drying with sodiumsulfate, evaporation of the solvent and chromatography over silicagel with ethylacetate yielded 0.21g (R)-1-[8-[(R)-2-benzoyloxycarbonyl-pyrrolidin-1-yl]-2,7-dimethoxy-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid benzyl ester as a colourless oil.

S-ISP: 609 (M+H)⁺.

b) (R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,7-dimethoxy-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid

182 mg (0.3 mmol) (R)-1-[8-[(R)-2-benzoyloxycarbonyl-pyrrolidin-1-yl]-2,7-dimethoxy-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid benzyl ester in 10 ml methanol were hydrogenated in the presence of 30 mg 5% Pd on carbon. Filtration and evaporation of the solvent yielded 109 mg (84%) (R)-1-[8-[(R)-2-carboxy-pyrrolidin-1-yl]-2,7-dimethoxy-8-oxo-octanoyl]-pyrrolidine-2-carboxylic

acid as colourless oil.

MS: 427 (M-H)⁺.

Example 7

5 (R)-1-[7-[(R)-2-Carboxy-pyrrolidin-1-yl]-7-oxo-heptanoyl]-pyrrolidine-2-carboxylic acid

- A mixture of 0.99 g (5 mmol) pimeloyl chloride, 1.15 g (10 mmol) D-proline and 1.4 ml (10 mmol) triethylamine in 100 ml dimethylformamide was warmed until a clear solution was obtained and then stirred over night at ambient temperature. The solvent was distilled off under vakuum. The residue was taken up with 2N hydrochloric acid and extracted with dichloromethane. Evaporation of the solvent and chromatography over silicagel with chloroform/acetone/formic acid 80/15/5 yielded 0.27 g (R)-1-[7-[(R)-2-carboxy-pyrrolidin-1-yl]-7-oxo-heptanoyl]-pyrrolidine-2-carboxylic acid as an oil.
- 15 MS: 353 (M-H)⁺.

Example 8

(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

20 a) (R)-1-[6-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl ester

- 0.97 g (10 mmol) D-proline-benzylester hydrochloride in 70 ml dichloromethane were stirred with 0.92 g (5 mmol) adipoyl chloride and 2.8 ml (20 mmol) triethylamine over the weekend under argon at room temperature.
- 25 Extraction with 2N hydrochloric acid and water, drying with sodiumsulfate, evaporation and chromatography over silicagel with acetoacetate to yielded 0.42 g (16%) (R)-1-[6-[(R)-2-benzoyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl ester as colourless oil.
- MS-ISP: 521 (M+H)⁺.

30 b) (R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

410 mg (0.79 mmol) (R)-1-[6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl ester in 100 ml methanol were hydrogenated in the presence of 50 mg 5% Pd on carbon. Filtration and evaporation of the solvent yielded 160 mg (59%) (R)-1-[6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid as colourless oil.
MS: 339 (M-H)⁺.

Example 9

(R)-1-[5-[(R)-2-Carboxy-pyrrolidin-1-yl]-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[5-[(R)-2-Benzyloxycarbonyl-pyrrolidin-1-yl]-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid benzyl ester

0.97 g (10 mmol) D-proline-benzylester hydrochloride in 70 ml dichloromethane were stirred with 0.85 g (5 mmol) glutaryl dichloride and 2.8 ml (20 mmol) triethylamine over night under argon at room temperature. Extraction with 2N hydrochloric acid and brine, drying with sodiumsulfate, evaporation and chromatography over silicagel with acetoacetate to yielded 0.44 g (17%) (R)-1-[5-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid benzyl ester as colourless oil.
MS-ISP: 507 (M+H)⁺.

b) (R)-1-[5-[(R)-2-Carboxy-pyrrolidin-1-yl]-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid

440 mg (0.87 mmol) (R)-1-[5-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid benzyl ester in 100 ml ethanol were hydrogenated in the presence of 40 mg 5% Pd on carbon. Filtration and evaporation of the solvent yielded 130 mg (46%) (R)-1-[5-[(R)-2-carboxy-pyrrolidin-1-yl]-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid as colourless oil.
MS-ISP: 327 (M+H)⁺.

Example 10

(R)-1-[4-[(R)-2-Carboxy-pyrrolidin-1-yl]-4-oxo-butryl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[4-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-4-oxo-butyryl]-pyrrolidine-2-carboxylic acid benzyl ester

To a solution of 300 mg (1.2 mmol) D-Proline benzyl ester hydrochloride and 0.35 ml (2.5 mmol) triethylamine in 9 ml dichloromethane at 0 °C was added
5 dropwise 68 ml (0.6 mmol) succinyl chloride and stirring continued for 24 h at room temperature. The reaction mixture was then washed sequentially with saturated ammonium chloride solution, saturated sodium bicarbonate solution and finally with water, and the aqueous phases back-extracted with dichloromethane. The combined organic extracts were dried over sodium
10 sulphate and concentrated in vacuo to afford 286 mg (94%) of the title compound as a pale yellow oil.

MS m/e (%): 510 (M + NH₄⁺, 20), 493 (M + H⁺, 100), 288 (80).

b) (R)-1-[4-[(R)-2-Carboxy-pyrrolidin-1-yl]-4-oxo-butyryl]-pyrrolidine-2-carboxylic acid

15 A solution of 256 mg (0.5 mmol) (R)-1-[4-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-4-oxo-butyryl]-pyrrolidine-2-carboxylic acid benzyl ester in 5 ml ethanol was stirred with 13 mg 10% Palladium on carbon under 1 atm of hydrogen for 16 h at room temperature. After filtration to remove the catalyst, concentration in vacuo afforded 170 mg (100%) of the title compound (R)-1-[4-
20 [(R)-2-carboxy-pyrrolidin-1-yl]-4-oxo-butyryl]-pyrrolidine-2-carboxylic acid as a colourless viscous oil.

MS m/e (%): 313 (M+H)⁺, 100).

Example 11

25 (R)-1-[[2-[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]ethoxy]-ethoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A mixture of 1.04 g (4 mmol) 2,2'-[oxybis(2,1-ethanedioxy)]bis acetyl chloride, 0.92 g (8 mmol) D-proline and 1.2 ml triethylamine in 200 ml dimethylformamide was stirred for three days at ambient temperature. The
30 solvent was distilled off under vakuum. The residue was chromatographed over silicagel with methanol to yield 0.42 g (R)-1-[[2-[2-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]ethoxy]-ethoxy]-acetyl]-pyrrolidine-2-carboxylic

acid as a beige hygroscopic solid.

MS-ISP: 417 (M+H)⁺.

Example 12

5 (R)-1-[3-[4-[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-phenyl]-propionyl]-pyrrolidine-2-carboxylic acid

A mixture of 1.30 g (5 mmol) 1,4-benzene dipropanoyl dichloride, 1.15 g (10 mmol) D-proline and 1.5 ml triethylamine in 100 ml dimethylformamide was stirred for 24 hours at ambient temperature. The suspension was filtered and
10 the solvent was distilled off under vakuum. The residue was taken up in acetoacetate, washed with 2N hydrochloric acid, dried over sodiumsulfate and chromatographed over silicagel with dichloromethane/ acetone/formic acid 80/5/15 to yield 0.48 g (R)-1-[3-[4-[3-[(R)-2-carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-phenyl]-propionyl]-pyrrolidine-2-carboxylic acid as colorless foam.
15 MS-ISP: 417 (M+H)⁺.

Example 13

20 (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]acetyl]-pyrrolidine-2-carboxylic acid

A mixture of 1.15 g (5 mmol) 1,4-benzene diacetyl dichloride, 1.15 g (10 mmol) D-proline and 1.5 ml triethylamine in 100 ml dimethylformamide was stirred for 20 hours at ambient temperature. The solvent was distilled off under vakuum. The residue was taken up in 30 ml 2N hydrochloric acid, treated
25 with ultrasound, filtered and dried to yield 1.19 g of a brown solid. This was stirred and refluxed for 30 minutes in 300 ml methanol. Filtration, evaporation and recrystallization from methanol/acetoacetate gave 0.18 g (R)-1-[[4-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]acetyl]-pyrrolidine-2-carboxylic acid as yellow crystals with melting point 210-214°C.
30 MS-ISP: 389 (M+H)⁺.

Example 14

(R)-1-[[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[2-[2-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]acetyl]-pyrrolidine-2-carboxylic acid benzyl ester

To 0.566 g (2.5 mmol) 1,2-phenylenedioxyacetic acid in 60 ml tetrahydrofuran was added a solution of 0.81 g (5 mmol) carbonyldiimidazole in 25 ml tetrahydrofuran. After stirring at room temperature for two hours 1.21 g (5 mmol) D-proline-benzylester hydrochloride in 30 ml dichloromethane and 1.4 ml triethylamine were added and stirring was continued over the weekend. The mixture was extracted with 2N hydrochloric acid and brine. Drying with sodium sulfate, evaporation of the solvent and chromatography over silicagel with ethylacetate yielded 0.25 g (R)-1-[[2-[2-[(R)-2-benzoyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]acetyl]-pyrrolidine-2-carboxylic acid benzyl ester as colorless oil.

MS m/e (%): 600 (1, M⁺), 509 (1), 368 (25), 246 (17), 217 (19), 204 (14), 91 (100).

b) (R)-1-[[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

230 mg (0.38 mmol) (R)-1-[[2-[2-[(R)-2-benzoyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]acetyl]-pyrrolidine-2-carboxylic acid benzyl ester in 100 ml ethanol were hydrogenated in the presence of 30 mg 5% Pd on carbon. Filtration and evaporation of the solvent yielded 0.2 g (R)-1-[[2-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid as colourless glass (that still contained small amounts of ethanol)

MS-ISP: 421 (M+H)⁺.

Example 15

(R)-1-[3-[6-[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-pyridin-2-yl]propionyl]-pyrrolidine-2-carboxylic acid

a) 3-[6-(2-Carboxy-ethyl)-pyridin-2-yl]-propionic acid

A mixture of 25.6 g (0.2 mol) naphthalin and 1.39 g (0.2 mol) lithium in 150 ml tetrahydrofurane was stirred for three hours at room temperature. After

cooling to -15°C a solution of 5.72 ml (0.1 mol) acetic acid in 10 ml tetrahydrofuran was added and stirring was continued for three hours at room temperature. Then 13.3 g (0.5 mol) 2,6-bis-(bromomethyl)pyridine in 65 ml tetrahydrofuran was added and stirring was continued over night at room temperature. 200 ml ether were added and the mixture was extracted with water. The water layers were filtered through 200ml BioRad AG1-X8 ion exchanger. The ion exchanger was washed with water until neutral and then eluated with acetic acid/water. Product containing fractions were evaporated, dissolved in water and lyophylised to yield 3.9 g (35%) 3-[6-(2-carboxy-ethyl)-pyridin-2-yl]-propionic acid as a light yellow powder.

MS m/e (%): 223 (M⁺,15), 178(100), 160(81), 132(68), 104(16), 77(13).

b) (R)-1-[3-[6-[3-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-pyridin-2-yl]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester

To 0.45 g (2.0 mmol) 3-[6-(2-carboxy-ethyl)-pyridin-2-yl]-propionic acid in a mixture of 25 ml tetrahydrofuran and 25 ml dichloromethane was added a solution of 0.65 g (4.0 mmol) carbonyldiimidazole in 20 ml tetrahydrofuran. After stirring at room temperature for two hours 0.97 g (4.0 mmol) D-proline-benzylester hydrochloride in 50 ml dichloromethane and 1.12 g triethylamine were added and stirring was continued for 18 hours. Then ethylacetate was added to the mixture followed by extraction with water. Drying with sodiumsulfate, evaporation of the solvent and chromatography over silicagel with ethylacetate/hexane 2/8, then ethylacetate, then ethylacetate/methanol 95/5 followed by a second chromatography of the product containing fractions on silicagel with acetone/hexane 6/4 yielded 0.18 g (15%) (R)-1-[3-[6-[3-[(R)-2-benzoyloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-pyridin-2-yl]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester as acolourless oil.

MS-ISP: 598 (M+H)⁺.

c) (R)-1-[3-[6-[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-pyridin-2-yl]propionyl]-pyrrolidine-2-carboxylic acid

0.17 g (0.29 mmol) (R)-1-[3-[6-[3-[(R)-2-benzoyloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-pyridin-2-yl]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester

in 100 ml ethanol were hydrogenated in the presence of 35 mg 5% Pd on carbon. Filtration and evaporation of the solvent yielded 0.11g (92%) (R)-1-[3-[6-[3-[(R)-2-carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-pyridin-2-yl]propionyl]-pyrrolidine-2-carboxylic acid as colorless oil.

5 MS-ISP: 418 (M+H)⁺.

Example 16

(R)-1-[3-[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid

10 a) (R)-1-Acryloyl-pyrrolidine-2-carboxylic acid benzyl ester

To a solution of 390 mg (1.6 mmol) D-Proline benzyl ester hydrochloride and 0.47 ml (3.4 mmol) triethylamine in 20 ml dichloromethane at 0 °C was added dropwise 0.2 ml (2.4 mmol) acryloyl chloride and stirring continued for 24 h at room temperature. The reaction mixture was then washed sequentially with
15 water, 1 M hydrochloric acid and once more with water, and the aqueous phases back-extracted with dichloromethane. The combined organic extracts were dried over sodium sulphate and concentrated in vacuo to afford 420 mg (100%) of the title compound (R)-1-acryloyl-pyrrolidine-2-carboxylic acid benzyl ester as a colourless oil.

20 MS m/e (%): 259 (M⁺, 25), 124 (100), 91 (25), 70 (21).

b) (R)-1-[3-[3-[(R)-2-Benzylloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester

A solution of 400 mg (1.5 mmol) (R)-1-acryloyl-pyrrolidine-2-carboxylic acid benzyl ester and 63 ml (0.75 mmol) propylamine in 5 ml acetonitrile was
25 stirred for 16 h at room temperature, then for 6 h at 45 °C, and finally for 16 h at 80 °C. Concentration in vacuo and flash chromatography (20% H₂O in acetone) afforded 84 mg (19%) of the title compound (R)-1-[3-[3-[(R)-2-benzylloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester as a pale yellow oil.

30 MS m/e (%): 578 (M+H⁺, 100).

c) (R)-1-[3-[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid

A solution of 84 mg (0.15 mmol) (R)-1-[3-[[3-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester in 3 ml ethanol was stirred with 10 mg 10% Palladium on carbon under 1 atm of hydrogen for 16 h at room temperature.

- 5 After filtration to remove the catalyst, concentration in vacuo afforded 58 mg (100%) of the title compound (R)-1-[3-[[3-[(R)-2-carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid as a white solid. MS m/e (%): 398 (M+H⁺, 100).

10

Example 17

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-ureido]-pyrrolidine-2-carboxylic acid

a) (R)-1-tert-Butoxycarbonylaminoacetyl-pyrrolidine-2-carboxylic acid benzyl ester

- 15 1.21 g (5 mmol) D-proline-benzylester hydrochloride were dissolved in 100 ml dichloromethane and stirred with 0.7 ml triethylamine. The mixture was extracted with water, dried with sodiumsulfate and evaporated. The residue was dissolved in a mixture of 100 ml tetrahydrofuran and 50 ml chloroform. 1.03 g (5 mmol) N,N'-dicyclohexylcarbodiimide and 0.88 g (5 mmol) BOC-glycin were added and stirring was continued for 18 hours at room
20 temperature. Five drops acetic acid were added and after 10 minutes at room temperature the mixture was filtered and the solvents were distilled off. The residue was taken up in acetoacetate, washed with aqueous citric acid, with aqueous sodiumbicarbonate and water, dried with sodiumsulfate and the
25 solvent was distilled off. Chromatography over silicagel with dichloromethane/methanol 99/1 yielded 1.43g (79%) (R)-1-tert-butoxycarbonylaminoacetyl-pyrrolidine-2-carboxylic acid benzyl ester as colorless oil.

MS m/e (%): 363 (M+H⁺, 1), 306 (29), 289 (10), 114 (44), 91 (76), 70 (100), 57
30 (64).

b) (R)-1-[[3-[2-[(R)-2-Benzylloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-ureido]-pyrrolidine-2-carboxylic acid benzyl ester

- 8.6 ml trifluoric acid were added dropwise at 0 °C to a solution of 1.57 g (4.34 mmol) (R)-1-tert-butoxycarbonylaminoacetyl-pyrrolidine-2-carboxylic acid benzyl ester in 8.6 ml dichloromethane and stirring was continued for half an hour at room temperature. The solution was washed with aqueous sodiumbicarbonate, dried with sodiumsulfate and evaporated. The residue was dissolved in 200 ml dichloromethane and stirred with 0.21 g (0.7 mmol) triphosgene and 1.8 ml (13 mmol) triethylamine for four hours at room temperature. The mixture was extracted with 1N hydrochloric acid, dried with sodiumsulfate and evaporated. The remaining 1.15 g residue were chromatographed over silicagel with dichloromethane/methanol 96/4 and the fractions containing product were chromatographed again over silicagel with dichloromethane/acetone/formic acid 80/15/5 to yield 0.23 g (33%) (R)-1-[[3-[2-[(R)-2-benzylloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-ureido]-pyrrolidine-2-carboxylic acid benzyl ester as an oil.

MS-ISP: 551 (M+H)⁺.

c) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-ureido]-pyrrolidine-2-carboxylic acid

- 0.14 g (0.36 mmol) (R)-1-[[3-[2-[(R)-2-benzylloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-ureido]-pyrrolidine-2-carboxylic acid benzyl ester in 60 ml ethanol were hydrogenated in the presence of 40 mg 5% Pd on carbon. Filtration, evaporation of the solvent and crystallization from methanol/ethylacetate yielded 0.07 g (52%) as white crystals with melting point 157-160 °C.

25

Example 18

(R)-1-[10-[(R)-2-Carboxy-pyrrolidin-1-yl]-10-oxo-decanoyl]-pyrrolidine-2-carboxylic acid Ca salt (1:1)

- A mixture of 1.20 g (5 mmol) sebacyl chloride, 1.15 g (10 mmol) D-proline and 1.4 ml (10 mmol) triethylamine in 100 ml dimethylformamide was stirred over the weekend at ambient temperature. The solvent was distilled off under vakuum. The residue was taken up with 40 ml aqueous citric acid and

30

extracted with ethylacetate. Evaporation of the solvent and chromatography over silicagel with chloroform/aceton/formic acid 80/5/15 yielded 1.21g (R)-1-[10-[(R)-2-carboxy-pyrrolidin-1-yl]-10-oxo-decanoyl]-pyrrolidine-2-carboxylic acid as an oil.

5 MS-ISP: 397 (M+H)⁺.

0.89 g (2.24 mmol) of this oil were dissolved in 50 ml ethanol and stirred with 0.175 g (2.24 mmol) calciumhydroxide for 48 hours. The suspension was filtered. The solid residue was taken up in 15 ml water, warmed to 80 °C, filtrated when hot and evaporated. The residue was suspended in ether,
10 filtered and washed with ether to yield 0.5 g (R)-1-[10-[(R)-2-carboxy-pyrrolidin-1-yl]-10-oxo-decanoyl]-pyrrolidine-2-carboxylic acid Ca salt (1:1) as a white solid.

C (theory) 55.28 H (theory) 6.96 N (theory) 6.45

C (found) 55.29 H (found) 7.11 N (found) 6.08

15

Example 19

(R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid Ca salt (1:1)

A mixture of 1.10 g (5 mmol) suberoyl chloride, 1.15 g (10 mmol) D-proline and
20 1.5 ml (10 mmol) triethylamine in 100 ml dimethylformamide was stirred over 20 hours at ambient temperature. The solvent was distilled off under vakuum. The residue was taken up with 40 ml aqueous citric caid acid and extracted with ethylacetate. Evaporation of the solvent and chromatography over silicagel with chloroform/aceton/formic acid 80/5/15 yielded 0.8 g (R)-1-[8-[(R)-
25 2-carboxy-pyrrolidin-1-yl]-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid as an oil.

MS-ISP: 367 (M-H)⁺. 0.79 g (2.15 mmol) of this oil were dissolved in 40 ml ethanol and stirred with 0.167 g (2.15 mmol) calciumhydroxide for 20 hours. The suspension was filtered and the solid residue was almost dissolved in 25
30 ml water. The solution was filtrated and evaporated to yield 0.5 g (R)-1-[8-[(R)-2-carboxy-pyrrolidin-1-yl]-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid Ca salt (1:1) as a white solid.

Example 20

(R)-1-[4'-[(R)-2-Carboxy-pyrrolidin-1-yl-carbonyl]-[2,2]-bithiazolyl-4-yl]pyrrolidine-2-carboxylic acid

a) (R)-1-[4'-[(R)-2-Carboxypyrrolidin-1-yl-carbonyl]-[2,2]-bithiazolyl-4-yl]pyrrolidine-2-carboxylic acid benzyl ester

Thionylchloride (2 ml) was added to a solution of 0.26 g(1 mmol) [2,2']bithiazolyl-4,4'-dicarboxylic acid in 20 ml tetramethylurea and the mixture was stirred for three days at room temperature. Excess thionylchloride and the solvent were distilled off under vakuum. The residue was three times taken up in dimethylformamide and evaporated and then dissolved in 50 ml pyridine. 0.28 g (1.1 mmol) D-proline-benzylester hydrochloride were added and stirring continued at room temperature for 24 hours. The solvent was distilled off, the residue taken up in acetoacetate and extracted with 2N HCl and brine. Drying with sodiumcarbonate, evaporation of the solvent and chromatography on silicagel with acetoacetate/hexane 1:1 yielded 0.094 g (R)-1-[4'-[(R)-2-carboxypyrrolidin-1-yl-carbonyl]-[2,2]-bithiazolyl-4-yl]pyrrolidine-2-carboxylic acid benzyl ester.

MS m/e(%): 630 (M⁺, 24), 539 (17), 495 (44), 449 (51), 380 (25), 329 (33), 313 (30), 223 (38), 194 (83), 180 (66), 145 (24), 137 (21), 91 (100).

b) (R)-1-[4'-[(R)-2-Carboxy-pyrrolidin-1-yl-carbonyl]-[2,2]-bithiazolyl-4-yl]pyrrolidine-2-carboxylic acid

0.51 g (0.08 mmol) (R)-1-[4'-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl][2,2]-bithiazol-4-yl]pyrrolidine-2-carboxylic acid benzyl ester in 50 ml methanol were stirred with 5 ml 2N sodiumhydroxide solution at room temperature for 64 hours. After addition of 2N hydrochloric acid until pH 1 the mixture was extracted with dichloromethane. The extracts were dried with sodiumsulfate and evaporated. Chromatography ocer silicagel with dichloromethane/acetone/formic acid 80/15/5 gave 0.04 g (R)-1-[4'-[(R)-2-Carboxy-pyrrolidin-1-yl-carbonyl]-[2,2]-bithiazolyl-4-yl]pyrrolidine-2-carboxylic acid as colorless solid.

MS-ISP: 451 (M+H)⁺.

Example 21

(R)-1-[[(R)-2-Carboxy-pyrrolidin-1-yl]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-Pyrrolidine-2-carboxylic acid tert-butyl ester

The title compound was prepared according to a literature procedure (M.

- 5 Thorsen, T.P. Andersen, U. Pedersen, B. Yde and S.-O. Lawesson, *Tetrahedron* 1985, 41, 5633 - 5636.

Starting from 25.0 g (217 mmol) D-proline, 27.52 g (74%) of (R)-pyrrolidine-2-carboxylic acid tert-butyl ester were obtained as a colorless oil.

b) (R)-(1)-Bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester

- 10 To a solution of 64.9 g (322 mmol) bromoacetyl bromide in 250 ml dichloromethane at 0°C was added dropwise a solution of 27.5 g (161 mmol) (R)-pyrrolidine-2-carboxylic acid tert-butyl ester and 30 ml (177 mmol) N-ethyl-diisopropylamine in 150 ml dichloromethane within 40 min. The reaction mixture was allowed to warm to room temperature overnight and was poured
- 15 into 600 ml of water. The organic phase was separated and the water phase was extracted with 600 ml dichloromethane. The combined organic phases were washed with saturated sodium bicarbonate solution and brine, dried (magnesium sulfate) and evaporated to yield 44.1 g (94%) of the title compound as a brown oil that crystallized upon standing at room temperature.
- 20 Melting point 51.5-53.2°C.

c) (R)-1-[[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- To a solution of 34.3 g (200 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 350 ml dichloromethane at 0°C were added dropwise
- 25 27.9 ml (200 mmol) triethylamine. After stirring for 45 min at this temperature, 17.8 g (200 mmol) bromoacetyl bromide were added dropwise. Stirring was continued at 0°C for 3 h and 250 ml 1 N hydrochloric acid solution were added. The organic phase was separated and was washed with saturated sodium bicarbonate solution and brine, dried (magnesium sulfate)
- 30 and evaporated to yield 45 g of a brown oil. Trituration with ethyl acetate and cooling to -78°C gave 7.1 g (9%) of a pale yellow solid.
- Melting point 75.0-76.0°C. MS m/e (%): 405 (M+Na⁺, 11), 383 (M+H⁺, 100).

d) (R)-1-[[4-(R)-2-Carboxy-pyrrolidin-1-yl]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 382 mg (1.0 mmol) (R)-1-[[4-(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 300 mg (quantitative) of RO-64-2799/000 as a pale yellow amorphous and hygroscopic solid which still contains trace amounts of trifluoroacetic acid.

MS m/e (%): 269 (M-H⁺, 4.5), 113 (CF₃CO₂⁻, 100).

10

Example 22

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[4-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

15

To a solution of 236 mg (2.1 mmol) potassium tert-butyrate in 2 ml dimethylformamide at room temperature was added dropwise a solution of 110 mg (1.0 mmol) hydroquinone in 2 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 584 mg (2.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 380 mg (71%) of the title compound as a colorless oil.

25 MS m/e (%): 550 (M+NH₄⁺, 100), 477 (23), 421 (65).

b) (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 350 mg (0.66 mmol) (R)-1-[[4-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 265 mg (96%) of the title compound as a white powder.

30

MS m/e (%): 443 (M+Na⁺, 48), 438 (M+NH₄⁺, 39), 421 (M+H⁺, 100).

Example 23

5 (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2,3,5,6-tetrafluoro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[4-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-2,3,5,6-tetrafluoro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 185 mg (1.65 mmol) potassium tert-butyrate in 1 ml
10 dimethylformamide at room temperature was added dropwise a solution of 137 mg (0.75 mmol) tetrafluorohydroquinone in 1 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 438 mg (1.50 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml dimethylformamide was added within 1-2 min. The reaction mixture was
15 stirred for additional 4 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 87 mg (19%) of the title compound as a white foam.

MS m/e (%): 622 (M+NH₄⁺, 100), 549 (32), 493 (57).

20 b) (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2,3,5,6-tetrafluoro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 80 mg (0.13 mmol) (R)-1-[[4-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-2,3,5,6-tetrafluoro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue
25 suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 64 mg (96%) of the title compound as a white powder.

MS m/e (%): 491 (M-H⁺, 100).

30

Example 24

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-chloro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[4-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-chloro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 185 mg (1.65 mmol) potassium tert-butyrate in 1 ml dimethylformamide at room temperature was added dropwise a solution of 108
5 mg (0.75 mmol) chlorohydroquinone in 1 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 438 mg (1.50 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 4 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 149 mg (35%)
10 of the title compound as a white foam.

MS m/e (%): 584 (M+NH₄⁺, 100), 511 (48), 455 (96).

b) (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-chloro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

15 A solution of 140 mg (0.25 mmol) (R)-1-[[4-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-chloro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight.
20 Filtration and drying gave 126 mg (quantitative) of the title compound as a white powder.

MS m/e (%): 453 (M-H, 100).

Example 25

25 (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[4-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 236 mg (2.1 mmol) potassium tert-butyrate in 2 ml
30 dimethylformamide at room temperature was added dropwise a solution of 140 mg (1.0 mmol) methoxyhydroquinone in 2 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 584 mg (2.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml

dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 280 mg (50%) of the title compound as a colorless oil.

5 MS m/e (%): 580 (M+NH₄⁺, 100), 563 (M+H⁺, 75), 507 (62), 451 (67).

b) (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 250 mg (0.44 mmol) (R)-1-[[4-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methoxy-phenoxy]-acetyl]-pyrrolidine-2-
10 carboxylic acid tert-butyl ester in 4 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 188 mg (94%) of the title compound as a white powder.

15 MS m/e (%): 473 (M+Na⁺, 45), 468 (M+NH₄⁺, 24), 451 (M+H⁺, 100).

Example 26

Mixture of (R)-1-[(4-hydroxy-3- and -2-methoxy-phenoxy)-acetyl]-pyrrolidine-2-carboxylic acid

20 a) Mixture of (R)-1-[(4-hydroxy-3- and -2-methoxy-phenoxy)-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

The title compounds were formed as side products during the preparation of (R)-1-[[4-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester.

25 Isolation and purification by flash-chromatography gave 80 mg (23%) of RO-64-2915/000 as a colorless oil.

¹H-NMR (CDCl₃, ppm): 1.41 (s, 2.4H), 1.45 (s, 6.6H), 1.81-2.32 (m, 4H), 3.55-3.82 (m, 2H), 3.75 (s, 3H), 4.39-4.78 (m, 3H), 6.16-6.26 (m, 1H) 6.40-6.43 (m, 1H), 6.67-6.74 (m, 1H).

30 b) Mixture of (R)-1-[(4-hydroxy-3- and -2-methoxy-phenoxy)-acetyl]-pyrrolidine-2-carboxylic acid

To a solution of 80 mg (0.23 mmol) mixture of (R)-1-[(4-hydroxy-3- and -2-methoxy-phenoxy)-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1 ml

dichloromethane were added 5 ml of a 4 N solution of hydrochloric acid in dioxane. After 24 h, the solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 65 mg (97%) of the title compound as a white powder.

MS m/e (%): 296 (M+H⁺, 100).

Example 27

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 561 mg (5.0 mmol) potassium tert-butate in 4 ml dimethylformamide at room temperature was added dropwise a solution of 275 mg (2.5 mmol) resorcinol in 4 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 1.46 mg (5.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 830 mg (62%) of the title compound as a colorless oil.

MS m/e (%): 550 (M+NH₄⁺, 100), 533 (M+H⁺, 95), 477 (48), 421 (95).

b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 750 mg (1.41 mmol) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 6 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 15 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 581 mg (98%) of the title compound as a white powder.

MS m/e (%): 443 (M+Na⁺, 32), 438 (M+NH₄⁺, 20), 421 (M+H⁺, 100).

Example 28

(R)-1-[(3-Hydroxy-phenoxy)-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[(3-Hydroxy-phenoxy)-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- 5 The title compound was formed as side product during the preparation of (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester. Isolation and purification by flash-chromatography gave 120 mg (37%) of RO-64-2802/000 as a colorless oil.
- 10 MS m/e (%): 344 (M+Na⁺, 9), 322 (M+H⁺, 73), 266 (100).

b) (R)-1-[(3-Hydroxy-phenoxy)-acetyl]-pyrrolidine-2-carboxylic acid

- To a solution of 120 mg (0.37 mmol) (R)-1-[(3-hydroxy-phenoxy)-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1 ml dichloromethane were added 5 ml of a 4 N solution of hydrochloric acid in dioxane. After 3 d, the
- 15 solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 95 mg (97%) of the title compound as a white powder.
- MS m/e (%): 264 (M-H⁺, 100).

Example 29

- 20 (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- To a solution of 236 mg (2.1 mmol) potassium tert-butyrate in 2 ml
- 25 dimethylformamide at room temperature was added dropwise a solution of 124 mg (1.0 mmol) 2,6-dihydroxytoluene in 2 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 584 mg (2.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at
- 30 room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 335 mg (61%) of the title compound as a colorless oil.
- MS m/e (%): 564 (M+NH₄⁺, 100), 491 (27), 435 (71).

b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 300 mg (0.55 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 226 mg (95%) of the title compound as a white powder.

MS m/e (%): 457 (M+Na⁺, 54), 452 (M+NH₄⁺, 55), 435 (M+H⁺, 100).

Example 30

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 236 mg (2.1 mmol) potassium tert-butyrate in 2 ml dimethylformamide at room temperature was added dropwise a solution of 140 mg (1.0 mmol) 5-methoxyresorcinol in 2 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 584 mg (2.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 407 mg (72%) of the title compound as a colorless oil.

MS m/e (%): 580 (M+NH₄⁺, 98), 563 (M+H⁺, 100), 507 (54), 451 (95).

b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 370 mg (0.66 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight.

Filtration and drying gave 287 mg (97%) of the title compound as a white powder.

MS m/e (%): 473 (M+Na⁺, 45), 468 (M+NH₄⁺, 30), 451 (M+H⁺, 100).

5

Example 31

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-methoxycarbonyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-methoxycarbonyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl

10 ester

To a solution of 595 mg (5.3 mmol) potassium tert-butyrate in 4 ml dimethylformamide at room temperature was added dropwise a solution of 420 mg (2.5 mmol) 3,5-dihydroxybenzoate in 4 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 1.46 mg (5.0 mmol) (R)-(1)-

15 bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml

dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 2 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 1.01 g (68%) of the title compound as a colorless oil.

20 MS m/e (%): 608 (M+NH₄⁺, 92), 591 (M+H⁺, 48), 535 (41), 479 (100).

b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-methoxycarbonyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 710 mg (1.2 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-methoxycarbonyl-phenoxy]-acetyl]-pyrrolidine-

25 2-carboxylic acid tert-butyl ester in 10 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 20 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 540 mg (94%) of the title compound as a white powder.

30 MS m/e (%): 477 (M-H⁺, 100).

Example 32

(R)-1-[[3-Carboxy-5-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

To 10 ml of a 0.5 N lithium hydroxide solution in methanol/water = 3:1 were
5 added 100 mg (0.2 mmol) of (R)-1-[[3-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-methoxycarbonyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid. The solution was allowed to stand at room temperature for 24 h. The mixture was adjusted to pH 6 by dropwise addition of hydrochloric acid solution and lyophilized to give 800 mg of a white powder. The product was isolated by
10 chromatography using an ion exchange resin (Dowex). Lyophilization gave 20 mg (22%) of the title compound as a white powder.

MS m/e (%): 487 (M+Na⁺, 61), 482 (M+NH₄⁺, 54), 465 (M+H⁺, 100).

Example 33

15 a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-cyano-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 1.35 g (10 mmol) 3,5-dihydroxybenzonitrile and 5.84 g (20 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 25 ml dimethylformamide at room temperature were added 7 g anhydrous
20 potassium carbonate. After stirring for additional 20 h, the potassium salts were filtered off and the solvent was removed *in vacuo*. The residue was purified by flash-chromatography to yield 4.77 g (86%) of the title compound as a colorless foam.

MS m/e (%): 575 (M+NH₄⁺, 100), 558 (M+H⁺, 42), 502 (35), 446 (85).

25 b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-cyano-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 280 mg (0.5 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-cyano-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml trifluoroacetic acid was stirred for 18 h
30 at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 280 mg (72%) of the title compound as a white powder.

MS m/e (%): 444 (M-H⁺, 100).

Example 34

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-1H-tetrazol-5-yl]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

- To a solution of 110 mg (0.2 mmol) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-cyano-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 10 ml 1,2-dimethoxyethane were added 200 mg (0.6 mmol) tributyltin azide. The mixture was heated at reflux for 3 days.
- After cooling to room temperature, 1.4 g of gaseous hydrochloric acid were bubbled into the solution to obtain a 4 N hydrochloric acid solution in 1,2-dimethoxyethane and stirring was continued for 12 h. The solvent was removed *in vacuo* and the oily residue was triturated with ether to give 61 mg (92%) of the title compound as a pale yellow amorphous solid.
- MS m/e (%): 511 (M+Na⁺, 41), 506 (M+NH₄⁺, 32), 489 (M+H⁺, 100).

Example 35

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-hydroxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-hydroxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- To a solution of 95 mg (0.75 mmol) phloroglucinol and 438 mg (1.5 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml dimethylformamide at room temperature were added 520 mg anhydrous potassium carbonate. After stirring overnight, the potassium salts were filtered off and the solvent was removed *in vacuo*. The residue was purified by flash-chromatography to yield 50 mg (12%) of the title compound as a white foam.

MS m/e (%): 566 (M+NH₄⁺, 96), 549 (M+H⁺, 88), 493 (47), 437 (100).

b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-hydroxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 46 mg (0.084 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-hydroxy-phenoxy]-acetyl]-pyrrolidine-2-

carboxylic acid tert-butyl ester in 1 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 5 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 35 mg (96%) of the title compound as a light brown powder.

MS m/e (%): 435 (M-H⁺, 100).

Example 36

(R)-1-[[3,5-Bis-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[3,5-Bis-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 95 mg (0.75 mmol) phloroglucinol and 220 mg (0.75 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml dimethylformamide at room temperature were added 520 mg anhydrous potassium carbonate. After 2 h and 6 h stirring at room temperature, 220 mg (0.75 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester were added (per addition) and stirring was continued for 18 h. The potassium salts were filtered off and the solvent was removed *in vacuo*. The residue was purified by flash-chromatography to yield 465 mg (82%) of the title compound as a colorless foam.

MS m/e (%): 777 (M+NH₄⁺, 100), 760 (M+H⁺, 4), 704 (11), 648 (22), 592 (22).

b) (R)-1-[[3,5-Bis-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 350 mg (0.47 mmol) (R)-1-[[3,5-bis-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 254 mg (92%) of the title compound as a light brown powder.

MS m/e (%): 614 (M+Na⁺, 73), 609 (M+NH₄⁺, 100), 592 (M+H⁺, 56).

Example 37

Mixture of (E)- and (Z)-(R)-1-[[3-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(N-hydroxycarbamimidoyl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

- 5 a) Mixture of (E)- and (Z)-(R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(N-hydroxycarbamimidoyl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 1.25 g (17.9 mmol) hydroxylamine hydrochloride in 6 ml dimethyl sulfoxide were added 1.82 g (18 mmol) triethylamine. An insoluble
10 material was filtered off and was washed with 5 ml tetrahydrofuran. The filtrate was concentrated *in vacuo* at 100 mbar to remove tetrahydrofuran and 2.0 g (3.59 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-cyano-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester were added. After stirring for 20 h at 75°C, the reaction mixture was diluted
15 with water and extracted with ethyl acetate. The organic solution was extracted with 1 N hydrochloric acid solution. The aqueous solution was adjusted to pH 10 with 1 N sodium hydroxide solution and extracted with ethyl acetate. The organic solution was washed with water, dried (sodium sulfate) and evaporated to give 1.74 g of the title compound as a colorless
20 foam.

MS m/e (%): 613 (M+Na⁺, 7), 591 (M+H⁺, 100), 535 (21), 479 (15).

b) Mixture of (E)- and (Z)-(R)-1-[[3-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(N-hydroxycarbamimidoyl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

- 25 A solution of 120 mg (0.2 mmol) mixture of (E)- and (Z)-(R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(N-hydroxycarbamimidoyl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting
30 suspension was stirred overnight. Filtration and drying gave 96 mg (quantitative) of the title compound as a light brown powder.

MS m/e (%): 501 (M+Na⁺, 31), 479 (M+H⁺, 100).

Example 38

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(5-oxo-4,5-dihydro-[1,2,4]oxadiazol-3-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

- 5 a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(5-oxo-4,5-dihydro-[1,2,4]oxadiazol-3-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To an ice-cooled solution of 300 mg (0.51 mmol) mixture of (E)- and (Z)-(R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(N-hydroxycarbamimidoyl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester and 43 mg (0.55 mmol) pyridine in 2 ml dimethylformamide were added dropwise 200 mg (0.51 mmol) 2-ethylhexyl chloroformate. The mixture was stirred at 0°C for 30 min, diluted with water and extracted with ethyl acetate. The organic phase was dried (sodium sulfate) evaporated and the oily residue was dissolved in 20 ml toluene. After 16 h stirring at room temperature, the toluene solution was washed with brine, dried (sodium sulfate) and filtered over silica gel to give 280 mg (89%) of the title compound as a viscous oil.

MS m/e (%): 639 (M+Na⁺, 51), 634 (M+NH₄⁺, 100), 617 (M+H⁺, 14), 561 (40), 505 (79).

- 20 b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(5-oxo-4,5-dihydro-[1,2,4]oxadiazol-3-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 220 mg (0.36 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(5-oxo-4,5-dihydro-[1,2,4]oxadiazol-3-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 135 mg (74%) of the title compound as a light brown powder.

MS m/e (%): 527 (M+Na⁺, 73), 522 (M+NH₄⁺, 75), 505 (M+H⁺, 100).

Example 39

(R)-1-[[3-[2-[(R)-2-Carbonxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(2-oxo-2,3-dihydro-[1,2,3,5]oxathiadiazol-4-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid (config. of S-oxide R:S=1:1)

- 5 a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(2-oxo-2,3-dihydro-[1,2,3,5]oxathiadiazol-4-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester (config. of S-oxide R:S=1:1)

To an ice-cooled solution of 300 mg (0.51 mmol) mixture of (E)- and (Z)-(R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(N-

- 10 hydroxycarbamimidoyl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester and 80 mg (1 mmol) pyridine in 2 ml dichloromethane were added dropwise 60 mg (0.51 mmol) thionyl chloride (dissolved in 0.3 ml dichloromethane). The mixture was stirred at 0°C for 45 min, diluted with dichloromethane, washed with water, dried (sodium sulfate) and evaporated.
- 15 Flash-chromatography gave 190 mg (59%) of the title compound as a semi-solid liquid.

MS m/e (%): 659 (M+Na⁺, 14), 654 (M+NH₄⁺, 100), 637 (M+H⁺, 11), 581 (14), 525 (74).

- b) (R)-1-[[3-[2-[(R)-2-Carbonxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(2-oxo-2,3-dihydro-[1,2,3,5]oxathiadiazol-4-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid (config. of S-oxide R:S=1:1)
- 20

A solution of 165 mg (0.26 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(2-oxo-2,3-dihydro-[1,2,3,5]oxathiadiazol-4-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester (config. of S-oxide R:S=1:1)

25

in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 115 mg (85%) of the title compound as a light brown powder.

- 30 MS m/e (%): 547 (M+Na⁺, 65), 542 (M+NH₄⁺, 86), 525 (M+H⁺, 100).

Example 40

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(5-tert-

5 butylsulfanyl-[1,2,4]oxadiazol-3-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(5-thioxo-4,5-dihydro-[1,2,4]oxadiazol-3-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

10 A mixture of 515 mg (0.87 mmol) mixture of (E)- and (Z)- (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(N-hydroxycarbamimidoyl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester, 177 mg (0.94 mmol) 1,1'-thiocarbonyldiimidazole and 530 mg (3.5 mmol) 1,8-diazabicyclo[5.4.0]undec-7-ene in 10 ml acetonitrile was stirred at
15 room temperature for 4 h. The mixture was concentrated in *in vacuo*, diluted with water, adjusted to pH 4-5 with 1 N hydrochloric acid solution and extracted with ethyl acetate. The extract was concentrated again, the residue was dissolved in 100 ml 1 N sodium hydroxide solution and washed with ether. The aqueous solution was adjusted to pH 4 with 1 N hydrochloric acid
20 solution and was extracted with ethyl acetate. The organic phase was washed with water, dried (sodium sulfate) and was evaporated to give 490 mg (89%) of the title compound as a pale brown foam.

MS m/e (%): 655 (M+Na⁺, 35), 650 (M+NH₄⁺, 100), 633 (M+H⁺, 14), 577 (31), 521 (52).

25 b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(5-tert-butylsulfanyl-[1,2,4]oxadiazol-3-yl)-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 315 mg (0.5 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-5-(5-thioxo-4,5-dihydro-[1,2,4]oxadiazol-3-yl)-
30 phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting

suspension was stirred overnight. Filtration and drying gave 160 mg (57%) of the title compound as a white powder.

MS m/e (%): 599 (M+Na⁺, 54), 594 (M+NH₄⁺, 71), 577 (M+H⁺, 100), 521 (37).

5

Example 41

(R)-1-[[2-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-3-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[2-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-3-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- 10 To a solution of 236 mg (2.1 mmol) potassium tert-butyrate in 2 ml dimethylformamide at room temperature was added dropwise a solution of 140 mg (1.0 mmol) pyrogallol-1-methyl ether in 2 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 584 mg (2.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml
- 15 dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 375 mg (67%) of the title compound as a colorless oil.

MS m/e (%): 580 (M+NH₄⁺, 14), 563 (M+H⁺, 100), 507 (14), 451 (12).

- 20 b) (R)-1-[[2-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-3-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

- A solution of 345 mg (0.61 mmol) (R)-1-[[2-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-3-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml trifluoroacetic acid was stirred for 4 h
- 25 at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 265 mg (96%) of the title compound as a white powder.

MS m/e (%): 473 (M+Na⁺, 54), 451 (M+H⁺, 100).

30

Example 42

(R)-1-[[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-3-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[2-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-3-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 236 mg (2.1 mmol) potassium tert-butyrate in 2 ml dimethylformamide at room temperature was added dropwise a solution of 124 mg (1.0 mmol) 3-methylcatechol in 2 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 584 mg (2.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 356 mg (65%) of the title compound as a colorless oil.

MS m/e (%): 547 (M+H⁺, 100), 491 (14), 435 (17).

b) (R)-1-[[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-3-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 325 mg (0.60 mmol) (R)-1-[[2-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-3-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 235 mg (91%) of the title compound as a white powder.

MS m/e (%): 457 (M+Na⁺, 59), 435 (M+H⁺, 100).

Beispiel 43

25 (R)-1-[[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-3,4,5,6-tetrachloro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[2-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-3,4,5,6-tetrachloro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 185 mg (1.65 mmol) potassium tert-butyrate in 1 ml dimethylformamide at room temperature was added dropwise a solution of 186 mg (0.75 mmol) tetrachlorocatechol in 1 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 438 mg (1.50 mmol) (R)-(1)-

bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 4 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 229 mg (45%) of the title compound as a white foam.

MS m/e (%): 671 (M+H⁺, 100), 615 (19), 559 (14).

b) (R)-1-[[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-3,4,5,6-tetrachloro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 220 mg (0.34 mmol) (R)-1-[[2-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-3,4,5,6-tetrachloro-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 172 mg (94%) of the title compound as a white powder.

MS m/e (%): 559 (M-H⁺, 100).

Example 44

R)-1-[[6-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[6-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 236 mg (2.1 mmol) potassium tert-butyrate in 2 ml dimethylformamide at room temperature was added dropwise a solution of 160 mg (1.0 mmol) 2,6-dihydroxynaphthalene in 2 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 584 mg (2.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 120 mg (21%) of the title compound as a white solid.

MS m/e (%): 605 (M+Na⁺, 14), 600 (M+NH₄⁺, 92), 583 (M+H⁺, 5), 527 (32), 471 (100).

b) (R)-1-[[6-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

- A solution of 75 mg (0.13 mmol) (R)-1-[[6-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 18 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 60 mg (98%) of the title compound as a white powder.
- 10 MS m/e (%): 493 (M+Na⁺, 58), 488 (M+NH₄⁺, 33), 471 (M+H⁺, 100).

Example 45

(R)-1-[[5-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

- 15 a) (R)-1-[[5-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- To a solution of 236 mg (2.1 mmol) potassium tert-butyrate in 2 ml dimethylformamide at room temperature was added dropwise a solution of 160 mg (1.0 mmol) 1,5-dihydroxynaphthalene in 2 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 584 mg (2.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 470 mg (81%) of the title compound as a pale yellow foam.
- 20
- 25

MS m/e (%): 605 (M+Na⁺, 10), 600 (M+NH₄⁺, 100), 527 (20), 471 (63).

b) (R)-1-[[5-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

- A solution of 290 mg (0.5 mmol) (R)-1-[[5-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml trifluoroacetic acid was stirred for 18 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight.
- 30

Filtration and drying gave 230 mg (98%) of the title compound as a white powder.

MS m/e (%): 493 (M+Na⁺, 45), 488 (M+NH₄⁺, 37), 471 (M+H⁺, 100).

5

Example 46

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- 10 To a solution of 236 mg (2.1 mmol) potassium tert-butyrate in 2 ml dimethylformamide at room temperature was added dropwise a solution of 160 mg (1.0 mmol) 2,3-dihydroxynaphthalene in 2 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 584 mg (2.0 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml
- 15 dimethylformamide was added within 1-2 min. The reaction mixture was stirred for additional 3 h at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 310 mg (53%) of the title compound as a colorless oil.

MS m/e (%): 605 (M+Na⁺, 27), 600 (M+NH₄⁺, 16), 583 (M+H⁺, 100), 527 (16),

20 471 (24).

b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

- A solution of 230 mg (0.4 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-
- 25 carboxylic acid tert-butyl ester in 5 ml trifluoroacetic acid was stirred for 18 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 180 mg (96%) of the title compound as a white powder.
- 30 MS m/e (%): 469 (M-H⁺, 100).

Example 47

(R)-1-[[2'-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-biphenyl-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[2'-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-

5 biphenyl-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 160 mg (1.44 mmol) potassium tert-butyrate in 1.6 ml dimethylformamide at room temperature was added dropwise a solution of 127 mg (0.69 mmol) 2,2'-dihydroxybiphenyl in 1 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 400 mg (1.37 mmol) (R)-(1)-

10 bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred overnight at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 84 mg (20%) of the title compound as a light brown oil.

15 MS m/e (%): 631 (M+Na⁺, 22), 626 (M+NH₄⁺, 100), 609 (M+H⁺, 16).

b) (R)-1-[[2'-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-biphenyl-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 87 mg (0.14 mmol) (R)-1-[[2'-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-biphenyl-2-yloxy]-acetyl]-pyrrolidine-2-

20 carboxylic acid tert-butyl ester

in 1 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 31 mg (44%) of the title compound as a light brown powder.

25 MS m/e (%): 519 (M+Na⁺, 62), 497 (M+H⁺, 100).

Example 48

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

30 a) (R)-1-[[4-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 160 mg (1.44 mmol) potassium tert-butyrate in 1.6 ml dimethylformamide at room temperature was added dropwise a solution of 116

mg (0.69 mmol) 1,4-naphthoquinone in 1 ml dimethylformamide. Stirring was continued for 2-3 min and a solution of 400 mg (1.37 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml dimethylformamide was added within 1-2 min. The reaction mixture was stirred overnight at room temperature. The solvent was removed *in vacuo* and the residue purified by flash-chromatography to yield 269 mg (67%) of the title compound as a light brown foam.

MS m/e (%): 600 (M+NH₄⁺, 100).

10 b) (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 172 mg (0.30 mmol) (R)-1-[[4-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 140 mg (quantitative) of the title compound as a light brown powder.

MS m/e (%): 469 (M-H⁻, 100).

20

Example 49

(R)-1-[[7-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[7-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

25 To a solution of 110 mg (0.69 mmol) 2,7-dihydroxynaphthalene and 400 mg (1.37 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 1.5 ml dimethylformamide at room temperature were added 480 mg anhydrous potassium carbonate. After stirring for additional 20 h, the potassium salts were filtered off and the solvent was removed *in vacuo*. The residue was purified by flash-chromatography to yield 296 mg (74%) of the title compound as a white foam.

30

MS m/e (%): 600 (M+NH₄⁺, 100), 527 (25), 471 (99).

b) (R)-1-[[7-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 272 mg (0.47 mmol) (R)-1-[[7-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-2-yloxy]-acetyl]-pyrrolidine-2-

5 carboxylic acid tert-butyl ester in 1.5 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 209 mg (95%) of the title compound as a white powder.

10 MS m/e (%): 493 (M+Na⁺, 48), 488 (M+NH₄⁺, 23), 471 (M+H⁺, 100).

Example 50

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:2)

15 a) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 75 mg (0.69 mmol) 1,3-phenylenediamine and 400 mg (1.37 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 3 ml dichloromethane at room temperature were added 0.21 ml (1.5 mmol)

20 triethylamine. The reaction mixture was stirred overnight at room temperature. The organic phase was washed with 1 N sodium carbonate solution and brine, dried (sodium sulfate) and evaporated. The residue was purified by flash-chromatography to yield 285 mg (78%) of the title compound as a light brown foam.

25 MS m/e (%): 553 (M+Na⁺, 15), 531 (M+H⁺, 100), 475 (20), 419 (10).

b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:2)

A solution of 170 mg (0.32 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-

30 carboxylic acid tert-butyl ester in 4 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight.

Filtration and drying gave 130 mg (63%) of the title compound as a brown foam.

MS m/e (%): 417 (M-H⁺, 100).

5

Example 51

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:2)

a) (R)-1-[[4-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- 10 To a solution of 75 mg (0.69 mmol) 1,4-phenylenediamine and 400 mg (1.37 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml tetrahydrofuran at room temperature were added 190 mg anhydrous potassium carbonate. The reaction mixture was stirred overnight at 40°C. The potassium salts were filtered off and the solvent was removed *in vacuo*. The residue was purified by flash-chromatography to yield 180 mg (50%) of the title compound as a brown foam.

MS m/e (%): 553 (M+Na⁺, 26), 531 (M+H⁺, 100).

b) (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:2)

- 20 A solution of 168 mg (0.32 mmol) (R)-1-[[4-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed, the residue suspended in toluene and the solvent removed again to eliminate excess trifluoroacetic acid.
- 25 The residue was re-suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 160 mg (78%) of the title compound as a brown foam.

MS m/e (%): 417 (M-H⁺, 100).

30

Example 52

(R)-1-[[[2-[(R)-2-Carbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-butyl-aminol]-acetyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

a) (R)-1-[[[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-butyl-amino]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 136 mg (1.86 mmol) butylamine and 995 mg (3.40 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml

5 dichloromethane at room temperature were added 0.52 ml (3.74 mmol)

triethylamine. The suspension was stirred overnight at room temperature.

The organic phase was washed with 1 N sodium carbonate solution and brine, dried (sodium sulfate) and evaporated. The residue was purified by flash-chromatography to yield 830 mg (98%) of the title compound as a light brown

10 oil.

MS m/e (%): 524 (2M+Ni⁺, 37), 496 (M+H⁺, 100), 468 (2), 440 (7).

b) (R)-1-[[[2-[(R)-2-Carbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-butyl-amino]-acetyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

A solution of 50 mg (0.1 mmol) (R)-1-[[[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-

15 1-yl]-2-oxo-ethyl]-butyl-amino]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 0.2 ml trifluoroacetic acid was stirred for 4 h at room temperature.

The solvent was removed *in vacuo* and the residue suspended in 10 ml ether.

The resulting suspension was stirred overnight. Filtration and drying gave 46 mg (92%) of the title compound as a light brown powder.

20 MS m/e (%): 382 (M-H⁺, 100).

Example 53

(R)-1-[[[2-[(R)-2-Carbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-(2-methoxy-ethyl)-amino]-acetyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

25 a) (R)-1-[[[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-(2-methoxy-ethyl)-amino]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 137 mg (1.83 mmol) 2-methoxyethylamine and 970 mg (3.33 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 4 ml dichloromethane at room temperature were added 0.51 ml (3.66 mmol)

30 triethylamine. The suspension was stirred overnight at room temperature.

The organic phase was washed with 1 N sodium carbonate solution and brine, dried (sodium sulfate) and evaporated. The residue was purified by flash-

chromatography to yield 397 mg (48%) of the title compound as a light brown oil.

MS m/e (%): 520 (M+Na⁺, 50), 498 (M+H⁺, 100).

b) (R)-1-[[[2-[(R)-2-Carbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-(2-methoxy-

5 ethyl)amino]-acetyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

A solution of 367 mg (0.74 mmol) (R)-1-[[[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-(2-methoxy-ethyl)-amino]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue
10 suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 350 mg (95%) of the title compound as a light brown foam.

MS m/e (%): 384 (M-H⁺, 100).

15

Example 54

(R)-1-[[Benzyl-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-amino]-acetyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

a) (R)-1-[[Benzyl-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-amino]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

20 To a solution of 104 mg (0.97 mmol) benzylamine and 514 mg (1.76 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml dichloromethane at room temperature were added 0.27 ml (1.94 mmol) triethylamine. The suspension was stirred overnight at room temperature. The organic phase was washed with 1 N sodium carbonate solution and brine,
25 dried (sodium sulfate) and evaporated. The residue was purified by flash-chromatography to yield 270 mg (58%) of the title compound as a light brown oil.

MS m/e (%): 552 (M+Na⁺, 12), 530 (M+H⁺, 100).

b) (R)-1-[[Benzyl-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-amino]-acetyl]-
30 pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

A solution of 200 mg (0.38 mmol) (R)-1-[[Benzyl-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-amino]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

in 2 ml trifluoroacetic acid was stirred for 3 h at room temperature. The solvent was removed *in vacuo* and the residue suspended in 10 ml ether. The resulting suspension was stirred overnight. Filtration and drying gave 189 mg (94%) of the title compound as white crystals.

5 MS m/e (%): 440 (M+Na⁺, 16), 418 (M+H⁺, 100).

Example 55

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-1,3-dibutyl-ureido]-acetyl]-pyrrolidine-2-carboxylic acid

10 a) (R)-1-Butylaminoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 1.58 ml (16 mmol) butylamine in 1 ml dichloromethane was added dropwise a solution of 470 mg (1.6 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml dichloromethane at room temperature. The suspension was stirred overnight. The organic phase was washed with 1 N sodium carbonate solution and brine, dried (sodium sulfate) and evaporated. The residue was purified by flash-chromatography to yield 370 mg (81%) of the title compound as a light brown oil.

MS m/e (%): 285 (M+H⁺, 100).

20 b) (R)-1-[[3-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-1,3-dibutyl-ureido]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

To a solution of 165 mg (0.58 mmol) (R)-1-butylaminoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml toluene were added 305 mg anhydrous sodium carbonate. The suspension was evaporated to dryness and the residue was re-suspended in 2 ml tetrahydrofuran. Then, 1.4 ml (2.88 mmol) of a phosgen solution (20 % in toluene) were added and stirring was continued for 6 h at 50°C. The sodium salts were filtered off and washed with tetrahydrofuran. The filtrate was evaporated and 165 mg (0.58 mmol) RO-64-2576/000 dissolved in 2 ml tetrahydrofuran and 305 mg anhydrous sodium carbonate were added again. After stirring overnight at 50°C, the organic phase was washed with 1 N hydrochloric acid solution, dried (magnesium sulfate) and evaporated to give 330 mg (91%) of the title compound as a light brown powder.

MS m/e (%): 617 (M+Na⁺, 19), 595 (M+H⁺, 100).

c) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-1,3-dibutyl-ureido]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 278 mg (0.47 mmol) (R)-1-[[3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-1,3-dibutyl-ureido]-acetyl]-pyrrolidine-2-

5 carboxylic acid tert-butyl ester

in 2 ml trifluoroacetic acid was stirred for 4 h at room temperature. The solvent was removed *in vacuo* and the residue was dried to give 167 mg (74%) of the title compound as brown foam.

MS m/e (%): 505 (M+Na⁺, 11), 487 (33), 483 (M+H⁺, 33), 465 (100).

10

Example 56

(R)-1-[[1,3-Dibenzyl-3-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-ureido]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-Benzylaminoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester

15 To a solution of 1.25 ml (11.4 mmol) benzylamine in 1 ml dichloromethane was added dropwise a solution of 335 mg (1.14 mmol) (R)-(1)-bromoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml dichloromethane at 0°C. The suspension was stirred for 4 h at this temperature. The organic phase was washed with 1 N sodium carbonate solution and brine, dried (sodium sulfate)

20 and evaporated. The residue was purified by flash-chromatography to yield 312 mg (86%) of the title compound as an amorphous solid.

MS m/e (%): 319 (M+H⁺, 100).

b) (R)-1-[[1,3-Dibenzyl-3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-ureido]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

25 To a solution of 233 mg (0.73 mmol) (R)-1-benzylaminoacetyl-pyrrolidine-2-carboxylic acid tert-butyl ester in 2 ml toluene at room temperature were added 0.20 ml (1.34 mmol) triethylamine. Over a period of 3 h, 0.30 ml (0.70 mmol) of a phosgen solution (20 % in toluene) were added in 6 portions and stirring was continued for 3 h at room temperature. The organic phase was

30 washed with 1 N hydrochloric acid solution, 1 N sodium carbonate solution and brine, dried (magnesium sulfate) and evaporated. The residue was purified by flash-chromatography to give 245 mg (50%) of the title compound as a light yellow oil.

MS m/e (%): 685 (M+Na⁺, 28), 680 (M+NH₄⁺, 100), 663 (M+H⁺, 30).

a) (R)-1-[[1,3-Dibenzyl-3-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-ureido]-acetyl]-pyrrolidine-2-carboxylic acid

A solution of 220 mg (0.33 mmol) (R)-1-[[1,3-Dibenzyl-3-[2-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-ureido]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml 4 N hydrochloric acid solution in dioxane was stirred for 5 h at room temperature. The product was extracted with 1 N sodium hydroxide solution, the basic extract was acidified with 1 N hydrochloric acid solution to pH 4 and the product extracted with ethyl acetate. The organic phase was dried (magnesium sulfate) and evaporated to give 74 mg (41%) of the title compound as a brown oil.

MS m/e (%): 549 (M-H⁺, 100).

Example 57

15 (R)-1-Benzylaminoacetyl-pyrrolidine-2-carboxylic acid

The title compound was formed via urea cleavage as side product during the preparation of (R)-1-[[1,3-Dibenzyl-3-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-ureido]-acetyl]-pyrrolidine-2-carboxylic acid. It was isolated from the acidified aqueous solution as follows. The water phase was concentrated and the residue was suspended in 2-propanol. Inorganic salts were filtered off over celite. Evaporation and drying gave 78 mg (45%) of (R)-1-benzylaminoacetyl-pyrrolidine-2-carboxylic acid as a colorless foam.

MS m/e (%): 261 (M-H⁺, 100).

25 In the following Examples the general procedures are used:

General Procedure A: EDC Coupling Reaction

To a stirred solution of D-proline benzyl ester hydrochloride (2 equiv.), a dicarboxylic acid (1 equiv.), N-methylmorpholine (6 equiv.) and hydroxybenzotriazole (2 equiv.) in dichloromethane at 0 °C was added N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (2 equiv.) and stirring continued at 0 °C for 2 h and then at room temperature for 16 h. The reaction mixture was then washed sequentially with 1 M hydrochloric acid, saturated sodium bicarbonate solution and finally with saturated brine, and

the aqueous phases back-extracted with dichloromethane. The combined organic extracts were dried over sodium sulphate and concentrated *in vacuo*. Purification by flash chromatography on kieselgel then afforded the title compound.

5 General Procedure B: Hydrogenolysis of benzyl ester

A solution of the benzyl ester in isopropanol was stirred with 5 wt% of 10% palladium on charcoal under 1 atm of hydrogen for 16 h at room temperature. After filtration to remove the catalyst, the reaction mixture was concentrated *in vacuo* and azeotroped three times with chloroform on a rotary evaporator to
10 remove last traces of isopropanol. The resulting product (often a viscous oil, semi-solid or foam) was triturated in ether and then dried *in vacuo* (10 mbar) at 50 °C for 16 h.

Example 58

15 (R)-1-[5-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,4-dimethyl-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

a) (R)-1-[5-[(R)-2-Benzylloxymethyl-pyrrolidin-1-yl]-2,4-dimethyl-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 3 diastereomers)

20 Using General Procedure A with 6.05 g (25 mmol) D-Proline benzyl ester hydrochloride and 2.0 g (12.5 mmol) 2,4-dimethylglutaric acid afforded, after flash chromatography (EtOAc), 4.3 g (64%) of the title compound as a light yellow oil. MS m/e (%): 535 (M+H⁺, 100).

b) (R)-1-[5-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,4-dimethyl-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

25 Using General Procedure B with 4.30 g (8.05 mmol) (R)-1-[5-[(R)-2-benzylloxymethyl-pyrrolidin-1-yl]-2,4-dimethyl-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 3 diastereomers) afforded 2.58 g (91%) of the title compound as a white foam. MS m/e (%): 355 (M+H⁺, 100).

30

Example 59

(R)-1-[4-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,3-dimethyl-4-oxo-butyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

a) (R)-1-[4-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-2,3-dimethyl-4-oxo-butyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 3 diastereomers)

Using General Procedure A with 2.0 g (8.2 mmol) D-Proline benzyl ester hydrochloride and 600 mg (4.1 mmol) 2,3-dimethylsuccinic acid afforded, after
5 flash chromatography (EtOAc), 1.45 g (69%) of the title compound as a colourless oil. MS m/e (%): 521 (M+H⁺, 100).

b) (R)-1-[4-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,3-dimethyl-4-oxo-butyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

Using General Procedure B with 1.45 g (2.78 mmol) (R)-1-[4-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-2,3-dimethyl-4-oxo-butyl]-pyrrolidine-2-
10 carboxylic acid benzyl ester (mixture of 3 diastereomers) afforded 780 mg (80%) of the title compound as a white foam. MS m/e (%): 341 (M+H⁺, 100).

Example 60

15

(R)-1-[trans-4-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[trans-4-[(R)-2-Benzoyloxycarbonyl-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid benzyl ester

20 Using General Procedure A with 2.0 g (8.2 mmol) D-Proline benzyl ester hydrochloride and 700 mg (4.1 mmol) *trans*-cyclohexane-1,4-dicarboxylic acid afforded, after flash chromatography (EtOAc), 1.67 g (76%) of the title compound as a colourless oil. MS m/e (%): 547 (M+H⁺, 100).

b) (R)-1-[trans-4-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid
25

Using General Procedure B with 1.61 g (2.95 mmol) (R)-1-[trans-4-[(R)-2-Benzoyloxycarbonyl-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 1.07 g (99%) of the title compound as a white foam. MS m/e (%): 367 (M+H⁺, 100).

30

Example 61

(R)-1-[cis-4-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[cis-4-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid benzyl ester

- 5 Using General Procedure A with 2.0 g (8.2 mmol) D-Proline benzyl ester hydrochloride and 700 mg (4.1 mmol) *cis*-cyclohexane-1,4-dicarboxylic acid afforded, after flash chromatography (EtOAc), 2.0 g (91%) of the title compound as a colourless oil. MS *m/e* (%): 547 ($M+H^+$, 100).

- b) (R)-1-[cis-4-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid
- 10 Using General Procedure B with 2.0 g (3.66 mmol) (R)-1-[cis-4-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 930 mg (69%) of the title compound as a white foam. MS *m/e* (%): 365 ($[M-H]^-$, 100).

15

Example 62

(R)-1-[3-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

- a) (R)-1-[3-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 3 diastereomers)
- 20

- Using General Procedure A with 5.6 g (23.2 mmol) D-Proline benzyl ester hydrochloride and 2.0 g (11.6 mmol) cyclohexane-1,3-dicarboxylic acid afforded, after flash chromatography (EtOAc), 4.4 g (70%) of the title compound as a colourless oil. MS *m/e* (%): 547 ($M+H^+$, 100).
- 25

- b) (R)-1-[3-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)
- Using General Procedure B with 4.4 g (8.05 mmol) (R)-1-[3-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 3 diastereomers) afforded 2.9 g (98%) of the title compound as a white foam. MS *m/e* (%): 367 ($M+H^+$, 100).
- 30

Example 63

Mixture of (R)-1-[(1R,2R)- and -(1S,2S)-2-[(R)-2-carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid

- 5 a) Mixture of (R)-1-[(1R,2R)- and -(1S,2S)-2-[(R)-2-benzyloxycarbonyl-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid benzyl

Using General Procedure A with 2.0 g (8.2 mmol) D-Proline benzyl ester hydrochloride and 700 mg (4.1 mmol) *trans*-cyclohexane-1,2-dicarboxylic acid afforded, after flash chromatography (EtOAc), 1.36 g (62%) of the title
10 compound as a colourless oil. MS m/e (%): 547 (M+H⁺, 100).

b) Mixture of (R)-1-[(1R,2R)- and -(1S,2S)-2-[(R)-2-carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 1.36 g (2.48 mmol) mixture of (R)-1-[(1R,2R)- and -(1S,2S)-2-[(R)-2-benzyloxycarbonyl-pyrrolidine-1-carbonyl]-
15 cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid benzyl afforded 900 mg (99%) of the title compound as a white foam. MS m/e (%): 367 (M+H⁺, 100).

Example 64

- 20 (R)-1-[[2,5-Dihydroxy-4-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[[4-[2-[(R)-2-Benzyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-2,5-dihydroxy-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 10.7 g (44.2 mmol) D-Proline benzyl ester hydrochloride and 5.0 g (22.1 mmol) 2,5-dihydroxy-1,4-phenylenediacetic acid
25 afforded, after flash chromatography (gradient: 70-100% EtOAc/hexane), 3.17 g (24%) of the title compound as a colourless foam. MS m/e (%): 601 (M+H⁺, 100).

b) (R)-1-[[2,5-Dihydroxy-4-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid

30 Using General Procedure B with 3.17 g (5.3 mmol) (R)-1-[[4-[2-[(R)-2-Benzyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-2,5-dihydroxy-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 2.2 g (99%) of the title compound as a white crystalline solid. MS m/e (%): 421 (M+H⁺, 100).

Example 65

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid

- 5 a) (R)-1-[[3-[2-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 2.0 g (8.2 mmol) D-Proline benzyl ester hydrochloride and 800 mg (4.1 mmol) 1,3-phenylenediacetic acid afforded, after flash chromatography (EtOAc), 1.93 g (84%) of the title compound as a
10 colourless oil. MS m/e (%): 586 (M+NH₄⁺, 100), 569 (M+H⁺, 60).

b) (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 1.84 g (3.2 mmol) (R)-1-[[3-[2-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-
15 carboxylic acid benzyl ester afforded 1.21 g (97%) of the title compound as a white foam. MS m/e (%): 421 (M+H⁺, 100).

Example 66

(R)-1-[4-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-benzoyl]-pyrrolidine-2-
20 carboxylic acid

a) (R)-1-[4-[(R)-2-Benzoyloxycarbonyl-pyrrolidine-1-carbonyl]-benzoyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 5.8 g (24.0 mmol) D-Proline benzyl ester hydrochloride and 2.0 g (12.0 mmol) benzene-1,4-dioic acid afforded, after flash
25 chromatography (EtOAc), 4.66 g (72%) of the title compound as a yellow oil. MS m/e (%): 558 (M+NH₄⁺, 100), 541 (M+H⁺, 95).

b) (R)-1-[4-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-benzoyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 4.66 g (8.62 mmol) (R)-1-[4-[(R)-2-Benzoyloxycarbonyl-pyrrolidine-1-carbonyl]-benzoyl]-pyrrolidine-2-carboxylic
30 acid benzyl ester afforded 3.05 g (95%) of the title compound as a colourless foam. MS m/e (%): 378 (M+NH₄⁺, 100), 361 (M+H⁺, 55).

Example 67

(R)-1-[3-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-benzoyl]-pyrrolidine-2-carboxylic acid

- 5 a) (R)-1-[3-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-benzoyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 5.8 g (24.0 mmol) D-Proline benzyl ester hydrochloride and 2.0 g (12.0 mmol) benzene-1,3-dioic acid afforded, after flash chromatography (EtOAc), 4.68 g (72%) of the title compound as a yellow oil.

- 10 MS m/e (%): 558 (M+NH₄⁺, 100), 541 (M+H⁺, 90).

b) (R)-1-[3-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-benzoyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 2.83 g (5.24 mmol) (R)-1-[3-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-benzoyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 1.9 g (100%) of the title compound as a colourless foam. MS m/e (%): 378 (M+NH₄⁺, 100), 361 (M+H⁺, 35).

15

Example 68

(R)-1-[6-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-pyridine-2-carbonyl]-pyrrolidine-2-carboxylic acid

20

a) (R)-1-[6-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-pyridine-2-carbonyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 5.8 g (24.0 mmol) D-Proline benzyl ester hydrochloride and 2.0 g (12.0 mmol) pyridine-2,6-dicarboxylic acid afforded, after flash chromatography (EtOAc), 5.0 g (77%) of the title compound as a yellow oil. MS m/e (%): 559 (M+NH₄⁺, 60), 542 (M+H⁺, 100).

25

b) (R)-1-[6-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-pyridine-2-carbonyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 3.2 g (5.9 mmol) (R)-1-[6-[(R)-2-

30

Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-pyridine-2-carbonyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 1.9 g (90%) of the title compound as a white foam. MS m/e (%): 379 (M+NH₄⁺, 100), 362 (M+H⁺, 65).

Example 69

(R)-1-[5-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-thiophene-2-carbonyl]-pyrrolidine-2-carboxylic acid

- 5 a) (R)-1-[5-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-thiophene-2-carbonyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 2.0 g (8.2 mmol) D-Proline benzyl ester hydrochloride and 700 mg (4.1 mmol) thiophene-2,5-dicarboxylic acid afforded, after flash chromatography (EtOAc), 1.84 g (84%) of the title compound as a
10 yellow crystalline solid. MS m/e (%): 564 (M+NH₄⁺, 70), 547 (M+H⁺, 100).

b) (R)-1-[5-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-thiophene-2-carbonyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 1.84 g (3.3 mmol) (R)-1-[5-[(R)-2-benzylloxycarbonyl-pyrrolidine-1-carbonyl]-thiophene-2-carbonyl]-pyrrolidine-
15 2-carboxylic acid benzyl ester afforded 770 mg (63%) of the title compound RO-64-2667/000 as a white crystalline solid. MS m/e (%): 365 ([M-H]⁻, 65).

Example 70

(R)-1-[5-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-furan-2-carbonyl]-pyrrolidine-2-carboxylic acid

- 20 a) (R)-1-[5-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-furan-2-carbonyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 2.0 g (8.2 mmol) D-Proline benzyl ester hydrochloride and 640 mg (4.1 mmol) furan-2,5-dicarboxylic acid afforded,
25 after flash chromatography (EtOAc), 1.7 g (78%) of the title compound as a yellow oil. MS m/e

b) (R)-1-[5-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-furan-2-carbonyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 1.7 g (3.2 mmol) (R)-1-[5-[(R)-2-benzylloxycarbonyl-pyrrolidine-1-carbonyl]-furan-2-carbonyl]-pyrrolidine-2-
30 carboxylic acid benzyl ester afforded 1.02 g (91%) of the title compound as a white foam. MS m/e (%): 351 (M+H⁺, 100).

Example 71

(S)-1-[6-[(S)-2-Carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

- 5 a) (S)-1-[6-[(S)-2-Benzylloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 1.0 g (4.1 mmol) L-Proline benzyl ester hydrochloride and 300 mg (2.1 mmol) adipic acid afforded, after flash chromatography (EtOAc), 1.07 g (100%) of the title compound as a colourless oil. MS m/e (%): 521 (M+H⁺, 100).

- 10 b) (S)-1-[6-[(S)-2-Carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 1.07 g (2.1 mmol) (S)-1-[6-[(S)-2-benzylloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 609 mg (87%) of the title compound as a white
15 crystalline solid. MS m/e (%): 341 (M+H⁺, 100).

Example 72

(S)-1-[[4-[2-[(S)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid

- 20 a) (S)-1-[[4-[2-[(S)-2-Benzylloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 1.0 g (4.1 mmol) L-Proline benzyl ester hydrochloride and 400 mg (2.1 mmol) 1,4-phenylenediacetic acid afforded, after flash chromatography (EtOAc), 1.07 g (91%) of the title compound as a
25 colourless oil. MS m/e (%): 586 (M+NH₄⁺, 100), 569 (M+H⁺, 97).

- b) (S)-1-[[4-[2-[(S)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 1.07 g (1.88 mmol) (S)-1-[[4-[2-[(S)-2-benzylloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-
30 carboxylic acid benzyl ester afforded 410 mg (56%) of the title compound as a white crystalline solid. MS m/e (%): 389 (M+H⁺, 100).

Example 73

(S)-1-[[2-[2-[(S)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

- 5 a) (S)-1-[[2-[2-[(S)-2-Benzyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 1.0 g (4.1 mmol) L-Proline benzyl ester hydrochloride and 790 mg (2.1 mmol) 1,2-phenylenedioxyacetic acid afforded, after flash chromatography (EtOAc), 1.08 g (87%) of the title compound as a colourless oil. MS m/e (%): 601 (M+H⁺, 100).

- 10 b) (S)-1-[[2-[2-[(S)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 1.08 g (1.8 mmol) (S)-1-[[2-[2-[(S)-2-Benzyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 717 mg (95%) of the title compound as a
15 white crystalline solid. MS m/e (%): 421 (M+H⁺, 100).

Example 74

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-naphthalen-1-yl]-acetyl]-pyrrolidine-2-carboxylic acid

- 20 a) 2-Diazo-1-(4-diazoacetyl-naphthalen-1-yl)-ethanone

To a stirred solution of 3.0 g (13.9 mmol) naphthalene-1,4-dicarboxylic acid and 4.4 ml (30.6 mmol) triethylamine in 200 ml THF at -15 °C was added dropwise 2.9 ml (30.6 mmol) ethyl chloroformate. After stirring for 15 min, 250 ml (approx. 0.3 M, approx. 75 mmol) of an ethereal solution of diazomethane
25 was added at 0 °C and stirring continued for 16 h at room temperature. The reaction mixture was then washed sequentially with saturated sodium bicarbonate solution, saturated ammonium chloride solution and finally with saturated brine, and the aqueous phases back-extracted with ether. The combined organic extracts were dried over magnesium sulphate and
30 concentrated *in vacuo*. Flash chromatography (gradient: 15-50% EtOAc/hexane) afforded 450 mg (12%) of the title compound as a yellow crystalline solid. MS m/e (%): 323 ([M+OAc]⁺, 100).

b) (4-Carboxymethyl-naphthalen-1-yl)-acetic acid

To a stirred solution of 450 mg (1.7 mmol) 2-diazo-1-(4-diazoacetyl-naphthalen-1-yl)-ethanone in 20 ml THF at -20 °C were added sequentially in the dark 1.5 ml water, 86 mg (375 mmol) silver benzoate (0.22 equiv.) and 687
5 ml (4.9 mmol) triethylamine (2.9 equiv.) and stirring continued for 2 h at room temperature. The reaction mixture was then diluted with 100 ml ether and extracted twice with saturated sodium bicarbonate solution. The combined aqueous phases were extracted three times with ether, then acidified with concentrated hydrochloric acid and extracted a further three times with ether.
10 The latter organic extracts were combined and dried over sodium sulphate and concentrated *in vacuo* to afford 330 mg (80%) of the title compound as a yellow crystalline solid. MS m/e (%): 244 (M^+ , 100), 199 ($M-CO_2H$, 80).

c) (R)-1-[[4-[2-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-naphthalen-1-yl]-acetyl]-pyrrolidine-2-carboxylic acid benzyl ester

15 Using General Procedure A with 200 mg (0.82 mmol) D-Proline benzyl ester hydrochloride and 100 mg (0.41 mmol) (4-carboxymethyl-naphthalen-1-yl)-acetic acid afforded, after flash chromatography (EtOAc), 188 mg (75%) of the title compound as a colourless oil. MS m/e (%): 636 ($M+NH_4^+$, 100), 619 ($M+H^+$, 75).

20 d) (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-naphthalen-1-yl]-acetyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 130 mg (0.21 mmol) (R)-1-[[4-[2-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-naphthalen-1-yl]-acetyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 79 mg (86%) of the title
25 compound as a white crystalline solid. MS m/e (%): 437 ($[M-H]^+$, 100).

Example 75

(R)-1-[[6-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-pyridin-2-yl]-acetyl]-pyrrolidine-2-carboxylic acid

30 a) (6-Cyanomethyl-pyridin-2-yl)-acetonitrile

To a stirred solution of 3.47 g (13.1 mmol) 2,6-bis(bromomethyl)pyridine in 50 ml dichloromethane was added dropwise a solution of 4.1 g (26.2 mmol) tetraethylammonium cyanide in 20 ml dichloromethane and the reaction

mixture then heated at 45 °C for 72 h. The reaction mixture was then cooled to room temperature and filtered. The filtrate was concentrated *in vacuo*, resuspended in ethyl acetate, filtered once again, and the second filtrate concentrated *in vacuo*. Flash chromatography (33% EtOAc/hexane) afforded
5 1.77 g (84%) of the title compound as a white crystalline solid. MS m/e (%): 157 (M⁺, 100), 130 ([M-HCN]⁺, 95), 90 (55).

b) (6-Carboxymethyl-pyridin-2-yl)-acetic acid

A solution of 3.0 g (19.1 mmol) (6-cyanomethyl-pyridin-2-yl)-acetonitrile in 30 ml concentrated hydrochloric acid was heated at 100 °C for 24 h. The reaction
10 mixture was then cooled to room temperature and concentrated *in vacuo*. The residue was redissolved in water, activated charcoal added, and the mixture heated at 50 °C for 30 min. After removal of the charcoal by filtration, the filtrate was concentrated *in vacuo* and the residue recrystallised from water at 4 °C to afford 2.48 g (100%) of the title compound as an off-white crystalline
15 solid. ¹H NMR d (250 MHz, D₂O) 8.42 (1H, t, *J* = 8 Hz), 7.80 (2H, d, *J* = 8 Hz).

c) (R)-1-[[6-[2-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-pyridin-2-yl]-acetyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 9.7 g (40 mmol) D-Proline benzyl ester hydrochloride and 3.9 g (20 mmol) (6-carboxymethyl-pyridin-2-yl)-acetic acid
20 afforded, after flash chromatography (gradient: 0-10% MeOH/EtOAc), 890 mg (8%) of the title compound as a yellow oil. MS m/e (%): 570 (M+H⁺, 100).

d) (R)-1-[[6-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-pyridin-2-yl]-acetyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 890 mg (1.56 mmol) (R)-1-[[6-[2-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-pyridin-2-yl]-acetyl]-
25 pyrrolidine-2-carboxylic acid benzyl ester afforded 410 mg (67%) of the title compound as a yellow crystalline solid. MS m/e (%): 390 (M+H⁺, 100).

Example 76

30 (R)-1-[[5-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-thiophen-2-yl]-acetyl]-pyrrolidine-2-carboxylic acid

a) 2,5-Bis-chloromethyl-thiophene

To 63.4 ml of a 37% aqueous solution of formaldehyde was added dropwise with ice-cooling 15.3 ml of concentrated hydrochloric acid and then 20 ml (0.25 mol) thiophene was added dropwise and stirring continued for 90 min at room temperature. The reaction mixture was then extracted with ether and the organic phases washed sequentially with water, saturated sodium bicarbonate solution and finally with saturated brine. The aqueous phases were back-extracted with ether and the combined organic extracts were dried over sodium sulphate and concentrated *in vacuo* to afford 38.8 g (86%) of the title compound RO-64-4463/000 as an amber-coloured oil. MS m/e (%): 184 (M^+ , 8), 182 (M^+ , 15), 180 (M^+ , 24), 147 ($[M-Cl]^+$, 44), 145 ($[M-Cl]^+$, 100), 110 (64).

b) (5-Cyanomethyl-thiophen-2-yl)-acetonitrile

To a stirred solution of 15 g (82.8 mmol) 2,5-bis-chloromethyl-thiophene in 500 ml dichloromethane was added dropwise a solution of 28.5 g (182 mmol) tetraethylammonium cyanide in 100 ml dichloromethane and the reaction mixture then heated at 45 °C for 24 h. The reaction mixture was then cooled to room temperature and washed twice with water. The organic phase was dried over sodium sulphate and concentrated *in vacuo*. Flash chromatography (20% EtOAc/hexane) afforded 4.88 g (36%) of the title compound as a yellow oil. MS m/e (%): 162 (M^+ , 44), 122 ($[M-CH_2CN]^+$, 100).

c) (5-Carboxymethyl-thiophen-2-yl)-acetic acid

To a stirred solution of 1.2 g (7.4 mmol) (5-cyanomethyl-thiophen-2-yl)-acetonitrile in 5 ml ethanol and 5 ml water was added 1.74 g (31.8 mmol) potassium hydroxide and the reaction mixture heated at 100 °C for 1 h. The reaction mixture was then cooled to room temperature and concentrated *in vacuo*. The residue was acidified with hydrochloric acid and extracted with three times with ether. The combined organic phases were dried over magnesium sulphate and concentrated *in vacuo* to afford 1.3 g (88%) of the title compound as a brown crystalline solid. MS m/e (%): 221 ($[M+Na-H]^+$, 50), 199 ($[M-H]^+$, 45), 155 ($[M-CO_2H]^+$, 100).

d) (R)-1-[[5-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-thiophen-2-yl]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

Using General Procedure A with 2.23 g (13 mmol) D-Proline benzyl ester hydrochloride and 1.3 g (6.5 mmol) (5-carboxymethyl-thiophen-2-yl)-acetic acid

afforded, after flash chromatography (gradient: 0-10% MeOH/EtOAc), 2.6 g (79%) of the title compound as a yellow oil. MS m/e (%): 524 (M+NH₄⁺, 90), 507 (M+H⁺, 10), 451 ([M+H-C₄H₉]⁺, 30), 395 ([M+H-2C₄H₉]⁺, 100).

e) (R)-1-[[5-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-thiophen-2-yl]-acetyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 600 mg (1.18 mmol) (R)-1-[[5-[2-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-2-oxo-ethyl]-thiophen-2-yl]-acetyl]-pyrrolidine-2-carboxylic acid tert-butyl ester afforded 100 mg (21%) of the title compound as a beige crystalline solid. MS m/e (%): 395 (M+H⁺, 100).

10

Example 77

(R)-1-[(2R,5S)-6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

a) (3R,6S)-3,6-Dimethoxy-cyclohexene

- 15 Lit. *J. Org. Chem.* **1988**, *53*, 5695. To a stirred solution of 0.70 g (3.1 mmol) palladium acetate, 16.9 g (156 mmol) benzoquinone and 0.4 ml (6.2 mmol) methanesulphonic acid in 200 ml methanol at room temperature was added via syringe pump over 4 h a solution of 5.95 ml (62 mmol) 1,3-cyclohexadiene in 5 ml methanol, and stirring continued for an additional 16 h. The reaction
- 20 mixture was extracted three times with ether and the combined organic extracts washed successively with water, 2 M sodium hydroxide solution and saturated brine. The organic phases were dried over sodium sulphate and concentrated *in vacuo*. Kugelrohr distillation (6 mbar, oven temp 120 °C) afforded 5.42 g (61%) of the title compound as a colourless oil. ¹H NMR d (250
- 25 MHz, CDCl₃) 5.92 (2H, s), 3.70 (2H, t, *J* = 5 Hz), 3.37 (6H, s), 1.90-1.65 (4H, m).

b) (2R,5S)-2,5-Dimethoxy-hexanedioic acid

- Lit. *J. Am. Chem. Soc.* **1983**, *105*, 5688. To a stirred solution of 5.0 g (35.2 mmol) (3R,6S)-3,6-dimethoxy-cyclohexene in 120 ml acetone and 120 ml water
- 30 were added 37.6 g (176 mmol) sodium periodate and 50 mg (0.24 mmol) ruthenium (III) chloride and stirring continued for 16 h at room temperature. 5 ml isopropanol was added and stirring continued for 30 min, then the reaction mixture filtered and the filtrate concentrated *in vacuo* to half-volume. 5 g

sodium bicarbonate was then added portionwise, and the mixture extracted three times with ethyl acetate. The combined organic phases were washed with saturated brine, dried over sodium sulphate, and concentrated *in vacuo* to afford 1.0 g (14%) of the title compound as an orange oil. MS m/e (%): 205

5 ([M-H]⁺, 100).

c) (R)-1-[(2R,5S)-6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 1.51 g (6.2 mmol) D-Proline benzyl ester hydrochloride and 643 mg (3.1 mmol) (2R,5S)-2,5-Dimethoxy-hexanedioic acid
10 afforded, after flash chromatography (gradient: 10-100% EtOAc/hexane then 10% MeOH/EtOAc), 643 mg (36%) of the title compound as a yellow oil. MS m/e (%): 581 (M+ H⁺, 100).

d) (R)-1-[(2R,5S)-6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

15 Using General Procedure B with 640 mg (1.10 mmol) (R)-1-[(2R,5S)-6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 440 mg (100%) of the title compound as a yellow solid. MS m/e (%): 399 [M-H]⁺, 100).

20

Example 78

(R)-1-[(2S,5S)- or -(2R,5R)-6-[(R)-2-Hydroxymethyl-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

a) Acetic acid (1RS,4RS)-4-acetoxy-cyclohex-2-enyl ester

Lit. *J. Org. Chem.* 1984, 49, 4619. To a stirred solution of 2.8 g (12.5 mmol)
25 palladium acetate, 27.2 g (267 mmol) lithium acetate, and 7.64 g (70.7 mmol) benzoquinone in 200 ml acetic acid at room temperature were added 26.1 g (300 mmol) manganese dioxide and a solution of 23.8 ml (250 mmol) 1,3-cyclohexadiene in 400 ml pentane, and stirring continued for 16 h. The two-phase reaction mixture was separated and the acetic acid phase extracted
30 twice with pentane. The combined organic extracts were washed successively with saturated brine, water, and 2 M sodium hydroxide solution, then the organic phases were dried over sodium sulphate and concentrated *in vacuo*. Recrystallisation from pentane afforded 16.1 g (33%) of the title compound as

an off-white crystalline solid. MS m/e (%): 138 ($[M-AcOH]^+$, 8), 96 ($[M-AcOH-COCH_3]^+$, 100), 43 (40).

b) (1RS,4RS)-Cyclohex-2-ene-1,4-diol

Lit. *J. Org. Chem.* 1984, 49, 4619. To a stirred solution of 19.8 g (99.8 mmol)
5 acetic acid (1RS,4RS)-4-acetoxy-cyclohex-2-enyl ester in 500 ml methanol was
added 120 ml 2 M sodium hydroxide solution and the reaction mixture
refluxed for 15 min. After cooling to room temperature, the reaction mixture
was concentrated *in vacuo* to 100 ml, and then saturated with sodium
hydroxide pellets. The mixture was extracted repeatedly with ethyl acetate
10 and the combined organic phases dried over sodium sulphate and concentrated
in vacuo to afford 10.2 g (90%) of the title compound as an off-white crystalline
solid. MS m/e (%): 113 ($[M-H]^+$, 13), 96 ($[M-H_2O]^+$, 36), 70 ($[M-H_2C=CHOH]^+$,
100).

c) (3RS,6RS)-3,6-Dimethoxy-cyclohexene

15 Lit. *J. Org. Chem.* 1988, 53, 5695. To 14.3 g (328 mmol) sodium hydride (55%
dispersion in oil) was added dropwise at 0 °C with stirring a solution of 10.1 g
(88.5 mmol) (1RS,4RS)-cyclohex-2-ene-1,4-diol in 100 ml THF. 34 ml (546
mmol) methyl iodide was then added and stirring continued for an additional
48 h at room temperature. The reaction was quenched with saturated
20 ammonium chloride solution and extracted with ether. The combined organic
extracts were washed successively with saturated ammonium chloride
solution and saturated brine, dried over sodium sulphate, and concentrated *in*
vacuo to afford 11.5 g (91%) of the title compound as a yellow oil. 1H NMR d
(250 MHz, $CDCl_3$) 5.90 (2H, s), 3.82 (2H, br t), 3.37 (6H, s), 2.10 (2H, m), 1.51
25 (2H, m).

d) (2RS,5RS)-2,5-Dimethoxy-hexanedioic acid

To a stirred solution of 6.0 g (42.2 mmol) (3RS,6RS)-3,6-dimethoxy-
cyclohexene in 140 ml acetone and 140 ml water were added 50.5 g (236 mmol)
sodium periodate and 67 mg (0.32 mmol) ruthenium (III) chloride and stirring
30 continued for 16 h at room temperature. 10 ml isopropanol was added and
stirring continued for 30 min, then the reaction mixture filtered and the
filtrate concentrated *in vacuo* to half-volume. 5 g sodium bicarbonate was then
added portionwise, and the mixture extracted three times with ether. The

aqueous phase was acidified with 25% hydrochloric acid and extracted repeatedly with ethyl acetate. The combined organic phases were dried over sodium sulphate, and concentrated *in vacuo* to afford 1.32 g (15%) of the title compound RO-64-5650/000 as an orange oil. Continuous extraction of the aqueous phase over 48 h afforded another 608 mg (7%) of product. MS m/e (%): 161 ([M-CO₂H]⁺, 20), 113 (58), 101 (69), 85 (50), 71 (100).

e) (R)-1-[(2S,5S)- or -(2R,5R)-6-[(R)-2-Benzylloxymethyl-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl and (R)-1-[(2R,5R)- or [(2S,5S)-6-[(R)-2-Benzylloxymethyl-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl

Using General Procedure A with 1.67 g (6.9 mmol) D-Proline benzyl ester hydrochloride and 250 mg (1.2 mmol) (2RS,5RS)-2,5-dimethoxy-hexanedioic acid afforded, after flash chromatography (gradient: 50-100% EtOAc/hexane then 10% MeOH/EtOAc), 95 mg (13%) of the title compound (mixture of 2 diastereomers) as a brown oil, and 172 mg (24%) of the title compound (single separated diastereomer) as a yellow oil. MS m/e (%): 581 (M+ H⁺, 100).

f) (R)-1-[(2S,5S)- or -(2R,5R)-6-[(R)-2-Hydroxymethyl-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 160 mg (0.28 mmol) (R)-1-[(2R,5R)- or [(2S,5S)-6-[(R)-2-benzylloxymethyl-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl afforded 90 mg (82%) of the title compound as a white foam. MS m/e (%): 399 ([M-H]⁻, 100), 355 ([M-H-CO₂]⁻, 61).

25

Example 79

(R)-1-[(2R,5R)- or -(2S,5S)-6-[(R)-2-Hydroxymethyl-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 95 mg (0.28 mmol) (R)-1-[(2S,5S)- or -(2R,5R)-6-[(R)-2-benzylloxymethyl-pyrrolidin-1-yl]-2,5-dimethoxy-6-oxo-

30 hexanoyl]-pyrrolidine-2-carboxylic acid benzyl afforded 70 mg (100%) of the title compound as a white foam. MS m/e (%): 399 ([M-H]⁻, 100), 355 ([M-H-CO₂]⁻, 20).

Example 80

(R)-1-[2,5-Dibenzyl-6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

- 5 a) Mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-dibenzyl-hex-3-enedioic acid diethyl ester

To a stirred solution of 688 mg (3.44 mmol) *trans*-2-butene-1,4-dicarboxylic acid diethyl ester in 35 ml THF was added 1.46 g (34.4 mmol) anhydrous lithium chloride and the resulting suspension cooled to -78 °C. 3.44 ml (6.88 mmol) of a 2 M solution of LDA in THF was added dropwise and stirring continued for 45 min. 0.82 ml (6.9 mmol) benzyl bromide was then added and stirring continued for 1 h at -78 °C and 10 min at 0 °C. The reaction was quenched at this temperature by addition of saturated ammonium chloride solution and the mixture extracted three times with ether. The combined organic phases were washed with saturated brine, dried over sodium sulphate, and concentrated *in vacuo*. Flash chromatography (5% EtOAc in hexane) afforded 927 mg (71%) of the title compound as a yellow oil. MS *m/e* (%): 398 ([M+NH₄]⁺, 100).

b) Mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-dibenzyl-hex-3-enedioic

- 20 To a stirred solution of 200 mg (0.53 mmol) mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-dibenzyl-hex-3-enedioic acid diethyl ester in 5 ml THF was added 4.3 ml (4.3 mmol) of 1 M sodium hydroxide solution. After stirring for 68 h at room temperature, the reaction mixture was acidified to pH 3 by addition of 1 M hydrochloric acid and extracted three times with ethyl acetate.
- 25 The combined organic phases were washed successively with water and with saturated brine, dried over sodium sulphate, and concentrated *in vacuo*. Flash chromatography (50% EtOAc in hexane containing 1% AcOH) afforded 127 mg (74%) of the title compound as a white crystalline solid. MS *m/e* (%): 342 ([M+NH₄]⁺, 100).

- 30 c) (E)-(R)-1-[2,5-Dibenzyl-6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 3 diastereomers)

Using General Procedure A with 1.04 g (4.30 mmol) D-Proline benzyl ester hydrochloride and 700 mg (2.16 mmol) mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-dibenzyl-hex-3-enedioic acid diethyl ester afforded, after flash chromatography (5% MeOH in EtOAc), 926 mg (61%) of the title compound as a yellow oil. MS m/e (%): 716 ($[M+NH_4]^+$, 100), 699 ($[M+H]^+$, 85).

d) (R)-1-[2,5-Dibenzyl-6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

Using General Procedure B with 926 mg (1.33 mmol) (E)-(R)-1-[2,5-dibenzyl-6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 3 diastereomers) afforded 610 mg (89%) of the title compound as a white crystalline solid. MS m/e (%): 519 ($[M-H]^-$, 100).

Example 81

15 (R)-1-[2,5-Dibutyl-6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (1 out of 3 possible diastereomers)

a) Mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-dibutyl-hex-3-enedioic acid diethyl ester

To a stirred solution of 2.0 g (9.99 mmol) *trans*-2-butene-1,4-dicarboxylic acid diethyl ester in 80 ml THF was added 2.54 g (59.9 mmol) anhydrous lithium chloride and the resulting suspension cooled to -78 °C. 10.0 ml (20.0 mmol) of a 2 M solution of LDA in THF was added dropwise and stirring continued for 45 min. 2.2 ml (20.4 mmol) butyl bromide was then added and stirring continued for 15 min at -78 °C, then 30 min at 0 °C, and then 4 h at room temperature.

25 The reaction was quenched by addition of saturated ammonium chloride solution and the mixture extracted three times with ether. The combined organic phases were washed successively with saturated ammonium chloride solution, water, and saturated brine, dried over sodium sulphate, and concentrated *in vacuo*. Successive flash chromatography (10% EtOAc in hexane for first column; then gradient: 10-100% toluene in cyclohexane for second column) afforded 509 mg (16%) of the title compound as a yellow oil. MS m/e (%): 330 ($[M+NH_4]^+$, 100).

b) Mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-dibutyl-hex-3-enedioic acid

To a stirred solution of 435 mg (1.39 mmol) mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-dibutyl-hex-3-enedioic acid diethyl ester in 10 ml THF was added 26 ml (26 mmol) of 1 M sodium hydroxide solution. After stirring for 72 h at room temperature, the reaction mixture was acidified to pH 3 by addition of 1 M hydrochloric acid and extracted three times with ethyl acetate. The combined organic phases were washed successively with water and with saturated brine, dried over sodium sulphate, and concentrated *in vacuo* to afford 346 mg (97%) of the title compound as a white crystalline solid. MS m/e (%): 255 ([M-H]⁻, 60), 211 ([M-H-CO₂]⁻, 100).

c) (E)-(R)-1-[2,5-Dibutyl-6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 2 out of 3 possible diastereomers)

and (E)-(R)-1-[2,5-Dibutyl-6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid benzyl ester (principally 1 out of 3 possible diastereomers)

Using General Procedure A with 653 mg (2.70 mmol) D-Proline benzyl ester hydrochloride and 346 mg (1.35 mmol) mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-dibutyl-hex-3-enedioic acid afforded, after flash chromatography (gradient: 33-50% EtOAc in hexane), 148 mg (17%) of the title compound RO-64-3271/000 (mixture of 2 diastereomers) as a yellow oil and 116 mg (14%) of the title compound (single diastereomer) as a yellow oil. MS m/e (%): 648 (M+NH₄⁺, 90), 631 (M+H⁺, 100).

d) (R)-1-[2,5-Dibutyl-6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (1 out of 3 possible diastereomers)

Using General Procedure B with 140 mg (0.22 mmol) (E)-(R)-1-[2,5-dibutyl-6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid benzyl ester (principally 1 out of 3 possible diastereomers) afforded 81 mg (81%) of the title compound RO-64-3273/000 (single diastereomer) as a colourless oil. MS m/e (%): 451 ([M-H]⁻, 100).

Example 82

(R)-1-[2,5-Dibutyl-6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (mixture of 2 out of 3 possible diastereomers)

- Using General Procedure B with 110 mg (0.22 mmol) (E)-(R)-1-[2,5-dibutyl-6-
5 [(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 2 out of 3 possible diastereomers) afforded 69 mg (88%) of the title compound (2 diastereomers) as a colourless oil. MS m/e (%): 451 ([M-H]⁺, 100).

10

Example 83

(R)-1-[2,5-Diisopropyl-6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

a) Mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-diisopropyl-hex-3-enedioic acid diethyl ester

- 15 To a stirred solution of 5.0 g (25.0 mmol) *trans*-2-butene-1,4-dicarboxylic acid diethyl ester in 120 ml THF was added 6.35 g (150 mmol) anhydrous lithium chloride and the resulting suspension cooled to -78 °C. 25.0 ml (50.0 mmol) of a 2 M solution of LDA in THF was added dropwise and stirring continued for 45 min. 4.7 ml (50 mmol) isopropyl bromide was then added and stirring
20 continued for 15 min at -78 °C, then 4 h at 0 °C, and then 48 h at room temperature. The reaction was quenched by addition of saturated ammonium chloride solution and the mixture extracted three times with ether. The combined organic phases were washed successively with saturated ammonium chloride solution, water, and saturated brine, dried over sodium sulphate, and
25 concentrated *in vacuo*. Successive flash chromatography (20% EtOAc in hexane for first column; 5% EtOAc in hexane for second column; 10% EtOAc in hexane for third column) afforded 259 mg (4%) of the title compound as a yellow oil. MS m/e (%): 302 ([M+NH₄]⁺, 100).

b) Mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-diisopropyl-hex-3-enedioic

- 30 To a stirred solution of 108 mg (0.38 mmol) mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-diisopropyl-hex-3-enedioic acid diethyl ester in 5 ml THF was added 10 ml (10 mmol) of 1 M sodium hydroxide solution. After stirring for 96 h at room temperature, the reaction mixture was acidified to pH 3 by addition

of 1 M hydrochloric acid and extracted three times with ethyl acetate. The combined organic phases were washed successively with water and with saturated brine, dried over sodium sulphate, and concentrated *in vacuo* to afford 70 mg (81%) of the title compound as a yellow oil. MS *m/e* (%): 227 ([M-H]⁺, 100).

c) (E)-(R)-1-[2,5-Diisopropyl-6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of the 3 diastereomers)

Using General Procedure A with 114 mg (0.47 mmol) D-Proline benzyl ester hydrochloride and 54 mg (0.24 mmol) mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-diisopropyl-hex-3-enedioic afforded, after flash chromatography (gradient: 33-50% EtOAc in hexane), 15 mg (11%) of the title compound as a colourless oil. MS *m/e* (%): 620 (M+NH₄⁺, 100).

d) (R)-1-[2,5-Diisopropyl-6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

Using General Procedure B with 41 mg (0.07 mmol) (E)-(R)-1-[2,5-diisopropyl-6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of the 3 diastereomers) afforded 28 mg (100%) of the title compound as a colourless oil. MS *m/e* (%): 423 ([M-H]⁺, 100).

Example 84

(R)-1-[5-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-7-methoxy-2-(2-methoxy-ethyl)-heptanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

a) Mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-bis-(2-methoxy-ethyl)-hex-3-enedioic acid diethyl ester

To a stirred solution of 5.0 g (25.0 mmol) *trans*-2-butene-1,4-dicarboxylic acid diethyl ester in 125 ml THF was added 6.35 g (150 mmol) anhydrous lithium chloride and the resulting suspension cooled to -78 °C. 25.0 ml (50.0 mmol) of a 2 M solution of LDA in THF was added dropwise and stirring continued for 45 min. 7.4 ml (78.7 mmol) 2-methoxyethyl bromide was then added and stirring continued for 15 min at -78 °C, then 1 h at 0 °C, and then 2 h at room temperature. The reaction was quenched by addition of saturated ammonium chloride solution and the mixture extracted three times with ether. The

combined organic phases were washed successively with saturated ammonium chloride solution, water, and saturated brine, dried over sodium sulphate, and concentrated *in vacuo*. Flash chromatography (50% toluene in EtOAc) afforded 1.29 g (16%) of the title compound as a yellow oil. MS *m/e* (%): 334 ($[M+NH_4]^+$, 100).

b) Mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-bis-(2-methoxy-ethyl)-hex-3-enedioic acid

To a stirred solution of 1.29 g (4.08 mmol) mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-bis-(2-methoxy-ethyl)-hex-3-enedioic acid diethyl ester in 10 ml THF was added 33 ml (33 mmol) of 1 M sodium hydroxide solution. After stirring for 18 h at room temperature, the reaction mixture was acidified to pH 3 by addition of 1 M hydrochloric acid and extracted three times with ethyl acetate. The combined organic phases were washed successively with water and with saturated brine, dried over sodium sulphate, and concentrated *in vacuo* to afford 894 mg (84%) of the title compound as a yellow oil. MS *m/e* (%): 259 ($[M-H]^+$, 100).

c) (E)-(R)-1-[5-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-7-methoxy-2-(2-methoxy-ethyl)-heptanoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 3 diastereomers)

Using General Procedure A with 1.86 g (7.69 mmol) D-Proline benzyl ester hydrochloride and 894 mg (3.84 mmol) mixture of (E)-(2R,5S)- and -(2RS,5SR)-2,5-bis-(2-methoxy-ethyl)-hex-3-enedioic acid afforded, after successive flash chromatography (gradient 50%-100% EtOAc in hexane then 10% MeOH in EtOAc for first column; 20% toluene in EtOAc for second column; 20% toluene in EtOAc for third column; gradient: 25-20% toluene in EtOAc for fourth column), 146 mg (7%) of the title compound as a light yellow oil. MS *m/e* (%): 652 ($M+NH_4^+$, 50), 635 ($M+H^+$, 100).

d) (R)-1-[5-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-7-methoxy-2-(2-methoxy-ethyl)-heptanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

Using General Procedure B with 146 mg (0.23 mmol) (E)-(R)-1-[5-[(R)-2-benzylloxycarbonyl-pyrrolidine-1-carbonyl]-7-methoxy-2-(2-methoxy-ethyl)-heptanoyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of 3

diastereomers) afforded 92 mg (88%) of the title compound as a colourless oil.
MS m/e (%): 455 ([M-H]⁺, 100).

Example 85

5 (R)-1-[2-[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-1-methyl-2-oxo-ethyl]-phenyl]-propionyl]-pyrrolidine-2-carboxylic acid (mixture of the 3 diastereomers)

a) (4-Benzyloxycarbonylmethyl-phenyl)-acetic acid benzyl ester

To a suspension of 10.0 g (51.5 mmol) benzene-1,4-diacetic acid, 0.94 g (7.73 mmol) 4-dimethylaminopyridine and 5.33 ml (51.5 mmol) benzyl alcohol in
10 150 ml dichloromethane at 0 °C was added 11.85 g (61.8 mmol) *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide and stirring continued at 0 °C for 2 h and then at room temperature for 16 h. The reaction mixture was then washed sequentially with 1 M hydrochloric acid, saturated sodium bicarbonate solution and finally with saturated brine, and the aqueous phases back-
15 extracted with dichloromethane. The combined organic extracts were dried over sodium sulphate and concentrated *in vacuo*. Flash chromatography (50% EtOAc in hexane) then afforded the title compound as a light yellow oil which crystallised on standing. MS m/e (%): 374 (M⁺, 10), 283 ([M-Bn]⁺, 16), 239 ([M-Bn-CO₂]⁺, 18), 91 (Bn⁺, 100).

20 b) Mixture of (R)-2-[4-[(S)- and (RS)-2-[4-[(RS)-1-benzyloxycarbonyl-ethyl]-phenyl]-propionic acid benzyl ester

To a stirred solution of 2.0 g (5.3 mmol) (4-benzyloxycarbonylmethyl-phenyl)-acetic acid benzyl ester in 80 ml THF was added 1.35 g (32 mmol) anhydrous lithium chloride and the resulting suspension cooled to -78 °C. 10.7 ml (21.4
25 mmol) of a 2 M solution of LDA in THF was added dropwise and stirring continued for 45 min. 1.33 ml (21.3 mmol) methyl iodide was then added and stirring continued for 15 min at -78 °C, then 20 min at 0 °C. The reaction was quenched at this temperature by addition of saturated ammonium chloride solution and the mixture extracted three times with ether. The combined
30 organic phases were dried over sodium sulphate, and concentrated *in vacuo*. Flash chromatography (EtOAc) afforded 2.1 g (100%) of the title compound as a brown oil. MS m/e (%): 420 ([M+NH₄]⁺, 100).

c) Mixture of (R)-2-[4-[(S)- and (RS)-2-[4-[(RS)-1-carboxy-ethyl)-phenyl]-propionyl]-phenyl]-propionic acid

A solution of 230 mg (0.57 mmol) mixture of (R)-2-[4-[(S)- and (RS)-2-[4-[(RS)-1-benzyloxycarbonyl-ethyl]-phenyl]-propionyl]-phenyl]-propionic acid benzyl ester in 20 ml

- 5 ethanol was stirred with 5 wt% of 10% palladium on charcoal under 1 atm of hydrogen for 16 h at room temperature. After filtration to remove the catalyst, the reaction mixture was concentrated *in vacuo* to afford 70 mg (55%) of the title compound as a white crystalline solid. MS m/e (%): 222 (M^+ , 23), 177 ($[M-CO_2H]^+$, 100), 131.

- 10 d) (R)-1-[2-[4-[2-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-1-methyl-2-oxo-ethyl]-phenyl]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of the 3 diastereomers)

Using General Procedure A with 370 mg (1.53 mmol) D-Proline benzyl ester hydrochloride and 170 mg (0.77 mmol) mixture of (R)-2-[4-[(S)- and (RS)-2-[4-

- 15 [(RS)-1-carboxy-ethyl)-phenyl]-propionyl]-phenyl]-propionyl]-pyrrolidine-2-carboxylic acid afforded, after flash chromatography (EtOAc), 170 mg (38%) of the title compound as a colourless oil. MS m/e (%): 614 ($M+NH_4^+$, 100), 597 ($M+H^+$, 60).

e) (R)-1-[2-[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-1-methyl-2-oxo-ethyl]-phenyl]-propionyl]-pyrrolidine-2-carboxylic acid (mixture of the 3 diastereomers)

- 20 Using General Procedure B with 170 mg (0.29 mmol) (R)-1-[2-[4-[2-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-1-methyl-2-oxo-ethyl]-phenyl]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester (mixture of the 3 diastereomers) afforded 50 mg (42%) of the title compound as a white foam. MS m/e (%): 415 ($[M-H]^+$, 100).

25

Example 86

(2E,4E)-(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexa-2,4-dienyl]-pyrrolidine-2-carboxylic acid

- 30 a) (2E,4E)-(R)-1-[6-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexa-2,4-dienyl]-pyrrolidine-2-carboxylic acid benzyl ester

To a stirred suspension of 0.91 g (5.32 mmol) 2,5-dimethyl-hex-2,4-dien-1,6-dioic acid in 80 ml dichloromethane containing two drops of pyridine was added dropwise at room temperature 0.77 ml (10.6 mmol) thionyl chloride and

the reaction mixture was then heated at 50 °C for 2 h. The resulting solution was cooled to 0 °C and added dropwise to a solution of 2.57 g (10.6 mmol) D-Proline benzyl ester hydrochloride and 3.0 ml (21.5 mmol) triethylamine in 50 ml dichloromethane at 0 °C. Stirring was continued for 2 h at 0 °C and then 24 h at room temperature. The reaction mixture was then washed sequentially with 1 M hydrochloric acid and with water, and the aqueous phases back-extracted with dichloromethane. The combined organic extracts were dried over sodium sulphate and concentrated *in vacuo*. Flash chromatography (EtOAc) afforded 2.79 g (96%) of the title compound as a colourless oil. MS m/e (%): 562 (M+NH₄⁺, 100), 545 (M+H⁺, 5).

b) (2E,4E)-(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexa-2,4-dienoyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 1.14 g (2.09 mmol) (2E,4E)-(R)-1-[6-[(R)-2-Benzyloxycarbonyl-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexa-2,4-dienoyl]-pyrrolidine-2-carboxylic acid benzyl ester and with dioxane as solvent afforded 127 mg (16%) of the title compound as a white solid. MS m/e (%): 365 (M+H⁺, 100).

Example 87

(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

A solution of 593 mg (1.09 mmol) (2E,4E)-(R)-1-[6-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexa-2,4-dienoyl]-pyrrolidine-2-carboxylic acid benzyl ester in 15 ml dioxane was stirred with 25 mg (0.11 mmol) platinum (IV) oxide under 1 atm of hydrogen for 72 h at room temperature. After filtration to remove the catalyst, the reaction mixture was concentrated *in vacuo* and azeotroped three times with chloroform on a rotary evaporator to remove last traces of dioxane, then triturated in ether to afford 400 mg (100%) of the title compound as a white foam. MS m/e (%): 369 (M+H⁺, 100).

30

Example 88

Mixture of (R)-1-[(3R,4R)- and -[(3S,4S)-3,4-dihydroxy-6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[6-[(R)-2-Benzylloxycarbonyl-pyrrolidine-1-carbonyl]-pyridine-2-carbonyl]-pyrrolidine-2-carboxylic acid benzyl ester

Using General Procedure A with 5.0 g (20.6 mmol) D-Proline benzyl ester hydrochloride and 1.5 g (10.3 mmol) *trans*-3-hexenedioic acid afforded, after
5 flash chromatography (EtOAc), 3.15 g (59%) of the title compound as a light yellow oil. MS *m/e* (%): 536 ($M+NH_4^+$, 100), 519 ($M+H^+$, 80).

b) Mixture of (R)-1-[(3R,4R)- and -(3S,4S)-6-[(R)-2-benzylloxycarbonyl-pyrrolidin-1-yl]-3,4-dihydroxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl ester

10 To a stirred solution of 3.13 g (6.04 mmol) (R)-1-[6-[(R)-2-benzylloxycarbonyl-pyrrolidine-1-carbonyl]-pyridine-2-carbonyl]-pyrrolidine-2-carboxylic acid benzyl ester in 10 ml acetone and 10 ml water were added 980 mg (7.25 mmol) 4-methylmorpholine-4-oxide and 0.6 ml of a 2.5% solution of osmium tetroxide in *tert*-butanol and stirring continued for 72 h at room temperature. 50 ml of
15 38% sodium hydrogensulphite solution was then added at 0 °C and stirring continued for a further 15 min. The reaction mixture was then filtered and extracted three times with ethyl acetate. The combined organic extracts were washed sequentially with 1 M hydrochloric acid and saturated brine, dried over sodium sulphate, and concentrated *in vacuo* to afford 3.34 g (100%) of the
20 title compound as a colourless oil. MS *m/e* (%): 575 ($M+Na^+$, 35), 570 ($M+NH_4^+$, 55), 553 ($M+H^+$, 100).

c) Mixture of (R)-1-[(3R,4R)- and -(3S,4S)-3,4-dihydroxy-6-[(R)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

Using General Procedure B with 506 mg (0.92 mmol) mixture of (R)-1-
25 [(3R,4R)- and -(3S,4S)-6-[(R)-2-benzylloxycarbonyl-pyrrolidin-1-yl]-3,4-dihydroxy-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid benzyl ester afforded 341 mg (100%) of the title compound as a white solid. MS *m/e* (%): 395 ($M+NH_4^+$, 55), 373 ($M+H^+$, 100).

30

Example 89

(E)-(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid

a) (E)-(R)-1-[6-[(R)-2-*tert*-Butoxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid *tert*-butyl ester

Using General Procedure A with 1.5 g (8.76 mmol) D-Proline *tert*-butyl ester and 630 mg (4.37 mmol) *trans*-3-hexenedioic acid afforded, after flash

- 5 chromatography (10% EtOH in EtOAc), 1.59 g (77%) of the title compound as a white crystalline solid. MS *m/e* (%): 568 ($M+NH_4^+$, 35), 451 ($M+H^+$, 100), 395 ($[M+H-C_3H_7]^+$, 32), 339 ($[M+H-2C_3H_7]^+$, 40).

b) (E)-(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid

- 10 To a stirred solution of 600 mg (1.33 mmol) (E)-(R)-1-[6-[(R)-2-*tert*-butoxycarbonyl-pyrrolidin-1-yl]-6-oxo-hex-3-enoyl]-pyrrolidine-2-carboxylic acid *tert*-butyl ester in 15 ml dichloromethane at 0 °C was added dropwise 4.4 ml (57.8 mmol) trifluoroacetic acid and stirring continued for 16 h at room temperature. Concentration *in vacuo* and azeotroping three times with
- 15 chloroform on a rotary evaporator afforded 402 mg (90%) of the title compound as a white foam. MS *m/e* (%): 339 ($M+H^+$, 100).

Example 90

(R)-1-[3-[[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-aminol]-propionyl]-pyrrolidine-2-carboxylic acid

- 20

a) (R)-1-Acryloyl-pyrrolidine-2-carboxylic acid benzyl ester

To a stirred solution of 390 mg (1.6 mmol) D-Proline benzyl ester hydrochloride and 0.47 ml (3.4 mmol) triethylamine in 20 ml dichloromethane at 0 °C was added dropwise 0.2 ml (2.4 mmol) acryloyl chloride and stirring

- 25 continued for 24 h at room temperature. The reaction mixture was then washed sequentially with water, 1 M hydrochloric acid and once more with water, and the aqueous phases back-extracted with dichloromethane. The combined organic extracts were dried over sodium sulphate and concentrated *in vacuo* to afford 420 mg (100%) of the title compound as a colourless oil. MS
- 30 *m/e* (%): 259 (M^+ , 25), 124 (100), 91 (25), 70 (21).

b) (R)-1-[3-[[3-[(R)-2-Benzoyloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-aminol]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester

A solution of 400 mg (1.5 mmol) (R)-1-acryloyl-pyrrolidine-2-carboxylic acid benzyl ester and 63 ml (0.75 mmol) propylamine in 5 ml acetonitrile was stirred for 16 h at room temperature, then for 6 h at 45 °C, and finally for 16 h at 80 °C. Concentration *in vacuo* and flash chromatography (20% H₂O in acetone) afforded 84 mg (19%) of the title compound as a pale yellow oil. MS m/e (%): 578 (M+H⁺, 100).

c) (R)-1-[3-[[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid

A solution of 84 mg (0.15 mmol) (R)-1-[3-[[3-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-propyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester in 3 ml ethanol was stirred with 10 mg 10% Palladium on charcoal under 1 atm of hydrogen for 16 h at room temperature. After filtration to remove the catalyst, concentration *in vacuo* afforded 58 mg (100%) of the title compound as a white solid. MS m/e (%): 398 (M+H⁺, 100).

15

Example 91

(R)-1-[3-[[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-cyclopropylmethyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid

a) (R)-1-[3-[[3-[(R)-2-Benzyloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-cyclopropylmethyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester

A solution of 444 mg (1.71 mmol) (R)-1-acryloyl-pyrrolidine-2-carboxylic acid benzyl ester and 74 ml (0.85 mmol) cyclopropylmethylamine in 5 ml acetonitrile was stirred for 1 h at room temperature, then for 16 h at 80 °C. Concentration *in vacuo* and flash chromatography (gradient: 0-100% MeOH in EtOAc) afforded 220 mg (44%) of the title compound as a yellow oil. MS m/e (%): 590 (M+H⁺, 100).

b) (R)-1-[3-[[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-cyclopropylmethyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid

A solution of 220 mg (0.37 mmol) (R)-1-[3-[[3-[(R)-2-benzyloxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-cyclopropylmethyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid benzyl ester in 20 ml isopropanol was stirred with 10 mg 10% Palladium on charcoal under 1 atm of hydrogen for 16 h at room temperature. After filtration to remove the catalyst, concentration *in*

30

vacuo afforded 153 mg (100%) of the title compound as a yellow solid. MS *m/e* (%): 408 ($[M-H]^-$, 100).

Example 92

5 (R)-1-[3-[(3,4-Dimethoxy-benzyl)-[3-[(R)-2-carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-amino]-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

a) (R)-1-Acryloyl-pyrrolidine-2-carboxylic acid tert-butyl ester

To a stirred solution of 5.0 g (29.2 mmol) D-Proline *tert*-butyl ester and 4.5 ml (32.1 mmol) triethylamine in 180 ml dichloromethane at 0 °C was added

10 dropwise 3.6 ml (43.8 mmol) acryloyl chloride and stirring continued for 48 h at room temperature. The reaction mixture was then washed sequentially with water, saturated ammonium chloride solution, once more with water and then with saturated brine, and the aqueous phases back-extracted with dichloromethane. The combined organic extracts were dried over sodium
15 sulphate and concentrated *in vacuo* to afford 6.6 g (100%) of the title compound as a yellow oil. MS *m/e* (%): 243 ($M+NH_4^+$, 33), 226 ($M+H^+$, 100).

b) (R)-1-[3-[[3-[(R)-2-*tert*-Butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-(3,4-dimethoxy-benzyl)-amino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

20 A solution of 1.0 g (4.44 mmol) (R)-1-acryloyl-pyrrolidine-2-carboxylic acid *tert*-butyl ester and 0.33 ml (2.22 mmol) veratrylamine in 25 ml acetonitrile was stirred for 16 h at 80 °C. Concentration *in vacuo* and flash chromatography (gradient: 0-10% MeOH in EtOAc) afforded 150 mg (10%) of the title compound as a light brown oil. MS *m/e* (%): 618 ($M+H^+$, 100).

25 c) (R)-1-[3-[(3,4-Dimethoxy-benzyl)-[3-[(R)-2-carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-amino]-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

To a stirred solution of 150 mg (0.24 mmol) (R)-1-[3-[[3-[(R)-2-*tert*-butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-(3,4-dimethoxy-benzyl)-amino]-propionyl]-pyrrolidine-2-carboxylic acid *tert*-butyl ester in 5 ml

30 dichloromethane at 0 °C was added dropwise 1.0 ml trifluoroacetic acid and stirring continued for 16 h at room temperature. Concentration *in vacuo* and azeotroping three times with chloroform on a rotary evaporator afforded, after

trituration in ether, 130 mg (87%) of the title compound as a yellow crystalline solid. MS m/e (%): 506 (M+H⁺, 100).

Example 93

5 (R)-1-[3-[[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-(2-methoxy-ethyl)-aminol-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

a) (R)-1-[3-[[3-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-(2-methoxy-ethyl)-aminol-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

A solution of 1.0 g (4.44 mmol) (R)-1-acryloyl-pyrrolidine-2-carboxylic acid tert-butyl ester and 0.19 ml (2.22 mmol) 2-methoxyethylamine in 25 ml
10 acetonitrile was stirred for 16 h at 80 °C. Concentration *in vacuo* and flash chromatography (gradient: 0-10% MeOH in EtOAc) afforded 300 mg (23%) of the title compound as a light brown oil. MS m/e (%): 526 (M+H⁺, 100).

b) (R)-1-[3-[[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-(2-methoxy-ethyl)-aminol-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)
15

To a stirred solution of 150 mg (0.29 mmol) (R)-1-[3-[[3-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-(2-methoxy-ethyl)-aminol-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml
dichloromethane at 0 °C was added dropwise 1.0 ml trifluoroacetic acid and
20 stirring continued for 16 h at room temperature. Concentration *in vacuo* and azeotroping three times with chloroform on a rotary evaporator afforded, after resuspension in water and subsequent lyophilisation, 100 mg (67%) of the title compound as a yellow oil. MS m/e (%): 436 (M+Na⁺, 35), 414 (M+H⁺, 100).

25

Example 94

(R)-1-[3-[Benzyl-[3-[(R)-2-carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-aminol-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

a) (R)-1-[3-[Benzyl-[3-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-aminol-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

30 A solution of 1.0 g (4.44 mmol) (R)-1-acryloyl-pyrrolidine-2-carboxylic acid tert-butyl ester and 0.24 ml (2.22 mmol) benzylamine in 25 ml acetonitrile was stirred for 16 h at 80 °C. Concentration *in vacuo* and flash chromatography

(gradient: 0-10% MeOH in EtOAc) afforded 470 mg (34%) of the title compound as a yellow oil. MS m/e (%): 558 (M+H⁺, 100).

b) (R)-1-[3-[Benzyl-[3-[(R)-2-carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-amino]-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

- 5 To a stirred solution of 200 mg (0.36 mmol) (R)-1-[3-[benzyl-[3-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-amino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml dichloromethane at 0 °C was added dropwise 1.0 ml trifluoroacetic acid and stirring continued for 16 h at room temperature. Concentration *in vacuo* and azeotroping three times with
- 10 chloroform on a rotary evaporator afforded, after trituration in ether, 160 mg (80%) of the title compound as a yellow crystalline solid. MS m/e (%): 468 (M+Na⁺, 30), 446 (M+H⁺, 100).

Example 95

- 15 (R)-1-[3-[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propylamino]-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

a) (R)-1-[3-[3-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-(4-trifluoromethyl-benzyl)-amino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- 20 A solution of 1.0 g (4.44 mmol) (R)-1-acryloyl-pyrrolidine-2-carboxylic acid tert-butyl ester and 0.32 ml (2.22 mmol) *para*-trifluoromethylbenzylamine in 25 ml acetonitrile was stirred for 16 h at 80 °C. Concentration *in vacuo* and flash chromatography (gradient: 0-10% MeOH in EtOAc) afforded 480 mg (31%) of the title compound as a light brown oil. MS m/e (%): 626 (M+H⁺, 100).

- 25 b) (R)-1-[3-[3-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propylamino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester
and (R)-1-[3-[3-[(R)-2-tert-Butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-ethyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester

- A solution of 290 mg (0.46 mmol) (R)-1-[3-[3-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-(4-trifluoromethyl-benzyl)-amino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 50 ml ethanol was stirred with
- 30 30 mg 10% Palladium on charcoal under 1 atm of hydrogen for 72 h at room temperature. After filtration to remove the catalyst, concentration *in vacuo*

and flash chromatography (gradient: 0-100% MeOH in EtOAc containing 1% Et₃N) afforded 36 mg (17%) of the title compound (R)-1-[3-[3-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propylamino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester as a yellow oil. MS m/e (%): 468 (M+H⁺, 100).

- 5 Further flash chromatography of the less polar fractions (gradient: 0-20% MeOH in EtOAc containing 1% Et₃N) afforded 87 mg (38%) of the by-product (R)-1-[3-[[3-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-ethyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester as a yellow oil. MS m/e (%): 496 (M+H⁺, 100).

10 c) (R)-1-[3-[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propylamino]-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

- To a stirred solution of 33 mg (0.07 mmol) (R)-1-[3-[3-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propylamino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml dichloromethane at 0 °C was added
15 dropwise 1.0 ml trifluoroacetic acid and stirring continued for 16 h at room temperature. Concentration *in vacuo* and azeotroping three times with chloroform on a rotary evaporator afforded, after resuspension in water and subsequent lyophilisation, 19 mg (76%) of the title compound as a yellow glassy solid. MS m/e (%): 354 ([M-H]⁻, 100).

20

Example 96

(R)-1-[3-[Ethyl-[3-[(R)-2-carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-amino]-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

- To a stirred solution of 85 mg (0.17 mmol) (R)-1-[3-[3-[(R)-2-tert-butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-ethyl-amino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml dichloromethane at 0 °C was added dropwise 1.0 ml trifluoroacetic acid and stirring continued for 16 h
25 at room temperature. Concentration *in vacuo* and azeotroping three times with chloroform on a rotary evaporator afforded, after resuspension in water and lyophilisation, 63 mg (97%) of the title compound as a yellow crystalline
30 solid. MS m/e (%): 442 ([M+OAc]⁻, 35), 382 ([M-H]⁻, 100).

Example 97

(R)-1-[3-[[3-[(R)-2-Carboxy-pyrrolidin-1-yl]-3-oxo-propyl]-(4-trifluoromethyl-benzyl)-amino]-propionyl]-pyrrolidine-2-carboxylic acid trifluoroacetate (1:1)

- To a stirred solution of 200 mg (0.32 mmol) (R)-1-[3-[[3-[(R)-2-tert-
5 butoxycarbonyl-pyrrolidin-1-yl]-3-oxo-propyl]-(4-trifluoromethyl-benzyl)-
amino]-propionyl]-pyrrolidine-2-carboxylic acid tert-butyl ester in 5 ml
dichloromethane at 0 °C was added dropwise 1.0 ml trifluoroacetic acid and
stirring continued for 16 h at room temperature. Concentration *in vacuo* and
azeotroping three times with chloroform on a rotary evaporator afforded, after
10 trituration in ether, 150 mg (75%) of the title compound as a yellow crystalline
solid. MS m/e (%): 536 (M+Na⁺, 20), 514 (M+H⁺, 100).

Example 98

- Mixture of (R)-1-[6-[(S)- and (RS)-1-[6-[(RS)-2-carboxy-pyrrolidin-1-yl]-6-oxo-
15 hexanoyl]-pyrrolidine-2-carboxylic acid

a) 1H-Pyrrole-2-carboxylic acid benzyl ester

- Lit. *J. Org. Chem.* **1979**, *44*, 975. To a stirred solution of 5.0 g (45.0 mmol)
pyrrole-2-carboxylic acid and 31.3 ml (225 mmol) triethylamine in 100 ml
DMF was added dropwise 26 ml (225 mmol) benzyl bromide and stirring
20 continued for 72 h at room temperature. The reaction mixture was then
concentrated *in vacuo* and the residue resuspended in dichloromethane and
washed twice with saturated sodium bicarbonate solution and twice with
water, and the aqueous phases back-extracted with dichloromethane. The
combined organic extracts were dried over sodium sulphate and concentrated
25 *in vacuo*. Flash chromatography (25% EtOAc in hexane) afforded 8.23 g (90%)
of the title compound as a yellow oil. MS m/e (%): 201 (M⁺, 33), 94 ([M-OBn]⁺,
22), 91 (Bn⁺, 100).

b) 1-[6-(2-Benzyloxycarbonyl-pyrrol-1-yl)-6-oxo-hexanoyl]-1H-pyrrole-2-
carboxylic acid benzyl ester

- 30 To a stirred solution of 1.0 g (4.97 mmol) 1H-pyrrole-2-carboxylic acid benzyl
ester, 0.06 g (0.50 mmol) 4-dimethylaminopyridine and 0.82 ml (5.47 mmol)
1,8-diazabicyclo[5.4.0]undec-7-ene in 40 ml dichloromethane at 0 °C was
added dropwise 0.36 ml (2.49 mmol) adipoyl chloride and stirring continued

for 1 h at room temperature. The reaction mixture was then washed sequentially with 1 M hydrochloric acid, saturated sodium bicarbonate solution and finally with saturated brine, and the aqueous phases back-extracted with dichloromethane. The combined organic extracts were dried
5 over sodium sulphate and concentrated *in vacuo*. Flash chromatography (25% EtOAc in hexane) afforded 450 mg (35%) of the title compound as a white crystalline solid. MS *m/e* (%): 530 ($M+NH_4^+$, 100).

c) Mixture of (R)-1-[6-[(S)- and (RS)-1-[6-[(RS)-2-carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid

- 10 Using General Procedure B with 800 mg (1.56 mmol) 1-[6-(2-benzyloxycarbonyl-pyrrol-1-yl)-6-oxo-hexanoyl]-1H-pyrrole-2-carboxylic acid benzyl ester afforded 470 mg (91%) of the title compound as a white crystalline solid. MS *m/e* (%): 339 ($[M-H]^+$, 100).

15

Example 99

1-[6-(2-Carboxy-pyrrol-1-yl)-6-oxo-hexanoyl]-1H-pyrrole-2-carboxylic acid

a) 1H-Pyrrole-2-carboxylic acid *tert*-butyl ester

- Lit. *Tetrahedron* 1985, 41, 5633. To a stirred solution of 10.0 g (90.0 mmol) pyrrole-2-carboxylic acid in 180 ml dioxane was added dropwise at 0 °C 18 ml
20 concentrated sulphuric acid 2-methylpropene was then condensed into the reaction flask over the course of 1 h, using a dry-ice condenser, and stirring continued for 16 h at 0 °C while the dry-ice condenser was periodically refilled so as to maintain a slow reflux of 2-methylpropene. The reaction mixture was then poured cautiously into an ice-cooled mixture of 400 ml ether and 150 ml 2
25 M sodium hydroxide solution. The phases were separated and the aqueous phase extracted twice more with ether. The combined organic phases were washed successively with 2 M sodium hydroxide solution, water and finally with saturated brine, then dried over sodium sulphate and concentrated *in vacuo* to afford 9.07 g (60%) of the title compound as a colourless oil which still
30 contained some dioxane. 1H NMR δ (250 MHz, $CDCl_3$) 9.60 (1H, br s), 6.90 (1H, m), 6.83 (1H, m), 6.22 (1H, m), 1.56 (9H, s).

b) 1-[6-(2-*tert*-Butoxycarbonyl-pyrrol-1-yl)-6-oxo-hexanoyl]-1H-pyrrole-2-carboxylic acid *tert*-butyl ester

- To a stirred solution of 1.3 g (7.77 mmol) 1H-pyrrole-2-carboxylic acid *tert*-butyl ester, 95 mg (0.78 mmol) 4-dimethylaminopyridine and 1.28 ml (8.56 mmol) 1,8-diazabicyclo[5.4.0]undec-7-ene in 40 ml dichloromethane at 0 °C was added dropwise 0.57 ml (3.91 mmol) adipoyl chloride and stirring
- 5 continued for 16 h at room temperature. A further 95 mg (0.78 mmol) 4-dimethylaminopyridine and 1.28 ml (8.56 mmol) 1,8-diazabicyclo[5.4.0]undec-7-ene were added and stirring continued for a further 4 h at room temperature. The reaction mixture was then washed sequentially with 1 M hydrochloric acid, saturated sodium bicarbonate solution and finally with saturated brine,
- 10 and the aqueous phases back-extracted with dichloromethane. The combined organic extracts were dried over sodium sulphate and concentrated *in vacuo*. Flash chromatography (gradient: 10-20% EtOAc in hexane) afforded 464 mg (13%) of the title compound as a light yellow crystalline solid. MS *m/e* (%): 462 ($M+NH_4^+$, 100).
- 15 c) 1-[6-(2-Carboxy-pyrrol-1-yl)-6-oxo-hexanoyl]-1H-pyrrole-2-carboxylic acid
- To a stirred solution of 55 mg (0.12 mmol) 1-[6-(2-*tert*-butoxycarbonyl-pyrrol-1-yl)-6-oxo-hexanoyl]-1H-pyrrole-2-carboxylic acid *tert*-butyl ester in 8 ml dichloromethane at 0 °C was added dropwise 0.15 ml (1.97 mmol) trifluoroacetic acid and stirring continued for 16 h at room temperature.
- 20 Concentration *in vacuo* and azeotroping three times with chloroform on a rotary evaporator afforded 39 mg (95%) of the title compound as an off-white crystalline solid. MS *m/e* (%): 350 ($M+NH_4^+$, 100).

25

Example 100

(R)-1-[6-[(R)-2-Carboxy-4,4-difluoro-pyrrolidin-1-yl]-6-oxo-hexanoyl]-4,4-difluoropyrrolidine-2-carboxylic acid

a) (2R)-4-Oxo-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-*tert*-butyl ester

- 30 A solution of 15.6 ml (0.22mol) dimethylsulfoxide in 50 ml dichloromethane was given to 9.60 ml oxalylchloride in 150 ml dichloromethane at minus 65 °C during a period of 10 minutes. After 5 minutes at -65 °C 32.1g (0.1mol) (2R,2R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-*tert*-butyl

ester in 100 ml dichloromethane were added and after additional 15 minutes 24.4 ml (0.18 mol) triethylamine. The cooling bath was removed, stirring was continued over night, and then the mixture was poured into ice-water.

Extraction with dichloromethane, followed by washing with 0.05 N HCl,

- 5 bicarbonate and water and chromatography over silicagel with dichloromethane/ ethylacetate 98:2 gave 11.1g (35%) (2R)-4-oxo-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-tert-butyl ester as a colorless liquid. MS m/e (%): 263(M-isobutylene, 7), 219 (8), 184 (24), 128 (14), 91 (58), 84 (40), 57 (100);

- 10 $[\alpha]_D = +1.1^\circ$ (c= 1% in methanol).

b) (2R)-4,4-Difluoro-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-tert-butyl ester

A solution of 0.64 g (0.002 mol) (2R)-4-oxo-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-tert-butyl ester in 3 ml dichloromethane was treated at 0°C with
15 0.79 ml (0.006 mol) diethylaminosulfur trifluoride, stirring was continued at room temperature for 32 hours and then the mixture was poured on ice.

Extraction with dichloromethane and filtration over silicagel with hexane followed by elution with dichloromethane gave 0.62 g (90%) (2R)-4,4-difluoro-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-tert-butyl ester as light

- 20 yellow oil.

MS m/e (%): 285(1), 206 (6), 106 (33), 91 (35), 57 (100); $[\alpha]_D = +43.0^\circ$ (c= 1% in methanol).

c) (2R)-4,4-Difluoro-pyrrolidine-2-carboxylic acid benzyl ester hydrochloride (1:1)

- 25 100 ml of dry HCl in diethylether was added to a solution of 7.51 g (0.022 mol) (2R)-4,4-difluoro-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-tert-butyl ester in a mixture 100 ml diethylether and 20 ml dichloromethane. After stirring for two days 5.84 g (96%) (2R)-4,4-difluoro-pyrrolidine-2-carboxylic acid benzyl ester hydrochloride (1:1) were isolated by filtration and dried
30 under reduced pressure. Mp.: 118-120°C; $[\alpha]_D = +29.3^\circ$ (c= 1% in methanol).

d) (R)-1-[6-[(R)-2-Benzoyloxycarbonyl-4,4-difluoro-pyrrolidin-1-yl]-6-oxo-hexanoyl]-4,4-difluoropyrrolidine-2-carboxylic acid benzyl ester

To a suspension of 1.11 g (0.004 mol) (2R)-4,4-difluoro-pyrrolidine-2-carboxylic acid benzyl ester hydrochloride (1:1) in 20 ml dichloromethane were added 1.17 ml (0.008 mol) triethylamine, and 0.29 ml (0.002 mol) adipoyldichloride in 5 ml dichloromethane. After stirring at room temperature over night the mixture was extracted with 1N HCl, water and aqueous sodiumbicarbonate and dried with sodiumsulfate. Chromatography over silicagel with dichloromethane/ethylacetate 8:2 gave 0.79 g (66%) (R)-1-[6-[(R)-2-benzyloxycarbonyl-4,4-difluoro-pyrrolidin-1-yl]-6-oxo-hexanoyl]-4,4-difluoropyrrolidine-2-carboxylic acid benzyl ester as colorless oil. ISP-MS: 593 (MH)⁺; $[\alpha]_D = +67.3^\circ$ (c= 1% in methanol).

e) (R)-1-[6-[(R)-2-Carboxy-4,4-difluoro-pyrrolidin-1-yl]-6-oxo-hexanoyl]-4,4-difluoropyrrolidine-2-carboxylic acid

0.59g (0.001mol) (R)-1-[6-[(R)-2-Benzyloxycarbonyl-4,4-difluoro-pyrrolidin-1-yl]-6-oxo-hexanoyl]-4,4-difluoropyrrolidine-2-carboxylic acid benzyl ester in 20 ml ethanol were hydrogenated at room temperature and atmospheric pressure in the presence of 0.12g 5% palladium/carbon. After completion of the reaction the catalyst was filtered off, the solvent was distilled off, and the residue was dissolved in dichloromethane. Evaporation gave 0.4g (97%) (R)-1-[6-[(R)-2-carboxy-4,4-difluoro-pyrrolidin-1-yl]-6-oxo-hexanoyl]-4,4-difluoropyrrolidine-2-carboxylic acid as a white foam. ISP-MS: 413 (MH)⁺; $[\alpha]_D = +54.2^\circ$ (c= 1% in dimethylsulfoxide).

Example 101

(R)-1-[[2-[2-[(R)-2-Carboxy-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethoxyl]-

phenoxy]acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid

a) (R)-1-[[2-[2-[(R)-2-Benzyloxycarbonyl-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethoxyl]-phenoxy]acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid benzyl ester

To a mixture of 1.11g (0.004 mol) (2R)-4,4-difluoro-pyrrolidine-2-carboxylic acid benzyl ester hydrochloride (1:1), 0.45g (0.002 mol) 1,2-

30 phenylenedioxydiacetic acid,

1.21g (0.012 mol) N-methylmorpholine, and 0.61g (0.004mol) 1-

hydroxybenzotriazole hydrate in 90 ml dichloromethane were added 0.77g

(0.004 mol) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimid hydrochloride. After

stirring at room temperature for 18 hours the mixture was extracted with 1N HCl, water, 10% aqueous sodium bicarbonate and again water.

Chromatography over silicagel with dichloromethane/ethylacetate 9:1 yielded 0.52g (39%) (R)-1-[[2-[2-[(R)-2-benzyloxycarbonyl-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid benzyl ester as colorless oil. ISP-MS: 673 (MH)⁺; [α]_D = +68.4° (c= 1% in methanol).

b) (R)-1-[[2-[2-[(R)-2-Carboxy-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid

0.47g (0.0007mol) (R)-1-[[2-[2-[(R)-2-Benzyloxycarbonyl-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid benzyl ester in 20 ml ethanol were hydrogenated at room temperature and atmospheric pressure in the presence of 0.09g 5% palladium/carbon. After completion of the reaction the catalyst was filtered off, the solvent was distilled off, and the residue was dissolved in dichloromethane. Evaporation gave 0.32g (94%) (R)-1-[[2-[2-[(R)-2-carboxy-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid as a light yellow foam. ISP-MS: 493 (MH)⁺; [α]_D = +46.8° (c= 1% in dimethylsulfoxide).

Example 102

20 (R)-1-[[4-[2-[(R)-2-Carboxy-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid

a) (R)-1-[[4-[2-[(R)-2-Benzyloxycarbonyl-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid benzyl ester

To a mixture of 1.11g (0.004 mol) (2R)-4,4-difluoro-pyrrolidine-2-carboxylic acid benzyl ester hydrochloride (1:1), 0.39g (0.002 mol) 1,4-phenylenediacetic acid, 1.21g (0.012 mol) N-methylmorpholine, and 0.61g (0.004mol) 1-hydroxybenzotriazole hydrate in 90 ml dichloromethane were added 0.77g (0.004 mol) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimid hydrochloride. After stirring at room temperature for 18 hours the mixture was extracted with 1N HCl, water, 10% aqueous sodium bicarbonate and again water.

Chromatography over silicagel with dichloromethane/ethylacetate 9:1 yielded 0.72g (56%) (R)-1-[[4-[2-[(R)-2-benzyloxycarbonyl-4,4-difluoropyrrolidin-1-yl]-2-

oxo-ethyl]-phenyl]-acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid benzyl ester as colorless oil. ISP-MS: 658 (MNH₄)⁺; [α]_D = +55.5° (c= 1% in methanol).

b) (R)-1-[[4-[2-[(R)-2-Carboxy-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid

- 5 0.64g (0.001mol) (R)-1-[[4-[2-[(R)-2-Benzoyloxycarbonyl-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid benzyl ester in 20 ml ethanol were hydrogenated at room temperature and atmospheric pressure in the presence of 0.13g 5% palladium/carbon. After completion of the reaction the catalyst was filtered off, the solvent was
- 10 distilled off, and the residue was dissolved in dichloromethane. Evaporation gave 0.28g (61%) (R)-1-[[4-[2-[(R)-2-carboxy-4,4-difluoropyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-4,4-difluoropyrrolidine-2-carboxylic acid as a light yellow foam. ISP-MS: 478 (MNH₄)⁺; [α]_D = +50.3° (c= 0.33 % in dimethylsulfoxide).

15

Example 103

(R)-1-[6-[(R)-2-Carboxy-2,5-dihydropyrrole-1-yl]-6-oxo-hexanoyl]-2,5-dihydropyrrole-2-carboxylic acid

- a) (2R,4R)-4-(Toluene-4-sulfonyloxy)-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-tert-butyl ester
- 20

A solution of 2.35g (0.007 mol) (2R,2R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-benzyl-ester 1-tert-butyl ester in 22 ml pyridine were treated at 5°C with 1.53g (0.008 mol) p-toluenesulfonyl chloride and kept in the refrigerator for 15 days. The pyridine was then distilled off in vacuo and the residue was

25 purified by chromatography on silicagel with dichloromethane/ethylacetate 95:5 to yield 2.81g (81%) (2R,4R)-4-(toluene-4-sulfonyloxy)-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-tert-butyl ester as colorless liquid. MS m/e (%): 376 (2), 340 (10), 284 (6), 240 (21), 91 (48), 68 (100), 57 (63); [α]_D = +24.5° (c= 1% in methanol).

- 30 b) (2R,4S)-4-Phenylselenanyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester

A solution of 7.99g (0.026 mol) diphenyl diselenide in 250 ml ethanol was treated with 1.59g (0.042 mol) sodium borohydride and stirring was continued

until the yellow solution turned colorless. After addition of 20.0g (0.042 mol) (2R,4R)-4-(toluene-4-sulfonyloxy)-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-tert-butyl ester the mixture was refluxed for 2.5 hours. A white precipitate was filtered off and the solvent was removed in vacuo.

- 5 Chromatography on silicagel with dichloromethane/methanol 98:2 gave 8.48g (51%)

(2R,4S)-4-Phenylselanyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester

- as a colorless oil. MS m/e (%): 399 (7), 326 (13), 270 (18), 226 (38), 186 (35), 68 (60), 57 (100), 41 (28). $[\alpha]_D = +40.4^\circ$ (c= 1% in methanol). In addition were isolated 0.6g (2R,4S)-4-phenylselanyl-pyrrolidine-1,2-dicarboxylic acid 2-benzyl ester 1-tert-butyl ester as colorless oil. MS m/e (%): 461(7), 326 (21), 270 (37), 248 (48), 226 (49), 209 (30), 91 (100), 68 (54), 57 (96). $[\alpha]_D = +33.2^\circ$ (c= 1% in methanol).

- 15 c) (R)-2,5-Dihydro-pyrrole-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester

- To a solution of 7.35g (0.016 mol) (2R,4S)-4-phenylselanyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester in 80 ml dichloromethane were added at 0-5°C 1.93 ml (0.024 mol) pyridine and 4.6 ml 30% hydrogen peroxide and stirring was continued for 1.5 hours. The mixture was extracted with 5% aqueous HCl, saturated aqueous sodium carbonate, and water. Chromatography on silicagel with ethylacetate/hexane 1:5 yielded 2.99g (77%) (R)-2,5-dihydro-pyrrole-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester as colorless oil. MS m/e (%): 186 (11), 168 (48), 140 (32), 112 (100), 68 (85), 57 (58). $[\alpha]_D = +242^\circ$ (c= 1% in chloroform).

- 25 d) (R)-2,5-Dihydro-1H-pyrrole-2-carboxylic acid ethyl ester trifluoroacetate (1:1)

- 1.5g (0.006mol) (R)-2,5-dihydro-pyrrole-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester were dissolved at 5°C in 10 ml trifluoroacetic acid and stirring was continued at room temperature for 3 hours. Evaporation of the solvent in vacuo gave 2.05g (quant.) of (R)-2,5-dihydro-1H-pyrrole-2-carboxylic acid ethyl ester trifluoroacetate (1:1) as a yellow oil. MS m/e (%): 142 (1), 68 (100), 45 (10), 41 (15). $[\alpha]_D = +95.4^\circ$ (c= 1% in methanol).

e) (R)-1-[6-[(R)-2-Ethylloxycarbonyl-2,5-dihydropyrrole-1-yl]-6-oxo-hexanoyl]-2,5-dihydropyrrole-2-carboxylic acid ethyl ester

To a suspension of 0.99 g (0.003 mol) (R)-2,5-dihydro-1H-pyrrole-2-carboxylic acid ethyl ester trifluoroacetate (1:1) in 20 ml dichloromethane were added
5 1.3 ml (0.009 mol) triethylamine, and 0.22 ml (0.0015 mol) adipoyldichloride in 5 ml dichloromethane. After stirring at room temperature over night the mixture was extracted with 1N HCl, water and aqueous sodiumbicarbonate and dried with sodiumsulfate. Chromatography over silicagel with ethylacetate gave 0.37 g (32%) (R)-1-[6-[(R)-2-ethylloxycarbonyl-2,5-dihydropyrrole-1-yl]-6-
10 oxo-hexanoyl]-2,5-dihydropyrrole-2-carboxylic acid ethyl ester as yellow oil. ISP-MS: 393 (MH)⁺.

f) (R)-1-[6-[(R)-2-Carboxy-2,5-dihydropyrrole-1-yl]-6-oxo-hexanoyl]-2,5-dihydropyrrole-2-carboxylic acid

0.09g (0.0002 mol) (R)-1-[6-[(R)-2-ethylloxycarbonyl-2,5-dihydropyrrole-1-yl]-6-
15 oxo-hexanoyl]-2,5-dihydropyrrole-2-carboxylic acid ethyl ester were stirred with aqueous HCl at 50°C for 3 hours. The solvent was evaporated and the residue dissolved in water and lyophilized to yield 0.08g (97%) (R)-1-[6-[(R)-2-carboxy-2,5-dihydropyrrole-1-yl]-6-oxo-hexanoyl]-2,5-dihydropyrrole-2-carboxylic acid as light yellow foam. ISP-MS: 335 (M-H)⁺.

20

Example 104

(R)-1-[[2-[2-[(R)-2-Carboxy-2,5-dihydropyrrole-1-yl]-2-oxo-ethoxyl]-phenoxy]acetyl]-2,5-dihydropyrrole-2-carboxylic acid

a) (R)-1-[[2-[2-[(R)-2-Ethylcarbonyl-2,5-dihydropyrrole-1-yl]-2-oxo-ethoxyl]-phenoxy]acetyl]-2,5-dihydropyrrole-2-carboxylic acid ethyl ester
25

To a mixture of 0.99g (0.003 mol) (R)-2,5-dihydro-1H-pyrrole-2-carboxylic acid ethyl ester trifluoroacetate (1:1), 0.34g (0.0015 mol) 1,2-phenylenedioxydiacetic acid,

1.3 ml (0.012 mol) N-methylmorpholine, and 0.46g (0.003mol) 1-
30 hydroxybenzotriazole hydrate in 80 ml dichloromethane were added 0.57g (0.003 mol) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimid hydrochloride. After stirring at room temperature for 18 hours the mixture was extracted with 1N HCl, water, 10% aqueous sodium bicarbonate and again water.

Chromatography over silicagel with ethylacetate yielded 0.45g (32%) (R)-1-[[2-[2-[(R)-2-ethylcarbonyl-2,5-dihydropyrrole-1-yl]-2-oxo-ethoxyl]-phenoxy]acetyl]-2,5-dihydropyrrole-2-carboxylic acid ethyl ester as colorless oil. ISP-MS: 473 (MH)⁺.

- 5 b) (R)-1-[[2-[2-[(R)-2-Carboxy-2,5-dihydropyrrole-1-yl]-2-oxo-ethoxyl]-phenoxy]acetyl]- 2,5-dihydropyrrole-2-carboxylic acid

0.17g (0.0004 mol) (R)-1-[[2-[2-[(R)-2-ethylcarbonyl-2,5-dihydropyrrole-1-yl]-2-oxo-ethoxyl]-phenoxy]acetyl]-2,5-dihydropyrrole-2-carboxylic acid ethyl ester were stirred with aqueous HCl at 50°C for 3 hours. The solvent was

- 10 evaporated and the residue dissolved in water and lyophilized to yield 0.13g (86%) (R)-1-[[2-[2-[(R)-2-carboxy-2,5-dihydropyrrole-1-yl]-2-oxo-ethoxyl]-phenoxy]acetyl]- 2,5-dihydropyrrole-2-carboxylic acid as white amorphous powder. ISP-MS: 417 (MH)⁺.

Example A

Tablets of the following composition were manufactured in the usual manner:

	<u>mg/tablet</u>
5 Active ingredient	100
Powd. lactose	95
White corn starch	35
Polyvinylpyrrolidone	8
Na carboxymethylstarch	10
10 Magnesium stearate	2
Tablet weight	250

Example B

Tablets of the following composition are manufactured in the usual manner:

	<u>mg/tablet</u>
Active ingredient	200
Powd. lactose	100
White corn starch	64
20 Polyvinylpyrrolidone	12
Na carboxymethylstarch	20
Magnesium stearate	4
Tablet weight	400

25

Example C

Capsules of the following composition are manufactured:

	<u>mg/capsule</u>
Active ingredient	50
Cryst. lactose	60
30 Microcrystalline cellulose	34

Talc	5
Magnesium stearate	1

Capsule fill weight 150


The active ingredient having a suitable particle size, the crystalline
5 lactose and the microcrystalline cellulose are homogeneously mixed with one
another, sieved and thereafter talc and magnesium stearate are admixed. The
finished mixture is filled into hard gelatine capsules of suitable size.

Claims

This Sheet Has Been Replaced




1. Compounds of the general formulae



I-A

or




I-B

5

wherein

R is SH, benzyl or phenyl, optionally substituted by hydroxy or lower alkoxy or



the group

;

THIS SHEET HAS BEEN REPLACED

10 R¹ is hydrogen or halogen;

X is $-(CH_2)_n-$; $-CH(R^2)(CH_2)_n-$; $-CH_2O(CH_2)_n-$; $-CH_2NH-$; benzyl, $-C(R^2)=CH-$; $-CH_2CH(OH)-$; or thiazol-2,5-diyl;

Y is $-S-S-$; $-(CH_2)_n-$; $-O-$; $-NH-$; $-N(R^3)-$; $-CH=CH-$; $-NHC(O)NH-$; $-N(R^3)C(O)N(R^3)-$; $-N[CH_2C_6H_3(OCH_3)_2]-$; $-N(CH_2C_6H_5)-$;

15 $-N(CH_2C_6H_5)C(O)N(CH_2C_6H_5)-$; $-N(alkoxyalkyl)-$; $-N(cycloalkyl-methyl)-$; 2,6-pyridyl; 2,5-furanyl; 2,5-thienyl; 1,2-cyclohexyl; 1,3-cyclohexyl; 1,4-cyclohexyl; 1,2-naphthyl; 1,4-naphthyl; 1,5-naphthyl; 1,6-naphthyl; biphenyl; or 1,2-phenylene, 1,3-phenylene and 1,4-phenylene, wherein the phenylene groups are optionally substituted by 1 - 4 substituents,

20 selected from halogen, lower alkyl, lower alkoxy, hydroxy, carboxy, $-COO-$ lower alkyl, nitrilo, 5-tetrazol, (2-carboxylic acid-pyrrolidin-1-yl)-2-oxo-ethoxy, N-hydroxycarbamimidoyl, 5-oxo-[1,2,4]oxadiazolyl, 2-oxo-[1,2,3,5]oxathiadiazolyl, 5-thioxo-[1,2,4]oxadiazolyl and 5-tert-butylsulfanyl-[1,2,4]oxadiazolyl;



X' is $-(CH_2)_n$; $-(CH_2)_nCH(R^2)$; $-(CH_2)_nOCH_2$; $-NHCH_2$; benzyl,
 $-CH=C(R^2)$; $-CH(OH)CH_2$; or thiazol-2,5-diyl;

R^2 is lower alkyl, lower alkoxy or benzyl and

n is 0-3,

- 5 and pharmaceutically acceptable salts or mono- and diesters thereof, with the exception of (R)-1-[(R)- and (R)-1-[(S)-3-mercapto-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid.

2. Compounds of general formula I-A in accordance with claim 1.

- 10 3. Compounds in accordance with claim 2, wherein X is $CH(R^2)(CH_2)_n$ and R^2 is methyl or methoxy and n is 0 or 1.

4. Compounds in accordance with claim 3, which are:

(R)-1-[(S)-3-[(S)-3-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-methyl-3-oxopropyl-disulfanyl]-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid,

- 15 (R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,7-dimethyl-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid,

(R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,7-dimethoxy-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid and

(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexanoyl]-

- 20 pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

5. Compounds in accordance with claim 2, wherein X is $-(CH_2)_n$ and n is 0 or 1.

- 25 6. Compounds in accordance with claim 5, which are:

(R)-1-[7-[(R)-2-Carboxy-pyrrolidin-1-yl]-7-oxo-heptanoyl]-pyrrolidine-2-carboxylic acid,

(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid,

30



This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

(R)-1-[5-[(R)-2-Carboxy-pyrrolidin-1-yl]-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid,

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]acetyl]-pyrrolidine-2-carboxylic acid,

- 5 (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-ureido]-pyrrolidine-2-carboxylic acid,

(R)-1-[[Benzyl-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-amino]-acetyl]-pyrrolidine-2-carboxylic acid,

- (R)-1-[cis-4-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid and
- 10

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid.

7. Compounds in accordance with claim 2, wherein X is $-\text{CH}_2\text{O}-$

- 15 8. Compounds in accordance with claim 7, which are:

(R)-1-[[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,

- (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,
- 20

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,

- 25 (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid and

(R)-1-[[5-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid.

- 30 9. Compounds in accordance with claim 2, wherein X is $-\text{CH}_2\text{NH}-$

10. Compounds in accordance with claim 9, which compound is



This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid.

11. Compounds in accordance with claim 2, wherein X is $-\text{CH}_2\text{CH}(\text{OH})-$.


5 12. Compounds in accordance with claim 11, which compound is

(2E,4E)-(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexa-2,4-dienoyl]-pyrrolidine-2-carboxylic acid.

13. A medicament, especially for the treatment or prevention of
10 amyloidosis, containing a compound in accordance with any one of claims 1-12 or (R)-1-[(R)- and (R)-1-[(S)-3-mercapto-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid or a pharmaceutically acceptable salt or mono- and diesters thereof as well as a therapeutically inert carrier material.

14. A process for the manufacture of compounds in accordance with any
15 one of claims 1-12, which process comprises

a) cleaving off the protecting group from a compound of formulae



20 wherein R, R¹, X, Y and X' are described in claim 1 and R³ is a protecting group,
to give a compound of formula I-A or I-B, and, if desired,

converting a compound of general formulae I-A and I-B into a pharmaceutically usable salt or into a mono- and diester thereof.

25



This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

15. Compounds according to any one of claims 1-12, whenever manufactured by the process defined in claim 14 or by an obvious chemical equivalent thereof.

16. Compounds according to any one of claims 1-12 or to (R)-1-[(R)- and
5 (R)-1-[(S)-3-mercapto-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid as well as pharmaceutically acceptable salts and mono-or diesters thereof for use as therapeutically active substances, especially against central and systemic forms of amyloidosis.

17. The use of compounds according to any one of claims 1-12 or of (R)-1-
10 [(R)- and (R)-1-[(S)-3-mercapto-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid and of pharmaceutically usable salts and mono-or diesters thereof, especially for the treatment or prevention of central and systemic forms of amyloidosis, and, respectively, for the production of corresponding medicaments.

15 18. The invention as hereinbefore described.



This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED



Abstract


The invention relates to D-prolines of the formula



wherein

R is SH, benzyl or phenyl, optionally substituted by hydroxy or lower alkoxy

or



the group

;

R¹ is hydrogen or halogen;

X is $-(CH_2)_n-$; $-CH(R^2)(CH_2)_n-$; $-CH_2O(CH_2)_n-$; $-CH_2NH-$; benzyl, $-C(R^3)=CH-$; $-CH_2CH(OH)-$; or thiazol-2,5-diyl;

Y is $-S-S-$; $-(CH_2)_n-$; $-O-$; $-NH-$; $-N(R^3)-$; $-CH=CH-$; $-NHC(O)NH-$; $-N(R^3)C(O)N(R^3)-$; $-N[CH_2C_6H_3(OCH_3)_2]-$; $-N(CH_2C_6H_5)-$; $-N(CH_2C_6H_5)C(O)N(CH_2C_6H_5)-$; $-N(alkoxyalkyl)-$; $-N(cycloalkyl-methyl)-$; 2,6-pyridyl; 2,5-furanyl; 2,5-thienyl; 1,2-cyclohexyl; 1,3-cyclohexyl; 1,4-cyclohexyl; 1,2-naphthyl; 1,4-naphthyl; 1,5-naphthyl; 1,6-naphthyl; biphenylen; or 1,2-phenylen, 1,3-phenylen and 1,4-phenylen, wherein the phenylen groups are optionally substituted by 1 - 4 substituents, selected from halogen, lower alkyl, lower alkoxy, hydroxy, carboxy, $-COO-$ lower alkyl, nitrilo, 5-tetrazol, (2-carboxylic acid-pyrrolidin-1-yl)-2-oxo-ethoxy, N-hydroxycarbamimidoyl, 5-oxo-[1,2,4]oxadiazolyl, 2-oxo-[1,2,3,5]oxathiadiazolyl, 5-thioxo-[1,2,4]oxadiazolyl and 5-tert-butylsulfanyl-[1,2,4]oxadiazolyl;

THIS SHEET HAS BEEN
REPLACED

X' is $-(CH_2)_n-$; $-(CH_2)_nCH(R^2)-$; $-(CH_2)_nOCH_2-$; $-NHCH_2-$; benzyl,
 $-CH=C(R^3)-$; $-CH(OH)CH_2-$; or thiazol-2,5-diyl;

R² is lower alkyl, lower alkoxy or benzyl and

n is 0-3,


and to pharmaceutically acceptable salts and mono- and diesters thereof.

The D-prolines of formula I-A and I-B can be used in the treatment or prevention of all forms of central and systemic amyloidosis.

This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED



D-PROLINE DERIVATIVES


9804381-3

The invention relates to D-prolines of the formula



I-A

or




I-B

wherein

- 5 R is SH, benzyl or phenyl, optionally substituted by hydroxy or lower alkoxy

or



the group ;


- R¹ is hydrogen or halogen;
- 10 X is $-(CH_2)_n-$; $-CH(R^2)(CH_2)_n-$; $-CH_2O(CH_2)_n-$; $-CH_2NH-$; benzyl, $-C(R^2)=CH-$; $-CH_2CH(OH)-$; or thiazol-2,5-diyl;
- Y is $-S-S-$; $-(CH_2)_n-$; $-O-$; $-NH-$; $-N(R^3)-$; $-CH=CH-$; $-NHC(O)NH-$; $-N(R^3)C(O)N(R^3)-$; $-N[CH_2C_6H_4(OCH_3)]_n-$; $-N(CH_2C_6H_4)-$; $-N(CH_2C_6H_4)C(O)N(CH_2C_6H_4)-$; $-N(alkoxyalkyl)-$; $-N(cycloalkyl-methyl)-$;
- 15 2,6-pyridyl; 2,5-furanyl; 2,5-thienyl; 1,2-cyclohexyl; 1,3-cyclohexyl; 1,4-cyclohexyl; 1,2-naphthyl; 1,4-naphthyl; 1,5-naphthyl; 1,6-naphthyl; biphenylen; or 1,2-phenylen, 1,3-phenylen and 1,4-phenylen, wherein the phenylen groups are optionally substituted by 1 - 4 substituents,

D-PROLINE DERIVATIVES

Abstract




The invention relates to D-prolines of the formula



I-A


or



I-B

wherein the definition of substituents is given in the description,
and to pharmaceutically acceptable salts and mono- and diesters thereof.


The D-prolines of formula I-A and I-B can be used in the treatment or prevention of all forms of central and systemic amyloidosis.



Claims


9804381-3

1. D-proline derivatives of the general formulae



I-A


or



I-B

wherein

R is SH, benzyl or phenyl, optionally substituted by hydroxy or lower alkoxy or



the group

- 10 R¹ is hydrogen or halogen;
- X is -(CH₂)_n-; -CH(R²)(CH₂)_n-; -CH₂O(CH₂)_n-; -CH₂NH-; benzyl, -C(R²)=CH-; -CH₂CH(OH)-; or thiazol-2,5-diyl;
- Y is -S-S-; -(CH₂)_n-; -O-; -NH-; -N(R³)-; -CH=CH-; -NHC(O)NH-; -N(R³)C(O)N(R³)-; -N[CH₂C₆H₃(OCH₃)₂]-; -N(CH₂C₆H₅)-;
- 15 -N(CH₂C₆H₅)C(O)N(CH₂C₆H₅)-; -N(alkoxyalkyl)-; -N(cycloalkyl-methyl)-; 2,6-pyridyl; 2,5-furanyl; 2,5-thienyl; 1,2-cyclohexyl; 1,3-cyclohexyl; 1,4-cyclohexyl; 1,2-naphthyl; 1,4-naphthyl; 1,5-naphthyl; 1,6-naphthyl;

9804381-3
5/4/00



biphenyl; or 1,2-phenylen, 1,3-phenylen and 1,4-phenylen, wherein the phenylen groups are optionally substituted by 1 - 4 substituents, selected from halogen, lower alkyl, lower alkoxy, hydroxy, carboxy, -COO-lower alkyl, nitrilo, 5-tetrazol, (2-carboxylic acid-pyrrolidin-1-yl)-
 5 2-oxo-ethoxy, N-hydroxycarbamimidoyl, 5-oxo-[1,2,4]oxadiazolyl, 2-oxo-[1,2,3,5]oxathiadiazolyl, 5-thioxo-[1,2,4]oxadiazolyl and 5-tert-butylsulfanyl-[1,2,4]oxadiazolyl;

X' is $-(CH_2)_n-$; $-(CH_2)_nCH(R^2)-$; $-(CH_2)_nOCH_2-$; $-NHCH_2-$; benzyl.
 $-CH=C(R^2)-$; $-CH(OH)CH_2-$; or thiazol-2,5-diyl;

10 R² is lower alkyl, lower alkoxy or benzyl and
 n is 0-3,

with the proviso, that when R is SH and R¹ is hydrogen, in formula I-B X cannot be $-CH(R^2)-CH_2-$ for R² hydrogen, lower alkyl or benzyl,


15 and pharmaceutically acceptable salts or mono- and diesters thereof

20 2. D-proline derivatives of general formula I-A in accordance with claim 1.

3. Compounds in accordance with claim 2, wherein X is $CH(R^2)(CH_2)_n-$ and R² is methyl or methoxy and n is 0 or 1.

4. Compounds in accordance with claim 3, which are:

(R)-1-[(S)-3-[(S)-3-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-methyl-3-oxopropyl-
 25 disulfanyl]-2-methyl-propionyl]-pyrrolidine-2-carboxylic acid,
 (R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,7-dimethyl-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid,
 (R)-1-[8-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,7-dimethoxy-8-oxo-octanoyl]-pyrrolidine-2-carboxylic acid and





(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid (mixture of 3 diastereomers)

5. Compounds in accordance with claim 2, wherein X is $-(CH_2)_n-$ and n is 0 or 1.


6. Compounds in accordance with claim 5, which are:

- (R)-1-[7-[(R)-2-Carboxy-pyrrolidin-1-yl]-7-oxo-heptanoyl]-pyrrolidine-2-carboxylic acid,
- (R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-6-oxo-hexanoyl]-pyrrolidine-2-carboxylic acid,
- (R)-1-[5-[(R)-2-Carboxy-pyrrolidin-1-yl]-5-oxo-pentanoyl]-pyrrolidine-2-carboxylic acid,
- (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]acetyl]-pyrrolidine-2-carboxylic acid,
- (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-ureido]-pyrrolidine-2-carboxylic acid,
- (R)-1-[[Benzyl-[2-[(R)-2-carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-amino]-acetyl]-pyrrolidine-2-carboxylic acid,
- (R)-1-[cis-4-[(R)-2-Carboxy-pyrrolidine-1-carbonyl]-cyclohexanecarbonyl]-pyrrolidine-2-carboxylic acid and
- (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethyl]-phenyl]-acetyl]-pyrrolidine-2-carboxylic acid.

7. Compounds in accordance with claim 2, wherein X is $-CH_2O-$

8. Compounds in accordance with claim 7, which are:

- (R)-1-[[2-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,
- (R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,





(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methoxy-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid,

(R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid.

5 (R)-1-[[3-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-2-methyl-phenoxy]-acetyl]-pyrrolidine-2-carboxylic acid and

(R)-1-[[5-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethoxy]-naphthalen-1-yloxy]-acetyl]-pyrrolidine-2-carboxylic acid.

10 9. Compounds in accordance with claim 2, wherein X is $-\text{CH}_2\text{NH}$:

10. Compounds in accordance with claim 9, which compound is

(R)-1-[[4-[2-[(R)-2-Carboxy-pyrrolidin-1-yl]-2-oxo-ethylamino]-phenylamino]-acetyl]-pyrrolidine-2-carboxylic acid.

15 11. Compounds in accordance with claim 2, wherein X is $-\text{CH}_2\text{CH}(\text{OH})-$.


12. Compounds in accordance with claim 11, which compound is


(2E,4E)-(R)-1-[6-[(R)-2-Carboxy-pyrrolidin-1-yl]-2,5-dimethyl-6-oxo-hexa-2,4-dienoyl]-pyrrolidine-2-carboxylic acid.

20 13. A medicament, especially for the treatment or prevention of amyloidosis, containing a compound in accordance with any one of claims 1-12 or a pharmaceutically acceptable salt or mono-and diesters thereof as well as a therapeutically inert carrier material.

25 14. A process for the manufacture of compounds in accordance with any one of claims 1-12, which process comprises

a) cleaving off the protecting group from a compound of formulae





wherein R, R¹, X, Y and X' are described in claim 1 and R¹ is a protecting group,

to give a compound of formula I-A or I-B, and, if desired,

5

converting a compound of general formulae I-A and I-B into a pharmaceutically usable salt or into a mono- and diester thereof.

15 15. Compounds according to any one of claims 1-12, whenever manufactured by the process defined in claim 14 or by an obvious chemical equivalent thereof.

16. Compounds according to any one of claims 1-12 as well as pharmaceutically acceptable salts and mono-or diesters thereof for use as
15 therapeutically active substances, especially against central and systemic forms of amyloidosis.

17. The use of compounds according to any one of claims 1-12 and of pharmaceutically usable salts and mono-or diesters thereof, for the production of corresponding medicaments for treating or preventing central and systemic forms of amyloidosis.

