

POTENTIATION OF ANTIMICROBIAL EFFECTS

Technical field

5

The present invention relates to a novel antimicrobial composition and to a process for the preparation thereof. The invention further relates to the composition for use as a pharmaceutical composition, to the use of the
10 composition for the preparation of a pharmaceutical preparation as well as to usages of said composition as a preservative additive in cosmetic, medical or food products. More specifically, the composition contains
15 antimicrobial properties of previously known antimicrobial compounds or impart antimicrobial properties to compounds which have not in themselves been regarded as antimicrobials.

20 Background to the invention

The use of antibacterial substances is common within, by way of example, the field of dermatology. The aim of this use is partly to treat the patient with a specific
25 antibacterial substance against some infection, e.g. antimycotics against fungal infections, partly to preserve the products against microbial contamination.

In cases where a patient is to be treated, specific-action
30 antimicrobial substances without general effects are often used. The disadvantage of such treatments is that "typing", a dermination in order to enable choice of the appropriate preparation, must be undertaken before the treatment. Additionally, there is the risk that other
35 microorganisms may invade during treatment against which the preparation has no effect.

On the other hand, if the antimicrobial substance is used as a preservative in the product, it is desirable that it has no effect on the patient, either of an antimicrobial or a toxic nature. The substances that are nowadays used
5 for this purpose are well documented as regards preservative effects but it is also known that many of them exhibit toxic effects in some form or other.

The present invention enables the elimination or at least
10 the drastic reduction of the above shortcomings or disadvantages.

General description of the invention

15 In the practice of the invention, it has surprisingly been found that the combination of particular monoglycerides with certain specific groups of chemical substances provides a synergistic effect with regard to antimicrobial properties. In this respect, it has been established that
20 the invention not only provides a synergistic effect in relation to compounds which were previously known to exhibit certain antimicrobial properties, but that it also leads to synergistic effects in relation to chemical substances which had previously not been known to exhibit
25 antimicrobial properties at all.

Aside from the monoglycerides used in accordance with the invention being tolerated particularly well by humans, especially due to their near relationship to substances
30 naturally occurring in the body, many of the substances with which the monoglycerides are combined in accordance with the invention are of a type which does not give rise to any toxic effects in humans. Accordingly, the present invention opens unsuspected possibilities to produce new
35 preparations of a medically active character as well as a preservative character and which do not give rise to any side-effects. Through the synergistic effect of the

invention, it additionally becomes possible to use the "active" component in a lower concentration than previously or to use an "active" component which has previously been out of the question altogether, which can also be an advantage from a non-toxicity viewpoint. Furthermore, the new combination in accordance with the invention has proven to give rise to antimicrobial properties of a very general character and therefore even the abovementioned disadvantages with typing and very specific-action antimicrobial properties can be eliminated or reduced.

Other advantages of the invention should be apparent to the man skilled in the art after reviewing the following more detailed description of the invention.

The antimicrobial composition in accordance with the present invention is more particularly distinguished in that it comprises an antimicrobially effective amount of a combination of components A) and B), where A) is the monoglyceride of lauric acid, the monoglyceride of myristic acid or a blend of both of these monoglycerides, and component B) is at least one chemical substance selected from the following groups:

25

- i) a local anaesthetic of the amide type;
- ii) carbamide;
- iii) an antimicrobial or antibacterial substance in the form of a steroid antibiotic, an imidazole derivative or a nitroimidazole derivative; and
- iv) a diol with 3 - 6 carbon atoms.

30

The composition can additionally contain, if so required, C) a conventional physiologically acceptable carrier and/or physiologically acceptable additives.

35

The invention is thus founded on the surprising

observation that the monoglyceride of lauric acid and the monoglyceride of myristic acid impart antimicrobial properties in the abovementioned groups of chemical substances where they have not previously been observed at all or potentiate the antimicrobial properties in the groups of chemical substances in question in those cases where antimicrobial properties, as such, were previously known. In this connection, it should be mentioned that John J. Kabara in the February 1984 JAOCs vol. 61:2 pp 397 - 403, "Antimicrobial Agents Derived from Fatty Acids", reports that the monoglycerides of, for example, lauric acid and myristic acid per se exhibit certain antimicrobial properties. However, Kabara has only demonstrated effects on Gram positive bacteria but above all else the practice of the present invention has proven that a strong synergistic effect is obtained which for example can mean that the minimum inhibitory concentration with regard to certain microorganisms is a considerable number of orders of magnitude lower than would be expected on the basis of purely accumulative effects.

Of the two monoglycerides, also known as monolaurin and monomyristin respectively, the monoglyceride of lauric acid, i.e. monolaurin, is especially preferred from an effectiveness viewpoint but also from a physiological viewpoint because for example upon application to the skin it is particularly well combinable or compatible with the lipids that are already present on the skin surface. Often, this means that monolaurin is the sole component A) in the composition or that it comprises the dominant component of the two monoglycerides, calculated by weight, but this is not necessarily always the case as the relevant monoglycerides can also function in certain preparations as carrier, vehicle or consistency agent, as will be reverted to further below. However, for use in skin products it is particularly convenient to blend monolaurin with monomyristin because the melting point for

monolaurin crystals in water is 36°C while the blend of monolaurin and monomyristin in the proportions 1:5 to 5:1, in water, melts at 31 - 32°C.

- 5 The especially preferred monoglycerides are the 1- or 3-monoglycerides of lauric acid and myristic acid, respectively, e.g. 1- or 3- monolaurin and 1-monomyristin, to a not insignificant degree because the corresponding 2-monoglyceride, if synthesized, is converted to the 1- or
10 3-monoglyceride in aqueous solution for example. In view of the above, the especially preferred monoglyceride is therefore 1-monolaurin or 3-monolaurin.

- As monolaurin and monomyristin respectively are generally
15 accepted nowadays as being completely non-toxic, it will be appreciated that additional commentary regarding the physiological acceptability of the composition of the invention need not be further elaborated here. Thus for applications in which the product comes into contact with
20 humans those grades of the relevant monoglyceride will, of course, be selected from those which are known to be non-toxic and acceptable.

- In relation to the especially preferred substances in the
25 various B) groups, the following should be noted.

Particularly preferred local anaesthetics of the amide type, i.e. component i), are lidocaine, prilocaine, mepivacaine, cinchocaine, bupivacaine and etidocaine.

30

The especially preferred substance amongst said local anaesthetics is lidocaine, to a not insignificant degree due to it being well known in medical connections.

- 35 Carbamide, often also known as urea, is of interest in relation to the invention as, per se, it has been previously known and generally accepted in dermatological

connections.

Fusidic acid and cephalosporin P can be mentioned as preferred antimicrobial substances of the steroid antibiotic form, of which fusidic acid is especially preferred.

Econazole nitrate and miconazole nitrate can be mentioned as preferred examples of antimicrobial substances of the imidazole derivative form, of which econazole is especially preferred.

Advantageous antimicrobial substances of the nitroimidazole derivative form are tinidazole and metronidazole.

As regards the diol with 3 - 6 carbon atoms, i.e. component iv), propanediols, butanediols, pentanediols and hexanediols are included, of which pentanediols are preferred. The particularly preferred pentanediol is 1,5-pentanediol.

The group B) chemical substance is, of course, also selected with regard to its physiological acceptability. The latter expression should thus be interpreted broadly as the composition of the invention is usable in medical connections as well as food and cosmetic connections, as will be described further below. In other words, the expression includes "pharmaceutically acceptable" in its conventional meaning in medical connections, and similarly "acceptable" or "allowed" in food and/or cosmetic products in accordance with the regulations applying in individual countries for these types of products.

The monoglycerides A), in accordance with the invention have proven to provide usable effects even at very low concentrations and therefore the lower limit for said component can be set as low as 0.2 ppm, calculated as

weight relative to the total weight of the entire composition. However, in many cases at least 1 ppm or at least 10 ppm can be required to achieve usable results in practice. On the other hand, the upper limit is virtually impossible to state in more specific figures, as the composition of the invention is often used, e.g. as a salve, cream or or lotion where the monoglyceride also functions as a carrier or consistency agent and due to this is used in concentrations of the order of several tens of percent, calculated on the total weight of the composition.

As will be apparent from the above, it is often a matter of very broad limits within which the preferred concentrations can be specified from 0.2 ppm to 30% of the total weight of the composition. Other suitable ranges within the above are from 1 ppm to 30% and from 10 ppm to 30%. However particularly preferred preparations of the salve, cream and lotion type often contain 5 - 30 weight percent, preferably 10 - 30 weight percent.

Component B) has also proven to give the above described synergistic effect together with component A) at very low concentrations. More particularly, it can often be a matter of a lower limit of 1 ppm, 10 ppm or 100 ppm, combined with an upper limit of 20 weight percent or even better 10 weight percent. Ranges of interest within the above are from 1 ppm to 10%, from 10 ppm to 10% and from 100 ppm to 10%, respectively. However, especially with products of the salve, cream and lotion type, it can often be a matter of preferred ranges of the order 0.1 - 20 weight percent, in particular 0.1 - 10 weight percent.

As is apparent from the above, the composition of the invention is usable firstly as an active component in pharmaceutical preparations to impart antimicrobial properties to such preparations, in particular

antibacterial and antifungal properties but also antiviral properties in certain cases. Secondly, as an additive, the composition of the invention acts as a preservative agent in food and cosmetic products. Additionally, they are, of course, usable in other connections where antimicrobial properties are desired, in which case the composition of the invention can be prepared in a form which is usable for the relevant application.

10 This means that the composition of the invention can often also contain component C), i.e. some form of physiologically acceptable carrier and/or one or more physiologically acceptable additives. Said carrier and additives are selected entirely according to conventional techniques and therefore need not be further commented upon here. However it should be stressed that the composition of the invention for use in medical connections is primarily intended for dermatological use which means that it is often prepared in the form of a salve, cream, gel, solution, lotion or emulsion to which any carrier and/or additive must be suited.

The present invention is further directed to a process for the preparation of the above defined antimicrobial composition, the above recited preferred embodiments of the composition as regards ingredients and amounts also applying in relation to the process. The process of the invention is more particularly distinguished in that: the composition is prepared in the form of a solid lipid crystal composition with surface-active hydrophilic crystals which are per se known, e.g. in GB patent publication 1,174,672.

The process in accordance with the invention for the preparation of such a lipid crystal composition includes heating the monoglyceride(s) A) to a temperature above the transition or conversion temperature of the lipid,

optionally in the presence of carrier and/or additive C),
adding substance(s) B) to the heated mixture and
maintaining said temperature until conversion has taken
place and cooling the mixture to approximately room
5 temperature or some other desired use-temperature to
produce said surface-active solid crystal composition.

A preferred embodiment of the process requires that the
cooling is carried out at a rate of around 1 - 3°C per
10 minute.

Other types of compositions in accordance with the
invention are prepared by mixing components A), B) and
optionally C), in per se known methods depending on the
15 desired form of preparation.

From the above it will further be clear that a special
aspect of the invention comprises the above defined
composition, including all the preferred variants thereof,
20 for use as an antimicrobial pharmaceutical composition,
especially an antibacterial or antifungal pharmaceutical
composition, conveniently in salve, cream, gel or lotion
form or particularly in a solid crystal composition of the
above mentioned type.

25 According to a further aspect of the invention, there is
provided the use of the above mentioned composition,
including the preferred embodiments thereof, for the
preparation of a pharmaceutical preparation for combatting
30 bacteria and fungi, which preparation is primarily
intended for dermatological use or application to the
skin.

Another use of the above mentioned composition, which also
35 includes the preferred embodiments thereof, in accordance
with the invention is as a preservative additive in
various types of products which are liable to attack by

bacteria or fungi, namely cosmetic, food and medical products.

Detailed description of the invention

5

The invention will now be further described in relation to the following non-limiting Examples which illustrate the preparation of compositions in accordance with the invention, the constituents of these compositions, and
10 their antimicrobial effects.

Example 1

Antimicrobial preparation containing lidocaine

15

A preparation with the following composition by weight was prepared:

20	1-glycerol monolaurate	5.5%
	1-glycerol monomyristate	16.5%
	Lidocaine	5 %
	Propylene glycol	5 %
	Purified water (Ph.Eur.)	to 100 %

25 Preparation was as follows: the monoglycerides were heated to 70°C together with the water and propylene glycol, and the lidocaine added. The mixture was then cooled at a rate of 1 - 3°C per minute to room temperature.

30 Conventional buffering substances can be added to the preparation to provide a pH of 6 - 7 so that the xylocaine stays in solution. Thickening agents, which can be organic or inorganic, can also be added to improve cosmetic properties, if so desired. If thickening agents are
35 utilized, these are added to the aqueous phase prior to heating or at 70°C. Buffering substances are conveniently added at 70°C. Other substances serving to modify the

properties of the product, for example in relation to taste, odour and consistency can be provided in an appropriate manner. This also applies to the remaining Examples.

5

The resulting preparation was tested in a Kelsey Test in which it proved to be very active against both bacteria and fungi. It was still active, from an antimicrobial viewpoint, after 6 inoculations with Staph. aureus,

10 Strept. faecalis, Ps. aeruginosa and Candida albicans. Effects on the replication of the HSV1 and 2 viruses have been demonstrated with the above preparation.

Example 2

15

Antimicrobial cream containing pentanediol

A cream with the following composition by weight was prepared:

20

1-glycerol monolaurate	7 %
1-glycerol monomyristate	21 %
1,5-pentanediol	5 %
Purified water	to 100 %

25

Additives such as consistency agents and buffering substances, for example, can be added to the product if so desired.

30 During preparation, the monoglycerides and water were heated to 70°C after which the pentanediol was added. After 15 minutes agitation at said temperature, the mixture was cooled at a rate of 1 - 3°C per minute to room temperature. Addition of consistency agents or buffering
 35 substances, if any, is conveniently done prior to cooling, i.e. at 70°C or prior to heating.

An MIC (minimum inhibitory concentration) determination showed the cream to have a 10,000 fold greater effect than a pure solution of 1,5-pentanediol on the microorganisms listed below:

5	<u>Microorganism</u>	<u>MIC value (mg/ml)</u>	
10		1,5-pentane- diol solution	Cream of the invention
	S. aureus	50	0.0013
	S. epidermis	50	0.00075
	C. albicans	40	0.0025
	T. rubrum	10	0.0013
15	P. ovale	20	0.00025

The above MIC values clearly show that the preparation can be utilized as an antimicrobial preparation at very low concentrations.

20

In addition to antibacterial and antifungal effects, the preparation also has an effect on the replication of HSV1 without damage to the host cell lines, e.g green monkey kidney cells, GMK AH 1.

25

Example 3

Antimicrobial gel containing pentanediol

30 A preparation with the following composition by weight was prepared:

35	1,5 pentanediol	5 %
	1-glycerol monolaurate	2 %
	Dipropylene glycol	20 %
	Carbopol	2 %
	NaOH	q.s.

Purified water to 100 %

The 1-glycerol monolaurate was dissolved in the propylene glycol and the 1,5-pentanediol added. The carbopol was dissolved in water and mixed with the pentanediol solution, after which the viscosity was adjusted with an aqueous solution of NaOH.

This Example shows the typical constituents for a composition in the form of a gel and additionally in which 1-glycerol monolaurate is used as the sole monoglyceride.

The effectiveness of the gel corresponds to the effectiveness of the cream in Example 2.

Example 4

Fusidic acid in crystalline monoglycerides with an enhanced antimicrobial spectrum.

A composition with the following composition by weight was prepared:

	Fusidic acid	2 %
25	1-glycerol monolaurate	7 %
	1-glycerol monomyristate	21 %
	Purified water	to 100 %

During preparation, the monoglycerides were mixed with water and heating carried out to 70°C. The fusidic acid was then added, after which mixing was effected for 15 minutes at said temperature after which the resultant mixture was cooled at a rate of 1 - 3°C per minute to room temperature. Optional buffering of the pH and addition of thickening agents is done prior to cooling of the mixture.

An MIC value determination showed that the composition had

a lower MIC value than a commercial preparation of fusidic acid and that the antimicrobial spectrum was enhanced with regard to effectiveness on fungi.

5 Example 5

Antibacterial preparation containing econazole nitrate

10 A composition with the following composition by weight was prepared:

	Econazole nitrate	1	%
	1-glycerol monolaurate	7	%
15	1-glycerol monomyristate	21	%
	Purified water	to 100	%

The composition was prepared in the same manner as the composition of Example 4. Investigation of the
20 antimicrobial properties of the composition showed it to have antimicrobial effectiveness in contrast to a commercial product containing only econazole nitrate in the same concentration as the above preparation.

25 Example 6

Acne preparation containing tinidazole

30 A composition with the following composition by weight was prepared:

	Tinidazole	2	%
	1-glycerol monolaurate	5	%
	1-glycerol monomyristate	15	%
35	Propylene glycol	30	%
	Carbomer	0.5	%
	Purified water	to 100	%

The composition was prepared as in Example 4. It proved to have a powerful killing effect on fungi and bacteria, both aerobic and anaerobic, in contrast to aqueous solutions of tinidazole which are not effective against aerobic bacteria. In particular, the product is usable for disease states such as acne and seborrhoea.

Example 7

Microbiologically stabilized preparation containing urea

The following composition was prepared:

Urea	10	%
Phosphate buffer to pH 6.0	q.s.	
1-glycerol monolaurate	7	%
1-glycerol monomyristate	21	%
Purified water	to 100	%

The composition was prepared as in Example 4.

In microbiological load tests, it proved to have bacteriocidal and fungistatic properties. Both the bactericidal and fungistatic effects are more powerful than that provided by either the monoglycerides or urea individually.

Monolaurin can also be added to the emulsion in order to enhance the antimicrobial effectiveness, either in solution, in an aqueous phase, in crystalline form in an aqueous phase or dissolved in a lipid phase. It is important that crystalline monolaurin is added at a temperature below 35°C in order to avoid effects on the formation of the emulsion. Smaller amounts of monolaurin can be added at the emulsifying temperature (around 70°C).

Lipid or water soluble medicaments are then dissolved in the phase in which their solubility is highest. This is exemplified in Examples 8 and 9 below.

5 Example 8

Emulsion containing urea

10 An emulsion of the water-in-oil type with the following composition was prepared:

	Liquid paraffin	20	g
	Monolaurin	1	g
	Sorbitan monooleate	1	g
15	Water	50	g
	Polyoxyethylene (20) stearate	1	g
	Carbomer	0.5	g

Components 1, 2 and 3 were mixed and heated to 70°C.
 20 Separately, components 4, 5 and 6 were blended and heated to 70°C. The mixtures were homogenized and cooled to room temperature. After cooling, urea was added in an amount corresponding to 1 - 10 g in 26.5 g of water.

25 Example 9

Emulsion containing monolaurin

30 An emulsion of the oil-in-water type was prepared from the following ingredients:

	<u>A</u>		
	Liquid paraffin	20	g
	Sorbitan monooleate	1	g
35	Water	20	g
	Polyoxyethylene (20) stearate	1	g
	Carbomer	0.5	g

B

	Monolaurin	5 g
	Monomyristin	15 g
5	Water	37.5 g

Emulsion A was prepared as earlier described and cooled to around 35°C. Lipid crystal dispersion B was prepared in the following manner. The components were mixed at 70°C and after 15 minutes the mixture was cooled at a rate of 1 - 3° per minute. When crystallisation was complete, A and B were mixed at 30 - 35°C.

Over and above what has been said earlier regarding the components forming part of the composition, it should be added that the practice of the invention has shown that it does not matter in what form the monoglycerides are present. In other words, it has proven that the monoglyceride has the same or a similar potentiating effect on antimicrobial properties whether it is present in solution, in amorphous form or in crystalline form. In this respect also, the invention thereby represents a valuable contribution to the art.

25

30

35



This Sheet Has Been Replaced



18

CLAIMS

1. An antimicrobial composition, characterized in that
it comprises an antimicrobially effective amount of a
5 combination of:

A) a monoglyceride of lauric acid, a monoglyceride
of monomyristic acid or a mixture of these
monoglycerides, and

10

B) at least one chemical substance selected from the
following groups:

i) a local anaesthetic of the amide type

15

ii) carbamide

iii) an antibacterial substance in the form of
a steroid antibiotic, an imidazole
derivative or a nitroimidazole derivative,
and

20

iv) a diol with 3 - 6 carbon atoms, and

C) optionally, a conventional physiologically
acceptable carrier and/or physiologically acceptable
additives.

25

2. A composition according to claim 1, characterized in
that said local anaesthetic of the amide type is selected
from the group consisting of lidocaine, prilocaine,
mepivacaine, cinchocaine, bupivacaine and etidocaine.

30

3. A composition according to claim 2, characterized in
that said local anaesthetic of the amide type is
lidocaine.

35 4. A composition according to any one of claims 1 to 3,
characterized in that the antibacterial substance in the
form of a steroid antibiotic is fusidic acid or



This Sheet Has Been Replaced



19

cephalosporin P.

5. A composition according to claim 4, characterized in that the antibacterial substance is fusidic acid.
- 5
6. A composition according to any one of claims 1 to 5, characterized in that the imidazole derivative is econazole nitrate or mikonazole nitrate.
- 10 7. A composition according to claim 6, characterized in that the imidazole derivative is econazole nitrate.
8. A composition according to any one of the preceding claims, characterized in that the nitroimidazole
- 15 derivative is tinidazole or metronidazole.
9. A composition according to any one of the preceding claims, characterized in that the diol with 3 - 6 carbon atoms is a pentanediol, preferably 1,5-pentanediol.
- 20 10 A composition according to any one of the preceding claims, characterized in that the monoglyceride A) is a monolaurate, preferably 1-glycerol monolaurate or 3-glycerol monolaurate.
- 25 11. A composition according to any one of the preceding claims, characterized in that it contains at least 0.2 ppm of the monoglyceride(s) A), calculated on the weight of the total composition.
- 30 12. A composition according to claim 11, characterized in that the concentration of A) is from 0.2 ppm to 30% calculated as weight on the weight of the total composition.
- 35 13. A composition according to any preceding claim, characterized in that it contains from 1 ppm to 20% of the



20

chemical substance(s) B), calculated as weight on the total weight of the composition, preferably from 1 ppm to 10%, alternatively from 0.1 to 20%, especially 0.1 to 10 weight percent.

5

14. A process for the preparation of an antimicrobial composition as defined in any one of claims 1 to 13 characterized by heating the monoglyceride(s) A) to a temperature over the transition or conversion temperature of the lipid, optionally in the presence of a carrier and/or additives C), adding the substance(s) B) to the heated mixture until conversion has taken place, and cooling the substance(s) B-containing mixture to around room temperature.

15

15. A process according to claim 13, characterized in that the cooling is carried out at a rate of around 1 - 3°C per minute.

20

16. A composition according to any one of claims 1 to 13 for use as an antimicrobial pharmaceutical composition.

17. Use of a composition as defined in any one of claims 1 to 13 for the preparation of a dermatological pharmaceutical preparation for combatting bacteria and fungi.

25

18. Use of a composition as defined in any one of claims 1 to 13 as a preservative additive in a cosmetic product, a food product, or a medical product.

30

35

ABSTRACT

An antimicrobial composition comprising an antimicrobially effective amount of a combination of:

5

A) a monoglyceride of lauric acid, a monoglyceride of monomyristic acid or a mixture of these monoglycerides, and

10

B) at least one chemical substance selected from the following group:

i) local anaesthetics of the amide type

ii) carbamide

15

iii) antibacterial substances in the form of steroid antibiotics, imidazole derivatives or nitroimidazole derivatives, and

iv) diols with 3 - 6 carbon atoms, and

20

C) optionally, a conventional physiologically acceptable carrier and/or physiologically acceptable additives.

A process for the preparation of this composition by
25 heating A) to the transition temperature of the lipid, adding B), and optionally C), and cooling the mixture to form a solid lipid crystal composition.

Use of the composition for the preparation of a
30 dermatological preparation for combatting bacteria and fungi or as a preservative additive in a cosmetic product, a food product, or a medical product.

35

ABSTRACT

POTENTIATION OF ANTIMICROBIAL EFFECTS

An antimicrobial composition comprising an antimicrobially effective amount of a combination of:

- 5
- A) a monoglyceride of lauric acid, a monoglyceride of monomyristic acid or a mixture of these monoglycerides, and
- 10 B) at least one chemical substance selected from the following group:
- i) local anaesthetics of the amide type
- ii) carbamide
- 15 iii) antibacterial substances in the form of steroid antibiotics, imidazole derivatives or nitroimidazole derivatives, and
- iv) diols with 3 - 6 carbon atoms, and
- 20 C) optionally, a conventional physiologically acceptable carrier and/or physiologically acceptable additives.

A process for the preparation of this composition by

25 heating A) to the transition temperature of the lipid, adding B), and optionally C), and cooling the mixture to form a solid lipid crystal composition.

Use of the composition for the preparation of a

30 dermatological preparation for combatting bacteria and fungi or as a preservative additive in a cosmetic product, a food product, or a medical product.

REPLACEMENT SHEET

30 AUG 1996



CLAIMS

1. An antimicrobial composition, characterized in that it comprises an antimicrobially effective amount of a combination of:

- A) a monoglyceride of lauric acid, a monoglyceride of myristic acid or a mixture of these monoglycerides, and
- B) at least one chemical substance selected from urea, fusidic acid, econazole nitrate (1-(2,4-dichloro-beta-(p-chlorobenzyloxy)-phenethyl)imidazol nitrate), mikonazole nitrate (1-((2,4-dichlorophenyl)-2-((2,4-dichlorophenyl)methoxy)-ethyl)-1H-imidazol nitrate), tinidazole (1-(2-(ethylsulphonyl)ethyl)-2-methyl-5-nitroimidazole), metronidazole (1-(2-hydroxyethyl)-2-methyl-5-nitroimidazole and pentanediol, and
- C) optionally, a conventional physiologically acceptable carrier and/or physiologically acceptable additives, the concentration of A) being from 0.2 ppm to 30% and of B) being from 1 ppm to 20%, in both cases calculated as weight of the total weight of the composition.

2. A composition according to claim 1, characterized in that the chemical substance B) is fusidic acid.

3. A composition according to claim 1, characterized in that the chemical substance B) is econazole nitrate.

4. A composition according to claim 1, characterized in that the chemical substance B) is tinidazole.

5. A composition according to claim 1, characterized in that the chemical substance B) is urea.



6. A composition according to claim 1, characterized in that the pentanediol is 1,5-pentanediol.

7. A composition according to any one of the preceding claims, characterized in that the monoglyceride A)
5 is a monolaurate, preferably 1-glycerol monolaurate or 3-glycerol monolaurate.

8. A composition according to any one of the preceding claims, characterized in that it contains from 1 ppm to 10% or alternatively from 0.1 to 20%, especially
10 from 0.1 to 10%, of the chemical substance(s) B), calculated as weight on the total weight of the composition.


9. A process for the preparation of an antimicrobial composition as defined in any one of claims 1 to 8, characterized by heating the monoglyceride(s) A) to a
15 temperature over the transition or conversion temperature of the lipid, optionally in the presence of a carrier and/or additives C), adding the substance(s) B) to the heated mixture until conversion has taken place, and cooling the substance(s) B-containing mixture to around
20 room temperature.

10. A process according to claim 9, characterized in that the cooling is carried out at a rate of around 1-3°C per minute.

11. A composition according to any one of claims 1
25 to 8 for use as an antimicrobial pharmaceutical composition.

12. Use of a composition as defined in any one of claims 1 to 8 for the preparation of a dermatological pharmaceutical preparation for combatting bacteria and
30 fungi.

13. Use of a composition as defined in any one of claims 1 to 8 as a preservative additive in a cosmetic product, a food product, or a medical product.



9603104-2

18

CLAIMS

1. An antimicrobial composition, characterized in that it comprises an antimicrobially effective amount of a combination of:

A) a monoglyceride of lauric acid, a monoglyceride of myristic acid or a mixture of these monoglycerides, and

10

B) at least one chemical substance selected from urea, fusidic acid, econazole nitrate (1-(2,4-dichloro-beta-(p-chlorobenzyloxy)-phenethyl)imidazol nitrate), mikonazole nitrate (1-((2(2,4-dichlorophenyl)-2((2,4-dichlorophenyl)methoxy)-ethyl)-1H-imidazol nitrate), tinidazole (1-(2-(ethylsulphonyl)ethyl)-2-methyl-5-nitroimidazole), metronidazole (1-(2-hydroxyethyl)-2-methyl-5-nitroimidazole and pentanediol, and

15

20

C) optionally, a conventional physiologically acceptable carrier and/or physiologically acceptable additives, the concentration of A) being from 0.2 ppm to 30% and of B) being from 1 ppm to 20%, in both cases calculated as weight of the total weight of the composition.

25

2. A composition according to claim 1, characterized in that the chemical substance B) is fusidic acid.

3. A composition according to claim 1, characterized in that the chemical substance B) is econazole nitrate.

30

4. A composition according to claim 1, characterized in that the chemical substance B) is tinidazole.

5. A composition according to claim 1, characterized in that the chemical substance B) is urea.

35



6. A composition according to claim 1, characterized in that the pentanediol is 1,5-pentanediol.

7. A composition according to any one of the preceding claims, characterized in that the monoglyceride A) is a monolaurate, preferably 1-glycerol monolaurate or 3-glycerol monolaurate..

8. A composition according to any one of the preceding claims, characterized in that it contains from 1 ppm to 10% or alternatively from 0.1 to 20%, especially from 0.1 to 10%, of the chemical substance(s) B), calculated as weight on the total weight of the composition.

9. A process for the preparation of an antimicrobial composition as defined in any one of claims 1 to 8, characterized by heating the monoglyceride(s) A) to a temperature over the transition or conversion temperature of the lipid, optionally in the presence of a carrier and/or additives C), adding the substance(s) B) to the heated mixture until conversion has taken place, and cooling the substance(s) B-containing mixture to around room temperature.

10. A process according to claim 9, characterized in that the cooling is carried out at a rate of around 1-3°C per minute.

11. A composition according to any one of claims 1 to 8 for use as an antimicrobial pharmaceutical composition.

12. Use of a composition as defined in any one of claims 1 to 8 for the preparation of a dermatological pharmaceutical preparation for combatting bacteria and fungi.

13. Use of a composition as defined in any one of claims 1 to 8 as a preservative additive in a cosmetic product, a food product, or a medical product.

