


NICOTINAMIDE DERIVATES USEFUL AS P38 INHIBITORS

This invention relates to novel compounds and their use as pharmaceuticals, particularly as p38 kinase inhibitors, for the treatment of conditions or disease states mediated by p38 kinase activity or mediated by cytokines produced by the activity of p38 kinase.

We have now found a group of novel compounds that are inhibitors of p38 kinase.

According to the invention there is provided a compound of formula (I):



(I)

wherein

R^1 is selected from hydrogen, C_{1-6} alkyl optionally substituted by up to three groups selected from C_{1-6} alkoxy, halogen and hydroxy, C_{2-6} alkenyl, C_{3-7} cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, phenyl optionally substituted by up to three groups selected from R^5 and R^6 , and heteroaryl optionally substituted by up to three groups selected from R^5 and R^6 ,

R^2 is selected from hydrogen, C_{1-6} alkyl and $-(CH_2)_q$ - C_{3-7} cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups,

or $(CH_2)_m$ R^1 and R^2 , together with the nitrogen atom to which they are bound, form a four- to six-membered heterocyclic ring optionally substituted by up to three C_{1-6} alkyl groups;

R^3 is chloro or methyl;

R^4 is the group $-NH-CO-R^7$ or $-CO-NH-(CH_2)_q$ - R^8 ;

R^5 is selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_q$ - C_{3-7} cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, $-CONR^9R^{10}$, $-NHCOR^{10}$, $-SO_2NHR^9$, $-(CH_2)_5$ $NHSO_2R^{10}$, halogen, CN, OH, $-(CH_2)_5$ $NR^{11}R^{12}$, and trifluoromethyl;

R^6 is selected from C_{1-6} alkyl, C_{1-6} alkoxy, halogen, trifluoromethyl and $-(CH_2)_5$ $NR^{11}R^{12}$;

R^7 is selected from hydrogen, C_{1-6} alkyl, $-(CH_2)_q-C_{3-7}$ cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, trifluoromethyl, $-(CH_2)_r$ heteroaryl optionally substituted by R^{13} and/or R^{14} , and $-(CH_2)_r$ phenyl optionally substituted by R^{13} and/or R^{14} ;

5 R^8 is selected from hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, $CONHR^9$, phenyl optionally substituted by R^{13} and/or R^{14} , and heteroaryl optionally substituted by R^{13} and/or R^{14} ;

R^9 and R^{10} are each independently selected from hydrogen and C_{1-6} alkyl, or R^9 and R^{10} , together with the nitrogen atom to which they are bound, form

10 a five- to six-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and $N-R^{15}$, wherein the ring may be substituted by up to two C_{1-6} alkyl groups;

R^{11} is selected from hydrogen, C_{1-6} alkyl and $-(CH_2)_q-C_{3-7}$ cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups,

15 R^{12} is selected from hydrogen and C_{1-6} alkyl, or R^{11} and R^{12} , together with the nitrogen atom to which they are bound, form a five or six-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and $N-R^{15}$;

20 R^{13} is selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_q-C_{3-7}$ cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, $-CONR^9R^{10}$, $-NHCOR^{10}$, halogen, CN, $-(CH_2)_sNR^{11}R^{12}$, trifluoromethyl, phenyl optionally substituted by one or more R^{14} groups and heteroaryl optionally substituted by one or more R^{14} groups;

R^{14} is selected from C_{1-6} alkyl, C_{1-6} alkoxy, halogen, trifluoromethyl and $-NR^{11}R^{12}$;

25 R^{15} is selected from hydrogen and methyl;

X and Y are each independently selected from hydrogen, methyl and halogen;

Z is halogen;

m is selected from 0, 1, 2, 3 and 4, wherein each carbon atom of the resulting carbon chain may be optionally substituted with up to two groups selected

30 independently from C_{1-6} alkyl and halogen;


n is selected from 0, 1 and 2;

q is selected from 0, 1 and 2;

r is selected from 0 and 1; and

s is selected from 0, 1, 2 and 3.

35 According to a further embodiment of the invention there is provided a compound of formula (IA):



(IA)

wherein R^1 , R^2 , R^3 , R^4 and m are as defined above.

According to one embodiment of the present invention, R^1 is selected from hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, phenyl optionally substituted by R^5 and/or R^6 , and heteroaryl optionally substituted by R^5 and/or R^6 , and R^2 is selected from hydrogen, C_{1-6} alkyl and $-(CH_2)_q-C_{3-7}$ cycloalkyl.

In a preferred embodiment, R^1 is selected from C_{1-6} alkyl, for example methyl, ethyl, n-propyl, isopropyl, 1-methylpropyl, 1,1-dimethylpropyl, 2,2-dimethylpropyl, 1-ethyl-1-methyl-propyl, n-butyl, isobutyl, 3-methylbutyl, 1,1-dimethylbutyl, 1,3-dimethylbutyl, 3,3-dimethylbutyl, 2-pentyl or 1-methylpentyl, optionally substituted by up to three groups selected from C_{1-6} alkoxy, in particular C_{1-4} alkoxy groups such methoxy or t-butoxy, halogen, in particular fluorine, and hydroxy; C_{2-6} alkenyl, for example C_{4-6} alkenyl such as 3-methylbut-2-enyl or 1,1-dimethylbut-2-enyl; C_{3-7} cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, for example cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, in particular cyclopropyl, optionally substituted by one or two C_{1-4} alkyl groups such as methyl or ethyl; phenyl optionally substituted by up to three groups selected from R^5 and R^6 , for example phenyl optionally substituted by up to three substituents, for example one or two substituents, such as C_{1-4} alkyl, in particular methyl, C_{1-4} alkoxy, in particular methoxy, halogen, in particular fluorine or chlorine, trifluoromethyl, $-(CH_2)_sNR^{11}R^{12}$ or $-(CH_2)_sNHSO_2R^{10}$, located on any position on the ring; heteroaryl optionally substituted by up to three groups selected from R^5 and R^6 , for example heteroaryl optionally substituted by one or two substituents, in particular a 5-membered heteroaryl such as furyl, thienyl or thiazolyl optionally substituted by C_{1-4} alkyl, in particular methyl. In a particularly preferred embodiment, R^1 is C_{1-6} alkyl, for example C_{2-5} alkyl, such as ethyl, n-propyl, isopropyl, 1-methylpropyl, 1,1-dimethylpropyl, 2,2-dimethylpropyl, n-butyl, isobutyl, 3-methylbutyl or 2-pentyl.

In another preferred embodiment, R^1 is selected from C_{3-7} cycloalkyl, phenyl optionally substituted by R^5 and/or R^6 , and heteroaryl optionally substituted by R^5 and/or R^6 . In a more preferred embodiment, R^1 is selected from C_{3-6} cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, in particular cyclopropyl, and phenyl optionally substituted by R^5 and/or R^6 . The phenyl may be optionally substituted by one or two substituents, located on any position on the phenyl ring.

Preferred substituents for the phenyl include C₁₋₄alkoxy, in particular methoxy, -(CH₂)₈NR¹¹R¹², and -(CH₂)₈NHSO₂R¹⁰.

In another preferred embodiment, R¹ is selected from C₁₋₆alkyl, for example n-propyl, 1-methylpropyl, isobutyl, 3-methylbutyl or 2,2-dimethylpropyl, and C₃₋₇cycloalkyl optionally substituted by one or more C₁₋₆alkyl groups, for example cyclopropyl optionally substituted by one or two methyl groups.

In a further preferred embodiment, R¹ is selected from C₁₋₆alkyl, for example methyl, ethyl, n-propyl, isopropyl, 1,1-dimethylpropyl, 1-ethyl-1-methyl-propyl, n-butyl, isobutyl, 1,1-dimethylbutyl, 1,3-dimethylbutyl, 3,3-dimethylbutyl, 2-pentyl or 1-methylpentyl, optionally substituted by up to three groups selected from C₁₋₆alkoxy, in particular C₁₋₄alkoxy groups such as methoxy or t-butoxy, halogen, in particular fluorine, and hydroxy; C₂₋₆alkenyl, for example C₄₋₆alkenyl such as 3-methylbut-2-enyl or 1,1-dimethylbut-2-enyl; C₃₋₇cycloalkyl optionally substituted by one or more C₁₋₆alkyl groups, for example, cyclopropyl, cyclopentyl or cyclohexyl, optionally substituted by one or two ethyl groups; phenyl optionally substituted by up to three groups selected from R⁵ and R⁶, for example phenyl optionally substituted by up to three substituents such as C₁₋₄alkyl, in particular methyl, C₁₋₄alkoxy, in particular methoxy, halogen, in particular fluorine or chlorine and trifluoromethyl, located on any position on the ring; heteroaryl optionally substituted by up to three groups selected from R⁵ and R⁶, in particular a 5-membered heteroaryl such as furyl, thienyl or thiazolyl optionally substituted by C₁₋₄alkyl, in particular methyl.

In a preferred embodiment, R² is selected from hydrogen; C₁₋₄alkyl, in particular methyl, ethyl, isopropyl or isobutyl; and -(CH₂)_q-C₃₋₆cycloalkyl, in particular cyclopropyl, -CH₂-cyclopentyl, -(CH₂)₂-cyclopentyl or cyclohexyl.

In another preferred embodiment, R² is selected from hydrogen, C₁₋₄alkyl and -CH₂-cyclopropyl. More preferably R² is hydrogen.

In a further preferred embodiment, (CH₂)_mR¹ and R², together with the nitrogen atom to which they are bound, form a four- to six-membered heterocyclic ring optionally substituted by up to three C₁₋₆alkyl groups, in particular an azetidiny, pyrrolidiny or piperidiny ring optionally substituted by one or two methyl, ethyl or propyl groups.

In a preferred embodiment, R³ is methyl.

In a preferred embodiment, R⁴ is the group -CO-NH-(CH₂)_q-R⁸.

In one embodiment of the present invention, R⁵ is selected from C₁₋₆alkyl, C₁₋₆alkoxy, -(CH₂)_q-C₃₋₇cycloalkyl, -CONR⁹R¹⁰, -NHCOR¹⁰, -SO₂NHR⁹, -(CH₂)₈NHSO₂R¹⁰, halogen, CN, OH, -(CH₂)₈NR¹¹R¹², and trifluoromethyl.

In a preferred embodiment, R⁵ is selected from C₁₋₄alkyl, in particular methyl; C₁₋₄alkoxy, in particular methoxy; -(CH₂)₈NHSO₂R¹⁰; halogen, in particular chlorine or fluorine; -(CH₂)₈NR¹¹R¹²; and trifluoromethyl.

In another preferred embodiment, R⁵ is selected from C₁₋₄alkoxy, in particular methoxy, -(CH₂)₈NR¹¹R¹², and -(CH₂)₈NHSO₂R¹⁰.

In a further preferred embodiment, R⁵ is selected from C₁₋₄alkyl, in particular methyl; C₁₋₄alkoxy, in particular methoxy; halogen, in particular chlorine or fluorine; and trifluoromethyl.

In a preferred embodiment, R^6 is selected from C_{1-4} alkyl, in particular methyl, ethyl or propyl; C_{1-4} alkoxy, in particular methoxy; halogen, in particular chlorine or fluorine; and trifluoromethyl.

In a further preferred embodiment, R^6 is C_{1-4} alkoxy, in particular methoxy.

5 In one embodiment of the present invention, R^7 is selected from hydrogen, C_{1-6} alkyl, $-(CH_2)_q-C_{3-7}$ cycloalkyl, trifluoromethyl, $-(CH_2)_r$ heteroaryl optionally substituted by R^{13} and/or R^{14} , and $-(CH_2)_r$ phenyl optionally substituted by R^{13} and/or R^{14} .

10 In a preferred embodiment, R^7 is selected from C_{1-6} alkyl, $-(CH_2)_q-C_{3-7}$ cycloalkyl, trifluoromethyl, $-(CH_2)_r$ heteroaryl optionally substituted by R^{13} and/or R^{14} , and $-(CH_2)_r$ phenyl optionally substituted by C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_q-C_{3-7}$ cycloalkyl, $-CONR^9R^{10}$, $-NHCOR^{10}$, halogen, CN, trifluoromethyl, phenyl optionally substituted by one or more R^{14} groups and/or heteroaryl optionally substituted by one or more R^{14} groups. In another preferred embodiment, R^7 is
15 selected from C_{1-4} alkyl, $-(CH_2)_q-C_{3-6}$ cycloalkyl, trifluoromethyl, $-(CH_2)_r$ heteroaryl optionally substituted by R^{13} and/or R^{14} , and $-(CH_2)_r$ phenyl optionally substituted by C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_q-C_{3-7}$ cycloalkyl, $-CONR^9R^{10}$, $-NHCOR^{10}$, halogen, CN, trifluoromethyl, phenyl optionally substituted by one or more R^{14} groups and/or heteroaryl optionally substituted by one or more R^{14} groups. In a more
20 preferred embodiment, R^7 is $-(CH_2)_r$ heteroaryl optionally substituted by R^{13} and/or R^{14} , in particular a five or six-membered heteroaryl containing at least one heteroatom selected from oxygen, nitrogen and sulfur, for example, pyridinyl optionally substituted by $-NR^{11}R^{12}$, furyl or thiophenyl.

25 In one embodiment of the present invention, R^8 is selected from hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, $CONHR^9$, phenyl optionally substituted by R^{13} and/or R^{14} , and heteroaryl optionally substituted by R^{13} and/or R^{14} .

In a preferred embodiment, R^8 is selected from C_{3-7} cycloalkyl, $CONHR^9$, heteroaryl optionally substituted by R^{13} and/or R^{14} , and phenyl optionally substituted by C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_q-C_{3-7}$ cycloalkyl, $-CONR^9R^{10}$, $-NHCOR^{10}$,
30 halogen, CN, trifluoromethyl, phenyl optionally substituted by one or more R^{14} groups and/or heteroaryl optionally substituted by one or more R^{14} groups. In another preferred embodiment, R^8 is selected from C_{3-7} cycloalkyl, heteroaryl optionally substituted by R^{13} and/or R^{14} , and phenyl optionally substituted C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_q-C_{3-7}$ cycloalkyl, $-CONR^9R^{10}$, $-NHCOR^{10}$, halogen, CN,
35 trifluoromethyl, phenyl optionally substituted by one or more R^{14} groups and/or heteroaryl optionally substituted by one or more R^{14} groups. In a more preferred embodiment, R^8 is selected from C_{3-6} cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, in particular cyclopropyl, heteroaryl optionally substituted by R^{13} and/or R^{14} , in particular a five or six-membered heteroaryl containing at least
40 one heteroatom selected from nitrogen and sulfur, for example, thiazolyl or thiadiazolyl, and phenyl optionally substituted by heteroaryl. In a particularly preferred embodiment, R^8 is selected from C_{3-6} cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, in particular cyclopropyl.

In a preferred embodiment, R^9 is selected from hydrogen and C_{1-4} alkyl.

In a preferred embodiment, R^{10} is selected from hydrogen and C_{1-4} alkyl, in particular methyl.

In one embodiment, R^{11} is selected from hydrogen, C_{1-6} alkyl and $-(CH_2)_q-C_{3-7}$ cycloalkyl optionally substituted by C_{1-6} alkyl.

5 In a preferred embodiment, R^{11} and R^{12} , together with the nitrogen atom to which they are bound, form a five or six-membered heterocyclic ring optionally further containing one additional heteroatom N- R^{15} .

In one embodiment of the present invention, R^{13} is selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_q-C_{3-7}$ cycloalkyl, $-CONR^9R^{10}$, $-NHCOR^{10}$, halogen, CN, -
10 $(CH_2)_sNR^{11}R^{12}$, trifluoromethyl, phenyl optionally substituted by one or more R^{14} groups and heteroaryl optionally substituted by one or more R^{14} groups;

In a preferred embodiment, R^{13} is selected from C_{1-4} alkyl, in particular methyl, C_{1-4} alkoxy, in particular methoxy, halogen, $-(CH_2)_sNR^{11}R^{12}$, phenyl
15 optionally substituted by one or more R^{14} groups and heteroaryl optionally substituted by one or more R^{14} groups. In a more preferred embodiment, R^{13} is selected from -
 $(CH_2)_sNR^{11}R^{12}$ and heteroaryl optionally substituted by one or more R^{14} groups, in particular a five or six-membered heteroaryl containing at least one nitrogen atom, for example, pyridyl.

In a preferred embodiment R^{14} is selected from from C_{1-4} alkyl, in particular
20 methyl, C_{1-4} alkoxy, in particular methoxy, and $-NR^{11}R^{12}$.

In a preferred embodiment, R^{15} is methyl.

In a preferred embodiment, X and Y are each independently selected from hydrogen, chlorine and fluorine. In a further preferred embodiment, X is fluorine. In
25 another preferred embodiment, Y is hydrogen.

In a preferred embodiment, Z is fluorine.

In one embodiment of the present invention, m is selected from 0, 1, 2, 3 and
4. In another embodiment of the present invention, m is selected from 0, 1, 2, 3 and 4, wherein each carbon atom of the resulting carbon chain may be optionally substituted with up to two groups selected independently from C_{1-6} alkyl.

30 In a preferred embodiment, m is selected from 0, 1, 2 and 3. In a further preferred embodiment, m is selected from 0, 1 and 2, in particular 0 and 1. When the carbon chain of m is substituted, these substituents are preferably one or two methyl groups or fluorine atoms. In one embodiment, the substituents are preferably one or two methyl groups. In another embodiment, the substituents are preferably one or two
35 fluorine atoms.

In a preferred embodiment, n is selected from 0 and 1. In particular, n is 0.

In a preferred embodiment, q is selected from 0 and 1. In particular, q is 0.

In a preferred embodiment, r is 0.

In a preferred embodiment, s is selected from 0 and 1.

40 It is to be understood that the present invention covers all combinations of particular and preferred groups described hereinabove.

Particular compounds according to the invention include those mentioned in the Examples. Specific examples which may be mentioned include:

6-(5-cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-cyclopropylmethyl-
45 nicotinamide;

- 6-(5-cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-(1-cyclopropylethyl)-nicotinamide;
 6-(5-cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-(2,2-dimethylpropyl)-nicotinamide;
 5 6-(5-cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-(2-methylpropyl)-nicotinamide; and
 6-(5-cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-(1-methylpropyl)-nicotinamide.
- Futher specific examples which may be mentioned include:
- 10 6-(5-cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-cyclobutylmethyl-nicotinamide;
 6-(5-cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-cyclobutyl-nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,4,5-trifluorobenzyl)nicotinamide;
 15 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,5-difluorobenzyl)nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3,4-difluorobenzyl)nicotinamide;
 N-(3-chlorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 20 N-(4-chlorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 N-(3-chloro-2-fluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 25 N-(2-chloro-3,6-difluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,3-difluoro-4-methylbenzyl)nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,3,5-trifluorobenzyl)nicotinamide;
 30 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3-fluoro-4-methylbenzyl)nicotinamide;
 N-(5-chloro-2-fluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 35 N-(2-chlorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(4-fluorobenzyl)nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,3,4-trifluorobenzyl)nicotinamide;
 40 N-benzyl-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[3-(trifluoromethyl)benzyl]nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(1,1-dimethylbutyl)nicotinamide;
 45

- N-(4-chloro-2-fluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[4-(trifluoromethyl)benzyl]nicotinamide;
 5 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[(5-methyl-2-furyl)methyl]nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,3-difluorobenzyl)nicotinamide;
 N-(3-chloro-4-fluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 10 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(4-methylbenzyl)nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[(3-methylthien-2-yl)methyl]nicotinamide;
 15 N-(3-chloro-2,6-difluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(1-ethyl-1-methylpropyl)nicotinamide;
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2-fluorobenzyl)nicotinamide;
 20 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(tert-pentyl)nicotinamide; and
 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3-methylbenzyl)nicotinamide.

25 As used herein, the term "alkyl" refers to straight or branched hydrocarbon chains containing the specified number of carbon atoms. For example, C₁₋₆alkyl means a straight or branched alkyl containing at least 1, and at most 6, carbon atoms. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, isobutyl, isopropyl and t-butyl. A C₁₋₄alkyl group is
 30 preferred, for example methyl, ethyl, isopropyl or t-butyl. The said alkyl groups may be optionally substituted with one or more fluorine atoms for example, trifluoromethyl.

As used herein, the term "alkenyl" refers to straight or branched hydrocarbon chains containing the specified number of carbon atoms and containing at least one
 35 double bond. For example, C₂₋₆alkenyl means a straight or branched alkenyl containing at least 2, and at most 6, carbon atoms and containing at least one double bond. Examples of "alkenyl" as used herein include, but are not limited to ethenyl, propenyl, 3-methylbut-2-enyl and 1,1-dimethylbut-2-enyl.

As used herein, the term "alkoxy" refers to a straight or branched chain alkoxy
 40 group, for example, methoxy, ethoxy, propoxy, prop-2-oxy, butoxy, but-2-oxy, 2-methylprop-1-oxy, 2-methylprop-2-oxy, pentoxy, or hexyloxy. A C₁₋₄alkoxy group is preferred, for example methoxy or ethoxy.

As used herein, the term "cycloalkyl" refers to a non-aromatic hydrocarbon ring containing the specified number of carbon atoms which may optionally contain
 45 up to one double bond. For example, C₃₋₇cycloalkyl means a non-aromatic ring

containing at least three, and at most seven, ring carbon atoms. Examples of "cycloalkyl" as used herein include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. A C₃₋₆cycloalkyl group is preferred, for example, cyclopropyl, cyclopentyl or cyclohexyl. The said cycloalkyl groups may be optionally substituted with one or more C₁₋₆alkyl groups, for example one or two methyl groups. In one embodiment, the cycloalkyl groups may be optionally substituted by up to four C₁₋₆alkyl groups, for example one or two C₁₋₆alkyl groups, in particular one or two C₁₋₄alkyl groups such as methyl or ethyl.

As used herein, the terms "heteroaryl ring" and "heteroaryl" refer to a monocyclic five- to seven-membered unsaturated hydrocarbon ring containing at least one heteroatom independently selected from oxygen, nitrogen and sulfur. Preferably, the heteroaryl ring has five or six ring atoms. Examples of heteroaryl rings include, but are not limited to, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, imidazolyl, pyrazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl and triazinyl. The said ring may be optionally substituted by one or more substituents independently selected from C₁₋₆alkyl and oxy.

As used herein, the terms "heterocyclic ring" or "heterocyclyl" refer to a monocyclic three- to seven-membered saturated hydrocarbon ring containing at least one heteroatom independently selected from oxygen, nitrogen and sulfur. Preferably, the heterocyclyl ring has five or six ring atoms. Examples of heterocyclyl groups include, but are not limited to, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, piperidyl, piperazinyl, morpholino, tetrahydropyranyl, tetrahydrofuranyl, and thiomorpholino. The said ring may be optionally substituted by one or more substituents independently selected from C₁₋₆alkyl and oxy.

As used herein, the terms "halogen" or "halo" refer to the elements fluorine, chlorine, bromine and iodine. Preferred halogens are fluorine, chlorine and bromine. A particularly preferred halogen is fluorine or chlorine.

As used herein, the term "optionally" means that the subsequently described event(s) may or may not occur, and includes both event(s) which occur and events that do not occur.

As used herein, the term "substituted" refers to substitution with the named substituent or substituents, multiple degrees of substitution being allowed unless otherwise stated.

As used herein, the term "solvate" refers to a complex of variable stoichiometry formed by a solute (in this invention, a compound of formula (I) or a salt thereof) and a solvent. Such solvents for the purpose of the invention may not interfere with the biological activity of the solute. Examples of suitable solvents include water, methanol, ethanol and acetic acid. Preferably the solvent used is a pharmaceutically acceptable solvent. Examples of suitable pharmaceutically acceptable solvents include water, ethanol and acetic acid. All such solvates are included within the scope of the present invention.

Certain compounds of formula (I) may exist in stereoisomeric forms (e.g. they may contain one or more asymmetric carbon atoms or may exhibit cis-trans isomerism). The individual stereoisomers (enantiomers and diastereomers) and


mixtures of these are included within the scope of the present invention. The present invention also covers the individual isomers of the compounds represented by formula (I) as mixtures with isomers thereof in which one or more chiral centres are inverted. Likewise, it is understood that compounds of formula (I) may exist in tautomeric forms other than that shown in the formula and these are also included within the scope of the present invention.

Salts of the compounds of the present invention are also encompassed within the scope of the invention and may, for example, comprise acid addition salts resulting from reaction of an acid with a basic nitrogen atom present in a compound of formula (I).

Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention. Representative salts include the following salts: Acetate, Benzenesulfonate, Benzoate, Bicarbonate, Bisulfate, Bitartrate, Borate, Bromide, Calcium Edetate, Camsylate, Carbonate, Chloride, Clavulanate, Citrate, Dihydrochloride, Edetate, Edisylate, Estolate, Esylate, Fumarate, Gluceptate, Gluconate, Glutamate, Glycolylarsanilate, Hexylresorcinate, Hydrabamine, Hydrobromide, Hydrochloride, Hydroxynaphthoate, Iodide, Isethionate, Lactate, Lactobionate, Laurate, Malate, Maleate, Mandelate, Mesylate, Methylbromide, Methylnitrate, Methylsulfate, Monopotassium Maleate, Mucate, Napsylate, Nitrate, N-methylglucamine, Oxalate, Pamoate (Embonate), Palmitate, Pantothenate, Phosphate/diphosphate, Polygalacturonate, Potassium, Salicylate, Sodium, Stearate, Subacetate, Succinate, Tannate, Tartrate, Teoclate, Tosylate, Triethiodide, Trimethylammonium and Valerate. Other salts which are not pharmaceutically acceptable may be useful in the preparation of compounds of this invention and these form a further aspect of the invention.


The compounds of this invention may be made by a variety of methods, including standard chemistry. Any previously defined variable will continue to have the previously defined meaning unless otherwise indicated. Illustrative general synthetic methods are set out below and then specific compounds of the invention are prepared in the working Examples.

A compound of formula (I) may be prepared by reacting a compound of (II)



(II)


in which R^1 , R^2 , Z, m and n are as hereinbefore defined and W is halogen, in particular bromine or chlorine, with a compound of formula (III)



(III)


in which R^3 , R^4 , X and Y are as hereinbefore defined,
in the presence of a catalyst, for example tetrakis(triphenylphosphine)palladium.

- 5 A compound of formula (II) may readily be prepared from a corresponding acid compound of formula (IV)



(IV)

- 10 in which Z, W and n are as hereinbefore defined,
by converting the acid to an activated form of the acid, for example the acid chloride,
by treatment with, for example, thionyl chloride, and then reacting the activated acid
thus formed with an amine compound of formula (V)




15

(V)

in which R^1 , R^2 and m are as hereinbefore defined,
under amide forming conditions.

- 20 Suitable amide forming conditions are well known in the art and include
treating a solution of the acid of formula (IV), or the activated form thereof, in for
example acetone or dichloromethane, with an amine of formula (V) in the presence of
sodium carbonate.

A compound of formula (III) may be prepared by reacting a compound of
formula (VI)




25

(VI)

in which R^3 , R^4 , X and Y are as hereinbefore defined and hal is halogen, in particular
iodine,

with bis(pinnacolato)diboron, PdCl₂dppf and potassium acetate in a solvent such as DMF.

Alternatively, when R⁴ is -CO-NH-(CH₂)_q-R⁸, a compound of formula (III) may be prepared by reacting an acid compound of formula (VII)




(VII)

in which R³, hal, X and Y are as hereinbefore defined,

with bis(pinnacolato)diboron, PdCl₂dppf and potassium acetate in a solvent such as DMF, and then forming an amide by reaction with an amine compound of formula (V) as hereinbefore defined.


A compound of formula (I) may also be prepared by reacting a compound of formula (VIII)



(VIII)

with a compound of formula (III) as hereinbefore defined and then reacting the acid thus formed with an amine of formula (V) as hereinbefore defined, under amide forming conditions.

Additionally, a compound of formula (I) may be prepared by reacting a compound of (II) as hereinbefore defined with a compound of formula (IX)




(IX)

in which R³, R⁴, X and Y are as hereinbefore defined,

in the presence of a catalyst, for example tetrakis(triphenylphosphine)palladium.


For example, one general method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 1 below.



Scheme 1

- i. $\text{R}^7\text{CO}_2\text{H}$, HATU, DIPEA, DMF.
- ii. Bis(pinacolato)diboron, PdCl_2dppf , KOAc, DMF.
- 5 iii. SOCl_2 .
- iv. $\text{R}^1(\text{CH}_2)_m\text{R}^2\text{NH}$, Na_2CO_3 , acetone.
- v. Na_2CO_3 , tetrakis(triphenylphosphine)palladium, propan-2-ol.


For example, another general method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 2 below.



Scheme 2

- i. SOCl_2 .
- ii. $\text{R}^8(\text{CH}_2)_q\text{NH}_2$, Na_2CO_3 , acetone.
- 5 iii. Bis(pinacolato)diboron, PdCl_2dppf , KOAc, DMF.
- iv. SOCl_2 .
- v. $\text{R}^1(\text{CH}_2)_m\text{R}^2\text{NH}$, Na_2CO_3 , acetone.
- vi. Na_2CO_3 , tetrakis(triphenylphosphine)palladium, propan-2-ol.


For example, another general method for preparing the compounds of formula
 10 (I) comprises the reactions set out in Scheme 3 below.



Scheme 3

- i. Bis(pinacolato)diboron, PdCl_2dppf , KOAc, DMF.
- ii. $\text{R}^8(\text{CH}_2)_q\text{NH}_2$, HATU, DIPEA, DMF.
- 5 iii. SOCl_2 .
- iv. $\text{R}^1(\text{CH}_2)_m\text{R}^2\text{NH}$, Na_2CO_3 , DCM.
- v. Na_2CO_3 , tetrakis(triphenylphosphine)palladium, propan-2-ol.


For example, another general method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 4 below.



Scheme 4

- i. NaHCO₃, tetrakis(triphenylphosphine)palladium, propan-2-ol.
- 5 ii. R¹(CH₂)_mR²NH, HATU, DIPEA, DMF.

For example, a further general method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 5 below.



Scheme 5

- i. $SOCl_2$.
 - 5 ii. $R^8(CH_2)_qNH_2$, Na_2CO_3 , DCM.
 - iii. NaH , $n-BuLi$, THF, $(iPrO)_3B$.
 - iv. $SOCl_2$.
 - v. $R^1(CH_2)_mR^2NH$, Na_2CO_3 , DCM.
 - vi. $NaHCO_3$, tetrakis(triphenylphosphine)palladium, propan-2-ol.
- 10 Whilst it is possible for the compounds of the present invention to be administered as the new chemical, the compounds of formula (I) are conveniently administered in the form of pharmaceutical compositions. Thus, in another aspect of the invention, we provide a pharmaceutical composition comprising a compound of formula (I), in admixture with one or more pharmaceutically acceptable carriers,
- 15 diluents or excipients.

The compounds of formula (I) may be formulated for administration in any suitable manner. They may, for example, be formulated for topical administration or administration by inhalation or, more preferably, for oral, transdermal or parenteral administration. The pharmaceutical composition may be in a form such that it can

20 effect controlled release of the compounds of formula (I). A particularly preferred method of administration, and corresponding formulation, is oral administration.

For oral administration, the pharmaceutical composition may take the form of, and be administered as, for example, tablets (including sub-lingual tablets) and capsules (each including timed release and sustained release formulations), pills, powders, granules, elixirs, tinctures, emulsions, solutions, syrups or suspensions prepared by conventional means with acceptable excipients.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Powders are prepared by comminuting the compound to a suitable fine size and mixing with a similarly comminuted pharmaceutical carrier such as an edible carbohydrate, as, for example, starch or mannitol. Flavoring, preservative, dispersing and coloring agent can also be present.

Capsules can be made by preparing a powder mixture as described above, and filling formed gelatin sheaths. Glidants and lubricants such as colloidal silica, talc, magnesium stearate, calcium stearate or solid polyethylene glycol can be added to the powder mixture before the filling operation. A disintegrating or solubilizing agent such as agar-agar, calcium carbonate or sodium carbonate can also be added to improve the availability of the medicament when the capsule is ingested.

Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like.

Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like. Tablets are formulated, for example, by preparing a powder mixture, granulating or slugging, adding a lubricant and disintegrant and pressing into tablets. A powder mixture is prepared by mixing the compound, suitably comminuted, with a diluent or base as described above, and optionally, with a binder such as carboxymethylcellulose, an aliginate, gelatin, or polyvinyl pyrrolidone, a solution retardant such as paraffin, a resorption accelerator such as a quaternary salt and/or an absorption agent such as bentonite, kaolin or dicalcium phosphate. The powder mixture can be granulated by wetting with a binder such as syrup, starch paste, acacia mucilage or solutions of cellulosic or polymeric materials and forcing through a screen. As an alternative to granulating, the powder mixture can be run through the tablet machine and the result is imperfectly formed slugs broken into granules. The granules can be lubricated to prevent sticking to the tablet forming dies by means of the addition of stearic acid, a stearate salt, talc or mineral oil. The lubricated mixture is then compressed into tablets. The compounds of the present invention can also be combined with free flowing inert carrier and compressed into tablets directly without going through the granulating or slugging steps. A clear or opaque protective coating consisting of a sealing coat of shellac, a coating of sugar or polymeric material and a polish coating of wax can be provided. Dyestuffs can be added to these coatings to distinguish different unit dosages.

Oral fluids such as solution, syrups and elixirs can be prepared in dosage unit form so that a given quantity contains a predetermined amount of the compound. Syrups can be prepared by dissolving the compound in a suitably flavored aqueous solution, while elixirs are prepared through the use of a non-toxic alcoholic vehicle. Suspensions can be formulated by dispersing the compound in a non-toxic vehicle. Solubilizers and emulsifiers such as ethoxylated isostearyl alcohols and polyoxyethylene sorbitol ethers, preservatives, flavor additives such as peppermint oil or saccharin, and the like can also be added.

Where appropriate, dosage unit formulations for oral administration can be microencapsulated. The formulation can also be prepared to prolong or sustain the release as for example by coating or embedding particulate material in polymers, wax or the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

The compounds of the present invention can also be administered in the form of liposome emulsion delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

The present invention includes pharmaceutical compositions containing 0.1 to 99.5%, more particularly, 0.5 to 90% of a compound of the formula (I) in combination with a pharmaceutically acceptable carrier.

Likewise, the composition may also be administered in nasal, ophthalmic, otic, rectal, topical, intravenous (both bolus and infusion), intraperitoneal, intraarticular, subcutaneous or intramuscular, inhalation or insufflation form, all using forms well known to those of ordinary skill in the pharmaceutical arts.

For transdermal administration, the pharmaceutical composition may be given in the form of a transdermal patch, such as a transdermal iontophoretic patch.

For parenteral administration, the pharmaceutical composition may be given as an injection or a continuous infusion (e.g. intravenously, intravascularly or subcutaneously). The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. For administration by injection these

may take the form of a unit dose presentation or as a multidose presentation preferably with an added preservative. Alternatively for parenteral administration the active ingredient may be in powder form for reconstitution with a suitable vehicle.

5 The compounds of the invention may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly
10 soluble salt.

Alternatively the composition may be formulated for topical application, for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to
15 assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

20 For administration by inhalation the compounds according to the invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, tetrafluoroethane, heptafluoropropane, carbon dioxide or other suitable gas. In the
25 case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

The pharmaceutical compositions generally are administered in an amount
30 effective for treatment or prophylaxis of a specific condition or conditions. Initial dosing in human is accompanied by clinical monitoring of symptoms, such symptoms for the selected condition. In general, the compositions are administered in an amount of active agent of at least about 100 $\mu\text{g/kg}$ body weight. In most cases they will be administered in one or more doses in an amount not in excess of about 20 mg/kg body weight per day.
35 Preferably, in most cases, dose is from about 100 $\mu\text{g/kg}$ to about 5 mg/kg body weight, daily. For administration particularly to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.1 mg/kg to 10 mg/kg and typically around 1 mg/kg . It will be appreciated that optimum dosage will be determined by standard methods for each treatment modality and indication, taking into
40 account the indication, its severity, route of administration, complicating conditions and the like. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The effectiveness of a selected actual dose can readily be determined, for example, by measuring clinical symptoms or standard anti-inflammatory
45 indicia after administration of the selected dose. The above dosages are exemplary of

the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention. For conditions or disease states as are treated by the present invention, maintaining consistent daily levels in a subject over an extended period of time, e.g., in a maintenance regime, can be particularly beneficial.

In another aspect, the present invention provides a compound of formula (I) for use in therapy.

The compounds of the present invention are generally inhibitors of the serine/threonine kinase p38 and are therefore also inhibitors of cytokine production which is mediated by p38 kinase. Within the meaning of the term "inhibitors of the serine/threonine kinase p38" are included those compounds that interfere with the ability of p38 to transfer a phosphate group from ATP to a protein substrate according to the assay described below.

It will be appreciated that the compounds of the invention may be selective for one or more of the isoforms of p38, for example p38 α , p38 β , p38 γ and/or p38 δ . In one embodiment, the compounds of the invention selectively inhibit the p38 α isoform. In another embodiment, the compounds of the invention selectively inhibit the p38 β isoform. In a further embodiment, the compounds of the invention selectively inhibit the p38 α and p38 β isoforms. Assays for determining the selectivity of compounds for the p38 isoforms are described in, for example, WO 99/61426, WO 00/71535 and WO 02/46158.

It is known that p38 kinase activity can be elevated (locally or throughout the body), p38 kinase can be incorrectly temporally active or expressed, p38 kinase can be expressed or active in an inappropriate location, p38 kinase can be constitutively expressed, or p38 kinase expression can be erratic; similarly, cytokine production mediated by p38 kinase activity can be occurring at inappropriate times, inappropriate locations, or it can occur at detrimentally high levels.

Accordingly, the present invention provides a method for the treatment of a condition or disease state mediated by p38 kinase activity, or mediated by cytokines produced by the activity of p38 kinase, in a subject which comprises administering to said subject a therapeutically effective amount of a compound of formula (I). The compound may be administered as a single or polymorphic crystalline form or forms, an amorphous form, a single enantiomer, a racemic mixture, a single stereoisomer, a mixture of stereoisomers, a single diastereoisomer or a mixture of diastereoisomers.

The present invention also provides a method of inhibiting cytokine production which is mediated by p38 kinase activity in a subject, e.g. a human, which comprises administering to said subject in need of cytokine production inhibition a therapeutic, or cytokine-inhibiting, amount of a compound of the present invention. The compound may be administered as a single or polymorphic crystalline form or forms, an amorphous form, a single enantiomer, a racemic mixture, a single stereoisomer, a mixture of stereoisomers, a single diastereoisomer or a mixture of diastereoisomers.

The present invention treats these conditions by providing a therapeutically effective amount of a compound of this invention. By "therapeutically effective amount" is meant a symptom-alleviating or symptom-reducing amount, a cytokine-

reducing amount, a cytokine-inhibiting amount, a kinase-regulating amount and/or a kinase-inhibiting amount of a compound. Such amounts can be readily determined by standard methods, such as by measuring cytokine levels or observing alleviation of clinical symptoms. For example, the clinician can monitor accepted measurement scores for anti-inflammatory treatments.

The compounds of the present invention can be administered to any subject in need of inhibition or regulation of p38 kinase or in need of inhibition or regulation of p38 mediated cytokine production. In particular, the compounds may be administered to mammals. Such mammals can include, for example, horses, cows, sheep, pigs, mice, dogs, cats, primates such as chimpanzees, gorillas, rhesus monkeys, and, most preferably, humans.

Thus, the present invention provides methods of treating or reducing symptoms in a human or animal subject suffering from, for example, rheumatoid arthritis, osteoarthritis, asthma, psoriasis, eczema, allergic rhinitis, allergic conjunctivitis, adult respiratory distress syndrome, chronic pulmonary inflammation, chronic obstructive pulmonary disease, chronic heart failure, silicosis, endotoxemia, toxic shock syndrome, inflammatory bowel disease, tuberculosis, atherosclerosis, neurodegenerative disease, Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, epilepsy, multiple sclerosis, aneurism, stroke, irritable bowel syndrome, muscle degeneration, bone resorption diseases, osteoporosis, diabetes, reperfusion injury, graft vs. host reaction, allograft rejections, sepsis, systemic cachexia, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome (AIDS), malaria, leprosy, infectious arthritis, leishmaniasis, Lyme disease, glomerulonephritis, gout, psoriatic arthritis, Reiter's syndrome, traumatic arthritis, rubella arthritis, Crohn's disease, ulcerative colitis, acute synovitis, gouty arthritis, spondylitis, and non articular inflammatory conditions, for example, herniated/ruptured/prolapsed intervertebral disk syndrome, bursitis, tendonitis, tenosynovitis, fibromyalgic syndrome and other inflammatory conditions associated with ligamentous sprain and regional musculoskeletal strain, pain, for example that associated with inflammation and/or trauma, osteopetrosis, restenosis, thrombosis, angiogenesis, cancer including breast cancer, colon cancer, lung cancer or prostatic cancer, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I).

A further aspect of the invention provides a method of treatment of a human or animal subject suffering from rheumatoid arthritis, asthma, psoriasis, chronic pulmonary inflammation, chronic obstructive pulmonary disease, chronic heart failure, systemic cachexia, glomerulonephritis, Crohn's disease, neurodegenerative disease, Alzheimer's disease, Parkinson's disease, epilepsy and cancer including breast cancer, colon cancer, lung cancer and prostatic cancer, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I).

A further aspect of the invention provides a method of treatment of a human or animal subject suffering from rheumatoid arthritis, asthma, psoriasis, chronic pulmonary inflammation, chronic obstructive pulmonary disease, chronic heart failure, systemic cachexia, glomerulonephritis, Crohn's disease and cancer including

breast cancer, colon cancer, lung cancer and prostatic cancer, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I).

5 A further aspect of the invention provides a method of treatment of a human or animal subject suffering from rheumatoid arthritis, asthma, chronic pulmonary inflammation, chronic obstructive pulmonary disease, neurodegenerative disease, Alzheimer's disease, Parkinson's disease and epilepsy which comprises administering to said subject a therapeutically effective amount of a compound of formula (I).

10 A further aspect of the invention provides a method of treatment of a human or animal subject suffering from any type of pain including chronic pain, rapid onset of analgesis, neuromuscular pain, headache, cancer pain, acute and chronic inflammatory pain associated with osteoarthritis and rheumatoid arthritis, post operative inflammatory pain, neuropathic pain, diabetic neuropathy, trigeminal neuralgia, post-hepatic neuralgia, inflammatory neuropathies and migraine pain which comprises
15 administering to said subject a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof.

The compounds of formula (I) may be employed alone or in combination with other therapeutic agents for the treatment of the above-mentioned conditions. In particular, in rheumatoid arthritis therapy, combination with other chemotherapeutic
20 or antibody agents is envisaged. Combination therapies according to the present invention thus comprise the administration of at least one compound of formula (I) and at least one other pharmaceutically active agent. The compound(s) of formula (I) and the other pharmaceutically active agent(s) may be administered together or separately and, when administered separately, this may occur separately or
25 sequentially in any order. The amounts of the compound(s) of formula (I) and the other pharmaceutically active agent(s) and the relative timings of administration will be selected in order to achieve the desired combined therapeutic effect. Examples of other pharmaceutically active agents which may be employed in combination with compounds of formula (I) for rheumatoid arthritis therapy include:
30 immunosuppressants such as amtolmetin guacil, mizoribine and rimexolone; anti-TNF α agents such as etanercept, infliximab, diacerein; tyrosine kinase inhibitors such as leflunomide; kallikrein antagonists such as subreum; interleukin 11 agonists such as oprelvekin; interferon beta 1 agonists; hyaluronic acid agonists such as NRD-101 (Aventis); interleukin 1 receptor antagonists such as anakinra; CD8 antagonists such
35 as amiprilose hydrochloride; beta amyloid precursor protein antagonists such as reumacon; matrix metalloprotease inhibitors such as cipemastat and other disease modifying anti-rheumatic drugs (DMARDs) such as methotrexate, sulphasalazine, cyclosporin A, hydroxychloroquine, auranofin, aurothioglucose, gold sodium thiomalate and penicillamine.

40

Examples

The following examples are illustrative embodiments of the invention, not limiting the scope of the invention in any way. Reagents are commercially available or are prepared according to procedures in the literature.

LCMS was conducted on a column (3.3cm x 4.6mm ID, 3um ABZ+PLUS), at a Flow Rate of 3ml/min, Injection Volume of 5µl, at room temperature and UV Detection Range at 215 to 330nm.

Intermediate 1: 6-Chloro-N-cyclopropylmethylnicotinamide

6-Bromonicotinic acid (200mg, 0.99mmol) was heated at reflux in thionyl chloride (2ml) for 2.5hrs. The reaction was allowed to cool to room temperature and the excess thionyl chloride evaporated under vacuum. The residue was dissolved in acetone (4ml), cyclopropylmethylamine (71mg, 0.10mmol) and sodium carbonate (500mg) were added to the solution. The reaction was stirred at room temperature for 4hrs, filtered and the filtrate reduced to dryness under vacuum to give 6-chloro-N-cyclopropylmethylnicotinamide as a cream solid. NMR: δ H [2 H₆]-DMSO 8.82,(2H, m), 8.23,(1H, dd), 7.63,(1H, d), 3.14,(2H, t), 1.01,(1H, m), 0.44,(2H, m), 0.22,(2H, m).

Intermediate 2: 6-Chloro-N-(4-methoxyphenyl)nicotinamide

6-Bromonicotinic acid (200mg, 0.99mmol) was heated at reflux in thionyl chloride (2ml) for 3hrs. The reaction was allowed to cool to room temperature and the excess thionyl chloride evaporated under vacuum. The residue was dissolved in DCM (2ml), p-anisidine (123mg, 0.10mmol) and sodium carbonate (500mg) were added to the solution. The reaction was stirred at room temperature for 4hrs, filtered and the filtrate reduced to dryness under vacuum to give 6-chloro-N-(4-methoxyphenyl)nicotinamide. NMR: δ H [2 H₆]-DMSO 10.37,(1H, b), 8.94,(1H, d), 8.34,(1H, dd), 7.70,(1H, d), 7.66,(2H, m), 6.95,(2H, m), 3.75,(3H, s).

Intermediate 3: 6-Chloro-N-(3-methoxybenzyl)nicotinamide

6-Bromonicotinic acid (200mg, 0.99mmol) was heated at reflux in thionyl chloride (2ml) for 3hrs. The reaction was allowed to cool to room temperature and the excess thionyl chloride evaporated under vacuum. The residue was dissolved in DCM (2ml), 3-methoxybenzylamine (137mg, 0.10mmol) and sodium carbonate (500mg) were added to the solution. The reaction was stirred at room temperature for 4hrs, filtered and the filtrate reduced to dryness under vacuum to give 6-chloro-N-(3-methoxybenzyl)nicotinamide. NMR: δ H [2 H₆]-DMSO 9.29,(1H, t), 8.88,(1H, d), 8.28,(1H, dd), 7.66,(1H, d), 7.25,(1H, t), 6.90,(2H, m), 6.83,(1H, m), 4.47,(2H, d), 3.74,(3H, s).

Intermediate 4: 6-Chloro-N-(3-methylsulphonylaminobenzyl)nicotinamide

6-Bromonicotinic acid (200mg, 0.99mmol) was heated at reflux in thionyl chloride (2ml) for 3hrs. The reaction was allowed to cool to room temperature and the excess thionyl chloride evaporated under vacuum. The residue was dissolved in DCM (2ml), 3-methylsulphonylaminobenzylamine (200mg, 0.10mmol) and sodium carbonate (500mg) were added to the solution. The reaction was stirred at room temperature for 4hrs, filtered and the filtrate reduced to dryness under vacuum to give 6-chloro-N-(3-methylsulphonylaminobenzyl)nicotinamide. NMR: δ H [2 H₆]-DMSO 9.30,(1H, t), 8.88,(1H, d), 8.28,(1H, dd), 7.67,(1H, d), 7.23,(1H, t), 7.10,(1H, s), 7.04,(1H, d), 6.97,(1H, d), 4.45,(2H, d), 2.90,(3H, s).

Intermediate 5: 6-Chloro-N-[2-(4-methylpiperazin-1-yl)phenyl]nicotinamide

6-Bromonicotinic acid (200mg, 0.99mmol) was heated at reflux in thionyl chloride (2ml) for 3hrs. The reaction was allowed to cool to room temperature and the excess thionyl chloride evaporated under vacuum. The residue was dissolved in DCM (2ml), 1-(2-aminobenzyl)-4-methylpiperazine (205mg, 0.10mmol) and sodium carbonate (500mg) were added to the solution. The reaction was stirred at room temperature for 4hrs, filtered and the filtrate reduced to dryness under vacuum to give 6-chloro-N-[2-(4-methylpiperazin-1-yl)phenyl]nicotinamide. NMR: δ H [2 H₆]-DMSO 11.62,(1H, s), 8.95,(1H, d), 8.32,(1H, dd), 8.25,(1H, d), 7.77,(1H, d), 7.34,(1H, m), 7.28,(1H, m), 7.10,(1H, m), 3.73,(2H, s), 2.56-2.20,(8H, b), 2.12,(3H, s).

Intermediate 6: 4-Methyl-N-(3-pyridin-2-yl-phenyl)-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide

3-Iodo-4-methyl-N-(3-pyridin-2-yl-phenyl)benzamide (Intermediate 7) (83mg, 0.20mmol), bis(pinnacolato)diboron (100mg, 0.39mmol), potassium acetate (97mg, 1.0mmol) and PdCl₂dppf (12mg) were heated at 80°C in DMF (2.5ml) for 4hrs. The cooled reaction was absorbed onto silica, applied to a bond-elut (10g, silica) and eluted with an ethylacetate / cyclohexane gradient (0 to 100%). The solvent was evaporated from the product fractions under vacuum and the residue triturated with ether to give 4-methyl-N-(3-pyridin-2-yl-phenyl)-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide as a white solid (31mg). LCMS: retention time 3.69min, MH⁺ 415.

Intermediate 7: 3-Iodo-4-methyl-N-(3-pyridin-2-yl-phenyl)benzamide

3-Iodo-4-methylbenzoic acid (154mg, 0.59mmol) was heated at 80°C in thionyl chloride (2ml) for 3hrs. The reaction was allowed to cool to room temperature and the excess thionyl chloride evaporated under vacuum. The residue was dissolved in acetone (3ml), 2-(3-aminophenyl)pyridine (100mg, 0.59mmol) and sodium carbonate (400mg) were added to the solution. The reaction was stirred at room temperature for 11 days, filtered and the filtrate reduced to dryness under vacuum. The residue was dissolved in ether and filtered through a bond-elut (1g, silica), washing with ether.

The solvent was evaporated from the combined filtrate and washings to give 3-iodo-4-methyl-N-(3-pyridin-2-yl-phenyl)benzamide as a cream foam. NMR: δ H CDCl₃ 8.70,(1H, dt), 8.33,(1H, d), 8.18,(1H, t), 7.93-7.89,(2H, m), 7.79-7.75,(4H, m), 7.50,(1H, t), 7.35,(1H, d), 7.26,(1H, m), 2.51,(3H, s).

Intermediate 8: N-Cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide

N-Cyclopropyl-3-iodo-4-methylbenzamide (Intermediate 9) (1.1g, 3.64mmol), bis(pinnacolato)diboron (1.85g, 7.28mmol), potassium acetate (1.79g, 18.2mmol) and PdCl₂dppf (55mg) were heated at 85°C in DMF (30ml) for 4.5hrs. The cooled reaction was absorbed onto silica, applied to a bond-elut (10g, silica) and eluted with an ethylacetate / cyclohexane gradient (0 to 100%). The solvent was evaporated from the product fractions under vacuum and the residue triturated with cyclohexane to give N-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide as a white solid (650mg). NMR: δ H [²H₆]-DMSO 8.40,(1H, d), 8.06,(1H, d), 7.76,(1H, dd), 7.23,(1H, d), 2.82,(1H, m), 2.48,(3H, s), 1.30,(12H, s), 0.66,(2H, m), 0.56,(2H, m).

Intermediate 9: N-Cyclopropyl-3-iodo-4-methylbenzamide

3-Iodo-4-methylbenzoic acid (1.0g, 3.8mmol) was heated at 80°C in thionyl chloride (10ml) for 2hrs. The reaction was allowed to cool to room temperature and the excess thionyl chloride evaporated under vacuum. The residue was dissolved in DCM (10ml), cyclopropylamine (0.32ml) and sodium carbonate (2.0g) were added to the solution. The reaction was stirred at room temperature for 18hrs, filtered and the filtrate reduced to dryness under vacuum. The residue was triturated with ether to give N-cyclopropyl-3-iodo-4-methylbenzamide as a white solid (1.1g). NMR: δ H [²H₆]-DMSO 8.46,(1H, d), 8.24,(1H, d), 7.74,(1H, dd), 7.38,(1H, d), 2.82,(1H, m), 2.38,(3H, s), 0.67,(2H, m), 0.55,(2H, m).

Intermediate 10: N-Cyclopropylmethyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide

4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzoic acid (Intermediate 17) (2.0g, 7.63mmol), DIPEA (4ml, 22.89mmol) and HATU (3.05g, 8.02mmol) were dissolved in DMF (20ml) and stirred at room temperature for 15mins. Cyclopropylmethylamine (568mg, 8.01mmol) was added and the reaction stirred at room temperature for 18hours. The solvent was evaporated under vacuum and the reaction partitioned between ethyl acetate (250ml) and water (50ml). The organic phase was washed with hydrochloric acid (2N, 50ml) and aqueous sodium bicarbonate (1M, 50ml), then dried (magnesium sulphate) and the solvent evaporated under vacuum. The residue was absorbed onto silica and purified by flash column chromatography eluting with cyclohexane / ethyl acetate (4:1). The solvent was evaporated from the product fractions under vacuum to give N-cyclopropylmethyl-4-

methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (1.73g). LCMS: retention time 3.47min, MH^+ 316. NMR: δH [2H_6]-DMSO 8.54,(1H, t), 8.11,(1H, d), 7.82,(1H, dd), 7.26,(1H, d), 3.12,(2H, t), 1.32,(12H, s), 1.03,(1H, m), 0.42,(2H, m), 0.22,(2H, m).

5

Intermediate 11: 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiazol-2-yl)-benzamide

4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzoic acid (2.0g, 7.63mmol), DIPEA (4ml, 22.89mmol) and HATU (3.05g, 8.02mmol) were dissolved in DMF (20ml) and stirred at room temperature for 15mins. 2-Aminothiazole (801mg, 8.01mmol) was added and the reaction stirred at room temperature for 18hours. The solvent was evaporated under vacuum and the reaction partitioned between ethyl acetate (250ml) and water (50ml). The organic phase was washed with hydrochloric acid (2N, 50ml) and aqueous sodium bicarbonate (1M, 50ml), then dried (magnesium sulphate) and the solvent evaporated under vacuum. The residue was absorbed onto silica and purified by flash column chromatography eluting with cyclohexane / ethyl acetate (4:1). The solvent was evaporated from the product fractions under vacuum to 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiazol-2-yl)-benzamide (1.72g). LCMS: retention time 3.66min, MH^+ 345. NMR: δH [2H_6]-DMSO 12.65,(1H, b), 8.32,(1H, d), 8.08,(1H, dd), 7.56,(1H, d), 7.35,(1H, d), 7.28,(1H, d), 2.54,(3H, s), 1.34,(12H, s).

Intermediate 12: 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiadiazol-2-yl)-benzamide

4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzoic acid (2.0g, 7.63mmol), DIPEA (4ml, 22.89mmol) and HATU (3.05g, 8.02mmol) were dissolved in DMF (20ml) and stirred at room temperature for 15mins. 2-Aminothiadiazole (810mg, 8.01mmol) was added and the reaction stirred at room temperature for 18hours. The solvent was evaporated under vacuum and the reaction partitioned between ethyl acetate (250ml) and hydrochloric acid (2N, 150ml). The aqueous was extracted with ethylacetate (2 x 250ml). The combined organic extracts were dried (magnesium sulphate) and the solvent evaporated under vacuum. The residue was absorbed onto silica and purified by flash column chromatography eluting with cyclohexane / ethyl acetate (4:1 then 1:1). The solvent was evaporated from the product fractions under vacuum to 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiadiazol-2-yl)-benzamide (0.95g). LCMS: retention time 3.34min, MH^+ 346. NMR: δH [2H_6]-DMSO 13.08,(1H, b), 9.22,(1H, s), 8.35,(1H, d), 8.11,(1H, dd), 7.38,(1H, d), 2.55,(3H, s), 1.34,(12H, s).

Intermediate 13: N-[4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide

- 5 N-(3-Iodo-4-methylphenyl)-3-furamide (Intermediate 15) (2.5g, 7.64mmol), bis(pinnacolato)diboron (2.13g, 8.41mmol), potassium acetate (825mg, 8.41mmol) and PdCl₂dppf (312mg, 0.38mmol) in DMF (20ml) were heated at 80°C for 20hrs. The cooled reaction was absorbed onto silica and applied to a bond-elut (silica, 10g) and eluted with a cyclohexane / ethyl acetate gradient. The product fractions were concentrated under vacuum, dissolved in DMF (40ml) and reacted with
- 10 bis(pinnacolato)diboron (7.76g, 30.57mmol), potassium acetate (3.0g, 30.57mmol) and PdCl₂dppf (249mg, 0.306mmol) at 80°C for 23 hrs. The cooled reaction was absorbed onto silica and applied to bond-eluts (silica, 2 x10g) and eluted with a cyclohexane / ethyl acetate gradient. The product fractions were concentrated under vacuum to give N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide. LCMS: retention time 3.55min, MH⁺ 328. NMR: δH [²H₆]-
- 15 DMSO 9.86,(1H, b), 8.36,(1H, m), 7.86-7.82,(2H, m), 7.77,(1H, t), 7.14,(1H, d), 6.99,(1H, m), 2.41,(3H, s), 1.30,(12H, s).

Intermediate 14: N-[4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]thiophene-3-amide

- 20 N-(3-Iodo-4-methylphenyl)thiophene-3-amide (Intermediate 16) (2.64g, 7.64mmol), bis(pinnacolato)diboron (2.13g, 8.41mmol), potassium acetate (825mg, 8.41mmol) and PdCl₂dppf (312mg, 0.38mmol) in DMF (20ml) were heated at 80°C for 20hrs. The cooled reaction was absorbed onto silica and applied to a bond-elut (silica, 10g) and eluted with a cyclohexane / ethyl acetate gradient. The product
- 25 fractions were concentrated under vacuum, dissolved in DMF (20ml) and reacted with bis(pinnacolato)diboron (1.77g, 7.0mmol), potassium acetate (687mg, 7.0mmol) and PdCl₂dppf (143mg, 0.175mmol) at 80°C for 16 hrs. The cooled reaction was absorbed onto silica and applied to a bond-elut (silica, 10g) and eluted with a cyclohexane / ethyl acetate gradient. The product fractions were concentrated under
- 30 vacuum to give N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]thiophene-3-amide. LCMS: retention time 3.65min, MH⁺ 344. NMR: δH [²H₆]-DMSO 9.99,(1H, b), 8.35,(1H, s), 7.90,(1H, d), 7.85,(1H, dd), 7.63,(2H, m), 7.14,(1H, d), 2.42,(3H, s), 1.30,(12H, s).

35 **Intermediate 15: N-(3-Iodo-4-methylphenyl)-3-furamide**

- 3-Furoic acid (2.4g, 21.45mmol) and HATU (8.15g, 21.45mmol) in DMF (25ml) were stirred at room temperature for 15mins. HOBt (2.9g, 21.45mmol), 3-iodo-4-methylaniline (5.0g, 21.45mmol) and DIPEA (11.2ml, 64.35mmol) were added and the reaction stirred at room temperature for 16hrs. The solvent was
- 40 evaporated under vacuum and the residue partitioned between ethyl acetate (100ml) and aqueous sodium carbonate (10%, 100ml). The aqueous layer was extracted with ethyl acetate (50ml) and the combined organic phases washed with hydrochloric acid

(2N, 75ml), water (75ml) and brine (75ml). The organic phase was dried (magnesium sulphate) and absorbed onto silica. The silica was applied to a flash silica column and eluted with cyclohexane / ethyl acetate (3:1). The solvent was evaporated from the product fractions under vacuum to give N-(3-iodo-4-methylphenyl)-3-furamide.

- 5 LCMS: retention time 3.52min, MH^+ 328. NMR: δH [2H_6]-DMSO 9.92,(1H, b), 8.36,(1H, d), 8.23,(1H, d), 7.80,(1H, t), 7.66,(1H, dd), 7.29,(1H, d), 6.98,(1H, d), 2.33,(3H, s).

Intermediate 16: N-(3-Iodo-4-methylphenyl)thiophene-3-amide

- 10 Thiophene-3-carboxylic acid (2.75g, 21.45mmol) and HATU (8.15g, 21.45mmol) in DMF (25ml) were stirred at room temperature for 15mins. HOBt (2.9g, 21.45mmol), 3-iodo-4-methylaniline (5.0g, 21.45mmol) and DIPEA (11.2ml, 64.35mmol) were added and the reaction stirred at room temperature for 16hrs. The solvent was evaporated under vacuum and the residue partitioned between ethyl acetate (100ml) and aqueous sodium carbonate (10%, 100ml). The aqueous layer was extracted with ethyl acetate (50ml) and the combined organic phases washed with hydrochloric acid (2N, 75ml), water (75ml) and brine (75ml). The organic phase was dried (magnesium sulphate) and absorbed onto silica. The silica was applied to a flash silica column and eluted with cyclohexane / ethyl acetate (4:1). The solvent was evaporated from the product fractions under vacuum to give N-(3-iodo-4-methylphenyl)thiophene-3-amide. LCMS: retention time 3.69min, MH^+ 344. NMR: δH [2H_6]-DMSO 10.06,(1H, b), 8.34,(1H, m), 8.29,(1H, d), 7.70,(1H, dd), 7.66,(1H, dd), 7.62,(1H, dd), 7.30,(1H, d), 2.34,(3H, s).
- 15
- 20

25 **Intermediate 17: 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)benzoic acid**

- 3-Iodo-4-methylbenzoic acid (10g, 38.16mmol), bis(pinnacolato)diboron (14.5g, 57.24mmol), potassium acetate (18.73g, 190.8mmol) and $PdCl_2dppf$ (3.12g, 3.8mmol) in DMF (200ml) were heated at 80°C for 21hrs. The solvent was evaporated from the cooled reaction under vacuum, the residue dissolved in ethyl acetate (300ml) and hydrochloric acid (2N, 300ml) and filtered through celite. The organic phase was separated and the aqueous extracted with ethyl acetate (2 x 300ml). The combined organic extracts were washed with brine (500ml) and dried (magnesium sulphate). The solvent was evaporated under vacuum and the residue absorbed onto silica and applied to a silica flash column. This was eluted with cyclohexane / ethyl acetate (5:1). The product fractions were concentrated under vacuum to give 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)benzoic acid. LCMS: retention time 3.65min. NMR: δH [2H_6]-DMSO 12.83,(1H, b), 8.23,(1H, d), 7.89,(1H, dd), 7.29,(1H, d), 2.51,(3H, s), 1.30,(12H, s).
- 30
- 35
- 40

Intermediate 18: N-[4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-2-pyrrolidin-1-yl-isonicotinamide

Bis(pinacolato)diborane (7.24g, 28.5mmol) was added to a mixture of N-(3-iodo-4-methylphenyl)-2-pyrrolidin-1-yl-isonicotinamide (Intermediate 19) (7.73g, 19mmol) in dimethylformamide (100ml) potassium acetate (9.32g, 95mmol) and PdCl₂dppf and the reaction was heated under an atmosphere of nitrogen at 80°C for 16 hours. The reaction was cooled and the solvent removed *in vacuo*. The residue was taken up in chloroform (150ml), washed with water (3x100ml) and brine (100ml), dried over magnesium sulfate, filtered and solvent removed *in vacuo*. The residue was purified by column chromatography (20:80 ethyl acetate:cyclohexane to 50:50 ethyl acetate:cyclohexane). To give N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-2-pyrrolidin-1-yl-isonicotinamide as a white solid (1.5g, 3.7mmol). LCMS: retention time 2.90 min MH⁺408. NMR: δ H – CDCl₃ 8.27 (1H, d), 7.99 (1H, dd), 7.76 (1H, b), 7.65 (1H, d), 6.20 (1H, d), 6.82 (1H, b), 6.77 (1H, b), 3.52 (4H, apparent t), 2.52 (3H, s), 2.25 (4H, m).

Intermediate 19: N-(3-Iodo-4-methylphenyl)-2-pyrrolidin-1-yl-isonicotinamide

A solution of N-(3-iodo-4-methylphenyl)-2-chloro-isonicotinamide (Intermediate 20) (7.00g, 18.8mmol) in pyrrolidine (20ml) was heated at 80°C under an atmosphere of nitrogen for 16 hours. Excess pyrrolidine was removed *in vacuo* and the residue was titrated with diethyl ether (20ml). The resulting solid was collected by filtration and dried *in vacuo* to give N-(3-iodo-4-methylphenyl)-2-pyrrolidin-1-yl-isonicotinamide as a pale yellow solid (7.73g, 18mmol). LCMS: retention time 2.77 min MH⁺408. NMR: δ H [²H₆] – DMSO 10.29 (1H, s), 8.29 (1H, d), 8.20 (1H, d), 7.71 (1H, dd), 7.72 (1H, dd), 6.97 (1H, brd), 6.88 (1H, b), 3.45 (2H, apparent t), 3.09 (2H, m), 2.35 (3H, s), 1.98 (2H, m), 1.82 (2H, m).

Intermediate 20: 2-Chloro-N-(3-iodo-4-methylphenyl)-isonicotinamide

2-Chloroisonicotinic acid (3.3g, 21mmol), HATU (8.75g, 23mmol), diisopropylethyl amine (10.9ml, 63mmol) and 4-iodo-3-methylaniline (5.00g, 21mmol) in dimethylformamide (50ml) were heated under nitrogen for 16 hours. The reaction was cooled, solvent removed *in vacuo* and the residue taken up in dichloromethane (150ml). The organic solution was washed with water (3x100ml) and brine (100ml), dried over magnesium sulfate, filtered and solvent removed *in vacuo*. The residue was purified by column chromatography (40:60 ethyl acetate:cyclohexane) to give 2-chloro-N-(3-iodo-4-methylphenyl)-isonicotinamide as a white solid (7.00g, 18.8mmol). LCMS: retention time 3.59 min MH⁺373. NMR: δ H [²H₆] – DMSO 10.52 (1H, s), 8.62 (1H, d), 8.29 (1H, d), 7.99 (1H, b), 7.87 (1H, dd), 7.70 (1H, dd), 7.34 (1H, d), 2.36 (3H, s).

Intermediate 21: 6-Chloro-N-cyclopropylmethylnicotinamide

6-Bromonicotinic acid (200mg, 0.99mmol) was heated at reflux in thionyl chloride (2ml) for 2.5hrs. The reaction was allowed to cool to room temperature and the excess thionyl chloride evaporated under vacuum. The residue was dissolved in acetone (4ml), cyclopropylmethylamine (71mg, 0.10mmol) and sodium carbonate (500mg) were added to the solution. The reaction was stirred at room temperature for 4hrs, filtered and the filtrate reduced to dryness under vacuum to give 6-chloro-N-cyclopropylmethylnicotinamide as a cream solid. NMR: δ H [2 H₆]-DMSO 8.82,(2H, m), 8.23,(1H, dd), 7.63,(1H, d), 3.14,(2H, t), 1.01,(1H, m), 0.44,(2H, m), 0.22,(2H, m).

General Method A

6-Bromonicotinic acid (100mg, 0.5mmol) was heated at 95°C in thionyl chloride (0.63ml) for 2hours. The excess thionyl chloride was evaporated under vacuum and the residue dissolved in DCM (2ml). To this solution, amine (0.5mmol) and sodium carbonate (100mg) were added and the reaction was stirred at room temperature for 2hours. The reaction was filtered and the residue washed with DCM. The combined filtrate and washings were reduced to dryness to give the desired 6-chloronicotinamide.

Compound	Amine	MH ⁺	Retention time (minutes)
Intermediate 22: 6-Chloro-N-(3-methylbutyl)nicotinamide	3-methylbutylamine	227	2.92
Intermediate 23: 6-Chloro-N-(1-cyclopropylethyl)nicotinamide	1-cyclopropylethylamine	225	2.65
Intermediate 24: 6-Chloro-N-(2,2-dimethylpropyl)nicotinamide	2,2-dimethylpropylamine	227	2.82
Intermediate 25: 6-Chloro-N-(2,2-dimethylcyclopropyl)nicotinamide	2,2-dimethylcyclopropylamine	225	2.67
Intermediate 26: 6-Chloro-N-cyclopropylnicotinamide	cyclopropylamine	197	2.19
Intermediate 27: 6-Chloro-N-cyclohexylmethylnicotinamide	cyclohexylmethylamine	253	3.18
Intermediate 28: 6-Chloro-N-cyclobutylnicotinamide	cyclobutylamine	211	2.51

Intermediate 29: 6-Chloro-N-(2-methylpropyl)nicotinamide	2-methylpropylamine	213	2.63
Intermediate 30: 6-Chloro-N-propylnicotinamide	propylamine	199	2.38
Intermediate 31: 6-Chloro-N-cyclopentylnicotinamide	cyclopentylamine	225	2.70

Intermediate 32: 6-Chloro-N-cyclobutylmethylnicotinamide

6-Chloro-N-cyclobutylmethylnicotinamide was prepared from cyclobutylmethylamine using General Method A.

- 5 NMR: δ H [2 H₆] – DMSO 8.81,(1H, d), 8.70,(1H, bt), 8.22,(1H, dd), 7.64,(1H, d), 3.30,(2H, t), 2.52,(1H, m), 1.99,(2H, m), 1.81,(2H, m), 1.73,(2H, m).

Intermediate 33: 6-Chloro-N-(1-methylpropyl)nicotinamide

- 10 6-Chloro-N-(1-methylpropyl)nicotinamide was prepared from 1-methylpropylamine using General Method A.

NMR: δ H [2 H₆] – DMSO 8.82,(1H, d), 8.42,(1H, d), 8.24,(1H, dd), 7.64,(1H, d), 3.91,(1H, m), 1.51,(2H, m), 1.15,(3H, d), 0.87,(3H, t).

- 15 **Intermediate 34: N-Cyclopropyl-5-fluoro-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide**

- 20 3-Bromo-N-cyclopropyl-5-fluoro-4-methylbenzamide (Intermediate 35, 900mg), bispinacolatodiboron (4.5g), potassium acetate (2.1g) and PdCl₂dppf (75mg) were mixed in DMF (40ml) and heated at 100°C for 18hours. The cooled reaction was absorbed onto silica and applied to SPE's (Si 2 x 10g). The SPE's were eluted with an ethylacetate / cyclohexane gradient (0-6.25% ethylacetate). The solvent was evaporated from the product fractions under vacuum and the residue recrystallised from cyclohexane to give N-cyclopropyl-5-fluoro-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (260mg).
LCMS: MH⁺ 320, retention time 3.39mins.

25

Intermediate 35: 3-Bromo-N-cyclopropyl-5-fluoro-4-methylbenzamide

- 30 3-Fluoro-4-methylbenzoic acid (462mg, 3.0mmol) was added to a stirred mixture of bromine (2.31ml, 45mmol) and iron powder (252mg, 4.5mmol) under nitrogen. The reaction was stirred at 20°C for 4 hours and then left to stand for 16 hours. Sodium thiosulphate solution (200ml) was added and the product was extracted into ethyl acetate (3 x 150ml). Ethyl acetate extracts were combined and

evaporated *in vacuo*. The crude product (mixture of isomers) was dissolved in dimethylformamide (7ml). Cyclopropylamine (208µl, 3.0mmol), HOBT (405mg, 3.0mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (575mg, 3.0mmol) and DIPEA (525µl, 3.0mmol) were added to the stirred solution. The reaction was stirred for 5 hours at 20°C. Solvent was removed *in vacuo* and the residue was partitioned between ethyl acetate and water. Combined ethyl acetate extracts were washed sequentially with aqueous sodium hydrogen carbonate and hydrochloric acid (0.5M), then dried (magnesium sulphate). The ethyl acetate was evaporated *in vacuo* and the residue was purified by silica biotage chromatography eluting with cyclohexane:ethyl acetate (6:1) to give 3-bromo-N-cyclopropyl-5-fluoro-4-methylbenzamide (359mg, 44%).
 NMR: δ H – CDCl₃ 7.68,(1H, s), 7.39,(1H, d), 6.19,(1H, bs), 2.88,(1H, m), 2.36,(3H, d), 0.88,(2H, m), 0.63,(2H, m). LCMS: MH⁺ 272.

15 **Intermediate 36: {5-[(Cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}boronic acid**

N-Cyclopropyl-5-fluoro-3-iodo-4-methylbenzamide (Intermediate 37, 5g) in THF (75ml) was cooled to 0°C and sodium hydride (60%, 1.23g) added portionwise over 10minutes. Once effervescence had ceased the reaction was cooled to -75°C and n-butyl lithium (1.6M in hexanes, 20ml) added over 25minutes maintaining a temperature of <-70°C. Triisopropyl borate (8ml) was added to the reaction over 10minutes and the reaction stirred at -70°C for 4hours. The reaction was quenched with water (20ml) and the mixture allowed to warm to 5°C. The reaction was concentrated under vacuum and the residue partitioned between saturated ammonium chloride and ethyl acetate. The organic phase was washed with saturated ammonium chloride, brine, dried (sodium sulphate) and reduced to dryness under vacuum. The residue was dissolved in DCM/ethyl acetate and purified by column chromatography on silica eluting with an ethyl acetate/ DCM gradient (5-100% ethyl acetate) and then methanol. The product fractions were combined and the solvent evaporated under vacuum to give {5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}boronic acid. LCMS MH⁺ 238, retention time 2.19min.

Intermediate 37: N-Cyclopropyl-5-fluoro-3-iodo-4-methylbenzamide

35 N-Iodosuccinimide (22.5g) was added in portions to a solution of 3-fluoro-4-methylbenzoic acid (15.4g) in trifluoromethanesulphonic acid (100ml) at 0°C over 3hours and the reaction then allowed to warm to room temperature overnight. The reaction mixture was poured into ice/water (400ml) and the precipitate filtered off and washed with water. The solid remaining was dissolved in ethyl acetate, washed with 40 aqueous sodium thiosulphate (x2), then brine, dried (magnesium sulphate) and the solvent evaporated under vacuum. The residue was mixed with thionyl chloride (30ml) and heated at 100°C for 2.5hours. The excess thionyl chloride was removed from the cooled reaction under vacuum and the residue dissolved in DCM (100ml). Sodium carbonate (25g) and cyclopropylamine (13ml) were added to the solution and

the reaction stirred at room temperature for 72 hours. The reaction was filtered and the residue washed with DCM and ethyl acetate. The solvent was evaporated from the combined filtrate and washings under vacuum. The residue was absorbed onto silica and chromatographed on a flash silica column eluting with an ethyl acetate /

- 5 cyclohexane gradient (22 – 28% ethyl acetate). Appropriate fractions were reduced to dryness under vacuum to give N-cyclopropyl-5-fluoro-3-iodo-4-methylbenzamide. LCMS; MH⁺ 320, retention time 3.16 minutes.

Intermediate 38: 6-{5-[(Cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinic acid

- 10 N-Cyclopropyl-5-fluoro-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (3.2g), methyl 6-chloronicotinate (1.73g), tetrakis(triphenylphosphine)palladium (210mg) and aqueous sodium hydrogen carbonate (1M, 30ml) were mixed in propan-2-ol (100ml) and heated at 90°C for 18 hours. The reaction was allowed to cool and the propan-2-ol removed under
- 15 vacuum. The residue was partitioned between ethyl acetate and aqueous sodiumhydrogen carbonate (1M). The aqueous phase was acidified with hydrochloric acid (2N) and extracted with ethyl acetate (x2). The organic extracts were washed with brine, dried (magnesium sulphate) and reduced to dryness under vacuum. The resulting foam was triturated with ether to give 6-{5-[(cyclopropylamino)carbonyl]-3-
- 20 fluoro-2-methylphenyl}nicotinic acid as a solid. LCMS: MH⁺ 315, retention time 2.87 mins.


Intermediate 39: (2-Ethylcyclopropyl)methylamine

- 25 A solution of 2-ethylcyclopropylcarboxamide (250mg, 2.2mmol) in THF was heated to reflux. Borane-dimethylsulphide (1M solution in DCM, 3.2ml, 3.2mmol) was added dropwise over 30 minutes and the reaction refluxed for 16 hours. Hydrochloric acid (6N, 0.5ml) was added dropwise and the mixture heated at reflux for 30 minutes. The cooled reaction mixture was diluted with water (20ml), washed with ether (50ml) and basified with sodium hydroxide (6N). The aqueous was extracted with ether
- 30 (50ml x 3) and ethyl acetate (50ml). The combined organic extracts were dried (magnesium sulphate), acidified with hydrogen chloride (3.3M in methanol) and reduced to dryness under vacuum to give (2-ethylcyclopropyl)methylamine (230mg). NMR: δ H [²H₆]-DMSO 7.85,(3H, b), 2.66,(2H, d), 1.30-1.13,(2H, m), 0.91,(3H, t), 0.77-0.66,(2H, m), 0.46,(1H, m), 0.33,(1H, m).

35

General Method B


The 2-chloropyridine (0.05mmol), phenyl pinacolborane (0.05mmol), tetrakis(triphenylphosphine) palladium (1mg) and aqueous sodium carbonate (0.25ml) in propan-2-ol (1ml) were heated at 85°C under nitrogen for 18 hours. The cooled reaction was diluted with ethyl acetate (4ml) and methanol (2ml) and filtered through an SCX bond-elut (1g). The product was eluted with 10% ammonia (s.g. 0.88) in methanol. The solvents were evaporated and the residue triturated with ether.

Example 1: N-(3-[5-(Cyclopropylmethyl-carbamoyl)-pyridin-2-yl]-4-methyl-phenyl)-2-pyrrolidin-1-yl-isonicotinamide

10


6-Chloro-N-cyclopropylmethylnicotinamide (Intermediate 1) (25mg, 0.098mmol) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-2-pyrrolidin-1-yl-isonicotinamide (Intermediate 18) (30mg, 0.074mmol), aqueous sodium carbonate (2N, 0.5ml) and tetrakis(triphenylphosphine)palladium (4mg) were heated at 80°C in DMF (1ml) for 18hours. The reaction was absorbed onto silica, applied to a bond-elut (10g, silica) and eluted with an ethylacetate/cyclohexane (0 to 100%), then acetone and methanol. The solvent was evaporated from the product fractions under vacuum and the residue triturated with ether to give N-(3-[5-(cyclopropylmethyl-carbamoyl)-pyridin-2-yl]-4-methyl-phenyl)-2-pyrrolidin-1-yl-isonicotinamide as a white solid (20mg). LCMS: retention time 2.42min, MH^+ 456. NMR: δ H [2H_6]-DMSO 10.32,(1H, s), 9.09,(1H, s), 8.82,(1H, t), 8.28,(1H, m), 8.19,(1H, m), 7.85,(1H, t), 7.76,(1H, m), 7.64,(1H, m), 7.31,(1H, m), 6.98,(1H, m), 6.88,(1H, s), 3.43,(4H, m), 3.18,(2H, m), 2.31,(3H, s), 1.95,(4H, m), 1.07,(1H, m), 0.45,(2H, m), 0.25,(2H, m).

25


Example 2: N-Cyclopropylmethyl-6-[2-methyl-5-(3-pyridin-2-yl-phenylcarbamoyl)-phenyl]-nicotinamide

6-Chloro-N-cyclopropylmethylnicotinamide (Intermediate 1) (18.5mg, 0.073mmol) and 4-methyl-N-(3-pyridin-2-yl-phenyl)-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 6) (30mg, 0.072mmol), aqueous sodium carbonate (2N, 0.5ml) and tetrakis(triphenylphosphine)palladium (4mg) were heated at 90°C in DMF (1ml) for 4hours. The reaction was absorbed onto silica, applied to a bond-elut (5g, silica) and eluted with an ethylacetate/cyclohexane (0 to 100%) and then acetone. The solvent was evaporated from the product fractions under vacuum and the residue triturated with ether to give N-cyclopropylmethyl-6-[2-methyl-5-(3-pyridin-2-yl-phenylcarbamoyl)-phenyl]-nicotinamide as a white solid (20mg). LCMS: retention time 3.18min, MH^+ 463. NMR: δH [2H_6]-DMSO 10.43,(1H, s), 9.14,(1H, s), 8.86,(1H, t), 8.69,(1H, s), 8.53,(1H, s), 8.34,(1H, d), 8.11,(1H, s), 8.01,(1H, d), 7.95-7.89,(3H, m), 7.81-7.78,(2H, m), 7.53-7.46,(2H, m), 7.38,(1H, t), 3.21,(2H, t), 2.44,(3H, s), 1.07,(1H, m), 0.47,(2H, m), 0.27,(2H, m).


Example 3: 6-(5-Cyclopropylcarbamoyl-2-methyl-phenyl)-N-cyclopropylmethyl-nicotinamide



6-Chloro-N-cyclopropylmethylnicotinamide (Intermediate 1) (25.5mg, 0.10mmol) and N-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 8) (30mg, 0.10mmol), aqueous sodium carbonate (2N, 0.5ml) and tetrakis(triphenylphosphine)palladium (4mg) were heated at 90°C in DMF (1ml) for 3hours. The reaction was absorbed onto silica, applied to a bond-elut (5g, silica) and eluted with an ethylacetate/cyclohexane (0 to 100%) and then acetone. The solvent was evaporated from the product fractions under vacuum and the residue triturated with ether to 6-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-N-cyclopropylmethyl-nicotinamide as a cream solid. LCMS: retention time 2.70min, MH^+ 350. NMR: δH [2H_6]-DMSO 9.11,(1H, s), 8.84,(1H, t), 8.48,(1H, d), 8.31,(1H, dd), 7.88,(1H, s), 7.81,(1H, d), 7.70,(1H, d), 7.41,(1H, d), 3.20,(1H, t), 2.86,(1H, m), 2.37,(3H, s), 1.06,(1H, m), 0.69,(2H, m), 0.57,(2H, m), 0.46,(2H, m), 0.26,(2H, m).


Example 4: N-Cyclopropylmethyl-6-[5-(thiadiazol-2-ylcarbamoyl)-2-methyl-phenyl]-nicotinamide

5 N-Cyclopropylmethyl-6-[5-(thiadiazol-2-ylcarbamoyl)-2-methyl-phenyl]-
 nicotinamide was prepared from 6-chloro-N-cyclopropylmethylnicotinamide
 (Intermediate 1) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-
 (thiadiazol-2-yl)-benzamide (Intermediate 12) using General Method B. LCMS:
 retention time 2.79min, MH^+ 394. NMR: δH [2H_6]-DMSO 13.14,(1H, b), 9.24,(1H,
 s), 9.14,(1H, s), 8.86,(1H, t), 8.35,(1H, d), 8.25,(1H, s), 8.10,(1H, d), 7.82,(1H, d),
 10 7.54,(1H, d), 3.21,(2H, t), 2.46,(3H, s), 1.07,(1H, m), 0.47,(2H, m), 0.27,(2H, m).

Example 5: N-Cyclopropylmethyl-6-[5-(thiazol-2-ylcarbamoyl)-2-methyl-phenyl]-nicotinamide


15 N-Cyclopropylmethyl-6-[5-(thiazol-2-ylcarbamoyl)-2-methyl-phenyl]-
 nicotinamide was prepared from 6-chloro-N-cyclopropylmethylnicotinamide
 (Intermediate 1) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-
 (thiazol-2-yl)-benzamide (Intermediate 11) using General Method B. LCMS: retention
 time 2.99min, MH^+ 393. NMR: δH [2H_6]-DMSO 12.71,(1H, b), 9.13,(1H, s),
 20 8.86,(1H, t), 8.34,(1H, d), 8.21,(1H, s), 8.07,(1H, d), 7.81,(1H, d), 7.57,(1H, d),
 7.52,(1H, d), 7.29,(1H, d), 3.21,(2H, t), 2.45,(3H, s), 1.07,(1H, m), 0.47,(2H, m),
 0.27,(2H, m).

Example 6: 6-[5-(Cyclopropylmethylcarbamoyl)-2-methyl-phenyl]-N-cyclopropylmethyl-nicotinamide




5 6-[5-(Cyclopropylmethyl)carbamoyl-2-methyl-phenyl]-N-cyclopropylmethyl-nicotinamide was prepared from 6-chloro-N-cyclopropylmethylnicotinamide (Intermediate 1) and N-(cyclopropylmethyl)-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 10) using General Method B. LCMS: retention time 2.87min, MH^+ 364. NMR: δH [2H_6]-DMSO 9.10,(1H, s), 8.83,(1H, t), 8.60,(1H, t), 8.30,(1H, dd), 7.92,(1H, s), 7.84,(1H, d), 7.71,(1H, d), 10 7.41,(1H, d), 3.19,(2H, t), 3.13,(2H, t), 2.37,(3H, s), 1.03,(2H, m), 0.44,(4H, m), 0.23,(4H, m).

Example 7: N-Cyclopropylmethyl-6-[5-(fur-3-ylcarboxylamino)-2-methyl-phenyl]-nicotinamide




15

N-Cyclopropylmethyl-6-[5-(fur-3-ylcarboxylamino)-2-methyl-phenyl]-nicotinamide was prepared from 6-chloro-N-cyclopropylmethylnicotinamide (Intermediate 1) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide (Intermediate 13) using General Method B. LCMS: retention time 20 2.96min, MH^+ 376. NMR: δH [2H_6]-DMSO 9.99,(1H, s), 9.10,(1H, s), 8.83,(1H, t), 8.38,(1H, s), 8.30,(1H, d), 7.80,(2H, s), 7.75,(1H, d), 7.66,(1H, d), 7.30,(1H, d), 3.20,(2H, t), 2.31,(3H, s), 1.06,(1H, m), 0.46,(2H, m), 0.27,(2H, m).


Example 8: N-Cyclopropylmethyl-6-[2-methyl-5-(thiophen-3-ylcarbonylamino)-phenyl]-nicotinamide

5 N-Cyclopropylmethyl-6-[2-methyl-5-(thiophen-3-ylcarbonylamino)-phenyl]-
 nicotinamide was prepared from 6-chloro-N-cyclopropylmethylnicotinamide
 (Intermediate 1) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-
 phenyl]thiophene-3-amide (Intermediate 14) using General Method B. LCMS:
 retention time 3.07min, MH^+ 392. NMR: δH [2H_6]-DMSO 10.11,(1H, s), 9.11,(1H,
 s), 8.83,(1H, t), 8.35,(1H, s), 8.30,(1H, dd), 7.85,(1H, s), 7.78,(1H, d), 7.67-7.63,(3H,
 10 m), 7.31,(1H, d), 3.20,(2H, t), 2.31,(3H, s), 1.06,(1H, m), 0.46,(1H, m), 0.27,(1H, m).

Example 9: 6-(5-Cyclopropylcarbamoyl-2-methyl-phenyl)-N-(4-methoxyphenyl)-nicotinamide


15 6-[5-Cyclopropylcarbamoyl-2-methyl-phenyl]-N-(4-methoxyphenyl)-
 nicotinamide was prepared from 6-chloro-N-(4-methoxyphenyl)nicotinamide
 (Intermediate 2) and N-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-
 [1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 8) using General Method B.
 LCMS: retention time 2.96min, MH^+ 402. NMR: δH [2H_6]-DMSO 10.38,(1H, s),
 20 9.20,(1H, s), 8.49,(1H, d), 8.40,(1H, dd), 7.91,(1H, s), 7.82,(1H, d), 7.76,(1H, d),
 7.71,(2H, d), 7.43,(1H, d), 6.96,(2H, d), 3.76,(3H, s), 2.87,(1H, m), 2.40,(3H, s),
 0.70,(2H, m), 0.58,(2H, m).

Example 10: N-(4-Methoxyphenyl)-6-[2-methyl-5-(thiadiazol-2-ylcarbamoyl)-phenyl]-nicotinamide




5 N-(4-Methoxyphenyl)-6-[2-methyl-5-(thiadiazol-2-ylcarbamoyl)-phenyl]-
nicotinamide was prepared from 6-chloro-N-(4-methoxyphenyl)nicotinamide
(Intermediate 2) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-
(thiadiazol-2-yl)-benzamide (Intermediate 12) using General Method B. LCMS:
retention time 3.05min, MH^+ 446. NMR: δH [2H_6]-DMSO 13.15,(1H, b), 10.41,(1H,
s), 9.24,(2H, m), 8.45,(1H, dd), 8.28,(1H, s), 8.11,(1H, d), 7.88,(1H, d), 7.71,(2H, d),
10 7.56,(1H, d), 6.97,(2H, d), 3.76,(3H, s), 2.48,(3H, s).

Example 11: N-(4-Methoxyphenyl)-6-[2-methyl-5-(thiazol-2-ylcarbamoyl)-phenyl]-nicotinamide



15 N-(4-Methoxyphenyl)-6-[2-methyl-5-(thiazol-2-ylcarbamoyl)-phenyl]-
nicotinamide was prepared from 6-chloro-N-(4-methoxyphenyl)nicotinamide
(Intermediate 2) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-
(thiazol-2-yl)-benzamide (Intermediate 11) using General Method B. LCMS: retention
time 3.22min, MH^+ 445. NMR: δH [2H_6]-DMSO 12.72,(1H, s), 10.40,(1H, s),
20 9.22,(1H, d), 8.44,(1H, dd), 8.24,(1H, s), 8.09,(1H, d), 7.87,(1H, d), 7.71,(2H, d),
7.58,(1H, d), 7.53,(1H, d), 7.30,(1H, d), 6.97,(2H, d), 3.76,(3H, s), 2.48,(3H, s).

Example 12: 6-(5-Cyclopropylmethylcarbamoyl-2-methyl-phenyl)-N-(4-methoxyphenyl)-nicotinamide




25

6-[5-Cyclopropylmethylcarbamoyl-2-methyl-phenyl]-N-(4-methoxyphenyl)-nicotinamide was prepared from 6-chloro-N-(4-methoxyphenyl)nicotinamide (Intermediate 2) and N-cyclopropylmethyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 10) using General Method B.

- 5 LCMS: retention time 3.12min, MH^+ 416. NMR: δH [2H_6]-DMSO 10.39,(1H, s), 9.21,(1H, d), 8.63,(1H, t), 8.41,(1H, dd), 7.96,(1H, s), 7.86,(1H, d), 7.79,(1H, d), 7.71,(2H, d), 7.44,(1H, d), 6.96,(2H, d), 3.76,(3H, s), 3.15,(2H, t), 2.41,(3H, s), 1.03,(1H, m), 0.43,(2H, m), 0.23,(2H, m).


10 **Example 13: 6-[5-(Fur-3-ylcarbamoylamino)-2-methyl-phenyl]-N-(4-methoxyphenyl)-nicotinamide**



6-[5-(Fur-3-ylcarbamoylamino)-2-methyl-phenyl]-N-(4-methoxyphenyl)-nicotinamide was prepared from 6-chloro-N-(4-methoxyphenyl)nicotinamide (Intermediate 2) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide (Intermediate 13) using General Method B. LCMS: retention time 3.19min, MH^+ 428. NMR: δH [2H_6]-DMSO 10.38,(1H, s), 10.00,(1H, s), 9.19,(1H, s), 8.38,(2H, m), 7.83,(1H, s), 7.80,(1H, s), 7.76,(1H, s), 7.73-7.69,(3H, m), 7.32,(1H, s), 7.01,(1H, s), 6.96,(2H, d), 3.76,(3H, s), 2.34,(3H, s).


20

Example 14: 6-(5-Cyclopropylcarbamoyl-2-methyl-phenyl)-N-(3-methoxybenzyl)-nicotinamide




- 25 6-[5-Cyclopropylcarbamoyl-2-methyl-phenyl]-N-(3-methoxybenzyl)-nicotinamide was prepared from 6-chloro-N-(3-methoxybenzyl)nicotinamide (Intermediate 3) and N-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 8) using General Method B. LCMS: retention time 2.94min, MH^+ 416. NMR: δH [2H_6]-DMSO 9.29,(1H, t), 9.15,(1H, s), 8.48,(1H, d), 8.35,(1H, d), 7.89,(1H, s), 7.81,(1H, d), 7.72,(1H, d), 7.41,(1H, d), 7.26,(1H, t), 6.93,(2H, m), 6.84,(1H, s), 4.51,(2H, d), 3.75,(3H, s), 2.86,(1H, m), 2.38,(3H, s), 0.69,(2H, m), 0.57,(2H, m).
- 30

Example 15: N-(3-Methoxybenzyl)-6-[2-methyl-5-(thiadiazol-2-ylcarbamoyl)-phenyl]-nicotinamide



5 N-(3-Methoxybenzyl)-6-[2-methyl-5-(thiadiazol-2-ylcarbamoyl)-phenyl]-nicotinamide was prepared from 6-chloro-N-(3-methoxybenzyl)nicotinamide (Intermediate 3) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiadiazol-2-yl)-benzamide (Intermediate 12) using General Method B. LCMS: retention time 3.02min, MH^+ 460. NMR: δH [2H_6]-DMSO 13.14,(1H, b), 9.32,(1H, t), 9.24,(1H, s), 9.18,(1H, d), 8.40,(1H, dd), 8.26,(1H, s), 8.10,(1H, d), 7.84,(1H, d),
10 7.55,(1H, d), 7.27,(1H, t), 6.94,(2H, m), 6.84,(1H, d), 4.53,(2H, d), 3.75,(3H, s), 2.46,(3H, s).


Example 16: N-(3-Methoxybenzyl)-6-[2-methyl-5-(thiazol-2-ylcarbamoyl)-phenyl]-nicotinamide



15


N-(3-Methoxybenzyl)-6-[2-methyl-5-(thiazol-2-ylcarbamoyl)-phenyl]-nicotinamide was prepared from 6-chloro-N-(3-methoxybenzyl)nicotinamide (Intermediate 3) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiazol-2-yl)-benzamide (Intermediate 11) using General Method B. LCMS: retention
20 time 3.20min, MH^+ 459. NMR: δH [2H_6]-DMSO 12.71,(1H, b), 9.31,(1H, t), 9.17,(1H, d), 8.39,(1H, dd), 8.22,(1H, s), 8.07,(1H, d), 7.83,(1H, d), 7.57,(1H, d), 7.52,(1H, d), 7.29-7.25,(2H, m), 6.94,(2H, m), 6.84,(1H, d), 4.52,(2H, d), 3.75,(3H, s), 2.45,(3H, s).

Example 17: 6-[5-Cyclopropylmethylcarbamoyl-2-methyl-phenyl]-N-(3-methoxybenzyl)-nicotinamide



5 6-[5-Cyclopropylmethylcarbamoyl-2-methyl-phenyl]-N-(3-methoxybenzyl)-nicotinamide was prepared from 6-chloro-N-(3-methoxybenzyl)nicotinamide (Intermediate 3) and N-cyclopropylmethyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 10) using General Method B. LCMS: retention time 3.07min, MH^+ 430. NMR: δH [2H_6]-DMSO 9.30,(1H, t), 9.16,(1H, d), 8.62,(1H, t), 8.36,(1H, dd), 7.94,(1H, s), 7.85,(1H, d), 7.74,(1H, d), 10 7.43,(1H, d), 7.27,(1H, t), 6.94-6.92,(2H, m), 6.84,(1H, d), 4.51,(2H, d), 3.75,(3H, s), 3.14,(2H, t), 2.39,(3H, s), 1.03,(1H, m), 0.43,(2H, m), 0.23,(2H, m).


Example 18: 6-[5-(Fur-3-ylcarbamoylamino)-2-methyl-phenyl]-N-(3-methoxybenzyl)-nicotinamide



15


6-[5-(Fur-3-ylcarbamoylamino)-2-methyl-phenyl]-N-(3-methoxybenzyl)-nicotinamide was prepared from 6-chloro-N-(3-methoxybenzyl)nicotinamide (Intermediate 3) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide (Intermediate 13) using General Method B. LCMS: retention time 20 3.17min, MH^+ 442. NMR: δH [2H_6]-DMSO 9.99,(1H, s), 9.29,(1H, t), 9.15,(1H, d), 8.38,(1H, s), 8.34,(1H, dd), 7.81,(2H, m), 7.75,(1H, d), 7.67,(1H, d), 7.31-7.25,(2H, m), 7.00,(1H, s), 6.94,(2H, m), 6.84,(1H, d), 4.51,(2H, d), 3.75,(3H, s), 2.32,(3H, s).

Example 19: N-(3-Methoxybenzyl)-6-[5-(thiophen-3-ylcarbonylamino)-2-methyl-phenyl]-nicotinamide




N-(3-Methoxybenzyl)-6-[5-(thiophen-3-ylcarbonylamino)-2-methyl-phenyl]-
 5 nicotinamide was prepared from 6-chloro-N-(3-methoxybenzyl)nicotinamide
 (Intermediate 3) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-
 phenyl]thiophene-3-amide (Intermediate 14) using General Method B. LCMS:
 retention time 3.27min, MH^+ 458. NMR: δH [2H_6]-DMSO 10.12,(1H, s), 9.29,(1H, t),
 9.15,(1H, d), 8.35-8.32,(2H, m), 7.86,(1H, s), 7.78,(1H, d), 7.68-7.65,(3H, m), 7.32-
 10 7.24,(2H, m), 6.94,(2H, m), 6.84,(2H, d), 4.51,(2H, d), 3.75,(3H, s), 2.32,(3H, s).

Example 20: 6-(5-Cyclopropylcarbamoyl-2-methyl-phenyl)-N-(3-methylsulphonylaminobenzyl)-nicotinamide




6-(5-Cyclopropylcarbamoyl-2-methyl-phenyl)-N-(3-
 15 methylsulphonylaminobenzyl)-nicotinamide was prepared from 6-chloro-N-(3-
 methylsulphonylaminobenzyl)nicotinamide (Intermediate 4) and N-cyclopropyl-4-
 methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 8)
 using General Method B. LCMS: retention time 2.71min, MH^+ 479. NMR: δH [2H_6]-
 20 DMSO 9.33,(1H, t), 9.15,(1H, s), 8.48-8.33,(3H, m), 7.89,(1H, s), 7.81,(1H, d),
 7.73,(1H, d), 7.41,(1H, d), 7.31,(1H, t), 7.21,(1H, s), 7.10,(2H, m), 4.51,(2H, d),
 2.99,(3H, s), 2.86,(1H, m), 2.38,(3H, s), 0.69,(2H, m), 0.57,(2H, m).

Example 21: N-(3-Methylsulphonylaminobenzyl)-6-[2-methyl-5-(thiadiazol-2-ylcarbamoyl)-phenyl]-nicotinamide




N-(3-Methylsulphonylaminobenzyl)-6-[2-methyl-5-(thiadiazol-2-ylcarbamoyl)-phenyl]-nicotinamide was prepared from 6-chloro-N-(3-methylsulphonylaminobenzyl)nicotinamide (Intermediate 4) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiadiazol-2-yl)-benzamide (Intermediate 12) using General Method B. LCMS: retention time 2.80min, MH^+ 523. NMR: δH [2H_6]-DMSO 9.35,(1H, t), 9.17,(2H, m), 8.38,(1H, d), 8.26,(1H, s), 8.09,(1H, d), 7.83,(1H, d), 7.52,(1H, d), 7.31,(1H, t), 7.22,(1H, s), 7.11,(2H, m), 4.52,(2H, d), 2.99,(3H, s), 2.46,(3H, s).

Example 22: N-(3-Methylsulphonylaminobenzyl)-6-[2-methyl-5-(thiazol-2-ylcarbamoyl)-phenyl]-nicotinamide




N-(3-Methylsulphonylaminobenzyl)-6-[2-methyl-5-(thiazol-2-ylcarbamoyl)-phenyl]-nicotinamide was prepared from 6-chloro-N-(3-methylsulphonylaminobenzyl)nicotinamide (Intermediate 4) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiazol-2-yl)-benzamide (Intermediate 11) using General Method B. LCMS: retention time 2.96min, MH^+ 522. NMR: δH [2H_6]-DMSO 10.19,(2H, b), 9.35,(1H, t), 9.17,(1H, s), 8.38,(1H, dd), 8.22,(1H, s), 8.07,(1H, d), 7.84,(1H, d), 7.57,(1H, d), 7.52,(1H, d), 7.31-7.28,(2H, m), 7.22,(1H, s), 7.11,(2H, m), 4.52,(2H, d), 2.99,(3H, s), 2.45,(3H, s).

Example 23: 6-(5-Cyclopropylmethylcarbamoyl-2-methyl-phenyl)-N-(3-methylsulphonylaminobenzyl)-nicotinamide



- 5 6-(5-Cyclopropylmethylcarbamoyl-2-methyl-phenyl)-N-(3-methylsulphonylaminobenzyl)-nicotinamide was prepared from 6-chloro-N-(3-methylsulphonylaminobenzyl)nicotinamide (Intermediate 4) and N-cyclopropylmethyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 10) using General Method B. LCMS: retention time 2.88min, MH^+ 493. NMR: δH [2H_6]-DMSO 9.34,(1H, t), 9.16,(1H, d), 8.96,(1H, b),
10 8.62,(1H, t), 8.35,(1H, dd), 7.94,(1H, s), 7.85,(1H, d), 7.75,(1H, d), 7.43,(1H, d), 7.31,(1H, t), 7.21,(1H, s), 7.11,(2H, m), 4.52,(2H, d), 3.14,(2H, t), 2.99,(3H, s), 2.39,(3H, s), 1.03,(1H, m), 0.43,(2H, m), 0.23,(2H, m).


15 **Example 24: 6-[5-(Fur-3-ylcarbonylamino)-2-methyl-phenyl]-N-(3-methylsulphonylaminobenzyl)-nicotinamide**



- 20 6-[5-(Fur-3-ylcarbonylamino)-2-methyl-phenyl]-N-(3-methylsulphonylaminobenzyl)-nicotinamide was prepared from 6-chloro-N-(3-methylsulphonylaminobenzyl)nicotinamide (Intermediate 4) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide (Intermediate 13) using General Method B. LCMS: retention time 2.93min, MH^+ 505. NMR: δH [2H_6]-DMSO 9.99,(1H, s), 9.32,(1H, t), 9.15,(1H, d), 8.95,(1H, b), 8.38,(1H, s), 8.33,(1H, dd), 7.81,(2H, d), 7.75,(1H, d), 7.68,(1H, d), 7.33-7.30,(2H, m), 7.21,(1H, s), 7.11,(2H, m), 7.01,(1H, s), 4.51,(2H, d), 2.99,(3H, s), 2.32,(3H, s).

25


Example 25: N-(3-Methylsulphonylaminobenzyl)-6-[5-(thiophen-3-ylcarbonylamino)-2-methyl-phenyl]-nicotinamide



5 N-(3-Methylsulphonylaminobenzyl)-6-[5-(thiophen-3-ylcarbonylamino)-2-methyl-phenyl]-nicotinamide was prepared from 6-chloro-N-(3-methylsulphonylaminobenzyl)nicotinamide (Intermediate 4) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]thiophene-3-amide (Intermediate 14) using General Method B. LCMS: retention time 3.03min, MH^+ 521. NMR: δH [2H_6]-DMSO 10.12,(1H, s), 9.33,(1H, t), 9.15,(1H, s), 8.78,(1H, b), 8.36-8.32,(2H, m), 7.86,(1H, s), 7.78,(1H, d), 7.69-7.65,(3H, m), 7.31,(2H, m), 7.21,(1H, s), 7.11,(2H, m), 4.51,(2H, d), 2.99,(3H, s), 2.32,(3H, s).

10

Example 26: 6-(5-Cyclopropylcarbamoyl-2-methyl-phenyl)-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]-nicotinamide




15

6-(5-Cyclopropylcarbamoyl-2-methyl-phenyl)-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]-nicotinamide was prepared from 6-chloro-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]nicotinamide (Intermediate 5) and N-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 8) using General Method B. LCMS: retention time 2.36min, MH^+ 484. NMR: δH [2H_6]-DMSO 11.70,(1H, b), 9.23,(1H, s), 8.50,(1H, d), 8.38,(1H, d), 8.33,(1H, d), 7.92,(1H, s), 7.83,(2H, m), 7.43,(1H, d), 7.36,(1H, t), 7.29,(1H, d), 7.11,(1H, t), 3.77,(2H, s), 2.87,(1H, m), 2.67-2.24,(11H, m), 2.13,(3H, s), 0.70,(2H, m), 0.58,(2H, m).


20

Example 27: N-[2-(4-Methylpiperazin-1-ylmethyl)phenyl]-6-[2-methyl-5-(thiadiazol-2-ylcarbamoyl)-phenyl]-nicotinamide



5 N-[2-(4-Methylpiperazin-1-ylmethyl)phenyl]-6-[2-methyl-5-(thiadiazol-2-ylcarbamoyl)-phenyl]-nicotinamide was prepared from 6-chloro-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]nicotinamide (Intermediate 5) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiadiazol-2-yl)-benzamide (Intermediate 12) using General Method B. LCMS: retention time 2.43min, MH^+ 528. NMR: δH [2H_6]-DMSO 13.07,(1H, b), 11.74,(1H, s), 9.26,(1H, s), 9.21,(1H, s),
10 8.43,(1H, d), 8.34,(1H, d), 8.29,(1H, s), 8.12,(1H, d), 7.93,(1H, d), 7.56,(1H, d), 7.36,(1H, t), 7.29,(1H, d), 7.11,(1H, t), 3.78,(2H, s), 2.67-2.26,(11H, m), 2.11,(3H, s).


Example 28: N-[2-(4-Methylpiperazin-1-ylmethyl)phenyl]-6-[2-methyl-5-(thiazol-2-ylcarbamoyl)-phenyl]-nicotinamide



15


20 N-[2-(4-Methylpiperazin-1-ylmethyl)phenyl]-6-[2-methyl-5-(thiazol-2-ylcarbamoyl)-phenyl]-nicotinamide was prepared from 6-chloro-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]nicotinamide (Intermediate 5) and 4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiazol-2-yl)-benzamide (Intermediate 11) using General Method B. LCMS: retention time 2.53min, MH^+ 527. NMR: δH [2H_6]-DMSO 12.73,(1H, b), 11.70,(1H, b), 9.26,(1H, d), 8.43,(1H, dd), 8.33,(1H, d), 8.25,(1H, s), 8.10,(1H, d), 7.93,(1H, d), 7.58,(1H, d), 7.54,(1H, d), 7.36,(1H, t), 7.30,(2H, m), 7.12,(1H, t), 3.78,(2H, s), 2.67-2.25,(11H, b), 2.14,(3H, s).

Example 29: 6-(5-Cyclopropylmethylcarbamoyl-2-methyl-phenyl)-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]-nicotinamide




6-(5-Cyclopropylmethylcarbamoyl-2-methyl-phenyl)-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]-nicotinamide was prepared from 6-chloro-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]nicotinamide (Intermediate 5) and N-cyclopropylmethyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 10) using General Method B. LCMS: retention time 2.46min, MH^+ 498. NMR: δH [2H_6]-DMSO 11.67,(1H, b), 9.24,(1H, s), 8.63,(1H, t), 8.39,(1H, d), 8.32,(1H, d), 7.97,(1H, s), 7.88-7.83,(2H, m), 7.45,(1H, d), 7.36,(1H, t), 7.30,(1H, d), 7.11,(1H, t), 3.77,(2H, s), 3.15,(2H, t), 2.70-2.21,(11H, m), 1.04,(1H, m), 0.43,(2H, m), 0.23,(2H, m).

Example 30: 6-[5-(Fur-3-ylcarbonylamino)-2-methyl-phenyl]-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]-nicotinamide



6-[5-(Fur-3-ylcarbonylamino)-2-methyl-phenyl]-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]-nicotinamide was prepared from 6-chloro-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]nicotinamide (Intermediate 5) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide (Intermediate 13) using General Method B. LCMS: retention time 2.53min, MH^+ 510. NMR: δH [2H_6]-DMSO 11.64,(1H, b), 10.02,(1H, s), 9.23,(1H, s), 8.38,(2H, m), 8.31,(1H, d), 7.86,(1H, s), 7.80,(1H, s), 7.76,(2H, m), 7.38-7.29,(3H, m), 7.11,(1H, t), 7.01,(1H, s), 3.77,(2H, s), 2.66-2.20,(11H, m), 2.16,(3H, s).






Example 31: N-[2-(4-Methylpiperazin-1-ylmethyl)phenyl]-6-[5-(thiophen-3-ylcarbonylamino)-2-methyl-phenyl]-nicotinamide











- N-[2-(4-Methylpiperazin-1-ylmethyl)phenyl]-6-[5-(thiophen-3-ylcarbonylamino)-2-methyl-phenyl]-nicotinamide was prepared from 6-chloro-N-[2-(4-methylpiperazin-1-ylmethyl)phenyl]nicotinamide (Intermediate 5) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]thiophene-3-amide (Intermediate 14) using General Method B. LCMS: retention time 2.58min, MH^+ 526. NMR: δH [2H_6]-DMSO 11.64,(1H, b), 10.14,(1H, s), 9.23,(1H, s), 8.38,(2H, m), 8.31,(1H, d), 7.91,(1H, s), 7.79-7.75,(2H, m), 7.65,(2H, m), 7.38-7.29,(3H, m), 7.11,(1H, t), 3.77,(2H, s), 2.67-2.24,(11H, m), 2.16,(3H, m).

General Method C

- The 6-chloronicotinamide (25mg), N-cyclopropyl-5-fluoro-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 34, 15mg), tetrakis(triphenylphosphino)palladium (2mg) and aqueous sodium hydrogen carbonate (1M, 0.5ml) were mixed in propan-2-ol (2ml) and heated at reflux for 18 hours. The propan-2-ol was evaporated and the residue diluted with ethylacetate / cyclohexane (1:2). The solution was applied to a SPE (Si, 2g) and eluted with ethylacetate / cyclohexane (1:2) and then ethylacetate. The solvent was evaporated from the ethylacetate fraction and the residue triturated with ether to give the desired product as a white solid.
- Examples 32 to 44 may also be prepared using {5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}boronic acid (Intermediate 36) in place of Intermediate 34.

Compound	Structure	6-Chloro-N-	MH ⁺	Retention time (minutes)
Example 32 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-cyclopropylmethyl-nicotinamide		6-Chloro-N-cyclopropylmethylnicotinamide (Intermediate 21)	368	2.78
Example 33 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-(3-methylbutyl)-nicotinamide		6-Chloro-N-(3-methylbutyl)nicotinamide (Intermediate 22)	384	3.10
Example 34 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-cyclobutylmethyl-nicotinamide		6-Chloro-N-cyclobutylmethylnicotinamide (Intermediate 32)	382	3.01
Example 35 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-(1-cyclopropylethyl)-nicotinamide		6-Chloro-N-(1-cyclopropylethyl)nicotinamide (Intermediate 23)	382	2.95
Example 36 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-(2,2-dimethylpropyl)-nicotinamide		6-Chloro-N-(2,2-dimethylpropyl)nicotinamide (Intermediate 24)	384	3.01

Example 37 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methylphenyl)-N-(2,2-dimethylcyclopropyl)-nicotinamide		6-Chloro-N-(2,2-dimethylcyclopropyl)nicotinamide (Intermediate 25)	382	2.90
Example 38 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methylphenyl)-N-cyclopropyl-nicotinamide		6-Chloro-N-cyclopropylnicotinamide (Intermediate 26)	354	2.60
Example 39 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methylphenyl)-N-cyclohexylmethyl-nicotinamide		6-Chloro-N-cyclohexylmethylnicotinamide (Intermediate 27)	410	3.22
Example 40 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methylphenyl)-N-cyclobutyl-nicotinamide		6-Chloro-N-cyclobutylnicotinamide (Intermediate 28)	368	2.79
Example 41 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methylphenyl)-N-(2-methylpropyl)-nicotinamide		6-Chloro-N-(2-methylpropyl)nicotinamide (Intermediate 29)	370	2.86
Example 42 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methylphenyl)-N-(1-methylpropyl)-nicotinamide		6-Chloro-N-(1-methylpropyl)nicotinamide (Intermediate 33)	370	2.84

Example 43 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methylphenyl)-N-propyl-nicotinamide		6-Chloro-N-propylnicotinamide (Intermediate 30)	356	2.72
Example 44 6-(5-Cyclopropylcarbamoyl-3-fluoro-2-methylphenyl)-N-cyclopentyl-nicotinamide		6-Chloro-N-cyclopentylnicotinamide (Intermediate 31)	382	2.92

General Method D

Intermediate 38 (40 μ mol) in DMF (0.5ml) was treated with HATU (1.12eq) and DIPEA (3eq). On shaking a solution was formed which was added to a solution of amine (1.2 – 2.0eq) in DMF (0.5ml). After shaking the reactions were left overnight at room temperature. The solvent was removed *in vacuo*, the residue dissolved in chloroform (1.0ml) and applied to an SPE (NH₂, 0.5g). The product was eluted with chloroform (1.5ml), ethyl acetate (1.5ml) and methanol/ethyl acetate (1:9, 1.5ml). The solvent was evaporated under vacuum from the product fraction.

10

Compound	Amine	MH ⁺	Retention time (minutes)
Example 45 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(1,3-thiazol-2-ylmethyl)nicotinamide	2-aminomethylthiazole	411	2.79
Example 46 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[2-(1,3-thiazol-2-yl)ethyl]nicotinamide	2-(2-aminoethyl)thiazole	425	2.78
Example 47	2-methylbenzylamine	418	3.26

6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2-methylbenzyl)nicotinamide			
Example 48 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,4,5-trifluorobenzyl)nicotinamide	2,4,5-trifluorobenzylamine	458	3.29
Example 49 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,5-difluorobenzyl)nicotinamide	2,5-difluorobenzylamine	440	3.21
Example 50 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3,4-difluorobenzyl)nicotinamide	3,4-difluorobenzylamine	440	3.24
Example 51 N-(3-chlorobenzyl)-6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl} nicotinamide	3-chlorobenzylamine	438	3.33
Example 52 N-(4-chlorobenzyl)-6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl} nicotinamide	4-chlorobenzylamine	438	3.34
Example 53 N-(3-chloro-2-fluorobenzyl)-6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-	3-chloro-2-fluorobenzylamine	456	3.36

methylphenyl}nicotinamide			
Example 54 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(5-fluoro-2-methylbenzyl)nicotinamide	5-fluoro-2-methylbenzylamine	436	3.30
Example 55 N-(2-chloro-3,6-difluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide	2-chloro-3,6-difluorobenzylamine	474	3.31
Example 56 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(1-methylbutyl)nicotinamide	2-pentylamine	384	3.14
Example 57 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,3-difluoro-4-methylbenzyl)nicotinamide	2,3-difluoro-4-methylbenzylamine	454	3.36
Example 58 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,3,5-trifluorobenzyl)nicotinamide	2,3,6-trifluorobenzylamine	458	3.29
Example 59 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(1,3-	1,3-dimethylbutylamine	398	3.28

dimethylbutyl)nicotinamide			
Example 60 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3-fluoro-4-methylbenzyl)nicotinamide	3-fluoro-4-methylbenzylamine	436	3.32
Example 61 N-(5-chloro-2-fluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide	5-chloro-2-fluorobenzylamine	456	3.36
Example 62 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2-phenylethyl)nicotinamide	2-phenylethylamine	418	3.20
Example 63 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2-methoxy-2-methylpropyl)nicotinamide	2-methoxy-2-methylpropylamine	400	2.79
Example 64 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2-methoxyethyl)nicotinamide	2-methoxyethylamine	372	2.63
Example 65 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3,3-	3,3-dimethylbutylamine	398	3.30

dimethylbutyl)nicotinamide			
Example 66 N-(2-tert-butoxyethyl)-6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide	2-t-butoxyethylamine	414	2.98
Example 67 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(tert-pentyl)nicotinamide	1,1-dimethylpropylamine	384	3.17
Example 68 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[(4-methyl-1,3-thiazol-2-yl)methyl]nicotinamide	2-(aminomethyl)-4-methylthiazole	425	2.88
Example 69 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2-hydroxy-1,1-dimethylpentyl)nicotinamide	1,1-dimethyl-2-hydroxypentylamine	428	3.08
Example 70 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[2-(trifluoromethyl)benzyl]nicotinamide	2-trifluoromethylbenzylamine	472	3.38
Example 71 N-(2-chlorobenzyl)-6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide	2-chlorobenzylamine	438	3.30

Example 72 N-cyclopropyl-3-fluoro-4-methyl-5-{5-[(4-methylpiperidin-1-yl)carbonyl]pyridin-2-yl}benzamide	4-methylpiperidine	396	3.09
Example 73 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-isobutyl-N-methylnicotinamide	N-isobutyl-N-methylamine	384	3.02
Example 74 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(4-fluorobenzyl)nicotinamide	4-fluorobenzylamine	422	3.19
Example 75 N-cyclopropyl-3-{5-[(3,3-diethylazetid-1-yl)carbonyl]pyridin-2-yl}-5-fluoro-4-methylbenzamide	3,3-diethylazetidine	410	3.24
Example 76 N-cyclopentyl-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-methylnicotinamide	N-cyclopropyl-N-methylamine	396	3.06
Example 77 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-ethyl-N-isopropylnicotinamide	N-ethyl-N-isopropylamine	384	2.98
Example 78	2,3,4-trifluorobenzylamine	458	3.30

6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,3,4-trifluorobenzyl)nicotinamide			
Example 79 N-benzyl-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide	benzylamine	404	3.14
Example 80 N-cyclopropyl-3-{5-[(2-ethylpiperidin-1-yl)carbonyl]pyridin-2-yl}-5-fluoro-4-methylbenzamide	2-ethylpiperidine	410	3.17
Example 81 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[3-(trifluoromethyl)benzyl]nicotinamide	3-trifluoromethylbenzylamine	472	3.40
Example 82 N-cyclopropyl-3-{5-[(2-ethyl-2-methylpiperidin-1-yl)carbonyl]pyridin-2-yl}-5-fluoro-4-methylbenzamide	2-ethyl-2-methylpiperidine	424	3.30
Example 83 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(1,1-dimethylbutyl)nicotinamide	1,1-dimethylbutylamine	398	3.30
Example 84 N-(4-chloro-2-fluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-	4-chloro-2-fluorobenzylamine	456	3.37

methyphenyl}nicotinamide			
Example 85 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3,3,3-trifluoropropyl)nicotinamide	3,3,3-trifluoropropylamine	410	3.00
Example 86 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[4-(trifluoromethyl)benzyl]nicotinamide	4-trifluoromethylbenzylamine	472	3.41
Example 87 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3-methoxyphenyl)nicotinamide	3-methoxyaniline	420	3.26
Example 88 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(4-methylphenyl)nicotinamide	4-methylaniline	404	3.34
Example 89 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3-fluorobenzyl)nicotinamide	3-fluorobenzylamine	422	3.20
Example 90 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[(5-	2-aminomethyl-5-methylfuran	408	3.09

methyl-2-furyl)methyl]nicotinamide			
Example 91 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3-methylphenyl)nicotinamide	3-methylaniline	404	3.36
Example 92 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,3-difluorobenzyl)nicotinamide	2,3-difluorobenzylamine	440	3.23
Example 93 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2,4-difluorobenzyl)nicotinamide	2,4-difluorobenzylamine	440	3.23
Example 94 N-(3-chloro-4-fluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide	3-chloro-4-fluorobenzylamine	456	3.37
Example 95 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(4-methylbenzyl)nicotinamide	4-methylbenzylamine	418	3.27
Example 96 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3,4,5-trifluorobenzyl)nicotinamide	3,4,5-trifluorobenzylamine	458	3.33

Example 97 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[(3-methylthien-2-yl)methyl]nicotinamide	2-aminomethyl-3-methylthiophene	424	3.21
Example 98 N-(3-chloro-2,6-difluorobenzyl)-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}nicotinamide	3-chloro-2,6-difluorobenzylamine	474	3.35
Example 99 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[(2-ethylcyclopropyl)methyl]nicotinamide	(2-ethylcyclopropyl)methylamine (Intermediate 39)	396	3.24
Example 100 N-cyclopropyl-3-fluoro-4-methyl-5-{5-[(2-propylpiperidin-1-yl)carbonyl]pyridin-2-yl}benzamide	2-propylpiperidine	424	3.33
Example 101 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[2-(4-methyl-1,3-thiazol-2-yl)ethyl]nicotinamide	2-(2-aminoethyl)-4-methylthiazole	439	2.65
Example 102 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2-fluoro-	2-fluoro-2-phenylethylamine	436	3.07

2-phenylethyl)nicotinamide			
Example 103 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-[1-methyl-2-(1,3-thiazol-2-yl)ethyl]nicotinamide	2-(2-aminopropyl)thiazole	439	2.70
Example 104 N-cyclopropyl-3-{5-[(2,4-dimethylpiperidin-1-yl)carbonyl]pyridin-2-yl}-5-fluoro-4-methylbenzamide	2,4-dimethylpiperidine	410	3.07
Example 105 N-cyclopropyl-3-{5-[(2,3-dimethylpiperidin-1-yl)carbonyl]pyridin-2-yl}-5-fluoro-4-methylbenzamide	2,3-dimethylpiperidine	410	3.02
Example 106 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3-methylbut-2-enyl)nicotinamide	3-methylbut-2-enylamine	382	2.97
Example 107 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-methyl-N-(1-methylcyclopentyl)nicotinamide	N-methyl-N-(1-methylcyclopentyl)amine	410	3.14
Example 108 N-(2-cyclopentylethyl)-6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-	N-(2-cyclopentylethyl)-N-methylamine	424	3.26

methylephenyl}-N-methylnicotinamide			
Example 109 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylephenyl}-N-[(2E)-1,1-dimethylbut-2-enyl]nicotinamide	1,1-dimethylbut-2-enylamine	396	3.06
Example 110 N-cyclopropyl-3-fluoro-4-methyl-5-{5-[(2,2,3-trimethylpyrrolidin-1-yl)carbonyl]pyridin-2-yl}benzamide	2,2-dimethyl-3-methyl-pyrrolidine	410	3.07
Example 111 N-cyclopropyl-3-{5-[(3-ethylpiperidin-1-yl)carbonyl]pyridin-2-yl}-5-fluoro-4-methylbenzamide	3-ethylpiperidine	410	3.10
Example 112 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylephenyl}-N-(3,3,3-trifluoro-2-methylpropyl)nicotinamide	2-methyl-3,3,3-trifluoropropylamine	424	3.00
Example 113 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylephenyl}-N-(1-ethyl-1-methylpropyl)nicotinamide	1-ethyl-1-methylpropylamine	398	3.15
Example 114 N-cyclopropyl-3-fluoro-4-methyl-5-{5-[(2-methylpiperidin-1-	2-methylpiperidine	396	2.90

yl)carbonyl]pyridin-2-yl} benzamide			
Example 115 N-cyclopropyl-3-{5-[(3,3-dimethylpiperidin-1-yl)carbonyl]pyridin-2-yl}-5-fluoro-4-methylbenzamide	3,3,-dimethylpiperidine	410	3.04
Example 116 N-cyclopropyl-3-fluoro-4-methyl-5-{5-[(3-methylpiperidin-1-yl)carbonyl]pyridin-2-yl} benzamide	3-methylpiperidine	396	2.95
Example 117 N-cyclohexyl-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-ethylnicotinamide	N-cyclohexyl-N-ethylamine	424	3.18
Example 118 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-ethylnicotinamide	ethylamine	342	2.58
Example 119 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-isopropyl-N-methylnicotinamide	N-isopropyl-N-methylamine	370	2.72
Example 120 6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2-hydroxy-1-methylpentyl)nicotinamide	3-hydroxy-1-methylpentylamine	414	2.88

Example 121 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(2-fluorobenzyl)nicotinamide	2-fluorobenzylamine	422	3.18
Example 122 6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(3-methylbenzyl)nicotinamide	3-methylbenzylamine	418	3.27
Example 123 N-(cyclopentylmethyl)-6-{5- [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl} nicotinamide	cyclopentylmethylamine	396	3.23

Abbreviations

	DCM	Dichloromethane
	DIPEA	N,N-Diisopropylethylamine
5	DME	Dimethoxyethane
	DMF	Dimethylformamide
	DMSO	Dimethylsulphoxide
	HATU	O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
10	HOBT	1-Hydroxybenzotriazole hydrate
	SPE	bond-elut (solid phase extraction column)

The activity of the compounds of the invention as p38 inhibitors may be demonstrated in the following assays:

15

p38 Kinase Assay

The peptide substrate used in the p38 assay was biotin-IPTSPITTTYFFFRRR-amide. The p38 and MEK6 proteins were purified to homogeneity from E.coli expression systems. The fusion proteins were tagged at the N-terminus with
 20 Glutathione-S-Transferase (GST). The maximum activation was achieved by incubating 20uL of a reaction mixture of 30nM MEK6 protein and 120nM p38 protein in the presence of 1.5uM peptide and 10mM Mg(CH₃CO₂)₂ in 100mM HEPES, pH

7.5, added to 15uL of a mixture of 1.5uM ATP with 0.08uCi [33 P]ATP, with or without 15uL of inhibitor in 6%DMSO. The controls were reactions in the presence (negative controls) or absence (positive controls) of 50 mM EDTA. Reactions were allowed to proceed for 60 min at room temperature and quenched with addition of 50uL of 250mM EDTA and mixed with 150uL of Streptavidin SPA beads (Amersham) to 0.5mg/reaction. The Dynatech Microfluor white U-bottom plates were sealed and the beads were allowed to settle overnight. The plates were counted in a Packard TopCount for 60 seconds. IC₅₀ values were obtained by fitting raw data to $\%I = 100 * (1 - (I - C2) / (C1 - C2))$, where I was CPM of background, C1 was positive control, and C2 was negative control.

α P38 Fluorescence Polarisation Method


α P38 was prepared in house. SB4777790-R Ligand was diluted in HEPES containing MgCl₂, CHAPS, DTT and DMSO. This was added to blank wells of a Black NUNC 384 well plate. α P38 was added to this ligand mixture then added to the remainder of the 384 well plate containing controls and compounds. The plates were read on an LJI Analyst and Fluorescence Anisotropy used to calculate the compound inhibition.

20 Results

The compounds described in the Examples were tested as described above and had IC₅₀ values of <10 μ M.

Claims:

1. A compound of formula (I):



(I)

wherein

- 10 R^1 is selected from hydrogen, C_{1-6} alkyl optionally substituted by up to three groups selected from C_{1-6} alkoxy, halogen and hydroxy, C_{2-6} alkenyl, C_{3-7} cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, phenyl optionally substituted by up to three groups selected from R^5 and R^6 , and heteroaryl optionally substituted by up to three groups selected from R^5 and R^6 ,

- 15 R^2 is selected from hydrogen, C_{1-6} alkyl and $-(CH_2)_q-C_{3-7}$ cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, or $(CH_2)_mR^1$ and R^2 , together with the nitrogen atom to which they are bound, form a four- to six-membered heterocyclic ring optionally substituted by up to three C_{1-6} alkyl groups;

R^3 is chloro or methyl;

- 20 R^4 is the group $-NH-CO-R^7$ or $-CO-NH-(CH_2)_q-R^8$;

R^5 is selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_q-C_{3-7}$ cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, $-CONR^9R^{10}$, $-NHCOR^{10}$, $-SO_2NHR^9$, $-(CH_2)_sNHSO_2R^{10}$, halogen, CN, OH, $-(CH_2)_sNR^{11}R^{12}$, and trifluoromethyl;

- 25 R^6 is selected from C_{1-6} alkyl, C_{1-6} alkoxy, halogen, trifluoromethyl and $-(CH_2)_sNR^{11}R^{12}$;

- R^7 is selected from hydrogen, C_{1-6} alkyl, $-(CH_2)_q-C_{3-7}$ cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, trifluoromethyl, $-(CH_2)_r$ heteroaryl optionally substituted by R^{13} and/or R^{14} , and $-(CH_2)_r$ phenyl optionally substituted by R^{13} and/or R^{14} ;


30 R^8 is selected from hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups, $CONHR^9$, phenyl optionally substituted by R^{13} and/or R^{14} , and heteroaryl optionally substituted by R^{13} and/or R^{14} ;

- R^9 and R^{10} are each independently selected from hydrogen and C_{1-6} alkyl,
 or R^9 and R^{10} , together with the nitrogen atom to which they are bound, form
 a five- to six-membered heterocyclic ring optionally containing one additional
 heteroatom selected from oxygen, sulfur and $N-R^{15}$, wherein the ring may be
 5 substituted by up to two C_{1-6} alkyl groups;
 R^{11} is selected from hydrogen, C_{1-6} alkyl and $-(CH_2)_q-C_{3-7}$ cycloalkyl
 optionally substituted by one or more C_{1-6} alkyl groups,
 R^{12} is selected from hydrogen and C_{1-6} alkyl,
 or R^{11} and R^{12} , together with the nitrogen atom to which they are bound,
 10 form a five or six-membered heterocyclic ring optionally containing one additional
 heteroatom selected from oxygen, sulfur and $N-R^{15}$;
 R^{13} is selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_q-C_{3-7}$ cycloalkyl
 optionally substituted by one or more C_{1-6} alkyl groups, $-CONR^9R^{10}$, $-NHCOR^{10}$,
 halogen, CN, $-(CH_2)_sNR^{11}R^{12}$, trifluoromethyl, phenyl optionally substituted by one
 15 or more R^{14} groups and heteroaryl optionally substituted by one or more R^{14} groups;
 R^{14} is selected from C_{1-6} alkyl, C_{1-6} alkoxy, halogen, trifluoromethyl and -
 $NR^{11}R^{12}$;
 R^{15} is selected from hydrogen and methyl;
 X and Y are each independently selected from hydrogen, methyl and halogen;
 20 Z is halogen;
 m is selected from 0, 1, 2, 3 and 4, wherein each carbon atom of the resulting
 carbon chain may be optionally substituted with up to two groups selected
 independently from C_{1-6} alkyl and halogen;
 n is selected from 0, 1 and 2;
 25 q is selected from 0, 1 and 2;
 r is selected from 0 and 1; and
 s is selected from 0, 1, 2 and 3, or
 a pharmaceutically acceptable salt or solvate thereof.
- 30 2. A compound according to claim 1 wherein R^1 is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-7} cycloalkyl optionally substituted by one or more C_{1-6} alkyl groups,
 phenyl optionally substituted by up to three substituents selected from R^5 and R^6 ,
 heteroaryl optionally substituted by up to three substituents selected from R^5 and R^6 .
- 35 3. A compound according to claim 1 or claim 2 wherein R^2 is hydrogen.
4. A compound according to any one of the preceding claims wherein R^3 is
 methyl.
- 40 5. A compound according to any one of the preceding claims wherein X is
 fluorine.
6. A compound according to any one of the preceding claims wherein R^7 is
 45 selected from C_{1-6} alkyl, $-(CH_2)_q-C_{3-7}$ cycloalkyl, trifluoromethyl, $-(CH_2)_r$ heteroaryl
 optionally substituted by R^{13} and/or R^{14} , and $-(CH_2)_r$ phenyl optionally substituted

by C₁₋₆alkyl, C₁₋₆alkoxy, -(CH₂)_q-C₃₋₇cycloalkyl, -CONR⁹R¹⁰, -NHCOR¹⁰, halogen, CN, trifluoromethyl, phenyl optionally substituted by one or more R¹⁴ groups and/or heteroaryl optionally substituted by one or more R¹⁴ groups.

- 5 7. A compound according to any one of the preceding claims wherein R⁸ is selected from C₃₋₇cycloalkyl, CONHR⁹, heteroaryl optionally substituted by R¹³ and/or R¹⁴, and phenyl optionally substituted by C₁₋₆alkyl, C₁₋₆alkoxy, -(CH₂)_q-C₃₋₇cycloalkyl, -CONR⁹R¹⁰, -NHCOR¹⁰, halogen, CN, trifluoromethyl, phenyl optionally substituted by one or more R¹⁴ groups and/or heteroaryl optionally substituted by one or more R¹⁴ groups.
- 10 8. A compound according to claim 1 as defined in any one of Examples 1 to 123.
- 15 9. A compound according to claim 1 which is 6-(5-cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-(2,2-dimethylpropyl)-nicotinamide, or a pharmaceutically acceptable salt or solvate thereof.
- 20 10. A compound according to claim 1 which is 6-(5-cyclopropylcarbamoyl-3-fluoro-2-methyl-phenyl)-N-(2,2-dimethylpropyl)-nicotinamide.
- 25 11. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 10, or a pharmaceutically acceptable salt or solvate thereof, in admixture with one or more pharmaceutically acceptable carriers, diluents or excipients.
- 30 12. A method for treating a condition or disease state mediated by p38 kinase activity or mediated by cytokines produced by the activity of p38 kinase comprising administering to a patient in need thereof a compound as claimed in any one of claims 1 to 10, or a pharmaceutically acceptable salt or solvate thereof.
- 35 13. A compound as claimed in any one of claims 1 to 10, or a pharmaceutically acceptable salt or solvate thereof, for use in therapy.
- 40 14. Use of a compound as claimed in any one of claims 1 to 10, or a pharmaceutically acceptable salt or solvate thereof, in the manufacture of a medicament for use in the treatment of a condition or disease state mediated by p38 kinase activity or mediated by cytokines produced by the activity of p38 kinase.
15. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 10, or a pharmaceutically acceptable salt or solvate thereof, which comprises
 - (a) reacting a compound of formula (II)


71



(II)

in which R^1 , R^2 , Z , m and n are as defined in claim 1 and W is halogen,
with a compound of formula (III)

5




(III)

in which R^3 , R^4 , X and Y are as defined in claim 1,
in the presence of a catalyst, or


10

(b) reacting a compound of formula (VIII)



(VIII)

15 with a compound of formula (III) as hereinbefore defined and then reacting the acid
thus formed with an amine of formula (V)




(V)

20 in which R^1 , R^2 and m are as defined in claim 1,
under amide forming conditions, or

(c) reacting a compound of formula (II) as hereinbefore defined with a compound
of formula (IX)

72



(IX)

in which R³, R⁴, X and Y are as defined in claim 1,
in the presence of a catalyst.

5