

PRINS REACTION AND COMPOUNDS USEFUL IN THE SYNTHESIS OF HALICHONDRIN MACROLIDES AND ANALOGS THEREOF

Background


5 The invention relates to intermediates useful in the synthesis of pharmaceutically active macrolide compounds and methods of synthesizing macrolide compounds. Halichondrin B is a potent anticancer agent originally isolated from the marine sponge *Halichondria okadai*, and subsequently found in *Axinella* sp., *Phakellia carteri*, and *Lissodendoryx* sp. A total synthesis of halichondrin B was published in 1992 (Aicher, T. D. et al., J. Am. Chem. Soc. 114:3162-3164). Further synthetic and structure-activity relationship studies
10 have been disclosed in U.S. Patent Nos. 5,338,865 and 5,436,238 and in Towle et al., *Annual Meeting of the American Association for Cancer Research*, April 6-10, 2002, 5721 and Wanget al., *Bioorg. Med. Chem. Lett.*, 10:1029-1032, 2000. Eribulin mesylate (also called Halaven®, E7389, and the mesylate salt of B1939), a nontaxane microtubule dynamics inhibitor, is a structurally simplified, synthetic analog of halichondrin B. Methods and intermediates for the synthesis of certain halichondrin B analogs and
15 intermediates are described in International Publication Nos. WO 2005/118565, WO 2009/046308, WO 2009/064029, and WO 2009/124237, WO 2015/066729, and WO 2016/179607; U.S. Patent No. 6,214,865; Austad et al., *Synlett* 24(3):333-337, 2013; Austad et al., *Synlett*, 24(3):327-332, 2013; and Chase et al., *Synlett* 24(3):323-326, 2013. New methods for the synthesis of halichondrin and its analogs (e.g., macrolide analogs) are desirable.

20

Summary of the Invention

In general, the present invention provides methods and compounds useful for the preparation of macrocyclic intermediates in the synthesis of a halichondrin macrolide or analog thereof. In particular, the methods disclosed herein may be useful in the preparation of a halichondrin macrolide or analog thereof through a
25 synthesis including a C.23-C.24 bond-forming Prins reaction. Prins reaction precursors may be prepared using a Sakurai reaction, as described herein.


In one aspect, the invention provides a method of preparing a compound of formula (A):



30

where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10
35 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, or A is a group of formula (1):



where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;


5 R_2 is H and P_1 is absent, H, alkyl, or a hydroxyl protecting group, or R_2 and P_1 combine to form a bond;

(i) R_3 is H, and P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

10 (ii) R_3 is $-(CH_2)_nNP_3P_4$, where P_3 is H or an N-protecting group, and (a) P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_4 is H or an N-protecting group, (b) P_2 and P_4 combine to form an alkylidene, or (c) each of P_2 and P_4 is H;

15 (iii) R_3 is $-(CH_2)_nOP_5$, where P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_5 is H, optionally substituted alkyl, or a hydroxyl protecting group; or P_2 and P_5 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

25 each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_B(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl,


alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

n, when present, is 0, 1, or 2;

k is 0 or 1;

5 R₁ is -OP₆, -CH(Y)₂, or -CH₂(Y), where P₆ is H or a hydroxyl protecting group;

R₄ is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl,



where

10 each P₇ is independently H or a hydroxyl protecting group; and

R₈ is -CH₂CH₂-COOR_C, -CH=CH-COOR_C, -CH₂CH₂-SO₂R_D, or -CH=CH-SO₂R_D;

R₆ is H, optionally substituted alkyl, or optionally substituted arylalkyl;


each Y is independently $-\text{COOR}_C$ or $-\text{SO}_2\text{R}_D$;

each R_C, when present, is independently optionally substituted alkyl, optionally substituted

15 aryl, or optionally substituted arylalkyl; and

each R_D, when present, is independently optionally substituted aryl or optionally substituted alkyl.


The method includes producing the compound of formula (A) from a compound of formula (B), a compound of formula (C), and R_5OH , where R_5 is optionally substituted acyl. The compound of formula (B) is:



where all variables are as described for the compound of formula (A).

25 The compound of formula (C) is:


R₄-R₇,
(C)



In some embodiments, the producing the compound of formula (A) includes reacting the compound of formula (B), the compound of formula (C), R_5OH , and a Lewis acid (e.g., an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)).

- 5 In certain embodiments, the method further includes producing the compound of formula (B) including reacting a compound of formula (D), a compound of formula (E), and a second Lewis acid.


The compound of formula (D) is:



10

where all variables are as described for the compound of formula (B).

The compound of formula (E) is:



15


where R_5 is silyl, and R_6 is as described for the compound of formula (B).

- In some embodiments, the second Lewis acid is an oxophilic second Lewis acid (e.g., boron trifluoride or a solvate thereof). In particular embodiments, the Lewis acid and the second Lewis acid are same. In further
20 embodiments, the Lewis acid and the second Lewis acid are different.

- In certain embodiments, the preparing the compound of formula (B) and the preparing the compound of formula (A) are performed as a single-pot transformation. In other embodiments, the method further includes
25 epimerizing the product of reacting the compound of formula (D), the compound of formula (E), and the second Lewis acid.


- In another aspect, the invention provides a method of preparing a compound of formula (B). The compound of formula (B) can be used as a reactant in the Prins reaction described herein. The method includes
30 producing the compound of formula (B) from a compound of formula (D) and a compound of formula (E).

The compound of formula (B) is:



where

- 5 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, or A is a group of formula (1):



where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;



15 R₂ is H and P₁ is absent, H, alkyl, or a hydroxyl protecting group, or R₂ and P₁ combine to form a bond;

(i) R₃ is H, and P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

(ii) R₃ is $-(CH_2)_nNP_3P_4$, where P₃ is H or an N-protecting group, and (a) P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₄ is H or an N-protecting group, (b) P₂ and P₄ combine to form an alkylidene, or (c) each of P₂ and P₄ is H;

(iii) R₃ is $-(CH_2)_nOP_5$, where P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₅ is H, optionally substituted alkyl, or a hydroxyl protecting group; or P₂ and P₅, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R₃ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;
 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,
 NR_B(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of
 R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl,
 alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or
 10 heterocyclic radical-alkyl;

n, when present, is 0, 1, or 2;

k is 0 or 1;

R₁ is -OP₆, -CH(Y)₂, or -CH₂(Y), where P₆ is H or a hydroxyl protecting group;


R₆ is H, optionally substituted alkyl, or optionally substituted aryl; and

15 Y is independently -COOR_C or -SO₂R_D;

R_C, when present, is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and

R_D, when present, is optionally substituted aryl or optionally substituted non-enolizable alkyl.


The compound of formula (D) is:



where all variables are as described for the compound of formula (B).

5

The compound of formula (E) is:



where R₅ is silyl, and R₆ is as described for the compound of formula (D).

10

In some embodiments, the producing the compound of formula (B) includes reacting the compound of formula (D), the compound of formula (E), and a Lewis acid (e.g., an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)).


15

In certain embodiments, the method further includes epimerizing the product of reacting the compound of formula (D), the compound of formula (E), and the Lewis acid.

In yet another aspect, the invention provides methods of preparing a halichondrin macrolide or analog thereof.

20

In particular embodiments, the halichondrin macrolide or an analogue thereof is:




a halichondrin macrolide or analog thereof,

where

25

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated

hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, or A is a group of formula (1):



5 where

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;


R₂ is H and P₁ is absent, H, alkyl, or a hydroxyl protecting group, or R₂ and P₁ combine to form a bond;

10 (i) R₃ is H, and P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;


(ii) R₃ is $-(\text{CH}_2)_n\text{NP}_3\text{P}_4$, where P₃ is H or an N-protecting group, and (a) P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₄ is H or an N-protecting group, (b) P₂ and P₄ combine to form an alkylidene, or (c) each of P₂ and P₄ is H;

15 (iii) R₃ is $-(\text{CH}_2)_n\text{OP}_5$, where P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₅ is H, optionally substituted alkyl, or a hydroxyl protecting group; or P₂ and P₅, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R₃ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



20



, where each P' is independently H or a hydroxyl protecting group;

each of A₁, A₂, and A₃ is independently H or OP'', where each P'' is independently H or a hydroxyl protecting group;

25 E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , $NR_B R_A$, $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_B(CO)NR_B R_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_B R_A$, or $O(CO)NR_B R_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

5 n, when present, is 0, 1, or 2;

k is 0 or 1; and

X_1 is $-CH_2-$ or $-O-$.


In certain embodiments, the method includes

10 (A) producing a compound of formula (IA) from a compound of formula (IIC), a compound of formula (III), and R_5OH , where R_5 is optionally substituted acyl;

and


(B) producing the halichondrin macrolide or analog thereof from the compound of formula (IA).

15 The compound of formula (IA) is:



where

20 R_1 is $-OP_6$, $-CH(Y)_2$, or $-CH_2(Y)$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-COOR_C$ or $-SO_2R_D$;



where

25 each P_7 is independently H or a hydroxyl protecting group;

R_8 is $-CH_2CH_2-COOR_C$, $-CH=CH-COOR_C$, $-CH_2CH_2-SO_2R_D$, or $-CH=CH-SO_2R_D$;


each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and

30 each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

R_5 is optionally substituted acyl; and

the remaining variables are as described for the halichondrin macrolide or analog thereof.


The compound of formula (IIC) is:



where all variables are as described for the compound of formula (IA).

The compound of formula (III) is:



where R_7 is $-\text{CHO}$ or $R_{7A}\text{O}$ , where each R_{7A} is independently an optionally substituted alkyl; and R_4 is as described for the compound of formula (IA).

In particular embodiments, the method includes:

(A) producing a compound of formula (IVB) from a compound of formula (IIA), a compound of formula (IIB), a compound of formula (IIIA), a compound of formula (IVA), and $R_5\text{OH}$, where R_5 is optionally substituted acyl;


(B) producing a compound of formula (IVC) from the compound of formula (IVB);

(C) producing a compound of formula (IVD) from the compound of formula (IVC);

and

(D) producing the halichondrin macrolide or analog thereof from the compound of formula (IVD).

The compound of formula (IIA) is:



where

R_1 is $-\text{OP}_6$, $-\text{CH}(\text{Y})_2$, or $-\text{CH}_2(\text{Y})$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-\text{COOR}_C$ or $-\text{SO}_2\text{R}_D$;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$,

where

m is 1, and R_F is a hydroxyl protecting group,

or


m is 0, and

(i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene;

and the remaining variables are as described for the halichondrin macrolide or analog thereof.

The compound of formula (IIB) is:




where R_S is silyl.

The compound of formula (IIIA) is:




where

R_{4A} is




where R_C is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl, and each P_7 is independently H or a hydroxyl protecting group;

R_7 is $-\text{CHO}$ or



where each R_{7A} is independently an optionally substituted alkyl.

The compound of formula (IVA) is:



where

each P_9 is independently a hydroxyl protecting group;

R_9 is $-\text{CHO}$ or $-\text{COOH}$;

(a1) R_{10} is H or a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;

(a2) R_{10} is H or a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond;

or

(a3) R_{10} and R_{11} combine to form a bond, and R_{12} is H;

(b1) A_1 and R_{14} combine to form oxo, R_{13} is H or a hydroxyl protecting group, and R_{15} is H;

or

(b2) A_1 is H or $-\text{OP}''$, and:

(i) R_{13} is H or a hydroxyl protecting group, and R_{14} and R_{15} combine to form a double bond;

or

(ii) R_{13} and R_{14} combine to form a bond, and R_{15} is H or $-\text{OP}''$; and

(c1) R_{16} is H, and P_8 is H or a hydroxyl protecting group;


or

(c2) R_{16} and P_8 combine to form a double bond;

and

each P'' , when present, is independently H or a hydroxyl protecting group.

The compound of formula (IVB) is:




(IVB)

where

X_1 is $-O-$, $-C(Y)_2-$, $-CH(Y)-$, or $-CH_2-$;

X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(CH(OP_{12}))-$, where P_{12} is H or a hydroxyl protecting group; and

5 R_G is




where

R_5 is an optionally substituted acyl; and


10 X_2 is H or $-CH_2-X_{2A}-CH_2-CH=CH_2$, where X_{2A} is $-O-$, $-C(R_H)_2-$, or $-NR_1-$, where each R_H is independently H or $-COOR_J$, R_1 is an *N*-protecting group, and R_J is C_{1-6} alkyl; and the remaining variables are as described for the compound of formula (IIA) and the compound of formula (IVA).

The compound of formula (IVC) is:



where


L_1 is



20 where L_2 and L_3 combine to form a bond;

and the remaining variables are as described for the compound of formula (IVB).

The compound of formula (IVD) is:




where each of A₁ and A₂ is independently H or –OPⁿ, and the remaining variables are as described for the compound of formula (IVC).

5

In further embodiments, step (A) includes:

(E) producing a compound of formula (IA) from the compound of formula (IIA), the compound of formula (IIIA), and R₅OH, where the compound of formula (IA) is:



10

where R_{4A} is as described for the compound of formula (IIIA), and the remaining variables are as described for the compound of formula (IA) above;


and

(F) producing the compound of formula (IVB) from the compound of formula (IF) and the compound of formula (IVA).


15

In yet further embodiments, step (E) includes reacting the compound of formula (IIA), the compound of formula (IIIA), R_5OH , and a Lewis acid (e.g., an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)).


- 5 In still further embodiments, step (F) includes reacting R_{4A} that is



with a glycol cleaving agent, and a Horner-Wadsworth-Emmons reaction with $(R_CO)_2P(O)-CH_2-COOR_C$ to produce R_{4A} that is



- 10 In other embodiments, step (F) includes reacting R_{4A} that is



with a 1,2-reducing agent and further reacting with an allyl halide or allyl pseudohalide.


- 15 In yet other embodiments, R_9 is $-CHO$, X_1 is $-C(Y)_2-$, $-CH(Y)-$, or $-CH_2-$, and step (A) includes reacting R_9 with R_1 that is $-CH(Y)_2$ or $-CH_2(Y)$ treated with a Brønsted base. In still other embodiments, X_1 is $-O-$, and step (A) includes esterifying R_1 that is $-OP_6$, where P_6 is H, with R_9 that is $-COOH$.

In some embodiments, step (B) includes reacting the compound of formula (IVB) with an olefin metathesis catalyst.

20

In particular embodiments, P_9 is H, R_{16} is H, and step (C) includes:


oxidizing the compound of formula (IVC) to produce a compound of formula (IVCa):



(IVCa)

- 25 where all variables are as described for the compound of formula (IVC);
and reacting the compound of formula (IVCa) with a 1,4-reducing agent to produce the compound of formula (IVD).

In certain embodiments, the compound of formula (IVB) is, or is converted to, a compound of formula (IVBa) prior to producing the compound of formula (IVC), where the compound of formula (IVBa) is:



(IVBa)

where all variables are as described for the compound of formula (IVB).

In further embodiments, the producing the halichondrin macrolide or analog thereof includes reacting the compound of formula (IVD) with a hydroxyl protecting group removing agent.

In yet further embodiments, the method includes:

(A) producing a compound of formula (VB) from a compound of formula (IIA), a compound of formula (IIB), a compound of formula (IIIB), a compound of formula (VA), and R_5OH ,


where R_5 is optionally substituted acyl;

(B) producing a compound of formula (VC) from the compound of formula (VB),

and

(C) producing the halichondrin macrolide or analog thereof from the compound of formula (VC).

The compound of formula (IIA) is:



(IIA)

where

R_1 is $-OP_6$, $-CH(Y)_2$, or $-CH_2(Y)$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-COOR_c$ or $-SO_2R_D$;

each R_c , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

R_D , when present, is optionally substituted aryl or optionally substituted non-enolizable alkyl;

R_E is $-CHO$ or $-CH_{(1+m)}(OR_F)_{(2-m)}$,

where

m is 1, and R_F is a hydroxyl protecting group,

or


m is 0, and

(i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene;

5 and the remaining variables are as described for the halichondrin macrolide or analog thereof.

The compound of formula (IIB) is:




10 where R_S is silyl.

The compound of formula (IIIB) is:



15 where


R_{4B} is but-3-en-1-yl,




and

R_7 is $-CHO$ or

20



The compound of formula (VA) is:



25

where

a designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or a designates (*S*)-stereogenic center, and Z is OR_{16} , where R_{16} is a hydroxyl protecting group;

30

(a1) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;

(a2) R_{10} is a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond;

or

(a3) R_{10} and R_{11} combine to form a bond, and R_{12} is H;

5

(b1) A_1 and R_{14} combine to form oxo, R_{13} is a hydroxyl protecting group, and R_{15} is H;

or

(b2) A_1 is H or $-OP''$, and:

10

(i) R_{13} is a hydroxyl protecting group, and R_{14} and R_{15} combine to form a double bond;

or

(ii) R_{13} and R_{14} combine to form a bond, and R_{15} is H or $-OP''$;


Y_1 is iodide, bromide, or trifluoromethanesulfonate;

R_9 is $-CHO$ or $-COOH$;

15

each P_9 is independently a hydroxyl protecting group, and X_3 is oxo; or both P_9 groups and X_3 , together with the atoms to which each is attached, combine to form a ketal; and each P'' is independently a hydroxyl protecting group.

The compound of formula (VB) is:



20

(VB)

where


X_1 is $-CH_2-$, $-CH(Y)-$, $-C(Y)_2-$, or $-O-$, where each Y is independently $-COOR_c$ or $-SO_2R_D$;

25

each R_c , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl; and

R_k is




30

R_5 is optionally substituted acyl; and


the remaining variables are as described for the compound of formula (VA) and the compound of formula (IIA).

The compound of formula (VC) is:



where

L₁ is



where L₂ and L₃ combine to form a bond;

b designates (*S*)-stereogenic center, if *a* designates (*R*)-stereogenic center;


and

b designates (*R*)-stereogenic center, if *a* designates (*S*)-stereogenic center; and

the remaining variables are as described for the compound of formula (VB).

In certain embodiments, step (A) includes:

(D) producing a compound of formula (ID) from the compound of formula (IIA), the compound of formula (IIB), the compound of formula (IIIB), and R₅OH, where R₅ is optionally substituted acyl, and where the compound of formula (ID) is:



where all variables are as described for the compound of formula (IIA), the compound of formula (IIB), the compound of formula (IIIB), and R₅OH;


and

(E) producing the compound of formula (VB) from the compound of formula (ID) and the compound of formula (VA).

In some embodiments, the producing the compound of formula (ID) includes reacting the compound of formula (IIC), the compound of formula (IIIB), R_5OH , and a Lewis acid (e.g., an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)).

In certain embodiments, step (D) includes a Sakurai reaction between the compound of formula (IIA) and the compound of formula (IIB) to produce a compound of formula (IIC), and a Prins reaction between the compound of formula (IIC), the compound of formula (IIIB), and R_5OH .



In further embodiments, the compound of formula (IIC) is:



where all variables are as described for the compound of formula (IIA).



In yet further embodiments, X_1 is $-CH_2-$, $-CH(Y)-$, or $-C(Y)_2-$, and step (A) includes reacting R_9 , where R_9 is $-CHO$, with R_1 that is $-CH(Y)_2$ or $-CH_2(Y)$ treated with a Brønsted base, to produce $-CO-X_1-$. In still further embodiments, X_1 is $-O-$, and step (A) includes esterifying R_9 that is $-COOH$ with R_1 that is $-OP_6$, where P_6 is H, to produce $-CO-X_1-$.

In other embodiments, step (B) includes reacting the compound of formula (VB) with a Cr(II) salt and a Ni(II) salt. In yet other embodiments, step (C) includes a step of nucleophilic ring-closing of the compound of formula (VC). In still other embodiments, both P_9 groups and X_3 , together with the atoms to which each is attached, combine to form a ketal. In some embodiments, R_{10} and R_{11} combine to form a bond, and R_{12} is H. In certain embodiments, R_{4B} is but-3-en-1-yl, and step (A) includes reacting the but-3-en-1-yl group with a dihydroxylating agent to produce R_{4B} that is





30

In particular embodiments, R_{4B} is




5 In further embodiments, R_{4B} is



In yet further embodiments, the method includes:

- 10 (A) producing a compound of formula (IF) from a compound of formula (IIA), a compound of formula (IIB), a compound of formula (IIIC), and R_5OH , where R_5 is optionally substituted acyl;
- (B) producing a compound of formula (IG) from the compound of formula (IF),
- and
- (C) producing the halichondrin macrolide or analog thereof from the compound of formula (IG) and
- 15 one or more intermediates.

The compound of formula (IF) is:




20 where

R_1 is $-OP_6$, $-CH(Y)_2$, or $-CH_2(Y)$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-COOR_C$ or $-SO_2R_D$;


each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

25 each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;



- 5 R_5 is optionally substituted acyl; and
 the remaining variables are as described for the halichondrin macrolide or analog thereof.

The compound of formula (IIA) is:



10

where

R_1 is $-\text{OP}_6$, $-\text{CH}(\text{Y})_2$, or $-\text{CH}_2(\text{Y})$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-\text{COOR}_C$ or $-\text{SO}_2\text{R}_D$;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

15

R_D , when present, is optionally substituted aryl or optionally substituted non-enolizable alkyl;

R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$,

where

m is 1, and R_F is a hydroxyl protecting group,

20

or

m is 0, and


(i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene; and

the remaining variables are as described for the compound of formula (IF).

25

The compound of formula (IIB) is:



where R_S is silyl.


30

The compound of formula (IIIC) is:




where R_7 is $-\text{CHO}$ or $\text{R}_{7A}\text{O}-\text{CH}(\text{OR}_{7A})-\text{CH}_3$, where each R_{7A} is independently an optionally substituted alkyl, and R_{4C} is as described for the compound of formula (IF).

The compound of formula (IG) is:




where
 R_{E1} is




the remaining variables are as described for the compound of formula (IF).


In some embodiments, step (B) includes reacting R_{4C} that is




with a glycol cleaving agent to produce an aldehyde, reacting the aldehyde with $(\text{R}_C\text{O})_2\text{P}(\text{O})\text{-CH}_2\text{-COOR}_C$ to produce R_{4C} that is





In certain embodiments, step (B) includes reducing R_{4C} that is



with a 1,4-reducing agent to produce R_{4C} that is




In further embodiments, step (B) includes reducing R_{4C} that is




5

In yet further embodiments, step (B) includes reducing R_{4C} that is




with a 1,2-reducing agent to produce a carbonyl group in R_{E1} .

10 In still further embodiments, the step reducing R_{4C} that is




In particular embodiments, step (B) includes reacting R_{4C} that is



15

with an oxidizing agent capable of converting an alcohol to a carbonyl group in R_{E1} .

In certain embodiments, the one or more intermediates is a compound of formula (VIA):



20

where

R_{10A} is $-\text{CH}_2-\text{CH}=\text{CH}_2$, $-(\text{CH}_2)_2-\text{CH}=\text{CH}_2$, or $-(\text{CH}_2)_3-\text{OP}_{10}$;

each R_{10} , R_{13} , P_9 and P_{10} is independently a hydroxyl protecting group;


A_1 is H or $-\text{OP}''$; and

Y_1 is chloro, bromo, iodo, trifluoromethanesulfonate, or trialkylsilane;

and step (C) includes:

- 5 (D) producing a compound of formula (VIB) from the compound of formula (IG) and the compound of formula (VIA).


The compound of formula (VIB) is:



(VIB)

where

L_4 is



R_1 is $-\text{OP}_6$ or $-\text{CH}(Y)$, where P_6 is H or a hydroxyl protecting group, and Y is $-\text{COOR}_C$ or $-\text{CHO}$;

A_1 and R_{14} combine to form oxo, R_{13} is H or a hydroxyl protecting group, and R_{15} is H;

or

A_1 is H or $-\text{OP}''$, and:

- (i) R_{13} is H or a hydroxyl protecting group, and R_{14} and R_{15} combine to form a double bond;

or

- (ii) R_{13} and R_{14} combine to form a bond, and R_{15} is H or $-\text{OP}''$;

each P_9 is independently a hydroxyl protecting group, and X_3 is oxo or X_3 , together with the carbon atom to which it is attached, is $-(\text{CH}(\text{OP}_{11}))$, where P_{11} is H or a hydroxyl protecting group; or both P_9 groups and X_3 , together with the atoms to which each is attached, combine to form a ketal; and

the remaining variables are as described for the compound of formula (VIA) and the compound of formula (IG).

In yet further embodiments, step (D) includes reacting the compound of formula (VIA) and the compound of formula (IG) with a Cr(II) salt and a Ni(II) salt.

In still further embodiments, R_{10A} is $-\text{CH}_2-\text{CH}=\text{CH}_2$, and step (C) includes:


(E) producing a compound of formula (VIC) from the compound of formula (VIB) and $R_{17}-\text{CH}_2\text{CH}=\text{CH}_2$, where R_{17} is $-\text{COOH}$ or a metallic or metalloid moiety;

(F) producing a compound of formula (VID) from the compound of formula (VIC);

and

(G) producing the halichondrin macrolide or analog thereof from the compound of formula (VID).


The compound of formula (VIC) is:



(VIC)

where

L_5 is




X_1 is $-\text{CH}_2-$ or $-\text{O}-$;

X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P_{12} is H or a hydroxyl protecting group; and

the remaining variables are as described for the compound of formula (VIB).


The compound of formula (VID) is:



(VID)

where

L₁ is



5

In some embodiments, R₁ is -CH₂(Y), X₁ is -CH₂-, X₄ is oxo or -(CH(OP₁₂))-, and step (E) includes reacting the compound of formula (VIB) with R₁₇-CH₂CH=CH₂, where R₁₇ is a metallic or metalloid moiety. In other embodiments, R₁ is -OP₆, P₆ is H, X₁ is -O-, X₄ is oxo, and step (E) includes esterifying the compound of formula (VIB) with R₁₇-CH₂CH=CH₂, where R₁₇ is -COOH. In yet other embodiments, the producing step (F) includes reacting the compound of formula (VIC) with an olefin metathesis catalyst. In still other

10

embodiments, step (G) includes reacting the compound of formula (VID) with a hydroxyl protecting group removing agent.

In certain embodiments, R_{10A} is -CH₂-CH₂-CH=CH₂, R₁ is -OP₆ or -CH₂(Y), where P₆ is H or a hydroxyl protecting group, and Y is -COOR_C, and step (C) includes:

15

(H) producing a compound of formula (VIE) from the compound of formula (VIB) and R₁₇-CH=CH₂, where R₁₇ is -COOH or a metallic or metalloid moiety;


(I) producing a compound of formula (VIF) from the compound of formula (VIE);

and

20


(J) producing the halichondrin macrolide or analog thereof from the compound of formula (VIF).

The compound of formula (VIE) is:



25

where




X_1 is $-\text{CH}_2-$ or $-\text{O}-$; and

X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P_{12} is H or a hydroxyl protecting group; and

the remaining variables are as described for the compound of formula (VIB).


5

The compound of formula (VIF) is:



(VIF)

where



10

L_1 is OR_5 Me or Me, where L_2 and L_3 combine to form a bond; and

the remaining variables are as described for the compound of formula (VIE).

In some embodiments, step (I) includes reacting the compound of formula (VIE) with an olefin metathesis

catalyst. In particular embodiments, step (J) includes reacting the compound of formula (VIF) with a hydroxyl

15 protecting group removing agent.

In certain embodiments, R_{10A} is $-(\text{CH}_2)_3-\text{OP}_{10}$, and step (C) includes:

(K) producing a compound of formula (VIG) from the compound of formula (VIB) and $(\text{R}_C\text{O})_2\text{P}(\text{O})-\text{CH}_2-\text{R}_P$, where each R_C is independently optionally substituted alkyl, optionally substituted aryl, or

20

optionally substituted arylalkyl, and R_P is H or $-\text{COOH}$;


(L) producing a compound of formula (VIF) from the compound of formula (VIG);

and

(M) producing the halichondrin macrolide or analog thereof from the compound of formula (VIF).

25


The compound of formula (VIG) is:



(VIG)

where

L₇ is




5

X₁ is $-\text{CH}_2-$ or $-\text{O}-$; and

the remaining variables are as described for the compound of formula (VIB) and for $(\text{R}_\text{C}\text{O})_2\text{P}(\text{O})-\text{CH}_2-\text{R}_\text{P}$.


10 The compound of formula (VIF) is:



(VIF)

where

L₁ is



15

X₄ is oxo; and

the remaining variables are as described for the compound of formula (VIG).

In further embodiments, the producing step (K) includes reacting the compound of formula (VIB) with $(R_cO)_2P(O)-CH_2-R_P$ under Claisen reaction conditions, where R_P is H and R_1 is $-CH_2(Y)$. In yet further embodiments, the producing step (K) includes esterifying the compound of formula (VIB) with $(R_cO)_2P(O)-CH_2-R_P$, where R_P is $-COOH$ and R_1 is $-OP_6$, where P_6 is H. In still further embodiments, step (M) includes

5 reacting the compound of formula (VIF) with a hydroxyl protecting group removing agent.

In some embodiments, step (C) includes:


(N) producing a compound of formula (VIIB) from the compound of formula (IG) and a compound of formula (VIIA);

10 (O) producing a compound of formula (VIIC) from the compound of formula (VIIB);

and

(P) producing the halichondrin macrolide or analog thereof from the compound of formula (VIIC).

The compound of formula (VIIA) is:



where

(i) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;

(ii) R_{10} is a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond;

20 or

(iii) R_{10} and R_{11} combine to form a bond, and R_{12} is H;

R_9 is $-CH_2-OP''$ or $-COOP''$;


R_{13} and each P_9 is independently a hydroxyl protecting group;

A_1 is H or $-OP''$;


25 each P'' is independently H or a hydroxyl protecting group; and

Y_1 is chloro, bromo, iodo, trifluoromethanesulfonate, or trialkylsilane.

The compound of formula (VIIB) is:



where



R_1 is $-\text{OP}_6$, $-\text{CH}_2(\text{Y})$, or $-\text{CH}(\text{Y})_2$, where P_6 is H or a hydroxyl protecting group, and Y is $-\text{COOR}_C$ or $-\text{CHO}$;

A_1 and R_{14} combine to form oxo, R_{13} is H or a hydroxyl protecting group, and R_{15} is H;

or

A_1 is H or $-\text{OP}''$, and:

(i) R_{13} is H or a hydroxyl protecting group, and R_{14} and R_{15} combine to form a double bond;

or


(ii) R_{13} and R_{14} combine to form a bond, and R_{15} is H or $-\text{OP}''$;

R_9 is $-\text{CHO}$ or $-\text{COOP}''$;

each P_9 is independently a hydroxyl protecting group, and X_3 is oxo or X_3 , together with the carbon atom to which it is attached, is $-(\text{CH}(\text{OP}_{11}))$, where P_{11} is H or a hydroxyl protecting group; or both P_9 groups and X_3 , together with the atoms to which each is attached, combine to form a ketal; and

the remaining variables are as described for the compound of formula (IG) and the compound of formula (VIIA).

The compound of formula (VIIC) is:



each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(CH(OP_{12}))_n-$, where P_{12} is H or a hydroxyl protecting group; and

5 the remaining variables are as described for the compound of formula (VIIB).

In certain embodiments, step (N) includes reacting the compound of formula (VIIA) and the compound of formula (IH) with a Cr(II) salt and a Ni(II) salt. In particular embodiments, step (O) includes an esterification reaction between R_1 that is $-OP_6$, where P_6 is H, and R_9 that is $-COOH$. In further embodiments, step (O) includes a reaction between R_1 that is $-CH(Y)_2$ or $-CH_2(Y)$, and R_9 that is $-CHO$. In yet further

10 embodiments, step (P) includes reacting the compound of formula (VIIC) with a hydroxyl protecting group removing agent.

In still further embodiments, the method includes:


15 (A) producing a compound of formula (VIID) from a compound of formula (IIA), a compound of formula (IIB), a compound of formula (IIIC), a compound of formula (VIIA), and R_5OH , where R_5 is optionally substituted acyl;

(B) producing a compound of formula (VIIE) from the compound of formula (VIID);

and

20 (C) producing the halichondrin macrolide or analog thereof from the compound of formula (VIIC).

The compound of formula (IIA) is:



25 where

R_1 is $-OP_6$, $-CH(Y)_2$, or $-CH_2(Y)$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-COOR_C$ or $-SO_2R_D$;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

30 each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

R_E is $-CHO$ or $-CH_{(1+m)}(OR_F)_{(2-m)}$,

where

m is 1, and R_F is a hydroxyl protecting group,

35 or


m is 0, and

(i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene; and

the remaining variables are as described for the halichondrin macrolide or analog thereof.

The compound of formula (IIB) is:




where R_S is silyl.

The compound of formula (IIIC) is:




where

R₇ is -CHO or , where each R_{7A} is independently an optionally substituted alkyl;
R_{4C} is



where each P₇, when present, is independently H or a hydroxyl protecting group.

The compound of formula (VIIA) is:



where

- (i) R₁₀ is a hydroxyl protecting group, R₁₁ is alkyl ether, and R₁₂ is H;
 - (ii) R₁₀ is a hydroxyl protecting group, and R₁₁ and R₁₂ combine to form a double bond;
- or

- (iii) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;


R₉ is -CHO or -COOH;

R₁₃ and each P₉ is independently a hydroxyl protecting group;

A₁ is H or -OPⁿ, where Pⁿ is H or a hydroxyl protecting group; and

Y₁ is chloro, bromo, iodo, trifluoromethanesulfonate, or trialkylsilane.

The compound of formula (VIID) is:




5 where

X₁ is $-\text{CH}_2-$, $-\text{CH}(\text{Y})-$, $-\text{C}(\text{Y})_2-$, or $-\text{O}-$, where each Y is independently $-\text{COOR}_c$ or $-\text{SO}_2\text{R}_d$;

X₄ is oxo, or X₄, together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P₁₂ is H or a hydroxyl protecting group;

R_L is



where


R₁₈ is $-\text{CHO}$ or $-\text{CH}_2\text{OP}_7$;

R₅ is optionally substituted acyl; and

the remaining variables are as described for the compound of formula (IIA).


15

The compound of formula (VIIE) is:



where

L₁ is



where L₂ and L₃ combine to form a bond;

each P₉ is independently a hydroxyl protecting group, and X₃ is oxo or X₃, together with the carbon atom to which it is attached, is $-(CH(OP_{11}))-$, where P₁₁ is H or a hydroxyl protecting group; and
 5 the remaining variables are as described for the compound of formula (VIID).

In some embodiments, step (A) includes:


(D) producing a compound of formula (IIC) from the compound of formula (IIA) and the compound of
 10 formula (IIB);

(E) producing a compound of formula (IF) from the compound of formula (IIC), the compound of formula (IIIC), and R₅OH;

and


(F) producing the compound of formula (VIIB) from the compound of formula (VIIA) and the
 15 compound of formula (IF).

The compound of formula (IIC) is:



20 where the variables are as described for the compound of formula (IIA).

The compound of formula (IF) is:



25 where the variables are as described for the compound of formula (IIC), the compound of formula (IIIC), and R₅OH.


In certain embodiments, the producing the compound of formula (IF) includes reacting the compound of formula (IIC), the compound of formula (IIIC), R_5OH , and a Lewis acid (e.g., an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)).

- 5 In particular embodiments, step (F) includes:

(G) producing a compound of formula (IH) from the compound of formula (IF); and

(H) producing the compound of formula (VIID) from the compound of formula (IH) and the compound of formula (VIIA).


- 10 The compound of formula (IH) is:



where the variables are as described for the compound of formula (IF).

- 15 In some embodiments, the producing step (G) includes reacting the compound of formula (IF) with an allylic reducing agent.


In certain embodiments, step (A) includes reducing R_{4C} that is



with a 1,2-reducing agent.


20

In particular embodiments, step (A) includes reducing R_{4C} that is




with a 1,4-reducing agent to produce a reduction product, and reducing the reduction product with a 1,2-reducing agent.

- 25 In further embodiments, step (A) includes reducing R_{4C} that is




with an allylic reducing agent to produce an acid product, and reducing the acid product with a 1,2-reducing agent capable of converting the carboxylic acid to an aldehyde.


In yet further embodiments, step (A) includes cleaving R_{4C} that is



with a glycol cleaving agent to produce an aldehyde, reacting the aldehyde $(R_CO)_2P(O)-CH_2-COOR_C$ to produce an enoate, reacting the enoate with a 1,4-reducing agent to produce R_{4C} that is



, and reacting R_{4C} that is



with a 1,2-reducing agent capable of converting the carboxylic acid to an aldehyde.


- 10 In still further embodiments, step (A) includes an esterification reaction between R_1 that is $-OP_6$, where P_6 is H, and R_9 that is $-COOH$. In other embodiments, step (A) includes a reaction between R_1 that is $-CH(Y)_2$ or $-CH_2(Y)$, and R_9 that is $-CHO$. In yet other embodiments, step (B) includes reacting the compound of formulas (VIID) with a Cr(II) salt and a Ni(II) salt. In still other embodiments, step (C) includes reacting the compound of formula (VIIE) with a hydroxyl protecting group removing agent.

15

In some embodiments, the method includes:

- (A) producing a compound of formula (VIII B) from a compound of formula (VIII A), a compound of formula (IIA), a compound of formula (IIB), and a compound of formula (IIIC), and R_5OH , where R_5 is optionally substituted acyl;
- 20 (B) producing a compound of formula (VIII C) from the compound of formula (VIII B); and
- (C) producing the halichondrin macrolide or analog thereof from the compound of formula (VIII C).

The compound of formula (IIA) is:



25

where

R_1 is $-OP_6$, $-CH(Y)_2$, or $-CH_2(Y)$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-COOR_C$ or $-SO_2R_D$;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

5 R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$,

where

m is 1, and R_F is a hydroxyl protecting group,

or

m is 0

10 (i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene; and

the remaining variables are as described for the halichondrin macrolide or analog thereof.

The compound of formula (IIB) is:




where R_S is silyl.

The compound of formula (IIIC) is:

20 $R_{4C}-R_7$,
(IIIC)

where

R_{4C} is




where

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each P_7 , when present, is independently H or a hydroxyl protecting group;;

30 R_7 is $-\text{CHO}$ or $R_{7A}\text{O}-\text{CH}(\text{OR}_{7A})-\text{CH}_2-$, where each R_{7A} is independently an optionally substituted alkyl.


The compound of formula (VIIIA) is:



where

- (i) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;
- 5 (ii) R_{10} is a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;
- or
- (iii) R_{10} and R_{11} combine to form a bond, and R_{12} is H;
- R_9 is $-\text{CHO}$ or $-\text{COOH}$;
- and
- 10 each P_9 and R_{13} is independently a hydroxyl protecting group.


The compound of formula (VIII B) is:



15 where

X_1 is $-\text{CH}_2-$, $-\text{CH}(\text{Y})-$, $-\text{C}(\text{Y})_2-$, or $-\text{O}-$, where each Y is independently $-\text{COOR}_c$ or $-\text{SO}_2\text{R}_b$;

R_M is




where

- 20 R_5 is optionally substituted acyl; and
- X_3 is oxo or X_3 , together with the carbon atom to which it is attached, is $-(\text{CH}(\text{OP}_{11}))-$, where P_{11} is H or a hydroxyl protecting group;
- X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P_{12} is H or a hydroxyl protecting group; and

the remaining groups are as described for the compound of formula (IIA) and the compound of formula (VIII A).


The compound of formula (VIIC) is:



(VIIC)

where

L₁ is



where L₂ and L₃ combine to form a bond;

A₁ is H or -OPⁿ, where Pⁿ is H or a hydroxyl protecting group; and

the remaining variables are as described for the compound of formula (VIII B).

In certain embodiments, step (A) includes:

(D) producing a compound of formula (IIC) from the compound of formula (IIA) and the compound of formula (IIB);


(E) producing a compound of formula (IF) from the compound of formula (IIC), the compound of formula (IIIC), and R₅OH;

(F) producing a compound of formula (IJ) from the compound of formula (IF);

and

(G) producing the compound of formula (VIII B) from the compound of formula (VIII A) and the compound of formula (IJ).


The compound of formula (IIC) is:



(IIC)

where the variables are as described for the compound of formula (IIA).


The compound of formula (IF) is:



5

where the variables are as described for the compound of formula (IIA), the compound of formula (IIIC), and R₅OH.

The compound of formula (IJ) is:




10

where the variables are as described for the compound of formula (VIIB).


In particular embodiments, step (D) includes reacting the compound of formula (IIB), the compound of formula (IIA), and a Lewis acid (e.g., the Lewis acid is an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)). In further embodiments, step (E) includes reacting the compound of formula (IIC), the compound of formula (IIIC), R₅OH and a Lewis acid (e.g., the Lewis acid is an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)). In yet further embodiments, the Lewis acid in step (D) and the Lewis acid in step (E) are different. In still further embodiments, the Lewis acid in step (D) and the Lewis acid in step (E) are same.

20

In some embodiments, step (F) includes converting




In particular embodiments, step (A) includes reacting R_{4C} that is



with a 1,2-reducing agent to produce a product, and reacting the product with a vinyl nucleophile.


In certain embodiments, step (A) includes reacting R_{4C} that is



with a 1,4-reducing agent to produce a product, reducing the product with a 1,2-reducing agent to produce a reduced product, and reacting the reduced product with a vinyl nucleophile.


5

In further embodiments, step (A) includes reacting R_{4C} that is



with an allylic reducing agent to produce an acid, reducing the acid with a 1,2-reducing agent to produce a reduced product, and reacting the reduced product with a vinyl nucleophile.

10 In yet further embodiments, step (A) includes cleaving R_{4C} that is



with a glycol cleaving agent to produce an aldehyde, reacting the aldehyde with $(R_CO)_2P(O)-CH_2-COOR_C$ to produce an enoate, reacting the enoate with a 1,4-reducing agent to produce a product, reducing the product with a 1,2-reducing agent to produce a reduced product, and reacting the reduced product with a vinyl nucleophile.

15

In still further embodiments, step (A) includes an esterification reaction between R_1 that is $-OP_6$, where P_6 is H, and R_9 that is $-COOH$.

In other embodiments, step (A) includes a reaction between R_1 that is $-CH(Y)_2$ or $-CH_2(Y)$, and R_9 , where R_9 is $-CHO$. In yet other embodiments, step (B) includes reacting the compound of formula (VIIIB) with an olefin metathesis catalyst. In still other embodiments, step (C) includes reacting the compound of formula (VIIC) with a hydroxyl protecting group removing agent.


20

In some embodiments, the method includes:

25 (A) producing a compound of formula (IXB) from a compound of formula (IXA), a compound of formula (IIB), a compound of formula (IIA), a compound of formula (IIIE), and R_5OH , where R_5 is optionally substituted acyl; and

(B) producing the halichondrin macrolide or analog thereof from the compound of formula (IXB).

The compound of formula (IXA) is:



where

5 R_9 is $-\text{CHO}$ or $-\text{COOP}''$;

(a1) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;

(a2) R_{10} is a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond;

or

(a3) R_{10} and R_{11} combine to form a bond, and R_{12} is H;

10 A_1 and R_{14} combine to form oxo, R_{13} is H or a hydroxyl protecting group, and R_{15} is H;

or

A_1 is H or $-\text{OP}''$, and:

(i) R_{13} is H or a hydroxyl protecting group, and R_{14} and R_{15} combine to form a double bond;

15 or

(ii) R_{13} and R_{14} combine to form a bond, and R_{15} is H or $-\text{OP}''$;


R_O is $-\text{CHO}$, $-\text{CH}_2\text{OP}''$, $-\text{CH}=\text{CH}_2$, $-\text{CH}(\text{OP}'')\text{CH}_2\text{OP}''$, $-\text{C}(\text{O})-\text{CH}_2\text{P}(\text{O})(\text{OR}_C)_2$, or halogen;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and

20 each P'' is independently H or a hydroxyl protecting group; and

each P_9 is independently a hydroxyl protecting group.

The compound of formula (IIA) is:



25

where

R_1 is $-\text{OP}_6$, $-\text{CH}(\text{Y})_2$, or $-\text{CH}_2(\text{Y})$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-\text{COOR}_D$ or $-\text{SO}_2\text{R}_D$;

30 each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$,

where

m is 1, and R_F is a hydroxyl protecting group,

or

m is 0, and

(i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene; and


the remaining variables are as described for the halichondrin macrolide or analog thereof.

5

The compound of formula (IIIE) is:

$R_{4E}-R_7$,
(IIIE)

where



each P_7 is independently H or a hydroxyl protecting group; and


R_8 is $-\text{CH}_2\text{CH}_2-\text{COOR}_C$, $-\text{CH}=\text{CH}-\text{COOR}_C$, $-\text{CH}_2\text{CH}_2-\text{SO}_2\text{R}_D$, or $-\text{CH}=\text{CH}-\text{SO}_2\text{R}_D$;

15

and


R_7 is $-\text{CHO}$ or , where each R_{7A} is independently an optionally substituted alkyl.

The compound of formula (IIB) is:




where R_5 is silyl.

The compound of formula (IXB) is:



(IXB)

where

X₁ is $-\text{CH}_2-$, $-\text{CH}(\text{Y})-$, $-\text{C}(\text{Y})_2-$, or $-\text{O}-$;R₅ is optionally substituted acyl;A₁ and R₁₄ combine to form oxo, R₁₃ is H or a hydroxyl protecting group, and R₁₅ is H;

or

A₁ is H or $-\text{OP}''$, and:(i) R₁₃ is H or a hydroxyl protecting group, and R₁₄ and R₁₅, together with the atoms to which each is attached, combine to form a double bond;

or

(ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or $-\text{OP}''$;

each P₉ is independently H or a hydroxyl protecting group, and X₃ is oxo, or X₃, together with the atom to which it is attached, combines to form $-(\text{CH}(\text{OP}_{11}))-$, where P₁₁ is H or a hydroxyl protecting group;


or

both P₉ and X₃, together with the atoms to which each is attached, combine to form a ketal;


R_{19A} is H, $-\text{OP}''$, or Y, and R_{19B} is H; or R_{19A} and R_{19B}, together with the atoms to which each is attached, combine to form a double bond; and

the remaining variables are as described for the compound of formula (IXA) and the compound of formula (IIA).


In some embodiments, step (A) includes reacting R_E that is $-\text{CHO}$ with the compound of formula (IIB) and a first Lewis acid to produce a group of the structure



In certain embodiments, step (A) includes reacting the group of the structure




with R₇, R₅OH, and a second Lewis acid to produce a group of the structure




In particular embodiments, the first Lewis acid and/or second Lewis acid is an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof). In further embodiments, the first Lewis acid is same as the second Lewis acid. In still further embodiments, the first Lewis acid is same as the second Lewis acid.

- 5 In other embodiments, step (A) includes converting R_{4E} that is




to a group of the structure



, and reacting this group with R_O that is $-C(O)-CH_2P(O)(OR_C)_2$ under Horner-Wadsworth-Emmons reaction conditions.


- 10 In yet other embodiments, step (A) includes reacting R_{4E} that is



with R_O that is $-CHO$ under Claisen reaction conditions, where R_8 is $-CH_2CH_2-COOR_C$ or $-CH_2CH_2-SO_2R_D$.

In still other embodiments, step (A) includes converting R_{4E} that is

- 15 , or
group of the structure




, and reacting this group with R_O that is a halogen under Nozaki-Hiyama-Kishi reaction conditions, where R_8 , when present, is $-CH_2CH_2-COOR_C$ or $-CH=CH-COOR_C$, and R_{14} and R_{15} combine to form a double bond.


20

In some embodiments, step (A) includes reacting R_1 that is $-OP_6$ and R_9 under esterification reaction conditions to produce a group of the structure $-X_1-C(O)-$, where P_6 is H, R_9 is $-COOH$, and X_1 is $-O-$. In certain embodiments, step (A) includes reacting R_1 that is $-CH(Y)_2$ or $-CH_2(Y)$ with R_9 under Claisen reaction conditions to produce a group of the structure $-X_1-C(O)-$, where R_9 is $-CHO$, and X_1 is $-C(Y)_2-$ or $-CH(Y)-$, respectively. In particular embodiments, step (B) includes reacting the compound of formula (IXB) with a hydroxyl protecting group removing agent. In further embodiments, D' is OP_1 , where P_1 is alkyl. In yet further embodiments, D is H. In still further embodiments, A is of the following structure:

25



In other embodiments, k is 0. In yet other embodiments, R₃ is $-(CH_2)_nNP_3P_4$ or $-(CH_2)_nOP_5$, where n is 0. In still other embodiments, D' is H. In some embodiments, A and D combine to form the following structure:




where, the bond to oxygen atom originates at the carbon atom, to which D is attached in formula (IA), and

where R₃ is $-(CH_2)_nNP_3P_4$ or $-(CH_2)_nOP_5$, where n is 2.

In certain embodiments, k is 1, and E is optionally substituted alkyl. In particular embodiments, G is O.

In still another aspect, the invention provides a method of preparing an intermediate in the synthesis of a halichondrin macrolide or analog thereof.


In some embodiments, the intermediate is:




where R₇ is $-CHO$ or $R_{7A}O$ , and each R_{7A} is independently alkyl or a hydroxyl protecting group; or both R_{7A} combine to form an optionally substituted alkylene.

In certain embodiments, the method includes:

(A) producing a compound of formula (Z2) from a compound of formula (Z1) and

, where each R_O is independently optionally substituted alkyl, where the compound of formula (Z1) is:



where


each R_{20A} is independently a hydroxyl protecting group, or both R_{20A}, together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo;

each R_{20B} is independently a hydroxyl protecting group, or both R_{20B} , together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo;

P_{12A} is H;


5 P_{12B} is H or a hydroxyl protecting group;

and the compound of formula (Z2) is:




(B) producing a compound of formula (Z3) from the compound of formula (Z2) and

10 $R_{21}O-P(=O)(OR_{21})_2$, where each R_{21} is independently optionally substituted alkyl, where the compound of formula (Z3) is:




15 (C) producing a compound of formula (Z4) from the compound of formula (Z3), where the compound of formula (Z4) is:



where X_5 is a halogen;

20 (D) producing a compound of formula (Z5) from the compound of formula (Z4), where the compound of formula (Z5) is:




where

R_{20A} is H or a hydroxyl protecting group; and


each R_{20B} is independently H or a hydroxyl protecting group, or both R_{20B} , together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo;

(E) producing a compound of formula (Z6) from the compound of formula (Z5) and $R_{7A}OH$



(F) producing a compound of formula (Z7) from the compound of formula (Z6) and

$R_{22}O-P(=O)(OR_{22})-CH_2-COOR_{22}$, where each R_{22} is independently optionally substituted alkyl, and the compound of formula (Z7) is:



and


(G) producing the compound of formula (Z) from the compound of formula (Z7).

15 In particular embodiments, step (E) includes reacting the compound of formula (Z5) with $R_{7A}OH$ and a Brønsted acid. In further embodiments, step (F) includes reacting the compound of formula (Z6) with a glycol cleaving agent to produce an aldehyde product, and reacting the aldehyde product with

$R_{22}O-P(=O)(OR_{22})-CH_2-COOR_{22}$ under Horner-Wadsworth-Emmons reaction conditions. In yet further embodiments, step (G) includes reacting the compound of formula (Z6) with a 1,4-reducing agent.


20

In still further embodiments, the intermediate is:



where

R_7 is $-\text{CHO}$ or



each R_{7A} is independently alkyl or a hydroxyl protecting group; or

both R_{7A} combine to form an optionally substituted alkylene;

5 R_8 is $-\text{CH}_2\text{CH}_2-\text{COOR}_C$, $-\text{CH}=\text{CH}-\text{COOR}_C$, $-\text{CH}_2\text{CH}_2-\text{SO}_2\text{R}_D$, or $-\text{CH}=\text{CH}-\text{SO}_2\text{R}_D$;


R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and

R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl.


10

In other embodiments, the method includes:

(A) producing a compound of formula (Z2) from a compound of formula (Z1) and



formula (Z1) is:



15

where

each R_{20A} is independently a hydroxyl protecting group, or both R_{20A} , together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo;

20


each R_{20B} is independently a hydroxyl protecting group, or both R_{20B} , together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo;

P_{12A} is H;


25


P_{12B} is H or a hydroxyl protecting group;

and the compound of formula (Z2) is:




(B) producing a compound of formula (Z3) from the compound of formula (Z2) and

, where each R₂₁ is independently optionally substituted alkyl, where the compound of formula (Z3) is:



5


(C) producing a compound of formula (Z4) from the compound of formula (Z3), where the compound of formula (Z4) is:



10

where X₅ is a halogen;

(D) producing a compound of formula (Z5) from the compound of formula (Z4), where the compound of formula (Z5) is:



15


where

R_{20A} is H or a hydroxyl protecting group; and

each R_{20B} is independently H or a hydroxyl protecting group, or both R_{20B}, together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo;

20

(E) producing a compound of formula (Z5A) from the compound of formula (Z5) and R_{7A}OH, where the compound of formula (Z5A) is:




where

R_{20A} is H or a hydroxyl protecting group; and

each R_{20B} is independently H or a hydroxyl protecting group, or both R_{20B} , together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo;


- 5 (F) producing a compound of formula (Z5B) from the compound of formula (Z5A), where the compound of formula (Z5B) is:



where R_{23} is H or $-OR_{20A}$;

- 10 and

(G) producing the compound of formula (Z8) from the compound of formula (Z5B) and



- 15 In further embodiments, step (E) includes reacting the compound of formula (Z5) with $R_{7A}OH$ and a Brønsted acid. In yet further embodiments, step (F) includes reacting the compound of formula (Z5A), in which both R_{20B} are H, with a glycol cleaving agent. In still further embodiments, step (E) or step (F) further includes reacting $-OR_{20A}$, where R_{20A} is optionally substituted acyl with an allylic reducing agent. In other embodiments, step (G) includes reacting the compound of formula (Z5B) with

- 20 under Horner-Wadsworth-Emmons reaction conditions to produce the compound of formula (Z8), in which R_8 is $-\text{CH}=\text{CH}-\text{COOR}_C$ or $-\text{CH}=\text{CH}-\text{SO}_2R_D$, respectively. In yet other embodiments, step (G) further includes reacting the compound of formula (Z8), in which R_8 is $-\text{CH}=\text{CH}-\text{COOR}_C$ or $-\text{CH}=\text{CH}-\text{SO}_2R_D$, with a 1,4-reducing agent to produce the compound of formula (Z8), in which R_8 is $-\text{CH}_2\text{CH}_2-\text{COOR}_C$ or $-\text{CH}_2\text{CH}_2-\text{SO}_2R_D$, respectively. In still other embodiments, the producing the compound of formula (Z2) includes reacting the compound of formula (Z1) with


- 25 under Horner-Wadsworth-Emmons reaction conditions. In some embodiments, step (B) includes reacting the compound of formula (Z2) with a 1,2-reducing agent to produce a reduction product, and reacting the reduction product with

- 30 under Horner-Wadsworth-Emmons reaction conditions. In certain embodiments, step (C) includes substituting free hydroxyl group in the compound of formula (Z3) with a halogen. In particular embodiments, step (D) includes subjecting the compound of formula (Z4) to reductive metal conditions.

In a further aspect, the invention provides the compounds of formula (II), (IID), (IIE), (IIF), (IIG), (IIH), (IIi), (IVAb), (VAb), (VIIAa), (VIAa), (VIID), (IXD), (IXF), (IXG), (Z), (Z1), (Z2), (Z3), (Z4), (Z5), (Z5A), (Z5B), or (Z6).


5

The compound of formula (II) is:



where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, or A is a group of formula (1):



15

where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₂ is H and P₁ is absent, H, alkyl, or a hydroxyl protecting group, or R₂ and P₁ combine to form a bond;

20

(i) R₃ is H, and P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;



(ii) R₃ is $-(CH_2)_nNP_3P_4$, where P₃ is H or an N-protecting group, and (a) P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₄ is H or an N-protecting group, (b) P₂ and P₄ combine to form an alkylidene, or (c) each of P₂ and P₄ is H;

25

(iii) R₃ is $-(CH_2)_nOP_5$, where P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₅ is H, optionally substituted alkyl, or a hydroxyl protecting group; or P₂ and P₅, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R₃ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:

30



group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,

NR_B(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or

10 heterocyclic radical-alkyl;

n, when present, is 0, 1, or 2;

k is 0 or 1;

R₁ is -OP₆, -CH(Y)₂, or -CH₂(Y), where P₆ is H or a hydroxyl protecting group;


R₆ is H, optionally substituted alkyl, or optionally substituted aryl; and

15 Y is independently -COOR_C or -SO₂R_D;

R_C, when present, is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and

R_D, when present, is optionally substituted aryl or optionally substituted non-enolizable alkyl.

20 The compound of formula (VIIAa) is:



(VIIAa)

where

R_7 is $-\text{CH}_2\text{OP}_A$, $-\text{CHO}$, or

5 $\text{R}_{7A}\text{O}-\text{C}(\text{OR}_{7A})_2$, where each R_{7A} is independently an optionally substituted alkyl;

R_9 is $-\text{CH}_2\text{OP}''$, $-\text{CHO}$, or $-\text{COOP}''$;

(i) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;

(ii) R_{10} is a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond;

or

10 (iii) R_{10} and R_{11} combine to form a bond, and R_{12} is H;

A_1 and R_{14} combine to form oxo, R_{13} is H or a hydroxyl protecting group, and R_{15} is H;

or

A_1 is H or $-\text{OP}''$, and:

(i) R_{13} is H or a hydroxyl protecting group, and R_{14} and R_{15} combine to form a double bond;

15

or


(ii) R_{13} and R_{14} combine to form a bond, and R_{15} is H or $-\text{OP}''$;

each P'' is independently H or a hydroxyl protecting group; and

each P_9 is independently a hydroxyl protecting group, and X_3 is oxo or X_3 , together with the carbon

20 atom to which it is attached, is $-(\text{CH}(\text{OP}_{11}))-$, where P_{11} is H or a hydroxyl protecting group; or both P_9 groups and X_3 , together with the atoms to which each is attached, combine to form a ketal.

The compound of formula (VIAa) is:



(VIAa)

25

where

R_7 is $-\text{CHO}$ or



R_{10A} is $-\text{CH}_2-\text{CH}=\text{CH}_2$, $-(\text{CH}_2)_2-\text{CH}=\text{CH}_2$, or $-(\text{CH}_2)_3-\text{OP}_{10}$;

each of R_{10} and P_{10} is independently a hydroxyl protecting group;

5 A_1 and R_{14} combine to form oxo, R_{13} is H or a hydroxyl protecting group, and R_{15} is H;

or

A_1 is H or $-\text{OP}''$, and:

(i) R_{13} is H or a hydroxyl protecting group, and R_{14} and R_{15} combine to form a double bond;

10 or


(ii) R_{13} and R_{14} combine to form a bond, and R_{15} is H or $-\text{OP}''$;

and

each P_9 is independently a hydroxyl protecting group, and X_3 is oxo or X_3 , together with the carbon atom to which it is attached, is $-(\text{CH}(\text{OP}_{11}))-$, where P_{11} is H or a hydroxyl protecting group; or both P_9

15 groups and X_3 , together with the atoms to which each is attached, combine to form ketal.

The A compound of formula (IID) or (IIF) is:



20 where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, a hydroxyl protecting group, and A is a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , or A is a group of formula (1):



where

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;


30 R_2 is H and P_1 is absent, H, alkyl, or a hydroxyl protecting group, or R_2 and P_1 combine to form a bond;

(i) R_3 is H, and P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

(ii) R_3 is $-(CH_2)_nNP_3P_4$, where P_3 is H or an N-protecting group, and (a) P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_4 is H or an N-protecting group, (b) P_2 and P_4 combine to form an alkylidene, or (c) each of P_2 and P_4 is H;

(iii) R_3 is $-(CH_2)_nOP_5$, where P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_5 is H, optionally substituted alkyl, or a hydroxyl protecting group; or P_2 and P_5 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_B(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

n , when present, is 0, 1, or 2;

k is 0 or 1;

X_1 is $-CH_2-$ or $-O-$;

X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(CH(OP_{12}))_n-$, where P_{12} is H or a hydroxyl protecting group; and

R_E is $-CHO$ or $-CH_{(1+m)}(OR_F)_{(2-m)}$, where


m is 1, and R_F is a hydroxyl protecting group,

or

m is 0, and


- (i) each R_F is independently an alkyl or hydroxyl protecting group, or
- (ii) both R_F combine to form an alkylene.

The compound of formula (IIE) or (IIG):



where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, a hydroxyl protecting group, and A is a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , or A is a group of formula (1):



where

15 L is $-(CH(OP_2))_n-$, $-(C(OH)(OP_2))_n-$, or $-C(O)-$;



R_2 is H and P_1 is absent, H, alkyl, or a hydroxyl protecting group, or R_2 and P_1 combine to form a bond;

(i) R_3 is H, and P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

20 (ii) R_3 is $-(CH_2)_nNP_3P_4$, where P_3 is H or an N-protecting group, and (a) P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_4 is H or an N-protecting group, (b) P_2 and P_4 combine to form an alkylidene, or (c) each of P_2 and P_4 is H;

(iii) R_3 is $-(CH_2)_nOP_5$, where P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_5 is H, optionally substituted alkyl, or a hydroxyl protecting group; or P_2 and P_5 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

25 (iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is H, optionally substituted alkyl, or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;
 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,
 NR_B(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of
 R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl,
 alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or
 10 heterocyclic radical-alkyl;


n, when present, is 0, 1, or 2;

k is 0 or 1;

X₁ is -CH₂- or -O-; and

15 X₄ is oxo, or X₄, together with the atom to which it is attached, is -(CH(OP₁₂))- , where P₁₂ is H or a hydroxyl protecting group.


The compound of formula (IIH) is:



20 where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated

hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , or A is a group of formula (1):



5 where

L is $-(CH(OP_2))_n-$, $-(C(OH)(OP_2))_n-$, or $-C(O)-$;


R_2 is H and P_1 is absent, H, alkyl, or a hydroxyl protecting group, or R_2 and P_1 combine to form a bond;

(i) R_3 is H, and P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;


(ii) R_3 is $-(CH_2)_nNP_3P_4$, where P_3 is H or an N-protecting group, and (a) P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_4 is H or an N-protecting group, (b) P_2 and P_4 combine to form an alkylidene, or (c) each of P_2 and P_4 is H;

(iii) R_3 is $-(CH_2)_nOP_5$, where P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_5 is H, optionally substituted alkyl, or a hydroxyl protecting group; or P_2 and P_5 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



20



, where each P' is independently H or a hydroxyl protecting group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_B(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of

25

R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

n , when present, is 0, 1, or 2;

5 k is 0 or 1;

each R_C is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

X_1 is $-\text{CH}_2-$ or $-\text{O}-$;

10 X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P_{12} is H or a hydroxyl protecting group; and

R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$, where

m is 1, and R_F is a hydroxyl protecting group,


or

m is 0, and

15 (i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene.


The compound of formula (Iii) is:



20

where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, a hydroxyl protecting group, and A is a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently
25 selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , or A is a group of formula (1):



where

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;


30 R_2 is H and P_1 is absent, H, alkyl, or a hydroxyl protecting group, or R_2 and P_1 combine to form a bond;

(i) R_3 is H, and P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

(ii) R_3 is $-(CH_2)_nNP_3P_4$, where P_3 is H or an N-protecting group, and (a) P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_4 is H or an N-protecting group, (b) P_2 and P_4 combine to form an alkylidene, or (c) each of P_2 and P_4 is H;

(iii) R_3 is $-(CH_2)_nOP_5$, where P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_5 is H, optionally substituted alkyl, or a hydroxyl protecting group; or P_2 and P_5 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_B(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;


n , when present, is 0, 1, or 2;

k is 0 or 1;

each R_C is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and


X_1 is $-CH_2-$ or $-O-$.

The compound of formula (VIID) is:



where

- 5 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, or A is a group of formula (1):





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₂ is H and P₁ is absent, H, alkyl, or a hydroxyl protecting group, or R₂ and P₁ combine to form a bond;

- 15 (i) R₃ is H, and P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;
- (ii) R₃ is $-(CH_2)_nNP_3P_4$, where P₃ is H or an N-protecting group, and (a) P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₄ is H or an N-protecting group, (b) P₂ and P₄ combine to form an alkylidene, or (c) each of P₂ and P₄ is H;
- 20 (iii) R₃ is $-(CH_2)_nOP_5$, where P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₅ is H, optionally substituted alkyl, or a hydroxyl protecting group; or P₂ and P₅, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or
- (iv) R₃ and P₂ combine to form an optionally substituted ethylene or a structure
- 25 selected from the group consisting of:





, where each P' is independently H or a hydroxyl protecting group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_B(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 n, when present, is 0, 1, or 2;

k is 0 or 1;

X₁ is -CH₂-, -CH(Y)-, -C(Y)₂-, or -O-, where each Y is independently -COOR_C or -SO₂R_D;

X₄ is oxo, or X₄, together with the atom to which it is attached, is -(CH(OP₁₂))- , where P₁₂ is H or a hydroxyl protecting group;


15 each P₉ and R₁₃ is independently a hydroxyl protecting group;

(i) R₁₀ is a hydroxyl protecting group, R₁₁ is alkyl ether, and R₁₂ is H;

(ii) R₁₀ is a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond; or

(iii) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;

R_{M1} is




where

R₅ is optionally substituted acyl;


P₁₃ is H or a hydroxyl protecting group; and

R_{4C} is




where R_C is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and P_7 , when present, is independently H or a hydroxyl protecting group.

The compound of formula (IVAb) is:



where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, a hydroxyl protecting group, and A is a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , or A is a group of formula (1):



where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;



R_2 is H and P_1 is absent, H, alkyl, or a hydroxyl protecting group, or R_2 and P_1 combine to form a bond;

(i) R_3 is H, and P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

(ii) R_3 is $-(CH_2)_nNP_3P_4$, where P_3 is H or an N-protecting group, and (a) P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_4 is H or an N-protecting group, (b) P_2 and P_4 combine to form an alkylidene, or (c) each of P_2 and P_4 is H;

(iii) R_3 is $-(CH_2)_nOP_5$, where P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_5 is H, optionally substituted alkyl, or a hydroxyl protecting group; or P_2 and P_5 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

- 5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;
 each Q_i is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,
 NR_B(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of
 R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl,
 alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or
 10 heterocyclic radical-alkyl;

n, when present, is 0, 1, or 2;

k is 0 or 1;

X₁ is -CH₂-, -CH(Y)-, -C(Y)₂-, or -O-, where each Y is independently -COOR_C or -SO₂R_D;

- 15 each R_C, when present, is independently optionally substituted alkyl, optionally substituted aryl, or
 optionally substituted arylalkyl;

each R_D, when present, is independently optionally substituted aryl or optionally substituted non-
 enolizable alkyl;

X₄ is oxo, or X₄, together with the atom to which it is attached, is -(CH(OP₁₂))- , where P₁₂ is H or a
 hydroxyl protecting group;

- 20 each P₉ is independently a hydroxyl protecting group;

(a1) R₁₀ is H or a hydroxyl protecting group, R₁₁ is alkyl ether, and R₁₂ is H;

(a2) R₁₀ is H or a hydroxyl protecting group, and R₁₁ and R₁₂ combine to form a double bond;

or

(a3) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;

- 25 (b1) A₁ and R₁₄ combine to form oxo, R₁₃ is H or a hydroxyl protecting group, and R₁₅ is H;

or

(b2) A₁ is H or -OP'', and:

(i) R_{13} is H or a hydroxyl protecting group, and R_{14} and R_{15} combine to form a double bond;

or

(ii) R_{13} and R_{14} combine to form a bond, and R_{15} is H or $-OP''$; and

5 (c1) R_{16} is H, and P_8 is H or a hydroxyl protecting group;

or


(c2) R_{16} and P_8 combine to form a double bond;

and

each P'' , when present, is independently H or a hydroxyl protecting group.


10

The compound of formula (VAb) is:



where

15 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, or A is a group of formula (1):



20

where

L is $-(CH(OP_2))_n-$, $-(C(OH)(OP_2))_n-$, or $-C(O)-$;

R_2 is H and P₁ is absent, H, alkyl, or a hydroxyl protecting group, or R_2 and P₁ combine to form a bond;

25


(i) R_3 is H, and P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

(ii) R_3 is $-(CH_2)_nNP_3P_4$, where P₃ is H or an N-protecting group, and (a) P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₄ is H or an N-protecting group, (b) P₂ and P₄ combine to form an alkylidene, or (c) each of P₂ and P₄ is H;

30

(iii) R_3 is $-(CH_2)_nOP_5$, where P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₅ is H, optionally substituted alkyl, or a hydroxyl protecting group; or P₂ and P₅, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,

$NR_B(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

n , when present, is 0, 1, or 2;

k is 0 or 1;

a designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or a designates (*S*)-stereogenic center, and Z is OR_{16} , where R_{16} is a hydroxyl protecting group;

X_1 is $-CH_2-$, $-CH(Y)-$, $-C(Y)_2-$, or $-O-$, where each Y is independently $-COOR_C$ or $-SO_2R_D$;

each P_9 is independently a hydroxyl protecting group, and X_3 is oxo; or both P_9 groups and X_3 , together with the atoms to which each is attached, combine to form ketal;

Y_1 is iodide, bromide, or trifluoromethanesulfonate;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and

each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

(a1) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;


(a2) R_{10} is a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond;

or

(a3) R_{10} and R_{11} combine to form a bond, and R_{12} is H;


- (b1) A₁ and R₁₄ combine to form oxo, R₁₃ is a hydroxyl protecting group, and R₁₅ is H;
or
(b2) A₁ is H or –OPⁿ, and:
(i) R₁₃ is a hydroxyl protecting group, and R₁₄ and R₁₅ combine to form a double bond;
or
(ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or –OPⁿ;
and
each Pⁿ is independently a hydroxyl protecting group.

10 The compound of formula (IXD) is:



where

- each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D
15 and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, or A is a group of formula (1):




20 where

L is –(CH(OP₂))–, –(C(OH)(OP₂))–, or –C(O)–;

R₂ is H and P₁ is absent, H, alkyl, or a hydroxyl protecting group, or R₂ and P₁ combine to form a bond;

- (i) R₃ is H, and P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting
25 group;
(ii) R₃ is –(CH₂)_nNP₃P₄, where P₃ is H or an N-protecting group, and (a) P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₄ is H or an N-protecting group, (b) P₂ and P₄ combine to form an alkylidene, or (c) each of P₂ and P₄ is H;
(iii) R₃ is –(CH₂)_nOP₅, where P₂ is absent, H, optionally substituted alkyl, or a
30 hydroxyl protecting group, and P₅ is H, optionally substituted alkyl, or a hydroxyl protecting group; or P₂ and P₅, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R₃ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



PO₃H₂, where each P' is independently H or a hydroxyl protecting group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,

NR_B(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

n, when present, is 0, 1, or 2;

k is 0 or 1;


X₁ is -CH₂-, -CH(Y)-, -C(Y)₂-, or -O-, each Y is independently -COOR_C or -SO₂R_D;

each R_C, when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D, when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

X₄ is oxo, or X₄, together with the atom to which it is attached, is $-(CH(OP_{12}))_n-$, where P₁₂ is H or a hydroxyl protecting group;

R₇ is -CHO, -CH₂OP_A or



, where each R_{7A} is independently an optionally substituted alkyl;

(i) R₁₀ is a hydroxyl protecting group, R₁₁ is alkyl ether, and R₁₂ is H;

(ii) R₁₀ is a hydroxyl protecting group, and R₁₁ and R₁₂ combine to form a double bond; or

(iii) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;

A₁ and R₁₄ combine to form oxo, R₁₃ is H or a hydroxyl protecting group, and R₁₅ is H;

or

A_1 is H or -OP", and

(i) R₁₃ is H or a hydroxyl protecting group, and R₁₄ and R₁₅, together with the atoms to which each is attached, combine to form a double bond;


or

(ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or -OPⁿ; each P₉ is independently H or a hydroxyl protecting group,

each P₉ is independently a hydroxyl protecting group, and X₃ is oxo, or X₃, together with the atom to which it is attached, combines to form $-(CH(OP_{11}))_n-$, where P₁₁ is H or a hydroxyl protecting group; or both P₉ and X₃, together with the atoms to which each is attached, combine to form a ketal; and


R_{19A} is H, -OP", or Y, and R_{19B} is H; or R_{19A} and R_{19B}, together with the atoms to which each is attached, combine to form a double bond.

15 The compound of formula (IXF) is:



where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, or A is a group of formula (1):



25 where

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;


R₂ is H and P₁ is absent, H, alkyl, or a hydroxyl protecting group, or R₂ and P₁ combine to form a bond:

(i) R₃ is H, and P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

(ii) R₃ is -(CH₂)_nNP₃P₄, where P₃ is H or an N-protecting group, and (a) P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₄ is H or an N-protecting group, (b) P₂ and P₄ combine to form an alkylidene, or (c) each of P₂ and P₄ is H;

(iii) R_3 is $-(CH_2)_nOP_5$, where P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_5 is H, optionally substituted alkyl, or a hydroxyl protecting group; or P_2 and P_5 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

5 (iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



10 E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_B(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or

15 heterocyclic radical-alkyl;

n, when present, is 0, 1, or 2;

k is 0 or 1;

X_1 is $-CH_2-$, $-CH(Y)-$, $-C(Y)_2-$, or $-O-$, where each Y is independently $-COOR_C$ or $-SO_2R_D$;

20 X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(CH(OP_{12}))-$, where P_{12} is H or a hydroxyl protecting group;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

25 each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

R_O is $-CHO$, $-CH_2OP''$, $-CH=CH_2$, $-CH(OP'')CH_2OP''$, $-C(O)-CH_2P(O)(OR_C)_2$, or halogen;

(i) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;

(ii) R_{10} is a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond; or

(iii) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;

A₁ and R₁₄ combine to form oxo, R₁₃ is H or a hydroxyl protecting group, and R₁₅ is H;

or

A₁ is H or -OP'', and

5 (i) R₁₃ is H or a hydroxyl protecting group, and R₁₄ and R₁₅, together with the atoms to which each is attached, combine to form a double bond;


or

(ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or -OP''; each P₉ is independently H or a hydroxyl protecting group;

10 and

each of P'', P₉, and P₁₃ is independently H or a hydroxyl protecting group.


The compound of formula (IXG) is:



(IXG)

where

R₇ is -CHO, -CH₂OP_A, or



, where each R_{7A} is independently an optionally substituted alkyl;

(i) R₁₀ is a hydroxyl protecting group, R₁₁ is alkyl ether, and R₁₂ is H;

20 (ii) R₁₀ is a hydroxyl protecting group, and R₁₁ and R₁₂ combine to form a double bond; or

(iii) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;

A₁ and R₁₄ combine to form oxo, R₁₃ is H or a hydroxyl protecting group, and R₁₅ is H;

or

A₁ is H or -OP'', and

25 (i) R₁₃ is H or a hydroxyl protecting group, and R₁₄ and R₁₅, together with the atoms to which each is attached, combine to form a double bond;

or

(ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or -OP''; each P₉ is independently H or a hydroxyl protecting group,

30 each P₉ is independently a hydroxyl protecting group, and X₃ is oxo, or X₃, together with the atom to which it is attached, combines to form -(CH(OP₁₁))-; where P₁₁ is H or a hydroxyl protecting group; or both P₉ and X₃, together with the atoms to which each is attached, combine to form a ketal; and

R_{19A} is H, -OPⁿ, or Y, and R_{19B} is H; or R_{19A} and R_{19B}, together with the atoms to which each is attached, combine to form a double bond.

In a yet further aspect, the invention provides a compound of formula (Z), (Z1), (Z2), (Z3), (Z4), (Z5), (Z5A), (Z5B), or (Z6), as described herein.

In the halichondrin macrolide and halichondrin macrolide analog structures herein, when P₁ is absent, each of D and D' is independently H or optionally substituted alkyl, and, when P₂ is absent, L is -C(O)-.

10

Definitions

Compounds useful in the invention may be isotopically labeled compounds. Useful isotopes include hydrogen, carbon, nitrogen, and oxygen (e.g., ²H, ³H, ¹³C, ¹⁴C, ¹⁵N, ¹⁸O, and ¹⁷O). Isotopically-labeled compounds can be prepared by synthesizing a compound using a readily available isotopically-labeled reagent in place of a non-isotopically-labeled reagent.

15

For any of the following chemical definitions, a number following an atomic symbol indicates that total number of atoms of that element that are present in a particular chemical moiety. As will be understood, other atoms, such as hydrogen atoms, or substituent groups, as described herein, may be present, as necessary, to satisfy the valences of the atoms. For example, an unsubstituted C₂ alkyl group has the formula -CH₂CH₃. When used with the groups defined herein, a reference to the number of carbon atoms includes the divalent carbon in acetal and ketal groups but does not include the carbonyl carbon in acyl, ester, carbonate, or carbamate groups. A reference to the number of oxygen, nitrogen, or sulfur atoms in a heteroaryl group only includes those atoms that form a part of a heterocyclic ring.

20

By "acetal" is meant -O-(CHR)-O-, where R is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl, or R group is a bond to an enumerated carbon atom, as shown in Chart 1, within the intermediate or within the halichondrin macrolide or analog thereof.

25

By "acetyl" is meant an acyl, in which R is -CX_nH_{3-n}, where n is 0, 1, 2, or 3, and each X is independently alkoxy or halogen, provided that, when n is 3, each X is independently halogen, and, when n is 2, either both of the X groups are independently halogen or both of the X groups are independently alkoxy. An acetyl group may be substituted (i.e., n is 1, 2, or 3) or unsubstituted (i.e., n is 0).

30

By "acyl" is meant -C(O)R, where R is H, alkyl, alkenyl, aryl, or arylalkyl. In exemplary acyl groups, R is H, C₁₋₁₂ alkyl (e.g., C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, or C₃₋₆ alkyl), C₂₋₁₂ alkenyl (e.g., C₂₋₈, C₂₋₆, C₂₋₄, C₃₋₁₂, or C₃₋₆ alkenyl), C₆₋₂₀ aryl (e.g., C₆₋₁₄, C₆₋₁₀, C₆₋₂₀, or C₈₋₁₄ aryl), monocyclic C₁₋₆ heteroaryl (e.g., monocyclic C₁₋₄ or C₂₋₆ heteroaryl), C₄₋₁₉ heteroaryl (e.g., C₄₋₁₀ heteroaryl), (C₆₋₁₄)aryl(C₁₋₆)alkyl, (C₁₋₆)heteroaryl(C₁₋₆)alkyl, or (C₄₋₉)heteroaryl(C₁₋₆)alkyl. As defined herein, any heteroaryl group present in an acyl group has from 1 to 4 heteroatoms selected independently from O, N, and S. An acyl group can be unsubstituted or substituted (e.g., optionally substituted acyl). In the optionally substituted acyl group, the substituent R is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl. In some embodiments, acyl is C₂₋₁₀ acyl.

35

40

By "acylating agent" is meant a compound that reacts with an amine or a hydroxyl group to produce an amide or an ester, respectively. An acylating agent has a formula R-LG, where R is acyl, and LG is halogen, carbonate, or -OR', where R' is acyl.

5

By "alkoxide" is meant an anionic compound RO⁻, where R is alkyl. A counterion for alkoxide can be an alkali metal cation, an alkali earth metal cation, or a tetraalkylammonium cation. Alkoxide can be optionally substituted in the same manner as alkyl.

10 By "alkoxy" is meant -OR, where R is alkyl. Alkoxy can be optionally substituted in the same manner as alkyl.

By "alkoxyalkyl" is meant -OR, where R is alkyl substituted by alkoxy. Each portion of the alkoxyalkyl can be optionally substituted in the same manner as alkyl.

15

By "alkoxyaryl" is meant -R'(R")_n, where n is 1 or 2, R' is arylene and R" is alkoxy, as defined herein. R' can be further optionally substituted in the same manner as aryl. R" can be optionally substituted in the same manner as alkyl.

20 By "alkoxyaryalkyl" is meant -R'(R"(R'''))_n, where n is an integer from 1 to 3, R' is alkylene, R" is arylene, and R''' is alkoxy, as defined herein. R' can be optionally substituted in the same manner as alkyl. R" can be further optionally substituted in the same manner as aryl. R''' can be optionally substituted in the same manner as alkyl.

25 By "alkyl" is meant a straight or branched chain saturated cyclic (i.e., cycloalkyl) or acyclic hydrocarbon group of from 1 to 12 carbons, unless otherwise specified. In some embodiments, alkyl is C₁₋₆ alkyl. Exemplary alkyl groups include C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, and C₃₋₆ alkyl. Specific examples include methyl, ethyl, 1-propyl, 2-propyl, 2-methyl-1-propyl, 1-butyl, 2-butyl, and the like. Alkyl group can be optionally substituted with 1, 2, 3, or 4 substituents selected from the group consisting of halogen, hydroxy, alkoxy, 30 aryloxy, arylalkyloxy, amino, oxo, alkylthio, alkylenedithio, alkylamino, [alkenyl]alkylamino, [aryl]alkylamino, [arylalkyl]alkylamino, dialkylamino, silyl, sulfonyl, cyano, nitro, carboxyl, and azido.

By "alkylamino" is meant -NHR, where R is alkyl. By "[alkenyl]alkylamino" is meant -NRR', where R is alkyl, and R' is alkenyl. By "[aryl]alkylamino" is meant -NRR', where R is alkyl, and R' is aryl. By

35 "[arylalkyl]alkylamino" is meant -NRR', where R is alkyl, and R' is arylalkyl. By "dialkylamino" is meant -NR₂, where each R is alkyl, selected independently.

By "alkylaryl" is meant -R'(R")_n, where n is an integer from 1 to 3, R' is arylene, and R" is alkyl. Alkylaryl can be optionally substituted in the same manner as defined for each R' and R" group.

40

By "alkylene" is meant a multivalent alkyl group. Alkylene groups can be optionally substituted in the same manner as alkyl groups. Alkylene may be a divalent alkylene. For example, a C₁ alkylene group is -CH₂-.

By "alkylenedithio" is meant $-S\text{-alkylene-S-}$. Alkylenedithio can be optionally substituted in the same manner as an alkylene group.

- 5 By "alkylhaloaryl" is meant $-R'(R'')_n-R'''$, where n is an integer from 1 to 5 and R' is arylene, R'' is halogen, and R''' is alkylene, as defined herein. R' can be further optionally substituted in the same manner as aryl. R''' can be further optionally substituted in the same manner as alkyl.

- 10 By "alkylthio" is meant $-SR$, where R is alkyl. Alkylthio can be optionally substituted in the same manner as an alkyl group.

- By "alkenyl" is meant a straight or branched chain cyclic or acyclic hydrocarbon group of, unless otherwise specified, from 2 to 12 carbons and containing one or more carbon-carbon double bonds. In some embodiments, alkenyl is C_{2-6} alkenyl. Exemplary alkenyl groups include C_{2-6} , C_{2-7} , C_{2-8} , C_{2-9} , C_{3-12} , and C_{3-6} alkenyl. Specific examples include ethenyl (i.e., vinyl), 1-propenyl, 2-propenyl (i.e., allyl), 2-methyl-1-propenyl, 1-butenyl, 2-butenyl (i.e., crotyl), and the like. Alkenyl group can be optionally substituted in the same manner as alkyl groups. Alkenyl groups, used in any context herein, may also be substituted with an aryl group.
- 15

- 20 By "amido" is meant $-NHR$, where R is acyl. Amido can be optionally substituted in the same manner as acyl.

- By "aminal" is meant $-O-CR_2-NR'$, where each R is independently H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl, or both R groups are together optionally substituted alkylene, and R' is H or an N -protecting group. In particular, R' can be an N -protecting group (e.g., Boc).
- 25

- By "amino" is meant $-NR_2$, where N and R_2 combine to form azido, or each R is independently H or an N -protecting group, or both R combine to form an N -protecting group. Amino can be unmasked, when each R is H, or masked, when at least one R is not H. Thus, optionally masked amino can be masked or unmasked amino.
- 30

- By "aminoalkyl" is meant $-R'(R'')_n$, where n is 1 or 2, R' is alkylene, and R'' is amino, as defined herein. R' can be optionally substituted in the same manner as an alkyl group.
- 35

- By "aryl" is meant a monocyclic or multicyclic ring system having one or more aromatic rings, where the ring system is carbocyclic. Exemplary aryl groups include C_{6-20} , C_{6-15} , C_{6-10} , C_{8-20} , and C_{8-15} aryl. A preferred aryl group is a C_{6-10} aryl group. Specific examples of carbocyclic aryl groups include phenyl, indanyl, indenyl, naphthyl, phenanthryl, anthracyl, and fluorenyl. Aryl group can be optionally substituted with 1, 2, 3, 4, or 5 substituents selected from the group consisting of alkyl, alkenyl, aryl, arylalkyl, halogen, alkoxy, aryloxy, arylalkyloxy, alkylthio, alkylenedithio, alkylamino, [alkenyl]alkylamino, [aryl]alkylamino, [arylalkyl]alkylamino, dialkylamino, silyl, sulfonyl, cyano, nitro, carboxyl, and azido.
- 40

By "arylalkyl" is meant $-R'R''$, where R' is alkylene, and R'' is aryl. Arylalkyl can be optionally substituted in the same manner as defined for each R' and R'' group.

- 5 By "arylalkyloxy" is meant $-OR$, where R is arylalkyl. Arylalkyloxy can be optionally substituted in the same manner as defined for arylalkyl.

By "arylene" is meant a multivalent aryl group. Arylene groups can be optionally substituted in the same manner as aryl groups. For example, a C_6 arylene group is phenylene.

10

By "aryloxy" is meant $-OR$, where R is aryl. Aryloxy can be optionally substituted in the same manner as aryl.

By "azido" is meant $-N_3$.

15

By "boronate" is meant $-OB(R)O-$, where R is alkyl, alkenyl, aryl, arylalkyl, alkoxy, or 2,6-diacetamidophenyl. Boronate can be substituted, when R is a substituted alkyl, substituted alkenyl, substituted aryl, substituted arylalkyl, or substituted alkoxy. Alternatively, boronate can be unsubstituted, when R is unsubstituted alkyl, unsubstituted alkenyl, aryl, unsubstituted arylalkyl, unsubstituted alkoxy, or 2,6-diacetamidophenyl.

20

By "carbamate" is meant a group, when a hydroxyl protecting group, having the formula $-OC(O)NR_2$, or, when an amine protecting group, having the formula $-NR'-C(O)OR$, where each R and R' is independently H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl.

25

By "carbonate" is meant $-OC(O)OR$, where R is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl.

By "carbonyl" is meant $-C(O)-$.

30

By "carboxyl" is meant $-C(O)OH$, in free acid, ionized, or salt form.

By "carboxylic acid" is meant $R-OH$, where R is optionally substituted acyl.

- 35 By "carboxylic acid anhydride" is meant $R-O-R$, where each R is independently optionally substituted acyl.

By "cyclic carbonate" is meant $-OC(O)O-$ that is part of a ring.

By "dicarbonyl" is meant $-C(O)-C(O)-$. Dicarbonyl-dioxo is $-OC(O)-COO-$.

40

By "ester" is meant $-OC(O)R$, where $-C(O)R$ is an optionally substituted acyl group.

By "ether" is meant -OR , where R is alkyl, alkenyl, arylalkyl, silyl, or 2-tetrahydropyranyl. Ether can be optionally substituted as defined for each R group.

By "halichondrin macrolide" is meant a lactone including the structure of carbons 1-30 as shown in Chart 1, where carbons 29 and 30 form part of a five- or six-membered ring.

By "haloalkyl" is meant -R'(R)_n , where n is an integer from 1 to 5 and R' is alkylene and R is halogen, as defined herein. R' can be further optionally substituted in the same manner as alkyl.


By "haloaryl" is meant -R'(R)_n , where n is an integer from 1 to 5 and R' is arylene and R is halogen, as defined herein. R' can be further optionally substituted in the same manner as aryl.

By "haloarylalkyl" is meant -R'(R''(R'''))_n , where n is an integer from 1 to 5 and R' is alkylene, R'' is arylene, and R''' is halogen, as defined herein. R' can be further optionally substituted in the same manner as alkyl. R'' can be further optionally substituted in the same manner as aryl.

By "halogen" is meant fluoro, chloro, bromo, or iodo.

By "heterocyclic radical" is meant a 5-, 6- or 7-membered ring, unless otherwise specified, containing one, two, three, or four heteroatoms independently selected from the group containing nitrogen, oxygen, and sulfur. The 5-membered ring has zero to one double bonds, and the 6- and 7-membered rings have zero to two double bonds. Certain heterocyclyl groups include from 1 to 9 carbon atoms. Other such groups may include up to 12 carbon atoms. The term "heterocyclyl" also represents a heterocyclic compound having a bridged multicyclic structure in which one or more carbons and/or heteroatoms bridges two non-adjacent members of a monocyclic ring, *e.g.*, a quinuclidinyl group. The term "heterocyclyl" includes bicyclic, tricyclic, and tetracyclic groups in which any of the above heterocyclic rings is fused to one, two, or three carbocyclic rings, *e.g.*, an aryl ring, a cyclohexane ring, a cyclohexene ring, a cyclopentane ring, a cyclopentene ring, or another monocyclic heterocyclic ring, such as indolyl, quinolyl, isoquinolyl, tetrahydroquinolyl, benzofuryl, benzothienyl and the like. Examples of fused heterocyclyls include tropanes and 1,2,3,5,8,8a-hexahydroindolizine. Heterocyclics include pyrrolyl, pyrrolinyl, pyrrolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, piperidinyl, homopiperidinyl, pyrazinyl, piperazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiomorpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, indolyl, quinolyl, isoquinolyl, benzimidazolyl, benzothiazolyl, benzoxazolyl, furyl, thienyl, thiazolidinyl, isothiazolyl, isoindazolyl, triazolyl, tetrazolyl, oxadiazolyl, purinyl, thiadiazolyl (*e.g.*, 1,3,4-thiadiazole), tetrahydrofuranlyl, dihydrofuranlyl, tetrahydrothienyl, dihydrothienyl, dihydroindolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, pyranlyl, dihydropyranlyl, dithiazolyl, benzofuranlyl, benzothienyl and the like. Still other exemplary heterocyclyls include: 2,3,4,5-tetrahydro-2-oxo-oxazolyl; 2,3-dihydro-2-oxo-1H-imidazolyl; 2,3,4,5-tetrahydro-5-oxo-1H-pyrazolyl (*e.g.*, 2,3,4,5-tetrahydro-2-phenyl-5-oxo-1H-pyrazolyl); 2,3,4,5-tetrahydro-2,4-dioxo-1H-imidazolyl (*e.g.*, 2,3,4,5-tetrahydro-2,4-dioxo-5-methyl-5-phenyl-1H-imidazolyl); 2,3-dihydro-2-thioxo-1,3,4-oxadiazolyl (*e.g.*, 2,3-dihydro-2-thioxo-5-phenyl-1,3,4-oxadiazolyl); 4,5-dihydro-5-oxo-1H-triazolyl (*e.g.*, 4,5-dihydro-3-methyl-4-amino 5-oxo-1H-triazolyl); 1,2,3,4-tetrahydro-2,4-dioxypyridinyl (*e.g.*, 1,2,3,4-tetrahydro-2,4-dioxo-

3,3-diethylpyridinyl); 2,6-dioxo-piperidinyl (e.g., 2,6-dioxo-3-ethyl-3-phenylpiperidinyl); 1,6-dihydro-6-oxopyrimidinyl; 1,6-dihydro-4-oxopyrimidinyl (e.g., 2-(methylthio)-1,6-dihydro-4-oxo-5-methylpyrimidin-1-yl); 1,2,3,4-tetrahydro-2,4-dioxopyrimidinyl (e.g., 1,2,3,4-tetrahydro-2,4-dioxo-3-ethylpyrimidinyl); 1,6-dihydro-6-oxo-pyridazinyl (e.g., 1,6-dihydro-6-oxo-3-ethylpyridazinyl); 1,6-dihydro-6-oxo-1,2,4-triazinyl (e.g., 1,6-dihydro-5-isopropyl-6-oxo-1,2,4-triazinyl); 2,3-dihydro-2-oxo-1*H*-indolyl (e.g., 3,3-dimethyl-2,3-dihydro-2-oxo-1*H*-indolyl and 2,3-dihydro-2-oxo-3,3'-spirop propane-1*H*-indol-1-yl); 1,3-dihydro-1-oxo-2*H*-iso-indolyl; 1,3-dihydro-1,3-dioxo-2*H*-iso-indolyl; 1*H*-benzopyrazolyl (e.g., 1-(ethoxycarbonyl)-1*H*-benzopyrazolyl); 2,3-dihydro-2-oxo-1*H*-benzimidazolyl (e.g., 3-ethyl-2,3-dihydro-2-oxo-1*H*-benzimidazolyl); 2,3-dihydro-2-oxo-benzoxazolyl (e.g., 5-chloro-2,3-dihydro-2-oxo-benzoxazolyl); 2,3-dihydro-2-oxo-benzoxazolyl; 2-oxo-2*H*-benzopyranlyl; 1,4-benzodioxanyl; 1,3-benzodioxanyl; 2,3-dihydro-3-oxo-4*H*-1,3-benzothiazinyl; 3,4-dihydro-4-oxo-3*H*-quinazolinyl (e.g., 2-methyl-3,4-dihydro-4-oxo-3*H*-quinazolinyl); 1,2,3,4-tetrahydro-2,4-dioxo-3*H*-quinazolyl (e.g., 1-ethyl-1,2,3,4-tetrahydro-2,4-dioxo-3*H*-quinazolyl); 1,2,3,6-tetrahydro-2,6-dioxo-7*H*-purinyl (e.g., 1,2,3,6-tetrahydro-1,3-dimethyl-2,6-dioxo-7*H*-purinyl); 1,2,3,6-tetrahydro-2,6-dioxo-1*H*-purinyl (e.g., 1,2,3,6-tetrahydro-3,7-dimethyl-2,6-dioxo-1*H*-purinyl); 2-oxobenz[*c,d*]indolyl; 1,1-dioxo-2*H*-naphth[1,8-*c,d*]isothiazolyl; and 1,8-naphthylenedicarboxamido. Heterocyclic groups also include groups of the formula



F' is selected from the group consisting of -CH₂-, -CH₂O- and -O-, and G' is selected from the group consisting of -C(O)- and -(C(R')(R''))_v-, where each of R' and R'' is, independently, selected from the group consisting of hydrogen or alkyl of one to four carbon atoms, and v is one to three and includes groups, such as 1,3-benzodioxolyl, 1,4-benzodioxanyl, and the like. Any of the heterocyclyl groups mentioned herein may be optionally substituted with one, two, three, four or five substituents independently selected from the group consisting of: (1) alkanoyl (e.g., formyl, acetyl, and the like); (2) alkyl (e.g., alkoxyalkylene, alkylsulfanylalkylene, aminoalkylene, azidoalkylene, acylalkylene, haloalkylene (e.g., perfluoroalkyl), hydroxyalkylene, nitroalkylene, or thioalkoxyalkylene); (3) alkenyl; (4) alkynyl; (5) alkoxy (e.g., perfluoroalkoxy); (6) alkylsulfanyl; (7) aryl; (8) amino; (9) aryl-alkylene; (10) azido; (11) cycloalkyl; (12) cycloalkyl-alkylene; (13) cycloalkenyl; (14) cycloalkenyl-alkylene; (15) halo; (16) heterocyclyl (e.g., heteroaryl); (17) (heterocyclyl)oxy; (18) (heterocyclyl)aza; (19) hydroxy; (20) oxo; (21) nitro; (22) sulfide; (23) thioalkoxy; (24) -(CH₂)_qCO₂R^A, where q is an integer from zero to four, and R^A is selected from the group consisting of (a) alkyl, (b) aryl, (c) hydrogen, and (d) aryl-alkylene; (25) -(CH₂)_qCONR^BR^C, where q is an integer from zero to four and where R^B and R^C are independently selected from the group consisting of (a) hydrogen, (b) alkyl, (c) aryl, and (d) aryl-alkylene; (26) -(CH₂)_qSO₂R^D, where q is an integer from zero to four and where R^D is selected from the group consisting of (a) alkyl, (b) aryl, and (c) aryl-alkylene; (27) -(CH₂)_qSO₂NR^ER^F, where q is an integer from zero to four and where each of R^E and R^F is, independently, selected from the group consisting of (a) hydrogen, (b) alkyl, (c) aryl, and (d) aryl-alkylene; (28) thiol; (29) aryloxy; (30) cycloalkoxy; (31) arylalkoxy; (31) heterocyclyl-alkylene (e.g., heteroaryl-alkylene); (32) silyl; (33) cyano; and (34) -S(O)R^H where R^H is selected from the group consisting of (a) hydrogen, (b) alkyl, (c) aryl, and (d) aryl-alkylene. In some embodiments, each of these groups can be further substituted as described herein. For example, the alkylene group of an aryl-C₁-alkylene or a heterocyclyl-C₁-alkylene can be further substituted with an oxo group to afford the respective aryloyl and (heterocyclyl)oyl substituent group. In addition, when a heterocyclyl group is present in a bioreversible group of the invention it may be substituted

with an ester, thioester, or disulfide group that is bound to a conjugating moiety, a hydrophilic functional group, or an auxiliary moiety as defined herein.

By "heterocyclic radical alkyl," as used herein, represents an alkyl group substituted with a heterocyclic radical. The heterocyclic radical and alkyl portions may be substituted as the individual groups as described herein.

By "hydroxyalkyl" is meant $-R'(R)_n$, where n 1 or 2, R' is alkylene and R is hydroxyl, as defined herein. R' can be further optionally substituted in the same manner as alkyl.

By "hydroxyaryl" is meant $-R'(R)_n$, where n is 1 or 2, R' is arylene and R is hydroxyl, as defined herein. R' can be further optionally substituted in the same manner as aryl.

By "hydroxyl" is meant $-OH$.

By "hydroxyl protecting group" is meant any group capable of protecting the oxygen atom to which it is attached from reacting or bonding. Hydroxyl protecting groups are known in the art, e.g., as described in Wuts, Greene's Protective Groups in Organic Synthesis, Wiley-Interscience, 4th Edition, 2006. Exemplary protecting groups (with the oxygen atom to which they are attached) are independently selected from the group consisting of esters, carbonates, carbamates, sulfonates, and ethers. In exemplary ester hydroxyl protecting groups, R of the acyl group is C_{1-12} alkyl (e.g., C_{1-8} , C_{1-6} , C_{1-4} , C_{2-7} , C_{3-12} , and C_{3-6} alkyl), C_{2-12} alkenyl (e.g., C_{2-8} , C_{2-6} , C_{2-4} , C_{3-12} , and C_{3-6} alkenyl), carbocyclic C_{6-20} aryl (e.g., C_{6-15} , C_{6-10} , C_{8-20} , and C_{8-15} aryl), monocyclic C_{1-6} heteroaryl (e.g., C_{1-4} and C_{2-6} heteroaryl), C_{4-19} heteroaryl (e.g., C_{4-10} heteroaryl), $(C_{6-15})aryl(C_{1-6})alkyl$, $(C_{4-19})heteroaryl(C_{1-6})alkyl$, or $(C_{1-6})heteroaryl(C_{1-6})alkyl$. Specific examples of acyl groups for use in esters include formyl, benzoyl/formyl, acetyl (e.g., unsubstituted or chloroacetyl, trifluoroacetyl, methoxyacetyl, triphenylmethoxyacetyl, and *p*-chlorophenoxyacetyl), 3-phenylpropionyl, 4-oxopentanoyl, 4,4-(ethylenedithio)pentanoyl, pivaloyl (Piv), vinylpivaloyl, crotonoyl, 4-methoxy-crotonoyl, naphthoyl (e.g., 1- or 2-naphthoyl), and benzoyl (e.g., unsubstituted or substituted, e.g., *p*-methoxybenzoyl, phthaloyl (including salts, such as triethylamine and potassium), *p*-bromobenzoyl, and 2,4,6-trimethylbenzoyl).

As defined herein, any heteroaryl group present in an ester group has from 1 to 4 heteroatoms selected independently from O, N, and S. In exemplary carbonate hydroxyl protecting groups, R is C_{1-12} alkyl (e.g., C_{1-8} , C_{1-6} , C_{1-4} , C_{2-7} , C_{3-12} , and C_{3-6} alkyl), C_{2-12} alkenyl (e.g., C_{2-8} , C_{2-6} , C_{2-4} , C_{3-12} , and C_{3-6} alkenyl), carbocyclic C_{6-20} aryl (e.g., C_{6-15} , C_{6-10} , C_{8-20} , and C_{8-15} aryl), monocyclic C_{1-6} heteroaryl (e.g., C_{1-4} and C_{2-6} heteroaryl), C_{4-19} heteroaryl (e.g., C_{4-10} heteroaryl), $(C_{6-15})aryl(C_{1-6})alkyl$, $(C_{4-19})heteroaryl(C_{1-6})alkyl$, or $(C_{1-6})heteroaryl(C_{1-6})alkyl$. Specific examples include methyl, 9-fluorenylmethyl, ethyl, 2,2,2-trichloroethyl, 2-(trimethylsilyl)ethyl, 2-(phenylsulfonyl)ethyl, vinyl, allyl, *t*-butyl, *p*-nitrobenzyl, and benzyl carbonates. As defined herein, any heteroaryl group present in a carbonate group has from 1 to 4 heteroatoms selected independently from O, N, and S. In exemplary carbamate hydroxyl protecting groups, each R is independently H, C_{1-12} alkyl (e.g., C_{1-8} , C_{1-6} , C_{1-4} , C_{2-7} , C_{3-12} , and C_{3-6} alkyl), C_{2-12} alkenyl (e.g., C_{2-8} , C_{2-6} , C_{2-4} , C_{3-12} , and C_{3-6} alkenyl), carbocyclic C_{6-20} aryl (e.g., C_{6-15} , C_{6-10} , C_{8-20} , and C_{8-15} aryl), monocyclic C_{1-6} heteroaryl (e.g., C_{1-4} and C_{2-6} heteroaryl), C_{4-19} heteroaryl (e.g., C_{4-10} heteroaryl), $(C_{6-15})aryl(C_{1-6})alkyl$, $(C_{4-19})heteroaryl(C_{1-6})alkyl$, or $(C_{1-6})heteroaryl(C_{1-6})alkyl$. Specific examples include *N*-phenyl and *N*-methyl-*N*-(*o*-

nitrophenyl) carbamates. As defined herein, any heteroaryl group present in a carbamate group has from 1 to 4 heteroatoms selected independently from O, N, and S. Exemplary ether hydroxyl protecting groups include C₁₋₁₂ alkyl (e.g., C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, and C₃₋₆ alkyl), C₂₋₁₂ alkenyl (e.g., C₂₋₈, C₂₋₆, C₂₋₄, C₃₋₁₂, and C₃₋₆ alkenyl), (C₆₋₁₅)aryl(C₁₋₆)alkyl, (C₄₋₁₉)heteroaryl(C₁₋₆)alkyl, (C₁₋₆)heteroaryl(C₁₋₆)alkyl, (C₁₋₆)alkoxy(C₁₋₆)alkyl, (C₁₋₆)alkylthio(C₁₋₆)alkyl, (C₆₋₁₀)aryl(C₁₋₆)alkoxy(C₁₋₆)alkyl, and silyl (e.g., tri(C₁₋₆ alkyl)silyl, tri(C₆₋₁₀ aryl or C₁₋₆ heteroaryl)silyl, di(C₆₋₁₀ aryl or C₁₋₆ heteroaryl)(C₁₋₆ alkyl)silyl, and (C₆₋₁₀ aryl or C₁₋₆ heteroaryl)di(C₁₋₆ alkyl)silyl). Specific examples of alkylethers include methyl and t-butyl, and an example of an alkenyl ether is allyl. Ether hydroxyl protecting groups can be used to protect a carboxyl group (e.g., with a C₁₋₁₂ alkyl (e.g., C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, and C₃₋₆ alkyl), (C₆₋₁₅)aryl(C₁₋₆)alkyl, (C₁₋₆)alkoxy(C₁₋₆)alkyl, (C₁₋₆)alkylthio(C₁₋₆)alkyl, or (C₆₋₁₀)aryl(C₁₋₆)alkoxy(C₁₋₆)alkyl). Examples of alkoxyalkyls and alkylthioalkyls that can be used as ether hydroxyl protecting groups include methoxymethyl, methylthiomethyl, (2-methoxyethoxy)methyl, and β-(trimethylsilyl)ethoxymethyl. Examples of arylalkyl groups that can be used as ether hydroxyl protecting groups include benzyl, p-methoxybenzyl (MPM), 3,4-dimethoxybenzyl, triphenylmethyl (trityl), o-nitrobenzyl, p-nitrobenzyl, p-halobenzyl, 2,6-dichlorobenzyl, p-cyanobenzyl, naphthylmethyl, and 2- and 4-picolyl ethers. Specific examples of silylethers include trimethylsilyl (TMS), triethylsilyl (TES), t-butyldimethylsilyl (TBS), t-butyldiphenylsilyl (TBDPS), triisopropylsilyl (TIPS), and triphenylsilyl (TPS) ethers. An example of an arylalkyloxyalkylether is benzyloxymethyl ether. As defined herein, any heteroaryl group present in an ether group has from 1 to 4 heteroatoms selected independently from O, N, and S. Vicinal or 1,3-diols may be protected with a diol protecting group (e.g., to produce a "cyclic protected diol"), such as acetal (e.g., containing C₁₋₆ alkylene), ketal (e.g., containing C₃₋₆ alkylene or C₃₋₆ cycloalkyl), cyclic silylene, cyclic carbonate, and cyclic boronate. Examples of acetal and ketal groups include methylene-dioxo, ethylidene-dioxo, benzylidene-dioxo, isopropylidene-dioxo, cyclohexylidene-dioxo, and cyclopentylidene-dioxo. An example of a cyclic silylene is di-t-butylsilylene. Another diol protecting group is 1,1,3,3-tetraisopropylsiloxanediyl. Examples of cyclic boronates include methyl, ethyl, phenyl, and 2,6-diacetamidophenyl boronates. Protecting groups may be substituted as is known in the art; for example, aryl and arylalkyl groups, such as phenyl, benzyl, naphthyl, or pyridinyl, can be substituted with C₁₋₆ alkyl, C₁₋₆ alkoxy, nitro, cyano, carboxyl, or halogen. Alkyl groups, such as methyl, ethyl, isopropyl, n-propyl, t-butyl, n-butyl, and sec-butyl, and alkenyl groups, such as vinyl and allyl, can also be substituted with oxo, arylsulfonyl, halogen, and trialkylsilyl groups. Preferred protecting groups are TBS and Piv. Protecting groups that are orthogonal are removed under different conditions, as is known in the art.

By "imido" is meant --NR_2 , where each R is independently optionally substituted acyl.

By "ketal" is meant $\text{--O--CR}_2\text{--O--}$, where each R is independently optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl, or both R groups are together optionally substituted alkylene, or each of the R groups is a bond to an enumerated carbon atom, as shown in Chart 1, within the intermediate or within the halichondrin macrolide or analog thereof.

By "macrocyclic" is meant a compound containing at least one *n*-membered ring, where *n* is equal to or greater than 10.

By "non-enolizable" is meant a group that, either alone or in combination with a group to which it is attached, cannot form an enol through a deprotonation/reprotonation sequence. For example, a "non-enolizable alkyl" can be bonded to a sulfone group or to a carbonyl group through a quaternary carbon atom (i.e., the carbon atom that is not bonded to a hydrogen atom).

5

By "*N*-protecting group" is meant a group protecting a nitrogen atom in a molecule from participating in one or more undesirable reactions during chemical synthesis (e.g., oxidation reactions, or certain nucleophilic and electrophilic substitutions). Commonly used *N*-protecting groups are disclosed in Wuts, Greene's Protective Groups in Organic Synthesis, Wiley-Interscience, 4th Edition, 2006. Exemplary *N*-protecting groups include acyl (e.g., formyl, acetyl, trifluoroacetyl, propionyl, pivaloyl, t-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, phthalyl, o-nitrophenoxycarbonyl, α-chlorobutyryl, benzoyl, 4-chlorobenzoyl, and 4-bromobenzoyl); sulfonyl-containing groups (e.g., benzenesulfonyl, p-toluenesulfonyl, o-nitrobenzenesulfonyl, and p-nitrobenzenesulfonyl); carbamate forming groups (e.g., benzyloxycarbonyl, p-chlorobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl, α,α-dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxy carbonyl, t-butyloxycarbonyl, diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, phenoxy carbonyl, 4-nitrophenoxy carbonyl, fluorenyl-9-methoxycarbonyl, cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl, and phenylthiocarbonyl), arylalkyl (e.g., triphenylmethyl); silyl groups (e.g., trimethylsilyl); and imine-forming groups (e.g., diphenylmethylenes). Preferred *N*-protecting groups are acetyl, benzoyl, phenylsulfonyl, p-toluenesulfonyl, p-nitrobenzenesulfonyl, o-nitrobenzenesulfonyl, t-butyloxycarbonyl (Boc), and benzyloxycarbonyl (Cbz).

25

By "oxo" or (O) is meant =O.

By "pharmaceutically acceptable salt" is meant a salt within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and animals without undue toxicity, irritation, allergic response and the like and commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, pharmaceutically acceptable salts are described in: Berge et al., J. Pharmaceutical Sciences 66:1-19, 1977 and in Pharmaceutical Salts: Properties, Selection, and Use, (Eds. P.H. Stahl and C.G. Wermuth), Wiley-VCH, 2008. Representative acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts and the like. A preferred salt is the mesylate salt.

40

By "pseudohalogen" is meant $\text{--O--SO}_2\text{R}$, where R is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl. Non-limiting examples of pseudohalogens include trifluoromethanesulfonate and nonaflate.

- 5 By "silyl" is meant --SiR_3 , where each R is independently alkyl, alkenyl, aryl, or arylalkyl. Examples of silyl groups include tri(C_{1-6} alkyl)silyl, tri(C_{6-10} aryl or C_{1-6} heteroaryl)silyl, di(C_{6-10} aryl or C_{1-6} heteroaryl)(C_{1-6} alkyl)silyl, and (C_{6-10} aryl or C_{1-6} heteroaryl)di(C_{1-6} alkyl)silyl. It will be understood that, when a silyl group includes two or more alkyl, alkenyl, aryl, heteroaryl, or arylalkyl groups, these groups are independently selected. As defined herein, any heteroaryl group present in a silyl group has from 1 to 4 heteroatoms
10 selected independently from O, N, and S. Silyl can be optionally substituted in the same manner as defined for each R group.

- By "silylene" is meant $\text{--SiR}_2\text{--}$, where each R is independently alkyl, alkenyl, aryl, arylalkyl, or alkoxy. By "dialkylsilylene" is meant a silylene, where each R is alkyl. Silylene can be optionally substituted in the same
15 manner as defined for each R group. Silylene-dioxo is a group having the formula $\text{--O--SiR}_2\text{--O--}$.

- By "strong base" is meant a Brønsted base, the conjugate acid of which has pKa that is greater than or equal to 13. Non-limiting examples of strong bases include alkyl alkali metals (e.g., butyl lithium or Schlosser's base), Grignard reagents (e.g., alkyl magnesium halide), alkali or alkali earth alkoxides, alkali or alkali earth
20 amides (e.g., diisopropylamide, tetramethylpiperidide, or bis(trimethylsilyl)amide), and phosphazene bases (e.g., Schwesinger base). Non-limiting examples of the alkali amides are lithium diisopropylamide, lithium tetramethylpiperidide, lithium bis(trimethylsilyl)amide, sodium bis(trimethylsilyl)amide, and potassium bis(trimethylsilyl)amide.

- 25 By "sulfonamide" is meant --NR , where R is sulfonyl.

- By "sulfonate" is meant $\text{--OS(O)}_2\text{R}$, where R is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl. In exemplary sulfonates, R is C_{1-12} alkyl (e.g., C_{1-8} , C_{1-6} , C_{1-4} , C_{2-7} , C_{3-12} , or C_{3-6} alkyl), C_{2-12} alkenyl (e.g., C_{2-8} , C_{2-6} , C_{2-4} , C_{3-12} , or C_{3-6} alkenyl), carbocyclic
30 C_{6-20} aryl (e.g., C_{6-15} , C_{6-10} , C_{8-20} , or C_{8-15} aryl), monocyclic C_{1-6} heteroaryl (e.g., C_{1-4} and C_{2-6} heteroaryl), C_{4-19} heteroaryl (e.g., C_{4-10} heteroaryl), (C_{6-15})aryl(C_{1-6})alkyl, (C_{4-19})heteroaryl(C_{1-6})alkyl, or (C_{1-6})heteroaryl(C_{1-6})alkyl. As defined herein, any heteroaryl group present in a sulfonate group has from 1 to 4 heteroatoms selected independently from O, N, and S.

- 35 By "sulfonyl" is meant $\text{--S(O)}_2\text{R}$, where R is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, optionally substituted arylalkyl, or silyl. Preferred R groups for sulfonyl are the same as those described above for sulfonates.

- By "thioacetal" is meant --S--(CHR)--S-- , where R is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl.
40

By "thioketal" is meant $-S-(CR_2)-S-$, where each R is independently optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl.

By "triflate" is meant trifluoromethanesulfonate.

5

The pKa values recited herein refer to the pKa values of a conjugate Brønsted acid in water at room temperature, unless stated otherwise.

References to compounds described herein encompass their salts, where applicable.


10

Detailed Description

The present invention provides compounds and methods that may be useful in the synthesis of halichondrin macrolides and analogs thereof (see Chart 1). Preferably, the halichondrin macrolide analog is eribulin or a salt thereof (e.g., eribulin mesylate). Preferably, the halichondrin macrolide is a halichondrin B macrolide.


15 The carbon-atom numbering schemes for a halichondrin macrolide and analog thereof are shown in Chart 1.

Chart 1



20 in which each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, a hydroxyl protecting group, and A is a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , or A is a group of formula (1):

25



where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_2 is H and P_1 is absent, H, alkyl, or a hydroxyl protecting group, or R_2 and P_1 combine to form a


30 bond;

(i) R_3 is H, and P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

(ii) R_3 is $-(CH_2)_nNP_3P_4$, where P_3 is H or an N-protecting group, and (a) P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_4 is H or an N-protecting group, (b) P_2 and P_4 combine to form an alkylidene, or (c) each of P_2 and P_4 is H;

(iii) R_3 is $-(CH_2)_nOP_5$, where P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_5 is H, optionally substituted alkyl, or a hydroxyl protecting group; or P_2 and P_5 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group; each of A_1 , A_2 , and A_3 is independently H or OP'' , where each P'' is independently H or a hydroxyl protecting group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_B(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

n , when present, is 0, 1, or 2; and


k is 0 or 1.

In the halichondrin macrolide and halichondrin macrolide analog structures, when P_1 is absent, each of D and D' is independently H or optionally substituted alkyl, and, when P_2 is absent, L is $-C(O)-$.

Prins Reaction


In one aspect, the invention provides a compound of formula (A) and methods of its preparation. The compound of formula (A) may be prepared from a compound of formula (B), a compound of formula (C), and R_5OH , where R_5 is optionally substituted acyl. For example, a compound of formula (B), a compound of formula (C), and R_5OH may be subjected to Prins reaction conditions, e.g., as known in the art. For example, the compound of formula (B), the compound of formula (C), and R_5OH may be reacted with a Lewis acid (e.g., an oxophilic Lewis acid (e.g., boron trifluoride or solvate thereof)).

The compound of formula (A) is:



where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, a hydroxyl protecting group, and A is a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , or A is a group of formula (1):



where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;



R_2 is H and P_1 is absent, H, alkyl, or a hydroxyl protecting group, or R_2 and P_1 combine to form a bond;

(i) R_3 is H, and P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

(ii) R_3 is $-(CH_2)_nNP_3P_4$, where P_3 is H or an N-protecting group, and (a) P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_4 is H or an N-protecting group, (b) P_2 and P_4 combine to form an alkylidene, or (c) each of P_2 and P_4 is H;

(iii) R_3 is $-(CH_2)_nOP_5$, where P_2 is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P_5 is H, optionally substituted alkyl, or a hydroxyl protecting group; or P_2 and P_5 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;


5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;
 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_B(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 n, when present, is 0, 1, or 2;

k is 0 or 1;

R₁ is -OP₆, -CH(Y)₂, or -CH₂(Y), where P₆ is H or a hydroxyl protecting group;

R₄ is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl,



15 or

where

each P₇ is independently H or a hydroxyl protecting group; and

R₃ is -CH₂CH₂-COOR_C, -CH=CH-COOR_C, -CH₂CH₂-SO₂R_D, or -CH=CH-SO₂R_D;

R₅ is optionally substituted acyl;

20 R₆ is H, optionally substituted alkyl, or optionally substituted arylalkyl;

each Y is independently -COOR_C or -SO₂R_D;


each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and

each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl.

5

In the compound of formula (A), when P_1 is absent, each of D and D' is independently H or optionally substituted alkyl, and, when P_2 is absent, L is $-C(O)-$.

The compound of formula (B) is:



10


where all variables are as described for formula (A).

The compound of formula (C) is:

15



where


R_7 is $-CHO$ or $R_{7A}O$ , where each R_{7A} is independently an optionally substituted alkyl, and R_4 is as described for the compound of formula (A).

20

The compound of formula (B) may be prepared from a compound of formula (D) and a compound of formula (E). For example, the compound of formula (D) and the compound of formula (E) may be subjected to Sakurai reaction conditions, e.g., as known in the art. In certain embodiments, the compound of formula (D) may be reacted with the compound of formula (E) and a Lewis acid (e.g., an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)) to produce the compound of formula (B).

25

The compound of formula (D) is:




30

where

all variables are as described for the compound of formula (B).

The compound of formula (E) is:




5 where

R₅ is silyl (e.g., Me₃Si-); and

R₆ is as described for the compound of formula (B).

The Sakurai reaction producing the compound of formula (B) introduces the secondary alcohol stereogenic center in the compound of formula (B). If the stereoselectivity of the Sakurai reaction is less than desirable, the Sakurai reaction product can be subjected to epimerization to enrich the product in the compound of formula (B) relative to its diastereomer. In a non-limiting example, the Sakurai reaction products (the compound of formula (B) and its C.27 diastereomer) can be reacted with an oxidizing agent capable of converting an alcohol to a carbonyl group (e.g., Dess-Martin periodinane) to give a compound of formula (F), and the compound (F) can then be subjected to enantioselective 1,2-reduction reaction conditions (e.g., Corey-Bakshi-Shibata reaction). Alternatively, if the diastereoselectivity of the Sakurai reaction favors the formation of the diastereomer of the compound of formula (B), the Mitsunobu reaction may be used to invert the secondary alcohol stereocenter in a diastereomeric Sakurai reaction product mixture.


20 The compound of formula (F) is:



where all variables are as described for formula (B).

25 Enantioselective 1,2-reduction reaction conditions are known in the art. In a non-limiting example, the enantioselective 1,2-reduction reaction is Corey-Bakshi-Shibata reaction. Corey-Bakshi-Shibata reduction can include reacting a ketone with CBS catalyst and borane or a solvate thereof. In a non-limiting example, (S)-CBS-oxazaborolidine catalyst can be used with borane or a solvate thereof (e.g., BH₃·THF) to control the stereochemistry of the resulting secondary alcohol in the preparation of the compound of formula (II) from the compound of formula (F).

Mitsunobu reaction conditions are known in the art. In a non-limiting example, the diastereomeric Sakurai reaction product mixture may be reacted with an azidicarboxylate compound (e.g., DIAD), phosphine (e.g., PPh₃), and a carboxylic acid (e.g., 3,5-dinitrobenzoic acid) to produce a compound of formula (G):



where

R_{Ac} is optionally substituted acyl; and

5 the remaining variables are as described for formula (B).

The compound of formula (G) may be converted to the compound of formula (B). In a non-limiting example, the compound of formula (G) may be subjected to alcoholysis or hydrolysis reaction conditions (e.g., $Mg(OMe)_2$ in MeOH) to produce the compound of formula (B). Alternatively, the compound of formula (G)

10 may be reacted with a 1,2-reducing agent to produce the compound of formula (B).

In the compounds described herein, R_6 may be optionally substituted alkyl (e.g., methyl).


In the compounds described herein, D may be H. In the compounds described herein, D' may be OP_1 (e.g., P_1 may be alkyl (e.g., methyl)). In the compounds described herein, G may be O.

15

In the compounds described herein, k may be 0, and R_1 may be $-CH_2(Y)$, where Y may be $-SO_2R_D$ where, R_D is optionally substituted aryl (e.g., phenyl) or optionally substituted non-enolizable alkyl.


20 In the compounds described herein, E may be optionally substituted alkyl (e.g., methyl).

In the compounds described herein, D may be H, and A may be:



25 In the compounds described herein, k may be 0, and R_1 may be $-CH(Y)_2$, or $-CH_2(Y)$. In the compounds described herein, R_3 may be $-(CH_2)_nNP_3P_4$ or $-(CH_2)_nOP_5$, where n may be 0.

In the compounds described herein, A and D may combine to form the following structure:



30 where the bond to the oxygen atom originates at the carbon atom to which D is attached in formula (A) or formula (B). In the compounds described herein, R_3 may be $-(CH_2)_nNP_3P_4$ or $-(CH_2)_nOP_5$, and n is 2.

In the compounds described herein, k may be 1, and E may be optionally substituted alkyl (e.g., methyl). In the compounds described herein, R₁ may be -CH₂(Y), where Y may be -COOR_D, where R_D may be optionally substituted alkyl (e.g., methyl).

5 Preparation of Halichondrin Macrolides and Analogs Thereof


A halichondrin macrolide or analog thereof may be prepared as described herein and using methods and reactions known in the art. Non-limiting examples of methods and reactions useful for the preparation of a halichondrin macrolide or analog thereof include U.S. patent application publication Nos. 2016/0264594, 2015/0158881, 2011/0184190, 2011/0054194, 2009/0203771, and 2009/0198074; International patent application publication No. WO 2016/179607; U.S. patent Nos. 5,338,865, 5,436,238, 6,214,865, and 8,445,701; and in Towle et al., *Annual Meeting of the American Association for Cancer Research*, April 6-10, 2002, 5721; Wang et al., *Bioorg. Med. Chem. Lett.*, 10:1029-1032, 2000; Aicher et al., *J. Am. Chem. Soc.*, 114:3162-3164, 1992; Ueda et al., *J. Am. Chem. Soc.*, 136:5171-5176; and Yamamoto et al., *J. Am. Chem. Soc.*, 134:893-896, 2012.

As described herein, one of skill in the art can identify the sequence of reactions involving hydroxyl protecting group removing agents and Brønsted acids to convert the compounds described below into a halichondrin macrolide or analog thereof. One of skill in the art can recognize that certain functional groups require protecting groups known in the art to reduce or prevent undesired reactivity. One of skill in the art can select appropriate protecting groups to be used in the syntheses described herein.

Chart 2 illustrates the macrocycle disconnections that may be useful for the preparation of a halichondrin macrolide or analog thereof.

25

Chart 2



a halichondrin macrolide or analog thereof

where X₁ is -O- or -CH₂-, and the remaining variables are as described in Chart 1.

Table 1 lists the macrocyclic retrosynthesis disconnections shown in Chart 2 and indicates corresponding exemplary bond-forming reactions as described herein. Any of these various reactions can be employed in the synthesis of halichondrin macrolide or analogs thereof. For example, these reactions may be employed to couple various fragments together or to form the macrocycle.


5

Table 1

Disconnection	Exemplary Reaction	Reference(s)
C.1-X ₁	Esterification (e.g., of the compound of formula (VIIB)) Claisen reaction (e.g., of the compound of formula (VIIB))	US 5,338,865 US 6,214,865 US 2016/0264594
C.2-C.3	Olefin Metathesis (e.g., of the compound of formula (VIE)) Horner-Wadsworth-Emmons (e.g., of the compound of formula (VIG))	US 2016/0264594 WO 2016/179607
C.3-C.4	Olefin Metathesis (e.g., of the compound of formula (VIC))	US 2016/0264594 WO 2016/179607
C.12-C.13	Olefin Metathesis (e.g., of the compound of formula (VIIB))	WO 2016/179607
C.13-C.14	Nozaki-Hiyama-Kishi (e.g., of the compound of formula (VIID))	US 2009/0203771
C.15-C.16	Olefin metathesis (e.g., of the compound of formula (IVB))	US 2016/0264594 WO 2016/179607
C.19-C.20	Nozaki-Hiyama-Kishi (e.g., of the compound of formula (VB))	US 2016/0264594 WO 2016/179607
C.23-C.24	Prins reaction (e.g., producing the compound of formula (IXB))	-

The methods of preparing a halichondrin macrolide or analog thereof disclosed herein include a C.23-C.24 bond-forming Prins reaction either as a macrocyclization step or as a step in the synthesis of non-macrocyclic intermediates. In some approaches, a halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)) may be prepared from the compound of formula (IA) using methods and reaction conditions known in the art.


A compound of formula (IA) may be produced from a compound of formula (IIA), a compound of formula (IIB), a compound of formula (III), and R₅OH. The compound of formula (IA) is:



where

R₁ is -OP₆, -CH(Y)₂, or -CH₂(Y), where P₆ is H or a hydroxyl protecting group, and each Y is independently -COOR_C or -SO₂R_D;

20



each P_7 is independently H or a hydroxyl protecting group;

R_8 is $-\text{CH}_2\text{CH}_2-\text{COOR}_C$, $-\text{CH}=\text{CH}-\text{COOR}_C$, $-\text{CH}_2\text{CH}_2-\text{SO}_2\text{R}_D$, or $-\text{CH}=\text{CH}-\text{SO}_2\text{R}_D$;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and


each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl; and

R_5 is optionally substituted acyl;

and the remaining variables are as described for the halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)).

In the compound of formula (IA), when P_1 is absent, each of D and D' is independently H or optionally substituted alkyl.

The compound of formula (IIA) is:



where

R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$,

where

m is 1, and R_F is a hydroxyl protecting group,

or

m is 0, and


(i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene;

and

the remaining variables are as described for the compound of formula (IA).

The compound of formula (IIB) is:




where R_S is silyl (e.g., Me_3Si-).

5

The compound of formula (III) is:




where



Preparation of the compound of formula (IA) may include reacting the compound of formula (IIA) (in which R_E is $-CHO$), the compound of formula (IIB), and a Lewis acid (e.g., an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)) to produce a compound of formula (IIC):

15



where the variables are as described for the compound of formula (IA).

20


The compound of formula (III) and the compound of formula (IIC) may be reacted under Prins reaction conditions to produce the compound of formula (IA). Prins reaction conditions are known in the art. In a non-limiting example, the compound of formula (III) and the compound of formula (IIC) may be reacted with a Lewis acid (e.g., an oxophilic Lewis acid (e.g., boron trifluoride or a solvate thereof)).

25

The following describes exemplary conditions for macrocyclization at the indicated positions. These reaction conditions may also be used to couple two or more fragments prior to macrocyclization.

C.1-X₁ Bond-forming Macrocyclization

30 A halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)) may be prepared through a C.0-C.1 or a O-C.1 bond-forming macrocyclization according to the following synthesis strategy from a compound of formula (VIIB):




(VIIB)

where


- (i) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;
 5 (ii) R_{10} is a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond;
 or
 (iii) R_{10} and R_{11} combine to form a bond, and R_{12} is H;

L_4 is




- 10 R_1 is $-\text{OP}_6$, $-\text{CH}_2(\text{Y})$, or $-\text{CH}(\text{Y})_2$, where P_6 is H or a hydroxyl protecting group, and Y is $-\text{COOR}_c$ or $-\text{CHO}$;
 each R_c , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and
 A_1 and R_{14} combine to form oxo, R_{13} is H or a hydroxyl protecting group, and R_{15} is H;
 or
 15 A_1 is H or $-\text{OP}''$, and:
 (i) R_{13} is H or a hydroxyl protecting group, and R_{14} and R_{15} combine to form a double bond;
 or
 (ii) R_{13} and R_{14} combine to form a bond, and R_{15} is H or $-\text{OP}''$;
 20 R_5 is optionally substituted acyl;
 R_9 is $-\text{CHO}$ or $-\text{COOP}''$;
 each P'' is independently H or a hydroxyl protecting group;
 each P_9 is independently a hydroxyl protecting group, and X_3 is oxo or X_3 , together with the carbon atom to which it is attached, is $-(\text{CH}(\text{OP}_{11}))$, where P_{11} is H or a hydroxyl protecting group; or both P_9 groups and
 25 X_3 , together with the atoms to which each is attached, combine to form a ketal;
 and the remaining variables are as described for the halichondrin macrolide or analog thereof.

The compound of formula (VIIB) is converted to a compound of formula (VIIC):



where

L_1 is



X_1 is $-\text{CH}_2-$, $-\text{CH}(\text{Y})-$, $-\text{C}(\text{Y})_2-$, or $-\text{O}-$;


X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P_{12} is H or a hydroxyl protecting group;

and the remaining variables are as described for the compound of formula (VIIB).

Preparation of the compound of formula (VIIC) may include an esterification reaction between R_1 that is $-\text{OP}_6$, where P_6 is H, and R_9 that is $-\text{COOH}$. Alternatively, preparation of the compound of formula (VIIC) may include a reaction between R_1 that is $-\text{CH}(\text{Y})_2$ or $-\text{CH}_2(\text{Y})$, and R_9 that is $-\text{CHO}$ (e.g., under Claisen reaction conditions). Oxidation of the Claisen product (X_4 , together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P_{12} is H) with an oxidizing agent capable of converting an alcohol to a carbonyl group may produce X_4 that is oxo. Further desulfonylation or decarboxylation of this compound of formula (VIIC) may produce the compound of formula (VIIC), in which X_1 is $-\text{CH}_2-$.

The compound of formula (VIIC) is then converted to the halichondrin macrolide or analog thereof using methods described herein and those known in the art, e.g., those described in US 5,338,865, US 2016/0264594, US 2009/0203771, and WO 2016/179607. For example, preparation of the halichondrin macrolide or analog thereof may include reacting the compound of formula (VIIC) with a hydroxyl protecting group removing agent.

The compound of formula (VIIB) may be prepared from a compound of formula (VIIA), the compound of formula (IIA), the compound of formula (IIB), a compound of formula (IIIC), and $R_5\text{OH}$. The compound of formula (VIIA) is:



where

- (i) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;
- 5 (ii) R_{10} is a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond;
- or
- (iii) R_{10} and R_{11} combine to form a bond, and R_{12} is H;
- R_9 is $-\text{CH}_2-\text{OP}''$, $-\text{CHO}$, or $-\text{COOP}''$;
- R_{13} and each P_9 is independently a hydroxyl protecting group;
- 10 A_1 is H or $-\text{OP}''$;
- each P'' is independently H or a hydroxyl protecting group; and
- Y_1 is chloro, bromo, iodo, trifluoromethanesulfonate, or trialkylsilane.

The compound of formula (IIIC) is:




where

R_7 is $-\text{CHO}$, $-\text{CH}_2\text{OP}_A$,

or $R_7\text{AO}-\text{CH}(\text{OR}_{7A})-\text{CH}_2-$, where each R_{7A} is independently an optionally substituted alkyl; and

20 R_{4C} is



where

- R_C is optionally substituted alkyl, optionally substituted aryl, or optionally substituted aryl/alkyl;
- 25 P_A is H or a hydroxyl protecting group; and
- P_7 , when present, is independently H or a hydroxyl protecting group.


Preparation of the compound of formula (VIIB) can be accomplished as follows. The compound of formula (IIIC), in which R_7 is $-\text{CH}_2\text{OP}_A$ or

30 $R_7\text{AO}-\text{CH}(\text{OR}_{7A})-\text{CH}_2-$, may be converted to a compound of formula (IIID):



where


R_{4D} is



all remaining variables are as described for the compound of formula (IIIC).


The compound of formula (IIID) may be reacted with the compound of formula (VIIA) under Nozaki-Hiyama-Kishi reaction conditions (e.g., with a Cr(II) salt and a Ni(II) salt as described herein) to form a compound of formula (VIIAa):

10



where

R_7 is $-\text{CH}_2\text{OP}_A$, $-\text{CHO}$, or



and all remaining variables are as described for the compound of formula (VIIB) and the compound of formula (IIID).


The compound of formula (VIIAa), in which R_7 is $-\text{CHO}$, may be reacted with a compound of formula (IIC) and $R_5\text{OH}$ under Prins reaction conditions to form the compound of formula (VIIB).

20

Alternatively, preparation of the compound of formula (VIIB) can be accomplished by reacting the compound of formula (IG) and the compound of formula (VIIA) under Nozaki-Hiyama-Kishi reaction conditions (e.g., with a Cr(II) salt and a Ni(II) salt as described herein).


25

The compound of formula (IG) is:



where


5 R_{E1} is



where R_5 is optionally substituted acyl;


and the remaining variables are as described for the compound of formula (VIIB).

10 The compound of formula (IG) may be formed from a compound of formula (IF):




where all variables are as described for the compound of formula (IG) and the compound of formula (IIIC).


15 R_{4C} that is




may be reacted with a glycol cleaving agent to produce an aldehyde, which may then be reacted with $(R_cO)_2P(O)-CH_2-COOR_c$ to produce R_{4C} that is




R_{4C} that is




may be reacted with a 1,4-reducing agent to produce R_{4C} that is



5 R_{4C} that is




may be reacted with an allylic reducing agent to produce R_{4C} that is




(e.g., producing this R_{4C} may further include an esterification reaction after the reaction with an allylic reducing agent).


10 R_{4C} that is




may be reacted with a 1,2-reducing agent to produce



R_{4C} that is




15 may be reacted with an oxidizing agent capable of converting an alcohol to a carbonyl group in



In general, the method described herein includes an allylic reduction at C.25. This reaction can be performed at any point prior to the formation of the halichondrin macrolide or analog thereof. For example,


20



C.2-C.3 and C.3-C.4 Bond-Forming Macrocyclizations

In another approach, a halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)) may be prepared through a C.2-C.3 or a C.3-C.4 bond-forming macrocyclization according to the following strategy.


In some embodiments, a halichondrin macrolide or analog thereof is prepared from a compound of formula (VIB) is:



(VIB)

where

L₄ is



R₁ is $-\text{OP}_6$ or $-\text{CH}(\text{Y})$, where P₆ is H or a hydroxyl protecting group, and Y is $-\text{COOR}_c$ or $-\text{CHO}$;

R₅ is optionally substituted acyl;

R_{10A} is $-\text{CH}_2-\text{CH}=\text{CH}_2$, $-(\text{CH}_2)_2-\text{CH}=\text{CH}_2$, or $-(\text{CH}_2)_3-\text{OP}_{10}$;

each of R₁₀ and P₁₀ is independently a hydroxyl protecting group;

A₁ and R₁₄ combine to form oxo, R₁₃ is H or a hydroxyl protecting group, and R₁₅ is H;

or

A₁ is H or $-\text{OP}''$, and:

(i) R₁₃ is H or a hydroxyl protecting group, and R₁₄ and R₁₅ combine to form a double bond;

or

(ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or $-\text{OP}''$;


each P₉ is independently a hydroxyl protecting group, and X₃ is oxo or X₃, together with the carbon atom to which it is attached, is $-(\text{CH}(\text{OP}_{11}))$, where P₁₁ is H or a hydroxyl protecting group; or both P₉ groups and X₃, together with the atoms to which each is attached, combine to form a ketal;

and

the remaining variables are as described for the halichondrin macrolide or analog thereof.


The compound of formula (VIB) may be converted to the halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)) using reaction conditions known in the art, e.g., as described herein.

In particular embodiments, preparation of the halichondrin macrolide or analog thereof from the compound of formula (VIB) through C.3-C.4 bond-forming macrocyclization includes producing a compound of formula (VIC) from the compound of formula (VIB). The compound of formula (VIC) is



where

L₅ is




X₁ is $-\text{CH}_2-$ or $-\text{O}-$;

X₄ is oxo, or X₄, together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P₁₂ is H or a hydroxyl protecting group; and

the remaining variables are as described for the compound of formula (VIB).


The compound of formula (VIC) may be prepared by reacting the compound of formula (VIB) with R₁₇-CH₂CH=CH₂, where R₁₇ is $-\text{COOH}$ or a metallic or metalloid moiety. The compound of formula (VIB), in which R₁ is $-\text{CH}_2(\text{Y})$, may be reacted with R₁₇-CH₂CH=CH₂, in which R₁₇ is a metallic or metalloid moiety, to produce the compound of formula (VIC), in which X₁ is $-\text{CH}_2-$, and X₄ is oxo or $-(\text{CH}(\text{OP}_{12}))-$. The compound of formula (VIB), in which R₁ is $-\text{OP}_6$, and P₆ is H, may be esterified with R₁₇-CH₂CH=CH₂, where R₁₇ is $-\text{COOH}$, to produce the compound of formula (VIC), in which X₁ is $-\text{O}-$, and X₄ is oxo.

The compound of formula (VIC) is then converted to a compound of formula (VID):



where

5 L₁ is



, where L₂ and L₃ combine to form a bond;
and the remaining variables are as described for the compound of formula (VIC).


10 The compound of formula (VID) is produced through a reaction of the compound of formula (VIC) with an olefin metathesis catalyst.

The compound of formula (VID) may be converted to the halichondrin macrolide or analog thereof using methods described herein and those known in the art, e.g., those described in US 2016/0264594, US 2009/0203771, and WO 2016/179607.

15

Preparation of the halichondrin macrolide or analog thereof may include reacting the compound of formula (VID) with a hydroxyl protecting group removing agent. A reaction of the compound of formula (VID) with a hydroxyl protecting group removing agent may lead to the C.3-C.4 double bond isomerization to give an enoate/enone described herein upon exposure to basic (e.g., isomerization mediated by a hydroxyl protecting group removing agent, such as a fluoride source) or acidic (e.g., isomerization mediated by a Brønsted acid) conditions. The resulting enoate/enone may undergo a reaction with a hydroxyl of -OR₁₀.


20 In certain embodiments, preparation of the halichondrin macrolide or analog thereof from the compound of formula (VIB) through C.2-C.3 bond-forming macrocyclization includes producing a compound of formula (VIE) from the compound of formula (VIB). The compound of formula (VIE) is:



(VIE)

where

L_6 is



X_1 is $-\text{CH}_2-$ or $-\text{O}-$;


X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P_{12} is H or a hydroxyl protecting group; and

the remaining variables are as described for the compound of formula (VIB).

The compound of formula (VIB) may be converted to a compound of formula (VIE) through a reaction with $\text{R}_{17}-\text{CH}=\text{CH}_2$, where R_{17} is $-\text{COOH}$ or a metallic or metalloid moiety. For example, the compound of formula (VIB), in which R_1 is $-\text{CH}_2(\text{Y})$, can be reacted with $\text{R}_{17}-\text{CH}_2\text{CH}=\text{CH}_2$, where R_{17} is a metallic or metalloid moiety, to give the compound of formula (VIE), in which X_1 is $-\text{CH}_2-$, and X_4 is oxo or $-(\text{CH}(\text{OP}_{12}))-$.

Alternatively, the compound of formula (VIB), in which R_1 is $-\text{OP}_6$, and P_6 is H, can be esterified with $\text{R}_{17}-\text{CH}_2\text{CH}=\text{CH}_2$, where R_{17} is $-\text{COOH}$, to give the compound of formula (VIE), in which X_1 is $-\text{O}-$, and X_4 is oxo.


The compound of formula (VIE) may be converted to a compound of formula (VIF):



(VIF)

where


L₁ is



, where L₂ and L₃ combine to form a bond;
and the remaining variables are as described for the formula (VIE).

- 5 The compound of formula (VIF) may be produced by a reaction of the compound of formula (VIE) with an olefin metathesis catalyst.

Alternatively, the compound of formula (VIF), in which X₄ is oxo, may be prepared from a compound of formula (VIG):




10

(VIG)

where

L₇ is



- 15 X₁ is -CH₂- or -O-;

each R_C is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

and the remaining variables are as described for the compound of formula (VIF).

- 20 The compound of formula (VIG) may be prepared by reacting the compound of formula (VIB) with (R_CO)₂P(O)-CH₂-R_P, where each R_C is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl, and R_P is H or -COOH. For example, the compound of formula (VIB), in which R₁ is -CH₂(Y), may be reacted with (R_CO)₂P(O)-CH₂-R_P, in which R_P is H, under Claisen reaction conditions to produce the compound of formula (VIG). Alternatively, the compound of formula (VIB), in

which R_1 is $-\text{OP}_6$, where P_6 is H, may be esterified with $(\text{R}_c\text{O})_2\text{P}(\text{O})-\text{CH}_2-\text{R}_p$, in which R_p is $-\text{COOH}$, to produce the compound of formula (VIG).


The compound of formula (VIG) is then converted to the compound of formula (VIF), e.g., through a Horner-Wadsworth-Emmons reaction, where all variables are as described for the compound of formula (VIG).

The compound of formula (VIF) is then converted to the halichondrin macrolide or analog thereof using methods described herein and those known in the art, e.g., those described in US 2016/0264594, US 2009/0203771, and WO 2016/179607.

10

The compound of formula (VIB) may be prepared from a compound of formula (VIA), a compound of formula (IIA), a compound of formula (IIB), a compound of formula (IIIC), and R_5OH .

The compound of formula (VIA) is:



15

where

each R_{10} , R_{13} , and P_9 is independently a hydroxyl protecting group;


A_1 is H or $-\text{OP}''$;

Y_1 is chloro, bromo, iodo, trifluoromethanesulfonate, or trialkylsilane; and

$\text{R}_{10\text{A}}$ is as described for the compound of formula (VIB).

20

The compound of formula (IIA) is:



25

where

R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$,

where

m is 1, and R_F is a hydroxyl protecting group,

30

or


m is 0, and

(i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene;

and the remaining variables are as described for the compound of formula (VIB).

The compound of formula (IIB) is:




5 where R_S is silyl;

The compound of formula (IIIC) is:




10 where

R_{4C} is




, where R_C is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl, and each P_7 is independently H or a hydroxyl protecting group; and

15 R_7 is $-CHO$ or




where each R_{7A} is independently an optionally substituted alkyl;


20 Preparation of the compound of formula (VIB) includes a Sakurai reaction between R_E that is $-CHO$ and the compound of formula (IIB) producing a group of the structure:



. This product may then be reacted with the compound of formula (IIIC) under Prins reaction conditions to produce a group of the structure:



. This group may be converted to




, as described herein.


If the stereochemistry of the secondary alcohol in the Sakurai reaction product differs from the desired stereochemistry, this group may be subjected to epimerization reaction conditions described herein (e.g., oxidation followed by Corey-Bakshi-Shibata reduction; alternatively, Mitsunobu reaction may be used to invert the stereogenic center).

If R_E is $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$, preparation of the compound of formula (IVB) may further include conversion of R_E to $-\text{CHO}$ under acetal deprotecting conditions, e.g., through a reaction with an aqueous Brønsted acid (e.g., when m is 0), or through oxidation using an oxidizing agent capable of converting an alcohol to a carbonyl group (e.g., when m is 1).


R_{4C} that is




may be reacted with a glycol cleaving agent to produce an aldehyde, which may then be reacted with $(\text{R}_C\text{O})_2\text{P}(\text{O})-\text{CH}_2-\text{COOR}_C$ to produce R_{4C} that is




R_{4C} that is




may be reacted with a 1,4-reducing agent to produce R_{4C} that is



R_{4C} that is




may be reacted with an allylic reducing agent to produce R_{4C} that is




(e.g., producing this R_{4C} may further include an esterification reaction after the reaction with an allylic reducing agent).


R_{4C} that is




may be reacted with a 1,2-reducing agent to produce




5 R_{4C} that is



may be reacted with an oxidizing agent capable of converting an alcohol to a carbonyl group in




10 In some embodiments, the compound of formula (VIB) is produced from the compound of formula (VIA) and a compound of formula (IG):



where

15 R_{E1} is



where R₅ is optionally substituted acyl;


and the remaining variables are as described for the compound of formula (VIB).

20 The compound of formula (VIA) is reacted with the compound of formula (IG) under Nozaki-Hiyama-Kishi reaction conditions to produce the compound of formula (VIB). For example, Nozaki-Hiyama-Kishi reaction on the compound of formula (VIA) and the compound of formula (IG) can include reacting the compound of formula (VIA) and the compound of formula (IG) with a Cr(II) salt and a Ni(II) salt.

The compound of formula (IG) may be prepared from the compound of formula (IIA), the compound of formula (IIB), the compound of formula (IIIC), and R_5OH , where R_5 is optionally substituted acyl.

In further embodiments, the compound of formula (VIB) may be prepared according to the following strategy.

- 5 The compound of formula (IIIC) may be converted to the compound of formula (IIID) as described herein. Further, the compound of formula (IIID) may be reacted with the compound of formula (VIA) under Nozaki-Hiyama-Kishi reaction conditions to produce a compound of formula (VIAa):




(VIAa)

- 10 where all variables are as described for the compound of formula (VIB) and the compound of formula (IIIC).

The compound of formula (VIAa), in which R_7 is $-CHO$, may be reacted with a compound of formula (IIC) and R_5OH under Prins reaction conditions to form the compound of formula (VIB).

- 15 In yet further embodiments, the compound of formula (VIC) may be prepared according to the following strategy. The compound of formula (IIA), in which R_E is $-CH_{(1+m)}(OR_F)_{(2-m)}$, may be reacted with $R_{17}-CH_2CH=CH_2$, where R_{17} is $-COOH$ or a metallic or metalloid moiety, to produce a compound of formula (IID):




(IID)

- 20 where all variables are as described for the compound of formula (VIC).


- For example, the compound of formula (IIA), in which R_1 is $-CH_2(Y)$, may be reacted with $R_{17}-CH_2CH=CH_2$, in which R_{17} is a metallic or metalloid moiety, to produce the compound of formula (VIC), in which X_1 is $-CH_2-$, and X_4 is oxo or $-(CH(OP_{12}))-$. The compound of formula (IIA), in which R_1 is $-OP_6$, and P_6 is H, may be esterified with $R_{17}-CH_2CH=CH_2$, where R_{17} is $-COOH$, to produce the compound of formula (IID), in which X_1 is $-O-$, and X_4 is oxo.

- 25 The compound of formula (IID), in which R_E is $-CHO$, may be reacted with the compound of formula (IIB) under Sakurai reaction conditions to produce a compound of formula (IIE):




where all variables are as described for the compound of formula (VIC).

- 5 The compound of formula (IIE) may be reacted with the compound of formula (VIAa) and R_5OH under Prins reaction conditions to produce the compound of formula (VIC), in which L_5 is:




- 10 In still further embodiments, the compound of formula (VIE) may be prepared according to the following strategy. The compound of formula (IIA) may be converted to a compound of formula (IIF) through a reaction with $R_{17}-CH=CH_2$, where R_{17} is $-COOH$ or a metallic or metalloid moiety. The compound of formula (IIF) is:



- 15 where all variables are as described for the compound of formula (VIE).


- For example, the compound of formula (IIA), in which R_1 is $-CH_2(Y)$, can be reacted with $R_{17}-CH_2CH=CH_2$, where R_{17} is a metallic or metalloid moiety, to give the compound of formula (IIF), in which X_1 is $-CH_2-$, and X_4 is oxo or $-(CH(OP_{12}))-$. Alternatively, the compound of formula (IIA), in which R_1 is $-OP_6$, and P_6 is H, can be esterified with $R_{17}-CH_2CH=CH_2$, where R_{17} is $-COOH$, to give the compound of formula (IIF), in which X_1 is $-O-$, and X_4 is oxo.
- 20

The compound of formula (IIF), in which R_E is $-CHO$, may be reacted with the compound of formula (IIB) under Sakurai reaction conditions to produce a compound of formula (IIG):




where all variables are as described for the compound of formula (VIE).

- 5 The compound of formula (IIF) may be reacted with the compound of formula (VIAa) and R_5OH under Prins reaction conditions to produce the compound of formula (VIE), in which L_6 is:



In some embodiments, the compound of formula (VIG) may be prepared according to the following strategy.

- 10 The compound of formula (IIA) may be converted to a compound of formula (IIH) through a reaction with $(R_CO)_2P(O)-CH_2-R_P$, where each R_C is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl, and R_P is H or $-COOH$. The compound of formula (IIH) is:




- 15 where

each R_C is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(CH(OP_{12}))_n-$, where P_{12} is H or a hydroxyl protecting group; and


- 20 all remaining variables are as described for the compound of formula (VIG).

The compound of formula (IIH), in which X_4 is oxo, may be converted to a compound of formula (III) through a reaction with the compound of formula (IIB) under Sakurai reaction conditions. The compound of formula (III) is:



where all variables are as described for the compound of formula (IIH).


- 5 The compound of formula (III) may be reacted with the compound of formula (VIAa), in which R_{10A} is $-\text{CH}_2-\text{CH}=\text{CH}_2$ or $-(\text{CH}_2)_3-\text{OP}_{10}$, to produce a compound of formula (VIGa):



where

- 10 R_{10A} is $-\text{CH}_2-\text{CH}=\text{CH}_2$ or $-(\text{CH}_2)_3-\text{OP}_{10}$,


L_7 is




and all remaining variables are as described for the compound of formula (VIG).

- 15 The compound of formula (VIGa), in which R_{10A} is $-\text{CH}_2-\text{CH}=\text{CH}_2$, may be converted to the compound of formula (VIGa), in which R_{10A} is $-(\text{CH}_2)_3-\text{OP}_{10}$, and P_{10} is H, by hydroboration/oxidation reaction. Hydroboration/oxidation reactions are known in the art. For example, 9-BBN or thexyl borane may be used for the hydroboration of R_{10A} , and sodium perborate or hydrogen peroxide and base may be used for the oxidation step providing $-(\text{CH}_2)_3-\text{OP}_{10}$, and P_{10} is H. The compound of formula (VIGa), in which R_{10A} is $-(\text{CH}_2)_3-\text{OP}_{10}$, and P_{10} is H, may be converted to the compound of formula (VIG).
- 20

In general, the method described herein includes an allylic reduction at C.25. This reaction can be performed at any point prior to the formation of the halichondrin macrolide or analog thereof. For example,




may be converted to



using an allylic reducing agent.

C.12-C.13 Bond-forming Macrocyclization

In still another approach, a halichondrin macrolide or analog thereof may be prepared from the compound of
5 of formula (VIII B):



(VIII B)

where

X_1 is $-CH_2-$, $-CH(Y)-$, $-C(Y)_2-$, or $-O-$, where each Y is independently $-COOR_C$ or $-SO_2R_D$;


10 X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(CH(OP_{12}))-$, where P_{12} is H or a hydroxyl protecting group; each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

- 15 (i) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;
 (ii) R_{10} is a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;
 or
 (iii) R_{10} and R_{11} combine to form a bond, and R_{12} is H;

each P_9 and R_{13} is independently a hydroxyl protecting group;

20 R_M is




where

R_5 is optionally substituted acyl; and

X₃ is oxo or X₃, together with the carbon atom to which it is attached, is $-(CH(OP_{11}))-$, where P₁₁ is H or a hydroxyl protecting group;

and the remaining variables are as described for the halichondrin macrolide or analog thereof.


- 5 The compound of formula (VIIIB) can be subjected to ring closing metathesis to produce a compound of formula (VIIIC):



where

- 10 A_1 is H or $-OP''$, where P'' is H or a hydroxyl protecting group;

L_1 is




OR_5 $\overset{\text{Me}}{\underset{|}{\text{C}}}$ or $\overset{\text{Me}}{\underset{|}{\text{C}}}$, where L_2 and L_3 combine to form a bond; and
 and the remaining variables are as described for the compound of formula (VIIb).

- 15 The halichondrin macrolide or analog thereof is then produced from the compound of formula (VIIC), e.g., according to the methods described in WO 2016/179607.

In general, the fragments may be coupled in any order when forming compound (VIII B). For example, the C.0-C.1 or O-C.1 bond between fragment (IIA) and fragment (VIII A) can be formed, as described herein, prior to or after the Sakurai and/or Prins reaction. Similarly, installation of the C.13 olefin may occur before or after any of the Sakurai, Prins, and C.0-C.1 or O-C.1 bond forming reactions.

In one embodiment, the compound of formula (VIII B) is produced from a compound of formula (VIII A), a compound of formula (II A), and a compound of formula (II B), a compound of formula (III C), and R_5OH . The compound of formula (II A) is:



where R_1 is $-\text{OP}_6$, $-\text{CH}(\text{Y})_2$, or $-\text{CH}_2(\text{Y})$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-\text{COOR}_C$ or $-\text{SO}_2\text{R}_D$;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

5 each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl; and

R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$, where

m is 1, and R_F is a hydroxyl protecting group,


or

10 m is 0, and

(i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene.

The compound of formula (VIII A) is:



15

where

(i) R_{10} is a hydroxyl protecting group, R_{11} is alkyl ether, and R_{12} is H;

(ii) R_{10} is a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond; or

20 (iii) R_{10} and R_{11} combine to form a bond, and R_{12} is H;

R_9 is $-\text{CHO}$ or $-\text{COOH}$; and

each P_9 and R_{13} is independently a hydroxyl protecting group.

The compound of formula (IIIC) is:


25

$R_{4C}-R_7$,

(IIIC)


where

R_7 is $-\text{CHO}$ or



, where each R_{7A} is independently an optionally substituted alkyl; and

30 R_{4C} is




where R_C is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;
and P_7 , when present, is independently H or a hydroxyl protecting group.

The compound of formula (IIB) is:




where R_S is silyl.


In certain embodiments, a compound of formula (IIC):



A compound of formula (IF):



A compound of formula (IJ):




25

The compound of formula (VIIIB) can be produced from the compound of formula (VIIIA) and the compound of formula (IJ) as described herein.


- Preparation of the compound of formula (VIIIB) may include an esterification reaction between R_1 that is –
 5 OP_6 , where P_6 is H, and R_9 that is $-COOH$. Alternatively, preparation of the compound of formula (VIIIB) may include a reaction between R_1 that is $-CH(Y)_2$ or $-CH_2(Y)$, and R_9 that is $-CHO$ (e.g., under Claisen reaction conditions). Oxidation of the Claisen product (X_4 , together with the atom to which it is attached, is –
 (CH(OP₁₂))–, where P_{12} is H) with an oxidizing agent capable of converting an alcohol to a carbonyl group may produce X_4 that is oxo. Further desulfonylation or decarboxylation of this compound of formula (VIIIB)
 10 may produce the compound of formula (VIIIB), in which X_1 is $-CH_2-$.

- Alternatively, the compound of formula (VIIIB) may be produced from the compound of formula (IF) and the compound of formula (VIIIA). For example, a compound of formula (VIIID) may be produced from the compound of formula (IF) and the compound of formula (VIIIA) using methods described herein. The
 15 compound of formula (VIIID) is:



where

R_{M1} is




R_5 is optionally substituted acyl;

P_{13} is H or a hydroxyl protecting group;

and the remaining variables are as described for the compound of formula (IF) and the compound of formula (VIIIB).

25


Alternatively, the compound of formula (VIIID) may be prepared from the compound of formula (IIC) and the compound of formula (VIIIA) as follows. The compound of formula (IIC) may be protected with a hydroxyl protecting group to give a compound of formula (IIJ):



where P_{13} is a hydroxyl protecting group, and the remaining variables are as described for the compound of formula (IIC).


5

The compound of formula (IIJ) may be reacted with the compound of formula (VIIIA) using methods described herein to produce the compound of formula (VIID), in which R_{M1} is:




, where P_{13} is a hydroxyl protecting group.

The compound of formula (VIID), in which R_{M1} is




10

, and P_{13} is H, may be reacted with the compound of formula (IIC) and R_5OH under Prins reaction conditions to produce the compound of formula (VIID), in which R_{M1} is



The compound of formula (VIID) may also be produced according to the following strategy. The compound of formula (IIA) and the compound of formula (VIIIA) may be converted to a compound of formula (VIII E) using methods described herein. The compound of formula (VIII E) is:



15



where all variables are as described for the compound of formula (VIID) and the compound of formula (IIA).



20

The compound of formula (VIID), in which R_{M1} is




 or , may be converted to the compound of formula (VIII B) as described herein.

In general, the method described herein includes an allylic reduction at C.25. This reaction can be performed at any point prior to the formation of the halichondrin macrolide or analog thereof.


In certain embodiments, the method includes converting


 to 
 using an allylic reducing agent.


Generally, the C.12 olefin can be introduced at any point prior to the metathesis reaction via several routes. For example, if R_{4C} is

, reaction with an allylic reducing agent can produce a compound in which R_{4C} is , which can be reduced with a 1,2-reducing agent (e.g., DIBAL-H) to produce the corresponding aldehyde, which can be reacted with a vinyl nucleophile.

If R_{4C} is

, the compound can be reduced with a 1,4-reducing agent (e.g., Stryker's reagent), and the product can be reduced with a 1,2-reducing agent (e.g., DIBAL-H) to produce the corresponding aldehyde, which can be reacted with a vinyl nucleophile.

If R_{4C} is

, the compound can be reacted with a glycol cleaving agent (e.g., NaIO₄), (R_{4C}O)₂P(O)-CH₂-COOR_{4C} (under Horner-Wadsworth-Emmons reaction conditions), a 1,4-reducing agent (e.g., Stryker's reagent), and a 1,2-reducing agent (e.g., DIBAL-H) to produce the corresponding aldehyde, which can be reacted with a vinyl nucleophile. Vinyl nucleophiles are known in the art. Vinyl


nucleophiles may be prepared in situ or in a separate reaction vessel. Reaction conditions for adding vinyl nucleophiles to carbonyl groups are known in the art.

The compound of formula (VIIC) can be formed by reacting the compound of formula (VIIB) with an olefin metathesis catalyst.

The halichondrin macrolide may be produced from the compound of formula (VIIC) using reaction conditions known in the art and those described herein. For example, the compound of formula (VIIC) may be reacted with a hydroxyl protecting group removing agent to produce the halichondrin macrolide.

10


In general, the method described herein includes an allylic reduction at C.25. This reaction can be performed at any point prior to the formation of the halichondrin macrolide or analog thereof. For example,



15

C.13-C.14 Bond-forming Macrocyclization

In still another approach, a halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)) may be prepared from the compound of formula (VIID):



20 where

- (i) R₁₀ is a hydroxyl protecting group, R₁₁ is alkyl ether, and R₁₂ is H;
- (ii) R₁₀ is a hydroxyl protecting group, and R₁₁ and R₁₂ combine to form a double bond;
- or
- (iii) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;

25 X₁ is -CH₂-, -CH(Y)-, -C(Y)₂-, or -O-,

where each Y is independently -COOR_C or -SO₂R_D;

each R_C, when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D, when present, is independently optionally substituted aryl or optionally substituted

30 non-enolizable alkyl;


X₄ is oxo, or X₄, together with the atom to which it is attached, is $-(CH(OP_{12}))-$, where P₁₂ is H or a hydroxyl protecting group;

A₁ is H or $-OP''$, where P'' is H or a hydroxyl protecting group;

Y₁ is chloro, bromo, iodo, trifluoromethanesulfonate, or trialkylsilane;

5 R₁₃ and each P₉ is independently a hydroxyl protecting group;

R_L is



where


R₁₈ is $-CHO$ or $-CH_2OP_7$;

10 R₅ is optionally substituted acyl; and

P₇, when present, is independently H or a hydroxyl protecting group;

and the remaining variables are as described for the halichondrin macrolide or analog thereof.


The compound of formula (VIID) is then converted to a compound of formula (VIIE):



15

where

L₁ is



20

where L₂ and L₃ combine to form a bond;

each P₉ is independently a hydroxyl protecting group, and X₃ is oxo or X₃, together with the carbon atom to which it is attached, is $-(CH(OP_{11}))-$, where P₁₁ is H or a hydroxyl protecting group;

and the remaining variables are as described for the compound of formula (VIID).


25 In certain embodiments, the compound of formula (VIID), in which R₁₈ is $-CHO$, is subjected to Nozaki-Hiyama-Kishi reaction conditions to produce the compound of formula (VIIE).

The compound of formula (VII E) may then be converted to a halichondrin macrolide or analog thereof using methods described herein and those known in the art, e.g., those described in US 2009/0203771, US 2016/0264594, and WO 2016/179607.

5

In general, the fragments may be coupled in any order when forming the compound of formula (VIID). For example, the C.0-C.1 or O-C.1 bond in the compounds of formula (VIID) can be formed, as described herein, prior to or after the Sakurai and/or Prins reaction.

- 10 In some embodiments, a compound of formula (VIID) is produced from a compound of formula (IIA), a compound of formula (IIB), a compound of formula (IIIC), and R_5OH , where R_5 is optionally substituted acyl. The compound of formula (VIIA) is:




15 where

R_9 is $-CHO$ or $-COOH$; and

the remaining variables are as described for the compound of formula (VIID).

The compound of formula (IIA) is:



20

where

R_1 is $-OP_6$, $-CH(Y)_2$, or $-CH_2(Y)$,

where P_6 is H or a hydroxyl protecting group, and each Y is independently $-COOR_C$ or $-SO_2R_D$;

25 each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

and

30 R_E is $-CHO$ or $-CH_{(1+m)}(OR_F)_{(2-m)}$,


where

m is 1, and R_F is a hydroxyl protecting group, or

m is 0, and (i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene.

The compound of formula (IIB) is:




5 where R_S is silyl

The compound of formula (IIIC) is:




10 where

R_{4C} is



or R_7 is $-\text{CHO}$ or




15


where each R_{7A} is independently an optionally substituted alkyl; and

the remaining variables are as described for the compound of formula (VIID).

20 Preparation of the compound of formula (VIID) includes a Sakurai reaction between R_E that is $-\text{CHO}$ and the compound of formula (IIB) producing a group of the structure:



. This product may then be reacted with the compound of formula (IIIC) under the Prins reaction conditions to produce a group of the structure:




. This group may be converted to R_L as described herein.

25 If the stereochemistry of the secondary alcohol in the Sakurai reaction product differs from the desired stereochemistry, this group may be subjected to epimerization reaction conditions described herein (e.g., oxidation followed by Corey-Bakshi-Shibata reduction; alternatively, Mitsunobu reaction may be used to invert the stereogenic center).

If R_E is $-\text{CH}_{(1+m)}(\text{ORF})_{(2-m)}$, preparation of the compound of formula (VIID) may further include conversion of R_E to $-\text{CHO}$ under acetal deprotecting conditions, e.g., through a reaction with an aqueous Brønsted acid (e.g., when m is 0), or through oxidation using an oxidizing agent capable of converting an alcohol to a carbonyl group (e.g., when m is 1).


Preparation of the compound of formula (VIID) may include an esterification reaction between R_1 that is $-\text{OP}_6$, where P_6 is H, and R_9 that is $-\text{COOH}$. Alternatively, preparation of the compound of formula (VIID) may include a reaction between R_1 that is $-\text{CH}(\text{Y})_2$ or $-\text{CH}_2(\text{Y})$, and R_9 that is $-\text{CHO}$ (e.g., under Claisen reaction conditions). Further desulfonylation or decarboxylation of this compound of formula (VIID) may produce the compound of formula (VIID), in which X_1 is $-\text{CH}_2-$.

In some embodiments, preparation of the compound of formula (VIID) includes producing a compound of formula (IIC) from the compound of formula (IIA) and the compound of formula (IIB) (e.g., under Sakurai reaction conditions). The compound of formula (IIC) is:



where all variables are as described for the compound of formula (IIA).


The compound of formula (IIC) may be reacted with the compound of formula (IIB), the compound of formula (IIIC), and $R_5\text{OH}$ to produce a compound of formula (IF):



where all variables are as described for the compound of formula (IIA), the compound of formula (IIIC), and $R_5\text{OH}$.

Preparation of the compound of formula (IF) may include a Prins reaction between the compound of formula (IIC), the compound of formula (IIIC), and $R_5\text{OH}$.

The compound of formula (IF) may be converted to a compound of formula (IH):




(IH)


where all variables are described for the compound of formula (IF).

5

In general, the method described herein includes an allylic reduction at C.25. This reaction can be performed at any point prior to the formation of the halichondrin macrolide or analog thereof. For example,



may be converted to



using an allylic reducing agent.

- 10 In certain embodiments, the compound of formula (IF) may be reacted with an allylic reducing agent to produce the compound of formula (IH).

In the reactions described herein, the requisite transformations of groups R_{4C} may be performed using reaction conditions known in the art. In some embodiments, the transformations of groups R_{4C} are as


- 15 described herein.

C.15-C.16 Bond-forming Macrocyclization

In one approach, the macrocyclization reaction is a carbon-carbon bond-forming reaction (e.g., olefin metathesis) that provides a C.15-C.16 bond in a halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)). A non-macrocyclic intermediate in the synthesis of the halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)) may be a compound of formula (IVB).


20

The compound of formula (IVB) is:



where

- 5 each P₉ is independently a hydroxyl protecting group;
- (a1) R₁₀ is H or a hydroxyl protecting group, R₁₁ is alkyl ether, and R₁₂ is H;
- (a2) R₁₀ is H or a hydroxyl protecting group, and R₁₁ and R₁₂ combine to form a double bond;
- or
- (a3) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;
- 10 (b1) A₁ and R₁₄ combine to form oxo, R₁₃ is H or a hydroxyl protecting group, and R₁₅ is H;
- or
- (b2) A₁ is H or –OP^{''}, and:
- (i) R₁₃ is H or a hydroxyl protecting group, and R₁₄ and R₁₅ combine to form a double bond;
- 15 or
- (ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or –OP^{''}; and
- (c1) R₁₆ is H, and P₈ is H or a hydroxyl protecting group;
- or
- (c2) R₁₆ and P₈ combine to form a double bond;
- 20 each P^{''}, when present, is independently H or a hydroxyl protecting group
- X₁ is –O–, –C(Y)₂–, –CH(Y)–, or –CH₂–, where each Y is independently –COOR_C or –SO₂R_D, where each R_C, when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl, and each R_D, when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;
- 25 R_G is



where

R₅ is an optionally substituted acyl, and


X₂ is H or $-\text{CH}_2-\text{X}_{2\text{A}}-\text{CH}_2-\text{CH}=\text{CH}_2$, where X_{2A} is $-\text{O}-$, $-\text{C}(\text{R}_\text{H})_2-$, or $-\text{NR}_\text{I}-$, where each R_H is independently H or $-\text{COOR}_\text{J}$, R_I is an *N*-protecting group, and R_J is C₁₋₆ alkyl;

5 and

X₄ is oxo, or X₄, together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P₁₂ is H or a hydroxyl protecting group;


and the remaining variables are as described for the halichondrin macrolide or analog thereof.

10 In some embodiments, the compound of formula (IVB) is a compound of formula (IVBa):




where all variables are as described in the compounds of formula (IVB).

15 The compound of formula (IVB) (e.g., the compound of formula (IVBa)) is converted to the compound of formula (IVC) through an olefin metathesis reaction. The compound of formula (IVC) is:



where


20 L₁ is



where L₂ and L₃ combine to form a bond;

and the remaining variables are as described for the compound of formula (IVB).

The compound of formula (IVC) is then converted to the compound of formula (IVD):




5

where each of A₁ and A₂ is independently H or -OPⁿ, and the remaining variables are as described for the compound of formula (IVC).

The compound of formula (IVC) may be converted to the compound of formula (IVD) using reaction conditions known in the art. For example, the compound of formula (IVC) may be reacted with a 1,4-reducing agent to give the compound of formula (IVD).

Preparation of the compound of formula (IVD) may further include reacting the compound of formula (IVA), (IVB), or (IVC), in which P₈ and R₁₆ are both H, with an oxidizing agent capable of converting an alcohol to a carbonyl group (e.g., Dess-Martin periodinane or a dimethylsulfonium compound) to give the compound of formula (IVA), (IVB), or (IVC), in which R₁₆ and P₈ combine to form a double bond (e.g., the compound of formula IVCa)). If P₈ is a hydroxyl protecting group, prior to oxidation, the compound of formula (IVA), (IVB), or (IVC) may be treated with a hydroxyl protecting group removing agent. In a non-limiting example, preparation of the compound of formula (IVD) includes oxidizing the compound of formula (IVC), in which P₉ is H, R₁₆ is H, with an oxidizing agent capable of converting an alcohol to a carbonyl group (e.g., Dess-Martin periodinane or a dimethylsulfonium compound) to give a compound of formula (IVCa):



where all variables are described for the compound of formula (IVC).

25


The compound of formula (IVD) is then converted to the halichondrin macrolide or analog thereof using methods described herein and those known in the art, e.g., those described in US 2016/0264594 and WO 2016/179607.

- 5 If the compound of formula (IVD) includes hydroxyl protecting groups as R_3 and/or P_5 , these hydroxyl protecting groups can be removed with a hydroxyl protecting group removing agent. For example, a hydroxyl protecting group removing agent may be a fluoride source, if the hydroxyl protecting group is a silyl group.
- 10 If, in the compound of formula (IVD), each P_9 is H, and X_3 is oxo, the synthesis may further involve a reaction with a Brønsted acid (e.g., a Brønsted acid having a pK_a of 5 ± 3), e.g., after the reaction of the compound of formula (IVD) with a hydroxyl protecting group removing agent (e.g., to convert P_9 from a hydroxyl protecting group into H).
- 15 If, in the compound of formula (IVD), X_3 is oxo, R_{10} is a hydroxyl protecting group, and R_{11} and R_{12} combine to form a double bond, treatment with a hydroxyl protecting group removing agent can provide the compound of formula (IVD), in which R_{10} and R_{11} combine to form a bond, and R_{12} is H.

- In general, the fragments may be coupled in any order when forming compound (IVB). For example, the
- 20 C.0-C.1 or O-C.1 bond between fragment (IIA) and fragment (IVA) can be formed, as described herein, prior to or after the Sakurai and/or Prins reaction. Similarly, installation of the C.15 olefin may occur before or after any of the Sakurai, Prins, and C.0-C.1 or O-C.1 bond forming reactions.


- In certain embodiments, the compound of formula (IVB) is produced from a compound of formula (IIA), a
- 25 compound of formula (IIB), a compound of formula (IIIA), a compound of formula (IVA), and R_5OH .

The compound of formula (IVA) is:



- 30 where
 R_9 is $-CHO$ or $-COOH$,
 and the remaining variables are as described for the compound of formula (IVB).

The compound of formula (IIA) is:



where

- 5 R_1 is $-\text{OP}_6$, $-\text{CH}(\text{Y})_2$, or $-\text{CH}_2(\text{Y})$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-\text{COOR}_C$ or $-\text{SO}_2\text{R}_D$;

R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$,


where

m is 1, and R_F is a hydroxyl protecting group, or

- 10 m is 0, and (i) each R_F is independently an alkyl or hydroxyl protecting group, or
(ii) both R_F combine to form an alkylene;

and the remaining variables are as described for the compound of formula (IVB).

The compound of formula (IIB) is:




where R_S is silyl.

The compound of formula (IIIA) is:




where

R_7 is $-\text{CHO}$ or




- 25 where each R_{7A} is independently an optionally substituted alkyl; and
 R_{4A} is




where R_C is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl, and each P_7 is independently H or a hydroxyl protecting group.

30

Preparation of the compound of formula (IVB) includes a Sakurai reaction between R_E that is $-\text{CHO}$ and the compound of formula (IIB) producing a group of the structure:




. This product may then be reacted with the compound of formula (IIIA) under Prins reaction conditions to produce a group of the structure:




. This group may be converted to R_G as described herein.

- 5 If the stereochemistry of the secondary alcohol in the Sakurai reaction product differs from the desired stereochemistry, this group may be subjected to epimerization reaction conditions described herein (e.g., oxidation followed by Corey-Bakshi-Shibata reduction; alternatively, Mitsunobu reaction may be used to invert the stereogenic center).
- 10 If R_E is $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$, preparation of the compound of formula (IVB) may further include conversion of R_E to $-\text{CHO}$ under acetal deprotecting conditions, e.g., through a reaction with an aqueous Brønsted acid (e.g., when m is 0), or through oxidation using an oxidizing agent capable of converting an alcohol to a carbonyl group (e.g., when m is 1).

- 15 Preparation of R_G may include reacting R_{4A} that is




with a glycol cleaving agent, and subjecting the resulting product to Horner-Wadsworth-Emmons reaction with $(\text{R}_C\text{O})_2\text{P}(\text{O})-\text{CH}_2-\text{COOR}_C$ to produce R_{4A} that is




. This reaction may be performed after Sakurai and Prins reactions.

- 20 Preparation of R_G may further include a reaction of R_{4A} that is



with a 1,2-reducing agent and reacting the resulting product with an allyl halide or allyl pseudohalide. This reaction may be performed after Sakurai and Prins reactions.

- 25 Group R_E may be converted to R_G before or after the reaction of the compound of formula (IIA) with the compound of formula (IVA). For example, conversion of R_E to R_G before the reaction of the compound of formula (IIA) with the compound of formula (IVA) may be carried out via compounds of formulae (IB) and (IC), as described herein. Conversion of R_E to R_G after the reaction of the compound of formula (IIA) with the compound of formula (IVA) may be carried out via a compound of formula (IVAA):



(IVAA)

where

X_1 is $-\text{CH}_2-$, $-\text{CH}(\text{Y})-$, $-\text{C}(\text{Y})_2-$, or $-\text{O}-$, where each Y is independently $-\text{COOR}_\text{C}$ or $-\text{SO}_2\text{R}_\text{D}$;

5 each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;


each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

X_4 is oxo, or X_4 , together with the atom to which it is attached, is $-(\text{CH}(\text{OP}_{12}))-$, where P_{12} is H or a hydroxyl

10 protecting group; and

the remaining variables are as described for the compound of formulae (IIA) and (IVA).


The compound of formula (IVAA), in which R_E is $-\text{CHO}$, may be reacted with the compound of formula (IIB) under Sakurai reaction conditions to produce a compound of formula (IVAB):



(IVAB)

where all variables are as described for the compound of formula (IVAA).


A compound of formula (IVAc) may be produced from the compound of formula (IVAB), the compound of formula (IIIA), and R_5OH . The compound of formula (IVAc) is:



(IVAc)

where


R_{G1} is



R₅ is optionally substituted acyl, and


the remaining variables are as described for the compound of formula (IVAb).

10 For example, the compound of formula (IVAb) may be reacted with the compound of formula (IIIA) and R₅OH under Prins reaction conditions to produce the compound of formula (IVAc), in which R_{G1} is




The compound of formula (IVAc) may be converted to the compound of formula (IVB) as described herein.

15 If the compound described herein (e.g., a Prins reaction product) includes:



the synthesis may further include reacting such compound with an allylic reducing agent to produce the group




20

A non-limiting example of an allylic reducing agent is a palladium complex (e.g., Pd(PPh₃)₄) in combination with a formic acid salt (e.g., trialkylammonium formate).


Alternatively, preparation of the compound of formula (IVB) may include producing a compound of formula (IA) from a compound of formula (IIA), a compound of formula (IIIA), and R_5OH . The compound of formula (IA) is:

5



where all variables are as described for the compound of formula (IIA), the compound of formula (IIIA), and R_5OH .


- 10 The compound of formula (IA) may be reacted with the compound of formula (IVA) to form the compound of formula (IVB). For example, the compound of formula (IA) may be converted to the compound of formula (IB), which may then be converted to the compound of formula (IC). The compound of formula (IB) is:



- 15 where all variables are as described for the compound of formula (IA).

Preparation of the compound of formula (IB) may include reacting the compound of formula (IA) with an allylic reducing agent.

- 20 The compound of formula (IC) is:



where X_2 is as described for the compound of formula (IVB), and the remaining variables are as described for the compound of formula (IB).

The compound of formula (IVB) may be prepared from the compound of formula (IC) and the compound of
5 formula (IVA) as described herein.


Preparation of the compound of formula (IVB), in which X_1 is $-C(Y)_2-$, $-CH(Y)-$, or $-CH_2-$, may include a reaction (e.g., Claisen reaction) between an intermediate (e.g., the compound of formula (IVA), in which R_9 is $-CHO$) and another intermediate (e.g., the compound of formula (IIA), (IA), (IB), or (IC), in which R_1 is $-CH(Y)_2$ or $-CH_2(Y)$) that was treated with a Brønsted base to produce an intermediate (e.g., the compound of
10 formula (IVB), in which X_1 is $-C(Y)_2-$, $-CH(Y)-$, or $-CH_2-$). If X_1 in the compound of formula (IVB) is $-CH_2-$, the preparation of the compound of formula (IVB) may further include a desulfonylation or decarboxylation reaction.

Alternatively, preparation of the compound of formula (IVB), in which X_1 is $-O-$, may include a reaction (e.g., an esterification reaction) between an intermediate (e.g., the compound of formula (IVA), in which R_9 is $-COOH$) and another intermediate (e.g., the compound of formula (IIA), (IA), (IB), or (IC), in which R_1 that is $-OP_6$, where P_6 is H) to produce an intermediate (e.g., the compound of formula (IVB), in which X_1 is $-O-$).


Preparation of certain compounds of formula (IVB) or (IVC) may further involve conversion of the compound of formula (IVB) or (IVC), in which A_1 is H, and R_{14} and R_{15} combine to form a double bond, into the compound of formula (IVB) or (IVC), respectively, in which R_{14} and A_1 combine to form oxo. In a non-limiting example, the enone in the compound of formula (IVB) or (IVC), in which R_{14} and R_{15} combine to form a double bond can be converted into a C.12-C.13 epoxide using a nucleophilic peroxide agent, e.g., *t*-butyl
25 hydroperoxide, which can then be converted into the compound of formula (IVB) or (IVC), in which A_1 and R_{14} combine to form oxo, using methods known in the art, e.g., by reacting with a bidentate phosphine ligand and a source of Pd(0) (see, e.g., Muzart, J., *Eur. J. Org. Chem.*, 4717-4741, 2011). Thus, the compound of formula (IVB) or (IVC), in which A_1 is OP'' , can be prepared. Other transformations may involve α -oxygenation to produce the compound of formula (IVB) or (IVC), in which R_{15} is OP'' .

30

In general, the method described herein includes an allylic reduction at C.25. This reaction can be performed at any point prior to the formation of the halichondrin macrolide or analog thereof. For example,



may be converted to



using an allylic reducing agent.


35

C.19-C.20 Bond-forming Macrocyclization

In another approach, the macrocyclization reaction can be a carbon-carbon bond-forming reaction (e.g., olefin metathesis) that provides a C.19-C.20 bond in a halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)). A non-macrocyclic intermediate in the synthesis of the

halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)) may be a compound of formula (VB).

The compound of formula (VB) is:



(VB)

where

a designates (*R*)-stereogenic center, and *Z* is a sulfonate, chloride, bromide, or iodide; or *a* designates (*S*)-stereogenic center, and *Z* is OR₁₆, where R₁₆ is a hydroxyl protecting group;

X₁ is -CH₂-, -CH(Y)-, -C(Y)₂-, or -O-, where each Y is independently -COOR_C or -SO₂R_D;


each P₉ is independently a hydroxyl protecting group, and X₃ is oxo; or both P₉ groups and X₃, together with the atoms to which each is attached, combine to form a ketal;

Y₁ is iodide, bromide, or trifluoromethanesulfonate;

each R_C, when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D, when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

R_K is



where R₅ is optionally substituted acyl;

(a1) R₁₀ is a hydroxyl protecting group, R₁₁ is alkyl ether, and R₁₂ is H;

(a2) R₁₀ is a hydroxyl protecting group, and R₁₁ and R₁₂ combine to form a double bond;

or

(a3) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;

(b1) A₁ and R₁₄ combine to form oxo, R₁₃ is a hydroxyl protecting group, and R₁₅ is H;

or

(b2) A₁ is H or -OP'', and:

(i) R₁₃ is a hydroxyl protecting group, and R₁₄ and R₁₅ combine to form a double bond;

or


(ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or -OP'';

each P'' is independently a hydroxyl protecting group;

and the remaining variables are as described for the halichondrin macrolide or analog thereof.

The compound of formula (VB) is then converted to the compound of formula (VC) using reaction conditions known in the art. For example, the compound of formula (VB) may be subjected to Nozaki-Hiyama-Kishi
 5 reaction conditions known in the art to produce the compound of formula (VC). Nozaki-Hiyama-Kishi reaction on the compound of formula (VB) can include reacting the compound of formula (VB) with a Cr(II) salt and a Ni(II) salt. Ancillary ligands can be used in combination with the metal salts. In a non-limiting example, a substituted 1,10-phenanthroline can be used in combination with a Ni(II) salt. Chiral ancillary
 10 (dihydrooxazolyl-phenyl)-sulfonamides can be used with a Cr(II) salt to control the stereochemistry of the carbonyl carbon, to which a vinyl nucleophile is added in the course of Nozaki-Hiyama-Kishi reaction.


The compound of formula (VC) is:



(VC)

where

L₁ is



where L₂ and L₃ combine to form a bond;


20 *b* designates (*S*)-stereogenic center, if *a* designates (*R*)-stereogenic center;
 and

b designates (*R*)-stereogenic center, if *a* designates (*S*)-stereogenic center;

and the remaining variables are as described for the compound of formula (IVB).

25 The compound of formula (VC) is then converted to the halichondrin macrolide or analog thereof using methods described herein and those known in the art, e.g., those described in US 2016/0264594, US 2009/0203771, and WO 2016/179607.

The compound of formula (VC), in which Z is OR₁₆, R₁₆ is a hydroxyl protecting group ester, *a* designates
 30 (*S*)-stereogenic center, and *b* designates (*R*)-stereogenic center, can be converted to a compound of formula (VD):



where *a* designates (*S*)-stereogenic center, *b* designates (*R*)-stereogenic center, Z is OR₁₆, in which R₁₆ is a hydroxyl protecting group (e.g., R₁₆, together with the atom to which it is attached, combine to form an ester),
 5 R_H is sulfonyl, and the remaining variables are as described for the compound of formula (VC).

The compound of formula (VC), in which Z is OR₁₆, R₁₆ is a hydroxyl protecting group ester, *a* designates (*S*)-stereogenic center, and *b* designates (*R*)-stereogenic center, may be converted to the compound of formula (VD), e.g., by reacting with a sulfonyl electrophile, such as a sulfonyl chloride or a sulfonyl
 10 anhydride. In the compound of formula (VD), a hydroxyl protecting group in Z may be removed, resulting in the formation of the C.16-C.20 furan ring in the structure of the halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)). In a non-limiting example, the concomitant removal of a hydroxyl protecting group in Z (e.g., if Z is an ester) and the formation of the C.16-C.20 furan ring in the structure of the halichondrin macrolide or analog thereof may be achieved by the treatment of the compound
 15 of formula (VD) with a strong base (e.g., a C₁₋₆ alkoxide (e.g., KOMe)).

The compound of formula (VB), in which Z is a sulfonate, chloride, bromide, or iodide, *a* designates (*S*)-stereogenic center, and *b* designates (*R*)-stereogenic center, may be converted to the halichondrin macrolide or analog thereof (e.g., eribulin or a salt thereof (e.g., eribulin mesylate)) directly upon formation of
 20 the compound of formula (VC), e.g., upon the reaction work up and/or purification (e.g., on silica gel).

If the compound of formula (VD) includes hydroxyl protecting groups as R₃ and/or P₅, these hydroxyl protecting groups can be removed with a hydroxyl protecting group removing agent. For example, a hydroxyl protecting group removing agent may be a fluoride source, if the hydroxyl protecting group is a silyl
 25 group.

If, in the compound of formula (VC), each P₉ is H, and X₃ is oxo, the synthesis may further involve a reaction with a Brønsted acid (e.g., a Brønsted acid having a pK_a of 5 ± 3), e.g., after the formation of the C.16-C.20 furan ring.
 30

If, in the compound of formula (VC), (VD), or a C.16-C.20 furan cyclization product thereof, X₃ is oxo, R₁₃ is a hydroxyl protecting group, and R₁₄ and R₁₅ combine to form a double bond, treatment with a hydroxyl protecting group removing agent can provide the compound of formula (VC), (VD), or a C.16-C.20 furan cyclization product thereof, in which R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H.

Preparation of the halichondrin macrolides or analogs thereof may further involve conversion of the compound of formula (VC), (VD), or a C.16-C.20 furan cyclization product thereof, in which A₁ is H, and R₁₄ and R₁₅ combine to form a double bond, into the compound of formula (VC), (VD), or a C.16-C.20 furan cyclization product thereof, respectively, in which R₁₄ and A₁ combine to form oxo. In a non-limiting example, the enone in the compound of formula (VC), (VD), or a C.16-C.20 furan cyclization product thereof, in which X₃ is oxo, and R₁₄ and R₁₅ combine to form a double bond, can be converted into a C.12-C.13 epoxide using a nucleophilic peroxide agent, e.g., *t*-butyl hydroperoxide, which can then be converted into the compound of formula (VC), (VD), or a C.16-C.20 furan cyclization product thereof, in which A₁ and R₁₄ combine to form oxo, using methods known in the art, e.g., by reacting with a bidentate phosphine ligand and a source of Pd(0) (see, e.g., Muzart, J., *Eur. J. Org. Chem.*, 4717-4741, 2011). Thus, the compound of formula (VC), (VD), or a C.16-C.20 furan cyclization product thereof, in which A₁ is OP⁺, can be prepared. Other transformations may involve α-oxygenation to produce the compound of formula (VC), (VD), or a C.16-C.20 furan cyclization product thereof, in which R₁₅ is OP⁺.


15

In general, the fragments may be coupled in any order when forming compound (VB). For example, the C.0-C.1 or O-C.1 bond between fragment (IIA) and fragment (VA) can be formed, as described herein, prior to or after the Sakurai and/or Prins reaction.

20

In some embodiments, the compound of formula (VB) is produced from a compound of formula (IIA), a compound of formula (IIB), a compound of formula (IIIB), and, R₅OH, where R₅ is optionally substituted acyl.

The compound of formula (VA) is:



(VA)

25


where

R₉ is -CHO or -COOH;

and the remaining variables are as described for the compound of formula (VB).

30

The compound of formula (IIA) is:



(IIA)

where

R_1 is $-\text{OP}_6$, $-\text{CH}(\text{Y})_2$, or $-\text{CH}_2(\text{Y})$, where P_6 is H or a hydroxyl protecting group, and each Y is independently $-\text{COOR}_C$ or $-\text{SO}_2\text{R}_D$;

each R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

R_E is $-\text{CHO}$ or $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$,

where


m is 1, and R_F is a hydroxyl protecting group, or

m is 0, and (i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene;

and the remaining variables are as described for the compound of formula (VB).

The compound of formula (IIB) is:




where R_S is silyl.

The compound of formula (IIIB) is:




where

R_7 is $-\text{CHO}$ or




where each R_{7A} is independently an optionally substituted alkyl, and

R_{4B} is but-3-en-1-yl,




where each P_7 is independently H or a hydroxyl protecting group.

Preparation of the compound of formula (VB) includes a Sakurai reaction between R_E that is $-\text{CHO}$ and the compound of formula (IIB) producing a group of the structure:




. This product may then be reacted with the compound of formula (IIIB) under Prins reaction conditions to produce a group of the structure:




. This group may be converted to R_K as described herein.

- 5 If the stereochemistry of the secondary alcohol in the Sakurai reaction product differs from the desired stereochemistry, this group may be subjected to epimerization reaction conditions described herein (e.g., oxidation followed by Corey-Bakshi-Shibata reduction; alternatively, Mitsunobu reaction may be used to invert the stereogenic center).
- 10 If R_E is $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$, preparation of the compound of formula (VB) may further include conversion of R_E to $-\text{CHO}$ under acetal deprotecting conditions, e.g., through a reaction with an aqueous Brønsted acid (e.g., when m is 0), or through oxidation using an oxidizing agent capable of converting an alcohol to a carbonyl group (e.g., when m is 1).
- 15 When R_{4B} is but-3-en-1-yl, preparation of the compound of formula (VB) includes conversion of the but-3-en-1-yl group to $-(\text{CH}_2)_2-\text{CHO}$. For example, the but-3-en-1-yl group may be reacted with a dihydroxylating agent to produce R_{4B} that is




, and cleaving with a glycol cleaving agent.

- 20 When R_{4B} is




, preparation of the compound of formula (VB) includes conversion of




to $-(\text{CH}_2)_2-\text{CHO}$, e.g., by reacting with an oxidizing agent capable of converting an alcohol into a carbonyl group.

- 25 When R_{4B} is




, preparation of the compound of formula (VB) includes conversion of



to $-(\text{CH}_2)_2-\text{CHO}$, e.g., using a glycol cleaving agent.

If at least one P_7 is a hydroxyl protecting group in the compound of formula (IIIB), preparation of the compound of formula (VB) may further include treating with a hydroxyl protecting group removing agent.


- 5 Group R_E may be converted to R_K before or after the reaction of the compound of formula (IIA) with the compound of formula (VA). For example, conversion of R_E to R_K before the reaction of the compound of formula (IIA) with the compound of formula (VA) may be carried out via compounds of formulae (ID) and (IE), as described herein. Conversion of R_E to R_K after the reaction of the compound of formula (IIA) with the compound of formula (VA) may be carried out via a compound of formula (VAa):



(VAa)

where all variables are as described for the compound of formulae (IIA) and (VA).


The compound of formula (VAa) may be reacted with the compound of formula (IIB) under Sakurai reaction conditions to produce a compound of formula (VAb):



(VAb)

where all variables are as described for the compound of formula (VA).


- 20 A compound of formula (VAc) may be prepared from the compound of formula (VAb), the compound of formula (IIIB), and R_5OH through Prins reaction. The compound of formula (VAc) is:



(VAc)

where


R_{K1} is



and the remaining variables are as described for the compound of formula (VA) and the compound of formula (IIIB).

The compound of formula (VAc) may be converted to the compound of formula (VB) as described herein.


In particular embodiments, the Sakurai reaction product is a compound of formula (IIC):



(IIC)

where all variables are as described for the compound of formula (IIA).


In certain embodiments, the Prins reaction product is a compound of formula (ID):



(ID)

where all variables are as described for the compound of formula (IIA), the compound of formula (IIIB), and R_5OH .

In further embodiments, a compound of formula (IE) may be prepared from the compound of formula (ID) (e.g., the compound of formula (ID) is reacted with an allylic reducing agent to produce the compound of formula (IE)). The compound of formula (IE) is:




where all variables are as described for the compound of formula (ID).

The compound of formula (IE) may be reacted with the compound of formula (VA), as described herein, to produce the compound of formula (VB).

Preparation of the compound of formula (VB), in which X_1 is $-C(Y)_2-$, $-CH(Y)-$, or $-CH_2-$, may include a reaction (e.g., Claisen reaction) between an intermediate (e.g., the compound of formula (VA), in which R_9 is $-CHO$) and another intermediate (e.g., the compound of formula (IIA), (ID), (IIC), or (IE), in which R_1 is $-CH(Y)_2$ or $-CH_2(Y)$) that was treated with a Brønsted base to produce an intermediate (e.g., the compound of formula (VB), in which X_1 is $-C(Y)_2-$, $-CH(Y)-$, or $-CH_2-$). If X_1 in the compound of formula (VB) is $-CH_2-$, the preparation of the compound of formula (VB) may further include a desulfonylation or decarboxylation reaction.


Alternatively, preparation of the compound of formula (VB), in which X_1 is $-O-$, may include a reaction (e.g., an esterification reaction) between an intermediate (e.g., the compound of formula (VA), in which R_9 is $-COOH$) and another intermediate (e.g., the compound of formula (IIA), (ID), (IIC), or (IE), in which R_1 is $-OP_6$, where P_6 is H) to produce an intermediate (e.g., the compound of formula (VB), in which X_1 is $-O-$).

In general, the method described herein includes an allylic reduction at C.25. This reaction can be performed at any point prior to the formation of the halichondrin macrolide or analog thereof. For example,



C.23-C.24 Macrocyclization

In another approach, a halichondrin macrolide or analog thereof may be prepared from the compound of formula (IXB):




(IXB)

where X₁ is -CH₂-, -CH(Y)-, -C(Y)₂-, or -O-, each Y is independently -COOR_C or -SO₂R_D;

each R_C, when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D, when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

L₁ is



, where L₂ and L₃ combine to form a bond;

R₅ is optionally substituted acyl;

A₁ and R₁₄ combine to form oxo, R₁₃ is H or a hydroxyl protecting group, and R₁₅ is H;

or

A₁ is H or -OP^{''}, and

(i) R₁₃ is H or a hydroxyl protecting group, and R₁₄ and R₁₅, together with the atoms to which each is attached, combine to form a double bond;

or


(ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or -OP^{''}; each P₉ is independently H or a hydroxyl protecting group,

X₃ is oxo, or X₃, together with the atom to which it is attached, combines to form -(CH(OP₁₁))- , where P₁₁ is

H or a hydroxyl protecting group; or both P₉ and X₃, together with the atoms to which each is attached, combine to form a ketal; and

R_{19A} is H, -OP^{''}, or Y, and R_{19B} is H; or R_{19A} and R_{19B}, together with the atoms to which each is attached, combine to form a double bond.

The compound of formula (IXB) can be produced from a compound of formula (IXA), a compound of formula (IIB), a compound of formula (IIA), a compound of formula (IIIE), and R₅OH. The compound of formula (IXA) is:



(IXA)

where

R₉ is -CHO or -COOPⁿ;

- 5 (a1) R₁₀ is a hydroxyl protecting group, R₁₁ is alkyl ether, and R₁₂ is H;
 (a2) R₁₀ is a hydroxyl protecting group, and R₁₁ and R₁₂ combine to form a double bond; or
 (a3) R₁₀ and R₁₁ combine to form a bond, and R₁₂ is H;

A₁ and R₁₄ combine to form oxo, R₁₃ is H or a hydroxyl protecting group, and R₁₅ is H;

or

- 10 A₁ is H or -OPⁿ, and:

(i) R₁₃ is H or a hydroxyl protecting group, and R₁₄ and R₁₅ combine to form a double bond;

or

(ii) R₁₃ and R₁₄ combine to form a bond, and R₁₅ is H or -OPⁿ;


R₀ is -CHO, -CH₂OPⁿ, -CH=CH₂, -CH(OPⁿ)CH₂OPⁿ, -C(O)-CH₂P(O)(OR_C)₂, or halogen;

- 15 each R_C, when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each Pⁿ is independently H or a hydroxyl protecting group; and

each P₉ is independently a hydroxyl protecting group.

- 20 The compound of formula (IIA) is:



(IIA)

where

R₁ is -OP₆, -CH(Y)₂, or -CH₂(Y), where P₆ is H or a hydroxyl protecting group, and each Y is independently

- 25 -COOR_C or -SO₂R_D;

each R_C, when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl;

each R_D, when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl;

- 30 R_E is -CHO or -CH_(1+m)(OR_F)_(2-m), where

m is 1, and R_F is a hydroxyl protecting group,

or

m is 0, and


(i) each R_F is independently an alkyl or hydroxyl protecting group, or

(ii) both R_F combine to form an alkylene.

The compound of formula (IIIE) is:




where R_{4E} is



where each P_7 is independently H or a hydroxyl protecting group; and R_8 is $-\text{CH}_2\text{CH}_2-\text{COOR}_C$, $-\text{CH}=\text{CH}-\text{COOR}_C$, $-\text{CH}_2\text{CH}_2-\text{SO}_2\text{R}_D$, or $-\text{CH}=\text{CH}-\text{SO}_2\text{R}_D$; and

10 R_7 is $-\text{CHO}$ or




, where each R_{7A} is independently an optionally substituted alkyl.

The compound of formula (IIB) is:



where R_S is silyl.

In some embodiments, the compound of formula (IXB) is produced according the following strategy. For example, a compound of formula (IXC) may be produced from the compound of formula (IXA), the compound of formula (IIA), in which R_E is $-\text{CH}_{(1+m)}(\text{OR}_F)_{(2-m)}$, the compound of formula (IIB), and $R_5\text{OH}$, as follows. The compound of formula (IXA) may be reacted with the compound of formula (IIA) to produce a compound of formula (IXC):



25 where

X_1 is $-\text{CH}_2-$, $-\text{CH}(\text{Y})-$, $-\text{C}(\text{Y})_2-$, or $-\text{O}-$;

X₄ is oxo, or X₄, together with the atom to which it is attached, is $-(CH(OP_{12}))-$, where P₁₂ is H or a hydroxyl protecting group; and


the remaining variables are as described for the compound of formula (IXA) and the compound of formula (IIA).

5

Preparation of the compound of formula (IXC) may include an esterification reaction between R₁ that is $-OP_6$, where P₆ is H, and R₉ that is $-COOP^*$, where P^{*} is H. Alternatively, preparation of the compound of formula (IXC) may include a reaction between R₁ that is $-CH(Y)_2$ or $-CH_2(Y)$, and R₉ that is $-CHO$ (e.g., under Claisen reaction conditions). Further, the compound of formula (IXC), in which X₄, together with the atom to which it is attached, is $-(CH(OP_{12}))-$, where P₁₂ is H, may be oxidized to produce the compound of formula (IXC), in which X₄ is oxo. Reaction conditions useful for this oxidation are known in the art. Further desulfonylation or decarboxylation of this compound of formula (IXC) may produce the compound of formula (IXC), in which X₁ is $-CH_2-$.


10

15 The compound of formula (IXC) may be converted to a compound of formula (IXD):



where

R₇ is $-CHO$, $-CH_2OP_A$ or




the remaining variables are as described for the compound of formula (IXB).

The compound of formula (IXD) may be prepared from the compound of formula (IXC) and the compound of formula (IIIE) as follows. The compound of formula (IIIE) may be converted to a compound of formula (IIIF):

25




where R_{4F} is






R_{4E} that is




may be converted to R_{4F} that is




5 using a glycol cleaving agent.

R₈ that is $-\text{CH}=\text{CH}-\text{COOR}_\text{C}$ or $-\text{CH}=\text{CH}-\text{SO}_2\text{R}_\text{D}$ may be reacted with a 1,4-reducing agent to produce R₈ that is $-\text{CH}_2\text{CH}_2-\text{COOR}_\text{C}$ or $-\text{CH}_2\text{CH}_2-\text{SO}_2\text{R}_\text{D}$, respectively.


10 R_{4E} that is




may be reacted with an allylic reducing agent to produce



, which, upon reduction with a 1,2-reducing agent may produce R_{4F} that is




15 A compound of formula (IXE) may be produced from the compound of formula (IIIF) and the compound of formula (IXC). The compound of formula (IXE) is:



where all variables are as described for the compound of formula (IXC) and the compound of formula (IXD).


R_{4F} that is



, where R₈ is $-\text{CH}_2\text{CH}_2-\text{COOR}_c$ or $-\text{CH}_2\text{CH}_2-\text{SO}_2\text{R}_b$, may be reacted with R₀ that is $-\text{CHO}$ under Claisen reaction conditions. This reaction may produce the compound of formula (IXE), in which R_{19A} is Y.

5


R_{4F} that is



may be reacted with R₀ that is halogen under Nozaki-Hiyama-Kishi reaction conditions. This reaction may produce the compound of formula (IXE), in which X₃, together with the atom to which it is attached, combines to form $-(\text{CH}(\text{OP}_{11}))-$, where P₁₁ is H.

10

R_{4F} that is




may be reacted with R₀ that is $-\text{C}(\text{O})-\text{CH}_2\text{P}(\text{O})(\text{OR}_c)_2$ under Horner-Wadsworth-Emmons reaction conditions. This reaction may produce the compound of formula (IXE), in which R_{19A} and R_{19B}, together with the atoms to which each is attached, combine to form a double bond.

15

The compound of formula (IXE) may be converted to the compound of formula (IXD) by reacting with the compound of formula (IIB) under Sakurai reaction conditions.


The compound of formula (IXD), in which R₇ is $-\text{CHO}$, may be reacted with R₅OH under Prins reaction conditions to produce the compound of formula (IXB), in which L₁ is:

20



In further embodiments, the compound of formula (IXD) may be prepared by reacting the compound of formula (IXC), in which R_E is $-\text{CHO}$, with a compound of formula (IIB) to produce a compound of formula (IXF):

25




(IXF)

where P₁₃ is H or a hydroxyl protecting group; and the remaining variables are as described for the compound of formula (IXC).

5

The compound of formula (IXD) may be produced from the compound of formula (IXF) and the compound of formula (IIIF) as described herein. Preparation of the compound of formula (IXD) may further include converting P₁₃ that is a hydroxyl protecting group to P₁₃ that is H.

- 10 In yet further embodiments, the compound of formula (IXD) may be prepared as follows. A compound of formula (IXG) may be prepared from the compound of formula (IXA) and the compound of formula (IIIF) using methods described herein. The compound of formula (IXG) is:




(IXG)

- 15 where all variables are as described for the compound of formula (IXD) and the compound of formula (IXA).

The compound of formula (IXE) may be prepared from the compound of formula (IXG) and the compound of formula (IIA) using methods described herein. Preparation of the compound of formula (IXD) from the compound of formula (IXE) is then carried out as described herein.

20

The compound of formula (IXD) may be prepared from the compound of formula (IXG) and the compound of formula (IIC). For example, the compound of formula (IIC) may be protected with a hydroxyl protecting group to produce a compound of formula (IIJ):




where P_{13} is a hydroxyl protecting group, and the remaining variables are as described for the compound of formula (IIC).

5


The compound of formula (IXD) may be produced from the compound of formula (IIJ) and the compound of formula (IXG) using methods described herein.

In general, the methods described herein can include reacting R_E that is $-\text{CHO}$ with the compound of formula (IIB) and a first Lewis acid (e.g., an oxophilic Lewis acid) to produce a group of the structure:

10



, which may be reacted with R_7 , $R_5\text{OH}$, and a second Lewis acid (e.g., an oxophilic Lewis acid) (same as or different from the first Lewis acid) to produce a group of the structure:




15 In other embodiments, R_1 that is $-\text{OP}_6$ is reacted with R_9 under esterification reaction conditions to produce a group of the structure $-\text{X}_1-\text{C}(\text{O})-$, where P_6 is H, R_9 is $-\text{COOH}$, and X_1 is $-\text{O}-$. Alternatively, R_1 that is $-\text{CH}(\text{Y})_2$ or $-\text{CH}_2(\text{Y})$ is reacted with R_9 under Claisen reaction conditions to produce a group of the structure $-\text{X}_1-\text{C}(\text{O})-$, where R_9 is $-\text{CHO}$, and X_1 is $-\text{C}(\text{Y})_2-$ or $-\text{CH}(\text{Y})-$.


20 The compound of formula (IXB) may also be accessed using methods and compounds described in PCT/US17/40401.

In general, the method described herein includes an allylic reduction at C.25. This reaction can be performed at any point prior to the formation of the halichondrin macrolide or analog thereof. In certain
25 embodiments, the compound of formula (IXB) is reacted with an allylic reducing agent to produce the halichondrin or analog thereof.

In general, the method described herein includes an allylic reduction at C.25. This reaction can be performed at any point prior to the formation of the halichondrin macrolide or analog thereof. For example,



may be converted to




using an allylic reducing agent.

Compounds of Formula (Z) and (Z8)

Compounds of formula (Z) and (Z8) may be used as the compound of formula (III) in the syntheses described herein.

Compound of Formula (Z)

The compound of formula (Z) is:




where R_7 is $-\text{CHO}$ or $\text{R}_{7\text{A}}\text{O}-\text{CH}(\text{OR}_{7\text{A}})-\text{CH}_3$, and
each $\text{R}_{7\text{A}}$ is independently alkyl or a hydroxyl protecting group; or
both $\text{R}_{7\text{A}}$ combine to form an optionally substituted alkylene.

The compound of formula (Z) may be prepared as described herein. For example, a compound of formula

(Z2) may be produced from a compound of formula (Z1) and $\text{R}_{19}\text{O}-\text{P}(=\text{O})(\text{OR}_{19})_2-\text{CH}_2-\text{COOR}_{19}$, where each R_{19} is independently optionally substituted alkyl.

The compound of formula (Z1) is:



where


each $\text{R}_{20\text{A}}$ is independently a hydroxyl protecting group, or both $\text{R}_{20\text{A}}$, together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo;

each $\text{R}_{20\text{B}}$ is independently a hydroxyl protecting group, or both $\text{R}_{20\text{B}}$, together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo;


P_{12A} is H;


P_{12B} is H or a hydroxyl protecting group.


The compound of formula (Z2) is:





where all variables are as described for the compound of formula (Z1).

10 The compound of formula (Z1) may be reacted with  under Horner-Wadsworth-Emmons reaction conditions to produce the compound of formula (Z2).


A compound of formula (Z3) may be produced from the compound of formula (Z2) and , in which each R₂₁ is independently optionally substituted alkyl. The compound of formula (Z3) is:



where all variables are as described for the compound of formula (Z2) and for .

20 The compound of formula (Z2) may be reacted with  under Horner-Wadsworth-Emmons reaction conditions to produce the compound of formula (Z3).

The compound of formula (Z3) may be converted to the compound of formula (Z4):




where X_5 is a halogen, and

the remaining variables are as described for the compound of formula (Z3).

The free hydroxyl in the compound of formula (Z3) may be substituted with a halogen (e.g., iodide) as described herein. Reaction conditions for halogen substitution with a halogen are known in the art. In a non-limiting example, the compound of formula (Z3) may be reacted with a sulfonyl anhydride (e.g., trifluoromethanesulfonic anhydride or methanesulfonic anhydride) to produce the corresponding sulfonate (e.g., trifluoromethanesulfonate or methanesulfonate), which, upon reaction with a halide source (e.g., alkali iodide), may produce the compound of formula (Z4).


The compound of formula (Z4) may be converted to the compound of formula (Z5):



where all variables are as described for the compound of formula (Z4).

The compound of formula (Z4) may be subjected to reductive metal conditions to produce the compound of formula (Z5). Reductive metal conditions are known in the art. In a non-limiting example, the compound of formula (Z4) may be reacted with a metal capable of inserting into carbon-halogen bonds (e.g., Zn(0) or an alkyl lithium (e.g., *t*-BuLi) in combination with a Brønsted acid (e.g., acetic acid) to produce the compound of formula (Z5). When the metal is Zn(0), the Brønsted acid is an ingredient of the reaction mixture. When the reaction is performed with an alkyl lithium, the Brønsted acid is added to quench the reaction.


The compound of formula (Z5) may be reacted with R_{7A}OH to produce the compound of formula (Z6):



where R_{7A} is as described for the compound of formula (Z).

The compound of formula (Z5) may be reacted with R_{7A}OH in the presence of a Brønsted acid to produce the compound of formula (Z6).


A compound of formula (Z7) may be produced from the compound of formula (Z6) and in which each R₂₂ is independently optionally substituted alkyl. The compound of formula (Z7) is:




where all variables are as described for the compound of formula (Z) and for

Chemical structure of a phosphonate group: R₂₂O-P(=O)(OR₂₂)-CH₂-COOR₂₂.

- 5 The compound of formula (Z6) may be reacted with a glycol cleaving agent to produce aldehyde (Z6A):



where all variables are as described for the compound of formula (Z5).

- 10 The aldehyde (Z6A) can be reacted with , in which each R₂₂ is independently optionally substituted alkyl, under Horner-Wadsworth-Emmons reaction conditions to produce the compound of formula (Z7).

- The compound of formula (Z) may be produced from the compound of formula (Z7). In a non-limiting example, the compound of formula (Z7) may be reacted with a 1,4-reducing agent to produce the compound of formula (Z).
- 15


Compound of Formula (Z8)

The compound of formula (Z8) is:



where

R₇ is -CHO or



- 25 each R_{7A} is independently alkyl or a hydroxyl protecting group; or both R_{7A} combine to form an optionally substituted alkylene;

R_8 is $-\text{CH}_2\text{CH}_2-\text{COOR}_C$, $-\text{CH}=\text{CH}-\text{COOR}_C$, $-\text{CH}_2\text{CH}_2-\text{SO}_2\text{R}_D$, or $-\text{CH}=\text{CH}-\text{SO}_2\text{R}_D$;

R_C , when present, is independently optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and


R_D , when present, is independently optionally substituted aryl or optionally substituted non-enolizable alkyl.

5

The compound of formula (Z8) may be prepared from the compound of formula (Z5) as follows.

The compound of formula (Z5) may be reacted with $R_{7A}OH$ in the presence of a Brønsted acid to produce the compound of formula (Z5A):

10




where

R_{20A} is H or a hydroxyl protecting group;

each R_{20B} is independently H or a hydroxyl protecting group, or both R_{20B} , together with the atoms to which each is attached, combine to form an acetal, ketal, cyclic carbonate, dicarbonyl-dioxo, or silylene-dioxo; and all remaining variables are as described for the compound of formula (Z8).

The compound of formula (Z5A) may be converted to a compound of formula (Z5B):

20



where

R_{23} is H or $-\text{OR}_{20A}$; and


all remaining variables are as described for the compound of formula (Z8). The compound of formula (Z5A), in which both R_{20B} are H, can be reacted with a glycol cleaving agent to produce the compound of formula (Z5B). Conversion of $-\text{OR}_{20A}$ in formula (Z5A) to H in the compound of formula (Z5B) (R_{23} is H) can be achieved using allylic deoxygenation methods known in the art, e.g., using allylic reducing agents, when R_{20A} is a hydroxyl protecting group that is an ester. Alternatively, when R_{20A} is H, $-\text{OR}_{20A}$ may be converted to H using deoxygenation reactions known in the art. Deoxygenation reactions are known in the art, e.g., Barton-McCombie deoxygenation and tin-free versions thereof (see, e.g., Cheneberg and Ollivier, *Chimia*, 70:67-76, 2016). The reaction replacing $-\text{OR}_{20A}$ with H can be performed before or after the glycol cleavage.

The compound of formula (Z5B) may be converted to the compound of formula (Z8), in which R₈ is –

CH=CH–COOR_C or –CH=CH–SO₂R_D, by a reaction with $\text{R}_{22}\text{O}-\text{P}(=\text{O})(\text{OR}_{22})-\text{COOR}_{\text{C}}$ or $\text{R}_{22}\text{O}-\text{P}(=\text{O})(\text{OR}_{22})-\text{SO}_2\text{R}_{\text{D}}$, in which each R₂₂ is independently optionally substituted alkyl, under Horner-Wadsworth-Emmons reaction conditions. The compound of formula (Z8), in which R₈ is –CH=CH–COOR_C or –CH=CH–SO₂R_D, can be converted to the compound of formula (Z8), in which R₈ is –CH₂CH₂–COOR_C or –CH₂CH₂–SO₂R_D, respectively, by a reaction with a 1,4-reducing agent. A reaction replacing –OR_{20A} with H can be performed before or after the Horner-Wadsworth-Emmons reaction and/or the reaction with a 1,4-reducing agent.


Further Compounds

The invention provides a compound of formula (II):



where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, a hydroxyl protecting group, and A is a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, or A is a group of formula (1):



where

L is –(CH(OP₂))–, –(C(OH)(OP₂))–, or –C(O)–;


R₂ is H and P₁ is absent, H, alkyl, or a hydroxyl protecting group, or R₂ and P₁ combine to form a bond;

(i) R₃ is H, and P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group;

(ii) R₃ is –(CH₂)_nNP₃P₄, where P₃ is H or an N-protecting group, and (a) P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₄ is H or an N-protecting group, (b) P₂ and P₄ combine to form an alkylidene, or (c) each of P₂ and P₄ is H;

(iii) R₃ is –(CH₂)_nOP₅, where P₂ is absent, H, optionally substituted alkyl, or a hydroxyl protecting group, and P₅ is H, optionally substituted alkyl, or a hydroxyl protecting group; or P₂ and P₅, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or

(iv) R_3 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is H, optionally substituted alkyl, or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,

$NR_B(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

n , when present, is 0, 1, or 2;

k is 0 or 1;

R_1 is $-OP_6$, $-CH(Y)_2$, or $-CH_2(Y)$, where P_6 is H or a hydroxyl protecting group;

R_6 is H, optionally substituted alkyl, or optionally substituted aryl; and

Y is independently $-COOR_C$ or $-SO_2R_D$;

R_C , when present, is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl; and

R_D , when present, is optionally substituted aryl or optionally substituted non-enolizable alkyl.

The invention also provides the compounds of formula (IID), (IIE), (IIF), (IIG), (IIH), (III), (IVAb), (VAb), (VIIAa), (VIAa), (VIID), (IXD), (IXF), (IXG), (Z), (Z1), (Z2), (Z3), (Z4), (Z5), (Z5A), (Z5B), or (Z6).

25 Amination

Compounds disclosed herein, in which R_3 is $-(CH_2)_nOP_5$, can be aminated to afford eribulin, as described herein. Amination conditions can be those known in the art. In a non-limiting example, C.35 hydroxyl in the compounds disclosed herein can be sulfonylated (e.g., by a reaction with a sulfonyl anhydride or a sulfonyl

chloride) and reacted with a nitrogen source (e.g., ammonia, azide, sulfamic acid, urea (H_2NCONH_2), or thiourea (H_2NCSNH_2)) to afford eribulin or a pharmaceutically acceptable salt thereof upon optional unmasking of the amino group (if the nitrogen source was azide, urea, or thiourea). In another non-limiting example, C.35 hydroxyl in the compounds disclosed herein can be halogenated (e.g., by Appel reaction or a reaction with thionyl chloride, sulfuryl chloride, phosphorus(III) chloride, or phosphorus(V) oxychloride) and reacted with a nitrogen source (e.g., ammonia, azide, sulfamic acid, a phthalimide salt, urea (H_2NCONH_2), or thiourea (H_2NCSNH_2)) to afford eribulin or a pharmaceutically acceptable salt thereof upon optional unmasking of the amino group (if the nitrogen source was azide, a phthalimide salt, urea, or thiourea).

Amino unmasking agents are further described herein. The amination reaction can provide a pharmaceutically acceptable salt of eribulin directly. Alternatively, the amination reaction can provide eribulin in a free base form. A pharmaceutically acceptable salt of eribulin can be prepared from eribulin through a salification reaction as described herein.

Masked Amines and Amine Unmasking Agents

The compounds used in the methods of the invention can contain a masked or unmasked amine (e.g., at C.35 carbon of the structure of the halichondrin macrolide analog, such as eribulin). An unmasked amine is $-\text{NH}_2$. An amine can be masked using methods known in the art, e.g., by protecting the amine with an *N*-protecting group. Alternatively, an amine can be masked as a nitrogen-containing moiety, which can be reacted with an amine unmasking agent to afford an amine. Non-limiting examples of the nitrogen-containing moieties include azide and an imide (e.g., phthalimide). Amine unmasking agents can be those known in the art for removing *N*-protecting groups from amines. In a non-limiting example, a Boc group can be removed using amine unmasking agents known in the art, e.g., a Brønsted acid (e.g., HCl in 1,4-dioxane or trifluoroacetic acid). When amine is masked as azide, the amine can be unmasked by subjecting the compound containing the masked amine to Staudinger reaction conditions (e.g., by contacting with a phosphine, such as trialkylphosphine, dialkylarylphosphine, alkylidiarylphosphine, or triarylphosphine) or by reacting the compound containing the masked amine with a reducing agent (e.g., LiAlH_4). When amine is masked as an imide (e.g., phthalimide), the amine can be unmasked by reacting with an amine unmasking agent known in the art, e.g., hydrazine.

Oxidizing Agents Capable of Converting an Alcohol to a Carbonyl Group

Oxidizing agents capable of converting an alcohol to a carbonyl group are known in the art. Non-limiting examples of these oxidizing agents include Dess-Martin periodinane, TEMPO (in the presence of bleach or BAIB), a dimethylsulfonium compound (e.g., dimethylchlorosulfonium chloride), aluminum trialkoxide with an excess of a ketone (e.g., acetone), and catalytic tetrapropylammonium perruthenate (TPAP) (in the presence of *N*-methylmorpholine oxide). The dimethylsulfonium compound can be prepared *in situ* under the conditions known for Parikh-Doering oxidation, Swern oxidation, Corey-Kim oxidation, or Pfitzner-Moffatt oxidation. Alternatively, the dimethylsulfonium compound can be prepared *in situ* by a reaction between trichloroacetic acid anhydride and dimethyl sulfoxide. An oxidation reaction of an alcohol to a carbonyl group (e.g., a ketone) can be performed using aluminum trialkoxide and an excess of a ketone (e.g., acetone) under the conditions known in the art for Oppenauer oxidation. Allylic and benzylic alcohols can also be oxidized with MnO_2 .

Reducing Agents

Reducing agents that can be used in the methods of the invention are those known in the art. A reducing agent can be an electron-transfer reducing agent, a metal hydride, or a metalloid hydride. Non-limiting examples of electron-transfer reducing agent include alkali metals in oxidation state (0), alkali earth metals in oxidation state (0), alkali arenides, lanthanide (II) salts (e.g., SmI_2), $\text{Zn}(0)$, $\text{Fe}(0)$, and $\text{Mn}(0)$. Non-limiting examples of metal hydrides and metalloid hydrides include boron hydride compounds (e.g., NaBH_4 , LiBH_4 , LiHBEt_3 , selectrides (e.g., L-selectride), and boranes (e.g., 9-BBN and alpine borane)), aluminum hydride compounds (e.g., LiAlH_4 , Red-Al®, and alanes (e.g., diisobutylaluminum hydride (DIBAL))), hydrosilanes (e.g., PMHS and Ph_2SiH_2), hydrostannanes (e.g., Bu_3SnH), copper hydride complexes (e.g., Stryker's reagent), palladium hydride complexes, platinum hydride complexes, iridium hydride complexes, rhodium hydride complexes, and ruthenium hydride complexes. Reducing agents can be formed *in situ*, e.g., a copper hydride complex can be formed by a reaction of a copper salt with, e.g., a boron hydride compound or a hydrosilane. Thus, some reducing reagents (e.g., boron hydride compounds, hydrosilanes, and hydrostannanes) can be used in combination with a catalytic quantity of a metal salt (e.g., Cu, Pd, Pt, Ir, Rh, or Ru salt). Alternatively, catalytic reducing agents can be metal salts (e.g., aluminum isopropoxide or a ruthenium complex) in combination with an alcohol, which undergo transfer hydrogenation of carbonyl-containing compounds without intermediacy of a metal hydride. Non-limiting examples of transfer hydrogenation reactions include Meerwein-Ponndorf-Verley reduction (e.g., using aluminum isopropoxide/isopropanol) and Ru-catalyzed transfer hydrogenation (e.g., Hashiguchi et al., *J. Am. Chem. Soc.*, 117:7562-7563, 1995).

When a substrate is an α,β -unsaturated carbonyl or sulfone compound (e.g., an α,β -enone or a vinyl sulfone), a reducing agent can be a 1,2-reducing agent or a 1,4-reducing agent. For example, a reaction between an α,β -unsaturated carbonyl compound and a 1,2-reducing agent can afford, e.g., an allylic alcohol (or an allylic amine, if the starting compound is an enamide), whereas a reaction between an α,β -unsaturated carbonyl compound and a 1,4-reducing agent can afford an α,β -saturated compound and can leave the carbonyl group intact after work up of the reaction mixture. Non-limiting examples of 1,2-reducing agents include metal hydrides and metalloid hydrides, e.g., aluminum hydride compounds, boron hydride compounds (e.g., CeCl_3 with NaBH_4), and ruthenium hydride complexes. Non-limiting examples of 1,4-reducing agents include boron hydride compounds (e.g., LiHBEt_3 and L-selectride), hydrostannanes, copper hydride complexes (e.g., Stryker's reagent), palladium hydride complexes, platinum hydride complexes, iridium hydride complexes, rhodium hydride complexes, and ruthenium hydride complexes. Non-limiting examples of the 1,4-reducing agents include copper (I) hydrides, which can be isolated (e.g., Stryker's reagent) or prepared *in situ* (e.g., from a copper (I) or copper (II) salt and a hydride source). Catalytic quantities of a copper salt (either copper (I) or copper (II) salt) in combination with stoichiometric or superstoichiometric quantities of a hydride source (e.g., a borohydride salt, borane, PMHS, or a hydrosilane (e.g., Ph_2SiH_2)). A non-limiting example of the reaction conditions that can be used for 1,4-reduction is described, e.g., in Baker et al., *Org. Lett.*, 10:289-292, 2008, the disclosure of which is incorporated herein by reference. Other metals can be used to catalyze 1,4-reduction, e.g., Ru, Pd, and Ir compounds.

40

Methods of the invention may include the use of selective reduction techniques to reduce some reactive groups with retention of other reactive groups. For example, a carboxylic acid may be reduced in the

presence of esters and/or olefins using sodium borohydride and iodine. Alternatively, a carboxylic acid may be reduced in the presence of esters and/or 1,1-disubstituted olefins by converting the carboxylic acid to a mixed anhydride (e.g., using N-methylmorpholine and *t*-butylchloroformate) and reducing the resulting mixed anhydride using borane reducing agents (e.g., 9-BBN).

5

A compound having an allylic leaving group (e.g., a carboxylate, a halide, or a sulfonate) can be treated with an allylic reducing agent to replace the leaving group with a hydrogen atom. A non-limiting example of allylic reducing agent is a palladium salt or complex (e.g., Pd(PPh₃)₄) in combination with a formic acid salt (e.g., trialkylammonium formate).

10

Hydroxyl Protecting Groups and Hydroxyl Protecting Group Removing Agents

Hydroxyl protecting groups can be as defined herein. In particular, a hydroxyl protecting group can be an acyl, a sulfonyl, an arylalkyl (e.g., benzyl or *p*-methoxybenzyl), an aryl (e.g., *p*-methoxyphenyl), or an optionally substituted silyl (e.g., TMS, TES, TBS, TIPS, TBDPS, or TPS). Hydroxyl protecting groups, hydroxyl protecting agents, and hydroxyl protecting reaction conditions can be selected to protect selectively certain hydroxyl groups in a compound, while leaving other hydroxyl groups unprotected. The choice of hydroxyl protecting groups for a compound can facilitate subsequent deprotection strategies, as some hydroxyl protecting groups can be removed in the presence of others using appropriate hydroxyl protecting group removing agents. Some of these strategies involving the choice of silyl hydroxyl protecting groups are discussed in, e.g., *Silicon-Based Blocking Agents*, Gelest, Inc., 2011.

20

Hydroxyl protecting group removing agents are those agents that can react with a compound having a protected hydroxyl group to afford the compound with a deprotected hydroxyl group. Hydroxyl protecting group removing agents and deprotection reaction conditions can be those known in the art. In a non-limiting example, hydroxyl masked as silyl ether can be unmasked by a reaction with a fluoride source (e.g., a fluoride salt, such as KF or TBAF). Alternatively, hydroxyl protected as TMS or TES ether can be deprotected by a reaction with a Brønsted acid (e.g., a carboxylic acid). In another non-limiting example, hydroxyl protected as an ester can be deprotected by a reaction with a base (e.g., alkali hydroxide (e.g., lithium hydroxide, sodium hydroxide, or potassium hydroxide) or C₁₋₆ alkoxide (e.g., alkali C₁₋₆ alkoxide or alkali earth C₁₋₆ alkoxide)). Alternatively, hydroxyl protected as an ester (e.g., pivaloyl ester) can be deprotected by a reaction with a 1,2-reducing agent (e.g., DIBAL-H). In yet another non-limiting example, hydroxyl protected as an arylalkyl ether (e.g., 1-arylalk-1-yl ether) can be deprotected using a reduction reaction, e.g., with Pd/C and H₂ or with Na/NH₃. Alternatively, hydroxyl protected as an alkoxy-arylalkyl ether (e.g., MPM ether) can be deprotected by a reaction with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ). In still another non-limiting example, hydroxyl protected as alkoxyalkyl ether (e.g., 1-alkoxyalk-1-yl) or THP ether can be deprotected by a reaction with a Brønsted acid. Cyclic protected diols, such as acetals or ketals (e.g., as 2-alkyl-1,3-dioxolane, 2,2-dialkyl-1,3-dioxolane, 2-alkyl-1,3-dioxane, or 2,2-dialkyl-1,3-dioxane), can be deprotected by a reaction with a Brønsted acid (e.g., a carboxylic acid).

35

Decarboxylation and Desulfonation

The conditions for the decarboxylation reaction can be those known in the art, e.g., Krapcho decarboxylation or a sequence including deprotection, if R_C is not H, by converting R_C to H and subsequent

protodecarboxylation. The conditions for the desulfonylation reaction can be those known in the art. For example, the desulfonylation reaction can include contacting the compound of formula (IA) or formula (IB) or an intermediate downstream of the compound of formula (IA) or formula (IB) with an electron-transferring reducing agent (e.g., SmI_2 ; Cr(III) salt and Mn(0) ; or Mg(0)). For exemplary desulfonylation conditions, see

5 WO 2009/064029.

Nozaki-Hiyama-Kishi Reaction

Nozaki-Hiyama-Kishi reaction conditions that may be used in transformation described herein can be those known in the art. Nozaki-Hiyama-Kishi reaction can include reacting substrates (an aldehyde and a vinyl

10 halide or pseudohalide) with a Cr(II) salt and a Ni(II) salt. Ancillary ligands can be used in combination with the metal salts. In a non-limiting example, a substituted 1,10-phenanthroline can be used in combination with a Ni(II) salt. Chiral ancillary ligands can be used to render the reaction stereoselective. In a non-limiting example, chiral *N*-(dihydrooxazolyl-phenyl)-sulfonamides can be used with a Cr(II) salt to control the stereochemistry of the carbonyl carbon, to which a vinyl nucleophile is added in the course of Nozaki-

15 Hiyama-Kishi reaction.

Olefin Metathesis

Olefin metathesis catalysts are known in the art and include Ru-carbene complexes (e.g., Grubbs and Hoveyda-Grubbs catalysts). Olefin metathesis-competent catalysts that may be used in the olefin

20 metathesis reactions described herein are known in the art (e.g., second generation Hoveyda-Grubbs-type catalysts, e.g., those in which the Ru-benzylidene moiety is modified to include electron-withdrawing and/or electron-donating groups). Non-limiting examples of the useful olefin metathesis reaction conditions are provided in, e.g., U.S. patent application publication No. 2016/0264594 and International patent application publication No. WO 2016/179607.

Salification

Eribulin mesylate can be produced by salification of eribulin, as describen herein. Salification reaction conditions are known in the art. Salification of eribulin can afford a pharmaceutically acceptable salt of eribulin (e.g., eribulin mesylate). In particular, salification reaction can involve contacting eribulin with a

30 Brønsted acid (e.g., a pharmaceutically acceptable Brønsted acid (e.g., methanesulfonic acid)) to afford a pharmaceutically acceptable salt of eribulin (e.g., *Handbook of Pharmaceutical Salts: Properties, Selection and Use*, ed.: Stahl and Wermuth, Wiley-VCH/VHCA, Weinheim/Zurich, 2002). Pharmaceutically acceptable salts of eribulin, e.g., eribulin mesylate, can be formed by methods known in the art, e.g., in situ during the final isolation and purification of the compound or separately by reacting the free base group with

35 a suitable organic acid. In one example, eribulin is treated with a solution of MsOH and NH_4OH in water and acetonitrile. The mixture is concentrated. The residue is dissolved in DCM -pentane, and the solution is added to anhydrous pentane. The resulting precipitate is filtered and dried under high vacuum to provide eribulin mesylate.


Epimerizations

Epimerization reactions can be used to invert a stereogenic center having an undesired stereochemical identity. For example, through epimerization, *R* stereogenic center can be converted to *S* stereogenic center

- and *vice versa*. Epimerization of a stereogenic sp^3 -carbon bonded to one hydrogen atom and to one hydroxyl group can be achieved through a reaction sequence involving oxidation of the hydroxyl group to a carbonyl group followed by a 1,2-reduction reaction. The 1,2-reduction reaction can provide the desired stereochemical identity diastereoselectively, or the reaction can be carried out using a chiral catalyst, chiral auxiliary, or a chiral reducing agent. Non-limiting examples of chiral reducing agents include alpine borane and propine borane. Non-limiting examples of 1,2-reduction reactions involving chiral catalysts are Corey-Bakshi-Shibata reduction, Noyori hydrogenation, and Noyori transfer hydrogenation. The oxidation/reduction reaction sequence can be carried out *in situ* using dynamic kinetic resolution. A dynamic kinetic resolution can further involve a reaction with a hydroxyl protecting agent, which removes the desired stereoisomer from the reduction/oxidation equilibrium. In a non-limiting example, a dynamic kinetic resolution of chiral secondary alcohols can involve reduction/oxidation equilibration using $\eta^5\text{-Ph}_5\text{CpRu(CO)}_2\text{H}$ in combination with enantioselective esterification using isopropenyl acetate catalyzed by a lipase enzyme (e.g., lipase B from *Candida Antarctica*, see, e.g., Martin-Matute et al., *J. Am. Chem. Soc.*, 127:8817-8825, 2005).
- The following examples are meant to illustrate the invention. They are not meant to limit the invention in any way.

Examples

Example 1



5 An exemplary compound of formula (IA) can be prepared as shown in the above scheme. This compound may be useful as an intermediate in the synthesis of a halichondrin macrolide. Compound 1 was treated with compound 2 in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ (exemplary Sakurai reaction conditions) to give compound 3.


10 Compound 5 can be prepared by treating compound 3 with compound 4 in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ and methoxyacetic acid (exemplary Prins reaction conditions). Compound 7 can be prepared by treating compound 3 with compound 6 in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ and methoxyacetic acid (exemplary Prins reaction conditions).

A halichondrin macrolide can be prepared from compound 5 or 7 using methods and intermediates disclosed, e.g., WO 2016/179607, the synthesis of which is hereby incorporated by reference and described


herein. For example, compounds **5** and **7** may be converted to a halichondrin macrolide, as described for the compound of formula (VIII C) in WO 2016/179607.


Example 2

- 5 An exemplary compound of formula (IA) for the synthesis of a halichondrin macrolide analog can be prepared as shown in this Example. Compound **8** was treated with compound **2** in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ (exemplary Sakurai reaction conditions) to give compound **9**.




- The scheme above illustrates two exemplary synthetic routes that can be used to access compound **11** from compound **9**. In one approach, compound **9** was reacted with Dess-Martin periodinane to afford enone **10**, which, upon reduction with either $(S)\text{-}(-)\text{-}2\text{-methyl-CBS-oxazaborolidine}$ and $\text{BH}_3 \cdot \text{THF}$ or $\text{Zn}(\text{BH}_4)_2$, provided compound **11** as the major diastereomer. $(S)\text{-}(-)\text{-}2\text{-methyl-CBS-oxazaborolidine}$ is:






Trimethyl(2-methylbuta-2,3-dien-1-yl)silane. Lithium chloride (4.0 g, 94 mmol) was charged a round bottom flask and dried with heating under vacuum. After cooling to room temperature, the flask was charged with diethyl ether (100 mL) and copper(I) cyanide (4.0 g, 45 mmol). The mixture was cooled to 0 °C and treated with 1 M TMSCH₂MgCl (45.0 mL, 45.0 mmol) over 10 min maintaining the internal temperature below 5 °C. The mixture was stirred at 0 °C for 1 h. After cooling to -78 °C, the mixture was treated with 2-butyne-1-yl p-toluenesulfonate (10.0 g, 44.5 mmol) in 3 portions. The mixture was slowly warmed up to room temperature over 18 h. The reaction mixture was filtered through a celite pad and rinsed with diethyl ether. The filtrate was concentrated with slight vacuum (bath: 10 °C) to give the title compound (4.09 g, 65.3%).

¹H NMR (400 MHz, CHLOROFORM-d) δ 4.51-4.58 (m, 2H), 1.70 (t, *J*=3.13 Hz, 3H), 1.33 (t, *J*=2.54 Hz, 2H), 0.06 (s, 9H).



(*S*)-3-((2*R*,3*R*,4*S*,5*S*)-5-((*S*)-2-hydroxy-4-methyl-3-methylenepent-4-en-1-yl)-3-methoxy-4-((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate. A mixture of (*S*)-3-((2*R*,3*R*,4*S*,5*S*)-3-methoxy-5-(2-oxoethyl)-4-((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate (1.61 g, 2.77 mmol) and trimethyl(2-methylbuta-2,3-dien-1-yl)silane (0.78 g, 5.5 mmol) in CH₂Cl₂ (32 mL) was cooled to -78 °C, treated with BF₃·OEt₂ (0.70 mL, 5.5 mmol) and stirred at -78 °C for 1 h. The reaction was quenched with sat. aq. NaHCO₃ (80 mL) and the mixture was extracted twice with MTBE (30 mL). The organic layers were combined, washed with brine (32.1 mL), and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate in *n*-heptane=10% to 50%) to give the title compound (960 mg, 53.5%) and a 1:1 mixture of the title compound and its epimer (631 mg, 35%).

¹H NMR (400 MHz, CHLOROFORM-d) δ 8.04-8.10 (m, 2H), 7.97-8.04 (m, 2H), 7.88-7.96 (m, 2H), 7.64-7.72 (m, 1H), 7.50-7.62 (m, 4H), 7.37-7.48 (m, 4H), 5.54-5.70 (m, 1H), 5.27 (s, 1H), 5.17 (s, 1H), 4.99 (s, 1H), 4.95 (s, 1H), 4.63 (br d, *J*=8.60 Hz, 1H), 4.57 (d, *J*=5.08 Hz, 2H), 3.87-3.97 (m, 2H), 3.76-3.85 (m, 1H), 3.41 (s, 3H), 3.01-3.19 (m, 2H), 2.52-2.63 (m, 1H), 2.19-2.35 (m, 2H), 1.93-2.03 (m, 1H), 1.88 (s, 3H), 1.68-1.79 (m, 1H).



(S)-3-((2R,3R,4S,5S)-3-methoxy-5-(4-methyl-3-methylene-2-oxopent-4-en-1-yl)-4-


((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate. A solution of (S)-3-

((2R,3R,4S,5S)-5-((S)-2-hydroxy-4-methyl-3-methylenepent-4-en-1-yl)-3-methoxy-4-

5 ((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate (0.63 g, 0.97 mmol) in CH_2Cl_2 (12.6 mL) was treated with sodium bicarbonate (0.16 g, 1.9 mmol) and Dess-Martin periodinane (0.495 g, 1.17 mmol). After stirring at room temperature for 1 h, the reaction was quenched with sat. NaHCO_3 (6.3 mL) and 20% (w/v) aq. Na_2SO_3 (6.3 mL), and extracted twice with MTBE (12.6 mL). The organic layers were combined, washed with brine (6.3 mL), and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate in *n*-heptane= 10% to 40%) to give the title compound (378 mg, 60%).

^1H NMR (400 MHz, CHLOROFORM-d) δ 7.98-8.11 (m, 4H), 7.89-7.97 (m, 2H), 7.62-7.72 (m, 1H), 7.50-7.62 (m, 4H), 7.35-7.49 (m, 4H), 5.54-5.69 (m, 3H), 5.08 (s, 1H), 4.97 (s, 1H), 4.57 (d, $J=5.08$ Hz, 2H), 3.97-4.05 (m, 2H), 3.90-3.96 (m, 1H), 3.49 (dd, $J=3.91, 14.07$ Hz, 1H), 3.42 (s, 3H), 3.05-3.24 (m, 2H), 2.98 (dd, $J=7.03, 17.58$ Hz, 1H), 2.47-2.58 (m, 1H), 2.18-2.35 (m, 2H), 1.82-1.91 (s, 3H).

15 $J=7.03, 17.58$ Hz, 1H), 2.47-2.58 (m, 1H), 2.18-2.35 (m, 2H), 1.82-1.91 (s, 3H).



(S)-3-((2R,3R,4S,5S)-5-((R)-2-hydroxy-4-methyl-3-methylenepent-4-en-1-yl)-3-methoxy-4-

((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate. (S)-CBS oxazaborolidine


20 (0.056 g, 0.20 mmol) was dissolved in THF (5.8 mL) and treated with 1 M $\text{BH}_3\cdot\text{THF}$ in THF (0.90 mL, 0.90 mmol). The mixture was stirred at room temperature for 1 h. After cooling to -40°C , the mixture was treated with a solution of (S)-3-((2R,3R,4S,5S)-3-methoxy-5-(4-methyl-3-methylene-2-oxopent-4-en-1-yl)-4-((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate (0.29 g, 0.45 mmol) in THF (5.8 mL) and stirred at a temperature between -30 and -15°C for 3 h. The reaction was quenched with methanol

25 (0.18 mL) and sat. aq. NH_4Cl (15 mL). The mixture was extracted twice with MTBE (29 mL), washed with brine, and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate in *n*-heptane= 10% to 50%) to give the title compound (120 mg, 41%).

¹H NMR (400 MHz, CHLOROFORM-d) δ 7.95-8.09 (m, 4H), 7.86-7.93 (m, 2H), 7.62-7.72 (m, 1H), 7.49-7.60 (m, 4H), 7.35-7.45 (m, 4H), 5.49-5.59 (m, 1H), 5.30-5.37 (m, 1H), 5.15-5.21 (m, 1H), 4.99 (s, 1H), 4.96 (s, 1H), 4.58-4.65 (m, 1H), 4.49-4.57 (m, 2H), 3.83-3.97 (m, 2H), 3.77-3.82 (m, 1H), 3.36 (s, 3H), 3.03-3.15 (m, 2H), 2.60-2.70 (m, 1H), 2.26 (t, $J=6.44$ Hz, 2H), 2.03-2.11 (m, 1H), 1.89 (s, 3H), 1.71-1.83 (m, 1H).

5

In another approach, compound **9** was subjected to Mitsunobu reaction with 3,5-dinitrobenzoic acid, DIAD, and PPh₃ to give compound **13**. Alcoholysis of compound **13** with Mg(OMe)₂ gave compound **11**.



(S)-3-((2R,3R,4S,5S)-5-((R)-2-((3,5-dinitrobenzoyl)oxy)-4-methyl-3-methylenepent-4-en-1-yl)-3-methoxy-4-


10 methoxy-4-((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate. A solution of (S)-3-((2R,3R,4S,5S)-5-((S)-2-hydroxy-4-methyl-3-methylenepent-4-en-1-yl)-3-methoxy-4-

((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate (0.33 g, 0.51 mmol) in THF (13.2 mL) was treated with 3,5-dinitrobenzoic acid (0.65 g, 3.1 mmol) and triphenylphosphine (0.80 g, 3.1 mmol).

The mixture was treated with DIAD (0.49 mL, 2.5 mmol) and stirred at room temperature for 20 h. The

15 reaction was quenched with sat. aq. NaHCO₃ (6.6 mL) and extracted twice with MTBE (9.9 mL). The organic layers were combined, washed with brine, and concentrated *in vacuo*. The residue was purified by column chromatography (ethyl acetate in *n*-heptane= 10% to 30%) to give the title compound (235 mg, 55%).

¹H NMR (400 MHz, CHLOROFORM-d) δ 9.13-9.17 (m, 2H), 9.10-9.13 (m, 1H), 7.89-7.99 (m, 6H), 7.65-7.72 (m, 1H), 7.50-7.62 (m, 4H), 7.35-7.45 (m, 4H), 5.98-6.08 (m, 1H), 5.36-5.46 (m, 1H), 5.31 (s, 1H), 5.29 (s, **20** 1H), 5.27 (s, 1H), 5.11 (s, 1H), 4.43-4.49 (m, 2H), 3.83-3.92 (m, 2H), 3.69-3.77 (m, 1H), 3.31 (s, 3H), 3.01-3.17 (m, 2H), 2.64-2.75 (m, 1H), 2.28-2.37 (m, 2H), 2.07-2.28 (m, 2H), 1.94 (s, 3H).




(S)-3-((2R,3R,4S,5S)-5-((R)-2-hydroxy-4-methyl-3-methylenepent-4-en-1-yl)-3-methoxy-4-

25 ((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate. A solution of (S)-3-((2R,3R,4S,5S)-5-((R)-2-((3,5-dinitrobenzoyl)oxy)-4-methyl-3-methylenepent-4-en-1-yl)-3-methoxy-4-


((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate (0.235 g, 0.279 mmol) in methanol (4.7 mL) and THF (0.24 mL) was treated with 6-10% Mg(OMe)₂ in methanol (0.22 g, 0.17 mmol) and stirred

at room temperature for 1 h. The reaction mixture was diluted with MTBE (23.5 mL) and washed with sat. aq. NaHCO₃ (4.7 mL) and brine. The aqueous layer was extracted with MTBE (5 mL). The organic layers were combined, concentrated *in vacuo*, and purified by silica gel column chromatography (ethyl acetate in *n*-heptane=10% to 40%) to give the title compound (40 mg, 22%).


- 5 ¹H NMR (400 MHz, CHLOROFORM-*d*) δ 7.95-8.09 (m, 4H), 7.86-7.93 (m, 2H), 7.62-7.72 (m, 1H), 7.49-7.60 (m, 4H), 7.35-7.45 (m, 4H), 5.49-5.59 (m, 1H), 5.30-5.37 (m, 1H), 5.15-5.21 (m, 1H), 4.99 (s, 1H), 4.96 (s, 1H), 4.58-4.65 (m, 1H), 4.49-4.57 (m, 2H), 3.83-3.97 (m, 2H), 3.77-3.82 (m, 1H), 3.36 (s, 3H), 3.03-3.15 (m, 2H), 2.60-2.70 (m, 1H), 2.26 (t, *J*=6.44 Hz, 2H), 2.03-2.11 (m, 1H), 1.89 (s, 3H), 1.71-1.83 (m, 1H).



- 10 **(*S*)-3-((2*R*,3*R*,4*S*,5*S*)-5-(((2*S*,6*R*)-6-((*R*)-3,4-bis(benzoyloxy)butyl)-3-((2-methoxyacetoxy)methyl)-4-methyl-5,6-dihydro-2*H*-pyran-2-yl)methyl)-3-methoxy-4-((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate.** A mixture of (*S*)-3-((2*R*,3*R*,4*S*,5*S*)-5-((*S*)-2-hydroxy-4-methyl-3-methylenepent-4-en-1-yl)-3-methoxy-4-((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate (0.035 g, 0.054 mmol) and (*R*)-5,5-dimethoxypentane-1,2-diyl dibenzoate (0.030 g, 0.081 mmol)
- 15 in CH₂Cl₂ (2.1 mL) was cooled to -30 °C and treated with methoxyacetic acid (0.062 mL, 0.81 mmol) and BF₃-OEt₂ (0.021 mL, 0.162 mmol). The mixture was stirred at a temperature between -30 and -20 °C for 1.5 h, quenched with sat. aq. NaHCO₃ (3.5 mL) and extracted twice with MTBE (3.5 mL). The organic layers were combined, washed with brine (1.8 mL), and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate in *n*-heptane= 20% to 50%) to give the title compound (21 mg,
- 20 37%).
- ¹H NMR (400 MHz, CHLOROFORM-*d*) δ 7.91-8.06 (m, 8H), 7.87 (d, *J*=7.42 Hz, 2H), 7.60 (m, 1H), 7.45-7.55 (m, 6H), 7.30-7.41 (m, 8H), 5.53-5.65 (m, 1H), 5.42-5.52 (m, 1H), 4.62-4.70 (m, 1H), 4.40-4.57 (m, 5H), 4.05-4.18 (m, 1H), 3.98 (s, 2H), 3.90 (s, 2H), 3.71-3.80 (m, 1H), 3.40 (s, 3H), 3.37 (s, 3H), 3.28-3.36 (m, 1H), 2.97-3.11 (m, 2H), 2.45-2.55 (m, 1H), 2.10-2.28 (m, 2H), 1.82-1.99 (m, 3H), 1.72-1.82 (m, 1H), 1.67 (s, 3H),
- 25 1.44-1.63 (m, 4H).



Compound **14** can be prepared by treating compound **11** with compound **4** in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ and methoxyacetic acid (exemplary Prins reaction conditions):



5


(S)-3-((2R,3R,4S,5S)-3-methoxy-5-(((2R,6S)-3-((2-methoxyacetoxymethyl)-4-methyl-6-(2-((2S,5S)-3-methylene-5-(3-(pivaloyloxy)propyl)tetrahydrofuran-2-yl)ethyl)-5,6-dihydro-2H-pyran-2-yl)methyl)-4-((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate. A mixture of **(S)-3-((2R,3R,4S,5S)-5-((R)-2-hydroxy-4-methyl-3-methylenepent-4-en-1-yl)-3-methoxy-4-((phenylsulfonyl)methyl)tetrahydrofuran-2-yl)propane-1,2-diyl dibenzoate** (71 mg, 0.11 mmol) and **3-((2S,5S)-4-methylene-5-(3-oxopropyl)tetrahydrofuran-2-yl)propyl pivalate** (43 mg, 0.15 mmol) in CH_2Cl_2 (3 mL) was cooled to -30°C and treated with methoxyacetic acid (0.13 mL, 1.6 mmol) and $\text{BF}_3 \cdot \text{OEt}_2$ (0.042 mL, 0.33 mmol). After stirring at -20 to -30°C for 2 h, the reaction was quenched with sat. NaHCO_3 (3.6 mL), and the resulting mixture was extracted twice with MTBE (3.6 mL). The organic layers were combined, dried over MgSO_4 , and concentrated *in vacuo*. The residue was purified by column chromatography (ethyl acetate in *n*-heptane= 10% to 40%) to give the title compound (4 mg, 4%). Mass ($M + \text{Na}^+$): 757.4

Compound **15** can be prepared by treating compound **14** with an allylic reducing agent (e.g., $\text{Pd}(\text{PPh}_3)_4/\text{HCO}_2\text{H}/\text{Et}_3\text{N}$).

- 5 Alternatively, compound **16** can be prepared by treating compound **11** with compound **6** in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ and methoxyacetic acid (exemplary Prins reaction conditions). Compound **17** can be prepared by treating compound **16** with an allylic reducing agent (e.g., $\text{Pd}(\text{PPh}_3)_4/\text{HCO}_2\text{H}/\text{Et}_3\text{N}$).

A halichondrin macrolide analog can be prepared from compound **15** or **17** using methods and intermediates disclosed, e.g., WO 2015/066729, the synthesis of which is hereby incorporated by reference and described herein. For example, compounds **15** and **17** may be converted to a halichondrin macrolide analog, as described for the compound of formula (VIIIE) in WO 2015/066729.

Example 3



15


An exemplary compound of formula (III) can be prepared as shown in the above scheme. Compound **18** was subjected to an aldol reaction with HCHO in the presence of K_2CO_3 to afford compound **19**. Compound **19** was treated with triethyl phosphonoacetate, which was deprotonated with Cs_2CO_3 , to afford compound **20**.

20

Compound **21** can be prepared from compound **20** by treating with a 1,2-reducing agent (e.g., DIBAL-H) followed by an alkoxyethyl phosphonate, e.g., diethyl-(methoxymethyl)-phosphonate, under Wittig reaction

conditions. Compound **22** can be prepared from compound **21** by treating with a sulfonyl halide or sulfonyl anhydride, e.g., Ts_2O , followed by a halide salt, e.g., NaI or KI. Compound **23** can be prepared from compound **22** by treating with a reducing metal, e.g., elemental zinc, and a Brønsted acid, e.g., acetic acid. Compound **24** can be prepared from compound **23** by a reaction with methanol in the presence of a Brønsted acid, e.g., HCl. Compound **25** can be prepared from compound **24** by a reaction with a glycol cleaving agent, e.g., NaIO_4 , followed by a Horner-Wadsworth-Emmons reaction with alkylated phosphonoacetate, e.g., trimethyl phosphonoacetate, that has been deprotonated with a base (e.g., NaH or Cs_2CO_3). Compound **26** can be prepared from compound **25** by treating with a 1,4-reducing agent (e.g., Stryker's reagent), followed by a cyclization reaction. The cyclization reaction may proceed spontaneously after the 1,4-reduction, e.g., on work-up or during purification.


Example 4



An exemplary intermediate of formula (IA) can be prepared from compounds **3** and **26**. Compound **3** can be treated with compound **26** in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ and methoxyacetic acid (exemplary Prins reaction conditions) to afford compound **27**. Compound **27** can be treated with an allylic reducing agent (e.g., $\text{Pd}(\text{PPh}_3)_4/\text{HCO}_2\text{H}/\text{Et}_3\text{N}$) to afford compound **28**. Compound **28** can be converted to compound **29** using methods known in the art, e.g., using NaBH_4/I_2 or by converting the carboxylic acid in compound **28** to a mixed anhydride (e.g., using N-methylmorpholine and *i*-butylchloroformate) reducing the resulting mixed anhydride using borane reducing agents (e.g., 9-BBN).

A halichondrin macrolide can be prepared from compound **29** using methods and intermediates disclosed, e.g., WO 2016/179607, the synthesis of which is hereby incorporated by reference and described herein. For example, compound **29** can be converted to a halichondrin macrolide, as described for the compound of formula (VIID) in WO 2016/179607.


Example 5




An exemplary intermediate of formula (IA) can be prepared from compounds **11** and **26**. Compound **11** can be treated with compound **26** in the presence of BF₃·OEt₂ and methoxyacetic acid (exemplary Prins reaction conditions) to afford compound **30**. Compound **30** can be treated with an allylic reducing agent (e.g., Pd(PPh₃)₄/HCO₂H/Et₃N) to afford compound **31**. Compound **31** can be converted to compound **15** using methods known in the art, e.g., using NaBH₄/I₂ or by converting the carboxylic acid in compound **31** to a mixed anhydride (e.g., using N-methylmorpholine and *i*-butylchloroformate) and reducing the resulting mixed anhydride using borane reducing agents (e.g., 9-BBN).

A halichondrin macrolide analog can be prepared from compound **15** using methods and intermediates disclosed, e.g., WO 2015/066729, the synthesis of which is hereby incorporated by reference. For example, compound **15** may be converted to a halichondrin macrolide analog, as described for the compound of formula (VIIIE) in WO 2015/066729.

Example 6




- 5 (4-methyl-2,6-*cis*-diphenyl-5,6-dihydro-2*H*-pyran-3-yl)methyl 2-methoxyacetate. A mixture of trimethyl(2-methylbuta-2,3-dien-1-yl)silane (0.050 g, 0.36 mmol) and benzaldehyde (0.11 mL, 1.1 mmol) in CH₂Cl₂ (4.0 mL) was cooled to -78 °C and treated with BF₃·OEt₂ (0.14 mL, 1.1 mmol). The mixture was stirred at -78 °C for 1.5 h. After addition of methoxyacetic acid (0.41 mL, 5.4 mmol), the mixture was warmed to -40 °C and then slowly warmed to -20 °C over 2 h with stirring. The reaction was quenched with
- 10 sat. aq. NaHCO₃ (5.0 mL) and extracted twice with MTBE (5.0 mL). The organic layers were combined, washed with brine (2.5 mL), and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate in *n*-heptane= 5% to 15%) to give the title compound (53 mg, 42%).
- 15 ¹H NMR (400 MHz, CHLOROFORM-*d*) δ 7.34-7.44 (m, 4H), 7.20-7.33 (m, 6H), 5.31 (br s, 1H), 4.65-4.76 (m, 2H), 4.13-4.23 (m, 1H), 3.87 (d, *J*=16.41 Hz, 1H), 3.72 (d, *J*=16.41 Hz, 1H), 3.36 (s, 3H), 2.48-2.63 (m, 1H), 2.16-2.31 (m, 1H), 1.85 (s, 3H).



(*R*)-4-((2*R*,6*S*)-5-((2-methoxyacetoxy)methyl)-4-methyl-6-phenyl-3,6-dihydro-2*H*-pyran-2-yl)butane-1,2-diyl dibenzoate.

A mixture of trimethyl(2-methylbuta-2,3-dien-1-yl)silane (0.057 g, 0.41 mmol) and benzaldehyde (0.034 mL, 0.34 mmol) in CH₂Cl₂ (2 mL) was cooled to -78 °C and treated with BF₃·OEt₂ (0.13 mL, 1.0 mmol). The mixture was stirred at -78 °C for 1.5 h. After addition of methoxyacetic acid (0.39 mL, 5.1 mmol) and a solution of (*R*)-5,5-dimethoxypentane-1,2-diyl dibenzoate (0.16 g, 0.44 mmol) in CH₂Cl₂ (1.2 mL), the mixture was stirred at -35 °C for 10 min and warmed up to -10 °C over 2 h. The reaction was quenched with sat. aq. NaHCO₃, extracted twice with MTBE (3.6 mL), washed with brine (1.8 mL), and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate in *n*-heptane= 10%, 20% and 30%) to give the title compound (82 mg, 42%).

¹H NMR (400 MHz, CHLOROFORM-*d*) δ 7.91-8.06 (m, 4H), 7.46-7.59 (m, 2H), 7.32-7.45 (m, 6H), 7.19-7.31 (m, 3H), 5.84-5.93 (m, 1H), 5.07-5.17 (m, 2H), 3.95-4.08 (m, 1H), 3.78-3.87 (m, 2H), 3.70-3.77 (m, 2H), 3.64-3.78 (m, 1H), 3.34 (s, 3H), 2.15-2.31 (m, 2H), 1.95-2.07 (m, 1H), 1.82-1.94 (m, 2H), 1.78 (s, 3H), 1.58-1.76 (m, 1H).



***N*-((4-methyl-2,6-*cis*-diphenyl-5,6-dihydro-2*H*-pyran-3-yl)methyl)acetamide.** A mixture of trimethyl(2-methylbuta-2,3-dien-1-yl)silane (0.098 g, 0.70 mmol) and benzaldehyde (0.21 mL, 2.1 mmol) in CH₂Cl₂ (2 mL) and acetonitrile (2 mL) was cooled to -40 °C and treated with BF₃·OEt₂ (0.27 mL, 2.1 mmol). The mixture was slowly warmed up to -10 °C over 3 h. The reaction was quenched with sat. aq. NaHCO₃ (3.9 mL) and extracted twice with MTBE (3.92 mL). The organic layers were combined, washed with brine (4.90 mL), and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate in *N*-heptane= 10% to 90%) to give the title compound (60 mg, 27%).

¹H NMR (400 MHz, CHLOROFORM-*d*) δ 7.28-7.49 (m, 10H), 5.21 (br s, 1H), 4.72 (dd, *J*=3.12, 10.93 Hz, 1H), 4.50 (br s, 1H), 3.93 (dd, *J*=6.44, 14.25 Hz, 1H), 3.41 (dd, *J*=3.51, 14.06 Hz, 1H), 2.44-2.64 (m, 1H), 2.15-2.28 (m, 1H), 1.83 (s, 3H), 1.64 (s, 3H).

The experiments described herein show that Sakurai and Prins reactions can be performed as a single-pot Sakurai-Prins cascade reaction.

Other Embodiments

Various modifications and variations of the described compositions and methods of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various
5 modifications of the described modes for carrying out the invention that are obvious to those skilled in the art are intended to be within the scope of the invention.

Other embodiments are in the claims.

10