TITLE OF THE INVENTION

Therapeutic Modulation of the Tumor Inflammatory Response

BACKGROUND OF THE INVENTION

Human (and other vertebrate) immune responses comprise complex and intricately regulated sequences of events, involving cells of several different types. An imm response can be triggered when an antigen in a human body encounters an antigen-presentin cell (APC). The APC can capture the antigen and display a portion of it on its surface in a f that can be recognized by a helper T (Th) lymphocyte. Upon binding of a Th lymphocyte w the APC-displayed antigen, the Th lymphocyte can become activated. An activated Th lymphocyte promotes activation of lymphocytes of other types, the particular type(s) depen on the identity of the Th lymphocyte and the context in which the antigen is displayed by th APC. Various types of Th lymphocytes can, for example, promote activation of cytotoxic T cells or proliferation or differentiation of antigen-specific B cells. Th lymphocytes can acti other lymphocytes by secreting one or more polypeptide hormones designated cytokines. T lymphocytes exhibit the CD4 antigen on their surface.

Functionally distinct types of Th cells have been described in humans. At le two types of Th cells have been characterized, based on the cytokines they produce. Type 1 cells (i.e., Th1 cells) can produce interleukin-2 (IL-2), interferon-gamma (IFN-g), and tum necrosis factor-beta (TNF-b). Type 2 Th cells (i.e., Th2 cells) can produce IL-4, IL-5, IL-6 IL-10.

Immune responses activated by Th1 cell activation (i.e., a 'type 1 immune response' or 'Th1-mediated response') are characterized by significant production of IFN-gamma and promotion of cytotoxic lymphocyte activity. Type 1 immune responses can be induced, for example, by the presence of bacteria in the human body. The cytotoxic lymphocytes activated in a Th1-mediated response are capable of recognizing and killing c that display the Th1-activating antigen on their surface. Thus, induction of a type 1 immu response can lead to elimination from the body of antigen-bearing cells. Excessive or othe inappropriate induction of a type 1 immune response can cause damage to normal (i.e., no diseased tissues) in a human.

Immune responses activated by Th2 cell activation (i.e., a 'type 2 immune response' or 'Th2-mediated response') are characterized by significant production of IL-4 and promotion of humoral immunity (e.g., production of immunoglobulins, particularly including IgE). Type 2 immune responses are commonly induced in response to chronic infections (e.g. parasitic infections), and tend to inhibit, prevent, or reverse type 1 immune responses. Type 2 immune responses will normally eliminate a pathogen from the body, and can thereby inhibit further infection by an infectious agent. However, type 2 immune responses generally do not eliminate all pathogen-infected or diseased cells from the body. Thus, cells which exhibit the antigen that induced the type 2 immune response may persist chronically in the body. This occurs particularly when antigen-bearing cells (e.g., virus-infected cells or tumor cells) induce inappropriate activation of a type 2 immune response, which can facilitate persistence of the antigen-bearing cells in a human body.

Inflammation is a normal localized immune response to invasion or injury caused by an infectious agent (e.g., a bacterium) or by a tumor. In a process analogous to the manner in which blood flow can increase the supply of glucose and oxygen to active muscle tissue during a period of exercise, an inflammatory response can increase the supply of elements of the immune system at a local disease (e.g., infection or tumor) site in order to mount an effective defensive immune response. An effective inflammatory response can be characterized by at least six events:

i) release of antigens from diseased or pathological cells at the disease site an secretion of chemotactic factors at the injured site;

ii) infiltration of the disease site by cells of the immune system;

iii) polarized type 1 or type 2 activation of the immune cells by the antigens released at the site;

iv) amplification of the inflammatory response over time, at least for a limite period;

v) elimination of the diseased or pathological cells by immune cells; and

vi) conversion of activated immune cells into memory cells which are capabl of providing long-term protection against the antigen or antigen-bearing cells.

Cancer is one of the foremost causes of mortality and morbidity among huma Many cancers are manifested by the existence of tumors, which are clumps or masses of can cells. In the past, cancer cells were generally believed to be non-immunogenic, since they a

derived from autologous (i.e., 'self') tissue, which normally does not induce an immune response. However, tumor-reactive lymphocytes can be isolated from patients afflicted with many types of cancer (Lee et al., 1997, Blood 90:1611-1617).

A significant portion of tumor mass is made up of lymphocytes. These cells, designated tumor-infiltrating lymphocytes (TILs), typically produce cytokines (e.g., IL-4) tha are characteristic of a type 2 inflammatory response (Roussel et al., 1996, Clin. Exp. Immuno 105:344-352). It has been postulated that predominance of type 2 TILs support a type 2 inflammation in tumors that inhibits tumoricidal cytotoxic immune responses. It has furthermore been suggested that modulation of the type of immune response exhibited by TI can have anti-cancer therapeutic effects (Gorelik et al., 1994, Cancer Immunol. Immunother. 39:117-126; Pellegrini et al., 1996, Cancer Immunol. Immunother. 42:1-8; Goedegebuure et a 1997, Cell Immunol. 175:150-156; Fujimoto et al., 1997, J. Immunol. 158:5619-5626; Okam et al., 1997, Int. J. Cancer 70:598-605; Stein et al., 1998, Eur. J. Med. Res. 3:194-202; Li et a 1998, J. Surg. Oncol. 67:221-227). However, despite this recent understanding regarding bo types of inflammatory response, no course of treatment has previously been identified where a type 2 inflammatory response (i.e., one conducive to tumor survival or growth) can be converted to or overcome by a type 1 inflammatory response (i.e., one in which tumor growt slows or halts and tumor regression is enhanced).

Proper organ function is important to overall health and prevention of cancer development. One theory relating to organ function is the 3,000 year old principle of Easter medicine, termed energy balance. Each of the twelve organs controlled by the brain has its c energy, or bioEnergy. The term bioEnergy is imperfectly defined as bioelectric charge, but i more than bioelectric charge because it can be manipulated by temperature alterations, such application of heat or cooling (e.g., using a water ice pack). Each organ has its own meridia or "electrical wire," and its bioEnergy can be assessed qualitatively. There are twenty-six different meridian "lines," including two central lines that separate the body into two hemicorpses and twelve organ lines on each of the two hemicorpses. The twelve organ meridians are heart, kidney, lung, pancreas, liver, pericardial, small intestine, bladder, large intestine, stomach, gall bladder, and triple warmer. Regulatory points are distributed along these meridian lines. These points can be manipulated by various stimuli (e.g., massage, he electricity), in order to stimulate under-active organs and sedate over-active organs. The

balance of energy among meridian lines and points is important to normal function of each organ. Health arises from the simultaneously well-balanced energy among all of the organs.

Current cancer therapies (e.g., surgery, radiotherapy, and chemotherapy) are relatively inefficient, and have very debilitating side effects that lead to relapses and death. I view of the overwhelming toll of human mortality and morbidity associated with cancer, an urgent need remains for therapeutic compositions, kits, and methods which can slow or rever tumor progression in humans while reducing morbidity and offering protection against tumor relapse. The present invention satisfies this need, at least in part, by providing therapeutic compositions, kits, and methods which can be used to reliably treat a variety of human cance reduce treatment-related morbidity (relative to prior art therapeutic methods), and offer protection against tumor relapse.

BRIEF SUMMARY OF THE INVENTION

The invention relates to a method of alleviating a tumor in a human patient. T method comprises locally administering to the tumor (i) an antigen-releasing agent, (ii) a leukocyte attractant, and (iii) interferon-gamma (IFN-g) and a second type 1 inflammatory response-(IR1-) promoting agent (e.g., one of interleukin-2 (IL-2), interleukin-12 (IL-12), tumor necrosis factor-alpha (TNF-a), and tumor necrosis factor-beta (TNF-b)).

The antigen-releasing agent induces release of one or more tumor antigens fr cells of the tumor, and can, for example, be one of a proteolytic enzyme, an apoptosis-induci agent, electrical current, a strong acid, and a strong base. The leukocyte attractant induces leukocytes to infiltrate the tumor. Together, the agents administered to the tumor induce a t 1 inflammatory response in the tumor and alleviate the tumor (i.e., cause it to shrink or disappear). An added benefit of this therapeutic method is that the incidence of tumor recurrence can be decreased, relative to other tumor alleviation methods.

In one embodiment, the method further comprises locally administering to th tumor a type 1 lymphocyte attractant (e.g., one of RANTES, IP-10, and Mig), in order to sustain the type 1 inflammatory response. The type 1 inflammatory response can also (or instead) be sustained by administering autologous leukocytes to the patient. These leukocyt can be isolated from the patient, expanded ex vivo, induced to differentiate, and returned to patient, preferably by local administration at the tumor site.

In order to reduce recurrence of the tumor, a memory cell-inducing agent (interleukin-15 (IL-15), interferon-alpha (IFN-a), or interferon-beta (IFN-b)) can be administered to the patient in order to enhance production of anti-tumor type 1 immune mem cells. When used, the memory cell-inducing agent is preferably administered locally to the tumor site after most (e.g., 90%) of the tumor mass has disappeared, but before the tumor has been completely ablated.

The therapeutic method can further comprise supplementing the patient's nutrition with a nutrient such as a vitamin (e.g., one or more of vitamins A, B, C, D, and E) o mineral (e.g., one or more of selenium, zinc, calcium, magnesium, iron, and copper). Additionally, the therapeutic method can further comprise assessing organ meridian status an administering organ balance meridian therapy. Both organ meridian status assessment and organ balance meridian therapy administration can be accomplished using an electronic devi including a microcurrent stimulating device such as the ACCU-O-MATIC™ microcurrent stimulator (available from Electro-Therapeutic Devices, Inc., Markham, Ontario, Canada).

The invention also includes compositions useful in performing a tumor therapeutic method described herein. For example, such a composition can comprise IFN-g a second IR1-promoting agent. Local administration of the composition to a tumor induces a type 1 inflammatory response in the tumor, and the tumor is thereby alleviated. The composition can further comprise one or more of a leukocyte attractant, an antigen-releasing agent, and a pharmaceutically acceptable carrier.

The invention also includes a kit for alleviating a tumor in a human patient. T kit can comprise one or more of the agents used in the methods described herein, equipment devices used in those methods, and instructional material which describes one or more of the methods.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

Figure 1 is an image that depicts the organ meridian end points for one side o the body. In the Figure, LU means lung, LI means large intestine, P means pancreas, TW means triple warmer, HT means heart, SI means small intestine, SP means pericardial, LV means liver, ST means stomach, GB means gall bladder, KI means kidney, and BL means bladder.

DETAILED DESCRIPTION OF THE INVENTION

The invention relates to compositions, kits, and methods for alleviation of can in human patients. The invention is based on inducing a type 1 inflammatory response at the site of a tumor. Induction (or enhancement) of a type 1 inflammatory response at a tumor sit (i.e., within and near the margins of a tumor) can cause the tumor to shrink or disappear, redu the invasiveness or metastatic capacity of the tumor, and enhance production of immune memory cells which specifically recognize tumor tissue and inhibit or prevent relapse (i.e., recurrence) of the tumor.

Although the desirability of exchanging a type 1 immune response in tumor tissue in place of the normally-prevalent type 2 response is becoming better understood, ther presently no treatment for reliably establishing a type 1 immune response in tumor tissue. Th present invention overcomes the deficiencies of the prior art by providing therapeutic compositions, kits, and methods which can reliably be used to activate Th1 cells in and arour tumor tissue, with the result that tumor-cytotoxic inflammation is induced in the tumor, leadi to its regression or elimination.

Prior art methods of inducing tumor cell death include administration of agen which are cytotoxic to tumor cells, but which either exhibit lesser cytotoxicity with respect t non-tumor cells or are selectively delivered to tumor cells. Although these methods have exhibited limited success at limiting the size and rate of progression of tumors, at least for limited periods, many of these methods also exhibit a critical drawback that limits their effectiveness. Delivery of a cytotoxic agent (e.g., radiation or a cytotoxic chemical compou to a tumor can kill not only tumor cells, but also any non-tumor cells which are present in th tumor. This cytotoxicity can be caused by the direct effect of the agent on the non-tumor cel by a 'by-stander' effect wherein the agent induces localized cytotoxicity that is not specific f tumor cells (e.g., by release of a cytotoxin such as ricin from an agent intended to deliver the cytotoxin specifically to tumor cells), or by other mechanisms. Tumors normally comprise a significant number of leukocytes which, if appropriately activated, can induce tumor cell de and lead to tumor regression. Many anti-tumor cytotoxic agents kill these lymphocytes in addition to tumor cells. Thus, although many prior art anti-tumor agents kill significant numbers of tumor cells, the agents also kill significant numbers of leukocytes in the tumor. Elimination or inactivation of these tumors reduces the body's ability to marshal its cytotoxi immune resources to combat the tumor. The net effect of anti-tumor treatment using many

these prior art anti-tumor agents is temporary reduction in tumor mass without enhancement o the body's ability to destroy the tumor remnants or prevent recurrence of the tumor. As a resu tumor recurrence frequently occurs following prior art anti-tumor treatment methods.

The methods described in this specification can be distinguished from prior ar anti-tumor therapeutic methods in several ways. For example, the methods described herein enhance activity and proliferation of the body's anti-tumor cytotoxic immune cells within and the immediate vicinity of the tumor, rather than killing these cells as in prior art methods. Furthermore, by enhancing activity and proliferation of anti-tumor immune cells, the method described herein can inhibit or prevent recurrence of the tumor in that patient's body.

The anti-tumor therapeutic methods described herein can be outlined as follov Enzymes or other compounds which cause release of antigens from tumor cells (and which preferably also induce tumor cell death) are delivered locally to a tumor. These compounds a selected such that they are not cytotoxic with respect to lymphocytes (or at least less cytotoxi with respect to lymphocytes than with regard to tumor cells). TILs are attracted to the tumor site by local administration of one or more chemokines to tumor tissue. Local administration type 1 lymphokines to the tumor polarizes the TILs to exhibit a type 1 inflammatory respons leading to tumor tissue destruction. Sustained amplification of the type 1 inflammatory response can be effected by repeated local delivery of the type 1-polarizing cytokines to the tumor site. The net effect of this treatment is that the body's own immune defenses are mobilized for destruction of the tumor. Continued survival of type 1-polarized lymphocytes converted into memory cells, in the patient's body can prevent recurrence of the tumor at the same site or at a different body location. These lymphocytes can also travel to and destroy related tumors which may exist in the patient's body (e.g., by metastasis of cells from the original tumor).

These methods, and compositions and kits for performing them, are set forth greater detail in the following sections.

Definitions

As used herein, each of the following terms has the meaning associated with this section.

The articles "a" and "an" are used herein to refer to one or to more than one (i.e to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

A "tumor" refers to a solid or semi-solid mass of tissue comprising cells which exhibit uncontrolled growth characteristic of cancer. Tumors include both benign tumors (i.e those which do not appear to be invading surrounding tissues or metastasizing to other body sites) and malignant (i.e., non-benign) tumors.

"Co-administration" of two or more agents means administration of the agents sufficiently close in time that the periods of significant activity of each of the agents in vivo overlaps. By way of example, co-administration includes administration of a first and a second agent sufficiently close in time that the level of activity, in vivo, that is attributable to the first agent is more than half its maximum post-administration level at a period of time when the level of activity, in vivo, that is attributable to the second agent is more than half its maximum post-administration level. Co-administered agents can be administered in the form of a single composition comprising each agent, in the form of individual agent preparations, or (i.e., where there are more than two agents) in combinations of such forms.

Description

Tumorigenesis is a largely continuous occurrence in animals such as humans. Every day, millions of cell divisions occur in a human, and mutations and other genetic lesions inevitably occur in at least some of these divisions, leading to generation of cells which exhibit uncontrolled growth (i.e., cancer cells). Recognition and destruction of cancer cells are among the normal functions of the human immune system. When appropriately activated, cells of the type 1 immune response (including, for example, cytotoxic T cells) are capable of specifically destroying cells which exhibit abnormal cell surface markers (e.g., cancerous or virus-infected cells). However, survival of cancer cells is sometimes promoted by activation of cells of the type 2 immune response, which are normally associated with inactivation of parasitic or chronic infective agents, but which are not normally associated with tumorotoxicity. The type 2 immune response can sometimes be induced by the cancerous cells themselves, such as where the cancerous cells secrete factors which promote induction of the type 2 immune response.

Among the characteristics of the type 2 immune response is that type 1 inflammatory responses are inhibited when it occurs. Because tumoricidal cytotoxicity is

associated with the type 1 inflammatory response, but not with the type 2 immune response, t type 2 immune state can prolong survival of tumor tissue and permit that tissue to evade the patient's normal immune defenses.

The anti-cancer therapeutic method described in this specification involves altering the type 2 immune response which normally occurs in tumor tissue such that a type 1 inflammatory response takes over (i.e., type 1 inflammation is initiated or de-repressed). Cel associated with the type 1 inflammatory response recognize and kill tumor cells, thereby achieving tumor reduction (or even elimination) in the absence of the often debilitating or disfiguring side effects associated with administration of prior art anti-tumor agents.

In a preferred embodiment, the treatment method comprises three elements, as set forth in the three ensuing paragraphs. Although the elements are designated 'first,' 'secon and 'third,' the actions corresponding to these elements need not be performed in this sequent order (although, in a preferred embodiment, the actions corresponding to the first element are performed prior to {e.g., at least two hours prior to} the actions corresponding to the second third elements).

First, one or more agents (e.g., tumor de-bulking agents such as proteases or mixtures of proteases) are administered directly to the tumor tissue (e.g., by injection into the tumor at one or more sites) in order to cause release of tumor antigens from tumor cells. Without being bound by any particular theory of operation, it is believed that release of antig from tumor cells permits those antigens (e.g., fragments of tumor cell surface proteins) to interact with immune cells in ways that are not possible when the antigens remain part of a tumor cell. Release of tumor antigens from tumor cells facilitates induction of immune responses which are specific for the tumor cells.

Second, TILs are recruited to the tumor site by local administration of one or more TIL chemoattractants. TILs that are recruited to the tumor site can be activated to secr chemokines which attract more TILs, induce the TILs to exhibit a type 1 inflammatory response, or both.

Third, leukocytes present at (or recruited to) the tumor site are induced to exh a type 1 inflammatory response by locally administering two or more type 1 inflammatory response promoting agents ("IR1-promoting agents") including interferon-gamma ("IFN-g") with others such as interleukin-2 ("IL-2"), and tumor necrosis factor beta ("TNF-b") to the tumor tissue. The IR1-promoting agents are preferably administered together in a single

composition; however, they can be administered separately, but closely in time (e.g., seconds minutes up to a few hours apart). The IR1-promoting agents can be locally administered to tl tumor, for example, by injecting them into one or more sites in the tumor, peritumorally, or both. Preferably, the IR1-promoting agents are injected into two or more sites in the tumor, t sites spaced approximately equally apart in the tumor mass.

In the presence of two or more of these agents, TILs exhibit characteristics of type 1 inflammatory response, such as the ability to kill tumor cells and virus-infected cells. Such TILs can destroy tumor tissue, leading to shrinkage, or even disappearance, of the tumc from the patient's body and alleviation of tumor-related symptoms (e.g., pain, weight loss, nausea, exhaustion, and symptoms associated with the presence of a tumor at a particular boc location).

Although the tumor treatment method can include only these three steps, preferred embodiments of the method include one or more of three additional steps, which ai designated the fourth, fifth, and sixth steps in the ensuing three paragraphs.

Fourth, the effectiveness (i.e., sustained potency) of the treatment can be enhanced by amplifying the type 1 inflammatory response induced in and/or around the tum tissue. This can be achieved, in one embodiment, by additional (e.g., sustained or repetitive) local administration of the same or different IR1-promoting agents to the tumor tissue or the tumor site. Alternatively, or in addition, autologous lymphocytes can be provided to or near tumor site, in order to provide a greater pool of lymphocytes from which TILs can be genera or separated. The autologous lymphocytes can have been induced to proliferate ex vivo, in order to enhance the number of cells available for administration to the tumor site. The additional lymphocytes can also have been induced to differentiate into Th1-VLA6^{+} or Th1-CD49f^{+} cells by treating them, ex vivo, with one or more IR1-promoting agents.

Fifth, generation, proliferation, or both, of immune memory cells can be indu by injecting a memory cell-inducing agent (e.g., interferon-alpha {IFN-a}, interferon-beta {IFN-b} or interleukin-15 {IL-15}) at or near the tumor site at one or more steps of the treatment. Immune memory cells (e.g., activated B or T lymphocytes) are normally formed the presence of an antigen, and normally are able to rapidly differentiate to form type 1 and 2 immune cells upon re-exposure to the antigen. However, during a type 2 immune respons formation of type 1 immune memory cells can be inhibited and their differentiation or activation in response to the presence of the antigen can be inhibited. Formation of these ty

memory cells is preferably enhanced during the period of type 1 inflammatory response brou about by the other steps of the treatment disclosed herein, resulting in formation of type 1 memory cells which can rapidly respond to recurrence of the tumor at the same or a different body site.

Sixth, the effectiveness of the anti-tumor treatment disclosed herein can be enhanced by enhancing the patient's general state of health, and particularly by enhancing the state of the patient's immune system. Nutritional supplementation methods are known in the art, and preferred regimens are described herein.

In the following sections, particular aspects of these six steps are disclosed. T anti-tumor treatment includes at least the steps of enhancing tumor antigen release, inducing tumor infiltration by lymphocytes, and induction of a type 1 inflammatory response. Of cour two or more of these steps can be combined, although the antigen-release step preferably precedes the others by a period of at least several hours (i.e., in order to generate a gradient o tumor antigen having its focus at the tumor site). One or more of the fourth, fifth, and sixth steps described above (and preferably all three) can be included in the method. The sixth ste can be performed at any time during the method, and is preferably performed throughout. Tl fourth and fifth steps preferably follow the antigen-release step by a period of at least several hours.

Enhancing Tumor Antigen Release

The method includes inducing release of antigens from tumor cells. Many methods which induce antigen release are also associated with significant cytotoxicity. Thus the antigen release step can effect de-bulking of the tumor as well. So long as more (prefera significantly more) tumor cells than lymphocytes are killed, the de-bulking can enhance or speed the anti-tumor treatment by reducing the tumor burden in the patient. Tumor de-bulki can be achieved by administering a tumor de-bulking agent to the tumor in order to cause de of at least a fraction of tumor cells and release of tumor antigens from the killed tumor cells. The antigen-release-inducing agent can be administered locally to the tumor tissue, systemically, or in any other manner in which the agent is brought into contact with the tum tissue.

The antigen-release-inducing agent is preferably one which exhibits relatively little or no cytotoxicity with regard to leukocytes, particularly with regard to TILs. Example

suitable antigen-release-inducing agents include compositions comprising one or more de-bulking agents, which include proteolytic enzymes such as trypsin, chymotrypsin, pepsin, and collagenase, apoptosis-inducing agents such as alkylphospholipids (e.g., alkylphosphocholine such as hexadecylphosphocholine or edelfosine), electrical current (e.g., delivered by way of electrodes inserted into tumor tissue), and strong acids and bases (e.g., concentrated solutions sodium and potassium hydroxides and hydrochloric acid). The agent is administered locally t the tumor tissue (e.g., by topical application to the tumor tissue or by intratumoral and/or peri tumoral injection) in order to decrease damage to non-tumor tissues.

Surgical tumor disruption and radiative tumor disruption methods can be used induce release of antigens from tumor cells. However, these methods are not preferred, owin to their relative non-specificity with respect to killing tumor cells and lymphocytes. Of cours it can be possible to overcome this limitation by providing lymphocytes to the tumor site fror an extracorporeal source (e.g., autologous lymphocytes which have been induced to prolifera differentiate, or both, ex vivo).

The following four paragraphs describe examples of methods of inducing rele of antigens from a tumor located in a patient's body.

An aqueous composition comprising one or more proteases (e.g., one or more trypsin, chymotrypsin, pepsin, and collagenase) is prepared, wherein the concentration of eac protease is about 2,000 to 10,000 units per cubic centimeter of tumor volume to be targeted. aliquot of this solution representing from about 1/50 to 1/100 of the volume of the tumor to b affected is injected into the tumor. For example, about 100 microliters of a suspension of several proteases, each having a concentration of about 50,000 units per milliliter can be injected into a tumor having a volume of about 10 milliliters. This solution is preferably injected into several locations in the same tumor, in order to distribute the antigen-releasing and/or tumor-de-bulking effects throughout the tumor mass. Using this treatment, about 10-20% of the tumor can be expected to be digested (i.e., about 10-20% of tumor cells will be killed). Of course, other concentrations and combinations of enzymes can be used.

In another embodiment, antigen release from a tumor is induced by injecting solution of one or more alkylphosphocholines into the tumor mass, preferably at spatially distinct locations. Alkylphosphocholines are phospholipids which have structural similarity naturally-occurring cell membrane phospholipids. Alkylphosphocholines such as hexadecylphosphocholine and edelfosine (1-O-octadecyl-2-O-methyl-rac-glycero-3-

phosphocholine) are relatively non-toxic with regard to lymphocytes, but can induce disruptic (i.e., apoptosis) of tumor cells, thereby effecting antigen release. A suspension containing ab 50-250 milligrams of each alkylphosphocholine per milliliter of tumor volume can be prepare and an aliquot having a volume of about 1/50 to 1/100 the tumor volume can be injected. For example, 200 microliters of a suspension comprising 25% (w/v) hexadecylphosphocholine or edelfosine can be injected into a tumor having a volume of about 10 milliliters. The treatmer described in this paragraph can be expected to disrupt about 10-20% of the tumor (i.e., to induce apoptosis of 10-20% of tumor cells).

In another embodiment, release of antigens from tumor cells can be induced b applying electrical potential across a portion of the tumor (i.e., by passing electrical current through a portion of the tumor). Electrochemical tumor disruption can be effected by insertii an electrode having a polarity (e.g., an anode) in one portion of a tumor mass, inserting an electrode having the opposite polarity (e.g., a cathode) in another portion of the tumor mass, and applying an electrical potential across the electrodes. Of course, multiple electrodes of e type can be used, so long as there is at least one electrode having a polarity opposite to that o another electrode. For example, an anode having a diameter of about 0.5 millimeter can be inserted in about the center of a tumor mass, and multiple cathodes can be inserted into the periphery of the tumor mass, about 3 centimeters apart from each other. A direct current potential of about 5-10 Volts is applied between the anode and cathodes, and about 40-100 milliamperes of current are delivered for a period of about 1-2 hours. The total charge delivered to the tumor during this period is estimated to be about 60-80 Coulombs per cubic centimeter of tumor. This treatment can be expected to induce tumor tissue destruction with about 3 centimeters from each electrode.

Disruption of tumor tissue and release of tumor antigens can also be effected direct administration to tumor tissue of a solution of concentrated acid or base. For example 10 molar solution of hydrochloric acid or a 10 molar solution of sodium hydroxide can be administered intra-tumorally. An aliquot of either of these solutions equal in volume to abo 1/50 to 1/100 of the tumor volume can be injected (for example, 100 microliters of one of th solutions can be injected into a tumor having a volume of 10 milliliters). The solution is preferably injected into two or more spatially distinct locations in the tumor mass, although single injection can be used for small tumors. Treatment as described in this paragraph can

expected to induce death of about 10-20% of tumor cells, and the concentration or identity of the acid or base can be adjusted to effect this level of tumor cell killing.

Inducing Tumor Infiltration by Lymphocytes

The anti-tumor therapeutic method includes a step wherein leukocytes are attracted to the tumor site, or into the tumor itself. Recruitment of leukocytes at a particular location can also be effected by causing leukocytes that are already at the location to prolifer It is known in the art that leukocytes of particular types can be attracted to a location in a boc (or in vitro) by the existence of one or more particular chemokines at the location. Of course the particular types of leukocytes vary with the particular chemokines. For example, monocy can be attracted to a site by the presence of MCP-1, MCP-2, MCP-3, or MCP-4. T cells are attracted by RANTES, IP-10, or Mig. Eosinophils can be attracted by the presence of eotaxi a body location. Furthermore, recruitment of type 1 inflammatory cells can be enhanced by inducing proliferation or activation of the cells at the site. For example, IL-2 is known to induce proliferation of lymphocytes which exhibit the CD4 antigen (including Th1 lymphocytes) and to induce activation of lymphocytes which exhibit the CD8 antigen (e.g., cytotoxic T lymphocytes).

Leukocyte infiltration into a tumor is preferably induced following a period o several (e.g., 6-8) hours following induction of tumor antigen release. Waiting for this perio prior to inducing leukocyte infiltration permits conditions used to disrupt tumor cells, or indi antigen release therefrom, to dissipate. This period also provides time for a gradient of tumc antigen to develop having the tumor at its most concentrated location. Formation of such a gradient can enhance the ability of lymphocytes and macrophages to localize specifically wi the tumor.

One embodiment of a method by which infiltration of lymphocytes into a tun is induced is as follows. A suspension comprising IFN-g and TNF-a is prepared, comprising about 20-100 units per milliliter of tumor volume of IFN-g and about 100-500 units per milliliter of tumor volume of TNF-a. The suspension further comprises IP-10 and Mig, eacl a concentration of about 1-100 nanograms per milliliter of tumor volume. An aliquot of this suspension is administered intra-tumorally, the volume of the aliquot being approximately 1 to 1/100 the volume of the tumor. The aliquot is preferably administered by injection into o or more sites within the tumor, multiple sites preferably being approximately equidistant fro

one another. Injection of this suspension activates monocytes such that they produce monocy chemoattractants and also attracts TILs to the tumor. In some instances, IP-10, Mig, or both, can be omitted from the suspension, because these TIL chemoattractants are normally produc by activated monocytes (which are attracted to the tumor site by IFN-g and TNF-a). The suspension can be repetitively administered to the tumor, or administered in a sustained-relea formulation; however, repetitive and sustained administration will often be unnecessary, owi to the self-sustaining nature of the type 1 inflammatory response induced by attracting monocytes to the tumor site (i.e., the monocytes attract more monocytes and TILs).

In another embodiment, granulocytes (e.g., neutrophils and basophils) are attracted to the tumor site by including one or more granulocyte-attractive agents in the suspension of chemokines that is administered to the tumor. Examples of such agents includ IL-8, granular component P-2 (GCP-2), growth-related oncogens 1, 2, and 3 (GROs), neutrophil-activating protein 2 (NAP-2), and others known in the art. The desirability of attracting granulocytes to a tumor can depend on the type of tumor being treated. For examp some tumors (e.g., certain brain tumors, such as gliomas) do not form solid tissue masses, bu instead have a gelatinous consistency. In gelatinous tumors, granulocytes can enhance inducement, endurance, or both, of a type 1 inflammatory response in the tumor tissue, presumably attributable to cytotoxic factor exhibited by granulocytes. Thus, it can be prefer to include administration of a granulocyte-attracting chemokine in a composition, kit, or met for treating a semi-solid tumor.

In addition, it is recognized that certain chemokines can be more conveniently provided to certain tissues than others. For example, the chemokine designated neurotactin (sometimes designated fractalkine or type 1 membrane protein) can be administered to brain tissue (i.e., for treatment of brain tumors).

The effectiveness of numerous individual chemokines for inducing infiltratior lymphocytes into tissues of various types is known, although not all of these characteristics a replicated in this specification.

Initiation of the type 1 inflammatory response can, alternatively, be achieved supplemented by injecting another inflammation-inducing agent into the tumor mass. For example, modified bacteria (e.g., Bacillus Calmest-Guerin), lipopolysaccharides, or mild tun de-bulking or tumor-antigen-releasing agents can be used to induce inflammation that will attract monocytes to the site. Once attracted to the site, the monocytes can secrete Th1-

attracting lymphokines. Because it is preferable to achieve rapid and efficient induction of a type 1 inflammatory response, administration of IP-10, Mig, or (preferably) both, is the preferred method of inducing inflammation.

Inducing a Type 1 Inflammatory Response

The anti-tumor therapeutic methods described in this specification also includ step in which the lymphocytes present within a tumor or at the tumor site are induced to exhi a phenotype and functional properties characteristic of a type 1 inflammatory response. Activation of Th1 cells and T cytotoxic1 (Tc-1) cells which have cell surface receptors that b specifically with tumor antigens is desirable, and contributes to tumor cell cytotoxicity. In some embodiments of the method, granulocytes can also be activated such that they contribu to the type 1 immune response and contribute to tumor cell cytotoxicity.

Promotion of a type 1 immune response within or in the vicinity of a tumor ca be achieved by local administration of lymphokines which are normally produced by lymphocytes associated with a type 1 immune response. For example, a suspension of IL-2, IFN-g, and TNF-b can be administered in order to achieve this effect. The suspension can al comprise TNF-a and IL-12. Any combination of these five lymphokines can be used. However, the combination of IL-2, IFN-g, and TNF-b is preferred, because it has been discovered that this combination exhibits better synergy and permits use of lower doses of ea of the three agents. For example, one useful suspension comprises 10-100 units of IL-2 per milliliter of tumor volume, 100-1000 units of IFN-g per milliliter of tumor volume, and 50-5 units of TNF-b per milliliter of tumor volume. An aliquot of this suspension having a volun from 1/50 to 1/100 of the volume of the tumor to be treated is administered intra-tumorally, one or more sites in the tumor mass. The lymphokine suspension should be administered ev 48-72 hours until the tumor mass becomes very small (e.g., less than 5% its original size) or undetectable (e.g., using digital imaging techniques such as CAT scanning). Alternatively, t lymphokine composition can be delivered in a sustained release form, so that fewer repetitic of the administration are necessary. For example, slowly-dissolving rods comprising a biodegradable matrix (e.g., PLGA) having the lymphokine(s) sequestered therein can be use effect delivery over a period of days or weeks.

Sustained Promotion of Type 1 Inflammatory Response

The effectiveness of the anti-tumor therapeutic method described in this specification can be enhanced by multiple (or sustained or continuous) local administration o one or more chemokines, followed by local administration of cytokines normally associated with a type 1 immune response. The effectiveness of the method can also be enhanced by single or multiple provision of leukocytes expanded or differentiated in vitro.

For example, Mig and IP-10, two chemokines that attract both Th1 and Tc1 lymphocytes are prepared in a suspension comprising 10-500 nanograms each per milliliter. This suspension can be injected into the tumor less than about an hour before administering autologous lymphocytes that have been expanded or differentiated in vitro. Providing these chemokines to diffuse in the tumor before providing the lymphocytes enhances infiltration of the lymphocytes into the tumor. Without being bound by any particular theory of operation, this enhancement of infiltration is believed to be attributable to the ability of these chemokin to activate avidity of the beta-1 integrin receptors on lymphocytes, causing them to migrate along the chemokine gradient. Thus, other compounds having this same effect can be used i place of these chemokines.

Lymphocyte populations can also be expanded and differentiated in vitro usir known methods. Preferably, the lymphocytes used in such in vitro procedures are obtained from the patient to whom they expanded or differentiated lymphocytes are to be administere so that the patient's immune system will not reject the lymphocytes and so that the administe lymphocytes will not attack healthy (e.g., non-tumor) patient tissues. Also, the lymphocytes should be treated (e.g., by exposing them to IL-12) in order to enhance expression of VLA-4 integrin necessary for tumor infiltration (Roussel et al., 1997, J. Leuk. Biol. 62:356).

Expansion of blood lymphocytes can be achieved by collecting lymphocytes from a patient. These lymphocytes will normally comprise a mixture of Th1-Tc1 and Th2-T cells. The lymphocytes are maintained for five days in the presence of a low dose if IL-2 (e 10-25 units per milliliter of medium) and a low dose (e.g., 10-100 units per milliliter of medium) of either IL-12 or IFN-g to promote Th1-Tc1 expansion. Following this incubatio non-adherent cells are collected and expanded from 0.25 million cells per milliliter to 1.0 million cells per milliliter in the same medium. The cells are expanded several fold over th course of about 10 days (e.g., in AIM-V serum-free medium {GIBCO} in the presence of th same cytokines). Differentiation of the cells is achieved by adding 10-100 units per millilit

IL-12 or IFN-a to the medium for 16-24 hours prior to injection of the cells into the patient. This IL-12 (or IFN-a) treatment enhances expression of integrin VLA-6. The resulting expanded type 1 cells are infused peri-tumorally into the patient's body at a ratio of about 10 million to 100 million cells per milliliter of targeted tumor. The cells are infused or injected or near the tumor site (e.g., within a void in the tumor mass formed by a tumor-antigen-relea enhancing agent, such as electrical current or a strong acid, or at a plurality of sites surroundi the tumor mass).

Alternatively, lymphocytes can be expanded by incubating blood lymphocyte obtained from a patient with T cell receptor-specific antibodies fixed to beads (e.g., polyacrylamide beads). After rinsing the beads, they can be suspended in a medium, at a concentration of about 1 million beads per million cells, in a medium such as AIM-V in the presence of 10-100 units per milliliter IL-12 or IFN-g. Cells in the suspension are expanded several fold over the course of about 10 days. Prior to injecting the cells into the patient, the cells are separated from the beads and differentiated by exposing them to 10-100 units per milliliter IFN-a for 16-24 hours in order to enhance expression of VLA-6. The resulting expanded type 1 cells are infused peri-tumorally into the patient's body at a ratio of about 10 million to 100 million cells per milliliter of targeted tumor. The cells are infused or injected or near the tumor site.

It is understood that in certain patients (e.g., those who mount an aggressive immune response to the tumor following the initial induction of type 1 inflammation and tho who are afflicted with only small tumors), promotion of a sustained type 1 inflammatory response will be unnecessary. Thus, the anti-tumor method described herein need not includ step in which promotion of such inflammation is sustained. Omission of this step may adversely affect production of immune memory cells, resulting in a less aggressive immune response in the event the tumor recurs. Thus, this step is preferably not omitted.

Generation and Proliferation of Immune Memory Cells

The human immune system will normally produce immune memory cells dur the course of a reaction to a pathogen-infected or tumor cell when a type 1 inflammatory response occurs in association with those cells. Production of immune memory cells can be insufficient to achieve an effective immune response upon recurrence of the tumor. Howeve generation and proliferation of immune memory cells can be enhanced during a type 1

inflammatory response by administering one or more appropriate agents (i.e., "memory cell-inducing agents" such as IL-15, IFN-a, and IFN-b) to the patient, preferably, at the site of inflammatory response.

In one embodiment, production of immune memory cells is enhanced by administering about 100-1000 units per milliliter of tumor volume of one or both of IL-15 an IFN-b at the tumor site. Administration of IL-15, IFN-b, or both, decreases the intensity of tl inflammatory response and promotes conversion of activated T cells into memory T cells. These memory T cells endure in the body and perform a 'patrolling' function for many years. Further growth of the tumor at the treatment site, or growth of the tumor at a distant site, can induce a rapid anti-tumor immune response, thereby providing protection against recurrence the tumor in the patient.

Of course, the anti-tumor treatment described herein can be performed withou enhancing generation and proliferation of immune memory cells. However, owing to the lon term protective effect that can be achieved if this step is performed, the method preferably includes a step of this type.

Patient Nutrition

In order to maximize the effectiveness of the anti-tumor treatment, the genera state of health of the patient's immune system should be supported to the greatest extent possible, so that the treatment is not limited by immunodeficiencies not related to the patient tumor load. For example, the patients nutritional intake should be monitored, and supplemented if necessary, in order to ensure that nutritional deficiencies do not limit the capacity of the patient's immune cells to mount a type 1 inflammatory response. Examples c suitable nutritional regimens have been described (e.g., Bendich, 1997, Nutrition, 13:154-15. Weber et al., 1997, Nutrition 13:450-460; Rayman, 2000, Lancet 356:233-241; Anura et al., 1998, Am. J. Clin. Nutr. 68(suppl):447S-463S). It is believed that particular nutritional requirements for which patient compliance should be monitored include a daily uptake of vitamin C (200-400 milligrams), vitamin E (200-400 IU), selenium (200-400 micrograms), a zinc (15-100 milligrams). In addition, a multi-vitamin formula that includes the vitamins A, and D, and minerals such as calcium, magnesium, iron, copper, and trace amounts of other vitamins and minerals normally found in a well-equilibrated formula should be included in t patient's daily diet in order to enhance the patient's immune function.

Organ Balance Meridian Therapy

The invention includes method of administering organ balance meridian thera that is administered independently or in conjunction with the immune modulatory methods o the invention.

Organ balance meridian therapy is administered using an electronic device, su as the ACCU-O-MATIC™ microcurrent stimulator (available from Electro-Therapeutic Devices, Inc., Markham, Ontario, Canada) or any other available transcutaneous electrical ne stimulator (TENS) device. The device sensitivity is calibrated on the subject by averaging th highest and the lowest meridian reading. The device is calibrated, for example by making meridian energy readings at the meridian end-points on the fingers and toes, as indicated in Figure 1, for each of the twelve pairs of organ meridians (i.e., polarized conductivity measurements for the meridian end point), averaging the highest and lowest meridian measurements, and zeroing the device at or near that average value.

After calibrating the device, organ meridian status is assessed by taking a meridian energy reading for each organ meridian at its end points. Organ meridian balance therapy is then administered to any organ in deficiency (i.e., an organ exhibiting hypofunctic an organ which exhibits or performs a normal function at a sub-normal rate or to a sub-norm degree) and any organ in excess (i.e., an organ exhibiting hyperfunction; an organ which exhibits or performs a normal function at a supra-normal rate or to a supra-normal degree). organ having a meridian that exhibits deficient energy is tonified by stimulating regulatory points on the corresponding organ meridian (e.g., by applying microcurrent with the negativ electrode contacting the meridian end point or tonifying point and the positive electrode elsewhere on the body). An organ having a meridian that exhibits excess energy is sedated using regulatory points on the corresponding organ meridian (e.g., by applying microcurrent with the positive electrode contacting the meridian end point or sedating point and the negat electrode elsewhere on the body).

The therapeutic agent used to stimulate or sedate the regulatory points is a polarized electric current that is administered at a specific frequency (i.e., about 80Hz) and a specific intensity (i.e., about 50 to 600 microamperes), and for a selected length of time (generally about 6 to 12 seconds). Administration of a positive current is used to sedate an organ in excess and administration of a negative current is used to tonify an organ in deficie

By administering organ balance meridian therapy, normal organ function is restored. When normal organ function occurs, the complementary conditions to alleviate tumors or prevent th development (or recurrence) are optimized. Thus, the organ balance meridian therapy metho described herein can be used to improve a patient's response to the tumor inflammatory response modulating methods described herein, to inhibit tumor recurrence, or both.

Compositions

The invention includes compositions which are useful in performance of the methods described herein. Such compositions can include individual agents which are packaged in a form convenient for one or more of storage, transportation, and administration human patients. These compositions can also include multiple agents which can be administered to patients in the form of a single composition (e.g., multiple IR1-promoting agents, multiple leukocyte attractants, multiple antigen releasing agents, multiple type 1 lymphocyte attractants, multiple nutritional supplements, and chemically compatible combinations of these ingredients). By way of example, a composition that is useful in the therapeutic methods described herein comprises both a leukocyte attractant and at least one IR1-promoting agent.

Kits

The invention also includes kits that include one or more of the compositions described herein. Various kits combine, in a single package or in a plurality of packages tha are sold, shipped, or promoted for complementary use, one or more of the compositions (e.g bulk packages or unit dosage forms), equipment or devices (e.g., electrodes or syringes), anc instructional material described herein. The instructional material can be a printed material, audio or visual material, a computer-readable document or presentation, or any other tangibl medium of expression whereby use of one or more of the methods and compositions descrit herein is explained to a medical practitioner, to a patient, or both. The instructional material need not be a single instructional material, but can be a series of pamphlets, videotapes, aud recordings, product package inserts, and the like. These materials can be provided together separately with various compositions and devices described herein, or they can be provided separately with the intention that the materials will be used in conjunction with composition devices described herein in order to perform one of the therapeutic methods described herei

Medical information obtained from patients undergoing a therapy described herein can be assembled in various ways to make a database. In this database, treatment parameters (e.g., type and dose of various agents, dosing schedule, and the like) can be correlated with the characteristics of the patient (e.g., age, gender, general state of health), the disease being treated (e.g., with the tumor type or stage of progression), with the outcome tha achieved (e.g., the rapidity with which the tumor is shrunk), and with any information relatin to recurrence. The resulting database can be consulted when therapy of a new patient is undertaken, so that the method used to treat that patient can be matched with therapies which resulted in a favorable outcome in similar patients. This database can be provided (e.g., in th form of a computer-readable database) to users of the compositions, kits, and methods described herein, or it can be consulted by experts who communicate with those users.

EXAMPLES

The invention is now described with reference to the following Examples. Th Examples are provided for the purpose of illustration only and the invention is not limited to these Examples, but rather encompasses all variations which are evident as a result of the teaching provided herein.

Example 1

A patient diagnosed with a breast tumor mass having a diameter of 5 centimeters is treated using the anti-tumor treatment described herein, as follows.

A 500 milliliter sample of the patient's peripheral blood is withdrawn, and the cells are expanded in vitro, using a method described in the Detailed Description or in the pr art. The patient is instructed to begin taking a multi-vitamin formula including daily amount of 300 milligrams of vitamin C, 300 IU of vitamin E, 200 micrograms of selenium, and 50 milligrams of zinc. The total dosage of the supplement is divided in three equal doses, one t be taken with each of the three daily meals.

After five days, the patient is prepared for tumor de-bulking by being immobilized in a bed and given a local anesthetic to desensitize the nerves of the breast tissu With the assistance of live digital imaging, one electrode (negative polarity) is inserted in th center of the tumor mass, and two electrodes (each positive polarity) are inserted, one in eac end of the tumor, about 2 centimeters from the center. Electric current is activated, and

performed as described in the Detailed Description, for a duration of 2 hours. Thereafter, the patient rests for six hours.

In order to promote tumor leukocyte infiltration, a suspension comprising hun IFN-g, TFN-a, IP-10, and Mig, each at a concentration specified in the Detailed Description. The suspension is injected at five points distributed equally at the tumor periphery, each injection containing about twenty microliters of the suspension. The patient remains at rest f 24 hours.

The patient is injected with an inflammation polarizing mixture comprising IF g, TNF-b, and IL-2, each at a concentration specified in the Detailed Description. This polarizing mixture is injected twice more, at 48 and 96 hours after the initial injection.

One hour later, about five billion expanded peripheral blood lymphocytes (expanded from the patient's blood sample) are injected peri-tumorally, at five different sites surrounding the tumor. An injection of the expanded cells is made into each of the three hol created by the electrical de-bulking treatment, and two more injections are made in the periphery of the tumor. An aliquot of the suspension comprising IFN-g, IL-2, and TNF-a is injected intra-tumorally (although this suspension can, alternatively, be combined with the c suspension prior to peri-tumoral infusion of the cells). This amplification step is repeated ev two or three days until the tumor has regressed to about 10 percent of its original size.

IL-15, IFN-b, or both, are injected at the tumor site, as described in the Detail Description, in order to terminate the type 1 inflammation by converting activated T cells int memory cells. The patient is thereafter discharged, and monthly follow-up sessions are scheduled.

Example 2

A patient in a poor nutritional state caused by a wide, thin stomach tumor covering the right wall of his stomach is treated using the anti-tumor treatment described he

The patient is placed on intravenous nutrition, including the daily nutritional requirements described in the Detailed Description. This treatment is continued for six days order to partially replenish the patient's nutritional state.

Hexadecylphosphocholine (in an amount described in the Detailed Descriptic is injected at a plurality of points within the tumor, the injections being guided by live digita imaging. A mixture of proteases, as described in the Detailed Description, is injected at a

plurality of points in the tumor. The inflammation at the tumor site is monitored after 24 hou and the tumor should exhibit much infiltration and inflammation, as is characteristic of stoma tumors.

A suspension comprising IFN-g, TNF-b, and IL-2, in amounts specified in the Detailed Description, is injected at several points in the tumor mass in order to polarize the inflammatory response to a type 1 response in the tumor tissue. The tumor is monitored dail and the polarizing suspension is injected into the tumor every 48 hours. After five days on th regimen, the tumor is anticipated to regress at a rate of about 10 percent of its total surface pe day. After 8 more days, the tumor is anticipated to have shrunk to about 15 percent of its original size, but the tissues surrounding the tumor are anticipated to have become inflamed, and the patient experiences stomach pain.

The polarization treatment is discontinued, and one or both of IFN-a and IFN- are injected into the tumor mass in order to halt the inflammatory response and induce memc cell production. One or both of IFN-a and IFN-b are also injected into the inflamed tissue surrounding the remaining tumor every two or three days, until the tumor has substantially disappeared, and the inflammation has diminished considerably.

Oral nutritional intake can then be re-established in the patient. Once this is r established, the patient is discharged and monthly follow-up monitoring sessions are schedul

The disclosure of every patent, patent application, and publication cited herei hereby incorporated herein by reference in its entirety.

While this invention has been disclosed with reference to specific embodimer it is apparent that other embodiments and variations of this invention can be devised by othe skilled in the art without departing from the true spirit and scope of the invention. The appended claims include all such embodiments and equivalent variations.

CLAIMS

I claim:

1. A method of inducing tumor cell death in a human patient by inducing a type 1 inflammatory response in a solid tumor, the method comprising

i) locally administering to the tumor an antigen-releasing agent, whereby a tumor antigen is released from cells of the tumor ;

iia) locally administering to the tumor a leukocyte attractant, whereby leukocytes are induced to infiltrate the tumor ; and

iib) locally administering to the tumor an agent having interferon-gamma (IFN-g) activity as a first type 1 inflammatory response promoting agent and a second type 1 inflammatory response- (IRI-) promoting agent selected from the group consisting of an agent having tumor necrosis factor-beta (TNF-b) activity, an agent having tumor necrosis factor-alpha (TNF-a) activity, an agent having interleukin-2 (IL-2) activity, an agent having interleukin-12 (IL-12) activity, and a mixture thereof, whereby a type 1 inflammatory response is induced in the tumor and tumor cell death is induced.

2. The method of claim 1, wherein the antigen-releasing agent is a tumor debulking agent.

3. The method of claim 1, wherein the antigen-releasing agent comprises an agent selected from the group consisting of a proteolytic enzyme, an apoptosis-inducing agent, electrical current, an acid, a base, and a mixture thereof.

4. The method of claim 3, wherein the antigen-releasing agent comprises a proteolytic enzyme selected from the group consisting of an agent having trypsin activity, an agent having chymotrypsin activity, an agent having pepsin activity, an agent having collagenase activity and a mixture thereof.

5. The method of claim 3, wherein the antigen-releasing agent comprises only one proteolytic enzyme.

6. The method of claim 3, wherein the antigen-releasing agent comprises at least two proteolytic enzymes.

7. The method of claim 3, wherein the antigen-releasing agent comprises an alkylphospholipid.

8. The method of claim 7, wherein thealkylphospholipid is an alkylphosphocholine.

9. The method of claim 7, wherein the alkylphosphocholine is selected from the group consisting of alkylphosphocholine having hexadecylphosphocholine activity, alkylphosphocholine having edelfosine activity, and a mixture thereof.

10. The method of claim 3, wherein the antigen-releasing agent is electrical current delivered by way of electrodes inserted into the tumor.

11. The method of claim 3, wherein the antigen-releasing agent comprises an acid selected from the group having hydrochloric acid, sulfuric acid activity, and a mixture thereof.

12. The method of claim 3, wherein the antigen-releasing agent comprises a base selected from the group having sodium hydroxide, concentrated potassium hydroxide activity, and a mixture thereof.

13. The method of claim1, wherein the antigen-releasing agent is administered to the tumor at least two hours before administering the leukocyte attractant to the tumor.

14. The method of claim 1, wherein the antigen-releasing agent and the leukocyte attractant are co-administered to the tumor.

15. The method of claim 1, wherein the antigen-releasing agent is administered to the tumor at least two hours before administering an agent having IFN-g activity to the tumor.

16. The method of claim1, wherein the antigen-releasing agent and the agent having IFN-g activity are co-administered to the tumor.

17. The method of claim1, wherein the leukocyte attractant comprises a monocyte attractant.

18. The method of claim 17, wherein the monocyte attractant is selected from the group of agents having MCP-1, MCP-2, MCP-3, MCP-4 activity, and a mixture thereof.

19. The method of claim 1, wherein the leukocyte attractant comprises a T cell attractant.

20. The method of claim 19, wherein the T cell attractant is selected from the group of agents having RANTES, IP-10, Mig activity, and a mixture thereof.

21. The method of claim 1, wherein the leukocyte attractant comprises a granulocyte attractant.

22. The method of claim 21, wherein the granulocyte attractant is selected from the group of agents having interleukin-8, granulocyte chemotactic protein-2, growth-related oncogen-1, growth-related oncogen-2, growth-related oncogen-3, neutrophil activated protein, neurotactin activity, and a mixture thereof.

23. The method of claim 21, wherein the granulocyte attractant is a eosinophil attractant.

24. The method of claim 23, wherein the eosinophil attractant is an agent having eotaxin activity.

25. The method of claim 1, wherein the leukocyte attractant is co-administered with at least one agent having IFN-g activity and the second IRl-promoting agent.

26. The method of claim1, wherein the leukocyte attractant and at least one of the agent having IFN-g activity and the second IRl-promoting agent are administered not more than two hours apart.

27. The method of claim 1, wherein the leukocyte attractant and at least one of the agent having IFN-g activity and the second IRl-promoting agent are administered more than two hours apart.

28. The method of claim 1, wherein the leukocyte attractant and at least one of the agent having IFN-g activity and the second IRl-promoting agent are co-administered.

29. The method of claim 1, wherein secondIR1-promoting agent is selected from the group agents having interleukin-2 (IL-2), interleukin-12 (IL-12), tumor necrosis factor alpha (TNF-a), tumor necrosis factor-beta (TNF-b) activity, and a mixture thereof.

30. The method of claim 29, wherein the second IRl-promoting agent comprises an agent having IL-2 activity.

31. The method of claim 29, wherein the second IRl-promoting agent comprises an agent having TNF-b activity.

32. The method of claim 29, wherein the second IRl-promoting agent comprises both agents having IL-2 and TNF-b activities.

33. The method of claim 1, wherein multiple aliquots of each of the agents having IFN-g activity,the second and additional IRl-promoting agents are administered to the patient, and wherein at least 48 hours elapse between aliquots.

34. The method of claim 1, wherein the agent having IFN-g activity ,second, and the additional IRl-promoting agents are co-administered.

35. The method of claim 1, wherein the agent having IFN-g activity, the second, and the additional IRl-promoting agents are separately administered not more than two hours apart.

36. The method of claim 1, wherein the agent having IFN-g activity and the second IRl-promoting agent are separately administered more than two hours apart.

37. The method of claim 1, further comprising iii) locally administering to the tumor a type 1 lymphocyte attractant in order to sustain the type 1 inflammatory response.

38. The method of claim 37, wherein the type 1 lymphocyte attractant is selected from the agents having RANTES, IP-10, Mig activities, and a mixture thereof.

39. The method of claim 37, wherein two type 1 lymphocyte attractants are locally administered and wherein the type 1 lymphocyte attractants are agents havingIP-10 and Mig activities.

40. The method of claim 37, further comprising iv) sustaining the type 1 inflammatory response by locally administering autologous leukocytes to the tumor.

41. The method of claim 37, further comprising iv) administering a memory cell-inducing agent to the patient after inducing the type 1 inflammatory response, whereby production of anti-tumor type 1 immune memory cells is enhanced.

42. The method of claim 41, further comprising v) supplementing the patient's nutrition with a nutrient selected from the group consisting of a vitamin, a mineral, and a mixture

thereof.

43. The method of claim 37, further comprising iv) supplementing the patient's nutrition with a nutrient selected from the group consisting of a vitamin, a mineral, and a mixture thereof.

44. The method of claim1, further comprising iii) locally administering autologous leukocytes to the tumor.

45. The method of claim 44, wherein the autologous leukocytes are obtained from the patient and expanded prior to locally administering them to the tumor.

46. The method of claim 44, wherein the autologous leukocytes are obtained from the patient and contacted with an IRl-promoting agent prior to locally administering them to the tumor.

47. The method of claim 44, wherein the autologous leukocytes are obtained from the patient, expanded ex vivo, and contacted with an IRl-promoting agent prior to locally administering them to the tumor.

48. The method of claim 47, wherein the leukocytes are contacted with both an IRl-promoting agent and with at least one of an agent having interferon-alpha (IFN-a) activity and an agent having IL-12 activity prior to locally administering them to the tumor.

49. The method of claim 44, further comprising iv) administering a memory cell-inducing agent to the patient after inducing the type 1 inflammatory response, whereby production of anti-tumor type 1 immune memory cells is enhanced.

50. The method of claim 49, further comprising v) supplementing the patient's nutrition with a nutrient selected from the group consisting of a vitamin, a mineral, and a mixture

thereof.

51. The method of claim 44, further comprising iv) supplementing the patient's nutrition with a nutrient selected from the group consisting of a vitamin, a mineral, and a mixture thereof.

52. The method of claim 1, further comprising iii) administering a memory cell-inducing agent to the patient after inducing the type 1 inflammatory response, whereby production of anti-tumor type 1 immune memory cells is enhanced.

53. The method of claim 52, wherein the memory cell-inducing agent is selected from the group consisting of an agent having interleukin-15 (IL-15) activity, an agent having IFN-a activity, an agent having IFN-b activity, and a mixture thereof.

54. The method of claim 52, wherein the memory cell-inducing agent is an agent having IL-15 activity.

55. The method of claim 52, wherein the memory cell-inducing agent is an agent having IFN-b activity.

56. The method of claim 52, wherein the memory cell-inducing agent is administered after the tumor shrinks to less than 10 percent of its size immediately prior to administration of the antigen-releasing agent.

57. The method of claim 1, further comprising supplementing the patient's nutrition with a nutrient selected from the group consisting of a vitamin, a mineral, and a mixture thereof.

58. The method of claim 57, wherein the vitamin is selected from the group consisting of vitamins A, B, C, D, E, and a mixture thereof.

59. The method of claim 58, wherein the vitamin is vitamin C and wherein the patient's nutrition is supplemented such that the patient receives from 200 to 400 milligrams of vitamin C daily.

60. The method of claim 58, wherein the vitamin is vitamin E and wherein the patient's nutrition is supplemented such that the patient receives from 200 to 400 international units of vitamin E daily.

61. The method of claim 57, wherein the mineral is selected from the group consisting of selenium, zinc, calcium, magnesium, iron, copper, and a mixture thereof.

62. The method of claim 61, wherein the mineral is selenium and wherein the patient's nutrition is supplemented such that the patient receives from 200 to 400 micrograms of selenium daily.

63. The method of claim 61, wherein the mineral is zinc and wherein the patient's nutrition is supplemented such that the patient receives from 15 to 100 milligrams of zinc daily.

64. The method of claim 57, wherein the patient's nutrition is supplemented beginning at least on the same day that the antigen-releasing agent is administered to the tumor, and continuing through at least the same day that an agent having IFN-g activity is administered to the tumor.

65. The method of claim 57, wherein the patient's nutrition is supplemented beginning at least five days before the antigen-releasing agent is administered to the tumor, and continuing through at least three days after the day that an agent having IFN-g activity is administered to the tumor.

66. A method of alleviating a tumor in a human patient by inducing a type 1 inflammatory response in a solid tumor, the method comprising

i) supplementing the patient's nutrition with a nutrient selected from the group consisting of a vitamin, a mineral, and a mixture thereof;

ii) locally administering to the tumor an antigen-releasing agent, whereby a tumor antigen is released from cells of the tumor; thereafter

iiia) locally administering to the tumor a leukocyte attractant, whereby leukocytes are induced to infiltrate the tumor; and

iiib) locally administering to the tumor a protein having interferon-gamma (IFN-g) activity and a second type 1 inflammatory response- (IRI-) promoting agent selected from the group consisting of an agent having tumor necrosis factor-beta (TNF-b) activity, an agent having tumor necrosis factor-alpha (TNF-a) activity, an agent having interleukin-2 (IL-2) activity, an agent having interleukin-12 (IL-12) activity, and a mixture thereof., whereby a type 1 inflammatory response is induced in the tumor; thereafter

iv) sustaining the type 1 inflammatory response by

iva) locally administering to the tumor a type 1 lymphocyte attractant, ivb) locally administering autologous leukocytes to the tumor, or ivc) both iva) and ivb); and thereafter

v) administering a memory cell-inducing agent to the patient after inducing the type 1 inflammatory response, whereby production of anti-tumor type 1 immune memory cells is enhanced and tumor cell death is induced.

67. A composition for alleviating a tumor in a human patient, the composition comprising an agent having IFN-g activity and a second IRI-promoting agent selected from the group consisting of an agent having tumor necrosis factor-beta (TNF-b) activity, an agent having tumor necrosis factor-alpha (TNF-a) activity, an agent having interleukin-2 (IL-2) activity, an agent having interleukin-12 (IL-12) activity, and a mixture thereof., whereby upon local administration of the composition to the tumor, a type 1 inflammatory response is induced in the tumor and tumor cell death is induced.

68. The composition of claim 67, wherein the second IRl-promoting agent is selected from the group of agents having IL-2, IL-12,TNF-a, and TNF-b activities.

69. The composition of claim 68, wherein the second IR1-promoting agent comprises an agent having IL-2 activity.

70. The composition of claim 69, further comprising an agent having TNF-b activity.

71. The composition of claim 67, further comprising a leukocyte attractant.

72. The composition of claim71, wherein the leukocyte attractant is selected from the group consisting of agents having MCP-1, MCP-2, MCP-3,MCP-4, RANTES, IP-10, Mig, interleukin-8, granulocyte chemotactic protein-2, growth-related oncogen1, growth-related oncogen-2, growth-related oncogen-3, neutrophil activated protein, neurotactin, eotaxin activities, and a mixture thereof.

73. The method of claim 1, further comprising iiia) assessing organ meridian status in the patient; and iiib) administering organ balance meridian therapy to the patient.

74. The method of claim 73, wherein an electronic device is used to assess organ meridian status and to administer organ balance meridian therapy.

75. The method of claim 74, wherein the device is a transcutaneous microcurrent stimulator.

76. A kit for alleviating a tumor in a human patient, the kit comprising an antigen-releasing agent; a leukocyte attractant; an agent having IFN-g activity; and a type 1 inflammatory response promoting agent selected from the group consisting of an agent having tumor necrosis factor-beta (TNF-b) activity, an agent having tumor necrosis

factor-alpha (TNF-a) activity, an agent having interleukin-2 (IL-2) activity, an agent having interleukin-12 (IL-12) activity, and a mixture thereof.

77. The kit of claim 76, further comprising an instructional material which describes administration of the antigen-releasing agent, the leukocyte attractant, the protein having IFN-g activity, and the type 1 inflammatory response promoting agent to the patient for inducing a type 1 inflammatory response in a solid tumor.

78. The kit of claim 76, wherein the antigen-releasing agent comprises an agent selected from the group consisting of a proteolytic enzyme, an apoptosis-inducing agent, electrical current, an acid, base; and a mixture thereof.

79. The kit of claim 78, wherein the antigen-releasing agent comprises a proteolytic enzyme selected from the group consisting of a protein having an agent having trypsin activity, an agent having chymotrypsin activity, an agent having pepsin activity, an agent having collagenase activity, and a mixture thereof.

80. The kit of claim 78, wherein the antigen-releasing agent comprises only one proteolytic enzyme.

81. The kit of claim 78, wherein the antigen-releasing agent comprises at least two proteolytic enzymes.

82. The kit of claim 78, wherein the antigen-releasing agent comprises an alkylphospholipid.

83. The kit of claim 82, wherein the alkylphospholipid is an alkylphosphocholine.

84. The kit of claim 82, wherein the alkylphosphocholine is selected from the group an alkylphosphocholine having hexadecylphosphocholine activity, an alkylphosphocholine having edelfosine activity, and a mixture thereof.

85. The kit of claim 78, wherein the antigen-releasing agent comprises an electrode for delivering electrical current into the tumor.

86. The kit of claim 78, wherein the antigen-releasing agent comprises an agent having activity of an acid selected from the group hydrochloric acid, sulfuric acid , and a mixture thereof.

87. The kit of claim 78, wherein the antigen-releasing agent comprises an agent having activity of a base selected from the group sodium hydroxide, potassium hydroxide, and a mixture thereof.

88. The kit of claim 76, wherein the leukocyte attractant comprises a monocyte attractant.

89. The kit of claim 88, wherein the monocyte attractant has the activity of an agent selected from the group MCP-1, MCP-2, MCP-3, MCP-4, and a mixture thereof.

90. The kit of claim 76, wherein the leukocyte attractant comprises a T cell attractant.

91. The kit of claim 90, wherein the T cell attractant has the activity of an agent selected from the group RANTES, IP-10, Mig, and a mixture thereof.

92. The kit of claim 76, wherein the leukocyte attractant comprises a granulocyte attractant.

93. The kit of claim 92 , wherein the granulocyte attractant has the activity of an agent selected from the group interleukin-8, granulocyte chemotactic protein-2, growth-related oncogen-1, growth-related oncogen-2, growth-related oncogen-3, neutrophil activated protein, neurotactin, and a mixture thereof.

94. The kit of claim 92, wherein the granulocyte attractant is a eosinophil attractant.

95. The kit of claim 94, wherein the eosinophil attractant is an agent having eotaxin activity.