
CLAIMS

1. An ~~liquid composition that is an~~ aqueous solution of a compound of formula (I)



I

wherein:

R^1 is selected from H, $-CH_2-O-R^3$, $-CH_2-S-R^3$, and $-CH_2-NR^3R^4$;

R^2 is selected from $-CH_2-O-R^3$, $-CH_2-S-R^3$, and $-CH_2-NR^3R^4$;

R^3 and R^4 are the same or different and are independently selected from H; substituted or non-substituted, unbranched or branched, saturated or unsaturated C3-C12 cycloalkyl or C1-C10 alkyl; substituted or non-substituted benzyl; substituted or non-substituted mono- or bicyclic aryl; substituted or non-substituted mono-, bi- or tricyclic C2-C10 heteroaryl or non-aromatic C2-C10 heterocyclyl containing one or several heteroatoms independently selected from N, O and S; or R^3 and R^4 in $-CH_2-NR^3R^4$ are bonded together and form, together with the nitrogen atom to which they are bonded, a substituted or non-substituted non-aromatic C2-C10 mono- or bicyclic heterocyclyl optionally containing one or several further heteroatoms independently selected from N, O and S and optionally comprising one or several cyclic keto groups;

wherein the substituents of the substituted groups are independently selected from unbranched or branched, saturated or unsaturated C3-C12 cycloalkyl or C1-C10 alkyl; halogen; halogen-substituted C1-C10 alkyl, mono- or bicyclic aryl; mono-, bi- or tricyclic C2-C10 heteroaryl or non-aromatic C2-C10 heterocyclyl containing one or several heteroatoms independently selected from N, O and S; C1-C10 alkoxy; amino; C1-C10 alkylamino;
or a pharmaceutically acceptable salt thereof, and
a pH regulating agent;
wherein the aqueous solution has a pH from 3.0 to 5.0, and wherein the compound is 2-(hydroxymethyl)-2-(methoxymethyl)quinuclidin-3-one.

~~2. The liquid composition according to claim 1, wherein R^1 is selected from H and $-CH_2-O-R^3$; and R^2 is $-CH_2-O-R^3$.~~

~~3. The liquid composition according to claim 1 or 2, wherein each R^3 is independently selected from H and substituted or non-substituted, unbranched or branched, saturated or unsaturated C1-C6 alkyl.~~

~~4. The liquid composition according to claim 3, wherein each R^3 is independently selected from H and methyl.~~

~~5. The liquid composition according to claim 4, wherein the compound is selected from 2-(hydroxymethyl)-2-(methoxymethyl)quinuclidin-3-one and 2,2-bis(hydroxymethyl)-quinuclidin-3-one.~~

~~6~~2. The ~~liquid composition~~aqueous solution according to ~~any one of the~~ claims 1-~~5~~, comprising at least one further therapeutically active agent.

~~7~~3. The ~~liquid composition~~aqueous solution according to any one of the claims 1-~~6~~2, having a pH between 3.0 and 4.5.

~~8~~4. The ~~liquid composition~~aqueous solution according to any one of the claims 1-~~7~~3, wherein the compound of formula (I) is present in the aqueous solution at a concentration of from 10 mg/mL to 250 mg/mL.

~~9~~5. The ~~liquid composition~~aqueous solution according to any one of the claims 1-~~8~~4, wherein the aqueous solution comprises NaCl at a concentration of from 0 to 3% weight by volume.

~~10~~6. The ~~liquid composition~~ aqueous solution according to any one of the claims 1-~~9~~5, for use in therapy.

~~11~~7. The ~~liquid composition~~ aqueous solution according to any one of the claims 1-~~9~~5, for use in the treatment of a disorder selected from hyperproliferative diseases, autoimmune diseases and heart diseases.

~~12~~8. Use of an aqueous solution ~~liquid composition~~ according to any one of the claims 1-~~9~~5, for the manufacturing of a medicament for the treatment of a disorder selected from hyperproliferative diseases, autoimmune diseases and heart diseases.

~~13~~9. A method for preparing an aqueous solution ~~liquid composition~~ according to any one of the claims 1-~~6~~2, by adding, to an aqueous phase, a compound of formula (I), or a pharmaceutically acceptable salt thereof; optionally at least one further therapeutically active agent; and a pH regulating agent in an amount so as to provide a pH in the composition of from 3.0 to 5.0.

~~14~~10. The method according to claim ~~13~~9, wherein the pH regulating agent is added in an amount so as to provide a pH in the composition of from 3.0 to 4.5.

~~15~~11. The method according to claim ~~13~~9 or claim ~~14~~10, wherein NaCl is added to the aqueous phase in an amount so as to provide a concentration of NaCl in the liquid composition of from 0 to 3% weight by volume.

~~16~~12. The method according to any one of the claims ~~13~~9 to ~~15~~11, wherein the compound of formula (I), or the pharmaceutically acceptable salt thereof, is added to the aqueous phase in an amount so as to provide a concentration of the compound of formula (I) in the liquid composition of from 10 mg/mL to 250 mg/mL