

CLAIMS**WHAT IS CLAIMED IS:**


1. An engineered host cell comprising:
 - a. a first nucleic acid molecule comprising an unnatural nucleotide; and
 - b. optionally, a second nucleic acid molecule encoding a modified transposition-associated protein or transposable element.
2. The engineered host cell of claim 1, further comprising:
 - a. a third nucleic acid molecule encoding a modified nucleoside triphosphate transporter, wherein the third nucleic acid molecule is incorporated in a genomic sequence of the engineer host cell, or comprises a plasmid encoding the modified nucleoside triphosphate transporter.
3. The engineered host cell of claim 1, wherein the modified nucleoside triphosphate transporter exhibits increased stability of expression in the engineered host cell as compared to an expression in an equivalent engineered host cell that does not comprise the second nucleic acid molecule encoding the modified transposition-associated protein.
4. The engineered host cell of claim 2, wherein the modified nucleoside triphosphate transporter comprises a deletion of an entire nucleic acid molecule encoding the nucleoside triphosphate transporter, an N-terminal truncation, a C-terminal truncation, or a truncation of both termini.
5. The engineered host cell of claim 2, wherein modified nucleoside triphosphate transporter comprises a nucleoside triphosphate transporter from *Phaeodactylum tricornutum* (*PtNTT₂*).
6. The engineered host cell of claim 5, wherein the *PtNTT₂* is under the control of a promoter selected from a pSC plasmid or a promoter from a lac operon.
7. The engineered host cell of claim 2, further comprising:
 - a. a Cas9 polypeptide or variants thereof; and
 - b. a single guide RNA (sgRNA) comprising a crRNA-tracrRNA scaffold, wherein the combination of Cas9 polypeptide or variants thereof and sgRNA modulates replication of the first nucleic acid molecule encoding the unnatural nucleotide.
8. The engineered host cell of claim 1, wherein the second nucleic acid molecule comprises a gene comprising *catalase* (*cat*), *IS1 protein insB-4* (*insB-4*), *IS1 protein insA-4* (*insA-4*), or a combination thereof.
9. The engineered host cell of claim 1, wherein the modified transposition-associated protein comprises Insertion element IS1 4 protein InsB, Insertion element IS1 4 protein

InsA, or a combination thereof; and wherein the modified transposable element comprises IS/.

10. The engineered host cell of claim 8, wherein the gene comprises one or more deletions, wherein the one or more deletions comprise an N-terminal deletion, a C-terminal deletion, a truncation at both termini, an internal deletion, and/or a deletion of the entire gene.
11. The engineered host cell of claim 1, further comprising a fifth nucleic acid molecule encoding a modified DNA repair response-associated protein, wherein the DNA repair response comprises recombinational repair, SOS response, nucleotide excision repair, or methyl-directed mismatch repair, or a combination thereof.
12. The engineered host cell of claim 11, wherein the modified DNA repair response-associated protein comprises RecA, Rad51, RadA, or LexA, or a combination thereof.
13. The engineered host cell of claim 1, wherein the engineered host cell is a prokaryotic cell comprising an *Escherichia coli* cell, *Escherichia coli* BL21 (DE3) cell.
14. The engineered host cell of claim 1, wherein the unnatural nucleotide comprises an unnatural base selected from the group consisting of 2-aminoadenin-9-yl, 2-aminoadenine, 2-F-adenine, 2-thiouracil, 2-thio-thymine, 2-thiocytosine, 2-propyl and alkyl derivatives of adenine and guanine, 2-amino-adenine, 2-amino-propyl-adenine, 2-aminopyridine, 2-pyridone, 2'-deoxyuridine, 2-amino-2'-deoxyadenosine 3-deazaguanine, 3-deazaadenine, 4-thio-uracil, 4-thio-thymine, uracil-5-yl, hypoxanthin-9-yl (I), 5-methyl-cytosine, 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 5-bromo, and 5-trifluoromethyl uracils and cytosines; 5-halouracil, 5-halocytosine, 5-propynyl-uracil, 5-propynyl cytosine, 5-uracil, 5-substituted, 5-halo, 5-substituted pyrimidines, 5-hydroxycytosine, 5-bromocytosine, 5-bromouracil, 5-chlorocytosine, chlorinated cytosine, cyclocytosine, cytosine arabinoside, 5-fluorocytosine, fluoropyrimidine, fluorouracil, 5,6-dihydrocytosine, 5-iodocytosine, hydroxyurea, iodouracil, 5-nitrocytosine, 5-bromouracil, 5-chlorouracil, 5-fluorouracil, and 5-iodouracil, 6-alkyl derivatives of adenine and guanine, 6-azapyrimidines, 6-azo-uracil, 6-azo cytosine, azacytosine, 6-azo-thymine, 6-thio-guanine, 7-methylguanine, 7-methyladenine, 7-deazaguanine, 7-deazaguanosine, 7-deaza-adenine, 7-deaza-8-azaguanine, 8-azaguanine, 8-azaadenine, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, and 8-hydroxyl substituted adenines and guanines; N4-ethylcytosine, N-2 substituted purines, N-6 substituted purines, O-6 substituted purines, those that increase the stability of duplex formation, universal nucleic acids, hydrophobic nucleic acids, promiscuous nucleic acids, size-expanded nucleic acids, fluorinated nucleic acids, tricyclic pyrimidines, phenoxazine cytidine(

[5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps, phenoxazine cytidine (9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido [3',2':4,5]pyrrolo [2,3-d]pyrimidin-2-one), 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid, wybutosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine and those in which the purine or pyrimidine base is replaced with a heterocycle.

15. The engineered host cell of claim 1, wherein the unnatural base is selected from the group consisting of:



16. The engineered host cell of claim 1, wherein the unnatural nucleotide further comprises an unnatural sugar moiety.
17. The engineered host cell of claim 16, wherein the unnatural sugar moiety is selected from the group consisting of a modification at the 2' position: OH; substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂F; O-alkyl, S-alkyl, N-alkyl; O-alkenyl, S-alkenyl, N-alkenyl; O-alkynyl, S-alkynyl, N-alkynyl; O-alkyl-O-alkyl, 2'-F, 2'-OCH₃,


2'-O(CH₂)₂OCH₃ wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁-C₁₀, alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, -O[(CH₂)_n O]_mCH₃, -O(CH₂)_nOCH₃, -O(CH₂)_nNH₂, -O(CH₂)_nCH₃, -O(CH₂)_n-ONH₂, and -O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10; and/or a modification at the 5' position: 5'-vinyl, 5'-methyl (R or S), a modification at the 4' position, 4'-S, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and any combination thereof.

18. A method of increasing the production of a nucleic acid molecule comprising an unnatural nucleotide, comprising:
 - a) incubating an engineered host cell with a plurality of unnatural nucleotides, wherein the engineered host cell comprises a modified nucleoside triphosphate transporter and optionally, a modified transposition-associated protein or transposable element; and
 - b) incorporating the plurality of unnatural nucleotides into one or more newly synthesized DNA strands, thereby generating the unnatural nucleic acid molecule;

wherein the modified transposition-associated protein or transposable element and the modified nucleoside triphosphate transporter increases retention of unnatural base pairs comprising the unnatural nucleotides in the one or more newly synthesized DNA strands.
19. The method of claim 18, wherein the modified transposition-associated protein comprises comprises Insertion element IS1 4 protein InsB, Insertion element IS1 4 protein InsA, or a combination thereof; and wherein the modified transposable element comprises IS1.
20. The method of claim 18, wherein the modified nucleoside triphosphate transporter comprises a codon optimized nucleoside triphosphate transporter from *Phaeodactylum tricornutum* (PtNTT₂).
21. The method of claim 18, wherein the unnatural nucleotide comprises an unnatural base selected from the group consisting of 2-aminoadenin-9-yl, 2-aminoadenine, 2-F-adenine, 2-thiouracil, 2-thio-thymine, 2-thiocytosine, 2-propyl and alkyl derivatives of adenine and guanine, 2-amino-adenine, 2-amino-propyl-adenine, 2-aminopyridine, 2-pyridone,

2'-deoxyuridine, 2-amino-2'-deoxyadenosine 3-deazaguanine, 3-deazaadenine, 4-thio-uracil, 4-thio-thymine, uracil-5-yl, hypoxanthin-9-yl (I), 5-methyl-cytosine, 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 5-bromo, and 5-trifluoromethyl uracils and cytosines; 5-halouracil, 5-halocytosine, 5-propynyl-uracil, 5-propynyl cytosine, 5-uracil, 5-substituted, 5-halo, 5-substituted pyrimidines, 5-hydroxycytosine, 5-bromocytosine, 5-bromouracil, 5-chlorocytosine, chlorinated cytosine, cyclocytosine, cytosine arabinoside, 5-fluorocytosine, fluoropyrimidine, fluorouracil, 5,6-dihydrocytosine, 5-iodocytosine, hydroxyurea, iodouracil, 5-nitrocytosine, 5-bromouracil, 5-chlorouracil, 5-fluorouracil, and 5-iodouracil, 6-alkyl derivatives of adenine and guanine, 6-azapyrimidines, 6-azo-uracil, 6-azo cytosine, azacytosine, 6-azo-thymine, 6-thio-guanine, 7-methylguanine, 7-methyladenine, 7-deazaguanine, 7-deazaguanosine, 7-deaza-adenine, 7-deaza-8-azaguanine, 8-azaguanine, 8-azaadenine, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, and 8-hydroxyl substituted adenines and guanines; N4-ethylcytosine, N-2 substituted purines, N-6 substituted purines, O-6 substituted purines, those that increase the stability of duplex formation, universal nucleic acids, hydrophobic nucleic acids, promiscuous nucleic acids, size-expanded nucleic acids, fluorinated nucleic acids, tricyclic pyrimidines, phenoxazine cytidine([5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps, phenoxazine cytidine (9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido [3',2':4,5]pyrrolo [2,3-d]pyrimidin-2-one), 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxoacetic acid, wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxoacetic acid methylester, uracil-5-oxoacetic acid, 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine and those in which the purine or pyrimidine base is replaced with a heterocycle.

22. The method of claim 19, wherein the unnatural base is selected from the group consisting of:



23. The method of claim 19, wherein the unnatural nucleotide further comprises an unnatural sugar moiety.
24. The method of claim 23, wherein the unnatural sugar moiety is selected from the group consisting of a modification at the 2' position: OH; substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂ CH₃, ONO₂, NO₂, N₃, NH₂F; O-alkyl, S-alkyl, N-alkyl; O-alkenyl, S-alkenyl, N-alkenyl; O-alkynyl, S-alkynyl, N-alkynyl; O-alkyl-O-alkyl, 2'-F, 2'-OCH₃, 2'-O(CH₂)₂OCH₃ wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁-C₁₀, alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, -O[(CH₂)_n O]_mCH₃, -O(CH₂)_nOCH₃, -O(CH₂)_n NH₂, -O(CH₂)_n CH₃, -O(CH₂)_n -ONH₂, and -O(CH₂)_nON[(CH₂)_n CH₃]₂, where n and m are from 1 to about 10; and/or a modification at the 5' position: 5'-vinyl, 5'-methyl (R or S), a modification at the 4' position, 4'-S, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and any combination thereof.
25. A method of preparing a modified polypeptide comprising an unnatural amino acid, comprising:
- incubating an engineered host cell with a plurality of unnatural nucleotides, wherein the engineered host cell comprises a modified nucleoside triphosphate transporter and, optionally, a modified transposition-associated protein or transposable element; and


d) incorporating the plurality of unnatural nucleotides into one or more newly synthesized DNA strands, thereby generating the unnatural nucleic acid molecule;

wherein the modified transposition-associated protein or transposable element and the modified nucleoside triphosphate transporter increases retention of unnatural base pairs which facilitates incorporation of the plurality of unnatural nucleotides into the newly synthesized polypeptide to generate the modified polypeptide.

26. The method of claim 25, wherein the modified transposition-associated protein comprises comprises Insertion element IS1 4 protein InsB, Insertion element IS1 4 protein InsA, or a combination thereof; and wherein the modified transposable element comprises IS1.
27. The method of claim 25, wherein the modified nucleoside triphosphate transporter comprises a codon optimized nucleoside triphosphate transporter from *Phaeodactylum tricorutum* (PtNTT₂).
28. The method of claim 25, wherein the unnatural nucleotide comprises an unnatural base selected from the group consisting of 2-aminoadenin-9-yl, 2-aminoadenine, 2-F-adenine, 2-thiouracil, 2-thio-thymine, 2-thiocytosine, 2-propyl and alkyl derivatives of adenine and guanine, 2-amino-adenine, 2-amino-propyl-adenine, 2-aminopyridine, 2-pyridone, 2'-deoxyuridine, 2-amino-2'-deoxyadenosine 3-deazaguanine, 3-dezaadenine, 4-thio-uracil, 4-thio-thymine, uracil-5-yl, hypoxanthin-9-yl (I), 5-methyl-cytosine, 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 5-bromo, and 5-trifluoromethyl uracils and cytosines; 5-halouracil, 5-halocytosine, 5-propynyl-uracil, 5-propynyl cytosine, 5-uracil, 5-substituted, 5-halo, 5-substituted pyrimidines, 5-hydroxycytosine, 5-bromocytosine, 5-bromouracil, 5-chlorocytosine, chlorinated cytosine, cyclocytosine, cytosine arabinoside, 5-fluorocytosine, fluoropyrimidine, fluorouracil, 5,6-dihydrocytosine, 5-iodocytosine, hydroxyurea, iodouracil, 5-nitrocytosine, 5-bromouracil, 5-chlorouracil, 5-fluorouracil, and 5-iodouracil, 6-alkyl derivatives of adenine and guanine, 6-azapyrimidines, 6-azo-uracil, 6-azo cytosine, azacytosine, 6-azo-thymine, 6-thio-guanine, 7-methylguanine, 7-methyladenine, 7-deazaguanine, 7-deazaguanosine, 7-deaza-adenine, 7-deaza-8-azaguanine, 8-azaguanine, 8-azaadenine, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, and 8-hydroxyl substituted adenines and guanines; N4-ethylcytosine, N-2 substituted purines, N-6 substituted purines, O-6 substituted purines, those that increase the stability of duplex formation, universal nucleic acids, hydrophobic nucleic acids, promiscuous nucleic acids, size-expanded nucleic acids,

fluorinated nucleic acids, tricyclic pyrimidines, phenoxazine cytidine([5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps, phenoxazine cytidine (9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido [3',2':4,5]pyrrolo [2,3-d]pyrimidin-2-one), 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid, wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine and those in which the purine or pyrimidine base is replaced with a heterocycle.

29. The method of claim 25, wherein the unnatural base is selected from the group consisting of:



30. The method of claim 25, wherein the unnatural nucleotide further comprises an unnatural sugar moiety selected from the group consisting of a modification at the 2' position: OH; substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂F; O-alkyl, S-alkyl, N-alkyl; O-alkenyl, S-alkenyl, N-alkenyl; O-alkynyl, S-alkynyl, N-alkynyl; O-alkyl-O-alkyl, 2'-F, 2'-OCH₃, 2'-O(CH₂)₂OCH₃ wherein the alkyl, alkenyl and alkynyl may be substituted or

unsubstituted C₁-C₁₀, alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, -O[(CH₂)_n O]_mCH₃, -O(CH₂)_nOCH₃, -O(CH₂)_nNH₂, -O(CH₂)_nCH₃, -O(CH₂)_n-ONH₂, and -O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10; and/or a modification at the 5' position: 5'-vinyl, 5'-methyl (R or S), a modification at the 4' position, 4'-S, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and any combination thereof.