


PYRIDINE DERIVATIVES WITH UMAMI FLAVOUR

This disclosure relates to chemical compounds and to their use in flavouring.

- 5 Umami is a flavour sensation generally associated with Asian cuisine. In addition, improved umami taste helps make low salt products more palatable. Umami flavour has traditionally been achieved by the addition of monosodium glutamate (MSG) to foodstuffs. However, the presence of MSG in foodstuffs is not universally welcome, and there is an interest in the achievement of umami taste without MSG.

10

It has been found that certain compounds confer desirable umami taste sensations on compositions intended to be taken orally. There is therefore provided a compound (including salts thereof) according to Formula I



15

in which

R¹ is selected from H, methyl and ethyl;

R² is selected from H, OH, fluorine, C₁-C₄ linear or branched alkyl, C₁-C₆ alkoxy wherein the alkyl group is linear or branched, or comprises or consists of a C₃-C₅ cycloalkyl moiety;

- 20 R³ is selected from H, methoxy, methyl and ethyl;

or R² and R³ together form a bridging moiety -O-CH₂-O- between the phenyl carbon atoms to which they are connected;

R⁴ is selected from OH and methoxy; and

R⁵ and R⁶ are independently selected from H and methyl;

- 25 R¹, R², R³, R⁴, R⁵ and R⁶ being selected such that,

- (i) when R^2 and R^3 together form a bridging moiety $-O-CH_2-O-$ between the phenyl carbon atoms to which they are connected, R^1 , R^5 , R^6 are H, and R^4 is OH; and
- (ii) when R^4 is OH and R^1 - R^3 are H, at least one of R^5 , R^6 is methyl.

5 In a particular embodiment, R^3 is selected from H and methoxy.

In a particular embodiment, R^2 is selected from H, OH, fluorine, methyl, C_1 - C_6 alkoxy wherein the alkyl group is linear or branched, or comprises or consists of a C_3 - C_5 cycloalkyl moiety.


10

In a further particular embodiment, R^2 is selected from methyl, methoxy and isobutyloxy, R^3 is H, R^4 is OH and R^5 and R^6 are H.


In a further particular embodiment, the compound (including salts thereof) of Formula I is

15 selected from the group consisting of:

1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one;




1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one;




20







1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one; and




1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one.



5 Other particular embodiments include (but are not limited to):





5 The compounds may be made by methods well known to the art. Non-limiting examples of suitable methods include:

1. Claisen-Schmidt condensation of an aromatic aldehyde with a substituted acetophenone. The base used for this condensation is sodium methoxide in methanol. The reactions were
10 performed in an organic solvent (in particular, THF). The obtained intermediate was hydrogenated in methanol with hydrogen and 10% palladium on carbon as a catalyst.

2. Friedel-Craft acylation of the substituted phenol with 2-pyridinyl-propionic acid derivatives.

15

3. A one-step reaction of 2-hydroxyacetophenone with pyridin-2-ylmethanol and $\text{RuCl}_2(\text{Me}_2\text{SO})_4$ as a catalyst in dioxane solvent.

4. A one-step reaction of 2-hydroxyacetophenone with isonicotinaldehyde was used with
20 $\text{RuCl}_2(\text{PPh}_3)_3$ as a catalyst in dioxane solvent.

Not all of these methods will work for all compounds, but the selection of a particular method for a given compound is well known within the skill of the art.

The compounds hereinabove described may be used individually or in combinations, either with other such compounds, or other umami flavourants, or both. Other umami flavorants include, but are not limited to MSG, GMP (guanosine monophosphate) and IMP (inosine mono phosphate). There is therefore also provided a method of adding umami flavour to a consumable composition, comprising the addition to the composition of at least one compound of the Formula I.

By "consumable composition" is meant any composition that is taken into the mouth for ultimate spitting out or ingestion. The composition may be in any physical form, solid, liquid or gaseous. Non-limiting examples include all food products, food additives, nutraceuticals, pharmaceuticals and any product placed in the mouth including (but not limited to) chewing gum, oral care products, and oral hygiene products including but not limited to, cereal products, rice products, tapioca products, sago products, baker's products, biscuit products, pastry products, bread products, confectionery products, dessert products, gums, chewing gums, flavor or flavor-coated food/beverage containers, yeast products, baking-powder, salt and spice products, snack foods, savoury products, mustard products, vinegar products, sauces (condiments), soups, seasonings, ready-to-eat meals, gravies, nuts & nut products, tobacco products, cigars, cigarettes, processed foods, vegetable products, meat and meat products, egg products, milk and dairy products, yoghourts, cheese products, butter and butter substitute products, milk substitute products, soy products, edible oils and fat products, medicaments, beverages, carbonated beverages, alcoholic drinks such as beers, wines and spirits, non-alcoholic drinks such as soft drinks, including forms requiring reconstitution including, without limitation, beverage powder, milk based beverage powder, sugar-free beverage powder, beverage syrup, beverage concentrate, coffee and tea, food extracts, plant extracts, meat extracts, condiments, gelatins, pharmaceutical and non-pharmaceutical gums, tablets, lozenges, drops, emulsions, elixirs, syrups and other preparations for making beverages, and combinations thereof.

There is therefore provided an umami-flavored consumable composition comprising a product base and at least one compound as hereinabove described. By "product base" is

meant the combination of all the usual art-recognised ingredients required for the particular consumable composition.

The concentrations at which the compounds hereinabove described may be used will naturally vary, depending on the taste desired, the nature of the other ingredients and the desired end-use. Typically the concentration will vary between 0.01-100ppm, but it is to be emphasised that this is only an indication of a typical useful concentration, and that there will be circumstances in which it may be possible to formulate outside this range. More particular ranges are 0.1-50ppm and 0.2-10ppm.

10

The form in which the compounds hereinabove described may be used may vary. In addition to powders, liquids, solutions, dispersions or emulsions, they may be used in spray-dried or encapsulated forms.

15 The umami-flavoured consumable compositions may also contain any of the standard ingredients used in making such compositions, in art-recognised quantities. Non-limiting examples include food additives such as sweeteners, emulsifiers, yeast extracts, hydrolyzed vegetable proteins, short chain peptides, amino acids, fermented products, enzyme digests, Maillard reaction products and botanical extracts.

20

In addition to at least one of the subject compounds, the umami flavor may contain flavoring materials well known to those skilled in the art, including the materials on the FEMA GRAS list and the EU positive list. Flavor materials commonly provide aroma which in itself, can help improve the overall umami profile of the product. However, some flavor materials also have taste effects which, when used with the compounds hereinabove described, improve or enhance the umami taste sensation or improve the overall profile.

In addition, other materials on the approved list of flavor materials provide a umami taste or umami-like character or may enhance the overall umami taste. Some examples from the FEMA GRAS which may be used with the subject compounds include items, such as,


30

N1-(2-methoxy-4-methylbenzyl)-N2-(2-(pyridin-2-yl)ethyl)oxalamide (FEMA # 4231), N-(heptan-4-yl)benzothiazolidine-5-carboxamide (FEMA# 4232), N1-(2,4-dimethoxybenzyl)-N2-(2-(pyridin-2-yl)ethyl)oxalamide (FEMA# 4233), N1-(2-methoxy-4-methylbenzyl)-N2-(2-(5-methylpyridin-2-yl)ethyl)oxalamide (FEMA # 4234), N-gluconyl ethanolamine (FEMA # 4254), N-gluconyl ethanolamine phosphate (FEMA # 4255), N-lactoyl ethanolamine (FEMA # 4256), N-lactoyl ethanolamine phosphate (FEMA # 4257), N-3,7-dimethyl-2,6-octadienylcyclopropylcarboxamide (FEMA #4267) and gamma amino-butyric acid.

There now follows a series of non-limiting examples, which describe particular embodiments.

Example 1:

Synthesis of 1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one.



To a stirring solution of picolinaldehyde (1.52g, 14.1 mmol) in THF (50 ml) and 7 ml 25% wt NaOMe was added 1-(2-hydroxy-4,5-dimethylphenyl)ethanone in 5 ml THF dropwise at room temperature (2.33g, 14.1 mmol). The solution was continuously stirred at room temperature for 5 hours, and then water was added to the reaction flask. The aqueous layer was acidified to pH=7 using 1N HCl and extracted three times with EtOAc. The organic layers combined and dried (Na₂SO₄). The solvent was removed *in vacuo* and the residue chromatographed on silica gel (15% EtOAc/Hex) to yield 1.17g of yellow product (32%).

¹H NMR (300 MHz, CDCl₃): δ 12.64 (s, 1H), 8.75 (d, *J*=5.1 Hz, 1H), 8.29 (d, *J*=15.9 Hz, 1H), 7.88-7.78 (m, 3H), 7.52 (d, *J*=8.4 Hz, 1H), 7.35 (d, d, d, *J*=1.5, 6.0, 9.0 Hz, 1H), 6.85 (s, 3H), 2.31 (s, 3H), 2.28 (s, 3H).


In a 250 ml round-bottom flask, (E)-1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)prop-2-en-1-one (1.17g, 4.62 mmol) and 0.18 g of 10% Pd/C were mixed together in wet

EtOAc. Hydrogenation was conducted at atmospheric pressure, then the catalyst was filtered off. The crude product was purified by chromatography on silica gel (20% EtOAc/hexane) to provide 0.58g (47.7%) of final compound.

¹H NMR (300 MHz, CDCl₃): δ 12.12 (s, 1H), 8.55 (d, *J*=5.1 Hz, 1H), 7.62 (d, d, d, *J*=2.1, 7.5, 9.6 Hz, 1H), 7.56 (s, 1H), 7.27 (d, *J*=7.5 Hz, 1H), 7.14 (d, d, *J*=4.5, 5.7 Hz, 1H), 6.77 (s, 1H), 3.52 (t, *J*=7.8 Hz, 2H), 3.25 (t, *J*=7.5 Hz, 2H), 2.26 (s, 3H), 2.21 (s, 3H).

Example 2:

Synthesis of 1-(2,4-dihydroxyphenyl)-3-(pyridine-2-yl)propan-1-one.



10

A mixture of resorcinol(2.2g, 20 mmol), 3-pyridin-2-yl-propionic acid sulphuric acid salt (5g, 20 mmol) and 50 ml boron trifluoride diethyl etherate (BF₃·Et₂O) was stirred at 90C for 5 hours under N₂ (monitored by TLC). The reaction mixture was poured into 10% aqueous NaOAc solution and allowed to stand for 30 min and the solution was neutralized to pH=7 by 1M HCl and was extracted with EtOAc (3X50ml). The combined EtOAc layer was washed with water, brine and dried over Na₂SO₄. The residue obtained after evaporation of the solvent was chromatographed over silica gel column using hexane-EtOAc mixture (1:1) as eluent to give mixture of products, which was then recrystallized in hot MTBE/Hex to afford final product (24%) with >97% purity.

20 ¹H NMR (300 mHz, CD₃OD) δ 8.46-8.44 (d,d,d , *J*=5.1, 1.8, 0.9 Hz, 1H), 7.78-7.72 (d, d, d, *J*=9.9, 7.8, 2.1 Hz, 1H), 7.78-7.75 (d, *J*=8.7 Hz, 1H), 7.40-7.37 (d, *J*=8.1 Hz, 1H), 7.29-7.23 (d,d, d, *J*=6.3, 4.8, 1.2 Hz, 1H), 6.38-6.34 (d, d, *J*=9.0, 2.4 Hz, 1H), 6.26-6.25 (d, *J*=2.4 Hz, 1H), 3.41 (t, *J*=7.5 Hz, 2H), 3.18 (t, *J*=8.4 Hz, 2H).

25 Example 3:

1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one.

The title compound was synthesized using the same procedure as described in example 1.

¹H NMR (300 MHz, CDCl₃): δ 12.29 (s, 1H), 8.53 (d, *J*=4.5 Hz, 1H), 7.63 (d, d, d, *J*=3.0, 9.6, 11.7 Hz, 1H), 7.26 (d, *J*=7.2 Hz, 1H), 7.14 (d, d, d, *J*=1.2, 5.4, 7.8 Hz, 1H), 6.79 (s, 1H), 6.71 (d, d, *J*=1.8, 8.4 Hz, 1H), 3.53 (t, *J*=7.2 Hz, 2H), 3.24 (t, *J*=7.2 Hz, 2H), 2.36 (s, 3H).

5 Example 4:

1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one.

The title compound was synthesized using the same procedure as described in example 1.

¹H NMR (300 MHz, CDCl₃): δ 12.77 (s, 1H), 8.53 (d, *J*=4.8 Hz, 1H), 7.72 (d, d, *J*=2.7, 8.7 Hz, 1H), 7.60 (d, d, d, *J*=2.1, 7.5, 9.6 Hz, 1H), 7.23 (d, *J*=8.1 Hz, 1H), 7.12 (d, d, d, *J*=1.5, 4.8, 6.3 Hz, 1H), 6.44-6.41 (m, 2H), 3.83 (s, 3H), 3.45 (t, *J*=7.5 Hz, 2H), 3.22 (t, *J*=7.8 Hz, 2H).

Example 5:

1-(2-hydroxyphenyl)-3-(5-methylpyridin-2-yl)propan-1-one.

15 The title compound was synthesized using the same procedure as described in example 1.

¹H NMR (300 MHz, CDCl₃): δ 12.25 (s, 1H), 8.36 (d, *J*=3.0 Hz, 1H), 7.85 (d, d, *J*=1.8, 8.4 Hz, 1H), 7.49-7.43 (m, 2H), 7.20 (d, *J*=9.0 Hz, 1H), 6.97 (d, d, *J*=2.1, 8.1 Hz, 1H), 6.89 (d, d, *J*=7.5, 8.1 Hz, 1H), 3.57 (t, *J*=6.9 Hz, 2H), 3.23 (t, *J*=6.9 Hz, 2H), 2.32 (s, 3H).

20 Example 6:

1-(4-ethoxy-2-hydroxyphenyl)-3-(pyridine-2-yl)propan-1-one.

The title compound was synthesized using the same procedure as described in example 1.

¹H NMR (300 MHz, CDCl₃): δ 12.71 (s, 1H), 8.53 (d, *J*=5.4 Hz, 1H), 7.73 (d, *J*=8.7 Hz, 1H), 7.63 (d, d, d, *J*=1.8, 7.5, 9.6 Hz, 1H), 7.28 (d, *J*=8.1 Hz, 1H), 7.15 (d, d, *J*=4.8, 6.3 Hz, 1H), 6.44-6.40 (m, 2H), 4.06 (q, *J*=7.5 Hz, 2H), 3.48 (t, *J*=7.8 Hz, 2H), 3.24 (t, *J*=7.8 Hz, 2H), 1.42 (t, *J*=7.5 Hz, 3H).

Example 7:

1-(2-hydroxy-3,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one.

30 The title compound was synthesized using the same procedure as described in example 1.

¹H NMR (300 MHz, CDCl₃): δ 12.40 (s, 1H), 8.55 (bd, *J*=5.2 Hz, 1H), 7.64 (d, d, d, *J*=1.8, 8.1, 10.2 Hz, 1H), 7.49 (s, 1H), 7.29 (d, *J*=8.1 Hz, 1H), 7.18-7.14 (m, 2H), 3.57 (t, *J*=7.5 Hz, 2H), 3.26 (t, *J*=7.8 Hz, 2H), 2.28 (s, 3H), 2.24 (s, 3H).

5 Example 8:

1-(2-hydroxy-5-methylphenyl)-3-(pyridine-2-yl)propan-1-one.

The title compound was synthesized using the same procedure as described in example 1.

¹H NMR (300 MHz, CDCl₃): δ 12.09 (s, 1H), 8.52 (d, *J*=4.8 Hz, 1H), 7.62-7.57 (m, 2H), 7.28-7.23 (m, 2H), 7.12 (d, d, *J*=4.8, 7.5 Hz, 1H), 6.86 (d, *J*=8.4 Hz, 1H), 3.54 (t, *J*=6.9 Hz, 2H), 3.23 (t, *J*=6.9 Hz, 2H), 2.28 (s, 3H).

Example 9:

1-(5-ethyl-2-hydroxyphenyl)-3-(pyridine-2-yl)propan-1-one.

The title compound was synthesized using the same procedure as described in example 1.

15 ¹H NMR (300 MHz, CDCl₃): δ 12.11 (s, 1H), 8.54 (d, *J*=3.3 Hz, 1H), 7.67-7.61 (m, 2H), 7.32 (d, d, *J*=2.1, 8.1 Hz, 1H), 7.16 (d, d, *J*=6.3, 8.4 Hz, 1H), 6.92 (d, *J*=8.7 Hz, 1H), 3.59 (t, *J*=6.6 Hz, 2H), 3.27 (t, *J*=6.6 Hz, 2H), 2.61 (q, *J*=8.7 Hz, 2H), 1.23 (t, *J*=7.5 Hz, 3H).

Example 10:

20 1-(2-hydroxyphenyl)-3-(3-methylpyridin-2-yl)propan-1-one.

The title compound was synthesized using the same procedure as described in example 1.

¹H NMR (300 MHz, CDCl₃): δ 12.31 (s, 1H), 8.34 (d, *J*=6.3 Hz, 1H), 7.91 (d, *J*=6.3 Hz, 1H), 7.49-7.44 (m, 2H), 7.09-7.04 (m, 1H), 6.98 (d, *J*=9.0 Hz, 1H), 6.91 (d, d, *J*=7.5, 7.5 Hz, 1H), 3.61 (t, *J*=7.5 Hz, 2H), 3.23 (t, *J*=7.5 Hz, 2H), 2.39 (s, 3H).

25

Example 11


1-(2-hydroxy-3,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one.

The title compound was synthesized using the same procedure as described in example 2.

30 ¹H NMR (300 MHz, CDCl₃): δ 12.40 (s, 1H), 8.55 (bd, *J*=5.2 Hz, 1H), 7.64 (d, d, d, *J*=1.8, 8.1, 10.2 Hz, 1H), 7.49 (s, 1H), 7.29 (d, *J*=8.1 Hz, 1H), 7.18-7.14 (m, 2H), 3.57 (t, *J*=7.5 Hz, 2H), 3.26 (t, *J*=7.8 Hz, 2H), 2.28 (s, 3H), 2.24 (s, 3H).

Example 12:

1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one.



5 Step1: To a stirred solution of 1-(2,4-dihydroxyphenyl)ethanone (3.2 g, 21.03 mmol) and K_2CO_3 in 100 mL of acetone at room temperature, the 1-iodo-2-methylpropane (3.6 ml, 21.03 mmol) was added dropwise. The mixture was refluxed for 8 hrs, and then the reaction was kept in 54°C for overnight. The reaction was cooled down to room temperature and quenched by 1N HCl to pH=7. The aqueous layer was extracted 3X with EtOAc, the organic
10 layers were combined, washed with water, brine and dried (Na_2SO_4). The solvent was removed *in vacuo* and the residue chromatographed on silica gel 5% EtOAc/Hex to give 1.1g (25%) of product.

1H NMR (300 MHz, $CDCl_3$): δ 12.67 (s, 1H), 7.55 (d, J =8.7 Hz, 1H), 6.36 (d, d, J =2.4, 8.7 Hz, 1H), 6.32 (d, J =2.4 Hz, 1H), 3.68 (d, J =6.6 Hz, 2H), 2.48 (s, 3H), 2.09-2.195 (m, 1H),
15 0.96 (s, 3H), 0.94 (s, 3H).

Step 2: To a stirred solution of picolinaldehyde (0.6 ml, 5.28 mmol) and NaOMe in Tetrahydrofuran (50 ml) and 1-(2-hydroxy-4-isobutoxyphenyl)ethanone in 5 ml THF was added at room temperature. The mixture was stirred at room temperature for 5 h, 1N HCl
20 was added until pH=7. The aqueous layer was extracted 3x with EtOAc. The organic layers combined, washed with water, brine and dried (Na_2SO_4). The solvent was removed *in vacuo* and the residue chromatographed on silica gel (25% EtOAc in Hex) to give 0.8g (51%) of yellow solid.

1H NMR (300 MHz, $CDCl_3$): δ 13.37 (s, 1H), 8.72 (d, J =5.1 Hz, 1H), 8.22 (d, J =15.0 Hz, 1H), 7.97 (d, J =9.0 Hz, 1H), 7.84 (d, J =15.0 Hz, 1H), 7.77 (d, d, J =1.8, 7.2 Hz, 1H), 7.50 (d, J =8.1 Hz, 1H), 7.33 (d, d, J =4.8, 7.8 Hz, 1H), 6.51 (d, d, J =2.4, 9.0 Hz, 1H), 6.47 (d, J =2.7 Hz, 1H), 3.80 (d, J =6.6 Hz, 2H), 2.20-2.07 (m, 1H), 1.60 (s, 3H), 1.05 (s, 3H).

Step 3: In a 200 mL round-bottomed flask, (E)-1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridin-2-yl)prop-2-en-1-one (1.08 g, 3.63 mmol) and 80mg of 10% Pd/C was mixed together in wet EtOAc (50 ml). Hydrogenation was conducted at atmospheric pressure room temperature until the starting material was gone. The catalyst was filtered and the crude was purified by chromatography on silica gel (25% EtOAc/hexane) to provide 0.92g of final compound (82%) as off white solid.

¹H NMR (300 MHz, CDCl₃): δ 12.73 (s, 1H), 8.52 (d, *J*=4.8 Hz, 1H), 7.72 (d, *J*=9.0 Hz, 1H), 7.61 (d, d, d, *J*=2.4, 7.8, 9.6 Hz, 1H), 7.25 (d, *J*=7.8 Hz, 1H), 7.13 (d, d, d, *J*=1.5, 4.8, 6.0 Hz, 1H), 6.43 (d, d, *J*=2.1, 8.4 Hz, 1H), 6.39 (d, *J*=2.7 Hz, 1H), 3.75 (d, *J*=7.2 Hz, 2H), 3.46 (t, *J*=7.2 Hz, 2H), 3.23 (t, *J*=6.9 Hz, 2H), 2.16-2.02 (m, 1H), 1.02 (d, *J*=7.2 Hz, 6H).

Example 13:

1-(4-cyclopropylmethoxy)-2-hydroxyphenyl)-3-(pyridine-2-yl)propan-1-one.

The title compound was synthesized using the same procedure as described in example 11.

¹H NMR (300 MHz, CDCl₃): δ 12.61 (s, 1H), 8.42 (d, *J*=5.7 Hz, 1H), 7.52 (d, d, d, *J*=1.8, 7.8, 9.6 Hz, 1H), 7.16 (d, *J*=7.5 Hz, 1H), 7.04 (d, d, *J*=4.8, 7.2 Hz, 1H), 6.34 (d, d, *J*=2.7, 9.0 Hz, 1H), 6.28 (d, *J*=2.4 Hz, 1H), 3.73 (d, *J*=6.9 Hz, 2H), 3.37 (t, *J*=7.2 Hz, 2H), 3.13 (t, *J*=7.5 Hz, 2H), 1.24-1.11 (m, 1H), 0.59-0.53 (m, 2H), 0.29-0.23 (m, 2H).

Example 14:


1-(4-(cyclopentyloxy)-2-hydroxyphenyl)-3-(pyridine-2-yl)propan-1-one.

The title compound was synthesized using the same procedure as described in example 11.

¹H NMR (300 MHz, CDCl₃): δ 12.74 (s, 1H), 8.53 (d, *J*=4.8 Hz, 1H), 7.74-7.70 (m, 1H), 7.62 (d, d, d, *J*=2.1, 8.1, 9.9 Hz, 1H), 7.26 (d, *J*=7.8 Hz, 1H), 7.14 (d, d, d, *J*=1.2, 4.8, 7.5 Hz, 1H), 6.41-6.37 (m, 2H), 4.80-4.76 (m, 1H), 3.47 (t, *J*=7.5 Hz, 2H), 3.23 (t, *J*=7.5 Hz, 2H), 2.01-1.78 (m, 8 H).

Example 15:

1-(4-hydroxybenzo[d][1,3]dioxol-5-yl)-3-(pyridin-2-yl)propan-1-one.



Step 1

- 1-(4-hydroxybenzo[d][1,3]dioxol-5-yl)ethanone (1.8 g, 9.99 mmol) was dissolved in THF (40 ml) to give a colorless solution. Sodium methanolate 30% (5 g, 27.8 mmol) was added.
- 5 Reaction mixture became dark green colored. After 15 min. stirring at room temperature 1.1 g (10 mmol) of picolinaldehyde dissolved in 10 ml THF was added dropwise during 10 min. The addition was started at 22 °C and was slightly exothermic. Stirring at room temperature was continued for another 3 hrs. Mixture was diluted with diethyl ether and the pH was adjusted to pH 6 with a concentrated hydrochloric acid solution while vigorously stirring.
- 10 The color changes from dark green to orange/red. After 5 minutes the formed solids were filtered. Solids were stirred with 100 mL of methanol and the remaining solids (mainly NaCl) were filtered again. The filtrate was evaporated and the residue was taken up in a mixture of diethylether and methanol. Solids were filtered and dried to yield 1.7 g of an orange powder. To remove the small amounts of impurities the orange product was stirred
- 15 with chloroform for 10 minutes, filtered and dried to yield 1.5 g (47%) of the (E)-1-(4-hydroxybenzo[d][1,3]dioxol-5-yl)-3-(pyridin-2-yl)prop-2-en-1-one.
- ¹H-NMR in DMSO-d₆: 6.17(2H, O-CH₂-O, s), 6.72(1H, CO-CH=, d), 7.60(1H, aromatic, m),
- 7.83(1H, aromatic, d), 7.94(1H, CO-CH=CH-, d), 8.09-8.13(2H, aromatic, m),
- 20 8.35(1H, aromatic, d), 8.75(1H, aromatic N-CH, d), 9.83(1H, OH, s)

The 1-(4-hydroxybenzo[d][1,3]dioxol-5-yl)ethanone was prepared by the reaction of trihydroxy acetophenone with diiodomethane.

25 Step 2

(E)-1-(4-hydroxybenzo[d][1,3]dioxol-5-yl)-3-(pyridin-2-yl)prop-2-en-1-one (0.5 g, 1.857 mmol) was added to methanol (50 ml) to give a yellow solution. Palladium on carbon (0.05 g) was added. Product was hydrogenated while stirring till conversion was completed (checked by TLC). The reaction mixture was filtered, evaporated and the solid residue was

washed with diethylether, and dried to yield 0.3 g (57%) of the title compound.

¹H-NMR in DMSO-d₆: 3.11(2H, CH₂-C=N, t), 3.49(2H, CO-CH₂-, t), 6.13(2H, O-CH₂-O, s),


6.64(1H, aromatic, d), 7.19(1H, aromatic, m), 7.33(1H, aromatic, d), 7.67-7.70(2H, aromatic,

5 m),

8.44(1H, aromatic N-CH, d), 9.83(1H, OH, s)

Example 16

1-(2-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one.



1-(2-methoxyphenyl)ethanone (3 g, 19.98 mmol), pyridin-2-ylmethanol (2.180 g, 19.98 mmol), and potassium hydroxide (1.121 g, 19.98 mmol) were added to dioxane (50 ml) to give a yellow suspension. RuCl₂(Me₂SO)₄ (0.2 g) was added. Reaction mixture was heated at 80°C for 1 hour.

15

Reaction mixture was filtered through Silicagel to remove catalyst and KOH. The filtrate was evaporated and the residue was purified by column chromatography.

Eluent: ethylacetate: heptane = 3:1. 0.2 g (4%) of the title compound was obtained.

¹H-NMR in chloroform, d: 3.20(2H, CO-CH₂-, t), 3.47(2H, -CH₂-Ar, t), 3.87(3H, -OCH₃, s),

20 6.95(2H, aromatic, m), 7.1(1H, aromatic, m), 7.23(1H, aromatic, d), 7.45(1H, aromatic, m)

7.58(1H, aromatic, m), 7.70(1H, aromatic, d), 8.50(1H, aromatic N-CH=, d)

Example 17

Testing of compounds

25 Various compounds prepared in the previous examples were tested.

Two solutions were prepared:

A: a solution of 0.3% NaCl,

B: a solution of 0.3% NaCl and 2 ppm compound as per Table 1.

The samples were tasted by a trained sensory panel composed of 20 women aged between 30 and 60. Table 1

Compound	Taste description
1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one	Umami, sweet, licorice, savoury
1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one	umami, sweet, salty, longlasting savouriness
1-(2-hydroxyphenyl)-3-(5-methylpyridin-2-yl)propan-1-one	Slightly umami, some green aftertaste, floral
1-(4-ethoxy-2-hydroxyphenyl)-3-(pyridine-2-yl)propan-1-one	Umami, floral
1-(4-cyclopropylmethoxy)-2-hydroxyphenyl)-3-(pyridine-2-yl)propan-1-one	Umami
1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one	Umami, sweet, salty, bouillon, lingering, intense
1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one	Umami, sweet, savoury, salty
1-(2-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one	Umami, slightly bitter
1-(2-hydroxyphenyl)-3-(3-methylpyridin-2-yl)propan-1-one	Slightly umami, savoury, meaty
1-(4-hydroxybenzo[d][1,3]dioxol-5-yl)-3-(pyridin-2-yl)propan-1-one	Clearly umami, slightly yeasty, bouillon
1-(2-hydroxy-3,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one	Umami
1-(4-(cyclopentoxo)-2-hydroxyphenyl)-3-(pyridine-2-yl)propan-1-one	Umami
1-(2-hydroxy-5-methylphenyl)-3-(pyridine-2-yl)propan-1-one	Umami
1-(2-methoxyphenyl)-3-(pyridine-2-yl)propan-1-one	Umami, slightly bitter
1-(5-ethyl-2-hydroxyphenyl)-3-(pyridine-2-yl)propan-1-one	Umami, slightly bitter
1-(2-hydroxy-3,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one.	Umami

Example 18

5 Comparative test with MSG

Two solutions were prepared:

A: a solution of 0.3% NaCl and 0.03% MSG,

B: a solution of 0.3% NaCl, 0.03% MSG and 5 ppm of 1-(4-hydroxybenzo[d][1,3]dioxol-5-yl)-3-(pyridin-2-yl)propan-1-one.

10

The samples were tasted by a trained sensory panel composed of 20 women aged between 30 and 60.

Solution A: salty, umami.

5 Solution B: salty, umami, more bouillon-like, more complex, less sweet.

Example 19

Tomato soup formulation

10 A tomato soup mix was prepared from 9.4 g sodium chloride, 1 g MSG, 0.08 g ribonucleotides (ex yeast), 32 g tomato powder (ex Sprea), 25.1 g glucose, 21 g starch (UltrasperseTM 5 ex National Starch), 5 g palm fat powder, 3 g yeast powder, 1 g onion powder, 0.15 g carrot powder, 0.05 g ground white pepper, 0.3 g celery extract powder, 0.05 g ground laurel leaf powder, and 1.85 g sucrose. 25 g of the well-mixed ingredients was
15 added to 250 g of boiling water and stirred until completely dissolved, to give a reference soup.

The reference soup was compared with a batch of the same soup containing 1.5 ppm of 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one. A small group of flavourists (2
20 male, 2 female) tasted the soups and agreed that the test soup was more umami, had a clear lingering effect, was more salty and was more complex than the reference soup.

Example 20

Potato chips

25

Plain potato chips were prepared.

One part was flavoured with 1.2 % sodium chloride (sample A),

One part was flavoured with 1.2 % sodium chloride and 0.3 % MSG (sample B),

One part was flavoured with 1.2 % sodium chloride and 2 ppm of 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one (sample C).
30

The samples were tasted by a trained sensory panel composed of 20 women aged between 30 and 60.

The panel preferred sample C over the other two samples.

Sample A was described as salty, sample B as salty and umami and sample C as salty, strong
5 umami, long lasting, savoury, bouillon.

Example 21

Tomato ketchup

10 A tomato ketchup was prepared from 19 % tomato paste (28-30 % dry weight), 8 % vinegar (15 %), 3 % sodium chloride, 20 % sugar and 50 % water.

To one-half of the batch was added 2 ppm of 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one.

A small group of flavourists (2 male, 2 female) tasted the ketchups and agreed that the test
15 ketchup tasted clearly more umami, more salty and had a pleasant savoury note compared with that of the ketchup without the compound.

Examples 22-24

The following compositions are made and tested. Using a bench-top tasting panel (consisting
20 of 10 panelists – flavorists, application experts), panelists were asked to record the sensory attribute differences (umami) between the samples. All samples were tasted blind and ranked on umami.

Example 22

25 Commercial Organic Chicken Broth (Swanson's Certified Organic Chicken Broth – No added MSG, contains about 0.4% sugars)

Control

<u>Ingredient</u>	<u>% (by weight)</u>
Commercial Organic Chicken Broth (550 mg Na/240 mL serving)	100

Reference

<u>Ingredient</u>	<u>% (by weight)</u>
MSG	0.05
Commercial Organic Chicken Broth (550 mg Na/240 mL serving)	Balance to 100

Ingredient 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one;	0.001
Commercial Organic Chicken Broth (550 mg Na/240 mL serving)	Balance to 100

- 5 *Sensory findings:* Panelists preferred 0.001% of 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one in this base. In a blind ranking of 6 different samples MSG was the highest umami with 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one finishing 2nd and Control finishing 5th.

10 Ingredient 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one	0.002
Commercial Organic Chicken Broth (550 mg Na/240 mL serving)	Balance to 100

Sensory findings: Panelists preferred 0.002% of 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one in this base. In a blind ranking of 6 different samples MSG was the highest umami with 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one finishing 3rd and Control finishing 5th.

Ingredient 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one	0.002
Commercial Organic Chicken Broth (550 mg Na/240 mL serving)	Balance to 100

Sensory findings: Panelists preferred 0.002% of 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one in this base. In a blind ranking of 6 different samples MSG was the highest umami with 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one finishing 5th and Control finishing 5th.

Example 23

Model Chicken Broth -- no MSG added, no sugar added

- 10 The model chicken broth powder was prepared from 11.9 g chicken fat, 23.81 g salt, 14.29 g sugar, 0.95 g onion powder, 49.01 g chicken powder, 0.04 g turmeric powder. To prepare the model chicken broth 2.0 g of powder was mixed 98.0 g hot water.

Control

<u>Ingredient</u>	<u>% (by weight)</u>
Model Chicken Broth (653 mg Na/240 mL serving)	100

15

Reference

<u>Ingredient</u>	<u>% (by weight)</u>
MSG	0.05
Model Chicken Broth (653 mg Na/240 mL serving)	Balance to 100

Ingredient 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one	0.0005
Model Chicken Broth (653 mg Na/240 mL serving)	Balance to 100

Sensory findings: Panelists preferred 0.0005% of 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one in this base. In a blind ranking of 6 different samples MSG was the highest umami with 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one finishing 3rd and Control finishing 5th.

Ingredient 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one	0.001
Model Chicken Broth (653 mg Na/240 mL serving)	Balance to 100

Sensory findings: Panelists preferred 0.001% of 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one in this base. In a blind ranking of 6 different samples MSG was the highest umami with 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one finishing 4th and Control finishing 5th.

Ingredient 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one	0.001
Model Chicken Broth (653 mg Na/240 mL serving)	Balance to 100

Sensory findings: Panelists preferred 0.001% of 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one in this base. In a blind ranking of 6 different samples MSG was the

highest umami with 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one finishing 2nd and Control finishing 5th.

Example 24

5 Model Chicken Broth – no MSG added, 0.3% sugar added

The model chicken broth powder was prepared from 11.9 g chicken fat, 23.81 g salt, 14.29 g sugar, 0.95 g onion powder, 49.01 g chicken powder, 0.04 g turmeric powder. To prepare the model chicken broth 2.0 g of powder was mixed 98.0 g hot water.

10

Control

<u>Ingredient</u>	<u>% (by weight)</u>
Model Chicken Broth – sugar added (653 mg Na/240 mL serving)	100

Reference

<u>Ingredient</u>	<u>% (by weight)</u>
MSG	0.05
Model Chicken Broth – sugar added (653 mg Na/240 mL serving)	Balance to 100

15 Ingredient 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one	0.0010
Model Chicken Broth – sugar added (653 mg Na/240 mL serving)	Balance to 100

Sensory findings: Panelists preferred 0.001% of 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one in this base. In a blind ranking of 6 different samples MSG was the highest umami with 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one finishing 5th and Control finishing 6th.

Ingredient: 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one	0.002
Model Chicken Broth – sugar added (653 mg Na/240 mL serving)	Balance to 100

Sensory findings: Panelists preferred 0.002% of compound 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one in this base. In a blind ranking of 6
5 different samples MSG was the highest umami with 1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one finishing 4th and Control finishing 6th.

Ingredient 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one	0.002
Model Chicken Broth – sugar added (653 mg Na/240 mL serving)	Balance to 100

Sensory findings: Panelists preferred 0.002% of 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-
10 2-yl)propan-1-one in this base. In a blind ranking of 6 different samples MSG was the highest umami with 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one finishing 3rd and Control finishing 6th.


Ingredient 1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one

<u>Ingredient</u>	<u>% (by weight)</u>
1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one	0.002
Model Chicken Broth – sugar added (653 mg Na/240 mL serving)	Balance to 100

15 *Sensory findings:* Panelists preferred 0.002% of 1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one in this base. In a blind ranking of 6 different samples MSG was

the highest umami with 1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one finishing 2nd and Control finishing 6th. – in this base at this level. 1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one performed well when evaluating for umami.

- 5 In certain embodiments there is provided a compound (including salts thereof) according to Formula I



in which

R¹ comprises H, methyl or ethyl;

- 10 R² comprises H, OH, fluorine, C₁-C₄ linear or branched alkyl, C₁-C₆ alkoxy wherein the alkyl group is linear or branched, or a C₃-C₅ cycloalkyl moiety;

R³ comprises H, methoxy, methyl or ethyl;

or R² and R³ together form a bridging moiety –O-CH₂-O- between the phenyl carbon atoms to which they are connected;

- 15 R⁴ comprises OH or methoxy; and

R⁵ and R⁶ independently comprise H or methyl;

R¹, R², R³, R⁴, R⁵ and R⁶ are such that,

- (i) when R² and R³ together form a bridging moiety –O-CH₂-O- between the phenyl carbon atoms to which they are connected, R¹, R⁵, R⁶ are H, and R⁴ is OH; and
 20 (ii) when R⁴ is OH and R¹-R³ are H, at least one of R⁵ or R⁶ is methyl.

In a particular embodiment, R³ is selected from H and methoxy.

- In a particular embodiment, R² comprises H, OH, fluorine, methyl, C₁-C₆ alkoxy wherein the
 25 alkyl group is linear or branched, or a C₃-C₅ cycloalkyl moiety.

In other particular embodiments, R^2 consists of a C_3 - C_5 cycloalkyl moiety.


In other particular embodiments, R^1 comprises H or methyl, R^2 comprises methyl, methoxy
5 or isobutyloxy, R^3 is H, R^4 is OH, and R^5 and R^6 are H.

These compounds confer desirable umami taste sensations on compositions intended to be taken orally, as disclosed above.

10 Although the embodiments have been described in detail through the above description and the preceding examples, these examples are for the purpose of illustration only and it is understood that variations and modifications can be made by one skilled in the art without departing from the scope of the disclosure. It should be understood that the embodiments described above are not only in the alternative, but can be combined.

Claims:

1. A compound (including salts thereof) according to Formula I



5 in which

R^1 comprises H, methyl or ethyl;

R^2 comprises H, OH, fluorine, C_1 - C_4 linear or branched alkyl, C_1 - C_6 alkoxy wherein the alkyl group is linear or branched, or a C_3 - C_5 cycloalkyl moiety;

R^3 comprises H, methoxy, methyl or ethyl;

10 or R^2 and R^3 together form a bridging moiety $-O-CH_2-O-$ between the phenyl carbon atoms to which they are connected;

R^4 comprises OH or methoxy; and

R^5 and R^6 independently comprise H or methyl;

R^1 , R^2 , R^3 , R^4 , R^5 and R^6 are such that,

15 (i) when R^2 and R^3 together form a bridging moiety $-O-CH_2-O-$ between the phenyl carbon atoms to which they are connected, R^1 , R^5 , R^6 are H, and R^4 is OH; and

(ii) when R^4 is OH and R^1 - R^3 are H, at least one of R^5 or R^6 is methyl.


2. The compound according to claim 1, in which R^3 is selected from H and methoxy.

20

3. The compound according to claim 1, in which R^2 comprises H, OH, fluorine, methyl, C_1 - C_6 alkoxy wherein the alkyl group is linear or branched, or a C_3 - C_5 cycloalkyl moiety.

4. The compound according to any one of claims 1 through 3, in which R² consists of a C₃-C₅ cycloalkyl moiety.
5. The compound according to any of claims 1 through 4, in which R¹ comprises H or methyl, R² comprises methyl, methoxy or isobutyloxy, R³ is H, R⁴ is OH, and R⁵ and R⁶ are H.
6. The compound according to claim 5, selected from the group consisting of:
1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one;
10 1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one;
1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one;
1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one;
and mixtures thereof.
- 15 7. A method of providing umami flavour to a consumable composition, comprising the addition to the composition of at least one compound according to any of claims 1 through 6.
8. A consumable composition having an umami flavour, comprising a product base and
20 at least one compound according to any of claims 1 through 6.
9. A consumable composition according to claim 8, in which the compound is present at a concentration of from about 0.01 to about 100ppm, optionally from about 0.1 to about 50ppm and further optionally from about 0.2 to about 10ppm.

10. A compound (including salts thereof) according to Formula I



in which

- 5 R^1 is selected from H, methyl and ethyl;
 R^2 is selected from H, OH, fluorine, C_1 - C_4 linear or branched alkyl, C_1 - C_6 alkoxy wherein the alkyl group is linear or branched, or comprises or consists of a C_3 - C_5 cycloalkyl moiety;
 R^3 is selected from H, methoxy, methyl and ethyl;
 10 or R^2 and R^3 together form a bridging moiety $-O-CH_2-O-$ between the phenyl carbon atoms to which they are connected;
 R^4 is selected from OH and methoxy; and
 R^5 and R^6 are independently selected from H and methyl;
 R^1 , R^2 , R^3 , R^4 , R^5 and R^6 being selected such that,
 15 (i) when R^2 and R^3 together form a bridging moiety $-O-CH_2-O-$ between the phenyl carbon atoms to which they are connected, R^1 , R^5 , R^6 are H, and R^4 is OH; and
 (ii) when R^4 is OH and R^1 - R^3 are H, at least one of R^5 , R^6 is methyl.
11. The compound according to claim 10, in which R^3 is selected from H and methoxy.
- 20
12. The compound according to claim 10, in which R^2 is selected from H, OH, fluorine, methyl, C_1 - C_6 alkoxy wherein the alkyl group is linear or branched, or comprises or consists of a C_3 - C_5 cycloalkyl moiety.

13. The compound according to claim 10, in which R¹ is selected from H and methyl, R² is selected from methyl, methoxy and isobutyloxy, R³ is H, R⁴ is OH and R⁵ and R⁶ are H.
- 5 14. The compound according to claim 13, selected from the group consisting of:
1-(2-hydroxy-4,5-dimethylphenyl)-3-(pyridine-2-yl)propan-1-one;
1-(2-hydroxy-4-methylphenyl)-3-(pyridin-2-yl)propan-1-one;
1-(2-hydroxy-4-methoxyphenyl)-3-(pyridin-2-yl)propan-1-one; and
1-(2-hydroxy-4-isobutoxyphenyl)-3-(pyridine-2-yl)propan-1-one.
- 10
15. A method of providing umami flavour to a consumable composition, comprising the addition to the composition of at least one compound according to claim 10.
16. A consumable composition having an umami flavour, comprising a product base and
15 at least one compound according to claim 10.
17. The consumable composition according to claim 16, in which the compound is present at a concentration of from 0.01-100ppm, particularly from 0.1-50ppm and more particularly from 0.2-10ppm.