FIGURE 1





Control
Metronidazole (10µg/ml)
Isoniazid (10µg/ml)
final compound 12 (10µg/ml)
final compound 12 (1µg/ml)
final compound 12 (0.1µg/ml)
Drug: Anaerobic
No drug: Aerobic
Log 10 [mRLU]
0
1
2
3
4
5
6
-1
0
1
2
3
4
5
Days after Anaerobiosis

FIGURE 2

A)


Day 2 after Anaerobiosis
Log 10 CFU/ml
0
1
2
3
4
5
6
7
8
Control
Metronidazole (10µg/ml)
Isoniazid (10µg/ml)
Final compound 12 (10µg/ml)
Final compound 12 (1µg/ml)
Final compound 12 (0.1µg/ml)

B)


Days 5 after Anaerobiosis
Log10 CFU/ml
0
1
2
3
4
5
6
7
8
9
10
Control
Metronidazole (10µg/ml)
Isoniazid (10µg/ml)
Final compound 12 (10µg/ml)
Final compound 12 (1µg/ml)
Final compound 12 (0.1µg/ml)



Effect of various drugs on dormant M. tuberculosis (Wayne model); n=2;
log10 CFU/ml
0
2
4
6
8
Control
Metronidazole 100 µg/ml
Isoniazid 10 µg/ml
Moxifloxacin 10 µg/ml
Moxifloxacin 1 µg/ml
Final compound 12 10 µg/ml
Final compound 12 1 µg/ml
Rifampicin 10 µg/ml
Rifampicin 1 µg/ml

FIGURE 3