

NOVEL PROCESSES FOR PRODUCING SOVAPREVRIR


CROSS REFERENCE TO RELATED APPLICATION

[0001] This application claims priority from U.S. Provisional Application No. 61/784,182, filed March 14, 2013, which is hereby incorporated by reference in its entirety.

BACKGROUND

[0002] Sovaprevir is a hepatitis C virus NS3 protease inhibitor, effective for treating HCV infection in humans.

[0003] Sovaprevir can be prepared by the method presented in US 7,906,619, Example 1.



SUMMARY

[0004] The disclosure provides methods of preparing Sovaprevir. These methods are designated Methods 1 and 2 in the EXAMPLES section of the application. A number of the steps in the methods are unique as are the overall methods for preparing Sovaprevir. The disclosure provides the steps occurring after the synthesis of Compound **13** in synthetic Method 1 to form Sovaprevir. The disclosure also provides the addition of Compound **F-1** to Compound **E** to form Sovaprevir. The formation of the product in Scheme IX, Method 2, (Compound IX) is provided, as is the formation of Sovaprevir from the steps occurring after the formation of the Scheme IX product.

[0005] The disclosure also provides intermediates useful for preparing Sovaprevir and close analogues of Sovaprevir. Particularly the disclosure provides, as useful intermediates, at least Compounds **13**, **14**, **15**, **C**, **C-1**, and **D** in Method 1, the compounds of Schemes IX, X, and XI, **F**, **F-1** (the first compound of Scheme XIII in Method 2), and **F-2**.

[0006] The disclosure provides a crystalline polymorph, Form F.

[0007] The disclosure further provides a bioavailable amorphous form of Sovaprevir and a method for making the amorphous form.

BRIEF DESCRIPTION OF THE DRAWINGS

[0008] FIG. 1. Amino acids residues P1', P1, P2, P2 Top, P3, and P4 as present in sovaprevir.

[0009] FIG. 2 is a graph of intensity (counts per second, CPS) versus scattering angle (degrees 2θ) showing the results of X-ray powder diffraction analysis of the Form F presented in Example 9.


[0010] FIG. 3 is a graph of heat flow (Watts per gram, W/g) versus temperature (°C) showing the results of differential scanning calorimetry analysis of the Form F polymorph, sample size 0.990 mg. Additional experimental details for the DSC analysis are provided in Example 9.

[0011] FIG. 4. is a graph of weight (percent) versus temperature (°C) showing the results of thermogravimetric analysis of the Form F polymorph. The analysis was performed on a 3.4990 mg sample. Analysis was performed using a TA Instruments 2950 thermogravimetric analyzer. Data were collected by heating under nitrogen at a rate of 10 °C/min to a maximum temperature of 350 °C.


DETAILED DESCRIPTION

[0012] Starting materials for sovaprevir production are prepared by methods known in the art. For example Compound **A** (tert-butyl ((1R)-1-((cyclopropylsulfonyl)carbamoyl)-2-vinylcyclopropyl)carbamate) is prepared by the method discussed in WO 2006/122188, page 34, Scheme III and pages 77-78. This procedure is illustrated in Scheme I.


Scheme I



[0013] Compound **B** ((S)-4-(tert-butoxy)-2-(tert-butyl)-4-oxobutanoic acid) is prepared by the method discussed in Evans et. al. *J. Org. Chem.* **1999**, 64, 6411-6417. The synthesis of Compound **B** is shown in Scheme II.




[0014] Compound **D** (4-chloro-7-methoxy-2-phenylquinoline) is prepared by the method reported in WO 2000/009543, page 51. This method is illustrated in Scheme III




[0015] Sovaprevir is a peptidomimetic compound comprised of amino acid like residues P1', P1, P2, P2-Top, P3, and P4. The presence of these residues in Sovaprevir is shown in FIG. 1. The sequence of steps utilized to attach the amino acid residues can vary. Examples 1 through 5 provide two methods by which amino acid residues P1', P1, P2, P2-Top, P3, and P4 can be attached to form sovaprevir. However, other sequences for attaching the amino acid residues to form sovaprevir are possible and will be readily apparent to those of skill in the art.

[0016] The disclosure provides a method of preparing Sovaprevir comprising adding Compound **F-1** to Compound **E** to provide Sovaprevir as follows:




[0017] This method may additionally comprise adding 4-chloro-7-methoxy-2-phenylquinoline to Compound **C-1**, to provide Compound **F-1** as follows:




prior to adding Compound **F-1** to Compound **E**.


[0018] The disclosure also provides a method of preparing a compound of formula (C) comprising at least Step 3 of the following 3 step process, wherein Step 1 comprises deprotecting Compound 13 with acid to provide Compound 14



Step 2 comprises coupling R_1H in base to Compound 14 to provide Compound 15;




Step 3 comprises demethylating Compound 15 with base to provide Compound **C**




where R_1 is a Nitrogen bound 4- to 7- membered heterocycloalkyl ring containing 0 to 2 additional heteroatoms independently chosen from N, O, and S which ring is optionally fused to a 5- or 6- membered heterocyclic ring, containing 1 or 2 heteroatoms independently chosen from N, O, and S, or 5- or 6-membered carbocyclic ring to form a bicyclic ring system, each of which 5- to 7- membered heterocycloalkyl ring or bicyclic ring system is optionally substituted with with 0 to 2 substituents independently chosen from fluoro, amino, hydroxyl, methyl, and trifluoromethyl. In certain embodiments R_1 is a 1-piperidinyl, 1-piperazinyl, or 1-pyrrolidinyl group, each of which is unsubstituted or substituted with 1 or 2 halogen substituents. In certain embodiments Step 2 includes coupling piperidine in base. In this case

Step 2 comprises coupling piperidine in base to Compound **14** to provide Compound **17**;



Step 3 comprises demethylating Compound **17** with base to provide Compound **C-1**




[0019] In certain embodiments R_1 is a 3,3-difluoropiperidin-1-yl group or R_1 is a 1-piperidinyl group.

[0020] The disclosure includes the above method for preparing Compound **C**, comprising Steps 2 and 3 of the method, where R_1 may carry any of the definitions set forth above for this variable.


[0021] The disclosure includes the above method for preparing Compound **C**, comprising Steps 1, 2, and 3 of the method, where R_1 may carry any of the definitions set forth above for this variable.

[0022] The disclosure includes a method of preparing Compound **F-2**, comprising adding 4-chloro-7-methoxy-2-phenylquinoline to Compound **C**, to provide Compound **F-2** as follows:



[0023] Where R_1 may carry any of the values set forth for this variable in the method for producing compound **C**, above. In certain embodiments of the method for producing Compound **F-2**, R_1 is a 1-piperidinyl, 1-piperazinyl, or 1-pyrrolidinyl group, each of which is unsubstituted or substituted with 1 or 2 halogen substituents, or R_1 is a 1-piperidinyl group.


[0024] The disclosure includes an intermediate useful in the processes described in this disclosure, of the Formula **C-2**




[0025] R_1 in Formula **C-2** may carry any of the definitions set forth above. In certain embodiments R_1 is C_1 - C_6 alkyl or hydroxyl, or R_1 is a 1-piperidinyl, 1-piperazinyl, or 1-pyrrolidinyl group, each of which is unsubstituted or substituted with 1 or 2 halogen substituents; and R_2 is hydrogen or C_1 - C_6 alkyl, or R_2 is methyl.

[0026] The disclosure includes intermediates of Formula **C** or **C-2**, in which R_1 is t-butoxy, hydroxyl, or 1-piperidine, and R_2 is methyl.

[0027] In certain embodiment the intermediate is a compound of Formula **C-1**.




[0028] The disclosure also includes intermediate of Formula **F** useful for preparing Sovaprevir.




[0029] R_1 and R_2 in Formula **F**, may carry any of the definitions set forth above for these variables. In certain embodiments R_1 is a 1-piperidinyl, 1-piperazinyl, or 1-pyrrolidinyl group, each of which is unsubstituted or substituted with 1 or 2 halogen substituents; and R_2 is hydrogen or methyl.


[0030] In certain embodiments the intermediate of Formula **F** is a compound of Formula **F-1**.




[0031] In other embodiments the disclosure includes compounds having the formula of any of Compounds IX, X, or XI:



Compound IX




Compound X



Compound XI.

[0032] The disclosure includes the method for making Sovaprevir shown above, comprising adding Compound **F-1** to Compound **E** to provide Sovaprevir, and, additionally comprising the step of hydrolyzing the ester in compound XI with base to form **F-1** as follows:



This step is performed prior to adding Compound **F-1** to Compound **E** to provide Sovaprevir.

[0033] The disclosure also provides a method of preparing a pure amorphous form of Sovaprevir, comprising: crystallizing Sovaprevir in isopropyl alcohol to form Sovaprevir crystalline Form F in greater than 98% purity; dissolving the Sovaprevir crystalline Form F in

6-12 volumes of a solvent such as acetone; adding 6-13 volumes water relative to volume acetone; and precipitating amorphous Sovaprevir.

[0034] The disclosure provides a method of preparing a pure amorphous form of Sovaprevir, comprising crystallizing Sovaprevir in isopropyl alcohol to form Sovaprevir crystalline Form F in greater than 98% purity; dissolving Form F in a solvent such as acetone to form dissolved Sovaprevir; and spray drying the dissolved Sovaprevir to form an amorphous Sovaprevir.

[0035] The disclosure provides a crystalline Sovaprevir form, comprising polymorph F.

[0036] The crystalline Sovaprevir of Form F has the characteristic 2θ values of FIG. 2.

[0037] The crystalline Sovaprevir form of Form F is characterized by an X-ray powder diffraction pattern obtained from a Cu K α source which comprises peaks at 2θ values of 4.2, 8.8, 13.0, and 19.9 +/- 0.2; or 9.7, 13.5, 14.9, 19.0, 19.6, 20.9, and 23.0 +/- 0.2; or 10.4, 17.1, 25.5, 26.1, 26.6, and 27.2 +/- 0.2.

[0038] The crystalline Sovaprevir of Form F has a melting point of 216 °C to 226 °C.

[0039] The crystalline Sovaprevir of Form F has a primary endotherm at 225 °C as determined by DSC.

EXAMPLES

ABBREVIATIONS


[0040] The following abbreviations may be useful when reviewing the schemes and examples of this disclosure.

DCM	Dichloromethane
DIEA/ DIPEA	N,N-diisopropylethylamine
DMF	Dimethyl formamide
DMSO	Dimethyl Sulfoxide
DSC	Differential Scanning Calorimetry
EDC	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
HATU	(1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate)
HOBt	Hydroxybenzotriazole
IPA	Isopropyl Alcohol
KO ^t Bu	Potassium Butoxide

MeOH	Methanol
MTBE	Methyl tert-Butyl Ether
TBTU	o-(Benzotriazol-1-yl)-N,N,N',N'-tetra methyluronium tetrafluoroborate
THF	Tetrahydrofuran
TFA	Trifluoroacetic Acid

EXAMPLE 1. SYNTHESIS OF (2S,4R)-1-((S)-2-(TERT-BUTYL)-4-OXO-4-(PIPERIDIN-1-YL)BUTANOYL)-4-HYDROXYPYRROLIDINE-2-CARBOXYLIC ACID (COMPOUND **C-1**)

[0041] The synthesis of Compound **C-1** is illustrated by Scheme IV. Examples 1-4 provide a complete method for synthesizing sovalprevir. This method is designated as Method 1.



[0042] HATU (1.3 equivalent) was added to a mixture of hydroxyproline, **16** (1.5 equivalent), diisopropyl ethyl amine (5 equivalent) and acid B (1 equivalent) in dimethylformamide at a temperature between 5 – 25 °C. The reaction was stirred for 15 h at room temperature, diluted with water and extracted with methyl t-butylether (MTBE). The organic layer was washed with 1M citric acid and brine. The organic layer was concentrated and residue crystallized from a mixture of MTBE/heptane to give compound **13**.


[0043] Compound **13** was treated with trifluoroacetic acid in dichloromethane (2.3:5 v/v) to remove the t-butyl group. All the volatiles were removed and the material (compound **14**) taken forward without purification to the next step.

[0044] Piperidine (1.2 equivalents) was added at $\sim 5^{\circ}\text{C}$ to a solution of Compound **14**, diisopropyl ethylamine (8.5 equivalents), EDC.HCl (1.3 equivalents), HOBt (0.14 equivalent) in dichloromethane. The reaction was stirred till completion at room temperature, diluted with dichloromethane, and washed with 1M citric acid followed by brine to give compound **17**. Other peptide coupling agents may be used in place of EDC.HCl. These include N,N'-Dicyclohexylcarbodiimide (DCC), triazole coupling agents such as N,N,N',N'-tetramethyl-O-(benzotriazol-1-yl)uronium tetrafluoroborate (TBTU), [Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate (HATU), and benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (PyBOP), or pyridium coupling agents such as 2-Chloro-1-methylpyridinium iodide (Mukaiyama's reagent) and other peptide coupling agents well known in the art.

[0045] Compound **17** (1 equivalent) was dissolved in a mixture of THF/MeOH/water (4/0.17/1 v/v) and lithium hydroxide (1.2 equivalents) was added. The reaction was stirred till hydrolysis was complete (by HPLC). The reaction was evaporated to dryness, diluted with water and extracted with MTBE. The aqueous layer was acidified with HCl and extracted with dichloromethane. The organic layer was dried and concentrated and the residue crystallized from MTBE to give compound C-1.

EXAMPLE 2. SYNTHESIS OF (1R)-1-AMINO-N-(CYCLOPROPYLSULFONYL)-2-VINYLCYCLOPROPANECARBOXAMIDE HYDROCHLORIDE (COMPOUND **E**)


[0046] Scheme V illustrates the synthesis of Compound **E**.



[0047] To a solution of compound A in 1,4-dioxane was added 4M HCl in dioxane and stirred for 3h at room temperature. The reaction mixture was evaporated to dryness to give Compound **E** which was used for the next step without further purification.


EXAMPLE 3. SYNTHESIS OF (2S,4R)-1-((S)-2-(TERT-BUTYL)-4-OXO-4-(PIPERIDIN-1-YL)BUTANOYL)-4-((7-METHOXY-2-PHENYLQUINOLIN-4-YL)OXY)PYRROLIDINE-2-CARBOXYLIC ACID (COMPOUND **F-1**)

[0048] Scheme VI illustrates the preparation of Compound **F-1**.



[0049] Potassium t-butoxide (2.5 equivalent) was added in portion to a solution of compound **C-1** (1 equivalent) and the chloroquinoline **D** (1.0 equivalent) in dimethylsulfoxide and stirred for 24 h at room temperature. The reaction mixture was diluted with dichloromethane and washed with 1M citric acid and concentrated. The residue was triturated with MTBE and the solid isolated by filtration. The solid was crystallized from isopropyl alcohol, cooled to ~3 °C and filtered to give Compound **F-1**.

EXAMPLE 4. PREPARATION OF SOVAPREVIR



[0050] To a solution of Compound **F-1** (1 equivalent), compound **E** (1 equivalent) and TBTU (1.3 equivalent) in dimethylformamide was added diisopropylethyl amine (5 equivalent) and stirred for 15 h at room temperature. The reaction mixture was diluted with

ethylacetate, washed with 1M citric acid, 1M LiOH followed by 1M citric acid and brine. The solution is treated with charcoal and concentrated. The residue was triturated in hot heptane cooled and filtered to give Sovaprevir Crude. Sovaprevir Crude is crystallized from isopropyl alcohol to obtain purified Sovaprevir. Crystallizations from isopropyl alcohol (IPA) will generally produce a crystalline form that is soluble in acetone and typically crystalline Form F is produced.


[0051] Sovaprevir is amorphous in nature and is produced by precipitating a solution of Sovaprevir Purified in acetone into water. Sovaprevir purified is dissolved in 6-12 volumes of acetone and this solution is added to water (6.7 to 12 volumes in relation to acetone used) with vigorous stirring. The precipitated solid is filtered and dried to obtain sovaprevir.

[0052] Sovaprevir Purified is dissolved in acetone and spray dried to obtain amorphous Sovaprevir.

EXAMPLE 5. PREPARATION OF SOVAPREVIR (METHOD 2)


[0053] Schemes VIII through XIII illustrated the preparation of Sovaprevir by Method B.

Scheme VIII




[0054] Lactone **18** is reacted in the presence of an acid catalyst such as amberlyst for a suitable time and temperature to obtain cis-hydroxy proline methyl ester, **19**, in high yield.

Scheme IX




[0055] Cis-hydroxyproline methylester, **19**, is reacted with compound **B** in presence of HATU and diisopropylethylamine in DMF to give compound **IX** as described in example 1.

Scheme X




[0056] Compound **IX** is treated with an acid such as trifluoroacetic acid in a solvent such as dichloromethane to give the acid, compound **20** which is then reacted with piperidine to give compound **X** as described in example 1.

Scheme XI




[0057] A solution of compound **X** is then reacted with Bromobenzene sulfonyl chloride (**21**) in presence of triethylamine and catalytic dimethylaminopyridine in solvent such as dichloromethane to obtain the brosylate (Compound **22**) (methanesulfonyl chloride can also be used to make the mesylate), which can be used for the next step. The brosylate is then reacted with compound **12** in a solvent such as dimethylformamide in presence of a base such as potassium carbonate or potassium hydroxide or bases of the like to obtain compound **XI**.

Scheme XII



[0058] The ester group of compound **XI** is hydrolyzed to the carboxylic acid using a base such as lithium hydroxide in a solvent such as tetrahydrofuran and water to obtain compound **F-1**.

Scheme XIII



[0059] Compound **F-1** is then treated with compound **E** as described in example 4.

EXAMPLE 6. CRYSTALLIZATION OF POLYMORPH F

[0060] Form F is crystallized from isopropanol. Form F was characterized by powder X-ray diffraction (pXRD), differential scanning calorimetry (DSC), thermogravimetric analysis (TG), and hot stage microscopy. The characterization results are presented in FIGs. 2-5.

[0061] XRPD patterns were collected using an Inel XRG-3000 diffractometer equipped with a curved position sensitive detector with a 2θ range of 120° . An incident beam of Cu K α radiation (40 kV, 30 mA) was used to collect data in real time at a resolution of $0.03^\circ 2\theta$. Prior to the analysis, a silicon standard (NIST SRM 640c) was analyzed to verify the Si 111 peak position. Samples were prepared for analysis by packing them into thin-walled glass capillaries. Each capillary was mounted onto a goniometer head and rotated during data acquisition. The monochromator slit was set at 5 mm by 160 μ m, and the samples were analyzed for 5 minutes. Characteristic peaks are observed at the following 2θ values: 4.2, 8.8, 9.7, 10.4, 13.0, 13.5, 14.9, 17.1, 19.0, 19.6, 19.9, 20.9, 23.0, 25.5, 26.1, 26.6, and 27.2 ± 0.2 .

[0062] Differential scanning calorimetry (DSC) was performed using a TA Instruments differential scanning calorimeter Q2000. The sample was placed into an aluminum DSC pan, and the weight accurately recorded. The pan was covered with a lid and then crimped. The sample cell was equilibrated at -50°C (or 25°C , depending on the sample) and heated under a nitrogen purge at a rate of $10^\circ\text{C}/\text{min}$, up to a final temperature of 250°C .

Indium metal was used as the calibration standard. Reported temperatures are at the transition maxima.

[0063] Hot stage microscopy was performed using a Linkam hot stage (model FTIR 600) mounted on a Leica DM LP microscope equipped with a SPOT Insight™ color digital camera. Temperature calibrations were performed using USP melting point standards. Samples were placed on a cover glass, and a second cover glass was placed on top of the sample. As the stage was heated, each sample was visually observed using a 20× 0.40 N.A. long working distance objective with crossed polarizers and a first order red compensator. Images were captured using SPOT software (v. 4.5.9).

[0064] Thermogravimetric analyses (TGA) were performed using a TA Instruments 2950 thermogravimetric analyzer. Each sample was placed in an aluminum sample pan and inserted into the TGA furnace. The furnace was heated under nitrogen at a rate of 10 °C/min, up to a final temperature of 350 °C. Nickely and ALUMEL were used as the calibration standards.

EXAMPLE 7. CHARACTERIZATION DATA FOR POLYMORPH FORM F

Sample Source	Technique	Analysis/Result
lot 146-181-2, LIMS 188807, crystallized from dry IPA	XRPD	Form F
	DSC ^a	very minor broad endo 158, endo 225 (57 J/g)
	TG ^b	1.59 @ 25 – 205 °C 1.34 @ 125 – 205 °C


a. endo = endotherm, temperatures (°C) reported are transition maxima. Temperatures are rounded to the nearest degree.

b. weight loss (%) at a certain temperature; weight changes (%) are rounded to 2 decimal places; temperatures are rounded to the nearest degree.


CLAIMS

What is claimed is:

1. A method of preparing sovalprevir comprising adding Compound **F-1** to Compound **E** to provide Sovaprevir as follows:




2. The method of Claim 1, additionally comprising adding 4-chloro-7-methoxy-2-phenylquinoline (Compound **D**) to Compound **C-1**, to provide Compound **F-1** as follows:




prior to adding Compound **F-1** to Compound **E**.

3. A method of preparing a compound of formula (C) comprising at least Step 3 of the following 3 step process, wherein


Step 1 comprises deprotecting Compound **13** with acid to provide Compound **14**



Step 2 comprises coupling R_1H in base to Compound **14** to provide Compound **15**;




Step 3 comprises demethylating Compound **15** with base to provide Compound **C**



where,

R_1 is a 1-piperidinyl, 1-piperazinyl, or 1-pyrrolidinyl group, each of which is unsubstituted or substituted with 1 or 2 halogen substituents.

4. The method of Claim 3, wherein R_1 is a 1-piperidinyl group.
5. The method of Claim 3, comprising Steps 2 and 3.
6. The method of Claim 3 comprising Steps 1, 2, and 3.
7. A method of preparing Compound **F-2**, comprising adding 4-chloro-7-methoxy-2-phenylquinoline (Compound **D**) to Compound **C**, to provide Compound **F-2** as follows:




where,

R_1 is a 1-piperidinyl, 1-piperazinyl, or 1-pyrrolidinyl group, each of which is unsubstituted or substituted with 1 or 2 halogen substituents.


8. The method of Claim 7 wherein R_1 is a 1-piperidinyl group.

9. The method of Claim 7, additionally comprising the following synthetic step, in which R_1 is a 1-piperidinyl group




10. The method of Claim 9, additionally comprising on or more of Steps 1, 2, and 3, where


Step 1 comprises deprotecting Compound **13** with acid to provide Compound **14**



Step 2 comprises coupling R_1H in base to Compound **14** to provide Compound **15**;




Step 3 comprises demethylating Compound **15** with base to provide Compound **C**



where,

R_1 is a 1-piperidinyl, 1-piperazinyl, or 1-pyrrolidinyl group, each of which is unsubstituted or substituted with 1 or 2 halogen substituents.


11. A compound having the formula:




R_1 is C_1 - C_6 alkyl or hydroxyl, or R_1 is a 1-piperidinyl, 1-piperazinyl, or 1-pyrrolidinyl group, each of which is unsubstituted or substituted with 1 or 2 halogen substituents; and R_2 is hydrogen or methyl.

12. A compound of Claim 11 in which R_1 is t-butoxy, hydroxyl, or 1-piperidine, and R_2 is methyl.

13. A compound of Claim 11 of the formula



14. A compound, having the Formula F




where,


R₁ is a 1-piperidiny, 1-piperazinyl, or 1-pyrrolidinyl group, each of which is unsubstituted or substituted with 1 or 2 halogen substituents;

R₂ is hydrogen or methyl.

15. A compound of Claim 14 of the formula




16. A compound of Claim 11, having the formula of Compound IX




Compound IX.

17. A compound of Claim 11, having the formula of Compound X




Compound X.

18. A compound of Claim 14, having the formula of Compound XI



Compound XI.

19. The process of Claim 1, additionally comprising the step of Demethylating Compound XI with base to form **F-1** as follows:



prior to adding Compound **F-1** to Compound **E** to provide Sovaprevir.

20. A method of preparing a pure amorphous form of Sovaprevir, comprising crystallizing Sovaprevir in isopropyl alcohol to form Sovaprevir crystalline Form F; dissolving the Sovaprevir crystalline Form F in 6-12 volumes acetone;

adding 6-13 volumes water relative to volume acetone; and
precipitating amorphous Sovaprevir of greater than 98% purity.

21. A method of preparing a pure amorphous form of Sovaprevir, comprising
crystallizing Sovaprevir in isopropyl alcohol to form Sovaprevir crystalline Form F;
dissolving Form F in an acetone to form dissolved Sovaprevir; and
spray drying the dissolved Sovaprevir to form an amorphous Sovaprevir of greater than 98%
purity.

22. A crystalline Sovaprevir form, comprising polymorph Form F.

23. The crystalline Sovaprevir form of Claim 22, having the characteristic 2θ values of
FIG. 2.

24. The crystalline Sovaprevir form of Claim 22, wherein the form is characterized by an
X-ray powder diffraction pattern obtained from a Cu $K\alpha$ source which comprises peaks at 2θ
values of 4.2, 8.8, 13.0, and 19.9 $\pm$ 0.2; or
9.7, 13.5, 14.9, 19.0, 19.6, 20.9, and 23.0 $\pm$ 0.2; or
10.4, 17.1, 25.5, 26.1, 26.6, and 27.2 $\pm$ 0.2.

25. The crystalline Sovaprevir form of Claim 22, wherein polymorph F has a melting
point of 216 °C to 226 °C.

26. The crystalline Sovaprevir form of Claim 22, wherein polymorph F has a primary
endotherm at 225 °C as determined by DSC.