CLAIMS

1. Use of a ~~A~~ botulinum neurotoxin in the manufacture of a medicament~~use~~ for~~in~~ treating a tremor of the upper limb ~~comprising~~ wherein said medicament is to be administered ~~the administration of a botulinum neurotoxin~~ to at least one muscle of each of the forearm/wrist, elbow and shoulder, wherein the botulinum neurotoxin is to be administered to at least one muscle of the forearm/wrist selected from the group of *M. extensor carpi ulnaris* (ECU), *M. extensor carpi radialis* (ECR), *M. pronator quadratus* (PQ) and *M.supinator* in a dosage in the range of 2 to 6 U and wherein the botulinum neurotoxin is to be administered to at least one muscle of the elbow in a dosage of about 20 U and to at least one muscle of the shoulder in a dosage of about 15 U.

2. The use~~A botulinum neurotoxin for use~~ according to claim 1, wherein the botulinum neurotoxin is to be administered in addition to at least one muscle of the wrist/forearm selected from the group of *M. flexor carpi radialis* (FCR), *M. flexor carpi ulnaris* (FCU) and *M. pronator teres* (PT) in a dose in the range of 4 to 16 U per muscle.

3. The use ~~A botulinum neurotoxin for use~~ according to claim 1 , wherein the botulinum neurotoxin is to be administered to at least one muscle of the forearm/wrist selected from the group of *M. extensor carpi ulnaris* (ECU), *M. extensor carpi radialis* (ECR), *M. pronator quadratus* (PQ) and *M. supinator* in a dosage of about 2.5 U.

4. The use ~~botulinum neurotoxin for use~~ according to claim 3, wherein the botulinum neurotoxin is to be administered in addition to at least one muscles of the wrist/forearm selected from the group of *M. flexor carpi radialis* (FCR), *M. flexor carpi ulnaris* (FCU) and *M. pronator teres* (PT) in a dose of about 10 U per muscle.

5. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 4, wherein the botulinum neurotoxin is not to be administered to the *M. exte~~r~~nsor digitorium communis*.

6. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 5, wherein the total dose of the botulinum neurotoxin to be administered to the forearm/wrist muscles does not exceed 65 U.

7. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 6, wherein the botulinum neurotoxin is to be administered to at least one muscle of the elbow selected from the group of *M. brachialis* and *M. triceps brachii*.

8. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 7, wherein the botulinum neurotoxin is not to be administered to the *M. biceps brachii*.

9. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 8, wherein the total dose~~does~~ of the botulinum neurotoxin to be administered to the muscles of the elbow does not exceed 40 U.

10. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 9, wherein the botulinum neurotoxin is to be administered to at least one muscle of the shoulder selected from the group of *M. latissimus dorsi*, *M. pectoralis major*, *M. supraspinatus* and *M. infraspinatus*.

11. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 10, wherein the botulinum neurotoxin is not to be administered to the *deltoid* and the *teres major* muscle.

12. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 11, wherein the total dose of the botulinum neurotoxin to be administered to muscles of the shoulder does not exceed 60 U.

13. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 12, wherein the total dose of the botulinum neurotoxin to be administered to the muscles of the forearm/wrist, elbow and shoulder does not exceed 165 U.

14. The use ~~botulinum neurotoxin for use~~ according to claim 1, wherein the botulinum neurotoxin is to be administered to 4, 5, 6, or 7 muscles of the forearm/wrist, 2 muscles of the elbow and 4 muscles of the shoulder in a dose according to the scheme:

M. extensor carpi ulnaris (ECU)	~~about~~ 2.5 U to 5 U
M. extensor carpi radialis (ECR)	~~about~~ 2.5 U to 5 U
M. flexor carpi radialis (FCR)	~~about~~ 5 U to 15 U
M. flexor carpi ulnaris (FCU)	~~about~~ 5 U to 15 U
M. pronator teres (PT)	~~about~~ 5 U to 15 U
M. pronator quadratus (PQ)	~~about~~ 2.5 U to 5 U
M. supinator	~~about~~ 2.5 U to 5 U
M. brachialis	about 20 U

M. triceps brachii	about 20 U
M. latissimus dorsi	about 15 U
M. pectoralis major	about 15 U
M. supraspinatus	about 15 U
M. infraspinatus	about 15 U

15. The use ~~botulinum neurotoxin for use~~ according to claim 1, wherein the botulinum neurotoxin is to be administered to 7 muscles of the forearm/wrist, 2 muscles of the elbow and 4 muscles of the shoulder in a dose according to the scheme:

M. extensor carpi ulnaris (ECU)	about 2.5 U
M. extensor carpi radialis (ECR)	about 2.5 U
M. flexor carpi radialis (FCR)	about 10 U
M. flexor carpi ulnaris (FCU)	about 10 U
M. pronator teres (PT)	about 10 U
M. pronator quadratus (PQ)	about 2.5 U
M. supinator	about 2.5 U
M. brachialis	about 20 U
M. triceps brachii	about 20 U
M. latissimus dorsi	about 15 U
M. pectoralis major	about 15 U
M. supraspinatus	about 15 U
M. infraspinatus	about 15 U

16. The ~~botulinum neurotoxin for~~ use according to any one of claims 1 to 15, wherein said botulinum neurotoxin is the neurotoxic component of a botulinum neurotoxin complex, wherein said neurotoxic component is devoid of any other protein component of the Clostridium botulinum neurotoxin complex.

17. The ~~botulinum neurotoxin for~~ use according to any one of claims 1 to 16, wherein said botulinum neurotoxin is selected from the group of serotypes consisting of A, B and E.

18. The use ~~botulinum neurotoxin for use~~ according to any one of claims 1 to 17, wherein the botulinum neurotoxin is to be administered together with at least one standard treatment selected from propranolol, primidone, any other antiepileptic or a calcium channel blocker, Deep Brain Stimulation (DBS), Magnetic Resonance guided High Frequency Ultrasound (MRgHiFUS), local electrical stimulation, biofeedback, kinematic assessment guided stimulation, anti-tremor appliances, anti-tremor smartphone apps or combinations thereof.

19. Use of a botulinum neurotoxin according to any one of claims 1 to 18 in the manufacture of a~~A~~ pharmaceutical composition ~~comprising a botulinum neurotoxin according to any one of claims 1 to 18 for use in~~ for treating a tremor of the upper limb.

~~20. A method of treating a tremor of the upper limb wherein the method comprises the administration of a therapeutically effective amount of a botulinum neurotoxin according to any one of claims 1 to 19.~~