

TRANSMEMBRANE PROTEASE, SERINE 6 (TMPRSS6) iRNA COMPOSITIONS AND METHODS OF USE THEREOF

Related Applications

5 This application claims the benefit of priority to U.S. Provisional Application No. 63/179,607, filed on April 26, 2021, and U.S. Provisional Application No. 63/278,227, filed on November 11, 2021. The entire contents of each of the foregoing applications are incorporated herein by reference.

Sequence Listing

10 The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on April 18, 2022, is named 121301_15420_SL.TXT and is 673,291 bytes in size.

Background of the Invention

15 TMPRSS6 (Transmembrane Protease, Serine 6), also known as matriptase-2, is a type II serine protease. It is primarily expressed in the liver, although high levels of TMPRSS6 mRNA are also found in the kidney, with lower levels in the uterus and much smaller amounts detected in many other tissues (Beliveau *et al.*, 2019, *Cell Chemical Biology* 26, 1559–1572). TMPRSS6 plays a key role in iron homeostasis *via* modulation of hepcidin expression. Hepcidin, a liver-derived peptide
20 hormone, is known as a central regulator of systemic iron homeostasis, and its unbalanced production contributes to the pathogenesis of a spectrum of iron disorders. Hepcidin functions by blocking the absorption of dietary iron from the intestine, and the release of iron from macrophages and hepatocytes (Ganz T. 2011, *Blood*, vol. 117, 17, 4425-4433). Hepcidin gene expression can be stimulated in response to iron through BMP/SMAD-dependent signal transduction cascade mediated
25 by the BMP-co-receptor hemojuvelin (HJV). TMPRSS6 inhibits BMP-mediated upregulation of hepcidin by cleaving the BMP co-receptor HJV, thus preventing BMP signaling, SMAD translocation to the nucleus, and hepcidin transcriptional activation, which causes downregulation of hepcidin levels (Finberg, K.E., *et al.*, 2010, *Blood* 115, 3817–3826; Wang, C.Y., *et al.*, 2014 *Front. Pharmacol.* 5, 114).

30 Therefore, inhibition of TMPRSS6 results in increased hepcidin levels, making it an attractive pharmacological target for disorders associated with iron overload and inappropriately low hepcidin or for disorders where iron restriction is desirable. Numerous disorders, such as thalassemias, hemochromatosis, and certain types of myelodysplastic syndromes (MDS), are associated with iron overload, a condition characterized by increased levels of iron. Iron overload can result in excess iron
35 deposition in various tissues and can eventually lead to tissue and organ damage. In addition, iron restriction is desirable in certain disorders such as polycythemia vera.

 Current treatments for disorders associated with iron overload and disorders where iron restriction is desirable (*e.g.* polycythemia vera) include phlebotomy or venesection, a treatment to remove iron-rich blood from the body; splenectomy; iron chelation therapy; and dieting. However,

these treatments are not always effective. Accordingly, there is a need in the art for alternative treatments for subjects having disorders associated with iron overload.

Summary of the Invention

5 The present invention provides iRNA compositions which affect the RNA-induced silencing complex (RISC)-mediated cleavage of RNA transcripts of a gene encoding Transmembrane protease, serine 6 (TMPRSS6). The TMPRSS6 gene may be within a cell, *e.g.*, a cell within a subject, such as a human subject. The present invention also provides methods of using the iRNA compositions of the invention for inhibiting the expression of a TMPRSS6 gene and/or for treating a subject who would
10 benefit from inhibiting or reducing the expression of a TMPRSS6 gene, *e.g.*, a subject suffering or prone to suffering from a TMPRSS6-associated disorder, *e.g.*, an iron overload associated disorder and/or a disorder of ineffective erythropoiesis, such as thalassemia, *e.g.*, β -thalassemia, hemochromatosis, myelodysplastic syndromes (MDS), or polycythemia vera.

 Accordingly, in an aspect, the present invention provides a double stranded ribonucleic acid
15 (dsRNA) for inhibiting expression of Transmembrane protease, serine 6 (TMPRSS6) in a cell, wherein said dsRNA comprises a sense strand and an antisense strand forming a double stranded region, wherein the antisense strand comprises a region of complementarity to an mRNA encoding TMPRSS6, and wherein the region of complementarity comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, or 20, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from any one of the
20 antisense nucleotide sequences in any one of Tables 2-7.

 In one embodiment, the dsRNA agent comprises a sense strand comprising a contiguous nucleotide sequence which has at least 85%, *e.g.*, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%, nucleotide sequence identity over its entire length to any one of the nucleotide sequences of the sense strands in any one of Tables 2-7 and an antisense strand
25 comprising a contiguous nucleotide sequence which has at least 85%, *e.g.*, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%, nucleotide sequence identity over its entire length to any one of the nucleotide sequences of the antisense strands in any one of Tables 2-7.

 In one embodiment, the dsRNA agent comprises a sense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three nucleotides from
30 any one of the nucleotide sequences of the sense strands in any one of Tables 2-7 and an antisense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three nucleotides from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-7.

 In one embodiment, the dsRNA agent comprises a sense strand comprising at least 15, *e.g.*,
35 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than two nucleotides from any one of the nucleotide sequences of the sense strands in any one of Tables 2-7 and an antisense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than two nucleotides from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-7.

In one embodiment, the dsRNA agent comprises a sense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than one nucleotide from any one of the nucleotide sequences of the sense strands in any one of Tables 2-7 and an antisense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than one nucleotide from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-7.

In one embodiment, the dsRNA agent comprises a sense strand comprising or consisting of a nucleotide sequence selected from the group consisting of any one of the nucleotide sequences of the sense strands in any one of Tables 2-7 and an antisense strand comprising or consisting of a nucleotide sequence selected from the group consisting of any one of the nucleotide sequences of the antisense strands in any one of Tables 2-7.

In one aspect, the present invention provides a double stranded ribonucleic acid (dsRNA) for inhibiting expression of Transmembrane protease, serine 6 (TMPRSS6) in a cell, wherein said dsRNA comprises a sense strand and an antisense strand forming a double stranded region, wherein the sense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three, *e.g.*, 3, 2, 1, or 0, nucleotides from any one of the nucleotide sequence of nucleotides 187-210; 227-254; 322-363; 362-390; 398-420; 404-429; 410-435; 439-461; 443-467; 448-474; 460-483; 466-488; 496-519; 519-542; 526-548; 557-593; 641-671; 652-676; 687-713; 725-762; 757-794; 886-908; 921-951; 956-987; 1051-1082; 1233-1269; 1279-1313; 1313-1341; 1327-1351; 1415-1439; 1447-1480; 1464-1486; 1486-1509; 1559-1589; 1571-1595; 1579-1609; 1707-1735; 1738-1764; 1806-1828; 1864-1886; 1934-1966; 1967-1991; 2008-2031; 2015-2043; 2042-2072; 2287-2311; 2297-2354; 2336-2361; 2360-2384; 2416-2438; 2481-2510; 2496-2527; 2526-2558; 2665-2693; 2693-2719; 2707-2729; 2799-2821; 2851-2874; 2971-2999; 2981-3006 and 3155-3195 of SEQ ID NO: 1, and the antisense strand comprises at least 15 contiguous nucleotides differing by no more than 3, *e.g.*, 3, 2, 1, or 0, nucleotides from the corresponding nucleotide sequence of SEQ ID NO:2.

In one aspect, the present invention provides a double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of Transmembrane protease, serine 6 (TMPRSS6) in a cell, wherein the dsRNA agent comprises a sense strand and an antisense strand forming a double stranded region, wherein the sense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three, *e.g.*, 3, 2, 1, or 0, nucleotides from any one of the nucleotide sequence of nucleotides 230-252, 324-346, 560-578, 560-582, 2338-2360, 3163-3185, 3169-3191, and 3172-3194 of SEQ ID NO: 1, and the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three, *e.g.*, 3, 2, 1, or 0, nucleotides from the corresponding nucleotide sequence of SEQ ID NO:2.

In one aspect, the present invention provides a double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of Transmembrane protease, serine 6 (TMPRSS6) in a cell, wherein the dsRNA agent comprises a sense strand and an antisense strand forming a double stranded region, wherein the sense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three, *e.g.*, 3, 2, 1, or 0, nucleotides from any one of the

nucleotide sequence of nucleotides 560-578, 2338-2360, and 3169-3191 of SEQ ID NO: 1, and the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than 3, *e.g.*, 3, 2, 1, or 0, nucleotides from the corresponding nucleotide sequence of SEQ ID NO:2.

5 In some embodiments, the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three, *e.g.*, 3, 2, 1, or 0, nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1556360, AD-1571158, AD-1571033, AD-1554875, AD-1571160, AD-1555117, AD-1554911, and AD-1556915.

10 In some embodiments, the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three, *e.g.*, 3, 2, 1, or 0, nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1556360, AD-1571158, and AD-1571033.

In one embodiment, the dsRNA agent comprises at least one modified nucleotide.

15 In one embodiment, substantially all of the nucleotides of the sense strand; substantially all of the nucleotides of the antisense strand comprise a modification; or substantially all of the nucleotides of the sense strand and substantially all of the nucleotides of the antisense strand comprise a modification.

In one embodiment, all of the nucleotides of the sense strand comprise a modification; all of the nucleotides of the antisense strand comprise a modification; or all of the nucleotides of the sense strand and all of the nucleotides of the antisense strand comprise a modification.

In one embodiment, at least one of the modified nucleotides is selected from the group consisting of a deoxy-nucleotide, a 3'-terminal deoxythymidine (dT) nucleotide, a 2'-O-methyl modified nucleotide, a 2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a locked nucleotide, an unlocked nucleotide, a conformationally restricted nucleotide, a constrained ethyl nucleotide, an abasic nucleotide, a 2'-amino-modified nucleotide, a 2'-O-allyl-modified nucleotide, 2'-C-alkyl-modified nucleotide, 2'-hydroxyl-modified nucleotide, a 2'-methoxyethyl modified nucleotide, a 2'-O-alkyl-modified nucleotide, a morpholino nucleotide, a phosphoramidate, a non-natural base comprising nucleotide, a tetrahydropyran modified nucleotide, a 1,5-anhydrohexitol modified nucleotide, a cyclohexenyl modified nucleotide, a nucleotide comprising a phosphorothioate group, a nucleotide comprising a methylphosphonate group, a nucleotide comprising a 5'-phosphate, a nucleotide comprising a 5'-phosphate mimic, a thermally destabilizing nucleotide, a glycol modified nucleotide (GNA), a nucleotide comprising a 2' phosphate, and a 2-O-(N-methylacetamide) modified nucleotide; and combinations thereof.

35 In one embodiment, the modifications on the nucleotides are selected from the group consisting of LNA, HNA, CeNA, 2'-methoxyethyl, 2'-O-alkyl, 2'-O-allyl, 2'-C-allyl, 2'-fluoro, 2'-deoxy, 2'-hydroxyl, and glycol; and combinations thereof.

In one embodiment, at least one of the modified nucleotides is selected from the group consisting of a deoxy-nucleotide, a 2'-O-methyl modified nucleotide, a 2'-fluoro modified nucleotide,

a 2'-deoxy-modified nucleotide, a glycol modified nucleotide (GNA), *e.g.*, Ggn, Cgn, Tgn, or Agn, a nucleotide with a 2' phosphate, *e.g.*, G2p, C2p, A2p or U2p, and, a vinyl-phosphonate nucleotide; and combinations thereof.

5 In some embodiments, the modified nucleotide comprises a short sequence of 3'-terminal deoxythymidine nucleotides (dT).

In some embodiments, the dsRNA agent further comprises at least one phosphorothioate internucleotide linkage. In some embodiments, the dsRNA agent comprises 6-8 phosphorothioate internucleotide linkages. In one embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the 3'-terminus of one strand. Optionally, the strand is the antisense strand. In another embodiment, the strand is the sense strand. In a related embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the 5'-terminus of one strand. Optionally, the strand is the antisense strand. In another embodiment, the strand is the sense strand. In another embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the both the 5'- and 3'-terminus of one strand. Optionally, the strand is the antisense strand. In another embodiment, the strand is the sense strand.

10
15

The double stranded region may be 19-30 nucleotide pairs in length; 19-25 nucleotide pairs in length; 19-23 nucleotide pairs in length; 23-27 nucleotide pairs in length; or 21-23 nucleotide pairs in length.

In one embodiment, each strand is independently no more than 30 nucleotides in length.

20 In one embodiment, the sense strand is 21 nucleotides in length and the antisense strand is 23 nucleotides in length.

The region of complementarity may be at least 17 nucleotides in length; between 19 and 23 nucleotides in length; or 19 nucleotides in length.

In one embodiment, at least one strand comprises a 3' overhang of at least 1 nucleotide. In another embodiment, at least one strand comprises a 3' overhang of at least 2 nucleotides.

25

In one embodiment, the dsRNA agent further comprises a ligand.

In one embodiment, the ligand is conjugated to the 3' end of the sense strand of the dsRNA agent.


In one embodiment, the ligand is conjugated to the 5' end of the sense strand of the dsRNA agent.

30


In one embodiment, the ligand is an N-acetylgalactosamine (GalNAc) derivative.

In one embodiment, the ligand is one or more GalNAc derivatives attached through a monovalent, bivalent, or trivalent branched linker.

In one embodiment, the ligand is



In one embodiment, the dsRNA agent is conjugated to the ligand as shown in the following schematic




5

and, wherein X is O or S.

In one embodiment, the X is O.

In one embodiment, the dsRNA agent is conjugated to the ligand as shown in the following schematic



10

In one embodiment, the dsRNA agent further comprises at least one phosphorothioate or methylphosphonate internucleotide linkage.

In one embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the 3'-terminus of one strand, *e.g.*, the antisense strand or the sense strand.

15 In another embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the 5'-terminus of one strand, *e.g.*, the antisense strand or the sense strand.

In one embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the both the 5'- and 3'-terminus of one strand. In one embodiment, the strand is the antisense strand.

In one embodiment, the base pair at the 1 position of the 5'-end of the antisense strand of the duplex is an AU base pair.

5 In one embodiment, the sense strand comprises at least 17 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-GACGCCACGCAUGCUGUGUGU -3' (SEQ ID NO: 119).

In one embodiment, the sense strand comprises at least 19 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-
10 GACGCCACGCAUGCUGUGUGU -3' (SEQ ID NO: 119).

In one embodiment, the sense strand comprises or consists of the nucleotide sequence of 5'-GACGCCACGCAUGCUGUGUGU -3' (SEQ ID NO: 119).

In one embodiment, the antisense strand comprises at least 17 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-
15 ACACACAGCAUGCUGGCGUCAC -3' (SEQ ID NO: 245).

In one embodiment, the antisense strand comprises at least 19 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-ACACACAGCAUGCUGGCGUCAC -3' (SEQ ID NO: 245).

In one embodiment, the antisense strand comprises at least 21 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-
20 ACACACAGCAUGCUGGCGUCAC -3' (SEQ ID NO: 245).

In one embodiment, the antisense strand comprises or consists of the nucleotide sequence of 5'-ACACACAGCAUGCUGGCGUCAC -3' (SEQ ID NO: 245).

In one embodiment, the sense strand comprises the nucleotide sequence of 5'-GACGCCACGCAUGCUGUGUGU -3' (SEQ ID NO: 119) and the antisense strand comprises the nucleotide sequence of 5'-ACACACAGCAUGCUGGCGUCAC -3' (SEQ ID NO: 245).
25

In one embodiment, the sense strand differs by no more than 3, e.g., 0, 1, 2, or 3, modified nucleotides from the nucleotide sequence of 5'-gsascgccacGfCfAfugcugugugu-3' (SEQ ID NO: 371) wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage.
30


In one embodiment, the antisense strand differs by no more than 3, e.g., 0, 1, 2, or 3, modified nucleotides from the nucleotide sequence of 5'-asdCsacdAcdAgcaudGcGfugcgucsasc -3' (SEQ ID NO: 497), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein dA, dG, and dC are 2'-deoxyadenosine-3'-phosphate, 2'-deoxyguanosine-3'-phosphate, and 2'-deoxycytidine-3'-phosphate respectively; and s is a phosphorothioate linkage.
35

In one embodiment, the sense strand comprises the nucleotide sequence of 5'-gsascgccacGfCfAfugcugugugu-3' (SEQ ID NO: 371) and the antisense strand comprises the nucleotide sequence of 5'-asdCsacdAcdAgcaudGcGfugcgucsasc -3' (SEQ ID NO: 497), wherein a,

g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein dA, dG, and dC are 2'-deoxyadenosine-3'-phosphate, 2'-deoxyguanosine-3'-phosphate, and 2'-deoxycytidine-3'-phosphate respectively; and s is a phosphorothioate linkage.

- 5 In one embodiment, the sense strand comprises the nucleotide sequence of 5'-gsascgccacGfCfAfugcuguguguL96 -3' (SEQ ID NO: 371) and the antisense strand comprises the nucleotide sequence of 5'-asdCsacdAc dAgcaudGcGfuggcgucsasc -3' (SEQ ID NO: 497), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein dA, dG, and dC are 2'-deoxyadenosine-3'-phosphate, 2'-deoxyguanosine-3'-phosphate, and 2'-deoxycytidine-3'-phosphate respectively; s is a phosphorothioate linkage, and L96 is N-[tris(GalNAc-alkyl)-amidodecanoyl]-4-hydroxyprolinol.

- 10 In one embodiment, the sense strand comprises the nucleotide sequence of 5'-gsascgccacGfCfAfugcugugugu -3' (SEQ ID NO: 371) and the antisense strand comprises the nucleotide sequence of 5'-asdCsacdAc dAgcaudGcGfuggcgucsasc -3' (SEQ ID NO: 497), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein dA, dG, and dC are 2'-deoxyadenosine-3'-phosphate, 2'-deoxyguanosine-3'-phosphate, and 2'-deoxycytidine-3'-phosphate respectively; and s is a phosphorothioate linkage, wherein the 3'-end of the sense strand is conjugated to the ligand as shown in the following schematic:



- 20 and, wherein X is O.

In one embodiment, the sense strand comprises at least 17 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-CCUUUGGAAUAAAGCUGCCUU -3' (SEQ ID NO: 844).

- 25 In one embodiment, the sense strand comprises at least 19 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-CCUUUGGAAUAAAGCUGCCUU -3' (SEQ ID NO: 844).

In one embodiment, the sense strand comprises or consists of the nucleotide sequence of 5'-CCUUUGGAAUAAAGCUGCCUU -3' (SEQ ID NO: 844).

In one embodiment, the antisense strand comprises at least 17 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-AAGGCAGCUUUAUCCAAAGGGC -3' (SEQ ID NO: 1868).

In one embodiment, the antisense strand comprises at least 19 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-AAGGCAGCUUUAUCCAAAGGGC -3' (SEQ ID NO: 1868).

In one embodiment, the antisense strand comprises at least 21 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-AAGGCAGCUUUAUCCAAAGGGC -3' (SEQ ID NO: 1868).

In one embodiment, the antisense strand comprises or consists of the nucleotide sequence of 5'-AAGGCAGCUUUAUCCAAAGGGC -3' (SEQ ID NO: 1868).

In one embodiment, the sense strand comprises the nucleotide sequence of 5'-CCUUUGGAAUAAAGCUGCCUU -3' (SEQ ID NO: 844) and the antisense strand comprises the nucleotide sequence of 5'-AAGGCAGCUUUAUCCAAAGGGC -3' (SEQ ID NO: 1868).

In one embodiment, the sense strand differs by no more than 3, e.g., 0, 1, 2, or 3, modified nucleotides from the nucleotide sequence of 5'-cscsuuugGfaAfUfAfaagcugccuu -3' (SEQ ID NO: 2095) wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage.


In one embodiment, the antisense strand differs by no more than 3, e.g., 0, 1, 2, or 3, modified nucleotides from the nucleotide sequence of 5'-asAfsggdCa(G2p)cuuuauUfcCfaaaggsgsc -3' (SEQ ID NO: 2324), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein G2p is guanosine-2'-phosphate; and s is a phosphorothioate linkage.

In one embodiment, the sense strand comprises the nucleotide sequence of 5'-cscsuuugGfaAfUfAfaagcugccuu -3' (SEQ ID NO: 2095) and the antisense strand comprises the nucleotide sequence of 5'-asAfsggdCa(G2p)cuuuauUfcCfaaaggsgsc -3' (SEQ ID NO: 2324), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein G2p is guanosine-2'-phosphate; and s is a phosphorothioate linkage.

In one embodiment, the sense strand comprises the nucleotide sequence of 5'-cscsuuugGfaAfUfAfaagcugccuuL96 -3' (SEQ ID NO: 2095) and the antisense strand comprises the nucleotide sequence of 5'-asAfsggdCa(G2p)cuuuauUfcCfaaaggsgsc -3' (SEQ ID NO: 2324), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein G2p is guanosine-2'-phosphate; s is a phosphorothioate linkage, and L96 is N-[tris(GalNAc-alkyl)-amidodecanoyl]-4-hydroxyprolinol.

In one embodiment, the sense strand comprises the nucleotide sequence of 5'-cscsuuugGfaAfUfAfaagcugccuu -3' (SEQ ID NO: 2095) and the antisense strand comprises the nucleotide sequence of 5'-asAfsggdCa(G2p)cuuuauUfcCfaaaggsgsc -3' (SEQ ID NO: 2324), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are

2'-fluoro A, G, C and U respectively; wherein G2p is guanosine-2'-phosphate, s is a phosphorothioate linkage, and wherein the 3'-end of the sense strand is conjugated to the ligand as shown in the following schematic:



5 and, wherein X is O.

In one embodiment, the sense strand comprises at least 17 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-UCACCUGCUUCUUCUGGUU-3' (SEQ ID NO: 1686).

10 In one embodiment, the sense strand comprises or consists of the nucleotide sequence of 5'-UCACCUGCUUCUUCUGGUU-3' (SEQ ID NO: 1686).

In one embodiment, the antisense strand comprises at least 17 contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of 5'-AACCAGAAGAAGCAGGUGA-3' (SEQ ID NO: 1790).

15 In one embodiment, the antisense strand comprises or consists of the nucleotide sequence of 5'-AACCAGAAGAAGCAGGUGA-3' (SEQ ID NO: 1790).

In one embodiment, the sense strand comprises the nucleotide sequence of 5'-UCACCUGCUUCUUCUGGUU-3' (SEQ ID NO: 1686) and the antisense strand comprises the nucleotide sequence of 5'-AACCAGAAGAAGCAGGUGA-3' (SEQ ID NO: 1790).

20 In one embodiment, the sense strand differs by no more than 3, e.g., 0, 1, 2, or 3, modified nucleotides from the nucleotide sequence of 5'-UfcAfcCfuGfcUfuCfuUfcUfgGfsusUf-3' (SEQ ID NO: 1974) wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage.


25 In one embodiment, the antisense strand differs by no more than 3, e.g., 0, 1, 2, or 3, modified nucleotides from the nucleotide sequence of 5'-asAfscCfaGfaAfgAfaGfcAfgGfusGfsa-3' (SEQ ID NO:), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage.

30 In one embodiment, the sense strand comprises the nucleotide sequence of 5'-UfcAfcCfuGfcUfuCfuUfcUfgGfsusUf-3' (SEQ ID NO: 2203) and the antisense strand comprises the nucleotide sequence of 5'-asAfscCfaGfaAfgAfaGfcAfgGfusGfsa-3' (SEQ ID NO:), wherein a,

g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage.

In one embodiment, the sense strand comprises the nucleotide sequence of 5'-Q191sUfcAfcCfuGfcUfuCfuUfcUfgGfsusUf -3' (SEQ ID NO: 1974) and the antisense strand comprises the nucleotide sequence of 5'-asApscCfaGfaAfgAfaGfcAfgGfusGfsa -3' (SEQ ID NO: 2203), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; s is a phosphorothioate linkage, and Q191 is N-[tris(GalNAc-alkyl)-amidododecanoyl]-(S)-pyrrolidin-3-ol-phosphorothioate (p-C12-(GalNAc-alkyl)3).

In one embodiment, the sense strand comprises the nucleotide sequence of 5'-UfcAfcCfuGfcUfuCfuUfcUfgGfsusUf -3' (SEQ ID NO: 1974) and the antisense strand comprises the nucleotide sequence of 5'-asApscCfaGfaAfgAfaGfcAfgGfusGfsa -3' (SEQ ID NO: 2203), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage, wherein the 5'-end of the sense strand is conjugated to the ligand as shown in the following schematic:



In another embodiment, the RNAi agent is a pharmaceutically acceptable salt thereof. "Pharmaceutically acceptable salts" of each of RNAi agents herein include, but are not limited to, a sodium salt, a calcium salt, a lithium salt, a potassium salt, an ammonium salt, a magnesium salt, a mixtures thereof. One skilled in the art will appreciate that the RNAi agent, when provided as a polycationic salt having one cation per free acid group of the optionally modified phosphodiester backbone and/or any other acidic modifications (*e.g.*, 5'-terminal phosphonate groups). For example, an oligonucleotide of "n" nucleotides in length contains n-1 optionally modified phosphodiester, so that an oligonucleotide of 21 nt in length may be provided as a salt having up to 20 cations (*e.g.*, 20 sodium cations). Similarly, an RNAi agents having a sense strand of 21 nt in length and an antisense strand of 23 nt in length may be provided as a salt having up to 42 cations (*e.g.*, 42 sodium cations). In the preceding example, where the RNAi agent also includes a 5'-terminal phosphate or a 5'-terminal vinylphosphonate group, the RNAi agent may be provided as a salt having up to 44 cations (*e.g.*, 44 sodium cations).

The present invention also provides cells containing any of the dsRNA agents of the invention and pharmaceutical compositions comprising any of the dsRNA agents of the invention.

The pharmaceutical composition of the invention may include dsRNA agent in an unbuffered solution, *e.g.*, saline or water, or the pharmaceutical composition of the invention may include the dsRNA agent is in a buffer solution, *e.g.*, a buffer solution comprising acetate, citrate, prolamine, carbonate, or phosphate or any combination thereof; or phosphate buffered saline (PBS).

5 In one aspect, the present invention provides a method of inhibiting expression of a Transmembrane protease, serine 6 (TMPRSS6) gene in a cell. The method includes contacting the cell with any of the dsRNAs of the invention or any of the pharmaceutical compositions of the invention, thereby inhibiting expression of the TMPRSS6 gene in the cell.

In one embodiment, the cell is within a subject, *e.g.*, a human subject, *e.g.*, a subject having a
10 Transmembrane protease, serine 6 (TMPRSS6)-associated disorder, such as a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic porphyria, Parkinson's Disease, Alzheimer's Disease or Friedreich's Ataxia.

15 In some embodiments, the TMPRSS6-associated disorder is β -thalassemia. In one embodiment, the TMPRSS6-associated disorder is β -thalassemia major. In another embodiment, the TMPRSS6-associated disorder is β -thalassemia intermedia. In some embodiments, the TMPRSS6-associated disorder is polycythemia vera.

In certain embodiments, the TMPRSS6 expression is inhibited by at least about 30%, 40%,
20 50%, 60%, 70%, 80%, 90%, or 95%. In one embodiment, inhibiting expression of TMPRSS6 decreases TMPRSS6 protein level in serum of the subject by at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 95%.

In certain embodiments, contacting the cell with the dsRNA agent increases the expression of hepcidin by at least 50%, 60%, 70%, 80%, 90%, or 95%. In one embodiment, increasing
25 expression of hepcidin increases hepcidin protein level in serum of the subject by at least 50%, 60%, 70%, 80%, 90%, or 95%.

In one aspect, the present invention provides a method of treating a subject having a disorder that would benefit from reduction in Transmembrane protease, serine 6 (TMPRSS6) expression. The method includes administering to the subject a therapeutically effective amount of any of the
30 dsRNAs of the invention or any of the pharmaceutical compositions of the invention, thereby treating the subject having the disorder that would benefit from reduction in TMPRSS6 expression.

In another aspect, the present invention provides a method of preventing at least one symptom in a subject having a disorder that would benefit from reduction in Transmembrane protease, serine 6 (TMPRSS6) expression. The method includes administering to the subject a
35 prophylactically effective amount of any of the dsRNAs of the invention or any of the pharmaceutical compositions of the invention, thereby preventing at least one symptom in the subject having the disorder that would benefit from reduction in TMPRSS6 expression.

In certain embodiments, the disorder is a Transmembrane protease, serine 6 (TMPRSS6)-associated disorder, *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective

erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic porphyria, Parkinson's Disease, Alzheimer's Disease or Friedreich's Ataxia .

5 In some embodiments, the TMPRSS6-associated disorder is β -thalassemia. In one embodiment, the TMPRSS6-associated disorder is β -thalassemia major. In another embodiment, the TMPRSS6-associated disorder is β -thalassemia intermedia. In some embodiments, the TMPRSS6-associated disorder is polycythemia vera.

10 In certain embodiments, administration of the dsRNA to the subject causes a decrease in the iron level, ferritin level and/or transferrin saturation level and/or a decrease in TMPRSS6 protein accumulation in the subject. In some embodiments, administration of the dsRNA to the subject causes an increase in the hemoglobin level and/or the hematocrit level in the subject.

15 In a further aspect, the present invention also provides methods of inhibiting the expression of TMPRSS6 in a subject. The methods include administering to the subject a therapeutically effective amount of any of the dsRNAs provided herein, thereby inhibiting the expression of TMPRSS6 in the subject.

 In one embodiment, the subject is human.

 In one embodiment, the dsRNA agent is administered to the subject at a dose of about 0.01 mg/kg to about 50 mg/kg.

20 In one embodiment, the dsRNA agent is administered to the subject subcutaneously or intravenously.

 In one embodiment, the methods of the invention include further determining the level of TMPRSS6 in a sample(s) from the subject.

25 In one embodiment, the level of TMPRSS6 in the subject sample(s) is a TMPRSS6 protein level in a blood, serum or liver sample(s).

 In one embodiment, the methods of the invention include further determining the level of iron and/or hepcidin in a sample(s) from the subject.

30 In certain embodiments, the methods of the invention further comprise administering to the subject an additional therapeutic agent. In one embodiment, the methods of the invention further comprise administering an iron chelator, *e.g.*, deferiprone, deferoxamine, and deferasirox, to a subject.

35 The present invention also provides kits comprising any of the dsRNAs of the invention or any of the pharmaceutical compositions of the invention, and optionally, instructions for use. In one embodiment, the invention provides a kit for performing a method of inhibiting expression of TMPRSS6 gene in a cell by contacting a cell with a double stranded RNAi agent of the invention in an amount effective to inhibit expression of the TMPRSS6 in the cell. The kit comprises an RNAi agent and instructions for use and, optionally, means for administering the RNAi agent to a subject.

 The present invention also provide an RNA-induced silencing complex (RISC) comprising an antisense strand of any of the dsRNA agents of the present invention.

Brief Description of the Drawings

Figure 1 is a schematic depicting the study plan to determine the efficacy of the dsRNA agents disclosed herein *in vivo* in Cynomolgus monkeys.

Figure 2 is a graph showing the percent of serum TMPRSS6 mRNA remaining in Cynomolgous monkeys (n=3 per group) subcutaneously administered a single 3 mg/kg or 10 mg/kg dose of the indicated dsRNA duplexes at Days 21, 22, 57, and 85 post-dose. TMPRSS6 mRNA levels are shown relative to control levels obtained from Cynomolgous monkeys administered PBS as a control.

Figure 3 is a graph showing the plasma iron levels, as a percent of predose levels, in Cynomolgous monkeys (n=3 per group) subcutaneously administered a single 3 mg/kg or 10 mg/kg dose of the indicated dsRNA duplexes at Days 1, 8, 15, 22, 29, 36, 43, 50, 57, 64, 71, 78, and 85 post-dose.

Figure 4 is a graph showing the percent transferrin saturation levels in Cynomolgous monkeys (n=3 per group) subcutaneously administered a single 3 mg/kg or 10 mg/kg dose of the indicated dsRNA duplexes at Days 1, 8, 15, 22, 29, 36, 43, 50, 57, 64, 71, 78, and 85 post-dose.

Detailed Description of the Invention

The present invention provides iRNA compositions which effect the RNA-induced silencing complex (RISC)-mediated cleavage of RNA transcripts of a Transmembrane protease, serine 6 (TMPRSS6) gene. The gene may be within a cell, *e.g.*, a cell within a subject, such as a human. The use of these iRNAs enables the targeted degradation of mRNAs of the corresponding gene (TMPRSS6) in mammals.

The iRNAs of the invention have been designed to target the human Transmembrane protease, serine 6 (TMPRSS6) gene, including portions of the gene that are conserved in the TMPRSS6 orthologs of other mammalian species. Without intending to be limited by theory, it is believed that a combination or sub-combination of the foregoing properties and the specific target sites or the specific modifications in these iRNAs confer to the iRNAs of the invention improved efficacy, stability, potency, durability, and safety.

Accordingly, the present invention provides methods for treating and preventing a Transmembrane protease, serine 6 (TMPRSS6)-associated disorder, *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, using iRNA compositions which effect the RNA-induced silencing complex (RISC)-mediated cleavage of RNA transcripts of a TMPRSS6 gene.

The iRNAs of the invention include an RNA strand (the antisense strand) having a region which is up to about 30 nucleotides or less in length, *e.g.*, 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-

21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 nucleotides in length, which region is substantially complementary to at least part of an mRNA transcript of a TMPRSS6 gene.

In certain embodiments, one or both of the strands of the double stranded RNAi agents of the invention is up to 66 nucleotides in length, *e.g.*, 36-66, 26-36, 25-36, 31-60, 22-43, 27-53 nucleotides
5 in length, with a region of at least 19 contiguous nucleotides that is substantially complementary to at least a part of an mRNA transcript of a TMPRSS6 gene. In some embodiments, such iRNA agents having longer length antisense strands may, for example, include a second RNA strand (the sense strand) of 20-60 nucleotides in length wherein the sense and antisense strands form a duplex of 18-30 contiguous nucleotides.

10 The use of iRNAs of the invention enables the targeted degradation of mRNAs of the corresponding gene (TMPRSS6 gene) in mammals. Using *in vitro* assays, the present inventors have demonstrated that iRNAs targeting a TMPRSS6 gene can potently mediate RNAi, resulting in significant inhibition of expression of a TMPRSS6 gene. Thus, methods and compositions including these iRNAs are useful for treating a subject having a TMPRSS6-associated disorder, *e.g.*, a disorder
15 associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic porphyria, Parkinson's Disease, Alzheimer's Disease or Friedreich's Ataxia.

20 Accordingly, the present invention provides methods and combination therapies for treating a subject having a disorder that would benefit from inhibiting or reducing the expression of a TMPRSS6 gene, *e.g.*, a Transmembrane protease, serine 6 (TMPRSS6)-associated disease, such as a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia),
25 polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, using iRNA compositions which effect the RNA-induced silencing complex (RISC)-mediated cleavage of RNA transcripts of a TMPRSS6 gene.

The present invention also provides methods for preventing at least one symptom in a subject having a disorder that would benefit from inhibiting or reducing the expression of a TMPRSS6 gene,
30 *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic porphyria, Parkinson's Disease, Alzheimer's Disease or Friedreich's Ataxia.

35 The following detailed description discloses how to make and use compositions containing iRNAs to inhibit the expression of a TMPRSS6 gene as well as compositions, uses, and methods for treating subjects that would benefit from inhibition and/or reduction of the expression of a TMPRSS6 gene, *e.g.*, subjects susceptible to or diagnosed with a TMPRSS6-associated disorder.

I. Definitions

In order that the present invention may be more readily understood, certain terms are first defined. In addition, it should be noted that whenever a value or range of values of a parameter are recited, it is intended that values and ranges intermediate to the recited values are also intended to be part of this invention.

The articles "a" and "an" are used herein to refer to one or to more than one (*i.e.*, to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element, *e.g.*, a plurality of elements.

The term "including" is used herein to mean, and is used interchangeably with, the phrase "including but not limited to".

The term "or" is used herein to mean, and is used interchangeably with, the term "and/or," unless context clearly indicates otherwise. For example, "sense strand or antisense strand" is understood as "sense strand or antisense strand or sense strand and antisense strand."

The term "about" is used herein to mean within the typical ranges of tolerances in the art. For example, "about" can be understood as about 2 standard deviations from the mean. In certain embodiments, about means $\pm 10\%$. In certain embodiments, about means $\pm 5\%$. When about is present before a series of numbers or a range, it is understood that "about" can modify each of the numbers in the series or range.

The term "at least", "no less than", or "or more" prior to a number or series of numbers is understood to include the number adjacent to the term "at least", and all subsequent numbers or integers that could logically be included, as clear from context. For example, the number of nucleotides in a nucleic acid molecule must be an integer. For example, "at least 19 nucleotides of a 21 nucleotide nucleic acid molecule" means that 19, 20, or 21 nucleotides have the indicated property. When at least is present before a series of numbers or a range, it is understood that "at least" can modify each of the numbers in the series or range.

As used herein, "no more than" or "or less" is understood as the value adjacent to the phrase and logical lower values or integers, as logical from context, to zero. For example, a duplex with an overhang of "no more than 2 nucleotides" has a 2, 1, or 0 nucleotide overhang. When "no more than" is present before a series of numbers or a range, it is understood that "no more than" can modify each of the numbers in the series or range. As used herein, ranges include both the upper and lower limit.

As used herein, methods of detection can include determination that the amount of analyte present is below the level of detection of the method.

In the event of a conflict between an indicated target site and the nucleotide sequence for a sense or antisense strand, the indicated sequence takes precedence.

In the event of a conflict between a sequence and its indicated site on a transcript or other sequence, the nucleotide sequence recited in the specification takes precedence.

As used herein, "Transmembrane protease, serine 6," used interchangeably with the term "TMPRSS6," refers to the type II plasma membrane serine protease (TTSP) gene or protein. TMPRSS6 is also known as matriptase-2, IRIDA (iron refractory iron-deficiency anemia),

transmembrane protease serine 6, type II transmembrane serine protease 6, and membrane-bound mosaic serine proteinase matriptase-2. TMPRSS6 is a serine protease Type II transmembrane protein of approximately 899 amino acids in length. TMPRSS6 contains multiple domains, *e.g.*, a short endo domain, a transmembrane domain, a sea urchin sperm protein/enteropeptidase domain/agrin (SEA) domain, two complement factor/urchin embryonic growth factor/BMP domains (CUB), three LDL-R class A domains (LDL_A), and a trypsin-like serine protease domain with conserved His-Asp-Ser triad (HDS).

The sequence of a human TMPRSS6 mRNA transcript can be found at, for example, GenBank Accession No. GI: 1755203660 (NM_153609.4; SEQ ID NO:1; reverse complement, SEQ ID NO: 2). The sequence of mouse TMPRSS6 mRNA can be found at, for example, GenBank Accession No. GI: 125656151 (NM_027902.2 ; SEQ ID NO:3; reverse complement, SEQ ID NO: 4). The sequence of rat TMPRSS6 mRNA can be found at, for example, GenBank Accession No. GI: 194474097 (NM_001130556.1 ; SEQ ID NO:5; reverse complement, SEQ ID NO: 6). The sequence of *Macaca fascicularis* TMPRSS6 mRNA can be found at, for example, GenBank Accession No. GI: 982272225 (XM_005567384.2; SEQ ID NO: 7; reverse complement, SEQ ID NO: 8). The sequence of *Macaca mulatta* TMPRSS6 mRNA can be found at, for example, GenBank Accession No. GI: 1622838152 (XM_015150283.2; SEQ ID NO: 9; reverse complement, SEQ ID NO: 10).

Additional examples of TMPRSS6 mRNA sequences are readily available through publicly available databases, *e.g.*, GenBank, UniProt, OMIM, and the *Macaca* genome project web site.

Further information on TMPRSS6 can be found, for example, at www.ncbi.nlm.nih.gov/gene/?term=TMPRSS6.

The entire contents of each of the foregoing GenBank Accession numbers and the Gene database numbers are incorporated herein by reference as of the date of filing this application.

The term TMPRSS6, as used herein, also refers to variations of the TMPRSS6 gene including variants provided in the SNP database. Numerous sequence variations within the TMPRSS6 gene have been identified and may be found at, for example, NCBI dbSNP and UniProt (see, *e.g.*, www.ncbi.nlm.nih.gov/snp/?term=TMPRSS6, the entire contents of which is incorporated herein by reference as of the date of filing this application.

As used herein, “target sequence” refers to a contiguous portion of the nucleotide sequence of an mRNA molecule formed during the transcription of a TMPRSS6 gene, including mRNA that is a product of RNA processing of a primary transcription product. In one embodiment, the target portion of the sequence will be at least long enough to serve as a substrate for iRNA-directed cleavage at or near that portion of the nucleotide sequence of an mRNA molecule formed during the transcription of a TMPRSS6 gene.

The target sequence may be from about 19-36 nucleotides in length, *e.g.*, about 19-30 nucleotides in length. For example, the target sequence can be about 19-30 nucleotides, 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 nucleotides in length. In certain embodiments, the target sequence is 19-23 nucleotides in length,

optionally 21-23 nucleotides in length. Ranges and lengths intermediate to the above recited ranges and lengths are also contemplated to be part of the disclosure.

As used herein, the term “strand comprising a sequence” refers to an oligonucleotide comprising a chain of nucleotides that is described by the sequence referred to using the standard nucleotide nomenclature.

“G,” “C,” “A,” “T,” and “U” each generally stand for a nucleotide that contains guanine, cytosine, adenine, thymidine, and uracil as a base, respectively. However, it will be understood that the term “ribonucleotide” or “nucleotide” can also refer to a modified nucleotide, as further detailed below, or a surrogate replacement moiety (see, *e.g.*, Table 1). The skilled person is well aware that guanine, cytosine, adenine, and uracil can be replaced by other moieties without substantially altering the base pairing properties of an oligonucleotide comprising a nucleotide bearing such replacement moiety. For example, without limitation, a nucleotide comprising inosine as its base can base pair with nucleotides containing adenine, cytosine, or uracil. Hence, nucleotides containing uracil, guanine, or adenine can be replaced in the nucleotide sequences of dsRNA featured in the invention by a nucleotide containing, for example, inosine. In another example, adenine and cytosine anywhere in the oligonucleotide can be replaced with guanine and uracil, respectively to form G-U Wobble base pairing with the target mRNA. Sequences containing such replacement moieties are suitable for the compositions and methods featured in the invention.

The terms “iRNA”, “RNAi agent,” “iRNA agent,” “RNA interference agent” as used interchangeably herein, refer to an agent that contains RNA as that term is defined herein, and which mediates the targeted cleavage of an RNA transcript *via* an RNA-induced silencing complex (RISC) pathway. iRNA directs the sequence-specific degradation of mRNA through a process known as RNA interference (RNAi). The iRNA modulates, *e.g.*, inhibits, the expression of a TMPRSS6 gene in a cell, *e.g.*, a cell within a subject, such as a mammalian subject.

In one embodiment, an RNAi agent of the invention includes a single stranded RNA that interacts with a target RNA sequence, *e.g.*, a TMPRSS6 target mRNA sequence, to direct the cleavage of the target RNA. Without wishing to be bound by theory it is believed that long double stranded RNA introduced into cells is broken down into siRNA by a Type III endonuclease known as Dicer (Sharp *et al.* (2001) *Genes Dev.* 15:485). Dicer, a ribonuclease-III-like enzyme, processes the dsRNA into 19-23 base pair short interfering RNAs with characteristic two base 3' overhangs (Bernstein, *et al.*, (2001) *Nature* 409:363). The siRNAs are then incorporated into an RNA-induced silencing complex (RISC) where one or more helicases unwind the siRNA duplex, enabling the complementary antisense strand to guide target recognition (Nykanen, *et al.*, (2001) *Cell* 107:309). Upon binding to the appropriate target mRNA, one or more endonucleases within the RISC cleave the target to induce silencing (Elbashir, *et al.*, (2001) *Genes Dev.* 15:188). Thus, in one aspect the invention relates to a single stranded RNA (siRNA) generated within a cell and which promotes the formation of a RISC complex to effect silencing of the target gene, *i.e.*, a TMPRSS6 gene. Accordingly, the term “siRNA” is also used herein to refer to an iRNA as described above.

In certain embodiments, the RNAi agent may be a single-stranded siRNA (ssRNAi) that is introduced into a cell or organism to inhibit a target mRNA. Single-stranded RNAi agents bind to the RISC endonuclease, Argonaute 2, which then cleaves the target mRNA. The single-stranded siRNAs are generally 15-30 nucleotides and are chemically modified. The design and testing of single-stranded siRNAs are described in U.S. Patent No. 8,101,348 and in Lima *et al.*, (2012) *Cell* 150:883-894, the entire contents of each of which are hereby incorporated herein by reference. Any of the antisense nucleotide sequences described herein may be used as a single-stranded siRNA as described herein or as chemically modified by the methods described in Lima *et al.*, (2012) *Cell* 150:883-894.

In certain embodiments, an “iRNA” for use in the compositions, uses, and methods of the invention is a double stranded RNA and is referred to herein as a “double stranded RNA agent,” “double stranded RNA (dsRNA) molecule,” “dsRNA agent,” or “dsRNA”. The term “dsRNA”, refers to a complex of ribonucleic acid molecules, having a duplex structure comprising two anti-parallel and substantially complementary nucleic acid strands, referred to as having “sense” and “antisense” orientations with respect to a target RNA, *i.e.*, a TMPRSS6 gene. In some embodiments of the invention, a double stranded RNA (dsRNA) triggers the degradation of a target RNA, *e.g.*, an mRNA, through a post-transcriptional gene-silencing mechanism referred to herein as RNA interference or RNAi.

In general, the majority of nucleotides of each strand of a dsRNA molecule are ribonucleotides, but as described in detail herein, each or both strands can also include one or more non-ribonucleotides, *e.g.*, a deoxyribonucleotide or a modified nucleotide. In addition, as used in this specification, an “iRNA” may include ribonucleotides with chemical modifications; an iRNA may include substantial modifications at multiple nucleotides. As used herein, the term “modified nucleotide” refers to a nucleotide having, independently, a modified sugar moiety, a modified internucleotide linkage, or modified nucleobase, or any combination thereof. Thus, the term modified nucleotide encompasses substitutions, additions or removal of, *e.g.*, a functional group or atom, to internucleoside linkages, sugar moieties, or nucleobases. The modifications suitable for use in the agents of the invention include all types of modifications disclosed herein or known in the art. Any such modifications, as used in a siRNA type molecule, are encompassed by “iRNA” or “RNAi agent” for the purposes of this specification and claims.

In certain embodiments of the instant disclosure, inclusion of a deoxy-nucleotide if present within an RNAi agent can be considered to constitute a modified nucleotide.

The duplex region may be of any length that permits specific degradation of a desired target RNA through a RISC pathway, and may range from about 19 to 36 base pairs in length, *e.g.*, about 19-30 base pairs in length, for example, about 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, or 36 base pairs in length, such as about 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 base pairs in length. In certain embodiments, the duplex region is 19-21 base pairs in length, *e.g.*, 21 base

pairs in length. Ranges and lengths intermediate to the above recited ranges and lengths are also contemplated to be part of the disclosure.

The two strands forming the duplex structure may be different portions of one larger RNA molecule, or they may be separate RNA molecules. Where the two strands are part of one larger molecule, and therefore are connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming the duplex structure, the connecting RNA chain is referred to as a "hairpin loop." A hairpin loop can comprise at least one unpaired nucleotide. In some embodiments, the hairpin loop can comprise at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 23 or more unpaired nucleotides. In some embodiments, the hairpin loop can be 10 or fewer nucleotides. In some embodiments, the hairpin loop can be 8 or fewer unpaired nucleotides. In some embodiments, the hairpin loop can be 4-10 unpaired nucleotides. In some embodiments, the hairpin loop can be 4-8 nucleotides.

Where the two substantially complementary strands of a dsRNA are comprised by separate RNA molecules, those molecules need not be, but can be covalently connected. Where the two strands are connected covalently by means other than an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming the duplex structure, the connecting structure is referred to as a "linker." The RNA strands may have the same or a different number of nucleotides. The maximum number of base pairs is the number of nucleotides in the shortest strand of the dsRNA minus any overhangs that are present in the duplex. In addition to the duplex structure, an RNAi may comprise one or more nucleotide overhangs. In one embodiment of the RNAi agent, at least one strand comprises a 3' overhang of at least 1 nucleotide. In another embodiment, at least one strand comprises a 3' overhang of at least 2 nucleotides, *e.g.*, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14, or 15 nucleotides. In other embodiments, at least one strand of the RNAi agent comprises a 5' overhang of at least 1 nucleotide. In certain embodiments, at least one strand comprises a 5' overhang of at least 2 nucleotides, *e.g.*, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14, or 15 nucleotides. In still other embodiments, both the 3' and the 5' end of one strand of the RNAi agent comprise an overhang of at least 1 nucleotide.

In certain embodiments, an iRNA agent of the invention is a dsRNA, each strand of which comprises 19-23 nucleotides, that interacts with a target RNA sequence, *e.g.*, a TMPRSS6 gene, to direct cleavage of the target RNA.

In some embodiments, an iRNA of the invention is a dsRNA of 24-30 nucleotides that interacts with a target RNA sequence, *e.g.*, a TMPRSS6 target mRNA sequence, to direct the cleavage of the target RNA.

As used herein, the term "nucleotide overhang" refers to at least one unpaired nucleotide that protrudes from the duplex structure of a double stranded iRNA. For example, when a 3'-end of one strand of a dsRNA extends beyond the 5'-end of the other strand, or *vice versa*, there is a nucleotide overhang. A dsRNA can comprise an overhang of at least one nucleotide; alternatively the overhang can comprise at least two nucleotides, at least three nucleotides, at least four nucleotides, at least five nucleotides or more. A nucleotide overhang can comprise or consist of a nucleotide/nucleoside

analog, including a deoxynucleotide/nucleoside. The overhang(s) can be on the sense strand, the antisense strand, or any combination thereof. Furthermore, the nucleotide(s) of an overhang can be present on the 5'-end, 3'-end, or both ends of either an antisense or sense strand of a dsRNA.

In one embodiment, the antisense strand of a dsRNA has a 1-10 nucleotide, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In one embodiment, the sense strand of a dsRNA has a 1-10 nucleotide, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In another embodiment, one or more of the nucleotides in the overhang is replaced with a nucleoside thiophosphate.

In certain embodiments, the antisense strand of a dsRNA has a 1-10 nucleotide, *e.g.*, 0-3, 1-3, 2-4, 2-5, 4-10, 5-10, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In one embodiment, the sense strand of a dsRNA has a 1-10 nucleotide, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In another embodiment, one or more of the nucleotides in the overhang is replaced with a nucleoside thiophosphate.

In certain embodiments, the antisense strand of a dsRNA has a 1-10 nucleotides, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In certain embodiments, the overhang on the sense strand or the antisense strand, or both, can include extended lengths longer than 10 nucleotides, *e.g.*, 1-30 nucleotides, 2-30 nucleotides, 10-30 nucleotides, 10-25 nucleotides, 10-20 nucleotides, or 10-15 nucleotides in length. In certain embodiments, an extended overhang is on the sense strand of the duplex. In certain embodiments, an extended overhang is present on the 3' end of the sense strand of the duplex. In certain embodiments, an extended overhang is present on the 5' end of the sense strand of the duplex. In certain embodiments, an extended overhang is on the antisense strand of the duplex. In certain embodiments, an extended overhang is present on the 3' end of the antisense strand of the duplex. In certain embodiments, an extended overhang is present on the 5' end of the antisense strand of the duplex. In certain embodiments, one or more of the nucleotides in the extended overhang is replaced with a nucleoside thiophosphate. In certain embodiments, the overhang includes a self-complementary portion such that the overhang is capable of forming a hairpin structure that is stable under physiological conditions.

“Blunt” or “blunt end” means that there are no unpaired nucleotides at that end of the double stranded RNA agent, *i.e.*, no nucleotide overhang. A “blunt ended” double stranded RNA agent is double stranded over its entire length, *i.e.*, no nucleotide overhang at either end of the molecule. The RNAi agents of the invention include RNAi agents with no nucleotide overhang at one end (*i.e.*, agents with one overhang and one blunt end) or with no nucleotide overhangs at either end. Most often such a molecule will be double-stranded over its entire length.

The term “antisense strand” or “guide strand” refers to the strand of an iRNA, *e.g.*, a dsRNA, which includes a region that is substantially complementary to a target sequence, *e.g.*, a TMPRSS6 mRNA.

As used herein, the term “region of complementarity” refers to the region on the antisense strand that is substantially complementary to a sequence, for example a target sequence, *e.g.*, a TMPRSS6 nucleotide sequence, as defined herein. Where the region of complementarity is not fully

complementary to the target sequence, the mismatches can be in the internal or terminal regions of the molecule. Generally, the most tolerated mismatches are in the terminal regions, *e.g.*, within 5, 4, or 3 nucleotides of the 5'- or 3'-end of the iRNA. In some embodiments, a double stranded RNA agent of the invention includes a nucleotide mismatch in the antisense strand. In some embodiments, the

5 antisense strand of the double stranded RNA agent of the invention includes no more than 4 mismatches with the target mRNA, *e.g.*, the antisense strand includes 4, 3, 2, 1, or 0 mismatches with the target mRNA. In some embodiments, the antisense strand double stranded RNA agent of the invention includes no more than 4 mismatches with the sense strand, *e.g.*, the antisense strand includes 4, 3, 2, 1, or 0 mismatches with the sense strand. In some embodiments, a double stranded

10 RNA agent of the invention includes a nucleotide mismatch in the sense strand. In some embodiments, the sense strand of the double stranded RNA agent of the invention includes no more than 4 mismatches with the antisense strand, *e.g.*, the sense strand includes 4, 3, 2, 1, or 0 mismatches with the antisense strand. In some embodiments, the nucleotide mismatch is, for example, within 5, 4, 3 nucleotides from the 3'-end of the iRNA. In another embodiment, the nucleotide mismatch is, for

15 example, in the 3'-terminal nucleotide of the iRNA agent. In some embodiments, the mismatch(s) is not in the seed region.

Thus, an RNAi agent as described herein can contain one or more mismatches to the target sequence. In one embodiment, an RNAi agent as described herein contains no more than 3 mismatches (*i.e.*, 3, 2, 1, or 0 mismatches). In one embodiment, an RNAi agent as described herein

20 contains no more than 2 mismatches. In one embodiment, an RNAi agent as described herein contains no more than 1 mismatch. In one embodiment, an RNAi agent as described herein contains 0 mismatches. In certain embodiments, if the antisense strand of the RNAi agent contains mismatches to the target sequence, the mismatch can optionally be restricted to be within the last 5 nucleotides from either the 5'- or 3'-end of the region of complementarity. For example, in such embodiments, for

25 a 23 nucleotide RNAi agent, the strand which is complementary to a region of a TMPRSS6 gene, generally does not contain any mismatch within the central 13 nucleotides. The methods described herein or methods known in the art can be used to determine whether an RNAi agent containing a mismatch to a target sequence is effective in inhibiting the expression of a TMPRSS6 gene. Consideration of the efficacy of RNAi agents with mismatches in inhibiting expression of a

30 TMPRSS6 gene is important, especially if the particular region of complementarity in a TMPRSS6 gene is known to have polymorphic sequence variation within the population.

The term "sense strand" or "passenger strand" as used herein, refers to the strand of an iRNA that includes a region that is substantially complementary to a region of the antisense strand as that term is defined herein.

35 As used herein, "substantially all of the nucleotides are modified" are largely but not wholly modified and can include not more than 5, 4, 3, 2, or 1 unmodified nucleotides.

As used herein, the term "cleavage region" refers to a region that is located immediately adjacent to the cleavage site. The cleavage site is the site on the target at which cleavage occurs. In some embodiments, the cleavage region comprises three bases on either end of, and immediately

adjacent to, the cleavage site. In some embodiments, the cleavage region comprises two bases on either end of, and immediately adjacent to, the cleavage site. In some embodiments, the cleavage site specifically occurs at the site bound by nucleotides 10 and 11 of the antisense strand, and the cleavage region comprises nucleotides 11, 12 and 13.

5 As used herein, and unless otherwise indicated, the term “complementary,” when used to describe a first nucleotide sequence in relation to a second nucleotide sequence, refers to the ability of an oligonucleotide or polynucleotide comprising the first nucleotide sequence to hybridize and form a duplex structure under certain conditions with an oligonucleotide or polynucleotide comprising the second nucleotide sequence, as will be understood by the skilled person. Such conditions can, for
10 example, be stringent conditions, where stringent conditions can include: 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50°C or 70°C for 12-16 hours followed by washing (see, *e.g.*, “Molecular Cloning: A Laboratory Manual, Sambrook, *et al.* (1989) Cold Spring Harbor Laboratory Press). Other conditions, such as physiologically relevant conditions as can be encountered inside an organism, can apply. The skilled person will be able to determine the set of conditions most
15 appropriate for a test of complementarity of two sequences in accordance with the ultimate application of the hybridized nucleotides.

Complementary sequences within an iRNA, *e.g.*, within a dsRNA as described herein, include base-pairing of the oligonucleotide or polynucleotide comprising a first nucleotide sequence to an oligonucleotide or polynucleotide comprising a second nucleotide sequence over the entire length of
20 one or both nucleotide sequences. Such sequences can be referred to as “fully complementary” with respect to each other herein. However, where a first sequence is referred to as “substantially complementary” with respect to a second sequence herein, the two sequences can be fully complementary, or they can form one or more, but generally not more than 5, 4, 3, or 2 mismatched base pairs upon hybridization for a duplex up to 30 base pairs, while retaining the ability to hybridize
25 under the conditions most relevant to their ultimate application, *e.g.*, inhibition of gene expression, in vitro or in vivo. However, where two oligonucleotides are designed to form, upon hybridization, one or more single stranded overhangs, such overhangs shall not be regarded as mismatches with regard to the determination of complementarity. For example, a dsRNA comprising one oligonucleotide 21 nucleotides in length and another oligonucleotide 23 nucleotides in length, wherein the longer
30 oligonucleotide comprises a sequence of 21 nucleotides that is fully complementary to the shorter oligonucleotide, can yet be referred to as “fully complementary” for the purposes described herein.

“Complementary” sequences, as used herein, can also include, or be formed entirely from, non-Watson-Crick base pairs or base pairs formed from non-natural and modified nucleotides, in so far as the above requirements with respect to their ability to hybridize are fulfilled. Such non-Watson-
35 Crick base pairs include, but are not limited to, G:U Wobble or Hoogsteen base pairing.

The terms “complementary,” “fully complementary” and “substantially complementary” herein can be used with respect to the base matching between the sense strand and the antisense strand of a dsRNA, or between two oligonucleotides or polynucleotides, such as the antisense strand of a double stranded RNA agent and a target sequence, as will be understood from the context of their use.

As used herein, a polynucleotide that is “substantially complementary to at least part of” a messenger RNA (mRNA) refers to a polynucleotide that is substantially complementary to a contiguous portion of the mRNA of interest (*e.g.*, an mRNA encoding a TMPRSS6 gene). For example, a polynucleotide is complementary to at least a part of a TMPRSS6 mRNA if the sequence
5 is substantially complementary to a non-interrupted portion of an mRNA encoding a TMPRSS6 gene.

Accordingly, in some embodiments, the antisense polynucleotides disclosed herein are fully complementary to the target TMPRSS6 sequence. In other embodiments, the antisense polynucleotides disclosed herein are substantially complementary to the target TMPRSS6 sequence and comprise a contiguous nucleotide sequence which is at least 80% complementary over its entire
10 length to the equivalent region of the nucleotide sequence of any one of SEQ ID NOs:1, 3, 5, 7, or 9, or a fragment of any one of SEQ ID NOs:1, 3, 5, 7, or 9, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% complementary.

In some embodiments, the antisense polynucleotides disclosed herein are substantially
15 complementary to a fragment of a target TMPRSS6 sequence and comprise a contiguous nucleotide sequence which is at least 80% complementary over its entire length to a fragment of SEQ ID NO: 1 selected from the group of nucleotides 187-210; 227-254; 322-363; 362-390; 398-420; 404-429; 410-435; 439-461; 443-467; 448-474; 460-483; 466-488; 496-519; 519-542; 526-548; 557-593; 641-671; 652-676; 687-713; 725-762; 757-794; 886-908; 921-951; 956-987; 1051-1082; 1233-1269; 1279-
20 1313; 1313-1341; 1327-1351; 1415-1439; 1447-1480; 1464-1486; 1486-1509; 1559-1589; 1571-1595; 1579-1609; 1707-1735; 1738-1764; 1806-1828; 1864-1886; 1934-1966; 1967-1991; 2008-2031; 2015-2043; 2042-2072; 2287-2311; 2297-2354; 2336-2361; 2360-2384; 2416-2438; 2481-2510; 2496-2527; 2526-2558; 2665-2693; 2693-2719; 2707-2729; 2799-2821; 2851-2874; 2971-2999; 2981-3006; and 3155-3195 of SEQ ID NO: 1, such as about 85%, about 90%, about 91%, about
25 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% complementary.

In some embodiments, the antisense polynucleotides disclosed herein are substantially complementary to a fragment of a target TMPRSS6 sequence and comprise a contiguous nucleotide sequence which is at least 80% complementary over its entire length to a fragment of SEQ ID NO: 1
30 selected from the group of nucleotides 230-252, 324-346, 560-578, 560-582, 2338-2360, 3163-3185, 3169-3191, and 3172-3194 of SEQ ID NO: 1, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% complementary.

In some embodiments, the antisense polynucleotides disclosed herein are substantially complementary to a fragment of a target TMPRSS6 sequence and comprise a contiguous nucleotide
35 sequence which is at least 80% complementary over its entire length to a fragment of SEQ ID NO: 1 selected from the group of nucleotides 560-578, 2338-2360, and 3169-3191 of SEQ ID NO: 1, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% complementary.

In other embodiments, the antisense polynucleotides disclosed herein are substantially complementary to the target TMPRSS6 sequence and comprise a contiguous nucleotide sequence which is at least about 80% complementary over its entire length to any one of the sense strand nucleotide sequences in any one of any one of Tables 2-7, or a fragment of any one of the sense strand nucleotide sequences in any one of Tables 2-7, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or 100% complementary.

In one embodiment, an RNAi agent of the disclosure includes a sense strand that is substantially complementary to an antisense polynucleotide which, in turn, is the same as a target TMPRSS6 sequence, and wherein the sense strand polynucleotide comprises a contiguous nucleotide sequence which is at least about 80% complementary over its entire length to the equivalent region of the nucleotide sequence of SEQ ID NOs: 2, 4, 6, 8, or 10, or a fragment of any one of SEQ ID NOs: 2, 4, 6, 8, or 10, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or 100% complementary.

In some embodiments, an iRNA of the invention includes a sense strand that is substantially complementary to an antisense polynucleotide which, in turn, is complementary to a target TMPRSS6 sequence, and wherein the sense strand polynucleotide comprises a contiguous nucleotide sequence which is at least about 80% complementary over its entire length to any one of the antisense strand nucleotide sequences in any one of any one of Tables 2-7, or a fragment of any one of the antisense strand nucleotide sequences in any one of Tables 2-7, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or 100% complementary.

In certain embodiments, the sense and antisense strands are selected from any one of duplexes AD-1556360, AD-1571158, AD-1571033, AD-1554875, AD-1571160, AD-1555117, AD-1554911, and AD-1556915.

In certain embodiments, the sense and antisense strands are selected from any one of duplexes AD-1556360, AD-1571158, and AD-1571033.

In general, an “iRNA” includes ribonucleotides with chemical modifications. Such modifications may include all types of modifications disclosed herein or known in the art. Any such modifications, as used in a dsRNA molecule, are encompassed by “iRNA” for the purposes of this specification and claims.

In certain embodiments of the instant disclosure, inclusion of a deoxy-nucleotide if present within an RNAi agent can be considered to constitute a modified nucleotide.

In an aspect of the invention, an agent for use in the methods and compositions of the invention is a single-stranded antisense oligonucleotide molecule that inhibits a target mRNA *via* an antisense inhibition mechanism. The single-stranded antisense oligonucleotide molecule is complementary to a sequence within the target mRNA. The single-stranded antisense oligonucleotides can inhibit translation in a stoichiometric manner by base pairing to the mRNA and physically obstructing the translation machinery, see Dias, N. *et al.*, (2002) *Mol Cancer Ther* 1:347-

355. The single-stranded antisense oligonucleotide molecule may be about 14 to about 30 nucleotides in length and have a sequence that is complementary to a target sequence. For example, the single-stranded antisense oligonucleotide molecule may comprise a sequence that is at least about 14, 15, 16, 17, 18, 19, 20, or more contiguous nucleotides from any one of the antisense sequences described herein.

The phrase “contacting a cell with an iRNA,” such as a dsRNA, as used herein, includes contacting a cell by any possible means. Contacting a cell with an iRNA includes contacting a cell *in vitro* with the iRNA or contacting a cell *in vivo* with the iRNA. The contacting may be done directly or indirectly. Thus, for example, the iRNA may be put into physical contact with the cell by the individual performing the method, or alternatively, the iRNA may be put into a situation that will permit or cause it to subsequently come into contact with the cell.

Contacting a cell *in vitro* may be done, for example, by incubating the cell with the iRNA. Contacting a cell *in vivo* may be done, for example, by injecting the iRNA into or near the tissue where the cell is located, or by injecting the iRNA into another area, *e.g.*, the bloodstream or the subcutaneous space, such that the agent will subsequently reach the tissue where the cell to be contacted is located. For example, the iRNA may contain or be coupled to a ligand, *e.g.*, GalNAc, that directs the iRNA to a site of interest, *e.g.*, the liver. Combinations of *in vitro* and *in vivo* methods of contacting are also possible. For example, a cell may also be contacted *in vitro* with an iRNA and subsequently transplanted into a subject.

In certain embodiments, contacting a cell with an iRNA includes “introducing” or “delivering the iRNA into the cell” by facilitating or effecting uptake or absorption into the cell. Absorption or uptake of an iRNA can occur through unaided diffusion or active cellular processes, or by auxiliary agents or devices. Introducing an iRNA into a cell may be *in vitro* or *in vivo*. For example, for *in vivo* introduction, iRNA can be injected into a tissue site or administered systemically. *In vitro* introduction into a cell includes methods known in the art such as electroporation and lipofection. Further approaches are described herein below or are known in the art.

The term “lipid nanoparticle” or “LNP” is a vesicle comprising a lipid layer encapsulating a pharmaceutically active molecule, such as a nucleic acid molecule, *e.g.*, an iRNA or a plasmid from which an iRNA is transcribed. LNPs are described in, for example, U.S. Patent Nos. 6,858,225, 6,815,432, 8,158,601, and 8,058,069, the entire contents of which are hereby incorporated herein by reference.

As used herein, a “subject” is an animal, such as a mammal, including a primate (such as a human, a non-human primate, *e.g.*, a monkey, and a chimpanzee), a non-primate (such as a cow, a pig, a horse, a goat, a rabbit, a sheep, a hamster, a guinea pig, a cat, a dog, a rat, or a mouse), or a bird that expresses the target gene, either endogenously or heterologously. In an embodiment, the subject is a human, such as a human being treated or assessed for a disease or disorder that would benefit from reduction in TMPRSS6 expression; a human at risk for a disease or disorder that would benefit from reduction in TMPRSS6 expression; a human having a disease or disorder that would benefit from reduction in TMPRSS6 expression; or human being treated for a disease or disorder that would

benefit from reduction in TMPRSS6 expression as described herein. In some embodiments, the subject is a female human. In other embodiments, the subject is a male human. In one embodiment, the subject is an adult subject. In another embodiment, the subject is a pediatric subject.

As used herein, the terms “treating” or “treatment” refer to a beneficial or desired result, such as reducing at least one sign or symptom of a TMPRSS6-associated disorder in a subject. Treatment also includes a reduction of one or more sign or symptoms associated with unwanted TMPRSS6 expression; diminishing the extent of unwanted TMPRSS6 activation or stabilization; amelioration or palliation of unwanted TMPRSS6 activation or stabilization. “Treatment” can also mean prolonging survival as compared to expected survival in the absence of treatment. The term “lower” in the context of the level of TMPRSS6 in a subject or a disease marker or symptom refers to a statistically significant decrease in such level. The decrease can be, for example, at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or more. In certain embodiments, a decrease is at least 20%. In certain embodiments, the decrease is at least 50% in a disease marker, *e.g.*, protein or gene expression level. “Lower” in the context of the level of TMPRSS6 in a subject is a decrease to a level accepted as within the range of normal for an individual without such disorder. In certain embodiments, “lower” is the decrease in the difference between the level of a marker or symptom for a subject suffering from a disease and a level accepted within the range of normal for an individual, *e.g.*, the level of decrease in bodyweight between an obese individual and an individual having a weight accepted within the range of normal.

As used herein, “prevention” or “preventing,” when used in reference to a disease, disorder or condition thereof, may be treated or ameliorated by a reduction in expression of a TMPRSS6 gene, refers to a reduction in the likelihood that a subject will develop a symptom associated with such a disease, disorder, or condition, *e.g.*, a symptom of unwanted or excessive TMPRSS6 expression, such as elevated iron levels or iron dysregulation. The likelihood of developing elevated iron levels or iron dysregulation is reduced, for example, when an individual having one or more risk factors for elevated iron levels or iron dysregulation either fails to develop elevated iron levels or iron dysregulation, or develops elevated iron levels or iron dysregulation with less severity relative to a population having the same risk factors and not receiving treatment as described herein. The failure to develop a disease, disorder or condition, or the reduction in the development of a symptom associated with such a disease, disorder or condition (*e.g.*, by at least about 10% on a clinically accepted scale for that disease or disorder), or the exhibition of delayed symptoms delayed (*e.g.*, by days, weeks, months or years) is considered effective prevention.

As used herein, the term “Transmembrane protease, serine 6-associated disease” or “TMPRSS6-associated disease,” is a disease or disorder that is caused by, or associated with TMPRSS6 gene expression or TMPRSS6 protein production. The term “TMPRSS6-associated disease” includes a disease, disorder or condition that would benefit from a decrease in TMPRSS6 gene expression, replication, or protein activity.

In some embodiments, the TMPRSS6-associated disease is a disorder associated with iron overload, a condition characterized by elevated iron levels, or iron dysregulation. Iron overload may

be caused, for example, by hereditary conditions, by elevated iron uptake from diet, or by excess iron administered parenterally that includes intravenous injection of excess iron, and transfusional iron overload.

In some embodiments, the TMPRSS6-associated disease is a disorder of ineffective erythropoiesis. Ineffective erythropoiesis is an abnormal expansion of the number of erythroid progenitor cells with unproductive synthesis of enucleated erythrocytes, leading to anemia and hypoxia. In particular, an increase in erythroid cells fails to produce a corresponding increase in red blood cells. As a consequence, iron absorption is still increased in response to stress, but the iron is deposited in the organs rather than being used to generate more erythrocytes.

In some embodiments, TMPRSS6-associated disorders include, but are not limited to, hereditary hemochromatosis, idiopathic hemochromatosis, primary hemochromatosis, secondary hemochromatosis, severe juvenile hemochromatosis, neonatal hemochromatosis, sideroblastic anemia, hemolytic anemia, dyserythropoietic anemia, sickle-cell anemia, hemoglobinopathy, thalassemia (*e.g.*, β -thalassemia and α -thalassemia), polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, chronic liver diseases, porphyria cutanea tarda, erythropoietic porphyria, atransferrinemia, hereditary tyrosinemia, cerebrohepatorenal syndrome, idiopathic pulmonary hemosiderosis, renal hemosiderosis.

In some embodiments, TMPRSS6 associated disorders include disorders associated with oral administration of excess iron, transfusional iron overload and intravenous injection of excess iron.

In other embodiments, TMPRSS6-associated disorders also include disorders with symptoms that are associated with or may be caused by iron overload. Such symptoms include increased risk for liver disease (cirrhosis, cancer), heart attack or heart failure, diabetes mellitus, osteoarthritis, osteoporosis, metabolic syndrome, hypothyroidism, hypogonadism, and in some cases premature death. In one embodiment, TMPRSS6-associated disorders include neurodegenerative disorders associated with iron overload and/or iron dysregulation, such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Friedreich's Ataxia, epilepsy and multiple sclerosis. Administration of an iRNA that targets TMPRSS6, *e.g.*, an iRNA described in any one of Tables 2-7 can treat one or more of these symptoms, or prevent the development or progression of a disease or disorder that is aggravated by increased iron levels.

In one embodiment, a TMPRSS6-associated disorder is a β -thalassemia. A β -thalassemia is any one of a group of hereditary disorders characterized by a genetic deficiency in the synthesis of beta-globin chains. In the homozygous state, beta thalassemia ("thalassemia major") causes severe, transfusion-dependent anemia. In the heterozygous state, the beta thalassemia trait ("thalassemia minor") causes mild to moderate microcytic anemia. "Thalassemia intermedia" is a β -thalassemia that results in subjects in whom the clinical severity of the disease is somewhere between the mild symptoms of β -thalassemia minor and the β -thalassemia major. Several laboratory tests may be used to help detect and diagnose thalassemia, for example, a complete blood count to determine the number of red blood cells and the number of hemoglobin, blood smear test, hemoglobin electrophoresis, gene sequencing, or iron tests to examine the level of iron, ferritin, unsaturated iron binding capacity, total

iron binding capacity, or the transferrin saturation level. The type and relative amounts of hemoglobin present in red blood cells are another indicator for thalassemia. β -thalassemia upsets the balance of beta and alpha hemoglobin chain formation and causes an increase in minor hemoglobin components. So individuals with the β -thalassemia major usually have larger percentages of Hb F. Those with β -thalassemia minor usually have elevated fraction of Hb A2.

In one embodiment, a β -thalassemia is thalassemia major. In another embodiment, a β -thalassemia is thalassemia intermedia.

In some embodiments, the TMPRSS6-associated disorder is polycythemia vera. Polycythemia vera is a type of blood cancer which causes the bone marrow to make excess red blood cells. These excess cells usually thicken the blood vessels, which make the patients more prone to develop blood clots, and other complications such as stroke or heart attack. Several tests may be performed to help detect and diagnose polycythemia vera, for example, a complete blood count, blood smear test, erythropoietin level test, bone marrow aspiration or biopsy, or gene sequencing.

"Therapeutically effective amount," as used herein, is intended to include the amount of an RNAi agent that, when administered to a subject having a TMPRSS6-associated disease, is sufficient to effect treatment of the disease (*e.g.*, by diminishing, ameliorating, or maintaining the existing disease or one or more symptoms of disease). The "therapeutically effective amount" may vary depending on the RNAi agent, how the agent is administered, the disease and its severity and the history, age, weight, family history, genetic makeup, the types of preceding or concomitant treatments, if any, and other individual characteristics of the subject to be treated.

"Prophylactically effective amount," as used herein, is intended to include the amount of an RNAi agent that, when administered to a subject having a TMPRSS6-associated disorder, is sufficient to prevent or ameliorate the disease or one or more symptoms of the disease. Ameliorating the disease includes slowing the course of the disease or reducing the severity of later-developing disease. The "prophylactically effective amount" may vary depending on the RNAi agent, how the agent is administered, the degree of risk of disease, and the history, age, weight, family history, genetic makeup, the types of preceding or concomitant treatments, if any, and other individual characteristics of the patient to be treated.

A "therapeutically-effective amount" or "prophylactically effective amount" also includes an amount of an RNAi agent that produces some desired effect at a reasonable benefit/risk ratio applicable to any treatment. The iRNA employed in the methods of the present invention may be administered in a sufficient amount to produce a reasonable benefit/risk ratio applicable to such treatment.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials (including salts), compositions, or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human subjects and animal subjects without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The phrase "pharmaceutically-acceptable carrier" as used herein means a pharmaceutically-acceptable material, composition, or vehicle, such as a liquid or solid filler, diluent, excipient, manufacturing aid (*e.g.*, lubricant, talc magnesium, calcium or zinc stearate, or steric acid), or solvent encapsulating material, involved in carrying or transporting the subject compound from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the subject being treated. Such carriers are known in the art. Pharmaceutically acceptable carriers include carriers for administration by injection.

The term "sample," as used herein, includes a collection of similar fluids, cells, or tissues isolated from a subject, as well as fluids, cells, or tissues present within a subject. Examples of biological fluids include blood, serum and serosal fluids, plasma, cerebrospinal fluid, ocular fluids, lymph, urine, saliva, and the like. Tissue samples may include samples from tissues, organs, or localized regions. For example, samples may be derived from particular organs, parts of organs, or fluids or cells within those organs. In certain embodiments, samples may be derived from the liver (*e.g.*, whole liver or certain segments of liver or certain types of cells in the liver, such as, *e.g.*, hepatocytes). In some embodiments, a "sample derived from a subject" refers to urine obtained from the subject. A "sample derived from a subject" can refer to blood or blood derived serum or plasma from the subject.

II. iRNAs of the Invention

The present invention provides iRNAs which inhibit the expression of a TMPRSS6 gene. In certain embodiments, the iRNA includes double stranded ribonucleic acid (dsRNA) molecules for inhibiting the expression of a TMPRSS6 gene in a cell, such as a cell within a subject, *e.g.*, a mammal, such as a human susceptible to developing a TMPRSS6-associated disorder, *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic porphyria, Parkinson's Disease, Alzheimer's Disease or Friedreich's Ataxia. The dsRNAi agent includes an antisense strand having a region of complementarity which is complementary to at least a part of an mRNA formed in the expression of a TMPRSS6 gene. The region of complementarity is about 19-30 nucleotides in length (*e.g.*, about 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, or 19 nucleotides in length).

Upon contact with a cell expressing the TMPRSS6 gene, the iRNA inhibits the expression of the TMPRSS6 gene (*e.g.*, a human, a primate, a non-primate, or a rat TMPRSS6 gene) by at least about 50% as assayed by, for example, a PCR or branched DNA (bDNA)-based method, or by a protein-based method, such as by immunofluorescence analysis, using, for example, western blotting or flow cytometric techniques. In certain embodiments, inhibition of expression is determined by the qPCR method provided in the examples herein with the siRNA at, *e.g.*, a 10 nM concentration, in an appropriate organism cell line provided therein. In certain embodiments, inhibition of expression *in*

vivo is determined by knockdown of the human gene in a rodent expressing the human gene, *e.g.*, a mouse or an AAV-infected mouse expressing the human target gene, *e.g.*, when administered as single dose, *e.g.*, at 3 mg/kg at the nadir of RNA expression.

A dsRNA includes two RNA strands that are complementary and hybridize to form a duplex structure under conditions in which the dsRNA will be used. One strand of a dsRNA (the antisense strand) includes a region of complementarity that is substantially complementary, and generally fully complementary, to a target sequence. The target sequence can be derived from the sequence of an mRNA formed during the expression of a TMPRSS6 gene. The other strand (the sense strand) includes a region that is complementary to the antisense strand, such that the two strands hybridize and form a duplex structure when combined under suitable conditions. As described elsewhere herein and as known in the art, the complementary sequences of a dsRNA can also be contained as self-complementary regions of a single nucleic acid molecule, as opposed to being on separate oligonucleotides.

Generally, the duplex structure is 15 to 30 base pairs in length, *e.g.*, 15-29, 15-28, 15-27, 15-26, 15-25, 15-24, 15-23, 15-22, 15-21, 15-20, 15-19, 15-18, 15-17, 18-30, 18-29, 18-28, 18-27, 18-26, 18-25, 18-24, 18-23, 18-22, 18-21, 18-20, 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 base pairs in length. In certain embodiments, the duplex structure is 18 to 25 base pairs in length, *e.g.*, 18-25, 18-24, 18-23, 18-22, 18-21, 18-20, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-25, 20-24, 20-23, 20-22, 20-21, 21-25, 21-24, 21-23, 21-22, 22-25, 22-24, 22-23, 23-25, 23-24 or 24-25 base pairs in length, for example, 19-21 basepairs in length. Ranges and lengths intermediate to the above recited ranges and lengths are also contemplated to be part of the disclosure.

Similarly, the region of complementarity to the target sequence is 15 to 30 nucleotides in length, *e.g.*, 15-29, 15-28, 15-27, 15-26, 15-25, 15-24, 15-23, 15-22, 15-21, 15-20, 15-19, 15-18, 15-17, 18-30, 18-29, 18-28, 18-27, 18-26, 18-25, 18-24, 18-23, 18-22, 18-21, 18-20, 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 nucleotides in length, for example 19-23 nucleotides in length or 21-23 nucleotides in length. Ranges and lengths intermediate to the above recited ranges and lengths are also contemplated to be part of the disclosure.

In some embodiments, the duplex structure is 19 to 30 base pairs in length. Similarly, the region of complementarity to the target sequence is 19 to 30 nucleotides in length.

In some embodiments, the dsRNA is about 19 to about 23 nucleotides in length, or about 25 to about 30 nucleotides in length. In general, the dsRNA is long enough to serve as a substrate for the Dicer enzyme. For example, it is well-known in the art that dsRNAs longer than about 21-23 nucleotides in length may serve as substrates for Dicer. As the ordinarily skilled person will also recognize, the region of an RNA targeted for cleavage will most often be part of a larger RNA molecule, often an mRNA molecule. Where relevant, a “part” of an mRNA target is a contiguous

sequence of an mRNA target of sufficient length to allow it to be a substrate for RNAi-directed cleavage (*i.e.*, cleavage through a RISC pathway).

One of skill in the art will also recognize that the duplex region is a primary functional portion of a dsRNA, *e.g.*, a duplex region of about 19 to about 30 base pairs, *e.g.*, about 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 base pairs. Thus, in one embodiment, to the extent that it becomes processed to a functional duplex, of *e.g.*, 15-30 base pairs, that targets a desired RNA for cleavage, an RNA molecule or complex of RNA molecules having a duplex region greater than 30 base pairs is a dsRNA. Thus, an ordinarily skilled artisan will recognize that in one embodiment, a miRNA is a dsRNA. In another embodiment, a dsRNA is not a naturally occurring miRNA. In another embodiment, an iRNA agent useful to target TMPRSS6 gene expression is not generated in the target cell by cleavage of a larger dsRNA.

A dsRNA as described herein can further include one or more single-stranded nucleotide overhangs, *e.g.*, 1-4, 2-4, 1-3, 2-3, 1, 2, 3, or 4 nucleotides. dsRNAs having at least one nucleotide overhang can have superior inhibitory properties relative to their blunt-ended counterparts. A nucleotide overhang can comprise or consist of a nucleotide/nucleoside analog, including a deoxynucleotide/nucleoside. The overhang(s) can be on the sense strand, the antisense strand, or any combination thereof. Furthermore, the nucleotide(s) of an overhang can be present on the 5'-end, 3'-end, or both ends of an antisense or sense strand of a dsRNA.

A dsRNA can be synthesized by standard methods known in the art. Double stranded RNAi compounds of the invention may be prepared using a two-step procedure. First, the individual strands of the double stranded RNA molecule are prepared separately. Then, the component strands are annealed. The individual strands of the siRNA compound can be prepared using solution-phase or solid-phase organic synthesis or both. Organic synthesis offers the advantage that the oligonucleotide strands comprising unnatural or modified nucleotides can be easily prepared. Similarly, single-stranded oligonucleotides of the invention can be prepared using solution-phase or solid-phase organic synthesis or both.

In an aspect, a dsRNA of the invention includes at least two nucleotide sequences, a sense sequence and an anti-sense sequence. The sense strand is selected from the group of sequences provided in any one of Tables 2-7, and the corresponding antisense strand of the sense strand is selected from the group of sequences of any one of Tables 2-7. In this aspect, one of the two sequences is complementary to the other of the two sequences, with one of the sequences being substantially complementary to a sequence of an mRNA generated in the expression of a TMPRSS6 gene. As such, in this aspect, a dsRNA will include two oligonucleotides, where one oligonucleotide is described as the sense strand in any one of Tables 2-7, and the second oligonucleotide is described as the corresponding antisense strand of the sense strand in any one of Tables 2-7.

In certain embodiments, the substantially complementary sequences of the dsRNA are contained on separate oligonucleotides. In other embodiments, the substantially complementary sequences of the dsRNA are contained on a single oligonucleotide.

In one embodiment, the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, or 20, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from any one of the antisense strand nucleotide sequences in any one of Tables 2-7.

It will be understood that, although the sequences in, for example, Tables 3 or 5, are not described as modified or conjugated sequences, the RNA of the iRNA of the invention *e.g.*, a dsRNA of the invention, may comprise any one of the sequences set forth in any one of Tables 2-7 that is unmodified, un-conjugated, or modified or conjugated differently than described therein. In other words, the invention encompasses dsRNA of Tables 2-7 which are un-modified, un-conjugated, modified, or conjugated, as described herein.

The skilled person is well aware that dsRNAs having a duplex structure of about 20 to 23 base pairs, *e.g.*, 21, base pairs have been hailed as particularly effective in inducing RNA interference (Elbashir *et al.*, *EMBO* 2001, 20:6877-6888). However, others have found that shorter or longer RNA duplex structures can also be effective (Chu and Rana (2007) *RNA* 14:1714-1719; Kim *et al.* (2005) *Nat Biotech* 23:222-226). In the embodiments described above, by virtue of the nature of the oligonucleotide sequences provided in any one of Tables 2-7, dsRNAs described herein can include at least one strand of a length of minimally 21 nucleotides. It can be reasonably expected that shorter duplexes having any one of the sequences in any one of Tables 2-7 minus only a few nucleotides on one or both ends can be similarly effective as compared to the dsRNAs described above. Hence, dsRNAs having a sequence of at least 19, 20, or more contiguous nucleotides derived from any one of the sequences of any one of Tables 2-7, and differing in their ability to inhibit the expression of a TMPRSS6 gene by not more than about 5, 10, 15, 20, 25, or 30 % inhibition from a dsRNA comprising the full sequence, are contemplated to be within the scope of the present invention.

In addition, the RNAs provided in Tables 2-7 identify a site(s) in a TMPRSS6 transcript that is susceptible to RISC-mediated cleavage. As such, the present invention further features iRNAs that target within one of these sites. As used herein, an iRNA is said to target within a particular site of an RNA transcript if the iRNA promotes cleavage of the transcript anywhere within that particular site. Such an iRNA will generally include at least about 19 contiguous nucleotides from any one of the sequences provided in any one of Tables 2-7 coupled to additional nucleotide sequences taken from the region contiguous to the selected sequence in a TMPRSS6 gene.

III. Modified iRNAs of the Invention

In certain embodiments, the RNA of the iRNA of the invention *e.g.*, a dsRNA, is unmodified, and does not comprise, *e.g.*, chemical modifications or conjugations known in the art and described herein. In other embodiments, the RNA of an iRNA of the invention, *e.g.*, a dsRNA, is chemically modified to enhance stability or other beneficial characteristics. In certain embodiments of the invention, substantially all of the nucleotides of an iRNA of the invention are modified. In other embodiments of the invention, all of the nucleotides of an iRNA or substantially all of the nucleotides of an iRNA are modified, *i.e.*, not more than 5, 4, 3, 2, or 1 unmodified nucleotides are present in a strand of the iRNA.

The nucleic acids featured in the invention can be synthesized or modified by methods well established in the art, such as those described in "Current protocols in nucleic acid chemistry," Beaucage, S.L. *et al.* (Edrs.), John Wiley & Sons, Inc., New York, NY, USA, which is hereby incorporated herein by reference. Modifications include, for example, end modifications, *e.g.*, 5'-end
5 modifications (phosphorylation, conjugation, inverted linkages) or 3'-end modifications (conjugation, DNA nucleotides, inverted linkages, *etc.*); base modifications, *e.g.*, replacement with stabilizing bases, destabilizing bases, or bases that base pair with an expanded repertoire of partners, removal of bases (abasic nucleotides), or conjugated bases; sugar modifications (*e.g.*, at the 2'-position or 4'-position) or replacement of the sugar; or backbone modifications, including modification or
10 replacement of the phosphodiester linkages. Specific examples of iRNA compounds useful in the embodiments described herein include, but are not limited to RNAs containing modified backbones or no natural internucleoside linkages. RNAs having modified backbones include, among others, those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified RNAs that do not have a phosphorus atom in their
15 internucleoside backbone can also be considered to be oligonucleosides. In some embodiments, a modified iRNA will have a phosphorus atom in its internucleoside backbone.

Modified RNA backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl
phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates,
20 phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5'-linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included. In some embodiments of the
25 invention, the dsRNA agents of the invention are in a free acid form. In other embodiments of the invention, the dsRNA agents of the invention are in a salt form. In one embodiment, the dsRNA agents of the invention are in a sodium salt form. In certain embodiments, when the dsRNA agents of the invention are in the sodium salt form, sodium ions are present in the agent as counterions for substantially all of the phosphodiester and/or phosphorothioate groups present in the agent. Agents
30 in which substantially all of the phosphodiester and/or phosphorothioate linkages have a sodium counterion include not more than 5, 4, 3, 2, or 1 phosphodiester and/or phosphorothioate linkages without a sodium counterion. In some embodiments, when the dsRNA agents of the invention are in the sodium salt form, sodium ions are present in the agent as counterions for all of the phosphodiester and/or phosphorothioate groups present in the agent.

35 Representative U.S. Patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S. Patent Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,195; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,316; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,625,050; 6,028,188; 6,124,445; 6,160,109; 6,169,170; 6,172,209; 6,239,265;

6,277,603; 6,326,199; 6,346,614; 6,444,423; 6,531,590; 6,534,639; 6,608,035; 6,683,167; 6,858,715; 6,867,294; 6,878,805; 7,015,315; 7,041,816; 7,273,933; 7,321,029; and U.S. Pat RE39464, the entire contents of each of which are hereby incorporated herein by reference.

Modified RNA backbones that do not include a phosphorus atom therein have backbones that
 5 are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatoms and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones;
 10 alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S, and CH₂ component parts.

Representative U.S. Patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Patent Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141;
 15 5,235,033; 5,64,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, the entire contents of each of which are hereby incorporated herein by reference.

Suitable RNA mimetics are contemplated for use in iRNAs provided herein, in which both the
 20 sugar and the internucleoside linkage, *i.e.*, the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound in which an RNA mimetic that has been shown to have excellent hybridization properties is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar backbone of an RNA is replaced with an amide containing backbone, in particular an
 25 aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative US patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Patent Nos. 5,539,082; 5,714,331; and 5,719,262, the entire contents of each of which are hereby incorporated herein by reference. Additional PNA compounds suitable for use in the iRNAs of the invention are described in, for
 30 example, in Nielsen *et al.*, *Science*, 1991, 254, 1497-1500.

Some embodiments featured in the invention include RNAs with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular --CH₂--NH--CH₂--, --CH₂--N(CH₃)--O--CH₂--[known as a methylene (methylimino) or MMI backbone], --CH₂--O--N(CH₃)--CH₂--, --CH₂--N(CH₃)--N(CH₃)--CH₂-- and --N(CH₃)--CH₂--CH₂-- of the above-referenced U.S.
 35 Patent No. 5,489,677, and the amide backbones of the above-referenced U.S. Patent No. 5,602,240. In some embodiments, the RNAs featured herein have morpholino backbone structures of the above-referenced U.S. Patent No. 5,034,506. The native phosphodiester backbone can be represented as O-P(O)(OH)-OCH₂--.

Modified RNAs can also contain one or more substituted sugar moieties. The iRNAs, *e.g.*, dsRNAs, featured herein can include one of the following at the 2'-position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl can be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Exemplary suitable modifications include O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. In other embodiments, dsRNAs include one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an iRNA, or a group for improving the pharmacodynamic properties of an iRNA, and other substituents having similar properties. In some embodiments, the modification includes a 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin *et al.*, *Helv. Chim. Acta*, 1995, 78:486-504) *i.e.*, an alkoxy-alkoxy group. Another exemplary modification is 2'-dimethylaminoethoxy, *i.e.*, a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples herein below, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), *i.e.*, 2'-O-CH₂-O-CH₂-N(CH₃)₂. Further exemplary modifications include: 5'-Me-2'-F nucleotides, 5'-Me-2'-OMe nucleotides, 5'-Me-2'-deoxynucleotides, (both R and S isomers in these three families); 2'-alkoxyalkyl; and 2'-NMA (N-methylacetamide).

Other modifications include 2'-methoxy (2'-OCH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications can also be made at other positions on the RNA of an iRNA, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked dsRNAs and the 5' position of 5' terminal nucleotide. iRNAs can also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative US patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Patent Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of which are commonly owned with the instant application. The entire contents of each of the foregoing are hereby incorporated herein by reference.

An iRNA can also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C), and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as deoxythymidine (dT), 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil


(pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo, particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-daazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in *Modified Nucleosides in Biochemistry, Biotechnology and Medicine*, Herdewijn, P. ed. Wiley-VCH, 2008; those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J. L., ed. John Wiley & Sons, 1990, those disclosed by Englisch *et al.*, *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y. S., Chapter 15, *dsRNA Research and Applications*, pages 289-302, Crooke, S. T. and Lebleu, B., Ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds featured in the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and 0-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., Eds., *dsRNA Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are exemplary base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative U.S. Patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Patent Nos. 3,687,808, 4,845,205; 5,130,30; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; 5,681,941; 5,750,692; 6,015,886; 6,147,200; 6,166,197; 6,222,025; 6,235,887; 6,380,368; 6,528,640; 6,639,062; 6,617,438; 7,045,610; 7,427,672; and 7,495,088, the entire contents of each of which are hereby incorporated herein by reference.

In some embodiments, an RNAi agent of the disclosure can also be modified to include one or more bicyclic sugar moieties. A "bicyclic sugar" is a furanosyl ring modified by a ring formed by the bridging of two carbons, whether adjacent or non-adjacent. A "bicyclic nucleoside" ("BNA") is a nucleoside having a sugar moiety comprising a ring formed by bridging two carbons, whether adjacent or non-adjacent, of the sugar ring, thereby forming a bicyclic ring system. In certain embodiments, the bridge connects the 4'-carbon and the 2'-carbon of the sugar ring, optionally, via the 2'-acyclic oxygen atom. Thus, in some embodiments an agent of the invention may include one or more locked nucleic acids (LNA). A locked nucleic acid is a nucleotide having a modified ribose moiety in which the ribose moiety comprises an extra bridge connecting the 2' and 4' carbons. In other words, an LNA is a nucleotide comprising a bicyclic sugar moiety comprising a 4'-CH₂-O-2' bridge. This structure effectively "locks" the ribose in the 3'-endo structural conformation. The addition of locked nucleic acids to siRNAs has been shown to increase siRNA stability in serum, and to reduce off-target effects (Elmen, J. *et al.*, (2005) *Nucleic Acids Research* 33(1):439-447; Mook, OR. *et al.*, (2007) *Mol Canc Ther* 6(3):833-843; Grunweller, A. *et al.*, (2003) *Nucleic Acids Research* 31(12):3185-3193). Examples of bicyclic nucleosides for use in the polynucleotides of the invention

include without limitation nucleosides comprising a bridge between the 4' and the 2' ribosyl ring atoms. In certain embodiments, the antisense polynucleotide agents of the invention include one or more bicyclic nucleosides comprising a 4' to 2' bridge.

A locked nucleoside can be represented by the structure (omitting stereochemistry),



5

wherein B is a nucleobase or modified nucleobase and L is the linking group that joins the 2'-carbon to the 4'-carbon of the ribose ring. Examples of such 4' to 2' bridged bicyclic nucleosides, include but are not limited to 4'-(CH₂)—O-2' (LNA); 4'-(CH₂)—S-2'; 4'-(CH₂)₂—O-2' (ENA); 4'-CH(CH₃)—O-2' (also referred to as "constrained ethyl" or "cEt") and 4'-CH(CH₂OCH₃)—O-2' (and analogs thereof; see, *e.g.*, U.S. Patent No. 7,399,845); 4'-C(CH₃)(CH₃)—O-2' (and analogs thereof; see *e.g.*, U.S. Patent No. 8,278,283); 4'-CH₂—N(OCH₃)-2' (and analogs thereof; see *e.g.*, U.S. Patent No. 8,278,425); 4'-CH₂—O—N(CH₃)-2' (see, *e.g.*, U.S. Patent Publication No. 2004/0171570); 4'-CH₂—N(R)—O-2', wherein R is H, C1-C12 alkyl, or a nitrogen protecting group (see, *e.g.*, U.S. Patent No. 7,427,672); 4'-CH₂—C(H)(CH₃)-2' (see, *e.g.*, Chattopadhyaya *et al.*, *J. Org. Chem.*, 2009, 74, 118-134); and 4'-CH₂—C(=CH₂)-2' (and analogs thereof; see, *e.g.*, U.S. Patent No. 8,278,426). The entire contents of each of the foregoing are hereby incorporated herein by reference.

Additional representative U.S. Patents and U.S. Patent Publications that teach the preparation of locked nucleic acid nucleotides include, but are not limited to, the following: U.S. Patent Nos. 6,268,490; 6,525,191; 6,670,461; 6,770,748; 6,794,499; 6,998,484; 7,053,207; 7,034,133; 7,084,125; 7,399,845; 7,427,672; 7,569,686; 7,741,457; 8,022,193; 8,030,467; 8,278,425; 8,278,426; 8,278,283; US 2008/0039618; and US 2009/0012281, the entire contents of each of which are hereby incorporated herein by reference.

Any of the foregoing bicyclic nucleosides can be prepared having one or more stereochemical sugar configurations including for example α-L-ribofuranose and β-D-ribofuranose (see WO 99/14226).

The RNA of an iRNA can also be modified to include one or more constrained ethyl nucleotides. As used herein, a "constrained ethyl nucleotide" or "cEt" is a locked nucleic acid comprising a bicyclic sugar moiety comprising a 4'-CH(CH₃)-O-2' bridge (*i.e.*, L in the preceding structure). In one embodiment, a constrained ethyl nucleotide is in the S conformation referred to herein as "S-cEt."

An iRNA of the invention may also include one or more "conformationally restricted nucleotides" ("CRN"). CRN are nucleotide analogs with a linker connecting the C2' and C4' carbons of ribose or the C3 and -C5' carbons of ribose. CRN lock the ribose ring into a stable conformation

and increase the hybridization affinity to mRNA. The linker is of sufficient length to place the oxygen in an optimal position for stability and affinity resulting in less ribose ring puckering.

Representative publications that teach the preparation of certain of the above noted CRN include, but are not limited to, U.S. Patent Publication No. 2013/0190383; and PCT publication WO 2013/036868, the entire contents of each of which are hereby incorporated herein by reference.

In some embodiments, an iRNA of the invention comprises one or more monomers that are UNA (unlocked nucleic acid) nucleotides. UNA is unlocked acyclic nucleic acid, wherein any of the bonds of the sugar has been removed, forming an unlocked "sugar" residue. In one example, UNA also encompasses monomer with bonds between C1'-C4' have been removed (*i.e.* the covalent carbon-oxygen-carbon bond between the C1' and C4' carbons). In another example, the C2'-C3' bond (*i.e.* the covalent carbon-carbon bond between the C2' and C3' carbons) of the sugar has been removed (see *Nuc. Acids Symp. Series*, 52, 133-134 (2008) and Fluiter *et al.*, *Mol. Biosyst.*, 2009, 10, 1039 hereby incorporated by reference).

Representative U.S. publications that teach the preparation of UNA include, but are not limited to, U.S. Patent No. 8,314,227; and U.S. Patent Publication Nos. 2013/0096289; 2013/0011922; and 2011/0313020, the entire contents of each of which are hereby incorporated herein by reference.

Potentially stabilizing modifications to the ends of RNA molecules can include N-(acetylaminocaproyl)-4-hydroxyprolinol (Hyp-C6-NHAc), N-(caproyl-4-hydroxyprolinol (Hyp-C6), N-(acetyl-4-hydroxyprolinol (Hyp-NHAc), thymidine-2'-O-deoxythymidine (ether), N-(aminocaproyl)-4-hydroxyprolinol (Hyp-C6-amino), 2-docosanoyl-uridine-3'-phosphate, inverted 2'-deoxy-modified ribonucleotide, such as inverted dT(idT), inverted dA (idA), and inverted abasic 2'-deoxyribonucleotide (iAb) and others. Disclosure of this modification can be found in WO 2011/005861.

In one example, the 3' or 5' terminal end of an oligonucleotide is linked to an inverted 2'-deoxy-modified ribonucleotide, such as inverted dT(idT), inverted dA (idA), or an inverted abasic 2'-deoxyribonucleotide (iAb). In one particular example, the inverted 2'-deoxy-modified ribonucleotide is linked to the 3'-end of an oligonucleotide, such as the 3'-end of a sense strand described herein, where the linking is via a 3'-3' phosphodiester linkage or a 3'-3'-phosphorothioate linkage.

In another example, the 3'-end of a sense strand is linked via a 3'-3'-phosphorothioate linkage to an inverted abasic ribonucleotide (iAb). In another example, the 3'-end of a sense strand is linked via a 3'-3'-phosphorothioate linkage to an inverted dA (idA).

In one particular example, the inverted 2'-deoxy-modified ribonucleotide is linked to the 3'-end of an oligonucleotide, such as the 3'-end of a sense strand described herein, where the linking is via a 3'-3' phosphodiester linkage or a 3'-3'-phosphorothioate linkage.

In another example, the 3'-terminal nucleotides of a sense strand is an inverted dA (idA) and is linked to the preceding nucleotide via a 3'-3'- linkage (*e.g.*, 3'-3'-phosphorothioate linkage).

Other modifications of the nucleotides of an iRNA of the invention include a 5' phosphate or 5' phosphate mimic, *e.g.*, a 5'-terminal phosphate or phosphate mimic on the antisense strand of an

iRNA. Suitable phosphate mimics are disclosed in, for example U.S. Patent Publication No. 2012/0157511, the entire contents of which are incorporated herein by reference.

A. Modified iRNAs Comprising Motifs of the Invention

5 In certain aspects of the invention, the double stranded RNA agents of the invention include agents with chemical modifications as disclosed, for example, in WO2013/075035, the entire contents of each of which are incorporated herein by reference. As shown herein and in WO2013/075035, one or more motifs of three identical modifications on three consecutive nucleotides may be introduced into a sense strand or antisense strand of a dsRNAi agent, particularly at or near the cleavage site. In
10 some embodiments, the sense strand and antisense strand of the dsRNAi agent may otherwise be completely modified. The introduction of these motifs interrupts the modification pattern, if present, of the sense or antisense strand. The dsRNAi agent may be optionally conjugated with a GalNAc derivative ligand, for instance on the sense strand.

More specifically, when the sense strand and antisense strand of the double stranded RNA
15 agent are completely modified to have one or more motifs of three identical modifications on three consecutive nucleotides at or near the cleavage site of at least one strand of a dsRNAi agent, the gene silencing activity of the dsRNAi agent was observed.

Accordingly, the invention provides double stranded RNA agents capable of inhibiting the expression of a target gene (*i.e.*, TMPRSS6 gene) *in vivo*. The RNAi agent comprises a sense strand
20 and an antisense strand. Each strand of the RNAi agent may be, for example, 17-30 nucleotides in length, 25-30 nucleotides in length, 27-30 nucleotides in length, 19-25 nucleotides in length, 19-23 nucleotides in length, 19-21 nucleotides in length, 21-25 nucleotides in length, or 21-23 nucleotides in length.

The sense strand and antisense strand typically form a duplex double stranded RNA
25 (“dsRNA”), also referred to herein as “dsRNAi agent.” The duplex region of a dsRNAi agent may be, for example, the duplex region can be 27-30 nucleotide pairs in length, 19-25 nucleotide pairs in length, 19-23 nucleotide pairs in length, 19- 21 nucleotide pairs in length, 21-25 nucleotide pairs in length, or 21-23 nucleotide pairs in length. In another example, the duplex region is selected from 19, 20, 21, 22, 23, 24, 25, 26, and 27 nucleotides in length.

30 In certain embodiments, the dsRNAi agent may contain one or more overhang regions or capping groups at the 3’-end, 5’-end, or both ends of one or both strands. The overhang can be, independently, 1-6 nucleotides in length, for instance 2-6 nucleotides in length, 1-5 nucleotides in length, 2-5 nucleotides in length, 1-4 nucleotides in length, 2-4 nucleotides in length, 1-3 nucleotides in length, 2-3 nucleotides in length, or 1-2 nucleotides in length. In certain embodiments, the
35 overhang regions can include extended overhang regions as provided above. The overhangs can be the result of one strand being longer than the other, or the result of two strands of the same length being staggered. The overhang can form a mismatch with the target mRNA or it can be complementary to the gene sequences being targeted or can be another sequence. The first and

second strands can also be joined, *e.g.*, by additional bases to form a hairpin, or by other non-base linkers.

In certain embodiments, the nucleotides in the overhang region of the dsRNAi agent can each independently be a modified or unmodified nucleotide including, but not limited to 2'-sugar modified, such as, 2'-F, 2'-O-methyl, thymidine (T), 2'-O-methoxyethyl-5-methyluridine (Teo), 2'-O-methoxyethyladenosine (Aeo), 2'-O-methoxyethyl-5-methylcytidine (m5Ceo), and any combinations thereof.

For example, TT can be an overhang sequence for either end on either strand. The overhang can form a mismatch with the target mRNA or it can be complementary to the gene sequences being targeted or can be another sequence.

The 5'- or 3'- overhangs at the sense strand, antisense strand, or both strands of the dsRNAi agent may be phosphorylated. In some embodiments, the overhang region(s) contains two nucleotides having a phosphorothioate between the two nucleotides, where the two nucleotides can be the same or different. In some embodiments, the overhang is present at the 3'-end of the sense strand, antisense strand, or both strands. In some embodiments, this 3'-overhang is present in the antisense strand. In some embodiments, this 3'-overhang is present in the sense strand.

The dsRNAi agent may contain only a single overhang, which can strengthen the interference activity of the RNAi, without affecting its overall stability. For example, the single-stranded overhang may be located at the 3'-end of the sense strand or, alternatively, at the 3'-end of the antisense strand. The RNAi may also have a blunt end, located at the 5'-end of the antisense strand (*i.e.*, the 3'-end of the sense strand) or *vice versa*. Generally, the antisense strand of the dsRNAi agent has a nucleotide overhang at the 3'-end, and the 5'-end is blunt. While not wishing to be bound by theory, the asymmetric blunt end at the 5'-end of the antisense strand and 3'-end overhang of the antisense strand favor the guide strand loading into RISC process.

In certain embodiments, the dsRNAi agent is a double blunt-ended of 19 nucleotides in length, wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides at positions 7, 8, 9 from the 5'end. The antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at positions 11, 12, and 13 from the 5'end.

In other embodiments, the dsRNAi agent is a double blunt-ended of 20 nucleotides in length, wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides at positions 8, 9, and 10 from the 5'end. The antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at positions 11, 12, and 13 from the 5'end.

In yet other embodiments, the dsRNAi agent is a double blunt-ended of 21 nucleotides in length, wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides at positions 9, 10, and 11 from the 5'end. The antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at positions 11, 12, and 13 from the 5'end.

In certain embodiments, the dsRNAi agent comprises a 21 nucleotide sense strand and a 23 nucleotide antisense strand, wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides at positions 9, 10, and 11 from the 5' end; the antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at positions 11, 12, and 13 from the 5' end, wherein one end of the RNAi agent is blunt, while the other end comprises a 2 nucleotide overhang. In one embodiment, the 2 nucleotide overhang is at the 3'-end of the antisense strand.

When the 2 nucleotide overhang is at the 3'-end of the antisense strand, there may be two phosphorothioate internucleotide linkages between the terminal three nucleotides, wherein two of the three nucleotides are the overhang nucleotides, and the third nucleotide is a paired nucleotide next to the overhang nucleotide. In one embodiment, the RNAi agent additionally has two phosphorothioate internucleotide linkages between the terminal three nucleotides at both the 5'-end of the sense strand and at the 5'-end of the antisense strand. In certain embodiments, every nucleotide in the sense strand and the antisense strand of the dsRNAi agent, including the nucleotides that are part of the motifs are modified nucleotides. In certain embodiments each residue is independently modified with a 2'-O-methyl or 3'-fluoro, *e.g.*, in an alternating motif. Optionally, the dsRNAi agent further comprises a ligand (such as, GalNAc₃).

In certain embodiments, the dsRNAi agent comprises a sense and an antisense strand, wherein the sense strand is 25-30 nucleotide residues in length, wherein starting from the 5' terminal nucleotide (position 1) positions 1 to 23 of the first strand comprise at least 8 ribonucleotides; the antisense strand is 36-66 nucleotide residues in length and, starting from the 3' terminal nucleotide, comprises at least 8 ribonucleotides in the positions paired with positions 1- 23 of sense strand to form a duplex; wherein at least the 3' terminal nucleotide of antisense strand is unpaired with sense strand, and up to 6 consecutive 3' terminal nucleotides are unpaired with sense strand, thereby forming a 3' single stranded overhang of 1-6 nucleotides; wherein the 5' terminus of antisense strand comprises from 10-30 consecutive nucleotides which are unpaired with sense strand, thereby forming a 10-30 nucleotide single stranded 5' overhang; wherein at least the sense strand 5' terminal and 3' terminal nucleotides are base paired with nucleotides of antisense strand when sense and antisense strands are aligned for maximum complementarity, thereby forming a substantially duplexed region between sense and antisense strands; and antisense strand is sufficiently complementary to a target RNA along at least 19 ribonucleotides of antisense strand length to reduce target gene expression when the double stranded nucleic acid is introduced into a mammalian cell; and wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides, where at least one of the motifs occurs at or near the cleavage site. The antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at or near the cleavage site.

In certain embodiments, the dsRNAi agent comprises sense and antisense strands, wherein the dsRNAi agent comprises a first strand having a length which is at least 25 and at most 29 nucleotides and a second strand having a length which is at most 30 nucleotides with at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at position 11, 12, 13 from the 5' end;

wherein the 3' end of the first strand and the 5' end of the second strand form a blunt end and the second strand is 1-4 nucleotides longer at its 3' end than the first strand, wherein the duplex region which is at least 25 nucleotides in length, and the second strand is sufficiently complementary to a target mRNA along at least 19 nucleotide of the second strand length to reduce target gene expression when the RNAi agent is introduced into a mammalian cell, and wherein Dicer cleavage of the dsRNAi agent results in an siRNA comprising the 3'-end of the second strand, thereby reducing expression of the target gene in the mammal. Optionally, the dsRNAi agent further comprises a ligand.

In certain embodiments, the sense strand of the dsRNAi agent contains at least one motif of three identical modifications on three consecutive nucleotides, where one of the motifs occurs at the cleavage site in the sense strand.

In certain embodiments, the antisense strand of the dsRNAi agent can also contain at least one motif of three identical modifications on three consecutive nucleotides, where one of the motifs occurs at or near the cleavage site in the antisense strand.

For a dsRNAi agent having a duplex region of 19-23 nucleotides in length, the cleavage site of the antisense strand is typically around the 10, 11, and 12 positions from the 5'-end. Thus the motifs of three identical modifications may occur at the 9, 10, 11 positions; the 10, 11, 12 positions; the 11, 12, 13 positions; the 12, 13, 14 positions; or the 13, 14, 15 positions of the antisense strand, the count starting from the first nucleotide from the 5'-end of the antisense strand, or, the count starting from the first paired nucleotide within the duplex region from the 5'-end of the antisense strand. The cleavage site in the antisense strand may also change according to the length of the duplex region of the dsRNAi agent from the 5'-end.

The sense strand of the dsRNAi agent may contain at least one motif of three identical modifications on three consecutive nucleotides at the cleavage site of the strand; and the antisense strand may have at least one motif of three identical modifications on three consecutive nucleotides at or near the cleavage site of the strand. When the sense strand and the antisense strand form a dsRNA duplex, the sense strand and the antisense strand can be so aligned that one motif of the three nucleotides on the sense strand and one motif of the three nucleotides on the antisense strand have at least one nucleotide overlap, *i.e.*, at least one of the three nucleotides of the motif in the sense strand forms a base pair with at least one of the three nucleotides of the motif in the antisense strand. Alternatively, at least two nucleotides may overlap, or all three nucleotides may overlap.

In some embodiments, the sense strand of the dsRNAi agent may contain more than one motif of three identical modifications on three consecutive nucleotides. The first motif may occur at or near the cleavage site of the strand and the other motifs may be a wing modification. The term "wing modification" herein refers to a motif occurring at another portion of the strand that is separated from the motif at or near the cleavage site of the same strand. The wing modification is either adjacent to the first motif or is separated by at least one or more nucleotides. When the motifs are immediately adjacent to each other then the chemistries of the motifs are distinct from each other, and when the motifs are separated by one or more nucleotide then the chemistries can be the same or different. Two or more wing modifications may be present. For instance, when two wing modifications are present,

each wing modification may occur at one end relative to the first motif which is at or near cleavage site or on either side of the lead motif.

Like the sense strand, the antisense strand of the dsRNAi agent may contain more than one motif of three identical modifications on three consecutive nucleotides, with at least one of the motifs occurring at or near the cleavage site of the strand. This antisense strand may also contain one or more wing modifications in an alignment similar to the wing modifications that may be present on the sense strand.

In some embodiments, the wing modification on the sense strand or antisense strand of the dsRNAi agent typically does not include the first one or two terminal nucleotides at the 3'-end, 5'-end, or both ends of the strand.

In other embodiments, the wing modification on the sense strand or antisense strand of the dsRNAi agent typically does not include the first one or two paired nucleotides within the duplex region at the 3'-end, 5'-end, or both ends of the strand.

When the sense strand and the antisense strand of the dsRNAi agent each contain at least one wing modification, the wing modifications may fall on the same end of the duplex region, and have an overlap of one, two, or three nucleotides.

When the sense strand and the antisense strand of the dsRNAi agent each contain at least two wing modifications, the sense strand and the antisense strand can be so aligned that two modifications each from one strand fall on one end of the duplex region, having an overlap of one, two, or three nucleotides; two modifications each from one strand fall on the other end of the duplex region, having an overlap of one, two or three nucleotides; two modifications one strand fall on each side of the lead motif, having an overlap of one, two or three nucleotides in the duplex region.

In some embodiments, every nucleotide in the sense strand and antisense strand of the dsRNAi agent, including the nucleotides that are part of the motifs, may be modified. Each nucleotide may be modified with the same or different modification which can include one or more alteration of one or both of the non-linking phosphate oxygens or of one or more of the linking phosphate oxygens; alteration of a constituent of the ribose sugar, *e.g.*, of the 2'-hydroxyl on the ribose sugar; wholesale replacement of the phosphate moiety with "dephospho" linkers; modification or replacement of a naturally occurring base; and replacement or modification of the ribose-phosphate backbone.

As nucleic acids are polymers of subunits, many of the modifications occur at a position which is repeated within a nucleic acid, *e.g.*, a modification of a base, or a phosphate moiety, or a non-linking O of a phosphate moiety. In some cases the modification will occur at all of the subject positions in the nucleic acid but in many cases it will not. By way of example, a modification may only occur at a 3'- or 5' terminal position, may only occur in a terminal region, *e.g.*, at a position on a terminal nucleotide or in the last 2, 3, 4, 5, or 10 nucleotides of a strand. A modification may occur in a double strand region, a single strand region, or in both. A modification may occur only in the double strand region of an RNA or may only occur in a single strand region of a RNA. For example, a phosphorothioate modification at a non-linking O position may only occur at one or both termini, may

only occur in a terminal region, *e.g.*, at a position on a terminal nucleotide or in the last 2, 3, 4, 5, or 10 nucleotides of a strand, or may occur in double strand and single strand regions, particularly at termini. The 5'-end or ends can be phosphorylated.

It may be possible, *e.g.*, to enhance stability, to include particular bases in overhangs, or to include modified nucleotides or nucleotide surrogates, in single strand overhangs, *e.g.*, in a 5'- or 3'-overhang, or in both. For example, it can be desirable to include purine nucleotides in overhangs. In some embodiments all or some of the bases in a 3'- or 5'-overhang may be modified, *e.g.*, with a modification described herein. Modifications can include, *e.g.*, the use of modifications at the 2' position of the ribose sugar with modifications that are known in the art, *e.g.*, the use of deoxyribonucleotides, 2'-deoxy-2'-fluoro (2'-F) or 2'-O-methyl modified instead of the ribosugar of the nucleobase, and modifications in the phosphate group, *e.g.*, phosphorothioate modifications. Overhangs need not be homologous with the target sequence.

In some embodiments, each residue of the sense strand and antisense strand is independently modified with LNA, CRN, cET, UNA, HNA, CeNA, 2'-methoxyethyl, 2'-O-methyl, 2'-O-allyl, 2'-C-allyl, 2'-deoxy, 2'-hydroxyl, or 2'-fluoro. The strands can contain more than one modification. In one embodiment, each residue of the sense strand and antisense strand is independently modified with 2'-O-methyl or 2'-fluoro.

At least two different modifications are typically present on the sense strand and antisense strand. Those two modifications may be the 2'-O-methyl or 2'-fluoro modifications, or others.

In certain embodiments, the N_a or N_b comprise modifications of an alternating pattern. The term "alternating motif" as used herein refers to a motif having one or more modifications, each modification occurring on alternating nucleotides of one strand. The alternating nucleotide may refer to one per every other nucleotide or one per every three nucleotides, or a similar pattern. For example, if A, B and C each represent one type of modification to the nucleotide, the alternating motif can be "ABABABABABAB...", "AABBAABBAABB...", "AABAABAABAAB...", "AAABAAABAAAB...", "AAABBBAAABBB...", or "ABCABCABCABC...", *etc.*

The type of modifications contained in the alternating motif may be the same or different. For example, if A, B, C, D each represent one type of modification on the nucleotide, the alternating pattern, *i.e.*, modifications on every other nucleotide, may be the same, but each of the sense strand or antisense strand can be selected from several possibilities of modifications within the alternating motif such as "ABABAB...", "ACACAC...", "BDBDBD..." or "CDCDCD...", *etc.*

In some embodiments, the dsRNAi agent of the invention comprises the modification pattern for the alternating motif on the sense strand relative to the modification pattern for the alternating motif on the antisense strand is shifted. The shift may be such that the modified group of nucleotides of the sense strand corresponds to a differently modified group of nucleotides of the antisense strand and *vice versa*. For example, the sense strand when paired with the antisense strand in the dsRNA duplex, the alternating motif in the sense strand may start with "ABABAB" from 5' to 3' of the strand and the alternating motif in the antisense strand may start with "BABABA" from 5' to 3' of the strand within the duplex region. As another example, the alternating motif in the sense strand may start with

“AABBAABB” from 5’ to 3’ of the strand and the alternating motif in the antisense strand may start with “BBAABBAA” from 5’ to 3’ of the strand within the duplex region, so that there is a complete or partial shift of the modification patterns between the sense strand and the antisense strand.

In some embodiments, the dsRNAi agent comprises the pattern of the alternating motif of 2'-O-methyl modification and 2'-F modification on the sense strand initially has a shift relative to the pattern of the alternating motif of 2'-O-methyl modification and 2'-F modification on the antisense strand initially, *i.e.*, the 2'-O-methyl modified nucleotide on the sense strand base pairs with a 2'-F modified nucleotide on the antisense strand and vice versa. The 1 position of the sense strand may start with the 2'-F modification, and the 1 position of the antisense strand may start with the 2'-O-methyl modification.

The introduction of one or more motifs of three identical modifications on three consecutive nucleotides to the sense strand or antisense strand interrupts the initial modification pattern present in the sense strand or antisense strand. This interruption of the modification pattern of the sense or antisense strand by introducing one or more motifs of three identical modifications on three consecutive nucleotides to the sense or antisense strand may enhance the gene silencing activity against the target gene.

In some embodiments, when the motif of three identical modifications on three consecutive nucleotides is introduced to any of the strands, the modification of the nucleotide next to the motif is a different modification than the modification of the motif. For example, the portion of the sequence containing the motif is “...N_aYYYN_b...,” where “Y” represents the modification of the motif of three identical modifications on three consecutive nucleotide, and “N_a” and “N_b” represent a modification to the nucleotide next to the motif “YYY” that is different than the modification of Y, and where N_a and N_b can be the same or different modifications. Alternatively, N_a or N_b may be present or absent when there is a wing modification present.

The iRNA may further comprise at least one phosphorothioate or methylphosphonate internucleotide linkage. The phosphorothioate or methylphosphonate internucleotide linkage modification may occur on any nucleotide of the sense strand, antisense strand, or both strands in any position of the strand. For instance, the internucleotide linkage modification may occur on every nucleotide on the sense strand or antisense strand; each internucleotide linkage modification may occur in an alternating pattern on the sense strand or antisense strand; or the sense strand or antisense strand may contain both internucleotide linkage modifications in an alternating pattern. The alternating pattern of the internucleotide linkage modification on the sense strand may be the same or different from the antisense strand, and the alternating pattern of the internucleotide linkage modification on the sense strand may have a shift relative to the alternating pattern of the internucleotide linkage modification on the antisense strand. In one embodiment, a double-stranded RNAi agent comprises 6-8 phosphorothioate internucleotide linkages. In some embodiments, the antisense strand comprises two phosphorothioate internucleotide linkages at the 5'-end and two phosphorothioate internucleotide linkages at the 3'-end, and the sense strand comprises at least two phosphorothioate internucleotide linkages at either the 5'-end or the 3'-end.

In some embodiments, the dsRNAi agent comprises a phosphorothioate or methylphosphonate internucleotide linkage modification in the overhang region. For example, the overhang region may contain two nucleotides having a phosphorothioate or methylphosphonate internucleotide linkage between the two nucleotides. Internucleotide linkage modifications also may be made to link the overhang nucleotides with the terminal paired nucleotides within the duplex region. For example, at least 2, 3, 4, or all the overhang nucleotides may be linked through phosphorothioate or methylphosphonate internucleotide linkage, and optionally, there may be additional phosphorothioate or methylphosphonate internucleotide linkages linking the overhang nucleotide with a paired nucleotide that is next to the overhang nucleotide. For instance, there may be at least two phosphorothioate internucleotide linkages between the terminal three nucleotides, in which two of the three nucleotides are overhang nucleotides, and the third is a paired nucleotide next to the overhang nucleotide. These terminal three nucleotides may be at the 3'-end of the antisense strand, the 3'-end of the sense strand, the 5'-end of the antisense strand, or the 5'-end of the antisense strand.

In some embodiments, the 2-nucleotide overhang is at the 3'-end of the antisense strand, and there are two phosphorothioate internucleotide linkages between the terminal three nucleotides, wherein two of the three nucleotides are the overhang nucleotides, and the third nucleotide is a paired nucleotide next to the overhang nucleotide. Optionally, the dsRNAi agent may additionally have two phosphorothioate internucleotide linkages between the terminal three nucleotides at both the 5'-end of the sense strand and at the 5'-end of the antisense strand.

In one embodiment, the dsRNAi agent comprises mismatch(es) with the target, within the duplex, or combinations thereof. The mismatch may occur in the overhang region or the duplex region. The base pair may be ranked on the basis of their propensity to promote dissociation or melting (*e.g.*, on the free energy of association or dissociation of a particular pairing, the simplest approach is to examine the pairs on an individual pair basis, though next neighbor or similar analysis can also be used). In terms of promoting dissociation: A:U is preferred over G:C; G:U is preferred over G:C; and I:C is preferred over G:C (I=inosine). Mismatches, *e.g.*, non-canonical or other than canonical pairings (as described elsewhere herein) are preferred over canonical (A:T, A:U, G:C) pairings; and pairings which include a universal base are preferred over canonical pairings.

In certain embodiments, the dsRNAi agent comprises at least one of the first 1, 2, 3, 4, or 5 base pairs within the duplex regions from the 5'-end of the antisense strand independently selected from the group of: A:U, G:U, I:C, and mismatched pairs, *e.g.*, non-canonical or other than canonical pairings or pairings which include a universal base, to promote the dissociation of the antisense strand at the 5'-end of the duplex.

In certain embodiments, the nucleotide at the 1 position within the duplex region from the 5'-end in the antisense strand is selected from A, dA, dU, U, and dT. Alternatively, at least one of the first 1, 2, or 3 base pair within the duplex region from the 5'-end of the antisense strand is an AU base pair. For example, the first base pair within the duplex region from the 5'-end of the antisense strand is an AU base pair.

In other embodiments, the nucleotide at the 3'-end of the sense strand is deoxythymidine (dT) or the nucleotide at the 3'-end of the antisense strand is deoxythymidine (dT). For example, there is a short sequence of deoxythymidine nucleotides, for example, two dT nucleotides on the 3'-end of the sense, antisense strand, or both strands.

5 In certain embodiments, the sense strand sequence may be represented by formula (I):



wherein:

i and j are each independently 0 or 1;

p and q are each independently 0-6;

10 each N_a independently represents an oligonucleotide sequence comprising 0-25 modified nucleotides, each sequence comprising at least two differently modified nucleotides;

each N_b independently represents an oligonucleotide sequence comprising 0-10 modified nucleotides;

each n_p and n_q independently represent an overhang nucleotide;

15 wherein N_b and Y do not have the same modification; and


XXX, YYY, and ZZZ each independently represent one motif of three identical modifications on three consecutive nucleotides. In one embodiment, YYY is all 2'-F modified nucleotides.

In some embodiments, the N_a or N_b comprises modifications of alternating pattern.

In some embodiments, the YYY motif occurs at or near the cleavage site of the sense strand.

20 For example, when the dsRNAi agent has a duplex region of 17-23 nucleotides in length, the YYY motif can occur at or the vicinity of the cleavage site (*e.g.*: can occur at positions 6, 7, 8; 7, 8, 9; 8, 9, 10; 9, 10, 11; 10, 11, 12; or 11, 12, 13) of the sense strand, the count starting from the first nucleotide, from the 5'-end; or optionally, the count starting at the first paired nucleotide within the duplex region, from the 5'-end.

25 In one embodiment, i is 1 and j is 0, or i is 0 and j is 1, or both i and j are 1. The sense strand can therefore be represented by the following formulas:



30 When the sense strand is represented by formula (Ib), N_b represents an oligonucleotide sequence comprising 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a independently can represent an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the sense strand is represented as formula (Ic), N_b represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a can independently represent an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

35 When the sense strand is represented as formula (Id), each N_b independently represents an oligonucleotide sequence comprising 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. In one

embodiment, N_b is 0, 1, 2, 3, 4, 5, or 6. Each N_a can independently represent an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

Each of X, Y and Z may be the same or different from each other.

In other embodiments, i is 0 and j is 0, and the sense strand may be represented by the

5 formula:

$5' n_p-N_a-YYY-N_a-n_q 3'$ (Ia).

When the sense strand is represented by formula (Ia), each N_a independently can represent an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

In one embodiment, the antisense strand sequence of the RNAi may be represented by

10 formula (II):

$5' n_q-N_a'-(Z'Z'Z')_k-N_b'-Y'Y'Y'-N_b'-(X'X'X')_l-N_a'-n_p' 3'$ (II)

wherein:

k and l are each independently 0 or 1;

p' and q' are each independently 0-6;

15 each N_a' independently represents an oligonucleotide sequence comprising 0-25 modified nucleotides, each sequence comprising at least two differently modified nucleotides;

each N_b' independently represents an oligonucleotide sequence comprising 0-10 modified nucleotides;

each n_p' and n_q' independently represent an overhang nucleotide;

20 wherein N_b' and Y' do not have the same modification; and

$X'X'X'$, $Y'Y'Y'$, and $Z'Z'Z'$ each independently represent one motif of three identical modifications on three consecutive nucleotides.

In some embodiments, the N_a' or N_b' comprises modifications of alternating pattern.

The $Y'Y'Y'$ motif occurs at or near the cleavage site of the antisense strand. For example, 25 when the dsRNAi agent has a duplex region of 17-23 nucleotides in length, the $Y'Y'Y'$ motif can occur at positions 9, 10, 11; 10, 11, 12; 11, 12, 13; 12, 13, 14; or 13, 14, 15 of the antisense strand, with the count starting from the first nucleotide, from the 5'-end; or optionally, the count starting at the first paired nucleotide within the duplex region, from the 5'-end. In one embodiment, the $Y'Y'Y'$ motif occurs at positions 11, 12, 13.

30 In certain embodiments, $Y'Y'Y'$ motif is all 2'-OMe modified nucleotides.

In certain embodiments, k is 1 and l is 0, or k is 0 and l is 1, or both k and l are 1.

The antisense strand can therefore be represented by the following formulas:

$5' n_q-N_a'-Z'Z'Z'-N_b'-Y'Y'Y'-N_a'-n_p' 3'$ (IIb);

$5' n_q-N_a'-Y'Y'Y'-N_b'-X'X'X'-n_p' 3'$ (IIc); or


35 $5' n_q-N_a'-Z'Z'Z'-N_b'-Y'Y'Y'-N_b'-X'X'X'-N_a'-n_p' 3'$ (IId).

When the antisense strand is represented by formula (IIb), N_b' represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the antisense strand is represented as formula (IIc), N_b' represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

- 5 When the antisense strand is represented as formula (IIId), each N_b' independently represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides. In one embodiment, N_b is 0, 1, 2, 3, 4, 5, or 6.

- 10 In other embodiments, k is 0 and l is 0 and the antisense strand may be represented by the formula:



When the antisense strand is represented as formula (IIa), each N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

Each of X' , Y' and Z' may be the same or different from each other.


- 15 Each nucleotide of the sense strand and antisense strand may be independently modified with LNA, CRN, UNA, cEt, HNA, CeNA, 2'-methoxyethyl, 2'-O-methyl, 2'-O-allyl, 2'-C-allyl, 2'-hydroxyl, or 2'-fluoro. For example, each nucleotide of the sense strand and antisense strand is independently modified with 2'-O-methyl or 2'-fluoro. Each X , Y , Z , X' , Y' , and Z' , in particular, may represent a 2'-O-methyl modification or a 2'-fluoro modification.


- 20 In some embodiments, the sense strand of the dsRNAi agent may contain YYY motif occurring at 9, 10, and 11 positions of the strand when the duplex region is 21 nt, the count starting from the first nucleotide from the 5'-end, or optionally, the count starting at the first paired nucleotide within the duplex region, from the 5'-end; and Y represents 2'-F modification. The sense strand may additionally contain XXX motif or ZZZ motifs as wing modifications at the opposite end of the duplex region; and XXX and ZZZ each independently represents a 2'-OMe modification or 2'-F modification.

- 25 In some embodiments the antisense strand may contain Y'Y'Y' motif occurring at positions 11, 12, 13 of the strand, the count starting from the first nucleotide from the 5'-end, or optionally, the count starting at the first paired nucleotide within the duplex region, from the 5'-end; and Y' represents 2'-O-methyl modification. The antisense strand may additionally contain X'X'X' motif or Z'Z'Z' motifs as wing modifications at the opposite end of the duplex region; and X'X'X' and Z'Z'Z' each independently represents a 2'-OMe modification or 2'-F modification.

- 30 The sense strand represented by any one of the above formulas (Ia), (Ib), (Ic), and (Id) forms a duplex with an antisense strand being represented by any one of formulas (IIa), (IIb), (IIc), and (IIId), respectively.

Accordingly, the dsRNAi agents for use in the methods of the invention may comprise a sense strand and an antisense strand, each strand having 14 to 30 nucleotides, the iRNA duplex represented by formula (III):





wherein:

i, j, k, and l are each independently 0 or 1;

5 p, p', q, and q' are each independently 0-6;

each N_a and N_a' independently represents an oligonucleotide sequence comprising 0-25 modified nucleotides, each sequence comprising at least two differently modified nucleotides;





each N_b and N_b' independently represents an oligonucleotide sequence comprising 0-10 modified nucleotides;

10 wherein each n_p' , n_p , n_q' , and n_q , each of which may or may not be present, independently represents an overhang nucleotide; and

XXX, YYY, ZZZ, $X'X'X'$, $Y'Y'Y'$, and $Z'Z'Z'$ each independently represent one motif of three identical modifications on three consecutive nucleotides.

In one embodiment, i is 0 and j is 0; or i is 1 and j is 0; or i is 0 and j is 1; or both i and j are 0; or both i and j are 1. In another embodiment, k is 0 and l is 0; or k is 1 and l is 0; k is 0 and l is 1; or both k and l are 0; or both k and l are 1.

Exemplary combinations of the sense strand and antisense strand forming an iRNA duplex include the formulas below:



When the dsRNAi agent is represented by formula (IIIa), each N_a independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the dsRNAi agent is represented by formula (IIIb), each N_b independently represents an oligonucleotide sequence comprising 1-10, 1-7, 1-5, or 1-4 modified nucleotides. Each N_a independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the dsRNAi agent is represented as formula (IIIc), each N_b , N_b' independently represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified

nucleotides. Each N_a independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the dsRNAi agent is represented as formula (III_d), each N_b , N_b' independently represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a , N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides. Each of N_a , N_a' , N_b , and N_b' independently comprises modifications of alternating pattern.

Each of X, Y, and Z in formulas (III), (III_a), (III_b), (III_c), and (III_d) may be the same or different from each other.

When the dsRNAi agent is represented by formula (III), (III_a), (III_b), (III_c), and (III_d), at least one of the Y nucleotides may form a base pair with one of the Y' nucleotides. Alternatively, at least two of the Y nucleotides form base pairs with the corresponding Y' nucleotides; or all three of the Y nucleotides all form base pairs with the corresponding Y' nucleotides.

When the dsRNAi agent is represented by formula (III_b) or (III_d), at least one of the Z nucleotides may form a base pair with one of the Z' nucleotides. Alternatively, at least two of the Z nucleotides form base pairs with the corresponding Z' nucleotides; or all three of the Z nucleotides all form base pairs with the corresponding Z' nucleotides.

When the dsRNAi agent is represented as formula (III_c) or (III_d), at least one of the X nucleotides may form a base pair with one of the X' nucleotides. Alternatively, at least two of the X nucleotides form base pairs with the corresponding X' nucleotides; or all three of the X nucleotides all form base pairs with the corresponding X' nucleotides.

In certain embodiments, the modification on the Y nucleotide is different than the modification on the Y' nucleotide, the modification on the Z nucleotide is different than the modification on the Z' nucleotide, or the modification on the X nucleotide is different than the modification on the X' nucleotide.

In certain embodiments, when the dsRNAi agent is represented by formula (III_d), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications. In other embodiments, when the RNAi agent is represented by formula (III_d), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications and $n_p' > 0$ and at least one n_p' is linked to a neighboring nucleotide *via* phosphorothioate linkage. In yet other embodiments, when the RNAi agent is represented by formula (III_d), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications, $n_p' > 0$ and at least one n_p' is linked to a neighboring nucleotide *via* phosphorothioate linkage, and the sense strand is conjugated to one or more GalNAc derivatives attached through a bivalent or trivalent branched linker (described below). In other embodiments, when the RNAi agent is represented by formula (III_d), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications, $n_p' > 0$ and at least one n_p' is linked to a neighboring nucleotide *via* phosphorothioate linkage, the sense strand comprises at least one phosphorothioate linkage, and the sense strand is conjugated to one or more GalNAc derivatives attached through a bivalent or trivalent branched linker.

In some embodiments, when the dsRNAi agent is represented by formula (IIIa), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications, $n_p' > 0$ and at least one n_p' is linked to a neighboring nucleotide *via* phosphorothioate linkage, the sense strand comprises at least one phosphorothioate linkage, and the sense strand is conjugated to one or more GalNAc derivatives
5 attached through a bivalent or trivalent branched linker.

In some embodiments, the dsRNAi agent is a multimer containing at least two duplexes represented by formula (III), (IIIa), (IIIb), (IIIc), and (IIId), wherein the duplexes are connected by a linker. The linker can be cleavable or non-cleavable. Optionally, the multimer further comprises a ligand. Each of the duplexes can target the same gene or two different genes; or each of the duplexes
10 can target same gene at two different target sites.

In some embodiments, the dsRNAi agent is a multimer containing three, four, five, six, or more duplexes represented by formula (III), (IIIa), (IIIb), (IIIc), and (IIId), wherein the duplexes are connected by a linker. The linker can be cleavable or non-cleavable. Optionally, the multimer further comprises a ligand. Each of the duplexes can target the same gene or two different genes; or each of
15 the duplexes can target same gene at two different target sites.


In one embodiment, two dsRNAi agents represented by at least one of formulas (III), (IIIa), (IIIb), (IIIc), and (IIId) are linked to each other at the 5' end, and one or both of the 3' ends, and are optionally conjugated to a ligand. Each of the agents can target the same gene or two different genes; or each of the agents can target same gene at two different target sites.

In certain embodiments, an RNAi agent of the invention may contain a low number of nucleotides containing a 2'-fluoro modification, *e.g.*, 10 or fewer nucleotides with 2'-fluoro modification. For example, the RNAi agent may contain 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 or 0 nucleotides with a 2'-fluoro modification. In a specific embodiment, the RNAi agent of the invention contains 10
20 nucleotides with a 2'-fluoro modification, *e.g.*, 4 nucleotides with a 2'-fluoro modification in the sense strand and 6 nucleotides with a 2'-fluoro modification in the antisense strand. In another specific embodiment, the RNAi agent of the invention contains 6 nucleotides with a 2'-fluoro modification, *e.g.*, 4 nucleotides with a 2'-fluoro modification in the sense strand and 2 nucleotides with a 2'-fluoro modification in the antisense strand.

In other embodiments, an RNAi agent of the invention may contain an ultra low number of nucleotides containing a 2'-fluoro modification, *e.g.*, 2 or fewer nucleotides containing a 2'-fluoro modification. For example, the RNAi agent may contain 2, 1 or 0 nucleotides with a 2'-fluoro modification. In a specific embodiment, the RNAi agent may contain 2 nucleotides with a 2'-fluoro modification, *e.g.*, 0 nucleotides with a 2-fluoro modification in the sense strand and 2 nucleotides with a 2'-fluoro modification in the antisense strand.

Various publications describe multimeric iRNAs that can be used in the methods of the invention. Such publications include WO2007/091269, U.S. Patent No. 7,858,769, WO2010/141511, WO2007/117686, WO2009/014887, and WO2011/031520 the entire contents of each of which are hereby incorporated herein by reference.

In certain embodiments, the compositions and methods of the disclosure include a vinyl phosphonate (VP) modification of an RNAi agent as described herein. In exemplary embodiments, a 5' vinyl phosphonate modified nucleotide of the disclosure has the structure:



5 wherein X is O or S;

R is hydrogen, hydroxy, fluoro, or C₁₋₂₀alkoxy (*e.g.*, methoxy or n-hexadecyloxy);

R^{5'} is =C(H)-P(O)(OH)₂ and the double bond between the C5' carbon and R^{5'} is in the *E* or *Z* orientation (*e.g.*, *E* orientation); and

B is a nucleobase or a modified nucleobase, optionally where B is adenine, guanine, cytosine,
10 thymine, or uracil.

A vinyl phosphonate of the instant disclosure may be attached to either the antisense or the sense strand of a dsRNA of the disclosure. In certain embodiments, a vinyl phosphonate of the instant disclosure is attached to the antisense strand of a dsRNA, optionally at the 5' end of the antisense strand of the dsRNA.

15 Vinyl phosphonate modifications are also contemplated for the compositions and methods of the instant disclosure. An exemplary vinyl phosphonate structure includes the preceding structure, where R^{5'} is =C(H)-OP(O)(OH)₂ and the double bond between the C5' carbon and R^{5'} is in the *E* or *Z* orientation (*e.g.*, *E* orientation).

As described in more detail below, the iRNA that contains conjugations of one or more
20 carbohydrate moieties to an iRNA can optimize one or more properties of the iRNA. In many cases, the carbohydrate moiety will be attached to a modified subunit of the iRNA. For example, the ribose sugar of one or more ribonucleotide subunits of a iRNA can be replaced with another moiety, *e.g.*, a non-carbohydrate (such as, cyclic) carrier to which is attached a carbohydrate ligand. A ribonucleotide subunit in which the ribose sugar of the subunit has been so replaced is referred to
25 herein as a ribose replacement modification subunit (RRMS). A cyclic carrier may be a carbocyclic ring system, *i.e.*, all ring atoms are carbon atoms, or a heterocyclic ring system, *i.e.*, one or more ring atoms may be a heteroatom, *e.g.*, nitrogen, oxygen, sulfur. The cyclic carrier may be a monocyclic ring system, or may contain two or more rings, *e.g.* fused rings. The cyclic carrier may be a fully saturated ring system, or it may contain one or more double bonds.

30 The ligand may be attached to the polynucleotide *via* a carrier. The carriers include (i) at least one “backbone attachment point,” such as, two “backbone attachment points” and (ii) at least one “tethering attachment point.” A “backbone attachment point” as used herein refers to a functional group, *e.g.* a hydroxyl group, or generally, a bond available for, and that is suitable for incorporation

of the carrier into the backbone, *e.g.*, the phosphate, or modified phosphate, *e.g.*, sulfur containing, backbone, of a ribonucleic acid. A “tethering attachment point” (TAP) in some embodiments refers to a constituent ring atom of the cyclic carrier, *e.g.*, a carbon atom or a heteroatom (distinct from an atom which provides a backbone attachment point), that connects a selected moiety. The moiety can be,

- 5 *e.g.*, a carbohydrate, *e.g.* monosaccharide, disaccharide, trisaccharide, tetrasaccharide, oligosaccharide, or polysaccharide. Optionally, the selected moiety is connected by an intervening tether to the cyclic carrier. Thus, the cyclic carrier will often include a functional group, *e.g.*, an amino group, or generally, provide a bond, that is suitable for incorporation or tethering of another chemical entity, *e.g.*, a ligand to the constituent ring.
- 10 The iRNA may be conjugated to a ligand *via* a carrier, wherein the carrier can be cyclic group or acyclic group. In one embodiment, the cyclic group is selected from pyrrolidinyl, pyrazolinyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, piperidinyl, piperazinyl, [1,3]dioxolane, oxazolidinyl, isoxazolidinyl, morpholinyl, thiazolidinyl, isothiazolidinyl, quinoxalinyl, pyridazinonyl, tetrahydrofuryl, and decalin. In one embodiment, the acyclic group is a serinol backbone or
- 15 diethanolamine backbone.

i. Thermally Destabilizing Modifications


- In certain embodiments, a dsRNA molecule can be optimized for RNA interference by incorporating thermally destabilizing modifications in the seed region of the antisense strand. As used
- 20 herein “seed region” means at positions 2-9 of the 5’-end of the referenced strand or at positions 2-8 of the 5’-end of the referenced strand. For example, thermally destabilizing modifications can be incorporated in the seed region of the antisense strand to reduce or inhibit off-target gene silencing.

- The term “thermally destabilizing modification(s)” includes modification(s) that would result with a dsRNA with a lower overall melting temperature (T_m) than the T_m of the dsRNA without
- 25 having such modification(s). For example, the thermally destabilizing modification(s) can decrease the T_m of the dsRNA by 1 – 4 °C, such as one, two, three or four degrees Celcius. And, the term “thermally destabilizing nucleotide” refers to a nucleotide containing one or more thermally destabilizing modifications.

- It has been discovered that dsRNAs with an antisense strand comprising at least one thermally
- 30 destabilizing modification of the duplex within the first 9 nucleotide positions, counting from the 5’ end, of the antisense strand have reduced off-target gene silencing activity. Accordingly, in some embodiments, the antisense strand comprises at least one (*e.g.*, one, two, three, four, five or more) thermally destabilizing modification of the duplex within the first 9 nucleotide positions of the 5’ region of the antisense strand. In some embodiments, one or more thermally destabilizing
- 35 modification(s) of the duplex is/are located in positions 2-9, such as, positions 4-8, from the 5’-end of the antisense strand. In some further embodiments, the thermally destabilizing modification(s) of the duplex is/are located at position 6, 7 or 8 from the 5’-end of the antisense strand. In still some further embodiments, the thermally destabilizing modification of the duplex is located at position 7 from the


5'-end of the antisense strand. In some embodiments, the thermally destabilizing modification of the duplex is located at position 2, 3, 4, 5 or 9 from the 5'-end of the antisense strand.

An iRNA agent comprises a sense strand and an antisense strand, each strand having 14 to 40 nucleotides. The RNAi agent may be represented by formula (L):




In formula (L), B1, B2, B3, B1', B2', B3', and B4' each are independently a nucleotide containing a modification selected from the group consisting of 2'-O-alkyl, 2'-substituted alkoxy, 2'-substituted alkyl, 2'-halo, ENA, and BNA/LNA. In one embodiment, B1, B2, B3, B1', B2', B3', and B4' each contain 2'-OMe modifications. In one embodiment, B1, B2, B3, B1', B2', B3', and B4' each contain 2'-OMe or 2'-F modifications. In one embodiment, at least one of B1, B2, B3, B1', B2', B3', and B4' contain 2'-O-N-methylacetamido (2'-O-NMA, 2'-O-CH₂C(O)N(Me)H) modification.


C1 is a thermally destabilizing nucleotide placed at a site opposite to the seed region of the antisense strand (*i.e.*, at positions 2-8 of the 5'-end of the antisense strand, or at positions 2-9 of the 5'-end of the antisense strand). For example, C1 is at a position of the sense strand that pairs with a nucleotide at positions 2-8 of the 5'-end of the antisense strand. In one example, C1 is at position 15 from the 5'-end of the sense strand. C1 nucleotide bears the thermally destabilizing modification which can include abasic modification; mismatch with the opposing nucleotide in the duplex; and sugar modification such as 2'-deoxy modification or acyclic nucleotide *e.g.*, unlocked nucleic acids (UNA) or glycerol nucleic acid (GNA), or 2'-5'-linked ribonucleotides ("3'-RNA").. In one embodiment, C1 has thermally destabilizing modification selected from the group consisting of: i) mismatch with the opposing nucleotide in the antisense strand; ii) abasic modification selected from the group consisting of:



selected from the group consisting of:



, wherein B is a modified or unmodified nucleobase, R^1 and R^2 independently are H, halogen, OR_3 , or alkyl; and R_3 is H, alkyl, cycloalkyl, aryl, aralkyl, heteroaryl or sugar. In one embodiment, the thermally destabilizing modification in C1 is a mismatch selected from the group consisting of G:G, G:A, G:U, G:T, A:A, A:C, C:C, C:U, C:T, U:U, T:T, and U:T; and optionally, at least one nucleobase in the mismatch pair is a 2'-deoxy nucleobase. In one example, the thermally



destabilizing modification in C1 is GNA or T1, T1', T2', and T3' each independently represent a nucleotide comprising a modification providing the nucleotide a steric bulk that is less or equal to the steric bulk of a 2'-OMe modification. A steric bulk refers to the sum of steric effects of a modification. Methods for determining steric effects of a modification of a nucleotide are known to one skilled in the art. The modification can be at the 2' position of a ribose sugar of the nucleotide, or a modification to a non-ribose nucleotide, acyclic nucleotide, or the backbone of the nucleotide that is similar or equivalent to the 2' position of the ribose sugar, and provides the nucleotide a steric bulk that is less than or equal to the steric bulk of a 2'-OMe modification. For example, T1, T1', T2', and T3' are each independently selected from DNA, RNA, LNA, 2'-F, and 2'-F-5'-methyl. In one embodiment, T1 is DNA. In one embodiment, T1' is DNA, RNA or LNA. In one embodiment, T2' is DNA or RNA. In one embodiment, T3' is DNA or RNA.

n^1 , n^3 , and q^1 are independently 4 to 15 nucleotides in length.
 n^5 , q^3 , and q^7 are independently 1-6 nucleotide(s) in length.
 n^4 , q^2 , and q^6 are independently 1-3 nucleotide(s) in length; alternatively, n^4 is 0.
 q^5 is independently 0-10 nucleotide(s) in length.
 n^2 and q^4 are independently 0-3 nucleotide(s) in length.

Alternatively, n^4 is 0-3 nucleotide(s) in length.

In one embodiment, n^4 can be 0. In one example, n^4 is 0, and q^2 and q^6 are 1. In another example, n^4 is 0, and q^2 and q^6 are 1, with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two

phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, n^4 , q^2 , and q^6 are each 1.

5 In one embodiment, n^2 , n^4 , q^2 , q^4 , and q^6 are each 1.

In one embodiment, C1 is at position 14-17 of the 5'-end of the sense strand, when the sense strand is 19-22 nucleotides in length, and n^4 is 1. In one embodiment, C1 is at position 15 of the 5'-end of the sense strand

10 In one embodiment, T3' starts at position 2 from the 5' end of the antisense strand. In one example, T3' is at position 2 from the 5' end of the antisense strand and q^6 is equal to 1.

In one embodiment, T1' starts at position 14 from the 5' end of the antisense strand. In one example, T1' is at position 14 from the 5' end of the antisense strand and q^2 is equal to 1.

15 In an exemplary embodiment, T3' starts from position 2 from the 5' end of the antisense strand and T1' starts from position 14 from the 5' end of the antisense strand. In one example, T3' starts from position 2 from the 5' end of the antisense strand and q^6 is equal to 1 and T1' starts from position 14 from the 5' end of the antisense strand and q^2 is equal to 1.

In one embodiment, T1' and T3' are separated by 11 nucleotides in length (*i.e.* not counting the T1' and T3' nucleotides).

20 In one embodiment, T1' is at position 14 from the 5' end of the antisense strand. In one example, T1' is at position 14 from the 5' end of the antisense strand and q^2 is equal to 1, and the modification at the 2' position or positions in a non-ribose, acyclic or backbone that provide less steric bulk than a 2'-OMe ribose.

25 In one embodiment, T3' is at position 2 from the 5' end of the antisense strand. In one example, T3' is at position 2 from the 5' end of the antisense strand and q^6 is equal to 1, and the modification at the 2' position or positions in a non-ribose, acyclic or backbone that provide less than or equal to steric bulk than a 2'-OMe ribose.

30 In one embodiment, T1 is at the cleavage site of the sense strand. In one example, T1 is at position 11 from the 5' end of the sense strand, when the sense strand is 19-22 nucleotides in length, and n^2 is 1. In an exemplary embodiment, T1 is at the cleavage site of the sense strand at position 11 from the 5' end of the sense strand, when the sense strand is 19-22 nucleotides in length, and n^2 is 1,

In one embodiment, T2' starts at position 6 from the 5' end of the antisense strand. In one example, T2' is at positions 6-10 from the 5' end of the antisense strand, and q^4 is 1.

35 In an exemplary embodiment, T1 is at the cleavage site of the sense strand, for instance, at position 11 from the 5' end of the sense strand, when the sense strand is 19-22 nucleotides in length, and n^2 is 1; T1' is at position 14 from the 5' end of the antisense strand, and q^2 is equal to 1, and the modification to T1' is at the 2' position of a ribose sugar or at positions in a non-ribose, acyclic or backbone that provide less steric bulk than a 2'-OMe ribose; T2' is at positions 6-10 from the 5' end of the antisense strand, and q^4 is 1; and T3' is at position 2 from the 5' end of the antisense strand, and

q^6 is equal to 1, and the modification to T3' is at the 2' position or at positions in a non-ribose, acyclic or backbone that provide less than or equal to steric bulk than a 2'-OMe ribose.

In one embodiment, T2' starts at position 8 from the 5' end of the antisense strand. In one example, T2' starts at position 8 from the 5' end of the antisense strand, and q^4 is 2.

5 In one embodiment, T2' starts at position 9 from the 5' end of the antisense strand. In one example, T2' is at position 9 from the 5' end of the antisense strand, and q^4 is 1.

In one embodiment, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 1, B3' is 2'-OMe or 2'-F, q^5 is 6, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense
10 strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 1, B3' is 2'-OMe or 2'-F, q^5 is 6, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two
15 phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense
25 strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 6, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 7, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1.
35

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 6, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 7, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1.

1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 1, B3' is 2'-OMe or 2'-F, q^5 is 6, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 1, B3' is 2'-OMe or 2'-F, q^5 is 6, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 5, T2' is 2'-F, q^4 is 1, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; optionally with at least 2 additional TT at the 3'-end of the antisense strand.


In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 5, T2' is 2'-F, q^4 is 1, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; optionally with at least 2 additional TT at the 3'-end of the antisense strand; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1.


In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end).

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).


In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

The RNAi agent can comprise a phosphorus-containing group at the 5'-end of the sense strand or antisense strand. The 5'-end phosphorus-containing group can be 5'-end phosphate (5'-P), 5'-end phosphorothioate (5'-PS), 5'-end phosphorodithioate (5'-PS₂), 5'-end vinylphosphonate (5'-VP), 5'-end methylphosphonate (MePhos), or 5'-deoxy-5'-C-malonyl (). When the 5'-end phosphorus-containing group is 5'-end vinylphosphonate (5'-VP), the 5'-VP can be either

5'-E-VP isomer (*i.e.*, *trans*-vinylphosphonate,



5'-Z-VP isomer (*i.e.*, *cis*-



vinylphosphonate, or mixtures thereof.

In one embodiment, the RNAi agent comprises a phosphorus-containing group at the 5'-end of the sense strand. In one embodiment, the RNAi agent comprises a phosphorus-containing group at the 5'-end of the antisense strand.

In one embodiment, the RNAi agent comprises a 5'-P. In one embodiment, the RNAi agent comprises a 5'-P in the antisense strand.

In one embodiment, the RNAi agent comprises a 5'-PS. In one embodiment, the RNAi agent comprises a 5'-PS in the antisense strand.

5 In one embodiment, the RNAi agent comprises a 5'-VP. In one embodiment, the RNAi agent comprises a 5'-VP in the antisense strand. In one embodiment, the RNAi agent comprises a 5'-E-VP in the antisense strand. In one embodiment, the RNAi agent comprises a 5'-Z-VP in the antisense strand.

10 In one embodiment, the RNAi agent comprises a 5'-PS₂. In one embodiment, the RNAi agent comprises a 5'-PS₂ in the antisense strand.

In one embodiment, the RNAi agent comprises a 5'-PS₂. In one embodiment, the RNAi agent comprises a 5'-deoxy-5'-C-malonyl in the antisense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, 15 T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-P. 20

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination 25 thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-PS₂.

30 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage 35

modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1. The RNAi agent also comprises a 5'-P.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1. The dsRNA agent also comprises a 5'-PS.

- 5 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1. The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.
- In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1. The RNAi agent also comprises a 5'-PS₂.

- 15 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1. The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

- In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-P.

- 25 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-PS.

- 30 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand
 5 (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F,
 10 q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-deoxy-5'-
 15 C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The RNAi agent also comprises a 5'-P.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The RNAi agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The dsRNAi RNA agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1;

with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The RNAi agent also comprises a 5' - P.

5 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The RNAi agent also comprises a 5' - PS.

10 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The RNAi agent also comprises a 5' - VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

15 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The RNAi agent also comprises a 5' - PS₂.

20 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

25 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5' - P.

30 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5' - PS.

35 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two

phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P and a targeting ligand. In one embodiment, the 5'-P is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS and a targeting ligand. In one embodiment, the 5'-PS is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP (*e.g.*, a 5'-E-VP, 5'-Z-VP, or combination thereof), and a targeting ligand.

In one embodiment, the 5'-VP is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂ and a targeting ligand. In one embodiment, the 5'-PS₂ is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl and a targeting ligand. In one embodiment, the 5'-deoxy-5'-C-malonyl is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-P and a targeting ligand. In one embodiment, the 5'-P is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand

5 (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-PS and a targeting ligand. In one embodiment, the 5'-PS is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

10 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at
15 positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-VP (*e.g.*, a 5'-E-VP, 5'-Z-VP, or combination thereof) and a targeting ligand. In one embodiment, the 5'-VP is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at
20 positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-PS₂ and a targeting ligand. In one embodiment, the 5'-PS₂ is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at
30 positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl and a targeting ligand. In one embodiment, the 5'-deoxy-5'-C-malonyl is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1;

with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P and a targeting ligand. In one embodiment, the 5'-P is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1;

with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS and a targeting ligand. In one embodiment, the 5'-PS is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP (e.g., a 5'-E-VP, 5'-Z-VP, or combination thereof) and a targeting ligand. In one embodiment, the 5'-VP is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂ and a targeting ligand. In one embodiment, the 5'-PS₂ is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide

linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl and a targeting ligand. In one embodiment, the 5'-deoxy-5'-C-malonyl is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P and a targeting ligand. In one embodiment, the 5'-P is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS and a targeting ligand. In one embodiment, the 5'-PS is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP (*e.g.*, a 5'-E-VP, 5'-Z-VP, or combination thereof) and a targeting ligand. In one embodiment, the 5'-VP is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications

within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂ and a targeting ligand. In one embodiment, the 5'-PS₂ is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl and a targeting ligand. In one embodiment, the 5'-deoxy-5'-C-malonyl is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In a particular embodiment, an RNAi agent of the present invention comprises:

- 15 (a) a sense strand having:
 - (i) a length of 21 nucleotides;
 - (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker; and
 - (iii) 2'-F modifications at positions 1, 3, 5, 7, 9 to 11, 13, 17, 19, and 21, and 2'-OMe modifications at positions 2, 4, 6, 8, 12, 14 to 16, 18, and 20 (counting from the 5' end);
- 20 and
- (b) an antisense strand having:
 - (i) a length of 23 nucleotides;
 - (ii) 2'-OMe modifications at positions 1, 3, 5, 9, 11 to 13, 15, 17, 19, 21, and 23, and 2'-F modifications at positions 2, 4, 6 to 8, 10, 14, 16, 18, 20, and 22 (counting from the 5' end); and
 - (iii) phosphorothioate internucleotide linkages between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);
- 30 wherein the dsRNA agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, an RNAi agent of the present invention comprises:

- (a) a sense strand having:
 - (i) a length of 21 nucleotides;
 - (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
 - (iii) 2'-F modifications at positions 1, 3, 5, 7, 9 to 11, 13, 15, 17, 19, and 21, and 2'-OMe modifications at positions 2, 4, 6, 8, 12, 14, 16, 18, and 20 (counting from the 5' end);
- 35 and

(iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);

and

(b) an antisense strand having:

- 5 (i) a length of 23 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 3, 5, 7, 9, 11 to 13, 15, 17, 19, and 21 to 23, and 2'-F modifications at positions 2, 4, 6, 8, 10, 14, 16, 18, and 20 (counting from the 5' end);
- and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between
- 10 nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);

wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- 15 (a) a sense strand having:
- (i) a length of 21 nucleotides;
- (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
- (iii) 2'-OMe modifications at positions 1 to 6, 8, 10, and 12 to 21, 2'-F modifications at
- 20 positions 7, and 9, and a deoxy-nucleotide (*e.g.* dT) at position 11 (counting from the 5' end); and
- (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- 25 (b) an antisense strand having:
- (i) a length of 23 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 3, 7, 9, 11, 13, 15, 17, and 19 to 23, and 2'-F modifications at positions 2, 4 to 6, 8, 10, 12, 14, 16, and 18 (counting from the 5' end);
- and
- 30 (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);

wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

35 In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
- (i) a length of 21 nucleotides;
- (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;

- (iii) 2'-OMe modifications at positions 1 to 6, 8, 10, 12, 14, and 16 to 21, and 2'-F modifications at positions 7, 9, 11, 13, and 15; and
- (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);

5 and

- (b) an antisense strand having:

- (i) a length of 23 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 5, 7, 9, 11, 13, 15, 17, 19, and 21 to 23, and 2'-F modifications at positions 2 to 4, 6, 8, 10, 12, 14, 16, 18, and 20 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);

10 wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a
15 blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
 - (i) a length of 21 nucleotides;
 - (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three
 - 20 GalNAc derivatives attached through a trivalent branched linker;
 - (iii) 2'-OMe modifications at positions 1 to 9, and 12 to 21, and 2'-F modifications at positions 10, and 11; and
 - (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);

25 and

- (b) an antisense strand having:

- (i) a length of 23 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 3, 5, 7, 9, 11 to 13, 15, 17, 19, and 21 to 23, and 2'-F modifications at positions 2, 4, 6, 8, 10, 14, 16, 18, and 20 (counting from the 5' end);
- 30 and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);

wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a
35 blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
 - (i) a length of 21 nucleotides;

- (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
- (iii) 2'-F modifications at positions 1, 3, 5, 7, 9 to 11, and 13, and 2'-OMe modifications at positions 2, 4, 6, 8, 12, and 14 to 21; and
- 5 (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:
- (i) a length of 23 nucleotides;
- 10 (ii) 2'-OMe modifications at positions 1, 3, 5 to 7, 9, 11 to 13, 15, 17 to 19, and 21 to 23, and 2'-F modifications at positions 2, 4, 8, 10, 14, 16, and 20 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);
- 15 wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
- (i) a length of 21 nucleotides;
- 20 (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
- (iii) 2'-OMe modifications at positions 1, 2, 4, 6, 8, 12, 14, 15, 17, and 19 to 21, and 2'-F modifications at positions 3, 5, 7, 9 to 11, 13, 16, and 18; and
- (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and
- 25 between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:
- (i) a length of 25 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 4, 6, 7, 9, 11 to 13, 15, 17, and 19 to 23, 2'-F
- 30 modifications at positions 2, 3, 5, 8, 10, 14, 16, and 18, and deoxy-nucleotides (*e.g.* dT) at positions 24 and 25 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);
- 35 wherein the RNAi agents have a four nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
- (i) a length of 21 nucleotides;

- (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
- (iii) 2'-OMe modifications at positions 1 to 6, 8, and 12 to 21, and 2'-F modifications at positions 7, and 9 to 11; and
- 5 (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:
- (i) a length of 23 nucleotides;
- 10 (ii) 2'-OMe modifications at positions 1, 3 to 5, 7, 8, 10 to 13, 15, and 17 to 23, and 2'-F modifications at positions 2, 6, 9, 14, and 16 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);
- 15 wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
- (i) a length of 21 nucleotides;
- 20 (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
- (iii) 2'-OMe modifications at positions 1 to 6, 8, and 12 to 21, and 2'-F modifications at positions 7, and 9 to 11; and
- (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- 25 and
- (b) an antisense strand having:
- (i) a length of 23 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 3 to 5, 7, 10 to 13, 15, and 17 to 23, and 2'-F
- 30 modifications at positions 2, 6, 8, 9, 14, and 16 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);
- wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a
- 35 blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
- (i) a length of 19 nucleotides;

- (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
 - (iii) 2'-OMe modifications at positions 1 to 4, 6, and 10 to 19, and 2'-F modifications at positions 5, and 7 to 9; and
 - (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:
 - (i) a length of 21 nucleotides;
 - (ii) 2'-OMe modifications at positions 1, 3 to 5, 7, 10 to 13, 15, and 17 to 21, and 2'-F modifications at positions 2, 6, 8, 9, 14, and 16 (counting from the 5' end); and
 - (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 19 and 20, and between nucleotide positions 20 and 21 (counting from the 5' end);
- wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In certain embodiments, the iRNA for use in the methods of the invention is an agent selected from agents listed in any one of Tables 2-7. These agents may further comprise a ligand.

III. iRNAs Conjugated to Ligands

Another modification of the RNA of an iRNA of the invention involves chemically linking to the iRNA one or more ligands, moieties or conjugates that enhance the activity, cellular distribution, or cellular uptake of the iRNA *e.g.*, into a cell. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 1989, 86: 6553-6556). In other embodiments, the ligand is cholic acid (Manoharan *et al.*, *Biorg. Med. Chem. Lett.*, 1994, 4:1053-1060), a thioether, *e.g.*, beryl-S-tritylthiol (Manoharan *et al.*, *Ann. N.Y. Acad. Sci.*, 1992, 660:306-309; Manoharan *et al.*, *Biorg. Med. Chem. Lett.*, 1993, 3:2765-2770), a thiocholesterol (Oberhauser *et al.*, *Nucl. Acids Res.*, 1992, 20:533-538), an aliphatic chain, *e.g.*, dodecandiol or undecyl residues (Saison-Behmoaras *et al.*, *EMBO J*, 1991, 10:1111-1118; Kabanov *et al.*, *FEBS Lett.*, 1990, 259:327-330; Svinarchuk *et al.*, *Biochimie*, 1993, 75:49-54), a phospholipid, *e.g.*, di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-phosphonate (Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36:3651-3654; Shea *et al.*, *Nucl. Acids Res.*, 1990, 18:3777-3783), a polyamine or a polyethylene glycol chain (Manoharan *et al.*, *Nucleosides & Nucleotides*, 1995, 14:969-973), or adamantane acetic acid (Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36:3651-3654), a palmityl moiety (Mishra *et al.*, *Biochim. Biophys. Acta*, 1995, 1264:229-237), or an octadecylamine or hexylamino-carbonyloxycholesterol moiety (Crooke *et al.*, *J. Pharmacol. Exp. Ther.*, 1996, 277:923-937).

In certain embodiments, a ligand alters the distribution, targeting, or lifetime of an iRNA agent into which it is incorporated. In some embodiments a ligand provides an enhanced affinity for a

selected target, *e.g.*, molecule, cell or cell type, compartment, *e.g.*, a cellular or organ compartment, tissue, organ or region of the body, as, *e.g.*, compared to a species absent such a ligand. In some embodiments, ligands do not take part in duplex pairing in a duplexed nucleic acid.

Ligands can include a naturally occurring substance, such as a protein (*e.g.*, human serum albumin (HSA), low-density lipoprotein (LDL), or globulin); carbohydrate (*e.g.*, a dextran, pullulan, chitin, chitosan, inulin, cyclodextrin, N-acetylglucosamine, N-acetylgalactosamine, or hyaluronic acid); or a lipid. The ligand can also be a recombinant or synthetic molecule, such as a synthetic polymer, *e.g.*, a synthetic polyamino acid. Examples of polyamino acids include polyamino acid is a polylysine (PLL), poly L-aspartic acid, poly L-glutamic acid, styrene-maleic acid anhydride copolymer, poly(L-lactide-co-glycolide) copolymer, divinyl ether-maleic anhydride copolymer, N-(2-hydroxypropyl)methacrylamide copolymer (HMPA), polyethylene glycol (PEG), polyvinyl alcohol (PVA), polyurethane, poly(2-ethylacrylic acid), N-isopropylacrylamide polymers, or polyphosphazene. Example of polyamines include: polyethylenimine, polylysine (PLL), spermine, spermidine, polyamine, pseudopeptide-polyamine, peptidomimetic polyamine, dendrimer polyamine, arginine, amidine, protamine, cationic lipid, cationic porphyrin, quaternary salt of a polyamine, or an alpha helical peptide.

Ligands can also include targeting groups, *e.g.*, a cell or tissue targeting agent, *e.g.*, a lectin, glycoprotein, lipid or protein, *e.g.*, an antibody, that binds to a specified cell type such as a kidney cell. A targeting group can be a thyrotropin, melanotropin, lectin, glycoprotein, surfactant protein A, Mucin carbohydrate, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-glucosamine multivalent mannose, multivalent fucose, glycosylated polyaminoacids, multivalent galactose, transferrin, bisphosphonate, polyglutamate, polyaspartate, a lipid, cholesterol, a steroid, bile acid, folate, vitamin B12, vitamin A, biotin, or an RGD peptide or RGD peptide mimetic. In certain embodiments, the ligand is a multivalent galactose, *e.g.*, an N-acetyl-galactosamine.

Other examples of ligands include dyes, intercalating agents (*e.g.* acridines), cross-linkers (*e.g.* psoralene, mitomycin C), porphyrins (TPPC4, texaphyrin, Sapphyrin), polycyclic aromatic hydrocarbons (*e.g.*, phenazine, dihydrophenazine), artificial endonucleases (*e.g.* EDTA), lipophilic molecules, *e.g.*, cholesterol, cholic acid, adamantane acetic acid, 1-pyrene butyric acid, dihydrotestosterone, 1,3-Bis-O(hexadecyl)glycerol, geranyloxyhexyl group, hexadecylglycerol, borneol, menthol, 1,3-propanediol, heptadecyl group, palmitic acid, myristic acid, O3-(oleoyl)lithocholic acid, O3-(oleoyl)cholenic acid, dimethoxytrityl, or phenoxazine) and peptide conjugates (*e.g.*, antennapedia peptide, Tat peptide), alkylating agents, phosphate, amino, mercapto, PEG (*e.g.*, PEG-40K), MPEG, [MPEG]₂, polyamino, alkyl, substituted alkyl, radiolabeled markers, enzymes, haptens (*e.g.* biotin), transport/absorption facilitators (*e.g.*, aspirin, vitamin E, folic acid), synthetic ribonucleases (*e.g.*, imidazole, bisimidazole, histamine, imidazole clusters, acridine-imidazole conjugates, Eu³⁺ complexes of tetraazamacrocycles), dinitrophenyl, HRP, or AP.

Ligands can be proteins, *e.g.*, glycoproteins, or peptides, *e.g.*, molecules having a specific affinity for a co-ligand, or antibodies *e.g.*, an antibody, that binds to a specified cell type such as a hepatic cell. Ligands can also include hormones and hormone receptors. They can also include non-

peptidic species, such as lipids, lectins, carbohydrates, vitamins, cofactors, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-glucosamine multivalent mannose, or multivalent fucose. The ligand can be, for example, a lipopolysaccharide, an activator of p38 MAP kinase, or an activator of NF- κ B.

5 The ligand can be a substance, *e.g.*, a drug, which can increase the uptake of the iRNA agent into the cell, for example, by disrupting the cell's cytoskeleton, *e.g.*, by disrupting the cell's microtubules, microfilaments, or intermediate filaments. The drug can be, for example, taxol, vincristine, vinblastine, cytochalasin, nocodazole, japlakinolide, latrunculin A, phalloidin, swinholide A, indanocine, or myoservin.

10 In some embodiments, a ligand attached to an iRNA as described herein acts as a pharmacokinetic modulator (PK modulator). PK modulators include lipophiles, bile acids, steroids, phospholipid analogues, peptides, protein binding agents, PEG, vitamins, *etc.* Exemplary PK modulators include, but are not limited to, cholesterol, fatty acids, cholic acid, lithocholic acid, dialkylglycerides, diacylglyceride, phospholipids, sphingolipids, naproxen, ibuprofen, vitamin E,
15 biotin. Oligonucleotides that comprise a number of phosphorothioate linkages are also known to bind to serum protein, thus short oligonucleotides, *e.g.*, oligonucleotides of about 5 bases, 10 bases, 15 bases, or 20 bases, comprising multiple of phosphorothioate linkages in the backbone are also amenable to the present invention as ligands (*e.g.* as PK modulating ligands). In addition, aptamers that bind serum components (*e.g.* serum proteins) are also suitable for use as PK modulating ligands
20 in the embodiments described herein.

 Ligand-conjugated iRNAs of the invention may be synthesized by the use of an oligonucleotide that bears a pendant reactive functionality, such as that derived from the attachment of a linking molecule onto the oligonucleotide (described below). This reactive oligonucleotide may be reacted directly with commercially-available ligands, ligands that are synthesized bearing any of a
25 variety of protecting groups, or ligands that have a linking moiety attached thereto.

 The oligonucleotides used in the conjugates of the present invention may be conveniently and routinely made through the well-known technique of solid-phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems® (Foster City, Calif.). Any other methods for such synthesis known in the art may additionally or alternatively be
30 employed. It is also known to use similar techniques to prepare other oligonucleotides, such as the phosphorothioates and alkylated derivatives.

 In the ligand-conjugated iRNAs and ligand-molecule bearing sequence-specific linked nucleosides of the present invention, the oligonucleotides and oligonucleosides may be assembled on a suitable DNA synthesizer utilizing standard nucleotide or nucleoside precursors, or nucleotide or
35 nucleoside conjugate precursors that already bear the linking moiety, ligand-nucleotide or nucleoside-conjugate precursors that already bear the ligand molecule, or non-nucleoside ligand-bearing building blocks.

 When using nucleotide-conjugate precursors that already bear a linking moiety, the synthesis of the sequence-specific linked nucleosides is typically completed, and the ligand molecule is then

reacted with the linking moiety to form the ligand-conjugated oligonucleotide. In some embodiments, the oligonucleotides or linked nucleosides of the present invention are synthesized by an automated synthesizer using phosphoramidites derived from ligand-nucleoside conjugates in addition to the standard phosphoramidites and non-standard phosphoramidites that are commercially available and routinely used in oligonucleotide synthesis.

A. Lipid Conjugates

In certain embodiments, the ligand or conjugate is a lipid or lipid-based molecule. In one embodiment, such a lipid or lipid-based molecule binds a serum protein, *e.g.*, human serum albumin (HSA). An HSA binding ligand allows for distribution of the conjugate to a target tissue, *e.g.*, a non-kidney target tissue of the body. For example, the target tissue can be the liver, including parenchymal cells of the liver. Other molecules that can bind HSA can also be used as ligands. For example, naproxen or aspirin can be used. A lipid or lipid-based ligand can (a) increase resistance to degradation of the conjugate, (b) increase targeting or transport into a target cell or cell membrane, or (c) can be used to adjust binding to a serum protein, *e.g.*, HSA.

A lipid based ligand can be used to inhibit, *e.g.*, control the binding of the conjugate to a target tissue. For example, a lipid or lipid-based ligand that binds to HSA more strongly will be less likely to be targeted to the kidney and therefore less likely to be cleared from the body. A lipid or lipid-based ligand that binds to HSA less strongly can be used to target the conjugate to the kidney.

In certain embodiments, the lipid based ligand binds HSA. In one embodiment, it binds HSA with a sufficient affinity such that the conjugate will be distributed to a non-kidney tissue. However, it is preferred that the affinity not be so strong that the HSA-ligand binding cannot be reversed.

In other embodiments, the lipid based ligand binds HSA weakly or not at all. In one embodiment, the conjugate will be distributed to the kidney. Other moieties that target to kidney cells can also be used in place of, or in addition to, the lipid based ligand.

In another aspect, the ligand is a moiety, *e.g.*, a vitamin, which is taken up by a target cell, *e.g.*, a proliferating cell. These are particularly useful for treating disorders characterized by unwanted cell proliferation, *e.g.*, of the malignant or non-malignant type, *e.g.*, cancer cells. Exemplary vitamins include vitamin A, E, and K. Other exemplary vitamins include are B vitamin, *e.g.*, folic acid, B12, riboflavin, biotin, pyridoxal or other vitamins or nutrients taken up by target cells such as liver cells. Also included are HSA and low density lipoprotein (LDL).

B. Cell Permeation Agents

In another aspect, the ligand is a cell-permeation agent, such as, a helical cell-permeation agent. In one embodiment, the agent is amphipathic. An exemplary agent is a peptide such as tat or antennopodia. If the agent is a peptide, it can be modified, including a peptidylmimetic, invertomers, non-peptide or pseudo-peptide linkages, and use of D-amino acids. In one embodiment, the helical agent is an alpha-helical agent, which has a lipophilic and a lipophobic phase.

The ligand can be a peptide or peptidomimetic. A peptidomimetic (also referred to herein as an oligopeptidomimetic) is a molecule capable of folding into a defined three-dimensional structure

similar to a natural peptide. The attachment of peptide and peptidomimetics to iRNA agents can affect pharmacokinetic distribution of the iRNA, such as by enhancing cellular recognition and absorption. The peptide or peptidomimetic moiety can be about 5-50 amino acids long, *e.g.*, about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 amino acids long.

- 5 A peptide or peptidomimetic can be, for example, a cell permeation peptide, cationic peptide, amphipathic peptide, or hydrophobic peptide (*e.g.*, consisting primarily of Tyr, Trp, or Phe). The peptide moiety can be a dendrimer peptide, constrained peptide or crosslinked peptide. In another alternative, the peptide moiety can include a hydrophobic membrane translocation sequence (MTS). An exemplary hydrophobic MTS-containing peptide is RFGF having the amino acid sequence
- 10 AAVALLPAVLLALLAP (SEQ ID NO: 14). An RFGF analogue (*e.g.*, amino acid sequence AALLPVLLAAP (SEQ ID NO:15) containing a hydrophobic MTS can also be a targeting moiety. The peptide moiety can be a “delivery” peptide, which can carry large polar molecules including peptides, oligonucleotides, and protein across cell membranes. For example, sequences from the HIV Tat protein (GRKKRRQRRRPPQ (SEQ ID NO:16) and the Drosophila Antennapedia protein
- 15 (RQIKIWFQNRRMKWKK (SEQ ID NO:17) have been found to be capable of functioning as delivery peptides. A peptide or peptidomimetic can be encoded by a random sequence of DNA, such as a peptide identified from a phage-display library, or one-bead-one-compound (OBOC) combinatorial library (Lam *et al.*, Nature, 354:82-84, 1991). Examples of a peptide or peptidomimetic tethered to a dsRNA agent *via* an incorporated monomer unit for cell targeting
- 20 purposes is an arginine-glycine-aspartic acid (RGD)-peptide, or RGD mimic. A peptide moiety can range in length from about 5 amino acids to about 40 amino acids. The peptide moieties can have a structural modification, such as to increase stability or direct conformational properties. Any of the structural modifications described below can be utilized.

- An RGD peptide for use in the compositions and methods of the invention may be linear or
- 25 cyclic, and may be modified, *e.g.*, glycosylated or methylated, to facilitate targeting to a specific tissue(s). RGD-containing peptides and peptidomimetics may include D-amino acids, as well as synthetic RGD mimics. In addition to RGD, one can use other moieties that target the integrin ligand, *e.g.*, PECAM-1 or VEGF.

- A “cell permeation peptide” is capable of permeating a cell, *e.g.*, a microbial cell, such as a
- 30 bacterial or fungal cell, or a mammalian cell, such as a human cell. A microbial cell-permeating peptide can be, for example, an α -helical linear peptide (*e.g.*, LL-37 or Ceropin P1), a disulfide bond-containing peptide (*e.g.*, α -defensin, β -defensin or bactenecin), or a peptide containing only one or two dominating amino acids (*e.g.*, PR-39 or indolicidin). A cell permeation peptide can also include a nuclear localization signal (NLS). For example, a cell permeation peptide can be a bipartite
- 35 amphipathic peptide, such as MPG, which is derived from the fusion peptide domain of HIV-1 gp41 and the NLS of SV40 large T antigen (Simeoni *et al.*, Nucl. Acids Res. 31:2717-2724, 2003).

C. Carbohydrate Conjugates

In some embodiments of the compositions and methods of the invention, an iRNA further comprises a carbohydrate. The carbohydrate conjugated iRNA is advantageous for the *in vivo* delivery of nucleic acids, as well as compositions suitable for *in vivo* therapeutic use, as described herein. As used herein, "carbohydrate" refers to a compound which is either a carbohydrate *per se* made up of one or more monosaccharide units having at least 6 carbon atoms (which can be linear, branched or cyclic) with an oxygen, nitrogen or sulfur atom bonded to each carbon atom; or a compound having as a part thereof a carbohydrate moiety made up of one or more monosaccharide units each having at least six carbon atoms (which can be linear, branched or cyclic), with an oxygen, nitrogen or sulfur atom bonded to each carbon atom. Representative carbohydrates include the sugars (mono-, di-, tri-, and oligosaccharides containing from about 4, 5, 6, 7, 8, or 9 monosaccharide units), and polysaccharides such as starches, glycogen, cellulose and polysaccharide gums. Specific monosaccharides include C5 and above (*e.g.*, C5, C6, C7, or C8) sugars; di- and trisaccharides include sugars having two or three monosaccharide units (*e.g.*, C5, C6, C7, or C8).

In certain embodiments, a carbohydrate conjugate for use in the compositions and methods of the invention is a monosaccharide.

In certain embodiments, the monosaccharide is an N-acetylgalactosamine (GalNAc). GalNAc conjugates, which comprise one or more N-acetylgalactosamine (GalNAc) derivatives, are described, for example, in US 8,106,022, the entire content of which is hereby incorporated herein by reference.

In some embodiments, the GalNAc conjugate serves as a ligand that targets the iRNA to particular cells. In some embodiments, the GalNAc conjugate targets the iRNA to liver cells, *e.g.*, by serving as a ligand for the asialoglycoprotein receptor of liver cells (*e.g.*, hepatocytes).

In some embodiments, the carbohydrate conjugate comprises one or more GalNAc derivatives. The GalNAc derivatives may be attached *via* a linker, *e.g.*, a bivalent or trivalent branched linker. In some embodiments the GalNAc conjugate is conjugated to the 3' end of the sense strand. In some embodiments, the GalNAc conjugate is conjugated to the iRNA agent (*e.g.*, to the 3' end of the sense strand) *via* a linker, *e.g.*, a linker as described herein. In some embodiments the GalNAc conjugate is conjugated to the 5' end of the sense strand. In some embodiments, the GalNAc conjugate is conjugated to the iRNA agent (*e.g.*, to the 5' end of the sense strand) *via* a linker, *e.g.*, a linker as described herein.

In certain embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a monovalent linker. In some embodiments, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a bivalent linker. In yet other embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a trivalent linker. In other embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a tetravalent linker.


In certain embodiments, the double stranded RNAi agents of the invention comprise one GalNAc or GalNAc derivative attached to the iRNA agent. In certain embodiments, the double stranded RNAi agents of the invention comprise a plurality (*e.g.*, 2, 3, 4, 5, or 6) GalNAc or GalNAc

derivatives, each independently attached to a plurality of nucleotides of the double stranded RNAi agent through a plurality of monovalent linkers.


In some embodiments, for example, when the two strands of an iRNA agent of the invention are part of one larger molecule connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming a hairpin loop comprising, a plurality of unpaired nucleotides, each unpaired nucleotide within the hairpin loop may independently comprise a GalNAc or GalNAc derivative attached *via* a monovalent linker. The hairpin loop may also be formed by an extended overhang in one strand of the duplex.

In some embodiments, for example, when the two strands of an iRNA agent of the invention are part of one larger molecule connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming a hairpin loop comprising, a plurality of unpaired nucleotides, each unpaired nucleotide within the hairpin loop may independently comprise a GalNAc or GalNAc derivative attached *via* a monovalent linker. The hairpin loop may also be formed by an extended overhang in one strand of the duplex.


In one embodiment, a carbohydrate conjugate for use in the compositions and methods of the invention is selected from the group consisting of:




Formula II,




Formula III,




Formula IV,




Formula V,




Formula VI,




Formula VII,




Formula VIII,




Formula IX,




Formula X,




Formula XI,




Formula XII,




Formula XIII,




Formula XIV,




Formula XV,




Formula XVI,




Formula XVII,




Formula XVIII,



Formula XIX,




Formula XX,




Formula XXI,


5




Formula XXII,




Formula XXIII;






, wherein Y is O or S and n is 3 -6 (Formula XXIV);




, wherein Y is O or S and n is 3-6 (Formula XXV);




Formula XXVI;




Formula XXVII; Formula XXIX;




Formula XXXI;



Formula XXX;




, and



Formula XXXII;


5 Formula XXXIII.



Formula XXXIV.


In another embodiment, a carbohydrate conjugate for use in the compositions and methods of the invention is a monosaccharide. In one embodiment, the monosaccharide is an N-

5 acetylgalactosamine, such as




Formula II.


In some embodiments, the RNAi agent is attached to the carbohydrate conjugate *via* a linker as shown in the following schematic, wherein X is O or S



In some embodiments, the RNAi agent is conjugated to L96 as defined in Table 1 and shown below:




Another representative carbohydrate conjugate for use in the embodiments described herein includes, but is not limited to,



(Formula XXXVI), when one of X or Y is an oligonucleotide, the other is a hydrogen.

In some embodiments, a suitable ligand is a ligand disclosed in WO 2019/055633, the entire contents of which are incorporated herein by reference. In one embodiment the ligand comprises the structure below:



In certain embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a monovalent linker. In some embodiments, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a bivalent linker. In yet other embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a trivalent linker.

In one embodiment, the double stranded RNAi agents of the invention comprise one or more GalNAc or GalNAc derivative attached to the iRNA agent. The GalNAc may be attached to any nucleotide *via* a linker on the sense strand or antisense strand. The GalNAc may be attached to the 5'-end of the sense strand, the 3' end of the sense strand, the 5'-end of the antisense strand, or the 3' –
 5 end of the antisense strand. In one embodiment, the GalNAc is attached to the 3' end of the sense strand, *e.g.*, *via* a trivalent linker.

In other embodiments, the double stranded RNAi agents of the invention comprise a plurality (*e.g.*, 2, 3, 4, 5, or 6) GalNAc or GalNAc derivatives, each independently attached to a plurality of nucleotides of the double stranded RNAi agent through a plurality of linkers, *e.g.*, monovalent linkers.

10 In some embodiments, for example, when the two strands of an iRNA agent of the invention is part of one larger molecule connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming a hairpin loop comprising, a plurality of unpaired nucleotides, each unpaired nucleotide within the hairpin loop may independently comprise a GalNAc or GalNAc derivative attached *via* a monovalent linker.

15 In some embodiments, the carbohydrate conjugate further comprises one or more additional ligands as described above, such as, but not limited to, a PK modulator or a cell permeation peptide.

Additional carbohydrate conjugates and linkers suitable for use in the present invention include those described in PCT Publication Nos. WO 2014/179620 and WO 2014/179627, the entire contents of each of which are incorporated herein by reference.

20 D. Linkers

In some embodiments, the conjugate or ligand described herein can be attached to an iRNA oligonucleotide with various linkers that can be cleavable or non-cleavable.

The term "linker" or "linking group" means an organic moiety that connects two parts of a compound, *e.g.*, covalently attaches two parts of a compound. Linkers typically comprise a direct
 25 bond or an atom such as oxygen or sulfur, a unit such as NR₈, C(O), C(O)NH, SO, SO₂, SO₂NH or a chain of atoms, such as, but not limited to, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, heterocyclalkyl, heterocyclalkenyl, heterocyclalkynyl, aryl, heteroaryl, heterocycl, cycloalkyl, cycloalkenyl, alkylarylalkyl, alkylarylalkenyl, alkylarylalkynyl, alkenylarylalkyl, alkenylarylalkenyl, alkenylarylalkynyl, alkynylarylalkyl, alkynylarylalkenyl, alkynylarylalkynyl, alkylheteroarylalkyl, alkylheteroarylalkenyl, alkylheteroarylalkynyl, alkenylheteroarylalkyl, alkenylheteroarylalkenyl, alkenylheteroarylalkynyl, alkynylheteroarylalkyl, alkynylheteroarylalkenyl, alkynylheteroarylalkynyl, alkylheterocyclalkyl, alkylheterocyclalkenyl, alkylheterocyclalkynyl, alkenylheterocyclalkyl, alkenylheterocyclalkenyl, alkenylheterocyclalkynyl, alkynylheterocyclalkyl, alkynylheterocyclalkenyl, alkynylheterocyclalkynyl, alkylaryl, alkenylaryl, alkynylaryl, alkylheteroaryl, alkenylheteroaryl, alkynylheteroaryl, which one or more methylenes can be interrupted or terminated by O, S, S(O), SO₂, N(R₈), C(O), substituted or unsubstituted aryl,

substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocyclic; where R8 is hydrogen, acyl, aliphatic, or substituted aliphatic. In one embodiment, the linker is about 1-24 atoms, 2-24, 3-24, 4-24, 5-24, 6-24, 6-18, 7-18, 8-18, 7-17, 8-17, 6-16, 7-17, or 8-16 atoms.

5 A cleavable linking group is one which is sufficiently stable outside the cell, but which upon entry into a target cell is cleaved to release the two parts the linker is holding together. In an exemplary embodiment, the cleavable linking group is cleaved at least about 10 times, 20, times, 30 times, 40 times, 50 times, 60 times, 70 times, 80 times, 90 times, or more, or at least 100 times faster in a target cell or under a first reference condition (which can, *e.g.*, be selected to mimic or represent intracellular conditions) than in the blood of a subject, or under a second reference condition (which
10 can, *e.g.*, be selected to mimic or represent conditions found in the blood or serum).

Cleavable linking groups are susceptible to cleavage agents, *e.g.*, pH, redox potential, or the presence of degradative molecules. Generally, cleavage agents are more prevalent or found at higher levels or activities inside cells than in serum or blood. Examples of such degradative agents include: redox agents which are selected for particular substrates or which have no substrate specificity,
15 including, *e.g.*, oxidative or reductive enzymes or reductive agents such as mercaptans, present in cells, that can degrade a redox cleavable linking group by reduction; esterases; endosomes or agents that can create an acidic environment, *e.g.*, those that result in a pH of five or lower; enzymes that can hydrolyze or degrade an acid cleavable linking group by acting as a general acid, peptidases (which can be substrate specific), and phosphatases.

20 A cleavable linkage group, such as a disulfide bond can be susceptible to pH. The pH of human serum is 7.4, while the average intracellular pH is slightly lower, ranging from about 7.1-7.3. Endosomes have a more acidic pH, in the range of 5.5-6.0, and lysosomes have an even more acidic pH at around 5.0. Some linkers will have a cleavable linking group that is cleaved at a selected pH, thereby releasing a cationic lipid from the ligand inside the cell, or into the desired compartment of
25 the cell.

A linker can include a cleavable linking group that is cleavable by a particular enzyme. The type of cleavable linking group incorporated into a linker can depend on the cell to be targeted. For example, a liver-targeting ligand can be linked to a cationic lipid through a linker that includes an ester group. Liver cells are rich in esterases, and therefore the linker will be cleaved more efficiently
30 in liver cells than in cell types that are not esterase-rich. Other cell-types rich in esterases include cells of the lung, renal cortex, and testis.

Linkers that contain peptide bonds can be used when targeting cell types rich in peptidases, such as liver cells and synoviocytes.

In general, the suitability of a candidate cleavable linking group can be evaluated by testing
35 the ability of a degradative agent (or condition) to cleave the candidate linking group. It will also be desirable to also test the candidate cleavable linking group for the ability to resist cleavage in the blood or when in contact with other non-target tissue. Thus, one can determine the relative susceptibility to cleavage between a first and a second condition, where the first is selected to be indicative of cleavage in a target cell and the second is selected to be indicative of cleavage in other

tissues or biological fluids, *e.g.*, blood or serum. The evaluations can be carried out in cell free systems, in cells, in cell culture, in organ or tissue culture, or in whole animals. It can be useful to make initial evaluations in cell-free or culture conditions and to confirm by further evaluations in whole animals. In certain embodiments, useful candidate compounds are cleaved at least about 2, 4, 5 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 times faster in the cell (or under *in vitro* conditions selected to mimic intracellular conditions) as compared to blood or serum (or under *in vitro* conditions selected to mimic extracellular conditions).

i. Redox cleavable linking groups

10 In certain embodiments, a cleavable linking group is a redox cleavable linking group that is cleaved upon reduction or oxidation. An example of reductively cleavable linking group is a disulphide linking group (-S-S-). To determine if a candidate cleavable linking group is a suitable “reductively cleavable linking group,” or for example is suitable for use with a particular iRNA moiety and particular targeting agent one can look to methods described herein. For example, a 15 candidate can be evaluated by incubation with dithiothreitol (DTT), or other reducing agent using reagents known in the art, which mimic the rate of cleavage which would be observed in a cell, *e.g.*, a target cell. The candidates can also be evaluated under conditions which are selected to mimic blood or serum conditions. In one, candidate compounds are cleaved by at most about 10% in the blood. In other embodiments, useful candidate compounds are degraded at least about 2, 4, 10, 20, 30, 40, 50, 20 60, 70, 80, 90, or about 100 times faster in the cell (or under *in vitro* conditions selected to mimic intracellular conditions) as compared to blood (or under *in vitro* conditions selected to mimic extracellular conditions). The rate of cleavage of candidate compounds can be determined using standard enzyme kinetics assays under conditions chosen to mimic intracellular media and compared to conditions chosen to mimic extracellular media.

25

ii. Phosphate-based cleavable linking groups

In other embodiments, a cleavable linker comprises a phosphate-based cleavable linking group. A phosphate-based cleavable linking group is cleaved by agents that degrade or hydrolyze the phosphate group. An example of an agent that cleaves phosphate groups in cells are enzymes such as 30 phosphatases in cells. Examples of phosphate-based linking groups are -O-P(O)(ORk)-O-, -O-P(S)(ORk)-O-, -O-P(S)(SRk)-O-, -S-P(O)(ORk)-O-, -O-P(O)(ORk)-S-, -S-P(O)(ORk)-S-, -O-P(S)(ORk)-S-, -S-P(S)(ORk)-O-, -O-P(O)(Rk)-O-, -O-P(S)(Rk)-O-, -S-P(O)(Rk)-O-, -S-P(S)(Rk)-O-, -S-P(O)(Rk)-S-, -O-P(S)(Rk)-S-, wherein Rk at each occurrence can be, independently, C1-C20 alkyl, C1-C20 haloalkyl, C6-C10 aryl, or C7-C12 aralkyl. Exemplary embodiments include -O-P(O)(OH)-O-, -O-P(S)(OH)-O-, -O-P(S)(SH)-O-, -S-P(O)(OH)-O-, -O-P(O)(OH)-S-, -S-P(O)(OH)-S-, -O-P(S)(OH)-S-, -S-P(S)(OH)-O-, -O-P(O)(H)-O-, -O-P(S)(H)-O-, -S-P(O)(H)-O-, -S-P(S)(H)-O-, -S-P(O)(H)-S-, and -O-P(S)(H)-S-. In certain embodiments a phosphate-based linking group is -O-

35

P(O)(OH)-O-. These candidates can be evaluated using methods analogous to those described above.

iii. Acid cleavable linking groups

In other embodiments, a cleavable linker comprises an acid cleavable linking group. An acid cleavable linking group is a linking group that is cleaved under acidic conditions. In certain
 5 embodiments acid cleavable linking groups are cleaved in an acidic environment with a pH of about 6.5 or lower (*e.g.*, about 6.0, 5.5, 5.0, or lower), or by agents such as enzymes that can act as a general acid. In a cell, specific low pH organelles, such as endosomes and lysosomes can provide a cleaving environment for acid cleavable linking groups. Examples of acid cleavable linking groups include but are not limited to hydrazones, esters, and esters of amino acids. Acid cleavable groups can have the
 10 general formula -C=NN-, C(O)O, or -OC(O). An exemplary embodiment is when the carbon attached to the oxygen of the ester (the alkoxy group) is an aryl group, substituted alkyl group, or tertiary alkyl group such as dimethyl pentyl or t-butyl. These candidates can be evaluated using methods analogous to those described above.


15 *iv. Ester-based linking groups*

In other embodiments, a cleavable linker comprises an ester-based cleavable linking group. An ester-based cleavable linking group is cleaved by enzymes such as esterases and amidases in cells. Examples of ester-based cleavable linking groups include, but are not limited to, esters of alkylene, alkenylene and alkynylene groups. Ester cleavable linking groups have the general formula -C(O)O-,
 20 or -OC(O)-. These candidates can be evaluated using methods analogous to those described above.


v. Peptide-based cleaving groups

In yet other embodiments, a cleavable linker comprises a peptide-based cleavable linking group. A peptide-based cleavable linking group is cleaved by enzymes such as peptidases and
 25 proteases in cells. Peptide-based cleavable linking groups are peptide bonds formed between amino acids to yield oligopeptides (*e.g.*, dipeptides, tripeptides *etc.*) and polypeptides. Peptide-based cleavable groups do not include the amide group (-C(O)NH-). The amide group can be formed between any alkylene, alkenylene or alkynylene. A peptide bond is a special type of amide bond formed between amino acids to yield peptides and proteins. The peptide based cleavage group is
 30 generally limited to the peptide bond (*i.e.*, the amide bond) formed between amino acids yielding peptides and proteins and does not include the entire amide functional group. Peptide-based cleavable linking groups have the general formula -NHCHRAC(O)NHCHRBC(O)-, where RA and RB are the R groups of the two adjacent amino acids. These candidates can be evaluated using methods analogous to those described above.


35 In some embodiments, an iRNA of the invention is conjugated to a carbohydrate through a linker. Non-limiting examples of iRNA carbohydrate conjugates with linkers of the compositions and methods of the invention include, but are not limited to,




(Formula XXXVII),



(Formula XXXVIII),




(Formula XXXIX),




5


(Formula XL),




(Formula XLI),



(Formula XLII),



(Formula XLIII), and




5

(Formula XLIV), when one of X or Y is an oligonucleotide, the other is a hydrogen.



In certain embodiments of the compositions and methods of the invention, a ligand is one or more “GalNAc” (N-acetylgalactosamine) derivatives attached through a bivalent or trivalent branched linker.

10 In one embodiment, a dsRNA of the invention is conjugated to a bivalent or trivalent branched linker selected from the group of structures shown in any of formula (XLV) – (XLVI):


Formula XXXXV



Formula XLVI



, or



;

5 Formula XLVII

Formula XLVIII






wherein:

q^{2A}, q^{2B}, q^{3A}, q^{3B}, q^{4A}, q^{4B}, q^{5A}, q^{5B} and q^{5C} represent independently for each occurrence 0-20 and wherein the repeating unit can be the same or different;

P^{2A}, P^{2B}, P^{3A}, P^{3B}, P^{4A}, P^{4B}, P^{5A}, P^{5B}, P^{5C}, T^{2A}, T^{2B}, T^{3A}, T^{3B}, T^{4A}, T^{4B}, T^{5A}, T^{5B}, T^{5C} are each

- 10 independently for each occurrence absent, CO, NH, O, S, OC(O), NHC(O), CH₂, CH₂NH or CH₂O; Q^{2A}, Q^{2B}, Q^{3A}, Q^{3B}, Q^{4A}, Q^{4B}, Q^{5A}, Q^{5B}, Q^{5C} are independently for each occurrence absent, alkylene, substituted alkylene wherein one or more methylenes can be interrupted or terminated by one or more of O, S, S(O), SO₂, N(R^N), C(R')=C(R''), C≡C or C(O);

R^{2A}, R^{2B}, R^{3A}, R^{3B}, R^{4A}, R^{4B}, R^{5A}, R^{5B}, R^{5C} are each independently for each occurrence absent, NH, O,


- 15 S, CH₂, C(O)O, C(O)NH, NHCH(R^a)C(O), -C(O)-CH(R^a)-NH-, CO, CH=N-O, , , , , , or heterocyclyl;

L^{2A}, L^{2B}, L^{3A}, L^{3B}, L^{4A}, L^{4B}, L^{5A}, L^{5B} and L^{5C} represent the ligand; *i.e.* each independently for each

occurrence a monosaccharide (such as GalNAc), disaccharide, trisaccharide, tetrasaccharide,

oligosaccharide, or polysaccharide; and R^a is H or amino acid side chain. Trivalent conjugating

- 20 GalNAc derivatives are particularly useful for use with RNAi agents for inhibiting the expression of a target gene, such as those of formula (XLIX):



wherein L^{5A} , L^{5B} and L^{5C} represent a monosaccharide, such as GalNAc derivative.

Examples of suitable bivalent and trivalent branched linker groups conjugating GalNAc derivatives include, but are not limited to, the structures recited above as formulas II, VII, XI, X, and XIII.

Representative U.S. Patents that teach the preparation of RNA conjugates include, but are not limited to, U.S. Patent Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928; 5,688,941; 6,294,664; 6,320,017; 6,576,752; 6,783,931; 6,900,297; 7,037,646; and 8,106,022, the entire contents of each of which are hereby incorporated herein by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications can be incorporated in a single compound or even at a single nucleoside within an iRNA. The present invention also includes iRNA compounds that are chimeric compounds.

“Chimeric” iRNA compounds or “chimeras,” in the context of this invention, are iRNA compounds, such as, dsRNAi agents, that contain two or more chemically distinct regions, each made up of at least one monomer unit, *i.e.*, a nucleotide in the case of a dsRNA compound. These iRNAs typically contain at least one region wherein the RNA is modified so as to confer upon the iRNA increased resistance to nuclease degradation, increased cellular uptake, or increased binding affinity for the target nucleic acid. An additional region of the iRNA can serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of iRNA inhibition of gene expression. Consequently, comparable results can often be obtained with shorter iRNAs when chimeric dsRNAs are used, compared to phosphorothioate deoxy dsRNAs hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

In certain instances, the RNA of an iRNA can be modified by a non-ligand group. A number of non-ligand molecules have been conjugated to iRNAs in order to enhance the activity, cellular

distribution or cellular uptake of the iRNA, and procedures for performing such conjugations are available in the scientific literature. Such non-ligand moieties have included lipid moieties, such as cholesterol (Kubo, T. *et al.*, *Biochem. Biophys. Res. Comm.*, 2007, 365(1):54-61; Letsinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 1989, 86:6553), cholic acid (Manoharan *et al.*, *Bioorg. Med. Chem. Lett.*, 1994, 4:1053), a thioether, *e.g.*, hexyl-S-tritylthiol (Manoharan *et al.*, *Ann. N.Y. Acad. Sci.*, 1992, 660:306; Manoharan *et al.*, *Bioorg. Med. Chem. Lett.*, 1993, 3:2765), a thiocholesterol (Oberhauser *et al.*, *Nucl. Acids Res.*, 1992, 20:533), an aliphatic chain, *e.g.*, dodecandiol or undecyl residues (Saison-Behmoaras *et al.*, *EMBO J.*, 1991, 10:111; Kabanov *et al.*, *FEBS Lett.*, 1990, 259:327; Svinarchuk *et al.*, *Biochimie*, 1993, 75:49), a phospholipid, *e.g.*, di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36:3651; Shea *et al.*, *Nucl. Acids Res.*, 1990, 18:3777), a polyamine or a polyethylene glycol chain (Manoharan *et al.*, *Nucleosides & Nucleotides*, 1995, 14:969), or adamantane acetic acid (Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36:3651), a palmityl moiety (Mishra *et al.*, *Biochim. Biophys. Acta*, 1995, 1264:229), or an octadecylamine or hexylamino-carbonyl-oxcholesterol moiety (Crooke *et al.*, *J. Pharmacol. Exp. Ther.*, 1996, 277:923). Representative United States patents that teach the preparation of such RNA conjugates have been listed above. Typical conjugation protocols involve the synthesis of RNAs bearing an aminolinker at one or more positions of the sequence. The amino group is then reacted with the molecule being conjugated using appropriate coupling or activating reagents. The conjugation reaction can be performed either with the RNA still bound to the solid support or following cleavage of the RNA, in solution phase. Purification of the RNA conjugate by HPLC typically affords the pure conjugate.

IV. Delivery of an iRNA of the Invention

The delivery of an iRNA of the invention to a cell *e.g.*, a cell within a subject, such as a human subject (*e.g.*, a subject in need thereof, such as a subject susceptible to or diagnosed with a TMPRSS6-associated disorder, *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis) can be achieved in a number of different ways. For example, delivery may be performed by contacting a cell with an iRNA of the invention either *in vitro* or *in vivo*. *In vivo* delivery may also be performed directly by administering a composition comprising an iRNA, *e.g.*, a dsRNA, to a subject. Alternatively, *in vivo* delivery may be performed indirectly by administering one or more vectors that encode and direct the expression of the iRNA. These alternatives are discussed further below.

In general, any method of delivering a nucleic acid molecule (*in vitro* or *in vivo*) can be adapted for use with an iRNA of the invention (see *e.g.*, Akhtar S. and Julian RL. (1992) *Trends Cell Biol.* 2(5):139-144 and WO94/02595, which are incorporated herein by reference in their entireties). For *in vivo* delivery, factors to consider in order to deliver an iRNA molecule include, for example, biological stability of the delivered molecule, prevention of non-specific effects, and accumulation of the delivered molecule in the target tissue. RNA interference has also shown success with local delivery to the CNS by direct injection (Dorn, G., *et al.* (2004) *Nucleic Acids* 32:e49; Tan, PH., *et al*

(2005) *Gene Ther.* 12:59-66; Makimura, H., *et al* (2002) *BMC Neurosci.* 3:18; Shishkina, GT., *et al* (2004) *Neuroscience* 129:521-528; Thakker, ER., *et al* (2004) *Proc. Natl. Acad. Sci. U.S.A.*

101:17270-17275; Akaneya, Y., *et al* (2005) *J. Neurophysiol.* 93:594-602). Modification of the RNA or the pharmaceutical carrier can also permit targeting of the iRNA to the target tissue and avoid

5 undesirable off-target effects. iRNA molecules can be modified by chemical conjugation to lipophilic groups such as cholesterol to enhance cellular uptake and prevent degradation. For example, an iRNA directed against ApoB conjugated to a lipophilic cholesterol moiety was injected systemically into mice and resulted in knockdown of apoB mRNA in both the liver and jejunum (Soutschek, J., *et al* (2004) *Nature* 432:173-178).

10 In an alternative embodiment, the iRNA can be delivered using drug delivery systems such as a nanoparticle, a dendrimer, a polymer, liposomes, or a cationic delivery system. Positively charged cationic delivery systems facilitate binding of an iRNA molecule (negatively charged) and also enhance interactions at the negatively charged cell membrane to permit efficient uptake of an iRNA by the cell. Cationic lipids, dendrimers, or polymers can either be bound to an iRNA, or induced to

15 form a vesicle or micelle (see *e.g.*, Kim SH, *et al* (2008) *Journal of Controlled Release* 129(2):107-116) that encases an iRNA. The formation of vesicles or micelles further prevents degradation of the iRNA when administered systemically. Methods for making and administering cationic- iRNA complexes are well within the abilities of one skilled in the art (see *e.g.*, Sorensen, DR, *et al* (2003) *J. Mol. Biol* 327:761-766; Verma, UN, *et al* (2003) *Clin. Cancer Res.* 9:1291-1300; Arnold, AS *et al*

20 (2007) *J. Hypertens.* 25:197-205, which are incorporated herein by reference in their entirety). Some non-limiting examples of drug delivery systems useful for systemic delivery of iRNAs include DOTAP (Sorensen, DR., *et al* (2003), *supra*; Verma, UN, *et al* (2003), *supra*), "solid nucleic acid lipid particles" (Zimmermann, TS, *et al* (2006) *Nature* 441:111-114), cardiolipin (Chien, PY, *et al* (2005) *Cancer Gene Ther.* 12:321-328; Pal, A, *et al* (2005) *Int J. Oncol.* 26:1087-1091),

25 polyethyleneimine (Bonnet ME, *et al* (2008) *Pharm. Res.* Aug 16 Epub ahead of print; Aigner, A. (2006) *J. Biomed. Biotechnol.* 71659), Arg-Gly-Asp (RGD) peptides (Liu, S. (2006) *Mol. Pharm.* 3:472-487), and polyamidoamines (Tomalia, DA, *et al* (2007) *Biochem. Soc. Trans.* 35:61-67; Yoo, H., *et al* (1999) *Pharm. Res.* 16:1799-1804). In some embodiments, an iRNA forms a complex with cyclodextrin for systemic administration. Methods for administration and pharmaceutical

30 compositions of iRNAs and cyclodextrins can be found in U.S. Patent No. 7,427,605, which is herein incorporated by reference in its entirety.

A. Vector encoded iRNAs of the Invention

iRNA targeting the TMPRSS6 gene can be expressed from transcription units inserted into

35 DNA or RNA vectors (see, *e.g.*, Couture, A, *et al.*, *TIG.* (1996), 12:5-10; Skillern, A, *et al.*, International PCT Publication No. WO 00/22113, Conrad, International PCT Publication No. WO 00/22114, and Conrad, U.S. Patent No. 6,054,299). Expression can be transient (on the order of hours to weeks) or sustained (weeks to months or longer), depending upon the specific construct used and the target tissue or cell type. These transgenes can be introduced as a linear construct, a circular

plasmid, or a viral vector, which can be an integrating or non-integrating vector. The transgene can also be constructed to permit it to be inherited as an extrachromosomal plasmid (Gassmann, *et al.*, *Proc. Natl. Acad. Sci. USA* (1995) 92:1292).

Viral vector systems which can be utilized with the methods and compositions described herein include, but are not limited to, (a) adenovirus vectors; (b) retrovirus vectors, including but not limited to lentiviral vectors, moloney murine leukemia virus, *etc.*; (c) adeno-associated virus vectors; (d) herpes simplex virus vectors; (e) SV 40 vectors; (f) polyoma virus vectors; (g) papilloma virus vectors; (h) picornavirus vectors; (i) pox virus vectors such as an orthopox, *e.g.*, vaccinia virus vectors or avipox, *e.g.* canary pox or fowl pox; and (j) a helper-dependent or gutless adenovirus. Replication-defective viruses can also be advantageous. Different vectors will or will not become incorporated into the cells' genome. The constructs can include viral sequences for transfection, if desired. Alternatively, the construct can be incorporated into vectors capable of episomal replication, *e.g.* EPV and EBV vectors. Constructs for the recombinant expression of an iRNA will generally require regulatory elements, *e.g.*, promoters, enhancers, *etc.*, to ensure the expression of the iRNA in target cells. Other aspects to consider for vectors and constructs are known in the art.

V. Pharmaceutical Compositions of the Invention

The present invention also includes pharmaceutical compositions and formulations which include the iRNAs of the invention. In one embodiment, provided herein are pharmaceutical compositions containing an iRNA, as described herein, and a pharmaceutically acceptable carrier. The pharmaceutical compositions containing the iRNA are useful for preventing or treating a TMPRSS6-associated disorder, *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis. Such pharmaceutical compositions are formulated based on the mode of delivery. One example is compositions that are formulated for systemic administration *via* parenteral delivery, *e.g.*, by subcutaneous (SC), intramuscular (IM), or intravenous (IV) delivery. The pharmaceutical compositions of the invention may be administered in dosages sufficient to inhibit expression of a TMPRSS6 gene.

In some embodiments, the pharmaceutical compositions of the invention are sterile. In another embodiment, the pharmaceutical compositions of the invention are pyrogen free.

The pharmaceutical compositions of the invention may be administered in dosages sufficient to inhibit expression of a TMPRSS6 gene. In general, a suitable dose of an iRNA of the invention will be in the range of about 0.001 to about 200.0 milligrams per kilogram body weight of the recipient per day, generally in the range of about 1 to 50 mg per kilogram body weight per day. Typically, a suitable dose of an iRNA of the invention will be in the range of about 0.1 mg/kg to about 5.0 mg/kg, such as, about 0.3 mg/kg and about 3.0 mg/kg. A repeat-dose regimen may include administration of a therapeutic amount of iRNA on a regular basis, such as every month, once every 3-6 months, or once a year. In certain embodiments, the iRNA is administered about once per month to about once per six months.

After an initial treatment regimen, the treatments can be administered on a less frequent basis. Duration of treatment can be determined based on the severity of disease.

In other embodiments, a single dose of the pharmaceutical compositions can be long lasting, such that doses are administered at not more than 1, 2, 3, or 4 month intervals. In some embodiments of the invention, a single dose of the pharmaceutical compositions of the invention is administered about once per month. In other embodiments of the invention, a single dose of the pharmaceutical compositions of the invention is administered quarterly (*i.e.*, about every three months). In other embodiments of the invention, a single dose of the pharmaceutical compositions of the invention is administered twice per year (*i.e.*, about once every six months).

The skilled artisan will appreciate that certain factors can influence the dosage and timing required to effectively treat a subject, including but not limited to mutations present in the subject, previous treatments, the general health or age of the subject, and other diseases present. Moreover, treatment of a subject with a prophylactically or therapeutically effective amount, as appropriate, of a composition can include a single treatment or a series of treatments.

The iRNA can be delivered in a manner to target a particular tissue (*e.g.*, hepatocytes).

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions can be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids, and self-emulsifying semisolids. Formulations include those that target the liver.

The pharmaceutical formulations of the present invention, which can conveniently be presented in unit dosage form, can be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers.

A. Additional Formulations

i. Emulsions

The compositions of the present invention can be prepared and formulated as emulsions.

Emulsions are typically heterogeneous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μm in diameter (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, L.V., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi *et al.*, in Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa., 1985, p. 301). Emulsions are often biphasic systems comprising two immiscible liquid phases intimately mixed and dispersed

with each other. In general, emulsions can be of either the water-in-oil (w/o) or the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase, the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase, the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions can contain additional components in addition to the dispersed phases, and the active drug which can be present as a solution either in the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants can also be present in emulsions as needed.

Pharmaceutical emulsions can also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous phase provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Other means of stabilizing emulsions entail the use of emulsifiers that can be incorporated into either phase of the emulsion. Emulsifiers can broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Rieger, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants can be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic, and amphoteric (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Rieger, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives, and antioxidants (Block, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

The application of emulsion formulations *via* dermatological, oral, and parenteral routes, and methods for their manufacture have been reviewed in the literature (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

ii. Microemulsions

In one embodiment of the present invention, the compositions of iRNAs and nucleic acids are formulated as microemulsions. A microemulsion can be defined as a system of water, oil, and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Rosoff, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: Controlled Release of Drugs: Polymers and Aggregate Systems, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215).

iii. Microparticles

An iRNA of the invention may be incorporated into a particle, *e.g.*, a microparticle. Microparticles can be produced by spray-drying, but may also be produced by other methods including lyophilization, evaporation, fluid bed drying, vacuum drying, or a combination of these techniques.

iv. Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly iRNAs, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs can cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to

aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers can be classified as belonging to one of five broad categories, *i.e.*, surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (see *e.g.*,

- 5 Malmsten, M. Surfactants and polymers in drug delivery, Informa Health Care, New York, NY, 2002; Lee *et al.*, Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p.92). Each of the above mentioned classes of penetration enhancers and their use in manufacture of pharmaceutical compositions and delivery of pharmaceutical agents are well known in the art.

10 *v. Excipients*

- In contrast to a carrier compound, a “pharmaceutical carrier” or “excipient” is a pharmaceutically acceptable solvent, suspending agent, or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient can be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Such agent are well known in the art.
- 15

vi. Other Components

- The compositions of the present invention can additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions can contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or can contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, *e.g.*, lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings, or aromatic substances, and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.
- 20
- 25
- 30

Aqueous suspensions can contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol, or dextran. The suspension can also contain stabilizers.

- In some embodiments, pharmaceutical compositions featured in the invention include (a) one or more iRNA and (b) one or more agents which function by a non-iRNA mechanism and which are useful in treating a Tmprss63-associated disorder, *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera, myelodysplastic syndrome,
- 35

congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic porphyria, Parkinson's Disease, Alzheimer's Disease or Friedreich's Ataxia.

Toxicity and prophylactic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose prophylactically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds that exhibit high therapeutic indices are preferred.

The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of compositions featured herein in the invention lies generally within a range of circulating concentrations that include the ED50, such as, an ED80 or ED90, with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the methods featured in the invention, the prophylactically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range of the compound or, when appropriate, of the polypeptide product of a target sequence (*e.g.*, achieving a decreased concentration of the polypeptide) that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) or higher levels of inhibition as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography.

In addition to their administration, as discussed above, the iRNAs featured in the invention can be administered in combination with other known agents used for the prevention or treatment of a TMPRSS6-associated disorder, *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis. In any event, the administering physician can adjust the amount and timing of iRNA administration on the basis of results observed using standard measures of efficacy known in the art or described herein.

VI. Methods For Inhibiting TMPRSS6 Expression

The present invention also provides methods of inhibiting expression of a TMPRSS6 gene in a cell. The methods include contacting a cell with an RNAi agent, *e.g.*, double stranded RNA agent, in an amount effective to inhibit expression of TMPRSS6 in the cell, thereby inhibiting expression of TMPRSS6 in the cell.

Contacting of a cell with an iRNA, *e.g.*, a double stranded RNA agent, may be done *in vitro* or *in vivo*. Contacting a cell *in vivo* with the iRNA includes contacting a cell or group of cells within a subject, *e.g.*, a human subject, with the iRNA. Combinations of *in vitro* and *in vivo* methods of contacting a cell are also possible. Contacting a cell may be direct or indirect, as discussed above. Furthermore, contacting a cell may be accomplished *via* a targeting ligand, including any ligand

described herein or known in the art. In some embodiments, the targeting ligand is a carbohydrate moiety, *e.g.*, a GalNAc₃ ligand, or any other ligand that directs the RNAi agent to a site of interest.

The term “inhibiting,” as used herein, is used interchangeably with “reducing,” “silencing,” “downregulating,” “suppressing”, and other similar terms, and includes any level of inhibition.

5 The phrase “inhibiting expression of a TMPRSS6” is intended to refer to inhibition of expression of any TMPRSS6 gene (such as, *e.g.*, a mouse TMPRSS6 gene, a rat TMPRSS6 gene, a monkey TMPRSS6 gene, or a human TMPRSS6 gene) as well as variants or mutants of a TMPRSS6 gene. Thus, the TMPRSS6 gene may be a wild-type TMPRSS6 gene, a mutant TMPRSS6 gene, or a transgenic TMPRSS6 gene in the context of a genetically manipulated cell, group of cells, or

10 organism.

“Inhibiting expression of a TMPRSS6 gene” includes any level of inhibition of a TMPRSS6 gene, *e.g.*, at least partial suppression of the expression of a TMPRSS6 gene. The expression of the TMPRSS6 gene may be assessed based on the level, or the change in the level, of any variable associated with TMPRSS6 gene expression, *e.g.*, TMPRSS6 mRNA level or TMPRSS6 protein level.

15 The expression of a TMPRSS6 may also be assessed indirectly based on the hepcidin mRNA level, hepcidin protein level, or iron levels in tissues or serum. This level may be assessed in an individual cell or in a group of cells, including, for example, a sample derived from a subject. It is understood that TMPRSS6 is expressed predominantly in the liver.

Inhibition may be assessed by a decrease in an absolute or relative level of one or more
20 variables that are associated with TMPRSS6 expression compared with a control level. The control level may be any type of control level that is utilized in the art, *e.g.*, a pre-dose baseline level, or a level determined from a similar subject, cell, or sample that is untreated or treated with a control (such as, *e.g.*, buffer only control or inactive agent control).

In some embodiments of the methods of the invention, expression of a TMPRSS6 gene is
25 inhibited by at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%, or to below the level of detection of the assay. In some embodiments, expression of a TMPRSS6 gene is inhibited by at least 70%. It is further understood that inhibition of TMPRSS6 expression in certain tissues, *e.g.*, in liver, without a significant inhibition of expression in other tissues, *e.g.*, brain, may be desirable. In some embodiments, expression level is determined using the assay method provided in Example 2
30 with a 10 nM siRNA concentration in the appropriate species matched cell line.

In certain embodiments, inhibition of expression *in vivo* is determined by knockdown of the human gene in a rodent expressing the human gene, *e.g.*, an AAV-infected mouse expressing the human target gene (*i.e.*, TMPRSS6), *e.g.*, when administered as a single dose, *e.g.*, at 3 mg/kg at the nadir of RNA expression. Knockdown of expression of an endogenous gene in a model animal
35 system can also be determined, *e.g.*, after administration of a single dose at, *e.g.*, 3 mg/kg at the nadir of RNA expression. Such systems are useful when the nucleic acid sequence of the human gene and the model animal gene are sufficiently close such that the human iRNA provides effective knockdown of the model animal gene. RNA expression in liver is determined using the PCR methods provided in Example 2.

Inhibition of the expression of a TMPRSS6 gene may be manifested by a reduction of the amount of mRNA expressed by a first cell or group of cells (such cells may be present, for example, in a sample derived from a subject) in which a TMPRSS6 gene is transcribed and which has or have been treated (*e.g.*, by contacting the cell or cells with an iRNA of the invention, or by administering an iRNA of the invention to a subject in which the cells are or were present) such that the expression of a TMPRSS6 gene is inhibited, as compared to a second cell or group of cells substantially identical to the first cell or group of cells but which has not or have not been so treated (control cell(s) not treated with an iRNA or not treated with an iRNA targeted to the gene of interest). In some embodiments, the inhibition is assessed by the method provided in Example 2 using a 10nM siRNA concentration in the species matched cell line and expressing the level of mRNA in treated cells as a percentage of the level of mRNA in control cells, using the following formula:

$$\frac{(\text{mRNA in control cells}) - (\text{mRNA in treated cells})}{(\text{mRNA in control cells})} \bullet 100\%$$

In other embodiments, inhibition of the expression of a TMPRSS6 gene may be assessed in terms of a reduction of a parameter that is functionally linked to TMPRSS6 gene expression, *e.g.*, TMPRSS6 protein level in blood or serum from a subject. TMPRSS6 gene silencing may be determined in any cell expressing TMPRSS6, either endogenous or heterologous from an expression construct, and by any assay known in the art.

Inhibition of the expression of a TMPRSS6 protein may be manifested by a reduction in the level of the TMPRSS6 protein that is expressed by a cell or group of cells or in a subject sample (*e.g.*, the level of protein in a blood sample derived from a subject). As explained above, for the assessment of mRNA suppression, the inhibition of protein expression levels in a treated cell or group of cells may similarly be expressed as a percentage of the level of protein in a control cell or group of cells, or the change in the level of protein in a subject sample, *e.g.*, blood or serum derived therefrom.

A control cell, a group of cells, or subject sample that may be used to assess the inhibition of the expression of a TMPRSS6 gene includes a cell, group of cells, or subject sample that has not yet been contacted with an RNAi agent of the invention. For example, the control cell, group of cells, or subject sample may be derived from an individual subject (*e.g.*, a human or animal subject) prior to treatment of the subject with an RNAi agent or an appropriately matched population control.

The level of TMPRSS6 mRNA that is expressed by a cell or group of cells may be determined using any method known in the art for assessing mRNA expression. In one embodiment, the level of expression of TMPRSS6 in a sample is determined by detecting a transcribed polynucleotide, or portion thereof, *e.g.*, mRNA of the TMPRSS6 gene. RNA may be extracted from cells using RNA extraction techniques including, for example, using acid phenol/guanidine isothiocyanate extraction (RNAzol B; Biogenesis), RNeasyTM RNA preparation kits (Qiagen®) or PAXgeneTM (PreAnalytixTM, Switzerland). Typical assay formats utilizing ribonucleic acid hybridization include nuclear run-on assays, RT-PCR, RNase protection assays, northern blotting, *in situ* hybridization, and microarray analysis.

In some embodiments, the level of expression of TMPRSS6 is determined using a nucleic acid probe. The term “probe”, as used herein, refers to any molecule that is capable of selectively binding to a specific TMPRSS6. Probes can be synthesized by one of skill in the art, or derived from appropriate biological preparations. Probes may be specifically designed to be labeled. Examples of molecules that can be utilized as probes include, but are not limited to, RNA, DNA, proteins, antibodies, and organic molecules.

Isolated mRNA can be used in hybridization or amplification assays that include, but are not limited to, Southern or northern analyses, polymerase chain reaction (PCR) analyses and probe arrays. One method for the determination of mRNA levels involves contacting the isolated mRNA with a nucleic acid molecule (probe) that can hybridize to TMPRSS6 mRNA. In one embodiment, the mRNA is immobilized on a solid surface and contacted with a probe, for example by running the isolated mRNA on an agarose gel and transferring the mRNA from the gel to a membrane, such as nitrocellulose. In an alternative embodiment, the probe(s) are immobilized on a solid surface and the mRNA is contacted with the probe(s), for example, in an Affymetrix® gene chip array. A skilled artisan can readily adapt known mRNA detection methods for use in determining the level of TMPRSS6 mRNA.

An alternative method for determining the level of expression of TMPRSS6 in a sample involves the process of nucleic acid amplification or reverse transcriptase (to prepare cDNA) of for example mRNA in the sample, *e.g.*, by RT-PCR (the experimental embodiment set forth in Mullis, 1987, U.S. Patent No. 4,683,202), ligase chain reaction (Barany (1991) *Proc. Natl. Acad. Sci. USA* 88:189-193), self sustained sequence replication (Guatelli *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi *et al.* (1988) *Bio/Technology* 6:1197), rolling circle replication (Lizardi *et al.*, U.S. Patent No. 5,854,033) or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers. In particular aspects of the invention, the level of expression of TMPRSS6 is determined by quantitative fluorogenic RT-PCR (*i.e.*, the TaqMan™ System). In some embodiments, expression level is determined by the method provided in Example 2 using, *e.g.*, a 10 nM siRNA concentration, in the species matched cell line.

The expression levels of TMPRSS6 mRNA may be monitored using a membrane blot (such as used in hybridization analysis such as northern, Southern, dot, and the like), or microwells, sample tubes, gels, beads or fibers (or any solid support comprising bound nucleic acids). See U.S. Patent Nos. 5,770,722, 5,874,219, 5,744,305, 5,677,195 and 5,445,934, which are incorporated herein by reference. The determination of TMPRSS6 expression level may also comprise using nucleic acid probes in solution.

In some embodiments, the level of mRNA expression is assessed using branched DNA (bDNA) assays or real time PCR (qPCR). The use of these methods is described and exemplified in

the Examples presented herein. In some embodiments, expression level is determined by the method provided in Example 2 using a 10nM siRNA concentration in the species matched cell line.

The level of TMPRSS6 protein expression may be determined using any method known in the art for the measurement of protein levels. Such methods include, for example, electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), hyperdiffusion chromatography, fluid or gel precipitin reactions, absorption spectroscopy, a colorimetric assays, spectrophotometric assays, flow cytometry, immunodiffusion (single or double), immunoelectrophoresis, western blotting, radioimmunoassay (RIA), enzyme-linked immunosorbent assays (ELISAs), immunofluorescent assays, electrochemiluminescence assays, and the like.

In some embodiments, the efficacy of the methods of the invention are assessed by a decrease in TMPRSS6 mRNA or protein level (*e.g.*, in a liver biopsy).

In some embodiments of the methods of the invention, the iRNA is administered to a subject such that the iRNA is delivered to a specific site within the subject. The inhibition of expression of TMPRSS6 may be assessed using measurements of the level or change in the level of TMPRSS6 mRNA or TMPRSS6 protein in a sample derived from fluid or tissue from the specific site within the subject (*e.g.*, liver or blood).

As used herein, the terms detecting or determining a level of an analyte are understood to mean performing the steps to determine if a material, *e.g.*, protein, RNA, is present. As used herein, methods of detecting or determining include detection or determination of an analyte level that is below the level of detection for the method used.

VII. Prophylactic and Treatment Methods of the Invention

The present invention also provides methods of using an iRNA of the invention or a composition containing an iRNA of the invention to inhibit expression of TMPRSS6, thereby preventing or treating a TMPRSS6-associated disorder, *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis. In the methods of the invention the cell may be contacted with the siRNA *in vitro* or *in vivo*, *i.e.*, the cell may be within a subject.

A cell suitable for treatment using the methods of the invention may be any cell that expresses a TMPRSS6 gene, *e.g.*, a liver cell. A cell suitable for use in the methods of the invention may be a mammalian cell, *e.g.*, a primate cell (such as a human cell, including human cell in a chimeric non-human animal, or a non-human primate cell, *e.g.*, a monkey cell or a chimpanzee cell), or a non-primate cell. In certain embodiments, the cell is a human cell, *e.g.*, a human liver cell. In the methods of the invention, TMPRSS6 expression is inhibited in the cell by at least 50, 55, 60, 65, 70, 75, 80, 85, 90, or 95, or to a level below the level of detection of the assay.

The *in vivo* methods of the invention may include administering to a subject a composition containing an iRNA, where the iRNA includes a nucleotide sequence that is complementary to at least a part of an RNA transcript of the TMPRSS6 gene of the mammal to which the RNAi agent is to be administered. The composition can be administered by any means known in the art including, but not

limited to oral, intraperitoneal, or parenteral routes, including intracranial (*e.g.*, intraventricular, intraparenchymal, and intrathecal), intravenous, intramuscular, subcutaneous, transdermal, airway (aerosol), nasal, rectal, and topical (including buccal and sublingual) administration. In certain
5 embodiments, the compositions are administered by intravenous infusion or injection. In certain
embodiments, the compositions are administered by subcutaneous injection. In certain embodiments,
the compositions are administered by intramuscular injection.

In one aspect, the present invention also provides methods for inhibiting the expression of a
TMPRSS6 gene in a mammal. The methods include administering to the mammal a composition
comprising a dsRNA that targets a TMPRSS6 gene in a cell of the mammal and maintaining the
10 mammal for a time sufficient to obtain degradation of the mRNA transcript of the TMPRSS6 gene,
thereby inhibiting expression of the TMPRSS6 gene in the cell. Reduction in gene expression can be
assessed by any methods known in the art and by methods, *e.g.* qRT-PCR, described herein, *e.g.*, in
Example 2. Reduction in protein production can be assessed by any methods known in the art, *e.g.*
ELISA. In certain embodiments, a puncture liver biopsy sample serves as the tissue material for
15 monitoring the reduction in the TMPRSS6 gene or protein expression. In other embodiments, a blood
sample serves as the subject sample for monitoring the reduction in the TMPRSS6 protein expression.

The present invention further provides methods of treatment in a subject in need thereof, *e.g.*,
a subject diagnosed with a TMPRSS6-associated disorder, such as a disorder associated with iron
overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -
20 thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera,
myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency,
erythropoietic porphyria, Parkinson's Disease, Alzheimer's Disease or Friedreich's Ataxia. In one
embodiment, a subject having a TMPRSS6-associated disorder has hereditary hemochromatosis. In
another embodiment, a subject having a TMPRSS6-associated disorder has β -thalassemia. In another
25 embodiment, a subject having a TMPRSS6-associated disorder has polycythemia vera.

The present invention further provides methods of prophylaxis in a subject in need thereof.
The treatment methods of the invention include administering an iRNA of the invention to a subject,
e.g., a subject that would benefit from a reduction of TMPRSS6 expression, in a prophylactically
effective amount of a dsRNA targeting a TMPRSS6 gene or a pharmaceutical composition
30 comprising a dsRNA targeting a TMPRSS6 gene.

In one aspect, the present invention provides methods of treating a subject having a disorder that
would benefit from reduction in TMPRSS6 expression, *e.g.*, a TMPRSS6-associated disease, such as a
disorder associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary
hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia
35 vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency,
erythropoietic porphyria, Parkinson's Disease, Alzheimer's Disease or Friedreich's Ataxia. Treatment of a
subject that would benefit from a reduction and/or inhibition of TMPRSS6 gene expression includes
therapeutic treatment (*e.g.*, a subject is having elevated iron levels) and prophylactic treatment (*e.g.*, the
subject is not having elevated iron levels or a subject may be at risk of developing elevated iron levels).

An iRNA of the invention may be administered as a “free iRNA.” A free iRNA is administered in the absence of a pharmaceutical composition. The naked iRNA may be in a suitable buffer solution. The buffer solution may comprise acetate, citrate, prolamine, carbonate, or phosphate, or any combination thereof. In one embodiment, the buffer solution is phosphate buffered saline (PBS). The pH and osmolarity of the buffer solution containing the iRNA can be adjusted such that it is suitable for administering to a subject.

Alternatively, an iRNA of the invention may be administered as a pharmaceutical composition, such as a dsRNA liposomal formulation.

Subjects that would benefit from an inhibition of TMPRSS6 gene expression are subjects susceptible to or diagnosed with a TMPRSS6-associated disorder, such as a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic porphyria, Parkinson’s Disease, Alzheimer’s Disease or Friedreich’s Ataxia. In an embodiment, the method includes administering a composition featured herein such that expression of the target a TMPRSS6 gene is decreased, such as for about 1, 2, 3, 4, 5, 6, 1-6, 1-3, or 3-6 months per dose. In certain embodiments, the composition is administered once every 3-6 months.

In one embodiment, the iRNAs useful for the methods and compositions featured herein specifically target RNAs (primary or processed) of the target TMPRSS6 gene. Compositions and methods for inhibiting the expression of these genes using iRNAs can be prepared and performed as described herein.

Administration of the iRNA according to the methods of the invention may result prevention or treatment of a TMPRSS6-associated disorder, *e.g.*, a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis, *e.g.*, hereditary hemochromatosis, β -thalassemia (*e.g.*, β -thalassemia major and β -thalassemia intermedia), polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic porphyria, Parkinson’s Disease, Alzheimer’s Disease or Friedreich’s Ataxia. Subjects can be administered a therapeutic amount of iRNA, such as about 0.01 mg/kg to about 200 mg/kg.

In one embodiment, the iRNA is administered subcutaneously, *i.e.*, by subcutaneous injection. In another embodiment, the iRNA is administered intravenously, *i.e.*, by intravenous injection. One or more injections may be used to deliver the desired dose of iRNA to a subject. The injections may be repeated over a period of time.

The administration may be repeated on a regular basis. In certain embodiments, after an initial treatment regimen, the treatments can be administered on a less frequent basis. A repeat-dose regimen may include administration of a therapeutic amount of iRNA on a regular basis, such as once per month to once a year. In certain embodiments, the iRNA is administered about once per month to about once every three months, or about once every three months to about once every six months.

The invention further provides methods and uses of an iRNA agent or a pharmaceutical composition thereof for treating a subject that would benefit from reduction and/or inhibition of

TMPRSS6 gene expression, *e.g.*, a subject having a TMPRSS6-associated disease, in combination with other pharmaceuticals and/or other therapeutic methods, *e.g.*, with known pharmaceuticals and/or known therapeutic methods, such as, for example, those which are currently employed for treating these disorders.

- 5 Accordingly, in some aspects of the invention, the methods which include either a single iRNA agent of the invention, further include administering to the subject one or more additional therapeutic agents.

- For example, in certain embodiments, an iRNA targeting TMPRSS6 is administered in combination with, *e.g.*, an agent useful in treating a disorder associated with iron overload. For
10 example, additional agents suitable for treating a subject that would benefit from reduction in TMPRSS6 expression, *e.g.*, a subject having a disorder associated with iron overload, may include iron chelators (*e.g.*, desferoxamine), folic acid, a blood transfusion, a phlebotomy, agents to manage ulcers, agents to increase fetal hemoglobin levels (*e.g.*, hydroxyurea), agents to control infection (*e.g.*, antibiotics and antivirals), agents to treat thrombotic state, or a stem cell or bone marrow transplant. A
15 stem cell transplant can utilize stem cells from an umbilical cord, such as from a relative, *e.g.*, a sibling. Exemplary iron chelators include desferoxamine, Deferasirox (Exjade), deferiprone, vitamin E, wheat germ oil, tocophersolan, and indicaxanthin.

- The iRNA agent and an additional therapeutic agent and/or treatment may be administered at the same time and/or in the same combination, *e.g.*, parenterally, or the additional therapeutic agent can be
20 administered as part of a separate composition or at separate times and/or by another method known in the art or described herein.

VIII. Kits

- In certain aspects, the instant disclosure provides kits that include a suitable container
25 containing a pharmaceutical formulation of a siRNA compound, *e.g.*, a double-stranded siRNA compound, or siRNA compound, (*e.g.*, a precursor, *e.g.*, a larger siRNA compound which can be processed into a siRNA compound, or a DNA which encodes an siRNA compound, *e.g.*, a double-stranded siRNA compound, or ssiRNA compound, or precursor thereof).

- Such kits include one or more dsRNA agent(s) and instructions for use, *e.g.*, instructions for
30 administering a prophylactically or therapeutically effective amount of a dsRNA agent(s). The dsRNA agent may be in a vial or a pre-filled syringe. The kits may optionally further comprise means for administering the dsRNA agent (*e.g.*, an injection device, such as a pre-filled syringe), or means for measuring the inhibition of TMPRSS6 (*e.g.*, means for measuring the inhibition of TMPRSS6 mRNA, TMPRSS6 protein, and/or TMPRSS6 activity). Such means for measuring the inhibition of
35 TMPRSS6 may comprise a means for obtaining a sample from a subject, such as, *e.g.*, a plasma sample. The kits of the invention may optionally further comprise means for determining the therapeutically effective or prophylactically effective amount.

 In certain embodiments the individual components of the pharmaceutical formulation may be provided in one container, *e.g.*, a vial or a pre-filled syringe. Alternatively, it may be desirable to

provide the components of the pharmaceutical formulation separately in two or more containers, *e.g.*, one container for a siRNA compound preparation, and at least another for a carrier compound. The kit may be packaged in a number of different configurations such as one or more containers in a single box. The different components can be combined, *e.g.*, according to instructions provided with the kit.

- 5 The components can be combined according to a method described herein, *e.g.*, to prepare and administer a pharmaceutical composition. The kit can also include a delivery device.

10 This invention is further illustrated by the following examples which should not be construed as limiting. The entire contents of all references, patents and published patent applications cited throughout this application, as well as the informal Sequence Listing and Figures, are hereby incorporated herein by reference.

15 EXAMPLES

Example 1. iRNA Synthesis

Source of reagents

Where the source of a reagent is not specifically given herein, such reagent can be obtained from any supplier of reagents for molecular biology at a quality/purity standard for application in
20 molecular biology.

siRNA Design

siRNAs targeting the human Transmembrane protease, serine 6 (TMPRSS6) gene (human: NCBI refseqID NM_153609.4, NCBI GeneID: 164656) were designed using custom R and Python
25 scripts. The human NM_153609.4 REFSEQ mRNA, has a length of 3197 bases.

Detailed lists of the unmodified TMPRSS6 sense and antisense strand nucleotide sequences are shown in Tables 2, 4 and 6. Detailed lists of the modified TMPRSS6 sense and antisense strand nucleotide sequences are shown in Tables 3, 5 and 7.

It is to be understood that, throughout the application, a duplex name without a decimal is
30 equivalent to a duplex name with a decimal which merely references the batch number of the duplex. For example, AD-959917 is equivalent to AD-959917.1.

siRNA Synthesis

siRNAs were designed, synthesized, and prepared using methods known in the art.
35 Briefly, siRNA sequences were synthesized on a 1 μ mol scale using a Mermade 192 synthesizer (BioAutomation) with phosphoramidite chemistry on solid supports. The solid support was controlled pore glass (500-1000 Å) loaded with a custom GalNAc ligand (3'-GalNAc conjugates), universal solid support (AM Chemicals), or the first nucleotide of interest. Ancillary synthesis reagents and standard 2-cyanoethyl phosphoramidite monomers (2'-deoxy-2'-fluoro, 2'-O-

methyl, RNA, DNA) were obtained from Thermo-Fisher (Milwaukee, WI), Hongene (China), or Chemgenes (Wilmington, MA, USA). Additional phosphoramidite monomers were procured from commercial suppliers, prepared in-house, or procured using custom synthesis from various CMOs. Phosphoramidites were prepared at a concentration of 100 mM in either acetonitrile or 9:1

5 acetonitrile:DMF and were coupled using 5-Ethylthio-1H-tetrazole (ETT, 0.25 M in acetonitrile) with a reaction time of 400 s. Phosphorothioate linkages were generated using a 100 mM solution of 3-((Dimethylamino-methylidene) amino)-3H-1,2,4-dithiazole-3-thione (DDTT, obtained from Chemgenes (Wilmington, MA, USA)) in anhydrous acetonitrile/pyridine (9:1 v/v). Oxidation time was 5 minutes. All sequences were synthesized with final removal of the DMT group ("DMT-Off").

10 Upon completion of the solid phase synthesis, solid-supported oligoribonucleotides were treated with 300 μ L of Methylamine (40% aqueous) at room temperature in 96 well plates for approximately 2 hours to afford cleavage from the solid support and subsequent removal of all additional base-labile protecting groups. For sequences containing any natural ribonucleotide linkages (2'-OH) protected with a tert-butyl dimethyl silyl (TBDMS) group, a second deprotection step was

15 performed using TEA.3HF (triethylamine trihydrofluoride). To each oligonucleotide solution in aqueous methylamine was added 200 μ L of dimethyl sulfoxide (DMSO) and 300 μ L TEA.3HF and the solution was incubated for approximately 30 mins at 60 °C. After incubation, the plate was allowed to come to room temperature and crude oligonucleotides were precipitated by the addition of 1 mL of 9:1 acetonitrile:ethanol or 1:1 ethanol:isopropanol. The plates were then centrifuged at 4 °C

20 for 45 mins and the supernatant carefully decanted with the aid of a multichannel pipette. The oligonucleotide pellet was resuspended in 20 mM NaOAc and subsequently desalted using a HiTrap size exclusion column (5 mL, GE Healthcare) on an Agilent LC system equipped with an autosampler, UV detector, conductivity meter, and fraction collector. Desalted samples were collected in 96 well plates and then analyzed by LC-MS and UV spectrometry to confirm identity and quantify

25 the amount of material, respectively.

Duplexing of single strands was performed on a Tecan liquid handling robot. Sense and antisense single strands were combined in an equimolar ratio to a final concentration of 10 μ M in 1x PBS in 96 well plates, the plate sealed, incubated at 100 °C for 10 minutes, and subsequently allowed to return slowly to room temperature over a period of 2-3 hours. The concentration and identity of

30 each duplex was confirmed and then subsequently utilized for in vitro screening assays.

Example 2. *In vitro* screening methods

Cell culture and 384-well transfections

For transfections, Hep3b cells (ATCC, Manassas, VA) were grown to near confluence at

35 37°C in an atmosphere of 5% CO₂ in Eagle's Minimum Essential Medium (Gibco) supplemented with 10% FBS (ATCC) before being released from the plate by trypsinization. Transfection was carried out by adding 7.5 μ L of Opti-MEM plus 0.1 μ L of Lipofectamine RNAiMax per well (Invitrogen, Carlsbad CA. cat # 13778-150) to 2.5 μ L of each siRNA duplex to an individual well in a 384-well plate. The mixture was then incubated at room temperature for 15 minutes. Forty μ L of complete

growth media without antibiotic containing $\sim 1.5 \times 10^4$ cells were then added to the siRNA mixture. Cells were incubated for 24 hours prior to RNA purification. Single dose experiments were performed at 10 nM, 1 nM, and 0.1 nM final duplex concentration.

5 Total RNA isolation using DYNABEADS mRNA Isolation Kit (Invitrogen™, part #: 610-12)

Cells were lysed in 75µl of Lysis/Binding Buffer containing 3 µL of beads per well and mixed for 10 minutes on an electrostatic shaker. The washing steps were automated on a Biotek EL406, using a magnetic plate support. Beads were washed (in 90µL) once in Buffer A, once in Buffer B, and twice in Buffer E, with aspiration steps in between. Following a final aspiration,

10 complete 10µL RT mixture was added to each well, as described below.

cDNA synthesis using ABI High capacity cDNA reverse transcription kit (Applied Biosystems, Foster City, CA, Cat #4368813)

A master mix of 1µl 10X Buffer, 0.4µl 25X dNTPs, 1µl Random primers, 0.5µl Reverse Transcriptase, 0.5µl RNase inhibitor and 6.6µl of H₂O per reaction were added per well. Plates were sealed, agitated for 10 minutes on an electrostatic shaker, and then incubated at 37 degrees C for 2 hours. Following this, the plates were agitated at 80 degrees C for 8 minutes.


Real time PCR




20 Two microlitre (µl) of cDNA were added to a master mix containing 0.5µl of human GAPDH TaqMan Probe (4326317E), 0.5µl human TMPRSS6, 2µl nuclease-free water and 5µl Lightcycler 480 probe master mix (Roche Cat # 04887301001) per well in a 384 well plates (Roche cat # 04887301001). Real time PCR was done in a LightCycler480 Real Time PCR system (Roche).

To calculate relative fold change, data were analyzed using the $\Delta\Delta C_t$ method and normalized to assays performed with cells transfected with 10nM AD-1955, or mock transfected cells. IC₅₀s were calculated using a 4 parameter fit model using XLFit and normalized to cells transfected with AD-1955 or mock-transfected. The sense and antisense sequences of AD-1955 are: sense: cuuAcGcuGAGuAcuucGAdTsdT (SEQ ID NO: 18) and antisense UCGAAGuACUcAGCGuAAGdTsdT (SEQ ID NO: 19).

30 The results of the single dose screen of the agents in Tables 2, 3, 6 and 7 in Hep3b cells are shown in Table 8.

Table 1. Abbreviations of nucleotide monomers used in nucleic acid sequence representation. It will be understood that these monomers, when present in an oligonucleotide, are mutually linked by 5'-3'-phosphodiester bonds; and it is understood that when the nucleotide contains a 2'-fluoro modification, then the fluoro replaces the hydroxy at that position in the parent nucleotide (*i.e.*, it is a 2'-deoxy-2'-fluoronucleotide).

Abbreviation	Nucleotide(s)
A	Adenosine-3'-phosphate
Ab	beta-L-adenosine-3'-phosphate
Abs	beta-L-adenosine-3'-phosphorothioate
Af	2'-fluoroadenosine-3'-phosphate
Afs	2'-fluoroadenosine-3'-phosphorothioate
As	adenosine-3'-phosphorothioate
C	cytidine-3'-phosphate
Cb	beta-L-cytidine-3'-phosphate
Cbs	beta-L-cytidine-3'-phosphorothioate
Cf	2'-fluorocytidine-3'-phosphate
Cfs	2'-fluorocytidine-3'-phosphorothioate
Cs	cytidine-3'-phosphorothioate
G	guanosine-3'-phosphate
Gb	beta-L-guanosine-3'-phosphate
Gbs	beta-L-guanosine-3'-phosphorothioate
Gf	2'-fluoroguanosine-3'-phosphate
Gfs	2'-fluoroguanosine-3'-phosphorothioate
Gs	guanosine-3'-phosphorothioate
T	5'-methyluridine-3'-phosphate
Tf	2'-fluoro-5-methyluridine-3'-phosphate
Tfs	2'-fluoro-5-methyluridine-3'-phosphorothioate
Ts	5-methyluridine-3'-phosphorothioate
U	Uridine-3'-phosphate
Uf	2'-fluorouridine-3'-phosphate
Ufs	2'-fluorouridine-3'-phosphorothioate
Us	uridine-3'-phosphorothioate
N	any nucleotide, modified or unmodified
a	2'-O-methyladenosine-3'-phosphate
as	2'-O-methyladenosine-3'-phosphorothioate
c	2'-O-methylcytidine-3'-phosphate
cs	2'-O-methylcytidine-3'-phosphorothioate
g	2'-O-methylguanosine-3'-phosphate
gs	2'-O-methylguanosine-3'-phosphorothioate
t	2'-O-methyl-5-methyluridine-3'-phosphate
ts	2'-O-methyl-5-methyluridine-3'-phosphorothioate
u	2'-O-methyluridine-3'-phosphate
us	2'-O-methyluridine-3'-phosphorothioate
s	phosphorothioate linkage
L10	N-(cholesterylcarboxamidocaproyl)-4-hydroxyprolinol (Hyp-C6-Chol)
L96	<p>N-[tris(GalNAc-alkyl)-amidodecanoyl]-4-hydroxyprolinol (Hyp-(GalNAc-alkyl)₃)</p> 

Abbreviation	Nucleotide(s)
Y34	2-hydroxymethyl-tetrahydrofuran-4-methoxy-3-phosphate (abasic 2'-OMe furanose) 
Y44	inverted abasic DNA (2-hydroxymethyl-tetrahydrofuran-5-phosphate) 
L10	N-(cholesterylcarboxamidocaproyl)-4-hydroxyprolinol (Hyp-C6-Chol) 
(Agn)	Adenosine-glycol nucleic acid (GNA) S-Isomer
(Cgn)	Cytidine-glycol nucleic acid (GNA) S-Isomer
(Ggn)	Guanosine-glycol nucleic acid (GNA) S-Isomer
(Tgn)	Thymidine-glycol nucleic acid (GNA) S-Isomer
P	Phosphate
VP	Vinyl-phosphonate
dA	2'-deoxyadenosine-3'-phosphate
dAs	2'-deoxyadenosine-3'-phosphorothioate
dC	2'-deoxycytidine-3'-phosphate
dCs	2'-deoxycytidine-3'-phosphorothioate
dG	2'-deoxyguanosine-3'-phosphate
dGs	2'-deoxyguanosine-3'-phosphorothioate
dT	2'-deoxythymidine-3'-phosphate
dTs	2'-deoxythymidine-3'-phosphorothioate
dU	2'-deoxyuridine
dUs	2'-deoxyuridine-3'-phosphorothioate
(C2p)	cytidine-2'-phosphate
(G2p)	guanosine-2'-phosphate
(U2p)	uridine-2'-phosphate
(A2p)	adenosine-2'-phosphate
(Chd)	2'-O-hexadecyl-cytidine-3'-phosphate
(Ahd)	2'-O-hexadecyl-adenosine-3'-phosphate
(Ghd)	2'-O-hexadecyl-guanosine-3'-phosphate
(Uhd)	2'-O-hexadecyl-uridine-3'-phosphate


Abbreviation	Nucleotide(s)
Q191s	<p data-bbox="483 277 1313 338">N-[tris(GalNAc-alkyl)-amidododecanoyl]-(S)-pyrrolidin-3-ol-phosphorothioate (p-C12-(GalNAc-alkyl)3)</p> 

Table 2. Unmodified Sense and Antisense Strand Sequences of TMPRSS6 dsRNA Agents

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1554875	GCCUGUGAGGACUCCCAAGAGU	20	232-252	ACUCTUGGAGUCCUCACAGGCCU	146	230-252
AD-1554909	GGUGCUACUCUGGUUUUUCU	21	324-344	AGGAAATACCAAGAGUAGCACCCC	147	322-344
AD-1554910	GUGCUACUCUGGUUUUUCU	22	325-345	AAGGAAUACCAAGAGUAGCACCCC	148	323-345
AD-1554911	UGCUACUCUGGUUUUUCU	23	326-346	ATAGGAAUACCAAGAGUAGCAC	149	324-346
AD-1554912	GCUACUCUGGUUUUUCU	24	327-347	ACUAGGAAUACCAAGAGUAGCAC	150	325-347
AD-1554913	CUACUCUGGUUUUUCU	25	328-348	ACCUAGGAAUACCAAGAGUAGCA	151	326-348
AD-1554914	UACUCUGGUUUUUCU	26	329-349	ACCCTAGGAAUACCAAGAGUAGC	152	327-349
AD-1554915	ACUCUGGUUUUUCU	27	330-350	AACCCUAGGAAUACCAAGAGUAG	153	328-350
AD-1554916	CUCUGGUUUUUCU	28	331-351	ATACCTAGGAAUACCAAGAGUA	154	329-351
AD-1554917	UCUGGUUUUUCU	29	332-352	AGUACCCUAGGAAUACCAAGAGU	155	330-352
AD-1554923	AUUUCCUAGGGUACAAGGCGU	30	338-358	ACGCCUTGUACCCUAGGAAUAC	156	336-358
AD-1554951	GGUCAGCCAGGUGUACUCAGU	31	366-386	ACUGAGTACACCUUGGUGACCAU	157	364-386
AD-1554955	AGCCAGGUGUACUCAGGCGU	32	370-390	ACUGCCTGAGUACACCUUGGUGA	158	368-390
AD-1554992	GCCACUUCUCCAGGAUCUUU	33	407-427	AAAGAUCCUGGGAGAAUGGGCGA	159	405-427
AD-1554997	UUCUCCAGGAUCUUACCCGU	34	412-432	ACGGGUAAGAUCCUGGGAGAAAGU	160	410-432
AD-1555000	UCCAGGAUCUUACCCGCCGU	35	415-435	ACGGCGGUAAGAUCCUGGGAGA	161	413-435
AD-1555030	GCCUCCCGCAGUGAAACCGCU	36	445-465	AGCGGUTUCAC'TGCGGAAGGCAC	162	443-465
AD-1555106	CAACUCCAGCUCGUCUUAUUU	37	522-542	AAAUAGACGGAGCUGGAGUUUGUA	163	520-542
AD-1555112	CAGCUCGUCUUAUUCUUUGU	38	528-548	ACAAAGGAAUAGACGGAGCUGGA	164	526-548
AD-1555114	CUCACCUUGCUUUCUGGUUUU	39	559-579	AAACCAGAAAGAACGAGGUGAGGG	165	557-579
AD-1555115	UCACCUUGCUUUCUGGUUUU	40	560-580	AGAACAGAAAGAACGAGGUGAGG	166	558-580
AD-1555117	ACCUGCUUUCUUGGUUCAUU	41	562-582	AAUGAACAGAAAGAACGAGGUGA	167	560-582
AD-1555118	CCUGCUUUCUUGGUUCAUUU	42	563-583	AAAUAGAACAGAAAGAACGAGGUG	168	561-583
AD-1555120	UGCUUUCUUGGUUCAUUU	43	565-585	AAGAAUGAACAGAAAGAACGAGG	169	563-585
AD-1555121	GCUUCUUCUGGUUCAUUUCU	44	566-586	AGAGAAUGAACAGAAAGAACGAG	170	564-586
AD-1555122	CUUCUUCUGGUUCAUUUCU	45	567-587	AGGAGAAUGAACAGAAAGAACGA	171	565-587

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1555123	UUUUUCUGGUUUAUUCUCCAU	46	568-588	ATGGAGAAUGAAACAGAAAGAGC	172	566-588
AD-1555128	CUGGUUUAUUCUCCAAUCCU	47	573-593	AGGATUTGGAGAAUGAACAGAA	173	571-593
AD-1555184	ACAGGGCCGAGUACGAAAGUGU	48	689-709	ACACTUCGUACTCGGCCCUAG	174	687-709
AD-1555185	CAGGGCCGAGUACGAAAGUGGU	49	690-710	ACCACUTCGUACUCGGCCCUGUA	175	688-710
AD-1555212	CCAGUGUGAAAGACAUAGCUU	50	737-757	AAGCTATGUCUTUCACACUGGCU	176	735-757
AD-1555213	CAGUGUGAAAGACAUAGCUGU	51	738-758	ACAGCUAUGUCTUUCACACUGGC	177	736-758
AD-1555234	AUUGAAUCCACGCUGGGUUU	52	759-779	AAACCCAGCGUGGAAUUCAAUGC	178	757-779
AD-1555235	UUUGAAUCCACGCUGGGUUGU	53	760-780	ACAACCCAGCGTGGAAUUCAAUG	179	758-780
AD-1555236	UGAAUCCACGCUGGGUUGUU	54	761-781	AACAACCCAGCGUGGAAUUCAAU	180	759-781
AD-1555238	AAUCCACGCUGGGUUGUUUAU	55	763-783	ATAACAAACCCAGCGUGGAAUUC	181	761-783
AD-1555241	UCCACGCUGGGUUGUUACCGU	56	766-786	ACGGTAACAACCCAGCGUGGAAU	182	764-786
AD-1555242	CCACGCUGGGUUGUUACCGCU	57	767-787	AGCGGUAACAACCCAGCGUGGAA	183	765-787
AD-1555243	CACGCUGGGUUGUUACCGCUU	58	768-788	AAGCGTTAACAACCCAGCGUGGA	184	766-788
AD-1555247	CUGGGUUGUUACCGCUACAGU	59	772-792	ACUGTAGCGGUAAACAACCCAGCG	185	770-792
AD-1555342	GGGACCGACUGGCCAUGUAUU	60	923-943	AAUACATGGCCAGUCGGUCCCGG	186	921-943
AD-1555343	GGACCGACUGGCCAUGUAUGU	61	924-944	ACAUACAUUGGCCAGUCGGUCCCG	187	922-944
AD-1555345	ACCGACUGGCCAUGUAUGACU	62	926-946	AGUCAUACAUUGGCCAGUCGGUCC	188	924-946
AD-1555346	CCGACUGGCCAUGUAUGACGU	63	927-947	ACGUCATACAUUGGCCAGUCGGUCC	189	925-947
AD-1555348	GACUGGCCAUGUAUGACGUGU	64	929-949	ACACGUCAUACAUUGGCCAGUCGG	190	927-949
AD-1555349	ACUGGCCAUGUAUGACGUGGU	65	930-950	ACCACGTCAUACAUUGGCCAGUCCG	191	928-950
AD-1555350	CUGGCCAUGUAUGACGUGGCU	66	931-951	AGCCACGUCAUACAUUGGCCAGUC	192	929-951
AD-1555366	AGGCUCAUACACCUCCGUGUAU	67	967-987	ATACACCGAGGTGAUGAGCCUCU	193	965-987
AD-1555428	GCCUGCACAGCUACUACGACU	68	1061-1081	AGUCGUAGUAGCUGUGCAGGCC	194	1059-1081
AD-1555429	CCUGCACAGCUACUACGACCU	69	1062-1082	AGGUCGTAGUAGCUGUGCAGGCC	195	1060-1082
AD-1555535	CCUCUCUGGACUACGGCCUUGU	70	1235-1255	ACAAGCCGUAGTCCAGAGAGGGC	196	1233-1255
AD-1555537	UCUCUGGACUACGGCCUUGGCU	71	1237-1257	AGCCAAGCCGUAGUCCAGAGAGG	197	1235-1257
AD-1555546	UACGGCUUGGCCCUUGGCUUU	72	1246-1266	AAACAGAGGGCCAAGCCGUAGU	198	1244-1266

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1555547	ACGGCUUGGCCCCUCUGGUUUU	73	1247-1267	AAAAACAGAGGGCCAAAGCCGUAG	199	1245-1267
AD-1555548	CGGCUUGGCCCCUCUGGUUUU	74	1248-1268	ACAAACAGAGGGCCAAAGCCGUA	200	1246-1268
AD-1555549	GGCUUGGCCCCUCUGGUUUU	75	1249-1269	ATCAAACAGAGGGCCAAAGCCGU	201	1247-1269
AD-1555581	GAGGAGGCAGAAAGUAUGAUUU	76	1281-1301	AAAUACATACUUCUGCCUCCAG	202	1279-1301
AD-1555583	GGAGGCAGAAAGUAUGAUUUU	77	1283-1303	ACAAAUCAUACTUCUGCCUCCUC	203	1281-1303
AD-1555584	GAGGCAGAAAGUAUGAUUUU	78	1284-1304	AGCAATCAUACUUCUGCCUCCU	204	1282-1304
AD-1555585	AGGCAGAAAGUAUGAUUUU	79	1285-1305	AGGCAAAUCAUACUUCUGCCUCC	205	1283-1305
AD-1555586	GGCAGAAAGUAUGAUUUU	80	1286-1306	ACGGCAAAUCATACUUCUGCCUC	206	1284-1306
AD-1555587	GCAGAAAGUAUGAUUUU	81	1287-1307	AACGGCAAAUCAUACUUCUGCCU	207	1285-1307
AD-1555588	CAGAAAGUAUGAUUUU	82	1288-1308	ACACGGCAAAUCAUACUUCUGCC	208	1286-1308
AD-1555589	AGAAAGUAUGAUUUU	83	1289-1309	AGCACGGCAAAATCAUACUUCUGC	209	1287-1309
AD-1555590	GAAGUAUGAUUUU	84	1290-1310	ATGCACGGCAAAUCAUACUUCUG	210	1288-1310
AD-1555615	CAGUGGACGAUCCAGAACAGU	85	1318-1338	ACUGTUCUGGATCGUCCACUGGC	211	1316-1338
AD-1555616	AGUGGACGAUCCAGAACAGGU	86	1319-1339	ACCUGTUCUGGAGUCCACUGG	212	1317-1339
AD-1555626	CCAGAAACAGGAGGCUGUGUGU	87	1329-1349	ACACACAGCCUCCUGUUCUGGAU	213	1327-1349
AD-1555628	AGAAACAGGAGGCUGUGUGGU	88	1331-1351	AGCCACACAGCCUCCUGUUCUGG	214	1329-1351
AD-1555706	UGUGCGGGUGCACUAUGGCUU	89	1449-1469	AAGCCATAGUGCACCCGCACACC	215	1447-1469
AD-1555707	GUUGCGGGUGCACUAUGGCUUU	90	1450-1470	AAAGCCAUAGUGCACCCGCACAC	216	1448-1470
AD-1555709	GCGGGUGCACUAUGGCUUUU	91	1452-1472	AACAAGCCAUAGUGCACCCGCAC	217	1450-1472
AD-1555711	GGGUGCACUAUGGCUUUU	92	1454-1474	AGUACAAGCCATAGUGCACCCGC	218	1452-1474
AD-1555717	ACUAUGGCUUGUACAACACAGU	93	1460-1480	ACUGGUTGUACAAGCCAUAGUGC	219	1458-1480
AD-1555723	GCUUGUACAACAGUCCGACU	94	1466-1486	AGUCCGACUGGTUGUACAAGCCA	220	1464-1486
AD-1555725	CUGCCUUGGAGAGUUCUCUU	95	1488-1508	AAGAGGAACUCTCCAGGGCAGGG	221	1486-1508
AD-1555768	GCCUGGAUGAGAGAAACUGCU	96	1565-1585	AGCAGUTUCUCTCAUCCAGGCCG	222	1563-1585
AD-1555771	UGGAUGAGAGAAACUGCGUUU	97	1568-1588	AAACGCAGUUUCUCUCAUCCAGG	223	1566-1588
AD-1555772	GGAUGAGAGAAACUGCGUUUU	98	1569-1589	AAAACGCAGUUTCUUCAUCCAG	224	1567-1589
AD-1555776	GAGAGAAACUGCGUUU	99	1573-1593	ACUGCAACGCAGUUUCUCUCAU	225	1571-1593

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1555789	UUUGCAGAGCCACAUUCCAGU	100	1586-1606	ACUGGAAUUGGUCUCUGCAAACG	226	1584-1606
AD-1555894	GUGGGACAUUACCUUCCAGU	101	1709-1729	ACUGGAAAGGUAAUGUCCCAU	227	1707-1729
AD-1555895	UGGGACAUUACCUUCCAGUU	102	1710-1730	AACUGGAAAGGUAAUGUCCACA	228	1708-1730
AD-1555897	GGACAUUACCUUCCAGUGUU	103	1712-1732	AACACUGGAAAGGUAAUGUCCCA	229	1710-1732
AD-1555898	GACAUUACCUUCCAGUGUGU	104	1713-1733	ACACACTGGAAGGUAAUGUCC	230	1711-1733
AD-1555899	ACAUUACCUUCCAGUGUGAU	105	1714-1734	ATCACACUGGAAAGGUAAUGUCC	231	1712-1734
AD-1555900	CAUUCACCUUCCAGUGUGAGU	106	1715-1735	ACUCACACUGGAAAGGUAAUGUC	232	1713-1735
AD-1556052	AUCGCUGACCGCUGGGUGAUU	107	1936-1956	AAUACCCACGCGGUCAGCGAUGA	233	1934-1956
AD-1556057	UGACCGCUGGGUGUAUACAGU	108	1941-1961	ACUGTUAUCACCCAGCGGUCAGC	234	1939-1961
AD-1556126	CGUGUUCUUGGGCAAGGUGUU	109	2010-2030	AACACCTUGCCCAAGGAACACGGU	235	2008-2030
AD-1556127	GUGUUCUUGGGCAAGGUGUGU	110	2011-2031	ACACACCUUGCCCAAGGAACACGG	236	2009-2031
AD-1556137	GCAAGGUGUGGCAGAACUCGU	111	2021-2041	ACGAGUTCUGCCACACCUUGCCC	237	2019-2041
AD-1556139	AAGGUGUGGCAGAACUCGCGU	112	2023-2043	ACGCGAGUUCUGCCACACCUUGC	238	2021-2043
AD-1556163	CUGGAGAGGUGUCCUUCAGU	113	2048-2068	ACUUGAAGGACACCUCCAGGC	239	2046-2068
AD-1556164	UGGAGAGGUGUCCUUCAGGU	114	2049-2069	ACCUUGAAGGACACCUCCAGG	240	2047-2069
AD-1556166	GAGAGGUGUCCUUCAGGUGU	115	2051-2071	ACACCUUGAAGGACACCUCCUCCA	241	2049-2071
AD-1556167	AGAGGUGUCCUUCAGGUGAU	116	2052-2072	ATCACCTUGAAGGACACCUCCUCC	242	2050-2072
AD-1556319	AUCCACACAGGACCUUGGCAGU	117	2299-2319	ACUGCACAGGUCCUGUGGGAUCA	243	2297-2319
AD-1556359	UGACGCCACGCAUGCUGUGUU	118	2339-2359	AACACAGCAUGCUGGCGGUCACC	244	2337-2359
AD-1556360	GACGCCACGCAUGCUGUGUGU	119	2340-2360	ACACACAGCAUGCUGGCGGUCAC	245	2338-2360
AD-1556382	GCUACCCGCAAGGGCAAGAAGU	120	2363-2383	ACUUCUTGCCCCUUGCGGUAGCCG	246	2361-2383
AD-1556383	CUACCCGCAAGGGCAAGAAGGU	121	2364-2384	ACCUUUTGCCCCUUGCGGUAGCC	247	2362-2384
AD-1556465	GGCCUAAACUACUUCGGCGUCU	122	2483-2503	AGACGCCGAAGTAGUAGGCCGG	248	2481-2503
AD-1556466	GCCUAAACUACUUCGGCGUCUU	123	2484-2504	AAGACGCCGAAGUAGUAGGCCG	249	2482-2504
AD-1556484	CUACACCCGCAUACAGGUGU	124	2502-2522	ACACCUUGGAGUCCGGGUGUAGAC	250	2500-2522
AD-1556510	GCUGGAUCCAGCAAGUGGUGU	125	2528-2548	ACACCACUUGCTGGAUCCAGCUG	251	2526-2548
AD-1556584	UGGCAGGAGGUGGCAUCUUGU	126	2670-2690	ACAAGATGCCACCUCCUGCCACC	252	2668-2690

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO.	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO.	Range in NM_153609.4
AD-1556585	GGCAGGAGGUGGCAUCUUGUU	127	2671-2691	AACAAAGUAGCCACCUCCUGCCAC	253	2669-2691
AD-1556586	GCAGGAGGUGGCAUCUUGUCU	128	2672-2692	AGACAAGUAGCCACCUCCUGCCA	254	2670-2692
AD-1556587	CAGGAGGUGGCAUCUUGUCUU	129	2673-2693	AAGACAAGUAGCCACCUCCUGCC	255	2671-2693
AD-1556613	UGAUGUCUGUCCAGUGAUGU	130	2699-2719	ACAUCACUGGAGCAGACAUACAGG	256	2697-2719
AD-1556677	CAAUUCUCUCUCCUCCGUCCU	131	2801-2821	AGGACGGAGGAGAGAAUUGGG	257	2799-2821
AD-1556709	GGCUCAGCAGCAAGAAUGCUU	132	2853-2873	AAGCAUTCUUGCUGGAGGCCAC	258	2851-2873
AD-1556710	GCUCAGCAGCAAGAAUGCUGU	133	2854-2874	ACAGCATUCUUGCUGGAGGCCA	259	2852-2874
AD-1556789	CUGGUCUAAACUUGGGAUCUGU	134	2973-2993	ACAGAUCCCAAGUAGACCAGGG	260	2971-2993
AD-1556790	UGGUCUAAACUUGGGAUCUGGU	135	2974-2994	ACCAGATCCCAAGUAGACCAGG	261	2972-2994
AD-1556791	GGUCUAAACUUGGGAUCUGGGU	136	2975-2995	ACCCAGAUCCTCAAGUAGACCAG	262	2973-2995
AD-1556795	UAAACUUGGGAUCUGGGAUUGU	137	2979-2999	ACAUTCCAGATCCCAAGUUAAGA	263	2977-2999
AD-1556799	UUUGGAUCUGGGAUUGGAAGU	138	2983-3003	ACUUCCAUCCCAAGAUCCCAAGU	264	2981-3003
AD-1556802	GGAUUCUGGGAUUGGAAGGUGU	139	2986-3006	ACACCTCCCAUCCCAAGAUCCCA	265	2984-3006
AD-1556908	UGAGCUCAGCUGCCCUUUGGU	140	3158-3178	ACCAAAGGCGAGCUGAGCUCACC	266	3156-3178
AD-1556909	GAGCUCAGCUGCCCUUUGGAU	141	3159-3179	ATCCAAAGGCGAGCUGAGCUCAC	267	3157-3179
AD-1556911	GCUCAGCUGCCCUUUGGAU	142	3161-3181	AAUCCAAAGGCGAGCUGAGCUC	268	3159-3181
AD-1556915	AGCUGCCCUUUGGAUAAAGU	143	3165-3185	ACUUTATUCCAAAGGCGAGCUGA	269	3163-3185
AD-1556917	CUGCCCUUUGGAUAAAGCUU	144	3167-3187	AAGCTUTAUUCCAAAGGCGAGCU	270	3165-3187
AD-1556918	UGCCCUUUGGAUAAAGCUGU	145	3168-3188	ACAGCUTAUUCCAAAGGCGAGC	271	3166-3188

Table 3. Modified Sense and Antisense Strand Sequences of TMRSS6 dsRNA Agents

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO.	Antisense Sequence 5' to 3'	SEQ ID NO.	mRNA target sequence 5' to 3'	SEQ ID NO.
AD-1554875	gscscugugaGfGfAfuccaagaguL96	272	asdCsuedTudGgagudCcUfcaaggcsesu	398	AGCCUGUGAGGACUCCAAAGAGA	524
AD-1554909	gsgsugcuacUfCfUfgguuuuuccuL96	273	asdGsgadAadTaccadGaGfuagcaccscsc	399	GGGGUGCUACUCUGGUUUUCCU	525
AD-1554910	gsusguacuCfUfGfguuuuuuccuL96	274	asdAsggdAadAuaacdAgAfguagcaccsc	400	GGGUGCUACUCUGGUUUUCCUA	526
AD-1554911	usgsuacuUfGfGfuuuuuuuccuL96	275	asdTsagdGadAuaacdCaGfuagcaccsc	401	GGUGCUACUCUGGUUUUCCUAG	527

AD-1554912	gscsuacucufGfGfUfUfaunucucaguuL96	276	asdCsuaadGgdAaauadCcAfagagagcsase	402	GUGCUACUCUGUGUAUUUCCUAGG	528
AD-1554913	csusacucufGfUfAfucucucaguuL96	277	asdCsuaadGgdAaauadCcAfagagagcsase	403	UGCUACUCUGUGUAUUUCCUAGGG	529
AD-1554914	usascucufGfUfUfucucucaguuL96	278	asdCsuaadGgdAaauadCcAfagagagcsase	404	GCUACUCUGUGUAUUUCCUAGGGU	530
AD-1554915	ascscucufGfUfUfucucucaguuL96	279	asdCsuaadGgdAaauadCcAfagagagcsase	405	CUACUCUGUGUAUUUCCUAGGGUA	531
AD-1554916	csuscucufGfUfUfucucucaguuL96	280	asdCsuaadGgdAaauadCcAfagagagcsase	406	UACUCUGUGUAUUUCCUAGGGUAC	532
AD-1554917	uscsucufGfUfUfucucucaguuL96	281	asdCsuaadGgdAaauadCcAfagagagcsase	407	ACUCUGUGUAUUUCCUAGGGUACA	533
AD-1554923	asusucufGfUfUfucucucaguuL96	282	asdCsuaadGgdAaauadCcAfagagagcsase	408	GUAUUCCUAGGGUACAAGGGCGG	534
AD-1554951	gsgsucufGfUfUfucucucaguuL96	283	asdCsuaadGgdAaauadCcAfagagagcsase	409	AUGGUCAGCCAGGUGUACUCAGG	535
AD-1554955	asgscucufGfUfUfucucucaguuL96	284	asdCsuaadGgdAaauadCcAfagagagcsase	410	UCAGCCAGGUGUACUCAGGCGAGU	536
AD-1554992	gscscucufGfUfUfucucucaguuL96	285	asdCsuaadGgdAaauadCcAfagagagcsase	411	UCGCCACUUCUCCAGGUAUCUUA	537
AD-1554997	ususcucufGfUfUfucucucaguuL96	286	asdCsuaadGgdAaauadCcAfagagagcsase	412	ACUUCUCCAGGUAUCUUAACCCGC	538
AD-1555000	uscsucufGfUfUfucucucaguuL96	287	asdCsuaadGgdAaauadCcAfagagagcsase	413	UCUCCAGGUAUCUUAACCCGCCGG	539
AD-1555030	gscscucufGfUfUfucucucaguuL96	288	asdCsuaadGgdAaauadCcAfagagagcsase	414	GUGCCUUCGCGAGUGAAACCGCC	540
AD-1555106	csasucufGfUfUfucucucaguuL96	289	asdCsuaadGgdAaauadCcAfagagagcsase	415	UACAAUCUCCAGGUAUCUUAUUC	541
AD-1555112	csasucufGfUfUfucucucaguuL96	290	asdCsuaadGgdAaauadCcAfagagagcsase	416	UCCAGCUCUCCAGGUAUCUUAUUC	542
AD-1555114	csuscucufGfUfUfucucucaguuL96	291	asdCsuaadGgdAaauadCcAfagagagcsase	417	CCCUCACUCCAGGUAUCUUAUUC	543
AD-1555115	uscsucufGfUfUfucucucaguuL96	292	asdCsuaadGgdAaauadCcAfagagagcsase	418	CCUACUCCAGGUAUCUUAUUCUUA	544
AD-1555117	ascscucufGfUfUfucucucaguuL96	293	asdCsuaadGgdAaauadCcAfagagagcsase	419	UCACUCCAGGUAUCUUAUUCUUA	545
AD-1555118	csuscucufGfUfUfucucucaguuL96	294	asdCsuaadGgdAaauadCcAfagagagcsase	420	CACCUCUCCAGGUAUCUUAUUCUUA	546
AD-1555120	usgscucufGfUfUfucucucaguuL96	295	asdCsuaadGgdAaauadCcAfagagagcsase	421	CCUGCUUCCAGGUAUCUUAUUCUUA	547
AD-1555121	gscscucufGfUfUfucucucaguuL96	296	asdCsuaadGgdAaauadCcAfagagagcsase	422	CUGCUUCCAGGUAUCUUAUUCUUA	548
AD-1555122	csusucufGfUfUfucucucaguuL96	297	asdCsuaadGgdAaauadCcAfagagagcsase	423	UGCUUCCAGGUAUCUUAUUCUUA	549
AD-1555123	ususcucufGfUfUfucucucaguuL96	298	asdCsuaadGgdAaauadCcAfagagagcsase	424	GCUUCCAGGUAUCUUAUUCUUA	550
AD-1555128	csusucufGfUfUfucucucaguuL96	299	asdCsuaadGgdAaauadCcAfagagagcsase	425	UUCUGGUAUCUUAUUCUUAUUCUUA	551
AD-1555184	ascscucufGfUfUfucucucaguuL96	300	asdCsuaadGgdAaauadCcAfagagagcsase	426	CUACAGGCGGAGUAUCUUAUUCUUA	552
AD-1555185	csasucufGfUfUfucucucaguuL96	301	asdCsuaadGgdAaauadCcAfagagagcsase	427	UACAGGCGGAGUAUCUUAUUCUUA	553
AD-1555212	csuscucufGfUfUfucucucaguuL96	302	asdCsuaadGgdAaauadCcAfagagagcsase	428	AGCCAGGUGGAAAGACUAUUCUUA	554

AD-1555213	csasuguggaAfAfGfacaugagugul96	303	asdCsagdCudAugncdTuUfcacagsgsc	429	GCCAGUGUGAAAGACAUAGCUGC	555
AD-1555234	asusugauuicFcfAfzgcuiggguuul96	304	asdAsacdCcdAgcgudGgAfauucausgsc	430	GCAUUGAAUUECCACGCUGGGUUG	556
AD-1555235	usugaaauccCfAfCfzgcuiggguuul96	305	asdCsaadCcdCagcgdTgGfauucaasug	431	CAUUGAAUUECCACGCUGGGUUGU	557
AD-1555236	usgsaaauccAfCfCfzgcuiggguuul96	306	asdAscadAcdCcgagdGnGfgaauacasu	432	AUUGAAUUECCACGCUGGGUUGUU	558
AD-1555238	asasuuecacGfCFUfzgguuuguuul96	307	asdTsaadCadAeccadGcGfuggausesa	433	UGAAUUECCACGCUGGGUUGUUA	559
AD-1555241	usescagcuGfGfGfhuuuaacegul96	308	asdCsggdTadAcaacdCcAfzgcuiggasau	434	AUUECCACGCUGGGUUGUUAACCGC	560
AD-1555242	csescagcuGfGfUfhuuuaacegul96	309	asdGsgcdGndAcaacdCcCfzgcuiggasasa	435	UUECCACGCUGGGUUGUUAACCGCU	561
AD-1555243	csascgcuGfUfUfhuuuaacegul96	310	asdAsgcdGgdTaaacdAcCfzgcuiggsgsa	436	UCCACGCUGGGUUGUUAACCGCUA	562
AD-1555247	csusggguugUfUfAfccgcuacagul96	311	asdCsugdTadGcgndAaCfaaccagsgsg	437	CGCUGGGUUGUUAACCGCUACAGC	563
AD-1555342	gsgsagccgaCfUfGfzgcuuuguuul96	312	asdAsuadCadTfgcccdAgUfzgcuiggcssgsg	438	CCGGGACCGACUGGGCCAUGUAUG	564
AD-1555343	gsgsagccgaCfUfGfzgcuuuguuul96	313	asdCsaudAcdAuggcdCaGfugcgucscsgsg	439	CGGGACCGACUGGGCCAUGUAUGA	565
AD-1555345	ascsagcuGfCfCfzgcuuuguuul96	314	asdGsudAcdAcaugdGcCfzgcuiggscsg	440	GGACCGACUGGGCCAUGUAUGACG	566
AD-1555346	csescagcuGfCfAfzgcuuuguuul96	315	asdCsugdCadTcaudGcCfzgcuiggsgscsg	441	GACCGACUGGGCCAUGUAUGACGU	567
AD-1555348	gsasagccgaCfUfGfhuuuaacegul96	316	asdCsacdGndCaucadAaUfzgcuiggsgsgsg	442	CCGACUGGGCCAUGUAUGACGUGG	568
AD-1555349	ascsugccgaCfUfGfhuuuaacegul96	317	asdCsacdCgdTcaudCaUfzgcuiggscsgsg	443	CGACUGGGCCAUGUAUGACGUGGC	569
AD-1555350	csusggcuuGfUfAfzgcuuuguuul96	318	asdGsacdAcdGueaudAcAfugcgcuiggscsg	444	GACUGGGCCAUGUAUGACGUGGCC	570
AD-1555366	asgsagcuuCaCfAfCfzgcuuuguuul96	319	asdTsacdAcdCgagdTgAfugagcuiggscsg	445	AGAGGCUCAUCCACCGCGGUGUAC	571
AD-1555428	gsescugcacAfGfCfhuuuaacegul96	320	asdGsudGndAguagdCuGfugcagcgscsg	446	GGGCUGCACAGCUACUACGACC	572
AD-1555429	csesugcacGfCFUfhuuuaacegul96	321	asdGsugdCgdTaguadGeUfzgcuiggscsgsg	447	GGCCUGCACAGCUACUACGACCC	573
AD-1555535	csesucucugGfAfCfhuuuaacegul96	322	asdCsaadGcdCguagdTcCfagagcgsgsg	448	GCCCUCUCUGGACUACGCGCUUGG	574
AD-1555537	uscsucuggaCfUfAfzgcuuuguuul96	323	asdGsacdAadGcgndAgUfzcagagcgsgsg	449	CCUCUCUGGACUACGCGCUUGGCC	575
AD-1555546	usascggcuuGfGfGfhuuuaacegul96	324	asdAsacdCadGagggdCcAfagcgcuuagsgu	450	ACUACGGCUUGGGCCCUUGGUUU	576
AD-1555547	ascsagcuuGfCfCfzgcuuuguuul96	325	asdAsaadCcdAgaggdGcCfaagcgcuuagsg	451	CUACGGCUUGGGCCCUUGGUUUUG	577
AD-1555548	csagscuuGfCfCfhuuuaacegul96	326	asdCsaadAcdCagagdGcCfcaagcgcuuagsg	452	UACGGCUUGGGCCCUUGGUUUUGA	578
AD-1555549	gsgscuuggcCfCFUfzgcuuuguuul96	327	asdTsacdAadCcaagdGgGfcaagcgcuuagsg	453	ACGGCUUGGGCCCUUGGUUUUGAU	579
AD-1555581	gsasagggcAfGfAfzgcuuuguuul96	328	asdAsaudCadTacuudCuGfzcucucuasg	454	CUGAGGAGGCAGAAGUAUGAUUU	580
AD-1555583	gsgsagggcAfAfGfhuuuaacegul96	329	asdCsaudAadCaucadTuCfugcuuucscsg	455	GAGGAGGCAGAGAUGAUUUUUGC	581

AD-1555584	gsasggcagaAfGfUfauuugcuL96	330	asdGscadAadTcauadCuUfucgucscsu	456	AGGAGGCAGAAAGUAUGAUUUUGCC	582
AD-1555585	asgsgeagaaGfUfAfugauugcuL96	331	asdGsgcdAadAucadAcUfucgucscsc	457	GGAGGCAGAAAGUAUGAUUUUGCCG	583
AD-1555586	gsascagaagUfAfUfauuugccguL96	332	asdCsugdCadAaucadTaCfucgucscsu	458	GAGGCAGAAAGUAUGAUUUUGCCGU	584
AD-1555587	gsascagaaguAfUfGfauuugccguL96	333	asdAsegdGcdAaucadAuAfcuucgucscsu	459	AGGCAGAAAGUAUGAUUUUGCCGUG	585
AD-1555588	csasgaaguaUfGfAfuuugccguL96	334	asdCsacdGgdCaaadCaUfaucucgscsc	460	GGCAGAAAGUAUGAUUUUGCCGUGC	586
AD-1555589	asgsaaguauGfAfUfugccguL96	335	asdGscadCgdGcaadTcAfuucucgsc	461	GCAGAAAGUAUGAUUUUGCCGUGCA	587
AD-1555590	gsasaguauGfUfUfugccguL96	336	asdTsgcdAcDgcaadAuCfuaucucscsu	462	CAGAAAGUAUGAUUUUGCCGUGCAC	588
AD-1555615	csasguugagGfAfUfcaagaacaguL96	337	asdCsugdTuDcugadTcGfucacugsc	463	GCCAGUGGACGAUCCAGAACAGG	589
AD-1555616	asgsuugagGfUfCfcaagaacaguL96	338	asdCsugdTuDcugadAuCfucacugsc	464	CCAGUGGACGAUCCAGAACAGGA	590
AD-1555626	csasgaacaGfGfAfugcugugcuL96	339	asdCsacdAcDAgcudCcUfucucgscsu	465	AUCCAGAACAGGAGGCUUGUGG	591
AD-1555628	asgsaacaggAfGfGfugcugugcuL96	340	asdGscddAcDAcagcdCuCfucucgsc	466	CCAGAACAGGAGGCUUGUGGCU	592
AD-1555706	usgsuugggUfGfCfauuugcuL96	341	asdAsgcdCadTagudCaCfucgacscsc	467	GGUGUGGGUGGACUUAUGGCUU	593
AD-1555707	gsusgcggguGfCfAfauuugcuL96	342	asdAsagdCcdAuagudGcAfucgacscsc	468	GUGUGCGGGUGGACUUAUGGCUU	594
AD-1555709	gsusggguGfCfUfauuugcuL96	343	asdAsagdCcdAuagudGcAfucgacscsc	469	GUGCGGGUGGACUUAUGGCUUUA	595
AD-1555711	gsusguGfUfAfUfugcugugcuL96	344	asdGsuadCadAgcudTaGfucgacscsc	470	GCGGGUGGACUUAUGGCUUUAACA	596
AD-1555717	ascsuugggUfUfGfuaaaccaguL96	345	asdCsugdGudTgucadAaGfucuaugsc	471	GCACUUAUGGCUUUAACAACAGU	597
AD-1555723	gsusuuGfUfAfCfagucgcuL96	346	asdGsuadCgdAcugdTgGfuaaaccscsa	472	UGGCUUUAACAACAGGAGGAC	598
AD-1555725	csusggcuGfUfGfagucgcuL96	347	asdAsagdGgdAcucdTcCfagggcagsg	473	CCCUGCCUGGAGAGUUAUCCUG	599
AD-1555768	gsusggcuGfUfGfagaaacugcuL96	348	asdGscadGudTucudTcAfucagggsc	474	CGGCCUGGAGAGAGAAACUUGCG	600
AD-1555771	usgsuugagAfGfAfauuugcuL96	349	asdAsacdGcdAgucdTgCfuaaaccscsa	475	CCUGGAGAGAGAGAAACUUGCUU	601
AD-1555772	gsusauGfUfAfUfagucgcuL96	350	asdAsacdCgdCagudTcUfucuaaccscsa	476	CUGGAGAGAGAGAAACUUGCUUUG	602
AD-1555776	gsasgagaaCfUfGfagucgcuL96	351	asdCsugdCadAcgudAgUfucucscsa	477	AUGAGAGAAACUGGCUUUGCAGA	603
AD-1555789	usugcagaGfCfCfauuuccaguL96	352	asdCsugdGadAugudGcUfucgaaaccsc	478	CGUUUGCAGAGCCACAUUCCAGU	604
AD-1555894	gsusgggacaUfUfCfauuuccaguL96	353	asdCsugdGadAugudGcUfucgaaaccsc	479	AUGUGGACAUUCCAUUCCAGU	605
AD-1555895	usgsuugcuUfCfAfucucgcuL96	354	asdAsacdGgdAggudGcUfucgaaaccsc	480	UGUGGACAUUCCAUUCCAGUG	606
AD-1555897	gsusauuGfUfCfUfagucgcuL96	355	asdAsacdCudGagudGcUfuaaaccscsa	481	UGGACAUUCCAUUCCAGUGUG	607
AD-1555898	gsasauuGfUfUfagucgcuL96	356	asdCsacdAcDtggaadGgUfuaaaccsc	482	GGGACAUUCCAUUCCAGUGUGA	608

AD-1555899	asasauuacCfUfUfccagugugauL96	357	asdTtsaadCadCuggadAgGfguauguscs	483	GGACAUUCACCUUCCAGUGUGAG	609
AD-1555900	esasuuaceUfUfCfcagugugauL96	358	asdCsuedAcadAucggdAaGfgugaaugus	484	GACAUUCACCUUCCAGUGUGAGG	610
AD-1556052	asuscgcugacCfCfGfcugggugauuL96	359	asdAsuedAcadCagcdGgUfcagcgaugsa	485	UCAUCGCUGACCCGUGGGUGAUA	611
AD-1556057	usgsaccgcuGfGfGfguguaacaguiL96	360	asdCsugdTuadAucacdCcAfgcggucasc	486	GCUGACCCGUGGGGUGUAUAACAGC	612
AD-1556126	csugsugnuceUfGfGfgcaaguguuL96	361	asdAsaadCcdTugccdCaGfgaacacsgsu	487	ACCGUGUUCUUGGGCAAGGUGUG	613
AD-1556127	gsusguuuccGfGfGfcaaguguguiL96	362	asdCsacdAcadCuugcdCcAfggaacacsg	488	CCGUGUUCUUGGGCAAGGUGUGG	614
AD-1556137	gsasaaaggugUfGfGfcaagaacuguiL96	363	asdCsagdGudTugcdCaCfaccuugscsc	489	GGCAAGGUGUGGCGAGAACUCGC	615
AD-1556139	asasggugugGfCfAfgaacucgcuL96	364	asdCsagdGadGuucudGcCfaccuugsc	490	GCAAGGUGUGGCGAGAACUCGCGC	616
AD-1556163	csusggagagGfUfGfuccuacaguiL96	365	asdCsuidGadAggaadAcCfuccuacsgsc	491	GCCUGGAGAGGUGUCCUUAAGG	617
AD-1556164	usgsagagaggUfGfUfuccuacaguiL96	366	asdCsuidTgdAaggadCaCfuccuacsgsg	492	CCUGGAGAGGUGUCCUUAAGGU	618
AD-1556166	gsasagaggugUfCfCfuccaagguguiL96	367	asdCsacdCudTgaagdGaCfaccuucscsa	493	UGGAGAGGUGUCCUUAAGGUGA	619
AD-1556167	asgsagaggugCfCfUfuccaagguguiL96	368	asdTtsaadCcdTugaadGgAfcaccuucsc	494	GGAGAGGUGUCCUUAAGGUGAG	620
AD-1556319	asuscccacaGfGfAfccugugcaguiL96	369	asdCsugdCadCaggudCcUfgugggauusca	495	UGAUCCACACAGGACCCUGUGCAGC	621
AD-1556359	usgsacgccaCfGfCfaguguguguiL96	370	asdAsaadCadGaaludCcUfgcggucasc	496	GGUGACGCCACGCAUGCUGUGUG	622
AD-1556360	gsasgcgcacGfCfAfugcuguguguiL96	371	asdCsacdAcadAgaudGcGfugcgucscsc	497	GUGACGCCACGCAUGCUGUGUGC	623
AD-1556382	gsusuaaccgcAfAfGfGfcaagaaguiL96	372	asdCsuidCudTgcccduGfGfguaugscsg	498	CGGCUACCCGCAAGGGCAAGAAAGG	624
AD-1556383	csusaccgcaAfGfGfGfcaagaaguiL96	373	asdCsuidTcdTugccdCuUfgcgguaugsc	499	GGCUACCCGCAAGGGCAAGAAAGGA	625
AD-1556465	gsusccuaacUfAfCfucggcguguiL96	374	asdGsacdGcdCgaagdTaGfuaagccsgsg	500	CCGGCCUAAACUACUUCGGCGUCU	626
AD-1556466	gsuscuacuuAfCfUfucggcguguiL96	375	asdAsgadCgdCgaadGuAfguuagccscsg	501	CGGCCUAAACUACUUCGGCGUCUA	627
AD-1556484	csusacacceGfCfAfucacaguguiL96	376	asdCsacdCudGugaudGcGfGfguaugsc	502	GUCUACACCCGCAUCACAGGUGU	628
AD-1556510	gsusuggaucCfAfGfcaaguguguiL96	377	asdCsacdCadCuugcdTgGfauccagcsusg	503	CAGCUGGAUCCAGCAAGUGGUGA	629
AD-1556584	usgsagcaggaGfGfUfGfcaucuuL96	378	asdCsaadGadTgcccduUfGfGfcaucsc	504	GGUGGACGAGGUGGCAUCUUGU	630
AD-1556585	gsusagagagGfUfGfGfcaucuuL96	379	asdAsaadAgaugcdAcCfuccuagccsc	505	GUGGCAGGAGGUGGCAUCUUGUC	631
AD-1556586	gsasagagaggUfGfGfcaucuuL96	380	asdGsacdAadGaugcdCaCfuccuagccsc	506	UGGCAGGAGGUGGCAUCUUGUCU	632
AD-1556587	csasggaggugGfGfCfcaucuuL96	381	asdAsaadCadAgaugcdCcAfcuccuagsc	507	GGCAGGAGGUGGCAUCUUGUCUC	633
AD-1556613	usgsauugucuGfCfUfccaaguguiL96	382	asdCsuidCadCuggadGcAfgaacuacsgsg	508	CCUGAUGUCUGUCUCCAGUGAUGG	634
AD-1556677	csasauucueUfCfUfuccuagucuiL96	383	asdGsgadCgdGaggadGaGfagaauugsgsg	509	CCCAAUUCUCUCUCCUCCGUCUCC	635

AD-1556709	gsgscucagc AfGfCfaagaugcuuL96	384	asdAsgedAudTcuugdCuGfcugagccsasc	510	GUGGUCACGACGACAAAGAAUGCUG	636
AD-1556710	gscsucagcaGfCfAfagaugcuL96	385	asdCsagdCadTucnuudGcUfugagccsca	511	UGGUCACGACGACAAAGAAUGCUGG	637
AD-1556789	csusggucaAfCfUfuggauncuL96	386	asdCsagdAudCccaadGuUfagaccagsgsg	512	CCCUGGUCUAAACUUUGGGAUCUCUGG	638
AD-1556790	usgsgucuaaCfUfUfuggauncuL96	387	asdCscadGadTcccadAgUfugagccagsgsg	513	CCUGGUCUAAACUUUGGGAUCUCUGGG	639
AD-1556791	gsgsucuaacUfUfGfggaucugguL96	388	asdCscadAgdAuuccdAaGfuaagaccsag	514	CUGGUCUAAACUUUGGGAUCUCUGGGA	640
AD-1556795	usasacuuGgGfAfUfcuuggaauL96	389	asdCsaudTcdCagadTcCfaaguuasgsa	515	UCUAAACUUUGGGAUCUCUGGGAUUGG	641
AD-1556799	ususgggaucUfGfGfagauggaauL96	390	asdCsnuudCcdAuuccdCaGfauccecaasgsu	516	ACUUGGGAUCUCUGGGAUUGGGAAGG	642
AD-1556802	gsgsaucugGfAfAfugaagguL96	391	asdCsacdCudTccaadTcCfagaucscsa	517	UGGGAUCUCUGGGAUUGGGAAGGUGC	643
AD-1556908	usgsagcucaGfCfUfcccunugguL96	392	asdCscadAadGgagadGcUfagagccasgc	518	GGUGAGCUCAGCUGCCCCUUUGGA	644
AD-1556909	gsasgcucagCfUfGfcccunugguL96	393	asdTscadAadAagggedAgCfugagcucscasc	519	GUGAGCUCAGCUGCCCCUUUGGAA	645
AD-1556911	gscsucagcuGfCfCfuuuggaauL96	394	asdAsnuudCcdAaaggdGcAfagagagcsusc	520	GAGCUCAGCUGCCCCUUUGGAAUA	646
AD-1556915	asgscugeccUfUfUfuggaauaaauL96	395	asdCsnuudTadTuccadAaGfaggagcugsgsa	521	UCAGCUGCCCCUUUGGGAUAAAGC	647
AD-1556917	csusgcccunUfGfGfaauaaagcuL96	396	asdAsgedTudTaanudCaAfaggagcagcsu	522	AGCUGCCCCUUUGGGAUAAAGCUG	648
AD-1556918	usgsccecuuGfGfAfauaagcugL96	397	asdCsagdCudTuanudCcAfaggagcagsgsc	523	GCUGCCCCUUUGGGAUAAAGCUGC	649

Table 4. Unmodified Sense and Antisense Strand Sequences of TMPRSS6 dsRNA Agent

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1557376	CGGAGGUGAUGGCGAGGAAGU	650	189-209	ACUUCCTCGCCCAUCACCCUCCGUC	848	187-209
AD-1557377	GGAGGUGATGGCGAGGAAGCU	651	190-210	AGCUUCCUGCCATCACCUCUGU	849	188-210
AD-1557396	AAGCCUGTGAGGACUCCAAU	652	229-249	ATUGGAGUCCUCACAGGCCUUGA	850	227-249
AD-1557398	GGCCUGAGGACUCCAAAGU	653	231-251	ATCUUGGAGUCCUCACAGGCCUU	851	229-251
AD-1557399	GCCUGAGGACUCCAAAGAU	20	232-252	ACUUGGAGUCCUCCACAGGCCU	852	230-252
AD-1557400	CCUGUAGGACUCCAAAGAU	654	233-253	ATCUCUTGGAGTCCUCACAGGCC	853	231-253
AD-1557401	CUGUAGGACTCCAAGAGAU	655	234-254	ATUCUTUGGAGUCCUCACAGGC	854	232-254
AD-1557437	CUACUCUGUAUUUCCUAGGU	25	328-348	ACCUAGGAAUACCCAGAGUAGCA	151	326-348
AD-1557440	CUCUGUAUTUCCUAGGGU	656	331-351	ATACCCTAGGAAATACCAGAGUA	855	329-351
AD-1557441	UCUGGUAUTUCCUAGGGUACU	657	332-352	AGUACCCUAGGAAUACCCAGAGU	155	330-352
AD-1557442	CUGGUAUUTCCUAGGGUACAU	658	333-353	ATGUACCCUAGGAAUACCCAGAG	856	331-353

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1557443	UGGUUUUCCCTAGGGUACAACU	659	334-354	ATUGUACCCUAGGAAAUACCCAGA	857	332-354
AD-1557444	GGUAAUUCCUAGGGUACAAGU	660	335-355	ACUUGUACCCUAGGAAAUACCCAG	858	333-355
AD-1557445	GUUUUCCCTAGGGUACAAGGU	661	336-356	ACCUUGTACCCCTAGGAAAUACCCA	859	334-356
AD-1557452	CUAGGGUACAAGGCGGAGGUU	662	343-363	AACCUCCGCCCTGTACCCUAGGA	860	341-363
AD-1557473	AUGGUACGCCAGGUGUACUCU	663	364-384	AGAGUACACCUUGGUGACCAUCA	861	362-384
AD-1557475	GGUACGCCAGGUGUACUCAGU	31	366-386	ACUGAGTACACCUUGGUGACCAU	157	364-386
AD-1557476	GUCAGCCAGGTGUACUCAGGU	664	367-387	ACCUGAGUACACCTGGCUGACCA	862	365-387
AD-1557477	UCAGCCAGGUGUACUCAGGCU	665	368-388	AGCCUGAGUACACCUUGGUGAC	863	366-388
AD-1557478	CAGCCAGGTGTACUCAGGCAU	666	369-389	ATGCCUGAGUACACCUUGGUGAC	864	367-389
AD-1557479	AGCCAGGUGUACUCAGGCAGU	32	370-390	ACUGCTGAGUACACCUUGGUGA	158	368-390
AD-1557509	CUCAAUCGCCACUUCUCCCAU	667	400-420	ATGGGAGAAUGGGCAUUGAGUA	865	398-420
AD-1557515	CGCCACUUCUCCAGGAGUUCU	668	406-426	AAGAUCCUGGGAGAGUGGCGAU	866	404-426
AD-1557516	GCCACUUCUCCAGGAGUUCUU	669	407-427	AAAGAUCUUGGGAGAGUGGCGA	159	405-427
AD-1557518	CACUUCUCCAGGAGUUCUACU	670	409-429	AGUAGAUCUUGGGAGAGUGGC	867	407-429
AD-1557522	UCUCCAGGATCUUACCCGCU	671	413-433	AGCGGTAAAGATCCUGGGAGAG	868	411-433
AD-1557523	CUCCAGGAGUUCUACCCGCCU	672	414-434	AGCGGGUAAAGAUCCUGGGAGAA	869	412-434
AD-1557524	UCCAGGATCTUACCCGCCU	673	415-435	ACGGCGGUAAAGATCCUGGGAGA	870	413-435
AD-1557550	UAGUGCCUUCGCGAGUGAAAU	674	441-461	ATUUCACUGCGGAAGGCACUAGA	871	439-461
AD-1557554	GCCUCCGCGAGUGAAACCGCU	36	445-465	AGCGGUTUACCTGCGGAAGGCAC	162	443-465
AD-1557555	CCUUCGCGAGTGAACCGGCCU	675	446-466	AGCGGTUUCACUGCGGAAGGCA	872	444-466
AD-1557556	CUUCGCGAGUGAAACCGCCAU	676	447-467	ATGGCGGUUUCACCTGCGGAAGGC	873	445-467
AD-1557559	CCGCGAGUGAAACCGCCAAAGU	677	450-470	ACUUUGGCGGUTUACUGCGGAA	874	448-470
AD-1557560	CGCAGUGAAACCGCCAAAGCU	678	451-471	AGCUUUGGCGGUTUACUGCGGGA	875	449-471
AD-1557561	GCAGUGAAACCGCCAAAGCCU	679	452-472	AGGCUUTGGCGGUTUACUGCGG	876	450-472
AD-1557562	CAGUGAAACCGCCAAAGCCCU	680	453-473	AGGCGUTUGGCGGUTUACUGCG	877	451-473
AD-1557563	AGUGAAACCGCCAAAGCCCAU	681	454-474	ATGGGCTUUGGCGGUTUACUGC	878	452-474
AD-1557571	CGCCAAAGCCAGAGAGUUCU	682	462-482	AGCAUUCUUGGCGUUGGCGGU	879	460-482
AD-1557572	GCCAAAGCCAGAGAGUUCUU	683	463-483	AAGCAUUCUUGGCGUUGGCGG	880	461-483
AD-1557577	AGCCAGAGAGAGUUCUACAGU	684	468-488	ACCUAGAGCAUCUUCUGGCGUU	881	466-488
AD-1557606	CAGCACCCGCCUGGGAACUUU	685	498-518	AAAGUCCCAAGCGGUGGUGGU	882	496-518
AD-1557607	AGCACCCGCTGGGAACUUU	686	499-519	ATAAGUTCCCAAGCGGUGGUGG	883	497-519
AD-1557629	ACAACUCCAGUCCGUCUUAUU	687	521-541	AAUAGACGGAGUGGAGUUGAG	884	519-541
AD-1557630	CAACUCCAGCTCCGUCUUAUU	688	522-542	AAUAGACGGAGCTGGAGUUGA	885	520-542
AD-1557639	UCACCUGCTUCUUCUGGUUCU	689	560-580	AGAACCAAGAAAGCAGGUGAGG	166	558-580
AD-1557640	CACCUGCUTCTUCUGGUUCAU	690	561-581	ATGAACCAAGAAAGCAGGUGAG	886	559-581

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1557642	CCUGCUUUCTUCUGGUUUAUUU	691	563-583	AAUUAACCAAGAAAGCAGGUG	168	561-583
AD-1557643	CUGCUUCUTCTGGUUAUUUCU	692	564-584	AGAAUUAACCAAGAAAGCAGGU	887	562-584
AD-1557644	UGCUUCUUCUGGUUAUUUCU	43	565-585	AAGAAUUAACCAAGAAAGCAGG	169	563-585
AD-1557646	CUUCUUCUGGUUAUUUCUCCU	693	567-587	AGGAGAAUUAACCAAGAAAGCA	171	565-587
AD-1557647	UUCUUCUGGUUAUUUCUCCAU	694	568-588	ATGGAGAAUUAACCAAGAAAGC	172	566-588
AD-1557648	UUCUUCUGGUUAUUUCUCCAU	695	569-589	ATUGGAGAAUUAACCAAGAAAG	888	567-589
AD-1557649	CUUCUGGUUAUUUCUCCAAU	696	570-590	ATUUGGAGAAUUAACCAAGAA	889	568-590
AD-1557650	UUCUGGUUAUUUCUCCAAU	697	571-591	AAUUUGGAGAAUUAACCAAGAA	890	569-591
AD-1557651	UCUGGUUAUUUCUCCAAUUCU	698	572-592	AGAUUUGGAGAAUUAACCAAG	891	570-592
AD-1557652	CUGGUUAUUUCUCCAAUUCU	699	573-593	AGGAUUTGGAGAAUUAACCAAG	892	571-593
AD-1557682	GUGGAGGAGCTGCUUCCACU	700	643-663	AGUGGACAGCAGCTCCUCCACCA	893	641-663
AD-1557685	GAGGAGCUGCTGUCCACAGUU	701	646-666	AACUGUGGACAGCAGCUCUCCA	894	644-666
AD-1557689	AGCUGCUGTCCACAGUCAACU	702	650-670	AGUUGACUGUGGACAGCAGCUC	895	648-670
AD-1557690	GCUGCUGUCCACAGUCAACAU	703	651-671	ATGUUGACUGUGGACAGCAGCUC	896	649-671
AD-1557693	GCUGUCCACAGUCAACAGCUU	704	654-674	AAGCUGTUGACTGTGGACAGCAG	897	652-674
AD-1557694	CUGUCCACAGTCAACAGCUCU	705	655-675	AGAGCUGUUGACUGUGGACAGCA	898	653-675
AD-1557695	UGUCCACAGUCAACAGCUCGU	706	656-676	ACGAGCTGUUGACTGUUGGACAGC	899	654-676
AD-1557708	ACAGGGCCGAGUACGAAAGUGU	48	689-709	ACACUUCGUACTCGGCCCUUGUAG	900	687-709
AD-1557711	GGGCCGAGTACGAAGUGGACU	707	692-712	AGUCCACUUCGTACUCGGCCCU	901	690-712
AD-1557712	GGCCGAGUACGAAGUGGACCU	708	693-713	AGGUCCACUUCGUACUCGGCCCU	902	691-713
AD-1557726	AUCCUGGAAAGCCAGUGUGAAU	709	727-747	ATUCACACUGGCGTCCAGGAUCA	903	725-747
AD-1557727	UCCUGGAAAGCCAGUGUGAAAU	710	728-748	ATUUCACACUGGCTUCCAGGAUC	904	726-748
AD-1557728	CCUGGAAGCCAGUGUGAAAGU	711	729-749	ACUUUCACACUGGCUUCCAGGAU	905	727-749
AD-1557729	CUGGAAGCCAGUGUGAAAGAU	712	730-750	ATCUUUCACACTGGCUUCCAGGA	906	728-750
AD-1557730	UGGAAGCCAGTUGUGAAAGACU	713	731-751	AGUCUUTCACACUGGCUUCCAGG	907	729-751
AD-1557731	GGAAAGCCAGUGUGAAAGACAU	714	732-752	ATGUCUTUCACACTGGCUUCCAG	908	730-752
AD-1557732	GAAGCCAGTGTGAAAGACAUU	715	733-753	AAUGUCTUUACACACUGGCUUCCA	909	731-753
AD-1557733	AAGCCAGUGUGAAAGACAUU	716	734-754	ATAUGUCUUUACACACUGGCUUCC	910	732-754
AD-1557734	AGCCAGUGTGAAGACAUAGU	717	735-755	ACUAUUGTUUUACACACUGGCUUC	911	733-755
AD-1557735	GCCAGUGUGAAAGACAUAGCU	718	736-756	AGCUAUGUCUUUACACACUGGCUU	912	734-756
AD-1557736	CCAGUGUGAAAGACAUAGCUU	50	737-757	AAGCUATGUCUTUCACACUGGCU	913	735-757
AD-1557738	AGUGUGAAAGACAUAGCUGCU	719	739-759	AGCAGTAUGUCUTUCACACUGG	914	737-759
AD-1557739	GUGUGAAAGACAUAGCUGCAU	720	740-760	ATGCAGCUAUGTCTUUCACACUG	915	738-760
AD-1557740	UGUGAAAGACAUAGCUGCAUU	721	741-761	AAUGCAGCUAUGUCUUUCCACACU	916	739-761
AD-1557741	GUGAAAGACATAGCUGCAUUU	722	742-762	AAAUGCAGCUATGTCTUUUCCACAC	917	740-762

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1557758	AUUUAAUCCACGCGUGGGUUU	52	759-779	AAACCCAGCGUGGAAUCAAUUGC	178	757-779
AD-1557762	AAUCCACGCTGGGUUUAU	723	763-783	ATAACAACCCAGCGUGGAAUUA	181	761-783
AD-1557767	CACGCGUGGUGUUAACCGCU	724	768-788	AAGCGGTAAACAACCCAGCGUGGA	184	766-788
AD-1557768	ACGCGUGGUGUUAACCGCUAU	725	769-789	ATAGCGGUAAACAACCCAGCGUGG	918	767-789
AD-1557769	CGCUGGGUTGTUACCGCUACU	726	770-790	AGUAGCGGUAAACAACCCAGCGGUG	919	768-790
AD-1557770	GCUGGGUUGUTACCGCUACAU	727	771-791	ATGUAGCGGUAAACAACCCAGCGU	920	769-791
AD-1557771	CUGGGUUGTUAACCGCUACAGU	728	772-792	ACUGAAGCGGUAAACAACCCAGCG	921	770-792
AD-1557772	UGGGUUGUTACCGCUACAGCU	729	773-793	AGCUGAAGCGGTAAACAACCCAGC	922	771-793
AD-1557773	GGGUUGUUAACCGCUACAGCUU	730	774-794	AAGCUGTAGCGGUAAACAACCCAG	923	772-794
AD-1557836	CAAAUCCGGGUGGAGUGGAU	731	888-908	ATCCACTCCAGCCGGAGUUUAG	924	886-908
AD-1557866	GGGACCGACUGGCCAUGUAUU	60	923-943	AAUACATGGCCAGTCGGUCCCGG	925	921-943
AD-1557871	CGACUGGCCATGUUAUGACGUU	732	928-948	AACGUCAUACATGGCCAGUCCGU	926	926-948
AD-1557881	CUGGAGAAGAGGCUCAUCACU	733	958-978	AGUGAUGAGCCTCTUCUCCAGGG	927	956-978
AD-1557882	UGGAGAAGAGGCUCAUCACCU	734	959-979	AGGUGATGAGGCUUUCUCCAGG	928	957-979
AD-1557883	GGAGAAGAGGCUCAUCACCUU	735	960-980	AAGUGAUGAGCCTCUUCCAG	929	958-980
AD-1557884	GAGAAAGAGGCTCAUCACCUU	736	961-981	AGAGGUGAUGAGCCUUCUCCCA	930	959-981
AD-1557886	GAAAGAGGCTCAUCACCUCCGU	737	963-983	ACCGAGGUGAUGAGCCUUCUC	931	961-983
AD-1557890	AGGCUCAUACCUCCGGUGUAU	67	967-987	ATACACCGAGGTGAUAGCCUUCU	193	965-987
AD-1557944	GAAGAAGGGCCUGCACAGCUU	738	1053-1073	AAGCUGTGCAGGCCUUCUCCCA	932	1051-1073
AD-1557945	AAGAAGGGCCTGCACAGCUAU	739	1054-1074	ATAGCUGUGCAGGCCUUCUCC	933	1052-1074
AD-1557948	AAGGGCCUGCACAGCUACU	740	1057-1077	ATAGUAGCUGUGCAGGCCUUCU	934	1055-1077
AD-1557949	AGGGCCUGCACAGCUACUACU	741	1058-1078	AGUAGUAGCUGTGCCAGGCCUUC	935	1056-1078
AD-1557953	CCUGCACAGTACUACGACCU	742	1062-1082	AGGUCGTAGUAGCTGUGCAGGCC	936	1060-1082
AD-1558059	CCUCUCUGGACUACGGCUUGU	70	1235-1255	ACAAGCCGUAGTCCAGAGAGGCC	196	1233-1255
AD-1558061	UCUCUGGACUACGGCUUGGCU	71	1237-1257	AGCCAAGCCGUAGTCCAGAGAGG	937	1235-1257
AD-1558065	UGGACUACGGCUUGGCCUUCU	743	1241-1261	AGAGGGCCAAGCCGUAGUCCAGA	938	1239-1261
AD-1558066	GGACUACGGCTUGGCCUUCU	744	1242-1262	AAGAGGGCCAAGCCGUAGUCCAG	939	1240-1262
AD-1558105	GAGGAGGCAGAAUGAUUUU	76	1281-1301	AAAUACATACUUCUGCCUCCAG	202	1279-1301
AD-1558106	AGGAGGCAGAAUGAUUUU	745	1282-1302	AAAUAUAUACUUCUCCUCCCA	940	1280-1302
AD-1558113	AGAAUGAUGATUUGCCGUGCU	746	1289-1309	AGCACGGCAATCAUACUUCUGC	209	1287-1309
AD-1558114	GAAGUAUGAUTUUGCCGUGCAU	747	1290-1310	ATGCACGGCAAAUUAUUCUUG	210	1288-1310
AD-1558115	AAGUAUGAUTUUGCCGUGCACU	748	1291-1311	AGUGACGGCAAAUUAUUCUUC	941	1289-1311
AD-1558116	AGUAUGAUTUUGCCGUGCACCU	749	1292-1312	AGGUGACGGCAAAUUAUUCUUC	942	1290-1312
AD-1558117	GUUAUGAUTUUGCCGUGCACCCU	750	1293-1313	AGGGUGACGGCAAAUUAUUCUUC	943	1291-1313
AD-1558136	GGCCAGUGGACGAUCCAGAAU	751	1315-1335	ATUCUGGAUCGTCACACUGGCCCU	944	1313-1335

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1558137	GCCAGUGGACGAUCCAGAACU	752	1316-1336	AGUUCUGGAUCGUCCACUGGCC	945	1314-1336
AD-1558138	CCAGUGGACGAUCCAGAACU	753	1317-1337	ATGUUCTGGAUCGTCCACUGGCC	946	1315-1337
AD-1558139	CAGUGGACGATCCAGAACAGU	754	1318-1338	ACUGUUCUGGATCGUCCACUGGCC	947	1316-1338
AD-1558142	UGGACGAUCCAGAACAGGAGU	755	1321-1341	ACUCCUGUUCUGGACUGGCCACU	948	1319-1341
AD-1558150	CCAGAACAGGAGGCGUGUGUGU	87	1329-1349	ACACACAGCCUCCUUCUUCUGAU	949	1327-1349
AD-1558152	AGAACAGGAGGCGUGUGUGGU	88	1331-1351	AGCCACACAGCCUCCUUCUUGG	214	1329-1351
AD-1558211	ACUUCACCTCCAGAUCCU	756	1415-1435	AGGAGATCUGGAGGUGAAGUUG	950	1413-1435
AD-1558215	CACCUCCAGAUCCUCCU	757	1419-1439	ATGAGGGAGAUCCUGGGAGGUGAA	951	1417-1439
AD-1558230	UGUGCGGTGCACUAUGGCUU	758	1449-1469	AAGCCATAGUGACCCGACACCC	215	1447-1469
AD-1558231	GUGCGGUGGCACUAUGGCUUU	90	1450-1470	AAAGCCAUAGUGCACCCGACAC	216	1448-1470
AD-1558232	UGCGGUGGCACUAUGGCUUGU	759	1451-1471	ACAAGCCAUAGTGACCCGACACA	952	1449-1471
AD-1558233	GCGGGUGCACTAUGGCUUGU	760	1452-1472	AACAAGCCAUAGUGCACCCGAC	217	1450-1472
AD-1558234	CGGGUGCACUAUGGCUUGUAU	761	1453-1473	ATACAAGCCAUAGTGACCCGACA	953	1451-1473
AD-1558235	GGGUGCACTATGGCUUGUAU	762	1454-1474	AGUACAAGCCATAGUGCACCCG	218	1452-1474
AD-1558236	GGUGCACUAUGGCUUGUAU	763	1455-1475	ATGUACAAGCCAUAGUGCACCCG	954	1453-1475
AD-1558238	UGCACUAUGGCUUGUAU	764	1457-1477	AGUUGUACAAGCCAUAGUGCAC	955	1455-1477
AD-1558239	GCACUAUGGCTUGUAACAACCU	765	1458-1478	AGGUUGTACAAAGCCAUAGUGCAC	956	1456-1478
AD-1558249	CUGCCUGGAGAGUCCUUCU	95	1488-1508	AAGAGGAACUCTCCAGGGCAGGG	221	1486-1508
AD-1558250	UGCCUUGGAGAGUCCUUCUGU	766	1489-1509	ACAGAGGAACUCCAGGGCAGG	957	1487-1509
AD-1558288	AACGGCCUGGAGAGAGAAU	767	1561-1581	ATUUCUCUACUCCAGGCCGUUGG	958	1559-1581
AD-1558289	ACGGCCUGGATGAGAGAAU	768	1562-1582	AGUUUCTCUCATCCAGGCCGUUG	959	1560-1582
AD-1558290	CGGCCUGGAGAGAGAAACU	769	1563-1583	AAGUUUCUCUACUCCAGGCCGUU	960	1561-1583
AD-1558292	GCCUGGAGAGAGAAACUGCU	96	1565-1585	AGCAGUTUUCUCTCAUCCAGGCCG	222	1563-1585
AD-1558293	CCUGGAGAGAGAAACUGCGU	770	1566-1586	ACGAGTUUUCUCUACUCCAGGCC	961	1564-1586
AD-1558301	AGAGAACTGCGUUUGCAGAU	771	1574-1594	ATCUGCAACCGCAGUUUCUCUCA	962	1572-1594
AD-1558302	GAGAAACUGCGUUUGCAGAGU	772	1575-1595	ACUCUGCAACCGCAGUUUCUCUC	963	1573-1595
AD-1558308	CUGCGUUUGCAGAGCCACAUU	773	1581-1601	AAUGUGGCUUGCAACCGCAGUU	964	1579-1601
AD-1558309	UGCGUUUGCAGAGCCACAUUU	774	1582-1602	AAUGUGGCUCTGCAACCGCAGU	965	1580-1602
AD-1558310	GCGUUUGCAGAGCCACAUUCU	775	1583-1603	AGAAUGTGGCUCUGCAACCGCAG	966	1581-1603
AD-1558311	CGUUUGCAGAGCCACAUUCCU	776	1584-1604	AGGAAUGUGGCTCTGCAACCGCA	967	1582-1604
AD-1558316	GCAGAGCCACAUUCCAGUGCU	777	1589-1609	AGCACUGGAUUGUGGCUCUGCAA	968	1587-1609
AD-1558419	UGGGACAUTACACCUUCCAGUU	778	1710-1730	AACUGGAAGGUGAAUUGCCACA	228	1708-1730
AD-1558420	GGGACAUCACCUUCCAGUGU	779	1711-1731	ACACUGGAAGGTGAUUGUCCAC	969	1709-1731
AD-1558421	GGACAUCACCUUCCAGUGUU	103	1712-1732	AACACUGGAAGGUGAAUUGCCCA	229	1710-1732
AD-1558423	ACAUUCACCUUCCAGUGUAU	780	1714-1734	ATCACACUGGAAGGUGAAUUGCC	231	1712-1734

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1558449	GAGCUGCGTGAAGAAGCCCAU	781	1740-1760	ATGGGCTUCUUCACGCAGCUCCG	970	1738-1760
AD-1558450	AGCUGCGUGAAGAAGCCCAU	782	1741-1761	ATUGGGCUUCUTACGCAGCUCC	971	1739-1761
AD-1558451	GCUGCGUGAAGAAGCCCAU	783	1742-1762	AGUUGGGUUCUUCACGCAGCUC	972	1740-1762
AD-1558452	CUGCGUGAAGAAGCCCAU	784	1743-1763	AGUUUGGGUUCUUCACGCAGCU	973	1741-1763
AD-1558453	UGCGUGAAGAAGCCCAU	785	1744-1764	AGGUUGGGUUCUUCACGCAGC	974	1742-1764
AD-1558508	AGCAGUGAGUUGGGCCU	786	1808-1828	AGAGCCACAGTCACAGUCCUCC	975	1806-1828
AD-1558546	CUCGAGGGUGAGUGGCCAU	787	1866-1886	AAUGGCCACUACCCUCCGAGGA	976	1864-1886
AD-1558576	AUCGUGACCGUGGGU	107	1936-1956	AAUACCCACAGCGTACGCAUGA	977	1934-1956
AD-1558577	UCGUGACCGUGGGU	788	1937-1957	ATAUACCCACAGCGGUCAGCGAUG	978	1935-1957
AD-1558578	CGCUGACCGTGGGUAU	789	1938-1958	ATUAUACCCACAGCGGUCAGCGAU	979	1936-1958
AD-1558579	GCUGACCGUGGGUAU	790	1939-1959	AGUUAUACCCACAGCGGUCAGCGA	980	1937-1959
AD-1558586	GCUGGGUGUAACAGCUGCCU	791	1946-1966	AGGACGUGUUAUACCCAGCGG	981	1944-1966
AD-1558609	UGCUUCCAGGAGGACAGCAU	792	1969-1989	AAUGCUGUCCUCCCTGGAAAGCAGU	982	1967-1989
AD-1558610	GCUCCAGGAGGACAGCAUGU	793	1970-1990	ACAUGTGUCCCTCCUGGAAGCAG	983	1968-1990
AD-1558611	CUUCCAGGAGGACAGCAUGU	794	1971-1991	ACCAUGCUGUCCUCCUGGAAGCA	984	1969-1991
AD-1558650	CGUGUCCCTGGGCAAGGUGU	795	2010-2030	AACACCTUGCCACAGGAACACGGU	235	2008-2030
AD-1558657	CUGGGCAAGGTUGGGCAGAAU	796	2017-2037	ATUCUGCCACACCTUGCCACAGGA	985	2015-2037
AD-1558658	UGGGCAAGGUGUGGCAGAAU	797	2018-2038	AGUUCUGCCACACCUUGCCACAG	986	2016-2038
AD-1558659	GGGCAAGGTGTGGCAGAAU	798	2019-2039	AAGUUCGTGCCACACCUUGCCACAG	987	2017-2039
AD-1558660	GGCAAGGUGUGGCAGAAU	799	2020-2040	AGAGUUCUGCCACACCUUGCCCA	988	2018-2040
AD-1558661	GCAAGGUGTGGCAGAAU	800	2021-2041	ACGAGUTCUGCCACACCUUGCC	237	2019-2041
AD-1558662	CAAGGUGUGGCAGAAU	801	2022-2042	AGCGAGTUCUGCCACACCUUGCC	989	2020-2042
AD-1558683	UGGCCUGGAGAGGUGUCCU	802	2044-2064	AAAGGACACCUUCCAGGCCAGC	990	2042-2064
AD-1558684	GGCCUGGAGAGGUGUCCU	803	2045-2065	AGAAGGACACCTCTCCAGGCCAG	991	2043-2065
AD-1558685	GCCUGGAGAGGUGUCCU	804	2046-2066	ATGAAGGACACCUUCCAGGCCA	992	2044-2066
AD-1558686	CCUGGAGAGGTGUCCU	805	2047-2067	ATUGAAGGACACCTCUCACAGGCC	993	2045-2067
AD-1558687	CUGGAGAGGUGUCCU	113	2048-2068	ACUUGAAGGACACCUUCCAGGC	239	2046-2068
AD-1558691	AGAGGUGUCCUCAAAGGUAU	806	2052-2072	ATCACCTUGAAGGACACCUUCC	242	2050-2072
AD-1558833	UGUGCAGUTGAUCCACAGGU	807	2289-2309	ACCUGUGGGAUCAACUGCACAUC	994	2287-2309
AD-1558835	UGCAGUUGAUCCACAGGACU	808	2291-2311	AGUCCUGUGGGAUCAACUGCACA	995	2289-2311
AD-1558843	AUCCACAGGACCUUGCAGU	117	2299-2319	ACUGCACAGGUCCCTGUGGGAUCA	996	2297-2319
AD-1558845	CCCACAGGACCUUGCAGCGU	809	2301-2321	ACGUGCACAGGUCCUUGGGGAU	997	2299-2321
AD-1558846	CCACAGGACCTGUGCAGCGAU	810	2302-2322	ATCGUGCACAGGTCUUGGGGA	998	2300-2322
AD-1558878	CCAGGUGACGCCACGCAUGCU	811	2334-2354	AGCAUGCGUGGCGTCAACCUUGUA	999	2332-2354
AD-1558882	GUGACGCCACGCAUGCUGUGU	812	2338-2358	ACACAGCAUGCGUGGCGUCACCU	1000	2336-2358

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1558883	UGACGCCACGCAUGCUGUGUU	118	2339-2359	AACACAGCAUGCGTGGCGUACCC	1001	2337-2359
AD-1558885	ACGCCACGCATGCGUGUGCU	813	2341-2361	AGCACACAGCATGCGUGGCGUCA	1002	2339-2361
AD-1558905	GGCUACCGCAAGGGCAAGAAU	814	2362-2382	ATUCUUGCCCTGCGGUAGCCGG	1003	2360-2382
AD-1558906	GCUACCCGCAAGGGCAAGAAU	120	2363-2383	ACUUCUTGCCCTGCGGUAGCCG	246	2361-2383
AD-1558907	CUACCGCAAGGGCAAGAAU	121	2364-2384	ACUUCUTGCCCTGCGGUAGCC	1004	2362-2384
AD-1558961	GUGCAAGGCACUCAGUGGCCU	815	2418-2438	AGGCCATUGAGTGCCUUGCACAC	1005	2416-2438
AD-1558992	CUAACUACTUCGGCGUCUACU	816	2486-2506	AGUAGACGCCGAAGUAGUAGGC	1006	2484-2506
AD-1558995	ACUACUUCGGCGUCUACACCU	817	2489-2509	AGGUGUAGACGCCGAAGUAGUUA	1007	2487-2509
AD-1558996	CUACUUCGGCGUCUACACCCU	818	2490-2510	AGGGUAGTAGACGCCGAAGUAGUU	1008	2488-2510
AD-1559004	GGGUUACACCCGCAUCACAU	819	2498-2518	ATGUGATGGGGUGUAGACGCCG	1009	2496-2518
AD-1559005	CGUUAACACCCGCAUCACAGU	820	2499-2519	ACUGUAGUGGGGTGUAGACGCC	1010	2497-2519
AD-1559008	CUACACCCGCAUCACAGGUGU	124	2502-2522	ACACCUGUGAUGCGGGUGUAGAC	250	2500-2522
AD-1559012	ACCCGCAUCACAGGUGUGAUU	821	2506-2526	AUACACACCUGTGAUGCGGGUGU	1011	2504-2526
AD-1559013	CCCGCAUCACAGGUGUGAUU	822	2507-2527	AGUACACCUUGUAGCGGGUGU	1012	2505-2527
AD-1559036	UGGAUCCAGCAAGUGGUGACU	823	2530-2550	AGUACACCUUGTGGAUCCAGC	1013	2528-2550
AD-1559038	GAUCCAGCAAGUGGUGACCUU	824	2532-2552	AAGGUCACCACTGCGGUAUCCA	1014	2530-2552
AD-1559039	AUCCAGCAAGTGGUGACCUU	825	2533-2553	ACAGGUCACCACTGCGGUAUCC	1015	2531-2553
AD-1559041	CCAGCAAGTGGUGACCUAGU	826	2535-2555	ACUAGGUCACCACTGCGGUAU	1016	2533-2555
AD-1559042	CAGCAAGUGGTGACCUAGGUGU	827	2536-2556	ACCUCAGGUCACCACTGCGGUA	1017	2534-2556
AD-1559044	GCAAGUGGTGACCUAGGAAU	828	2538-2558	ATUCCUCAGGUCACCACTGCGU	1018	2536-2558
AD-1559105	UGGUGGCAGGAGGUGGCAUCU	829	2667-2687	AGAUGCCACCTGCTGCCACACA	1019	2665-2687
AD-1559106	GGUGGCAGGAGGUGGCAUCUU	830	2668-2688	AAGAUGCCACCTGCTGCCACCA	1020	2666-2688
AD-1559107	GUGGCAGGAGGUGGCAUCUUU	831	2669-2689	AAAGAUGCCACCTGCTGCCACCA	1021	2667-2689
AD-1559109	GGCAGGAGGUGGCAUCUUGUU	127	2671-2691	AACAAGAUGCCACCTGCTGCCAC	253	2669-2691
AD-1559133	UCCUGAUGUCUGCUCACAGUU	832	2695-2715	AACUGGAGCAGACAUAGGGACG	1022	2693-2715
AD-1559136	CUGAUGUCTGCUCCAGUGAUU	833	2698-2718	AUACACTGGAGCAGACAUAGGG	1023	2696-2718
AD-1559147	UCCAGUGATGGCAGGAGGAAU	834	2709-2729	AUCCUCCUGCCATCAGUGGAGC	1024	2707-2729
AD-1559233	GGCUCAGCAGCAAGAAUGCUU	132	2853-2873	AAGCAUTCUUGCUGGUGAGCCAC	258	2851-2873
AD-1559318	CUAACUUGGGAUCUGGGAAUU	835	2978-2998	AUUCGCCAGAUCCCAAGUAGAC	1025	2976-2998
AD-1559323	UUGGGAUCTGGGAUUGGAAU	836	2983-3003	ACUUCCAUCCCAAGAUCCCAAGU	264	2981-3003
AD-1559431	GUGAGCUCAGCUGCCCUUUGU	837	3157-3177	ACAAAGGGCAGCUGAGCUCACCU	1026	3155-3177
AD-1559436	CUCAGCUGCCCUUUGGAAUUAU	838	3162-3182	ATAUCCAAAGGGCAGCUGAGCU	1027	3160-3182
AD-1559437	UCAGCUGCCCTUUGGAAUAAU	839	3163-3183	ATUAUCCAAAGGGCAGCUGAGC	1028	3161-3183
AD-1559438	CAGCUGCCCTUUGGAAUAAU	840	3164-3184	ATUAUTCCAAAGGGCAGCUGAG	1029	3162-3184
AD-1559441	CUGCCCUUTGGAAUAAAGCUU	841	3167-3187	AAGCUUTAUUCCAAAGGGCAGCU	1030	3165-3187

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4	Antisense Sequence 5' to 3'	SEQ ID NO:	Range in NM_153609.4
AD-1559443	GCCUUUUGGAAUAAAGCUGCU	842	3169-3189	AGCAGTUUUAUTCCAAAGGGCAG	1031	3167-3189
AD-1559444	CCUUUUGGAATAAAGCUGCCU	843	3170-3190	AGGCAGCUUUAUCCAAAGGGCA	1032	3168-3190
AD-1559445	CCUUUGGAAUAAAGCUGCCUU	844	3171-3191	AAGGCAGCUUUAUCCAAAGGGC	1033	3169-3191
AD-1559447	UUUGGAAUAAAGCUGCCUGAU	845	3173-3193	ATCAGGCAGCUTUAUCCAAAGG	1034	3171-3193
AD-1559448	UUGGAAUAAAGCUGCCUGAUU	846	3174-3194	AUACAGGCAGTUTUAUCCAAAG	1035	3172-3194
AD-1559449	UGGAAUAAAGCUGCCUGAUCU	847	3175-3195	AGAUCAGGCAGCUTUAUCCAAA	1036	3173-3195

Table 5. Modified Sense and Antisense Strand Sequences of TMPRSS6 dsRNA Agents

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO	Antisense Sequence 5' to 3'	SEQ ID NO	mRNA target sequence 5' to 3'	SEQ ID NO
AD-1557376	csGsgaggugdAudGgegaggagauL96	1037	asdCsuuuc(Tgn)cgccAudCacucuccgsu	1264	GACGGAGGUGAUGGCGAGGAAGC	1491
AD-1557377	gsGsgaggugdTgdGegaggagauL96	1038	asdGseuuc(C2p)uegcdCadTcaccuccgsu	1265	ACGGAGGUGAUGGCGAGGAAGCG	1492
AD-1557396	asagsgccugdTgdAggacuccauL96	1039	asdTsuag(C2p)uuccudCadCagccuucgsa	1266	UCAAGGCCUGAGGACUCCAAG	1493
AD-1557398	gsgscucugdTgdGacuccaagauL96	1040	asdTseuug(C2p)agucdCudCacaggcccsu	1267	AAGGCCUGAGGACUCCAAGAG	1494
AD-1557399	gscscugugdTgdAcuccaagagauL96	1041	asdCsuuuc(C2p)ggagdTgdTcacaggccsu	1268	AGGCCUGAGGACUCCAAGAGA	524
AD-1557400	cscsugugdTgdGadCuccaagagauL96	1042	asdTseuuc(Tgn)ggagdTgdCucacaggccsc	1269	GGCCUGAGGACUCCAAGAGAA	1495
AD-1557401	csusugugdTgdAcTccaagagauL96	1043	asdTseuuc(Tgn)ugagdTgdCucacaggccsc	1270	GCCUGAGGACUCCAAGAGAAA	1496
AD-1557437	csusacucdTgdGudAnuuccuagguL96	1044	asdCsuag(C2p)aaudAcCagagugscsa	1271	UGCUACUCUGGUUUUCCUAGGG	529
AD-1557440	csuscugguadTudTccuaggguauL96	1045	asdTsaccc(Tgn)aggadAadTaccagagcsa	1272	UACUCUGGUUUUCCUAGGGUAC	532
AD-1557441	uscusugguadTudCcuaggguacuL96	1046	asdGsuacc(C2p)uagdTgdAadAuaaccagsgu	1273	ACUCUGGUUUUCCUAGGGUACA	533
AD-1557442	csusugguadTgdCuaaggguacuL96	1047	asdTsguac(C2p)cuagdTgdAuaaccagsgu	1274	CUCUGGUUUUCCUAGGGUACAA	1497
AD-1557443	usgsuauuudCcdTaggguacuL96	1048	asdTsuag(C2p)ccuadGgdAaauaccagsgu	1275	UCUGGUUUUCCUAGGGUACAAG	1498
AD-1557444	gsgsuauuudCudAgggguacuL96	1049	asdCsuuug(Tgn)ccuadAgdTgaauaccagsgu	1276	CUGGUUUUCCUAGGGUACAAGG	1499
AD-1557445	ggsuauuudCcdTadGggguacuL96	1050	asdCsuuug(Tgn)acccdTadGgaauaccagsgu	1277	UGGUUUUCCUAGGGUACAAGGC	1500
AD-1557452	csusagguadCcdAgggcgagguuL96	1051	asdAsceuc(C2p)gcccudTgdTaccuagsgu	1278	UCCUAGGGUACAAGGGCGGAGGUG	1501
AD-1557473	asusggucagdTgdAgguacuL96	1052	asdGsuag(C2p)accudGgdCugaccuagsgu	1279	UGAUGGUCAGCCAGGUGUACUCA	1502
AD-1557475	gsgsucagccdTgdGugguacuL96	1053	asdCsuuug(Tgn)acacdTgdGgcuagccagsgu	1280	AUGGUACGCCAGGUGUACUCAGG	535
AD-1557476	ggsucagccdTgdTguacuagguL96	1054	asdCsuuug(C2p)uacacdTgdTggcuagccagsgu	1281	UGGUACGCCAGGUGUACUCAGGC	1503
AD-1557477	uscusagccagdTgdGuaucagguL96	1055	asdGseuuc(Tgn)guacdTgdCugcuagccagsgu	1282	GGUCAGCCAGGUGUACUCAGGCA	1504
AD-1557478	csasgcccagdTgdTaccuagguL96	1056	asdTsgccu(C2p)agudCadCugcuagccagsgu	1283	GUCAGCCAGGUGUACUCAGGCAG	1505

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO	Antisense Sequence 5' to 3'	SEQ ID NO	mRNA target sequence 5' to 3'	SEQ ID NO
AD-1557479	asgscacagcudGudAucacagcagcagL96	1057	asdCsuagcc(Tgn)gagudAcdAccuagcagcgsa	1284	UCAGCCAGGUGUACUCAGGCAGU	536
AD-1557509	csuscauucgdCcdAucuccccaL96	1058	asdTsgggg(G2p)aagudGgcGauuagagusa	1285	UACUCAAUUCGCCACUUCUCCAG	1506
AD-1557515	csagccacudCudCecaggaucuuL96	1059	asdAsgauc(C2p)uagggdAgaAagugcgcgsasu	1286	AUGCCACUUCUCCCCAGGAUCUU	1507
AD-1557516	gscscauuecdTcdCaggaucuuL96	1060	asdAsgauc(C2p)uagggdGadGagugcgcgsa	1287	UCGCCACUUCUCCCCAGGAUCUU	537
AD-1557518	csascuuecdCcdAgaucuuL96	1061	asdGsuag(Agn)uccudGgdGagagugcgsa	1288	GCCACUUCUCCCCAGGAUCUUACC	1508
AD-1557522	uscuccacgdGadTcuuacccgcL96	1062	asdGscggg(Tgn)aagadTcdCugggagagasc	1289	CUUCUCCAGGAUCUUACCCGCC	1509
AD-1557523	csuscccaggdAudCuuacccgcL96	1063	asdGsgcgg(G2p)uagadAudCugggagagasa	1290	UUUCCCCAGGAUCUUACCCGCCG	1510
AD-1557524	uscscacaggdTcdTuuacccgcL96	1064	asdCsgcgg(G2p)guaadGadTccugggagasa	1291	UCUCCCCAGGAUCUUACCCGCCGG	539
AD-1557550	usasugccudTcdCgcagugaaL96	1065	asdTsuuca(C2p)uagcdGadAagcacuagasa	1292	UCUAGUGCCUCCGAGUGAAAC	1511
AD-1557554	gscscuuecdCcdGugaacgcL96	1066	asdGscggg(Tgn)uaccdTcdCggaagcgsasc	1293	GUGCCUUCGCGAGUGAAACCGCC	540
AD-1557555	csusuuecdAgaTgaacccgcL96	1067	asdGsgcgg(Tgn)uucadCudCcggaagcgsa	1294	UGCCUUCGCGAGUGAAACCGCCA	1512
AD-1557556	csusucegcdGudGaaacccgcL96	1068	asdTsgcgg(G2p)uucudAcdTgcggagcgsa	1295	GCCUUCGCGAGUGAAACCGCCAA	1513
AD-1557559	csccgcagugdAadAccgccaaL96	1069	asdCsuuug(G2p)cggdTudCacugcgsasa	1296	UUCGCGAGUGAAACCGCCAAAGC	1514
AD-1557560	csagcagugadAadCcgccaaL96	1070	asdGscuuu(C2p)cggdTudTcacugcgsa	1297	UCCGCGAGUGAAACCGCCAAAGCC	1515
AD-1557561	gscsagugadAcdCgcaaaL96	1071	asdGsgcgg(Tgn)ggcgGudTucacugcgsa	1298	CCGCGAGUGAAACCGCCAAAGCCC	1516
AD-1557562	csasugaaadCcdGccaaL96	1072	asdGsgcgg(Tgn)uagcdGgdTuuacugcgsa	1299	CGCAGUGAAACCGCCAAAGCCCA	1517
AD-1557563	asgsuagaaadCgdCcaagccL96	1073	asdTsgggc(Tgn)uagcdCgdGuuacugcgsa	1300	GCAGUGAAACCGCCAAAGCCCG	1518
AD-1557571	csagccaaagCcdCagaagL96	1074	asdGscuac(Tgn)uagcdGgdCuuuagcgsa	1301	ACCGCCAAAGCCCGAGAGAGUCU	1519
AD-1557572	gscsaaagcdCcdAgaagL96	1075	asdAsgcau(C2p)uucudGgdGuuugcgsa	1302	CCGCCAAAGCCCGAGAGAGUCUC	1520
AD-1557577	asgscaccagadAgaAugcuaL96	1076	asdCsuug(Agn)gcaudCudTcuugcgsa	1303	AAAGCCAGAGAGAGAGAGAGGA	1521
AD-1557606	csasgaccdGcdCugggaaL96	1077	asdAsgauu(C2p)ccagdGgdGgugcgsa	1304	ACCAGCACCCCGCCUGGGAACUUA	1522
AD-1557607	asgscaccdCcdTgggaL96	1078	asdTsaagu(Tgn)ccagdGgdCggugcgsa	1305	CCAGCACCCCGCCUGGGAACUUA	1523
AD-1557629	ascsaaccdAgaCuccuL96	1079	asdAauaga(C2p)ggagdCudGgaugcgsa	1306	CUACAACUCCAGCUCGUCUUAU	1524
AD-1557630	csasacccadGcdTccgcuL96	1080	asdAauag(Agn)cgagdGcdTggagcgsa	1307	UACAACUCCAGCUCGUCUUAUUC	541
AD-1557639	uscscaccdTudCuucggL96	1081	asdGsaacc(Agn)gaagdAadCagagcgsa	1308	CCUCACCCUGCUUCUUCUGGUUA	544
AD-1557640	csasccugcdTcdTucggL96	1082	asdTsgaac(C2p)agaadGadAagcagcgsa	1309	CUCACCCUGCUUCUUCUGGUUAU	1525
AD-1557642	csccugcudTudCugggL96	1083	asdAauaga(Agn)ccagdAadGagcagcgsa	1310	CACCUGCUUCUUCUGGUUAUUC	546
AD-1557643	csugcucudTcdTgguL96	1084	asdGsaug(Agn)accadGadAgaagcgsa	1311	ACCGCUUCUUCUGGUUAUUCU	1526
AD-1557644	uscgcuucudCudGguuL96	1085	asdAsgaa(C2p)aaacdAgaAgaagcgsa	1312	CCUGCUUCUUCUGGUUAUUCUC	547
AD-1557646	csusucucudGgdTucuuL96	1086	asdGsgaga(Agn)uagadCcdAgaagcgsa	1313	UGCUUCUUCUGGUUAUUCUCCA	549
AD-1557647	usucucugdGudTcauL96	1087	asdTsggag(Agn)auagdAadCagaagcgsa	1314	GUUCUUCUGGUUAUUCUCCAA	550
AD-1557648	uscscuuegdTudCaucL96	1088	asdTsuag(C2p)aaagdAadCagaagcgsa	1315	GUUCUUCUGGUUAUUCUCCAA	1527
AD-1557649	csusucugdCcdAucuccaaL96	1089	asdTsuugg(Agn)gaagdGadAccagagcgsa	1316	UUUCUUCUGGUUAUUCUCCAAA	1528
AD-1557650	usucugudCadTucuaaL96	1090	asdAsuuug(G2p)agaadTgdAaccagagcgsa	1317	UCUUCUGGUUAUUCUCCAAAUC	1529

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO	Antisense Sequence 5' to 3'	SEQ ID NO	mRNA target sequence 5' to 3'	SEQ ID NO
AD-1557651	uscsggguucdAudTcucaaauculL96	1091	asdGsaunuu(G2p)gagadAudGaaccagagasa	1318	CUUCUGGUUUAUUCUCCAAAUCC	1530
AD-1557652	csusggguucadTudCucaaauculL96	1092	asdGsaunuu(Tgn)ggagadAadTgaaccagasa	1319	UUCUGGUUUAUUCUCCAAAUCCC	551
AD-1557682	gusugggagadGedTguguccaculL96	1093	asdGugugai(C2p)agcadGedTcuuccacsa	1320	UGGUGGAGGAGGUCUGUCUCCACA	1531
AD-1557685	gsasggagcadGedTguccacaculL96	1094	asdAseuuiu(G2p)gaeadGedAgnuucacsa	1321	UGGAGGAGGUCUGUCUCCACAGUC	1532
AD-1557689	asgscugcugdTcdCacagucaculL96	1095	asdGsunua(C2p)ugugadGadCagcagucsc	1322	GGAGCUGCUGUCCACAGUCAACA	1533
AD-1557690	gscsugcugudCodAcagucaculL96	1096	asdTsgung(Agn)cugudGgdAcagcagcusc	1323	GAGCUGCUGUCCACAGUCAACAG	1534
AD-1557693	gscsugucadCadGucaacagcuculL96	1097	asdAsgcug(Tgn)ugadTgdTggacagcusc	1324	CUGCUGUCCACAGUCAACAGCUC	1535
AD-1557694	csusguccadAadTcaacagcuculL96	1098	asdGsgacui(G2p)uugadCudGuggacagcsc	1325	UGCUGUCCACAGUCAACAGCUCG	1536
AD-1557695	usgsuaccadGudCaacagcuculL96	1099	asdCsgagc(Tgn)gugudAadTgugacagcsc	1326	GCUGUCCACAGUCAACAGCUCGG	1537
AD-1557708	ascsagggccadGadGuacgaaugulL96	1100	asdCsacuu(C2p)guadTcdGgcccugusag	1327	CUACAGGGCCGAGUACGAAAGUGG	552
AD-1557711	gsgsgccgagdTadCgaaguggaculL96	1101	asdGsucau(C2p)uucgdTadCucggccusc	1328	CAGGGCCGAGUACGAAAGUGGACC	1538
AD-1557712	gsgscggagudAcdGaaguggaculL96	1102	asdGsguuc(Agn)cuuudGudAcueggccscsu	1329	AGGGCCGAGUACGAAAGUGGACCC	1539
AD-1557726	asusccuggadAcdCagugugaaulL96	1103	asdTsucau(C2p)uucgdCudTccaggaucsa	1330	UGAUCCUGGAAAGCCAGUGUGAAA	1540
AD-1557727	uscscuggaadGcdCagugugaaulL96	1104	asdTsucau(C2p)uucgdGedTuccaggaucsc	1331	GAUCCUGGAAAGCCAGUGUGAAAAG	1541
AD-1557728	cscsuggaaadCcdAguugugaaulL96	1105	asdCsaunuc(Agn)cacudGgdCuuccaggsasu	1332	AUCCUGGAAAGCCAGUGUGAAAAGA	1542
AD-1557729	csusggaaadCcdGugugaaulL96	1106	asdTscuuu(C2p)uucgdTgdGcuuccaggsa	1333	UCCUGGAAAGCCAGUGUGAAAAGAC	1543
AD-1557730	usgsaagccadAadTgugaaagaculL96	1107	asdGsucau(Tgn)cacudCudGcuuccaggsag	1334	CCUGGAAAGCCAGUGUGAAAAGACA	1544
AD-1557731	gsgsaagccadGudGugaaagaculL96	1108	asdTsguuc(Tgn)uucadAadTggcuuccasag	1335	CUGGAAAGCCAGUGUGAAAAGACAU	1545
AD-1557732	gsasagccagdTgdTgaagacaculL96	1109	asdAsguuc(Tgn)uucadCadCuuggcuuccsa	1336	UGGAAAGCCAGUGUGAAAAGACAU	1546
AD-1557733	asasgccaadGudGaaagacaculL96	1110	asdTsaugui(C2p)uucudAcdAcuuggcuuccsc	1337	GGAAGCCAGUGUGAAAAGACAUAG	1547
AD-1557734	asgscagugdTgdAaagacaculL96	1111	asdCsuauu(Tgn)cuuudCadCacuggcucusc	1338	GAAGCCAGUGUGAAAAGACAUAGC	1548
AD-1557735	gscscagugudGadAagacaculL96	1112	asdGsucau(G2p)uucudTcdAcacuggcucusu	1339	AAGCCAGUGUGAAAAGACAUAGCU	1549
AD-1557736	cscsagugudAadAagacaculL96	1113	asdAsgcuu(Tgn)uucudTudCacacuggcusc	1340	AGCCAGUGUGAAAAGACAUAGCUG	554
AD-1557738	asgsugugaadAadAacacaculL96	1114	asdGscagc(Tgn)uugudCudTucacacusc	1341	CCAGUGUGAAAAGACAUAGCUGCA	1550
AD-1557739	gusugugaaadGadCauagcugcuculL96	1115	asdTsgcag(C2p)uugudTcdTucacacusc	1342	CAGUGUGAAAAGACAUAGCUGCAU	1551
AD-1557740	usgsugaaadAcdAuagcugcuculL96	1116	asdAsguca(G2p)uucudGudCuuccacacscsu	1343	AGUGUGAAAAGACAUAGCUGCAU	1552
AD-1557741	gusugaaagadCadTgucugcuculL96	1117	asdAaagc(Agn)gcuadTgdTcuuccacacsc	1344	GUGUGAAAAGACAUAGCUGCAUUG	1553
AD-1557758	asusugaaadCcdAcgucggguculL96	1118	asdAaacc(Agn)gcuudGgdAauncacacsc	1345	GCAUUGAAUCCACGCGGGGUUG	556
AD-1557762	asasuuccadGcdTggguguguculL96	1119	asdTsaaca(Agn)cccadGcdGuggaucacsa	1346	UGAAUUCACGCGGGGUUGUUAAC	559
AD-1557767	csascgucggdTgdTguuacguculL96	1120	asdAsgcgg(Tgn)uucadAcdCacgucggsa	1347	UCCACGCGGGGUUGUUAACCGCUA	562
AD-1557768	ascsgucggdTgdTguuacguculL96	1121	asdTsaagc(G2p)uucadAadCccagcuggsag	1348	CCACGCGGGGUUGUUAACCGCUAC	1554
AD-1557769	csgscugggdTgdTuaacgucaculL96	1122	asdGsuuag(C2p)uucadAcdAaccagcuggsu	1349	CACGCGGGGUUGUUAACCGCUACA	1555
AD-1557770	gscsugggudGudTaccgucaculL96	1123	asdTsguag(C2p)gguadAcdAaccagcugsu	1350	ACGCGGGGUUGUUAACCGCUACAG	1556
AD-1557771	csusgggugdTudAccgucaculL96	1124	asdCsuua(G2p)cgguadAadCaaccagcsg	1351	CGCUGGGGUUGUUAACCGCUACAGC	563

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO	Antisense Sequence 5' to 3'	SEQ ID NO	mRNA target sequence 5' to 3'	SEQ ID NO
AD-1557772	usgsaguuuTadCgcuaacagcuL96	1125	asdGscuu(Agn)gcggdTadAcaaccasgsc	1352	GCUGGGUUGUACCGCUACAGCU	1557
AD-1557773	gsaguuuAdCgcuaacagcuL96	1126	asdAsgcu(Tgn)agcgGudAcaaccasg	1353	CUGGGUUGUACCGCUACAGCUA	1558
AD-1557836	csasaacuccGgdCuggaguggauL96	1127	asdTscac(Tgn)ccagCcdGgaguuuagsag	1354	CUCAACUCCGGCUGGAGUGGAC	1559
AD-1557866	gsagaccGadCudGgcuaugauuL96	1128	asdAsuaac(Tgn)ggccAdTgcguccasgsg	1355	CCGGACCCAGCUGGCCAUGUAG	564
AD-1557871	csagacugcdCadTgnaugacguL96	1129	asdAscgu(Agn)uacdTgdGccagucgsgsu	1356	CCGGACUGGCCAUGUAGACGUG	1560
AD-1557881	csusgagagadGadGgcuaacauL96	1130	asdGsuag(G2p)agcdTcdTuccacagsgsg	1357	CCUGGAGAAAGAGGCUCAUCACC	1561
AD-1557882	usgsagagadAgdGeuaucacuL96	1131	asdGsguag(Tgn)gagcdCudCuuuccasgsg	1358	CCUGGAGAAAGAGGCUCAUCACCU	1562
AD-1557883	gsagagagadGgdCueaucacuL96	1132	asdAsggu(Agn)ugadCcdTcnuccasgsg	1359	CUGGAGAAAGAGGCUCAUCACCU	1563
AD-1557884	gsagagagadGgdTcaucacuL96	1133	asdGaggu(G2p)augadGedCcuuuccesca	1360	UGGAGAAAGAGGCUCAUCACCU	1564
AD-1557886	gsasagagagcdTcdAucacucguL96	1134	asdCscgag(G2p)ugadGadGccuuccesuc	1361	GAGAAAGAGGCUCAUCACCU	1565
AD-1557890	asgsacuaudCadCucgugguauL96	1135	asdTsacac(C2p)gaggdTgdAugagcucsu	1362	AGAGGCUCAUCACCU	571
AD-1557944	gsasagagagcdGcdCugcagacuL96	1136	asdAsgcu(Tgn)gagcdGedCcuuuccesca	1363	UGGAGAAAGAGGCUCAUCACCU	1566
AD-1557945	asasagagagcdCcdTgcacagcuL96	1137	asdTsagcu(G2p)ugcdGgdCcuuuccesuc	1364	GGAGAAAGAGGCUCAUCACCU	1567
AD-1557948	asasgagccudGcdAacagacuL96	1138	asdTsaguu(G2p)ccgudGedAagccuuccesuc	1365	AGAAGGCUCAUCACCU	1568
AD-1557949	asgsagccuudCadCagcuacuL96	1139	asdGsuag(Tgn)gcuadTgdCagccuuccesuc	1366	GAGGCUCAUCACCU	1569
AD-1557953	cscsuagacGcdTactaacgacuL96	1140	asdGsguag(Tgn)aguuadGedTgucagagsgsc	1367	GGCUGCACAGCUACUACGACCC	573
AD-1558059	cscsuagacGadCuaagcguuL96	1141	asdCsaag(C2p)guagdTcdCagagagsgsg	1368	GCCUCUCUGGACUACGGCUUGG	574
AD-1558061	uscucuggadCudAegcguuL96	1142	asdGscac(G2p)ccgudAgdTccagagagsgsg	1369	CCUCUCUGGACUACGGCUUGGCC	575
AD-1558065	usgsagacudGgdCuuagcccuL96	1143	asdGsaag(G2p)ccagdTcdGuaucacsgsa	1370	UCUGGACUACGGCUUGGCCCU	1570
AD-1558066	gsagacuagcdGcdTuggcccuL96	1144	asdAsgag(G2p)ccadGedCgucagccasg	1371	CUGGACUACGGCUUGGCCCU	1571
AD-1558105	gsasgagagcdAgdAguuaguuuL96	1145	asdAsaaca(Tgn)acuadCudGccuuccasg	1372	CUGAGGAGGCAGAAAGUAUGAUU	580
AD-1558106	asgsagagcdGadAguuaguuuL96	1146	asdAsaaca(Tgn)acuadTcdTgcuccesca	1373	UGAGGAGGCAGAAAGUAUGAUU	1572
AD-1558113	asgsaaguuadGadTuuagccguL96	1147	asdGscag(G2p)ccadTcdAucuuuccesuc	1374	GCAGAAAGUAUGAUUUGCCGUGCA	587
AD-1558114	gsasaguuagadAudTugccguL96	1148	asdTsgcac(G2p)gcaadAudCauuuccesuc	1375	CAGAAAGUAUGAUUUGCCGUGCAC	588
AD-1558115	asasguuagadTudTgcccguL96	1149	asdGsuag(C2p)gagcdAadTcauuccesuc	1376	AGAAGUAUGAUUUGCCGUGCAC	1573
AD-1558116	asgsuagauudTudGccgucacuL96	1150	asdGsguag(C2p)gagcdAadAucuuuccesuc	1377	GAAGUAUGAUUUGCCGUGCAC	1574
AD-1558117	gsusaaguuudTgdCcgucacccuL96	1151	asdGsguag(C2p)gagcdCadAucuuuccesuc	1378	AAGUAUGAUUUGCCGUGCAC	1575
AD-1558136	gsagccagudGadCgaucagauL96	1152	asdTsucag(G2p)aucgdTcdCacugcccsu	1379	AGGCCAGUGGACGACUCCAGAAC	1576
AD-1558137	gscsaguggdAcdGaucagauL96	1153	asdGsuuag(G2p)gagcdGudCcacugccsc	1380	GGCCAGUGGACGACUCCAGAAC	1577
AD-1558138	cscsaguggadCgdAuccagauL96	1154	asdTsguu(Tgn)ggaudTcdTccuagsgsc	1381	GGCCAGUGGACGACUCCAGAAC	1578
AD-1558139	csasuguggadGadTccagacauL96	1155	asdCsuuag(G2p)uggdTcdGuccacugsgsc	1382	GCCAGUGGACGACUCCAGAAC	589
AD-1558142	usgsagacudCadAgaacagauL96	1156	asdCsuuag(G2p)uucdTgdAucgucacsu	1383	AGGCCAGUCCAGAACAGG	1579
AD-1558150	cscsagacudGgdAggcuugguL96	1157	asdCsaaca(Tgn)gagcdTgucuggsu	1384	AUCCAGAACAGGAGGCUUGG	591
AD-1558152	asgsaacagadAgdGeuugugguL96	1158	asdGscac(Tgn)gagcdCudCcuuuccesuc	1385	CCAGAACAGGAGGCUUGG	592

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO	Antisense Sequence 5' to 3'	SEQ ID NO	mRNA target sequence 5' to 3'	SEQ ID NO
AD-1558211	ascsmuaccdTcdCagauucuccuL96	1159	asdGsgaga(Tgn)cuagdgGadGguagaugusg	1386	CAACUUCACCUCCAGAUUCCUCC	1580
AD-1558215	csascucccdAgaAuccuccuauL96	1160	asdTsgagG(C2p)agauCudGggagguGsasa	1387	UUCACCUCCAGAUUCCUCCAC	1581
AD-1558230	usgsugcgggdTgdCacuauugguuL96	1161	asdAsgcca(Tgn)aguGdCadCccgcacasc	1388	GGUGGCGGGUGCAGAUUGGCUU	1593
AD-1558231	gsusgcgggudGedAciuagguuuL96	1162	asdAsagcc(Agn)uagudGedAaccgcacsc	1389	GUGUGCGGGUGCAGAUUGGCUU	1594
AD-1558232	usgscgggudCadCuauugguuuL96	1163	asdCsaagc(C2p)auagdTgdAaccgcacsc	1390	UGUGCGGGUGCAGAUUGGCUU	1582
AD-1558233	gscsgggugcdAcdTaugguuuL96	1164	asdAscaag(C2p)cuaudGudGcaccgcsasc	1391	GUGCGGGUGCAGAUUGGCUU	1595
AD-1558234	csrgsgggucadCudAugguuuL96	1165	asdTsacaa(G2p)ccauAgaTgacccgcsca	1392	UGCGGGUGCAGAUUGGCUU	1583
AD-1558235	gsrgugcadcTadTggguuuL96	1166	asdGsuaca(Agn)gcccTadGugcaccgsc	1393	GCGGGUGCAGAUUGGCUU	1596
AD-1558236	gsrgugcadcAuaGguuuL96	1167	asdTsguac(Agn)agccAuaGguuuL96	1394	CGGGUGCAGAUUGGCUU	1584
AD-1558238	usgscacuandGgdCuuguaacuL96	1168	asdGsuuu(Agn)cagcdCedAuaugcasc	1395	GGUGCAGAUUGGCUU	1585
AD-1558239	gscsacuauGdGduuuaacuL96	1169	asdGsguuu(Tgn)acaadGedCauaagucsc	1396	GUGCAGAUUGGCUU	1586
AD-1558249	csusgcccudGadGaguuuuL96	1170	asdAsgagG(Agn)acuTcdCaggcgagsg	1397	CCUGCCCCUGGAGAGUUCUUG	1599
AD-1558250	usgscuccudAgaAguuuuuL96	1171	asdCsagG(C2p)aaudCudCaggcgagsg	1398	CCUGCCCCUGGAGAGUUCUUG	1587
AD-1558288	asascggccudGgdAugagaaauL96	1172	asdTsuuu(C2p)uauCudAagccguusg	1399	CCACGGCCUGGAGAGAGAAAC	1588
AD-1558289	ascsgggccudGadTgagagaaauL96	1173	asdGsuuu(Tgn)uauCudCaggcgagsg	1400	CAACGGCCUGGAGAGAGAAAC	1589
AD-1558290	csrgsgccudAuaGagagaaauL96	1174	asdAsguuu(C2p)uauCudCaggcgagsg	1401	AACGGCCUGGAGAGAGAAAC	1590
AD-1558292	gscscugaudGadGagaaacuL96	1175	asdGscagG(Tgn)uauCudAuccagcscg	1402	CGGCCUGGAGAGAGAGAAAC	1600
AD-1558293	csusgguuGadAgaacuuL96	1176	asdCsgcag(Tgn)uauCudCauccagcsc	1403	GGCCUGGAGAGAGAGAGAAAC	1591
AD-1558301	asgsagaaadTgdCguuugcagauL96	1177	asdTscugc(Agn)aaagCudGmuuucscsa	1404	UGAGAGAAACUGCGUUGCAGAG	1592
AD-1558302	gsasgaacuGdGguuugcagauL96	1178	asdCsucG(C2p)aaagCudAguuucscsu	1405	GAGAGAAACUGCGUUGCAGAGC	1593
AD-1558308	csusgguuudGedAgaagcauuL96	1179	asdAsugG(C2p)uauCudAaacgcagsg	1406	AACUGCGUUGCAGAGCCACA	1594
AD-1558309	usgscguuudCadGagcauuL96	1180	asdAsauG(C2p)gcuTgdCaaacgcagsg	1407	ACUGCGUUGCAGAGCCACA	1595
AD-1558310	gscgguuudGadAgaacuuL96	1181	asdGsaauG(Tgn)gcuCudCaaacgcagsg	1408	CUGCGUUGCAGAGCCACA	1596
AD-1558311	csrgsuuudGadGcccauuL96	1182	asdGsgauG(C2p)uagGdTgdTgcaaacgscsa	1409	UGCGUUGCAGAGCCACA	1597
AD-1558316	gscsagagcdAcdAuuccaguuL96	1183	asdGscacu(C2p)gaudGudGgcuuGcsasa	1410	UUGCAGAGCCACA	1598
AD-1558419	usgsagacudTadAauuccaguuL96	1184	asdAscugG(Agn)agudGadAuguuccscsa	1411	UGUGGGACA	1606
AD-1558420	gsrgacauudCadCuuccaguuL96	1185	asdCsacug(C2p)aaagdTgdAauguccscsa	1412	GUGGGACA	1599
AD-1558421	gsrgacauudAcdCuuccaguuL96	1186	asdAscacu(C2p)gaagGudGauuuccscsa	1413	UGGGACA	607
AD-1558423	ascsauuacudCudTccaguuL96	1187	asdTscacu(C2p)uagGdTgdGuaugucsc	1414	GGACA	609
AD-1558449	gsasgcugcdTgdAagaagccauL96	1188	asdTsgggG(Tgn)uauCudCgagcucscg	1415	CGGAGCUGGUGAAGAGGCCAA	1600
AD-1558450	asgsucgGadAgaagccauL96	1189	asdTsuagG(C2p)uauCudAagcagucsc	1416	GGAGCUGGUGAAGAGGCCAA	1601
AD-1558451	gscsugcGadAadGaagcccauL96	1190	asdGsuuG(Tgn)uauCudAagcagucsc	1417	GAGCUGGUGAAGAGGCCAA	1602
AD-1558452	csusgcgGadAgaagcccauL96	1191	asdGsguuG(C2p)gcuCudTcacgcagscsu	1418	AGCUGCGUGAAGAGGCCAA	1603
AD-1558453	usgscguGadAgaagcccauL96	1192	asdGsgguu(C2p)gcuCudTcacgcagscsu	1419	GCUGCGUGAAGAGGCCAA	1604

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO	Antisense Sequence 5' to 3'	SEQ ID NO	mRNA target sequence 5' to 3'	SEQ ID NO
AD-1558508	asgscacugudGadCuguggcuccuL96	1193	asdGsaagc(C2p)acagdTcdAacagucscsc	1420	GGAGCACUGUGACUGUGGCCUCC	1605
AD-1558546	csusccagagGdGadGaguggccauL96	1194	asdAsuggc(C2p)acudAcCucgggsgsa	1421	UCCUCCGAGGGUGAGUGGCCAUG	1606
AD-1558576	asuscgeugadCcdGcugggugauL96	1195	asdAsueac(C2p)ecagdTcagcgaugsgsa	1422	UCAUCGUGACCCGCGUGGUGAUA	611
AD-1558577	uscsugcagcdCcdCugggugauL96	1196	asdTsuauc(C2p)ecagdTcdGucagcagsg	1423	CAUCGUGACCCGUGGUGGUAUA	1607
AD-1558578	csugscugaccdCcdTgggugauL96	1197	asdTsuauc(Agn)ccadCcdGucagcagsg	1424	CAUCGUGACCCGUGGUGGUAUA	1608
AD-1558579	gscsuagaccGdGdGgugauL96	1198	asdGsuuac(C2p)accadAcdCgucagcsgsa	1425	UCGUGACCCGUGGUGGUAUA	1609
AD-1558586	gscsuagggudAudAacagugccuL96	1199	asdGsaagc(C2p)ugudAudCaccagcsgsg	1426	CCGUGGGUGAUACAGCUGCCC	1610
AD-1558609	usgscuuecadGgdAggacagcauL96	1200	asdAsugcu(G2p)ueudCcdTggagcagsgu	1427	ACUGCUUCCAGGAGGACAGCAUG	1611
AD-1558610	gscsuuecadGdGgacagcauL96	1201	asdCsaugc(Tgn)gucdTcdCugagcagsg	1428	CUGCUUCCAGGAGGACAGCAUGG	1612
AD-1558611	csusuccagGdAgdGacagcauL96	1202	asdCsaug(C2p)ugudCudCugagcagsgsa	1429	UGCUUCCAGGAGGACAGCAUGGC	1613
AD-1558650	csugsuuuccdTgdGgcaagggguL96	1203	asdAsacc(Tgn)ugcdCadGgaacagsgsu	1430	ACCGUUGUCCUGGCCAAGGUGUG	613
AD-1558657	csusgggcaadGgdTguggcagaaL96	1204	asdTsueug(C2p)ecadCcdTgucceagsgsa	1431	UCCUGGGCAAGGUGUGGCGAGAAC	1614
AD-1558658	usgsgcaagGdGdGgagcauL96	1205	asdGsuuuc(G2p)ccadAcCcuagcagsgsg	1432	CCUGGGCAAGGUGUGGCGAGAACU	1615
AD-1558659	gsgsgcaagGdTgdTggcagaaL96	1206	asdGsuuuc(Tgn)gccadCcdCcuagcagsgsg	1433	CUGGGCAAGGUGUGGCGAGAACUC	1616
AD-1558660	gsgscagggudGdGgcaagaaL96	1207	asdGsaugu(C2p)ugcdAcCcuagcagsgsa	1434	UGGGCAAGGUGUGGCGAGAACUCG	1617
AD-1558661	gscsaagggudTgdGagaaucguL96	1208	asdCsaggu(Tgn)ueudCadCacuuagcscsc	1435	GGGCAAGGUGUGGCGAGAACUCGC	615
AD-1558662	csasagggudGdGgcaagaaL96	1209	asdGsgcag(Tgn)ueudCcdAcacuuagcsc	1436	GGCAAGGUGUGGCGAGAACUCGCG	1618
AD-1558683	usgsgcugGdAgdAgguguccuL96	1210	asdAsagga(C2p)acudCudCagcagcagsg	1437	GCUGGCCUGGAGAGGUGUCCUUC	1619
AD-1558684	gsgscugggadGdGguguccuL96	1211	asdGsaagc(Agn)ccadCcdTccagcagsg	1438	CUGGCCUGGAGAGGUGUCCUUA	1620
AD-1558685	gscscugggadAgdGuguccuL96	1212	asdTsgaag(G2p)acacCudCuccagcagsgsa	1439	UGGCCUGGAGAGGUGUCCUCAA	1621
AD-1558686	cscsuaggadGgdTgucuucauL96	1213	asdTsugaal(G2p)gacacCcdTcuccagcagsg	1440	GGCCUGGAGAGGUGUCCUCAAAG	1622
AD-1558687	csusggagadGdGgucuucauL96	1214	asdCsuuga(Agn)ggacAcCucuccagsgsc	1441	GCCUGGAGAGGUGUCCUCAAAGG	617
AD-1558691	asgsagggudCcdTuccagggauL96	1215	asdTscac(Tgn)ugadGdAcacuuagcsc	1442	GGAGAGGUGUCCUCAAAGGUGAG	620
AD-1558833	usgsugcagdTgdAuuccagguL96	1216	asdCscugu(G2p)ggadCadAcuagcagsg	1443	GAUGGCAGUUGAUUCCACAGGA	1623
AD-1558835	usgscaguuGdAudCccacaggaL96	1217	asdGsuuuc(G2p)ugggadAudCacucagcagsg	1444	UGUGCAGUUGAUUCCACAGGACC	1624
AD-1558843	asuscccacGdAcuugugcauL96	1218	asdCsuaga(C2p)aggadCcdTgugggauL96	1445	UGAUCCACAGGACCGUGGACGC	621
AD-1558845	cscscacagGdAcCugugcagguL96	1219	asdCsugcu(G2p)acagGdCcuugggsgsu	1446	AUCCACAGGACCGUGGACGCGA	1625
AD-1558846	cscsacagGdCcdTgugcaggaL96	1220	asdTscgu(G2p)ccadGdTccuugggsgsa	1447	UCCACAGGACCGUGGACGCGAG	1626
AD-1558878	cscsagggudCgdCcagcauL96	1221	asdGsaug(C2p)guggdCgdTccuugggsgsa	1448	UACCAGGUGAGCGCACGCAUGCU	1627
AD-1558882	gscsagcccdAcCgagcugguL96	1222	asdCsacag(C2p)augdGdGgucacscsu	1449	AGGUGACCGCACGCAUGCUGUGU	1628
AD-1558883	usgsaccccdCgdCaugcugguL96	1223	asdAsacac(C2p)caudCgdTgggucacsc	1450	GGUGACCGCACGCAUGCUGUGUG	622
AD-1558885	ascsgccacGdCadTgucugguL96	1224	asdGsaaga(C2p)agcdTgdCgugggsgsa	1451	UGACGCCACGCAUGCUGUGGCC	1629
AD-1558905	gsgscuaccGdCadAgggcagaaL96	1225	asdTsucuu(G2p)cccdTgdCgugagcsgsg	1452	CCGGCUACCGCAAGGGCAAGAAG	1630
AD-1558906	gscsuaccGdAadGggcagaaL96	1226	asdCsuuuc(Tgn)gcccdTudGgugagcsgsg	1453	CGGCUACCGCAAGGGCAAGAAGG	624

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO	Antisense Sequence 5' to 3'	SEQ ID NO	mRNA target sequence 5' to 3'	SEQ ID NO
AD-1558907	csusaccgcadAgdGgcagagagguL96	1227	asdCseuuc(Tgn)ugcccdCudTgcgguagscsc	1454	GGCUACCCGCAAGGGCAAGGAAGGA	625
AD-1558961	gsusgcagaggdCadCucaaguggcucL96	1228	asdGsgcca(C2p)ugagdTgdCcuugcacsasc	1455	GUGUGCAAAGGCACUCAGUGGGCCG	1631
AD-1558992	csusaacuadTudCggcgucucacL96	1229	asdGsuaga(C2p)gccegdAdCguaguuagscsc	1456	GCCUAAACUACUUCGGGUCUACACA	1632
AD-1558995	ascstaeuadGgdCgucuaacacL96	1230	asdGsgguu(Tgn)gacgdGdGaguuagutsa	1457	UAACUACUUCGGGUCUACACCC	1633
AD-1558996	csusacuucgdGdCgucuaacacL96	1231	asdGsgguu(Tgn)agacdGdCgaguuagsusu	1458	AACUACUUCGGGUCUACACCCG	1634
AD-1559004	gscgcuacacAcCgcgaucacacL96	1232	asdTsguga(Tgn)ggcgGdGuaagacscscg	1459	CGGCGUCUACACCCGCAUCACAG	1635
AD-1559005	csgsuacuacAcCgcgaucacacL96	1233	asdCsuug(Tgn)ugcgGdTguagacscsc	1460	GGCGUCUACACCCGCAUCACAGG	1636
AD-1559008	csusacaccdGcdAucacagguL96	1234	asdCsacu(C2p)ugaudGdGgguuagscsc	1461	GUCUACACCCGCAUCACAGGU	628
AD-1559012	ascscgcacudCadCagguugauL96	1235	asdAsuac(Agn)ccuGdTgdAugcgguusgu	1462	ACACCCGCAUCACAGGUGUAUC	1637
AD-1559013	cscscgcacudAcCagguugauL96	1236	asdGsauc(C2p)accudGdGaugcgguusgu	1463	CACCCGCAUCACAGGUGUAUCA	1638
AD-1559036	usgsuacacGdCagguugauL96	1237	asdGsuac(C2p)acuudGdTggaucacscsc	1464	GCUGAUCCAGCAAGUGGUGACC	1639
AD-1559038	gsasucacGdAadGugguagacuuL96	1238	asdAsgguc(Agn)ccacdTudCcuugacscsa	1465	UGGAUCCAGCAAGUGGUGACCUG	1640
AD-1559039	asuscacacGdAgdTggugacuuL96	1239	asdCsagu(C2p)accadCudTgcuugacscsc	1466	GGAUCCAGCAAGUGGUGACCUGA	1641
AD-1559041	cscsagcaagdTgdGugacuuL96	1240	asdCsacag(C2p)uacCdCcuugacscsa	1467	AUCCAGCAAGUGGUGACCUGAGG	1642
AD-1559042	csasgcaagdTgdTgacuuL96	1241	asdCsucac(C2p)gucadCcdAcuugcugscsa	1468	UCCAGCAAGUGGUGACCUGAGGA	1643
AD-1559044	gsasagugdTgdAccuagagauL96	1242	asdTsuc(C2p)agguCdCacuugcugsc	1469	CAGCAAGUGGUGACCUGAGGAAC	1644
AD-1559105	usgsuggcadGdAgguugacuuL96	1243	asdGsaug(C2p)accuCdTgcccacscsa	1470	UGUGGUGGAGGAGGUGGCAUCU	1645
AD-1559106	gsusuggcadGdGugugacuuL96	1244	asdAsagu(C2p)cccdTcdCugccacscsc	1471	GUGGUGGAGGAGGUGGCAUCUU	1646
AD-1559107	gsusggcagdTgdGugugacuuL96	1245	asdAsagu(C2p)cccdCudCcuugcscsa	1472	UGGUGGAGGAGGUGGCAUCUUG	1647
AD-1559109	gsusggcagdTgdGugugacuuL96	1246	asdAsagu(C2p)cccdCudCcuugcscsa	1473	GUGGAGGAGGUGGCAUCUUGUC	631
AD-1559133	uscscuagdTgdCugcucacuuL96	1247	asdAscug(Agn)gcagdTudCcuugcscsc	1474	CGUCCUGAUGUGUGUCCUCCAGUG	1648
AD-1559136	csusgaugdTgdCuccagaguuL96	1248	asdAsuac(Tgn)ggagdTudCacuugcscsc	1475	CCUGAUGUGUGUCCUCCAGUGAUG	1649
AD-1559147	uscscuagdTgdCuccagaguuL96	1249	asdAsuac(C2p)cccdCudCcuugcscsc	1476	GUCCAGUGAUGGAGGAGGAGUG	1650
AD-1559233	gsusgcagdTgdCuccagaguuL96	1250	asdAsuac(Tgn)cccdCudCcuugcscsc	1477	GUGGUCACAGCAAGAAUGCUG	636
AD-1559318	csusaacuGdTgdAucugggauL96	1251	asdAsuac(C2p)agadCcdCacuugcscsc	1478	GUCUAAACUUGGGAUCUGGGAUUG	1651
AD-1559323	usugggaudTgdGgaugaguuL96	1252	asdCsucc(Agn)uuccCdGauccacscsc	1479	ACUUGGGAUCUGGGAUUGGAAGG	642
AD-1559431	gsusgcagdTgdCugcucuuL96	1253	asdCsaaag(C2p)gcagdTudCagcucacscsc	1480	AGGUGAGUCAGCUGCCCUUUGG	1652
AD-1559436	csuscagcudCcdCcuugaguuL96	1254	asdTsauuc(C2p)aaagdTgdCagcugcscsc	1481	AGCUCAGCUGCCCUUUGGAUAA	1653
AD-1559437	uscsagcudCcdCcuugaguuL96	1255	asdTsauuc(C2p)aaagdTgdCagcugcscsc	1482	GCUCAGCUGCCCUUUGGAUAAA	1654
AD-1559438	csasgcudCudTuggaauaaL96	1256	asdTsauuc(Tgn)ccadAgdGgagcugcscsc	1483	CUCAGCUGCCCUUUGGAUAAAAG	1655
AD-1559441	csusgcudTgdGauaaaguuL96	1257	asdAsguu(Tgn)uuccCdAagcagcscsc	1484	AGCUGCCCUUUGGAUAAAAGCUG	648
AD-1559443	gscscuuuGdGadAuaagcucL96	1258	asdGscag(Tgn)uuccCdCaaagcscsc	1485	CUGCCCUUUGGAUAAAAGCUGCC	1656
AD-1559444	cscscuuuGdAadTaaagcucL96	1259	asdGscag(C2p)uuccCdCaaagcscsc	1486	UGCCCUUUGGAUAAAAGCUGCCU	1657
AD-1559445	cscscuuuGdAadTaaagcucL96	1260	asdAsggca(C2p)uuccAudTccaaagcscsc	1487	GCCCUUUGGAUAAAAGCUGCCUG	1658

Duplex Name	Sense Sequence 5' to 3'	SEQ ID NO	Antisense Sequence 5' to 3'	SEQ ID NO	mRNA target sequence 5' to 3'	SEQ ID NO
AD-1559447	ususuggaudAadAgcugccugauL96	1261	asdTscagg(C2p)agcudTudAuuccaaasg	1488	CCUUUGGAAUAAAGCUGCCUGAU	1659
AD-1559448	ususggaauadAadGeugccugauL96	1262	asdAsucag(G2p)ecagdTudTuuccaasag	1489	CUUUGGAAUAAAGCUGCCUGAUC	1660
AD-1559449	usgsaauaadAadCugccugauL96	1263	asdGsauc(a(G2p)gcagdTudTuuuccasasa	1490	UUUGGAAUAAAGCUGCCUGAUCC	1661

Table 6. Unmodified Sense and Antisense Strand Sequences of TMPRSS6 dsRNA Agents

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO	Range in NM_153609.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO	Range in NM_153609.4	SEQ ID NO
AD-1570929.1	CGGAGGUGAUGGCGAGGAAGU	650	189-209	ACUUCCTCGCCAUACACCUCCGUC	848	187-209	848
AD-1570930.1	CCUGUGAGGACUCCAAAGAGAU	654	233-253	AUCUCUTGGAGUCCUCACAGGCC	1726	231-253	1726
AD-1570931.1	CUGUGAGGACUCCAAAGAGAAU	1662	234-254	AUUCTCTUGGAGUCCUCACAGGC	1727	232-254	1727
AD-1570932.1	CUCUGGUAUUUCCUAGGGUAU	28	331-351	AUACCTTAGGAAUACACAGAGUA	1728	329-351	1728
AD-1570933.1	GGUAUUUCCUAGGGUACAAGU	660	335-355	ACUUGUACCCUAGGAAUACACAG	858	333-355	858
AD-1570934.1	GUUUUUUCCUAGGGUACAAGGU	1663	336-356	ACCUAGTACCCUAGGAAUACCA	1729	334-356	1729
AD-1570935.1	GGUCAGCCAGGUGUACUCAGU	31	366-386	ACUGAGTACACCUAGGUGACCAU	157	364-386	157
AD-1570936.1	UCAGCCAGGUGUACUCAGGCU	665	368-388	AGCCTGAGUACACCUAGGUGACC	1730	366-388	1730
AD-1570937.1	AGCCAGGUGUACUCAGGCAGU	32	370-390	ACUGCCTGAGUACACCUAGGUGA	158	368-390	158
AD-1570938.1	CACUUCUCCAGGAGUACUUAU	670	409-429	AGUAGAUCUUGGGAGAGUGGC	867	407-429	867
AD-1570939.1	UCUCCAGGAGUUAUACCCGCU	1664	413-433	AGCGGTAAGAUUCCUGGGAGAG	1731	411-433	1731
AD-1570940.1	GCCUUCCGCAGUGAAACCCGCU	36	445-465	AGCGGUTUACUCUGCGGAAGGCAC	1732	443-465	1732
AD-1570941.1	CCUUCCGCAGUGAAACCCGCU	1665	446-466	AGCGGUTUACUCUGCGGAAGGCA	872	444-466	872
AD-1570942.1	GCAGUGAAACCGCCAAAGCCU	679	452-472	AGGCTUTGGCGGUUUCACUGCG	1733	450-472	1733
AD-1570943.1	CAGUGAAACCGCCAAAGCCCU	680	453-473	AGGCGUTUUGCGGUUUCACUGCG	1734	451-473	1734
AD-1570944.1	AGUGAAACCGCCAAAGCCCAU	681	454-474	AUGGGCTUUGCGGUUUCACUGC	1735	452-474	1735
AD-1570945.1	CGCCAAAGCCCAAGAGUAGCU	682	462-482	AGCATCTUCUGGGCUUUGGCGGU	1736	460-482	1736
AD-1570946.1	AGCCCAAGAGAGUAGCUAAGGU	684	468-488	ACCUTGAGCAUCUUCUGGGCUU	1737	466-488	1737
AD-1570947.1	AGCACCCGCGGGAACUUAU	1666	499-519	AUAGUTCCCAGGCGGGUGCUGG	1738	497-519	1738

Duplex Name	Sense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:
AD-1570948.1	CAACUCCAGCUCGUCUAUUU	522-542	37	AAAUAGACGGAGCUGGAGUUGUA	520-542	163
AD-1570949.1	UCACCUUGCUCUUCUGGUUCU	560-580	40	AGAACCCAGAAAGCAGGUGAGG	558-580	166
AD-1570950.1	CCUGCUUCUUCUGGUUCAUUU	563-583	42	AAAUAGAACCCAGAAAGCAGGUG	561-583	168
AD-1570951.1	CUGCUUCUUCUGGUUCAUUU	564-584	1667	AGAATGAACCCAGAAAGCAGGUG	562-584	1739
AD-1570952.1	CUUCUUCUGGUUCAUUCUCCU	567-587	45	AGGAGAAUAGAACCCAGAAAGCA	565-587	171
AD-1570953.1	UUCUUCUGGUUCAUUCUCCU	568-588	46	AUGGAGAAUAGAACCCAGAAAGC	566-588	1740
AD-1570954.1	CUUCUGGUUCAUUCUCCAAU	570-590	1668	AUUUGGAGAAUAGAACCCAGAAAG	568-590	1741
AD-1570955.1	CUGGUUCAUUCUCCAAAUCCU	573-593	47	AGGATUTGGAGAAUAGAACCCAGAA	571-593	173
AD-1570956.1	GCUGCUGUCCACAGUCAACAU	651-671	703	AUGUTGACUGUGGACAGCAGCUC	649-671	1742
AD-1570957.1	GCUGUCCACAGUCAACAGCUU	654-674	704	AAGCTGTUGACUGUGGACAGCAG	652-674	1743
AD-1570958.1	UGUCCACAGUCAACAGCUCGU	656-676	706	ACGAGCTGUUGACUGUGGACAGC	654-676	1744
AD-1570959.1	GGCCGAGUACGAAGUGGACCU	693-713	708	AGGUCCACUUCGUACUCGGGCCU	691-713	902
AD-1570960.1	AUCCUGGAAGCCAGUGUGAAU	727-747	709	AUUCACACUGGCUUCCAGGAUCA	725-747	1745
AD-1570961.1	CCUGGAAGCCAGUGUGAAAGU	729-749	711	ACUUTCACACUGGCUUCCAGGAU	727-749	1746
AD-1570962.1	UGGAAGCCAGUGUGAAAGACU	731-751	1669	AGUCTUTCACACUGGCUUCCAGG	729-751	1747
AD-1570963.1	GGAAAGCCAGUGUGAAAGACAU	732-752	714	AUGUCUTUCACACUGGCUUCCAG	730-752	1748
AD-1570964.1	GAAAGCCAGUGUGAAAGACAUU	733-753	1670	AAUGTCTUUCACACUGGCUUCCA	731-753	1749
AD-1570965.1	AGCCAGUGUGUGAAAGACAUAGU	735-755	1671	ACUATGTCTUUCACACUGGCUUC	733-755	1750
AD-1570966.1	CCAGUGUGUGAAAGACAUAGCUU	737-757	50	AAGCTATGUCUUCACACUGGCU	735-757	1751
AD-1570967.1	AGUGUGUGAAAGACAUAGCUGCU	739-759	719	AGCAGCTAUGUCUUCACACUGG	737-759	1752
AD-1570968.1	GUGAAAGACAUAGCUGCAUUU	742-762	1672	AAAUAGCAGUAUGUCUUCACAC	740-762	1753
AD-1570969.1	AUUGAAUUCACACGUGGGUUU	759-779	52	AAACCCAGCUGGAAUUCACUUGC	757-779	178
AD-1570970.1	AAUUCACGUGGGUUGUUAU	763-783	55	AUAACAACCCAGCUGGAAUUA	761-783	1754
AD-1570971.1	CACGUGGGUUGUUAACCGCUU	768-788	58	AAGCGGTAAACAACCCAGCUGGA	766-788	184
AD-1570972.1	UGGGUUGUUAACCGCUACAGCU	773-793	1673	AGCUGUAGCGGUAAACAACCCAGC	771-793	1755
AD-1570973.1	GGGUUGUUAACCGCUACAGCUU	774-794	730	AAGCTGTAGCGGUAAACAACCCAG	772-794	1756

Duplex Name	Sense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:
AD-1570974.1	CAAAACUCCGGCUGGAGUGGAU	888-908	731	AUCCACTCCAGCCGGAGUUUGAG	886-908	1757
AD-1570975.1	GGGACCGACUGGCCAUGUAUU	923-943	60	AAUACATGGCCAGUCGGUCCCGG	921-943	186
AD-1570976.1	CGACUGGCCAUGUAUGACGUU	928-948	1674	AACGTCAUACAUGGCCAGUCCGU	926-948	1758
AD-1570977.1	UGGAGAAGAGGCUCUACACCU	959-979	734	AGGUGATGAGCCUUCUUCUCCAGG	957-979	928
AD-1570978.1	GGAGAAGAGGCUCUACACCUU	960-980	735	AAGGTGAUGAGCCUUCUUCUCCAG	958-980	1759
AD-1570979.1	GAAGAAGGGCCUGCACAGCUU	1053-1073	738	AAGCTGTGCAGGCCCUUCUUCUCCA	1051-1073	1760
AD-1570980.1	AGGGCCUGCACAGCUACUACU	1058-1078	741	AGUAGUAGCUGUGCAGGCCCUUC	1056-1078	1761
AD-1570981.1	CCUGCACAGCUACUACGACCU	1062-1082	69	AGGUCGTAGUAGCUGUGCAGGCC	1060-1082	195
AD-1570982.1	GAGGAGGCAGAAAGUAUGAUUU	1281-1301	76	AAAUACATACUUCUGCCUCCUACAG	1279-1301	202
AD-1570983.1	AGGAGGCAGAAAGUAUGAUUUU	1282-1302	745	AAAATCAUACUUCUGCCUCCUCCA	1280-1302	1762
AD-1570984.1	AGUAUGAUUUGCCUGGACCU	1292-1312	1675	AGGUGCAGGCAAAUCAAUACUUC	1290-1312	942
AD-1570985.1	CCAGUGGACGAUCCAGAAACAU	1317-1337	753	AUGUTCTGGAUCGUCACUUGGCC	1315-1337	1763
AD-1570986.1	CCAGAAACAGGAGGCUGUGUGU	1329-1349	87	ACACACAGCCUCCUGUUCUGGAU	1327-1349	213
AD-1570987.1	AGAACAGGAGGCUGUGUGGCU	1331-1351	88	AGCCACACAGCCUCCUGUUCUGG	1329-1351	214
AD-1570988.1	ACUUCACCCUCCAGAUUCUCCU	1415-1435	1676	AGGAGATCUGGGAGGUGAAGUUG	1413-1435	950
AD-1570989.1	UGUGCGGGUGCACUAUGGCUU	1449-1469	89	AAGCCATAGUGCACCCGCACACC	1447-1469	215
AD-1570990.1	GUGCGGGUGCACUAUGGCUUU	1450-1470	90	AAAGCCAUAGUGCACCCGCACAC	1448-1470	216
AD-1570991.1	GGGUGCACUAUGGCUUGUACU	1454-1474	92	AGUACAAAGCCAUAGUGCACCCCGC	1452-1474	1764
AD-1570992.1	GGUGCACUAUGGCUUGUACAU	1455-1475	763	AUGUACAAAGCCAUAGUGCACCCCG	1453-1475	1765
AD-1570993.1	UGCACUAUGGCUUGUACAAACU	1457-1477	764	AGUUGUACAAGCCAUAGUGCACCC	1455-1477	955
AD-1570994.1	GCACUAUGGCUUGUACAAACCU	1458-1478	1677	AGGUTGTACAAGCCAUAGUGCAC	1456-1478	1766
AD-1570995.1	CUGCCUUGGAGAGUUCUCCU	1488-1508	95	AAGAGGAAACUCCAGGGCAGGG	1486-1508	1767
AD-1570996.1	ACGGCCUGGAGAGAGAAACU	1562-1582	1678	AGUUTCTCUCAUCCAGGCCGUUG	1560-1582	1768
AD-1570997.1	GCCUGGAGAGAGAAACUGCU	1565-1585	96	AGCAGUTUCUCUCAUCCAGGCCG	1563-1585	1769
AD-1570998.1	CCUGGAGAGAGAAACUGCGU	1566-1586	770	ACGCAGTUUCUCUCAUCCAGGCC	1564-1586	961
AD-1570999.1	AGAGAAACUGCGUUUUGCAGAU	1574-1594	1679	AUCUGCAAACGCAGUUUUCUCUCA	1572-1594	1770

Duplex Name	Sense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:
AD-1571000.1	GCGUUUGCAGAGCCACAUUCU	1583-1603	775	AGAATGTGGCUCUCGAAAGCGCAG	1581-1603	1771
AD-1571001.1	UGGGACAUUACACCUUCCAGUU	1710-1730	102	AACUGGAAGGUGAAUGUCCACACA	1708-1730	228
AD-1571002.1	GAGCUGCGUGAAGAAGCCCAU	1740-1760	1680	AUGGGCTUUCUACGACGCUCCG	1738-1760	1772
AD-1571003.1	CGCUGACCGCUGGGUGUAU	1938-1958	1681	AUATCACCACGCGGUCAGCGAU	1936-1958	1773
AD-1571004.1	GCUUCCAGGAGGACAGCAUGU	1970-1990	793	ACAUGCTGUCCUCCUGGAAGCAG	1968-1990	1774
AD-1571005.1	CGUGUCCUGGGCAAGGUGUU	2010-2030	109	AACACCTUGCCCCAGGAACACGGU	2008-2030	235
AD-1571006.1	GGGCAAGGUGUGGCAGAACUU	2019-2039	1682	AAGUTCTGCCACACCUUGCCCCAG	2017-2039	1775
AD-1571007.1	GCAAGGUGUGGCAGAACUCGU	2021-2041	111	ACGAGUTCUGCCACACCUUGGCC	2019-2041	237
AD-1571008.1	CAAGGUGUGGCAGAACUCGU	2022-2042	801	AGCGAGTUCUGCCACACCUUGCC	2020-2042	989
AD-1571009.1	GGCCUGGAGAGGUGUCCUUCU	2045-2065	803	AGAAGGACACCUUCCAGGCCAG	2043-2065	1776
AD-1571010.1	CUGGAGAGGUGUCCUUCAGU	2048-2068	113	ACUUGAAGGACACCUUCCAGGC	2046-2068	239
AD-1571011.1	AGAGGUGUCCUUCAGGUGAU	2052-2072	116	AUCACCTUGAAGGACACCUUCC	2050-2072	1777
AD-1571012.1	GCUACCGCAAGGGCAAGAGU	2363-2383	120	ACUUCUTGCCCCUUGCGGUAGCCG	2361-2383	1778
AD-1571013.1	CUACCGCAAGGGCAAGAGGU	2364-2384	121	ACCUTCTUGCCCCUUGCGGUAGCC	2362-2384	247
AD-1571014.1	ACUACUUCGGCGUCUACACCU	2489-2509	817	AGGUGUAGACGCCGGAAGUAGUUA	2487-2509	1007
AD-1571015.1	CUACUUCGGCGUCUACACCU	2490-2510	818	AGGGTGTAGACGCCGGAAGUAGUU	2488-2510	1779
AD-1571016.1	GCGUCUACACCCGCAUCACAU	2498-2518	819	AUGUGATGCGGGUGUAGACGCCG	2496-2518	1780
AD-1571017.1	CGUCUACACCCGCAUCACAGU	2499-2519	820	ACUGTGAUGCGGGUGUAGACGCC	2497-2519	1781
AD-1571018.1	ACCCGCAUCACAGGUGUGAUU	2506-2526	821	AAUCACACCUUGUGAUGCGGGUGU	2504-2526	1782
AD-1571019.1	GAUCCAGCAAGUGGUGACCUU	2532-2552	824	AAGGTCACCAUUGCUGGAUCCA	2530-2552	1783
AD-1571020.1	GGCAGGAGGUGGCAUCUUGUU	2671-2691	127	AACAAGUAGCCACCUCCUGCCAC	2669-2691	253
AD-1571021.1	UCCCGAUGUCUGCUCCAGUU	2695-2715	832	AACUGGAGCAGACAUCAGGGACG	2693-2715	1022
AD-1571022.1	CUGAUGUCUGCUCCAGUAGUU	2698-2718	1683	AAUACCTGGAGCAGACAUCAGGG	2696-2718	1023
AD-1571023.1	GGCUCAGCAGCAAGAAUGCUU	2853-2873	132	AAGCAUTCUUGCUGCUGAGCCAC	2851-2873	258
AD-1571024.1	UUGGGAUCUGGGGAUUGGAAGU	2983-3003	138	ACUUCCAUUCGCCAGAUCCCAAGU	2981-3003	264
AD-1571025.1	CAGCUGCCCCUUGGAAUAAU	3164-3184	1684	AUUUAUTCCAAAGGGCAGCUGAG	3162-3184	1784

Duplex Name	Sense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:
AD-1571026.1	CUGCCCUUUGGAAUAAAGCUU	3167-3187	144	AAGCTUTAUUCCAAAGGGCAGCU	3165-3187	270
AD-1571027.1	GCCCUUUGGAAUAAAGCUGCU	3169-3189	842	AGCAGCTUUUUUCCAAAGGGCAG	3167-3189	1785
AD-1571028.1	CCUCACCUUGCUUUCUUGGUU	558-578	1685	AACCAGAAAGCAGGUGAGGGG	556-578	1786
AD-1571029.1	CCUCACCUUGCUUUCUUGGUU	558-578	1685	AACCAGAAAGCAGGUGAGGGCU	556-578	1787
AD-1571030.1	UCACCUUGCUUUCUUGGUU	560-578	1686	AACCAGAAAGCAGGUGAGG	558-578	1788
AD-1571031.1	UCACCUUGCUUUCUUGGUU	560-578	1686	AACCAGAAAGCAGGUGACU	558-578	1789
AD-1571032.1	ACCUUGCUUUCUUGGUU	562-578	1687	AACCAGAAAGCAGGUGA	560-578	1790
AD-1571033.1	UCACCUUGCUUUCUUGGUU	560-578	1686	AACCAGAAAGCAGGUGA	560-578	1790
AD-1571034.1	GGAGGUGAUGGCGAGGAAGCU	190-210	1688	AGCUTCCUCGCCAUCACCCUGU	188-210	1791
AD-1571035.1	AAGGCCUGAGGACUCCAAU	229-249	1689	AUUGGAGUCCUCACAGGCCUUGA	227-249	1792
AD-1571036.1	GGCCUGAGGACUCCAAAGAU	231-251	653	AUCUTGGAGUCCUCACAGGCCUU	229-251	1793
AD-1571037.1	GCCUGAGGACUCCAAAGAGU	232-252	20	ACUTUGGAGUCCUCACAGGCCU	230-252	146
AD-1571038.1	CUACUCUGGUUUUUCUAGGU	328-348	25	ACCUAGGAAAUACCCAGAGUAGCA	326-348	151
AD-1571039.1	UCUGGUUUUUCUAGGGUACU	332-352	29	AGUACCCUAGGAAAUACCCAGAGU	330-352	155
AD-1571040.1	CUGGUUUUUCUAGGGUACAU	333-353	1690	AUGUACCCUAGGAAAUACCCAGAG	331-353	1794
AD-1571041.1	UGGUUUUUCUAGGGUACAAU	334-354	1691	AUUGTACCCUAGGAAAUACCCAG	332-354	1795
AD-1571042.1	CUAGGGUACAAGGCGGAGGUU	343-363	662	AACCTCCGCCUUGUACCCUAGGA	341-363	1796
AD-1571043.1	AUGGUCAGCCAGGUGUACUCU	364-384	663	AGAGTACACCCUGGCUGACCAUCA	362-384	1797
AD-1571044.1	GUCAGCCAGGUGUACUCAGGU	367-387	1692	ACCUGAGUACACCCUGGCUGACCA	365-387	1798
AD-1571045.1	CAGCCAGGUGUACUCAGGCAU	369-389	1693	AUGCCUGAGUACACCCUGGCUGAC	367-389	1799
AD-1571046.1	CUCAUUGGCCACUUCUCCCAU	400-420	667	AUGGGAGAAGUGGCGAUUGAGUA	398-420	1800
AD-1571047.1	CGCCACUUCUCCCAAGGAUCUU	406-426	668	AAGATCCUGGGAGAAAGUGGCGAU	404-426	1801
AD-1571048.1	GCCACUUCUCCCAAGGAUCUUU	407-427	33	AAAGAUCCUGGGAGAAAGUGGCGA	405-427	159
AD-1571050.1	UCCCAGGAUCUUACCCGCCGU	415-435	35	ACGGCGGUUAGAUCUCCUGGGAGA	413-435	161
AD-1571051.1	UAGUGCCUUCUCCGACAGUAGAAU	441-461	1694	AUUUCACUGCGGAAGGCACUAGA	439-461	1802
AD-1571052.1	CUUCCGCAGUGAUAACCCGCCAU	447-467	676	AUGGCGGUUUUCACUGCGGAAGGC	445-467	1803

Duplex Name	Sense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:
AD-1571053.1	CCGCAGUGAAACCGCCAAAGU	450-470	677	ACUUTGGCGGUUUUACUGCGGAA	448-470	1804
AD-1571054.1	CGCAGUGAAACCGCCAAAGCU	451-471	678	AGCUTUGGCGGUUUUACUGCGGA	449-471	1805
AD-1571055.1	GCCAAAGCCCAAGAAAGUCUU	463-483	683	AAGCAUCUUCUGGGCUUUGCGGG	461-483	880
AD-1571056.1	CAGCACCGCGCUGGGAAACUUU	498-518	685	AAAGTUUCCAGCGGGUGUGGU	496-518	1806
AD-1571057.1	ACAAACUCCAGCUCGCCGUCUAUU	521-541	687	AAUAGACGGAGCUGGAGUUGUAG	519-541	884
AD-1571058.1	CACCUUGCUUUCUGGUUCAU	561-581	1695	AUGAACCCAGAAAGCAGGUGAG	559-581	1807
AD-1571059.1	UGCUUCUUCUGGUUCAUUCUU	565-585	43	AAGAAUGAACCCAGAAAGCAGG	563-585	169
AD-1571060.1	UCUUCUGGUUCAUUCUCCAAU	569-589	1696	AUUGGAGAAUGAACCCAGAAAG	567-589	1808
AD-1571061.1	UUCUGGUUCAUUCUCCAAAUU	571-591	1697	AAUUTGGAGAAUGAACCCAGAA	569-591	1809
AD-1571062.1	UCUGGUUCAUUCUCCAAAUUCU	572-592	1698	AGAUTUGGAGAAUGAACCCAGAA	570-592	1810
AD-1571063.1	GUUGAGGAGCUGCUGUCCACU	643-663	1699	AGUGGACAGCAGCUCUCCACCA	641-663	1811
AD-1571064.1	GAGGAGCUGCUGUCCACAGUU	646-666	1700	AACUGUGGACAGCAGCUCUCCCA	644-666	894
AD-1571065.1	AGCUGCUGUCCACAGUCAACU	650-670	1701	AGUUGACUGUGGACAGCAGCUC	648-670	895
AD-1571066.1	CUGUCCACAGUCAACAGCUCU	655-675	1702	AGAGCUGUUGACUGUGGACAGCA	653-675	898
AD-1571067.1	ACAGGGCCGAGUACGAAGUGU	689-709	48	ACACTUCGUACUCGGGCCUUGUAG	687-709	1812
AD-1571068.1	GGGCCGAGUACGAAGUGGACU	692-712	1703	AGUCCACUUCGUACUCGGCCUUG	690-712	1813
AD-1571069.1	UCCUGGAAGCCAGUGUGAGAAU	728-748	710	AUUUCACACUGGGCUUCCAGGAUC	726-748	1814
AD-1571070.1	CUGGAAGCCAGUGUGAGAAAGAU	730-750	712	AUCUTUCACACUGGGCUUCCAGGA	728-750	1815
AD-1571071.1	AAGCCAGUGUGAGAAAGACAUAU	734-754	716	AU AUGUCUUUACACACUGGCUUCC	732-754	1816
AD-1571072.1	GCCAGUGUGAGAAAGACAUAGCU	736-756	718	AGCUAUGUCUUUACACACUGGCUU	734-756	1817
AD-1571074.1	UGUGAAAGACAUAGCUGCAUU	741-761	721	AAUGCAGCUAUGUCUUUUCACACU	739-761	916
AD-1571075.1	ACGCUUGGUUGUUACCGCUAU	769-789	1704	AUAGCGUAAACAAACCCAGCGUGG	767-789	1818
AD-1571076.1	CGCUGGGUUGUUACCGCUACU	770-790	1705	AGUAGCGGUAAACAAACCCAGCGUG	768-790	919
AD-1571077.1	GCUGGGUUGUUACCGCUACAU	771-791	1706	AUGUAGCGGUAAACAAACCCAGCGU	769-791	1819
AD-1571078.1	CUGGGUUGUUACCGCUACAGU	772-792	59	ACUGTAGCGGUAAACAAACCCAGCG	770-792	185
AD-1571079.1	CUGGAGAAAGAGGCUCAUCACU	958-978	733	AGUGAUGAGGCCUCUCUCCAGGG	956-978	1820

Duplex Name	Sense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:
AD-1571080.1	GAGAAAGGGCUCAUACCCUCU	961-981	1707	AGAGGUGAUGAGCCUUCUCCCA	959-981	930
AD-1571081.1	GAAAGAGGCUCACACCCUCGGU	963-983	1708	ACCGAGGUGAUGAGCCUUCUCUC	961-983	931
AD-1571082.1	AGGCUCAUCACCCUCGGUGUAU	967-987	67	AUACACCGAGGUGAUGAGCCUCU	965-987	1821
AD-1571083.1	AAGAAGGGCCUGCACAGCUAU	1054-1074	1709	AUAGCUGUGCAGGCCUUCUUC	1052-1074	1822
AD-1571084.1	AAGGGCCUGCACAGCUACUAU	1057-1077	740	AUAGTAGCUGUGCAGGCCUUCU	1055-1077	1823
AD-1571085.1	CCUCUCUGGACUACGGCUUGU	1235-1255	70	ACAAGCCGUAUCCAGAGAGGGC	1233-1255	1824
AD-1571086.1	UCUCUGGACUACGGCUUGGCU	1237-1257	71	AGCCAAGCCGUAUCCAGAGAGG	1235-1257	197
AD-1571087.1	UGGACUACGGCUUGGCCUCUCU	1241-1261	743	AGAGGGCCAAGCCGUAUCCAGA	1239-1261	938
AD-1571088.1	GGACUACGGCUUGGCCUCUCU	1242-1262	1710	AAGAGGGCCAAGCCGUAUCCAG	1240-1262	939
AD-1571089.1	AGAAUAUGAUUUUUGCCGUGCU	1289-1309	83	AGCACGGCAAAUCAAUACUUCGC	1287-1309	1825
AD-1571090.1	GAAUAUGAUUUUUGCCGUGCAU	1290-1310	84	AUGCACGGCAAAUCAAUACUUCUG	1288-1310	1826
AD-1571091.1	AAGUAUGAUUUUUGCCGUGCACU	1291-1311	1711	AGUGCACGGCAAAUCAAUACUUCU	1289-1311	1827
AD-1571092.1	GUUAUAUUUUGCCGUGCACCCU	1293-1313	1712	AGGGTGACGGCAAAUCAAUACUU	1291-1313	1828
AD-1571093.1	GGCCAGUGGACGAUCCAGAAU	1315-1335	751	AUUCTGGAUCGUCCACUGGCCCU	1313-1335	1829
AD-1571094.1	GCCAGUGGACGAUCCAGAAACU	1316-1336	752	AGUUCUGGAUUGUCCACUGGCC	1314-1336	945
AD-1571096.1	UGGACGAUCCAGAACAGGAGU	1321-1341	755	ACUCCUGUUCUGGAUUGUCCACU	1319-1341	948
AD-1571097.1	CACCUCCAGAUUCCUCCAU	1419-1439	757	AUGAGGGAGAUUUGGGAGGUGAA	1417-1439	1830
AD-1571098.1	UGCGGGUGCACUAUGGCUUGU	1451-1471	759	ACAAGCCAUAGUGCACCCGCACA	1449-1471	1831
AD-1571099.1	GCGGGUGCACUAUGGCUUGUU	1452-1472	91	AACAAGCCAUAGUGCACCCGCAC	1450-1472	217
AD-1571100.1	CGGGUGCACUAUGGCUUGUAU	1453-1473	761	AUACAAGCCAUAGUGCACCCGCA	1451-1473	1832
AD-1571102.1	AACGGCCUGGAUGAGAGAAAU	1561-1581	767	AUUUCUCUACUCCAGGCCGUUGG	1559-1581	1833
AD-1571103.1	CGGCCUGGAUGAGAGAAACUU	1563-1583	769	AAGUTUCUCUCAUCCAGGCCGUU	1561-1583	1834
AD-1571104.1	GAGAAACUGCGUUGCAGAGU	1575-1595	772	ACUCTGCAACGCAGUUUCUCUC	1573-1595	1835
AD-1571105.1	CUGCGUUUGCAGAGCCACAUU	1581-1601	773	AAUGTGGCUCUGCAACGCAGUU	1579-1601	1836
AD-1571106.1	UGCGUUUGCAGAGCCACAUUU	1582-1602	774	AAAUGUGGCUUGCAACGCAGU	1580-1602	1837
AD-1571107.1	CGUUUGCAGAGCCACAUUCCU	1584-1604	776	AGGAAUGUGGCUUGCAACGCACA	1582-1604	1838

Duplex Name	Sense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:
AD-1571108.1	GCAGAGCCACAUUCCAGUGCU	1589-1609	777	AGCACUGGAAUGUGGCUCUGCAA	1587-1609	968
AD-1571109.1	GGGACAUUCCACCUUCCAGUGU	1711-1731	779	ACACTGGAAGGUGAAUGUCCAC	1709-1731	1839
AD-1571110.1	GGACAUUCCACCUUCCAGUGUU	1712-1732	103	AACACUGGAAGGUGAAUGUCCCA	1710-1732	229
AD-1571111.1	ACAUUCCACCUUCCAGUGUGAU	1714-1734	105	AUCACACUGGAAGGUGAAUGUCC	1712-1734	1840
AD-1571112.1	AGCUGCGUGAAGAAAGCCCAAU	1741-1761	782	AUUGGGCUUUCUACACGCAGCUCC	1739-1761	1841
AD-1571113.1	GCUGCGUGAAGAAGCCCAACU	1742-1762	783	AGUUUGGCUUUCUACACGCAGCUC	1740-1762	1842
AD-1571114.1	CUGCGUGAAGAAGCCCAACCU	1743-1763	784	AGGUTGGGCUUUCUACACGCAGCU	1741-1763	1843
AD-1571115.1	UGCGUGAAGAAGCCCAACCCU	1744-1764	785	AGGGTUGGGCUUUCUACACGCAGC	1742-1764	1844
AD-1571116.1	AGCACUGUGACUGUGGCCUCU	1808-1828	786	AGAGGCCACAGUCACAGUCCUCC	1806-1828	1845
AD-1571117.1	CUCCGAGGUGAGUGGCCAUU	1866-1886	787	AAUGGCCACUCACCCUCGGAGGA	1864-1886	976
AD-1571118.1	AUCGUGACCGCUGGGUGAUU	1936-1956	107	AAUACCCACAGCGGUACAGGAUGA	1934-1956	233
AD-1571119.1	UCGUGACCGCUGGGUGAUU	1937-1957	788	AUAUACCCACAGCGGUACAGGAUG	1935-1957	1846
AD-1571120.1	GCUGACCGCUGGGUGAUU	1939-1959	790	AGUUUACACCCACAGCGGUACGCGA	1937-1959	980
AD-1571121.1	GCUGGGUGAUUACAGCUGCCU	1946-1966	791	AGGACGUGUUUACACCCACGCGG	1944-1966	981
AD-1571122.1	UGCUUCCAGGAGGACAGCAUU	1969-1989	792	AAUGCUGUCCUCCUGGAAAGCAGU	1967-1989	1847
AD-1571123.1	CUUCCAGGAGGACAGCAUGGU	1971-1991	794	ACCATGCUGUCCUCCUGGAAAGCA	1969-1991	1848
AD-1571124.1	CUGGGCAAGGUGUGGCAGAAU	2017-2037	1713	AUUCTGCCACACCUUUGCCCAGGA	2015-2037	1849
AD-1571125.1	UGGGCAAGGUGUGGCAGAAU	2018-2038	797	AGUUCUGCCACACCUUUGCCCAGG	2016-2038	986
AD-1571126.1	GGCAAGGUGUGGCAGAAACUCU	2020-2040	799	AGAGTUCUGCCACACCUUUGCCCA	2018-2040	1850
AD-1571127.1	UGGCCUGGAGAGGUGUCCUUU	2044-2064	802	AAAGGACACCUUCCAGGCCAGC	2042-2064	990
AD-1571128.1	GCCUGGAGAGGUGUCCUUCAU	2046-2066	804	AUGAAGGACACCUUCCAGGCCA	2044-2066	1851
AD-1571129.1	CCUGGAGAGGUGUCCUUCAAU	2047-2067	1714	AUUGAAGGACACCUUCCAGGCC	2045-2067	1852
AD-1571130.1	UGUGCAGUUGAUCCACAGGU	2289-2309	1715	ACCUUGGGGAUCAACUGCACAU	2287-2309	994
AD-1571131.1	UGCAGUUGAUCCACAGGACU	2291-2311	808	AGUCCUGUGGGGAUCAACUGCACA	2289-2311	995
AD-1571132.1	AUCCACAGGACCUUGUGCAGU	2299-2319	117	ACUGCACAGGUCCUGUGGGAUCA	2297-2319	243
AD-1571133.1	CCCACAGGACCUUGUGCAGCGU	2301-2321	809	ACGCTGCACAGGUCCUGUGGGAU	2299-2321	1853

Duplex Name	Sense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:
AD-1571134.1	CCACAGGACCUUGUGCAGCGAU	2302-2322	1716	AUCGUGCAGCAGGUCUUGGGGA	2300-2322	1854
AD-1571135.1	CCAGGUGACGCCACGCAUGCU	2334-2354	811	AGCATGCGUGGCGUACCCUGGUA	2332-2354	1855
AD-1571136.1	GUGACGCCACGCAUGCUGUGU	2338-2358	812	ACACAGCAUGCGUGGCGUCACCU	2336-2358	1000
AD-1571137.1	UGAGGCCACGCAUGCUGUGUU	2339-2359	118	AACACAGCAUGCGUGGCGUACCC	2337-2359	244
AD-1571138.1	ACGCCACGCAUGCUGUGUGCU	2341-2361	1717	AGCACACAGCAUGCGUGGCGUCA	2339-2361	1856
AD-1571139.1	GGCUACCGCAAGGGCAAGAAU	2362-2382	814	AUUCTUGCCCUUUGCGGUAGCCGG	2360-2382	1857
AD-1571140.1	GUGCAAGGCACUCAGUGGCCU	2418-2438	815	AGCCACUGAGUGCCUUGCACCAC	2416-2438	1858
AD-1571141.1	CUAAACUACUUCGGCGUCUACU	2486-2506	1718	AGUAGACGCCGAAGUAGUAGGC	2484-2506	1006
AD-1571142.1	CUACACCCGCAUCACAGGUGU	2502-2522	124	ACACUGUGAUGCGGGUGUAGAC	2500-2522	250
AD-1571143.1	CCCGCAUCACAGGUGUGAUCU	2507-2527	822	AGAUCACACCUUGAUGCGGGUG	2505-2527	1012
AD-1571144.1	UGGAUCCAGCAAGUGGUGACU	2530-2550	823	AGUACCAUUGCUGGUAUCCAGC	2528-2550	1859
AD-1571145.1	AUCCAGCAAGUGGUGACCUGU	2533-2553	1719	ACAGGUCACCAUUGCUGGAUCC	2531-2553	1860
AD-1571146.1	CCAGCAAGUGGUGACCUGAGU	2535-2555	1720	ACUCAGGUCACCAUUGCUGGAU	2533-2555	1016
AD-1571147.1	CAGCAAGUGGUGACCUGAGGU	2536-2556	1721	ACCUCAGGUCACCAUUGCUGGA	2534-2556	1017
AD-1571148.1	GCAAUGUGGUGACCUGAGGAAU	2538-2558	1722	AUUCUACAGGUCACCAUUGCUG	2536-2558	1861
AD-1571149.1	UGGUGGCAGGAGGUGGCAUCU	2667-2687	829	AGAUGCCACCUCCUGCCACCACA	2665-2687	1862
AD-1571150.1	GGUGGCAGGAGGUGGCAUCUU	2668-2688	830	AAGATGCCACCUCCUGCCACCAC	2666-2688	1863
AD-1571151.1	GUGGCAGGAGGUGGCAUCUUU	2669-2689	831	AAAGAUGCCACCUCCUGCCACCA	2667-2689	1021
AD-1571152.1	UCCAGUGAUGGCAGGAGGAUU	2709-2729	1723	AAUCCUCCUGCCAUACUGGAGC	2707-2729	1864
AD-1571153.1	CUAAUUGGGAUCUGGGAAUU	2978-2998	835	AAUUCUCCAGAUCCCAAGUUAGAC	2976-2998	1025
AD-1571154.1	GUGAGCUCAGCUGCCCCUUUGU	3157-3177	837	ACAAAGGCAGCUGAGCUCACCU	3155-3177	1026
AD-1571155.1	CUCAGCUGCCCCUUUGGAAU	3162-3182	838	AUAUTCCAAAGGCAGCUGAGCU	3160-3182	1865
AD-1571156.1	UCAGCUGCCCCUUUGGAAUAAU	3163-3183	1724	AUUATUCCAAAGGCAGCUGAGC	3161-3183	1866
AD-1571157.1	CCUUUUGGAAUAAAGCUGCCU	3170-3190	1725	AGGCAGUUUAUCCAAAGGGCA	3168-3190	1867
AD-1571158.1	CCUUUGGAAUAAAGCUGCCUU	3171-3191	844	AAGGCAGCUUUAUCCAAAGGGC	3169-3191	1868
AD-1571159.1	UUUGGAAUAAAGCUGCCUGAU	3173-3193	845	AUCAGGCAGCUUUAUCCAAAGG	3171-3193	1869

Duplex Name	Sense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	Range in NM_153609.4	SEQ ID NO:
AD-1571160.1	UUGGAAUAAAGCUGCCUGAUU	3174-3194	846	AAUCAGGCAGCUUUAUUCCAAAG	3172-3194	1870
AD-1571161.1	UGGAAUAAAGCUGCCUGAUCU	3175-3195	847	AGAUCAGGCAGCUUUAUUCCAA	3173-3195	1871

Table 7. Modified Sense and Antisense Strand Sequences of TMPRSS6 dsRNA Agents

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1570929.1	csgsgaggUfgAfUfgfcgaggaagL96	1872	asCfsuudCc(Tgn)cgccauCfaCfucecgsc	2099	GACGGAGGUGAUGGCGAGGA	1491
AD-1570930.1	cscsugugAfgGfAfCfucaagagauL96	1873	asUfscudCu(Tgn)ggagucCfuCfacaggscsc	2100	GGCCUGUGAGGACUCCAAAGAG	1495
AD-1570931.1	csusgugaGfgAfCfUfccaaagaaL96	1874	asUfsuedTfc(Tgn)uggaguCfcUfcacagsgsc	2101	GCUGUGAGGACUCCAAAGAGA	1496
AD-1570932.1	csuseuggUfaUfUfUfccuaggguauL96	1875	asUfsaedCCc(Tgn)aggaaaUfaCfcagagsusa	2102	UACUCUGGUUUUCCUAGGG	532
AD-1570933.1	gsgsuauUfcCfUfAfggguacaagL96	1876	asCfsuudGu(Agn)cccuagGfaAfauiaccsag	2103	CUGGUUUUCCUAGGGUACA	1499
AD-1570934.1	gsusauuuCfcUfAfGfgguacaagL96	1877	asCfscudTgc(Tgn)accuaGfgAfauiaccsa	2104	UGGUUUUCCUAGGGUACAA	1500
AD-1570935.1	gsgsucagCfcAfGfGfuguaucagL96	1878	asCfsugdAg(Tgn)acaccuGfgCfugaccsas	2105	AUGGUCAGCCAGGUACUCA	535
AD-1570936.1	uscsagccAfgGfUfGfuaucaaggcuL96	1879	asGfscddTgc(Agn)guacaeCfuGfgeugascsc	2106	GGUCAGCCAGGUUACUCAGG	1504
AD-1570937.1	asgsccagGfuGfUfAfcuacaggcuL96	1880	asCfsugdCc(Tgn)gaguacAfcCfuggcuagsga	2107	UCAGCCAGGUUACUCAGGCA	536
AD-1570938.1	csaseuuUfcCfCfAfGgaucuuacuL96	1881	asGfsuadAg(Agn)uccuggGfaGfaagugsgs	2108	GCCACUUCUCCCAAGGAUCUUA	1508
AD-1570939.1	uscsuuccAfgGfAfUfUfcuuaccgcguL96	1882	asGfscgdGg(Tgn)aagaucCfuGfaggagasag	2109	CUUCUCCCAAGGAUCUUAACCCG	1509
AD-1570940.1	gseseuuCfGfCfAfGfugaaccgcguL96	1883	asGfscgdGu(Tgn)uacugCfGfaggacsasc	2110	GUGCCUUCGCGAGUGAAACCG	540

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1570941.1	cscsuuccGfcAfGfUfgaaaccgccuL96	1884	asGfsgcdGg(Tgn)uucacuGfcGfgaaggscsa	2111	UGCCUUCCGCAGUGAAAAACCGC CA	1512
AD-1570942.1	gscsagugAfaAfCfCfccaagaagccuL96	1885	asGfsgcdTu(Tgn)ggggguUfuCfacuagcsgsg	2112	CCGCAGUGAAAAACCGCCAAAGC CC	1516
AD-1570943.1	csasgugaAfaCfCfGfccaaagccuL96	1886	asGfsggdCu(Tgn)ugggggUfuUfcaugscsg	2113	CGCAGUGAAAAACCGCCAAAGCC CA	1517
AD-1570944.1	asgsugaaAfcCfGfCfcaaaagccuL96	1887	asUfsggdGc(Tgn)uuggcgGfuUfuaacugsc	2114	GCAGUGAAAAACCGCCAAAGCCC AG	1518
AD-1570945.1	csgsccaaAfgCfCfCfagaagaugcuL96	1888	asGfscadTc(Tgn)ucugggCfuUfugggcgsgsu	2115	ACCGCCAAAGCCAGAAAGAUG CU	1519
AD-1570946.1	asgscccaGfaAfGfAfugcucaagguL96	1889	asCfscudTg(Agn)gcaucuUfcUfggcgususu	2116	AAAGCCCAGAAAGAUGCUCAAG GA	1521
AD-1570947.1	asgsccaccCfGfCfCfUfgggaacuuuL96	1890	asUfsgaadGu(Tgn)cccaggCfGfGfugcugsgsg	2117	CCAGCACCCGCCUGGGAAACUU AC	1523
AD-1570948.1	csasacucCfaGfCfUfcgucuaauuL96	1891	asAfsaudAg(Agn)cggagcUfgGfaguugsus a	2118	UACAAUCCAGCUCCGUCUCU UC	541
AD-1570949.1	uscsaccuGfcUfUfCfuucugguuL96	1892	asGfsgaadCc(Agn)gaagaaGfcAfggugagsgsg	2119	CCUCACCUGCUUCUUCUGGUU CA	544
AD-1570950.1	cscsugcuUfcUfUfCfugguucauuuL96	1893	asAfsaudGa(Agn)ccagaaGfaAfgcaggsgsg	2120	CACCUGCUUCUUCUGGUUUCU UC	546
AD-1570951.1	csusgcuUfUfCfUfgguucauuL96	1894	asGfsgaadTg(Agn)accagaAfgAfgcagsgsu	2121	ACCUGCUUCUUCUGGUUUCU CU	1526
AD-1570952.1	csusucuuCfuGfGfUfucuuuL96	1895	asGfsggadGa(Agn)uugaaccAfgAfgaagsgsa	2122	UGCUCUUCUGGUUUCUUCUC CA	549
AD-1570953.1	ususeuuUfgGfUfUfcauuuL96	1896	asUfsggdAg(Agn)augaaacCfaGfaagaagsg	2123	GCUUCUUCUGGUUUCUUCUCC AA	550
AD-1570954.1	csusucugGfuUfCfAfuuccaaauL96	1897	asUfsguudGg(Agn)gaugaAfcCfagaagsgasa	2124	UUCUUCUGGUUUCUUCUCCAA AU	1528
AD-1570955.1	csusggguCfaUfUfCfucaaaauL96	1898	asGfsggadTu(Tgn)ggagaaUfgAfacagsgasa	2125	UUCUGGUUUCUUCUCCAAUUC CC	551
AD-1570956.1	gscsugcuGfuCfCfAfcagucacauL96	1899	asUfsguudTg(Agn)cuguggAfcAfgcagcsusc	2126	GAGCUGCUGUCCACAGUCAAC AG	1534
AD-1570957.1	gscsugucCfaCfAfGfuaacaaguuL96	1900	asAfsgcdTg(Tgn)ugacugUfgGfacaagsgsg	2127	CUGCUGUCCACAGUCAACAGC UC	1535

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1570958.1	usgsuuccaCfaGfUfCfaacagcugcuL96	1901	asCfsgadGc(Tgn)guugacUfgUfggacagsc	2128	GCUGUCCACAGUCAACAGCUC	1537
AD-1570959.1	gsgsccgaGfuAfCfCfagaggaccuL96	1902	asGfsgudCc(Agn)cucucguAfcUfcggccscsu	2129	AGGGCCGAGUACGAAGUGGA	1539
AD-1570960.1	asusccugGfaAfAfGfCfcagugagaauL96	1903	asUfsuudAc(Agn)cuggcuUfcCfaggaucesca	2130	UGAUCCUGGAAAGCCAGUGUG	1540
AD-1570961.1	cscsuggaAfgCfCfAfagugagaaguL96	1904	asCfsuudTc(Agn)cacuggCfuUfcagggsasu	2131	AUCCUGGAAGCCAGUGUGAA	1542
AD-1570962.1	usgsaaagCfcAfGfUfgugaagaagacuL96	1905	asGfsgudTu(Tgn)cacacuGfgCfuuccasgsg	2132	CCUGGAAGCCAGUGUGAAAAAG	1544
AD-1570963.1	gsgsaagcCfaGfUfGfGfagaagacaauL96	1906	asUfsgudCu(Tgn)ucacacUfgGfueuccasag	2133	CUGGAAGCCAGUGUGAAAAAGA	1545
AD-1570964.1	gsasagccAfgUfGfUfgaagaagacaauL96	1907	asAfsugdTc(Tgn)uucacaCfuGfgcuuccesca	2134	UGGAAGCCAGUGUGAAAAAGAC	1546
AD-1570965.1	asgsccagUfgUfGfAfagaagacaauL96	1908	asCfsuadTg(Tgn)cuuucaCfaCfuggcususc	2135	GAAGCCAGUGUGAAAAAGACAU	1548
AD-1570966.1	cscsagugUfgAfAfAfagacaauL96	1909	asAfsugdTa(Tgn)gucuuuCfaCfagugscsu	2136	AGCCAGUGUGAAAAAGACAUAG	554
AD-1570967.1	asgsugugAfaAfGfAfafcauagcuguL96	1910	asGfseadGc(Tgn)auguauUfuCfacausgsg	2137	GCA	1550
AD-1570968.1	gsusgaaaGfaCfAfUfagcugcauuuL96	1911	asAfsaudGc(Agn)gcuauuUfcUfuuacacsac	2138	GUGUGAAAGACAUAGCUGCA	1553
AD-1570969.1	asusugaaUfuCfCfAfagcugggguuL96	1912	asAfsacdCc(Agn)gcguggaAfaUfuaaungsc	2139	GCAUUGAAUUCCACGCUGGGU	556
AD-1570970.1	asasuuccAfcGfCfUfaggguuguuuauL96	1913	asUfsaadCa(Agn)cccagcGfuCfagaauuesca	2140	UGAAUUCCACGCUGGGGUUGU	559
AD-1570971.1	csasegcuGfgGfUfUfguuaccgcuL96	1914	asAfsgcdGg(Tgn)aaacaacCfcAfagcuggsa	2141	UACCACGCUGGGGUUGUUACCCG	562
AD-1570972.1	usgsgguuGfuUfAfCfegcuacagcuL96	1915	asGfseudGu(Agn)gggguaAfcAfaccacsagc	2142	CU	1557
AD-1570973.1	gsgsguuuUfuAfCfCfagcuacagcuL96	1916	asAfsgcdTg(Tgn)agcggguAfaCfaaccacsag	2143	CUGGGUUGUUACCCGCUACAGC	1558
AD-1570974.1	csasaacuCfcGfGfCfagagagcuL96	1917	asUfsecdAc(Tgn)ccagccGfgAfguuungasag	2144	CUCAAACUCCGGCUGGAGUGG	1559

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1570975.1	gsgsgaacGfaCfaUfGfccaugauuL96	1918	asAfsuadCa(Tgn)ggccagUfcGfguccesgsg	2145	CCGGACCGACUGGCCAUGUA	564
AD-1570976.1	csgsacugGfcCfAfUfguaagacguL96	1919	asAfsagdTc(Agn)uacaugGfcCfagucgsgsu	2146	ACCGACUGGCCAUGUAUGACG	1560
AD-1570977.1	usgsagaaAfgAfGfGfcauacacuuL96	1920	asGfsgudGa(Tgn)ggcccuCfuUfuccasgsg	2147	CCUGGAGAAAGAGGCUCAUCAC	1562
AD-1570978.1	gsgsagaaGfaGfGfCfcauacacuuL96	1921	asAfsggdTg(Agn)ugagccUfcUfuccesasg	2148	CUGGAGAAAGAGGCUCAUCACC	1563
AD-1570979.1	gsasagaaGfgCfCfGfcagacacuuL96	1922	asAfsagdTg(Tgn)gcaggcCfcUfuccesasa	2149	UGGAAGAAGGGCCUGCACACAGC	1566
AD-1570980.1	asgsggcccUfcCfAfCfagcuacuL96	1923	asGfsuadGu(Agn)gcugugCfaGfgccucusc	2150	GAAGGGCCUGCACAGCUACUA	1569
AD-1570981.1	cscsugcaCfaGfCfUfcauacgacuuL96	1924	asGfsgudCg(Tgn)aguagcUfgUfagcagcsc	2151	GGCCUGCACAGCUACUACGAC	573
AD-1570982.1	gsasgggagGfcAfGfAfaguaugauuL96	1925	asAfsaudCa(Tgn)acuuccUfcCfuccuesag	2152	CUGAGGAGGCAGAAAGUAUGA	580
AD-1570983.1	asgsgagggCfaGfAfAfuaugauuuuL96	1926	asAfsaadTc(Agn)uacuuCfGfCfuccucasa	2153	UGAGGAGGCAGAAAGUAUGAU	1572
AD-1570984.1	asgsuauGfAfUfUfGfcegugcacuuL96	1927	asGfsgudGc(Agn)cggcaaAfuCfuaucususc	2154	GAAGUAUGAUUUUGCCGUGCA	1574
AD-1570985.1	cscsagugGfaGfGfAfuccagacauL96	1928	asUfsgudTc(Tgn)ggauccUfcCfagucgscsc	2155	GGCCAGUGGACGAUCCAGAAC	1578
AD-1570986.1	cscsagaaCfaGfGfAfagcuguguuL96	1929	asCfsacdAc(Agn)gcuccUfgUfucuggsasu	2156	AUCCAGAACAGGAGGCUGUG	591
AD-1570987.1	asgsaacaGfgAfGfGfcegugugguL96	1930	asGfscddAc(Agn)cagccuCfcUfguuucsgsg	2157	CCAGAACAGGAGGCUGUGUG	592
AD-1570988.1	asesuucaCfcUfCfCfcauuccuuL96	1931	asGfsgadGa(Tgn)cuagggaGfgUfgaagusug	2158	CAACUUCACCUCCAGACUUCUC	1580
AD-1570989.1	usgsugcgGfgUfGfCfcauugggcuL96	1932	asAfsagdCa(Tgn)agugcaCfcCfcaacascsc	2159	GGUGUGCGGGUGCACUAUGG	593
AD-1570990.1	gsusgcggGfuGfCfAfcauagggcuuL96	1933	asAfsagdCc(Agn)uagugcAfcCfcaacascsc	2160	GUGUGCGGGUGCACUAUGGC	594
AD-1570991.1	gsgsgugcAfcUfAfAfUfggcuuquacL96	1934	asGfsuadCa(Agn)gccaaGfuGfcaaccsgsc	2161	GCGGGUGCACUAUGGCUUGU	596

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1570992.1	gsgsugcaCfuAfUfGfgcuuguaacaul96	1935	asUfsgudAc(Agn)agcccaAfuUfGfcaaccscg	2162	CGGGUGCACUAUGGCUUGUACAA	1584
AD-1570993.1	usgseacuAfuGfGfCfuuguaacaul96	1936	asGfssuudGu(Agn)caagccAfuAfuGfcaaccsc	2163	GGUGCACUAUGGCUUGUACAACC	1585
AD-1570994.1	gscsacuaUfgGfCfUfuguaacaul96	1937	asGfsgudTg(Tgn)acaagcCfaUfagugcsasc	2164	GUGCACUAUGGCUUGUACAACCA	1586
AD-1570995.1	csusgcccUfgGfAfGfaguuccuacaul96	1938	asAfsagadGg(Agn)acucucCfaGfagcagsgsg	2165	CCUGCCCCUGGAGAGAUUCCUCUG	599
AD-1570996.1	ascsgggcUfgGfAfUfagagaaacaul96	1939	asGfssuudTc(Tgn)cucaucCfaGfagcagsgsg	2166	CAACGGCCUGGAGAGAGAAACU	1589
AD-1570997.1	gscsugggAfuGfAfGfagagaaacaul96	1940	asGfscadGu(Tgn)ucucucAfuCfagcagsgsg	2167	CGCCUGGAGAGAGAGAAACUGCG	600
AD-1570998.1	cscsuggaUfgAfuGfAfgaaacugcual96	1941	asCfsgadAg(Tgn)uucucuCfaUfcaagsgsc	2168	GGCCUGGAGAGAGAGAAACUGCGU	1591
AD-1570999.1	asgsagaaAfcUfGfCfguuugcagaul96	1942	asUfscudGc(Agn)aacgcaGfuUfucucscsa	2169	UGAGAGAAACUGGCUUUGCAGGAG	1592
AD-1571000.1	gscsguuuGfcAfGfAfgccacauucul96	1943	asGfssuudTg(Tgn)ggcucuGfcAfaagcsasg	2170	CUGCGUUUGCAGAGCCACAUUCC	1596
AD-1571001.1	usgsagacAfuUfCfAfcuuccagaul96	1944	asAfsuudGg(Agn)aggugaAfuGfucuccasca	2171	UGUGGGACAUUACCUUCCAGUG	606
AD-1571002.1	gsasgcugCfGfUfGfAfgaagcccaul96	1945	asUfsgudGc(Tgn)uucucaCfGfagcagsgsg	2172	CGGAGCUGCGUGAAGAAAGCCC	1600
AD-1571003.1	csgscugaCfcGfCfUfGfGfagaaacaul96	1946	asUfssuudTc(Agn)cccagcGfGfagcagsgsas	2173	AUCGUGACCGCUGGGUGUAUAC	1608
AD-1571004.1	gscsuuccAfuGfAfGfagagcaugul96	1947	asCfssuudGc(Tgn)gucucCfuGfagagcsasg	2174	CUGCUUCCAGGAGGACAGCAU	1612
AD-1571005.1	csgsuguuCfcUfGfGfcaagguugul96	1948	asAfsuudCc(Tgn)ugcccaGfGfagcagsgsu	2175	ACCGUGUCCUGGGCAAGGUGUG	613
AD-1571006.1	gsgsgcaaGfGfUfGfUfGfagaaacaul96	1949	asAfsuudTc(Tgn)gcccacaCfcUfugcccsasg	2176	CUGGGCAAGGUGUGGCGAGAA	1616
AD-1571007.1	gscsaaggUfgUfGfGfagaaacugul96	1950	asCfsgadGu(Tgn)cuugccaCfaCfuuugscsc	2177	GGGCAAGGUGUGGCGAGAAACU	615
AD-1571008.1	csasagguGfuGfCfGfagaaacugul96	1951	asGfscgAdAg(Tgn)uucugcAfcAfcuuugscsc	2178	GGCAAGGUGUGGCGAGAAACUCGCG	1618

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1571009.1	gsgscugGfaGfAfGfguguccuulL96	1952	asGfsaadGg(Agn)caccucUfcCfagggccsasg	2179	CUGGCCUGGAGAGGUGUCCUU CA	1620
AD-1571010.1	csusggagAfGfUfGfuccuucaguuL96	1953	asCfsuudGa(Agn)ggagacCfuCfuccagsgc	2180	GCUGGAGAGGUGUCCUUCAA GG	617
AD-1571011.1	asgsagguGfuCfCfUfucagggugauL96	1954	asUfscadCc(Tgn)ugaaggAfcAfcucuscsc	2181	GGAGAGGUGUCCUUCAAAGGU GAG	620
AD-1571012.1	gscsuaccGfcAfAfGfggcaagaaguuL96	1955	asCfsuudCu(Tgn)gcccuuGfcGfugagcscsg	2182	CGCUACCGCAAGGGCAAGAA GG	624
AD-1571013.1	csusaccgCfaAfGfGfgcaagaaguuL96	1956	asCfsuudTc(Tgn)ugcccuUfgCfugagsgsc	2183	GGCUACCGCAAGGGCAAGAA GG	625
AD-1571014.1	ascsuacuUfcGfGfCfucuaacuuL96	1957	asGfsugudGu(Agn)gacgccGfaAfugauususa	2184	UAACUACUUCGGCGUACAC CC	1633
AD-1571015.1	csusacuUfcGfGfCfucuaacuuL96	1958	asGfsugdTg(Tgn)agacgcCfGfAfguaguuuu	2185	AACUACUUCGGCGUACAC CG	1634
AD-1571016.1	gscsguacuAfcAfCfCfcaucacuuL96	1959	asUfsgudGa(Tgn)ggggguGfuAfgacgscsg	2186	CGCGUCUACACCCCGCAUCAC AG	1635
AD-1571017.1	csgsueuaCfaCfCfCfcaucacuuL96	1960	asCfsugdTg(Agn)ugcgggUfgUfagagsgsc	2187	GGCGUCUACACCCCGCAUCACA GG	1636
AD-1571018.1	ascscgcAfUfCfAfCfagugugauuL96	1961	asAfsuudAc(Agn)ccugugAfuGfaggugsgsu	2188	ACACCCGCAUCACAGGUGUGA UC	1637
AD-1571019.1	gsasuecaGfcAfAfGfugugagaccuuL96	1962	asAfsugdTc(Agn)ccacuGfcUfggaucscsa	2189	UGGAUCCAGCAAGUGGUGACC UG	1640
AD-1571020.1	gsgscaggAfGfUfGfcaucunguuL96	1963	asAfsuudAc(Agn)ugccacCfuCfucgcsasc	2190	GUGGCAGGAGGUGGCAUCUU GUC	631
AD-1571021.1	uscscugAfUfUfCfuguccaguuL96	1964	asAfsuudCc(Agn)gcaagAfuCfagggascsg	2191	CGUCCUGAUGUCUGUCUCCAG UG	1648
AD-1571022.1	csusgaugUfcUfGfCfuccagugauuL96	1965	asAfsuudAc(Tgn)ggagcaGfaCfaucagsgsg	2192	CCUGAUGUCUGUCUCCAGUGA UG	1649
AD-1571023.1	gsgseucaGfcAfGfCfaagauguuL96	1966	asAfsugdTc(Tgn)cuugcuGfcUfagccscasc	2193	GUGGCUCAGCAGCAAGAAUGC UG	636
AD-1571024.1	ususgggaUfcUfGfGfcauggaaguuL96	1967	asCfsuudCc(Agn)uuuccaGfaUfccaasgsu	2194	ACUUGGAUCUUGGGAUUGGA AGG	642
AD-1571025.1	csasgcugCfcCfUfUfuggaauuuL96	1968	asUfauudAu(Tgn)ccaaagGfgCfagcugsasg	2195	CUCAGCUGCCCUUUGGAUAA AG	1655

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1571026.1	csusgccUfuUfGfGfaaagaacuuL96	1969	asAfsgeU(Tgn)auuccaAfaGfggcagsesu	2196	AGCUGCCCUUUGGAAUAAAAGCUG	648
AD-1571027.1	gsesccuuUfgGfAfAfuaaagcugcuL96	1970	asGfscadGc(Tgn)uuauucCfaAfagggcsasg	2197	CUGCCCUUUGGAAUAAAAGCUGCC	1656
AD-1571028.1	csesueacCfuGfCfUfucnuucgguuL96	1971	asAfsecaGfaAfGfaagcAfgGfugaggsgsg	2198	CCCUACACCGUCUUCUCUGG	2328
AD-1571029.1	csesueacCfuGfCfUfucnuucgguuL96	1971	asAfsecaGfaAfGfaagcAfgGfugaggsgsu	2199	CCCUACACCGUCUUCUCUGG	2328
AD-1571030.1	uscsacCfuGfCfUfucnuucgguuL96	1972	asAfsecaGfaAfGfaagcAfgGfugaggsg	2200	CCUACACCGUCUUCUCUGGUU	2329
AD-1571031.1	uscsacCfuGfCfUfucnuucgguuL96	1972	asAfsecaGfaAfGfaagcAfgGfugaggsgsu	2201	CCUACACCGUCUUCUCUGGUU	2329
AD-1571032.1	ascsCfuGfCfUfucnuucgguuL96	1973	asAfsecaGfaAfGfaagcAfgGfugaggsgsu	2202	UCACCGUCUUCUCUGGUU	2330
AD-1571033.1	Q191stUfcAfcCfuGfcUfuCfuUfcUfgGfsusUf	1974	asAfscCfaGfaAfGfaGfAfaGfCfAfgGfugGfsa	2203	UCACCGUCUUCUCUGGUU	2330
AD-1571034.1	gsgsagguGfaUfGfGfcgagggaagcuL96	1975	asGfscudTc(C2p)uegcccaUfcAfcuccsgsu	2204	ACGGAGGUGAUGGCGAGGAA	1492
AD-1571035.1	asasggccUfgUfGfAfaggacuccaauL96	1976	asUfsugdGa(G2p)uccucaCfaGfgecuusgsa	2205	UCAAGGCCUGUGAGGACUCCA	1493
AD-1571036.1	gsgsccugUfgAfGfGfaiuccaagauL96	1977	asUfscudTg(G2p)aguccuCfaCfaggcccsusu	2206	AAGGCCUGUGAGGACUCCAAG	1494
AD-1571037.1	gsescugUfGfGfAfceuaagaguL96	1978	asCfsucdTu(G2p)gaguccUfcAfcaggcsesu	2207	AGGCCUGUGAGGACUCCAAGA	524
AD-1571038.1	csusacucUfgGfUfAfuuuuccuagguL96	1979	asCfscudAg(G2p)aaaauacCfaGfaguagcsa	2208	UGCUACUCUGGUUUUCCUAG	529
AD-1571039.1	uscsugguAfuUfUfCfcuaggguacuL96	1980	asGfsuadCc(C2p)uagggaaAfuAfcagagsgsu	2209	ACUCUGGUUUUCCUAGGGU	533
AD-1571040.1	csusggguUfuUfCfCfuaggguuacuL96	1981	asUfsugdAc(C2p)cuagggaAfaUfaccagsasg	2210	CUCUGGUUUUCCUAGGGUAC	1497
AD-1571041.1	usgsuguUfuCfCfUfaggguacuauL96	1982	asUfsugdTa(C2p)ccuaggAfaAfuaccasgsa	2211	UCUGGUUUUCCUAGGGUAC	1498
AD-1571042.1	csusagggUfaCfAfagggcgagguuL96	1983	asAfscedTc(C2p)gccuugUfaCfcuagsgsa	2212	UCCUAGGGUACAAAGCGGAG	1501
AD-1571043.1	asusggucAfcCfCfAfsgguacuL96	1984	asGfsagdTa(C2p)accuaggCfuGfaccauscsa	2213	UGAUGGUCAGCCAGGUGUAC	1502
AD-1571044.1	gsuscagCfaGfGfUfguacuagguL96	1985	asCfscudGa(G2p)uacaccUfgGfcuagcsesa	2214	UGGUCAGCCAGGUGUACUCAG	1503

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1571045.1	csasgcccaGfgUfGfUfacucaggcauL96	1986	asUfsgcdCu(G2p)aguacaCfcUfsgcugsasc	2215	GUCAGCCAGGUGUACUCAGGC	1505
AD-1571046.1	csusecauCfcCfCfAfcuueucecauL96	1987	asUfsggdGa(G2p)aaguggCfGfAfuugagsus	2216	AG	1506
AD-1571047.1	csgsceacUfuCfUfCfccaggaucauL96	1988	asAfsagdTc(C2p)ugggagAfaGfugggcsasu	2217	UACUCAAUCCGCCACUUCUCCC	1507
AD-1571048.1	gsescacuUfcUfCfCfcaggaucauL96	1989	asAfsagdAu(C2p)cuagggaGfaAfguggcsgs	2218	UU	537
AD-1571050.1	uscsceagGfaUfCfUfuacccgccguL96	1990	asCfsggdCg(G2p)guuagaUfcCfugggasgsa	2219	UCGCCACUUCUCCCAGGAUC	539
AD-1571051.1	usasgugcCfuUfCfCfcagugaaauL96	1991	asUfsgndCa(C2p)ugcggaAfgGfcacuaggsa	2220	UA	1511
AD-1571052.1	csusuecgCfaGfUfGfaaacgcccauL96	1992	asUfsggdCg(G2p)uuuacUfcCfugggasgsa	2221	GG	1513
AD-1571053.1	csescagUfgAfAfAfcgcccaagauL96	1993	asCfsgndTg(G2p)cgguuuCfaCfuggggsasa	2222	AC	1514
AD-1571054.1	csgscaguGfaAfAfCfcgcccaagauL96	1994	asCfsgndTu(G2p)gcgguuUfcAfcugcggsa	2223	GC	1515
AD-1571055.1	gsescaaaGfcCfCfAfAgaagaugcuL96	1995	asAfsagdAu(C2p)uuuugGfcUfuugcsgs	2224	CC	1520
AD-1571056.1	csasgcacCfcGfCfCfcfugggaucauL96	1996	asAfsagdTu(C2p)ccaggeGfGfugcugsgsu	2225	UCCGCCAGUGAAACCGCCAAA	1522
AD-1571057.1	ascsaacuCfcAfGfCfuccgucuaauL96	1997	asAfsuadGa(C2p)ggagcuGfGfAfuugusas	2226	UA	1524
AD-1571058.1	csasccugCfuUfCfUfucugguucauL96	1998	asUfsgadAc(C2p)agaagaAfgCfaggugsasg	2227	UU	1525
AD-1571059.1	usgscuueUfuCfUfGfguuecauL96	1999	asAfsagdAu(G2p)aacagAfaGfaagcagsg	2228	CUACACUCCAGCUCUCCGUCUA	547
AD-1571060.1	uscsuueuGfGfUfUfCfaucuecauL96	2000	asUfsgudGa(G2p)aaugaCfcAfgaagagasg	2229	CUUCUUCUGGUUUCAUUCUCCA	1527
AD-1571061.1	ususcuggUfuCfAfUfucuecaauL96	2001	asAfsuadTg(G2p)agaagAfaCfcagaasgsa	2230	AA	1529
AD-1571062.1	uscsugguUfcAfUfUfucuecaauL96	2002	asCfsgaudTu(G2p)gagaauGfaAfcagagasg	2231	UC	1530
					CC	

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1571063.1	gsusggagGfaGfCfUfgcuguccaculL96	2003	asGfsugdGa(C2p)agcagcUfcCfuaccscsa	2232	UGGUGGAGGAGCUGCUGUCC	1531
AD-1571064.1	gsasggagCfuGfCfUfguccacaguuL96	2004	asAfsudGu(G2p)gacagcAfgCfuuccscsa	2233	UGGAGGAGCUGCUGUCCACAG	1532
AD-1571065.1	asgscugcUfgUfCfCfacagucaculL96	2005	asGfsuudGa(C2p)uguggaCfaGfcaagcscsc	2234	GGAGCUGCUGUCCACAGUCAA	1533
AD-1571066.1	csusguccAfcAfcGfUfcaacaguculL96	2006	asGfsagdCu(G2p)uugacuGfuGfagcagcsa	2235	UGCUGUCCACACAGUCAACAGCU	1536
AD-1571067.1	ascsagggCfcGfAfcGfuacgagugulL96	2007	asCfsacdTu(C2p)guacucGfgCfcugusag	2236	CUACAGGGCCGAGUACGAAAGU	552
AD-1571068.1	gsusggcgAfgUfAfcfgaagugagaculL96	2008	asGfsudCa(C2p)uucgnaCfuCfagcccsusg	2237	CAGGGCCGAGUACGAAAGUGG	1538
AD-1571069.1	uscsuaggAfaGfCfCfagugugaaulL96	2009	asUfsuudCa(C2p)acuggcUfuCfcaggasusc	2238	GAUCCUGGAAAGCCAGUGUGA	1541
AD-1571070.1	csusggaaGfcCfAfcGfugugaaagaulL96	2010	asUfscudTu(C2p)acacugGfcUfuaccagsgsa	2239	UCCUGGAAGCCAGUGUGUAAA	1543
AD-1571071.1	asasgccaGfuGfUfGfaaagacauulL96	2011	asUfsaudGu(C2p)uuuacAfcUfaggcuuscsc	2240	GGAAGCCAGUGUGUAAAGACA	1547
AD-1571072.1	gscscaguGfuGfAfcAfcagacauagaculL96	2012	asGfsudAu(G2p)uuuucAfcAfcugcsus	2241	AAGCCAGUGUGUAAAGACAU	1549
AD-1571074.1	usgsugaaAfgAfcAfcfuagcugcaulL96	2013	asAfsugdCa(C2p)cuuanguCfuUfuacacscsu	2242	AGUGUGAAAGACAUAAGCUGC	1552
AD-1571075.1	ascsugcugGfgUfUfGfuuaccgcuulL96	2014	asUfsagdCg(G2p)uaacaaCfcCfagcugsgsg	2243	CCAGCUGGGUUGUUACCCGU	1554
AD-1571076.1	csusguggGfuUfGfUfuaccgcuaculL96	2015	asGfsuadGc(G2p)guuacaAfcCfcagcugsgsg	2244	CACGCUGGGUUGUUACCCGCUA	1555
AD-1571077.1	gscsugggUfuGfUfUfUfaccgcuaculL96	2016	asUfsugdAg(C2p)gguuacAfaCfcagcugsgsu	2245	ACGCUGGGUUGUUACCCGUAC	1556
AD-1571078.1	csusggguUfgUfUfAfcgcuacagulL96	2017	asCfsugdTa(G2p)cgguaaCfaAfcacagsgsg	2246	CGCUGGGUUGUUACCCGUACA	563
AD-1571079.1	csusggagAfaGfAfcfgcuacaculL96	2018	asGfsugdAu(G2p)agccucUfuCfuaccagsgsg	2247	CCUUGGAGAAGAGGCUCAUCA	1561
AD-1571080.1	gsasgaagAfgGfCfUfcauacaculL96	2019	asGfsagdGu(G2p)augagcCfuCfuuccscsa	2248	UGGAGAAGAGGCUCAUCACCU	1564

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1571081.1	gsasagagGfcUfCfAfucacucggulL96	2020	asCfscgdAg(G2p)ugaugaGfcCfucucusc	2249	GAGAAAGAGGCUCAUACACCCUGG	1565
AD-1571082.1	asgsgeucAfuCfAfCfucgguguanL96	2021	asUfsaedAc(C2p)gagugAfuGfagcucsesu	2250	AGAGGCUCAUACACCCUGGUGU	571
AD-1571083.1	asasgaagGfgCfCfUfgcacaguanL96	2022	asUfsagdCu(G2p)ugcaggCfcCfucucusc	2251	GGAAGAAGGGCCUGCACAGCU	1567
AD-1571084.1	asasgggcCfuGfCfAfCfagcucuanL96	2023	asUfsagdTa(G2p)ecugucAfgGfcccucsesu	2252	AGAAGGGCCUGCACAGCUACU	1568
AD-1571085.1	csesucucUfgGfAfCfucggcuugulL96	2024	asCfsaadGc(C2p)guagucCfaGfagaggsgc	2253	GCCCUCUCUGGACUACGGCUU	574
AD-1571086.1	usesucugGfaCfUfAfCfagcucggculL96	2025	asGfscddAa(G2p)ecguagUfcCfagagaggsg	2254	CCUCUCUGGACUACGGCUUUGG	575
AD-1571087.1	usgsagucAfcGfGfCfucggccuculL96	2026	asGfscgdGg(C2p)caagccGfuAfgucaggsa	2255	UCUGGACUACGGCUUUGGCCCU	1570
AD-1571088.1	gsasacuaCfGfCfUfugggccuculL96	2027	asAfsagdGg(G2p)ccaagcCfGfagucssasg	2256	CUGGACUACGGCUUUGGCCCU	1571
AD-1571089.1	asgsaaguAfuGfAfUfugccgugculL96	2028	asGfscgdCg(G2p)caaaucAfuAfcucucgsc	2257	GCAGAAGUAUGAUUUGGCCGU	587
AD-1571090.1	gsasaguaUfgAfuUfugccgugcualL96	2029	asUfsagdAc(G2p)gcaauCfaUfacucucsg	2258	CAGAAGUAUGAUUUGGCCGUG	588
AD-1571091.1	asasguauGfaUfUfUfgccgugcualL96	2030	asGfscgdCa(G2p)ggcaaaUfcAfuacucsesu	2259	AGAAGUAUGAUUUGGCCGUGC	1573
AD-1571092.1	gsusaugaUfuUfGfCfCfugccaccucL96	2031	asCfsggdTg(C2p)acggcaAfaUfcauacsesu	2260	AAGUAUGAUUUGGCCGUGCACC	1575
AD-1571093.1	gsgeccagUfgGfAfCfCfaguccagaaL96	2032	asUfsugdTa(G2p)aucgucCfaCfugccscsu	2261	AGGGCCAGUGGACGAUCCAGA	1576
AD-1571094.1	gsccagagGfgAfcGfauccagaaL96	2033	asGfsuudCu(G2p)gcuugCfcAfcugccscsc	2262	GGCCAGUGGACGAUCCAGAA	1577
AD-1571096.1	usgsagagAfuCfCfAfgaacaggagulL96	2034	asCfsugdCu(G2p)uucuggAfuCfugcucsesu	2263	AGUGGACGAUCCAGAACAGG	1579
AD-1571097.1	csascucCfcAfcGfAfucuccucuaL96	2035	asUfsagdGg(G2p)agauucGfgGfaggugsas	2264	UUCACCUCCAGAUUCUCCUC	1581
AD-1571098.1	usgsccggUfgCfAfCfCfuaugggcuugulL96	2036	asCfsaadGc(C2p)auagucCfaCfcccagcsa	2265	UGUGCGGGUGCACUAUUGGCU	1582

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1571099.1	gscsgggguGfcAfCfUfauugcuuugulL96	2037	asAfscaadAg(C2p)cauaguGfcAfcgccgsase	2266	GUGCGGGUGCACUAUGGCUU GUA	595
AD-1571100.1	csgsgggugCfaCfUfAfuugcuuugulL96	2038	asUfsaedAat(G2p)ccauagUfgCfaceccgsca	2267	UAC	1583
AD-1571102.1	asascgggcCfuGfGfAfugagagaaauL96	2039	asUfsuudCu(C2p)ucauccAfgGfccguusgsg	2268	CCAACGGCCUGGAUGAGAGAA AC	1588
AD-1571103.1	csgsgccuGfgAfUfGfagagaaacuuL96	2040	asAfsaudTu(C2p)ucueauCfcAfggccgsusu	2269	AACGGCCUGGAUGAGAGAGAA CUG	1590
AD-1571104.1	gsasgaaaCfuGfCfGfuuugcagagulL96	2041	asCfsuudTg(C2p)aaacgcAfgUfuucucsu	2270	GAGAGAAACUGCGUUUUGCAG AGC	1593
AD-1571105.1	csusgcguUfuGfCfAfgagccacuuL96	2042	asAfsaudTg(G2p)ucucgcAfaAfcgcagsusu	2271	AACUGCGUUUGCAGAGAGCCACA UU	1594
AD-1571106.1	usgscguUfgCfAfGfagccacuuL96	2043	asAfsaudGu(G2p)gcucugCfaAfacgcagsu	2272	ACUGCGUUUGCAGAGAGCCACA UC	1595
AD-1571107.1	csgsuuugCfaGfAfGfccacacuuL96	2044	asGfsaudAu(G2p)uggcucUfgCfaaacgsca	2273	UGCGUUUGCAGAGAGCCACA CA	1597
AD-1571108.1	gscsagagCfcAfCfAfuuccagugulL96	2045	asGfsaudCu(G2p)gauguGfgCfucugcsasa	2274	UUGCAGAGCCACAUAUCCAGUG CA	1598
AD-1571109.1	gsgsgacaUfuCfAfCfuuccagugulL96	2046	asCfsaudTg(G2p)aaagugAfaUfguuccsase	2275	GUGGACAUUCCACCUUCCAGU GU	1599
AD-1571110.1	gsgsacauUfcAfCfCfuuccagugulL96	2047	asAfscauCu(G2p)gaagguGfaAfuugccsca	2276	UGGACAUUCCACCUUCCAGUG UG	607
AD-1571111.1	ascsauucAfcCfUfUfccagugugulL96	2048	asUfscauCa(C2p)uggaagGfuGfaaunguscsc	2277	GGACAUUCCACCUUCCAGUG AG	609
AD-1571112.1	asgseuigeGfuGfAfAfgaagcccauL96	2049	asUfsaudGg(C2p)uuuucAfcGfcagcuscsc	2278	GGAGCUGCGUGAAAGAGGCCCA AC	1601
AD-1571113.1	gscsugcgUfgAfAfGfagcccaacuL96	2050	asGfsuudGg(G2p)uuuucCfaCfcagcuscsc	2279	GAGCUGCGUGAAAGAGGCCCA CC	1602
AD-1571114.1	csusgcguGfaAfGfAfagcccaacuL96	2051	asGfsaudTg(G2p)gcuuucUfcAfcgcagsesu	2280	AGCUGCGUGAAAGAGGCCCA CC	1603
AD-1571115.1	usgscguUfgAfAfAfgcccaacuL96	2052	asGfsaudTg(G2p)ggcuucUfuCfacgcagsc	2281	GCUGCGUGAAAGAGGCCCA CC	1604
AD-1571116.1	asgseacuGfuGfAfCfugggccuulL96	2053	asGfsaudGg(C2p)acagucAfcAfgugcuscsc	2282	GGAGCACUGUGACUGUGGCCU CC	1605

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1571117.1	csuscegaGfgGfUfGfagugggccaauL96	2054	asAfsugdGc(C2p)acucacCfcUfcccagsgsa	2283	UCCUCCGAGGGUGAGUGGCCA UG	1606
AD-1571118.1	asusgecuGfaCfCfGfuggggugaauL96	2055	asAfsuedAc(C2p)cagcggUfcAfcggausgsa	2284	UCAUCGCUGACCCGCUUGGGUGA UA	611
AD-1571119.1	uscsgeugAfcCfGfCfugggugaauL96	2056	asUfsaudCa(C2p)ccagcgGfuCfagegasusg	2285	CAUCGCUGACCCGCUUGGGUGAU AA	1607
AD-1571120.1	gscsugacCfGcUfGfGfuggaauaauL96	2057	asGfsuudAu(C2p)accacgCfGfGfucagcsgsa	2286	UCGUGACCCGCUUGGGUGAUAA CA	1609
AD-1571121.1	gscsugggUfgAfUfAfacagcugccuL96	2058	asGfsgcdAg(C2p)uguuauCfaCfcccagsgsg	2287	CCGCUUGGGUGAUAAACAGCUGC CC	1610
AD-1571122.1	usgsuucCfaGfGfAfcgacagcauL96	2059	asAfsugdCu(G2p)uccuccUfgGfaagcasgsu	2288	ACUGCUUCCAGGAGGACAGCA UG	1611
AD-1571123.1	csusucaGfgAfGfGfGfagcaugguL96	2060	asCfscadTg(C2p)uguccuCfcUfGgaagscsa	2289	UGCUUCCAGGAGGACAGCAUG GC	1613
AD-1571124.1	csusgggcAfaGfGfUfGfugacagaauL96	2061	asUfsuedTg(C2p)ccacacUfuGfcccagsgsa	2290	UCCUGGGCAAGGUGUGGCAG AAC	1614
AD-1571125.1	usgsggcaAfgGfUfGfugggcagaauL96	2062	asGfsuudCu(G2p)ccacacCfuUfcccagsgsg	2291	CCUGGGCAAGGUGUGGCAGA ACU	1615
AD-1571126.1	gsgscaagGfuUfGfGfGfagcauauL96	2063	asGfsagdTu(C2p)ugccacAfcCfuugccscsa	2292	UGGGCAAGGUGUGGCAGAAC UCG	1617
AD-1571127.1	usgsgecuGfgAfGfAfcguguccuuuL96	2064	asAfsagdGat(C2p)accuccCfcAfcgcccagsgc	2293	GCUGGCCUGGAGAGGUGUCCU UC	1619
AD-1571128.1	gscscuggAfgAfGfGfuguccuauL96	2065	asUfsagdAg(C2p)acaccuCfuCfcccagsgcsa	2294	UGGCCUGGAGAGGUGUCCUUC AA	1621
AD-1571129.1	cscsuggaGfaGfGfUfGfuccuauL96	2066	asUfsugdAa(C2p)gacaccUfcUfcccagsgsc	2295	GGCCUGGAGAGGUGUCCUUCA AG	1622
AD-1571130.1	usgsuugaGfuUfGfAfucccagagguL96	2067	asCfscudGu(G2p)ggaucaAfcUfcccagsgc	2296	GAUGGCAGUUGAUUCCACACAG GA	1623
AD-1571131.1	usgscauUfgAfUfCfcccagaggaauL96	2068	asGfsuedCu(G2p)ugggauCfaAfcugcascsa	2297	UGUGCAGUUGAUUCCACACAGGA CC	1624
AD-1571132.1	asuscccaCfaGfGfAfcgugugcaguL96	2069	asCfsugdCa(C2p)agguccUfgUfggggauscsa	2298	UGAUCCACAGGACCUGUGCA GC	621
AD-1571133.1	cscscacaGfgAfCfCfugugcagguL96	2070	asCfsgcdTg(C2p)acagguCfcUfGggggsasu	2299	AUCCACAGGACCUGUGGCAGC GA	1625

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1571134.1	cscacagGfaCfCfUfgucagcgaL96	2071	asUfscgdCu(G2p)cacaggUfcCfugggsgsa	2300	UCCCACAGGACCUGUGCAGCG AG	1626
AD-1571135.1	cscsagguGfaCfGfCfcaugcgaucL96	2072	asGfscadTg(C2p)gugcgUfcAfcuuggsusa	2301	UACCAGGUGACGCCACGCCAUG CU	1627
AD-1571136.1	gsusgacGfCfAfcGfcaugcugugL96	2073	asCfscadAg(C2p)augcgUfgCfucacscsu	2302	AGGUGACGCCACGCCAUGCUGU GU	1628
AD-1571137.1	usgsacGfCfAfcGfCfcaugcugugL96	2074	asAfcscadCa(G2p)caugcgUfgGfucacscsc	2303	GGUGACGCCACGCCAUGCUGUG UG	622
AD-1571138.1	ascsgccaCfGfCfAfcUfgcugugugL96	2075	asGfscadCa(C2p)agcagUfgUfgcugscsa	2304	UGACGCCACGCCAUGCUGUG CC	1629
AD-1571139.1	gsuscuacCfGfCfAfcGfcaagcaL96	2076	asUfscudTu(G2p)ccuugCfGfuaagcsgsg	2305	CCGGUACCCGCAAGGGCAAGA AG	1630
AD-1571140.1	gsusgcaaGfGfCfAfcUfucagugcL96	2077	asGfsgcdCa(C2p)ugagugCfCfUfgcacsasc	2306	GUGUGCAAGGCACUCACUGGC CG	1631
AD-1571141.1	csusaacuAfcUfUfCfGcgcucuaL96	2078	asGfscudGa(C2p)gcgaaGfuAfcuaagsgsc	2307	GCCUAAUACUUCGGCGUCUA CA	1632
AD-1571142.1	csusacacCfcGfCfAfcUfucagugL96	2079	asCfscadCu(G2p)ugaugCfGfGfuaagasc	2308	GUCUACACCCGCAUCACAGGU GU	628
AD-1571143.1	cscscgcaUfcAfcUfCfGcugugauL96	2080	asGfscudCa(C2p)accuGfUfgcgggsgsg	2309	CACCCGCAUCACAGGUGUGAU CA	1638
AD-1571144.1	usgsaucCfaGfCfAfcUfgugugacL96	2081	asGfscudAc(C2p)acuuGfUfgGfauccagsc	2310	GCUGGAUCCAGCAAGUGGUG ACC	1639
AD-1571145.1	asusccagCfaAfcUfGfUfgugaccuL96	2082	asCfscadGu(C2p)accacuUfgCfugcuscsc	2311	GGAUCCAGCAAGUGGUGACCU GA	1641
AD-1571146.1	cscsagcaAfcUfGfGfUfgugaccuL96	2083	asCfscadAg(C2p)uacccaCfuUfgcuggsasu	2312	AUCCAGCAAGUGGUGACCUGA GG	1642
AD-1571147.1	csasgcaaGfuGfGfUfgaccugagL96	2084	asCfscudCa(G2p)aucaccAfcUfugcuggsa	2313	UCCAGCAAGUGGUGACCUGAG GA	1643
AD-1571148.1	gsusaaguGfGfUfgGfAfcuagagauL96	2085	asUfscudCu(C2p)agguaCfcAfcuugcscusg	2314	CAGCAAGUGGUGACCUGAGG AAC	1644
AD-1571149.1	usgsugGfCfaGfGfAfcgugcgaucL96	2086	asGfscudGc(C2p)accuUfgCfaccacsa	2315	UGUGGUGCAGGAGGUGGCA UCU	1645
AD-1571150.1	gsusgagCfGfGfAfcUfgugcgaucL96	2087	asAfcsgadTg(C2p)caucUfcUfgcaccasc	2316	GUGGUGCAGGAGGUGGCAU CUU	1646

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA Target Sequence 5' to 3'	SEQ ID NO:
AD-1571151.1	gsusggcaGfgAfGfGfuggcaucuuL96	2088	asAfsagdAu(G2p)ccacuuCfcUfgccacscsa	2317	UGGUGGCAGGAGGUGGCAUC UUG	1647
AD-1571152.1	uscsaaguGfaUfGfGfcagaggaauL96	2089	asAfsuedCu(C2p)ugccaUfcAfcugagsgsc	2318	GUCCAGUGAUGGCAGGAGG AUG	1650
AD-1571153.1	csusaacuUfgGfGfAfcuugggaaauL96	2090	asAfsuudCc(C2p)agauccCfaAfguuagsasc	2319	GUCUACUUUGGAUCUGGGA AUG	1651
AD-1571154.1	gsusgagcUfcAfGfCfcugccuuuugL96	2091	asCfsaadAg(G2p)gcagcuGfaGfcuacacscsu	2320	AGGUGAGCUCAGCUGGCCUUU GG	1652
AD-1571155.1	csuscagcUfgCfcCfcfuuggaauuL96	2092	asUfsaudTc(C2p)aaagggCfaGfcugagsgscu	2321	AGCUCAGCUGCCCCUUUGGAU AA	1653
AD-1571156.1	uscsagcuGfcCfcUfuuggaauaauL96	2093	asUfsuadTu(C2p)caaaaggGfcAfcugagsgsc	2322	GCUCAGCUGCCCCUUUGGAUA AA	1654
AD-1571157.1	cscscuuuGfgAfAfUfaaagcugccuL96	2094	asGfsgcdAg(C2p)uuuuuuCfcAfaagggscsa	2323	UGCCCUUUGGAUAAAAGCUGC CU	1657
AD-1571158.1	cscsuuugGfaAfUfAfaagcugccuL96	2095	asAfsaggdCa(G2p)cuuuuuUfcCfaaaggsgsc	2324	GCCCUUUGGAUAAAAGCUGCC UG	1658
AD-1571159.1	ususuggaAfAfAfAfcugcugccuL96	2096	asUfscadGg(C2p)ageuuuAfUfccaasgsg	2325	CCUUUGGAUAAAAGCUGCCUG AU	1659
AD-1571160.1	ususggaaUfaAfAfGfCfcugccuL96	2097	asAfsuedAg(G2p)cagcuUfaUfuccaasag	2326	CUUUGGAUAAAAGCUGCCUG AUC	1660
AD-1571161.1	usgsaaUfaAfGfCfcugcugauL96	2098	asGfscadCa(G2p)gcagcuUfuAfuuccasasa	2327	UUUGGAUAAAAGCUGCCUGA UCC	1661

Table 8. Single Dose Screen in Hep3b Cells

	10 nM		1 nM		0.1 nM	
Duplex	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev
AD-1570929.1	55	8	73	7	55	8
AD-1571034.1	76	5	94	7	129	8
AD-1571035.1	62	15	73	11	82	8
AD-1571036.1	53	8	75	11	92	4
AD-1554875.1	14	3	21	4	30	5
AD-1571037.1	29	7	44	12	98	9
AD-1570930.1	15	3	22	3	30	2
AD-1570931.1	11	2	14	1	21	8
AD-1554909.1	22	6	39	1	44	8
AD-1554910.1	21	4	30	3	39	7
AD-1554911.1	21	3	32	10	36	5
AD-1554912.1	21	3	46	3	51	5
AD-1554913.1	50	6	71	15	66	13
AD-1571038.1	95	21	96	9	121	9
AD-1554914.1	47	8	74	9	70	9
AD-1554915.1	28	3	51	9	50	7
AD-1554916.1	34	5	54	8	64	8
AD-1570932.1	17	3	34	5	55	8
AD-1554917.1	25	5	47	4	52	8
AD-1571039.1	31	3	55	13	89	10
AD-1571040.1	37	8	43	11	86	11
AD-1571041.1	36	9	61	16	97	32
AD-1570933.1	92	14	109	22	97	3
AD-1570934.1	80	11	103	9	71	9
AD-1554923.1	41	6	79	16	72	7
AD-1571042.1	69	19	70	4	93	5
AD-1571043.1	56	11	81	16	107	8
AD-1554951.1	32	5	59	2	56	7
AD-1570935.1	60	12	79	12	73	7
AD-1571044.1	78	9	64	14	122	21
AD-1570936.1	103	24	105	13	102	22
AD-1571045.1	76	15	99	15	122	22
AD-1554955.1	31	6	48	4	51	6
AD-1570937.1	27	5	54	3	61	3
AD-1571046.1	37	9	60	16	87	16
AD-1571047.1	23	3	28	7	45	9
AD-1554992.1	85	6	99	9	75	2
AD-1571048.1	74	14	98	10	111	20
AD-1570938.1	36	4	71	12	70	11
AD-1554997.1	24	6	43	3	50	4
AD-1570939.1	111	16	117	11	84	8

Duplex	10 nM		1 nM		0.1 nM	
	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev
AD-1555000.1	30	5	51	4	64	12
AD-1571050.1	51	10	87	6	88	8
AD-1571051.1	44	7	68	18	77	15
AD-1555030.1	30	6	61	7	57	9
AD-1570940.1	27	4	62	6	70	6
AD-1570941.1	103	16	113	11	79	10
AD-1571052.1	23	4	38	1	40	7
AD-1571053.1	31	2	58	14	76	5
AD-1571054.1	28	5	46	5	56	6
AD-1570942.1	47	4	70	5	76	8
AD-1570943.1	27	7	42	3	68	4
AD-1570944.1	38	6	36	4	62	6
AD-1570945.1	52	8	87	7	67	4
AD-1571055.1	43	6	68	12	83	10
AD-1570946.1	80	11	89	11	82	4
AD-1571056.1	44	3	70	13	87	13
AD-1570947.1	54	9	80	14	84	9
AD-1571057.1	43	3	62	6	67	14
AD-1555106.1	16	5	17	2	35	8
AD-1570948.1	26	7	34	6	53	7
AD-1555112.1	33	5	61	4	64	8
AD-1571028.1	65	8	87	6	105	10
AD-1571029.1	69	12	83	4	112	19
AD-1555114.1	25	6	36	3	43	12
AD-1555115.1	26	5	38	4	40	6
AD-1570949.1	29	5	45	3	56	8
AD-1571030.1	37	2	61	12	74	12
AD-1571031.1	46	11	64	14	79	10
AD-1571058.1	34	4	44	3	53	5
AD-1555117.1	23	6	27	5	38	2
AD-1571032.1	54	1	80	14	86	7
AD-1571033.1	44	5	80	11	101	23
AD-1555118.1	30	7	33	3	47	5
AD-1570950.1	31	6	44	5	63	5
AD-1570951.1	28	6	33	7	46	3
AD-1555120.1	24	6	33	4	53	11
AD-1571059.1	28	5	44	6	55	5
AD-1555121.1	36	5	55	4	69	7
AD-1555122.1	22	4	32	5	49	6
AD-1570952.1	25	4	45	6	52	8
AD-1555123.1	35	7	43	1	70	5
AD-1570953.1	93	6	102	14	101	12

Duplex	10 nM		1 nM		0.1 nM	
	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev
AD-1571060.1	25	4	42	9	53	11
AD-1570954.1	23	6	32	3	72	17
AD-1571061.1	22	3	35	3	43	3
AD-1571062.1	44	7	68	5	87	15
AD-1555128.1	36	7	41	6	63	15
AD-1570955.1	31	8	35	2	48	9
AD-1571063.1	80	10	88	14	89	6
AD-1571064.1	87	8	94	6	123	6
AD-1571065.1	68	4	80	9	93	8
AD-1570956.1	48	9	76	7	93	19
AD-1570957.1	50	11	66	3	82	14
AD-1571066.1	35	5	43	9	82	26
AD-1570958.1	69	12	102	7	92	3
AD-1555184.1	87	13	100	12	99	5
AD-1571067.1	80	18	77	11	93	12
AD-1555185.1	63	15	88	15	100	13
AD-1571068.1	71	10	55	6	73	15
AD-1570959.1	104	13	106	9	85	6
AD-1570960.1	48	9	62	18	79	16
AD-1571069.1	57	5	41	10	81	9
AD-1570961.1	73	12	101	2	94	13
AD-1571070.1	48	5	44	11	78	8
AD-1570962.1	57	11	88	6	82	13
AD-1570963.1	33	6	52	4	50	8
AD-1570964.1	52	10	83	7	92	19
AD-1571071.1	59	4	65	6	85	16
AD-1570965.1	86	17	109	12	100	16
AD-1571072.1	72	6	75	4	120	8
AD-1555212.1	42	11	56	7	71	13
AD-1570966.1	32	5	39	9	58	6
AD-1555213.1	33	6	36	5	47	7
AD-1570967.1	35	8	58	10	52	4
AD-1571074.1	19	3	31	6	33	4
AD-1570968.1	30	6	41	4	44	7
AD-1555234.1	30	6	41	6	56	5
AD-1570969.1	42	8	62	10	61	8
AD-1555235.1	51	9	77	12	72	5
AD-1555236.1	59	7	67	15	68	3
AD-1555238.1	45	8	55	9	58	1
AD-1570970.1	77	10	88	32	74	6
AD-1555241.1	41	6	57	10	39	9
AD-1555242.1	47	6	83	6	71	3

Duplex	10 nM		1 nM		0.1 nM	
	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev
AD-1555243.1	41	8	65	7	67	4
AD-1570971.1	93	11	108	8	92	14
AD-1571075.1	25	5	37	5	38	3
AD-1571076.1	15	4	33	9	41	8
AD-1571077.1	39	9	43	13	46	11
AD-1555247.1	42	4	51	4	78	8
AD-1571078.1	16	3	40	14	49	4
AD-1570972.1	53	15	67	39	40	18
AD-1570973.1	45	5	35	8	55	8
AD-1570974.1	76	12	81	16	81	9
AD-1555342.1	73	16	69	3	78	18
AD-1570975.1	108	21	84	15	103	13
AD-1555343.1	80	12	92	5	91	9
AD-1555345.1	84	10	97	6	103	13
AD-1555346.1	54	12	71	7	86	3
AD-1570976.1	71	11	70	9	93	5
AD-1555348.1	57	22	64	6	84	12
AD-1555349.1	36	8	50	2	66	7
AD-1555350.1	57	9	58	10	77	10
AD-1571079.1	71	12	77	10	65	11
AD-1570977.1	34	8	68	10	92	8
AD-1570978.1	30	13	53	4	86	5
AD-1571080.1	63	11	70	3	71	14
AD-1571081.1	76	12	79	3	94	16
AD-1555366.1	42	4	48	2	78	2
AD-1571082.1	37	3	54	5	56	11
AD-1570979.1	31	8	54	12	72	10
AD-1571083.1	45	4	54	6	56	8
AD-1571084.1	34	1	53	11	58	11
AD-1570980.1	82	13	81	14	92	9
AD-1555428.1	48	12	75	8	96	4
AD-1555429.1	47	7	66	8	90	4
AD-1570981.1	34	14	66	3	92	6
AD-1555535.1	41	2	65	5	71	6
AD-1571085.1	48	6	77	6	69	8
AD-1555537.1	52	1	63	4	115	12
AD-1571086.1	40	2	54	6	61	3
AD-1571087.1	69	15	76	4	97	13
AD-1571088.1	39	7	63	8	60	7
AD-1555546.1	20	4	30	4	56	7
AD-1555547.1	24	3	47	4	73	11
AD-1555548.1	41	5	55	5	79	7

Duplex	10 nM		1 nM		0.1 nM	
	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev
AD-1555549.1	61	10	89	7	84	11
AD-1555581.1	35	5	60	9	95	12
AD-1570982.1	55	1	80	11	95	11
AD-1570983.1	61	5	84	10	100	13
AD-1555583.1	40	4	65	3	89	9
AD-1555584.1	50	5	78	11	102	8
AD-1555585.1	49	4	74	13	86	9
AD-1555586.1	48	11	70	5	86	18
AD-1555587.1	34	9	60	6	89	11
AD-1555588.1	40	7	56	7	91	10
AD-1555589.1	34	3	52	11	83	13
AD-1571089.1	32	3	42	6	60	3
AD-1555590.1	46	6	68	16	87	5
AD-1571090.1	40	8	54	10	69	12
AD-1571091.1	39	8	52	7	56	7
AD-1570984.1	77	11	100	10	110	8
AD-1571092.1	39	9	76	6	86	16
AD-1571093.1	71	9	76	7	86	10
AD-1571094.1	66	7	73	18	104	13
AD-1570985.1	25	5	43	10	60	4
AD-1555615.1	43	2	60	7	82	12
AD-1555616.1	60	10	84	22	91	9
AD-1571096.1	90	10	95	12	96	14
AD-1555626.1	69	15	67	11	99	11
AD-1570986.1	71	6	90	10	93	5
AD-1555628.1	81	7	85	11	102	15
AD-1570987.1	119	16	99	14	126	8
AD-1570988.1	82	7	96	8	116	10
AD-1571097.1	43	3	65	11	61	6
AD-1555706.1	60	10	78	16	101	18
AD-1570989.1	59	17	83	12	96	12
AD-1555707.1	34	8	57	5	81	9
AD-1570990.1	63	9	67	8	93	9
AD-1571098.1	48	3	73	3	82	10
AD-1555709.1	44	3	72	12	89	14
AD-1571099.1	50	11	79	12	92	7
AD-1571100.1	24	5	44	3	64	10
AD-1555711.1	49	5	78	4	97	15
AD-1570991.1	77	8	122	9	114	11
AD-1570992.1	78	6	127	24	97	12
AD-1570993.1	28	4	51	1	77	6
AD-1570994.1	67	1	85	18	102	11

Duplex	10 nM		1 nM		0.1 nM	
	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev
AD-1555717.1	42	2	57	1	73	8
AD-1555723.1	48	5	70	11	100	13
AD-1555725.1	42	3	71	11	98	17
AD-1570995.1	90	13	110	20	129	15
AD-1571102.1	24	4	37	4	58	5
AD-1570996.1	47	8	87	24	112	13
AD-1571103.1	43	6	68	8	92	14
AD-1555768.1	37	8	66	14	92	18
AD-1570997.1	43	6	85	17	89	20
AD-1570998.1	61	7	91	16	90	23
AD-1555771.1	17	3	34	6	44	5
AD-1555772.1	23	3	43	10	66	17
AD-1555776.1	52	12	82	12	117	23
AD-1570999.1	120	18	120	23	154	33
AD-1571104.1	70	9	56	7	91	16
AD-1571105.1	20	1	40	5	40	5
AD-1571106.1	31	2	47	7	74	14
AD-1571000.1	38	5	94	12	112	16
AD-1571107.1	25	1	52	5	70	6
AD-1555789.1	27	2	55	7	72	9
AD-1571108.1	65	9	87	6	92	21
AD-1555894.1	52	13	65	9	115	5
AD-1555895.1	37	7	58	8	78	17
AD-1571001.1	59	11	96	15	96	16
AD-1571109.1	62	4	83	7	93	10
AD-1555897.1	57	15	88	21	125	13
AD-1571110.1	79	11	109	10	118	10
AD-1555898.1	47	7	87	24	114	24
AD-1555899.1	78	6	109	14	104	4
AD-1571111.1	88	5	95	8	107	17
AD-1555900.1	45	4	99	12	86	5
AD-1571002.1	19	8	61	5	69	6
AD-1571112.1	27	3	50	6	65	11
AD-1571113.1	41	2	64	8	82	16
AD-1571114.1	39	5	62	5	77	15
AD-1571115.1	54	7	70	7	74	13
AD-1571116.1	41	4	70	8	75	13
AD-1571117.1	110	3	108	23	102	14
AD-1556052.1	19	3	42	5	73	17
AD-1571118.1	24	5	60	6	79	3
AD-1571119.1	30	3	55	10	83	13
AD-1571003.1	42	4	87	7	94	11

Duplex	10 nM		1 nM		0.1 nM	
	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev
AD-1571120.1	44	7	58	11	77	15
AD-1556057.1	33	5	69	12	71	7
AD-1571121.1	69	11	77	6	87	8
AD-1571122.1	46	4	62	13	81	19
AD-1571004.1	106	5	115	12	111	10
AD-1571123.1	90	10	103	6	102	9
AD-1556126.1	43	2	103	18	100	16
AD-1571005.1	40	16	99	10	88	10
AD-1556127.1	38	3	75	14	77	7
AD-1571124.1	44	6	84	11	102	15
AD-1571125.1	54	6	95	16	107	19
AD-1571006.1	35	0	76	9	80	2
AD-1571126.1	49	11	70	10	72	12
AD-1556137.1	40	1	85	17	86	3
AD-1571007.1	66	12	117	20	104	14
AD-1571008.1	55	5	101	25	107	8
AD-1556139.1	48	7	84	15	101	21
AD-1571127.1	60	5	76	6	79	8
AD-1571009.1	23	6	76	19	66	8
AD-1571128.1	42	5	71	11	95	15
AD-1571129.1	47	9	71	11	87	18
AD-1556163.1	27	6	81	14	85	11
AD-1571010.1	61	5	94	11	73	5
AD-1556164.1	52	5	41	7	77	2
AD-1556166.1	55	10	88	7	89	14
AD-1556167.1	43	7	93	13	114	8
AD-1571011.1	44	12	99	12	101	14
AD-1571130.1	48	3	82	12	83	15
AD-1571131.1	54	7	78	10	99	21
AD-1556319.1	34	5	47	15	62	9
AD-1571132.1	75	15	100	25	114	14
AD-1571133.1	96	24	110	24	126	31
AD-1571134.1	52	14	87	14	108	10
AD-1571135.1	47	12	65	4	138	37
AD-1571136.1	93	7	105	14	112	13
AD-1556359.1	31	6	36	1	81	0
AD-1571137.1	59	9	81	10	100	13
AD-1556360.1	26	2	49	15	48	12
AD-1571138.1	85	18	93	18	91	16
AD-1571139.1	51	10	92	15	100	24
AD-1556382.1	38	8	63	6	40	8
AD-1571012.1	58	6	71	10	54	15

Duplex	10 nM		1 nM		0.1 nM	
	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev
AD-1556383.1	44	7	81	10	70	22
AD-1571013.1	58	12	90	6	86	10
AD-1571140.1	117	32	120	16	131	16
AD-1556465.1	36	2	70	8	68	2
AD-1556466.1	8	2	24	4	41	3
AD-1571141.1	52	11	88	15	97	12
AD-1571014.1	63	13	45	10	91	24
AD-1571015.1	49	6	83	9	80	19
AD-1571016.1	47	4	67	1	59	7
AD-1571017.1	55	5	90	13	90	14
AD-1556484.1	49	13	87	2	79	16
AD-1571142.1	84	11	94	20	97	11
AD-1571018.1	48	9	83	10	94	16
AD-1571143.1	49	5	73	12	95	5
AD-1556510.1	34	6	57	3	68	10
AD-1571144.1	28	7	53	10	74	9
AD-1571019.1	28	2	54	2	75	9
AD-1571145.1	38	7	51	4	77	8
AD-1571146.1	39	3	63	3	81	11
AD-1571147.1	38	6	48	9	77	5
AD-1571148.1	25	1	46	6	68	4
AD-1571149.1	59	7	68	8	72	5
AD-1571150.1	41	11	65	8	88	5
AD-1571151.1	59	1	74	13	94	13
AD-1556584.1	67	5	102	17	89	17
AD-1556585.1	54	3	92	17	91	21
AD-1571020.1	86	13	118	16	114	9
AD-1556586.1	57	7	93	3	103	13
AD-1556587.1	47	8	75	9	94	7
AD-1571021.1	72	1	95	4	117	8
AD-1571022.1	47	3	84	8	97	8
AD-1556613.1	48	9	62	7	88	12
AD-1571152.1	52	4	67	8	92	13
AD-1556677.1	40	5	80	18	94	12
AD-1556709.1	66	16	92	5	91	9
AD-1571023.1	56	8	94	13	85	10
AD-1556710.1	51	6	69	9	91	12
AD-1556789.1	57	4	97	5	93	12
AD-1556790.1	75	5	113	21	107	13
AD-1556791.1	77	13	101	22	99	19
AD-1571153.1	53	9	65	9	95	11
AD-1556795.1	43	4	82	5	99	17

Duplex	10 nM		1 nM		0.1 nM	
	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev	Avg % message remaining	St. Dev
AD-1556799.1	64	3	87	11	104	7
AD-1571024.1	85	11	113	13	115	7
AD-1556802.1	62	10	95	14	92	14
AD-1571154.1	47	6	67	7	96	7
AD-1556908.1	37	5	82	10	93	11
AD-1556909.1	70	16	101	11	114	24
AD-1556911.1	20	1	39	3	42	8
AD-1571155.1	11	3	24	3	45	6
AD-1571156.1	8	1		0		
AD-1571025.1	40	8	49	4	56	8
AD-1556915.1	29	8	37	8	58	12
AD-1556917.1	22	4	39	6	55	9
AD-1571026.1	30	6	52	6	57	12
AD-1556918.1	18	4	33	9	46	5
AD-1571027.1	45	4	66	6	86	3
AD-1571157.1	18	7	37	8	57	10
AD-1571158.1	10	2	17	3	20	5
AD-1571159.1	18	0	22	5	42	4
AD-1571160.1	16	1	26	4	35	5
AD-1571161.1	27	3	31	10	55	12

Example 3. *In vivo* Efficacy of dsRNA Duplexes in Non-Human Primates (NHP)

Selected duplexes of interest, identified from the above *in vitro* studies, were evaluated *in vivo* in non-human primates. Figure 1 provides a depiction of the study design.

In particular, 15 male Cynomolgus monkeys were divided into 5 groups of 3 each and were subcutaneously administered a single 3 mg/kg dose of AD-1556360, a single 10 mg/kg dose of AD-1556360, a single 3 mg/kg dose of AD-1571158, or a single 3 mg/kg dose of AD-1571033, or PBS as a control (see Table 9). For each animal, two liver biopsy samples (one per lobe) of about 100 mg each were collected following 12 hours of fasting on Day 22, Day 57, and/or Day 85. Liver biopsy and serum samples were also collected from the animals 21 days prior to dosing. One mL of blood was collected into tubes without anticoagulant weekly from Day 1 for hepcidin level, iron level, transferrin saturation level, and red blood cell (RBC) count determinations. Following clotting, serum was aliquoted and stored at -80°C.

Tissue mRNA was extracted and analyzed by the RT-QPCR method. TMPRSS6 mRNA levels were compared to the levels of the housekeeping gene, GAPDH. The values were then normalized to the average of PBS vehicle control group. The data were expressed as percent of baseline value, and presented as mean plus standard deviation.

Iron and transferrin saturation levels were determined using commercially available kits from Roche.

The results, shown in Figures 2-4, demonstrate that all three exemplary duplexes, AD-1556360, AD-1571158, and AD-1571033, potently and durably inhibit the expression of TMPRSS6 messenger RNA *in vivo* (Figure 2), potently and durably lower plasma iron levels (Figure 3), and potently and durably lower transferrin saturation levels (Figure 4). Transferrin saturation is a measure of the amount of iron bound to serum transferrin, and corresponds to the ratio of serum iron and total iron-binding capacity.

10

Table 9. Treatment Groups

Group No.	Duplex	Dose Level (mg/kg)	No. of males
1	PBS (control)	0	3
2	AD-1556360	3	3
3	AD-1556360	10	3
4	AD-1571158	3	3
5	AD-1571033 (benchmark comparator duplex)	3	3

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments and methods described herein. Such equivalents are intended to be encompassed by the scope of the following claims.

We claim:

1. A double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of
5 Transmembrane protease, serine 6 (TMPRSS6) in a cell, wherein the dsRNA agent comprises a sense strand and an antisense strand forming a double stranded region, wherein the antisense strand comprises a region of complementarity to an mRNA encoding TMPRSS6, and wherein the region of complementarity comprises at least 15 contiguous nucleotides differing by no more than 3 nucleotides from any one of the antisense nucleotide sequences in any one of Tables 2-7.
- 10 2. The dsRNA agent of claim 1, wherein the dsRNA agent comprises a sense strand comprising at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the nucleotide sequences of the sense strands in any one of Tables 2-7 and an antisense strand comprising at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the
15 nucleotide sequences of the antisense strands in any one of Tables 2-7.
3. The dsRNA agent of claim 1, wherein the dsRNA agent comprises a sense strand comprising at least 15 contiguous nucleotides differing by no more than two nucleotides from any one of the nucleotide sequences of the sense strands in any one of Tables 2-7 and an antisense strand comprising
20 at least 15 contiguous nucleotides differing by no more than two nucleotides from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-7.
4. The dsRNA agent of claim 1, wherein the dsRNA agent comprises a sense strand comprising at least 15 contiguous nucleotides differing by no more than one nucleotide from any one of the
25 nucleotide sequences of the sense strands in any one of Tables 2-7 and an antisense strand comprising at least 15 contiguous nucleotides differing by no more than one nucleotide from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-7.
5. The dsRNA agent of claim 1, wherein the dsRNA agent comprises a sense strand comprising
30 a nucleotide sequence selected from the group consisting of any one of the nucleotide sequences of the sense strands in any one of Tables 2-7 and an antisense strand comprising a nucleotide sequence selected from the group consisting of any one of the nucleotide sequences of the antisense strands in any one of Tables 2-7.
- 35 6. A double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of Transmembrane protease, serine 6 (TMPRSS6) in a cell, wherein the dsRNA agent comprises a sense strand and an antisense strand forming a double stranded region, wherein the sense strand comprises at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the

nucleotide sequence of nucleotides 187-210; 227-254; 322-363; 362-390; 398-420; 404-429; 410-435; 439-461; 443-467; 448-474; 460-483; 466-488; 496-519; 519-542; 526-548; 557-593; 641-671; 652-676; 687-713; 725-762; 757-794; 886-908; 921-951; 956-987; 1051-1082; 1233-1269; 1279-1313; 1313-1341; 1327-1351; 1415-1439; 1447-1480; 1464-1486; 1486-1509; 1559-1589; 1571-1595; 1579-1609; 1707-1735; 1738-1764; 1806-1828; 1864-1886; 1934-1966; 1967-1991; 2008-2031; 2015-2043; 2042-2072; 2287-2311; 2297-2354; 2336-2361; 2360-2384; 2416-2438; 2481-2510; 2496-2527; 2526-2558; 2665-2693; 2693-2719; 2707-2729; 2799-2821; 2851-2874; 2971-2999; 2981-3006; and 3155-3195 of SEQ ID NO: 1, and the antisense strand comprises at least 15 contiguous nucleotides from the corresponding nucleotide sequence of SEQ ID NO:2.

10

7. A double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of Transmembrane protease, serine 6 (TMPRSS6) in a cell, wherein the dsRNA agent comprises a sense strand and an antisense strand forming a double stranded region, wherein the sense strand comprises at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the nucleotide sequence of nucleotides 230-252, 324-346, 560-578, 560-582, 2338-2360, 3163-3185, 3169-3191, and 3172-3194 of SEQ ID NO: 1, and the antisense strand comprises at least 15 contiguous nucleotides from the corresponding nucleotide sequence of SEQ ID NO:2.

15

8. A double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of Transmembrane protease, serine 6 (TMPRSS6) in a cell, wherein the dsRNA agent comprises a sense strand and an antisense strand forming a double stranded region, wherein the sense strand comprises at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the nucleotide sequence of nucleotides 560-578, 2338-2360, and 3169-3191 of SEQ ID NO: 1, and the antisense strand comprises at least 15 contiguous nucleotides from the corresponding nucleotide sequence of SEQ ID NO:2.

20

9. The dsRNA agent of any one of claims 1-8, wherein the antisense strand comprises at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1556360, AD-1571158, AD-1571033, AD-1554875, AD-1571160, AD-1555117, AD-1554911, and AD-1556915.

30

10. The dsRNA agent of any one of claims 1-9, wherein the antisense strand comprises at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1556360, AD-1571158, and AD-1571033.

35

11. The dsRNA agent of any one of claims 1-10, wherein the dsRNA agent comprises at least one modified nucleotide.
12. The dsRNA agent of any one of claims 1-11, wherein substantially all of the nucleotides of the sense strand; substantially all of the nucleotides of the antisense strand comprise a modification; or substantially all of the nucleotides of the sense strand and substantially all of the nucleotides of the antisense strand comprise a modification.
13. The dsRNA agent of any one of claims 1-12, wherein all of the nucleotides of the sense strand comprise a modification; all of the nucleotides of the antisense strand comprise a modification; or all of the nucleotides of the sense strand and all of the nucleotides of the antisense strand comprise a modification.
14. The dsRNA agent of any one of claims 11-13, wherein at least one of the modified nucleotides is selected from the group a deoxy-nucleotide, a 3'-terminal deoxythymidine (dT) nucleotide, a 2'-O-methyl modified nucleotide, a 2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a 2'-5'-linked ribonucleotide (3'-RNA), a locked nucleotide, an unlocked nucleotide, a conformationally restricted nucleotide, a constrained ethyl nucleotide, an abasic nucleotide, a 2'-amino-modified nucleotide, a 2'-O-allyl-modified nucleotide, 2'-C-alkyl-modified nucleotide, a 2'-methoxyethyl modified nucleotide, a 2'-O-alkyl-modified nucleotide, a morpholino nucleotide, a phosphoramidate, a non-natural base comprising nucleotide, a tetrahydropyran modified nucleotide, a 1,5-anhydrohexitol modified nucleotide, a cyclohexenyl modified nucleotide, a nucleotide comprising a 5'-phosphorothioate group, a nucleotide comprising a 5'-methylphosphonate group, a nucleotide comprising a 5' phosphate or 5' phosphate mimic, a nucleotide comprising vinyl phosphonate, a glycol nucleic acid (GNA), a glycol nucleic acid S-Isomer (S-GNA), a nucleotide comprising 2-hydroxymethyl-tetrahydrofuran-5-phosphate, a nucleotide comprising 2'-deoxythymidine-3' phosphate, a nucleotide comprising 2'-deoxyguanosine-3'-phosphate, and a terminal nucleotide linked to a cholesteryl derivative and a dodecanoic acid bisdecylamide group; and combinations thereof.
15. The dsRNA agent of any one of claims 11-13, wherein the modifications on the nucleotides are selected from the group consisting of LNA, HNA, CeNA, 2'-methoxyethyl, 2'-O-alkyl, 2'-O-allyl, 2'-C-allyl, 2'-fluoro, 2'-deoxy, 2'-hydroxyl, and glycol; and combinations thereof.
16. The dsRNA agent of any one of claims 11-13, wherein at least one of the modified nucleotides is selected from the group consisting of a deoxy-nucleotide, a 2'-O-methyl modified nucleotide, a 2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a glycol modified

nucleotide (GNA), a nucleotide comprising a 2' phosphate, and, a vinyl-phosphonate nucleotide; and combinations thereof.

17. The dsRNA agent of any one of claims 1-16, wherein the double stranded region is 19-30
5 nucleotide pairs in length.
18. The dsRNA agent of claim 17, wherein the double stranded region is 19-25 nucleotide pairs
in length.
- 10 19. The dsRNA agent of claim 17, wherein the double stranded region is 19-23 nucleotide pairs
in length.
20. The dsRNA agent of claim 17, wherein the double stranded region is 23-27 nucleotide pairs
in length.
15
21. The dsRNA agent of claim 17, wherein the double stranded region is 21-23 nucleotide pairs
in length.
22. The dsRNA agent of any one of claims 1-21, wherein each strand is independently no more
20 than 30 nucleotides in length.
23. The dsRNA agent of any one of claims 1-22, wherein the sense strand is 21 nucleotides in
length and the antisense strand is 23 nucleotides in length.
- 25 24. The dsRNA agent of any one of claims 1-23, wherein the region of complementarity is at
least 17 nucleotides in length.
25. The dsRNA agent of any one of claims 1-24, wherein the region of complementarity is
between 19 and 23 nucleotides in length.
30
26. The dsRNA agent of any one of claims 1-25, wherein the region of complementarity is 19
nucleotides in length.
27. The dsRNA agent of any one of claims 1-26, wherein at least one strand comprises a 3'
35 overhang of at least 1 nucleotide.

28. The dsRNA agent of any one of claims 1-26, wherein at least one strand comprises a 3' overhang of at least 2 nucleotides.

29. The dsRNA agent of any one of claims 1-28, further comprising a ligand.

5

30. The dsRNA agent of claim 29, wherein the ligand is conjugated to the 3' end of the sense strand of the dsRNA agent.

31. The dsRNA agent of claim 29, wherein the ligand is conjugated to the 5' end of the sense strand of the dsRNA agent.


10

32. The dsRNA agent of any one of claims 29-31, wherein the ligand is an N-acetylgalactosamine (GalNAc) derivative.

33. The dsRNA agent of any one of claims 29-32, wherein the ligand is one or more GalNAc derivatives attached through a monovalent, bivalent, or trivalent branched linker.


15

34. The dsRNA agent of claim 32 or 33, wherein the ligand is



20

35. The dsRNA agent of claim 34, wherein the dsRNA agent is conjugated to the ligand as shown in the following schematic




and, wherein X is O or S.

36. The dsRNA agent of claim 35, wherein the X is O.

5

37. The dsRNA agent of claim 34, wherein the dsRNA agent is conjugated to the ligand as shown in the following schematic



10 38. The dsRNA agent of any one of claims 1-37, wherein the dsRNA agent further comprises at least one phosphorothioate or methylphosphonate internucleotide linkage.

39. The dsRNA agent of claim 38, wherein the phosphorothioate or methylphosphonate internucleotide linkage is at the 3'-terminus of one strand.

15

40. The dsRNA agent of claim 39, wherein the strand is the antisense strand.

41. The dsRNA agent of claim 39, wherein the strand is the sense strand.

20 42. The dsRNA agent of claim 39, wherein the phosphorothioate or methylphosphonate internucleotide linkage is at the 5'-terminus of one strand.

43. The dsRNA agent of claim 42, wherein the strand is the antisense strand.
44. The dsRNA agent of claim 42, wherein the strand is the sense strand.
- 5 45. The dsRNA agent of claim 38, wherein the phosphorothioate or methylphosphonate internucleotide linkage is at both the 5'- and 3'-terminus of one strand.
46. The dsRNA agent of claim 45, wherein the strand is the antisense strand.
- 10 47. The dsRNA agent of any one of claims 1-46, comprising 6-8 phosphorothioate or methylphosphonate internucleotide linkages.
48. The dsRNA agent of any one of claims 1-47, wherein the base pair at the 1 position of the 5'-end of the antisense strand of the duplex is an AU base pair.
- 15 49. The dsRNA agent of any one of claims 1-48, wherein the sense strand comprises at least 17 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'-GACGCCACGCAUGCUGUGUGU -3'(SEQ ID NO: 119).
- 20 50. The dsRNA agent of any one of claims 1-49, wherein the sense strand comprises at least 19 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'-GACGCCACGCAUGCUGUGUGU -3'(SEQ ID NO: 119).
51. The dsRNA agent of any one of claims 1-50, wherein the sense strand comprises the
- 25 nucleotide sequence of 5'- GACGCCACGCAUGCUGUGUGU -3'(SEQ ID NO: 119).
52. The dsRNA agent of any one of claims 1-51, wherein the sense strand consists of the nucleotide sequence of 5'- GACGCCACGCAUGCUGUGUGU -3'(SEQ ID NO: 119).
- 30 53. The dsRNA agent of any one of claims 1-52, wherein the antisense strand comprises at least 17 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'-ACACACAGCAUGCUGGCGUCAC -3' (SEQ ID NO: 245).
54. The dsRNA agent of any one of claims 1-53, wherein the antisense strand comprises at least
- 35 19 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'-ACACACAGCAUGCUGGCGUCAC -3' (SEQ ID NO: 245).

55. The dsRNA agent of any one of claims 1-54, wherein the antisense strand comprises at least 21 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'-ACACACAGCAUGCGUGGCGUCAC -3' (SEQ ID NO: 245).

5 56. The dsRNA agent of any one of claims 1-55, wherein the antisense strand comprises the nucleotide sequence of 5'-ACACACAGCAUGCGUGGCGUCAC -3' (SEQ ID NO: 245).

57. The dsRNA agent of any one of claims 1-56, wherein the antisense strand consists of the nucleotide sequence of 5'-ACACACAGCAUGCGUGGCGUCAC -3' (SEQ ID NO: 245).

10

58. The dsRNA agent of any one of claims 1-57, wherein the sense strand comprises the nucleotide sequence of 5'-GACGCCACGCAUGCUGUGUGU -3' (SEQ ID NO: 119) and the antisense strand comprises the nucleotide sequence of 5'-ACACACAGCAUGCGUGGCGUCAC -3' (SEQ ID NO: 245).

15

59. The dsRNA agent of any one of claims 1-58, wherein the sense strand differs by no more than 3 modified nucleotides from the nucleotide sequence of 5'-gsascgccacGfCfAfugcugugugu-3' (SEQ ID NO: 371) wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage.

20

60. The dsRNA agent of any one of claims 1-59, wherein the antisense strand differs by no more than 3 modified nucleotides from the nucleotide sequence of 5'-asdCsacdAcdAgcaudGcGfugcgucsasc -3' (SEQ ID NO: 497), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein dA, dG, and dC are 2'-deoxyadenosine-3'-phosphate, 2'-deoxyguanosine-3'-phosphate, and 2'-deoxycytidine-3'-phosphate respectively; and s is a phosphorothioate linkage.

25

61. The dsRNA agent of any one of claims 1-60, wherein the sense strand comprises the nucleotide sequence of 5'-gsascgccacGfCfAfugcugugugu-3' (SEQ ID NO: 371) and the antisense strand comprises the nucleotide sequence of 5'-asdCsacdAcdAgcaudGcGfugcgucsasc -3' (SEQ ID NO: 497), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein dA, dG, and dC are 2'-deoxyadenosine-3'-phosphate, 2'-deoxyguanosine-3'-phosphate, and 2'-deoxycytidine-3'-phosphate respectively; and s is a phosphorothioate linkage.


30

35

62. The dsRNA agent of any one of claims 1-61, wherein the sense strand comprises the nucleotide sequence of 5'-gsascgccacGfCfAfugcuguguguL96 -3' (SEQ ID NO: 371) and the

antisense strand comprises the nucleotide sequence of 5'- asdCsacdAc dAgcaudGcGfugcgucsasc -3' (SEQ ID NO: 497), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein dA, dG, and dC are 2'-deoxyadenosine-3'-phosphate, 2'-deoxyguanosine-3'-phosphate, and 2'-deoxycytidine-3'-phosphate respectively; s is a phosphorothioate linkage, and L96 is N-[tris(GalNAc-alkyl)-amidodecanoyl]-4-hydroxyprolinol.

63. The dsRNA agent of any one of claims 1-61, wherein the sense strand comprises the nucleotide sequence of 5'- gsascgccacGfCfAfugcugugugu -3' (SEQ ID NO: 371) and the antisense strand comprises the nucleotide sequence of 5'- asdCsacdAc dAgcaudGcGfugcgucsasc -3' (SEQ ID NO: 497), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein dA, dG, and dC are 2'-deoxyadenosine-3'-phosphate, 2'-deoxyguanosine-3'-phosphate, and 2'-deoxycytidine-3'-phosphate respectively; and s is a phosphorothioate linkage, wherein the 3'-end of the sense strand is conjugated to the ligand as shown in the following schematic:



and, wherein X is O.

64. The dsRNA agent of any one of claims 1-48, wherein sense strand comprises at least 17 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'- CCUUUGGAAUAAAGCUGCCUU -3' (SEQ ID NO: 844).

65. The dsRNA agent of any one of claims 1-48 and 64, wherein the sense strand comprises at least 19 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'- CCUUUGGAAUAAAGCUGCCUU -3' (SEQ ID NO: 844).

66. The dsRNA agent of any one of claims 1-48, 64 and 65, wherein the sense strand comprises the nucleotide sequence of 5'- CCUUUGGAAUAAAGCUGCCUU -3' (SEQ ID NO: 844).

67. The dsRNA agent of any one of claims 1-48 and 64-66, wherein the sense strand consists of the nucleotide sequence of 5'-CCUUUGGAAUAAAGCUGCCUU-3' (SEQ ID NO: 844).
68. The dsRNA agent of any one of claims 1-48 and 64-67, wherein the antisense strand
5 comprises at least 17 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'-AAGGCAGCUUUAUUCCAAAGGGC-3' (SEQ ID NO: 1868).
69. The dsRNA agent of any one of claims 1-48 and 64-68, wherein the antisense strand
10 comprises at least 19 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'-AAGGCAGCUUUAUUCCAAAGGGC-3' (SEQ ID NO: 1868).
70. The dsRNA agent of any one of claims 1-48 and 64-69, wherein the antisense strand
comprises at least 21 contiguous nucleotides differing by no more than 3 nucleotides from the
nucleotide sequence of 5'-AAGGCAGCUUUAUUCCAAAGGGC-3' (SEQ ID NO: 1868).
15
71. The dsRNA agent of any one of claims 1-48 and 64-70, wherein the antisense strand
comprises the nucleotide sequence of 5'-AAGGCAGCUUUAUUCCAAAGGGC-3' (SEQ ID NO:
1868).
- 20 72. The dsRNA agent of any one of claims 1-48 and 64-71, wherein the antisense strand consists
of the nucleotide sequence of 5'-AAGGCAGCUUUAUUCCAAAGGGC-3' (SEQ ID NO: 1868).
73. The dsRNA agent of any one of claims 1-48 and 64-72, wherein the sense strand comprises
the nucleotide sequence of 5'-CCUUUGGAAUAAAGCUGCCUU-3' (SEQ ID NO: 844) and the
25 antisense strand comprises the nucleotide sequence of 5'-AAGGCAGCUUUAUUCCAAAGGGC-
3' (SEQ ID NO: 1868).
74. The dsRNA agent of any one of claims 1-48 and 64-73, wherein the sense strand differs by
no more than 3 modified nucleotides from the nucleotide sequence of 5'-
30 cscsuuugGfaAfUfAfaagcugccuu-3' (SEQ ID NO: 2095) wherein a, g, c and u are 2'-O-methyl (2'-
OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s
is a phosphorothioate linkage.
75. The dsRNA agent of any one of claims 1-48 and 64-74, wherein the antisense strand differs
35 by no more than 3 modified nucleotides from the nucleotide sequence of 5'-
asAfsggdCa(G2p)cuuuauUfcCfaagsgsc-3' (SEQ ID NO: 2324), wherein a, g, c and u are 2'-O-

methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein G2p is guanosine-2'-phosphate; and s is a phosphorothioate linkage.


76. The dsRNA agent of any one of claims 1-48 and 64-75, wherein the sense strand comprises the nucleotide sequence of 5'- cscsuuugGfaAfUfAfaagcugccuu -3' (SEQ ID NO: 2095) and the antisense strand comprises the nucleotide sequence of 5'- asAfsggdCa(G2p)cuuuauUfcCfaaaggsgsc -3' (SEQ ID NO: 2324), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein G2p is guanosine-2'-phosphate; and s is a phosphorothioate linkage.

10

77. The dsRNA agent of any one of claims 1-48 and 64-76, wherein the sense strand comprises the nucleotide sequence of 5'- cscsuuugGfaAfUfAfaagcugccuuL96 -3' (SEQ ID NO: 2095) and the antisense strand comprises the nucleotide sequence of 5'- asAfsggdCa(G2p)cuuuauUfcCfaaaggsgsc -3' (SEQ ID NO: 2324), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein G2p is guanosine-2'-phosphate; s is a phosphorothioate linkage, and L96 is N-[tris(GalNAc-alkyl)-amidodecanoyl]-4-hydroxyprolinol.

78. The dsRNA agent of any one of claims 1-48 and 64-76, wherein the sense strand comprises the nucleotide sequence of 5'- cscsuuugGfaAfUfAfaagcugccuu -3' (SEQ ID NO: 2095) and the antisense strand comprises the nucleotide sequence of 5'- asAfsggdCa(G2p)cuuuauUfcCfaaaggsgsc -3' (SEQ ID NO: 2324), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; wherein G2p is guanosine-2'-phosphate, s is a phosphorothioate linkage, and wherein the 3'-end of the sense strand is conjugated to the

25 ligand as shown in the following schematic:




and, wherein X is O.

79. The dsRNA agent of any one of claims 1-48, wherein the sense strand comprises at least 17 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'-UCACCUGCUUCUUCUGGUU-3' (SEQ ID NO: 1686).
- 5 80. The dsRNA agent of any one of claims 1-48 and 79, wherein the sense strand comprises the nucleotide sequence of 5'-UCACCUGCUUCUUCUGGUU-3' (SEQ ID NO: 1686).
81. The dsRNA agent of any one of claims 1-48, 79 and 80, wherein the sense strand consists of the nucleotide sequence of 5'-UCACCUGCUUCUUCUGGUU-3' (SEQ ID NO: 1686).
- 10 82. The dsRNA agent of any one of claims 1-48 and 79-81, wherein the antisense strand comprises at least 17 contiguous nucleotides differing by no more than 3 nucleotides from the nucleotide sequence of 5'-AACCAGAAGAAGCAGGUGA-3' (SEQ ID NO: 1790).
- 15 83. The dsRNA agent of any one of claims 1-48 and 79-82, wherein the antisense strand comprises the nucleotide sequence of 5'-AACCAGAAGAAGCAGGUGA-3' (SEQ ID NO: 1790).
84. The dsRNA agent of any one of claims 1-48 and 79-83, wherein the antisense strand consists of the nucleotide sequence of 5'-AACCAGAAGAAGCAGGUGA-3' (SEQ ID NO: 1790).
- 20 85. The dsRNA agent of any one of claims 1-48 and 79-84, wherein the sense strand comprises the nucleotide sequence of 5'-UCACCUGCUUCUUCUGGUU-3' (SEQ ID NO: 1686) and the antisense strand comprises the nucleotide sequence of 5'-AACCAGAAGAAGCAGGUGA-3' (SEQ ID NO: 1790).
- 25 86. The dsRNA agent of any one of claims 1-48 and 79-85, wherein the sense strand differs by no more than 3 modified nucleotides from the nucleotide sequence of 5'-UfcAfcCfuGfcUfuCfuUfcUfgGfsusUf-3' (SEQ ID NO: 1974), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage.
- 30 87. The dsRNA agent of any one of claims 1-48 and 79-86, wherein the antisense strand differs by no more than 3 modified nucleotides from the nucleotide sequence of 5'-asAfscCfaGfaAfgAfaGfcAfgGfusGfsa-3' (SEQ ID NO: 2203), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage.
- 35

88. The dsRNA agent of any one of claims 1-48 and 79-87, wherein the sense strand comprises the nucleotide sequence of 5'-UfcAfcCfuGfcUfuCfuUfcUfgGfsusUf-3' (SEQ ID NO: 1974) and the antisense strand comprises the nucleotide sequence of 5'-asAfscCfaGfaAfgAfaGfcAfgGfusGfsa-3' (SEQ ID NO: 2203), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage.

89. The dsRNA agent of any one of claims 1-48 and 79-88, wherein the sense strand comprises the nucleotide sequence of 5'-Q191sUfcAfcCfuGfcUfuCfuUfcUfgGfsusUf-3' (SEQ ID NO: 1974) and the antisense strand comprises the nucleotide sequence of 5'-asAfscCfaGfaAfgAfaGfcAfgGfusGfsa-3' (SEQ ID NO: 2203), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; s is a phosphorothioate linkage, and Q191 is N-[tris(GalNAc-alkyl)-amidododecanoyl]-(S)-pyrrolidin-3-ol-phosphorothioate (p-C12-(GalNAc-alkyl)3).

90. The dsRNA agent of any one of claims 1-48 and 79-88, wherein the sense strand comprises the nucleotide sequence of 5'-UfcAfcCfuGfcUfuCfuUfcUfgGfsusUf-3' (SEQ ID NO: 1974) and the antisense strand comprises the nucleotide sequence of 5'-asAfscCfaGfaAfgAfaGfcAfgGfusGfsa-3' (SEQ ID NO: 2203), wherein a, g, c and u are 2'-O-methyl (2'-OMe) A, G, C, and U respectively; Af, Gf, Cf and Uf are 2'-fluoro A, G, C and U respectively; and s is a phosphorothioate linkage, wherein the 5'-end of the sense strand is conjugated to the ligand as shown in the following schematic:



91. A cell containing the dsRNA agent of any one of claims 1-90.

92. A pharmaceutical composition for inhibiting expression of a gene encoding Transmembrane protease, serine 6 (TMPRSS6) comprising the dsRNA agent of any one of claims 1-90.

93. The pharmaceutical composition of claim 92, wherein dsRNA agent is in an unbuffered solution.

94. The pharmaceutical composition of claim 93, wherein the unbuffered solution is saline or water.
95. The pharmaceutical composition of claim 92, wherein said dsRNA agent is in a buffer
5 solution.
96. The pharmaceutical composition of claim 95, wherein the buffer solution comprises acetate, citrate, prolamine, carbonate, or phosphate or any combination thereof.
- 10 97. The pharmaceutical composition of claim 96, wherein the buffer solution is phosphate buffered saline (PBS).
98. A method of inhibiting expression of a Transmembrane protease, serine 6 (TMPRSS6) gene in a cell, the method comprising contacting the cell with the dsRNA agent of any one of claims 1-90,
15 or the pharmaceutical composition of any one of claims 92-97, thereby inhibiting expression of the TMPRSS6 gene in the cell.
99. The method of claim 98, wherein the cell is within a subject.
- 20 100. The method of claim 98, wherein the subject is a human.
101. The method of claim 98 or 100, wherein the subject has a TMPRSS6-associated disorder.
102. The method of claim 101, wherein the TMPRSS6-associated disorder is a disorder associated
25 with iron overload and/or a disorder of ineffective erythropoiesis.
103. The method of claim 101, wherein the TMPRSS6-associated disorder is selected from the group consisting of hereditary hemochromatosis, β -thalassemia, polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic
30 porphyria, Parkinson's Disease, Alzheimer's Disease and Friedreich's Ataxia.
104. The method of claim 101, wherein the TMPRSS6-associated disorder is β -thalassemia.
105. The method of claim 104, wherein the β -thalassemia is thalassemia major.
35
106. The method of claim 104, wherein the β -thalassemia is thalassemia intermedia.
107. The method of claim 101, wherein the TMPRSS6-associated disorder is polycythemia vera.

108. The method of any one of claims 98-107, wherein contacting the cell with the dsRNA agent inhibits the expression of TMPRSS6 by at least 50%, 60%, 70%, 80%, 90%, or 95%.

109. The method of any one of claims 98-108, wherein inhibiting expression of TMPRSS6
5 decreases TMPRSS6 protein level in serum of the subject by at least 50%, 60%, 70%, 80%, 90%, or 95%.

110. The method of any one of claims 98-109, wherein contacting the cell with the dsRNA agent increases the expression of hepcidin by at least 50%, 60%, 70%, 80%, 90%, or 95%.

10

111. The method of any one of claims 98-110, wherein increasing expression of hepcidin increases hepcidin protein level in serum of the subject by at least 50%, 60%, 70%, 80%, 90%, or 95%.

112. A method of treating a subject having a disorder that would benefit from reduction in Transmembrane protease, serine 6 (TMPRSS6) expression, comprising administering to the subject a therapeutically effective amount of the dsRNA agent of any one of claims 1-58, or the pharmaceutical composition of any one of claims 50-55, thereby treating the subject having the disorder that would benefit from reduction in TMPRSS6 expression.

20

113. A method of preventing at least one symptom in a subject having a disorder that would benefit from reduction in Transmembrane protease, serine 6 (TMPRSS6) expression, comprising administering to the subject a prophylactically effective amount of the dsRNA agent of any one of claims 1-48, or the pharmaceutical composition of any one of claims 50-55, thereby preventing at
25 least one symptom in the subject having the disorder that would benefit from reduction in TMPRSS6 expression.

114. The method of claim 112 or 113, wherein the disorder is a TMPRSS6-associated disorder.

115. The method of claim 114, wherein the TMPRSS6-associated disorder is a disorder associated with iron overload and/or a disorder of ineffective erythropoiesis.

116. The method of claim 115, wherein the TMPRSS6-associated disorder is selected from the group consisting of hereditary hemochromatosis, β -thalassemia, polycythemia vera, myelodysplastic syndrome, congenital dyserythropoietic anemias, pyruvate kinase deficiency, erythropoietic porphyria, Parkinson's Disease, Alzheimer's Disease and Friedreich's Ataxia.

35

117. The method of claim 114, wherein the TMPRSS6-associated disorder is β -thalassemia.
118. The method of claim 117, wherein the β -thalassemia is thalassemia major.
- 5 119. The method of claim 117, wherein the β -thalassemia is thalassemia intermedia.
120. The method of claim 114, wherein the TMPRSS6-associated disorder is polycythemia vera.
121. The method of any one of claims 112-120, wherein the subject is a human.
- 10 122. The method of any one of claims 112-121, wherein the administration of the agent to the subject causes a decrease in iron level, a decrease in ferritin level, a decrease in a transferrin saturation level, an increase in hemoglobin level, an increase in hematocrit level, and/or a decrease in TMPRSS6 protein accumulation.
- 15 123. The method of any one of claims 112-122, wherein the dsRNA agent is administered to the subject at a dose of about 0.01 mg/kg to about 50 mg/kg.
124. The method of any one of claims 112-123, wherein the dsRNA agent is administered to the
- 20 subject subcutaneously.
125. The method of any one of claims 112-123, wherein the dsRNA agent is administered to the subject intravenously.
- 25 126. The method of any one of claims 112-125, further comprising determining the level of TMPRSS6 in a sample(s) from the subject.
127. The method of claim 126, wherein the level of TMPRSS6 in the subject sample(s) is a TMPRSS6 protein level in a blood, serum or liver sample(s).
- 30 128. The method of any one of claims 112-127, further comprising determining the level of iron and/or hepcidin in a sample(s) from the subject.
129. The method of any one of claims 112-128, further comprising administering to the subject an
- 35 additional therapeutic agent for treatment of a TMPRSS6-associated disorder.
130. The method of claim 129, wherein the additional therapeutic agent is an iron chelator.

131 The method of claim 130, wherein the iron chelator is selected from the group consisting of deferiprone, deferoxamine, and deferasirox.

132. A kit comprising the dsRNA agent of any one of claims 1-90 or the pharmaceutical
5 composition of any one of claims 92-97.

133. A vial comprising the dsRNA agent of any one of claims 1-90 or the pharmaceutical composition of any one of claims 92-97.

10 134. A syringe comprising the dsRNA agent of any one of claims 1-90 or the pharmaceutical composition of any one of claims 92-97.

135. An RNA-induced silencing complex (RISC) comprising an antisense strand of the dsRNA agent of any one of claims 1-90.
15