

Description

This invention relates to fungicidal compositions containing cyclopentene, cyclohexanone and cyclohexenone derivatives, some of which are novel, a process for preparing such compounds and their use as fungicides.

Chem. Abs. 108: 131507s (1988) discloses 2-benzylcyclohex-2-en-1-one. However there is no suggestion in this abstract that this compound possesses any fungicidal activity.


Bull. Chem. Soc. Jap., 43(7), (1970), pp. 2204-8 discloses 1-benzyl-2-carboxycyclopent-1-ene and Bull. Chem. Soc. Jap., 60(2), (1987), pp. 836-8 discloses 1-benzyl-2-methoxycarbonylcyclopent-1-ene. However, there is no indication in these references that either of these compounds exhibits any fungicidal activity.

Chem. Abs. 62, (1965), p. 2728 discloses 2-(4-chlorobenzyl)dihydroresorcinol and 2-(4-bromobenzyl)dihydroresorcinol. However, this abstract gives no evidence that either of these compounds exhibits significant fungicidal activity against phytopathogenic fungi.

Moreover, Chem. Abs. 52: 13701i (1958) discloses 2-benzyl-3-ethoxy-5,5-dimethylcyclohex-2-en-1-one but, again, there is no indication that this compound has any fungicidal properties.


It has now been found that certain cyclopentene, cyclohexanone and cyclohexenone derivatives exhibit fungicidal activity in specific screens.

According to the present invention there is therefore provided a fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula




in which n represents an integer from 0 to 5; each R represents a halogen atom, nitro, cyano, hydroxyl, alkyl, haloalkyl, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, alkoxy carbonyl, carboxyl, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, carbamoyl, alkylamido, cycloalkyl or phenyl group; R^1 and R^2 independently represent a hydrogen atom or an alkyl group; and R^5 represents a hydrogen atom or an alkyl or cycloalkyl group.

Also according to the present invention there is provided a fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula




in which n , R , R^1 , R^2 and R^5 are as defined above.

Further according to the present invention there is provided a fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula




in which n, R, R¹ and R² are as defined above and X and Y independently represent a halogen atom.

According to another aspect of the present invention there is provided a compound of the general formula



25 in which n, R, R¹ and R² are as defined above.

According to a further aspect of the present invention there is provided a fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula



in which n, R, R¹ and R² are as defined above and R⁴ represents an alkyl group.

40 When any of the foregoing substituents represents or contains an alkyl substituent group, this may be linear or branched and may contain up to 12, preferably up to 6, and especially up to 4, carbon atoms. A cycloalkyl substituent group may contain 3 to 8, preferably 3 to 6, carbon atoms.

It is preferred that R¹ and R² independently represent a hydrogen atom or a C₁₋₄alkyl, particularly a methyl, group.

45 Preferably, R represents a halogen, especially a chlorine, atom.

It is also preferred that R⁵ represents a hydrogen atom or a C₁₋₆alkyl group. Most preferably, R⁵ represents a methyl group.

It is also preferred that R⁴ represents a primary or secondary alkyl group, primary alkyl groups being especially preferred. Preferably, R⁴ represents a C₁₋₄alkyl group, especially a primary C₁₋₄alkyl group. 50 Most preferably, R⁴ represents a methyl or butyl, especially an isobutyl, group.

It is further preferred that X and Y independently represent a chlorine or bromine atom. Most preferably, X and Y both represent a bromine atom.

A particularly preferred sub-group of compounds of formulae I, II, III, VI and VII is that in which n is 1, R represents a chlorine atom, preferably substituted at the 4-position of the phenyl ring, R¹ and R² both 55 represent a hydrogen atom or both represent a methyl group.

A method of making fungicidal compositions according to the invention is also provided which comprises bringing a compound of formula I, II, III, VI or VII into association with at least one carrier. Such compositions may contain a single compound or a mixture of several compounds of formulae I, II, III, VI or

VII.

A composition according to the invention preferably contains from 0.5 to 95% by weight of active ingredient.

A carrier, in a composition according to the invention is any material with which the active ingredient is formulated to facilitate application to the locus to be treated, which may for example be a plant, seed or soil, or to facilitate storage, transport or handling. A carrier may be a solid or a liquid, including a material which is normally gaseous but which has been compressed to form a liquid, and any of the carriers normally used in formulating fungicidal compositions may be used.

Suitable solid carriers include natural and synthetic clays and silicates, for example natural silicas such as diatomaceous earths; magnesium silicates, for example talcs; magnesium aluminium silicates, for example attapulgites and vermiculites; aluminium silicates, for example kaolinites, montmorillonites and micas; calcium carbonate; calcium sulphate; ammonium sulphate; synthetic hydrated silicon oxides and synthetic calcium or aluminium silicates; elements, for example carbon and sulphur; natural and synthetic resins, for example coumarone resins, polyvinyl chloride, and styrene polymers and copolymers; solid polychlorophenols; bitumen; waxes, for example beeswax, paraffin wax, and chlorinated mineral waxes; and solid fertilisers, for example superphosphates.

Suitable liquid carriers include water; alcohols, for example isopropanol and glycols; ketones, for example acetone, methyl ethyl ketone, methyl isobutyl ketone and cyclohexanone; ethers; aromatic or araliphatic hydrocarbons, for example benzene, toluene and xylene; petroleum fractions, for example, kerosine and light mineral oils; chlorinated hydrocarbons, for example carbon tetrachloride, perchloroethylene and trichloroethane. Mixtures of different liquids are often suitable.

Fungicidal compositions are often formulated and transported in a concentrated form which is subsequently diluted by the user before application. The presence of small amounts of a carrier which is a surface-active agent facilitates this process of dilution. Thus preferably at least one carrier in a composition according to the invention is a surface-active agent. For example the composition may contain at least two carriers, at least one of which is a surface-active agent.

A surface-active agent may be an emulsifying agent, a dispersing agent or a wetting agent; it may be nonionic or ionic. Examples of suitable surface-active agents include the sodium or calcium salts of polyacrylic acids and lignin sulphonic acids; the condensation products of fatty acids or aliphatic amines or amides containing at least 12 carbon atoms in the molecule with ethylene oxide and/or propylene oxide; fatty acid esters of glycerol, sorbitol, sucrose or pentaerythritol; condensates of these with ethylene oxide and/or propylene oxide; condensation products of fatty alcohol or alkyl phenols, for example p-octylphenol or p-octylcresol, with ethylene oxide and/or propylene oxide; sulphates or sulphonates of these condensation products; alkali or alkaline earth metal salts, preferably sodium salts, of sulphuric or sulphonic acid esters containing at least 10 carbon atoms in the molecule, for example sodium lauryl sulphate, sodium secondary alkyl sulphates, sodium salts of sulphonated castor oil, and sodium alkylaryl sulphonates such as dodecylbenzene sulphonate; and polymers of ethylene oxide and copolymers of ethylene oxide and propylene oxide.

The compositions of the invention may for example be formulated as wettable powders, dusts, granules, solutions, emulsifiable concentrates, emulsions, suspension concentrates and aerosols. Wettable powders usually contain 25, 50 or 75% w of active ingredient and usually contain in addition to solid inert carrier, 3-10% w of a dispersing agent and, where necessary, 0-10% w of stabiliser(s) and/or other additives such as penetrants or stickers. Dusts are usually formulated as a dust concentrate having a similar composition to that of a wettable powder but without a dispersant, and may be diluted in the field with further solid carrier to give a composition usually containing $\frac{1}{2}$ -10% w of active ingredient. Granules are usually prepared to have a size between 10 and 100 BS mesh (1.676 - 0.152 mm), and may be manufactured by agglomeration or impregnation techniques. Generally, granules will contain $\frac{1}{2}$ -75% w active ingredient and 0-10% w of additives such as stabilisers, surfactants, slow release modifiers and binding agents. The so-called "dry flowable powders" consist of relatively small granules having a relatively high concentration of active ingredient. Emulsifiable concentrates usually contain, in addition to a solvent and, when necessary, co-solvent, 1-50% w/v active ingredient, 2-20% w/v emulsifiers and 0-20% w/v of other additives such as stabilisers, penetrants and corrosion inhibitors. Suspension concentrates are usually compounded so as to obtain a stable, non-sedimenting flowable product and usually contain 10-75% w active ingredient, 0.5-15% w of dispersing agents, 0.1-10% w of suspending agents such as protective colloids and thixotropic agents, 0-10% w of other additives such as defoamers, corrosion inhibitors, stabilisers, penetrants and stickers, and water or an organic liquid in which the active ingredient is substantially insoluble; certain organic solids or inorganic salts may be present dissolved in the formulation to assist in preventing sedimentation or as anti-freeze agents for water.

Aqueous dispersions and emulsions, for example compositions obtained by diluting a wettable powder or a concentrate according to the invention with water, also lie within the scope of the invention. The said emulsions may be of the water-in-oil or of the oil-in-water type, and may have a thick 'mayonnaise' like consistency.

5 The composition of the invention may also contain other ingredients, for example other compounds possessing herbicidal, insecticidal or fungicidal properties.

Of particular interest in enhancing the duration of the protective activity of the compounds of this invention is the use of a carrier which will provide a slow release of the fungicidal compounds into the environment of the plant which is to be protected. Such slow-release formulations could, for example, be
10 inserted in the soil adjacent to the roots of a plant, or could include an adhesive component enabling them to be applied directly to the stem of a plant.

According to the present invention there is also provided a compound of the general formula II or III in which n, R, R¹, R², R⁵, X and Y are as defined above.


According to another aspect of the present invention there is provided a compound of the general
15 formula I in which n, R, R¹, R² and R⁵ are as defined above with the proviso that, when n is 0 and R¹ and R² both represent a hydrogen atom, R⁵ does not represent a hydrogen atom or a methyl group.

Also according to the present invention there is provided a compound of the general formula VI in which n, R, R¹ and R² are as defined above with the proviso that, when n is 0, R¹ and R² do not both represent a hydrogen atom.


20 According to a further aspect of the present invention there is provided a compound of the general formula VII in which n, R, R¹, R² and R⁴ are as first defined above with the provisos that, when n is 1, R¹ and R² both represent a hydrogen atom and R⁴ represents a methyl group then R does not represent a chlorine or bromine atom substituted at the 4-position of the phenyl ring, and, when n is 0 and R¹ and R² both represent a methyl group, R⁴ does not represent an ethyl group.

25 The invention still further provides the use as a fungicide of a composition as defined above or a compound of formula I, II, III, VI or VII as defined above, and a method for combating fungus at a locus, which comprises treating the locus, which may be for example plants subject to or subjected to fungal attack, seeds of such plants or the medium in which such plants are growing or are to be grown, with such a compound or composition.

30 The present invention also provides a process for the preparation of a compound of formula I as defined above which comprises heating a compound of the general formula



40 or the general formula



in which n, R, R¹, R² and R⁵ are as first defined above with the proviso that, when n is 0 and R¹ and R²
55 both represent a hydrogen atom, R⁵ does not represent a hydrogen atom or a methyl group, and X and Y are as defined above with a compound of the general formula

MOR⁵ (IV)

in which R^5 is as first defined above and M represents an alkali metal, preferably a sodium, atom, in the presence of a polar solvent.

Preferably, the polar solvent is a compound of the general formula




in which R^5 is as first defined above, dimethylformamide or dimethylsulphoxide.

If a compound of formula V is used as solvent, preferably, R^5 has the same meaning in formula IV and formula V. For instance, if the compound of formula IV is sodium methoxide, it is preferred that the solvent of formula V is methanol.

The reaction is conveniently carried out at a temperature from 80 °C to the reflux temperature of the solvent. Preferably, an excess of the compound of formula IV is used.

The present invention further provides a process for the preparation of a compound of formula II as defined above which comprises reacting a compound of the general formula



in which n, R, R^1 and R^2 are as first defined above and X and Y are as defined above with a compound of the general formula



in which R^5 is as first defined above and M is as defined above in the presence of a solvent of the general formula




in which R^5 is as first defined above.

Preferably, R^5 has the same meaning in formula IV and formula V. For instance, if the compound of formula IV is sodium methoxide, it is preferred that the solvent of formula V is methanol.

The reaction is conveniently carried out at a temperature from 0 °C to the reflux temperature of the solvent V, preferably, using an excess of the compound of formula IV.

The present invention still further provides a process for the preparation of a compound of formula III as defined above which comprises reacting a compound of the general formula



in which n, R, R^1 and R^2 are as first defined above, with a compound XY, in which X and Y are as defined above. Alternatively, compounds of formula III may be generated *in situ* and then treated with a compound of formula IV as described above to form compounds of formula I in a one-pot synthesis.


The process may be carried out in the presence of a solvent. Suitable solvents include petroleum, lower alcohols, such as methanol, chlorinated hydrocarbons, such as carbon tetrachloride, ethers and acetic acid.

The reaction is suitably carried out at a temperature from -10°C to room temperature, depending on the nature of the solvent, if present. The preferred temperature is from 0°C to room temperature.

The present invention also provides a process for the preparation of a compound of formula VI as defined above which comprises reacting a compound of the general formula

5

10



(VII)

15

in which n , R , R^1 , R^2 and R^4 are as first defined above, with a reducing agent and subsequently hydrolysing the reaction mixture.

20

The reducing agent is preferably a strong reducing agent, such as a complex metal hydride or hydrogen in combination with a catalyst. Complex metal hydrides, such as lithium aluminium hydride, sodium aluminium hydride and modified derivatives thereof such as "REDAL" (Trade Mark: sodium bis (2-methoxyethoxy) aluminium hydride in toluene), are especially preferred.

The process is conveniently carried out in the presence of a solvent. Suitable solvents include ethers, such as diethyl ether, tetrahydrofuran and oligoethers, and hydrocarbons.

25

The reaction is suitably carried out at a temperature from 20°C to 80°C , depending on the nature of the solvent, if present.

It is advisable to destroy any excess reducing agent remaining at the end of the reduction process to prevent further reaction. If a complex metal hydride, such as lithium aluminium hydride, is used as reducing agent, any excess may be destroyed by the addition of water and sodium hydroxide to the reaction mixture.


30

This process for preparing compounds of formula VI proceeds via an enol ether intermediate. Accordingly, to obtain compounds of formula VI, it is necessary to include a hydrolysis work-up in the above process. This may comprise the addition of a dilute mineral acid, such as hydrochloric acid, to the reaction mixture after the reduction step and serves to convert any remaining intermediate enol ether into the desired cyclohexenone derivative of formula VI.

35

The present invention further provides a process for the preparation of a compound of formula VII as defined above which comprises reacting a compound of the general formula

40



(VIII)

45

in which n , R , R^1 and R^2 are as first defined above, with a first compound of the general formula



50

in which R^4 is as first defined above, with the provisos that, when n is 1, R^1 and R^2 both represent a hydrogen atom and R^4 represents a methyl group then R does not represent a chlorine or bromine atom substituted at the 4-position of the phenyl ring, and, when n is 0 and R^1 and R^2 both represent a methyl group, R^4 does not represent an ethyl group, in the presence of an acid; and, if desired, converting the compound of formula VII so obtained into another compound of formula VII by reaction with a second different compound of formula IX in the presence of an acid.

55

The acid may be any substance which acts as a source of protons. Suitable acids include inorganic acids such as sulphuric acid, organic acids such as p-toluenesulphonic acid and ion exchange resins.


The process may be carried out in the presence of a solvent. Suitable solvents include petroleum, toluene and benzene. Solvents which form azeotropic mixtures with the reactants are particularly preferred.

The reaction is suitably carried out at a temperature from 70°C to 130°C, depending on the nature of the solvent, if present. The preferred temperature is from 80°C to 120°C.

Compounds of formula VIII may be conveniently prepared by reacting a compound of the general formula




in which R¹ and R² are as first defined above, with a compound of the general formula



in which R and n are as first defined above and L represents a suitable leaving group, in the presence of a suitable base, such as potassium hydroxide.



Compounds of formula IV, V, IX, X and XI and the compounds XY are known compounds or can be prepared by processes analogous to known processes.

The compounds of formulae I, II, III, VI and VII are also useful as intermediates in the preparation of fungicidally active cyclopentane derivatives of the general formula



in which n, R, R¹ and R² are as first defined above and A represents a nitrogen atom or a CH group. Certain compounds of formula XII are the subject of co-pending patent applications GB-A1-2180236 and EP-A2-0267778.


The compounds disclosed in EP-A2-0267778 and GB-A1-2180236 exist in two stereoisomeric forms which have the following structures:-




30 The letters A and B will be used hereinafter to denote compounds having the same stereochemical configuration as isomers A and B above.

Isomers A and B can be separated by, for instance, chromatography and exhibit different fungicidal activity. Generally, isomers of formula XIIA exhibit greater fungicidal activity than isomers of formula XIIB. The process used to synthesise compounds of formula XIIA from compounds of formulae I, II, III, VI and VII is set out in the following reaction scheme:-

35





In the above reaction scheme, n , R , R^1 , R^2 , R^4 , R^5 , X , Y , M and A are as previously defined, R^3 represents an optionally substituted alkyl or aryl group, preferably a C_{1-4} alkyl or a phenyl group each optionally substituted by one or more substituents selected from halogen atoms, nitro, cyano, hydroxyl, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkoxy, amino, C_{1-4} alkylamino, di- C_{1-4} alkylamino, C_{1-4} alkoxycarbonyl, carboxyl, C_{1-4} alkanoyl, C_{1-4} alkylthio, C_{1-4} alkylsulphinyl, C_{1-4} alkylsulphonyl, carbamoyl, C_{1-4} alkylamido, C_{3-8} cycloalkyl and phenyl groups, and Q represents a hydrogen or alkali metal, preferably sodium, atom. The intermediate compounds and process steps in the above reaction scheme which are not part of the present invention are the subject of copending patent applications EP-A-0361560, EP-A-0359305 and EP-A-0357404.

The invention is further illustrated by the following Examples.

Example 1

Preparation of 2-(4-chlorobenzyl)-3-(2-methylpropoxy)-5,5-dimethylcyclohex-2-en-1-one
(Formula VII: $n = 1$, $R = 4\text{-Cl}$, $R^1 = R^2 = \text{CH}_3$, $R^3 = (\text{CH}_3)_2\text{CH-CH}_2\text{O}$)

(a) Preparation of 2-(4-chlorobenzyl)-5,5-dimethylcyclohexane-1,3-dione

449g (3.21 mols) dimedone (5,5-dimethylcyclohexane-1,3-dione) were added to a solution of aqueous potassium hydroxide comprising 166g of 85% potassium hydroxide (2.52 moles) in 700 ml of water. The mixture was then warmed and a clear orange solution was obtained at 47°C. The solution was then heated to 59°C and 544g (3.21 mols) molten 4-chlorobenzyl chloride were added over a period of 1 hour with further heating to 85°C. Heating was continued for a further 2½ to 3 hours up to a temperature of 100°C. The mixture was then cooled, the solid product filtered off, washed with water and dried in a vacuum oven at 50°C. The crude solid (815g) was then dissolved in 2400ml methanol at reflux and 200ml water added to produce a permanent cloudiness. The mixture was then allowed to cool to room temperature overnight with stirring. The solid so obtained was filtered, washed with about 400ml cold methanol and dried in a vacuum oven to produce 340g 2-(4-chlorobenzyl)-5,5-dimethylcyclohexane-1,3-dione as a white solid, m.pt. 188-190°C. Yield: 42%.

(b) Preparation of 2-(4-chlorobenzyl)-3-(2-methylpropoxy)-5,5-dimethylcyclohex-2-en-1-one

325g (1.23 mol) of the 2-(4-chlorobenzyl)-5,5-dimethylcyclohexane-1,3-dione obtained in (a), 1.6 litres toluene, 182g (2.5 mol) isobutanol and 5g p-toluenesulphonic acid were stirred together at reflux under a Dean-Stark apparatus. The temperature of the reaction mixture was approximately 90°C. As water distilled off, the reaction mixture changed from a thin slurry to a yellow solution. After 14 hours reflux, the reaction mixture was cooled and shaken twice with 500ml aliquots of 10% aqueous sodium hydroxide. The toluene layer was then flashed to give 389g yellow/orange oil which crystallised on standing. Recrystallisation of the solid from 60/80 petroleum produced 331g 2-(4-chlorobenzyl)-3-(2-methylpropoxy)-5,5-dimethylcyclohex-2-en-1-one as a white crystalline solid, m.pt. 60-61°C. Yield: 84%.

Example 2

Preparation of 2-(4-chlorobenzyl)-3-methoxy-5,5-dimethylcyclohex-2-en-1-one
(Formula VII: $n = 1$, $R = 4\text{-Cl}$, $R^1 = R^2 = \text{CH}_3\text{O-}$)

A solution of 154g of the 2-(4-chlorobenzyl)-3-(2-methylpropoxy)-5,5-dimethylcyclohex-2-en-1-one obtained in Example 1 in 1200 ml methanol containing 3g p-toluenesulphonic acid was refluxed for 2 hours. The reaction mixture was then extracted with 3 litres water and 1 litre diethyl ether and re-extracted with a further 1 litre diethyl ether. The organic phases were then back-washed first with 200 ml 10% aqueous sodium hydroxide and then with 100 ml saturated sodium chloride solution, dried over anhydrous magnesium sulphate and flashed. The residue was then crystallised in 60/80 petroleum, filtered and air-dried to give 98g 2-(4-chlorobenzyl)-3-methoxy-5,5-dimethylcyclohex-2-en-1-one as a white solid, m.pt. 62-63°C. Yield: 73%.

Example 3

Preparation of 2-(4-chlorobenzyl)-5,5-dimethylcyclohex-2-en-1-one
(Formula VI: $n = 1$, $R = 4\text{-Cl}$, $R^1 = R^2 = \text{CH}_3$)

98g (0.35 mol) of the 2-(4-chlorobenzyl)-3-methoxy-5,5-dimethylcyclohex-2-en-1-one obtained in Example 2 were added to a slurry of 6.65g (0.175 mol) lithium aluminium hydride in 490 mls diethyl ether at a rate sufficient to maintain reflux and the final reaction mixture refluxed for a further 30 minutes. 5ml water were then added, followed by 5ml 15% aqueous sodium hydroxide and a further 15ml water and the resulting precipitate was filtered off. The filtrate was then shaken in 200ml 5M hydrochloric acid for five minutes and the organic layer then separated, washed twice with 100ml aliquots of saturated sodium bicarbonate solution, dried over anhydrous magnesium sulphate and stripped. The resulting oil was then dissolved in 430mls dichloromethane, 18g (0.085 mols) pyridinium chlorochromate were added, and the reaction mixture stirred for 3 hours. 600ml diethyl ether were added and the solid was then filtered off. The

filtrate was washed three times with 10% sodium hydroxide, once with 2.5M hydrochloric acid and once with saturated sodium bicarbonate solution. It was then dried over anhydrous magnesium sulphate and stripped to give 82g of crude product. Distillation of the crude product under reduced pressure (0.15mm mercury) gave 79g 2-(4-chlorobenzyl)-5,5-dimethylcyclohex-2-en-1-one, b.pt. 130 °C at 0.15mm mercury.

5 Yield: 91%.

Example 4

10 Preparation of 2-(4-chlorobenzyl)-2,3-dibromo-5,5-dimethylcyclohexan-1-one
(Formula III: $n = 1$, $R = 4\text{-Cl}$, $R^1 = R^2 = \text{CH}_3$, $X = Y = \text{Br}$)

10g (40.2mmols) of the 2-(4-chlorobenzyl)-5,5-dimethylcyclohex-2-en-1-one obtained in Example 3 were dissolved in 50ml 30/40 petroleum at 0 °C. 6.72g (40.2mmols) bromine were then added to the solution. After 5 - 10 minutes the solution decolourised and a precipitate formed. The solution was then cooled
15 further and the precipitate filtered off to give 12.4g 2-(4-chloro-benzyl)-2,3-dibromo-5,5-dimethylcyclohexan-1-one as a solid, m.pt. 82 - 84 °C. Yield: 75%.

Example 5

20 Preparation of 1-(4-chlorobenzyl)-3,3-dimethyl-2-methoxycarbonylcyclopent-1-ene
(Formula I: $n = 1$, $R = 4\text{-Cl}$, $R^1 = R^2 = \text{CH}_3$, $R^5 = \text{CH}_3$)

A solution of sodium methoxide was prepared by adding 2.8g (121 mmol) sodium to 50ml methanol. A slurry of the 2-(4-chlorobenzyl)-2,3-dibromo-5,5-dimethylcyclohexan-1-one obtained in Example 4 in methanol was then prepared and added to the sodium methoxide solution at reflux. Reflux was continued
25 overnight. The reaction mixture was then quenched with 200ml water, extracted twice with 100ml aliquots of diethyl ether, backwashed with water, dried over anhydrous magnesium sulphate and flashed to give 8g of a yellow oil. By gas chromatography analysis, it was established that 6.6g 1-(4-chlorobenzyl)-3,3-dimethyl-2-methoxycarbonyl-cyclopent-1-ene were produced as an oil. The structure of the product was established by
30 n.m.r. spectroscopy. Yield: 78%.

Example 6

35 Preparation of 2-(4-chlorobenzyl)-2,3-dibromo-5,5-dimethylcyclohexan-1-one
(Formula III: $n = 1$, $R = 4\text{-Cl}$, $R^1 = R^2 = \text{CH}_3$, $X = Y = \text{Br}$)

5g of the 2-(4-chlorobenzyl)-5,5-dimethylcyclohex-2-en-1-one obtained in Example 3 were dissolved in 25ml tetrachloromethane at 5-10 °C. 3.2g bromine were then added to the solution over a period of 10 minutes. The solution decolourised and 2-(4-chlorobenzyl)-2,3-dibromo-5,5-dimethylcyclohexan-1-one
40 formed in solution.

Example 7

45 Preparation of 2-(4-chlorobenzyl)-4,4-dimethyl-3-methoxycarbonylcyclopent-1-ene
(Formula II: $n = 1$, $R = 4\text{-Cl}$, $R^1 = R^2 = \text{CH}_3$, $R^5 = \text{CH}_3$)

A solution of sodium methoxide was prepared by adding 1.1g sodium to 10ml methanol. 50ml methanol was then added to the reaction mixture obtained in Example 6 followed by the solution of sodium methoxide, keeping the temperature of the reaction mixture at 10-15 °C. After 2 hours, the mixture was
50 partitioned between dichloromethane and water, backwashed with water, dried over anhydrous magnesium sulphate and flashed to give 5.65g crude 2-(4-chlorobenzyl)-4,4-dimethyl-3-methoxycarbonylcyclopent-1-ene as an oil. The structure of the product was established by n.m.r. spectroscopy.

Example 8

Preparation of 1-(4-chlorobenzyl)-3,3-dimethyl-2-methoxycarbonylcyclopent-1-ene
(Formula I: $n=1$, $R=4\text{-Cl}$, $R^1=R^2=\text{CH}_3$, $R^5=\text{CH}_3$)

The crude 2-(4-chlorobenzyl)-4,4-dimethyl-3-methoxycarbonylcyclopent-1-ene obtained in Example 7 was refluxed for 15 hours in methanol containing one equivalent of sodium methoxide to give 1-(4-chlorobenzyl)-3,3-dimethyl-2-methoxycarbonylcyclopent-1-ene as an oil.

10 Example 9

The fungicidal activity of compounds of the invention was investigated in specific screens and by means of the following tests.

(a) Antisporulant activity against vine downy mildew (*Plasmopara viticola*; Pva)

The test is a direct antisporulant one using a foliar spray. The lower surfaces of leaves of whole vine plants (cv Cabernet Sauvignon) are inoculated by spraying with an aqueous suspension containing 10^4 zoospores/ml 2 days prior to treatment with the test compound. The inoculated plants are kept for 24 hours in a high humidity compartment, then 24 hours at glasshouse ambient temperature and humidity. Infected leaves are sprayed on their lower surfaces with a solution of active material in 1:1 water/acetone containing 0.04% "TWEEN 20" (Trade Mark; a polyoxyethylene sorbitan ester surfactant). The spraying is carried out with a moving track sprayer giving an application rate of 1kg/ha. After spraying, the plants are returned to normal glasshouse conditions for 96 hours and are then transferred to the high humidity compartment for 24 hours to induce sporulation, prior to assessment. Assessment is based on the percentage of the leaf area covered by sporulation compared with that on control leaves.

(b) Direct protectant activity against vine downy mildew (*Plasmopara viticola*; Pvp)

The test is a direct protectant one using a foliar spray. The lower surfaces of leaves of whole vine plants (cv Cabernet Sauvignon) are sprayed with the test compound at a dosage of 1 kilogram of active material per hectare using a track sprayer as described under (a), and after a subsequent 24 hours under normal glasshouse conditions the lower surfaces of the leaves are inoculated by spraying with an aqueous solution containing 10^4 zoospores/ml. The inoculated plants are kept for 24 hours in a high humidity compartment, 5 days under normal glasshouse conditions and then returned for a further 24 hours to high humidity. Assessment is based on the percentage of leaf area covered by sporulation compared with that on control leaves.

(c) Activity against barley powdery mildew (*Erysiphe graminis* f.sp. *hordei*; Eg)

The test is a direct therapeutic one, using a foliar spray. Leaves of barley seedlings, (cv. Golden Promise) are inoculated by dusting with mildew conidia one day prior to treatment with the test compound. The inoculated plants are kept overnight at glasshouse ambient temperature and humidity prior to treatment. The plants are sprayed with the test compound at a dosage of 1 kilogram of active material per hectare using a track sprayer as described under (a). After drying, plants are returned to a compartment at 20-25 °C and moderate humidity for up to 7 days, followed by assessment. Assessment is based on the percentage of leaf area covered by sporulation compared with that on leaves of control plants.

(d) Activity against wheat brown rust (*Puccinia recondita*; Pr)

The test is a direct protectant one using a foliar spray. Wheat seedlings (cv Brigand) are grown to the 1-1½ leaf stage. The plants are then sprayed with the test compound at a dosage of 1 kg/ha using a track sprayer as described under (a). Test compounds are applied as solutions or suspensions in a mixture of acetone and water (50:50 v/v) containing 0.04% surfactant ("TWEEN 20" - Trade Mark).

18-24 hours after treatment, the seedlings are inoculated by spraying the plants from all sides with an aqueous spore suspension containing about 10^5 spores/ml. For 18 hours after inoculation, the plants are kept in high humidity conditions at a temperature of 20-22 °C. Thereafter, the plants are kept in ambient glasshouse conditions, that is, in moderate relative humidity and at a temperature of 20 °C.

The disease is assessed 10 days after inoculation on the basis of the percentage of the plant covered by sporulating pustules compared with that on the control plants.

(e) Activity against *Fusarium* in-vitro (*Fusarium* species; Fsl)

This test measures the in vitro activity of compounds against a species of Fusarium that causes stem and root rots.

Compound is dissolved or suspended in acetone and added to molten half strength Potato Dextrose Agar to give a final concentration of 100ppm compound and 3.5% acetone. After the agar has set, plates

are inoculated with 6mm diameter plugs of agar and mycelium taken from a 7 day old culture of *Fusarium* sp..

Plates are incubated at 20 °C for 5 days and radial growth from the plug is measured.

The extent of disease control in all the above tests is expressed as a rating compared with either an untreated control or a diluent-sprayed-control, according to the criteria:-

0 = less than 50% disease control

1 = about 50-80% disease control


2 = greater than 80% disease control

The results of these tests are set out in the Table below:-

Compound Example No.	Fungicidal Activity				
	Pva	Pvp	Eg	Pr	Fsl
2 4/6	1	2	1	2	1


Claims

1. A fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula



in which n represents an integer from 0 to 5; each R represents a halogen atom, nitro, cyano, hydroxyl, alkyl, haloalkyl, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, alkoxycarbonyl, carboxyl, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, carbamoyl, alkylamido, cycloalkyl or phenyl group; R¹ and R² independently represent a hydrogen atom or an alkyl group; and R⁵ represents a hydrogen atom or an alkyl or cycloalkyl group.

2. A fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula



in which n, R, R¹, R² and R⁵ are as defined in claim 1.

3. A fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula



are inoculated with 6mm diameter plugs of agar and mycelium taken from a 7 day old culture of *Fusarium* sp..

Plates are incubated at 20 °C for 5 days and radial growth from the plug is measured.

The extent of disease control in all the above tests is expressed as a rating compared with either an untreated control or a diluent-sprayed-control, according to the criteria:-


- 0 = less than 50% disease control
- 1 = about 50-80% disease control
- 2 = greater than 80% disease control

The results of these tests are set out in the Table below:-

Compound Example No.	Fungicidal Activity				
	Pva	Pvp	Eg	Pr	Fsl
2 4/6	1	2	1	2	1


Claims

1. A fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula




in which n represents an integer from 0 to 5; each R represents a halogen atom, nitro, cyano, hydroxyl, alkyl, haloalkyl, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, alkoxycarbonyl, carboxyl, alkanoyl, alkylthio, alkylsulphanyl, alkylsulphonyl, carbamoyl, alkylamido, cycloalkyl or phenyl group; R¹ and R² independently represent a hydrogen atom or an alkyl group; and R⁵ represents a hydrogen atom or an alkyl or cycloalkyl group.

2. A fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula




in which n, R, R¹, R² and R⁵ are as defined in claim 1.

3. A fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula




in which n, R, R¹ and R² are as defined in claim 1 and X and Y independently represent a halogen atom.

- 15 4. A fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula



25 in which n, R, R¹ and R² are as defined in claim 1.

- 30 5. A fungicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula




40 in which n, R, R¹ and R² are as defined in claim 1 and R⁴ represents an alkyl group.

- 45 6. A compound of the general formula II or III as defined in claim 2 or claim 3.
7. A compound of the general formula I in which n, R, R¹, R² and R⁵ are as defined in claim 1 with the proviso that, when n is 0 and R¹ and R² both represent a hydrogen atom, R⁵ does not represent a hydrogen atom or a methyl group.
- 50 8. A compound of the general formula VI in which n, R, R¹ and R² are as defined in claim 4 with the proviso that, when n is 0, R¹ and R² do not both represent a hydrogen atom.
9. A compound of the general formula VII in which n, R, R¹, R² and R⁴ are as defined in claim 5 with the provisos that, when n is 1, R¹ and R² both represent a hydrogen atom and R⁴ represents a methyl group then R does not represent a chlorine or bromine atom substituted at the 4-position of the phenyl ring, and, when n is 0 and R¹ and R² both represent a methyl group, R⁴ does not represent an ethyl group.
- 55 10. A method of combating fungus at a locus which comprises treating the locus with a composition as defined in any one of claims 1 to 5 or a compound of formula I, II, III, VI or VII as defined in any one of




claims 6 to 9.

11. The use as a fungicide of a composition as defined in any one of claims 1 to 5 or a compound of formula I, II, III, VI or VII as defined in any one of claims 6 to 9.
12. A process for the preparation of a compound of formula I as defined in claim 7 which comprises heating a compound of the general formula



or the general formula




in which n , R , R^1 , R^2 and R^5 are as defined in claim 7 and X and Y are as defined in claim 3, with a compound of the general formula



in which R^5 is as defined in claim 1 and M represents an alkali metal atom, in the presence of a polar solvent.

13. A process for the preparation of a compound of formula II as defined in claim 6 which comprises reacting a compound of the general formula



in which n , R , R^1 and R^2 are as defined in claim 1 and X and Y are as defined in claim 3 with a compound of the general formula




in which R^5 is as defined in claim 1 and M is as defined in claim 11, in the presence of a solvent of the general formula






in which R^5 is as defined in claim 1.

14. A process for the preparation of a compound of formula III as defined in claim 6 which comprises reacting a compound of the general formula




in which n , R , R^1 and R^2 are as defined in claim 1, with a compound XY , in which X and Y are as defined in claim 3.

15. A process for the preparation of a compound of formula VI as defined in claim 8 which comprises reacting a compound of the general formula




in which n , R , R^1 and R^2 are as defined in claim 8 and R^4 is as defined in claim 5 with a reducing agent and subsequently hydrolysing the reaction mixture.

16. A process for the preparation of a compound of formula VII as defined in claim 9 which comprises reacting a compound of the general formula




in which n , R , R^1 and R^2 are as defined in claim 9, with a first compound of the general formula



in which R^4 is as defined in claim 9, in the presence of an acid, and, if desired, converting the compound of formula VII so obtained into another compound of formula VII by reaction with a second different compound of formula IX in the presence of an acid.


Patentansprüche

1. Eine fungizide Zusammensetzung, die ein Trägermaterial und als aktiven Bestandteil eine Verbindung der allgemeinen Formel




umfaßt, in der n eine ganze Zahl von 0 bis 5 darstellt; jeder Rest R ein Halogenatom, eine Nitro-, Cyano-, Hydroxyl-Alkyl-, Halogenalkyl-, Alkoxy-, Halogenalkoxy-, Amino-, Alkylamino-, Dialkylamino-, Alkoxycarbonyl-, Carboxyl-, Alkanoyl-, Alkylthio-, Alkylsuffinyl-, Alkylsulfonyl-, Carbamoyl-, Alkylamido-, Cycloalkyl- oder Phenyl-Gruppe darstellt; R¹ und R² unabhängig voneinander ein Wasserstoffatom oder eine Alkylgruppe darstellen; und R⁵ ein Wasserstoffatom oder eine Alkyl- oder Cycloalkyl-Gruppe darstellt.

2. Eine fungizide Zusammensetzung, die ein Trägermaterial und als aktiven Bestandteil eine Verbindung der allgemeinen Formel




umfaßt, in der n, R, R¹, R² und R⁵ wie in Anspruch 1 definiert sind.

3. Eine fungizide Zusammensetzung, die ein Trägermaterial und als aktiven Bestandteil eine Verbindung der allgemeinen Formel



umfaßt, in der n, R, R¹ und R² wie in Anspruch 1 definiert sind und X und Y unabhängig voneinander ein Halogenatom darstellen.


4. Eine fungizide Zusammensetzung, die ein Trägermaterial und als aktiven Bestandteil eine Verbindung der allgemeinen Formel



umfaßt, in der n, R, R¹ und R² wie in Anspruch 1 definiert sind.




5. Eine fungizide Zusammensetzung, die ein Trägermaterial und als aktiven Bestandteil eine Verbindung der allgemeinen Formel




umfaßt, in der n, R, R¹ und R² wie in Anspruch 1 definiert sind und R⁴ eine Alkylgruppe darstellt.

- 15 6. Eine Verbindung der allgemeinen Formel II oder III wie in Anspruch 2 oder Anspruch 3 definiert.
7. Eine Verbindung der allgemeinen Formel I, in der n, R, R¹, R² und R⁵ wie in Anspruch 1 definiert sind unter der Bedingung, daß wenn n 0 ist und R¹ und R² Wasserstoffatome darstellen, R⁵ kein Wasserstoffatom oder keine Methylgruppe darstellt.
- 20 8. Eine Verbindung der allgemeinen Formel VI, in der n, R, R¹ und R² wie in Anspruch 4 definiert sind unter der Bedingung, daß wenn n 0 ist, R¹ und R² keine Wasserstoffatome darstellen.
- 25 9. Eine Verbindung der allgemeinen Formel VII, in der n, R, R¹, R² und R⁴ wie in Anspruch 5 definiert sind unter der Bedingung, daß wenn n gleich 1 ist, R¹ und R² Wasserstoffatome darstellen und R⁴ eine Methylgruppe darstellt, dann R kein in der 4-Stellung des Phenylrings substituiertes Chlor- oder Bromatom darstellt, und daß, wenn n gleich 0 ist und R¹ und R² Methylgruppen darstellen, R⁴ keine Ethylgruppe darstellt.
- 30 10. Ein Verfahren zum Bekämpfen von Fungi an einem Ort, das umfaßt ein Behandeln des Ortes mit einer Zusammensetzung wie in irgendeinem der Ansprüche 1 - 5 definiert oder mit einer Verbindung der Formel I, II, III, VI oder VII wie in irgendeinem der Ansprüche 6 - 9 definiert.
- 35 11. Die Verwendung als ein Fungizid von einer Zusammensetzung wie in irgendeinem der Ansprüche 1 - 5 definiert oder von einer Verbindung der Formel I, II, III, VI oder VII wie in irgendeinem der Ansprüche 6 - 9 definiert.
- 40 12. Ein Verfahren zur Herstellung einer Verbindung der Formel I wie in Anspruch 7 definiert, das umfaßt ein Erhitzen einer Verbindung der allgemeine Formel



oder der allgemeinen Formel






in der n , R , R^1 , R^2 und R^5 wie in Anspruch 5 definiert sind und X und Y wie in Anspruch 3 definiert sind, mit einer Verbindung der allgemeinen Formel

MOR⁵ (IV)

in der R^5 wie in Anspruch 1 definiert ist und M ein Alkalimetallatom darstellt, in Gegenwart eines polaren Lösungsmittels.

13. Ein Verfahren zur Herstellung einer Verbindung der Formel II wie in Anspruch 6 definiert, das umfaßt ein Umsetzen einer Verbindung der allgemeinen Formel



in der n , R , R^1 und R^2 wie in Anspruch 1 definiert sind und X und Y wie in Anspruch 3 definiert sind, mit einer Verbindung der allgemeinen Formel


MOR⁵ (IV)

in der R^5 wie in Anspruch 1 definiert ist und M wie in Anspruch 13 definiert ist, in Gegenwart eines Lösungsmittels der allgemeinen Formel

R⁵OH (V)


in der R^5 wie in Anspruch 1 definiert ist.

14. Ein Verfahren zur Herstellung einer Verbindung der Formel III wie in Anspruch 6 definiert, das umfaßt eine Umsetzen einer Verbindung der allgemeinen Formel



in der n , R , R^1 und R^2 wie in Anspruch 1 definiert sind, mit einer Verbindung XY , in der X und Y wie in Anspruch 3 definiert sind.


15. Ein Verfahren zur Herstellung einer Verbindung der Formel VI wie in Anspruch 8 definiert, das umfaßt eine Umsetzen einer Verbindung der allgemeinen Formel





in der n, R, R¹ und R² wie in Anspruch 8 definiert sind und R⁴ wie in Anspruch 5 definiert ist, mit einem Reduktionsmittel und anschließend ein Hydrolysieren des Reaktionsgemisches.

16. Ein Verfahren zur Herstellung einer Verbindung der Formel VII wie in Anspruch 9 definiert, das umfaßt eine Umsetzen einer Verbindung der allgemeinen Formel




in der n, R, R¹ und R² wie in Anspruch 9 definiert sind, mit einer ersten Verbindung der allgemeinen Formel



in der R⁴ wie in Anspruch 9 definiert ist, in Gegenwart einer Säure und gegebenenfalls Umwandeln der so erhaltenen Verbindung der Formel VII in eine andere Verbindung der Formel VII durch Umsetzen mit einer zweiten, unterschiedlichen Verbindung der Formel IX in Gegenwart einer Säure.


25 Revendications

1. Une composition fongicide qui comporte un support, et, comme ingrédient actif, un composé de la formule générale



dans laquelle n représente un nombre entier de 0 à 5 ; chaque R représente un atome d'halogène, un groupe nitro, cyano, hydroxyle, alkyle, haloalkyle, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, alkoxy-carbonyl, carboxyle, alkanoyl, alkylthio, alkylsulfinyle, alkylsulfonyl, carbamoyl, alkylamido, cycloalkyle ou phényle ; R¹ et R² représentent indépendamment un atome d'hydrogène ou un groupe alkyle et R⁵ représente un atome d'hydrogène ou un groupe alkyle ou cycloalkyle.


2. Une composition fongicide qui comporte un support, et, comme ingrédient actif, un composé de la formule générale



dans laquelle n, R, R¹, R² et R⁵ sont tels que définis dans la revendication 1.




3. Une composition fongicide qui comporte un support et, comme ingrédient actif, un composé de la formule générale




15 dans laquelle n, R, R¹ et R² sont tels que définis dans la revendication 1 et X et Y représentent indépendamment un atome d'halogène.

4. Une composition fongicide qui comporte un support et, comme ingrédient actif, un composé de la formule générale



dans laquelle n, R, R¹ et R² sont tels que définis dans la revendication 1.

- 30 5. Une composition fongicide qui comporte un support, et, comme ingrédient actif, un composé de la formule générale




dans laquelle n, R, R¹ et R² sont tels que définis dans la revendication 1 et R⁴ représente un groupe alkyle.

- 45 6. Un composé de la formule générale II ou III tel que défini dans la revendication 2 ou la revendication 3.
7. Un composé de la formule générale I dans lequel n, R, R¹, R² et R⁵ sont tels que définis dans la revendication 1 sous la condition que, lorsque n est 0 et R¹ et R² représentent chacun un atome d'hydrogène, R⁵ ne représente pas un atome d'hydrogène ni un groupe méthyle.
- 50 8. Un composé de la formule générale VI, dans lequel n, R, R¹ et R² sont tels que définis dans la revendication 4, sous la condition que, lorsque n est 0, R¹ et R² ne représentent pas chacun un atome d'hydrogène.
- 55 9. Un composé de la formule générale VII dans lequel n, R, R¹, R² et R⁴ sont tels que définis dans la revendication 5 sous les conditions que, lorsque n est 1, R¹ et R² représentent chacun un atome d'hydrogène et R⁴ représente un groupe méthyle tandis que R ne représente pas un atome de chlore ni un atome de brome substitué en position 4 du noyau phénylique, et, lorsque n est 0 et R¹ et R²




représentent chacun un groupe méthyle, R^4 ne représente pas un groupe éthyle.

10. Une méthode pour combattre les mycètes en un endroit, méthode selon laquelle on traite l'endroit avec une composition telle que définie dans l'une quelconque des revendications 1 à 5 ou un composé de formules I, II, III, VI ou VII tel que défini dans l'une quelconque des revendications 6 à 9.
11. L'utilisation comme fongicide d'une composition telle que définie dans l'une quelconque des revendications 1 à 5 ou d'un composé de formules I, II, III, VI ou VII tel que défini dans l'une quelconque des revendications 6 à 9.
12. Un procédé pour la préparation d'un composé de formule I tel que défini dans la revendication 7, procédé selon lequel on chauffe un composé de formule générale



ou de formule générale




dans lesquelles n , R , R^1 , R^2 et R^5 sont tels que définis dans la revendication 7 et X et Y sont tels que définis ci-dessus, avec un composé de formule générale

MOR^5 (IV)

dans laquelle R^5 est tel que défini dans la revendication 1 et M représente un atome de métal alcalin, en présence d'un solvant polaire.

13. Un procédé pour la préparation d'un composé de formule II tel que défini dans la revendication 6, procédé selon lequel on fait réagir un composé de la formule générale



dans laquelle n , R , R^1 et R^2 sont tels que définis dans la revendication 1 et X et Y sont tels que définis dans la revendication 3 avec un composé de la formule générale

MOR^5 (IV)




dans laquelle R^5 est tel que défini dans la revendication 1 et M est tel que défini dans la revendication 11, en présence d'un solvant de la formule générale

R^5OH (V)


dans laquelle R^5 est tel que défini dans la revendication 1.

14. Un procédé pour la préparation d'un composé de formule III tel que défini dans la revendication 6, procédé selon lequel on fait réagir un composé de la formule générale




dans laquelle n, R, R^1 et R^2 sont tels que définis dans la revendication 1, avec un composé XY, dans lequel X et Y sont tels que définis dans la revendication 3.

15. Un procédé pour la préparation d'un composé de la formule VI tel que défini dans la revendication 8, procédé selon lequel on fait réagir un composé de la formule générale



dans laquelle n, R, R^1 et R^2 sont tels que définis dans la revendication 8 et R^4 est tel que défini dans la revendication 5 avec un agent réducteur et, par la suite, on hydrolyse le mélange réactionnel.

16. Un procédé pour la préparation d'un composé de formule VII tel que défini dans la revendication 9, procédé selon lequel on fait réagir un composé de formule générale



dans laquelle n, R, R^1 et R^2 sont tels que définis dans la revendication 9, avec un premier composé de formule générale

R^4OH (IX)

dans laquelle R^4 est tel que défini dans la revendication 9, en présence d'un acide ; et, si on le souhaite, on convertit le composé de formule VII ainsi obtenu en un autre composé de formule VII par réaction avec un second composé différent de formule IX en présence d'un acide.